

Strategies for Regional Modelling of Surface Mass Balance at the Monte Sarmiento Massif, Tierra del Fuego

Franziska Temme¹, David Farías-Barahona^{2,1}, Thorsten Seehaus¹, Ricardo Jaña³, Jorge Arigony-Neto^{4,5},
Inti Gonzalez^{6,7}, Anselm Arndt⁸, Tobias Sauter⁸, Christoph Schneider⁸, Johannes J. Fürst¹

5 ¹Institut für Geographie, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, 91058, Germany

²Departamento de Geografía, Universidad de Concepción, Concepción, 4030000, Chile

³Departamento Científico, Instituto Antártico Chileno, Punta Arenas, 6200000, Chile

⁴Instituto de Oceanografía, Universidade Federal do Rio Grande, Rio Grande, 96203, Brazil

⁵Instituto Nacional de Ciência e Tecnologia da Criosfera, Brazil

10 ⁶Centro de Estudios del Cuaternario de Fuego-Patagonia y Antártica, Punta Arenas, 6200000, Chile

⁷Programa Doctorado Ciencias Antárticas y Subantárticas, Universidad de Magallanes, Punta Arenas, 6200000, Chile

⁸Geography Department, Humboldt-Universität zu Berlin, Berlin, 10099, Germany

Correspondence to: Franziska Temme (franziska.temme@fau.de)

Abstract. This study investigates strategies for ~~melt model~~ calibration of surface mass balance (SMB) models in the Monte
15 Sarmiento Massif (MSM), Tierra del Fuego, with the goal to achieve realistic simulations of the regional ~~surface mass balance~~
(SMB). Applied calibration strategies range from a local single-glacier calibration to a regional calibration with the inclusion
of a snowdrift parametrization. We apply four SMB models of different complexity. This way, we examine the model
transferability in space, the benefit of regional mass change observations and the advantage of increasing the complexity level
20 regarding included processes. Measurements include ablation and ice thickness observations at Schiaparelli Glacier as well as
elevation changes and flow velocity from satellite data for the entire study site. Performance of simulated SMB is validated
against geodetic mass changes and stake observations of surface melting. Results show that transferring SMB models in space
is a challenge, and common practices can produce distinctly biased estimates. Model performance can be significantly
improved by the use of remotely sensed regional observations. Furthermore, we have shown that snowdrift does play an
important role for the SMB in the Cordillera Darwin, where strong and consistent winds prevail. The massif-wide average
25 annual SMB between 2000 and 2022 falls between ~~-0.28 and -0.07~~ ~~-0.25 and -0.07~~ m w.e. yr⁻¹, depending on the applied
model. The SMB is mainly controlled by surface melting and snowfall. The model intercomparison does not indicate one
obviously best-suited model for SMB simulations in the MSM.

1. Introduction

Together with the Northern and the Southern Patagonian Icefield, the Cordillera Darwin Icefield (CDI) in Tierra del Fuego
30 experienced strong losses during the last decades (Rignot et al., 2003; Willis et al., 2012; Melkonian et al., 2013; Braun et al.,
2019; Dussaillant et al., 2019). The glaciers of Tierra del Fuego have contributed to about 5% of the total glacier mass loss in
South America between 2000-2011/14 with a mean annual mass balance (MB) rate of -0.29 ± 0.03 m water equivalent per


year (w.e. yr⁻¹) (Braun et al., 2019). However, the difficult accessibility of Patagonian glaciers and the harsh conditions result in scarce in-situ observations of glacier MB (Lopez et al., 2010). Especially the Cordillera Darwin remains poorly explored
35 (Lopez et al., 2010; Gacitúa et al., 2021).

The CDI is the third largest temperate icefield in the Southern Hemisphere with an area of 2606 km² (state in 2014) including neighboring ice bodies that are not directly connected to the main ice body (Bown et al., 2014). It is located in the southernmost part of the Andes in Tierra del Fuego (Fig. 1) spanning about 200 km in zonal (71.8-68.5°W) and 50 km in meridional (54.9-54.2°S) direction. The two most prominent peaks are Monte Darwin (also known as Monte Shipton) (2568 m above sea level
40 (a.s.l.)) and Monte Sarmiento (2207 m a.s.l.) (Rada and Martinez, 2022). The climate in the Cordillera Darwin is strongly influenced by the year-round prevailing westerlies, which reach a maximum intensity in austral summer. Within the so-called storm track of the westerly belt, frontal systems pass over the region inducing abundant precipitation (Garreaud et al., 2009). The interaction of these moist air masses with the topography causes intense precipitation over the western side and rain-shadow effects and decreasing precipitation amounts towards the east (Porter and Santana, 2003; Strelin et al., 2008). In the
45 westernmost region of the Cordillera Darwin lies Monte Sarmiento. The study site comprises two icefields of around 70 km² and 39 km² (Barcaza et al., 2017), respectively (Fig. 1), which we group together as the Monte Sarmiento Massif (MSM) in this study. Similar to the large icefields in Patagonia, studies show that most of the glaciers in this region have experienced glacier thinning and retreat in the last decades as well (Strelin et al., 2008; Melkonian et al., 2013; Meier et al., 2019).

Many glaciers in southern Patagonia, including the Cordillera Darwin, largely advanced during the Little Ice Age cold interval
50 with maximum advances in the 16th to 19th century (Villalba et al., 2003; Glasser et al., 2004; Strelin et al., 2008; Masiokas et al., 2009; Koch, 2015; Meier et al., 2019). In the last decades, most glaciers in Patagonia and Tierra del Fuego have been strongly losing mass (Rignot et al., 2003; Strelin and Iturraspe, 2007; Strelin et al., 2008; Willis et al., 2012; Melkonian et al., 2013; Braun et al., 2019; Dussaillant et al., 2019). Thinning rates in the Cordillera Darwin have been analyzed the first time by Melkonian et al. (2013) with average annual thinning of -1.6 ± 0.7 m yr⁻¹ (2001-2011). More recent studies focused on the
55 Andes estimate average annual thinning in Tierra del Fuego around -0.32 ± 0.02 m yr⁻¹ (2000-2011/14) (Braun et al. 2019) and -0.56 ± 0.32 m yr⁻¹ (2000-2016) (Dussaillant et al. 2019). Estimates of average annual thinning rates in this area range from -0.32 ± 0.02 m yr⁻¹ (2000-2011/14) by Braun et al. (2019) to -1.6 ± 0.7 m yr⁻¹ (2001-2011) by Melkonian et al. (2013). Average thinning rates are found to be distinctly higher in the northeastern compared to the southwestern part due to the strong precipitation gradient across the mountain range (Melkonian et al., 2013). For individual glaciers in the south (e.g., Garibaldi
60 Glacier), Melkonian et al. (2013) even noticed slight thickening.

Simulating glacier melt ranges from empirical approaches to complex energy-balance models including many physical details. The former ~~assume a linear relationship between temperature and~~ relates melt rates to air temperature, requiring little input. Energy-balance models compute all relevant energy fluxes at the glacier surface, thus rely on numerous meteorological and surface input variables (Gabbi et al., 2014). In between, there is a wide range of different complex implementations. To
65 improve the representation of the spatial and diurnal variability of melt, radiation has been included to temperature-index

models (e.g., Hock, 1999; Pellicciotti et al., 2005). -Previous studies (e.g., Six et al., 2009; Gabbi et al., 2014; Réveillet et al., 2017) have shown that physically-based models can give accurate results when local high-quality meteorological measurements exist, however, when remote meteorological data or reanalysis data are used, the performance decreases rapidly (Gabbi et al., 2014). Thus, more complex models might not be the optimal choice for areas with limited in-situ meteorological measurements, like the Cordillera Darwin. As Patagonian glacier evolution is highly correlated with air temperature (Strelin and Iturraspe, 2007; Weidemann et al., 2020; Mutz and Aschauer, 2022), it is likely that a temperature-based model is able to sufficiently reproduce glacier behavior in the Cordillera Darwin.



75 **Fig. 1: Overview of the study site and a subset of the available in-situ measurements at Schiaparelli Glacier. The inset map displays Patagonia and its icefields: the Northern (NPI) and Southern (SPI) Patagonian Icefield and the Cordillera Darwin Icefield (CDI). The numbers inside the catchment areas refer to the respective glacier ID. Glacier outlines are taken from the Chilean glacier inventory (Bareaza et al., 2017) mark 2004, 2013 and 2019 extents. The satellite image is from Sentinel-2 (2019-02-04) with coordinates in UTM projection, zone 19S.**

80 ~~In the westernmost region of the Cordillera Darwin lies Monte Sarmiento. The main pyramidal summit arises from the correspondent Monte Sarmiento Massif (MSM) and reaches 2207 m a.s.l. Several glaciers descend from all sides of the MSM,~~

together covering $\sim 70 \text{ km}^2$ (Bareaza et al., 2017). To the south of MSM, another glacierized area of 39 km^2 is centered around Pico Marumby (1253 m a.s.l.). Both ice bodies together represent the MSM study area in this study (Fig. 1). Similar to the large icefields in Patagonia, studies show that most of the glaciers in this region have experienced glacier thinning and retreat in the last decades as well (Strelin et al., 2008; Melkonian et al., 2013; Meier et al., 2019).

In order to answer the question, which models are able to reproduce the MB in these unique climatic conditions, we apply four surface mass balance (SMB) models of different complexity at the MSM: a) a positive degree-day model (PDD) (Braithwaite and Olesen, 1989/1995), b) a simplified energy balance (SEB) model (Oerlemans, 2001) using potential insolation, c) a SEB model using the actual insolation (accounting for cloud cover, shading effects and diffuse radiation) and d) the physically-based “Coupled Snowpack and Ice surface energy and mass balance model in PYthon” (COSIPY) (Sauter et al., 2020).

The SMB is given by surface ablation and accumulation. Accumulation is typically considered to equal solid precipitation. Yet, it also depends on deposition, melt-water percolation and subsequent refreezing as well as on avalanching and snow redistribution by wind. The latter can play a decisive role for the spatial heterogeneity in accumulation of mountain glaciers and can reduce or increase the amount by a large factor (Winstral and Marks, 2002; Lehning et al., 2008; Mott et al., 2008; Dadic et al., 2010; Warscher et al., 2013). In southern Patagonia, where strong winds prevail all year round, we hypothesize that snowdrift has a crucial impact on accumulation and with it on the SMB.

Essential for model performance is an appropriate calibration of model parameters, requiring reliable observations. Parameter tuning has been accomplished with different types of data, ranging from in-situ observations of surface ablation and snow properties (e.g., Six et al. 2009; van Pelt et al., 2012; Gabbi et al. 2014; Réveillet et al., 2017; Zolles et al., 2019) to satellite products, e.g., snow line altitudes or mass changes (e.g., Schaefer et al., 2013; Rounce et al, 2020; Barandun et al., 2021). As continuous SMB monitoring is challenging over larger spatial scales covering multiple glaciers, regional modelling attempts often rely on short-term monitoring efforts on single or few glaciers (e.g., Schaefer et al., 2013; Schaefer et al., 2015; Ziemen et al., 2016; Groos et al., 2017; Bown et al., 2019). Though effective, this strategy is in stark contrast with our knowledge that relations between the atmospheric conditions and the surface melt are highly variable in space and time (Pellicciotti et al., 2005, 2008; MacDougall and Flowers, 2011; Gurgiser et al., 2013; Sauter and Galos, 2016; Réveillet et al., 2017; Zolles et al., 2019). Thus, this common approach inherently implies important uncertainties in the SMB estimate and decreases model performance. Discrepancies become evident when modelling results are compared to independent values on specific mass loss from glaciological or geodetic observations. Such comparisons are often inherent in glacier or ice-sheet mass budgeting using various techniques (e.g., Bentley, 2009; Minowa et al., 2021).

The overall goal in this study is therefore to assess and give advice on various strategies for SMB model calibration in the Cordillera Darwin with the aim to achieve reliable simulations of the regional SMB. This objective entails several more specific questions that we want to answer:

Q1. Does a single-glacier calibration guarantee an appropriate regional SMB model transferability ensure transferability of the model producing appropriate regional SMB estimates?

Q2. Is it beneficial to ingest regional geodetic mass change observations into the melt-SMB model calibration?

Q3. Can the performance of the SMB model be improved by increasing the complexity level regarding included processes?

The study is structured as follows: Sect. 2 and 3 describe the study site and data as well as methods and the experimental design. In Sect. 4, we describe the model performance using different calibration strategies and the main characteristics of the SMB in the MSM together with the differences between the employed SMB model types. Sect. 5 provides a discussion of these results and assesses the main limitations and challenges. In Sect. 6, we summarize the main conclusions.

2. Study site and data

2.1 The Monte Sarmiento Massif

~~The MSM lies in the westernmost region of the Cordillera Darwin. The study site comprises two icefields of around 70 km² and 39 km² (Barcaza et al., 2017), respectively (Fig. 1), which we group together as the MSM in this study. The main pyramidal summit, Monte Sarmiento, reaches 2207 m a.s.l. Several glaciers descend from all sides of the MSM, together covering ~70 km² (Barcaza et al., 2017). To the south of MSM, another glacierized area of 39 km² is centered around Pico Marumby (1253 m a.s.l.). Both ice bodies together represent the MSM study area in this study (Fig. 1).~~ The larger icefield includes both land- and lake-terminating glaciers, whereas the smaller one consists entirely of land-terminating glaciers. Schiaparelli Glacier is the largest glacier of the MSM with an area of 24.3 km² in 2016 (Meier et al., 2018). It descends Monte Sarmiento to the north-west almost to sea level and calves into a moraine-dammed proglacial lake, which was formed after strong recession in the 1940s (Meier et al., 2019). Meier et al. (2019) found a continuous average glacier retreat of approximately 5 m yr⁻¹ from 1973 to 2018. Analysis of the surface energy and mass balance of Schiaparelli Glacier with a physically based energy-balance model revealed a glacier-wide mean annual climatic mass balance of -1.8 ± 0.36 m w.e. yr⁻¹ in 2000-2017 (Weidemann et al., 2020). The mass balance is dominated by surface melt and precipitation (Weidemann et al., 2020).

The largest glaciers in the study site after Schiaparelli are Pagels, Lovisato, Conway and Emma Glacier. Emma Glacier was the target for studying Holocene glaciation in the MSM, which indicated that the Holocene glacier behavior in Tierra del Fuego and southern Patagonia responds synchronously to the same regional climate change (Strelin et al., 2008). The other glaciers in the MSM are largely un-surveyed, except by remote sensing (Melkonian et al., 2013; Braun et al., 2019; Dussaillant et al., 2019). From the geodetic MB data from previous studies, different patterns are observed. Despite the rather small study site and proximity of the glaciers, the characteristics of the geodetic MB in 2000-2013 (see Section 2.4) are very heterogenous (Fig. 2). Lovisato Glacier shows by far the highest mass loss. Satellite images (see Fig. 1) reveal large amounts of icebergs in the proglacial lake, indicating significant calving losses for this glacier. A clear contrast between lake- and land-terminating glaciers is not visible. There are several lake-terminating glaciers in the northern part of the study site (Schiaparelli, Conway, Lovisato), however, land-terminating glaciers in this area show similar MBs. ~~Specific~~ Geodetic MBs are also heterogenous in the southern part of the study site, despite all glaciers being land-terminating.

2.2 In-situ observations at Schiaparelli Glacier

We use observational data of two automatic weather stations (AWSs) at Schiaparelli Glacier (Fig. 1). AWS Rock (92 m a.s.l.) is located on rock close to the glacier front. Since the installation in 09/2015, it has been measuring air temperature T , relative humidity RH , global radiation G , wind speed U , wind direction DIR and precipitation RRR in hourly resolution. Bucket-based precipitation measurements often show under-catch due to wind and snow (Rasmussen et al., 2012; Buisán et al., 2017), specifically if the bucket is not heated as in this case. Due to the high wind velocities, precipitation measurements are known to be specifically error-prone in Patagonia ~~due to the high wind velocities~~ (Schneider et al., 2003; Weidemann et al., 2018b; Temme et al., 2020). We therefore assume that the measurement instrument only records a fraction of the total precipitation and ~~increase the recorded values, thus, the annual amount needs to be increased~~ by 20%. AWS Glacier (140 m a.s.l. in 09/2016) is located on ice in the ablation area of Schiaparelli Glacier. It has been measuring T , RH , G , U , DIR and air pressure $PRES$ in hourly resolution since 08/2013 to present with some interruptions. Since this AWS is subject to tilting due to melting of the ice surface, we do not use measurements that require a horizontal sensor orientation from this station. In addition, we identified a step change and a multi-annual drift in the RH measurements. These measurements were therefore discarded. RH values at AWS Glacier were inferred from AWS Rock assuming identical specific humidity. Values of T , corrected RH and $PRES$ from AWS Glacier ~~together with G and RRR from AWS Rock~~ are used to inform the statistical downscaling. G and RRR from AWS Rock serve to ~~and~~ evaluate modelled radiation and precipitation.

Several ablation stakes, ~~spread over~~ concentrated on the lowest part of the ablation area, deliver information about surface melt. Stakes have been installed on varying locations and for irregular time spans ranging from few months to almost one year. The largest number of stakes was installed in the period 11/2018 – 04/2019 with 6 stakes at the same time (see Fig. 1). In the other periods the number of measured stakes ranges from one to four (see Fig. S4). The stake located next to AWS Glacier has been in use for the longest period between 08/2013 and ~~034/2019~~ 03/2019. Additionally, an automatic ablation sensor has been measuring each -150 mm of melt (recording the time point when 150 mm have melted) from 09/2016 to 11/2017, giving a temporally higher resolved information on surface melt.


In April 2016 the ice thickness was measured with a ground-penetrating radar in the ablation area approximately parallel to the glacier front of Schiaparelli Glacier (Fig. 1). Measurements reveal a maximum ice thickness of 324 m with an estimated uncertainty of around 10% (Gacitúa et al., 2021).

2.3 Reanalysis data

The ERA5 reanalysis dataset is the latest global climate reanalysis product of the European Centre for Medium-Range Weather Forecasts (ECMWF). Being a global data set, ERA5 shows high temporal and spatial resolution with an hourly time step and an approximately 31-km horizontal grid over 137 vertical levels (Hersbach et al., 2020). ERA5 and its previous versions have

been successfully applied in modelling studies in Patagonia (e.g., Lenaerts et al., 2014; Bravo et al., 2019; Sauter, 2020; Temme et al., 2020; Weidemann et al., 2020).

180 ERA5 data is required to extend the time period for our modelling framework beyond the AWS records. Therefore, we infer the local surface conditions near AWS Glacier from the spatially coarse ERA5 data, averaging the four closest grid points to the AWSs. U and cloud cover fraction N are directly taken from ERA5. For T , RH and $PRES$, Quantile Mapping (Gudmundsson et al., 2012) was used to relate ERA5 to AWS data (see Sect. 3.1). For downscaling of precipitation, we use a model of orographic precipitation (see Sect. 3.1). It requires the large-scale precipitation and upwind information on geopotential height, 185 air temperature, wind vectors and relative humidity between 850 and 500 hPa which were extracted in a rectangular domain upstream of the study site (54.0-55.0°S, 72.0-71.25°W).



190 **Fig. 2: Specific geodetic mass balance (MB) from elevation change rates for the individual glaciers (> 3 km²) in the MSM study site in 2000-2013. Blue outlines highlight the lake-terminating glaciers. Grey shading indicates glaciers with an area < 3 km². Glacier outlines mark the 2004 extent.**

2.4 Remotely sensed data

Glacier outlines used in this study rely on the Chilean glacier inventory realized by the water directorate of Chile (DGA). It is the first comprehensive glacier inventory of Chile, created from Landsat TM and ETM+ images acquired between 2000–2003 (Barcaza et al., 2017).

195 We use the elevation changes data from Braun et al. (2019), specifically for the period between 2000–2013 (full coverage of the study area). In general, the elevation changes data was derived from two digital elevation models (DEMs). The DEM for the year 2000 corresponds to the Shuttle Radar Topography Mission (SRTM). The 2013 DEM was derived using differential synthetic aperture radar (SAR) interferometry techniques from the TerraSAR-X add-on for the Digital Elevation Measurement mission (TanDEM-X). Postprocessing details can be found in Braun et al. (2019). To estimate the geodetic MB for this study,
200 glacier outlines were taken from the Chilean glacier inventory (Barcaza et al., 2017), and a density factor of 900 kg m^{-3} is used to convert volume to mass changes. Errors and uncertainties from the geodetic MBs were calculated using a standard error propagation equation from Braun et al. (2019), including error sources such as DEM differencing (considering spatial autocorrelation), glacier outlines, and uncertainties from volume to mass conversion ($\pm 60 \text{ kg m}^{-3}$), radar penetration signal, and hypsometric gap filling. The specific MBs calculated from the elevation change rates (2000–2013) are presented in Fig. 2.
205 The annual MB is negative throughout the region, however, with a rather wide range from -1.30 to $-0.09 \text{ m w.e. yr}^{-1}$ (Fig. 2). Glacier outlines from the Monte Sarmiento Massif and their surrounding glaciers are extracted from the two national Chilean inventories generated by the water directorate of Chile (DGA). The first comprehensive glacier inventory of Chile was created from Landsat TM and ETM+ images acquired in 2004 (Barcaza et al., 2017) and later updated using Sentinel-2 images acquired in 2019 (DGA, 2022). The latter inventory presents an improvement in the glacier catchment areas, however, we observed
210 small inconsistencies in both inventories (ice-covered areas not included in the inventory). Hence, we homogenized both inventories and corrected them. To do so, we have used the original and close-date satellite images of both inventories to manually correct these inconsistencies. Moreover, we also generated glacier outlines of 2013 using composite band of Landsat OLI images.

We calculate the geodetic MB using digital elevation models (DEMs) from 2000 and 2013. The DEMs from the 2013
215 TerraSAR-X add-on for the Digital Elevation Measurement mission (TanDEM-X) correspond to a part of the dataset generated in the study of Braun et al. (2019), which presents a complete coverage of the study area. Braun et al. (2019) calculated the elevation changes for the entire Tierra del Fuego region from synthetic aperture radar (SAR) DEMs between 2000 and 2011/2015. For this study, the elevation changes were derived from the Shuttle Radar Topography Mission (SRTM) in 2000 and the 2013 TanDEM-X DEMs. In general, the TanDEM-X DEMs were derived using differential SAR interferometry
220 techniques. Details regarding the SAR approach can be found in Braun et al. (2019).

To obtain precise elevation changes fields, the TanDEM-X DEMs are horizontally and vertically coregistered to the SRTM (reference) DEM using stable areas (Braun et al., 2019; Sommer et al., 2020). Subsequently, the elevation changes differencing is estimated. Data gaps are filled by applying an elevation change versus altitude function by calculating the mean elevation

change within 100 m height bins across the glacier area. Finally, we remove steep slopes ($>50^\circ$) to avoid artificial biases introduced by outliers and filter each elevation band by applying a quantile filter (1%–99%) (Seehaus et al., 2019; Sommer et al., 2020).

To estimate the geodetic MB between 2000 and 2013, we use the two corresponding abovementioned glacier inventories to take into account the glacier area loss (Sommer et al., 2020). To convert volume to mass changes a density factor of 900 kg m^{-3} is applied.

Errors and uncertainties from the geodetic MB ($\frac{M}{\Delta t}$) were calculated using a standard error propagation equation (1) from Braun et al. (2019), which considers the following factors:

- Accuracy of the elevation change rates (considering spatial autocorrelation and hypsometric gap filling) ($\delta_{\Delta h/\Delta t}$)
- Accuracy of the glacier areas (δ_A) (for this study we will include the accuracy of the two glacier inventories)
- Uncertainty from volume to mass conversion using a fixed density (δ_ρ)
- Potential bias due to different SAR signal penetration ($\frac{V_{pen}}{\Delta t}$) (details in Braun et al., 2019)

$$dM = \sqrt{\left(\frac{M}{\Delta t}\right)^2 * \left(\left(\left[\frac{\delta_{\Delta h/\Delta t}}{\Delta t} \right]^2 + \left[\frac{\delta_{A1}}{A1} \right]^2 + \left[\frac{\delta_{A2}}{A2} \right]^2 + \left[\frac{\delta_\rho}{\rho} \right]^2 \right) + \left(\left(\frac{V_{pen}}{\Delta t} \right) * \rho \right)^2} \quad (1)$$

The geodetic MB calculated from the elevation change rates (2000-2013) are presented in Fig. 2. The annual MB is negative throughout the region, however, with a rather wide range from -1.30 to -0.09 m w.e. yr⁻¹ (Fig. 2).

Multi-mission ~~synthetic aperture radar~~SAR remote sensing data was employed to obtain information on glacier speeds. The database covers the period 2001 – 2021 (ERS-1/2 IM SAR 07/2001-08/2001; ENVISAT ASAR, 03/2007-07/2007; ALOS PALSAR 08/2007-09/2010; TerraSAR-X/TanDEM-X 05/2011-02/2021). More detailed information on the sensor specifications can be found in Seehaus et al. (2015).

3. Methods

3.1 Atmospheric forcing

Atmospheric forcing for the SMB models requires T , RH , U , $PRES$, G , RRR and N . T , RH and $PRES$ are statistically downscaled to the AWS location and subsequently extrapolated over the study site, while U and N are directly taken from ERA5. G and RRR are produced using additional models for radiation and orographic precipitation, respectively. The statistical performance of all input variables compared to AWS measurements is summarized in Table S1.

Statistical downscaling of T , RH and $PRES$ is performed via Quantile Mapping. Quantile Mapping is a technique for statistical bias correction of climate-model outputs by transferring the cumulative distribution function of the model to the cumulative distribution function of the observation (Gudmundsson et al., 2012; Cannon et al., 2015). This technique has been successfully applied in Patagonia before (e.g., Weidemann et al., 2018a, 2020). Statistically downscaled T and $PRES$ are spatially

extrapolated from AWS Glacier over the topography using a linear temperature lapse rate (TLR) and the barometric equation, respectively.

255 G over the glacier surface is modelled based on the radiation scheme of Mölg et al. (2009a). It calculates both the direct and diffuse part of the incoming solar radiation from N , T , RH and $PRES$. Radiation is corrected for slope and aspect of the respective grid cell. Furthermore, both self-shading and topographic shading are considered, thus shaded grid cells only receive the diffuse component of the incoming solar radiation (Mölg et al., 2009a, b).

As precipitation events can be short-lived and highly variable in space, it is challenging to infer reliable distributions over
260 complex terrain from coarse global datasets. Furthermore, a direct extrapolation of the sparse AWS measurement network in the CDI using altitudinal lapse rates is critical because measurements in this region are error-prone (Schneider et al., 2003; Schneider et al., 2007; Temme et al., 2020). Therefore, we follow a physically motivated approach using an orographic precipitation model (~~OPM~~), which has been successfully used in glaciological studies before (e.g., Schuler et al., 2008; Weidemann et al., 2018a, 2020). The ~~OPM~~ model is based on the linear steady-state theory of orographic precipitation and
265 includes airflow dynamics, cloud time scales and advection and downslope evaporation (Smith and Barstad, 2004; Barstad and Smith, 2005). This way, the precipitation resulting from forced orographic uplift over a mountain is calculated (Weidemann et al., 2018a). For a more detailed description see Smith and Barstad (2004), Barstad and Smith (2005) and Sauter (2020).

The ~~OPM orographic precipitation model~~ assumes stable and saturated conditions, thus time intervals that do not fulfill these
270 constraints need to be excluded (Smith and Barstad, 2004; Weidemann et al., 2018a). We use relative humidity, Brunt-Väisälä-frequency and Froude number as model constraints in order to ensure saturated, stable airflow without flow blocking. A positive zonal wind component guarantees that airflow crosses the mountains from west to east. ~~The total precipitation is calculated adding the large-scale precipitation (after removing the orographic component from the ERA5 precipitation) to the orographic precipitation calculated in the model. Based on the annual precipitation amounts from AWS Rock, we are able to constrain the relative humidity threshold (90%) above which orographic precipitation can occur and the large-scale precipitation from ERA5. This way, we guarantee that the annual total precipitation at the AWS location agrees with the observed amounts. Sensitivity tests suggest an optimal relative humidity threshold of 90%. eC~~onversion (τ_e) and fallout (τ_f) timescales of hydrometeors (τ) are varied within the range of previous studies (Table 1) in the latter model calibration (Jiang and Smith, 2003; Barstad and Smith, 2005; Smith and Evans, 2007; Weidemann et al., 2013, 2018a, 2020; Sauter, 2020).

280 ~~3.2 Surface mass balance models~~

~~We use four types of SMB models with different complexity. This way, we can understand which type of model is well suited and which processes are essential for SMB simulations in the MSM. Calibration parameters for each model are summarized in Table 1. The calibration approach is described in detail in Sect. 3.5.1. For calibration, simulations were limited to the period in which observations are available (04/1999-03/2019). The final SMB simulations have been extended to the period 04/1999-~~

285 ~~03/2022 at the end to produce the most comprehensive and updated results possible. In the following we explain the different models.~~

Table 1: Overview of the calibration parameters for the atmospheric forcing as well as for the SMB model-variants. Best values are given for calibration Strategy C (see Sect. 3.5.1). Asterisk (*) indicates deviating values for calibration Strategy A. The specific parameter ranges were inferred from the given references.

<u>Model</u>	Parameter	Sampled values (value1, value2, ...) or range (min to max by step)	Optimal S setting	Reference
Atmospheric forcing	TLR ($\text{K } 100 \text{ m}^{-1}$)	-0.60, -0.65, -0.7	-0.7* / -0.6	Buttstädt et al. (2009); Koppes et al. (2009); Schaefer et al. (2015); Bown et al. (2019); Weidemann et al. (2020)
	τ (s)	850, 1000, 1200, 1400	1200 / 850*	Smith (2003); Barstad and Smith (2005); Schuler et al. (2008); Jarosch et al. (2012); Sauter (2020)
PDD	DDF_{ice} ($\text{mm d}^{-1} \text{ } ^\circ\text{C}^{-1}$)	3.0 to 10.0 by 0.5	5.0	Gabbi et al. (2014); Réveillet et al. (2017)
	DDF_{snow} ($\text{mm d}^{-1} \text{ } ^\circ\text{C}^{-1}$)	3.0 to 7.0 by 0.5	3.0	
SEB_Gpot	C_0 (W m^{-2})	-80 to 0 by 5	-2 / 10	Oerlemans (2001); Machguth et al. (2006); Gabbi et al. (2014); Réveillet et al. (2017)
	C_1 ($\text{W m}^{-2} \text{ K}^{-1}$)	2 to 30 by 2	10 / 2	
SEB_G	C_0 (W m^{-2})	-80 to 0 by 5	-20	Oerlemans (2001); Machguth et al. (2006); Gabbi et al. (2014); Réveillet et al. (2017)
	C_1 ($\text{W m}^{-2} \text{ K}^{-1}$)	2 to 30 by 2	10	
COSIPY	α_{ice}	0.300, 0.333, 0.367, 0.400 0.300 to 0.467 by 1/3	0.400	Oerlemans (2001); Schaefer et al. (2015); Weidemann et al. (2020)
	z_{ice} (mm)	0.3 to 2.4 by 0.7	0.3	Brock et al. (2006); Cullen et al. (2007); Mölg et al. (2012)
	$\alpha_{firnsalbedo}$ (days)	0.50 to 0.65 by 0.056, 14, 22	0.5022	Oerlemans (1998); Mölg et al. (2012); Arndt (2021) Oerlemans and Knap (1998); Mölg et al. (2012)
Snowdrift	D_{max} (mm)	4 to 12 by 2	8 / 6	Warscher et al. (2013)
	K	-0.2 to +0.2 by 0.1	0 / 0.1	Warscher et al. (2013)

290 **3.2 Surface mass balance models**

We use four types of SMB models with different complexity. This way, we can understand which type of model is well-suited and which processes are essential for SMB simulations in the MSM. Calibration parameters for each model are summarized

in Table 1. The calibration approach is described in detail in Sect. 3.5.1. For calibration, simulations were limited to the period in which observations are available (04/1999-03/2019). The final SMB simulations have been extended to the period 04/1999-03/2022 at the end to produce the most comprehensive and updated results possible. A complete overview of the model set-up and fixed parameters values is given in the supplement (Table S2). In the following we explain the different models.

3.2.1 PDD

Positive degree day (PDD) models (Braithwaite and Olesen, 1989) assume a linear relationship between air temperature and to surface melt by melt factors that distinguishes between ice and snow surfaces. The melt M is calculated by:

$$M = \begin{cases} \frac{1}{n} * DDF_{ice/snow} * T_a, & T_a > T_T \\ 0, & T_a \leq T_T \end{cases} \quad (1)$$

$DDF_{ice/snow}$ (mm d⁻¹ °C⁻¹) are the degree-day factors for ice and snow, respectively, n is the number of timesteps per day (here $n = 8$), T_a is the air temperature and T_T is the temperature threshold above which melt occurs (here $T_T = 1.0$ °C) (Pellicciotti et al., 2005; Gabbi et al., 2014). The model keeps track of the snow depth in each grid cell to decide whether snow or ice is melted. Accumulation occurs as snowfall at locations where air temperature lies below 1.0 °C.

3.2.2 SEB_Gpot

In the simplified energy-balance (SEB) melt model of Oerlemans (2001), the available melt energy Q_M is calculated by parametrizing the temperature-dependent energy fluxes with the empirical factors C_0 (W m⁻²) and C_1 (W m⁻² K⁻¹):

$$Q_M = (1 - \alpha)I + C_0 + C_1 T_a \quad (2)$$

If snow-free, α is set to $\alpha_{ice} = 0.3$. The snow albedo is not a fixed value but it considers the aging and densification processes using a parameterization via air temperatures since the last snowfall event:

$$\alpha_{snow} = a_1 - a_2 \log(T_a), \quad (3)$$

where a_1 is the albedo of fresh snow (0.9) and $a_2 = 0.155$ (Pellicciotti et al., 2005).

In this first model variant (SEB_Gpot), the incoming solar radiation I is directly related to the potential insolation I_{pot} via the atmospheric transmissivity τ_{atm} ($\tau_{atm} = 0.384$) giving $I = I_{pot} * \tau_{atm}$. Accumulation occurs as snowfall at locations where air temperature is below 1.0 °C.

3.2.3 SEB_G

In order to have a more accurate representation of the incoming solar radiation at each location of the glacier basin, a radiation model is employed. We use the incoming shortwave radiation which we computed with the radiation model based on M \ddot{o} lg et al. (2009a) (see Sect 3.1) as input for a second model variant of the SEB model (SEB_G) ($I = G$). Using the SEB model with

320 two differently complex radiation information, we are able to analyze the importance of accurate radiation input for SMB modelling.

3.2.4 COSIPY

The open-source **e**COupled Snowpack and Ice surface energy and mass balance model in PYthon (COSIPY) (Sauter et al., 2020) is an updated version of the preceding model COSIMA (**e**COupled Snowpack and Ice surface energy and **m**MAss balance model) by Huintjes et al. (2015). COSIPY is based on the concept of energy and mass conservation. It combines a surface energy balance with a multi-layer subsurface snow and ice model, where the computed surface melt water serves as input for the subsurface model (Sauter et al., 2020). In comparison to the previous model types, the primary difference is that the energy fluxes are treated explicitly. Moreover, snow-densification as well as melt-water percolation and refreezing in the snow-cover are possible. Strictly speaking, COSIPY calculates the climatic mass balance following the definition in (Cogley et al., 2011), giving the surface plus the internal mass balance. To maintain readability, we will stick to the term ‘surface mass balance’ also for COSIPY although we underrate the included processes in COSIPY this way.

The energy balance model combines all energy fluxes F at the glacier surface:

$$F = SW_{in}(1 - \alpha) + LW_{in} + LW_{out} + Q_{sen} + Q_{lat} + Q_g + Q_R \quad (4)$$

where SW_{in} is the incoming shortwave radiation taken from the radiation model (G) (see Sect 3.1), α is the surface albedo, LW_{in} and LW_{out} are the incoming and outgoing longwave radiation, Q_{sen} and Q_{lat} are the turbulent sensible and latent heat flux, Q_g is the ground heat flux and Q_R the rain heat flux. Melt only occurs if the surface temperature is at the melting point (0.0 °C) and F is positive. Under this condition, the available energy for surface melt Q_M equals F . Otherwise, this energy is used for changing the near-surface ice or snow temperature. The total ablation not only comprises surface melting but also sublimation and subsurface melting. Mass gain by accumulation is possible via snowfall, deposition and refreezing. A logistic transfer function is applied to derive snowfall from precipitation scaling around a threshold temperature of 1.0 °C.

The albedo is parameterized based on the approach by Oerlemans (1998) where the snow albedo depends on the time since the last snowfall and the snow depth. The turbulent heat fluxes are parameterized using a bulk approach. COSIPY offers two options to correct the flux-profile relationship by adding a stability correction; we confined ourselves to the Monin-Obukhov similarity theory (Sauter et al., 2020). With sensitivity testing, we found the ice albedo (α_{ice}), the roughness length of ice (z_{ice}) and the ~~albedo time constant (t_{albedo})~~ firm albedo (α_{firm}) as the most important tuning parameters in the MSM.

3.3 Snowdrift

Redistribution of snow caused by wind plays a key role for the spatial heterogeneity in accumulation. In Tierra del Fuego and Patagonia, where strong winds prevail throughout the year, we hypothesize that snowdrift has a crucial impact on accumulation and the SMB. Thus, a simple parametrization to capture wind-driven snow redistribution based on Warscher et al. (2013) was

350 slightly modified and added to the SMB model types. The scheme determines locations that are sheltered from or exposed to wind by an analysis of the topography and corrects the solid precipitation accordingly:

$$P_{snow,SD} = P_{snow} * (1 + C_{wind}) \quad (5)$$

The correction factor C_{wind} for each grid cell is calculated by

$$C_{wind} = U * E * (D_{max}(1 - dSVF) - 1) + K \quad (6)$$

355 D_{max} gives the maximum deposition in millimeters, $dSVF$ is the directed sky-view factor and K is a calibration constant, which was set to 0.1 by Warscher et al. (2013). We vary it in a range from -0.2 to +0.2 in the snowdrift calibration together with the D_{max} (Table 1). E is a factor for weighting with elevation (linearly) ranging from 0 to 1, assuming that lower wind speeds prevail at lower elevations which reduces the snow redistribution. In this study, we additionally include a weighting with prevailing wind speed U to further improve the performance because we observe different wind directions with different
360 velocities and we suppose that more (less) snow is redistributed also during periods of higher (lower) wind velocities. For a more detailed description of the snowdrift scheme refer to Warscher et al. (2013).

3.4 Ice flux and mass budgeting

Ice surface velocity fields are derived from the SAR imagery database by applying intensity offset tracking on co-registered image pairs (Strozzi et al., 2002). Tracking parameters were adjusted depending on sensor specification and acquisition
365 intervals. The tracking is done using multiple tracking patch sizes in order to account for different glacier flow speeds and a subsequent stacking of the results was applied. More details on the processing, including filtering and error estimation, can be found in Seehaus et al. (2018).

In order to estimate the ice flux, a flux gate was defined along a cross-profile following the thickness surveys in 2016 (Gacitúa et al., 2021). By combining the obtained surface velocity information with the ice thickness measurements along this flux gate,
370 the ice flux was computed following the approach of Seehaus et al. (2015) and Rott et al. (2011). In order to account for ice thickness changes at the flux gate throughout the observation period, the measured ice thickness values were corrected by a surface lowering rate of -2.8 m yr^{-1} derived from the annual average elevation change rate between 2000 and 2019 (see Sect. 2.4) near the flux gate. The resulting ice flux through the flux gate is summarized in Table S23.

The combination of the SMB integrated over the glacier area above the flux gate and the mass lost through the flux gate allows
375 us to determine the total mass budget of Schiaparelli Glacier. This value is comparable to the specific-geodetic MB from elevation changes in the area above the flux gate and will be used as one calibration constraint in this study.

3.5 Experimental design

3.5.1 Calibration strategies

We use three different strategies for model calibration that are summarized in Fig. 3. The calibration strategies are based on calculations of model skill. The choice for which parameters enter the calibration was preceded by sensitivity studies on an exhaustive set of parameters with the aim to cover all relevant contributions to SMB. The TLR and τ give control on the amount of snowfall and on temperature-dependent melting. For the PDD and the two SEB variants, we calibrate the model-specific parameters (DDF_{ice} and DDF_{snow} , and C_0 and C_1 , respectively). For COSIPY we must constrain the number of calibrated parameters to limit the modelling effort. Therefore, we decided for the ice albedo (α_{ice}) and the roughness length of ice (z_{ice}), which constrain ice melting addressing both, the radiative and turbulent energy fluxes. To have a control also on the higher elevated, firn-covered parts of the glaciers, we include the firn albedo (α_{firn}). Sensitivity testing revealed that those parameters impact the SMB results the strongest. Other parameters we had tested are the temperature of snow/rain transfer, the albedo of snow, the albedo time constant which gives the effect of ageing on the snow albedo, and the method of stability correction. Those parameters were fixed at the end because they either were interdependent with other parameters, had minor impact on the overall results or showed clear advantage of the one method.


In Strategy A, calibration is focused only on Schiaparelli Glacier where we have in-situ observations. These include ablation stake measurements (see Section 2.2) and estimation of total glacier mass budget using a combination of elevation changes and mass flux through a flux gate parallel to the glacier front (see Sect. 3.4). Measured melt at each ablation stake is compared to modeled average melt at all grid cells of same altitude (± 5 m) at Schiaparelli Glacier for the respective same period. Ablation measurements give control on the processes of melting in the ablation area whereas the mass budget gives an additional control on the basin-wide mass overturning and with it on the amount of accumulation. After this glacier-specific calibration, the model is transferred to regional scales, i.e., the surrounding glaciers in the study site, with the parameter setting we found in the calibration.

In Strategy B, we use regional specific geodetic MB observations from MSM elevation changes (2000-2013). This way we calibrate the SMB model towards the massif-wide average in order to guarantee that the total net amount of accumulation and ablation on a regional scale is close to observations. Since dynamical losses at calving fronts are not considered in the SMB but are included in the geodetic MB, we exclude glaciers that have significant calving losses (Lovisato Glacier). However, we include glaciers in the average value that are lake-terminating but known to have only minor calving losses. Furthermore, only glaciers larger than 3 km² are considered because small glaciers involve larger uncertainties. The average annual MB of this subset of glaciers is referred to as B_{MSMnc} in the following and comprises around 71% of the total glaciated area. The annual mean-average SMB of the according glaciers is then calibrated towards this observed value.

In Strategy C, we follow Strategy B, but additionally activate a snowdrift module that needs to be calibrated in this step. After defining the regional massif-wide amount of accumulation in Strategy B, we now optimize the distribution of snowfall on the local scale with the inclusion of the snowdrift. As calibration constraints we again rely on the B_{MSMnc} , but additionally consider

410 the total mass budget of Schiaparelli Glacier to incorporate information about local distribution of snow. We ensure mass conservation by keeping the total amount of snowfall nearly ($\pm 10\%$) constant.

We use the PDD for calibration of the climate- and snowdrift-related parameters. These include the TLR and τ as well as D_{max} and K . Additional model-specific parameters are DDF_{ice} and DDF_{snow} . The number of varied parameters and their ranges are based on what has been used in previous studies (see Table 1). For the other three models, we fix the temperature and precipitation field and the snowdrift parameters based on the results of the PDD calibration. This way, we guarantee consistency in the atmospheric forcing and save computational cost. Thus, only model-specific parameters are calibrated for those following Strategy C (see Fig. 3).



420 **Fig. 3: Overview of the three calibration strategies used for the calibration of the PDD model, the snowdrift module and the final calibration of the SEB_Gpot, SEB_G and COSIPY.**

The model skill is calculated using different combinations of observations, depending on the respective calibration strategy. These include the ablation stake measurements at Schiaparelli Glacier, the total mass budget of Schiaparelli Glacier and the specific-geodetic MB derived from elevation changes on a massif-wide average (glaciers $> 3 \text{ km}^2$) excluding significantly calving glaciers (B_{MSMnc}). To calculate the model skill for each run, the simple averaging method of Pollard et al. (2016) is used applying full-factorial sampling. Taking the misfit between model and observation, an objective aggregate score is determined (Pollard et al., 2016; Albrecht et al., 2020). The misfits are calculated by mean squared errors between observation and model. Thereby, the individual, normalized score $S_{i,j}$ is obtained for each considered measurement type i and each parameter sample j (see Table 1):

$$S_{i,j} = e^{\frac{-M_{i,j}}{\overline{M}_i}} \quad (7)$$

430 Here, \overline{M}_i is the median of all misfits of one measurement type (for all parameter combinations). The unweighted, aggregated score for each run is the product of the individual scores

$$S_j = \prod_{i=1}^3 S_{i,j}. \quad (8)$$

The run with the highest aggregated score S_j implies the optimal parameter combination.

3.5.2 Model evaluation and intercomparison

435 To investigate the model performance, we compare modelled surface and observed ~~specific-geodetic~~ MB of the individual land-terminating glaciers basins (> 3 km²) in the study site (2000-2013). To determine the agreement, we compute the area-weighted root mean square error (RMSE). ~~Since we are able to estimate the ice flux along the measured thickness profile at Schiaparelli Glacier, we directly compare the integrated surface and specific MB above the flux gate. In this way, Schiaparelli Glacier is included in the RMSE estimation.~~ Furthermore, we assess the agreement between modelled and observed ablation
 440 at the ablation stakes, ~~where we have measurements between 2013 and 2019 in the observation period between 2013 and 2019.~~ In order to investigate the performance of SMB models with a different degree of complexity, we compare the results of four model types. After calibrating the model-specific parameters of each model individually, the best guess SMB characteristics and uncertainties of each model can be compared with each other in respect to the observed geodetic MBs. Uncertainties and sensitivity to the calibration parameters are discussed in the supplementary material.

445 ~~3.5.3 Uncertainty assessment~~

~~Uncertainties in the SMB estimation come from three different sources in this study: Climatic forcing related uncertainties, model inherent uncertainties and model type related uncertainties. Uncertainties in the climatic forcing mainly stem from snowfall and air temperatures. Model inherent uncertainties relate to process parameterizations and limitations in the model calibration. Uncertainties due to model type are linked to model capability to realistically represent the controlling factors on~~
 450 ~~SMB and are outlined by the use of different SMB models. With our comprehensive model calibration, we are able to assess all three types of uncertainties. Climatic uncertainties are given by the range of TLRs and τ used for calibration of the PDD. To determine these uncertainty ranges, we use all runs with different combinations of TLR and τ out of the 10 best ranked runs (see scores in Sect. 3.5.1). Model inherent uncertainties are assessed by considering the 10 best ranked parameter combinations for each model (5 best ranked for the PDD due to the significantly smaller sample size) after setting the climatic~~
 455 ~~forcing. Model type related uncertainties are quantified by comparing the results of the four different models.~~

4. Results


4.1 Strategies for model calibration

4.1.1 Strategy A: Single-glacier calibration

460 Results of model calibration show that ablation stake measurements give a control on melting only since almost no snowfall
occurs at the stake locations. Thus, the ability to reproduce ablation at the stakes depends principally on the DDF_{ice} (see
Fig. S1a,b). The total mass budget, additionally, depends strongly on the TLR and τ , thus both the ablation and the distribution
and amount of snowfall over elevation (see Fig. S1c). Based on this information, we are able to narrow down the amount of
solid precipitation. The combination of both datasets allows an accurate calibration of ablation at Schiaparelli Glacier as well
as a profound-well-informed estimate of precipitation amounts over its catchment area.

465 An overview of the calibration scores for Strategy A is presented in Fig. S1d. This strategy suggests a TLR of
-0.70 °C 100 m⁻¹ at Schiaparelli Glacier. The requirement for a higher TLR tells us that a steeper SMB gradient with respect
to elevation is needed a large amount of precipitation is needed to be in solid form in order to meet the observations, resulting
in reduced ablation with altitude and increased snowfall. A similar signal comes from the τ , where a value of 850+000 s is
most suitable producing a precipitation field with rather a high amounts of orographic precipitation. The degree-day-factors
470 DDF_{ice} and DDF_{snow} are set to 56.0 and 3.0 mm d⁻¹ °C⁻¹, respectively.

After the local calibration at Schiaparelli Glacier, the model is transferred to the surrounding glaciers. The results are given in
Table 2. Comparing the surface (-0.51-62 m w.e. yr⁻¹) and the specific-geodetic MB (-0.79 m w.e. yr⁻¹) delivers an estimated
calving flux of 0.28-17 m w.e. yr⁻¹ (6.944.26 Mt yr⁻¹) at Schiaparelli Glacier. However, the application on the regional scale
shows that the SMB is consistently overestimated compared to the geodetic observations (Fig. 4). The observed value for the
475 B_{MSMc} of
-0.51 m w.e. yr⁻¹ is even positive with +0.16-09 m w.e. yr⁻¹ in the model (Table 2). Furthermore, several land-terminating
glaciers, where no dynamical-calving losses are involved, have a clearly positive annual SMB which differs distinctly from
the observations. The poor agreement is reflected in a high RMSE of 0.65-56 m w.e. yr⁻¹ (0.52 m w.e. yr⁻¹ including
Schiaparelli Glacier cut along the flux gate) (Table 2).



480

Fig. 4: Difference of modelled surface to observed specific-geodetic MB (2000-2013) for the three calibration strategies. Dotted areas indicate lake termination precluding a direct comparison of the two datasets. Grey shading indicates glaciers with an area < 3 km². Glacier outlines mark the 2004 extent.

4.1.2 Strategy B: Regional calibration

485

In a second stepstrategy, we use the regional specific-geodetic MB as the sole calibration target. Therefore, we rely on the massif-wide average annual specific-geodetic MB obtained from satellite observations, excluding glaciers with major calving losses (B_{MSMnc}) (see Fig. S1e). Following the approach in Strategy A, the model calibration is performed with the PDD calibrating the same parameters again. All-Whereas the DDF_{ice} updates to 5.0 mm d⁻¹ °C⁻¹, the DDF_{snow} parameters remains unchanged. _____ apart _____ from _____ thThe TLR, _____ which _____ becomes _____ changes _____ to -0.60 °C 100 m⁻¹ and the τ to 1200 s. Accordingly, the precipitation field itself stays the same, but the amount of precipitation and the ratio of solid and liquid precipitation are shifted towards less snowfall.

490

Using regional observations of specific-geodetic MB from satellite data, the calibration for a regional application can be improved. With this strategy the observedAs it is the sole calibration target, the value of the B_{MSMnc} is reproduced perfectly with a modelled value of -0.51 m w.e. yr⁻¹ (Table 2). Individual glaciers show a loss of performance, e.g., at Schiaparelli
 495 Glacier the simulated SMB gets more negative. However, looking at several land-terminating glaciers of the MSM (glaciers 149, 152 and 159), the agreement has considerably increased (Fig. 4). This is also reflected in a strong decrease of the RMSE to 0.28-30 m w.e. yr⁻¹ (0.45 m w.e. yr⁻¹ including Schiaparelli Glacier cut along the flux gate) (Table 2). The negative-positive SMB bias from calibration Strategy A is no longer discernible.

Table 2: Comparison of modelled surface to observed specific-geodetic MB (m w.e. yr⁻¹) (2000-2013) from the PDD using three different calibration strategies as well as from the SEB_Gpot, SEB_G and COSIPY for the glaciers in the study site (> 3km²). The results of Strategy C equal the final results of the PDD model. B_{MSMnc} gives the massif-wide annual average MB excluding glaciers with major calving losses. The root mean square error (RMSE) is weighted by area and calculated from the land-terminating glaciers only as well as including Schiaparelli cut along the flux gate. Asterisk marks lake termination.

Name/ID	Area (km ²)	geodetic MB (m w.e. yr ⁻¹)	SMB (m w.e. yr ⁻¹)					
			PDD			SEB_Gpot	SEB_G	COSIPY
			Strategy A	Strategy B	Strategy C			
<u>133 - Conway*</u>	<u>8.45</u>	<u>-0.18</u>	<u>1.59</u>	<u>0.77</u>	<u>0.29</u>	<u>0.28</u>	<u>0.29</u>	<u>0.87</u>
<u>136 - Schiaparelli*</u>	<u>25.03</u>	<u>-0.79</u>	<u>-0.62</u>	<u>-1.11</u>	<u>-1.02</u>	<u>-0.79</u>	<u>-0.79</u>	<u>-1.20</u>
<u>136 - Schiaparelli_FG</u>	<u>23.15</u>	<u>-0.59</u>	<u>-0.67</u>	<u>-1.32</u>	<u>-1.22</u>	<u>-1.11</u>	<u>-1.10</u>	<u>-1.27</u>
<u>138*</u>	<u>3.89</u>	<u>-0.50</u>	<u>1.68</u>	<u>0.92</u>	<u>2.03</u>	<u>2.39</u>	<u>2.30</u>	<u>2.45</u>
<u>139 - Lovisato*</u>	<u>12.57</u>	<u>-1.30</u>	<u>0.62</u>	<u>-0.15</u>	<u>0.61</u>	<u>0.92</u>	<u>0.95</u>	<u>0.80</u>
<u>142 - Emma</u>	<u>7.28</u>	<u>-0.21</u>	<u>0.04</u>	<u>-0.68</u>	<u>-0.40</u>	<u>-0.43</u>	<u>-0.30</u>	<u>-0.45</u>
<u>144</u>	<u>3.83</u>	<u>-0.74</u>	<u>-0.24</u>	<u>-0.78</u>	<u>-0.88</u>	<u>-0.95</u>	<u>-0.73</u>	<u>-0.83</u>
<u>149</u>	<u>3.91</u>	<u>-0.65</u>	<u>0.04</u>	<u>-0.52</u>	<u>-0.64</u>	<u>-0.55</u>	<u>-0.64</u>	<u>-0.78</u>
<u>152</u>	<u>3.60</u>	<u>-0.33</u>	<u>0.96</u>	<u>0.22</u>	<u>0.05</u>	<u>-0.14</u>	<u>-0.04</u>	<u>0.26</u>
<u>157</u>	<u>3.55</u>	<u>-0.09</u>	<u>0.31</u>	<u>-0.22</u>	<u>0.13</u>	<u>0.36</u>	<u>0.33</u>	<u>0.32</u>
<u>159 - Pagels</u>	<u>18.69</u>	<u>-0.45</u>	<u>-0.06</u>	<u>-0.65</u>	<u>-0.54</u>	<u>-0.47</u>	<u>-0.52</u>	<u>-0.74</u>
<u>B_{MSMnc}</u>	<u>78.23</u>	<u>-0.51</u>	<u>0.09</u>	<u>-0.51</u>	<u>-0.43</u>	<u>-0.33</u>	<u>-0.32</u>	<u>-0.45</u>
<u>RMSE</u>			<u>0.56</u>	<u>0.30</u>	<u>0.17</u>	<u>0.19</u>	<u>0.16</u>	<u>0.31</u>

4.1.3 Strategy C: Regional calibration including snowdrift


Adding snowdrift delivers additional parameters that need to be calibrated with the PDD. Therefore, we fix the model parameters as determined in Strategy B. Afterwards, the snowdrift parameters are calibrated, suggesting a D_{max} of 68.0 mm and K of +0.40. The snowdrift scheme redistributes snow on average from the northwest to the southeast of the massif due to prevailing northwesterly flow. Subsequently, the southeastern glaciers obtain higher snowfall amounts, whereas from the northwestern glaciers snow is on average removed. With this procedure the agreement between modelled and observed MBs is further improved (Fig. 4), although the resulting simulated B_{MSMnc} (-0.43 m w.e. yr⁻¹) is slightly overestimated (Table 2). At Schiaparelli Glacier, where a large part of the accumulation area is located east of the prominent Monte Sarmiento, snow is on average deposited producing a slightly less negative SMB, which is closer to observations. For the land-terminating glaciers,

the difference between model and observations now lies within close to the uncertainty of the observation (Table S5), with a total RMSE of ~~0.18-17~~ m w.e. yr⁻¹. Therefore, further tuning is neither required nor justifiable. ~~The RMSE including Schiaparelli Glacier cut along the flux gate can also be reduced further with Strategy C (0.39 m w.e. yr⁻¹).~~

The other three SMB models are limited to calibration Strategy C for sake of computational cost. We use the TLR, τ and snowdrift parameters as found by the PDD, and calibrate the model-specific parameters only (see Table 1, Fig. S2). For the SEB_Gpot/SEB_G we get a C_0 and C_1 of ~~-10/-20~~ W m⁻² and 12/10 W m⁻² °C⁻¹, respectively. COSIPY calibration reveals an α_{ice} of 0.4, an α_{firm} of 0.5 and a z_{ice} of 0.3 mm ~~and a τ_{albedo} of 22 days.~~

520 4.2 Surface mass balance of the Monte Sarmiento Massif

Generally, all four models give very similar results of SMB (Fig. 5). The spatial distribution and seasonal/interannual patterns are captured by all models in a similar way. We will summarize the main characteristics of SMB in the MSM in the following, and highlight differences between the models. For this analysis, we include all glaciers in the study site (no area limit) to produce the most comprehensive results possible.




525 **Fig. 5: Mean annual surface MB (SMB) for the four SMB models (2000–2022). Dotted lines mark altitude in 300 m intervals with intensity decreasing with height. Glacier outlines represent 2004 (black), 2013 (darkgrey) and 2019 (lightgrey) extents.**

The massif-wide average annual SMB lies just below equilibrium, with the PDD and COSIPY producing a more negative value (~~-0.285~~ and ~~-0.2049~~ m w.e. yr⁻¹, respectively) than the SEB_Gpot and SEB_G (~~-0.07~~ and ~~-0.1309~~ m w.e. yr⁻¹, respectively). For all models, the SMB is mainly influenced by snowfall (average of +1.6674 to +1.7987 m w.e. yr⁻¹) and melt (average of ~~-1.9387~~ to ~~-2.6455~~ m w.e. yr⁻¹). Snowfall is almost zero at the lowest parts of the glaciers indicating melt all year round (Fig. S3a). The distribution of snow reflects the topography, increasing strongly towards the summits and showing largest snow deposition south-east of the mountain peaks and ridges. The highest amounts are found in the wind-sheltered

slopes of the Monte Sarmiento summit. For all four SMB models, we see high mass gain due to snowfall on the elevated areas of the massif (up to around 10 m w.e. yr⁻¹) and extreme mass loss at the glacier tongues (up to around -10 m w.e. yr⁻¹) (Fig. 5). Several glaciers have large ablation areas (Schiaparelli, Lovisato, Pagels). However, Schiaparelli Glacier stands out due to its large size, the range of altitude and its ~~very~~ huge glacier tongue causing a much larger area of intense ablation compared to the other glaciers in the region. Depending on the model type, the massif-wide equilibrium line altitude is on average between 770 and ~~794~~83 m a.s.l. during the study period. Equilibrium line altitudes tend to be lower in the east of the massif compared to the west, which can be confirmed by snow line altitudes from satellite observations ~~as well in the region~~ (Table S4). Due to the location in the higher mid-latitudes, the seasonal variations are huge. In summer, the average ~~mass balance~~SMB is negative up to around 900-1000 m a.s.l., which leaves (almost) no area of mass gain for several glaciers in the region (see Fig. S3~~e,d~~). This applies in particular to the southern, lower elevated massif. In winter, the ~~largest part~~majority of the MSM area is characterized by a positive SMB. The cooler temperatures cause higher snowfall amounts, and we observe snowfall also over lower altitudes (see Fig. S3~~e~~). More than ~~6533~~% of the total snow accumulates in winter (June to August) and spring (September to November), only 13% in summer (December to February).

Over the course of the 22-year~~s~~ study period, we see a phase of more negative and more positive annual SMB that all four models agree on (Fig. 6). Massif-wide more positive MB values are prevailing between 2009 and 2015/16, more negative before and after this phase. More negative MBs coincide with over-average temperatures and decreased snowfall and vice versa. All models agree that the most negative MBs likely occurred in 2003/04, 2005/06, 2016/17, 2019/20 and 2020/21. However, the amplitude of annual mass balances differs significantly between the models. Overall, the PDD and COSIPY tend to simulate more negative MBs, however, COSIPY also simulates more positive MB in several positive years and the SEB_Gpot and SEB more positive ones (Fig. 6).



555 Fig. 6: Annual massif-wide average SMB for the four SMB models (left axis) together with the anomaly of temperature (black line) and snowfall (black star) from the 2000-2022 average (right axis).

4.3 Uncertainty quantification

Uncertainties in the SMB results stem from the climatic forcing, the model-specific parameters and the type of model. We want to quantify those in the following. To assess the uncertainty related to the climatic forcing used, we compare the results of the 10 best ranked PDD runs with different combination of TLR and τ . This ranking differs depending on the calibration strategy applied. Using Strategy A, the maximum range around the best guess in the SMB is -0.38 to $+0.17$ m w.e. yr $^{-1}$ for the individual glaciers and -0.32 to $+0.04$ m w.e. yr $^{-1}$ for the B_{MSMne} (see Table S3). With Strategy B, the uncertainties are overall smaller with ± 0.29 m w.e. yr $^{-1}$ for the individual glacier MBs and below ± 0.10 m w.e. yr $^{-1}$ for the B_{MSMne} .

The model-specific uncertainty stems from the calibration of the model-specific parameters (see Table S4). For the PDD, the SMB of individual glaciers differs up to $-0.45/+0.40$ m w.e. yr $^{-1}$ from the best estimate. Also, the range around the B_{MSMne} is large with -0.34 to $+0.25$ m w.e. yr $^{-1}$. For the SEB_Gpot and the SEB_G, the ranges in SMB of the individual glaciers are smaller with $-0.14/+0.28$ m w.e. yr $^{-1}$ and $-0.13/+0.23$ m w.e. yr $^{-1}$, respectively. The B_{MSMne} can be determined accurately for both models in a range of roughly ± 0.10 m w.e. yr $^{-1}$, respectively. For COSIPY, the best ranked run equals the maximum estimated SMB for each individual glacier. The range of uncertainty is up to -0.26 m w.e. yr $^{-1}$ for the individual glaciers and -0.20 m w.e. yr $^{-1}$ for the B_{MSMne} . The RMSEs can be determined rather accurately within a range of ± 0.15 m w.e. yr $^{-1}$ for all models, apart from the PDD where the range is up to $+0.25$ m w.e. yr $^{-1}$.

Comparing the results of the best ranked run of each model (Table S4), we observe that the minimum estimate of each glacier comes either from the PDD or COSIPY, whereas the maximum estimate comes often from the SEB_Gpot. The estimates for the individual glaciers differ between 0.16 and 0.63 m w.e. yr⁻¹. The B_{MSMnc} is simulated between 0.46 and -0.29 m w.e. yr⁻¹.

4.34 Model intercomparison

We can compare modeled and observed MB for the individual glacier catchments to assess the performance of the individual models (Fig. 7). The area-weighted RMSE (Table 2) is similar for the PDD, SEB_Gpot and SEB_G (0.178, 0.19 and 0.16 m w.e. yr⁻¹) and largest for COSIPY (0.310 m w.e. yr⁻¹) comparing land-terminating glaciers only. However, if we include Schiaparelli Glacier (cut along the flux gate), the ranking changes distinctly. Since with Strategy B and C, the performance at Schiaparelli Glacier is known to decrease slightly compared to Strategy A, also the RMSEs are getting larger. The clearly best performance is now achieved with COSIPY (0.25 m w.e. yr⁻¹) followed by both SEB variants, and the least with the PDD (0.39 m w.e. yr⁻¹). The range of uncertainty of the RMSEs is very similar for all four models (see Table S46), with RMSEs lying between 0.167 and 0.43-34 m w.e. yr⁻¹. Only the PDD stands out with a maximum RMSE of 0.64-75 m w.e. yr⁻¹ if we include Schiaparelli Glacier, again. The B_{MSMnc} range of the 10 best ranked runs are very similar (range below 0.20-24 m w.e. yr⁻¹) for the SEB_Gpot, SEB_G and COSIPY. For the PDD, this range is distinctly larger with 0.59-51 m w.e. yr⁻¹ taking the 5 (due to the smaller sample size) best ranked runs.

In order to answer the question about investigate the importance of including accurate information about incoming radiation, we can directly compare the performance of the SEB_Gpot with the SEB_G. The former relies on the potential radiation whereas the latter accurately calculates the direct and diffuse part of incoming shortwave radiation taking into account cloud cover and shading. Generally, both models tend to overestimate the SMB in the MSM (Fig. 7), which is reflected also in the overestimation of the B_{MSMnc} (-0.33 m w.e. yr⁻¹ and -0.32 m w.e. yr⁻¹ for the SEB_Gpot and SEB_G, respectively) (Table 2). The differences between both models are overall minor, although the RMSE for the SEB_G is smaller (Table 2). Thus, for this study site, the improvement by using more accurate instead of potential radiation appears insignificant. This finding further agrees with the fact that the PDD also produces satisfying results.




Fig. 7: Difference of modelled surface to observed specific-geodetic MB (2000-2013) for the four SMB models. Dotted areas indicate lake termination precluding a direct comparison of the two datasets. The displayed results for the PDD are those from Strategy C in Fig. 3. Grey shading indicates glaciers with an area < 3 km². Glacier outlines mark the 2004 extent.

The second observation available for model evaluation are the stake measurements. However, the agreement between measured and modelled ablation at the stakes is poor for all considered SMB models (see Fig. S4 and S5). Mean RMSEs are in the range between 4.023.92 and 4.90.78 m w.e. yr⁻¹ (Table 3), which is about 33% of the observed melt. The model bias ranges between -0.77 and 3.51 m w.e. yr⁻¹. The best results are achieved with COSIPY followed by the PDD regarding both, RMSE and bias followed by COSIPY. At the individual ablation stakes (Fig. S4), COSIPY behaves distinctly different to the other models, for which melt rates are more similar for most of the time. Subsequently, COSIPY meets the observations better for the first half of the time where the other models often underestimate the melting. However, after 2018 COSIPY clearly overestimates the melt rates which causes the overall poor RMSE value degrading the overall statistics. In general, we consider the ablation measurements as error-prone looking at individual observations. Thus, we consider the large RMSE and bias values as caused only partly by poor model performance.

Table 3: Comparison (RMSE and mean bias in m w.e. yr⁻¹) between observed and modeled melt at the stakes. S1to5 includes all individual stake observations in 2013-2019, Sauto comprises the measurements by the automatic ablation sensor. and Smean gives the average RMSE of both variables.

Stake	PDD		SEB_Gpot		SEB_G		COSIPY	
	RMSE	bias	RMSE	bias	RMSE	bias	RMSE	bias
S1to5	<u>3.92</u>	<u>2.07</u>	<u>4.78</u>	<u>3.51</u>	<u>4.56</u>	<u>3.23</u>	<u>3.99</u>	<u>-1.00</u>
Sauto	<u>4.12</u>	<u>-1.02</u>	<u>3.94</u>	<u>1.90</u>	<u>3.90</u>	<u>2.04</u>	<u>3.76</u>	<u>-0.77</u>

5. Discussion

615 5.1 Strategies for model calibration

The single-glacier model calibration at Schiaparelli Glacier (Strategy A) results in a TLR of $-0.70\text{ }^{\circ}\text{C } 100\text{ m}^{-1}$ which is slightly stronger compared to previously reported annual values that vary from -0.60 to $-0.67\text{ }^{\circ}\text{C } 100\text{ m}^{-1}$ in the southern Patagonia region (Strelin and Iturraspe, 2007; Buttstädt et al., 2009; Koppes et al., 2009; Schaefer et al., 2015; Weidemann et al., 2018a, 2020). Furthermore, calibration suggests a τ of ~~1000~~850 s, which differs significantly from the value used at Schiaparelli
620 Glacier in a recent SMB study (Weidemann et al., 2020). However, it agrees well with values reported in various other applications, including southern Patagonia (Smith and Barstad, 2004; Barstad and Smith, 2005; Smith and Evans, 2007; Schuler et al., 2008; Jarosch et al., 2012; Sauter, 2020). ~~Furthermore, we calculated a rough estimate of τ from ERA5 data, which gave us a similar value of 1050 ± 350 s.~~ The degree-day-factors DDF_{ice} and DDF_{snow} of 5.6.0 and $3.0\text{ mm d}^{-1}\text{ }^{\circ}\text{C}^{-1}$, respectively, lie within the range of previously reported values (Stuefer et al., 2007; Gabbi et al., 2014; Réveillet et al., 2017).
625 At Gran Campo Nevado, Schneider et al. (2007) found a value of $7.6\text{ mm d}^{-1}\text{ }^{\circ}\text{C}^{-1}$ for ice in summertime. Calculating the average DDF_{ice} directly from measured ablation and positive degree-day sum at the stake locations (Groos et al., 2017) delivers a value very close to the calibrated one with 5.50 $\text{mm d}^{-1}\text{ }^{\circ}\text{C}^{-1}$ at the automatic ablation sensor and $6.0\text{ mm d}^{-1}\text{ }^{\circ}\text{C}^{-1}$ at the individual stakes.

Going from a single-glacier calibration (Strategy A) to a regional calibration (Strategy B), ~~only~~ the TLR and τ needs changing
630 and the DDF_{ice} decreases slightly to $5.0\text{ mm d}^{-1}\text{ }^{\circ}\text{C}^{-1}$. Here, a lapse rate A TLR of $-0.60\text{ }^{\circ}\text{C } 100\text{ m}^{-1}$ is required, which is distinctly lower than the result of Strategy A. However, this value is close to values used in the Cordillera Darwin before ranging
from -0.60 to $-0.63\text{ }^{\circ}\text{C } 100\text{ m}^{-1}$ (Strelin and Iturraspe, 2007; Koppes et al., 2009). The τ of 1200 s produces a precipitation field with less orographic contribution compared to Strategy A. The value is still distinctly smaller than in Weidemann et al. (2020). This
635 Both changes significantly reduces the snowfall amounts and results in a better match with observed specific-geodetic MBs. ~~One reason, why the other model parameters did not change, may be the fact that Schiaparelli Glacier covers a large part of the study site. However, in general we must not assume that a single glacier calibration is able to tune all parameters correctly except for the TLR.~~

The results suggest that the exclusive use of ablation stakes (Weidemann et al., 2020), which have been installed on the lowest
640 part of Schiaparelli Glacier, for model calibration shows limited utility because no information about accumulation is included. Thus, adding the total mass budget of Schiaparelli Glacier by a flux gate approach brings significant benefit to constrain the drainage basin-wide mass input. Still, the transfer of a SMB model, which has been calibrated to a single glacier, to a regional study site (Strategy A) can imply severe biases in the overall mass budget. This demonstrates that ~~the~~ model parameters is are not transferable from one single glacier to the surrounding. This shortcoming has been reported similarly in previous studies
645 with various melt models and at many locations (e.g., MacDougall and Flowers, 2010; Gurgiser et al., 2013; Zolles et al., 2019). In general, the SMB in the MSM is excessively overestimated which indicates that the SMB model either produces too

little melt or receives excessive snowfall. The latter seems more likely, since melt is well constrained by the stake measurements at least at Schiaparelli Glacier whereas the precipitation amounts are generally more uncertain. By the use of a regional calibration strategy (Strategy B) the agreement between the observed geodetic and modelled surface MB can be significantly improved. This highlights the importance of including regional observations for realistic simulations of regional surface mass balance in the Cordillera Darwin.

Considering the regional distribution of the difference of SMB to the geodetic observations (Fig. 4), the model tends to overestimate the MB on the land-terminating glaciers in the northwest (e.g., 149, 152) and underestimate it in the southeast (e.g., Emma, Pagels) of the massifs. This pattern indicates that snowfall amounts are overestimated on the northwestern slopes and underestimated on the southeastern slopes, which may be associated with neglect of climatic gradients, e.g., in temperature or precipitation. Mass transfer by snowdrift due to the consistent westerlies has been neglected so far. With the addition of a basic snowdrift scheme (Strategy C), the agreement between modelled and observed mass balance can be improved further. Thus, the results express that snowdrift plays an important role for the SMB in the MSM.

The calibration of the SEB models and COSIPY reveals realistic parameter values within the range of previous applications as well (see Table 1). For COSIPY, the calibrated parameters z_{ice} and α_{firn} are lie on the margin of the range, implying that a larger range may be beneficial, or a parameter not considered in calibration is not chosen optimally. However, we decided not to extend the limits of these parameters would result further into physically unrealistic ranges values. We have not been able to find a parameter that was neglected in the calibration and would solve the issue. Apart from the model-inherent parameters, the difficulties with the calibration of COSIPY might alternatively lie in the input dataset. Variables that are only considered in COSIPY and not in the other models are for example wind speed and relative humidity, which both affect turbulent heat fluxes and thereby impact the choice of ice roughness length.

A high discrepancy between modelled and observed mass balance is obtained for two lake-terminating glaciers south of Monte Sarmiento (138 and Lovisato) (Fig. 7). Due to the lake termination, it is expected that the modelled SMB is higher than the specific-geodetic MB. However, the difference is extremely large, especially when considering snow-redistribution due to snowdrift. In Sect. 5.45, we will discuss possible explanations for this discrepancy.

5.2 Surface mass balance of the Monte Sarmiento Massif

The mean annual SMB of ~~-0.795~~ to ~~-1.205~~ m w.e. yr⁻¹ (2000-2013) at Schiaparelli Glacier is distinctly less negative than the previous estimate for the period 2000-2017 (-1.8 ± 0.36 m w.e. yr⁻¹) by Weidemann et al. (2020), however in much better agreement with the satellite observations (-0.79 ± 0.190 m w.e. yr⁻¹). The massif-wide average SMB over the full study period (2000-2022) is estimated between ~~-0.285~~ and ~~-0.07~~ m w.e. yr⁻¹ depending on the model choice. In the eastern part of the CDI, an average SMB of ~~-0.53~~ m w.e. yr⁻¹ has been simulated between 2000 and 2006 using a PDD model (Buttstädt et al., 2009). Similarly large accumulation amounts over the highest parts of the glaciers as well as the extreme ablation over the glacier tongues, that we see at our study site, have been reported for the Southern Patagonian Icefield (Schaefer et al., 2015). We can confirm that the SMB of the MSM is controlled by winter ablation-accumulation and summer temperature as has been observed

680 in the Cordillera Darwin before (Weidemann et al., 2020; Mutz and Aschauer, 2022). The orientation of the individual glaciers does not seem to dictate a particular pattern. Glaciers that receive more direct solar radiation (e.g., Schiaparelli, Conway, Pagels) do not show more negative MBs than glaciers with stronger shading (e.g., Lovisato, 138).

We simulate an average ELA at ~~770-79583~~ m for the MSM. This is close to the mean ELA at 730 ± 50 m simulated at Schiaparelli Glacier in 2000-2017 (Weidemann et al., 2020), however higher than the ELA suggested by Bown et al. (2014)
685 for Ventisquero Glacier at the southwestern edge of the CDI at around 650 m in 2004. In the CDI's northern edge at Marinelli Glacier and the eastern edge at Martial Este Glacier, average ELAs have been reported at around 1100 m (Buttstädt et al., 2009; Bown et al., 2014). The altitude difference can be explained by the more continental conditions due to lee-side effects that reduce the precipitation in the east of the CDI (Strelin and Iturraspe, 2007). Whereas the MSM is located on the western edge of the CDI, directly exposed to the moist westerly winds causing abundant precipitation and, thus, higher accumulation
690 amounts (Bown et al., 2014), which results in lower equilibrium lines.

Ice loss due to dynamical adjustment and calving are assumed to play an important role only for few glaciers in the CDI (Koppes et al., 2009; Bown et al., 2014; Weidemann et al., 2020), like Marinelli Glacier (Porter and Santana, 2003). Weidemann et al. (2020) conclude that mass loss due to SMB processes is the main reason for the recent areal changes of Schiaparelli Glacier. Based on our results, we can confirm that the SMB contributes the largest amount to the ice loss at
695 Schiaparelli Glacier. However, calving is not negligible. Using calibration Strategy A, where the PDD model is tuned to the Schiaparelli Glacier conditions directly, we assess a resulting calving flux of ~~0.28-17~~ m w.e. yr⁻¹, which equals a mass loss of ~~4.266.94~~ Mt yr⁻¹ at Schiaparelli Glacier. The average specific-geodetic MB estimated from elevation changes for the whole study site is with ~~-0.62-55~~ m w.e. yr⁻¹ (2000-2013) distinctly more negative than the SMB (~~-0.2819~~ m w.e. yr⁻¹ by the PDD, in the same period) indicating that dynamical-calving losses are not insignificant in the region. However, in order to determine
700 the calving flux more accurately, detailed information about the ice thickness and velocities at the glacier fronts are required.

5.3 Uncertainty quantification

~~The uncertainties related to the climatic forcing are larger if we rank the runs following Strategy A than following Strategy B (see Table S3). This shows that with a single glacier calibration we are able to represent the local climate at this single glacier, however, we face large difficulties to accurately calibrate the climate variables for an entire region.~~

~~The model-inherent uncertainties show model distinct magnitudes. However, apart from the PDD, all models produce rather stable results looking at the B_{MSMmc} as well as the individual glaciers (Table S4). The increased uncertainties of the PDD model may be due to the reduced sample size of the model specific parameters in this case. To compensate for this disparity, we calculate the uncertainties of the PDD taking only the top 5 instead of the top 10 ranked model runs as we did for the other models. Yet, a direct comparison with the other models might not be straightforward. However, this also shows that the model-specific uncertainty depends strong on the range and sample size of applied parameters.~~

710

Comparing the results of the four model types, the differences are moderate for the B_{MSMnc} , however significant for several individual glaciers. The estimated SMBs of the SEB_Gpot and SEB_G are on average similar and more positive than the PDD and COSIPY.

Overall, the largest uncertainties are related to the climate forcing and the applied model type for values of both, the individual glaciers and the B_{MSMnc} . The model inherent uncertainties are generally smaller. This shows that the model choice is of large importance and highlights the significance of accurate downscaling of climatic forcing data.

5.34 Model intercomparison

Overall, we achieve a very good agreement between the modelled surface and the observed specific-geodetic MB. For most glaciers, the RMSEs are in a similar range as the uncertainties in geodetic MB (Table S5). We want to highlight the remarkable performance of all four models used under these challenging conditions with very sparse observations leading to over-parametrization issues.

Previous studies (e.g., Six et al., 2009; Gabbi et al., 2014; Réveillet et al., 2017) of melt model comparison have come to the conclusion that more complex, physically-based models can achieve more realistic SMB results in case they are based on high-quality and well-distributed in-situ observations. If observations are limited or inferred from distant weather stations, the performance decreases rapidly, and less complex, empirical models produce better results (Gabbi et al., 2014; Réveillet et al., 2017). Since we focus on a study area where in-situ measurements are extremely limited and, thus, need to infer model input from reanalysis data via downscaling, and furthermore glacier SMB is known to be highly correlated with precipitation and air temperature (Weidemann et al., 2020), we strongly challenge the question of which melt-SMB model can produce the most realistic SMB.

Results are validated against the individual specific-geodetic MBs and the stake measurements. The results of this study show that less complex model types overall outperform COSIPY, although the simulated melt at the ablation stakes is best represented by COSIPY, as long as we ignore Schiaparelli Glacier. When including the total mass budget of Schiaparelli Glacier, the performance of COSIPY is higher compared to the other models. Both SEB model variants tend to overestimate the SMB in the MSM on average, however the SEB_G achieves the smallest RMSE compared to geodetic observations of the land-terminating glaciers. The MB of Schiaparelli Glacier, the largest glacier of the massif, is also simulated well by both SEB variants. Comparison of the measured against modelled melt at the stakes delivers similar results for all models with large RMSEs (Table 3). Although the PDDCOSIPY achieves the overall smallest RMSE and bias, the difference to the other models is small compared to the difference to the measurements, we do not see a clear advantage over the other models looking at the measurements individually (Fig. S4 and S5). Instead, COSIPY gets the melt rates very well for the S1to5 during the first years (Fig. S4). Only after 2018, it overestimates the ablation which decreases the overall performance.

Gabbi et al. (2014) concluded that models considering the temperature- and radiation-induced melt separately are more suitable for long-term simulation periods because they are less sensitive to temperature. However, shorter time periods might not be able to bring issues like parameter instability to light (Gabbi et al., 2014), which might apply to our study period. The

importance of correct radiation information cannot be confirmed even by comparing the two SEB model variants we used.

745 Although the agreement with observations can be increased (see Table S1), including accurate radiation calculation (SEB_G) instead of potential radiation (SEB_Gpot) only produces a minor ~~changes~~ improvement in the glacier wide SMBs. Interestingly, the SEB_G produces always slightly larger melt rates at the individual stake locations (Fig. S4), whereas at the automatic ablation sensor we do not see this consistent pattern (Fig. S5).

Overall, due to the small sample size of glaciers ~~and the large impact of Schiaparelli Glacier on the RMSE~~, it is not possible
750 to point out the one best-suited melt-SMB model for the MSM. The ~~very~~ strong correlation with air temperature as well as precipitation make the PDD a good predictor of the SMB. Both SEB model variants show convincing performance as well, although they tend to produce a less negative B_{MSMnc} . Still, the highest agreement with geodetic MB is achieved using the SEB_G. However, COSIPY ~~might overall~~ delivers more accurate and confident results (smaller uncertainty) (Table S6) and can best reproduce the melt at the stakes. Both SEB model variants show convincing performance as well, although they tend
755 to overestimate the B_{MSMnc} . As in this study, in Schneider et al. (2007) the applied PDD and energy balance model at the Gran Campo Nevado showed very similar results. In order to better understand the interaction between the atmosphere and the glacier surface, a physically-based energy and mass balance model like COSIPY is advantageous.

5.45 Challenges and limitations

A large discrepancy between the surface and specific-geodetic MB is modelled for glaciers 138 and Lovisato. Both glaciers
760 are calving, thus, a positive anomaly in SMB is to be expected, however, the difference seems very high. Including the snowdrift parameterization (Strategy C), the discrepancy gets even larger due to the mainly prevailing north-westerlies during snowfall events. The results from the four different SMB models imply a mass loss through calving of 2.50 to 2.95 m w.e. yr⁻¹ and 1.92-2 to 2.25 m w.e. yr⁻¹ for glaciers 138 and Lovisato, which equals an ice mass of 9.736-57 to 11.288-22 Mt yr⁻¹ and 23.8828-75 to 27.6533-83 Mt yr⁻¹, respectively. The question, if these values are realistic, will be discussed in the following
765 We will discuss in the following if these values are realistic.

Assessing satellite images of the last years, it can be confirmed that Lovisato Glacier has significant calving losses, seen through large amounts of icebergs in the proglacial lake (see Fig. 1). Lovisato Glacier has a frontal width of around 500 m. Satellite observations suggest surface velocities of around 400-2 m yr⁻¹ in the recent years. In order to obtain the suggested ice mass loss of 28.75 to 33.83 23.88 to 27.65 Mt yr⁻¹, an ice thickness of around 13059-15087 m would be necessary. The 2019
770 consensus estimate gives an ice thickness of up to ~190 m at this area of Lovisato Glacier (Farinotti et al., 2019). Other ice thickness reconstructions estimate a thickness between 144 to 200 m (Carrivick et al., 2016; Millan et al., 2022). Subsequently, the high calving rates suggested by our results are not-unrealistic realistic for Lovisato Glacier.

For glacier 138, however, we do not see any major icebergs that would indicate a significant calving flux. Surface velocities are below 20 m yr⁻¹ and maximum ice thickness between 50 and 70 m (Farinotti et al., 2019; Millan et al., 2022). This would
775 result in a calving flux 3-4 times magnitudes smaller than implied by our results. Therefore, we reject the calving explanation for glacier 138. It is one of the smallest glaciers that we included in the comparison with satellite observations. Due to the

small size, the uncertainty in the observed elevation change rate is large. Furthermore, the DEMs used for the calculations have large gaps over this glacier, specifically in the accumulation area. Thus, we assume ~~an~~-that in reality increased the uncertainty for glacier 138 ~~is even larger~~, which could cause the large difference between model and observation in this case.

780 Another factor that could explain the large discrepancy between geodetic and surface MB are limitations in the snowdrift parametrization. The snowdrift scheme does not track the snow on its way from one location to another, but identifies locations sheltered from or exposed to wind and, subsequently, corrects the snowfall amounts based on that. Looking at the study site, the question can be asked where the snow deposited at glacier 138 should come from. The main snowdrift direction is towards the south-east. There is no area directly north-west from glacier 138, where we would expect much snowfall that could be
785 blown to and deposited at glacier 138. This highlights one limitation of the snowdrift parametrization. However, even without snowdrift (Strategy B), our results require a calving flux of more than 1.4220 m w.e. yr⁻¹ (Table 2). Thus, limitations are given by the ~~melt-SMB~~ model itself and the climatic forcing as well.

Using one TLR throughout the whole study site and throughout the year is a major simplification. ~~Melt-SMB~~ models are highly sensitive to the air temperature field. It is known that the TLR over mountainous terrain does not only vary temporally but also
790 locally (Gardner and Sharp, 2009; Gardner et al., 2009; Petersen et al., 2013; Ayala et al., 2015; ~~Hanna et al., 2015~~; Heynen et al., 2016; Shaw et al., 2016; Shen et al., 2016; ~~Hanna et al., 2017~~). Bravo et al. (2019a) found that the observed lapse rates at the SPI are steeper in the east compared to the west, and that differences exist between the lower and upper section of glaciers. Thus, it is possible that a northwest to southeast gradient in temperature (lapse rate) prevails in the MSM, affecting the SMB. However, since we do not have any measurements of TLR at the study site that would allow a more realistic estimate,
795 a constant and linear lapse rate is applied.

6. Conclusion

We investigated strategies for ~~melt-SMB~~ model calibration in the Cordillera Darwin in order to achieve realistic simulations of the regional SMB. Therefore, we applied three different calibration strategies, ranging from a local single-glacier calibration transferred to the regional scale (Strategy A), to a regional calibration without (Strategy B) and with (Strategy C) the inclusion
800 of a snowdrift parametrization. This way, we examined the model transferability in space, the advantage of regional mass change observations and the benefit of increasing the complexity level regarding included processes. Furthermore, we constrained the main characteristics of SMB in the MSM. We considered the following measurements: ablation and ice thickness measurements at Schiaparelli Glacier as well as elevation changes and flow velocities from satellite data for the entire study site. Performance of simulated MB is validated against geodetic mass changes and stake observations.
805 Our analysis suggests that the exclusive use of ablation stakes from the lowest part of Schiaparelli Glacier for model calibration shows limited utility because no information about accumulation is included. Adding the total mass budget of Schiaparelli Glacier by a flux gate approach brings significant benefit to constrain the drainage basin-wide mass input. Still, calibration at one single glacier and subsequent transfer to regional scales (Strategy A) resulted in a clearly biased SMB. Such an important

bias implies that spatial model transfers are critical even on such small scales as the MSM. Model performance can be significantly improved by the use of remotely sensed regional observations (Strategy B), e.g., here the annual massif-wide average ~~specific-geodetic~~ MB. Such observations are available on global scales, often dating back to 2000 (e.g., Hugonnet et al., 2021). Including a snowdrift parametrization (Strategy C) can further increase the agreement between modelled and observed MB of individual glacier basins. This demonstrates that snowdrift has an important influence on the accumulation in the MSM where strong and consistent westerly winds prevail.

To answer the main study questions, we can summarize that this study has shown that transferring SMB models in space is a challenge, and common practices can produce distinctly biased estimates (Q1). Thus, we advise to incorporate regional observations for a regional application of SMB models (Q2). Furthermore, we have shown that snowdrift does play an important role for the SMB in the Cordillera Darwin, and thus the inclusion of this process is beneficial (Q3). However, increasing the complexity level of the SMB models from an empirical approach to a physically-based model, did not result in an improvement.

The main characteristics of SMB in the MSM are reproduced in similar way by all four models applied in this study. The massif-wide average annual SMB between 2000 and 2022 ranges between -0.285 and -0.07 m w.e. yr⁻¹ with an average ELA between 770 and 79583 m, depending on the exact model. The SMB is mainly controlled by melt and snowfall as has been observed similarly in southern Patagonia. The spatial pattern is characterized by high amounts of snowfall over the high-altitude areas up to 10 m w.e. yr⁻¹ and extreme surface melt over the glacier tongues down to -10 m w.e. yr⁻¹. The model intercomparison did not indicate one clear best-suited model for SMB simulations in the MSM. Thus, the performance of the SMB cannot generally be improved by increasing the complexity level of the model. The PDD delivered surprisingly unexpectedly good results considering the simplicity of the model. Yet, the physically-based model COSIPY, which is much more challenging to calibrate, did produce convincing results as well, and might produce slightly more stable values (smaller uncertainty and range of values in the 10 top ranked simulations). Both SEB model variants show reasonable results as well, although they tend to overestimate the average SMB in the MSM. Overall, the SEB_G achieves the best agreement with geodetic observations.

The main limitation of this study are the sparse observations in the Cordillera Darwin, which cause over-parametrization and preclude extensive model calibration and validation. We particularly missed information about precipitation amounts in mountainous areas. Moreover, measurements of TLR are missing, which we have shown to be essential for the SMB simulations. With the combination of in-situ and satellite observations, we have been able to appropriately calibrate both fields. However, the uncertainties linked to the climatic forcing are still large. Including snowdrift and solely considering regional calibration targets together with mass budgeting of the most prominent Schiaparelli Glacier, we succeeded to reduce the RMSE with respect to the geodetic measurements below-to their associated errors.

Code and data availability. ERA5 reanalysis data is available via the Copernicus Climate Data Store (<https://cds.climate.copernicus.eu/cdsapp#!/home>). Ice thickness measurements in 2016 are accessible at

<https://doi.org/10.1594/PANGAEA.919331>. Meteorological and ablation stake observations are available ~~on request on the~~ [Pangaea Database \(https://doi.org/10.1594/PANGAEA.956569\)](https://doi.org/10.1594/PANGAEA.956569). The code for the COSIPY model (version 1.4) is available at <https://github.com/cryotools/cosipy>. ~~The code for the PDD model is available at~~ <https://github.com/FAU-glacier-systems/positive-deg-day-model>. ~~The code for the two SEB model variants is available at~~ <https://github.com/FAU-glacier-systems/simplified-energy-balance-model>. ~~Modelling Model forcing and SMB model~~ output of this study is available ~~from the~~ ~~corresponding author on request at~~ <https://doi.org/10.5281/zenodo.7798666>.

845

Author contribution. The concept of this study was developed by Fürst, Temme and Schneider. Temme implemented the simulations with support of Sauter, Arndt and Fürst. In-situ observational data were collected and provided by Schneider, Jaña, Arigony-Neto and Gonzalez. Satellite observations were processed and provided by Farías-Barahona and Seehaus. Temme led the writing process with the support of all authors.

850

Competing interests. The authors declare that they have no conflict of interest.

Disclaimer. The presented content only reflects the authors' views and the European Research Council Executive Agency is not responsible for any use that may be made of the information it contains.

855

Acknowledgements. This research was funded by the German Research Foundation (DFG) within the MAGIC project (FU 1032/5-1). Fürst has received funding from the European Union's Horizon 2020 research and innovation programme via the European Research Council (ERC) as a Starting Grant (StG) under grant agreement No 948290. Farías-Barahona acknowledges the Vicerrectoría de Investigación y Desarrollo, Universidad de Concepción, Postdoctorado VRID. Seehaus received support by the ESA Living Planet Fellowship Programme (Project MIT-AP). Arigony-Neto received funding from the Rio Grande do Sul state Research Support Foundation (FAPERGS nos. 17/25510000518-0 and 21/2551-0002034-2). The authors want to thank Thomas Mölg who provided the model code for the radiation module. The authors gratefully acknowledge the scientific support and HPC resources provided by the Erlangen National High Performance Computing Center (NHR@FAU) of the Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU). NHR funding is provided by federal and Bavarian state authorities. NHR@FAU hardware is partially funded by the German Research Foundation (DFG) – 440719683. The authors want to thank the Chilean National Forest Corporation (CONAF) for enabling and supporting the field work in the Monte Sarmiento Massif, Parque Nacional Alberto de Agostini.

860

865

References

- Albrecht, T., Winkelmann, R., and Levermann, A.: Glacial-cycle simulations of the Antarctic Ice Sheet with the Parallel Ice Sheet Model (PISM)-Part 2: Parameter ensemble analysis, *The Cryosphere*, 14, 633–656, <https://doi.org/10.5194/tc-14-633-2020>, 2020.
- [Arndt, A.; Scherer, D.; Schneider, C.: Atmosphere Driven Mass-Balance Sensitivity of Halji Glacier, Himalayas. Atmosphere 12, 426. https://doi.org/10.3390/atmos12040426, 2021.](https://doi.org/10.3390/atmos12040426)
- Ayala, A., Pellicciotti, F., and Shea, J. M.: Modeling 2 m air temperatures over mountain glaciers: Exploring the influence of katabatic cooling and external warming, *Journal of Geophysical Research: Atmospheres*, 120, 3139–3157, <https://doi.org/10.1002/2015JD023137>, 2015.
- Barcaza, G., Nussbaumer, S. U., Tapia, G., Valdés, J., García, J. L., Videla, Y., Albornoz, A., and Arias, V.: Glacier inventory and recent glacier variations in the Andes of Chile, South America, *Ann Glaciol*, 58, 166–180, <https://doi.org/10.1017/aog.2017.28>, 2017.
- Barstad, I. and Smith, R. B.: Evaluation of an orographic precipitation model, *J Hydrometeorol*, 6, 85–99, <https://doi.org/10.1175/JHM-404.1>, 2005.
- Bentley, C. R.: Mass balance of the Antarctic ice sheet: observational aspects, *Mass Balance of the Cryosphere*, 459–490, <https://doi.org/10.1017/cbo9780511535659.014>, 2009.
- Bown, F., Rivera, A., Zenteno, P., Bravo, C., and Cawkwell, F.: First Glacier Inventory and Recent Glacier Variation on Isla Grande de Tierra Del Fuego and Adjacent Islands in Southern Chile, *Global Land Ice Measurements from Space*, 661–674, https://doi.org/10.1007/978-3-540-79818-7_28, 2014.
- Bown, F., Rivera, A., Pełlicki, M., Bravo, C., Oberreuter, J., and Moffat, C.: Recent ice dynamics and mass balance of Jorge Montt Glacier, Southern Patagonia Icefield, *Journal of Glaciology*, 65, 732–744, <https://doi.org/10.1017/jog.2019.47>, 2019.
- [Braithwaite, R.: Positive degree-day factors for ablation on the Greenland ice sheet studied by energy-balance modelling, J Glaciol, 41 \(137\), 153-160, doi:10.3189/S0022143000017846, 1995.](https://doi.org/10.1017/S0022143000017846)
- [Braithwaite, R. J. and Olesen, O. B.: Calculation of glacier ablation from air temperature, West Greenland, in: Glacier fluctuations and climatic change, Kluwer Academic Publishers, Dordrecht, 219–233, 1989.](https://doi.org/10.1017/S0022143000017846)
- Braun, M. H., Malz, P., Sommer, C., Farías-Barahona, D., Sauter, T., Casassa, G., Soruco, A., Skvarca, P., and Seehaus, T. C.: Constraining glacier elevation and mass changes in South America, *Nat Clim Change*, 9, 130–136, <https://doi.org/10.1038/s41558-018-0375-7>, 2019.
- Bravo, C., Quincey, D. J., Ross, A. N., Rivera, A., Brock, B., Miles, E., and Silva, A.: Air Temperature Characteristics, Distribution, and Impact on Modeled Ablation for the South Patagonia Icefield, *Journal of Geophysical Research: Atmospheres*, 124, 907–925, <https://doi.org/10.1029/2018JD028857>, 2019a.

- Bravo, C., Bozkurt, D., Gonzalez-Reyes, Á., Quincey, D. J., Ross, A. N., Farías-Barahona, D., and Rojas, M.: Assessing snow accumulation patterns and changes on the Patagonian Icefields, *Front Environ Sci*, 7, 1–18, <https://doi.org/10.3389/fenvs.2019.00030>, 2019b.
- [Buisán, S., Earle, M., Collado, J., Kochendorfer, J., Alastrué, J., Wolff, M., Smith, C.G., and López-Moreno, J.I.: Assessment of snowfall accumulation underestimation by tipping bucket gauges on the Spanish operational network. *Atmos. Meas. Tech.*, 10, 1079–1091. <https://doi.org/10.5194/amt-10-1079-2017>, 2017.](#)
- Buttstädt, M., Möller, M., Iturraspe, R., and Schneider, C.: Mass balance evolution of Martial Este Glacier, Tierra del Fuego (Argentina) for the period 1960–2009, *Advances in Geosciences*, 22, 117–124, <https://doi.org/10.5194/adgeo-22-117-2009>, 2009.
- Cannon, A. J., Sobie, S. R., and Murdock, T. Q.: Bias correction of GCM precipitation by quantile mapping: How well do methods preserve changes in quantiles and extremes?, *J Climate*, 28, 6938–6959, <https://doi.org/10.1175/JCLI-D-14-00754.1>, 2015.
- Carrivick, J. L., Davies, B. J., James, W. H. M., Quincey, D. J., and Glasser, N. F.: Distributed ice thickness and glacier volume in southern South America, *Global Planet Change*, 146, 122–132, <https://doi.org/10.1016/j.gloplacha.2016.09.010>, 2016.
- [Cogley, J. C., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser, G., Möller, M., Nicholson, M., and Zemp, M.: Glossary of Glacier Mass Balance and Related Terms, *IACS Contrib. No. 2*, 2011.](#)
- Dadic, R., Mott, R., Lehning, M., and Burlando, P.: Parameterization for wind-induced preferential deposition of snow, *Hydrological Process*, 24, 1994–2006, <https://doi.org/10.1002/hyp.7776>, 2010.
- [DGA: Metodología de inventario público de glaciares, SDT N°447, 2022. Ministerio de Obras Públicas, Dirección General de Aguas Unidad de Glaciología y Nieves. Realizado por: Casassa, G., Espinoza, A., Segovia, A., Huenante, J., 2022.](#)
- Dussailant, I., Berthier, E., Brun, F., Masiokas, M., Hugonnet, R., Favier, V., Rabatel, A., Pitte, P., and Ruiz, L.: Two decades of glacier mass loss along the Andes, *Nat Geosci*, 12, 802–808, <https://doi.org/10.1038/s41561-019-0432-5>, 2019.
- Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth, *Nat Geosci*, 12, 168–173, <https://doi.org/10.1038/s41561-019-0300-3>, 2019.
- Gabbi, J., Carenzo, M., Pellicciotti, F., Bauder, A., and Funk, M.: A comparison of empirical and physically based glacier surface melt models for long-term simulations of glacier response, *Journal of Glaciology*, 60, 1199–1207, <https://doi.org/10.3189/2014JG14J011>, 2014.
- Gacitúa, G., Schneider, C., Arigony, J., González, I., Jaña, R., and Casassa, G.: First ice thickness measurements in Tierra del Fuego at Schiaparelli Glacier, Chile, *Earth Syst Sci Data*, 13, 231–236, <https://doi.org/10.5194/essd-13-231-2021>, 2021.
- Gardner, A. S. and Sharp, M.: Sensitivity of net mass-balance estimates to near-surface temperature lapse rates when employing the degree-day method to estimate glacier melt, *Ann Glaciol*, 50, 80–86, <https://doi.org/10.3189/172756409787769663>, 2009.

- Gardner, A. S., Sharp, M. J., Koerner, R. M., Labine, C., Boon, S., Marshall, S. J., Burgess, D. O., and Lewis, D.: Near-surface temperature lapse rates over arctic glaciers and their implications for temperature downscaling, *J Climate*, 22, 4281–4298, <https://doi.org/10.1175/2009JCLI2845.1>, 2009.
- Garreaud, R. D., Vuille, M., Compagnucci, R., and Marengo, J.: Present-day South American climate, *Palaeogeogr Palaeoclimatol Palaeoecol*, 281, 180–195, <https://doi.org/10.1016/j.palaeo.2007.10.032>, 2009.
- 935 Glasser, N. F., Harrison, S., Winchester, V., and Aniya, M.: Late Pleistocene and Holocene palaeoclimate and glacier fluctuations in Patagonia, *Glob Planet Change*, 43, 79–101, <https://doi.org/10.1016/j.gloplacha.2004.03.002>, 2004.
- Groos, A. R., Mayer, C., Smiraglia, C., Diolaiuti, G., and Lambrecht, A.: A first attempt to model region-wide glacier surface mass balances in the Karakoram: Findings and future challenges, *Geografia Fisica e Dinamica Quaternaria*, 40, 137–159, 940 <https://doi.org/10.4461/GFDQ2017.40.10>, 2017.
- Gudmundsson, L., Bremnes, J. B., Haugen, J. E., and Engen-Skaugen, T.: Technical Note: Downscaling RCM precipitation to the station scale using statistical transformations - A comparison of methods, *Hydrol Earth Syst Sci*, 16, 3383–3390, <https://doi.org/10.5194/hess-16-3383-2012>, 2012.
- Gurgiser, W., Molg, T., Nicholson, L., and Kaser, G.: Mass-balance model parameter transferability on a tropical glacier, 945 *Journal of Glaciology*, 59, 845–858, <https://doi.org/10.3189/2013JoG12J226>, 2013.
- Hanna, E., Mernild, S., de Villiers, S., and Yde, J.: Surface air temperature variations and lapse rates on Olivares Gamma Glacier, Rio Olivares Basin, Central Chile, from a novel meteorological sensor network, *Meteorological Applications*, 2017, 12–14, 2017.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, 950 D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., de Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 955 Heynen, M., Miles, E., Ragetti, S., Buri, P., Immerzeel, W. W., and Pellicciotti, F.: Air temperature variability in a high-elevation Himalayan catchment, *Ann Glaciol*, 57, 212–222, <https://doi.org/10.3189/2016AoG71A076>, 2016.
- [Hock, R.: A distributed temperature-index ice- and snowmelt model including potential direct solar radiation, *J Glaciol*, 45, 101-111, 1999.](#)
- Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussailant, I., Brun, F., 960 and Kääb, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- Huintjes, E., Sauter, T., Schroter, B., Maussion, F., Yang, W., Kropaček, J., Buchroithner, M., Scherer, D., Kang, S., and Schneider, C.: Evaluation of a Coupled Snow and Energy Balance Model for Zhadang Glacier, Tibetan Plateau, Using

- Glaciological Measurements and Time-Lapse Photography, *Arct Antarct Alp Res*, 47, 573–590, 1965 <https://doi.org/10.1657/AAAR0014-073>, 2015.
- Jarosch, A. H., Anslow, F. S., and Clarke, G. K. C.: High-resolution precipitation and temperature downscaling for glacier models, *Clim Dynam*, 38, 391–409, <https://doi.org/10.1007/s00382-010-0949-1>, 2012.
- [Jiang, Q. and Smith, R. B.: Cloud timescales and orographic precipitation, *J Atmos Sci*, 60, 1543–1559, <https://doi.org/10.1175/2995.1>, 2003.](https://doi.org/10.1175/2995.1)
- 970 Koch, J.: Little Ice Age and recent glacier advances in the Cordillera Darwin, Tierra del Fuego, Chile, *Anales del Instituto de la Patagonia*, 43, 127–136, <https://doi.org/10.4067/s0718-686x2015000100011>, 2015.
- Koppes, M., Hallet, B., and Anderson, J.: Synchronous acceleration of ice loss and glacial erosion, *Glacier Marine* [Marine](https://doi.org/10.3189/002214309788608796) [37](https://doi.org/10.3189/002214309788608796) [olocene](https://doi.org/10.3189/002214309788608796) [li](https://doi.org/10.3189/002214309788608796), Chilean Tierra del Fuego, *Journal of Glaciology*, 55, 207–220, <https://doi.org/10.3189/002214309788608796>, 2009.
- Lehning, M., Löwe, H., Ryser, M., and Raderschall, N.: Inhomogeneous precipitation distribution and snow transport in steep 975 terrain, *Water Resour Res*, 44, 1–19, <https://doi.org/10.1029/2007WR006545>, 2008.
- Lenaerts, J. T. M., van den Broeke, M. R., van Wessem, J. M., van de Berg, W. J., van Meijgaard, E., van Uft, L. H., and Schaefer, M.: Extreme precipitation and climate [gradients](https://doi.org/10.1175/JCLI-D-13-00579.1) [-37](https://doi.org/10.1175/JCLI-D-13-00579.1) [olocene](https://doi.org/10.1175/JCLI-D-13-00579.1) [37](https://doi.org/10.1175/JCLI-D-13-00579.1) [n](https://doi.org/10.1175/JCLI-D-13-00579.1) patagonia revealed by high-resolution regional atmospheric climate modeling, *J Climate*, 27, 4607–4621, <https://doi.org/10.1175/JCLI-D-13-00579.1>, 2014.
- Lopez, P., Chevallier, P., Favier, V., Pouyaud, B., Ordenes, F., and Oerlemans, J.: A regional view of fluctuations in glacier 980 length in southern South America, *Global Planet Change*, 71, 85–108, <https://doi.org/10.1016/j.gloplacha.2009.12.009>, 2010.
- MacDougall, A. H. and Flowers, G. E.: Spatial and temporal transferability of a distributed energy-balance glacier melt model, *J Climate*, 24, 1480–1498, <https://doi.org/10.1175/2010JCLI3821.1>, 2011.
- Masiokas, M. H., Rivera, A., Espizua, L. E., Villalba, R., Delgado, S., and Aravena, J. C.: Glacier fluctuations in extratropical South America during the past 1000 years, *Palaeogeogr Palaeoclimatol—Palaeoecol*, 281, 242–268, 985 <https://doi.org/10.1016/j.palaeo.2009.08.006>, 2009.
- Meier, W. J. H., Griebinger, J., Hochreuther, P., and Braun, M. H.: An updated multi-temporal glacier inventory [for](https://doi.org/10.3389/feart.2018.00062) [the](https://doi.org/10.3389/feart.2018.00062) [37](https://doi.org/10.3389/feart.2018.00062) [olocene](https://doi.org/10.3389/feart.2018.00062) [37](https://doi.org/10.3389/feart.2018.00062) [n](https://doi.org/10.3389/feart.2018.00062) patagonian andes with changes between the little ice age and 2016, *Front Earth Sci—(Lausanne)*, 6, <https://doi.org/10.3389/feart.2018.00062>, 2018.
- Meier, W. J. H., Aravena, J. C., Griebinger, J., Hochreuther, P., Soto-Rogel, P., Zhu, H., de Pol-Holz, R., Schneider, C., and 990 Braun, M. H.: [La](https://doi.org/10.3390/geosciences9080340) [37](https://doi.org/10.3390/geosciences9080340) [olocene](https://doi.org/10.3390/geosciences9080340) [te](https://doi.org/10.3390/geosciences9080340) holocene glacial fluctuations of Schiaparelli Glacier at Monte Sarmiento Massif, Tierra del Fuego (54°24's), *Geosciences—(Switzerland)*, 9, 3–540, <https://doi.org/10.3390/geosciences9080340>, 2019.
- Melkonian, A. K., Willis, M. J., Pritchard, M. E., Rivera, A., Bown, F., and Bernstein, S. A.: Satellite-derived volume loss rates and glacier speeds for the Cordillera Darwin Icefield, Chile, *The Cryosphere*, 7, 823–839, <https://doi.org/10.5194/tc-7-823-2013>, 2013.
- 995 Millan, R., Mouginot, J., Rabatel, A., and Morlighem, M.: Ice velocity and thickness of the world's glaciers, *Nat Geosci*, 15, 124–129, <https://doi.org/10.1038/s41561-021-00885-z>, 2022.

- Minowa, M., Schaefer, M., Sugiyama, S., Sakakibara, D., and Skvarca, P.: Frontal ablation and mass loss of the Patagonian icefields, *Earth Planet Sci Lett*, 561, 116811, <https://doi.org/10.1016/j.epsl.2021.116811>, 2021.
- Mölg, T., Cullen, N. J., Hardy, D. R., Winkler, M., and Kaser, G.: Quantifying climate change in the tropical midtroposphere over East Africa from glacier shrinkage on Kilimanjaro, *J Clim*, 22, 4162–4181, <https://doi.org/10.1175/2009JCLI2954.1>, 2009a.
- Mölg, T., Cullen, N. J., and Kaser, G.: Solar radiation, cloudiness and longwave radiation over low-latitude glaciers: Implications for mass-balance modelling, *Journal of Glaciology*, 55, 292–302, <https://doi.org/10.3189/002214309788608822>, 2009b.
- Mott, R., Faure, F., Lehning, M., Löwe, H., Hynek, B., Michlmayer, G., Prokop, A., and Schöner, W.: Simulation of seasonal snow-cover distribution for glacierized sites on Sonnblick, Austria, with the Alpine3D model, *Ann Glaciol*, 49, 155–160, <https://doi.org/10.3189/172756408787814924>, 2008.
- Mutz, S. G. and Aschauer, J.: Empirical glacier mass-balance models for South America, *Journal of Glaciology*, 1–15, <https://doi.org/10.1017/jog.2022.6>, 2022.
- Nuth, C. and Kääb, A.: Co-registration and bias corrections of satellite elevation data sets for quantifying glacier thickness change, *The Cryosphere*, 5, 271–290, <https://doi.org/10.5194/tc-5-271-2011>, 2011.
- Oerlemans, J.: *Glaciers and climate change*, 2nd edition., Routledge, London, 2001.
- Oerlemans, J. and Knap, W. H.: A 1 year record of global radiation and albedo in the ablation zone of Morteratschgletscher, Switzerland, *Journal of Glaciology*, 44, 231–238, <https://doi.org/10.1017/S0022143000002574>, 1998.
- Pellicciotti, F., Brock, B., Strasser, U., Burlando, P., Funk, M., and Corripio, J.: An enhanced temperature-index glacier melt model including the shortwave radiation balance: Development and testing for Haut Glacier d’Arolla, Switzerland, *Journal of Glaciology*, 51, 573–587, <https://doi.org/10.3189/172756505781829124>, 2005.
- Pellicciotti, F., Helbing, J., Rivera, A., Favier, V., Corripio, J., Araos, J., Sicart, J.-E., and Carenzo, M.: A study of the energy balance and melt regime on Juncal Norte Glacier, semi-arid Andes of central Chile, using melt models of different complexity, *Hydrol Process*, 22, 3980–3997, <https://doi.org/10.1002/hyp.7085>, 2008.
- Petersen, L., Pellicciotti, F., Juszak, I., Carenzo, M., and Brock, B.: Suitability of a constant air temperature lapse rate over an Alpine glacier: testing the Greuell and Böhm model as an alternative, *Ann Glaciol*, 54, 120–130, <https://doi.org/10.3189/2013AoG63A477>, 2013.
- Pollard, D., Chang, W., Haran, M., Applegate, P., and DeConto, R.: Large ensemble modeling of the last deglacial retreat of the West Antarctic Ice Sheet: Comparison of simple and advanced statistical techniques, *Geosci Model Dev*, 9, 1697–1723, <https://doi.org/10.5194/gmd-9-1697-2016>, 2016.
- Porter, C. and Santana, A.: Rapid 20th Century Retreat of Ventisquero Marinelli in the Cordillera Darwin Icefield, *Anales del Instituto de la Patagonia*, 31, 17–26, 2003.
- Rada, C. and Martinez, N.: UNCHARTED: Cordillera Darwin v0.98, <https://doi.org/https://doi.org/10.6084/m9.figshare.19246140.v1>, 2022.

- [Rasmussen, R., Baker, B., Kochendorfer, J., Meyers, T., Landolt, S., Fischer, A.P., Black, J., Thériault, J.M., Kucera, P., Gochis, D., Smith, C., Nitu, R., Hall, M., Ikeda, K., and Gutmann, E.: How Well Are We Measuring Snow: The NOAA/FAA/NCAR Winter Precipitation Test Bed, *B Am Meteorol Soc*, 93, 811–829, <https://doi.org/10.1175/BAMS-D-11-00052.1>, 2012.](https://doi.org/10.1175/BAMS-D-11-00052.1)
- 035 Réveillet, M., Vincent, C., Six, D., and Rabatel, A.: Which empirical model is best suited to simulate glacier mass balances?, *Journal of Glaciology*, 63, 39–54, <https://doi.org/10.1017/jog.2016.110>, 2017.
- Rignot, E., Rivera, A., and Casassa, G.: Contribution of the Patagonia Icefields of South America to Sea Level Rise, *Science* (1979), 302, 434–437, <https://doi.org/10.1126/science.1087393>, 2003.
- Rott, H., Müller, F., Nagler, T., and Floricioiu, D.: The imbalance of glaciers after disintegration of Larsen-B ice shelf, Antarctic Peninsula, *The Cryosphere*, 5, 125–134, <https://doi.org/10.5194/tc-5-125-2011>, 2011.
- 040 [Rounce D.R., Khurana T., Short M.B., Hock R., Shean D.E., Brinkerhoff D.J.: Quantifying parameter uncertainty in a largescale glacier evolution model using Bayesian inference: application to High Mountain Asia. *J Glaciol* 66 \(256\), 175–187, <https://doi.org/10.1017/jog.2019.91>, 2020.](https://doi.org/10.1017/jog.2019.91)
- Sauter, T.: Revisiting extreme precipitation amounts over southern South America and implications for the Patagonian Icefields, *Hydrol Earth Syst Sci*, 24, 2003–2016, <https://doi.org/10.5194/hess-24-2003-2020>, 2020.
- 045 Sauter, T. and Galos, P.: Effects of local advection on the spatial sensible heat flux variation on a mountain glacier, *The Cryosphere*, 10, 2887–2905, <https://doi.org/10.5194/tc-10-2887-2016>, 2016.
- Sauter, T., Arndt, A., and Schneider, C.: COSIPY v1.3 – an open-source coupled snowpack and ice surface energy and mass balance model, *Geosci Model Dev*, 13, 5645–5662, <https://doi.org/10.5194/gmd-13-5645-2020>, 2020.
- 1050 Schaefer, M., Machguth, H., Falvey, M., and Casassa, G.: Modeling past and future surface mass balance of the Northern Patagonia Icefield, *J Geophys Res—Earth Surf*, 118, 571–588, <https://doi.org/10.1002/jgrf.20038>, 2013.
- Schaefer, M., Machguth, H., Falvey, M., Casassa, G., and Rignot, E.: Quantifying mass balance processes on the Southern Patagonia Icefield, *The Cryosphere*, 9, 25–35, <https://doi.org/10.5194/tc-9-25-2015>, 2015.
- Schneider, C., Glaser, M., Kilian, R., Santana, A., Butorovic, N., and Casassa, G.: Weather Observations Across the Southern Andes at 53°S, *Phys Geogr*, 24, 97–119, <https://doi.org/10.2747/0272-3646.24.2.97>, 2003.
- 1055 Schneider, C., Kilian, R., and Glaser, M.: Energy balance in the ablation zone during the summer season at the Gran Campo Nevado Ice Cap in the Southern Andes, *Global and Planetary Change*, 59, 175–188, <https://doi.org/10.1016/j.gloplacha.2006.11.033>, 2007.
- Schuler, T. v., Crochet, P., Hock, R., Jackson, M., Barstad, I., and Jóhannesson, T.: Distribution of snow accumulation on the Svartisen ice cap, Norway, assessed by a model of orographic precipitation, *Hydrol Process*, 22, 3998–4008, <https://doi.org/10.1002/hyp.7073>, 2008.
- 1060 Seehaus, T., Marinsek, S., Helm, V., Skvarca, P., and Braun, M.: Changes in ice dynamics, elevation and mass discharge of Dinsmoor–Bombardier–Edgeworth glacier system, Antarctic Peninsula, *Earth Planet Sci Lett*, 427, 125–135, <https://doi.org/10.1016/j.epsl.2015.06.047>, 2015.

- 065 Seehaus, T., Cook, A. J., Silva, A. B., and Braun, M.: Changes in glacier dynamics in the northern Antarctic Peninsula since 1985, *The Cryosphere*, 12, 577–594, <https://doi.org/10.5194/tc-12-577-2018>, 2018.
- Shaw, T. E., Brock, B. W., Fyffe, C. L., Pellicciotti, F., Rutter, N., and Diotri, F.: Air temperature distribution and energy-balance modelling of a debris-covered glacier, *Journal of Glaciology*, 62, ~~1–14~~[185–198](https://doi.org/10.1017/jog.2016.31), <https://doi.org/10.1017/jog.2016.31>, 2016.
- 070 Shen, Y., Shen, Y., Goetz, J., and Brenning, A.: Spatial-temporal variation of near-surface temperature lapse rates over the Tianshan Mountains, central Asia, *Journal of Geophysical Research: Atmospheres*, 121, 14,006–14,017, <https://doi.org/10.1002/2016JD025711>, 2016.
- Six, D., Wagon, P., Sicart, J. E., and Vincent, C.: Meteorological controls on snow and ice ablation for two contrasting months on Glacier de Saint-Sorlin, France, *Ann Glaciol*, 50, 66–72, <https://doi.org/10.3189/172756409787769537>, 2009.
- 075 Smith, R. B. and Barstad, I.: A linear theory of orographic precipitation, *J Atmos Sci*, 61, 1377–1391, [https://doi.org/10.1175/1520-0469\(2004\)061<1377:ALTOOP>2.0.CO;2](https://doi.org/10.1175/1520-0469(2004)061<1377:ALTOOP>2.0.CO;2), 2004.
- Smith, R. B. and Evans, J. P.: Orographic precipitation and water vapor fractionation over the southern Andes, *J Hydrometeorol*, 8, 3–19, <https://doi.org/10.1175/JHM555.1>, 2007.
- Strelin, J. and Iturraspe, R.: Recent evolution and mass balance of Cordón Martial glaciers, Cordillera Fueguina Oriental, *Global Planet Change*, 59, 17–26, <https://doi.org/10.1016/j.gloplacha.2006.11.019>, 2007.
- 080 Strelin, J., Casassa, G., Rosqvist, G., and Holmlund, P.: Holocene glaciations in the Ema Glacier valley, Monte Sarmiento Massif, Tierra del Fuego, *Palaeogeogr Palaeoclimatol Palaeoecol*, 260, 299–314, <https://doi.org/10.1016/j.palaeo.2007.12.002>, 2008.
- Strozzi, T., Luckman, A., Murray, T., Wegmuller, U., and Werner, C. L.: Glacier motion estimation using SAR offset-tracking procedures, *IEEE Transactions on Geoscience and Remote Sensing*, 40, 2384–2391, <https://doi.org/10.1109/TGRS.2002.805079>, 2002.
- 085 Stuefer, M., Rott, H., and Skvarca, P.: Glacier Perito Moreno, Patagonia: Climate sensitivities and glacier characteristics preceding the 2003/04 and 2005/06 damming events, *Journal of Glaciology*, 53, 3–16, <https://doi.org/10.3189/172756507781833848>, 2007.
- 090 Temme, F., Turton, J. v., Mölg, T., and Sauter, T.: Flow regimes and Föhn types characterize the local climate of Southern Patagonia, *Atmosphere (Basel)*, 11, [899](https://doi.org/10.3390/ATMOS11090899), <https://doi.org/10.3390/ATMOS11090899>, 2020.
- [van Pelt, W. J. J., Oerlemans, J., Reijmer, C. H., Pohjola, V. A., Pettersson, R., and van Angelen, J. H.: Simulating melt, runoff and refreezing on Nordenskiöldbreen, Svalbard, using a coupled snow and energy balance model, *The Cryosphere*, 6, 641–659, <https://doi.org/10.5194/tc-6-641-2012>, 2012.](https://doi.org/10.5194/tc-6-641-2012)
- 095 Villalba, R., Lara, A., Boninsegna, J. A., Masiokas, M., Delgado, S., Aravena, J. C., Roig, F. A., Schmelter, A., Wolodarsky, A., and Ripalta, A.: Large-Scale Temperature Changes Across the Southern Andes: 20th-Century Variations in the Context of the Past 400 Years, 177–232, https://doi.org/10.1007/978-94-015-1252-7_10, 2003.

- Warscher, M., Strasser, U., Kraller, G., Marke, T., Franz, H., and Kunstmann, H.: Performance of complex snow cover descriptions in a distributed hydrological model system: A case study for the high Alpine terrain of the Berchtesgaden Alps, *Water Resour Res*, 49, 2619–2637, <https://doi.org/10.1002/wrcr.20219>, 2013.
- Weidemann, S., Sauter, T., Schneider, L., and Schneider, C.: Impact of two conceptual precipitation downscaling schemes on mass-balance modeling of Gran Campo Nevado ice cap, Patagonia, *Journal of Glaciology*, 59, 1106–1116, <https://doi.org/10.3189/2013JG13J046>, 2013.
- Weidemann, S. S., Sauter, T., Malz, P., Jaña, R., Arigony-Neto, J., Casassa, G., and Schneider, C.: Glacier mass changes of lake-terminating grey and tyndall glaciers at the southern patagonia icefield derived from geodetic observations and energy and mass balance modeling, *Front Earth Sci (Lausanne)*, 6, 1–16, <https://doi.org/10.3389/feart.2018.00081>, 2018a.
- [Weidemann, S., T. Sauter, R. Kilian, D. Steger, N. Butorovic and C. Schneider: A 17-year Record of Meteorological Observations Across the Gran Campo Nevado Ice Cap in Southern Patagonia, Chile, Related to Synoptic Weather Types and Climate Modes, Front Earth Sci, 6, https://doi.org/10.3389/feart.2018.00053, 2018b.](#)
- Weidemann, S. S., Arigony-Neto, J., Jaña, R., Netto, G., Gonzalez, I., Casassa, G., and Schneider, C.: Recent climatic mass balance of the schiaparelli glacier at the monte sarmiento massif and reconstruction of little ice age climate by simulating steady-state glacier conditions, *Geosciences (Switzerland)*, 10, 1–17, <https://doi.org/10.3390/geosciences10070272>, 2020.
- Willis, M. J., Melkonian, A. K., Pritchard, M. E., and Rivera, A.: Ice loss from the Southern Patagonian Ice Field, South America, between 2000 and 2012, *Geophys Res Lett*, 39, 1–6, <https://doi.org/10.1029/2012GL053136>, 2012.
- Winstral, A. and Marks, D.: Simulating wind fields and snow redistribution using terrain-based parameters to model snow accumulation and melt over a semi-arid mountain catchment, *Hydrol Process*, 16, 3585–3603, <https://doi.org/10.1002/hyp.1238>, 2002.
- Ziemen, F. A., Hock, R., Aschwanden, A., Khroulev, C., Kienholz, C., Melkonian, A., and Zhang, J.: Modeling the evolution of the Juneau Icefield between 1971 and 2100 using the Parallel Ice Sheet Model (PISM), *Journal of Glaciology*, 62, 199–214, <https://doi.org/10.1017/jog.2016.13>, 2016.
- Zolles, T., Maussion, F., Galos, P., Gurgiser, W., and Nicholson, L.: Robust uncertainty assessment of the spatio-temporal transferability of glacier mass and energy balance models, *The Cryosphere*, 13, 469–489, <https://doi.org/10.5194/tc-13-469-2019>, 2019.