

Late summer transition from a free-tropospheric to boundary layer source of Aitken mode aerosol in the high Arctic

Ruth Price¹, Andrea Baccarini^{2,*}, Julia Schmale², Paul Zieger^{3,4}, Ian M. Brooks¹, Paul Field^{1,5}, and Ken S. Carslaw¹

¹School of Earth and Environment, University of Leeds, Leeds, UK

²Extreme Environments Research Laboratory, École Polytechnique Fédérale de Lausanne, Sion, Switzerland

³Department of Environmental Science, Stockholm University, Stockholm, Sweden

⁴Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

⁵Met Office, Exeter, UK

* now at: Laboratory of Atmospheric Processes and their Impacts, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

Correspondence: Ruth Price (eersp@leeds.ac.uk)

Abstract. In the Arctic, the aerosol budget plays a particular role in determining the behaviour of clouds, which are important for the surface energy balance and thus for the region's climate. A key question is the extent to which cloud condensation nuclei in the high Arctic summertime boundary layer are controlled by local emission and formation processes as opposed to transport from outside. Each of these sources is likely to respond differently to future changes in ice cover. Here we use a global model and observations from ship and aircraft field campaigns to understand the source of high Arctic aerosol in late summer. We find that particles formed remotely, i.e. ~~at lower latitudes~~, at latitudes outside the Arctic, are the dominant source of boundary layer Aitken mode particles during the sea ice melt period up to the end of August. Particles from such remote sources, entrained into the boundary layer from the free troposphere, account for nucleation and Aitken mode particle concentrations that are otherwise underestimated by the model. This source from outside the high Arctic declines as photochemical rates decrease towards the end of summer, and is largely replaced by local new particle formation driven by iodic acid ~~emitted from the surface and associated with created during~~ increases the simulated Aitken mode particle concentrations by two orders of magnitude during sea ice freeze-up and is consistent with strong fluctuations in nucleation mode concentrations that occur in September. Our results suggest a high-Arctic aerosol regime shift in late summer, and only after this shift do cloud condensation nuclei become sensitive to local aerosol processes.

15 1 Introduction

The Arctic is a key component of the global climate system. Over the past five decades, the mean surface temperature of the Arctic has increased 3–4 times faster than the global average (AMAP, 2021; Rantanen et al., 2022). Regional changes in the Arctic can have global impacts, such as climate feedbacks from albedo changes due to loss of sea ice, land ice and snow (Notz and Stroeve, 2018; Zhang et al., 2019; Flanner et al., 2011); changes in carbon sources due to increased wildfires (Walker et al.,

20 2019; Randerson et al., 2006) and melting permafrost (Hugelius et al., 2014; Biskaborn et al., 2019); and global sea level rise from melting of the Greenland ice sheet (Cazenave et al., 2018; Shepherd et al., 2012).

Clouds are a major control on the surface energy balance in the Arctic. Due to the low solar insolation and high albedo of sea ice in the high Arctic (i.e. pack ice regions north of 80 °), the shortwave cooling effect of clouds is less important than at lower latitudes. Instead, longwave effects dominate such that the net radiative effect of low-level Arctic clouds is surface warming for 25 all but a few weeks in the middle of summer (Curry et al., 1993; Shupe and Intrieri, 2004; Sedlar et al., 2011; Kay et al., 2016).

Low clouds are common and can persist in the mixed-phase for several days (Shupe, 2011). Aerosol particles act as cloud condensation nuclei (CCN) and ice-nucleating particles (INP) and therefore influence the behaviour of clouds. Observational Arctic case studies have shown that perturbations to aerosol concentrations can change the radiative effect of clouds (Lubin and Vogelmann, 2006; Garrett and Zhao, 2006; Mauritzen et al., 2011). Modelling studies reproduce this behaviour, while also 30 highlighting the difficulty in creating models that can accurately simulate the complex behaviour of Arctic clouds (Alterskjaer et al., 2010; Birch et al., 2012; Stevens et al., 2018).

The aerosol budget of the high Arctic is a balance of different processes, including natural-and-anthropogenic primary emissions, new particle formation, condensational growth from vapours, long-range transport of anthropogenic emissions, and wet 35 scavenging (Willis et al., 2018; Schmale et al., 2021). Transport of aerosol from lower latitudes is more efficient during the winter and spring because it is thermodynamically easier in winter for air to enter the Arctic region from the south than it is in summer (Stohl, 2006).

Cold air masses sitting over ice-covered portions of the Arctic Ocean create sharp north-south temperature gradients, described as the polar dome, that act as a barrier for air moving towards the Arctic lower troposphere from the south.

In the winter, several factors make it easier for air from south to penetrate the polar dome, including a more southward extent of the dome, and cooling of air during transport due to proximity to snow and ice surfaces over land.

Also, the removal of aerosol 40 by frozen precipitation in winter is less effective than removal by drizzle in summer (Browse et al., 2012; Korhonen et al., 2008b). These seasonal cycles of aerosol transport and removal efficiency results in a transition from Arctic haze in the spring to more pristine conditions in the summer with fewer accumulation mode particles, less anthropogenic influence, and more, smaller nucleation and Aitken mode particles (Engvall et al., 2008; Ström et al., 2003; Stohl, 2006; Tunved et al., 2013; Schmale et al., 2022).

Such a transition is evident in measurements e.g. from Svalbard (Ström et al., 2003; Engvall et al., 2008; Tunved et al., 2013; Karl et al., 2014), from pan-Arctic observatories (Schmale et al., 2022), or from the recent year-long MOSAiC campaign in the pack ice region (Boyer et al., 2023).

Unlike the winter, when long-range transport is the dominant source of Arctic aerosols, summertime particles are thought to be strongly controlled by new particle formation (NPF) and growth from precursor vapours, which takes place in many Arctic locations. However, the large variety of vapours that can play a role, as well as the strong seasonal variation in key processes, 50 have made it difficult to understand the main drivers of Arctic NPF. Observations of Arctic NPF are discussed in Schmale and Baccarini (2021) and will be briefly summarised here. Sulphuric acid (H_2SO_4) and methanesulphonic acid (MSA) have been shown to drive NPF and growth in regions close to or influenced by open water such as Svalbard and the Canadian Arctic archipelago (Beek et al., 2020; Chang et al., 2011; Willis et al., 2017) (Heintzenberg et al., 2017; Beck et al., 2020; Chang et al., 2011; Willis et al., 2017).

The marine biogenic precursor dimethylsulphide (DMS) is the main source of H_2SO_4 and MSA to these regions. Open water

55 is also a source of organic vapours which are observed to contribute to condensational growth of small particles (Willis et al., 2017). Ammonia has biogenic sources in the Arctic, for example seabird colonies, and has been observed to contribute to NPF events in Svalbard and Greenland (Croft et al., 2016; Beck et al., 2020). ~~Iodide-acid-()~~Iodine is known to be emitted from kelp in coastal areas and ~~has iodine-containing compounds have~~ been observed to drive NPF events near the coast of Greenland (Sipila et al., 2016; Allan et al., 2015). Of relevance to this study, Baccarini et al. (2020a) recently observed ~~iodide-acid-driven NPF~~NPF driven by iodic acid (HIO_3) in the pack ice region of the Arctic Ocean during the sea ice freeze-up, suggesting an ~~iodide-acid-iodine~~ source from snow, ice or ocean water. Recent laboratory results elucidate the chemical pathway for the creation of iodic acid from iodine (Finkenzeller et al., 2022).

60 Modelling studies have demonstrated the importance of NPF for the budget of Arctic aerosol. Merikanto et al. (2009) and Gordon et al. (2017) used the global aerosol model GLOMAP to show the importance of NPF on a global scale. Both studies indicate that a high fraction (greater than 80%) of particles and CCN in the Arctic are derived from new particle formation. While the model configuration in Merikanto et al. (2009) only included parameterisations based on H_2SO_4 , the model used by Gordon et al. (2017) included parameterisations for neutral and ion-induced binary (H_2SO_4 -water) and ternary (H_2SO_4 -ammonia-water) NPF, NPF from organic molecules and H_2SO_4 , and pure organic NPF driven by highly oxygenated molecules (HOMs). The authors found that a significant fraction (greater than 40%) of Arctic CCN originate from secondary organic aerosol, including HOMs. Korhonen et al. (2008b) and Browse et al. (2014), both using GLOMAP to investigate Arctic aerosol, found that boundary layer NPF driven by H_2SO_4 was required to explain measured size distributions or CCN concentrations in the high Arctic during summer. Karl et al. (2012) used an aerosol dynamics model to study NPF events that were observed during the summers of 1996, 2001 and 2008. H_2SO_4 and organic vapours were used in the model to drive NPF events. Simulations of NPF driven by H_2SO_4 followed by growth from condensation of organic vapours were able to reproduce the particle size distributions observed during NPF events. The inclusion of organic vapours as a driver of NPF led to an overprediction in the concentration of particles, though the authors note significant uncertainty in the sources and concentrations of organic vapours in the high Arctic. Croft et al. (2019) used a chemical transport and aerosol microphysics model to show that condensation of secondary organic vapour played a key role in particle formation events in the Canadian Arctic during the summer. The vapours were assumed to have a marine, biogenic source. Browse et al. (2014), using GLOMAP to investigate the response of Arctic aerosol to sea ice loss, found that boundary layer NPF driven by was required to explain measured CCN concentrations in the high Arctic during summer.

75 Boundary layer NPF as a source of Arctic aerosol implies a potentially high sensitivity of the local aerosol budget to the changing climate. Any future increase in the extent of open water and the marginal ice zone in summer will affect the emission of aerosol and precursor gases to the atmosphere. Dall’Osto et al. (2017, 2018) have found a correlation between frequency of NPF events and the extent of open water near Svalbard and Greenland, respectively. Such an increase in occurrence of NPF with sea ice loss could be expected to increase CCN concentrations in the Arctic under future warmer conditions, though this is far from certain since changing sea ice extent is not the only controlling factor. Gilgen et al. (2018) found that reduction of sea ice in the year 2050 leads to increased emissions of DMS in a global aerosol-climate model. The increased DMS emissions, along with increased sea spray aerosol and meteorological changes, cause higher cloud drop number concentrations over the Arctic

90 Ocean. This is in line with results from another global model study, Struthers et al. (2011), which used an atmospheric climate model to investigate the response of sea spray aerosol to sea ice loss. They found a strong increase in sea salt emissions and thus higher cloud drop concentrations, but the effect on clouds and energy budget was uncertain due to poor model representations of aerosol-cloud interactions. In contrast to these studies, the results from Browse et al. (2014) suggest that interactions between aerosol particles of different sizes could lead to a suppression of NPF in an ice-free Arctic summer. Their results showed an
95 increase in the sink of condensable vapours due to stronger emission of sea spray aerosol from the open water, resulting in a decrease in the concentration of smaller particles from NPF. In addition, the growth of sea spray particles from condensation of vapours shifted the size distribution to larger sizes, leading to enhanced scavenging by precipitation and a decrease in drop concentrations.

100 The conflicting results from Browse et al. (2014), Gilgen et al. (2018) and Struthers et al. (2011) highlight the difficulty in modelling the Arctic aerosol budget and how it might change in future. Uncertainties in model parameterisations stem from the knowledge gaps in processes controlling the aerosol budget, and cause differences in climate projections from different model set-ups. For example, the sea spray parameterisations used in the Gilgen et al. (2018) and Struthers et al. (2011) studies include empirical representations of the effect of temperature on the sea spray aerosol size distribution, while the parameterisation used in the Browse et al. (2014) study does not. This could be a cause of the discrepancy in their predictions of sea spray aerosol
105 response to sea ice loss, though this has not been studied. The studies also differ in their treatment of primary marine organic emissions, with only the Browse et al. (2012) study including such a source. Some field studies from the [Arctic high Arctic pack ice region](#) have indicated the importance of primary marine organics in the region ([Leck and Bigg, 2010; Karl et al., 2013](#))
[\(Leck and Bigg, 2010; Karl et al., 2013\)](#), as well as raising questions about possible recycling mechanisms of particles after they are emitted ([i.e. through ageing or the particle break-up theory, Lawler et al., 2021](#))
110 [\(i.e. through ageing or the particle break-up theory, Lawler et al., 2021\)](#). The open questions surrounding primary marine emissions complicate modelling of Arctic NPF due to the the sink of condensable vapours from larger particles. Models with size-resolved aerosol microphysics and chemistry are better equipped to study these questions than bulk, single-moment models considering total mass only.

115 Aitken mode particles can act as CCN in the Arctic, which increases the influence of NPF over the Arctic aerosol budget. Karlsson et al. (2022) used measurements of cloud residual particles (i.e. particles obtained by drying cloud droplets or ice crystals) from the high Arctic to show that Aitken mode particles were acting as CCN during a period of frequent boundary layer new particle formation. Observations of aerosol size distributions in and out of cloud in sub-Arctic Finland showed that on average 30% of Aitken particles (defined as 25-95 nm) were activated to form droplets (Komppula et al., 2005). In Svalbard, measurements of cloud residuals (Karlsson et al., 2021) and cloud drop number concentrations (Koike et al., 2019)
120 have been used to show the activation of particles smaller than 50 nm diameter during periods of high supersaturation, such as when updraft speeds are high or when accumulation mode concentrations are low. Results from parcel models and large-eddy simulation (LES) models are in agreement with the observations that Aitken particles can act as CCN (Pöhlker et al., 2021; Bulatovic et al., 2021), [though Bulatovic et al. \(2021\) find that two sets of aerosol conditions under which Aitken activation is favourable in their model have real-world occurrence probabilities of 5 and 17%, raising questions about how widespread the phenomenon could be](#). Activation of Aitken particles means that particles formed by NPF and subsequent growth may only

125 need to grow up 20-50 nm diameter to act as CCN in Arctic clouds, making it more plausible that such particles could survive long enough in the atmosphere to be important for cloud formation.


130 A further complicating factor that affects our understanding of summertime Arctic aerosol processes is the decoupling of the surface, where most aerosol measurements have been made in the high Arctic, and the cloud layer, where the aerosols act as CCN. Such decoupling is caused by the thermodynamic structure of the high Arctic summer boundary layer. Regimes of turbulence at the surface and in the cloud mixed layer have previously been shown to occur in configurations where the two regimes do not interact, inhibiting transport of moisture or particles vertically between the different layers (Shupe et al., 2013; Sotiropoulou et al., 2016). Decoupling of the surface in this way has implications for aerosol-cloud interactions because it implies that the aerosol sources, concentrations and size distributions at the surface may only be relevant for the cloud layer during sporadic events of mixing. This is very different from dynamics typical of lower latitudes, where strong convection can promote mixing of heat, moisture and aerosols from the surface up to higher altitudes. Models have struggled to capture decoupling in the Arctic, showing a tendency to become coupled too often (Birch et al., 2012; Sotiropoulou et al., 2016).

140 Recent observations from the high Arctic provide a new opportunity to explore the questions surrounding the source of summertime aerosol. The Arctic Ocean 2018 (AO2018) expedition took place in August and September 2018 between Svalbard and the North Pole (Vüllers et al., 2020) (Vüllers et al., 2020; Leck et al., 2020). The observed properties and behaviour of the aerosol challenge our current understanding of aerosol sources and sinks. Firstly, mass spectrometry measurements clearly show that iodine acid is the main driver of NPF events, which occur primarily during sea ice freeze-up (Baccarini et al., 2020a). This process was not included in previous large-scale modelling studies. Secondly, the long-time series of particle concentrations and size distributions, spanning several weeks over the end of the sea ice melt period and transition to freeze-up, provide an opportunity to understand how sources and sinks are related to the melting/freezing cycle and changes in photochemistry. 145 In particular, the observations show a distinct transition in aerosol behaviour in late summer, when the iodine acid NPF events begin to take place. Here we aim to interpret this change in behaviour in terms of changes in the dominant aerosol sources.

150 In this study we used measurements from AO2018 to evaluate the Arctic aerosol budget in the global climate model UKESM1. We compared the accuracy of simulations with NPF in the boundary layer to that of simulations with NPF in the free troposphere. We also introduced an iodine acid NPF scheme to the model to investigate iodine acid as an Arctic aerosol source compared to other components of the Arctic aerosol budget. The observational datasets are introduced in Sect. 2. The model is described in Sect. 3, including the different NPF schemes and our approach to the inclusion of iodine acid. Results are presented in Sect. 4 and discussed in Sect. 5.

2 Observations

155 We use data from three campaigns in different years: Arctic Ocean 2018 (AO2018), The Arctic Summer Cloud Ocean Study (ASCOS) in 2008, and the Atmospheric Tomography Mission (ATom) in 2016. We only use model output from a simulation of the year 2018 because the Arctic Ocean 2018 dataset is the main focus of this study. The interannual variability in aerosol concentration may limit how representative observations from 2008 (ASCOS) or 2016 (ATom 1) can be to assess model output



Images/fig01.pdf

Figure 1. Map showing location of where observations were measured. Co-ordinates on map are for *Oden* during AO2018 (blue line), aircraft during leg 1 of ATom (orange line) and helicopter during ASCOS (pink-purple line). Note only ATom 1 co-ordinates with a latitude greater than 60°N are shown here since we discarded data from further south in this study.

from 2018. However, as we will show in later sections, the difference in particle concentration from different simulations in this study can be several orders of magnitude and therefore is likely to exceed the range of concentrations that would be measured

160 Simulated values of N_{50} (number concentration of particles larger than 50 nm diameter) over a 30 period suggest that the interannual variability of N_{50} in the Arctic does not account for such large differences in our simulated particle concentrations (?).

2.1 Arctic Ocean 2018

The Arctic Ocean 2018 (AO2018) expedition took place in August and September 2018 aboard the Swedish icebreaker *Oden*.

165 The ship travelled from Svalbard to the North Pole and then drifted, moored to an ice floe, for 4 weeks before travelling back to Svalbard (Fig. 1). Further details and meteorological conditions of the campaign are presented in Vüllers et al. (2020) and Leck et al. (2020). Here, we compare model output to aerosol and gas-phase measurements, which were measured during AO2018 as part of the Microbiology-Ocean-Cloud Coupling in the High Arctic (MOCCHA) campaign.

We use two aerosol datasets from the AO2018 campaign, involving three different instruments (Baccarini et al., 2020a). A 170 differential mobility particle sizer (DMPS) measured particles in the size range 10–959 nm (Karlsson and Zieger, 2020). We integrate this size distribution over the ranges 15–100 nm and 100–500 nm. Total particle concentration for diameters greater than 2.5 nm was measured by an ultrafine condensation particle counter (UCPC). The UCPC and integrated DMPS data were

used to calculate the concentration of all particles in the size range 2.5–15 nm (Baccarini and Schmale, 2020). This 2.5–15 nm time series was also supplemented by a particle size magnifier during periods when the UCPC was not in operation. All 175 data has been selected for "clean" periods, i.e. excluding periods where the ship's exhaust might influence the measurements. In addition to the aerosol data, iodic acid concentrations were measured using a nitrate chemical ionization mass spectrometer (Baccarini et al., 2020b).

The measurements of nucleation mode particle concentration at the surface during AO2018 show marked difference in aerosol behaviour linked to the onset of local sea ice freeze-up. During the freeze period, peaks in the nucleation mode concentration occur during NPF events, lasting on the order of hours (see section 4.2). Thus the nucleation mode concentration is 180 on average higher in the freeze period than the melt period, and also fluctuates more.

2.2 Arctic Summer Cloud Ocean Study 2008

The Arctic Summer Cloud Ocean Study (ASCOS) campaign also took place on the icebreaker *Oden*, at roughly the same time of year as AO2018 but a decade earlier (2008). The ASCOS drift period was from 12th August until 2nd September 2008 and 185 took place close to 87°N. A full description of the campaign is given in Tjernström et al. (2014).

A helicopter was used during ASCOS to take measurements above the surface, with two condensation particle counters (CPCs) and one optical particle counter (OPC) used to measure aerosol concentrations ([Leck et al., 2022](#)). The CPCs and OPC detected particles larger than 3 nm, 14 nm, and 300 nm respectively, giving an overall aerosol size distribution in the ranges 3–14, 14–300 and >300 nm. These do not exactly match the aerosol size ranges available from the AO2018 data, but still give 190 us valuable information about the aerosol size distribution from this campaign. The aerosol measurements are presented in Kupiszewski et al. (2013).

2.3 Atmospheric Tomography Mission

The Atmospheric Tomography Mission (ATom) was a multi-year flight campaign that used the NASA DC-8 aircraft to study the effects of air pollution on the chemistry of the atmosphere. There were four legs of ATom, each carried out in a different season 195 in different years. Here, we use measurements taken during the first leg, which took place in summer 2016. Measurements were taken over a wide range of latitudes and altitudes. We only use measurements taken north of 60°N.

Aerosol size distributions were measured during ATom using a nucleation-mode aerosol size spectrometer (NMASS), an ultra-high-sensitivity aerosol size spectrometer (UHSAS) and a laser aerosol spectrometer (LAS). The full set of ATom aerosol measurements are presented in Brock et al. (2019). Aerosol size distributions are available for particles between approximately 200 3 nm and 3.5 μm diameters.

Note that the AO2018, ASCOS and ATom campaigns use instruments with different ways of sizing particles. The DMPS measures the electrical mobility diameter of particles. The CPCs and NMASS measure particles based on the critical diameter for droplet nucleation at the instrument's operating supersaturation. The OPC, UHSAS and LAS all rely on optical methods to measure particles, meaning they measure the optical equivalent diameter of particles. No attempt has been made to convert

205 from optical equivalent diameter to mobility diameter. Such a conversion would require information about the refractive indices of the particles measured by the optical instruments, which is not available.

3 Model description

We used the UK Earth System Model version 1 (UKESM1, Mulcahy et al., 2020; Sellar et al., 2019) in its atmosphere-only configuration, which uses output from a fully-coupled run of UKESM1 to prescribe sea surface temperatures (SSTs) and some 210 biogenic emissions, such as DMS, from the ocean. The dynamical core of the model is the UK Met Office Unified Model (UM) in global atmosphere configuration version 7.1 (GA7.1, Walters et al., 2019) with a horizontal resolution of 1.875° longitude by 1.25° latitude and 85 vertical levels. The vertical resolution is approximately 50 m at the surface, 150 m at 1 km altitude and on the order of km at the highest model level. We ran the model in its "nudged" configuration, which means horizontal winds and potential temperature are relaxed to ERA-Interim values on a 6-hourly time scale above approximately 1 km. The model 215 simulates gas and aerosol chemistry using the UK chemistry and aerosols model (UKCA, Morgenstern et al., 2009; O'Connor et al., 2014). Aerosol microphysics is simulated by the Global Model of Aerosol Processes (GLOMAP-Mode, Mann et al., 2010) using 5 log-normal aerosol size modes (4 soluble and 1 insoluble). [Mode sizes and geometric widths are given in Table 1](#). The model uses 2-moment aerosol microphysics, meaning that the number and mass in each mode are prognostic variables. There are four aerosol species: sulphate, organic carbon, black carbon and sea salt. Processes handled by GLOMAP include 220 primary emissions, coagulation within and between modes, condensational growth and ageing, new particle formation, dry deposition, wet deposition within and below clouds, and aqueous sulphate production in cloud droplets.

Emissions of SO₂, black carbon (BC) and organic carbon (OC) are for the year 2014 for all emissions sectors except 225 biomass burning, which are from a climatology of the years 1995-2004. We use a climatology for biomass burning to minimise any potential bias from boreal forest fire emissions not being from the same year as the simulation time. The emissions datasets used for aerosols and precursor vapours are Hoesly et al. (2018) (SO₂, anthropogenic OC and BC), Marle et al. (2017) (biomass burning OC and BC) and Sindelarova et al. (2014) (monoterpenes). Sea salt emissions are calculated using the Gong (2003) parameterisation. Primary marine organic emissions are then calculated using the sea spray flux, 10 m wind speed and chlorophyll-a concentration, using the Gantt et al. (2015) parameterisation. Note that because we use an atmosphere-only configuration in this study, the chlorophyll-a concentration is taken from an ancillary file, produced using model output from 230 the fully-coupled model. Marine emissions are scaled by grid box open water fraction for sea ice regions.

3.1 New particle formation schemes

UKESM includes binary homogeneous nucleation of water and sulphuric acid (H₂SO₄) using the parameterisation of Vehkamäki et al. (2002). [In this study, we included two additional NPF formation schemes in the model. We](#) [The Vehkamäki et al. \(2002\) scheme creates particles mostly in the cold free and upper troposphere. Here, we run simulations with NPF schemes that also](#) [create particles in the boundary layer \(BL\). We](#) [do this so that we can compare NPF driven by iodic acid, observed during AO2018, to other nucleation mechanisms known to be important in the Arctic and extra-Arctic. We](#) simulated nucleation of

Table 1. Description of aerosol modes and parameterisations in the model. Y/N=yes/no i.e. if Y, this species is allowed to exist in this mode. J96=?.

| | Mode | | | | | Water uptake |
|----------------------------------|--|--------|--------------|----------|-----------|-----------------------------|
| | Nucleation | Aitken | Accumulation | Coarse | Insoluble | |
| <u>Size (nm)</u> | 1–10 | 10–100 | 100–500 | 500–1000 | 10–100 | |
| <u>Width</u> | 1.59 | 1.59 | 1.4 | 2 | 1.59 | |
| <u>Sulphate</u> | Y | Y | Y | Y | Y | As for SO_4 in J96 |
| <u>Organic carbon</u> | Y | Y | Y | Y | Y | 65% that of sulphate |
| <u>Black carbon</u> | N | Y | Y | Y | Y | None |
| <u>Sea salt</u> | N | N | Y | Y | N | As for Cl in J96 |
| <u>Hygroscopic growth</u> | Following ZSR theory as described in Mann et al. (2010). | | | | | |
| <u>Aerosol activation scheme</u> | Abdul-Razzak and Ghan (2000) | | | | | |

H_2SO_4 in the boundary layer (BL) BL by cluster activation as described by Kulmala et al. (2006) (hereafter K06). We simulated organically mediated H_2SO_4 nucleation using the parameterisation of Metzger et al. (2010) (hereafter M10). Simulations are labelled by their NPF scheme and the region of. We refer to simulations by the atmosphere the scheme is applied to (e.g. BL for a scheme switched on in the boundary layer only or FT for a scheme switched on in the free troposphere only) precursor vapour used to drive new particle formation in each simulation, i.e. SA for the use of sulphuric acid in the Kulmala et al. (2006) scheme, SOA for secondary organic vapours in Metzger et al. (2010), or IA for iodic acid.

The formation rates of 1.5 nm clusters, J_* , used by M10 and K06 are given in $\text{cm}^{-3} \text{ s}^{-1}$ by

$$J_{*,K06} = k_{K06} C_{SA} \quad (1)$$

$$245 \quad J_{*,M10} = k_{M10} C_{SA} C_{secorg SOA} \quad (2)$$

where k_{K06} is 10^{-6} s^{-1} , k_{M10} is $10^{-13} \text{ cm}^3 \text{ s}^{-1}$, and C_x are the concentrations of the precursor vapours in cm^{-3} . The 3 nm particle formation rate is calculated from the 1.5 nm cluster formation rates using the method from Kerminen and Kulmala (2002), which accounts for growth of clusters and loss to existing particles.

It should be noted that these two BL NPF schemes use empirical parameterisations that have not been developed with or 250 tested against data from the Arctic. Moreover, the parameterisations only consider the influence of H_2SO_4 and secondary organic vapours. MSA and ammonia are not included in UKESM and thus are not modelled as NPF precursors in this study. These omissions are potential sources of bias given the importance of MSA and ammonia in the Arctic (Willis et al., 2018; Beck et al., 2020).

As we show below, entrainment of nucleation mode aerosol from the FT is an important process a key surface aerosol source in the central Arctic in UKESM. To investigate the role of FT entrainment as a source of surface particles in the high Arctic, we ran an additional sensitivity test with an imposed NPF rate in the low FT (simulation M10_PrcSOA_PRSC). A particle

formation rate of 10^{-2} ~~particles~~ $\text{cm}^{-3} \text{ s}^{-1}$ was used for altitudes above the top of the BL and below 7.5 km, and for latitudes north of 80°N . Since this rate is higher than what the ~~M10~~ [Metzger et al. \(2010\)](#) scheme typically produces, it allows us to investigate the sensitivity of the surface concentrations to a strong free troposphere source.

260 **3.2 Iodic acid**

An empirical model for the steady-state concentration of iodic acid has been produced from observations taken in the high Arctic (Baccarini et al., 2020a). The concentration in cm^{-3} , C_{IA} , is given by

$$C_{IA} = \frac{E}{v_{dep} + h \cdot CS} \quad (3)$$

265 where E is the emission rate of iodine atoms in $\text{cm}^{-2} \text{ s}^{-1}$, v_{dep} is the dry deposition velocity of iodic acid in cm s^{-1} , h is the surface mixed layer height in cm and CS is the condensation sink due to ~~existing aerosols~~ [all aerosols \(including the nucleation mode formed through this process\)](#), given in s^{-1} . [E can be considered a net emission rate, which also accounts for the conversion of iodine into HIO₃ \(the rate is assumed to be constant\). This parameterisation does not include the HIO₃ chemical formation mechanism described in Finkenzeller et al. \(2022\). Still, it can reproduce the observed iodic acid concentration, as shown in Figure 2 and already reported in Baccarini et al. \(2020a\). Hence, it serves well the scope of this work.](#) It was observed 270 during AO2018 that fog and cloud droplets acted as a strong sink of the iodic acid, however we do not account for that here since the parameterisation of clouds in the coarse resolution model gridboxes is unlikely to be representative of conditions at the ship.

We used Eq. (3) to diagnose the steady-state concentration of iodic acid in model gridboxes at each timestep. The iodic acid 275 was then used as a precursor to drive NPF, [a process which also depletes the iodic acid gas concentration](#). The dry deposition velocity of iodic acid was calculated by the model assuming a diffusion constant equal to that of H₂SO₄. The condensation sink due to existing aerosols was calculated during model runs using the aerosol size distribution produced by the model. We use the modelled dry aerosol diameter in the condensation sink calculation, meaning that the effects of water uptake by aerosol are not included. For the surface mixed layer height, we take the BL height calculated by the UM. Finally, we use a value of 5.21 $\times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$ for E , which is approximately equal to the median of the distribution of E measured by Baccarini et al. 280 (2020a).

The concentration of iodic acid was observed to increase towards the end of summer, during the sea ice freeze-up (Baccarini et al., 2020a). It is thought that the freezing of sea water can trigger the emission of iodine from the surface. To incorporate this behaviour into UKESM, we calculate the concentration of iodic acid in gridboxes where sea ice fraction is non-zero and where the surface temperature is less than -5°C . Although the observed freeze-up date is defined as the day when the 14-day running 285 mean of surface temperature reaches -2°C (the temperature at which sea water freezes), we found that -5°C acted as a better threshold for ice freeze-up in the model. This is because there is a cold bias in the model such that when the observed 14-day running mean temperature reaches -2°C , the modelled surface temperature is closer to -5°C (see Fig. A1). The calculated

concentration of iodic acid is then equally distributed from the surface the BL height. This is equivalent to assuming that the lifetime of iodic acid is long enough for the gas to be mixed throughout the BL.

290 It is important to note that a direct/causal mechanism linking the freeze-up to enhanced iodine emissions has not yet been identified. However, the results of our study would remain valid even if the two processes were not directly related. In fact, the surface temperature threshold used in the model is a good tracer for the summer to autumn transition, which has been associated with higher iodine concentration in the Arctic (Baccarini et al., 2020a; Sharma et al., 2019)

295 The particle formation rate at 3 nm from iodic acid is calculated using the method from Kerminen and Kulmala (2002) as for the H_2SO_4 activation and organically mediated schemes, using a kinetic rate of cluster formation from iodic acid. This is in line with recent results from cloud chamber experiments (He et al., 2021). The cluster formation rate in $\text{cm}^{-3} \text{ s}^{-1}$ is given by

$$J_{*,IA} = k_{IA} C_{IA}^2 \quad (4)$$

where $k_{IA} = 10^{-13} \text{ cm}^3 \text{ s}^{-1}$. The mass created from HIO_3 -driven new particle formation is added to the sulphate model component.

300 3.3 Secondary organic vapours

Assumptions about the production of secondary organic aerosol (**SOA**) material have a large effect on modelled Arctic aerosol. **SOA** Secondary organic aerosol is created in the model by the oxidation of monoterpenes by ozone, the hydroxyl radical and the nitrate radical (Spracklen et al., 2006). These reactions produce **SOA** secondary organic aerosol on the timescale of hours. This is a simplification of the process, since in reality there are many chemical species and reaction pathways involved in the 305 production of **SOA** secondary organics, which operate on a range of timescales and produce a range of volatilities. In particular, some **SOA** organic aerosol precursor species have a longer lifetime than monoterpenes and can therefore be transported further in the atmosphere before forming **SOA** aerosol and condensing on existing aerosol. In the model, most monoterpenes are oxidized close to the source region (e.g. boreal forests) and quickly condense onto existing particles, therefore the concentration of **SOA** organic aerosol precursor gases is very low in the central Arctic; as a consequence, NPF involving organic vapours is extremely 310 weak. To investigate the effect of these assumptions on Arctic NPF, we ran sensitivity simulations where the oxidation rate of monoterpenes was reduced by a factor of 100. The labels of these simulations use the suffix _SeeOrgOXID.

We alter the oxidation rate of monoterpenes to promote transport of monoterpenes north, to account for missing species and reactions that create organic aerosol precursors. This approach allows us to test the effect of neglected organic species with oxidation rates different from monoterpenes.

315 3.4 Ageing of insoluble particles

The default assumptions in UKESM about particle ageing have a very substantial effect on Arctic aerosol, particularly where NPF is a major source of Aitken mode particles. Particle ageing in a model is the transfer of insoluble particles into the soluble particle modes after condensation of water-soluble material. In UKESM, sources of insoluble carbonaceous particles

Table 2. Description of simulations.

| Simulation | Description |
|----------------------------|--|
| Main body | |
| CONTROL | Model set-up based on UKESM1 atmosphere-only configuration. Binary H_2SO_4 -water vapour NPF parameterised as per Vehkamäki et al. (2002). |
| K06_BL_SOA | Additional BL NPF parameterised by Kulmala et al. (2006) M10_ALL Organically-mediated NPF in all model levels (driven by secondary organic aerosol precursors) in the BL , using Metzger et al. (2010) |
| M10_BL_SOA_PRSC | Organically-mediated NPF in the BL all model levels , using Metzger et al. (2010) M10_Prcs_AS M10_ALL, but with, and a prescribed NPF rate of $10^{-2} \text{ cm}^{-3} \text{ s}^{-1}$ between the BL top and 7.5 km and north of 80°N |
| IA_BL | Additional BL NPF driven by IA |
| IA_BL_M10_ALL_SOA | As M10_All but Organically-mediated NPF in all model levels, using Metzger et al. (2010) , with additional BL NPF driven by IA |
| M10_BL_SOA_85N | Additional NPF in the BL using Metzger et al. (2010) for gridboxes north of 85°N only |
| XXX_SeeOrg | |
| Appendices | |
| SA | Additional BL H_2SO_4 NPF parameterised by Kulmala et al. (2006) |
| SOA_ALL_LEVELS | Organically-mediated NPF in all model levels, using Metzger et al. (2010) |
| XXX_OXID | Additional change to oxidation of monoterpenes as described in section 3.3 |

are biofuel and biomass burning emissions (mean diameter 150 nm), fossil fuel burning emissions (mean diameter 60 nm) and primary marine organic carbon emissions (mean diameter 160 nm). By default, the aged mass from the insoluble mode is moved into the soluble Aitken mode (Mulcahy et al., 2020). However, the mean diameter of the insoluble mode does not usually correspond to the size limits of the soluble Aitken mode, and this can lead to undesirable behaviour in the model. Thus, when mass from the insoluble mode is moved into the soluble Aitken mode, it will typically increase the mean diameter of the soluble mode beyond its upper limit (100 nm). When small particles enter the Aitken mode following growth of nucleation mode particles, they are artificially strongly depleted because "mode merging" (Mann et al., 2010) requires that their mass is averaged with the larger particles already existing in the Aitken mode. This combination of assumptions (the size of aged particles and mode merging) is adequate for reproducing size distributions in the mid-latitudes, where anthropogenic and fire emissions are a more dominant source. However, our early simulations showed that this method has a very substantial effect in the Arctic where NPF is occurring in air that has aged during long-range transport from low latitudes to the Arctic (see appendix C). In particular, we found that the particle size distribution was extremely insensitive to the NPF rate. We therefore altered the model such that aged carbonaceous particles are moved directly into the soluble accumulation mode (100 – 500 nm), as is appropriate for their diameter. Our CONTROL simulation uses this altered ageing scheme. Simulations using the model's default ageing scheme are presented in appendix C.

A description of all model simulations is given in Table 2.

We organise the results as follows. First we examine the behaviour of our empirical iodic acid scheme by comparing modelled surface iodic acid concentrations to observations in Sect. 4.1. Then in Sect. 4.2 we consider the effect of ~~different BL NPF schemes~~ ~~BL NPF~~ in the model ~~(including iodic acid)~~ on the surface aerosol concentration during AO2018. Since we show that BL NPF is not taking place locally during the melt period of AO2018, we consider FT NPF as a source of surface particles in Sect. 4.3, using surface observations from AO2018 and aerosol profiles from ASCOS in 2008 and ATom in 2016. Finally, we consider a combination of ~~local (BL) and non-local (FT) NPF~~ ~~BL and FT sources~~ in Sect. 4.4.

When comparing model output to observations of surface aerosol concentration, we use the overlap index defined in Pastore and Calcagnì (2019) to quantify the similarity between the distribution in aerosol concentration from observations and different model simulations. ~~An overlap index of 0 indicates no overlap between the distributions, while an overlap index of 1 indicates that the two probability density functions $A(x)$ and $B(x)$ are identical.~~ For two probability density functions $A(x)$ and $B(x)$, the overlap index $\eta(A, B)$ is defined as

$$\eta(A, B) = \int \min[A(x), B(x)] dx \quad (5)$$

~~which can be thought of as integrating the area where the distributions overlap. If the two distributions overlap completely, the entire area under the distribution will be integrated so the index will be 1. Whereas if they do not overlap at all, no area will be integrated and the index will be 0.~~ Overlap indices closer to 1 therefore indicate simulations with good model-observation agreement in terms of the magnitude and variability of the two time series even if individual peaks and troughs do not match temporally. ~~In our case, we apply the overlap index to discrete distributions of binned aerosol concentration, so the integral becomes a sum.~~

4.1 Iodic acid concentration

Figure 2 shows a time series and PDF of surface HIO_3 concentration from observations and model output. To collocate the model output with the ship, we take the model gridbox nearest the ship's position. In the observations, the surface HIO_3 concentration is lower in the melt period (before day 239) than in the freeze period (after day 239). The surface concentration in the freeze period has a baseline of approximately 10^6 cm^{-3} and peak values 5–6 times higher lasting on the order of hours, whereas in the melt period the concentration reaches 10^6 cm^{-3} only for brief periods, such as on day 228. The periods of peak surface concentration in the freeze period correspond to observed NPF events.

In the model, HIO_3 is only emitted when the surface temperature is -5°C or less. The modelled temperature reaches this threshold on approximately day 236 at the ship's position, such that the model starts to emit HIO_3 at a similar time to the observed freeze-up onset (day 239), when HIO_3 concentrations were observed to increase at the ship. After HIO_3 starts to be emitted in the model, the empirical scheme we use for HIO_3 production consistently calculates surface HIO_3 concentration to be the same order of magnitude as the observations. The PDF in Fig. 2(b) is for the freeze season only, i.e. day 239 onward. The observed distribution of surface HIO_3 concentration is broadly captured, though the model does not reproduce the highest or lowest observed concentrations. This is likely to be because the model does not resolve the spatially heterogeneous sea ice

Images/fig02.pdf

Figure 2. Times series and PDF of surface iodic acid concentration during AO2018 from observations and model output. Model output is from simulation IA BL (orange lines). Observations are shown as 3-hourly mean (black lines). Red dashed lines in (a) show periods of observed NPF events. PDF of concentration is for freeze season only, i.e. after 27th August 2018 (day 239).

state (which controls the emissions of iodic acid) nor the variability in clouds and fog, which control iodic acid scavenging as well as influencing the condensation sink due to existing aerosols.

Table 3. Overlap indices calculated for the PDFs of nucleation, Aitken and accumulation aerosol concentrations from observations and each simulation, measured at the surface. PDFs are separated into the melt and freeze periods before overlap indices are calculated. Bold text indicates the greatest overlap index in each mode for the melt and freeze periods.

| Simulation | Nucleation | | Aitken | | Accumulation | |
|---------------------------|----------------------------------|--|-------------|-------------|--------------|-------------|
| | Melt | Freeze | Melt | Freeze | Melt | Freeze |
| CONTROL | 0.01 | 0.00 | 0.31 | 0.12 | 0.57 | 0.64 |
| K06_BL_SOA | 0.73 | M10_ALL M10_ALL_SeeOrg M10_BL 0.05 | 0.73 | 0.34 | 0.50 | 0.55 |
| M10_BL_SOA_85N | 0.01 | 0.00 | 0.33 | 0.13 | 0.56 | 0.67 |
| M10_Prcs_SOA_PRSC | 0.35 | 0.52 | 0.75 | 0.37 | 0.51 | 0.55 |
| M10_Prcs_SeeOrg_IA | IA_BL 0.25 | 0.47 | 0.35 | 0.70 | 0.58 | 0.58 |
| IA_BL_SeeOrg_SOA | IA_BL_M10_AL 0.51 0.63 | | 0.69 | 0.77 | 0.55 | 0.56 |

4.2 Effect of NPF in the boundary layer

370 4.2.1 Time series of particle concentrations

Figure 3 shows time series and PDFs of surface aerosol concentration in three particle size ranges from the AO2018 observations and for simulations CONTROL, **K06_BL**, **M10_BL**, **IA_BL** and **IA_BL_SeeOrg** **SOA** and **IA**. Observations are from the ship (approximately 15 m above the ground) and model output is for the first model level, between approximately 0 and 37 m.

375 A time series of the measured nucleation mode (2.5–15 nm diameter) particle concentration during AO2018 is shown in Fig. 3 (a). Periods where iodic acid NPF events were observed are marked with red dashed lines. There is a marked difference in the behaviour of the nucleation mode concentration before and after the onset of sea ice freeze-up (27th August 2018, day 239). In the melt period (up to day 239), the nucleation particle concentration rarely exceeds 100 cm^{-3} and is usually between $1\text{--}10 \text{ cm}^{-3}$. The NPF events, which occurred after the freeze-up began in the vicinity of the ship on day 239, are associated with peaks in the nucleation mode particle concentration lasting a few hours, causing fluctuations in particle concentration between 380 approximately 10 and 10^4 cm^{-3} , consistent with a strong local source.

The modelled nucleation mode particle concentration is underestimated in simulations CONTROL, **IA_BL** and **IA_BL_SeeOrg** **and IA** in the melt period, while the inclusion of HIO_3 NPF means that **IA_BL** and **IA_BL_SeeOrg** **performs** **performs** better in the freeze period than during the melt. The nucleation mode concentration in CONTROL varies from roughly 10^{-6} to 10^{-1} cm^{-3} and is usually at least an order of magnitude lower than observed throughout the whole period. In contrast, 385 simulations with **(IA_BL and IA_BL_SeeOrg)** **produce** **IA produces** nucleation mode concentrations of roughly the correct

Images/fig03.pdf

Figure 3. Times series and PDFs of surface aerosol concentration during AO2018 from observations and model simulations with various boundary layer NPF mechanisms. Model output is from simulations CONTROL (pink lines), **K06_BL_SOA** (grey lines), **M10_BL** (blue lines), **IA_BL** (thick orange lines) and **IA_BL_SeeOrg** (thin orange lines). Observations are shown as 3-hourly mean (black lines) and standard deviation (grey shading). Aerosol concentrations are shown for particles with diameter (a-c) 2.5-15 nm, (d-f) 15-100 nm and (g-i) 100-500 nm. Red dashed lines in (a, d, g) show observed NPF events. PDFs are separated by observed sea ice freeze-up date, 27th August 2018 (day 239).

order of magnitude during the freeze period (day 239 onwards), with a distinct change in concentration and behaviour around day 235. However, the lack of HIO_3 emissions during the melt period means that the model continues to underestimate the concentration by several orders of magnitude. The model underprediction of nucleation mode particle concentration in the melt period is an indication that HIO_3 NPF is not the only part of the regional aerosol budget that needs consideration to produce an accurate simulation. ~~Simulations K06_BL and M10_BL both use BL NPF schemes and have Simulation SOA uses a BL NPF scheme and has~~ higher nucleation mode concentrations than CONTROL as a result. In the melt period, ~~K06_BL and M10_BL both consistently simulate SOA consistently simulates~~ nucleation mode concentrations of the same order of magnitude as the observations. ~~Interestingly, the two simulations have very similar nucleation mode concentrations despite the differences in the NPF schemes. K06_BL occasionally simulates higher nucleation mode concentrations than M10_BL, most notably on days 239–243.~~ In the freeze period, ~~K06_BL and M10_BL still underestimate SOA still underestimates~~ the nucleation mode concentration despite simulating higher concentrations than CONTROL.

Overlap indices for the PDFs of observed surface aerosol concentration and modelled concentration from all simulations are given in Table 3. The underestimation of nucleation mode concentration by simulation CONTROL is highlighted by the fact that its overlap indices are close to 0 in both the melt and the freeze periods. The ~~K06_BL and M10_BL SOA~~ PDFs of nucleation mode concentration for the melt period match the observations ~~well, producing overlap indices better than CONTROL, producing an overlap index~~ of 0.73 ~~for M10_BL and for K06_BL, which is the highest value achieved for the nucleation mode in the melt period. In the~~ ~~However, in the~~ freeze period, the overlap ~~indices are also low (index is low at 0.05 for M10_BL and for K06_BL).~~ IA performs moderately better than CONTROL in both periods, giving overlap indices of 0.25 ~~in the melt period and 0.47 in the freeze period.~~

Observed Aitken mode concentrations (diameter 15–100 nm) lie between about 1 and 1000 cm^{-3} , with a mean of 60 cm^{-3} over the whole period. In contrast, concentrations in the CONTROL simulation are around 1 cm^{-3} , frequently fall below 0.1 cm^{-3} , and never exceed 10 cm^{-3} . The ~~simulations simulation~~ with HIO_3 again ~~show shows~~ a sharp increase in particle concentration around day 235 when HIO_3 starts being emitted during the freeze period. Model Aitken mode concentrations then vary between being comparable to the observations and being 1–2 orders of magnitude too low. ~~The Aitken mode concentration in simulation IA_BL_SeeOrg behaves similarly to that of IA_BL, although there are some periods where the concentration in IA_BL_SeeOrg exceeds those in the IA_BL simulation by a factor of more than 10 and is in better agreement with observed concentrations (notably days 244–247). This is likely because of the extra condensable vapour in IA_BL_SeeOrg, which promotes more particle growth from the nucleation mode.~~ As in the nucleation mode, the Aitken mode concentrations from ~~K06_BL and M10_BL SOA~~ are higher than CONTROL, on the same order of magnitude as the observations in the melt period but underestimating observations in the freeze period.

The observed accumulation mode concentrations (diameter 100–500 nm) are typically around $10\text{--}100 \text{ cm}^{-3}$, with brief periods of less than a day where they fall to 1 cm^{-3} or lower, which is characteristic of the central Arctic (Bigg et al., 1996; Bigg and Leck, 2001; Mauritzen et al., 2011; Leck and Svensson, 2015). All simulations capture the observations well, except for the periods of extremely low concentration. The good agreement shows that NPF is not an important source of accumulation mode aerosol. As shown in e.g. Stevens et al. (2018) and Loewe et al. (2017), the periods of very low concentration are associated

with efficient scavenging in drizzle on smaller spatial scales than represented in a global model. Nevertheless, the generally good model-observation agreement means that the aerosol surface area, and hence condensation sink for nucleating vapours, is reasonable in the model, and therefore not a cause of the biases in nucleation and Aitken mode particle concentrations.

4.2.2 Aerosol vertical profiles

425 In Sect. 4.2.1, we showed that the use of ~~BL-NPF schemes~~ (organically-mediated BL NPF in the model (simulation SOA) instead of only the default ~~upper tropospheric scheme~~ NPF scheme (simulation CONTROL) increased nucleation and Aitken particle concentrations at the surface during the melt period of AO2018. In this section, we examine the effect of ~~these schemes~~ ~~this scheme~~ on particle concentrations aloft, ~~and show that switching on BL NPF increases particle concentrations in the Arctic FT, perhaps counter-intuitively.~~

430 Figure 4 shows simulated nucleation mode aerosol vertical profiles for the AO2018 campaign period. As discussed in Sect. 4.2.1, the nucleation mode concentrations at the surface in CONTROL are typically less than 0.1 cm^{-3} and never above 1 cm^{-3} . We showed that this is an underestimation of the measurements taken at the ship. Figure 4 shows that despite the low concentrations at the surface, the model produces nucleation mode concentrations of up to 1000 cm^{-3} in the low FT, for example on day 244 above approximately 4 km. These higher concentrations aloft suggest that NPF is being simulated in the 435 FT, but that those particles do not reach the surface.

440 The vertical profile for simulation ~~M10_BL_SOA~~ shows that the nucleation mode concentration is higher than CONTROL above the BL as well as at the surface, even though this simulation is using the same NPF mechanism as CONTROL in the FT. We previously showed that ~~M10_BL_SOA~~ had nucleation mode concentrations at the surface that were 1–3 orders of magnitude higher than that of CONTROL. In the FT, the increase is up to an order of magnitude. The white lines in Fig. 4 show the ~~modelled BL height~~ ~~top of the BL in the model~~, which is used by the NPF schemes to separate the BL from the FT. The increased nucleation mode concentrations in ~~M10_BL_SOA~~ above this BL height suggest that particles are being created in the BL at lower latitudes and then transported north in the FT. To examine this effect further, we ran a simulation where the ~~M10~~ organically-mediated NPF scheme was used in the BL only for latitudes north of 85° ~~(as opposed to globally, as it is in SOA)~~. The nucleation mode vertical ~~profile~~ ~~profiles~~ for this simulation, ~~M10_BL_SOA~~ 85N, ~~is~~ ~~are~~ also shown in Fig. 4 and 445 ~~shows~~ ~~show~~ lower concentrations than ~~M10_BL_SOA~~ in both the BL and the FT. In fact, the nucleation mode concentration in ~~M10_BL_SOA~~ 85N is mostly the same as that of CONTROL. The output from ~~M10_BL_SOA~~ 85N therefore shows that the higher concentrations in ~~M10_BL_SOA~~ are from latitudes south of 85° . The concentration of organic vapours north of 85°N is much lower than at lower latitudes (see the steep latitudinal gradient in Fig. E4), which potentially accounts for the low particle concentrations in ~~M10_BL_SOA~~ 85N. Note that the smaller spatial extent of the NPF scheme could lead to less 450 depletion of precursors than when the scheme is used globally, thereby increasing the NPF rate in ~~M10_BL_SOA~~ 85N relative to ~~M10_BL_SOA~~. However, this non-linear behaviour would have the opposite effect to what we describe here so we can disregard it.

Images/fig04.pdf

Figure 4. Nucleation mode aerosol profiles simulated for AO2018 campaign period. Model output is from simulations (a) CONTROL, (b) **M10_BL_SOA** and (c) **M10_SOA_BL_85N**. Model output was colocated with the position of the *Oden*. White lines show BL height.

4.3 Effect of free-tropospheric NPF

In Sect. 4.2 we showed that the model default NPF scheme is insufficient to produce an accurate simulation of the aerosol
455 concentrations measured at the surface during AO2018. The inclusion of iodic acid NPF improves the concentration in the

nucleation and Aitken modes in the freeze period. Nevertheless, substantial model underestimations in particle concentration remain during the melt period, indicating some other missing source. ~~Simulations~~ A simulation with BL NPF driven by ~~or a combination of and~~ secondary organic vapours ~~produce~~ (simulation SOA) produced more accurate surface concentrations in the melt period of AO2018 compared to observations. However, as we showed in Sect. 4.2.2, ~~these simulations do using~~

460 ~~output from simulation SOA_85N, the SOA simulation does~~ not produce more NPF in the BL in the high Arctic. Rather, small particles produced at lower latitudes are transported north, increasing nucleation and Aitken concentrations aloft as well as at the surface. This behaviour in the model ~~suggests~~ ~~raises the question~~ that the FT could be a source of particles to the surface, ~~either from particles created by NPF aloft, or in the BL at lower latitudes before being transported into the high Arctic FT~~. Such a source has been considered before, for example by Korhonen et al. (2008a), who showed that entrainment of secondary 465 particles from the FT is an important source of CCN over the Southern Ocean in GLOMAP, and ~~?Igel et al. (2017)~~ who used a high resolution LES model to show that particles can be transported from the free troposphere to the surface under conditions typical of the high Arctic. Also of relevance to entrainment processes in Arctic clouds, Solomon et al. (2011) used an LES model to show that a humidity inversion and entrainment of water vapour at cloud top helps to maintain the cloud by supplying moisture.

470 In this section we explore the role of the FT as a source of particles at the surface in both periods. We ran a simulation with ~~M10~~ ~~the Metzger et al. (2010) organically-mediated NPF scheme~~ switched on at all model levels ~~. We also ran a simulation and~~ with a fixed NPF rate for model levels between the top of the BL and 7.5 km ~~.(simulation SOA_PRSC)~~. The use of an ~~idealised, constant NPF rate above the top of the BL in SOA_PRSC tests the sensitivity of the surface aerosol concentration to a source of aerosols from the FT. We use the output from SOA_PRSC to examine whether particles from the FT are being~~ 475 ~~entrained into the BL in the model.~~

4.3.1 Aerosol vertical profiles

Figure 5 compares simulated aerosol vertical profiles against observations from ASCOS (see Sect. 2.2). Model particle concentrations in these size ranges were calculated using August 2018 monthly mean data, colocated with the ASCOS flights which took place in August 2008. Concentrations of 3–14 nm diameter particles were consistently in the range 10^2 – 10^3 cm $^{-3}$ in the 480 FT, with lower values recorded below 1 km. The CONTROL simulation fails to capture these concentrations, underestimating the ASCOS mean concentration throughout the profile by 2 orders of magnitude in the FT and 3 orders of magnitude at the surface. Inclusion of organic ~~NPF (M10_ALL_BL_NPF (SOA))~~ substantially increases particle concentrations, but they remain at least 1 order of magnitude lower than observed. ~~Even when we allow much higher concentrations of organic vapours to be transported into the Arctic (M10_ALL_SeeOrg), 3–14 nm diameter particle concentration is still underestimated by up to 1 order of magnitude throughout the profile.~~ The model captures the 14–300 nm concentration better than the smaller particles. This is consistent with the results from the AO2018 comparisons, where the accumulation mode was captured much better than the nucleation or Aitken modes. All simulations shown here have 14–300 nm diameter particle concentrations of the same 485 order as those measured in the ASCOS flights.

Images/fig05.pdf

Figure 5. Aerosol vertical profiles from model output and ASCOS campaign observations from 2008. Model output is from colocated 2018 monthly mean values from simulations CONTROL (pink), **M10_ALL_SOA** (thick orange blue lines), **M10_ALL_SeeOrg** (thin orange lines) and **M10_SOA_Prcs-PRSC** (dark green lines). Observed values are given as the mean profile from each ASCOS flight (grey lines), overall mean (solid black) and overall median (dashed black). Profiles are for particles with size (a) 3-14 nm, measured during ASCOS using a UCPC and (b) 14-300 nm, measured using a CPC (particles greater than 14 nm) and a CLASP instrument (particles greater than 300 nm).

Images/fig06.pdf

Figure 6. Aerosol vertical profiles from model output and ATom campaign observations from 2016. Model output is from colocated 2018 monthly mean values from simulations CONTROL (pink), M10_ALL_SOA (thick orange blue lines), M10_ALL_SeeOrg (thin orange lines) and M10_SOA_Prcs_PRSC (dark green lines). ATom observations are taken from leg 1 of the campaign and restricted to measurements that were taken north of 60°N. Observations correspond to mean profiles from different days (grey lines), the overall mean (black solid lines) and overall median (black dashed lines). Profiles are for particles with size (a) 5–10 nm, (b) 10–100 nm and (c) 100–500 nm. Observations were recorded at standard temperature and pressure, model output has been adjusted to account for this.

Aerosol vertical profiles measured in 2016 during the ATom campaign are shown in Fig. 6 with model output for the 490 year 2018. Consistent with the ASCOS and AO2018 datasets, the CONTROL simulation predicts 5–10 nm diameter particle concentrations up to 3 orders of magnitude lower than the ATom measurements in the lowest 5 km of the atmosphere, with the largest model-observations discrepancies at the surface. In contrast to ASCOS, the ~~M10_ALL_SOA~~ simulations captures the nucleation mode profile better, producing concentrations that are within the range of the ATom measurements. The Aitken mode is underestimated by all simulations, though simulation ~~M10_ALL_SOA~~ is closer to the observations than CONTROL in 495 the lowest part of the FT. All simulations capture the accumulation mode well at the surface but underestimate particle number aloft by roughly 50%.

~~M10_ALL underpredicts the profile of SOA underpredicts the nucleation mode aerosol concentrations measured during the ASCOS campaign, even after perturbing the model monoterpenes to increase the availability of precursors in the Arctic (M10_ALL_SeeOrg) and ATom campaigns. However, SOA simulates higher nucleation mode concentrations than CONTROL.~~ 500 ~~As shown in section 4.2.2, the higher concentrations in the central Arctic in SOA occur as a result of transport of particles from further south, and concentrations are also increased in the FT by this transport. We therefore artificially tested what NPF rate would be required to explain these observed profiles. We tested various prescribed nucleation rates applied above the boundary layer. A rate of 10^{-2} s^{-1} produced a nucleation mode profile in best agreement with ASCOS and ATom (M10_Prse). tested the sensitivity of aerosol concentrations at the surface to a strong source of particles in the FT in simulation SOA_PRSC, to 505 see if the particles transported into the FT could be entrained into the Arctic BL.~~

In the ASCOS region (the central Arctic), ~~M10_Prse SOA_PRSC~~ is the only simulation to produce 3–14 nm particle concentrations close to the observed profiles (Fig. 5). ~~The nucleation mode concentration in SOA_PRSC is approximately an order of magnitude greater than that of SOA, including at the surface, suggesting that FT entrainment of secondary particles is taking place in the high Arctic in the model.~~ In the ATom region over continental North America, ~~M10_ALL and M10_Prse both behave the same~~ nucleation mode concentrations from SOA_PRSC are greater than that of SOA in most of the FT, with the greatest difference being nearly an order of magnitude at 4 km. We prescribed the nucleation rate in the model for latitudes north of 80 °N, which ~~means it has little effect over~~ is further north than the ATom area. We also tested the prescribed rate in gridboxes north of 60 °N and still found good agreement with the ATom and ASCOS profiles (Figs. D1 and D2).

~~The model higher nucleation mode concentrations at the surface in SOA_PRSC relative to SOA (Fig. 5) confirms that 515 particles that have been created in the FT can be entrained into the Arctic boundary layer in the model. Since we prescribed a stronger FT NPF rate for all latitudes north of 80° in SOA_PRSC, the vertical transport of the secondary particles could be occurring in any location over such a region and then be transported through the BL to the location of the helicopter. In the next section, we will examine the effect of this FT source of particles on the particle size distribution at the surface during the AO2018 campaign.~~

~~The model~~ output shown in Figs. 5 and 6 is for the year 2018, while the ~~observational~~ observational data is from 2008 (ASCOS) and 2016 (ATom). However, the large underestimations of the nucleation mode by simulation CONTROL (up to 3 orders of magnitude at the surface) for both campaigns is unlikely to be because of interannual variability alone. The CONTROL simulation also underestimates the particle concentrations from the AO2018 campaign, which was in the same year as our 520

model output. We will show in the next section that simulation SOA_PRSC performs better than CONTROL when evaluated with AO2018 measurements, consistent with the conclusions of this section using older measurements.

4.3.2 Time series of particle concentrations

Time series and PDFs of particle number concentration at the surface from AO2018 observations and simulations M10_ALL and M10_Prse CONTROL and SOA_PRSC are shown in Fig. 7 for nucleation, Aitken and accumulation mode particles.

In the nucleation mode, simulations M10_ALL and M10_Prse both increase simulation SOA_PRSC increases the simulated particle concentration by 2–3 orders of magnitude in the melt period relative to CONTROL (Fig. 2). Like M10_BL, M10_ALL captures the observed nucleation mode concentration well in the melt period. The 7), and by an order of magnitude relative to SOA (Fig. 3). In the freeze period, the higher FT NPF rate in M10_Prse SOA_PRSC produces a nucleation mode concentration that is 1–2 orders of magnitude higher than M10_ALL SOA. This higher concentration in M10_Prse further demonstrates the way the SOA_PRSC demonstrates that the FT is acting as a source of particles to the surface in the model, although the fixed NPF rate in M10_Prse overestimates the nucleation mode concentration in the melt period. The nucleation mode concentration declines from the melt to the freeze period in M10_ALL. The particle concentration decreases from above 1 in the melt period to approximately 1, occasionally reaching 0.1. M10_Prse produces SOA_PRSC produces nucleation mode concentrations close to 100 cm^{-3} in the both both the freeze and melt periods. As such, M10_Prse SOA_PRSC overestimates the nucleation mode concentration in the melt period and captures the lower end of the observed distribution of nucleation concentration in the freeze periods period, but does not capture the highest concentrations seen during the NPF events.

In the Aitken mode, particle concentrations are also higher in M10_ALL and M10_Prse SOA_PRSC than in CONTROL. Despite differences between M10_ALL and M10_Prse SOA and SOA_PRSC in the nucleation mode, they produce the same concentration of Aitken particles. Melt-period Aitken particle concentrations in M10_ALL and M10_Prse SOA_PRSC are 1–3 orders of magnitude higher than in CONTROL, taking them to within 1 order of magnitude of the observed particle concentration for most of the melt period. As in the nucleation mode, Aitken concentrations decrease in M10_ALL and M10_Prse SOA_PRSC in the freeze period such that the observations are underestimated by 1–2 orders of magnitude in the freeze period. The declining Aitken concentrations in both of these simulations suggests a decrease in the particle growth rates towards late summer, possibly driven by declining photochemical production of precursor vapours (see appendix E). The Aitken mode concentration decreases even in M10_Prse, which has a despite the fixed NPF rate above the BL in SOA_PRSC, highlighting that particle growth by condensation is an important process as well as the particle formation rate itself.

The accumulation mode is well captured by both M10_ALL and M10_Prse SOA_PRSC, like in SOA and CONTROL. As for IA_BL, the changes to the model NPF scheme in the M10_SOA simulations have little effect on the accumulation mode concentration relative to CONTROL.

Local formation processes in M10_Prse are dominated by the fixed nucleation rate above the BL. However, local particle formation is also affected by the availability of condensable vapours (if new particles grow faster, they are more likely to survive in the atmosphere for longer rather than being lost to coagulation with other particles). Therefore, since growth is important, we tested the sensitivity of NPF in the M10_Prse simulation to the availability of secondary organic vapour by allowing more

vapour to be transported to the Arctic in simulation M10_Prse_SeeOrg. Time series and PDFs of surface particle concentration during AO2018 is shown on Fig. 7. In the nucleation mode, M10_Prse_SeeOrg produces a concentration of approximately 100, behaving similarly to M10_Prse.

In the Aitken mode, there are periods where M10_Prse_SeeOrg behaves the same as M10_Prse and brief periods where it produces higher concentrations. For example, on days 223–229, the concentration in M10_Prse_SeeOrg is nearly an order of magnitude greater than that from M10_Prse, resulting in an overestimation of the observed Aitken concentration.

4.4 Combining local and non-local NPF

Overall, M10_ALL_SOA performs well in the melt period of AO2018 but still underestimates particle concentration in the freeze period, when simulation IA performs well. This suggests that a combination of secondary particles from aloft transported secondary particles and local BL NPF involving iodic acid is required to capture the full aerosol time series. To test this, we combined the use of the organically-mediated NPF scheme with our iodic acid NPF scheme. The combined simulation of M10_ALL and IA_BL_SOA and IA is shown as simulation IA_BL_M10_ALL_SOA in Fig. 7. IA_BL_M10_ALL_SOA captures both the baseline of the nucleation mode in the melt period and the peaks in the freeze period. In the Aitken mode, IA_BL_M10_ALL_SOA is within 1 order of magnitude of the observations except for a few brief periods, such as days 253–255 when the observed concentration is overestimated in the model.

The behaviour of the simulations can be summarised by comparing the overlap of PDFs of simulated particle concentration with those from observations (Table 3). Simulations with or organically-mediated NPF in the BL (K06_BL, M10_BL, M10_ALL, IA_BL_M10_ALL) have higher overlap indices for the nucleation (0.73) and Aitken modes (0.73) than CONTROL in the melt period, when the source of secondary particles from the FT has the greatest influence, but lower indices shows less of an improvement in the freeze period. IA_SOA also shows improvement relative to CONTROL in the melt period. During freeze-up, IA_BL_M10_ALL_SOA slightly overestimates the nucleation mode concentration, meaning that its nucleation mode overlap index in this period (0.51) is lower than that of M10_Prse_SOA_PRSC (0.52) and M10_Prse_SeeOrg (), which do, which does not overestimate the concentration in the same way. However, it is clear from the time series that the simulation simulations with iodic acid NPF (IA_BL, IA_BL_SeeOrg, IA_BL_M10_ALL_SOA) capture the nucleation concentration peaks from NPF events in a way that the M10_Prse simulations do SOA_PRSC simulation does not. Simulation IA_BL_M10_ALL_SOA has the highest overlap index of all simulations for the Aitken mode in the freeze period (0.77), as it is the only simulation to capture the highest values of Aitken concentration during that time.

5 Discussion and conclusion

We have used field observations and a global aerosol-climate model with an empirical iodic acid nucleation scheme to investigate sources of aerosol in the high Arctic summer. Our results point to a regime transition occurring in late summer from a free-tropospheric source of secondary particles to local boundary layer *in situ* new particle formation in the boundary layer, driven by iodic acid. The onset of iodic acid new particle formation (triggered by sea ice freeze-up) coincides with a decline

| Vapour species | Change in concentration from Aug to Sep 2018 [%] |
|--------------------------------|--|
| DMS | -28.8 |
| SO ₂ | 135.2 |
| H ₂ SO ₄ | -85.2 |
| OH | -87.7 |
| Monoterpenes | 31.2 |
| Secondary organics | -5.9 |

Table 4. Table of change in simulated concentrations of different vapours from August to September 2018 in simulation CONTROL. Concentration changes are given as the change in the mean value over gridboxes at the surface with latitude 80–90 N.

590 in the free tropospheric source rate brought on by declining rates of photochemical production of precursor vapours in the free troposphere. The net effect of the transition from free troposphere to boundary layer nucleation is a fairly constant nucleation and Aitken mode at the surface.

There are several key conclusions we can draw from the simulations we have presented here. They are listed below.

595 – The default settings of the UKESM1 model cannot capture Arctic aerosol concentrations. The nucleation mode particle concentration at the surface is at least an order of magnitude too low in CONTROL compared to AO2018, ASCOS and ATom observations, sometimes underestimating observed concentrations by as much as 5 orders of magnitude. The surface Aitken mode concentration is also underestimated by up to 3 orders of magnitude. The model performs better at simulating the surface accumulation mode concentration. The accumulation mode concentration in CONTROL is generally within the range of the observations, but does not capture the lowest concentrations (less than 1 cm⁻³) seen in the time series of AO2018 observations. These periods of low accumulation mode concentration are important in controlling the behaviour and radiative effects of low-level Arctic clouds (Mauritsen et al., 2011; Loewe et al., 2017; Birch et al., 2012; Stevens et al., 2018).

600 – ~~Simulations with or organically-mediated boundary layer~~ Our simulation with new particle formation ~~produce in the boundary layer produces~~ more accurate nucleation and Aitken mode concentrations at the surface during the AO2018 melt period than the CONTROL simulation. However, modelled aerosol concentrations aloft ~~as well as output from a simulation where~~, ~~combined with results of a simulation with~~ boundary layer new particle formation ~~is switched on only~~ north of 85°N ~~show~~, ~~shows~~ that little to no boundary layer new particle formation takes place in situ during the AO2018 before the iodic acid new particle formation events occur. This ~~result~~ is in contrast to results from Browne et al. (2014), who found that new particle formation from H₂SO₄ in the Arctic boundary layer was an important source of high-Arctic CCN in GLOMAP. Instead, ~~the UKESM simulations with boundary layer new particle formation schemes cause more particles to be created at lower latitudes. These small particles are then transported north through the free troposphere, and provide a source of~~ in these UKESM simulations, nucleation and Aitken mode particles ~~to~~ ~~at~~ the surface in the high Arctic ~~were~~ created at lower latitudes and then transported northwards through the free troposphere. Higher

concentrations at the surface from a simulation with a prescribed new particle formation rate in the low free troposphere
615 supports the hypothesis of a ~~free-tropospheric~~free-tropospheric source to the surface (Fig. 8, left-hand side).

- Our simulations suggest that a seasonal regime shift triggered by changes in photochemistry coincides with the beginning of the iodic acid season triggered by the sea ice freeze-up. This is portrayed in Fig. 8. The net effect of these two changes is a ~~consistent~~fairly constant source of particles controlled by new particle formation from two different mechanisms.
620 Photochemistry declines towards the end of the AO2018 campaign period as a result of the reduction in incoming solar radiation at the end of summer. In the model output from simulation CONTROL, surface concentrations of the OH radical decline by 87.7% over 80–90°N from August 2018, causing H_2SO_4 to decline by 85.2% while SO_2 increases in concentration in the region by 135.2% over the same period (Table 4). The reduction in H_2SO_4 inhibits new particle formation and particle growth (Fig. E5). In simulation ~~M10_PrcSOA_PRSC~~, the Aitken mode particle concentration declines from August to September 2018 even though the ~~main source of these particles~~ (new particle formation rate
625 in the free troposphere) is prescribed to be constant. This behaviour highlights the importance of particle growth rates, which can vary alongside the new particle formation rate itself. Following this decline in the extra-Arctic aerosol source, the local iodic acid-driven particle formation begins.

Our results have implications for the future Arctic aerosol budget. Iodic acid new particle formation in the Arctic boundary layer is strongly coupled to the surface and therefore sensitive to changes in the sea ice. However, our simulations show
630 that entrainment of secondary particles from the free troposphere is also an important source of surface aerosol at nucleation and Aitken mode sizes. Free-tropospheric new particle formation is unlikely to have such a strong sensitivity to local sea ice changes, since the precursor vapours and background aerosol in the free troposphere are likely to have been transported from other regions. Further work will be required to understand how the balance of boundary layer versus free-tropospheric nucleation will evolve in the changing Arctic, since predictions of cloud behaviour and surface energy balance in the future
635 Arctic depend on knowledge of the Arctic aerosol budget. ~~It is also Previous studies have shown that models struggle to simulate the thermodynamic structure of the boundary layer in the high Arctic, for example by failing to produce the decoupled conditions that are common (Birch et al., 2012; Sotiropoulou et al., 2016). Such biases could inhibit accurate predictions of free-tropospheric aerosol sources, since the entrainment of particles in the boundary layer will rely on accurate representation of the turbulent mixing created by the structure of the cloud layer.~~

640 This study shows that it is important to assess the influence of boundary layer versus free-tropospheric sources compared to other uncertain Arctic aerosol processes and their different representations in models, for example sea spray parameterisations and primary marine sources. We did not consider primary sources for the Aitken mode during AO2018, such as biogenic marine particles. Although our model includes primary marine aerosol emissions, these are mostly limited to the accumulation mode. Thus, our model does not include direct marine emissions of smaller particles. Previous field studies in the Arctic pack ice region have found evidence that organics in Aitken and accumulation mode aerosols and cloud water were related to polymer gels found in the surface microlayer of sea water, suggesting a primary marine aerosol source (??Leck and Bigg, 2010; ?; Karl et al., 2013; ?), although our earlier modelling results suggest that to account for

the observed Aitken mode concentrations would require an unrealistically high surface source (Korhonen et al., 2008b). Also proposed in the literature is an atmospheric processing pathway where larger primary particles break-up to form more, smaller particles. We cannot use our model results to exclude such a source from the aerosol size distributions measured during AO2018. The balance of primary and secondary aerosol sources in this region merits further work, and would likely require improvements to existing model parameterisations (i.e. new particle formation and growth rates, primary marine emission fluxes) and the development of new parameterisations to test in the model (i.e. particle emissions from open leads independent of wind speed, and size-resolved break-up rates for primary marine particles).

A greater understanding of aerosol conditions in the Arctic free troposphere is needed. It is not possible to distinguish from the model output presented here whether particles coming from the free troposphere have been created there **by new particle formation** or have first been transported from lower latitudes, where they were created in the boundary layer. Our results indicate the importance of obtaining measurements of aerosol size distributions and precursor vapours from above the boundary layer in the high Arctic, which have previously been sparse. Moreover, the Kulmala et al. (2006) H_2SO_4 and Metzger et al. (2010) organically-mediated new particle formation schemes we use produce very similar surface particle concentrations in the region of study, such that we cannot use these datasets to evaluate the accuracy of one over the other. This highlights an open question in modelling of Arctic aerosol. It is crucial to understand which precursor species are important for Arctic aerosol in order to understand how changes to different parts of the climate system will affect the formation and behaviour of Arctic clouds. The production of secondary organic vapour in UKESM1 is crude, accounting only for the oxidation of monoterpenes, and we have not included any effects of ammonia, which has been shown in laboratory and Arctic field studies to contribute to new particle formation. The inclusion of ammonia in the model could lead to more new particle formation in the boundary layer since ammonia can stabilise H_2SO_4 clusters. However, before these potential sources of model bias can be improved, more observational data is needed about what drives new particle formation throughout the Arctic, both in the free troposphere and at the surface.

The accumulation mode in the model is relatively unaffected by the different new particle formation schemes we used, and is better simulated than the smaller modes. This shows that accumulation mode aerosol and the contribution of the particles to CCN is not affected by new particle formation **in these simulations**. However, comparison with AO2018 observations shows that the model cannot capture periods of up to a day of very low accumulation mode concentration (approximately 1 cm^{-3} and below). Given that the accumulation mode affects new particle formation via the sink of condensable vapours, it will be important to consider this model bias in future work. Moreover, observations indicate the importance of fog as a sink of the iodic acid, so more work is needed to assess the role of fog in controlling the frequency of iodic acid new particle formation events. A higher resolution model with better cloud parameterisations would be better suited to such studies than the coarse, global model we have used here.

Code availability. The code used to analyse model output and produce the figures is available on Github at <https://github.com/ruthprice/price-acp-figures>.

Data availability. Data from the Arctic Ocean 2018 campaign is available on the Bolin Data Centre, <https://bolin.su.se/data/oden-ao-2018-expedition-2>. The data used here from the ASCOS campaign is available on the Bolin Centre Database, <https://doi.org/10.17043/oden-ascos-2008-aerosol-stratification-1>. All data from the Atmospheric Tomography mission are openly available and archived in the Oak Ridge National Laboratory Distributed Active Archive Center (ORNL DAAC) at <https://doi.org/10.3334/ORNLDAAAC/1925>.

Images/fig07.pdf

Figure 7. Time series and PDFs of aerosol concentration at the surface during AO2018 from observations and model output. Model output is from simulations **M10_ALL_CONTROL** (orange-pink lines), **M10_SOA_Prse_PRSC** (thick green lines), **M10_Prse_SeeOrg** (thin green lines) and **IA_BL_M10_ALL_SOA** (red lines). Other than **IA_BL_M10_ALL_SOA** and the observations, lines have been made slightly transparent on this figure so that it is easier to view the results from **IA_BL_M10_ALL_SOA**. Observations are shown as 3-hourly mean (black lines) and standard deviation (grey shading). Aerosol concentrations are shown for particles with diameter (a–c) 2.5–15 nm, (d–f) 15–100 nm and (g–i) 100–500 nm. Red dashed lines in (a, d, g) show observed NPF events. PDFs are separated by observed sea ice freeze-up date, 27th August 2018 (day 239).

Images/fig08.pdf

Figure 8. Schematic of processes controlling the concentration of nucleation mode particles at the surface in the high Arctic. The summer sea ice melt period is on the left and the freeze-up period in late summer/early autumn is on the right. A [eartoon schematic](#) of the nucleation mode particle concentration is shown at the bottom.

Images/figA1.pdf

Figure A1. Time series of surface temperature during the AO2018 campaign periods from observations and simulation CONTROL. Observations are shown as 30-min mean values (grey line) and 14-day running mean values (black line). Model output is shown as 3-hourly mean values (pale pink line) and 14-day running mean values (dark pink line). The observed sea ice freeze-up point (when the 14-day mean reaches -2°C) is marked with a red dashed line.

Figure A1 shows the surface temperature during AO2018 from ship measurements and CONTROL output. The sea ice freeze-up point is defined to be when the 14-day running mean of the surface temperature reaches -2°C . During AO2018, this occurred on 27th August 2018 (day 239). The model output at this time gives a surface temperature of approximately -6°C . We use -5°C as the proxy for sea ice freezing in the iodic acid scheme instead of -2°C , so that the iodic acid is triggered on 690 roughly the observed freeze-up day in the model.

Appendix B: Choice of new particle formation parameterisation

In section 4 we show that switching on NPF in the BL in the model creates more particles at the surface than using the Vehkamäki et al. (2002) scheme only (simulation SOA relative to CONTROL). To test the sensitivity of the results to the choice of BL NPF scheme, we ran simulations using the K06 scheme (simulation SA) and using the M10 scheme in the FT 695 as well as the BL (SOA_ALL_LEVELS). Aerosol number concentrations at the surface from these simulations are shown in figure B1. Interestingly, simulations SA, SOA and SOA_ALL_LEVELS have very similar aerosols concentrations at the surface, despite the differences in the NPF schemes. SA occasionally simulates higher nucleation mode concentrations than SOA or SOA_ALL_LEVELS, most notably on days 239–243.

Images/figB1.pdf

Figure B1. Time series and PDFs of surface aerosol concentration during AO2018 from observations and model output. Model output is from simulations SA (grey lines), SOA (blue lines) and SOA_ALL_LEVELS (orange lines). Observations are shown as 3-hourly mean (black lines) and standard deviation (grey shading). Aerosol concentrations are shown for particles with diameter (a–c) 2.5–15 nm, (d–f) 15–100 nm and (g–i) 100–500 nm. Red dashed lines in (a, d, g) show observed NPF events. PDFs are separated by observed sea ice freeze-up date, 27th August 2018 (day 239).

Appendix C: Particle ageing

700 In Sect. 4.2 we show that ~~K06_BL_and_M10_BL_SA_and_SOA~~ simulate more accurate concentrations of the nucleation and Aitken modes during AO2018 than simulation CONTROL (3). Figure C1 shows the surface aerosol concentrations for simulations UKESM_default and CONTROL, ~~K06_BL_default_and_K06_BL_and_M10_BL_default_and_M10_BL_SA_default_and_SA_and_SOA_default_and_SOA~~. The simulations with the ageing change produce higher concentrations in the nucleation and Aitken modes than the DEFAULT simulations, and this difference is greater in the simulations with BL NPF than it is between UKESM_DEFAULT and CONTROL. This is because the mode merging described in Sect. 3.4 affects the Aitken mode concentration more when there is a stronger source of smaller particles. In the Aitken mode, the concentrations from simulations ~~K06_BL_and_M10_BL_SA_and_SOA~~ are typically higher than ~~K06_BL_DEFAULT_and_M10_BL_SA_DEFAULT_and_SOA_DEFAULT~~ by a factor of 2–10. The model-observation agreement is therefore improved in simulations with the ageing change.

705

710 Figure C2 shows mean aerosol profiles simulated for the AO2018 period for the soluble nucleation, Aitken and accumulation modes and the insoluble mode. Model output is shown for simulations CONTROL, UKESM_default, ~~K06_BL_SA(_default)~~ ~~M10_BL_SOA(_default)~~, ~~M10_FT_SOA_ALL_LEVELS(_default)~~, ~~M10_ALL_SOA_ALL_LEVELS(_default)~~ and ~~M10_Prse_SOA_PRSC(_default)~~. The perturbed ageing scheme produces up to an order of magnitude more Aitken mode particles than the default scheme in the low FT and at the surface, independently of which NPF scheme is used. The insoluble mode is not significantly affected by the 715 changes to the ageing scheme.

Appendix D: Latitude limit for prescribed FT NPF rate

720 In simulation ~~M10_Prse_SOA_PRSC~~, we impose a NPF rate of $10^{-2} \text{ cm}^{-3} \text{ s}^{-1}$ in model levels between the top of the boundary layer and 7.5 km altitude. The rate is imposed in gridboxes north of 80°N . In Sect. 4.3.1 we compare the ~~M10_Prse_SOA_PRSC~~ output to aerosol profiles from the ASCOS and ATom campaigns to test how realistic such a NPF rate may be. However, the data we use from ATom was taken further south, between approximately 60 – 80°N (see Fig. 1). To test our prescribed FT NPF rate using ATom observations we therefore ran another simulation, ~~M10_Prse_SOA_PRSC_60N~~, where the rate is imposed north of 60°N .

725 Figures D1 and D2 show output from ~~M10_Prse_and_M10_Prse_SOA_PRSC_and_SOA_PRSC_60N~~ with observations from ASCOS and ATom. Aerosol concentrations from the two simulations are within an order of magnitude of each other. While the prescribed rate increases aerosol concentrations relative to CONTROL in the ASCOS region (Sect. 4.3.1), it makes little difference in the ATom region even when we extend the region in which the rate is applied. This is likely because NPF rates are already higher in the ATom region than the ASCOS region due to the relative proximity of ATom to open water and boreal forests, both of which supply precursor vapours to the atmosphere.

Images/figC1.pdf

Figure C1. Time series and PDFs of surface aerosol concentration during AO2018 from observations and model output. Model output is from simulations UKESM_DEFAULT (thin pink lines), CONTROL (thick pink lines), **K06SA_BL_DEFAULT** (think grey lines), **K06_BL_SA** (thick grey lines), **M10SOA_BL_DEFAULT** (thin blue lines) and **M10_BL_SOA** (thick blue lines). Observations are shown as 3-hourly mean (black lines) and standard deviation (grey shading). Aerosol concentrations are shown for particles with diameter (a–c) 2.5–15 nm, (d–f) 15–100 nm and (g–i) 100–500 nm. Red dashed lines in (a, d, g) show observed NPF events. PDFs are separated by observed sea ice freeze-up date, 27th August 2018 (day 239).

Images/figC2.pdf

Figure C2. Mean simulated aerosol profiles for AO2018 campaign period. Concentrations shown are for (a–b) soluble nucleation, (c–d) soluble Aitken, (e–f) soluble accumulation and (g–h) insoluble Aitken modes. Model output is from simulations UKCA_DEFAULT and CONTROL (pink lines), **K06SA_BL**_default and **K06_BL_SA** (grey lines), **M10SOA_ALL_LEVELS**_default and **M10SOA_ALL_LEVELS** (red-orange lines), **M10SOA_BL**_default and **M10_BL_SOA** (dark-blue lines) and **M10SOA_PrcsPRSC**_default and **M10SOA_Prcs-PRSC** (green lines).

Images/figD1.pdf

Figure D1. Aerosol vertical profiles from model output and ASCOS campaign observations. Model output is from colocated monthly mean values from simulations [M10SOA_Prse_PRSC](#) (thick dark green) and [M10SOA_PrsePRSC_60N](#) (thin dark green). Observed values are given as the mean profile from each ASCOS flight (grey lines), overall mean (solid black) and overall median (dashed black). Profiles are for particles with size (a) 3–14 nm, measured during ASCOS using a UCPC and (b) 14–300 nm, measured using a CPC (particles greater than 14 nm) and a CLASP instrument (particles greater than 300 nm).

Images/figD2.pdf

Figure D2. Aerosol vertical profiles from model output and ATom campaign observations. Model output is from colocated monthly mean values from simulations [M10SOA_Prcs-PRSC](#) (thick dark green) and [M10SOA_PrcsPRSC_60N](#) (thin dark green). ATom observations are taken from leg 1 of the campaign and restricted to measurements that were taken north of 60°N. Observations correspond to mean profiles from different days (grey lines), the overall mean (black solid lines) and overall median (black dashed lines). Profiles are for particles with size (a) 5–10 nm, (b) 10–100 nm and (c) 100–500 nm. Observations were recorded at standard temperature and pressure, model output has been adjusted to account for this.

Appendix E: Aerosol precursor vapours and growth rate

Images/figE1.pdf

Figure E1. Time series and PDFs of surface aerosol concentration during AO2018 from observations and model output. Model output is from simulations SOA (thick blue lines) and SOA_OXID (thin blue lines). Observations are shown as 3-hourly mean (black lines) and standard deviation (grey shading). Aerosol concentrations are shown for particles with diameter (a–c) 2.5–15 nm, (d–f) 15–100 nm and (g–i) 100–500 nm. Red dashed lines in (a, d, g) show observed NPF events. PDFs are separated by observed sea ice freeze-up date, 27th August 2018 (day 239).

Images/figE2.pdf

Figure E2. Time series and PDFs of surface aerosol concentration during AO2018 from observations and model output. Model output is from simulations SOA_PRSC (thick green lines) and SOA_PRSC_OXID (thin green lines). Observations are shown as 3-hourly mean (black lines) and standard deviation (grey shading). Aerosol concentrations are shown for particles with diameter (a–c) 2.5–15 nm, (d–f) 15–100 nm and (g–i) 100–500 nm. Red dashed lines in (a, d, g) show observed NPF events. PDFs are separated by observed sea ice freeze-up date, 27th August 2018 (day 239).

730 In our simulations we have tested the effect of different NPF rates on the modelled aerosol concentrations. However, particle formation is also affected by the availability of condensable vapours (if new particles grow faster, they are more likely to survive in the atmosphere for longer rather than being lost to coagulation with other particles). Therefore, since growth is important, we tested the sensitivity of NPF in the SOA and SOA_PRSC simulations to the availability of secondary organic vapour by allowing more vapour to be transported to the Arctic in simulations SOA_OXID and SOA_PRSC_OXID. Time series and PDFs of surface particle concentration during AO2018 are shown on Fig. E1 and Fig. E2. In the nucleation mode, SOA_PRSC_OXID produces a concentration of approximately 100 cm^{-3} , behaving similarly to SOA_PRSC. SOA_OXID produces nucleation mode concentrations up to 3 orders of magnitude greater than SOA for brief periods between 22nd August and 1st September 2018, but overall this effect is not enough to account for the underestimation of the observed nucleation mode concentrations by SOA in the freeze period.


740 In the Aitken mode, there are periods where SOA_PRSC_OXID behaves the same as SOA_PRSC and brief periods where it produces higher concentrations. For example, on days 223–229, the concentration in SOA_PRSC_OXID is nearly an order of magnitude greater than that from SOA_PRSC, resulting in an overestimation of the observed Aitken concentration. The Aitken mode concentrations in SOA_OXID are similar to that of SOA.

745 Figures E3 and E4 show maps and zonal means of H_2SO_4 and secondary organic vapour concentration from simulation CONTROL. In August, Arctic H_2SO_4 concentration peaks just above the surface and at approximately 8 km (Fig. E3(a)). The concentration decreases throughout most of the troposphere from August to September by at least an order of magnitude. In August, the secondary organics have a maximum at the surface from $60\text{--}70^\circ\text{N}$ and some of this plume spreads north to the high Arctic. Unlike H_2SO_4 , concentrations do not significantly reduce from August to September. The mean surface concentration of H_2SO_4 for $80\text{--}90^\circ\text{N}$ reduces by 85.2% from August to September, while for secondary organics the reduction is only 5.9%.

750 Percentage changes for other vapours are given in Table 4.

The decline in vapour concentration from August to September drives a reduction in the growth rate of aerosols from condensation of vapour. Figure E5 shows aerosol growth rates calculated from model output from simulation CONTROL. Arctic growth rates are lower than in the mid-latitudes and tropics, where H_2SO_4 and secondary organics have higher concentrations. In the central Arctic, the decreasing growth rate is driven by the decrease in H_2SO_4 . Secondary organic vapours contribute little 755 to the growth rate in the central Arctic region, but dominate in the continental Arctic in North America and northern Eurasia, where concentrations are high near the boreal forest source regions.

In Sect. 4.3.2 we showed that the source of aerosols from FT NPF weakens towards the end of summer. This occurs even when the FT NPF rate is held constant in time as in simulation M10_PresSOA_PRSC. The slower aerosol growth rate shown in Fig. E5 curbs FT NPF in September. This highlights that it is important to understand the growth of new particles as well as 760 their formation.




Images/figE3.pdf

Figure E3. Zonal means and maps of simulated monthly mean H_2SO_4 concentration from simulation CONTROL. Maps are taken from model level (c–d) at surface and (e–f) with altitude 2 km.

Author contributions. RP ran model simulations, assisted by PF, and analysed output with guidance from KSC. JS, AB and PZ performed aerosol measurements during AO2018. RP, PF, IMB and KSC contributed towards the design of the study. All authors were involved in discussion of results and writing of the manuscript.

Competing interests. KSC is an executive editor of ACP. The peer-review process was guided by an independent editor, and the authors have
765 also no other competing interests to declare.

Acknowledgements. RP was supported by a Natural Environment Research Council (NERC) Ph.D. studentship through the SPHERES Doctoral Training Partnership (NE/L002574/1). PZ was supported by the Swedish Research Council (VR starting grant 2018-05045). JS




Images/figE4.pdf

Figure E4. Zonal means and maps of simulated monthly mean secondary organic vapour concentration from simulation CONTROL. Maps are taken from model level (c–d) sat surface and (e–f) with altitude 2 km.

and AB received funding from the Swiss National Science Foundation (Grant no. 169090) and the Swiss Polar Institute. JS holds the Ingvar Kamprad Chair for Extreme Environments Research sponsored by Ferring Pharmaceuticals. KSC was funded by the NERC project A-CURE 770 (NE/P013406/1). IMB was funded by NERC (grant No. NE/R009686/1)

We thank Hamish Gordon for his advice on new particle formation schemes in the model. We acknowledge use of the Monsoon2 system, a collaborative facility supplied under the Joint Weather and Climate Research Programme, a strategic partnership between the Met Office and the Natural Environment Research Council. This work used JASMIN, the UK's collaborative data analysis environment (<https://jasmin.ac.uk>).

775 This work is part of the ASCOS and Arctic Ocean (AO) 2018 expeditions. The Swedish Polar Research Secretariat (SPRS) provided access to the icebreaker Oden and logistical support. We are grateful to the Chief Scientists Caroline Leck, Patricia Matrai and Michael Tjernström for planning and coordination of ASCOS and AO2018, to the SPRS logistical staff and to I/B Oden's Captain Mattias Peterson and his crew.



Images/figE5.pdf

Figure E5. Zonal means and maps of aerosol growth rates from simulation CONTROL. Growth rates are calculated offline using model output of temperature and concentrations of H_2SO_4 and secondary organic vapour. Maps are taken from model level (c–d) at surface and (e–f) with altitude 2 km.

We thank the two reviewers and the community commenters for their time spent giving feedback on this paper.

Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation: 2. Multiple aerosol types, *Journal of Geophysical Research: Atmospheres*, 105, 6837–6844, <https://doi.org/10.1029/1999JD901161>, 2000.

Allan, J. D., Williams, P. I., Najera, J., Whitehead, J. D., Flynn, M. J., Taylor, J. W., Liu, D., Darbyshire, E., Carpenter, L. J., Chance, R., Andrews, S. J., Hackenberg, S. C., and McFiggans, G.: Iodine observed in new particle formation events in the Arctic atmosphere during 785 ACCACIA, *Atmospheric Chemistry and Physics*, 15, 5599–5609, <https://doi.org/10.5194/acp-15-5599-2015>, 2015.

Alterskjær, K., Kristjánsson, J. E., and Hoose, C.: Do anthropogenic aerosols enhance or suppress the surface cloud forcing in the Arctic?, *Journal of Geophysical Research Atmospheres*, 115, <https://doi.org/10.1029/2010JD014015>, 2010.

AMAP: Arctic Climate Change Update 2021: Key Trends and Impacts. Summary for Policy-makers, Arctic Monitoring and Assessment Programme (AMAP), Trømso, Norway, 2021.

790 Baccarini, A. and Schmale, J.: Ultrafine particle concentration measured during the Arctic Ocean 2018 expedition [data set], <https://doi.org/10.17043/oden-ao-2018-aerosol-ufp-1>, 2020.

Baccarini, A., Karlsson, L., Dommen, J., Duplessis, P., Vüllers, J., Brooks, I. M., Saiz-Lopez, A., Salter, M., Tjernström, M., Baltensperger, U., Zieger, P., and Schmale, J.: Frequent new particle formation over the high Arctic pack ice by enhanced iodine emissions, *Nature Communications*, 11, 4924, <https://doi.org/10.1038/s41467-020-18551-0>, 2020a.

795 Baccarini, A., Schmale, J., and Dommen, J.: Iodic acid, sulfuric acid and methanesulfonic acid collected during the Arctic Ocean 2018 expedition [data set], <https://doi.org/10.17043/oden-ao-2018-aerosol-cims-1>, 2020b.

Beck, L. J., Sarnela, N., Junninen, H., Hoppe, C. J. M., Garmash, O., Bianchi, F., Riva, M., Rose, C., Per akyl a, O., Wimmer, D., Kausiala, O., Jokinen, T., Ahonen, L., Mikkil a, J., Hakala, J., He, X., Kontkanen, J., Wolf, K. K. E., Cappelletti, D., Mazzola, M., Traversi, R., Petroselli, C., Viola, A. P., Vitale, V., Lange, R., Massling, A., Nojgaard, J. K., Krejci, R., Karlsson, L., Zieger, P., Jang, S., Lee, K., 800 Vakkari, V., Lampilahti, J., Thakur, R. C., Leino, K., Kangasluoma, J., Duplissy, E., Siivola, E., Marbouti, M., Tham, Y. J., Saiz-Lopez, A., Pet aj a, T., Ehn, M., Worsnop, D. R., Skov, H., Kulmala, M., Kerminen, V., and Sipil a, M.: Differing mechanisms of new particle formation at two Arctic sites., *Geophysical Research Letters*, <https://doi.org/10.1029/2020gl091334>, 2020.

Bigg, E. K. and Leck, C.: Cloud-active particles over the central Arctic Ocean, *Journal of Geophysical Research: Atmospheres*, 106, 32 155–32 166, <https://doi.org/10.1029/1999JD901152>, 2001.

805 Bigg, K. E., Leck, C., and Nilsson, D. E.: Sudden changes in arctic atmospheric aerosol concentrations during summer and autumn, *Tellus B: Chemical and Physical Meteorology*, 48, 254–271, <https://doi.org/10.3402/tellusb.v48i2.15890>, 1996.

Birch, C. E., Brooks, I. M., Tjernström, M., Shupe, M. D., Mauritsen, T., Sedlar, J., Lock, A. P., Earnshaw, P., Persson, P. O. G., Milton, S. F., and Leck, C.: Modelling atmospheric structure, cloud and their response to CCN in the central Arctic: ASCOS case studies, *Atmospheric Chemistry and Physics*, 12, 3419–3435, <https://doi.org/10.5194/acp-12-3419-2012>, 2012.

810 Biskaborn, B. K., Smith, S. L., Noetzli, J., Matthes, H., Vieira, G., Streletschi, D. A., Schoeneich, P., Romanovsky, V. E., Lewkowicz, A. G., Abramov, A., Allard, M., Boike, J., Cable, W. L., Christiansen, H. H., Delaloye, R., Diekmann, B., Drozdov, D., Etzelm uller, B., Grosse, G., Guglielmin, M., Ingeman-Nielsen, T., Isaksen, K., Ishikawa, M., Johansson, M., Johannsson, H., Joo, A., Kaverin, D., Kholodov, A., Konstantinov, P., Kr oger, T., Lambiel, C., Lanckman, J. P., Luo, D., Malkova, G., Meiklejohn, I., Moskalenko, N., Oliva, M., Phillips, M., Ramos, M., Sannel, A. B. K., Sergeev, D., Seybold, C., Skryabin, P., Vasiliev, A., Wu, Q., Yoshikawa, K., Zheleznyak, M., and Lantuit, H.: Permafrost is warming at a global scale, *Nature Communications*, 10, <https://doi.org/10.1038/s41467-018-08240-4>, 2019.

Boyer, M., Aliaga, D., Pernov, J. B., Angot, H., Quélérver, L. L. J., Dada, L., Heutte, B., Dall’Osto, M., Beddows, D. C. S., Brasseur, Z., Beck, I., Bucci, S., Duetsch, M., Stohl, A., Laurila, T., Asmi, E., Massling, A., Thomas, D. C., Nøjgaard, J. K., Chan, T., Sharma, S., Tunved, P., Krejci, R., Hansson, H. C., Bianchi, F., Lehtipalo, K., Wiedensohler, A., Weinhold, K., Kulmala, M., Petäjä, T., Sipilä, M., Schmale, J., and Jokinen, T.: A full year of aerosol size distribution data from the central Arctic under an extreme positive Arctic Oscillation: insights from the Multidisciplinary drifting Observatory for the Study of Arctic Climate (MOSAiC) expedition, *Atmospheric Chemistry and Physics*, 23, 389–415, <https://doi.org/10.5194/acp-23-389-2023>, 2023.

Brock, C. A., Williamson, C., Kupc, A., Froyd, K. D., Erdesz, F., Wagner, N., Richardson, M., Schwarz, J. P., Gao, R. S., Katich, J. M., Campuzano-Jost, P., Nault, B. A., Schroder, J. C., Jimenez, J. L., Weinzierl, B., Dollner, M., Bui, T., and Murphy, D. M.: Aerosol size distributions during the Atmospheric Tomography Mission (ATom): Methods, uncertainties, and data products, *Atmospheric Measurement Techniques*, 12, 3081–3099, <https://doi.org/10.5194/amt-12-3081-2019>, 2019.

Brooks, I. M., Tjernström, M., Persson, P. O. G., Shupe, M. D., Atkinson, R. A., Canut, G., Birch, C. E., Mauritsen, T., Sedlar, J., and Brooks, B. J.: The Turbulent Structure of the Arctic Summer Boundary Layer During The Arctic Summer Cloud-Ocean Study, *Journal of Geophysical Research: Atmospheres*, 122, 9685–9704, <https://doi.org/10.1002/2017JD027234>.

Browse, J., Carslaw, K. S., Arnold, S. R., Pringle, K., and Boucher, O.: The scavenging processes controlling the seasonal cycle in Arctic sulphate and black carbon aerosol, *Atmospheric Chemistry and Physics*, 12, 6775–6798, <https://doi.org/10.5194/acp-12-6775-2012>, 2012.

Browse, J., Carslaw, K. S., Mann, G. W., Birch, C. E., Arnold, S. R., and Leck, C.: The complex response of Arctic aerosol to sea-ice retreat, *Atmospheric Chemistry and Physics*, 14, 7543–7557, <https://doi.org/10.5194/acp-14-7543-2014>, 2014.

Bulatovic, I., Igel, A. L., Leck, C., Heintzenberg, J., Riipinen, I., and Ekman, A. M. L.: The importance of Aitken mode aerosol particles for cloud sustenance in the summertime high Arctic – a simulation study supported by observational data, *Atmospheric Chemistry and Physics*, 21, 3871–3897, <https://doi.org/10.5194/acp-21-3871-2021>, 2021.

Cazenave, A., Meyssignac, B., Ablain, M., Balmaseda, M., Bamber, J., Barletta, V., Beckley, B., Benveniste, J., Berthier, E., Blazquez, A., Boyer, T., Caceres, D., Chambers, D., Champollion, N., Chao, B., Chen, J., Cheng, L., Church, J. A., Chuter, S., Cogley, J. G., Dangendorf, S., Desbruyères, D., D'oll, P., Domingues, C., Falk, U., Famiglietti, J., Fenoglio-Marc, L., Forsberg, R., Galassi, G., Gardner, A., Groh, A., Hamlington, B., Hogg, A., Horwath, M., Humphrey, V., Husson, L., Ishii, M., Jaeggi, A., Jevrejeva, S., Johnson, G., Kolodziejczyk, N., Kusche, J., Lambeck, K., Landerer, F., Leclercq, P., Legresy, B., Leuliette, E., Llovel, W., Longuevergne, L., Loomis, B. D., Luthcke, S. B., Marcos, M., Marzeion, B., Merchant, C., Merrifield, M., Milne, G., Mitchum, G., Mohajerani, Y., Monier, M., Monselesan, D., Nerem, S., Palanisamy, H., Paul, F., Perez, B., Piecuch, C. G., Ponte, R. M., Purkey, S. G., Reager, J. T., Rietbroek, R., Rignot, E., Riva, R., Roemmich, D. H., Sørensen, L. S., Sasgen, I., Schrama, E. J., Seneviratne, S. I., Shum, C. K., Spada, G., Stammer, D., van de Wal, R., Velicogna, I., von Schuckmann, K., Wada, Y., Wang, Y., Watson, C., Wiese, D., Wijffels, S., Westaway, R., Woppelmann, G., and Wouters, B.: Global sea-level budget 1993–present, *Earth System Science Data*, 10, 1551–1590, <https://doi.org/10.5194/essd-10-1551-2018>, 2018.

Chang, R. Y., Sjostedt, S. J., Pierce, J. R., Papakyriakou, T. N., Scarratt, M. G., Michaud, S., Levasseur, M., Leaitch, W. R., and Abbatt, J. P.: Relating atmospheric and oceanic DMS levels to particle nucleation events in the Canadian Arctic, *Journal of Geophysical Research Atmospheres*, 116, <https://doi.org/10.1029/2011JD015926>, 2011.

Croft, B., Wentworth, G. R., Martin, R. V., Leaitch, W. R., Murphy, J. G., Murphy, B. N., Kodros, J. K., Abbatt, J. P., and Pierce, J. R.: Contribution of Arctic seabird-colony ammonia to atmospheric particles and cloud-albedo radiative effect, *Nature Communications*, 7, <https://doi.org/10.1038/ncomms13444>, 2016.

Croft, B., Martin, R. V., Leaitch, W. R., Burkart, J., Chang, R. Y.-W., Collins, D. B., Hayes, P. L., Hodshire, A. L., Huang, L., Kodros, J. K., Moravek, A., Mungall, E. L., Murphy, J. G., Sharma, S., Tremblay, S., Wentworth, G. R., Willis, M. D., Abbatt, J. P. D., and Pierce,

855 J. R.: Arctic marine secondary organic aerosol contributes significantly to summertime particle size distributions in the Canadian Arctic Archipelago, *Atmospheric Chemistry and Physics*, 19, 2787–2812, <https://doi.org/10.5194/acp-19-2787-2019>, 2019.

Curry, J. A., Schramm, J. L., and Ebert, E. E.: Impact of clouds on the surface radiation balance of the Arctic Ocean, *Meteorology and Atmospheric Physics*, 51, 197–217, <https://doi.org/10.1007/BF01030494>, 1993.

860 Dall’Osto, M., Beddows, D. C., Tunved, P., Krejci, R., Strom, J., Hansson, H. C., Yoon, Y. J., Park, K. T., Becagli, S., Udisti, R., Onasch, T., O’dowd, C. D., Simó, R., and Harrison, R. M.: Arctic sea ice melt leads to atmospheric new particle formation, *Scientific Reports*, 7, <https://doi.org/10.1038/s41598-017-03328-1>, 2017.

Dall’Osto, M., Geels, C., Beddows, D. C., Boertmann, D., Lange, R., Nojgaard, J. K., Harrison, R. M., Simo, R., Skov, H., and Massling, A.: Regions of open water and melting sea ice drive new particle formation in North East Greenland, *Scientific Reports*, 8, <https://doi.org/10.1038/s41598-018-24426-8>, 2018.

Engvall, A.-C., Krejci, R., Ström, J., Minikin, A., Treffeisen, R., Stohl, A., and Herber, A.: In-situ airborne observations of the microphysical 865 properties of the Arctic tropospheric aerosol during late spring and summer, *Tellus B: Chemical and Physical Meteorology*, 60, 392–404, <https://doi.org/10.1111/j.1600-0889.2008.00348.x>, 2008.

Finkenzeller, H., Iyer, S., He, X.-C., Simon, M., Koenig, T. K., Lee, C. F., Valiev, R., Hofbauer, V., Amorim, A., Baalbaki, R., Baccarini, A., Beck, L., Bell, D. M., Caudillo, L., Chen, D., Chiu, R., Chu, B., Dada, L., Duplissy, J., Heinritzi, M., Kemppainen, D., Kim, C., Krechmer, J., Kürten, A., Kvashnin, A., Lamkaddam, H., Lee, C. P., Lehtipalo, K., Li, Z., Makhmutov, V., Manninen, H. E., Marie, G., Marten, R., 870 Mauldin, R. L., Mentler, B., Müller, T., Petäjä, T., Philippov, M., Ranjithkumar, A., Rörup, B., Shen, J., Stolzenburg, D., Tauber, C., Tham, Y. J., Tomé, A., Vazquez-Pufleau, M., Wagner, A. C., Wang, D. S., Wang, M., Wang, Y., Weber, S. K., Nie, W., Wu, Y., Xiao, M., Ye, Q., Zauner-Wieczorek, M., Hansel, A., Baltensperger, U., Brioude, J., Curtius, J., Donahue, N. M., Haddad, I. E., Flagan, R. C., Kulmala, M., Kirkby, J., Sipilä, M., Worsnop, D. R., Kurten, T., Rissanen, M., and Volkamer, R.: The gas-phase formation mechanism of iodic acid as an atmospheric aerosol source, *Nature Chemistry*, <https://doi.org/10.1038/s41557-022-01067-z>, 2022.

875 Flanner, M. G., Shell, K. M., Barlage, M., Perovich, D. K., and Tschudi, M. A.: Radiative forcing and albedo feedback from the Northern Hemisphere cryosphere between 1979 and 2008, *Nature Geoscience*, 4, 151–155, <https://doi.org/10.1038/ngeo1062>, 2011.

Gantt, B., Johnson, M. S., Crippa, M., Prévôt, A. S., and Meskhidze, N.: Implementing marine organic aerosols into the GEOS-Chem model, *Geoscientific Model Development*, 8, 619–629, <https://doi.org/10.5194/gmd-8-619-2015>, 2015.

Garrett, T. J. and Zhao, C.: Increased Arctic cloud longwave emissivity associated with pollution from mid-latitudes, *Nature*, 440, 787–789, 880 <https://doi.org/10.1038/nature04636>, 2006.

Gilgen, A., Huang, W. T. K., Ickes, L., Neubauer, D., and Lohmann, U.: How important are future marine and shipping aerosol emissions in a warming Arctic summer and autumn?, *Atmospheric Chemistry and Physics*, 18, 10 521–10 555, <https://doi.org/10.5194/acp-18-10521-2018>, 2018.

Gong, S. L.: A parameterization of sea-salt aerosol source function for sub- and super-micron particles, *Global Biogeochemical Cycles*, 17, 885 <https://doi.org/10.1029/2003gb002079>, 2003.

Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., K urten, A., Lehtipalo, K., Makhmutov, V., Molteni, U., Rissanen, M. P., Stozkhov, Y., Tr ostl, J., Tsagkogeorgas, G., Wagner, R., Williamson, C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K. S.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres, *Journal of Geophysical Research: Atmospheres*, 122, 8739–8760, <https://doi.org/10.1002/2017JD026844>, 2017.

He, X.-C., Tham, Y. J., Dada, L., Wang, M., Finkenzeller, H., Stolzenburg, D., Iyer, S., Simon, M., Kürten, A., Shen, J., Rörup, B., Rissanen, M., Schobesberger, S., Baalbaki, R., Wang, D. S., Koenig, T. K., Jokinen, T., Sarnela, N., Beck, L. J., Almeida, J., Amanatidis, S., Amorim, A., Ataei, F., Baccarini, A., Bertozi, B., Bianchi, F., Brilke, S., Caudillo, L., Chen, D., Chiu, R., Chu, B., Dias, A., Ding, A., Dommen, J., Duplissy, J., Haddad, I. E., Carracedo, L. G., Granzin, M., Hansel, A., Heinritzi, M., Hofbauer, V., Junninen, H., Kangasluoma, J., 895 Kemppainen, D., Kim, C., Kong, W., Krechmer, J. E., Kvashin, A., Laitinen, T., Lamkaddam, H., Lee, C. P., Lehtipalo, K., Leiminger, M., Li, Z., Makhmutov, V., Manninen, H. E., Marie, G., Marten, R., Mathot, S., Mauldin, R. L., Mentler, B., Möhler, O., Müller, T., Nie, W., Onnela, A., Petäjä, T., Pfeifer, J., Philippov, M., Ranjithkumar, A., Saiz-Lopez, A., Salma, I., Scholz, W., Schuchmann, S., Schulze, B., Steiner, G., Stozhkov, Y., Tauber, C., Tomé, A., Thakur, R. C., Väisänen, O., Vazquez-Pufleau, M., Wagner, A. C., Wang, Y., Weber, S. K., Winkler, P. M., Wu, Y., Xiao, M., Yan, C., Ye, Q., Ylisirniö, A., Zauner-Wieczorek, M., Zha, Q., Zhou, P., Flagan, R. C., Curtius, J., Baltensperger, U., Kulmala, M., Kerminen, V.-M., Kurtén, T., Donahue, N. M., Volkamer, R., Kirkby, J., Worsnop, D. R., and Sipilä, M.: Role of iodine oxoacids in atmospheric aerosol nucleation, <http://science.sciencemag.org/>, 2021.

Heintzenberg, J., Tunved, P., Galí, M., and Leck, C.: New particle formation in the Svalbard region 2006–2015, *Atmospheric Chemistry and Physics*, 17, 6153–6175, <https://doi.org/10.5194/acp-17-6153-2017>, 2017.

Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, 900 T. C., Dawidowski, L., Kholod, N., Kurokawa, J. I., Li, M., Liu, L., Lu, Z., Moura, M. C., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.

Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A., Ping, C. L., Schirrmeyer, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks of circumpolar permafrost 910 carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573–6593, <https://doi.org/10.5194/bg-11-6573-2014>, 2014.

Igel, A. L., Ekman, A. M. L., Leck, C., Tjernström, M., Savre, J., and Sedlar, J.: The free troposphere as a potential source of arctic boundary layer aerosol particles, *Geophysical Research Letters*, 44, 7053–7060, <https://doi.org/10.1002/2017GL073808>, 2017.

Karl, M., Leck, C., Gross, A., and Pirjola, L.: A study of new particle formation in the marine boundary layer over the central 915 Arctic Ocean using a flexible multicomponent aerosol dynamic model, *Tellus, Series B: Chemical and Physical Meteorology*, 64, <https://doi.org/10.3402/tellusb.v64i0.17158>, 2012.

Karl, M., Leck, C., Coz, E., and Heintzenberg, J.: Marine nanogels as a source of atmospheric nanoparticles in the high Arctic, *Geophysical Research Letters*, 40, 3738–3743, <https://doi.org/10.1002/grl.50661>, 2013.

Karl, M., Leck, C., Rad, F. M., Bäcklund, A., Lopez-Aparicio, S., and Heintzenberg, J.: New insights in sources of the sub-micrometre 920 aerosol at Mt. Zeppelin observatory (Spitsbergen) in the year 2015, *Tellus, Series B: Chemical and Physical Meteorology*, 71, 1–29, <https://doi.org/10.1080/16000889.2019.1613143>, 2019.

Karlsson, L. and Zieger, P.: Aerosol particle number size distribution data collected during the Arctic Ocean 2018 expedition [data set], <https://doi.org/10.17043/oden-ao-2018-aerosol-dmps-1>, 2020.

Karlsson, L., Krejci, R., Koike, M., Ebelt, K., and Zieger, P.: A long-term study of cloud residuals from low-level Arctic clouds, *Atmospheric 925 Chemistry and Physics*, 21, 8933–8959, <https://doi.org/10.5194/acp-21-8933-2021>, 2021.

Karlsson, L., Baccarini, A., Duplessis, P., Baumgardner, D., Brooks, I. M., Chang, R. Y., Dada, L., D allenbach, K. R., Heikkinen, L., Krejci, R., Leaitch, W. R., Leck, C., Partridge, D. G., Salter, M. E., Wernli, H., Wheeler, M. J., Schmale, J., and Zieger, P.: Physical and Chemical

Properties of Cloud Droplet Residuals and Aerosol Particles During the Arctic Ocean 2018 Expedition, *Journal of Geophysical Research: Atmospheres*, 127, <https://doi.org/10.1029/2021jd036383>, 2022.

930 Kay, J. E., L'Ecuyer, T., Chepfer, H., Loeb, N., Morrison, A., and Cesana, G.: Recent Advances in Arctic Cloud and Climate Research, <https://doi.org/10.1007/s40641-016-0051-9>, 2016.

Kerminen, V.-M. and Kulmala, M.: Analytical formulae connecting the "real" and the "apparent" nucleation rate and the nuclei number concentration for atmospheric nucleation events, *Aerosol Science*, 33, 609–622, www.elsevier.com/locate/jaerosci, 2002.

Koike, M., Ukita, J., Strom, J., Tunved, P., Shiobara, M., Vitale, V., Lupi, A., Baumgardner, D., Ritter, C., Hermansen, O., Yamada, K., and 935 Pedersen, C. A.: Year-Round In Situ Measurements of Arctic Low-Level Clouds: Microphysical Properties and Their Relationships With Aerosols, *Journal of Geophysical Research: Atmospheres*, 124, 1798–1822, [https://doi.org/https://doi.org/10.1029/2018JD029802](https://doi.org/10.1029/2018JD029802), 2019.

Komppula, M., Lihavainen, H., Kerminen, V.-M., Kulmala, M., and Viisanen, Y.: Measurements of cloud droplet activation of aerosol particles at a clean subarctic background site, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/https://doi.org/10.1029/2004JD005200>, 2005.

940 Korhonen, H., Carslaw, K. S., Spracklen, D. V., Mann, G. W., and Woodhouse, M. T.: Influence of oceanic dimethyl sulfide emissions on cloud condensation nuclei concentrations and seasonality over the remote Southern Hemisphere oceans: A global model study, *Journal of Geophysical Research Atmospheres*, 113, <https://doi.org/10.1029/2007JD009718>, 2008a.

Korhonen, H., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., and Ström, J.: A global model study of processes controlling aerosol size distributions in the Arctic spring and summer, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009114>, 945 2008b.

Kulmala, M., Lehtinen, K. E. J., and Laaksonen, A.: Atmospheric Chemistry and Physics Cluster activation theory as an explanation of the linear dependence between formation rate of 3 nm particles and sulphuric acid concentration, *Atmos. Chem. Phys.*, 6, 787–793, www.atmos-chem-phys.net/6/787/2006/, 2006.

Kupiszewski, P., Leck, C., Tjernström, M., Sjogren, S., Sedlar, J., Graus, M., Müller, M., Brooks, B., Swietlicki, E., Norris, S., and Hansel, A.: 950 Vertical profiling of aerosol particles and trace gases over the central Arctic Ocean during summer, *Atmospheric Chemistry and Physics*, 13, 12 405–12 431, <https://doi.org/10.5194/acp-13-12405-2013>, 2013.

Lawler, M. J., Saltzman, E. S., Karlsson, L., Zieger, P., Salter, M., Baccarini, A., Schmale, J., and Leck, C.: New Insights Into the Composition and Origins of Ultrafine Aerosol in the Summertime High Arctic, *Geophysical Research Letters*, 48, e2021GL094395, <https://doi.org/https://doi.org/10.1029/2021GL094395>, e2021GL094395 2021GL094395, 2021.

955 Leck, C. and Bigg, E. K.: Aerosol production over remote marine areas - A new route, *Geophysical Research Letters*, 26, 3577–3580, <https://doi.org/10.1029/1999gl010807>, 1999.

Leck, C. and Bigg, E. K.: New particle formation of marine biological origin, vol. 44, pp. 570–577, <https://doi.org/10.1080/02786826.2010.481222>, 2010.

Leck, C. and Svensson, E.: Importance of aerosol composition and mixing state for cloud droplet activation over the Arctic pack ice in 960 summer, *Atmospheric Chemistry and Physics*, 15, 2545–2568, <https://doi.org/10.5194/acp-15-2545-2015>, 2015.

Leck, C., Mattrai, P., Perttu, A.-M., and Gårdfeldt, K.: Expedition report: SWEDARTIC Arctic Ocean 2018, Tech. rep., 2020.

Leck, C., Sedlar, J., Swietlicki, E., Sjögren, S., Brooks, B., and Norris, S.: Vertical stratification of submicrometer aerosol particles measured during the high-Arctic ASCOS expedition 2008 [dataset], <https://doi.org/10.17043/oden-ascos-2008-aerosol-stratification-1>, 2022.

965 Loewe, K., Ekman, A. M. L., Paukert, M., Sedlar, J., Tjernström, M., and Hoose, C.: Modelling micro- and macrophysical contributors to the dissipation of an Arctic mixed-phase cloud during the Arctic Summer Cloud Ocean Study (ASCOS), *Atmospheric Chemistry and Physics*, 17, 6693–6704, <https://doi.org/10.5194/acp-17-6693-2017>, 2017.

970 Lubin, D. and Vogelmann, A. M.: A climatologically significant aerosol longwave indirect effect in the Arctic, *Nature*, 439, 453–456, <https://doi.org/10.1038/nature04449>, 2006.

975 Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J., and Johnson, C. E.: Description and evaluation of GLOMAP-mode: a modal global aerosol microphysics model for the UKCA composition-climate model, *Geoscientific Model Development*, 3, 519–551, <https://doi.org/10.5194/gmd-3-519-2010>, 2010.

980 Marle, M. J. V., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A. L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and Werf, G. R. V. D.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), *Geoscientific Model Development*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.

985 Mauritsen, T., Sedlar, J., Tjernström, M., Leck, C., Martin, M., Shupe, M., Sjogren, S., Sierau, B., Persson, P. O. G., Brooks, I. M., and Swietlicki, E.: An Arctic CCN-limited cloud-aerosol regime, *Atmospheric Chemistry and Physics*, 11, 165–173, <https://doi.org/10.5194/acp-11-165-2011>, 2011.

990 Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Atmospheric Chemistry and Physics Impact of nucleation on global CCN, *Atmos. Chem. Phys.*, 9, 8601–8616, www.atmos-chem-phys.net/9/8601/2009/, 2009.

Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S., Weingartner, E., Riipinen, I., Kulmala, M., Spracklen, D. V., Carslaw, K. S., and Baltensperger, U.: Evidence for the role of organics in aerosol particle formation under atmospheric conditions, *Proceedings of the National Academy of Sciences of the United States of America*, 107, 6646–6651, <https://doi.org/10.1073/pnas.0911330107>, 2010.

995 Morgenstern, O., Braesicke, P., O’Connor, F. M., Bushell, A. C., Johnson, C. E., Osprey, S. M., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model-Part 1: The stratosphere, *Geosci. Model Dev.*, 2, 43–57, www.geosci-model-dev.net/2/43/2009/, 2009.

Mulcahy, J. P., Johnson, C., Jones, C. G., Povey, A. C., Scott, C. E., Sellar, A., Turnock, S. T., Woodhouse, M. T., Abraham, N. L., Andrews, M. B., Bellouin, N., Browse, J., Carslaw, K. S., Dalvi, M., Folberth, G. A., Glover, M., Grosvenor, D. P., Hardacre, C., Hill, R., Johnson, B., Jones, A., Kipling, Z., Mann, G., Molland, J., O’Connor, F. M., Palmiéri, J., Reddington, C., Rumbold, S. T., Richardson, M., Schutgens, N. A., Stier, P., Stringer, M., Tang, Y., Walton, J., Woodward, S., and Yool, A.: Description and evaluation of aerosol in UKESM1 and HadGEM3-GC3.1 CMIP6 historical simulations, *Geoscientific Model Development*, 13, 6383–6423, <https://doi.org/10.5194/gmd-13-6383-2020>, 2020.

Notz, D. and Stroeve, J.: The Trajectory Towards a Seasonally Ice-Free Arctic Ocean, *Current Climate Change Reports*, 4, 407–416, <https://doi.org/10.1007/s40641-018-0113-2>, 2018.

990 O’Connor, F. M., Johnson, C. E., Morgenstern, O., Abraham, N. L., Braesicke, P., Dalvi, M., Folberth, G. A., Sanderson, M. G., Telford, P. J., Voulgarakis, A., Young, P. J., Zeng, G., Collins, W. J., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model-Part 2: The troposphere, *Geoscientific Model Development*, 7, 41–91, <https://doi.org/10.5194/gmd-7-41-2014>, 2014.

Pastore, M. and Calcagnì, A.: Measuring distribution similarities between samples: A distribution-free overlapping index, *Frontiers in Psychology*, 10, <https://doi.org/10.3389/fpsyg.2019.01089>, 2019.

1000 Pöhlker, M. L., Zhang, M., Campos Braga, R., Krüger, O. O., Pöschl, U., and Ervens, B.: Aitken mode particles as CCN in aerosol- and updraft-sensitive regimes of cloud droplet formation, *Atmospheric Chemistry and Physics*, 21, 11 723–11 740, <https://doi.org/10.5194/acp-21-11723-2021>, 2021.

Randerson, J. T., Liu, H., Flanner, M. G., Chambers, S. D., Jin, Y., Hess, P. G., Pfister, G., Mack, M. C., Treseder, K. K., Welp, L. R., Chapin, F. S., Harden, J. W., Goulden, M. L., Lyons, E., Neff, J. C., Schuur, E. A., and Zender, C. S.: The impact of boreal forest fire on climate warming, *Science*, 314, 1130–1132, <https://doi.org/10.1126/science.1132075>, 2006.

1005 Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Communications Earth & Environment*, 3, 168, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.

Schmale, J. and Baccarini, A.: Progress in Unraveling Atmospheric New Particle Formation and Growth Across the Arctic, *Geophysical Research Letters*, 48, <https://doi.org/https://doi.org/10.1029/2021GL094198>, 2021.

1010 Schmale, J., Zieger, P., and Ekman, A. M.: Aerosols in current and future Arctic climate, *Nature Climate Change*, 11, 95–105, <https://doi.org/10.1038/s41558-020-00969-5>, 2021.

Schmale, J., Sharma, S., Decesari, S., Pernov, J., Massling, A., Hansson, H.-C., von Salzen, K., Skov, H., Andrews, E., Quinn, P. K., Upchurch, L. M., Eleftheriadis, K., Traversi, R., Gilardoni, S., Mazzola, M., Laing, J., and Hopke, P.: Pan-Arctic seasonal cycles and long-term trends of aerosol properties from 10 observatories, *Atmospheric Chemistry and Physics*, 22, 3067–3096, <https://doi.org/10.5194/acp-22-3067-2022>, 2022.

1015 Sedlar, J., Tjernström, M., Mauritsen, T., Shupe, M. D., Brooks, I. M., Persson, P. O. G., Birch, C. E., Leck, C., Sirevaag, A., and Nicolaus, M.: A transitioning Arctic surface energy budget: the impacts of solar zenith angle, surface albedo and cloud radiative forcing, *Climate Dynamics*, 37, 1643–1660, <https://doi.org/10.1007/s00382-010-0937-5>, 2011.

Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, 1020 S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahaan, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *Journal of Advances in Modeling Earth Systems*, 11, 4513–4558, 1025 <https://doi.org/https://doi.org/10.1029/2019MS001739>, 2019.

Sharma, S., Barrie, L. A., Magnusson, E., Brattstrom, G., Leaitch, W. R., Steffen, A., and Landsberger, S.: A Factor and Trends Analysis of Multidecadal Lower Tropospheric Observations of Arctic Aerosol Composition, Black Carbon, Ozone, and Mercury at Alert, Canada, *Journal of Geophysical Research: Atmospheres*, 124, 14 133–14 161, <https://doi.org/10.1029/2019JD030844>, 2019.

1030 Shepherd, A., Ivins, E. R., A, G., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Jougin, I., King, M. A., Lenaerts, J. T. M., Li, J., Ligtenberg, S. R. M., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R., Glenn, M., Mouginot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, E., Rott, H., Sørensen, L. S., Scambos, T. A., Scheuchl, B., Schrama, E. J. O., Smith, B., Sundal, A. V., van Angelen, J. H., van de Berg, W. J., van den Broeke, M. R., Vaughan, D. G., Velicogna, I., Wahr, J., Whitehouse, P. L., Wingham, D. J., Yi, D., Young, D., and Zwally, H. J.: A Reconciled Estimate of Ice-Sheet Mass Balance, *Science*, 338, <https://doi.org/10.1126/science.1228102>, 2012.

1035 Shupe, M. D.: Clouds at Arctic Atmospheric Observatories. Part II: Thermodynamic Phase Characteristics, *Journal of Applied Meteorology and Climatology*, 50, 645–661, <https://doi.org/10.1175/2010JAMC2468.1>, 2011.

Shupe, M. D. and Intrieri, J. M.: Cloud Radiative Forcing of the Arctic Surface: The Influence of Cloud Properties, Surface Albedo, and Solar Zenith Angle, *Journal of Climate*, 17, 616 – 628, [https://doi.org/10.1175/1520-0442\(2004\)017<0616:CRFOTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0616:CRFOTA>2.0.CO;2), 2004.

Shupe, M. D., Persson, P. O. G., Brooks, I. M., Tjernström, M., Sedlar, J., Mauritzen, T., Sjogren, S., and Leck, C.: Cloud and boundary layer interactions over the Arctic sea ice in late summer, *Atmospheric Chemistry and Physics*, 13, 9379–9399, <https://doi.org/10.5194/acp-13-9379-2013>, 2013.

Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrakou, T., Muller, J.-F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmospheric Chemistry and Physics*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014.

1045 Sipila, M., Sarnela, N., Jokinen, T., Henschel, H., Junninen, H., Kontkanen, J., Richters, S., Kangasluoma, J., Franchin, A., Perakyla, O., Rissanen, M. P., Ehn, M., Vehkamaki, H., Kurten, T., Berndt, T., Petaja, T., Worsnop, D., Ceburnis, D., Kerminen, V. M., Kulmala, M., and O'Dowd, C.: Molecular-scale evidence of aerosol particle formation via sequential addition of HIO_3 , *Nature*, 537, 532–534, <https://doi.org/10.1038/nature19314>, 2016.

Solomon, A., Shupe, M. D., Persson, P. O. G., and Morrison, H.: Moisture and dynamical interactions maintaining decoupled Arctic mixed-phase stratocumulus in the presence of a humidity inversion, *Atmospheric Chemistry and Physics*, 11, 10127–10148, <https://doi.org/10.5194/acp-11-10127-2011>, 2011.

1050 Sotiropoulou, G., Sedlar, J., Tjernström, M., Shupe, M. D., Brooks, I. M., and Persson, P. O. G.: The thermodynamic structure of summer Arctic stratocumulus and the dynamic coupling to the surface, *Atmospheric Chemistry and Physics*, 14, 12573–12592, <https://doi.org/10.5194/acp-14-12573-2014>, 2014.

1055 Sotiropoulou, G., Sedlar, J., Forbes, R., and Tjernström, M.: Summer Arctic clouds in the ECMWF forecast model: an evaluation of cloud parametrization schemes, *Quarterly Journal of the Royal Meteorological Society*, 142, 387–400, <https://doi.org/10.1002/qj.2658>, 2016.

Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Mann, G. W., and Sihto, S.-L.: *Atmospheric Chemistry and Physics*, European Geosciences Union, 6, 5631–5648, www.atmos-chem-phys.net/6/5631/2006/, 2006.

1060 Stevens, R. G., Loewe, K., Dearden, C., Dimitrelos, A., Possner, A., Eirund, G. K., Raatikainen, T., Hill, A. A., Shipway, B. J., Wilkinson, J., Romakkaniemi, S., Tonttila, J., Laaksonen, A., Korhonen, H., Connolly, P., Lohmann, U., Hoose, C., Ekman, A. M. L., Carslaw, K. S., and Field, P. R.: A model intercomparison of CCN-limited tenuous clouds in the high Arctic, *Atmospheric Chemistry and Physics*, 18, 11041–11071, <https://doi.org/10.5194/acp-18-11041-2018>, 2018.

Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006888>, 2006.

1065 Ström, J., Umegård, J., Tørseth, K., Tunved, P., Hansson, H.-C., Holmén, K., Wismann, V., Herber, A., and König-Langlo, G.: One year of particle size distribution and aerosol chemical composition measurements at the Zeppelin Station, Svalbard, March 2000–March 2001, *Physics and Chemistry of the Earth, Parts A/B/C*, 28, 1181 – 1190, <https://doi.org/https://doi.org/10.1016/j.pce.2003.08.058>, 2003.

Struthers, H., Ekman, A. M. L., Glantz, P., Iversen, T., Kirkevåg, A., Mårtensson, E. M., Seland, Ø., and Nilsson, E. D.: The effect of sea ice loss on sea salt aerosol concentrations and the radiative balance in the Arctic, *Atmospheric Chemistry and Physics*, 11, 3459–3477, <https://doi.org/10.5194/acp-11-3459-2011>, 2011.

1070 Tjernström, M., Leck, C., Birch, C. E., Bottenheim, J. W., Brooks, B. J., Brooks, I. M., Bäcklin, L., Chang, R. Y., Leeuw, G. D., Liberto, L. D., Rosa, S. D. L., Granath, E., Graus, M., Hansel, A., Heintzenberg, J., Held, A., Hind, A., Johnston, P., Knulst, J., Martin, M., Matrai, P. A., Mauritzen, T., Müller, M., Norris, S. J., Orellana, M. V., Orsini, D. A., Paatero, J., Persson, P. O., Gao, Q., Rauschenberg, C., Ristovski, Z., Sedlar, J., Shupe, M. D., Sierau, B., Sirevaag, A., Sjogren, S., Stetzer, O., Swietlicki, E., Szczodrak, M., Vaattovaara, P., Wahlberg, N., Westberg, M., and Wheeler, C. R.: The Arctic Summer Cloud Ocean Study (ASCOS): Overview and experimental design, *Atmospheric Chemistry and Physics*, <https://doi.org/10.5194/acp-14-2823-2014>, 2014.

Tunved, P., Ström, J., and Krejci, R.: Arctic aerosol life cycle: linking aerosol size distributions observed between 2000 and 2010 with air mass transport and precipitation at Zeppelin station, Ny-Ålesund, Svalbard, *Atmospheric Chemistry and Physics*, 13, 3643–3660, <https://doi.org/10.5194/acp-13-3643-2013>, 2013.

1080 Vehkamäki, H., Kulmala, M., Napari, I., Lehtinen, K. E. J., Timmreck, C., Noppel, M., and Laaksonen, A.: An improved parameterization for sulfuric acid–water nucleation rates for tropospheric and stratospheric conditions, *Journal of Geophysical Research*, 107, 4622, <https://doi.org/10.1029/2002JD002184>, 2002.

Vüller, J., Achtert, P., Brooks, I., Tjernström, M., Prytherch, J., and III, R. N.: Meteorological and cloud conditions during the Arctic Ocean 2018 expedition, *Atmospheric Chemistry and Physics*, pp. 1–43, <https://doi.org/10.5194/acp-2020-219>, 2020.

1085 Walker, X. J., Baltzer, J. L., Cumming, S. G., Day, N. J., Ebert, C., Goetz, S., Johnstone, J. F., Potter, S., Rogers, B. M., Schuur, E. A., Turetsky, M. R., and Mack, M. C.: Increasing wildfires threaten historic carbon sink of boreal forest soils, *Nature*, 572, 520–523, <https://doi.org/10.1038/s41586-019-1474-y>, 2019.

1090 Walters, D., Baran, A. J., Boutle, I., Brooks, M., Earnshaw, P., Edwards, J., Furtado, K., Hill, P., Lock, A., Manners, J., Morcrette, C., Mulcahy, J., Sanchez, C., Smith, C., Stratton, R., Tennant, W., Tomassini, L., Weverberg, K. V., Vosper, S., Willett, M., Browse, J., Bushell, A., Carslaw, K., Dalvi, M., Essery, R., Gedney, N., Hardiman, S., Johnson, B., Johnson, C., Jones, A., Jones, C., Mann, G., Milton, S., Rumbold, H., Sellar, A., Ujjie, M., Whitall, M., Williams, K., and Zerroukat, M.: The Met Office Unified Model Global Atmosphere 7.0/7.1 and JULES Global Land 7.0 configurations, *Geoscientific Model Development*, 12, 1909–1963, <https://doi.org/10.5194/gmd-12-1909-2019>, 2019.

Willis, M. D., Kollner, F., Burkart, J., Bozem, H., Thomas, J. L., Schneider, J., Aliabadi, A. A., Hoor, P. M., Schulz, H., Herber, A. B., 1095 Leaitch, W. R., and Abbatt, J. P.: Evidence for marine biogenic influence on summertime Arctic aerosol, *Geophysical Research Letters*, 44, 6460–6470, <https://doi.org/10.1002/2017GL073359>, 2017.

Willis, M. D., Leaitch, W. R., and Abbatt, J. P.: Processes Controlling the Composition and Abundance of Arctic Aerosol, *Reviews of Geophysics*, 56, 621–671, <https://doi.org/10.1029/2018RG000602>, 2018.

1100 Zhang, R., Wang, H., Fu, Q., Rasch, P. J., and Wang, X.: Unraveling driving forces explaining significant reduction in satellite-inferred Arctic surface albedo since the 1980s, *Proceedings of the National Academy of Sciences of the United States of America*, 116, 23 947–23 953, <https://doi.org/10.1073/pnas.1915258116>, 2019.