

Differential Temperature Sensitivity of Intracellular Metabolic Processes and Extracellular Soil Enzyme Activities.

Adetunji Alex Adekanmbi^{1,2}, Laurence Dale¹, Liz Shaw¹, and Tom Sizmur¹

5 ¹Department of Geography and Environmental Science, University of Reading, Reading, RG6 6DW, UK

²Department of Soil Science and Land Management, Federal University of Technology, PMB 65, 920001, Minna, Nigeria.

Correspondence to: Tom Sizmur (t.sizmur@reading.ac.uk)

10 **Abstract.** Predictions concerning the feedback of soil heterotrophic respiration to a warming climate often do not differentiate between the extracellular and intracellular steps involved in soil organic matter decomposition. This study examined the temperature sensitivities of intracellular metabolic processes and extracellular soil enzyme activities and how they are influenced by previous temperatures. We pre-incubated soils at 5 °C, 15 °C or 26 °C to acclimatise the microbial communities to different thermal regimes for 60 days before measuring potential activities of β -glucosidase and chitinase (extracellular enzymes), glucose-induced respiration (intracellular metabolic processes), and basal respiration at a range of assay temperatures (5 °C, 15 °C, 26 °C, 37 °C, and 45 °C). A higher pre-incubation temperature decreased soil pH and C/N ratio and decreased β -glucosidase potential activity and respiration, but not chitinase potential activity. It is likely that this legacy effect on β -glucosidase and respiration is an indirect effect of substrate depletion rather than physiological acclimation or genetic adaptation. Pre-incubation temperature effects on temperature sensitivity were subtle and restricted to extracellular activities, perhaps because of the short (60 day) duration of the pre-incubation at temperatures that were below the initial optimum (~30 °C) for the mesophilic soil community. However, we found that the intracellular and extracellular steps differ in their temperature sensitivity and this observation differs depending on the range of temperature used for Q10 estimates of temperature sensitivity. Between 5 °C and 15 °C intracellular and extracellular enzyme processes show equal temperature sensitivity, but between 15 °C and 26 °C intracellular metabolic processes were more temperature sensitive than extracellular enzyme activity and between 26 °C and 37 °C extracellular enzyme activity was more temperature sensitive than intracellular metabolic processes. This result implies that depolymerisation of higher molecular weight carbon is more sensitive to temperature changes at higher temperatures (e.g. higher temperatures on extremely warm days) but the respiration of the generated monomers is more sensitive to temperature changes at moderate temperatures (e.g. mean daily maximum soil temperature). However, studies using multiple soil types and a greater range of pre-incubation temperatures are required to generalize our results. Nevertheless, since climate change predictions currently indicate that there will be a greater frequency and severity of hot summers and heatwaves, it is possible that global warming may reduce the importance of extracellular depolymerisation relative to intracellular metabolic processes as the rate limiting step of soil organic matter mineralization. We conclude that extracellular and intracellular steps are not equally sensitive to changes in soil temperature and that the

35 previous temperature a soil is exposed to may influence the potential activity, but not temperature sensitivity, of extracellular and intracellular processes.

1 Introduction

40 Understanding the temperature sensitivity of soil organic matter (SOM) decomposition will help predict how soils might respond to climate change. There are two major enzymatically-mediated steps involved in the decomposition of SOM to produce CO₂ (Bárta et al., 2013; Maire et al., 2013; Blagodatskaya et al., 2016). The first step, extracellular depolymerisation, requires extracellular enzymes of microbial (and also plant and animal) origin to depolymerize macromolecular constituents of SOM and produce soluble low molecular weight microbial substrates (Maire et al., 2013). The second step, intracellular metabolism, results in the release of CO₂ after substrates are absorbed and catabolised by microbial cells, involving a multitude of intracellular processes.

45 Many ecological studies have examined the temperature sensitivity of SOM decomposition, but most of them measure the end product as respired CO₂ (e.g. Wang et al., 2013) or mass loss of C substrate (e.g. Kirwan et al., 2014), which does not differentiate between the temperature sensitivity of contributing extracellular and intracellular processes. Temperature sensitivity, defined as the rate of change in reaction rate with respect to temperature, is the first derivative of the relationship between temperature and reaction rate (Alster et al., 2020). Temperature-rate relationships are typically unimodal, reflecting rising reaction rates with temperature due to thermodynamic effects and then a decline in rate with further increase in temperature related to thermal effects on enzyme activation and ultimately, denaturation (Alster et al., 2020). Parameters describing the temperature-rate relationship have been shown to vary both with respect to extracellular enzyme type and, between microbial taxa for the same enzyme type (Alster et al., 2016). Extracellular enzyme activity, rather than intracellular metabolic processes, is widely thought to be the rate-limiting step for respiration of organic matter in soils (Jan et al., 2009; Bradford, 2013), but very few studies have explicitly compared the temperature sensitivity of extracellular and intracellular processes to understand how each step might respond to increases in temperature and whether the magnitude of dependence of intracellular catabolism and CO₂ respiration on extracellular enzyme activities for supply of substrate increases or decreases with increasing temperature. Ultimately, this lack of information limits our predictive understanding of how the soil carbon cycle will respond to future global temperature changes (Blagodatskaya et al., 2016).

As already stated, decomposition of SOM is a function of the heterotrophic microbial community and the extracellular enzymes it produces. If the microbial community and its enzyme production adapts to warming (or cooling), this might result in variation in the size of microbial enzyme (and biomass) pools (Fanin et al., 2022). In addition, thermal adaptation of the microbial community may, for a given enzyme-catalysed reaction, modulate the temperature- reaction rate relationship (e.g.

manifest as a shift in the temperature optimum of reaction rates) and thus temperature sensitivity of extra- and intra-cellular processes depending on soil thermal history (Wallenstein et al., 2010). Adaptation at the level of the microbial community may be through acclimation (phenotypic or physiological change to respond to thermal regime, including production of different isozymes within taxa), evolutionary changes within taxa leading to novel isozymes, or, species sorting where taxa (including their enzyme systems) already better adapted to a certain temperature competitively exclude those less adapted (Birgander et al., 2013). Whether adaptive processes modulate the activity and temperature response relationship to the same extent for intracellular and extracellular processes is not known. Since extracellular enzymes catalyse what is believed to be the rate limiting step in SOM decomposition (Duly and Nannipieri, 1998; Alvarez et al., 2018), any thermal adaptation of extracellular enzymes will then determine how much substrate is available for subsequent uptake and respiration and represents an important control on the response of ecosystems to warming (Bradford, 2013).

In this study, we measured potential extracellular enzyme activity and glucose-induced respiration, as a proxy for intracellular metabolic processes, at 5 assay temperatures (5 °C, 15 °C, 26 °C, 37 °C and 45 °C) following pre-incubation for 60 days at 5 °C, 15 °C, or 26 °C. The aim was to compare the temperature sensitivity of the extra- and intracellular steps of organic matter decomposition in soils that have previously been incubated at different thermal regimes, alongside measurements of key soil properties that we consider may lead to changes in potential enzyme activity. The pre-incubation temperatures were selected to be realistic for the site where the soil was sampled. We hypothesise that (i) extracellular and intracellular processes are not equally sensitive in their response to increasing temperature, given the involvement of different enzymes and (for intracellular catabolism) biochemical networks, and that (ii) extracellular and intracellular processes, and their temperature sensitivity, are influenced by pre-incubation temperature due to thermal adaptation of the soil microbial community.

2 Methodology

2.1 Soil sampling and pre-incubation

Soils samples were collected from a depth of 3 – 10 cm from a permanent grassland field at Sonning UK (latitude 51° 28.564', longitude 000° 54.198'), sieved with a 4 mm sieve, mixed, and homogenised before randomly allocating to replicates. Four 'field moist' (soil moisture content = 0.13 g H₂O g soil⁻¹) replicate sub-samples (750 g) were pre-incubated at 5 °C, 15 °C (similar to the mean daily minimum (6.8 °C) and the mean daily maximum (15.0 °C) temperatures measured at the University of Reading Atmospheric Observatory, close to the sampling location, between 2009 and 2019), and 26 °C (typical of a temperature measured on a warm summer day at the University of Reading Atmospheric Observatory; Figure S-1) for a period of 60 days in plastic containers with the cover of each container loosely closed. Soil moisture content was adjusted to the initial field moist condition every two weeks for soils incubated at 5 °C and 15 °C and weekly for soils incubated at 26 °C. The soil is a slightly acidic loamy soil, classified as Chromic Endoskeletal Luvisol. A detailed description of the site is provided by Adekanmbi et al., (2020).

2.2 Experimental design

100 The experimental design included 3 pre-incubation temperatures (5 °C, 15 °C, and 26 °C), replicated 4 times, resulting in 12 experimental units. At the end of the 60-day pre-incubation period, soils were subsampled for determination of basal respiration and substrate induced respiration using glucose as the substrate (Section 2.3), and the potential activity of β -glucosidase (β -1,4-glucosidase) and chitinase (N-acetyl β – D – glycosaminidase) extracellular enzymes (Section 2.4) all measured at 5 assay temperatures (5 °C, 15 °C, 26 °C, 37 °C and 45 °C). Assays were performed on all experimental units within the same week
105 to minimise variability due to time of assay. Incubation temperatures were randomised to prevent systematic bias in the results. A portion of the soil from each replicate sample was also analysed for total C, total N, pH, and microbial biomass carbon (Section 2.5).

2.3 Basal and substrate-induced respiration

For each replicate sample, 15 g (13.31 g dry weight equivalent) of soil was weighed into a 50 ml centrifuge tube. Glucose
110 solution (2 ml) was added at concentrations of 0 (deionized water only) or 10 mg g⁻¹ soil (an initially saturating concentration of glucose as determined in a preliminary experiment; supplementary information), thus bringing the soil to 58 % of its water holding capacity. The soil was then mixed to distribute the solution throughout. Following soil-substrate mixing, the tube was ventilated by blowing in lab air with a 20 ml syringe. The tubes were then sealed with septum stoppers and 15 ml of lab air was injected. The headspace was flushed by moving the syringe plunger up and down several times before sampling 15 ml of
115 head space gas (as the T0 sample) and injecting into an evacuated 12 ml exetainer vial, creating overpressure, using a tap and needle attached to the syringe. Soil samples were incubated for one hour at either 5 °C, 15 °C, 26 °C, 37 °C or 45 °C. It is likely that during the first few minutes of the assay the soils were changing temperature from room temperature to the assay temperature. At the end of the incubation, the process of injecting air, flushing, and sampling was repeated (T1 sample). Headspace gas samples were stored at 20 °C prior to analysis by an Agilent 7890B gas chromatograph. After calibrating with
120 CO₂ gas standards, the concentration of CO₂ in mg L⁻¹ was converted to C-CO₂ mg C g⁻¹ h as described by (Salazar-Villegas et al., 2016):

$$\text{CO}_2 \text{ (mg C g}^{-1} \text{ h}^{-1}\text{)} = \frac{V (T1 - T0)}{Wt}$$

Where V = volume of headspace in the centrifuge tube; T1 is CO₂ concentration after a 1 hour incubation in mg L⁻¹; T0 is CO₂
125 concentration before 1 hour incubation in mg L⁻¹, W is the dry weight of the soil, and t is the time between T0 and T1 measurements in hours.

2.4 Extracellular enzyme assays

Extracellular enzyme assay methods were based on Eivazi and Tabatabai, (1988) and Parham and Deng, (2000) for β -1,4-glucosidase (β -glucosidase) and N-acetyl β – D – glycosaminidase (chitinase), respectively. For each experimental replicate,

130 1 g of soil was weighed into a 50 ml centrifuge tube and mixed with 4ml of room temperature pre-incubated 4-methylumbelliferone (MUB) buffer (pH 6) and either 1ml 25mM *p*-nitrophenyl- β -D-glucopyranoside or 10 mM *p*-nitrophenyl-N-acetyl-b-D-glucosaminide solution, to assess β -glucosidase and chitinase activity, respectively. Samples were incubated at 5 °C, 15 °C, 26 °C, 37 °C, or 45°C for 30 minutes, after which 1 ml 0.5 M CaCl₂ and 4 ml Tris buffer (pH 12) was added to stop the reaction. It is likely that during the first few minutes of the assay the soils were changing temperature from room
135 temperature to the assay temperature. Samples were mixed by swirling, then filtered with Whatman No. 2 filter paper. Additionally, 2 blanks (for each run) were created by adding substrate to tubes containing the mixture after the reaction had stopped. Colour intensity of the filtrate was measured using a spectrophotometer at 400 nm and blank-corrected sample absorbance converted to $\mu\text{g } p\text{-nitrophenol reaction}^{-1}$ using *p*-nitrophenol standard solutions (0, 10, 20, 30, 40 and 50 $\mu\text{g } p\text{-nitrophenol}$). Potential enzyme activities were expressed as $\mu\text{g } p\text{-nitrophenol g}^{-1}$ dry soil h^{-1} . The 30-minute assay incubation
140 time was within the time range where product accumulation was linear with incubation time, according to a preliminary experiment (see supplementary material).

2.5 Measurement of total carbon, total nitrogen, pH, and microbial biomass carbon

Microbial biomass carbon was measured using the fumigation/extraction method described by Vance et al. (1987). Four replicates from each pre-incubation temperature were weighed to the moist mass equivalent to 50 g oven-dried soil in beakers
145 and placed in a vacuum desiccator lined with damp paper towel to ensure high humidity, along with a beaker containing about 50 ml ethanol-free chloroform and several anti- bumping granules. The desiccator was evacuated, and the chloroform allowed to boil for two minutes before the valve was closed and the desiccator kept in the dark for 24 hours. Before extraction, the chloroform was removed, the desiccator evacuated three times and the samples left to vent to ensure no chloroform remained in the soil.

150 Extraction was carried out on both fumigated soil and non-fumigated duplicates. Samples of both were placed into 350 ml polypropylene bottles, to which 200 ml 0.5 M K₂SO₄ was added, before being placed on an oscillating shaker for 30 minutes. The suspension was then filtered into polypropylene universal tubes before being stored in a freezer prior to analysis. After removal from the freezer, samples were diluted by a factor of 10, and filtered to remove CaSO₄ that had precipitated, before
155 analysis for total organic carbon (TOC) using a Shimadzu TOC 5000. Also analysed were method blanks consisting of K₂SO₄ that had not been used to extract soil, to correct for any part of the reading not due to organic carbon content. TOC extracted from fumigated and non-fumigated samples was converted to a biomass carbon value by multiplying the difference (E_c) by 2.64, following Vance et al., (1987). The TOC of the non-fumigated soil before conversion represents the K₂SO₄ extractable carbon.

160 Total C and N were determined using the dry combustion method. 2 mm sieved soil samples were ground for three minutes in an agate ball mill. From the residue, 10 mg duplicates were weighed out using a five-point balance and placed in tin foil

capsules for measurement. C and N concentrations were analysed using a C/N Elemental Analyser (Thermo Flash 2000 EA). The C/N ratio was calculated from total C and N.

165

pH was determined in water (10 g air-dried soil: 25ml deionised water) following end-over-end shaking (30 rpm, 15 mins) and using a calibrated (pH 4.0 and pH 7.0) pH meter.

2.6 Temperature sensitivity

170 Temperature sensitivity (Q10) of both the intra (glucose induced respiration) and extra-cellular (chitinase and β -glucosidase) enzyme activities was calculated using the equal time measurement method, as described by Karhu et al., (2014). Q10 was calculated at three temperature ranges (Q10_{5-15°C}, Q10_{15-26°C}, and Q10_{26-37°C}). The primary reason why we calculated Q10 at different temperatures is because we found that temperature sensitivity was different at different temperature ranges. This meant that there was not a good linear relationship between the natural log of enzyme intracellular metabolic processes or
175 extracellular enzyme activity and temperature, apart for β -glucosidase, as demonstrated in the supplementary material. Similar range of incubation temperatures was used by Wang et al., (2013). Arrhenius enzyme activation energy (Ea) was calculated from the slope of the relationship between $-1/R_0T$ and the natural logarithm of rate of enzyme activity (R_0 = the gas universal constant: 8.314 J mol⁻¹; T = temperature in Kelvin), as described by Li et al., (2015). Ea was calculated using assays within the 5 °C – 26 °C temperature range for all four assays to ensure that the data used to calculate Ea conformed to the Arrhenius
180 functional form (Schulte, 2015).

2.7 Statistical analysis

Two-way Analysis of Variance (ANOVA) was carried out to assess the effects of pre-incubation temperature and assay temperature on basal respiration, glucose-induced respiration, and extracellular enzyme activities. We also assessed whether
185 intracellular and extracellular steps were equally sensitive to temperature, and whether this was influenced by pre-incubation temperature, by performing a two-way ANOVA on the Ea and Q10 values using assay type, and pre-incubation temperature as factors. One way ANOVA was carried out to assess the effect of pre-incubation temperature on soil properties. ANOVA was performed in Minitab version 18. Tukey pairwise comparisons were used to assess the significance of differences between individual treatment means.

190 3 Results

3.1 Impact of pre-incubation temperature on selected soil properties

The effects of soil pre-incubation temperature on soil total C, total N, C/N ratio, pH and microbial biomass carbon are presented in Figure 1. Pre-incubation temperature did not have a statistically significant impact on C ($P = 0.641$) or N ($P = 0.439$). However, soil C/N ratio was significantly ($P < 0.05$) higher in soil pre-incubated at 15 °C and 5 °C, compared to soil pre-
195 incubated at 26 °C. Also, pre-incubation temperature significantly ($P < 0.05$) influenced soil pH which decreased in the order 5 °C > 15 °C > 26 °C. There was no statistically significant effect of soil pre-incubation temperature on soil microbial biomass ($P = 0.206$).

3.2 Responses of intracellular and extracellular processes to pre-incubation temperature and assay temperature.

200 The influence of pre-incubation temperature on the potential activities of β -glucosidase (Figure 2A) and chitinase (Figure 2B) extracellular enzymes, the rate of glucose-induced respiration (representing intracellular metabolic processes) (Figure 2C), and the basal respiration rate (Figure 2D) across the full range of assay temperatures (5 °C to 45 °C) are presented in Figure 2.

Both the pre-incubation temperature ($P < 0.0001$), assay temperature ($P < 0.0001$) and their interaction ($P = 0.001$) significantly
205 influenced potential β -glucosidase activity in soil. Soils pre-incubated at 26 °C had a lower potential β -glucosidase activity in soil compared to those pre-incubated at 15 °C or 5 °C. Increasing assay temperature increased β -glucosidase activity up to the maximum assay temperature of 45 °C (Figure 2A). Pre-incubating soils at 15 °C resulted in significantly greater potential β -glucosidase activity at the higher assay temperatures (45 °C and 37 °C) than by pre-incubating soils at 5 °C or 26 °C.

210 Both assay temperature ($P < 0.0001$) and interaction between assay and pre-incubation temperatures ($P < 0.001$) significantly influenced potential chitinase activity, but pre-incubation temperature ($P = 0.077$) did not. Chitinase activity increased with increasing assay temperature, reaching maximum when assayed at 37 °C, but was then lower when assayed at 45 °C (Figure 2B). Pre-incubating soil at 26 °C and assaying at 37 °C resulted in a significantly ($P = 0.001$) greater chitinase activity than pre-incubating at 5 °C assaying at 37 °C. When assayed at 5 °C or 15 °C, pre-incubation at 26 °C resulted in lower chitinase
215 activities than pre-incubation at 15 °C or 5 °C.

Both the pre-incubation temperature ($P < 0.0001$), and assay temperature ($P < 0.0001$) significantly influenced glucose-induced respiration, but not their interaction ($P = 0.130$). Similarly, the pre-incubation temperature ($P = 0.001$), and assay temperature ($P < 0.0001$) significantly influenced basal respiration, but not their interactions ($P = 0.250$). With or without glucose addition,
220 pre-incubating soil at 26 °C resulted in lower soil respiration compared to pre-incubating soil at 5 °C or 15 °C (Figure 2C and 2D). Glucose-induced respiration increased with increasing assay temperature, reaching maximum between 26 °C and 37 °C, but was significantly lower at 45 °C. Basal respiration increased with increasing assay temperature up to 26 °C then declined only slightly. The addition of 10 mg g⁻¹ soil of glucose led to about a 4-fold increase in CO₂ respired, compared to no addition of glucose substrate.

225

3.3 Effect of pre-incubation temperature and assay type (intra- or extracellular) on temperature sensitivity of metabolic processes

3.3.1 Temperature coefficient (Q10)

230 The effects of pre-incubation temperature and enzyme type on $Q_{10_{5-15^{\circ}\text{C}}}$, $Q_{10_{15-26^{\circ}\text{C}}}$, and $Q_{10_{26-37^{\circ}\text{C}}}$, are presented in Figure 3. There was no overall significant effect ($P > 0.05$) of pre-incubation temperature on Q10 calculated using all three temperature intervals. There was also no significant effect of assay type ($P = 0.393$), or the interaction between assay type and pre-incubation temperature ($P = 0.700$), on $Q_{10_{5-15^{\circ}\text{C}}}$. However, $Q_{10_{15-26^{\circ}\text{C}}}$ significantly differed with assay type ($P < 0.0001$), but not for the assay type*pre-incubation interaction ($P = 0.160$). The $Q_{10_{15-26^{\circ}\text{C}}}$ was significantly lower for both extracellular

235 enzymes (chitinase and β -glucosidase) than for glucose-induced respiration or basal respiration, irrespective of pre-incubation temperature. This result indicates that intracellular metabolic processes are more temperature sensitive than extracellular enzymes in this soil between 15 °C and 26 °C. Furthermore, $Q_{10_{26-37^{\circ}\text{C}}}$ was significantly affected by assay type ($P < 0.0001$) but exhibited the opposite pattern to $Q_{10_{15-26^{\circ}\text{C}}}$. $Q_{10_{26-37^{\circ}\text{C}}}$ for chitinase activity and β -glucosidase activity were significantly ($P < 0.05$) greater than the $Q_{10_{26-37^{\circ}\text{C}}}$ for glucose-induced respiration and basal respiration. This finding indicates that

240 extracellular enzymes are more temperature sensitive than intracellular metabolic processes in this soil between 26 °C and 37 °C. $Q_{10_{26-37^{\circ}\text{C}}}$ for chitinase activity was also significantly ($P < 0.05$) greater than the $Q_{10_{26-37^{\circ}\text{C}}}$ for β -glucosidase activity. There was also a significant interaction between enzyme type and pre-incubation temperature ($P = 0.018$). Chitinase activity was less temperature sensitive when soil was pre-incubated at 26 °C compared to when pre-incubated at 15 °C or 5 °C.

245 3.3.2 Arrhenius activation energy (Ea)

The activation energy (Ea), derived from the fit of the Arrhenius equation (Figure 4) to assays performed between 5 °C and 26 °C (basal respiration and glucose-induced respiration) and between 5°C and 37°C (extracellular enzymes), differed significantly with assay type ($P < 0.0001$) and pre-incubation temperature ($P = 0.002$) and there was a significant interaction between assay type and pre-incubation temperature ($P = 0.029$). Ea increased with increasing pre-incubation temperature, with

250 soils pre-incubated at 26 °C exhibiting the highest Ea and soils pre-incubated at 5 °C exhibiting the lowest. β -glucosidase activity and chitinase activity had a significantly ($P < 0.05$) lower Ea than intracellular metabolic activity, and basal respiration, though chitinase had similar Ea with intracellular metabolic activity when soils were pre-incubated at 26 °C.

4 Discussion

Understanding whether soil intracellular and extracellular processes, which each play a distinct role in SOM decomposition,

255 are equally sensitive to temperature changes was the major motivation for this study. We also examined whether pre-incubation

temperature drives thermal adaption of the soil microbial community and results in differential alteration of temperature sensitivity of intracellular and extracellular enzyme activity. Therefore, we pre-incubated soil samples at three different temperatures to expose the soil microbial community to a particular thermal regime and then assayed intracellular processes (as respiration induced by a saturating concentration of glucose) and potential extracellular enzyme activity. Because
260 extracellular enzymes were assayed in soil slurries and also in the presence of excess substrate, it was assumed that substrate diffusion or substrate concentration did not limit reaction rates and that the observed potential reaction rate was thus a function of enzyme properties and enzyme concentration (Wallenstein and Weintraub, 2008). Alongside intracellular and extracellular enzyme steps we measured basal respiration as a reference. We assume that the rate of basal respiration will represent the intracellular metabolic processes as supplied by substrate from extracellular enzyme activity, and that the rate of respiration
265 may be limited by substrate availability (to extracellular processes) and its supply (to intracellular processes) by extracellular activity and diffusion. Thus, any differential effects of pre-incubation temperature on temperature sensitivity of basal respiration cannot be interpreted solely as a function of differences in the cellular physiology of the microbial communities present.

270 Examining the general shape of the response of potential activity to assay temperature, we found that activity of β -glucosidase increased with increasing incubation temperature to our highest assay temperature of 45 °C. Our result is consistent with the increase in β -glucosidase activity with temperature reported in other studies using assay temperatures as low as 2 °C and as high as 65 °C (Steinweg et al., 2013) or 70 °C (Trasar-Cepeda et al., 2007) and showing increases in activity up to and beyond 45 °C. The potential activity of chitinase also increased with temperature, but, in contrast to β -glucosidase, the response, over
275 the range of assay temperatures, was non-monotonic, reaching a maximum activity between 37 °C and 45 °C. This observed non-monotonic response to increasing temperature is interpreted in terms of three distinct phases: (i) a rising phase where temperature increases lead to increasing reaction rate due to thermodynamic effects, (ii) a plateau which represents the optimum temperature and (iii) a steep falling phase where rate declines beyond the optimum temperature (Schulte, 2015) attributed to thermal denaturation of proteins. Our optimum for chitinase (37 to 45 °C) is relatively consistent with the report
280 of a maximum activity for soil chitinase of 45.5 °C (as assayed through quantification of N-acetyl glucosamine released from added chitin; (Rodriguez-Kabana, et al., 1983)) but contrasts to the optimum of ~63 °C reported in the study by (Parham and Deng, 2000) using the same p-nitrophenol-based assay as used here. Differences in these optimum temperature-activity responses between soils may be due to differences in microbial composition (and thus microbial-produced chitinase isozymes) between soils. Optimum temperatures varying between 40 °C and 60 °C have been recorded for chitinases (partially) purified
285 from soil microorganisms (Gao et al., 2008; Alster et al., 2016; Du et al., 2021; Thakur et al., 2021). Additionally, soil-type dependent stabilization of enzyme structure against thermal denaturation through interaction with soil surfaces might also mediate differential temperature responses (Sarkar et al., 1989). It is presumed that β -glucosidase activity in our study soil had a temperature optimum beyond the maximum tested and our finding that the optimum temperature for chitinase activity was lower than that of β -glucosidase is likely due to between-enzyme family differences in protein structural properties

290 conferring thermal stability, resulting in differential susceptibility of different enzyme families to thermal denaturation or
degree of stabilization in soil. Our finding that intracellular catabolism increased with increasing assay temperature up to an
optimal temperature between 26 °C and 37 °C, followed by a significant decline thereafter, is very likely due to the inability
of the microbial population to function optimally above 37 °C due to impairments in their physiological processes (Todd-
Brown et al., 2012; Maire et al., 2013) and uncoupling of relative rates of constituent enzymes leading to regulatory
295 compromise (Prentice et al. 2020). The optimum temperature recorded here is greater than the annual average temperature but
is within the range of the maximum soil temperature experienced for this soil. These findings are in agreement with other
studies on temperate soils recording an optimum temperature for microbial growth of ~30 °C (Bárcenas-Moreno et al., 2009),
although basal respiration rate has been shown to increase with increasing temperature to 45 °C and not to be coupled to
microbial growth (Pietikäinen et al., 2005). Prentice et al., (2020) identify a relationship between the inflection point of the
300 rising phase of the thermal response of intracellular enzyme activities and the growth rate of the organism. This inflection
seems to occur between 15 °C and 26 °C for intracellular metabolic processes and between 26 °C and 37 °C for extracellular
enzymes in our experiment and pre-incubation temperature does not consistently affect the temperature at which this inflection
point occurs (Figure 2). However, the chitinase enzyme activity does not clearly demonstrate the bell-shaped thermal response
expected by macromolecular rate theory. This response may have been observed if assays were undertaken at more
305 temperatures between 26 °C and 45 °C.

We found that the intracellular metabolic processes and extracellular enzyme activities differ in their temperature sensitivity
and this observation differs depending on the range of temperature used for Q10 estimates of temperature sensitivity.
Intracellular metabolic processes were more sensitive to temperature changes at a moderate temperature range (15 °C and 26
310 °C) than extracellular enzymes. Conversely, extracellular enzymes were more sensitive than intracellular processes to
temperature changes at a higher temperature range (26 °C and 37 °C). These results imply that, in the soil we studied,
extracellular depolymerase activity was more temperature sensitive at higher temperatures and intracellular catalytic processes
were more temperature sensitive at moderate temperatures. At the site where the soil was collected for this experiment the
annual mean daily maximum soil temperature was approximately 15 °C, whereas 26 °C reflected a typical hot summer day.
315 Therefore, assuming the absence of any thermal adaptation, we might expect intracellular metabolic processes to be more
sensitive to global warming-induced increases in the mean daily maximum soil temperature, but extracellular enzymes might
be more sensitive to increased maximum temperatures on extremely warm days. The findings described above support our
first hypothesis that the potential rate of extracellular depolymerisation and intracellular catabolism are not equally temperature
sensitive steps in the mineralisation of organic matter in soils. As far as we are aware, only one other study (Blagodatskaya et
320 al., 2016) has considered intra- and extracellular steps involved in organic matter decomposition and their responses to
temperature separately. Our finding that intracellular metabolic processes are more temperature sensitive at moderate
temperatures is in agreement with (Blagodatskaya et al., 2016) who calculated a $Q_{10_{10-20}}$ for intracellular glucose oxidation of
5.1 and $Q_{10_{10-20}}$ for chitinase and β -glucosidase activity of 1.9 and 2, respectively. Other previous research (Trasar-Cepeda et

al., 2007) has compared intracellular metabolic processes (via dehydrogenase assay) and extracellular enzyme activity
325 responses to a wider range of temperatures (5-70 °C), but, not necessarily under substrate-excess conditions for intracellular
metabolic processes, as we have done here. Thus, further experiments are required to evaluate the applicability of our finding
of a greater temperature sensitivity of extracellular activities at higher (26 °C and 37 °C) temperature ranges to other soil types.

Climate change predictions currently indicate that there will be a greater frequency and severity of hot summers and heatwaves
330 in Europe (Meehl and Tebaldi, 2004; Christidis et al., 2015), including Southeast England, where the soil was collected for
this study. Therefore, our findings imply that, in the absence of substrate availability (or other, e.g. moisture) limitations to
activity, the rate of extracellular depolymerase-catalysed reactions will increase during heatwaves to a greater extent than the
rate of intracellular metabolic processes. Depending on the relative sizes of the intra- and extracellular enzyme pools and
substrate availability, it is possible that global warming may reduce the importance of extracellular depolymerisation relative
335 to intracellular metabolic processes as the rate limiting step of SOM mineralization under in situ conditions. Such a switch in
rate limitation, if applicable generally across all extra- and intracellular reactions, would result in a build-up of low molecular
weight substrates in the soil and thus potential for greater losses of C from the soil profile as dissolved organic carbon, an often
overlooked component of terrestrial carbon budgets (Evans et al., 2014; Cook et al., 2018).

340 The temperature sensitivity of C mineralisation is generally found to decrease with temperature (Niklińska and Klimek, 2007;
Wang et al., 2013) and this trend has been observed in a synthesis of soil respiration measurements from laboratory studies
which revealed that Q10 correlates negatively with the range of temperatures used to generate the Q10 value below 25 °C
(Hamdi et al., 2013). This is consistent with kinetic theory of temperature dependence of reaction rates that explains that the
fraction of molecules with sufficient energy to react decreases in relative terms as temperature increases (Davidson and
345 Janssens, 2006). However, similar results to our study have been reported for mineralization of (labile) C where calculated
Q10 values were lower in the 0-10 or 5-15 °C range than 10-20 or 15-25 °C range, respectively (Howard and Howard, 1993;
Wang et al., 2013). These findings possibly reflect that CO₂ production is not a function of a single non-enzyme catalysed
chemical reaction but is subject to moderation by the temperature sensitivity of other components in the involved biochemical
network, for example, reduced membrane fluidity at lower temperature with implications for substrate uptake and function of
350 membrane-embedded proteins (Schulte, 2015). Also, based on kinetic theory, it is suggested that substrates that are more
recalcitrant should have higher temperature sensitivities (Davidson and Janssens, 2006). It might be initially supposed that the
substrates that are hydrolysed by chitinase and β -glucosidase enzymes in depolymerization reactions might be more recalcitrant
than glucose and other lower molecular weight substrates for intracellular respiration and, in consequence, the extracellular-
catalysed reactions should have higher temperature sensitivities. This supposition is supported by the Q10_{26-37°C} data but not
355 for Q10 calculated using the other temperature ranges. However, it should be recognised that chitinase and β -glucosidase have
relatively simple dimeric or trimeric substrates in nature and are assayed using artificial and simple substrates that may not be
more recalcitrant than those used in intracellular metabolism. In addition, the theoretical predictions refer to chemical

decomposition reactions and not necessarily those involving enzyme catalysis (Blagodatskaya et al., 2016). Indeed, comparison of intracellular versus extracellular estimated activation energies (Figure 4) suggested that the extracellular enzyme substrates had similar or lower (for β -glucosidase) recalcitrance. The activation energy values we obtained were in broad correspondence with those reported in other studies (Trasar-Cepeda et al., 2007).

In addition to the differences between the temperature sensitivity of extra- and intra-cellular processes, the extracellular activities were not equally temperature sensitive to each other according to Q_{10}^{26-37} and activation energy (integrating the temperature response between 5 and 26 °C), with chitinase being more sensitive than β -glucosidase. Previous studies have also shown that the temperature sensitivity of particular classes of enzyme differs within the same soil environment (Wallenstein et al., 2010), although specific comparisons between β -glucosidase and chitinase have not always revealed significant differences between these enzyme classes (e.g. Nottingham et al., 2016 ; Min et al., 2014; Min et al., 2019; Wei et al., 2021) and therefore the sign and magnitude of within-soil differences may not be consistent across soil types. In the case of differential temperature sensitivity with respect to assay type, there are implications for temperature-dependent variation in the quality of monomeric SOM constituents supplying respiration (Wallenstein et al. 2011). In the chitinase vs β -glucosidase example here, the relative activity of these enzymes would change with temperature (assuming no change in enzyme or substrate concentration) altering the relative production of glucose and N-acetyl-glucosamine monomers and thus C and N resource availability to soil microbial communities (Min et al., 2014).

In respect of the second hypothesis, the observation that pre-incubation at 26 °C resulted in significantly lower activity, when considered across all assay temperatures, for β -glucosidase and intracellular catalytic enzymes (as well as basal respiration), compared to pre-incubation at 5 °C or 15 °C, suggests possible adaptation of these processes to the direct or indirect effects of temperature. The indirect effects could be due to temperature-induced changes in soil properties during pre-incubation, with consequences for soil microbial activities (Sinsabaugh, 1994; Sinsabaugh et al., 1991; Adeli et al., 2005; Sinsabaugh et al., 2008; Puissant et al., 2019). It was evident in our results that pre-incubating soils at 26 °C reduced the C/N ratio when compared to pre-incubation at 5 °C or 15 °C. This probably reflects enhanced decomposition of organic matter at the warmer pre-incubation temperature and the resulting mass loss of CO₂-C and enrichment of N (on a mass basis) leading to the statistically significant effect when expressed in C/N ratio from. Temperature-induced changes in C/N ratio have been reported previously (Bárta et al., 2013; Souza and Billings, 2022). The lower intracellular enzyme activity after two months exposure to a higher pre-incubation temperature is likely due to a lower (indicated, but not statistically significant) microbial biomass (and thus a reduced intracellular enzyme pool) responding to depleted relative C availability. The lower activity of β -glucosidase for 26 °C-pre-incubated soil most likely also reflects a lower enzyme pool size, given the nature of the potential assay used to measure reaction rate and its relationship to enzyme concentration (Wallenstein and Weintraub, 2008). It is likely that such indirect effects of pre-incubation temperature on the microbial enzyme pool size masks any direct thermal acclimatation or genetic adaptation of the soil microbial community and subsequent change in the temperature sensitivity of the enzymes it produces.

It is often found that substrate depletion plays a greater role in the response of soil microbial communities to warming than physiological or genetic shifts (Domeignoz-Horta et al., 2022). Compared to β -glucosidase, there was less evidence of an effect of pre-incubation temperature on the potential enzyme activity of chitinase (no significance of pre-incubation as a main effect).

395 The potential activity of an enzyme in soil is a function of production versus turnover rate. Accordingly, the balance between these two processes, for β -glucosidase, must have been differentially influenced by pre-incubation temperature, probably both directly and indirectly via, for example, reduced microbial biomass, and complex enzyme regulation in response to altered C availability relative to other nutrients (Allison and Vitousek, 2005; Ferraz de Almeida et al., 2015).

400 Whilst pre-incubation at 26 °C reduced the rate of intracellular metabolic processes, it did not lead to an alteration of community intracellular temperature response traits (i.e. the shape of the temperature response) as evidenced by the non-significant interaction between pre-incubation and assay temperature or a pre-incubation effect on temperature sensitivity as evaluated by calculation of Q10s (Figure 3) or Ea (Figure 4). This result agrees with another study that showed minimal adaptation of the temperature response (of microbial growth) to pre-incubation temperature when the temperature was below

405 the initial optimum (~30 °C) for the mesophilic soil community (Bárcenas-Moreno et al., 2009), although pre-incubation above the optimum led to corresponding increases in the optimum for microbial growth. Minimal adaptive response to pre-incubation substantially below the initial optimum (i.e. 5 and 15 °C in our study) is explained in terms of a rate of species sorting (ultimately favouring a community better adapted to the pre-incubation conditions) being too slow to manifest within the 60-day pre-incubation period due to slow microbial generation times at colder temperatures (Bárcenas-Moreno et al., 2009). In

410 contrast to intracellular metabolic processes, there was some evidence (significant pre-incubation \times assay temperature interaction) of a pre-incubation effect on the temperature response for potential extracellular enzyme activity, although effects were quite subtle and only systematic with pre-incubation temperature for chitinase where activity assayed at 37 °C decreased in order of decreasing pre-incubation temperature (also discernible in effects of pre-incubation on Q10_{26-37°C} and Ea). The lower Ea for chitinase for soil samples pre-incubated at the lower temperatures is consistent with the concept that cold

415 adaptation of microorganisms leads to the production of cold adapted enzymes, by adjustment of chemical structure of the active site, with lower activation energies (Wallenstein et al., 2011). However, the few previous experimental studies examining the temperature response/sensitivity of extracellular enzyme potentials in soils exposed to differing thermal regimes have suggested no difference in temperature sensitivity (Schindlbacher et al., 2015; Jing et al., 2019) and therefore an absence of thermal adaptation of temperature sensitivity. However, these experiments involved long-term field-based warming

420 treatments and it is suggested that the effects of the experimental warming were negligible against the effects of wide seasonal temperature variations (Jing et al., 2019). Other studies, however, have demonstrated seasonal changes in temperature sensitivity of extracellular enzymes (Wallenstein et al. 2011; Wallenstein et al. 2009). These changes likely result from temporal changes in production of isoenzymes (by different organisms or within the same organism transcribing alternative

enzyme-encoding genes) but whether these patterns represent an adaptation to seasonally-varying temperature, or are driven
425 by other factors that change seasonally (e.g. substrate availability) is not clear (Wallenstein et al., 2011).

5 Conclusion

Our results advance understanding of how SOM decomposition will change under future global warming conditions. We show that the potential rates of the intracellular and extracellular steps of SOM decomposition are not equally sensitive to changes in temperature and that individual extracellular enzymes have different temperature sensitivities. Specifically, for our
430 individual grassland soil pre-incubated at just three representative temperatures, we have demonstrated that potential activities of extracellular depolymerase enzymes (β -glucosidase and chitinase) have greater sensitivity to increases in temperature in the range of temperatures experienced on extremely warm days (between 26 °C and 37 °C) than the rate of intracellular metabolic processes involved in catabolism of monomeric (e.g. glucose) substrates to CO₂. Since a greater prevalence of extremely hot days and heatwaves are predicted, the importance of intracellular metabolic processes may increase and the importance of
435 extracellular enzyme activity may decrease as the rate limiting step in SOM decomposition.

For the extracellular enzymes studied, we found differential temperature sensitivity with respect to enzyme class. Here, the implications are for temperature-dependent variation in the quality of monomeric SOM substrates supplying respiration and potential feedbacks to soil microbial community composition given taxa-specific competitive utilization of substrates
440 (Wallenstein et al. 2011). Whilst interpretation should be within the context of the pre-incubation conditions (60 days at temperatures less than the optimum for activity of the mesophilic community), we have also shown that the thermal history (i.e. pre-incubation temperature) of a soil might modulate the relative responses in reaction rates to current temperature. This is both through enzyme-dependent reduction of potential activity across assay temperatures in 26 °C pre-incubated soil (for intracellular enzymes and β -glucosidase but not chitinase) and also subtle adaptation of the temperature response trait to pre-
445 incubation temperature (for extracellular enzymes but not intracellular metabolic processes). Measurements of CO₂ alone as a response variable while studying the effect of warming may obscure our understanding of the temperature sensitivity of the intracellular and extracellular steps of SOM decomposition.

450 Data Availability

Data associated with this publication has been uploaded to the Mendeley Data repository and is embargoed until 13th October 2023:

Adekanmbi, Adetunji Alex; Dale, Laurence; Shaw, Liz; Sizmur, Tom (2022), "Data for: Differential Temperature Sensitivity of Intracellular and Extracellular Soil Enzyme Activities", Mendeley Data, V1, doi: 10.17632/xvr3dzvdcw.1

455

Author Contributions

AAA, LD, LS, and TS conceptualised the research and designed the experiment. AAA and LD carried out the laboratory work. AAA and LD analysed the data, with support from TS and LS. AAA and LD undertook data visualisation. AAA wrote the original draft of the paper. TS, LS, LD and AAA undertook subsequent reviewing and editing. TS supervised the project and
460 LS co-supervised the project.

Competing Interests

The authors declare that they have no conflict of interest

465 **Acknowledgements**

The authors gratefully acknowledge Technical Services staff within the Environmental Science Research Division at the University of Reading for technical support & assistance in this work

References


- 470 Adekanmbi, A. A., Shaw, L. J., and Sizmur, T.: Effect of Sieving on Ex Situ Soil Respiration of Soils from Three Land Use Types, *J. Soil Sci. Plant Nutr.*, 20, 912–916, <https://doi.org/https://doi.org/10.1007/s42729-020-00177-2>, 2020.
- Adeli, A., Sistani, K. R., Bal'a, M. F., and Rowe, D. E.: Phosphorus Dynamics in Broiler Litter-Amended Soils, *Commun. Soil Sci. Plant Anal.*, 36, 1099–1115, <https://doi.org/10.1081/CSS-200056876>, 2005.
- Allison, S. D. and Vitousek, P. M.: Responses of extracellular enzymes to simple and complex nutrient inputs, *Soil Biol. Biochem.*, 37, 937–944, <https://doi.org/10.1016/j.soilbio.2004.09.014>, 2005.
- 475 Alster, C. J., Baas, P., Wallenstein, M. D., Johnson, N. G., and von Fischer, J. C.: Temperature sensitivity as a microbial trait using parameters from macromolecular rate theory, *Front. Microbiol.*, 7, 1–10, <https://doi.org/10.3389/fmicb.2016.01821>, 2016.
- Alster, C. J., von Fischer, J. C., Allison, S. D., and Treseder, K. K.: Embracing a new paradigm for temperature sensitivity of soil microbes., *Glob. Chang. Biol.*, 26, 3221–3229, <https://doi.org/https://doi.org/10.1111/gcb.15053>, 2020.
- 480 Alvarez, G., Shahzad, T., Andanson, L., Bahn, M., Wallenstein, M. D., and Fontaine, S.: Catalytic power of enzymes decreases with temperature: new insights for understanding soil C cycling and microbial ecology under warming, *Glob. Chang. Biol.*, 0–2, <https://doi.org/10.1111/gcb.14281>, 2018.
- Bárcenas-Moreno, G., Brandón, M. G., Rousk, J., and Bååth, E.: Adaptation of soil microbial communities to temperature: Comparison of fungi and bacteria in a laboratory experiment, *Glob. Chang. Biol.*, 15, 2950–2957, <https://doi.org/10.1111/j.1365-2486.2009.01882.x>, 2009.
- 485 Bárta, J., Šlajsová, P., Tahovska, K., Pícek, T., and Santruckova, H.: Different temperature sensitivity and kinetics of soil enzymes indicate seasonal shifts in C, N and P nutrient stoichiometry in acid forest soil, *Biogeochemistry*, 117, 525–537, <https://doi.org/10.1007/s10533-013-9898-1>, 2014.
- Birgander, J., Reischke, S., Jones, D. L., and Rousk, J.: Temperature adaptation of bacterial growth and ¹⁴C-glucose mineralisation in a laboratory study, *Soil Biol. Biochem.*, 65, 294–303, <https://doi.org/10.1016/j.soilbio.2013.06.006>, 2013.
- 490 Blagodatskaya, E., Blagodatsky, S., Khomyakov, N., Myachina, O., and Kuzyakov, Y.: Temperature sensitivity and enzymatic mechanisms of soil organic matter decomposition along an altitudinal gradient on Mount Kilimanjaro, *Sci. Rep.*, 6, 1–11, <https://doi.org/10.1038/srep22240>, 2016.
- 495 Bradford, M. A.: Thermal adaptation of decomposer communities in warming soils, *Front. Microbiol.*, 4, 1–16, <https://doi.org/10.3389/fmicb.2013.00333>, 2013.

- Christidis, N., Jones, G. S., and Stott, P. A.: Dramatically increasing chance of extremely hot summers since the 2003 European heatwave, *Nat. Clim. Chang.*, 5, 46–50, <https://doi.org/10.1038/nclimate2468>, 2015.
- Cook, S., Whelan, M. J., Evans, C. D., Gauci, V., Peacock, M., Garnett, M. H., Kho, L. K., Teh, Y. A., and Page, S. E.: Fluvial organic carbon fluxes from oil palm plantations on tropical peatland, *Biogeosciences*, 15, 7435–7450, <https://doi.org/10.5194/bg-15-7435-2018>, 2018.
- Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, <https://doi.org/10.1038/nature04514>, 2006.
- Du, J., Duan, S., Miao, J., Zhai, M., and Cao, Y.: Purification and characterization of chitinase from *Paenibacillus* sp., *Biotechnol. Appl. Biochem.*, 68, 30–40, <https://doi.org/10.1002/bab.1889>, 2021.
- Duly, O. and Nannipieri, P.: Intracellular and extracellular enzyme activity in soil with reference to elemental cycling, *Zeitschrift für Pflanzenernährung und Bodenkd.*, 161, 243–248, <https://doi.org/10.1002/jpln.1998.3581610310>, 1998.
- Eivazi, F. and Tabatabai, M. A.: Glucosidases and Galactosidases in Soils, *Soil Biol. Biochem.*, 20, 601–606, [https://doi.org/https://doi.org/10.1016/0038-0717\(88\)90141-1](https://doi.org/https://doi.org/10.1016/0038-0717(88)90141-1), 1988.
- Evans, C. D., Page, S. E., Jones, T., Moore, S., Gauci, V., Laiho, R., Hruska, J., Allot, T. E. ., Billet, M. F., Tipping, E., Freeman, C., and Garnett, M. H.: Contrasting vulnerability of drained tropical and high-latitude peatlands to fluvial loss of stored carbon, *Global Biogeochem. Cycles*, 28, 1215–1234, <https://doi.org/10.1002/2013GB004782>, 2014.
- Fanin, N., Mooshammer, M., Sauvadet, M., Meng, C., Alvarez, G., Bernard, L., Bertrand, I., Blagodatskaya, E., Bon, L., Fontaine, S., Niu, S., Lashermes, G., Maxwell, T. L., Weintraub, M. N., Wingate, L., Moorhead, D., and Nottingham, A. T.: Soil enzymes in response to climate warming: Mechanisms and feedbacks, *Funct. Ecol.*, 36, 1378–1395, <https://doi.org/10.1111/1365-2435.14027>, 2022.
- Gao, X. A., Ju, W. T., Jung, W. J., and Park, R. D.: Purification and characterization of chitosanase from *Bacillus cereus* D-11, *Carbohydr. Polym.*, 72, 513–520, <https://doi.org/10.1016/j.carbpol.2007.09.025>, 2008.
- Hamdi, S., Moyano, F., Sall, S., Bernoux, M., and Chevallier, T.: Synthesis analysis of the temperature sensitivity of soil respiration from laboratory studies in relation to incubation methods and soil conditions, *Soil Biol. Biochem.*, 58, 115–126, <https://doi.org/10.1016/j.soilbio.2012.11.012>, 2013.
- Howard, D. M. and Howard, P. J. A.: Relationships between co₂ evolution, moisture content and temperature for a range of soil types, *Soil Biol. Biochem.*, 25, 1537–1546, [https://doi.org/10.1016/0038-0717\(93\)90008-Y](https://doi.org/10.1016/0038-0717(93)90008-Y), 1993.
- Jan, M. T., Roberts, P., Tonheim, S. K., and Jones, D. L.: Protein breakdown represents a major bottleneck in nitrogen cycling in grassland soils, *Soil Biol. Biochem.*, 41, 2272–2282, <https://doi.org/10.1016/j.soilbio.2009.08.013>, 2009.


- Jing, X., Wang, Y., Chung, H., Mi, Z., Wang, S., Jing, X., Wang, Y., and Chung, H.: No temperature acclimation of soil extracellular enzymes to experimental warming in an alpine grassland ecosystem on the Tibetan Plateau Stable URL : <https://www.jstor.org/stable/24716843> No temperature acclimation of soil extracellular enzymes to experim, *Biogeochemistry*, 117, 39–54, <https://doi.org/10.1007/s10533-013-9844-2>, 2019.
- 530 Karhu, K., Auffret, M. D., Dungait, J. A. J., Hopkins, D. W., Prosser, J. I., Singh, B. K., Subke, J. A., Wookey, P. A., Agren, G. I., Sebastià, M. T., Gouriveau, F., Bergkvist, G., Meir, P., Nottingham, A. T., Salinas, N., and Hartley, I. P.: Temperature sensitivity of soil respiration rates enhanced by microbial community response., *Nature*, 513, 81–84, <https://doi.org/10.1038/nature13604>, 2014.
- Kirwan, M. L., Guntenspergen, G. R., and Langley, J. A.: Temperature sensitivity of organic-matter decay in tidal marshes, *Biogeosciences*, 11, 4801–4808, <https://doi.org/10.5194/bg-11-4801-2014>, 2014.
- 535 Li, J., He, N., Wei, X., Gao, Y., and Zuo, Y.: Changes in temperature sensitivity and activation energy of soil organic matter decomposition in different Qinghai-Tibet Plateau grasslands, *PLoS One*, 10, 1–14, <https://doi.org/10.1371/journal.pone.0132795>, 2015.
- Maire, V., Alvarez, G., Colombet, J., Comby, A., Despinasse, R., Dubreucq, E., Joly, M., Lehours, A. C., Perrier, V., Shahzad, T., and Fontaine, S.: An unknown oxidative metabolism substantially contributes to soil CO₂ emissions, *Biogeosciences*, 10, 1155–1167, <https://doi.org/10.5194/bg-10-1155-2013>, 2013.
- 540 Meehl, G. A. and Tebaldi, C.: More intense, more frequent, and longer lasting heat waves in the 21st century, *Science* (80-.), 305, 994–997, <https://doi.org/10.1126/science.1098704>, 2004.
- Min, K., Lehmeier, C. A., Ballantyne, F., Tatarko, A., and Billings, S. A.: Differential effects of pH on temperature sensitivity of organic carbon and nitrogen decay, *Soil Biol. Biochem.*, 76, 193–200, <https://doi.org/10.1016/j.soilbio.2014.05.021>, 2014.
- 545 Min, K., Bagchi, S., Buckeridge, K., Billings, S. A., Ziegler, S. E., and Edwards, K. A.: Temperature sensitivity of biomass - specific microbial exo - enzyme activities and CO₂ efflux is resistant to change across short - and long - term timescales, *Glob. Chang. Biol.*, 25, 1793–1807, <https://doi.org/10.1111/gcb.14605>, 2019.
- Niklińska, M. and Klimek, B.: Effect of temperature on the respiration rate of forest soil organic layer along an elevation gradient in the Polish Carpathians, *Biol. Fertil. Soils*, 43, 511–518, <https://doi.org/10.1007/s00374-006-0129-y>, 2007.
- 550 Nottingham, A. T., Turner, B. L., Whitaker, J., Ostle, N., Bardgett, R. D., McNamara, N. P., Salinas, N., and Meir, P.: Temperature sensitivity of soil enzymes along an elevation gradient in the Peruvian Andes, *Biogeochemistry*, 127, 217–230, <https://doi.org/10.1007/s10533-015-0176-2>, 2016.
- Parham, J. A. and Deng, S. P.: Detection, quantification and characterization of b- glucosaminidase activity in soil, *Soil Biol.*

- 555 Biochem., 32, 1183–1190, [https://doi.org/https://doi.org/10.1016/S0038-0717\(00\)00034-1](https://doi.org/https://doi.org/10.1016/S0038-0717(00)00034-1), 2000.
- Pietikäinen, J., Pettersson, M., and Bååth, E.: Comparison of temperature effects on soil respiration and bacterial and fungal growth rates, *FEMS Microbiol. Ecol.*, 52, 49–58, <https://doi.org/10.1016/j.femsec.2004.10.002>, 2005.
- Prentice, E. J., Hicks, J., Ballerstedt, H., Blank, L. M., Liang, L. L., Schipper, L. A., and Arcus, V. L.: The inflection point hypothesis: The relationship between the temperature dependence of enzyme-catalyzed reaction rates and microbial growth rates, *Biochemistry*, 59, 3562–3569, <https://doi.org/10.1021/acs.biochem.0c00530>, 2020.
- 560 Puissant, J., Jones, B., Goodall, T., Mang, D., Blaud, A., Gweon, H. S., Malik, A., Jones, D. L., Clark, I. M., Hirsch, P. R., and Griffiths, R.: The pH optimum of soil exoenzymes adapt to long term changes in soil pH, *Soil Biol. Biochem.*, 138, 107601, <https://doi.org/10.1016/j.soilbio.2019.107601>, 2019.
- Rodriguez-Kabana, R., Godoy, G., Morgan-Jones, G., and Shelby, R. A.: The determination of soil chitinase activity :
565 Conditions for assay and ecological studies, *Plant Soil*, 75, 95–106, 1983.
- Salazar-Villegas, A., Blagodatskaya, E., and Dukes, J. S.: Changes in the size of the active microbial pool explain short-term soil respiratory responses to temperature and moisture, *Front. Microbiol.*, 7, 1–10, <https://doi.org/10.3389/fmicb.2016.00524>, 2016.
- Sarkar, J. M., Leonowicz, A., and Bollag, J. M.: Immobilization of enzymes on clays and soils, *Soil Biol. Biochem.*, 21, 223–
570 230, [https://doi.org/10.1016/0038-0717\(89\)90098-9](https://doi.org/10.1016/0038-0717(89)90098-9), 1989.
- Schindlbacher, A., Schneckler, J., Takriti, M., Borcken, W., and Wanek, W.: Microbial physiology and soil CO₂ efflux after 9 years of soil warming in a temperate forest - no indications for thermal adaptations, *Glob. Chang. Biol.*, 21, 4265–4277, <https://doi.org/10.1111/gcb.12996>, 2015.
- Schulte, P. M.: The effects of temperature on aerobic metabolism: Towards a mechanistic understanding of the responses of
575 ectotherms to a changing environment, *J. Exp. Biol.*, 218, 1856–1866, <https://doi.org/10.1242/jeb.118851>, 2015.
- Sinsabaugh, R. L., Antibus, R. K., and Linkins, A. E.: An enzymic approach to the analysis of microbial activity during plant litter decomposition, *Agric. Ecosyst. Environ.*, 34, 43–54, [https://doi.org/10.1016/0167-8809\(91\)90092-C](https://doi.org/10.1016/0167-8809(91)90092-C), 1991.
- Sinsabaugh, R. L., Lauber, C. L., Weintraub, M. N., Ahmed, B., Allison, S. D., Crenshaw, C., Contosta, A. R., Cusack, D., Frey, S., Gallo, M. E., Gartner, T. B., Hobbie, S. E., Holland, K., Keeler, B. L., Powers, J. S., Stursova, M., Takacs-Vesbach, C., Waldrop, M. P., Wallenstein, M. D., Zak, D. R., and Zeglin, L. H.: Stoichiometry of soil enzyme activity at global scale,
580 *Ecol. Lett.*, 11, 1252–1264, <https://doi.org/10.1111/j.1461-0248.2008.01245.x>, 2008.
- Sinsabaugh, R. S.: Enzymic analysis of microbial pattern and process, *Biol. Fertil. Soils*, 17, 69–74, <https://doi.org/10.1007/BF00418675>, 1994.


- Souza, L. F. T. and Billings, S. A.: Temperature and pH mediate stoichiometric constraints of organically derived soil nutrients, *Glob. Chang. Biol.*, 28, 1630–1642, <https://doi.org/10.1111/gcb.15985>, 2022.
- Steinweg, M. J., Jagadamma, S., Frerichs, J., and Mayes, M. A.: Activation Energy of Extracellular Enzymes in Soils from Different Biomes, *PLoS One*, 8, e59943, <https://doi.org/10.1371/journal.pone.0059943>, 2013.
- Thakur, N., Nath, A. K., Chauhan, A., and Gupta, R.: Purification, characterization, and antifungal activity of *Bacillus cereus* strain NK91 chitinase from rhizospheric soil samples of Himachal Pradesh, India, *Biotechnol. Appl. Biochem.*, 69, 1830–1842, <https://doi.org/10.1002/bab.2250>, 2021.
- Todd-Brown, K., Hopkins, F. M., Kivlin, S. N., Talbot, J. M., and Allison, S. D.: A framework for representing microbial decomposition in coupled climate models, *Biogeochemistry*, 109, 19–33, <https://doi.org/https://doi.org/10.1007/s10533-011-9635-6>, 2012.
- Trasar-Cepeda, C., Gil-Sotres, F., and Leirós, M. C.: Thermodynamic parameters of enzymes in grassland soils from Galicia, NW Spain, *Soil Biol. Biochem.*, 39, 311–319, <https://doi.org/10.1016/j.soilbio.2006.08.002>, 2007.
- Vance, E. D., Brookes, P. C., and Jenkinson, D. S.: An Extraction method for measuring Soil Microbial Biomass C, *Soil Biol. Biochem.*, 19, 703–707, [https://doi.org/10.1016/0038-0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6), 1987.
- Wallenstein, M., Allison, S. D., Ernakovich, J., Steinweg, J. M., and Sinsabaugh, R.: Controls on the Temperature Sensitivity of Soil Enzymes: A Key Driver of In Situ Enzyme Activity Rates., in: *Soil Enzymology*, edited by: Shukla, G. and Varma, A., Springer Berlin Heidelberg, Berlin, Heidelberg, 245–258, https://doi.org/10.1007/978-3-642-14225-3_13, 2010.
- Wallenstein, M. D. and Weintraub, M. N.: Emerging tools for measuring and modeling the in situ activity of soil extracellular enzymes, *Soil Biol. Biochem.*, 40, 2098–2106, <https://doi.org/10.1016/j.soilbio.2008.01.024>, 2008.
- Wang, G., Zhou, Y., Xu, X., Ruan, H., and Wang, J.: Temperature Sensitivity of Soil Organic Carbon Mineralization along an Elevation Gradient in the Wuyi Mountains, China, *PLoS One*, 8, e53914, <https://doi.org/10.1371/journal.pone.0053914>, 2013.
- Wei, L., Zhu, Z., Liu, S., Xiao, M., Wang, J., Deng, Y., Kuzyakov, Y., Wu, J., and Ge, T.: Temperature sensitivity (Q₁₀) of stable, primed and easily available organic matter pools during decomposition in paddy soil, *Appl. Soil Ecol.*, 157, 103752, <https://doi.org/10.1016/j.apsoil.2020.103752>, 2021.




615 **Figure 1: Effects of pre-incubation temperature on soil total carbon, total nitrogen, carbon-to-nitrogen (C/N) ratio, pH and microbial biomass carbon. Each bar and error bar represents mean and standard error of 4 replicate samples at each pre-incubation temperature. Means with the same letter are not significantly different ($P > 0.05$).**



620 **Figure 2: Response of β -glucosidase activity (a), chitinase activity (b), glucose-induced respiration (c), and basal respiration (d) to five assay temperatures (5 °C, 15 °C, 26 °C, 37 °C and 45 °C) undertaken on soils pre-incubated at three different temperatures (5 °C, 15 °C, and 26 °C). Each symbol and error bar represent mean and standard error of 4 replicate samples.**



625 **Figure 3: Effects of pre-incubation temperature on temperature sensitivity (Q10_{5-15°C}, Q10_{15-26°C}, and Q10_{26-37°C}) of basal respiration rate, glucose-induced respiration, and extracellular (chitinase and β-glucosidase) enzyme activity. Each bar and error bar represent mean and standard error of 4 replicate samples each pre-incubated at one of three different pre-incubation temperatures (5 °C, 15 °C, or 26 °C). Assay types sharing the same upper case letters are not significantly different (P > 0.05).**



635 **Figure 4: Effects of pre-incubation temperature (5 °C, 15 °C, or 26 °C) on Arrhenius enzyme activation energy (Ea) for basal respiration rate, intracellular (glucose substrate induced respiration) enzyme activity, and extracellular (chitinase and β-glucosidase) enzyme activity. Ea was calculated based on assays undertaken between (5 °C and 26 °C for all four assays. Each bar and error bar represent mean and standard error of 4 replicates. Enzyme types sharing the same upper case letters are not significantly different ($P > 0.05$).**