Revealing the sources and sinks of negative cluster ions in an urban environment through quantitative analysis

3	Rujing Yin ^{1,2#} , Xiaoxiao Li ^{1,3#} , Chao Yan ^{2,4} , Runlong Cai ² , Ying Zhou ⁴ , Juha Kangasluoma ² , Nina Sarnela ² , Janne
4	Lampilahti ² , Tuukka Petäjä ² , Veli-Matti Kerminen ² , Federico Bianchi ² , Markku Kulmala ^{2,4} , Jingkun Jiang ^{1*}

- ¹State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua
 University, Beijing, 100084, China
- ²Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, 00014
 Helsinki, Finland
- 9 ³School of Resources and Environmental Sciences, Wuhan University, 430072 Wuhan, China
- ¹⁰ ⁴Aerosol and Haze Laboratory, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing
- 11 University of Chemical Technology, 100029 Beijing, China
- 12 [#]These authors contributed equally to this work.
- 13 *: Correspondence to: J. Jiang (jiangjk@tsinghua.edu.cn)
- 14

15	This	PDF	file	includes:
----	------	-----	------	-----------

- 16 Main text
- 17 Figures 1 to 10
- 18
- 19

Abstract. Atmospheric cluster ions are important constituents in the atmosphere, and their concentrations and 20 21 compositions govern their role in atmospheric chemistry. However, there is currently limited quantitative research on 22 atmospheric ion compositions, sources, and sinks, especially in the urban atmosphere where pollution levels and human 23 populations are intense. In this study, we measured the compositions of negative cluster ions (mobility diameter smaller 24 than 1.6 nm) and neutral molecules using an atmospheric pressure interface high-resolution time-of-flight mass spectrometer (APi-TOF) and a chemical ionization mass spectrometer in urban Beijing. Quantitative analysis of cluster 25 ions was performed by their comparison with condensation sink (CS), reagent ions, and neutral molecules. We 26 demonstrate the feasibility of quantifying cluster ions with different compositions using *in-situ* measured ion mobility 27 distributions from a neutral cluster and air ion spectrometer (NAIS). The median concentration of negative cluster ions 28 was 85 (61-112 for 25-75%) cm⁻³ during the measurement period, which was negatively correlated with CS. The negative 29 cluster ions mainly consisted of inorganic nitrogen-containing ions, inorganic sulfur-containing ions, and organic ions in 30 the form of adducts with NO₃⁻ or HSO4⁻. The CHON-related organic ions accounted for over 70% of the total organic 31 32 ions. Although the molecules clustered with NO₃⁻ and HSO₄⁻ had similar compositions, we found that HSO₄⁻ clustered more efficiently with CHO and CHON_{nonNPs} species (CHON excluding nitrated phenols), while NO3⁻ clustered more 33 efficiently with nitrated phenols (CHON_{nonNPs}). Additionally, most organic ions were positively correlated with neutral 34 35 molecules, resulting in similar diurnal cycles of organic ions and neutral molecules. However, an exception was found for CHON_{NPs}, the concentration of which is also significantly influenced by the reagent ions NO₃. The charge fractions 36 are generally higher for molecules with higher molecular weight and oxidation state, and the opposite diurnal variation 37 of charging fractions between H₂SO₄ and organic species indicates a charging competition between them. Finally, we 38 39 choose HSO_4^- and $C_3H_3O_4^-$ as representatives to calculate the contribution of different formation and loss pathways. We found their losses are condensational loss onto aerosol particles (73-75%), ion-molecule reaction losses (19%), and ion-40 ion recombination losses (6-8%). 41

42 Keywords: atmospheric cluster ions, ion composition, quantitative analysis, urban atmosphere, condensation sink

43

44 **1 Introduction**

Atmospheric cluster ions are electrically charged atoms, molecules, and molecular clusters. Their mobility is usually 45 larger than 0.5 cm²·V⁻¹·s⁻¹ (Hõrrak et al., 2000). The primary ions, such as N₂⁺, O₂⁺, NO⁺, O⁻, and O₂⁻, are initially formed 46 via cosmic radiation, gamma radiation, and other near-ground local sources, such as radon decay, lightning, plant 47 48 emissions, waterfall and seashore generation, combustion, and high-voltage transmission lines emission (Carslaw et al., 49 2002; Eisele, 1989; Hirsikko et al., 2007; Tammet et al., 2006). Primary ions will subsequently undergo ion-molecule 50 reactions, ion-ion recombination, or deposition onto particles, accompanied by the evolution of their electrical mobility, concentration, and composition (Curtius et al., 2006; Harrison, 2003; Kontkanen et al., 2013; Shuman et al., 2015). 51 52 Atmospheric ions play important roles in atmospheric chemistry (Bates, 1982; Luts and Salm, 1994), ion-induced 53 nucleation (Charlson et al., 1992; Lee et al., 2003; Lovejoy, 2004), atmosphere conductivity (G. Baumgaertner et al., 54 2013) and air quality (Jiang et al., 2018). Moreover, atmospheric ions can benefit human health at certain levels of concentration, while their low concentrations may cause headaches and insomnia (Chu et al., 2019; Malcolm et al., 2009).
 The effects of atmospheric cluster ions are highly related to their mobility, concentration, and composition.

57 There are substantial differences in the characteristics of atmospheric ions and their dynamic variations among diverse 58 environments. Globally, atmospheric ion concentrations span a wide range from 100 to 5000 cm⁻³ (Hirsikko et al., 2011; 59 Usoskin et al., 2004), determined mainly by differences in ion production and loss rates. On one hand, ion production rates vary with geographical location, land cover type, and weather conditions (Chen et al., 2016; Ling et al., 2010; 60 61 Usoskin et al., 2004). On the other hand, ion loss rates are largely related to ions carrying opposite charges, aerosol surface area concentrations, and deposition rates (Harrison, 2003; Tammet et al., 2006). For ion compositions, hundreds 62 of new ions can be formed through ion-molecule reactions after the primary ions are initially formed. It has been found 63 that besides the primary ions, NO_3^- , HSO_4^- , and their clusters were predominant ions due to their high acidities (Davidson 64 et al., 1977; Viggiano et al., 1980). Oxygenated organic molecules (OOMs) were observed clustering with NO₃⁻ and 65 HSO₄⁻ naturally in the atmosphere and the composition of these organic atmospheric ions has been reported to be similar 66 to those of neutral vapor molecules (Bianchi et al., 2017; Ehn et al., 2010; Yin et al., 2021). Specifically, ion compositions 67 were found very different between NPF and non-NPF events, and ion compositions during NPF imply different 68 69 nucleation pathways in clean and polluted atmospheres (Bianchi et al., 2016; Ehn et al., 2010; Eisele et al., 2006; Kirkby 70 et al., 2016; Yin et al., 2021). Thus, the measurement of ion characteristics requires high temporal measurements in 71 various environments.

72 Ion characteristics at polluted urban sites, where the ion production and loss processes may be very different from the 73 clean sites, are relatively poorly understood. At many polluted urban sites, intense pollution emissions led to high 74 concentrations of neutral gas species and high aerosol concentrations. For the ion production processes, the neutral gas 75 molecules at urban sites are significantly different from those at clean sites. High NO_x concentrations were found to be 76 accompanied by high concentrations of nitrogen-containing organic molecules at polluted urban sites (Li et al., 2022; 77 Qiao et al., 2021). For ion loss processes, condensation sink (CS) at urban sites can reach up to two orders of magnitude 78 higher values compared with clean sites (Cai et al., 2017), which would cause a significant loss of ions onto large particles in urban air. These may lead to big differences in ion concentrations and compositions between clean and urban sites. 79 80 For instance, Iida et al. (2006) have shown that the ion concentration near a polluted site in Boulder might be substantially 81 suppressed by high ion loss rates, resulting in only a minor contribution from ion-induced nucleation to NPF. Although some recent studies have revealed the characteristics of ions in clean environments (Bianchi et al., 2016; Chen et al., 82 2016; Ehn et al., 2010), only a few studies have focused on such characteristics in an urban atmosphere (Yin et al., 2021). 83

Simultaneous analysis of ion concentrations and compositions is rare in the field campaign. Various ion counters have been designed to measure ion concentrations in the last century. Among these, Gerdien counters (Gerdien, 1905), the balance scanning mobility analyzer (BSMA) (Tammet, 2006), the air ion spectrometer (AIS) (Mirme et al., 2007), and a neutral cluster and air ion spectrometer (NAIS) (Manninen et al., 2016; Manninen et al., 2009) are the most widely used ion counters or ion mobility spectrometers. However, these electrical mobility-based instruments do not provide composition information of ions. Mass spectrometers have been widely used to measure ion compositions. Quadrupole mass spectrometers with approximate unit mass resolution have been applied in a series of ion composition measurements (Arnold et al., 1978; Arnold and Viggiano, 1982; Eisele, 1989; Eisele et al., 2006; Eisele and Tanner, 1990;
Viggiano, 1993). With the development of high-resolution mass spectrometer techniques, such as an atmospheric
pressure interface time-of-flight mass spectrometer (APi-TOF, Tofwerk AG), more species of atmospheric ions were
identified (Ehn et al., 2010; Eisele et al., 2006; Junninen et al., 2010; Kirkby et al., 2016), but the concentration measured
by the APi-TOF remains to be quantified.

It is needed to develop a robust and easy-to-use method to quantify concentrations of different ions in APi-TOF in field 96 97 measurements. The detection efficiency of an APi-TOF to different ions is mainly determined by the mass-dependent transmission efficiency. The calibration of transmission efficiency often requires complex laboratory calibration settings, 98 99 including the generation, classification, and counting of ions with different m/z ratios (Heinritzi et al., 2016; Junninen et al., 2010). Alternatively, the relative transmission efficiency of an APi-TOF compared to primary ions can be obtained 100 using a depletion method with the help of a chemical ionization inlet (Heinritzi et al., 2016). But the settings are still 101 102 complex and only relative transmission efficiency is obtained. Ehn et al. (2011) have shown that the ion concentration 103 measured by NAIS and the size-segregated ion signal measured by APi-TOF have good correlations in Hyytiälä. Then 104 by converting the ion mobility into the m/z of the ions (Hõrrak et al., 2000), the *in-situ* calibration of the mass 105 spectrometers can be achieved using the synchronous measurements of well-calibrated ion spectrometers (e.g. NAIS). However, the feasibility of this method still needs to be examined, especially in polluted urban atmospheres. 106

107 In this study, we apply an *in-situ* quantification method of atmospheric ion compositions measured by an APi-TOF and 108 reveal the governing factors of atmospheric cluster ion concentration and composition at polluted urban sites. We perform 109 field measurements of atmospheric negative cluster ions at an urban site using the APi-TOF and NAIS simultaneously. 110 By comparing the two *in-situ* measurements, we quantify the concentration of ions measured by the APi-TOF using an 111 improved method. The concentration and composition of ions are compared with the clean sites, and the reasons for the 112 observed differences are explored. The origins and composition variations of ions are characterized and compared with 113 the neutral clusters measured by a chemical ionization mass spectrometer with NO3⁻ as the reagent ions (abbreviated as 114 (nitrate) CI-APi-TOF). The driving factors for the variations in ion characteristics at the urban site are revealed, including CS, reagent ions, and neutral molecules. The charge fractions of various organic species were determined. To better 115 116 quantify the formation and loss scheme of atmospheric ions, the ions HSO_4^- and $C_3H_3O_4^-$ are taken as representatives in 117 the mass balance equation and their variations are well-captured.

118 2 Methods

119 **2.1 Field measurement**

A field measurement is conducted at the Aerosol and Haze Laboratory of Beijing University of Chemical Technology Station (AHL/BUCT station), Beijing, China (Liu et al., 2020; Yin et al., 2021). The AHL/BUCT station is a typical urban site surrounded by three traffic roads and some residential buildings. The station is ~18 m above the ground with no higher buildings around it within 50 m. It is equipped with many state-of-the-art instruments and routine instruments for air quality measurements. Besides, the measurements at a boreal forest station SMEAR II (System for Measurement of forest Ecosystem and Atmospheric Relationships) located at Hyytiälä, Finland were used for comparison. Details of the

126 instruments and their measurement periods in these two stations are given in Table S1.

127 The atmospheric negative cluster ion compositions were measured with an atmospheric pressure interface time-of-flight high-resolution mass spectrometers (APi-HTOF) from Feb 14 to Feb 27, 2018, in Beijing, and from April 7 to June 8, 128 2013 in Hyytiälä (Yan et al., 2018). The ambient air was sampled from the window through a 1.4-m long 1/4 in, stainless 129 130 steel tubing. The sampling flow rate of the APi-HTOF is ~0.8 LPM, and an extra sheath flow of 3 LPM was sampled to 131 minimize the loss of ions in the sampling tubes. Thus the total sampling flow rate was 3.8 LPM. The APi-HTOF was operated in the negative ion mode to measure negative ions and its voltages were adjusted to reduce cluster 132 fragmentations. The mass resolution of the APi-HTOF is ~4500 at m/z 200. The elements C, H, O, N, and S were used 133 134 for peak assignment. Peaks that cannot be assigned with these elements within 10 ppm are labeled as "unknown" or 135 "others". The relative mass-dependent transmission efficiency in the HTOF was calibrated through the depletion method when the APi-HTOF was operated with a nitrate chemical ionization (CI) inlet in front, for which the voltages were kept 136 137 the same as in the APi-mode (Heinritzi et al., 2016). Detailed descriptions of the measurement and data analysis have been described in our previous study (Yin et al., 2021). In Hyytiälä, ions were sampled through a core sampling method. 138 It has a thick and long tube outside and a thin and short coaxial tube inside to minimize the sampling losses with a total 139 140 flow rate of 5 LPM. The details of the APi-TOF measurements in Hyytiälä can be found in a previous study (Yan et al., 2018). For further analysis in this study, we quantified the negative cluster ions in both Beijing and Hyytiälä using the 141 *in-situ* quantification method, as will be discussed in Section 2.2. 142

- 143 The neutral gas molecules were measured by a (nitrate-) CI-APi-LTOF (Bertram et al., 2011; Jokinen et al., 2012) from Jan 23 to April 14, 2018, in urban Beijing. It was operated side by side with the APi-TOF to achieve simultaneous 144 145 measurements of neutral molecules and cluster ions. NO3⁻ and its adducts (HNO3)1-2NO3⁻ were used as the reagent ions 146 to chemically ionize the gas molecules with a laminar flow ionization source mounted in front of the APi-LTOF (Eisele 147 and Tanner, 1993). Nitric acid is volatilized and carried by a total of 20 LPM sheath flow, then it is exposed to soft X-148 ray to produce NO₃⁻ and its adducts (HNO₃)₁₋₂NO₃⁻. Mostly the species that can be ionized by NO₃⁻ and its adducts $(HNO_3)_{1-2}NO_3^-$ are highly oxygenated molecules, such as H_2SO_4 and OOMs. The mass resolution of the (nitrate-) CI-149 APi-LTOF is ~8000 at m/z 200. The peak fitting and assignment of the molecular compositions followed our previous 150 151 studies where binPMF solutions were used as assistants for the peak assignment (Nie et al., 2022). The sensitivity for 152 H_2SO_4 was calibrated via the controlled reaction between SO_2 and OH at the end of the campaign (Kürten et al., 2012; 153 Li et al., 2019). The relative mass-dependent transmission efficiency of the CI-APi-LTOF was calibrated using the 154 depletion method (Heinritzi et al., 2016). OOMs were quantified assuming they have the same collision-limit charging efficiency as H₂SO₄ with reagent ions, and the mass-dependent transmission efficiency is also used for the quantification 155 156 of OOMs. The calibration results have been shown in our previous study (Yin et al., 2021).
- The ion mobility distribution was measured by a NAIS (NAIS, Airel Ltd.) (Mirme and Mirme, 2013) from Jan 12th to Dec 31st, 2018 in urban Beijing, and from Jan 1st to Dec 31st, 2013 in Hyytiälä (Kontkanen et al., 2013). In both stations,
- the measurement periods of NAIS covered the measurement periods of APi-TOF. NAIS can simultaneously measure positive and negative ions with ion mobility of 3.2-0.0013 cm²·V⁻¹·s⁻¹, corresponding to the mobility diameter range of
- 161 0.8-40 nm. In practice, the NAIS switched between detecting naturally charged ions and total particles (including neutral
- 101 0.6-40 mil. In practice, the WARS switched between detecting naturally enarged fors and total particles (including neural

162 and charged particles) by switching off/on a unipolar corona charger. The measurement cycle period was 2 min for the ion mode, 2 min for the particle mode, and 30 s for offset. For the ion measurement, two cylindrical mobility 163 spectrometers were operated in parallel to measure positive and negative ions simultaneously. In this study, we focus on 164 the negative cluster ions measured in the ion mode, whose ion mobility is larger than 0.8 cm²·V⁻¹·s⁻¹ and diameter smaller 165 than ~ 1.6 nm. A total of 109 mobility bins in the negative ion mode were measured in urban Beijing, while a total of 28 166 mobility bins were measured in Hyytiälä. In urban Beijing, ambient air is drawn into the system through a 1.8 m long 167 copper tube with a diameter of 4 cm, positioned at the window. The sample flow rate is maintained at 54 LPM. The 168 inversion kernel of NAIS was calibrated based on the method described by Wagner et al. (2016) and sampling losses 169 were further considered in the data inversion. The results of NAIS were calibrated considering both the transfer functions 170 and sampling losses (Mirme and Mirme, 2013; Wagner et al., 2016). 171

172 **2.2** The quantification of negative cluster ion

Although we have calibrated the relative mass-dependent transmission efficiency in the APi-HTOF, there are still two problems for quantification: (1) the absolute transmission efficiency is unknown; (2) the relative transmission efficiency curve is calibrated by adding a CI-inlet in front of the APi-HTOF, and the influences of CI-inlet remain unknown. For example, a previous study using a high-resolution differential mobility analyzer and electrometer to calibrate the transmission efficiencies under the APi-mode and CI-mode found that the transmission efficiency under the CI-mode shifts to larger m/z compared to APi-mode (Heinritzi et al., 2016).

- To address these two problems in quantifying APi-HTOF, we used an improved method to obtain the absolute transmission efficiency of the APi-HTOF through the comparison with the synchronous *in-situ* measurement from the NAIS. The key to linking the measurement results of ions by the APi-HTOF with those by the NAIS is to convert the m/z of ions to ion mobilities. As the resolving power of APi-HTOF is higher than that of NAIS, the ion signals in APi-HTOF in specific m/z ranges were summed up to compare with the ion concentrations measured by the NAIS.
- For some specific ions with known peak assignments and structures, like NO_3^- and HSO_4^- , the ion mobility can be directly measured by ion mobility spectrometers (IMS). The results reported in previous literature (Jen et al., 2015; Liang et al., 2013; Spangler and Collins, 1975; Stano et al., 2008) are summarized in Table S1 and were applied in this study. For the remaining ions, like the hundreds of complex organic ions, the Stokes-Millikan equation was applied to convert their m/z to mobilities (Ehn et al., 2011; Friedlander, 2000; Tammet, 1995), as in Eq. 1.

189
$$Z = \frac{1}{\sqrt{1 + m_g / m}} \frac{Q}{3\pi\mu} \frac{1 + Kn(1.257 + 0.4e^{-1.1/Kn})}{d_m + d_g}$$
Eq.1

where Z is the electrical mobility, m_g the mass of a carrier gas molecule, m the mass of ions, Q the charge of ions, μ the viscosity of the carrier gas, Kn the Knudsen number, d_m the mobility diameter of ions, and d_g the diameter of the carrier gas molecule. Here, the carrier gas is ambient air, and m_g , d_g , and μ are adopted as 29.0 g/mol, 0.3 nm, and 1.73×10^{-5} kg·m⁻¹·s⁻¹ at 278 K and 101.3 kPa (Ku and De La Mora, 2009). *Kn* and d_m are calculated as Eq. 2-3.

194
$$K_n = \frac{2\lambda}{d_m + d_g}$$
 Eq.2

$$d_m = \sqrt[3]{\frac{6m}{\pi\rho}}$$
 Eq.3

where λ is the mean free path of carrier gas (62 nm at 278 K and 101.3 kPa) and ρ the density of ions. The appropriate 196 density for calculating the mobility of ions is controversial. Ehn et al. (2011) used the bulk density of aconitic acid (1.66 197 198 g·cm⁻³) for the conversion, but molecular clusters may have different densities with bulk substances. To address this problem, we compared the measured mobility of an IMS for different types of OOMs with the calculated mobilities 199 according to the Stokes-Millikan method using 1.1, 1.3, and 1.6 g·cm⁻³ as the densities, respectively (Fig. S1). The 200 experimental results of the IMS are from Krechmer et al. (2016). By comparing the measured and calculated mobilities 201 202 for organic ions with different compositions, densities of 1.1 and 1.3 g·cm⁻³ were ultimately used in the mass-mobility 203 conversion of ions with m/z smaller than 150 or larger than 200, respectively, whereas between m/z of 150 and 200, a linear curve of density is used. Though some ions with m/z larger than 400 fall into the region of 1.1 g cm^3 in Fig. S1, 204 they were mostly OOM dimers that did not exist in Beijing (Qiao et al., 2021). Thus a density of 1.3 g·cm⁻³ was applied 205 for large organic ions in Beijing. Finally, the mobilities of all ions were adjusted according to the ambient temperature 206 and pressure before being applied in the calibration of APi-HTOF (Tammet, 1995). 207

208 Using the synchronous *in-situ* measurement results of the NAIS as the reference, we obtained the detection efficiency of 209 the APi-HTOF for ions with different m/z. Here, since the ion concentrations measured by the NAIS have already 210 considered sampling losses and represent atmospheric concentrations, the obtained detection efficiency of the APi-HTOF 211 should represent a combined result of the transmission efficiency of the APi-HTOF and the sampling efficiency in the 212 sampling line of the APi-HTOF. The ions used in the APi-HTOF are in the m/z range from 32 to 908, corresponding to the mobility diameters of 0.8-1.6 nm and ion mobilities of 3.16 to 0.77 cm²·V⁻¹·s⁻¹, covering 12 mobility bins of the 213 214 NAIS measurements. The number concentrations and signal intensities measured by the NAIS and APi-HTOF had good consistencies (r \ge 0.66) in the 12 bins, and the correlations were the best over the *m/z* range of 331-791 (r > 0.94) (Fig. 215 216 S2). The good correlation between the APi-HTOF and NAIS indicates that both instruments captured the variation of the atmospheric negative cluster ions. Then the detection efficiency of a certain m/z in the APi-HTOF was calculated as the 217 slope of the linear fitting line between signal intensities acquired in the APi-HTOF and number concentrations in the 218 219 NAIS. Compared with the detection efficiencies calculated from the mass-dependent density, using a fixed density of 1.66 g·cm⁻³ would overestimate the detection efficiency for small ions and underestimate that for large ions, with a 220 deviation of up to 40% (Fig. S3). 221

222 After excluding the effect of sampling losses in the APi-HTOF, the absolute transmission efficiency of the APi-HTOF was derived, and the obtained pattern agreed well with the relative transmission efficiency as shown in Fig. 1. The 223 sampling losses of ions in the APi-TOF is assumed the same as those of neutral particles with the same sizes (Mahfouz 224 225 and Donahue, 2021). As the mass-dependent density was applied, the absolute and relative transmission efficiency curves 226 showed similar patterns, i.e., both peaked at $\sim m/z$ 400 and the maximum absolute transmission efficiency reached ~4.9%. The discrepancy between the absolute and relative transmission efficiency at small values of m/z may be caused by the 227 uncertainty of small ion measurements in the NAIS, sampling efficiency calculation, or the influences of CI-inlet, as the 228 229 voltage settings remain the same during the calibration experiment. It would result in an uncertainty of ~28% for the 230 quantification of small ions. The absolute transmission efficiency obtained by our calibration is roughly the same order of magnitude as that in previous studies, and the shape is similar as well (Heinritzi et al., 2016; Junninen et al., 2010). Finally, the fit line of the detection efficiency was applied to quantify the ion compositions measured by the APi-HTOF, considering both the sampling efficiency and the transmission efficiency (Fig. 1).

234

Fig. 1 The detection and transmission efficiency curves

235 After the quantification, the total negative cluster ion concentrations measured by the APi-HTOF were consistent with 236 those measured by the NAIS (Fig. 2). However, when the total ion concentration was lower than ~70 cm⁻³, the total ion concentration measured by the NAIS was higher than that measured by the APi-HTOF. Specifically, the total cluster ion 237 concentration measured by the NAIS was still higher than 25 cm⁻³ even when ions detected by APi-HTOF approached 238 zero. These signals detected by the NAIS are mainly from small cluster ions (3.16-1.28 cm²·V⁻¹·s⁻¹, 0.8~1.27 nm), the 239 fraction of which rises to >80% of the total counts when the total cluster ion concentration measured by the APi-HTOF 240 241 is lower than ~70 cm⁻³. As the APi-HTOF has a much lower detection limit than the NAIS (Fig. S4), these signals could be caused by the high noise of small ions in the NAIS. The detection limit of the APi-HTOF is the lowest at a diameter 242 of 1.4 nm, which is ~ 0.02 cm⁻³, and it increases when the detected ions move to lower or higher sizes due to the decrease 243 in detection efficiency. Under a minimum diameter of 0.9 nm and a maximum of 1.7 nm, the detection limits of APi-244 HTOF were ~ 0.4 and ~ 4 cm⁻³, respectively. The detection limit of the NAIS decreases with an increase in ion diameter. 245 246 The detection limit of the NAIS for cluster ions varies from 300 to 30 cm⁻³ at diameters ranging from 0.8 to 2 nm. Thus, 247 the ion concentration measured by the APi-HTOF is more reliable than that measured by the NAIS when the ion 248 concentration is extremely low. As the influence of background noise would diminish when signals increase and the 249 detection efficiency is calculated using data during the whole period, the background noise in the NAIS has limited 250 influence on the calibration we have done above.

251

Fig. 2 Ion concentrations measured by APi-HTOF and NAIS

252 **2.3** The simulation of HSO₄⁻ and C₃H₃O₄⁻ concentration

253 To quantitatively address the sources and sinks of air ions, we simulate the concentration of ions using the mass balance equation and identify their dominant production and loss pathways. The simulated concentrations are compared with the 254 255 measured concentrations. HSO_4^- and $C_3H_3O_4^-$ were chosen as they are one of the most abundant negative ions in different atmospheric environments and their reaction rates with neutral molecules are determined. The production pathways 256 considered in this simulation are the ion-molecular reactions between NO3⁻ and neutral molecules H₂SO₄ and C₃H₄O₄. 257 Previously, it was reported that NO₃⁻ was formed in the earlier stage of atmospheric ion formation (Beig and Brasseur, 258 259 2000; Luts, 1995) and it continues to ionize neutral gaseous that are more acidic than HNO₃, such as H₂SO₄ and C₃H₄O₄ 260 (Eisele, 1989; Tanner and Eisele, 1991). The loss pathways in the atmosphere mainly include condensational loss onto the particles, ion-molecule reactions, and ion-ion recombination with ions of opposite charge. The formation of H_2SO_4 261 262 dimer ions could be an important loss pathway for HSO_4^- (Beck et al., 2022). $C_3H_3O_4^-$ could further react with H_2SO_4 263 and change back to its neutral form $C_3H_4O_4$ (Beig and Brasseur, 2000). Thus, the following ion-molecule reactions are considered in the dynamic models to simulate the formation of HSO_{4^-} and $C_3H_3O_{4^-}$ in urban Beijing. 264

$$NO_3^- + H_2SO_4 \xrightarrow{\kappa_1} HSO_4^- + HNO_3$$

$$NO_3^- + C_3H_4O_4 \xrightarrow{\lambda_2} C_3H_3O_4^- + HNO_3$$
R.2

267
$$C_3H_3O_4^- + H_2SO_4 \xrightarrow{k_3} HSO_4^- + C_3H_4O_4$$

$$HSO_{4}^{-} + H_{2}SO_{4} \xrightarrow{k_{4}} H_{2}SO_{4} \cdot HSO_{4}^{-} R.4$$

R.3

where k_1 to k_4 are the rate constants of these reactions, which are 2.32×10^{-9} , 2.5×10^{-9} , 2.0×10^{-9} , and 2.0×10^{-9} cm³·s⁻ respectively (Beig and Brasseur, 2000; Lovejoy and Curtius, 2001; Viggiano et al., 1997). The ion-molecule reactions between HSO₄⁻ and organic molecules are not considered in this model due to the lack of reaction rate constants.

272 The mass balance equations for HSO_4^- and $C_3H_3O_4^-$ can be written as Eq.4 and Eq.5, respectively.

273
$$\frac{d[\text{HSO}_{4}^{-}]}{dt} = k_{1}[\text{NO}_{3}^{-}][\text{H}_{2}\text{SO}_{4}] - k_{4}[\text{HSO}_{4}^{-}][\text{H}_{2}\text{SO}_{4}] - \text{CS}[\text{HSO}_{4}^{-}] - \alpha n^{+}[\text{HSO}_{4}^{-}]$$
Eq.4

274
$$\frac{d[C_{3}H_{3}O_{4}^{-}]}{dt} = k_{2}[NO_{3}^{-}][C_{3}H_{4}O_{4}] - k_{3}[C_{3}H_{3}O_{4}^{-}][H_{2}SO_{4}] - CS[C_{3}H_{3}O_{4}^{-}] - \alpha n^{+}[C_{3}H_{3}O_{4}^{-}]$$
Eq.5

where CS[HSO₄⁻] and CS[C₃H₃O₄⁻] are loss rates caused by the condensational loss onto particles; αn^+ [HSO₄⁻] and αn^+ [C₃H₃O₄⁻] are loss rates caused by the ion-ion recombination between ions of opposite charge; the value of α is assumed to be 1.6×10⁻⁶ cm³·s⁻¹ (Bates, 1982; Zauner-Wieczorek et al., 2022); n^+ is the total concentration of positive ions and is measured by NAIS. Assuming a steady state, the concentration of HSO₄⁻ and C₃H₃O₄⁻ are calculated as Eq.6 and Eq.7, respectively. HSO₄⁻ formed through the ion-molecule reaction between C₃H₃O₄⁻ and H₂SO₄ is neglected considering the low concentration of C₃H₃O₄⁻ in Beijing.

281
$$[HSO_{4}^{-}] = \frac{k_{1}[NO_{3}^{-}][H_{2}SO_{4}]}{CS + k_{3}[H_{2}SO_{4}] + \alpha n^{+}}$$
Eq.6

282
$$[C_{3}H_{3}O_{4}^{-}] = \frac{k_{5}[NO_{3}^{-}][C_{3}H_{4}O_{4}]}{CS + k_{2}[H_{2}SO_{4}] + \alpha n^{+}}$$
Eq.7

283 **3. Results and discussion**

284 **3.1 Negative cluster ion concentration and composition in urban Beijing**

The median concentration of negative cluster ions is 85 cm⁻³ in Beijing during the measurement period, and its 25th and 75th percentiles are 61 cm⁻³ and 112 cm⁻³, respectively. These values are significantly lower than concentrations reported at typical rural, forest, and marine sites (Fig. 3a). The total concentration of air ions ranges between 14 cm⁻³ and 467 cm⁻ ³ in urban Beijing, similar to that in the Indian urban city, Pune, which is also a highly polluted site (Gautam et al., 2017). Ion concentrations in these two polluted urban cities are so far the lowest, significantly lower than at other urban, rural and marine, and forest sites, which are mostly above 200 cm⁻³ (Chen et al., 2017; Dos Santos et al., 2015; Hirsikko et al., 2011; Tammet, 2015a; Tammet, 2015b). It should be noted that although Paris is an urban site, its air is relatively
clean in terms of aerosol concentration compared to the Chinese and Indian polluted urban sites. Similarly, cluster ion
concentrations at some urban sites can also be as high as those at rural and remote sites due to the relatively clean
atmosphere (Hirsikko et al., 2011).

295

Fig. 3 Ion concentration comparison between Beijing and other sites.

296 The ion concentrations are negatively correlated with CS at both the urban site Beijing and the clean forest site Hyytiälä. 297 In urban Beijing, the median ion concentration decreases from 181 to 51 ions cm⁻³ when CS increases from 0.0032 to 298 0.12 s⁻¹ (Fig. 3b). The negative correlation between the ion concentration and CS exists for all the mobility ranges and 299 is more intensive for ions with mobilities between 1.72 and 0.90 cm²·V⁻¹·s⁻¹, corresponding to a diameter range of about 300 1.03-1.44 nm (Fig. S5). A similar negative correlation is also observed in Hyytiälä, where the median ion concentration decreases from 606 to 369 ions cm⁻³ when CS increases from 0.00017 to 0.012 s⁻¹. Whereas, the inverse relation between 301 302 the negative cluster ion concentration and CS is much stronger in Beijing than in Hyytiälä. Combining data from these 303 two sites, a negative correlation can still be seen over a wide CS range from 0.00017 to 0.12 s⁻¹. The influence of CS on 304 the positive cluster ions also exists, as positive and negative ions are closely related to each other (Fig. S6). Thus, CS is 305 an important influencing factor for the total cluster ion concentrations in both Beijing and Hyytiälä. This is very likely 306 the reason for the lowest ion concentrations at the two polluted sites (Beijing and Pune, Fig. 3a) having the highest 307 aerosol mass loadings.

308 In urban Beijing, the estimated lifetime of cluster ions due to the combined effect of condensation loss and ion-ion recombination of cluster ions is 0.02-0.4 min during haze periods ($PM_{2.5} > 75 \ \mu g/m^3$, average CS = 0.091 s⁻¹) and 0.1-309 310 1.6 min during clean periods ($PM_{2.5} < 75 \ \mu g/m^3$, average CS = 0.023 s⁻¹) (Fig. S7). The lifetime of cluster ions in Hyytiälä is between 0.8-14 min (average $CS = 0.0018 \text{ s}^{-1}$), much longer than that in Beijing. Such distinct CS-dependence of total 311 312 cluster ions in urban Beijing is rarely reported in previous studies despite that the condensational loss of ions on large 313 particles has been widely recognized as important. On the contrary, most of the studies have found that weather 314 conditions and ionizing radiation are the controlling factors for ion concentrations. For example, ion production rates in 315 Hyytiälä were reported to be largely affected by variations in seasonal radiation and wind speed (Chen et al., 2016). This 316 indicates that CS may be the driving factor of cluster ion concentration at highly polluted sites with high aerosol mass 317 loadings and relatively constant ion production rates, while ion production rates may be the driving factor at relatively 318 clean sites where aerosol mass loadings are low and ion production rate varies significantly (Hirsikko et al., 2007).

The composition of negative cluster ions is mainly comprised of inorganic nitrogen-containing ions, inorganic sulfurcontaining ions, and organic ions in urban Beijing, accounting for 20-22%, 8-15%, and 37-43%, respectively (Fig. 4 & Fig. S8). NO_2^- , NO_3^- , and $HNO_3NO_3^-$ are the most abundant among the detected inorganic nitrogen-containing ions. They possibly exist in the forms of $NO_2^-(H_2O)_n$, $NO_3^-(H_2O)_n$, and $NO_3^-(H_2O)_n$ in the atmosphere (Luts, 1995), 323 with the loosely bounded water molecules being evaporated from the cluster ions when passing through the mass 324 spectrometer. The concentration of NO₂⁻ and NO₃⁻ were observed to be well correlated with each other in the atmosphere 325 of Beijing, and the concentration of NO_2^- is ~20 % of that of NO_3^- (Fig. S9a). This is consistent with the ion chemical 326 models suggesting that NO_2^- and NO_3^- can be stably formed through a series of reactions between primary ions, NO, NO₂, and HNO₃ in the atmosphere (Beig and Brasseur, 2000; Kawamoto and Ogawa, 1984). HNO₃NO₃⁻ is subsequently 327 328 formed by adding an HNO₃ molecule to NO₃⁻. Inorganic sulfur-containing ions are mainly in the form of HSO₄⁻, SO₅⁻, 329 and $H_2SO_4HSO_4^-$. The ion HSO_4^- is mainly produced by the ion-molecule reaction between NO_3^- and H_2SO_4 which will 330 be discussed hereinafter. $H_2SO_4HSO_4^-$ is formed by the further addition of an H_2SO_4 molecule, which is proposed as the 331 first step of ion-induced nucleation (Lovejoy, 2004). SO_5^{-1} is likely to be generated through the reaction between O_2 and 332 SO_3^- (Möhler et al., 1992), and its signal variation is similar to that of HSO_4^- (Fig. S9b). Similar compositions and 333 variations of inorganic sulfur-containing ions were also observed at Hyytiälä (Ehn et al., 2010).

334

Fig. 4 Ion composition in Beijing and Hyytiälä

335 The negative organic ions are mainly CHO-related or CHON-related organic ions in the form of the adduct with NO_3^- or 336 HSO₄⁻ (Fig. 4). Only minor fractions (~4%) are in the deprotonated form of CHO⁻ or CHON⁻. These indicate that the 337 ionization schematic of organic ions is mainly through ion-molecular reaction with NO_3^- or HSO_4^- in Beijing. As few neutral sulfur-containing organics were observed, it is unlikely that the identified CHON-HSO4⁻ is a cluster of sulfur-338 339 containing organics and NO₃-, but rather a cluster of nitrogen-containing organics and HSO₄-. Among all negative organic 340 ions, CHON organic ions adducted with NO_3^{-} (CHON·NO₃⁻) are the most abundant and account for 56% and 69% during clean and haze periods, respectively. Here the haze periods were identified based on whether PM_{2.5} concentration is 341 higher than 75 μ g·m⁻³. Though the fraction of CHON·NO₃⁻ does not change much, their compositions are significantly 342 343 different between clean and haze periods, mainly influenced by the differences in neutral gaseous molecules. During the haze periods where the average NO_x concentration is 30.1 ± 14.6 ppb, the organic ions are dominated by nitrated phenols 344 345 (NPs), such as $C_6H_5NO_3 \cdot NO_3^-$ and $(C_6H_5NO_3)_2 \cdot NO_3^-$. During the clean periods when the average NO_x decreases to 15.0 ± 11.5 ppb, a series of organic ion peaks were observed for m/z above 300. The highest peaks in the series include 346 C₁₀H₁₅NO₁₀·NO₃⁻, C₁₀H₁₄O₁₁·NO₃⁻, and C₁₀H₁₆NO₁₁·NO₃⁻, which are possibly the adducts of NO₃⁻ and highly 347 348 oxygenated products from monoterpenes under high NO_x conditions (Yan et al., 2020). NPs have high signals in urban Beijing and their properties are different from most other CHON compounds, such as volatilities and charging 349 350 efficiencies as will be discussed below. Thus, CHON species were divided into CHON_{NPs} and CHON_{nonNPs} groups in the 351 following analysis (Nie et al., 2022).

The negative cluster ion spectra in urban Beijing are significantly different from that in Hyytiälä, possibly due to the high NO_x. Contrary to Beijing, CHO-related organic ions are more abundant than CHON-related in Hyytiälä, with fractions of 67% and 33%, respectively. The higher CHON ion fractions in urban Beijing are consistent with previous studies showing that neutral OOMs are composed of more nitrogen-containing species in urban Beijing compared to 356 Hyytiälä due to the high NO_x concentration (Li et al., 2022; Qiao et al., 2021). In addition, the higher CHON_{NPs} ion fractions in urban Beijing are also the result of high anthropogenic emissions of aromatics and high NO_x concentrations 357 358 (Cheng et al., 2021). Another difference is that the monoterpene-related OOM dimer ions are abundant in Hyytiälä 359 (Isidorov et al., 1985) but are rarely observed in Beijing. This is also related to the neutral molecules as the formation of neutral OOM dimers would be suppressed under high NO_x conditions (Nie et al., 2022). Moreover, a larger fraction of 360 361 the organic ions are in the form of adducts with HSO_4^- in Beijing than in Hyytiälä. This is related to the higher HSO_4^- 362 /NO₃⁻ ratio in Beijing (~0.36) compared with Hyytiälä (~0.05). The fraction of HSO₄⁻ adducted ions would increase during clear days in Hyytiälä but its fraction is still lower than those in Beijing (Bianchi et al., 2017). Despite the different 363 compositions between urban Beijing and forest Hyytiälä, the ion mobility distributions are generally quite similar (Fig. 364 S10). The mode mobility peaks of ions for both sites are $\sim 2 \text{ cm}^2 \cdot \text{V}^{-1} \cdot \text{s}^{-1}$, corresponding to a mobility range dominated 365 366 by HSO₄- dimer, trimer, and organic ions. These indicate that similar ion mobility distributions do not necessarily indicate 367 similar ion compositions and direct measurements should be performed.

368 Comparing the organic ions and neutral molecules measured by the APi-HTOF and the (nitrate-) CI-APi-LTOF, we found 369 that the composition of organic ions is similar to that of neutral molecules after subtracting the reagent ions (Fig. S11). 370 The element number distributions of organic compounds detected in the APi-HTOF and CI-APi-LTOF are similar, in which the overlapping region has C atoms in the range of 3 to 14, O atoms in the range of 1 to 15, and N atoms in the 371 372 range of 0 to 2 (Fig. S12). There are in total 293 organic species measured both in organic ions and neutral molecules, and in addition, inorganic sulfur-containing ions such as (H₂SO₄)₀₋₃HSO₄⁻ were measured in both instruments as well. 373 The neutral molecules detected by CI-APi-TOF was identified by subtracting one NO3. Such similar elemental 374 375 distributions and abundant common ions in APi-HTOF and CI-APi-LTOF indicate that naturally charging processes in 376 the atmospheres are quite similar to the NO3⁻ charging processes in the CI-APi-TOF, despite that HSO4⁻ also has a 377 considerable concentration in the atmosphere and would charge neutral molecules as efficiently as NO₃⁻. Here the reagent 378 ions such as NO₃⁻ and HSO₄⁻ were already subtracted and only the neutral molecular formulas were compared. Organic 379 compounds with carbon atom numbers less than 3 were not considered in our analysis. In addition, most molecules 380 detected only by CI-APi-TOF contain more C atoms of up to 20. This may be because the concentrations and signals of 381 organic ions with such high carbon numbers are too low to be assigned with a certain formula in the APi-HTOF.

382 **3.2 Formation of negative cluster ions**

To further characterize the formation of negative cluster ions, the influences from the reagent ions (i.e., NO_3^- and HSO_4^-) and the neutral molecules were explored in this section. Ions in form of adducts with NO_3^- and HSO_4^- were compared to explore their ionization selectivity towards different compounds. The concentration variations of cluster ions were compared with that of the neutral molecules and the reagent ions to explore their driving factors. Then the charge fractions for different compositions were compared.

388 **3.2.1 The influence of reagent ions**

389 The two main reagent ions, NO₃⁻ and HSO₄⁻, have different concentrations and variations in urban Beijing (Fig. S13). 390 Firstly, NO₃⁻ is about 3 times higher than HSO₄⁻. During the measurement period, the concentration of NO₃⁻ is 9.6 ± 6.0 391 cm⁻³, and that of HSO₄⁻ is 3.2 ± 3.4 cm⁻³ in urban Beijing. Similarly, NO₃⁻ is also more abundant than HSO₄⁻ in Hyytiälä, indicating that NO_3^- has greater chances to collide with neutral molecules than HSO_4^- in the atmosphere. Secondly, both 392 393 NO_3^- and HSO_4^- increase with the increase of total negative cluster ions in urban Beijing, while it is reversed in Hyytiälä. 394 This illustrates the significant impact of CS on the concentration of cluster ions in urban Beijing, that is, with the decrease 395 of CS, both the concentrations of reagent ions and product ions increase to different degrees. Moreover, the variation of 396 HSO_4^- is more significant than that of NO_3^- in urban Beijing. NO_3^- maintains a relatively constant proportion (15%±5%) 397 in the total negative cluster ions during the whole sampling period in urban Beijing, while the fraction of HSO_4^- varies significantly due to the strong diurnal variation of H₂SO₄. The reason for the relatively constant NO₃⁻ fraction in total 398 399 negative cluster ions needs further investigation, and this phenomenon might offer a chance of combining NO_3^- and organic ions attached to NO₃⁻ to predict the neutral molecules. 400

The compositions of molecules that cluster with NO₃⁻ and HSO₄⁻ are generally similar, with NO₃⁻ clustering with a few 401 402 more compounds with high carbon numbers and low oxygen numbers. Fig. 5a shows the carbon atom numbers and 403 average carbon oxidation states (\overline{OS}_{C}) of molecules that are attached to NO₃⁻ and HSO₄⁻. For molecules with less than 8 404 carbon atoms, molecules attached to NO₃⁻ and HSO₄⁻ has similar \overline{OS}_{C} between -3 and 3. For molecules with more than 405 8 carbon atoms, molecules with lower \overline{OS}_{C} (<-0.5) are only attached to NO₃. The average \overline{OS}_{C} of the molecules attached to NO₃⁻ and HSO₄⁻ are -0.3 and 0.22, respectively, and the former is close to the molecules detected in CI-APi-LTOF (-406 407 0.45). Similarly, the average O/C ratio for molecules adducting with NO_3^- and neutral molecules detected in CI-APi-408 LTOF are 0.85 and 0.82, respectively, slightly lower than the average O/C ratio of molecules adducting with HSO₄-(1.03). These indicate that the selectivity of HSO_4^- and NO_3^- towards organic molecules are similar, with only slight 409 410 differences for compounds with high carbon numbers and low oxygen numbers. The slight difference may be partly 411 because the concentrations of these adductes of these compounds and HSO₄⁻ are too low to be accurately distinguished. Overall, O/C ratio of compounds adducted with both NO₃⁻ and HSO₄⁻ are both significantly higher than those detected 412 413 by the (iodide)CI-APi-TOF and (oxygen)CI-APi-TOF (Li et al., 2021; Riva et al., 2019), indicating their higher 414 selectivity for oxidized compounds, which is consistent with previous theoretical simulations (Nadykto et al., 2018). In 415 addition, when only CHO and CHONnonNPs molecules clustering with both NO3- and HSO4- are considered, their volatility 416 distributions are very similar when clustering with NO_3^- or HSO_4^- , which further indicates that the selectivity difference 417 between NO₃⁻ and HSO₄⁻ towards organic molecules is very small. (Fig. 5b & Fig. S14).

418

Fig. 5 The species of organic ions adducting with NO₃⁻ and HSO₄⁻

419 For the 88 species that form clusters with both NO_3^- and HSO_4^- , the ratio of the two ionization forms (M·HSO₄⁻/M·NO₃⁻)

420 is positively related to the ratio of reagent ions (HSO_4^{-}/NO_3^{-}). In addition, HSO_4^{-} forms clusters with CHO and CHON_{nonNPs} more efficiently, while NO₃⁻ forms clusters with CHON_{NPs} more efficiently, despite the above-mentioned 421 422 similarity in their selectivity towards different organic molecules. There are an overall of 33 CHO, 40 CHON_{nonNPs}, and 423 15 CHON_{NPs} species that both form ions with NO₃⁻ and HSO₄⁻. The average ratio of HSO₄⁻/NO₃⁻ is 0.36, and the relative 424 ratio of M·HSO₄⁻/M·NO₃⁻ are 0.74, 0.73, and 0.23 for CHO, CHON_{nonNPs}, and CHON_{NPs} species, respectively. Generally, 425 M·HSO₄⁻/M·NO₃⁻ ratio is positively related to HSO₄⁻/NO₃⁻ ratio, and the dots are close to the 1:1 line as shown in Fig. 6. 426 Specifically, the dots for CHON_{NPs} are mostly lower than 1:1 line, especially under high H₂SO₄ concentration. Similar 427 decreasing trends were also observed for the M·HSO₄-/M·NO₃⁻ ratio for CHO and CHON_{nonNPs}. This is possibly due to 428 the competition between different neutral compounds for the reagent ions, as the atmospheric concentration of ions is up 429 to 7 orders of magnitudes lower than neutral molecules. For example, gaseous H₂SO₄ may compete with the organic 430 molecules to react with HSO4-. As a highly electronegative substance, H2SO4 prefers to be negatively charged and form 431 strongly bounded clusters with alkaline substances such as HSO₄⁻ or amines in the atmosphere, thus HSO₄⁻ would prefer 432 to cluster with H₂SO₄ during daytime and its efficiency of clustering with organic molecules drops as a result. During 433 the nighttime, when the competition from H₂SO₄ weakens, the formation of M·HSO₄⁻ increases.

434

Fig. 6 The ratio between organic ions adducting with NO3⁻ and HSO4⁻

435 **3.2.2** The influence of neutral molecules

436 The diurnal cycles of negative cluster ions are mainly determined by their corresponding neutral molecules, and 437 exceptions were found for CHON_{NPs}-related ions, the diurnal cycles of which are more related to the reagent ions (Fig. 7). The reagent ions and their inorganic clusters ((HNO₃)₀₋₂NO₃⁻ and (H₂SO₄)₀₋₂HSO₄⁻) both reach their maximum at 438 439 13:00 but show different diurnal patterns. (H_2SO_4)₀₋₂HSO₄ has a single peak pattern that resembled neutral H_2SO_4 , while 440 (HNO₃)₀₋₂NO₃⁻ has a three-peak pattern that matched the total concentration of negative cluster ions. For the organic ions, the CHO and CHONnonNPs-related ions both peak in the afternoon, which resemble the diurnal pattern of 441 442 corresponding neutral molecules. However, the diurnal patterns of CHON_{NPs}-related ions and neutral CHON_{NPs} show opposite variations, and that of CHON_{NPs}-related ions is closer to the diurnal variation of NO₃⁻. As CHON_{NPs}-related 443 444 ions reach their maximum at nighttime, neutral CHON_{NPs} reach their maximum at both noon and evening, indicating 445 that the formation of CHON_{NPs}-related ions may be different from other organic ions. Besides, although the diurnal variations of CHO and CHON_{nonNPs}-related ions roughly follow those of neutral molecules, it was found that their peak 446 times coincided with that of (HNO₃)₀₋₂NO₃⁻ at 13:00 and 17:00. Differently, the diurnal patterns of CHO and 447 448 CHON_{nonNPs}-related ions tended to peak during nighttime in Hyytiälä (Bianchi et al., 2017). This could be due to the different diurnal patterns of neutral molecules as those of the reagent ions are similar between Beijing and Hyytiälä. 449

450

Fig. 7 The diurnal patterns of negative cluster ions, neutral molecules, and charge fractions

451 Consistently, the cluster ion concentrations were found to be positively correlated with that of neutral molecules under

similar CS for CHO and $CHON_{nonNPs}$ species during the whole sampling period. As shown in Fig. 8, generally, the concentrations of all organic ions decrease with increasing CS as indicated in Section 3.1. When CS is larger than $0.01s^{-1}$, the concentrations of CHO and $CHON_{nonNPs}$ -related ions rise with their corresponding neutral gas concentration under similar CS levels, while that of $CHON_{NPs}$ -related ions remains constant or slightly declines. When CS is smaller than 0.01 s⁻¹, which is usually accompanied by low concentrations of neutral molecules, the relationships between organic ions and the corresponding neutral molecules are not obvious, this may be due to a combined effect of neutral molecules and the reagent ions.

459

Fig. 8 The variation of organic ions with neutral molecules under different CS

460 The charge fraction between organic ions and neutral molecules reflects a charging competition between different neutral 461 compounds. Here, we calculate charge fraction as the proportion between the concentrations of cluster ions and their corresponding neutral molecules. As shown in Fig. 7c, different organic molecules have similar diurnal variations of 462 463 charge fractions, which are highest at 5:00 and lowest at 15:00. While the HSO₄-/H₂SO₄ ratio has the opposite pattern 464 and is highest during the daytime. The opposite diurnal patterns between organic ions and HSO_4^- indicates that H_2SO_4 and organic molecules would compete for being charged. The median charge fractions of CHO and CHONnonNPs 465 molecules with NO₃⁻ and HSO₄⁻ are close to each other and range from 2×10^{-7} to 2×10^{-6} , while the charge fractions of 466 CHON_{NPs}-related ions are much lower than these organic ions. This suggests that CHON_{NPs} are less likely to form cluster 467 ions with NO_3^- and HSO_4^- than CHO and CHON_{nonNPs}. As the charge fraction for organic ions attached to NO_3^- and 468 469 HSO₄⁻ has similar diurnal patterns, it is deduced that the diurnal variation of charge fractions is mainly driven by the 470 variation of neutral molecules instead of reagent ions.

471 For CHO and CHON_{nonNPs}, the charge fractions between organic ions and neutral molecules generally increase with the 472 carbon atom numbers and \overline{OS}_{C} of the molecules. As shown in Fig. 9, the average charge fractions of different organic ions varie between 3×10⁻⁸ and 8×10⁻⁶, and species with larger carbon atom numbers and higher oxidation states tend to 473 474 have higher charge fractions, either when clustering with NO3⁻ or HSO4⁻ (Fig. S15). The charge fraction difference 475 between molecules with different carbon atom numbers and \overline{OS}_{C} is a reflection of the differences in reaction rates 476 between reagent ions and the neutral molecules or the stabilities of the cluster ions. In summary, through the 477 comprehensive quantitative analysis of atmospheric cluster ions, reagent ions, and neutral molecules, we can roughly 478 predict the composition and concentration variations of atmospheric negative cluster ions from measured neutral 479 molecules and CS, as indicated by Fig. 8. However, as the actual charging fractions for different species differ in a wide 480 range as shown in Fig. 9, the prediction of actual concentrations of atmospheric ions would require more work.

481

Fig. 9 The influence of molecular characteristics on the charge fraction

482 **3.3** Quantification of the sources and sinks of representative negative cluster ions

483 We found that the ionization of H_2SO_4 and $C_3H_4O_4$ by NO_3^- and condensational loss to particles are the main formation

484 and loss pathways for HSO_4^- and $C_3H_3O_4^-$ respectively. As shown in Fig. 10, the simulated concentrations of both $HSO_4^$ and C₃H₃O₄⁻ by Eqs.6-7 have a good consistency with the measured concentrations, indicating that the considered 485 486 production and loss pathways can reproduce the measured ion concentration in urban Beijing. The ionization of H₂SO₄ 487 and $C_3H_4O_4$ by NO₃⁻ is the main production pathway. For both HSO₄⁻ and $C_3H_3O_4^-$, the condensational loss is the main loss pathway, contributing to more than 70% of the total loss rate, which explains why CS is a driving factor for the 488 489 concentration of cluster ions in urban Beijing. Ion-ion recombination process only accounts for 6% and 8% of the total 490 loss rates for HSO_4^- and $C_3H_3O_4^-$, which is due to the low ion concentrations in the polluted urban atmosphere. Notably, the ion-molecular reactions between the ions with H_2SO_4 could contribute to 19% of the loss of both HSO_4^- and $C_3H_3O_4^-$. 491 The transformation from $C_3H_3O_4^-$ back to $C_3H_4O_4$ indicates that the conversion of ions back to neutral molecules is also 492 493 significant.

494

Fig. 10 Simulation of the sources and sinks of HSO₄⁻ and C₃H₃O₄⁻.

495 4 Conclusions

496 We quantified the composition-resolved ion concentrations in the atmosphere by combining mass spectrometry and 497 electrical mobility measurements. The absolute transmission efficiency in the mass spectrometry APi-HTOF obtained 498 with the ion mobility spectrometer (NAIS in this study) agreed well with the relative transmission efficiency obtained in 499 the chemical ionization mode of the APi-HTOF. The calibrated ion concentrations with different m/z ranges agreed well with the concentration of ions with corresponding mobility ranges of the ion mobility spectrometer. These indicate that 500 501 an APi-HTOF can be well calibrated by running an APi-HTOF and an ion mobility spectrometer side by side in ambient 502 measurements. Furthermore, we propose that the transmission efficiency of a CI-APi-TOF can also be obtained through 503 *in-situ* comparison with an ion mobility spectrometer, as long as the X-ray and voltages of a chemical ionization inlet 504 are turned off and atmospheric ions are directly measured.

The cluster ion concentrations and composition in urban Beijing are largely affected by the high condensation sink and nitrogen oxides. Firstly, ion concentration decreased significantly with an increasing CS for cluster ions in all the mobility ranges. Median cluster ion concentrations in urban Beijing were only ~85 cm⁻³, much lower than those reported at clean and rural sites due to the high CS in urban Beijing. Due to high concentrations of nitrogen oxides, the organic ion compositions were composed of more nitrogen-containing species in urban Beijing than those in Hyytiälä. This is consistent with the higher neutral nitrogen-containing molecular fractions in urban Beijing as previously reported.

The formation of cluster ions was analyzed by comparing their concentrations, volatility distributions, and charge fractions with CS, reagent ions, and their corresponding neutral molecules. The organic cluster ions are mainly in the form of adducts with NO_3^- or HSO_4^- . The ratio of the two ionization forms (M·HSO₄⁻/M·NO₃⁻) is positively related to the ratio of reagent ions (HSO₄⁻/NO₃⁻). Although the molecules clustered with HSO₄⁻ and NO₃⁻ are similar in composition,

515	we found that M·HSO ₄ ⁻ formed more efficiently for CHO and CHON _{nonNPs} , while M·NO ₃ ⁻ formed more efficiently with
516	$CHON_{NPs}$ when targeting the compounds that can be ionized by both HSO_4 - and NO_3 The concentrations of organic
517	ions are positively correlated with that of neutral molecules, resulting in their similar diurnal cycles. However, an
518	exception was found for $CHON_{NPs}$, the concentration of which is also significantly influenced by the reagent ions NO_3
519	The charge fractions tend to be higher for organic molecules with higher molecular weight and oxidation state. We also
520	observed a charging competition between different neutral organic and inorganic compounds, such as the competition
521	between H_2SO_4 and organic ions. Through this quantitative analysis, it is possible to infer the variation of atmospheric
522	negative cluster ions from the measured neutral molecules and CS in the urban atmosphere. The formation pathway of
523	HSO ₄ ⁻ and C ₃ H ₃ O ₄ ⁻ were well characterized by the ionization of H ₂ SO ₄ and C ₃ H ₄ O ₄ by NO ₃ ⁻ , and their loss processes
524	are dominated by condensation loss, with minor contributions from ion-molecular reactions and ion-ion recombination.
525	Inspired by the high dependence of charge fraction on the molecule species, we can take the ambient atmosphere as a
526	natural ion-molecule reaction chamber, and the charge fraction observed for different neutral molecules may provide
527	some insights into the charging efficiency of organic species measured with the reagent ions of NO_3^- and HSO_4^- in the
528	chemical ionization mass spectrometers and help better quantify them.
529	
530	Data availability
530 531	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average
530531532	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from
530531532533	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785.
 530 531 532 533 534 	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785.
 530 531 532 533 534 535 	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785. Author contributions
 530 531 532 533 534 535 536 	 Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785. Author contributions RY and JK designed the study. RY, XL, CY, RC, YZ, JK, NS, and JL participated in data collection and performed the data
 530 531 532 533 534 535 536 537 	 Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785. Author contributions RY and JK designed the study. RY, XL, CY, RC, YZ, JK, NS, and JL participated in data collection and performed the data analysis. RY and XL prepared the first version of the manuscript with contributions from all co-authors. All authors approved
 530 531 532 533 534 535 536 537 538 	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785. Author contributions RY and JK designed the study. RY, XL, CY, RC, YZ, JK, NS, and JL participated in data collection and performed the data analysis. RY and XL prepared the first version of the manuscript with contributions from all co-authors. All authors approved the final version of the manuscript.
 530 531 532 533 534 535 536 537 538 539 	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785. Author contributions RY and JK designed the study. RY, XL, CY, RC, YZ, JK, NS, and JL participated in data collection and performed the data analysis. RY and XL prepared the first version of the manuscript with contributions from all co-authors. All authors approved the final version of the manuscript.
 530 531 532 533 534 535 536 537 538 539 540 	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785. Author contributions RY and JK designed the study. RY, XL, CY, RC, YZ, JK, NS, and JL participated in data collection and performed the data analysis. RY and XL prepared the first version of the manuscript with contributions from all co-authors. All authors approved the final version of the manuscript. Competing interests
 530 531 532 533 534 535 536 537 538 539 540 541 	Data availability The detection efficiency of APi-TOF, the time series of total negative cluster ions and CS in urban Beijing, and the average spectrums of negative clusters ions measured by APi-TOF during haze and clean periods in urban Beijing are available from https://zenodo.org/record/7791785. Author contributions RY and JK designed the study. RY, XL, CY, RC, YZ, JK, NS, and JL participated in data collection and performed the data analysis. RY and XL prepared the first version of the manuscript with contributions from all co-authors. All authors approved the final version of the manuscript. Competing interests The authors declare that they have no conflict of interest.

543 Acknowledgments

- 544 This research has been supported by National Natural Science Foundation of China (grant no. 22188102, 92044301, and
- 545 22106083), Samsung PM_{2.5} SRP, the Academy of Finland (grant no. 337549, 302958, 337549, 1325656, 316114, 325647,
- and 332547), and European Research Council (grant no. 742206). The authors gratefully acknowledge the support of the
- 547 research teams in AHL/BUCT laboratory and SMEAR II station.
- 548
- 549

550 **References**

- 551 Arnold, F. and Viggiano, A. A.: Combined mass spectrometric composition measurements of positive and negative ions in the
- 552 lower ionosphere—I. Positive ions, Planet. Space Sci., 30, 1295-1305, 10.1016/0032-0633(82)90103-9, 1982.
- Arnold, F., Böhringer, H., and Henschen, G.: Composition measurements of stratospheric positive ions, Geophys. Res. Lett.,
 5, 653-656, 10.1029/GL005i008p00653, 1978.
- Bates, D. R.: Recombination of Small Ions in the Troposphere and Lower Stratosphere, Planet. Space Sci., 30, 1275-1282,
 10.1016/0032-0633(82)90101-5, 1982.
- 557 Beck, L. J., Schobesberger, S., Sipilä, M., Kerminen, V. M., and Kulmala, M.: Estimation of sulfuric acid concentration using 558 ambient ion composition and concentration data obtained with atmospheric pressure interface time-of-flight ion mass 559 spectrometer, Atmos. Meas. Tech., 15, 1957-1965, 10.5194/amt-15-1957-2022, 2022.
- Beig, G. and Brasseur, G. P.: Model of tropospheric ion composition: A first attempt, J. Geophys. Res.: Atmos., 105, 2267122684, Doi 10.1029/2000jd900119, 2000.
- Bertram, T. H., Kimmel, J. R., Crisp, T. A., Ryder, O. S., Yatavelli, R. L. N., Thornton, J. A., Cubison, M. J., Gonin, M., and
 Worsnop, D. R.: A field-deployable, chemical ionization time-of-flight mass spectrometer, Atmos. Meas. Tech., 4, 1471-1479,
 10.5194/amt-4-1471-2011, 2011.
- Bianchi, F., Garmash, O., He, X., Yan, C., Iyer, S., Rosendahl, I., Xu, Z., Rissanen, M. P., Riva, M., Taipale, R., Sarnela, N.,
 Petäjä, T., Worsnop, D. R., Kulmala, M., Ehn, M., and Junninen, H.: The role of highly oxygenated molecules (HOMs) in
 determining the composition of ambient ions in the boreal forest, Atmos. Chem. Phys., 17, 13819-13831, 10.5194/acp-1713819-2017, 2017.
- 569 Bianchi, F., Trostl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov, A., Bukowiecki,
- 570 N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kurten, A., Manninen, H. E., Munch, S.,
- Perakyla, O., Petaja, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., Kulmala, M., Dommen, J.,
 and Baltensperger, U.: New particle formation in the free troposphere: A question of chemistry and timing, Science, 352, 1109-
- 573 1112, 10.1126/science.aad5456, 2016.
- Cai, R. L., Yang, D. S., Fu, Y. Y., Wang, X., Li, X. X., Ma, Y., Hao, J. M., Zheng, J., and Jiang, J. K.: Aerosol surface area
 concentration: a governing factor in new particle formation in Beijing, Atmos. Chem. Phys., 17, 12327-12340, 10.5194/acp17-12327-2017, 2017.

- 577 Carslaw, K. S., Harrison, R. G., and Kirkby, J.: Cosmic Rays, Clouds, and Climate, Science, 298, 1732-1737,
 578 doi:10.1126/science.1076964, 2002.
- Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley, J. A., Jr., Hansen, J. E., and Hofmann, D. J.: Climate
 forcing by anthropogenic aerosols, Science, 255, 423-430, 10.1126/science.255.5043.423, 1992.
- 581 Chen, X., Kerminen, V.-M., Paatero, J., Paasonen, P., Manninen, H. E., Nieminen, T., Petäjä, T., and Kulmala, M.: How do air
- ions reflect variations in ionising radiation in the lower atmosphere in a boreal forest?, Atmos. Chem. Phys., 16, 14297-14315,
 10.5194/acp-16-14297-2016, 2016.
- 584 Chen, X., Virkkula, A., Kerminen, V.-M., Manninen, H. E., Busetto, M., Lanconelli, C., Lupi, A., Vitale, V., Del Guasta, M.,
- 585 Grigioni, P., Vaananen, R., Duplissy, E.-M., Petaja, T., and Kulmala, M.: Features in air ions measured by an air ion
- 586 spectrometer (AIS) at Dome C, Atmos. Chem. Phys., 17, 13783-13800, 10.5194/acp-17-13783-2017, 2017.
- Cheng, X., Chen, Q., Jie Li, Y., Zheng, Y., Liao, K., and Huang, G.: Highly oxygenated organic molecules produced by the
 oxidation of benzene and toluene in a wide range of OH exposure and NOx conditions, Atmos. Chem. Phys., 21, 12005–12019,
 10.5194/acp-21-12005-2021, 2021.
- 590 Chu, C.-H., Chen, S.-R., Wu, C.-H., Cheng, Y.-C., Cho, Y.-M., and Chang, Y.-K.: The effects of negative air ions on cognitive 591 function: an event-related potential (ERP) study, International Journal of Biometeorology, 63, 1309-1317, 10.1007/s00484-592 019-01745-7, 2019.
- Curtius, J., Lovejoy, E. R., and Froyd, K. D.: Atmospheric ion-induced aerosol nucleation, Space Sci. Rev., 125, 159-167,
 10.1007/s11214-006-9054-5, 2006.
- Davidson, J. A., Fehsenfeld, F. C., and Howard, C. J.: The heats of formation of NO₃⁻ and NO₃⁻ association complexes with
 HNO₃ and HBr, Int. J. Chem. Kinet., 9, 17-29, 10.1002/kin.550090104, 1977.
- Dos Santos, V. N., Herrmann, E., Manninen, H. E., Hussein, T., Hakala, J., Nieminen, T., Aalto, P. P., Merkel, M., Wiedensohler,
 A., Kulmala, M., Petaja, T., and Hameri, K.: Variability of air ion concentrations in urban Paris, Atmos. Chem. Phys., 15,
 13717-13737, 10.5194/acp-15-13717-2015, 2015.
- 600 Ehn, M., Junninen, H., Petaja, T., Kurten, T., Kerminen, V. M., Schobesberger, S., Manninen, H. E., Ortega, I. K., Vehkamaki,
- H., Kulmala, M., and Worsnop, D. R.: Composition and temporal behavior of ambient ions in the boreal forest, Atmos. Chem.
- 602 Phys., 10, 8513-8530, 10.5194/acp-10-8513-2010, 2010.
- Ehn, M., Junninen, H., Schobesberger, S., Manninen, H. E., Franchin, A., Sipila, M., Petaja, T., Kerminen, V. M., Tammet, H.,
- Mirme, A., Mirme, S., Hõrrak, U., Kulmala, M., and Worsnop, D. R.: An Instrumental Comparison of Mobility and Mass
- 605 Measurements of Atmospheric Small Ions, Aerosol Sci. Technol., 45, 522-532, 10.1080/02786826.2010.547890, 2011.
- Eisele, F. L.: Natural and Anthropogenic Negative-Ions in the Troposphere, J. Geophys. Res.: Atmos., 94, 2183-2196,
 10.1029/JD094iD02p02183, 1989.
- Eisele, F. L. and Tanner, D. J.: Identification of Ions in Continental Air, J. Geophys. Res.: Atmos., 95, 20539-20550,
 10.1029/JD095iD12p20539, 1990.
- 610 Eisele, F. L. and Tanner, D. J.: Measurement of the gas phase concentration of H2SO4 and methane sulfonic acid and estimates
- of H2SO4 production and loss in the atmosphere, J. Geophys. Res.: Atmos., 98, 9001-9010, 10.1029/93JD00031, 1993.

- 612 Eisele, F. L., Lovejoy, E. R., Kosciuch, E., Moore, K. F., Mauldin III, R. L., Smith, J. N., McMurry, P. H., and Iida, K.: Negative
- atmospheric ions and their potential role in ion-induced nucleation, 111, 10.1029/2005JD006568, 2006.
- 614 Friedlander, S. K.: Smoke, dust, and haze, Oxford university press New York2000.
- 615 G. Baumgaertner, A. J., Thayer, J. P., Neely III, R. R., and Lucas, G.: Toward a comprehensive global electric circuit model:
- 616 Atmospheric conductivity and its variability in CESM1(WACCM) model simulations, J. Geophys. Res.: Atmos., 118, 9221-
- 617 9232, 10.1002/jgrd.50725, 2013.
- Gautam, A. S., Siingh, D., and Kamra, A. K.: Statistical analysis of the atmospheric ion concentrations and mobility
 distributions at a tropical station, Pune, Q. J. R. Meteorol. Soc., 143, 2116-2128, 10.1002/qj.3071, 2017.
- Gerdien, H.: Die absolute Messung der specifischen Leitfähigkeit und der Dichte des verticalen Leitungsstromes in der
 Atmosphäre, Terr. Magn. Atmos. Electr., 10, 65-74, 10.1029/TE010i002p00065, 1905.
- Harrison, R. G.: Ion-aerosol-cloud processes in the lower atmosphere, Reviews of Geophysics, 41, 1012,
 10.1029/2002rg000114, 2003.
- Heinritzi, M., Simon, M., Steiner, G., Wagner, A. C., Kürten, A., Hansel, A., and Curtius, J.: Characterization of the massdependent transmission efficiency of a CIMS, Atmos. Meas. Tech., 9, 1449-1460, 10.5194/amt-9-1449-2016, 2016.
- Hirsikko, A., Paatero, J., Hatakka, J., and Kulmala, M.: The 222Rn activity concentration, external radiation dose and air ion
 production rates in a boreal forest in Finland between March 2000 and June 2006, 12, 265-278, 2007.
- Hirsikko, A., Nieminen, T., Gagné, S., Lehtipalo, K., Manninen, H. E., Ehn, M., Hõrrak, U., Kerminen, V. M., Laakso, L.,
 McMurry, P. H., Mirme, A., Mirme, S., Petäjä, T., Tammet, H., Vakkari, V., Vana, M., and Kulmala, M.: Atmospheric ions and
- 630 nucleation: a review of observations, Atmos. Chem. Phys., 11, 767-798, 10.5194/acp-11-767-2011, 2011.
- Hõrrak, U., Salm, J., and Tammet, H.: Statistical characterization of air ion mobility spectra at Tahkuse Observatory:
 Classification of air ions, J. Geophys. Res.: Atmos., 105, 9291-9302, 10.1029/1999JD901197, 2000.
- Iida, K., Stolzenburg, M., McMurry, P., Dunn, M. J., Smith, J. N., Eisele, F., and Keady, P.: Contribution of ion-induced
 nucleation to new particle formation: Methodology and its application to atmospheric observations in Boulder, Colorado, J.
 Geophys. Res.: Atmos., 111, 10.1029/2006JD007167, 2006.
- Isidorov, V. A., Zenkevich, I. G., and Ioffe, B. V.: Volatile organic compounds in the atmosphere of forests, Atmos. Environ.,
 19, 1-8, 10.1016/0004-6981(85)90131-3, 1985.
- 638 Jen, C. N., Hanson, D. R., and McMurry, P. H.: Toward Reconciling Measurements of Atmospherically Relevant Clusters by
- Chemical Ionization Mass Spectrometry and Mobility Classification/Vapor Condensation, Aerosol Sci. Technol., 49, i-iii,
 10.1080/02786826.2014.1002602, 2015.
- Jiang, S.-Y., Ma, A., and Ramachandran, S.: Negative air ions and their effects on human health and air quality improvement,
 Int. J. Mol. Sci., 19, 2966, 10.3390/ijms19102966, 2018.
- Jokinen, T., Sipilä, M., Junninen, H., Ehn, M., Lönn, G., Hakala, J., Petäjä, T., Mauldin, R. L., Kulmala, M., and Worsnop, D.
 R.: Atmospheric sulphuric acid and neutral cluster measurements using CI-APi-TOF, Atmos. Chem. Phys., 12, 4117-4125, 10.5194/acp-12-4117-2012, 2012.

- Junninen, H., Ehn, M., Petäjä, T., Luosujärvi, L., Kotiaho, T., Kostiainen, R., Rohner, U., Gonin, M., Fuhrer, K., Kulmala, M.,
 and Worsnop, D. R.: A high-resolution mass spectrometer to measure atmospheric ion composition, Atmos. Meas. Tech., 3,
 1039-1053, 10.5194/amt-3-1039-2010, 2010.
- Kawamoto, H. and Ogawa, T.: A steady state model of negative ions in the lower stratosphere, Planet. Space Sci., 32, 12231233, 10.1016/0032-0633(84)90066-7, 1984.
- Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J.,
 Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner,
- M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle,
 C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov,
- V., Mathot, S., Molteni, U., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A.
 D., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G.,
- 657 Stozhkov, Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D.,
- 658 Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M.,
- Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, Nature, 533, 521-526, 10.1038/nature17953,
 2016.
- Kontkanen, J., Lehtinen, K. E. J., Nieminen, T., Manninen, H. E., Lehtipalo, K., Kerminen, V. M., and Kulmala, M.: Estimating
 the contribution of ion–ion recombination to sub-2 nm cluster concentrations from atmospheric measurements, Atmos. Chem.
 Phys., 13, 11391-11401, 10.5194/acp-13-11391-2013, 2013.
- Krechmer, J. E., Groessl, M., Zhang, X., Junninen, H., Massoli, P., Lambe, A. T., Kimmel, J. R., Cubison, M. J., Graf, S., Lin,
 Y.-H., Budisulistiorini, S. H., Zhang, H., Surratt, J. D., Knochenmuss, R., Jayne, J. T., Worsnop, D. R., Jimenez, J.-L., and
- 666 Canagaratna, M. R.: Ion mobility spectrometry–mass spectrometry (IMS–MS) for on- and offline analysis of atmospheric gas
- 667 and aerosol species, Atmos. Meas. Tech., 9, 3245-3262, 10.5194/amt-9-3245-2016, 2016.
- Ku, B. K. and de la Mora, J. F.: Relation between Electrical Mobility, Mass, and Size for Nanodrops 1–6.5 nm in Diameter in
 Air, Aerosol Sci. Technol., 43, 241-249, 10.1080/02786820802590510, 2009.
- Kürten, A., Rondo, L., Ehrhart, S., and Curtius, J.: Calibration of a chemical ionization mass spectrometer for the measurement
 of gaseous sulfuric acid, J. Phys. Chem. A, 116, 6375-6386, 10.1021/jp212123n, 2012.
- Lee, S.-H., Reeves, J. M., Wilson, J. C., Hunton, D. E., Viggiano, A. A., Miller, T. M., Ballenthin, J. O., and Lait, L. R.: Particle
 formation by ion nucleation in the upper troposphere and lower stratosphere, Science, 301, 1886-1889,
 doi:10.1126/science.1087236, 2003.
- Li, X., Chee, S., Hao, J., Abbatt, J. P. D., Jiang, J., and Smith, J. N.: Relative humidity effect on the formation of highly oxidized
 molecules and new particles during monoterpene oxidation, Atmos. Chem. Phys., 19, 1555-1570, 10.5194/acp-19-1555-2019,
 2019.
- Li, X., Li, Y., Lawler, M. J., Hao, J., Smith, J. N., and Jiang, J.: Composition of Ultrafine Particles in Urban Beijing:
 Measurement Using a Thermal Desorption Chemical Ionization Mass Spectrometer, Environ. Sci. Technol., 55, 2859-2868,
 10.1021/acs.est.0c06053, 2021.
- Li, X., Li, Y., Cai, R., Yan, C., Qiao, X., Guo, Y., Deng, C., Yin, R., Chen, Y., Li, Y., Yao, L., Sarnela, N., Zhang, Y., Petäjä, T.,
 Bianchi, F., Liu, Y., Kulmala, M., Hao, J., Smith, J. N., and Jiang, J.: Insufficient Condensable Organic Vapors Lead to Slow

- 683 Growth of New Particles in an Urban Environment, Environ. Sci. Technol., 55, 9936-9946, 10.1021/acs.est.2c01566, 2022.
- Liang, X., Zhou, Q., Wang, W., Wang, X., Chen, W., Chen, C., Li, Y., Hou, K., Li, J., and Li, H.: Sensitive detection of black
 powder by a stand-alone ion mobility spectrometer with an embedded titration region, Anal. Chem., 85, 4849-4852,
 10.1021/ac400337s, 2013.
- Ling, X., Jayaratne, R., and Morawska, L.: Air ion concentrations in various urban outdoor environments, Atmos. Environ.,
 44, 2186-2193, 10.1016/j.atmosenv.2010.03.026, 2010.
- 689 Liu, Y., Yan, C., Feng, Z., Zheng, F., Fan, X., Zhang, Y., Li, C., Zhou, Y., Lin, Z., Guo, Y., Zhang, Y., Ma, L., Zhou, W., Liu,
- 690 Z., Dada, L., Daellenbach, K., Kontkanen, J., Cai, R., Chan, T., and Kulmala, M.: Continuous and comprehensive atmospheric
- observations in Beijing: a station to understand the complex urban atmospheric environment, Big Earth Data, 4, 295-321,
 10.1080/20964471.2020.1798707, 2020.
- Lovejoy, E. R.: Atmospheric ion-induced nucleation of sulfuric acid and water, J. Geophys. Res., 109, D08204,
 10.1029/2003jd004460, 2004.
- 695 Lovejoy, E. R. and Curtius, J.: Cluster ion thermal decomposition (II): Master equation modeling in the low-pressure limit and 696 fall-off regions. Bond energies for $HSO_4^{-}(H_2SO_4)(_x)(HNO_3)(_y)$, J. Phys. Chem. A, 105, 10874-10883, 10.1021/jp012496s, 2001.
- Luts, A.: Evolution of negative small ions at enhanced ionization, J. Geophys. Res.: Atmos., 100, 1487-1496,
 10.1029/94JD01836, 1995.
- Luts, A. and Salm, J.: Chemical-composition of small atmospheric ions near the ground, J. Geophys. Res.: Atmos., 99, 10781 10785, 10.1029/93jd03225, 1994.
- Mahfouz, N. G. A. and Donahue, N. M.: Technical note: The enhancement limit of coagulation scavenging of small charged
 particles, Atmos. Chem. Phys., 21, 3827-3832, 10.5194/acp-21-3827-2021, 2021.
- Malcolm, C. P., Cowen, P. J., and Harmer, C. J.: Research Letter: High-density negative ion treatment increases positive
 affective memory, Psychological Medicine, 39, 1930-1932, 10.1017/S0033291709990444, 2009.
- Manninen, H. E., Mirme, S., Mirme, A., Petaja, T., and Kulmala, M.: How to reliably detect molecular clusters and nucleation
 mode particles with Neutral cluster and Air Ion Spectrometer (NAIS), Atmos. Meas. Tech., 9, 3577-3605, 10.5194/amt-9 3577-2016, 2016.
- Manninen, H. E., Petäjä, T., Asmi, E., Riipinen, I., Nieminen, T., Mikkilä, J., Hõrrak, U., Mirme, A., Mirme, S., and Laakso,
 L.: Long-term field measurements of charged and neutral clusters using Neutral cluster and Air Ion Spectrometer (NAIS),
 Boreal Environ. Res., 2009.
- Mirme, A., Tamm, E., Mordas, G., Vana, M., Uin, J., Mirme, S., Bernotas, T., Laakso, L., Hirsikko, A., and Kulmala, M.: A
 wide-range multi-channel air ion spectrometer, Boreal. Environ. Res., 12, 247-264, 2007.
- 713 Mirme, S. and Mirme, A.: The mathematical principles and design of the NAIS a spectrometer for the measurement of cluster
- ion and nanometer aerosol size distributions, Atmos. Meas. Tech., 6, 1061-1071, 10.5194/amt-6-1061-2013, 2013.
- 715 Möhler, O., Reiner, T., and Arnold, F.: The formation of SO5- by gas phase ion-molecule reactions, The Journal of Chemical
- 716 Physics, 97, 8233-8239, 10.1063/1.463394, 1992.

- 717 Nadykto, A. B., Herb, J., Yu, F., and Nazarenko, K. M.: Clustering of highly oxidized organic acid with atmospheric NO₃⁻ and
- 718 HSO₄⁻ ions and neutral species: Thermochemistry and implications to new particle formation, Chem. Phys. Lett., 706, 175-
- 719 181, 10.1016/j.cplett.2018.05.080, 2018.
- 720 Nie, W., Yan, C., Huang, D. D., Wang, Z., Liu, Y., Qiao, X., Guo, Y., Tian, L., Zheng, P., Xu, Z., Li, Y., Xu, Z., Qi, X., Sun, P.,
- Wang, J., Zheng, F., Li, X., Yin, R., Dallenbach, K. R., Bianchi, F., Petäjä, T., Zhang, Y., Wang, M., Schervish, M., Wang, S.,
- 722 Qiao, L., Wang, Q., Zhou, M., Wang, H., Yu, C., Yao, D., Guo, H., Ye, P., Lee, S., Li, Y. J., Liu, Y., Chi, X., Kerminen, V.-M.,
- Ehn, M., Donahue, N. M., Wang, T., Huang, C., Kulmala, M., Worsnop, D., Jiang, J., and Ding, A.: Secondary organic aerosol
- formed by condensing anthropogenic vapours over China's megacities, Nature Geoscience, 15, 255-261, 10.1038/s41561-022-
- 725 00922-5, 2022.
- Qiao, X., Yan, C., Li, X., Guo, Y., Yin, R., Deng, C., Li, C., Nie, W., Wang, M., Cai, R., Huang, D., Wang, Z., Yao, L., Worsnop,
 D. R., Bianchi, F., Liu, Y., Donahue, N. M., Kulmala, M., and Jiang, J.: Contribution of Atmospheric Oxygenated Organic
 Compounds to Particle Growth in an Urban Environment, Environ. Sci. Technol., 55, 13646-13656, 10.1021/acs.est.1c02095,
 2021.
- Riva, M., Rantala, P., Krechmer, J. E., Peräkylä, O., Zhang, Y., Heikkinen, L., Garmash, O., Yan, C., Kulmala, M., Worsnop,
 D., and Ehn, M.: Evaluating the performance of five different chemical ionization techniques for detecting gaseous oxygenated
 organic species, Atmos. Meas. Tech., 12, 2403-2421, 10.5194/amt-12-2403-2019, 2019.
- Shuman, N. S., Hunton, D. E., and Viggiano, A. A.: Ambient and Modified Atmospheric Ion Chemistry: From Top to Bottom,
 Chem. Rev., 115, 4542-4570, 10.1021/cr5003479, 2015.
- Spangler, G. E. and Collins, C. I.: Reactant ions in negative ion plasma chromatography, Anal. Chem., 47, 393-402,
 10.1021/ac60353a019, 1975.
- Stano, M., Safonov, E., Kucera, M., and Matejcik, S.: Ion Mobility Spectrometry Study of Negative Corona Discharge in
 Oxygen/Nitrogen Mixtures, Chem. Listy, 102, S1414-S1417, 2008.
- Tammet, H.: Size and mobility of nanometer particles, clusters and ions, Journal of Aerosol Science, 26, 459-475,
 10.1016/0021-8502(94)00121-E, 1995.
- Tammet, H.: Continuous scanning of the mobility and size distribution of charged clusters and nanometer particles in
 atmospheric air and the Balanced Scanning Mobility Analyzer BSMA, Atmos. Res., 82, 523-535,
 10.1016/j.atmosres.2006.02.009, 2006.
- Tammet, H.: Nanoion 2010-11, University of Tartu [dataset], <u>http://dx.doi.org/10.15155/repo-2</u>, 2015a.
- Tammet, H., Hõrrak, U., Laakso, L., and Kulmala, M.: Factors of air ion balance in a coniferous forest according to
 measurements in Hyytiälä, Finland, Atmos. Chem. Phys., 6, 3377-3390, 10.5194/acp-6-3377-2006, 2006.
- Tammet, H. H., Urmas: FinEstIon2003_06. Dataset of air ion and atmospheric nanoparticle measurements, University of Tartu
 [dataset], <u>http://dx.doi.org/10.15155/repo-4</u>, 2015b.
- Tanner, D. J. and Eisele, F. L.: Ions in oceanic and continental air masses, J. Geophys. Res.: Atmos., 96, 1023-1031,
 10.1029/90JD02131, 1991.
- 751 Usoskin, I. G., Gladysheva, O. G., and Kovaltsov, G. A.: Cosmic ray-induced ionization in the atmosphere: spatial and temporal

- changes, Journal of Atmospheric and Solar-Terrestrial Physics, 66, 1791-1796, 10.1016/j.jastp.2004.07.037, 2004.
- Viggiano, A. A.: IN-SITU MASS-SPECTROMETRY AND ION CHEMISTRY IN THE STRATOSPHERE AND
 TROPOSPHERE, Mass Spectrometry Reviews, 12, 115-137, 10.1002/mas.1280120203, 1993.

Viggiano, A. A., Perry, R. A., Albritton, D. L., Ferguson, E. E., and Fehsenfeld, F. C.: The role of H2SO4 in stratospheric negative-ion chemistry, 85, 4551-4555, 10.1029/JC085iC08p04551, 1980.

Viggiano, A. A., Seeley, J. V., Mundis, P. L., Williamson, J. S., and Morris, R. A.: Rate constants for the reactions of XO_3^- (H₂O)(*n*) (X = C, HC, and N) and NO₃⁻(HNO₃)(*n*) with H₂SO₄: Implications for atmospheric detection of H₂SO₄, J. Phys. Chem. A, 101, 8275-8278, 10.1021/jp971768h, 1997.

Wagner, R., Manninen, H. E., Franchin, A., Lehtipalo, K., Mirme, S., Steiner, G., Petäjä, T., and Kulmala, M.: On the accuracy
of ion measurements using a Neutral cluster and Air Ion Spectrometer, Boreal Environ. Res., 21, 230-241, 2016.

762 Yan, C., Dada, L., Rose, C., Jokinen, T., Nie, W., Schobesberger, S., Junninen, H., Lehtipalo, K., Sarnela, N., Makkonen, U.,

Garmash, O., Wang, Y., Zha, Q., Paasonen, P., Bianchi, F., Sipilä, M., Ehn, M., Petäjä, T., Kerminen, V.-M., Worsnop, D. R.,

and Kulmala, M.: The role of H₂SO₄-NH₃ anion clusters in ion-induced aerosol nucleation mechanisms in the boreal forest,
 Atmos. Chem. Phys., 18, 13231-13243, 10.5194/acp-18-13231-2018, 2018.

766 Yan, C., Nie, W., Vogel, A. L., Dada, L., Lehtipalo, K., Stolzenburg, D., Wagner, R., Rissanen, M. P., Xiao, M., Ahonen, L.,

767 Fischer, L., Rose, C., Bianchi, F., Gordon, H., Simon, M., Heinritzi, M., Garmash, O., Roldin, P., Dias, A., Ye, P., Hofbauer,

768 V., Amorim, A., Bauer, P. S., Bergen, A., Bernhammer, A. K., Breitenlechner, M., Brilke, S., Buchholz, A., Mazon, S. B.,

- Canagaratna, M. R., Chen, X., Ding, A., Dommen, J., Draper, D. C., Duplissy, J., Frege, C., Heyn, C., Guida, R., Hakala, J.,
- Heikkinen, L., Hoyle, C. R., Jokinen, T., Kangasluoma, J., Kirkby, J., Kontkanen, J., Kurten, A., Lawler, M. J., Mai, H., Mathot,
- S., Mauldin, R. L., 3rd, Molteni, U., Nichman, L., Nieminen, T., Nowak, J., Ojdanic, A., Onnela, A., Pajunoja, A., Petaja, T.,
- Piel, F., Quelever, L. L. J., Sarnela, N., Schallhart, S., Sengupta, K., Sipila, M., Tome, A., Trostl, J., Vaisanen, O., Wagner, A.
- C., Ylisirnio, A., Zha, Q., Baltensperger, U., Carslaw, K. S., Curtius, J., Flagan, R. C., Hansel, A., Riipinen, I., Smith, J. N.,
- 774 Virtanen, A., Winkler, P. M., Donahue, N. M., Kerminen, V. M., Kulmala, M., Ehn, M., and Worsnop, D. R.: Size-dependent
- influence of NO_x on the growth rates of organic aerosol particles, Sci. Adv., 6, eaay4945, 10.1126/sciadv.aay4945, 2020.
- Yin, R., Yan, C., Cai, R., Li, X., Shen, J., Lu, Y., Schobesberger, S., Fu, Y., Deng, C., Wang, L., Liu, Y., Zheng, J., Xie, H.,
 Bianchi, F., Worsnop, D. R., Kulmala, M., and Jiang, J.: Acid–Base Clusters during Atmospheric New Particle Formation in
 Urban Beijing, Environ. Sci. Technol., 55, 10994-11005, 10.1021/acs.est.1c02701, 2021.
- 779 Zauner-Wieczorek, M., Curtius, J., and Kürten, A.: The ion-ion recombination coefficient α: comparison of temperature- and
- 780 pressure-dependent parameterisations for the troposphere and stratosphere, Atmos. Chem. Phys., 22, 12443-12465,
- 781 10.5194/acp-22-12443-2022, 2022.
- 782



Figure 1. The detection efficiency and absolute transmission efficiency of APi-HTOF determined through the *in-situ* comparison with NAIS and the relative transmission efficiency of APi-HTOF determined through the depletion method. The relative transmission efficiency is obtained by dividing the detection efficiency by the sampling efficiency of cluster ions. The voltage settings of APi-HTOF remain the same during the experiment.



Figure 2. The total concentration of negative cluster ions ($n_{sub-1.6 \text{ nm}}$) measured by APi-HTOF is well correlated with that measured by NAIS. Colors represent the ratio of negative ions smaller than 1.3 nm ($n_{sub-1.3 \text{ nm}}$) to negative cluster ions ($n_{sub-1.6 \text{ nm}}$).



Figure 3. (a) Negative cluster ions observed in global urban, rural, forest, and marine sites. The circles represent the
median values, and the error bars represent the 25%-75% ranges. Ion mobility ranges for different sites were between sub1.6 and sub-2 nm. (b) The decreasing trends of the negative cluster ion concentrations with the increase of CS were
observed in both Beijing and Hyytiälä. The markers and error bars are median values and 25%-75% ranges, respectively.
For comparison, NAIS data were used for both urban Beijing and the clean forest site Hyytiälä.



Figure 4. The negative cluster ion compositions in Beijing and Hyytiälä. The bar plots were average mass spectra of negative cluster ions during clean and haze periods in Beijing and Hyytiälä measured by APi-HTOF. Signals for m/z>300were multiplied by 10 times in Beijing for clearer views. Haze periods were identified based on whether PM_{2.5} concentration is higher than 75 µg·m⁻³. The average NO_x concentrations for the three situations are 30.1, 15.0, and 1.5 ppb, respectively. The pie charts were species distribution of organic ions, including the deprotonated form, adducts to NO₃⁻ and HSO₄⁻ forms of CHO and CHON species.



Figure 5. (a) The carbon atom number and average carbon oxidation state (\overline{OS}_C) of molecules that adduct to NO₃⁻ and HSO₄⁻ respectively. (b) The volatility distribution of CHO and CHON_{nonNPs} species in the forms of adducts with both NO₃⁻ and HSO₄⁻. Only species that attached to both NO₃⁻ and HSO₄⁻ were considered for calculating the volatility

815 distribution. The volatilities were calculated using the method of Qiao et al. (2021).



Figure 6. The ratio of CHO·HSO4⁻/CHO·NO3⁻ (a), CHON_{nonNPs}·HSO4⁻/CHON_{nonNPs}·NO3⁻ (b), and CHON_{NPs}·HSO4⁻ $/CHON_{NPs} \cdot NO_3^-$ (c) as a function of the ratio of HSO_4^-/NO_3^- . The color represents the concentration of sulfuric acid.





Figure 7. Diurnal cycles of negative cluster ions, corresponding neutral molecules, and the charge fractions between them in urban Beijing. (a) The diurnal cycles of reagent ions and their inorganic clusters, together with those of organic ions. The organic ions in the forms of adducts with NO₃⁻ and HSO₄⁻ were summed into CHO-, CHON_{nonNPs}⁻, and CHON_{NPs}-related ions. (b) The diurnal cycles of sulfuric acid and the corresponding neutral molecules of organic ions measured by the (nitrate-) CI-APi-LTOF. The concentrations of sulfuric acid, CHO, and CHONnonNPs were multiplied by a factor of 100, 20, and 20, respectively, to match that of CHON_{NPs} in the figure. (c) The diurnal cycles of charge fractions between negative cluster ions and their corresponding neutral molecules. The charge fractions for CHO, CHON_{nonNPs}, and CHON_{NPs} in the form of adducts with NO₃⁻ and HSO₄⁻ were calculated and compared with the HSO₄⁻/H₂SO₄ ratio.



Neutral molecules measured by CI-APi-TOF (cm⁻³)

Figure 8. Scatter plots of the concentrations of organic ions measured by APi-TOF and their corresponding neutral molecules measured by CI-APi-TOF in urban Beijing, colored by CS. The cluster ions were CHO·NO₃⁻ (a), CHO·HSO₄⁻ (b), CHON_{nonNPs}·NO₃⁻ (c), CHON_{nonNPs}·HSO₄⁻ (d), CHON_{NPs}·NO₃⁻ (e), and CHON_{NPs}·HSO₄⁻ (f). The neutral molecules were CHO (a and b), CHON_{nonNPs} (c and d), and CHON_{NPs} (e and f), respectively. The data was divided into 5 groups based on CS in the range of 0.001-0.01, 0.01-0.02, 0.02-0.06, 0.06-0.1, and 0.1-0.2 s⁻¹, respectively.



Figure 9. The carbon atom number and average carbon oxidation state (\overline{OS}_C) of molecules in CHO·NO₃⁻ (a) and CHON_{nonNPs}·NO₃⁻ (b). The color represents their charge fraction during the measurement periods.





Figure 10. (a) The measured and simulated concentrations of HSO_4^- and $C_3H_3O_4^-$ in Beijing. The simulation is performed according to dynamic models as described in Section 2.3 and four formation and loss pathways were considered. (b) The fractions of the formation or loss rates of HSO_4^- and $C_3H_3O_4^-$ contributed by different pathways. The average values of the results were shown here.