

Documentary evidence of urban droughts and their impact in the eastern Netherlands: the cases of Deventer and Zutphen, 1500–1795

Dániel Johannes Moerman¹

¹Faculty of Humanities, Department of Art and Culture, History, Antiquity, Vrije Universiteit Amsterdam, De Boelelaan 1105, 1081 HV, Amsterdam

Correspondence to: Dániel Moerman d.j.moerman@vu.nl

Gewijzigde veldcode

Abstract: Compared to other parts of Europe, very little is known about pre-instrumental drought periods in the Netherlands. Existing reconstructions are based primarily on data from England, France, and Germany, while more local studies on drought and its impact are still absent. This article thus aims to expand our knowledge of droughts in the Netherlands between 1500 and 1795, by focusing specifically on drought in an urban context to provide a more precise and local idea of the impact and severity of drought. The main case studies are cities in the eastern part of the country, Deventer and Zutphen. Both cities lay in relative close proximity to each other and share similar geological and hydrological conditions, as well as extensive archives that can be used to gather documentary data regarding historical drought periods. The three primary aims of the article are: 1) to examine the potential use of documentary data from the city archives of Deventer and Zutphen for historical drought reconstruction; 2) to establish droughts for both cities on the basis of the year, month/season in which they took place, as well as ranking the droughts according to the impact-based Historical Severity Drought Scale (HSDS) and 3) to compare the data from this analysis with that of other indices. In the end, the article strengthens the need to focus on documentary data from local case studies regarding drought, not only to provide more precise local reconstructions of drought-severity compared to regional studies, but also to take into account the long-term effects on urban waterscapes and the provisioning of fresh water.

1. Introduction

In recent years, droughts have become a more pressing topic of research. Worldwide, droughts of varying severity affect societies, whether on an agricultural, hydrological, or on wider socio-economic level, which is expected to increase within the current trends of climatic change (Kchouk et. al., 2021; Savelli et. al., 2022; Spinoni et. al., 2018). The study of past droughts for the pre-instrumental period on the basis of documentary evidence and natural proxies, such as dendroclimatology, has displayed the possibility to reconstruct drought-events and their societal impact in Europe, which has led to the development of several historical drought reconstructions and indices. (Bauch et. al., 2020; Brázdil et. al., 2016/2018/2019/2020; Camenisch et. al., 2020; Garnier, 2019; Kiss, 2017/2020; Leijonhufvud and Retsö, 2021; Piervitali and Colacino, 2001; Pribyl and Cornes, 2020; Stangl and Foelsche, 2022). However, very little to no historical drought data exists for the Netherlands. The limited data available from the voluminous works of Buisman (1995/1996/1998/2000/2006/2015) is based primarily on reconstructions and sources from England, France and Germany, and sporadic sources from across the Netherlands. A recent study by Camenisch and Salvisberg (2020), has emphasised the need to analyse regional

38 and local aspects of droughts by studying geographically limited source samples, such as municipal data from city
39 archives. Compared with other, supra-regional drought indices, this can lead to a more detailed understanding of
40 the extent and severity of certain droughts on a local level, while also providing insights into previously unknown
41 droughts. Even droughts with a larger geographical footprint, such as the infamous 1540 ‘Megadrought’ (Wetter
42 et. al. 2014), can thus demonstrate a greater temporal diversity if more localised data is included in the analysis
43 (Maughan et. al. 2022). As such, the data provided by Buisman cannot suffice to study the local or regional severity
44 and impact of drought for the Netherlands, and, as follows, further research is needed.

45 This article aims to expand our knowledge of pre-instrumental droughts in the Netherlands between 1500 and
46 1795, focusing on two cities in the eastern part of the country – Deventer and Zutphen. Both have rich municipal
47 archives, relatively similar geohydrological, and are located in close proximity to one another. The focus on the
48 eastern Netherlands also has a climatological reason, as a recent study has indicated that the eastern inland parts
49 of the Netherlands could be more prone to future droughts compared to the western coastal regions. While the
50 western parts also receive ample discharge from the rivers Rhine and Meuse, the eastern regions generally depend
51 more on precipitation for drought mitigation, given that their elevation above the level of the two rivers makes it
52 impossible for water to reach these areas without pumping. As such, the possibilities for drought mitigation in the
53 eastern regions are regarded as more limited compared to the west. A comparative analysis has also shown that
54 the differences in precipitation between the east and western parts are accompanied by differences in solar radiation
55 and temperature, which influence potential evapotranspiration. This trend has been visible since the 1950s, and is
56 expected to continue with stronger drying trends in the inland regions due to an increase in temperatures as a result
57 of global warming (Phillip et. al., 2020).

58 The focus on more specific urban contexts also moves away from the focus on agricultural drought, which is
59 dominant in historiography, shifting the emphasis to the wider hydrological and socio-economic impact of drought
60 within a city’s walls. This implies a focus on sources from city archives that describe the specific effects of
61 droughts on urban water provisioning, the accessibility of canals and harbours, and sanitary issues. Common
62 factors to denominate drought severity according to the Palmer Drought Severity Index, or PDSI, such as
63 temperature, precipitation levels and soil-moisture deficits, are not enough to determine the impact of droughts on
64 urban environments. Urbanisation, and other large-scale influences of human actions on the distribution and use
65 of water, have often been ignored in many classical drought indices that focused primarily on precipitation and
66 temperature data (Briffa, Van Der Schrier and Jones, 2009; Savelli et. al., 2022). Many previous studies into past
67 droughts worked in relative isolation, without taking into account the complex interactions between natural and
68 human processes in the hydrological sphere (AghaKouchak et. al., 2021; Van Loon et. al. 2016; Maughan et. al.
69 2022; Mukherjee, Mishra and Trenberth, 2018; Vörösmarty et. al., 2004)). These factors are more present in
70 another index, the Historical Severity Drought Scale (HSDS). This index allows for a reconstruction of droughts
71 based on a systemic inventory of the different hydrological and socio-economic impacts to determine levels of
72 drought severity (Garnier, 2014/2019; Metger and Jacob Rousseau, 2020). Urban documentary data provide more
73 precise local reconstructions of drought-severity, as they describe the variety of responses to droughts, allowing
74 for the creation of indices along the HSDS. As such, urban droughts refer to specific effects of drought on the
75 urban environment, which can be reconstructed with the use of urban archives to provide a the long-term
76 perspective on the effects of droughts on urban water systems. This is primarily relevant given the rising interest

77 in the effects of drought on urban environments for the present as well as the future (Machairas and Van de Ven,
78 2022; Szalinska, Otop and Tokarczyk, 2021).

79 This article has three primary aims: 1) examining the potential use of documentary data from the city archives of
80 Deventer and Zutphen for historical drought reconstruction; 2) to establish droughts for both cities on the basis of
81 the year, month/season in which they took place, as well as ranking the droughts according to the impact-based
82 Historical Drought Severity Scale; and 3) to compare the data from this analysis with that of other indices, such as
83 the [Van Engelen, Buisman, and IJnsen](#) temperature series for the Netherlands, the supra-regional drought index,
84 ~~of~~ (SDI), which comprises data from Switzerland, France, the Netherlands and Germany, (Camenisch and
85 Salvisberg, 2020), and the Old World Drought Atlas (OWDA), which provides an overview of
86 dendrochronological drought data on a regional scale (Cook et.al., 2015).

87

88 The article is divided in six sections. The first ~~section~~ provides a detailed overview of the sources used in the
89 reconstruction of drought for Deventer and Zutphen. Section two will present outcomes from the study of these
90 sources, by which the drought years are presented via a chronological HSDS. Section three discusses a specific set
91 of examples from the sources, providing a more detailed analysis of the data and their respective values. Sections
92 four, five, and six compare the data gathered in this study with other indices, followed by a final discussion and
93 conclusion.

94

95 **2. The data**

96 To reconstruct past weather and climatic phenomena, historical climatologists draw from a large amount of
97 documentary sources that provide either direct or indirect (proxy) data about changes in weather or abnormal
98 patterns of precipitation and temperatures (Brázdil et. al., 2010; Pfister, 2018). As for drought reconstructions, the
99 documentary evidence often consists of annals, chronicles, and diaries, in which people recorded daily or
100 extraordinary weather situations, or more institutional sources, such as tax and harvest records, and religious data
101 with regard to rogation ceremonies (Brazdil et. al. 2013/2019/2020; Dominguez-Castro et. al., 2012; Kiss and
102 Nolic, 2015). Throughout most parts of Europe, municipal records, from cities, towns and villages, became more
103 systematised from the end of the fifteenth century onward, often containing deliberations and resolutions that
104 indicate means by which local or state governments aimed to alleviate the effects of drought or other weather
105 extremes (Garnier, 2019; Gorostiza, Escayol and Barriendos, 2021; Grau Satorras et. al., 2021). Therefore,
106 municipal archives qualify as a reliable *Fundgrube* for (proxy) evidence of urban droughts during the pre-
107 instrumental period.

108 For this study, the municipal archives of two cities in the eastern Netherlands, Deventer and Zutphen, have been
109 studied extensively in search of references to drought-related phenomena. Deventer and Zutphen are both situated
110 along the IJssel river on sandy river dunes from the Holocene and relied on surface water from the rivers and clean
111 groundwater for everyday use (Vogelzang, 1956). The primary sources that have been studied were primarily
112 official municipal records, such as daily resolutions from the city government, ordinance books, and petitions. For
113 Deventer, a long-running series of sources, including daily resolutions, decrees from the magistracy (*buurspraken*)
114 and citizen petitions are available from 1459 until 1795. Both the daily resolutions and books of concordances

115 come with alphabetical reference books from eighteenth and nineteenth-century authors, which provide a useful,
116 yet also limited tool to find certain relevant entries regarding drought. In the case of Zutphen, the extensive series
117 of daily resolutions and can be studied from 1573 until the start of the nineteenth century. These series, including
118 the digitised reference books provided the primary source for Zutphen. In this regard, it must be noted that for
119 certain periods, particularly the seventeenth century, the amount of sources regarding Zutphen was generally less
120 extensive compared to Deventer.

121

122 3. Methodology

123 In this section, I discuss several indices and explain the choice for the HSDS as the preferred method to rank the
124 severity of the droughts for Deventer and Zutphen. Many historical drought reconstructions have been done on the
125 basis of natural proxy-data from dendroclimatological reconstructions. These focus on tree-ring analysis to
126 reconstruct tree growth that provides insights into precipitation and temperature levels. This can be expressed
127 along the PDSI as an estimate of relative dryness based on reconstructions of temperature and precipitation
128 (Brázdil et. al. 2018). Certain long-term dendroclimatological reconstructions, such as the OWDA for Europe and
129 parts of North-Africa, use a self-calibrating PDSI (scPDSI) to create year-by-year maps of reconstructed summer
130 droughts on a 5414-point half-degree longitude-by-latitude grid. The scPDSI has a high degree of spatial
131 comparability across a broad range of climatological regions, which allows for comparisons with other pre-
132 instrumental droughts, for example in North-America (Cook et. al. 2015).

133 One of the most commonly used indices to categorise drought-severity in Europe is based on the seven-point
134 ordinal index devised by Pfister during the 1980s, also named ‘Pfister Indices’ (Brázdil 2020; Nash et. al., 2021;
135 Pfister, Camenisch and Dobrovolný, 2018). These indices can indicate both temperature differences and variations
136 in precipitation. In the seven-point index for precipitation, values ranging from rather wet to extremely wet (+1 to
137 +3) and rather dry to extremely dry (-1 to -3) are used to typify periods on the basis of direct or proxy-based
138 information regarding precipitation within a certain area. Such an index cannot be built on descriptive documentary
139 evidence alone, ~~but~~ and should also include proxy-data, such as evidence from plant-phenology and
140 dendroclimatological analysis. A merely descriptive index would only be able to use a three-point scale, only
141 taking into account the extraordinary (-1 or +1) as a deviation from the average (0). Every seven-point index also
142 ~~needs~~ requires a reference period to denote the deviations from the average, which often consists of a series of
143 instrumental measurements from the period prior to the full onset of global warming, most commonly ~~from~~ 1906
144 to 1960 (Pfister, Camenisch and Dobrovolný, 2018).

145 Several studies into historical droughts within Europe have applied the seven-point index as a means to indicate
146 the severity of past droughts (Bauch et. al., 2020; Brázdil et. al. 2013; Camenisch and Salvisberg, 2020;
147 Leijonhufvud and Retsö, 2021). However, there are also certain limits to the seven-point index. Kiss and Nikolić
148 (2015), for example, remarked that the requirements for the index can hardly be met for the European Middle
149 Ages, where the amount of available documentary evidence is often insufficient to estimate the severity of drought
150 on a month-by-month basis. In their attempt to create a 400-year long drought-index for the cities of Bern and
151 Rouen, Camenisch and Salvisberg (2020) similarly argue that, given the available data from both cities – primarily
152 chronicles and municipal records from the fourteenth to the early eighteenth century – did not allow for all three

153 index values (-1 to -3) to be used. The sources from both city's only provide instances of extreme drought events,
 154 which left a significant mark on inhabitant's memory and prompted city governments to take action. Therefore,
 155 instead of using all three values, only extremely dry (-3) and very dry (-2) were used in their analysis, considering
 156 that the more frequent and less impactful droughts (-1) were usually not recorded. For both cities, most droughts
 157 during the 400-year period were characterised as very dry (-2), and only a few instances were classified as
 158 extremely dry (-3). The survey also led to the identification of specific accumulations of droughts, for instance, at
 159 the end of the fourteenth, second half of the sixteenth, and the 1670s and early, as seasonal difference was
 160 discovered as the droughts in Bern often ~~took place~~ occurred during the summer, while those in Rouen were more
 161 prevalent during their spring season.
 162

163 The previous conclusions can also be applied for the corpus of municipal sources that have been investigated for
 164 Deventer and Zutphen. However, the documentary data from Deventer and Zutphen does not allow for a precise
 165 month-by-month reconstruction, as the duration of the droughts is not mentioned in the primarily descriptive cannot
 166 be accurately reconstructed from the primarily descriptive data. Monthly records of precipitation are required, to
 167 categorise such droughts into a seven-point index. In this case, a drought can only be denoted as very dry (-2) after
 168 at least a one-and-a-half months of reduced precipitation, while the value of extremely dry (-3) is reserved for two
 169 or more months without rainfall (Camenisch and Salvisberg, 2020). As the archival data from Deventer and
 170 Zutphen do not provide insights into the length of certain droughts, only referring to 'long' or 'prolonged' periods
 171 of drought, which do not indicate without indicating a specific timeframe, the seven-point index cannot be applied.
 172 However, the primary references to drought concern descriptions of its human and economic impact on a societal
 173 level, which are also more accurate representations of past perceptions of drought than modern conceptions of
 174 precipitation and evaporation (Garnier, 2015). This data can be used according to the HSDS to delineate droughts
 175 on an impact-centred scale. The HSDS distinguishes droughts on the basis of societal reactions that can be found
 176 in various sources, which are classified in categories on a 1 to 5 scale (see table 1) from an absence of precipitation
 177 to full-scale social crisis. An additional category, -1, denotes instances where both qualitative and quantitative data
 178 are considered insufficient, but a drought reference is kept solely for the purpose chronological reconstruction
 179 (Garnier, 2014). This additional category does not apply to any of the cases discussed in this article.

180 **Table 1:** Historical Severity Drought Scale (for the sixteenth to nineteenth centuries), from Garnier (2014)

Index	Description
5	exceptional drought: no possible supply, shortage, sanitary problems, very high prices of wheat, forest fires
4	severe low-water mark: navigation impossible, lay-off of wheatmills, search for new springs, forest fires, death of cattle
3	general low-water (difficulties for navigation) and water reserves
2	local low-water in rivers, first effects on vegetation
1	absence of rainfall: rogations, evidences in texts
-1	insufficient qualitative and quantitative information but the event is kept in the chronological reconstruction

181
 182 In order to identify periods of drought, an extensive study of the above-mentioned sources was carried out. When
 183 available, reference books were used as an additional tool for finding entries connected to drought-related issues.

184 These concerned aspects like water provisioning, fires, watermills, and other matters related to waterworks and
185 shipping, as well as a dearth in foodstuffs and other items as a result of drought. Firstly, the sources for Deventer
186 were studied on a year-by-year basis, in which all entries were searched for direct or indirect references to drought.
187 This yielded ~~many a steady base of~~ results that formed the ~~basis-foundation~~ of the following archival research.
188 Second in line were petition books, which were also studied on a year-by-year basis. The daily resolutions were
189 not studied on a year-by-year basis because of the density of the source material, ~~which as this~~ would render an
190 extensive page-by-page study too time-consuming. Instead, the daily resolutions were studied primarily on the
191 basis of reference books and findings from other sources. In all instances, not only the drought years found in the
192 other sources were consulted in the daily resolutions, but also two years before and after a ~~drought~~
193 ~~reference~~~~reference to drought~~. This was deemed relevant given the insidious nature of drought and possibility that
194 source might display certain developments of a drought on an earlier basis. After the study for Deventer was
195 completed, the study of Zutphen ~~started off~~~~began~~ with an analysis of the largely digitised reference works
196 regarding the daily resolutions. The earlier discovered drought years for Deventer were used as reference points,
197 and were used to study specific years, including the years before and after.

198 For each city, the rough data was first copied into separate databases ~~for each city~~, after which the data were filtered
199 by setting aside references that did not directly relate to drought. These included references to future measures to
200 be taken when severe droughts would occur, or measures where the relation to a drought-event was less
201 ~~clear~~~~obvious~~. Secondly, the remaining drought-events were filtered for each city according to drought-type
202 (meteorological, agricultural, hydrological, socio-economic) and season. Hereafter, a chronological database was
203 created combining the data from Deventer and Zutphen in a chronological overview of the specific drought events
204 for each year. This specific overview was also used for the next step: ranking the severity of each drought per year
205 according to the HSDS.


206

207 4. Outcomes

208 The most common types of drought mentioned in documentary sources refer to instances of meteorological
209 drought, referring to a deficiency of precipitation over a specific period of time. This is usually followed by
210 agricultural drought, which refers to the effects of meteorological drought on agricultural production. Hydrological
211 drought takes into account the consequences of water shortages in rivers, streams, lakes, and underground water
212 tables, while socio-economic drought describes the effects of drought when the former causes widespread
213 economic and societal disruption, most commonly in the form of subsistence crises (Brázdil et. al., 2018; Wilhite
214 and Pulwarty, 2017). As municipal records usually only contain references to extreme weather events, the
215 descriptions of drought in the sources refer almost exclusively to extremities (Camenisch and Salvisberg, 2020;
216 Garnier, 2019).


217 Based on the indicators of drought and its severity in the studied sources, an HSDS index has been constructed
218 including ~~both the~~ data from Deventer and Zutphen (see fig.1). The index ranks droughts on an annual basis using
219 the five-point scale, although instances of purely meteorological droughts (scale 1) and its effects (rogation
220 ceremonies and public prayer) have not been ~~discovered~~~~found~~. In total, 33 years with drought have been
221 reconstructed. ~~This includes~~ 26 drought years for Deventer, 16 for Zutphen, and only nine coinciding years ~~for~~

222 both cities. Hydrological droughts with a significant impact on the city's waterway's and the availability of water
 223 (scale 3) are amongst the most common forms of drought described in the sources, occurring 24 times. More
 224 extreme hydrological conditions, those within scale 4, are less common but still make up a significant part of the
 225 recorded droughts, namely nine instances. Scale 5, denoting full-scale societal crisis and critical shortages of food
 226 and water, has not been identified.



227
 228 **Figure 1: Chronology and severity levels of droughts within Deventer and Zutphen according to the Historical**
 229 **Severity Drought Scale (HSDS), 1500–1795.**

230
 231 With regard to both Deventer and Zutphen (see fig. 2), hydrological drought is by far the most common type of
 232 drought described in the sources. In most cases, this-These refers to low water levels or a complete lack of water
 233 in certain rivers and canals, as well as a shortage of water in wells and pumps. Meteorological drought is more
 234 prevalent in sources from Deventer, although in general the descriptions refer exclusively to 'excessive', 'strong',
 235 'prolonged', or 'long-lasting' periods of drought, often accompanied with a reference to the hydrological effects,
 236 such as dried up waterways and wells. Agricultural drought is mentioned very rarely in the sources. There is only
 237 one reference from Deventer that explicitly mentions negative agricultural yields in the city's hinterlands as a
 238 result of a drought, and the fact that this led to increased prices for certain foodstuffs. Last but not least, socio-
 239 economic drought only occurs during very strong droughts, usually the result of an accumulation of events leading
 240 to a severe lack of water and a shortage of food and other goods. This specific factor seems absent in the sources
 241 from Deventer and Zutphen.



242

243

244

245

Figure 2: Difference in drought types per year for Deventer and Zutphen in terms of meteorological (Met), hydrological (Hyd), agricultural (Agr) and Socio-economic (Soc), during the period 1500-1795

246

247

248

249

250

251

252

253

254

255

256

257

While there are a number of different drought years for both cities (see fig. 2), there are ~~specific years that coincide~~ several coinciding years, although ~~this does~~ not always occur in terms of severity. The year 1615, for example, is ranked 3 for Deventer, yet 2 for Zutphen. The sources for Deventer for 1615 indicate both a period of drought and lack of water, while Zutphen did not seem to suffer from the low water levels on the IJssel river. Explanations for such differences in hydrological drought can be found in the geohydrological differences between both cities. Apart from the IJssel river, the groundwater tables of Deventer and Zutphen were also influenced by the influx of water from two other streams coming in from the east: the Schipbeek for Deventer and Berkel for Zutphen (see fig. 3). These streams fed the surrounding moats and canals of the cities, which determined the availability of water for milling, or the water level in the wells and pumps. The Schipbeek was a man-made stream, which since its creation in the fifteenth century often suffered from silting due to increased amounts of sediment, human pollution, and poor management. Hence, the Schipbeek was considered an unreliable source of water, in particular during periods of drought (Schutten, 1981). As a natural river, the Berkel suffered less from such


258 problems, and it was known as a relatively reliable supplier of water to the groundwater tables below Zutphen.
259 This could explain different impacts of hydrological drought between both cities. Nevertheless, many coinciding
260 drought years, such as 1733, 1753, 1772, 1779, 1781, and 1783, indicate similar levels of hydrological drought
261 for both cities. which points out similar effects of the rivers.—



262 **Figure 3: The locations of Deventer and Zutphen on a modern map of the Netherlands, indicating the IJssel river and**
263 **the Schipbeek and Berkel substreams (map by Bert Brouwenstijn, VU Amsterdam).**
264

265 A notable level of difference between the two cities is that of seasonality (see fig. 4). Deventer seems to have a
266 much higher rate of spring droughts – recorded between March and May – and summer droughts – recorded

267 between June and August –, while Zutphen displays a larger amount of winter droughts – recorded between
 268 December and February. It must be noted that this difference is also due to the higher density in data for Deventer.
 269 However, both cities seem to have witnessed an equal amount of autumn droughts – recorded between September
 270 and November –, which, together with summer droughts constitute the most common category of droughts based
 271 on seasonality.



272 **Figure 4: The number of droughts according to season for Deventer and Zutphen, 1500-1795.**

273
 274
 275 Similar to the research by Camenisch and Salvisberg, the results for Deventer and Zutphen also display specific
 276 clusters or accumulations of drought years that took place within a span of several, sometimes subsequent years.
 277 Droughts with a moderate to severe impact, ranking 3 or 4 on the HSDS, occurred during the years 1630–1640,
 278 1650–1652, 1662–1669, 1731–1733, 1781–1783, and 1790–1794. This does not include years in which references
 279 are made to the damaging effects of previous droughts, often a year or even multiple years after a severe drought
 280 occurred. Most of the severe droughts ranking 4 on the HSDS occurred during the second half of the eighteenth
 281 century, between 1753 and 1783.

282

283 **5. Examples from the sources**

284 It would go beyond the scope of this article to dive into the details of each specific drought year discovered for
 285 both cities. A brief overview of these can be found in appendix 1 at the end of the article. Nevertheless, to make
 286 sense of the otherwise rather abstract notions mentioned in the HSDS, it is necessary to provide a number of
 287 detailed examples. The number of examples has been restricted to the most extreme and detailed examples, some of
 288 which coincide for both Deventer and Zutphen. These are 1669, ~~1733~~-1753, ~~1781~~, and 1783.

289 **5.1. The year 1669**

290 Deventer witnessed a period of severe drought in September 1669, which, according to municipal documents, led
 291 to extraordinarily low water levels on the IJssel river. As a result, many of the wells and pumps in the city were
 292 rendered dry and unusable. The inhabitants described the lack of water as an inconvenience and public clamour

293 regarding the scarcity of water was heard throughout the city. One of the main concerns was the risk of fire, which
294 was worsened by the shortage of water. As for Zutphen, references to the shortage of water are less explicit for
295 September that year. Here, no explicit mention of water scarcity is made in the city governments documentation,
296 but the fear of fire becomes apparent in a resolution that directed the city crier to call upon all inhabitants to store
297 water in case of an uneventful fire. While the impact of the drought is very explicit for Deventer (scale 4), the
298 reference to compulsory storing of water for Zutphen (scale 3) also implicitly links to hydrological drought but
299 less to a direct societal impact or near-crisis situation.

300 ~~5.2. The year 1733~~

301 ~~The year 1733 seems to show the opposite in terms of references. As for Deventer, the impact of the drought was~~
302 ~~felt primarily during the summer, which led to a lack of water in the Schipbeek river that supplied water to the~~
303 ~~city's harbour and canals. However, whether this had an impact on the water levels in the city's wells and pumps~~
304 ~~is not mentioned. In Zutphen, the 1733 drought was first mentioned in October, when a genever distillery petitioned~~
305 ~~to the city government that their capacity to produce suffered due to the great shortage of water within the city. In~~
306 ~~this case, the effects of the hydrological drought are more explicit for Zutphen (rank 4) than for Deventer (rank 3).~~
307 ~~Nevertheless, it can be assumed that the lack of water in the Schipbeek hampered navigation and the supply of~~
308 ~~water power to Deventer's watermills. ———~~

309 5.3. The year 1753

310 During the year 1753, equally severe droughts are mentioned for both Deventer and Zutphen in terms of impact.
311 In Deventer, the effects of drought were first felt in June, when an 'excessive drought' (*excessive droogte*) led to
312 a shortage of water in the city's wells. This lack of water led to a general shortage of water that prompted the city
313 government to take action. In Zutphen, the impact of the drought was reported in September, which mentioned the
314 low water levels on both the IJssel and Berkel rivers that led to the 'paralysis' (*verlamminge*) of most wells and
315 pumps. This displays a similarity in drought severity (rank 4), which refers to societal setbacks, for example by
316 limiting water use, rather than a full socio-economic crisis, although the potential for the latter could have been
317 present.

318 ~~5.4. The year 1781~~

319 ~~In 1781, the severity of drought is indexed equally on the HSDS for both cities (rank 4). In July that year, the water~~
320 ~~level in the Schipbeek was reported to have once again reached an absolute low point, which was detriment to the~~
321 ~~city, although no further details of the negative impacts were recorded. It can be assumed, however, that the drying~~
322 ~~up of the Schipbeek must have been felt, as it would have certainly paralysed the watermills. The impact of drought~~
323 ~~in Zutphen was already felt in February, implying that the drought started in the winter. Here, the drought and low~~
324 ~~water levels resulted in a lack of navigation via the Berkel river and a limited operation of the city's watermills.~~
325 ~~However, no effects on the availability of water in both cities' wells and pumps is mentioned.~~

326 5.5. The year 1783

327 The most detailed drought year (rank 4) recorded for both cities occurred in 1783. In Deventer, the strong and
328 excessive drought led to a lack of water in most of the wells during around the beginning of August. Later during
329 that month, a rare instance of agricultural drought is also mentioned as the a great spring drought, which led to a

330 reduced yield in buckwheat. This implies that the prolonged drought probably set in during the spring-months,
331 while its effects did not become detriment until the end of the summer when the prices of cereals increased
332 significantly. In Zutphen, the effects were primarily felt by the drying up of the Berkel river, which led to a
333 standstill of all watermills at the beginning of August. Another likely effect of the drought of 1783 was an epidemic
334 of dysentery in both Zutphen and Deventer. In Zutphen, the onset of the epidemic in towns and villages around
335 the city was noticed in early August, while the first case within the city walls was recorded on the fourth day of
336 that month. The disease spread rapidly during the following months, and the epidemic must have lasted until the
337 end of October. The spread of water-borne diseases like dysentery can be attributed to a lack of clean, fresh water
338 as a result of drought, which prompted people to use polluted water, or to seek water from unsafe sources (Brázdil
339 et. al. 2020; Camenisch et. al. 2020; Garnier, 2019; Pribyl, 2020).

340 In general, the source material often refers to similar indicators of hydrological drought, which often hindered
341 socio-economic life, but rarely resulted in a widespread disruption of daily life. Instances of agricultural drought
342 and its effects on food prices or general subsistence are very rare and only account for one particular case; the year
343 1783, when the prolonged drought led to a shortage of water, shutdown of watermills, dearth in cereals, and an
344 outbreak of dysentery in both cities. However, the sources do not suggest that this led to a moment of crisis. There
345 were also notable differences in the responses to drought, which do not correspond one-on-one for both cities
346 during most years, despite the relative proximity and similarity of both cities in terms of geological and
347 hydrological circumstances and the systems of water provisioning.

348

349 6. Comparison with [the Van Engelen, Buisman, and IJnsen Temperature Series](#)

350 Compared to other countries, very little concrete data with regard to temperature and/or precipitation exist for the
351 Netherlands prior to the instrumental period after 1850. The Royal Netherlands Meteorological Institute (KNMI),
352 founded in 1854, has a collection of ‘antique data’, consisting of early instrumental observations from the
353 eighteenth and early nineteenth century. These datasets are comprised of observations from several weather
354 stations across the Netherlands. Most of the stations from which eighteenth century records exist are located in the
355 province of Holland – such as Amsterdam, Alkmaar, Bergen (North-Holland), Delft, Haarlem, Leiden Rijnsburg,
356 and Zwanenburg – leading to rather regional measurements more typical for the precipitation-rich western
357 provinces along the North Sea coast, not the inland provinces that are more susceptible to strong droughts. The
358 early records for the eighteenth century also contain very few consistent records regarding precipitation (Geurts
359 and Van Engelen, 1992). Most data ~~known for the pre-instrumental~~from this period consists ~~primarily~~of
360 reconstructions regarding winter and summer temperatures.

361


362 The longest list of pre-instrumental, and partially instrumental, estimations of winter and summer temperatures
363 ~~available via the KNMI was~~is compiled by Buisman~~d~~, in collaboration with Van Engelen, and IJnsen. Despite its
364 incredible length, running from ~~the year~~751 CE until 2000, this data~~set~~ is generally not well-known outside of
365 Dutch-speaking academia (Van Engelen, Buisman and IJnsen, 2001; Pfister, Camenisch and Dobrovolný, 2018).
366 ~~The~~is data-series was constructed with the use of various proxy-data from the early modern period, such as the
367 weather diary of German pastor David Fabricius for the larger Frisian area in the north of the Netherlands, a set of

368 frost-day notes from the German city of Kassel, the 'tow barge' records from De Vries and the Manley (1974)
369 records of monthly temperatures in central England. ~~Buisman and IJnsen also included~~It also includes data from
370 the ~~forementioned records of the~~ aforementioned weather stations (1706-1905). The winter – from November to
371 March – and summer – from May to September – temperatures in this series have been categorised along an annual
372 nine-point scale from 1 (extremely soft/cool) to 9 (extremely harsh/warm) (IJnsen, 2010). In addition to the
373 categorization of annual values, the series also contains annual temperature averages in degrees Celsius. This is
374 divided between average summer (JJA), winter (DJF) and annual (November-October) mean temperature.
375

376 For the comparison with the HSDS for Deventer and Zutphen, only values from 7/~~7~~ to 9/~~9~~, implying above
377 average summer and winter temperatures have been taken into account as relevant for possible correspondence
378 between drought and above or below average temperatures. Overall, the result of the comparison was rather
379 meagre. Only a handful of years displayed a correspondence between cases of moderate to strong and ~~extremely~~
380 ~~very~~ strong droughts – those ranking 3, 4 or 5 on the HSDS – and above or below average summer or winter
381 temperatures. Correspondences between droughts and high summer temperatures were found for the years 1534,
382 1556, 1669, 1733, 1779, 1781, and 1783. Only three years, 1556, 1781, and 1783, were ranked as extremely warm
383 (9). Only for 1672 there was a correspondence between drought below average winter temperatures (7). When
384 looking at the annual average temperature (in °C) for the summer months (JJA), a statistical comparison shows a
385 rather weak Pearson correlation ($r=0.17$) for the Deventer and moderate correlation ($r=0.45$) for Zutphen. This
386 suggest very weak to moderate correlations between the annual average summer temperatures and the HSDS for
387 either city. Comparing the average annual temperature series with the HSDS led to an even weaker ($r=0.04$)
388 correlation for Deventer, and a moderate correlation ($r=0.52$) for Zutphen. However, it must be noted that due to
389 the small set of years, these results only bear a rather low level of statistical significance.
390

Met opmaak: Lettertype: Niet cursief

Met opmaak: Lettertype: Niet cursief



398 **Figure 5: Scatterplots of the comparison between average annual summer and average annual temperatures**
 399 **with the HSDS for Deventer.**

Met opmaak: Lettertype: Vet

401 The low number of correspondence with the drought years for Deventer and Zutphen can indicate two aspects; 1)
 402 drought periods did not necessarily coincide with periods of above average or extreme heat (or winter droughts
 403 with extreme cold); 2) the series of temperatures provided by [Van Engelen, Buisman, and IJnsen](#) ~~do not~~ might also
 404 not provide precise enough information, given the reliance on non-local sources for the reconstruction of pre-
 405 instrumental temperature records. While modern data mentioned earlier show a trend of rising temperatures since
 406 the 1950s contributing to increased drought-risk in the eastern regions of the Netherlands (Phillip et. al., 2020),
 407 this is not ~~in line with the findings of necessarily in line with~~ the data presented in this article. ~~Similar historical~~
 408 ~~studies~~ A similar study with regard to northwestern Europe ~~suggested higher correlations between temperature and~~
 409 ~~droughts than for temperature and precipitation, which might indicate that drought indices refer primarily to above-~~
 410 ~~average temperatures and evapotranspiration, also suggest a lower influence of temperature on the severity of~~
 411 ~~drought compared to precipitation during the early modern period~~ (Leijonhufvud and Retsö, 2021). ~~Given the~~
 412 ~~relatively low correlation between the Van Engelen, Buisman, and IJnsen temperature series and HSDS, the latter~~
 413 ~~cannot be concluded for Deventer or Zutphen.~~ As such, aspect one can be supported ~~for Deventer and Zutphen~~ on
 414 the basis of the comparison with ~~Van Engelen, Buisman, and IJnsen~~ ~~Buisman-IJnsen~~. Aspect two can be used to
 415 prove that the reliance on data from various distant locations is not always useful when studying specific territories
 416 and localities. This can also be tested by using a large compiled index of drought-years for multiple nearby
 417 territories, which is the case with the SDI.

418

419 7. Comparison with the SDI

420 The SDI was created by Camenisch and Salvisberg (2020) with the use of pre-existing precipitation reconstructions
421 from documentary sources for the Netherlands and Belgium, Germany, France, and Switzerland between 1315
422 and 1715, applying the seven-point scale index. ~~When the data from Bern and Rouen was compared with the SDI,~~
423 ~~only the years 1556, 1567, and 1681, were present in all three indices. The comparison between Bern and Rouen~~
424 ~~also displayed a deviation in the data regarding certain ‘megadroughts’, as the extreme droughts of 1473 and 1540~~
425 ~~were only reported in Bern.~~ Because the SDI is based on years when a drought was reported ~~somewhere within a~~
426 ~~specific country across different territories,~~ the amount of drought-years is significantly higher than in more local
427 indices. When comparing their data from Bern and Rouen with the SDI, the number of corresponding droughts
428 was relatively low, namely a total of seventeen corresponding cases out of the 87 drought-years in the SDI.

429

430 When comparing the data between 1500 and 1715, there ~~were are~~ only eight corresponding drought-years, out of
431 52 instances mentioned in the SDI for this period. These concern ten instances in total; eight specifically with
432 regard to Deventer (1534, 1556, 1615, 1630, 1634, 1645, 1666, and 1669), two concerning both Deventer and
433 Zutphen (1615 and 1669), and none specifically for Zutphen. This indicates that 44 droughts recorded in the SDI
434 were not found ~~in the sources~~ for Deventer and Zutphen, while 14 instances of drought (1588, 1589, 1597, 1612,
435 1629, 1633, 1638, 1650, 1652, 1662, 1667, 1672, 1690, 1696) were documented specifically for Deventer and/or
436 Zutphen during this period, but do not occur in the SDI. ~~Comparing the HSDS values for Deventer with the SDI~~
437 ~~led to a rather weak negative correlation ($r=-0.36$).~~ Such a rather low degree in correspondence supports the
438 conclusions regarding Bern and Rouen, that generalised drought data cannot easily be applied to reconstruct or
439 strengthen knowledge of the specific local droughts. In fact, it shows that local sources can provide ~~better~~ insights
440 into droughts that may not appear in compiled data-sets, ~~which This~~ prompts the need to do more in-depth research
441 for multiple regions and localities to minimise faulty generalisations about the widespread effects of drought on
442 different parts of society.

443

444 8. Comparison with the OWDA

445 Camenisch and Salvisberg (2020) also compared their findings ~~for Bern, Rouen, and the SDI~~ with the OWDA. ~~The~~
446 ~~OWDA is~~ a freely accessible online database that provides year-by-year data – either via a dataset or an interactive
447 map – of drought severity throughout Europe and certain parts of North Africa and the Middle East on a 0.5 degrees
448 latitude/longitude grid, going back as far as 0 CE and coming to a halt in 2012. The OWDA displays drought-
449 severity on a scPDSI scale from extremely dry (-6) to extremely wet (6). It is based on a vast amount of
450 dendrochronological data for Europe, completed with additional information historical data on hydroclimatic
451 extremes, but only with regard to spring and summer drought conditions (Cook et. al., 2015). This is also the main
452 setback of the OWDA, as it can only be used to compare drought conditions from June to August. Another pitfall
453 is the scPDSI ranking-system, which has to be calibrated to other forms of indices, such as the seven-point Pfister
454 index or the HSDS. Camenisch and Salvisberg tested the OWDA against the data from individual indices of Bern
455 and Rouen, as well as the SDI. They used the censure of -2.5 on the scPDSI scale as the mark of moderate to severe

456 and extreme droughts. As expected, the comparisons with the drought indices for Bern and Rouen led showed low
457 similarities between the OWDA ($r=0.32$ and $r=0.22$) for the respective indices. The wider SDI yielded a more
458 moderate similarity ($r=0.42$) with the data from the OWDA, which was also the only statistically significant
459 outcome given difference in sample size. (the most results that can be regarded as statistically significant using the
460 Pearson correlation ($r=0.42$)).

462 For the comparison with the HSDS for Deventer and Zutphen, grid snapshots were generated for each
463 reconstructed drought year, using the area which includes Deventer and Zutphen (52.34 to 52.°N, and 6 to 6.48
464 °E) (see figure 8). Following the example of Camenisch and Salvisberg (2020), only values of -2.5 or lower were
465 taken into account for relevant comparisons, and no usable data was available for the years 1638 and 1662. The
466 outcome of the comparison was rather meagre, as from eleven drought years corresponding to relevant outcomes
467 of the OWDA survey (1534, 1615, 1630, 1634, 1652, 1666, 1669, 1753, 1790, 1793, and 1794), only one year,
468 1666, was relevant as it fell within the range of summer (JJA) drought. No usable data was also available for the
469 years 1638 and 1662. Another interesting aspect is that some of the major summer drought-years in the HSDS,
470 such as 1783, only received a ranking of -2 on the sePDSI scale of the OWDA. However, the OWDA data
471 for certain years, such as 1615, 1630, 1669, and 1793, which indicate indicating autumn and winter droughts, could
472 perhaps indicate that the effects of the summer droughts was still felt during the following seasons. A quantitative
473 comparison between the HSDS for Deventer and the OWDA has shown a moderate negative correlation ($r=0.44$),
474 while for Zutphen this yielded a mildly positive correlation ($r=0.37$). Once again, though there seems to be a
475 moderate correlation between the two datasets, it must be taken into account that the sample size for this
476 comparison remained rather small.

477 Perhaps the reconstructions using the OWDA are susceptible to the same criticism as the comparisons to the Van
478 Engelen, Buisman, and Jansen series, as well as the SDI. Individual Comparisons between the HSDS and these
479 datasets sometimes show strong deviation from the drought years reconstructed for Deventer and
480 Zutphen, which. This could indicate the more localised character of most droughts, focusing specifically on their
481 local effects and how these worked out on society. Yet it also shows the limits of dendroclimatological analysis
482 on the basis of tree rings as a proxy for drought, which highlights the value of using documentary sources as a
483 means to verify the occurrence of historic droughts (Bothe et. al., 2019; Maughan, 2022; Pribyl, 2020).

484 9. Discussion and Conclusion

485 This article aimed to provide the first documentary evidence-based look at pre-instrumental droughts in the eastern
486 Netherlands between 1500 and 1795, focusing on two case studies: the cities of Deventer and Zutphen. This was
487 done by 1) examining the possibility of urban municipal archives to reconstruct past droughts; 2) creating drought
488 indices for both cities; and 3) by comparing the gathered data with other indices to spot possible correspondence.

489 The archives of Deventer and Zutphen contain plenty of municipal records that provided impact-based instances
490 of drought from the early sixteenth to the late eighteenth century. As for Deventer, slightly longer-running and a
491 larger amount records are available compared to Zutphen, where consistent records, such as daily resolutions date
492 back from the second half of the sixteenth century. Nevertheless, similar examples of drought-related measures
493 were found that indicate how droughts affected both cities primarily in terms of hydrological circumstances. The

494 most common issues are related to low water levels in the rivers and canals around the city hampering navigation
495 and low groundwater tables leading to a lack of water in wells and pumps. The main problem with the information
496 from the documentary evidence from both archives is that although it provides a good view on the impact of
497 drought in cities like Deventer and Zutphen, it remains difficult to establish the exact duration of droughts. The
498 extent of droughts is only mentioned in terms of general wordings like 'prolonged' and 'extraordinary. As of such,
499 the seven-point index, in which drought-severity is measured according to monthly thresholds, cannot be applied
500 the data found for Deventer and Zutphen.

501 The alternative, creating an index along the HSDS, applies better to the source-material, yet it is less precise as
502 the seven-point index, which is also calibrated using an instrumental reference-period. Nevertheless, using the
503 HSDS for Deventer and Zutphen has led to an index with a total of 33 droughts of varying severity on the scale of
504 1 (deficiency of precipitation) to 5 (widespread societal disruption/erisis) for the period 1500–1795. As is the case
505 with municipal records, only extreme instances of drought are reported, most of which appeared to fall within the
506 range of scale 3 and 4, denoting primarily hydrological droughts in the forms of dried up waterways, wells, and
507 pumps. Widespread societal disruption in terms of scale 5 was not discovered in the sources, which indicates that
508 the droughts had a disturbing rather than a crippling effect on society. The data from both cities also suggests a
509 difference in seasonality, as there seems to be an unequal distribution between spring and summer droughts. There
510 were also notable differences between similar indexed drought years for both cities, by which the effects of drought
511 were reported differently to indicate similar levels of severity, for example by referring to dried up wells in
512 Deventer and shut-down watermills in Zutphen. Although both instances indicate a scale 4 drought on the HSDS,
513 referring to hydrological circumstances leading to socio-economic drought, it can be questioned whether both
514 examples were considered as equally severe by contemporaries. Was a low-water mark in wells and pumps, for
515 instance, considered just as bad as a period without the ability to employ watermills? The descriptive nature of the
516 HSDS makes it a valuable index for the study of qualitative data from municipal records, although the next step
517 should be to calibrate such data according to a more precise scale. This scale should be based on different
518 conceptions from contemporary records to determine drought-severity more precisely. This can be done by
519 extending the categories into different levels of, for example, hydrological drought. For instance, a lack of
520 navigation and lay-off of watermills can be regarded as more critical or disastrous compared to a general shortage
521 of water for domestic purposes like cooking and washing, while the need for a stable availability of water for
522 firefighting purposes could be regarded as more important regarding the wide-ranging socio-economic effects a
523 major fire could have on the city as a whole (Garrioch, 2018).

524 Comparison with other indices, such as the Van Engelen, Buisman, and-IJnsen temperature series, the SDI, and
525 the OWDA, have yielded different insights ~~with regard to the~~when compared to the ~~data from this study~~HSDS.
526 This was both the case in terms of quantitative and direct comparisons between the different datasets. The
527 comparisons with the Van Engelen, Buisman, and-IJnsen temperature series, yielded weak to moderate results for
528 average annual summer temperatures, displaying no strong correlation between droughts and temperature for the
529 HSDS regarding Deventer and Zutphen. -turned out to be unfruitful, probably because temperature was of less
530 influence on these droughts, and. The latter could also be influenced by the fact that because the dataset compiled
531 with input from multiple areas outside of the Netherlands cannot be used to create regional or localaccurately
532 reconstructions ~~of~~ extreme temperatures on a local scale. The comparison with the SDI for the sixteenth and
533 seventeenth centuries led to a similar limited number of corresponding drought years, ~~which indicates also~~

534 ~~indicating~~ that ~~such~~ supra-regional indices ~~do not correspond~~ often have little correspondence ~~one-on-one~~ with
535 more localised documentary-based drought reconstructions. The same can be said of the comparison with the data
536 gathered from single-year based snapshots from the OWDA. In this case the correspondence was even lower
537 regarding the sole focus on summer droughts, although the indications for certain years could point towards
538 possible long-lasting effects of summer droughts during consecutive months. For each comparison, however, the
539 limited size of the dataset for the HSDS concerning Deventer and Zutphen made quantitative analysis and
540 comparisons difficult to render on a high degree of statistical significance. To enhance this, more data from several
541 locations could be added to the existing dataset to create a more encompassing series along the HSDS for the
542 eastern Netherlands, or the country as a whole.——

543 All in all, Nevertheless, the data for Deventer and Zutphen display ~~both~~ evidence for a small number of ~~wider~~
544 supra-regional droughts ~~as well as,~~ but the sources primarily indicate a larger number of local droughts specifically
545 mentioned in the documentary sources for the period under study. These concern ~~primarily~~ moderate to severe
546 instances of drought that impacted society and prompted responses from the city government to avert possible
547 negative outcomes, such as food and water shortages. As such, the source ~~materials~~ to reconstruct droughts ~~is~~ are
548 closely connected to the societal responses to drought, which indicates that specific instances of drought, primarily
549 hydrological drought, impacted society not necessarily by causing a widespread crisis but by limiting the use of
550 water and waterways. The urban sources also record very little instances of agricultural drought, of which only
551 once instance was found for a 300-year period. Remarkable is also that, at least for Deventer, the 'megadrought'
552 of 1540 is entirely absent in the sources. As Camenisch and Salvisberg (2020) demonstrated, however, this is not
553 rare with regard to more localised reconstructions. Although major European drought events, as in 1540, feature
554 widely in supra-regional indices, ~~which are~~ comprised of documentary and natural proxy data from across different
555 regions (Wetter et al., 2014), they are less likely to show in more local, urban ~~documentary analyses~~ evidence.
556 Drought reconstructions for specific locations, whether cities or villages with adequate data density, therefore
557 should be taken into account when compiling large-scale drought reconstructions, to gain a more accurate picture
558 of the regional and local spread of drought and its severity in terms of societal impact.

559
560 However, comparisons between specific, localities is another aspect that requires more attention. Deventer and
561 Zutphen, for example, despite their similarities and close proximity to one another yield a number of different
562 drought years. This can be explained, in part, by a difference in source-density for specific periods. More and
563 longer-running series of sources were available for Deventer, but considering the relative consistency and duration
564 of the municipal records for both cities it could also be argued that droughts were not always perceived as equally
565 menacing. Explanations for this can be found in the source-type, municipal records, which mostly refer only to
566 high-impact drought-events that required a governmental response, but also at the local level, for example by
567 studying the hydrological, geological, and socio-economic aspects of each city. This would include the dependence
568 of specific water sources for a city's economy, such as the need to operate watermills, or the general system of
569 water provisioning and how this was impacted across different areas within a city. Differing hydrological or socio-
570 political means that strengthened or helped to alleviate the effects of past drought could thus play an important
571 part in determining the severity of drought on a local level (Metger and Jacob Rousseau, 2020). This could provide
572 a better image of droughts through human actions and natural circumstances that have an influence on the local
573 impact and severity of drought and other climatic hazards, which counts not only for the past but also the future

574 (Degroot et. al., 2021; Kchouk et. al., 2021; Savelli et. al., 2022; Van Loon et. al., 2016). More research is needed
575 in order to draw broader conclusions on the specific local impacts of urban droughts, and how this was influenced
576 by local natural or human factors over time.

577

578 **Data availability**

579 The data used in this article is included in two supplements attached to this article. The archival sources used for
580 the research of this paper are publicly and/or digitally accessible via the websites of the HCO
581 (<https://collectieoverijssel.nl/>) and ZuRAZ. (<https://erfgoedcentrumzutphen.nl/>) and can be found in appendix 1.

582 [The Van Engelen, Buisman, and IJnsen temperature series is available via the website of the Royal Netherlands
583 Meteorological Institute \(<https://www.knmi.nl/nederland-nu/klimatologie/daggegevens/antieke-waarnemingen>\).](https://www.knmi.nl/nederland-nu/klimatologie/daggegevens/antieke-waarnemingen)

584 The SDI is available as a supplement to the article by Camenisch and Salvisberg ([https://doi.org/10.5194/cp-16-
585 2173-2020](https://doi.org/10.5194/cp-16-2173-2020)). The OWDA can be freely consulted via the project website (<http://drought.memphis.edu/OWDA/>).

586 **Supplement**

587 The supplement related to this article is available via: <https://doi.org/10.17026/dans-x3p-camy>

588 **Competing Interests**

589 The authors declare that they have no conflict of interest.

590 **Acknowledgments**

591 This article was written as part of the research project: Coping with drought. An environmental history of drinking
592 water and climate adaptation in the Netherlands. Funding and necessary support for this research was provided by
593 the Dutch Research Council (Nederlandse Organisatie voor Wetenschappelijk Onderzoek, NWO).

594 **Financial support**

595 This research has been fully supported by Dutch Research Council (NWO) under file number 406.18.HW.015.

596

597

598

599

600

601

602

603

604

605

606

607 **Appendix 1: Archival sources**

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Met opmaak: Nederlands (standaard)

608 Historisch Centrum Overijssel (HCO) (Regional Archives of Overijssel), Deventer, Stad Deventer, periode
609 Middeleeuwen, 1241-1591 (ID 0690), Edicta magistratus die buyspraecht genoemtp or Dat boick der
610 buyspraiken, 1459-1538, 1555-1596, 135.1, 3.

611 Historisch Centrum Overijssel, Deventer, Schepenen en Raad van de stad Deventer, periode Republiek 1591-1795
612 (ID 0691), Prothocoll des Rades van dagelicken resolutien, or Liber quotidianarum resolutionum civitatis
613 Daventriensis, 1591-1795, 4.14,

614 Historisch Centrum Overijssel, Deventer, Schepenen en Raad van de stad Deventer, periode Republiek 1591-1795
615 (ID 0691), Register van resolutien van Schepenen en Raad en Gezworen Gemeente (Concordaten), 1600-1794,
616 6a-m.

617 Historisch Centrum Overijssel, Deventer, Schepenen en Raad van de stad Deventer, periode Republiek 1591-1795
618 (ID 0691), Register van verordeningen en bekendmakingen van het stedelijk bestuur (Buurspraakboek) or Liber
619 publicationum, 7a-g.

620 Erfgoed Centrum Zutphen (ZuRAZ) (Regional Archives of Zutphen and surrounding areas), Zutphen, Oud-
621 Archief van de stad Zutphen, 1206-1815 (ID 0001), Memorien- en resolutieboek van de stad Zutphen, registers
622 van resoluties van de magistraat, 1573-1808, 2, 3, 6, 8, 18, 32, 35, 37, 46, 49, 50, 52.

623 Erfgoed Centrum Zutphen, Zutphen, Oud-Archief van de stad Zutphen, 1206-1815 (ID 0001), Repertoria op de
624 resoluties van de magistraat, 1573-1620, 110.

625 Erfgoed Centrum Zutphen, Zutphen, Oud-Archief van de stad Zutphen, 1206-1815 (ID 0001), Repertoria op de
626 resoluties van de magistraat, 1620-1660, 111.

627 Erfgoed Centrum Zutphen, Zutphen, Oud-Archief van de stad Zutphen, 1206-1815 (ID 0001), Repertoria op de
628 resoluties van de magistraat, 1661-1700, 112.

629 Erfgoed Centrum Zutphen, Zutphen, Oud-Archief van de stad Zutphen, 1206-1815 (ID 0001), Repertoria op de
630 resoluties van de magistraat, 1701-1740, 113.

631 Erfgoed Centrum Zutphen, Zutphen, Oud-Archief van de stad Zutphen, 1206-1815 (ID 0001), Repertoria op de
632 resoluties van de magistraat, 1741-1780, 114.

633 Erfgoed Centrum Zutphen, Zutphen, Oud-Archief van de stad Zutphen, 1206-1815 (ID 0001), Repertoria op de
634 resoluties van de magistraat, 122.

635

636 References

637 Aghakouchak A. et. al., Anthropogenic Drought: Definition, Challenges, and Opportunities, *Reviews of*
638 *Geophysics*, 59, 1-23, <https://doi.org/10.1029/2019RG000683>, 2021.

639 Bauch, M. et. al., A prequel to the Dantean Anomaly: the precipitation seesaw and droughts of 1302 to 1307 in
640 Europe, *Clim. Past*, 16, <https://doi.org/10.5194/cp-16-2343-2020>, 2020.

641 Bothe O. et. al., Inconsistencies between observed, reconstructed, and simulated precipitation indices for England
642 since the year 1650 CE, *Clim. Past*, 15, 307-334, <https://doi.org/10.5194/cp-15-307-2019>, 2019.

643 Brázdil, R. et. al., European climate of the past 500 years: new challenges for historical climatology, *Clim. Change*,
644 101, 7-40 DOI 10.1007/s10584-009-9783-z, 2010

645 Brázdil, R. et. al., Droughts in the Czech Lands, 1090-2012 AD, *Clim. Past*, 9, 1985-2002. doi:10.5194/cp-9-
646 1985-2013, 2013

647
648 Brázdil, R. et. al., Documentary and Instrumental-based drought indices for the Czech Lands back to AD1501,
649 *Clim. Research.*, 70, 103-117, doi: 10.3354/cr01380, 2016.

650

Gewijzigde veldcode

Gewijzigde veldcode

651 Brázdil, R. et. al. Documentary data and the study of past droughts: a global state of the art, *Clim. Past*, 14, 1915-
652 1960, <https://doi.org/10.5194/cp-14-1915-2018>, 2018.

653

654 Brázdil, R. et. al., Extreme Droughts and Human Responses to them: the Czech Lands in the Pre-Instrumental
655 Period. *Clim. Past*, 15, 1-24 <https://doi.org/10.5194/cp-15-1-2019>, 2019.

656 Brázdil, R. et. al.: Droughts in Historical Times in Europe, as derived from Documentary Evidence, in:
657 Palaeohydrology. Traces, Tracks and Trails of Extreme Events, edited by. Herget J. and Fontana, A., Springer,
658 Cham 65-96. <https://doi.org/10.5194/cp-16-2125-2020>, 2020.

659 Briffa, K. van der Schier G. and Jones P., Wet and dry summers in Europe since 1750: evidence of increasing
660 drought, *International Journal of Climatology*, DOI: 10.1102/joc. 1836, 2009.

661

662 Buisman, J. Duizend jaar weer, wind en water in de Lage Landen, Van Wijnen, Franeker,
663 1995/1996/1998/2000/2006/2015.

664

665 Camenisch, C. and Salvisberg M. Droughts in Bern and Rouen from the 14th to the beginning of the 18th century
666 derived from documentary evidence, *Clim. Past*, 16, 2173-2182. <https://doi.org/10.5194/cp-16-2173-2020>, 2020.

667

668 Camenisch, C. et. al. Extreme Heat and Drought in 1473 and their Impacts in Europe in the Context of the early
669 1470s, *Reg. Environ. Change*, 20, <https://doi.org/10.1007/s10113-020-01601-0> 2020, 2020

670

671 Cook, E. et. al., Old World megadroughts and pluvials during the Common Era, *Science Advances*, 1, DOI:
672 10.1126/sciadv.150056, 2015.

673

674 Degroot D. et. al., Towards a rigorous understanding of societal responses to climate change, *Nature*, 591, 539-
675 550, <https://doi.org/10.1038/s41586-021-03190-2>, 2021

676

677 Dominguez-Castro F. et. al., Assessing extreme droughts in Spain during 1750-1850 from rogation ceremonies,
678 *Clim. Past*, 8, 705-722, 2012.

679 Garnier E., Strengthened Resilience from Historic Experience. European Societies Confronted with
680 Hydrometeor in the Sixteenth to Twentieth Centuries, in: *Hydrometeorological Hazards. Interfacing Science and
681 Policy*, edited by Quevaeviller Ph., Wiley & Sons, New York, 2014

682

683 Garnier E., European historic droughts beyond the modern instrumental records 16th-20th centuries, in: *Drought:
684 Research and Science-Policy Interfacing*, edited by Andreu J. et. al., CRC Press, Boca Raton, 2015.

685

686 Garnier, E. Historic Drought from Archives: Beyond the Instrumental Record, in: *Drought. Science and Policy*,
687 edited by Iglesias, A., Assimakopoulos, D., and Van Lanen, H.A.J., Wiley-Blackwell, New York, 45-67, 2019.

688

689 Garrioch D., Towards a fire history of European cities (late Middle Ages to late nineteenth century), *Urban
690 History*, 46, 202-224, <https://doi.org/10.1017/S0963926818000275>, 2018.

691

692 Gorostiza, S. Escayol M. and Barriendos M., Controlling water infrastructure and codifying water knowledge:
693 institutional responses to severe drought in Barcelona (1620-1650), *Clim. Past*, 17, 913-927,
694 <https://doi.org/10.5194/cp-17-913-2021> , 2021.

695

696 Grau-Sattoras M. et. al., Prudent Peasantries: Multilevel Adaptation to Drought in Early Modern Spain (1600-
697 1715), *Environment and History*, 27, 3-36, [https://doi.org.vu-
698 nl.idm.oclc.org/10.3197/096734019X15463432086964](https://doi.org/vu-nl.idm.oclc.org/10.3197/096734019X15463432086964) 2021.

699

700 Geurts H. and Van Engelen A., Beschrijving antieke meetreeksen, Koninklijk Nederlands Meteorologisch
701 Instituut: Historische weerkundige waarnemingen, KNMI, De Bilt, part V, 1992.

702

703 Kchouk, S. et. al., A review of drought indices: predominance of drivers over impact and the importance of local
704 context, *Nat. Haz. Syst. Sci.*, [preprint], <https://doi.org/10.5194/nhess-2021-152>, 17 June 2021.

705

706 Kiss, A. and Nikolić Z., Droughts, Dry Spells and Low Water Levels in Medieval Hungary (and Croatia) I: The
707 Great Droughts of 1362, 1474, 1479, 1494 and 1507, *Journal of Environmental Geography*, 8, 11-22, Doi:
708 10.1515/jengeo-2015-0002, 2015.

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

709
710 Kiss, A., Droughts and Low Water Levels in Late Medieval Hungary II: 1361, 1439, 1443-4, 1455, 1473, 1480,
711 1482(?), 1502-3, 1506: Documentary versus Three-ring (OWDA) Evidence, *Journal of Environmental*
712 *Geography*, 10, 43-56, DOI: 10.1515/jengeo-2017-0012, 2017.
713
714 Kiss, A., The great (1506-)1507 drought and its consequences in Hungary in a (Central) European context, *Reg.*
715 *Environ. Change*, 20, <https://doi.org/10.1007/s10113-020-01634-5> 2020
716
717 Leijonhufvud L. and Retsö D., Documentary evidence of droughts in Sweden between the Middle Ages and ca.
718 1800 CE, *Clim. Past*, 17, 2015-2029, <https://doi.org/10.5194/cp-17-2015-2021>, 2021.
719
720 Machairas, I. and Van de Ven, F., An urban drought categorization framework and the vulnerability of a lowland
721 city to groundwater urban droughts, *Nat. Hazards*, 116, <https://doi.org/10.1007/s11069-022-05767-0>
722
723 Maughan, N. et. al., Societal impacts of historical droughts in a warming world, *Reg. Environ. Change*, 22,
724 <https://doi.org/10.1007/s10113-022-01935-x>, 2022.
725
726 Mukherjee S. Mishra A. and Trenberth K., Climate Change and Drought: a Perspective on Drought Indices,
727 *Current Climate Change Reports*, 4, 145-163, <https://doi.org/10.1007/s40641-018-0098-x>, 2018.
728
729 Nash D. et. al., Climate indices in historical climate reconstructions: A global state-of-the-art, *Clim. Past*, 17,
730 1273-1314, Doi [10.5194/cp-17-1273-2021](https://doi.org/10.5194/cp-17-1273-2021), 2021.
731 Pfister, C. Evidence from the Archives of Societies: Institutional Sources, in: *The Palgrave Handbook of Climate*
732 *History*, edited by White S. et. al., Palgrave Macmillan, London, 67-81, [https://doi.org/10.1057/978-1-137-](https://doi.org/10.1057/978-1-137-43020-5_4)
733 [43020-5_4](https://doi.org/10.1057/978-1-137-43020-5_4), 2018,
734
735 Pfister C. Camenisch C. and Dobrovolný P., Analysis and Interpretation: Temperature and Precipitation Indices,
736 in: *The Palgrave Handbook of Climate History*, edited by White S. et. al., Palgrave Macmillan, London, 115-
737 129, https://doi.org/10.1057/978-1-137-43020-5_11, 2018.
738
739 Phillip, S. et. al., Regional differentiation in climate change induced drought trends in the Netherlands, *Environ.*
740 *Res. Lett.*, 15, <https://doi.org/10.1088/1748-9326/ab97ca>, 2020.
741
742 Piervitali E. and Colacino M., Evidence from Drought in Western Sicily During the Period 1565-1915 From
743 Liturgical Offices, *Clim. Change*, 49, 225-235, 2001.
744
745 Pribyl K. and Cornes R., Droughts in Medieval and Early Modern England, part 1: the evidence, *Weather*, 75,
746 168-172, doi:10.1002/wea.3599, 2020.
747
748 Pribyl, K., A survey of the impact of summer droughts in southern and eastern England, 1200-1700, *Clim. Past*,
749 16, 1027-1041. <https://doi.org/10.5194/cp-16-1027-2020>, 2020.
750
751 Savelli, E. et. al., Drought and society: scientific progress, blind spots, and future prospects, *WIREs Clim.*
752 *Change* 761, DOI: 10.1002/wcc.761, 2022.
753
754 Schutten, G., Varen waar geen water is. Reconstructie van een verdwenen wereld. Broekhuis, 1981.
755
756 Spinoni J. et. al., Will drought events become more frequent and severe in Europe?, *International Journal of*
757 *Climatology*, 38, 1718-1736, DOI: 10.1002/joc.5291, 2018.
758
759 Stangl, M. and Foelsche U., Climate History of the Principality of Transylvania during the Maunder Minimum
760 (MM) Years (1645-1715 CE) Reconstructed from German Language Sources, *Climate*, 10,
761 <https://doi.org/10.3390/cli10050066>, 2022.

762 Szalinska W., Otop I. and Tokarczyk T., Local urban risk assessment of dry and hot hazards for planning
763 mitigation issues, *Climate Risk Management* 34, doi:10.1016/j.crm.2021.100371, 2021.

764

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Met opmaak: nova-legacy-e-list_item, Patroon: Doorzichtig (Wit)

765
766 Van Engelen A., Buisman J. and IJnsen F., A millenium of weather, winds and water in the Low Countries, in:
767 History and Climate, Memories of the Future? Edited by Jones et. al., Kluwer Academic/Plenum Publishers,
768 101-124, 2001
769
770 Van Loon, A. et. al., Drought in a human-modified world: reframing drought definitions, understanding, and
771 analysis approaches, Hydrol. Earth Syst. Sci., 20, 3631-3650, doi:10.5194/hess-20-3631-2016, 2016.
772
773 Vogelzang, I. De drinkwatervoorziening van Nederland voor de aanleg van de drinkwaterleidingen. Joh. Mulder,
774 Gouda, 1956.

775 Vörösmarty C. et. al., Humans Transforming the Global Water System, Eos, Transactions American Geophysical
776 Union, 85, 509-514,

777 Wetter, O. et. al., The year-long unprecedented European heat and drought of 1540 – a worst case, Clim. Change,
778 125, 349-363, DOI 10.1007/s10584-014-1184-2, 2014.

779 Wilhite D. and Pulwarty R., Drought as Hazard: Understanding the Natural and Social Context, in: Drought and
780 Water Crisis: Integrating Science, Management, and Policy, 2nd edition, edited by Wilhite D. and Pulwarty R.,
781 CRC Press: Boca Raton FL, 3-20, 2018.
782

783 IJnsen, F., Methoden van onderzoek naar 12 eeuwen temperatuur in Nederland. Verslag van de gevolgde
784 statistische aanpak, Stiens, 2010.

785
786
787
788