

Multi-proxy assessment of brachiopod shell calcite as a potential archive of seawater temperature and oxygen isotope composition

Thomas Letulle¹, Danièle Gaspard², Mathieu Daëron³, Florent Arnaud-Godet¹, Arnaud Vinçon-Laugier¹, Guillaume Suan¹, Christophe Lécuyer¹.

¹ Univ Lyon, UCBL, ENSL, UJM, CNRS, LGL-TPE, F-69622, Villeurbanne, France

² UMR 7207, Centre de Recherche en Paléontologie, Paris (CR2P), CNRS, MNHN, Sorbonne- Université. Muséum national d'Histoire naturelle, 8 Rue Buffon, CP 38, 75005 Paris, France.

³ Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Orme des Merisiers, F-91191 Gif-sur-Yvette Cedex, France.

Correspondence to: Thomas Letulle (thomas.letulle@univ-lyon1.fr)

Abstract

Most of our knowledge of past ~~seawaterocean~~ temperature history is based on the $\delta^{18}\text{O}$ ~~valuesmeasurements~~ of calcium carbonate fossil shells. However, the determination of past temperature using this proxy requires the knowledge of past ocean $\delta^{18}\text{O}$, which is generally poorly constrained. Other ~~carbonate-based~~ paleothermometers ~~using carbonate archives~~, such as Mg/Ca ratios, and clumped isotopes (Δ_{47}), have been developed to estimate independently paleotemperatures, and allow past ocean $\delta^{18}\text{O}$ ~~values~~ to be calculated using various groups of calcifying organisms. Articulated brachiopods are some of the most commonly used ~~archives~~ in studies of past oceanic geochemistry and temperature. They are abundant in the fossil record since the Cambrian, ~~for decades~~, and their low Mg-calcite mineralogy has been considered relatively resistant to diagenetic alteration ~~for decades~~. Here, we investigate the potential of brachiopod shells as recorders of ~~seawatergrowing~~ temperatures and seawater $\delta^{18}\text{O}$ ~~values~~ using new brachiopod shell geochemical data, by testing multiple ~~well~~ established or ~~suggested paleothermometers applied to carbonate archivessupposed carbonate-based paleothermometers~~.

Modern articulated brachiopod shells covering a wide range of temperatures (-1.9 to 25.5°C), depths (5 to 3431 m) and salinities (33.4 to 37.0 PSU), were analysed for their stable isotope compositions ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$ and Δ_{47}), and elemental ratios (Mg/Ca, Sr/Ca, Na/Ca and Li/Ca). Our data allowed us to propose a revised oxygen isotope fractionation equation between modern brachiopod shell calcite and seawater:

$$T = -5.0 (\pm 0.2) (\delta^{18}\text{O}_c - \delta^{18}\text{O}_{\text{sw}}) + 19.4 (\pm 0.4) \quad T = -5.2 (\pm 0.3) (\delta^{18}\text{O}_c - \delta^{18}\text{O}_{\text{sw}}) + 19.9 (\pm 0.8) \\ R^2 = 0.95 \quad (n=53)$$

Where $\delta^{18}\text{O}_c$ is in ‰VPDB, $\delta^{18}\text{O}_{\text{sw}}$ is in ‰VSMOW and T is in °C. ~~The measured Δ_{47} values show a strong correlation with growing temperatures~~ Our results strongly support the use of clumped isotopes as an alternative temperature proxy, but ~~is confirm~~ significantly offsets relative to the canonical relationship established for other biogenic and abiogenic calcium carbonate minerals. ~~Our results strongly support the use of clumped isotopes as an alternative temperature proxy and indicate that brachiopod Δ_{47} values can be used with $\delta^{18}\text{O}$ to estimate past $\delta^{18}\text{O}_{\text{sw}}$ with a precision of about $\pm 1\%$ VSMOW. The obtained Brachiopod shell~~ Mg/Ca ratios show no relationship with temperatures, indicating that this ratio is a poor recorder of past changes in temperatures, an observation at variance with several previous studies. ~~Brachiopod shell Sr/Ca, Na/Ca and Li/Ca display relatively good and~~ Despite significant correlations with brachiopod living temperature, ~~but data brachiopod shell Sr/Ca, Na/Ca and~~

Li/Ca values indicate the influence of environmental and biological factors unrelated to temperature, that undermines their potential as alternative temperature proxy. Kinetic effects (growth rates) could explain most of the deviation of brachiopod shell calcite from expected isotopic equilibrium with seawater, and part of the distribution of Sr/Ca, Na/Ca and Li/Ca ratios. Our proposed revision of marine temperature and water $\delta^{18}\text{O}$ proxies based on brachiopod shell geochemistry is promising to refine the record of these oceanic parameters in the Phanerozoic.

1. Introduction

Oxygen isotope ratios ($\delta^{18}\text{O}$) of brachiopod shell calcite constitute the most extensive record of marine temperatures over the Phanerozoic eon (Prokoph et al., 2008; Veizer and Prokoph, 2015). Nevertheless, our limited knowledge of $\delta^{18}\text{O}$ values of past seas and oceans prevent confident interpretation of this record, as carbonate $\delta^{18}\text{O}$ values ($\delta^{18}\text{O}_\text{c}$) is dependent on both shell growth temperature and growing-living water $\delta^{18}\text{O}$ ($\delta^{18}\text{O}_\text{sw}$) (Epstein et al., 1953; Kim and O'Neil, 1997; Kim et al., 2007; Brand et al., 2013, 2019). Other paleotemperature proxies, such as trace element ratios and clumped isotopes (Δ_{47}), may be the key to build a more confident record of Phanerozoic ocean temperatures. Owing to the determination over the last decades of reliable oxygen isotope fractionation equations between calcite and seawater, such independent temperature estimates can be combined with $\delta^{18}\text{O}_\text{c}$ measurements to reconstruct $\delta^{18}\text{O}_\text{sw}$ values and hence characterise oceanic or local hydrography.

Alternative carbonate archive thermometers have been developed for different calcifying organisms. and have been proposed in the literature as potentially applicable to brachiopod shells. Element/Ca ratios such as Mg/Ca, Sr/Ca, Li/Ca, are the most common in the literature. Mg/Ca thermometry of foraminiferal calcite (Nürnberg, 1995; Anand et al., 2003) is a widely used proxy that is the basis for reconstruction of changes in $\delta^{18}\text{O}_\text{sw}$ and ice volumes over the Cenozoic era (Lear, 2000; Billups and Schrag, 2003; Miller et al., 2020). Building on the pioneering work of Lowenstam (1961), Brand et al., (2013, 2019) documented a positive correlation between seawater temperature and the MgCO_3 content of brachiopod shells sampled at a worldwide scale, and, Plus, following conclusions of Jiménez-López et al. (2004), Brand et al. (2013) developed an isotopic fractionation equation considering the amount of Mg^{2+} in substitution to Ca^{2+} in the crystal lattice of the low-Mg calcitic shell (Brand et al., 2013). This MgCO_3 -corrected isotopic fractionation equation which was abandoned in their more recent works (Brand et al., 2019). MgCO_3 -temperature relationship is described by a curve named the “global brachiopod Mg line” (GBMgL), yet Brand et al., (2013, 2019) acknowledge species-specific deviations from this concept falling well above the said “line”. Brachiopod shell Mg content has been found to have also shows a strong taxonomic control trend, with shells of thecideid and craniid being made of high-Mg calcite, in contrast with the low-Mg calcite of rhynchonellide and terebratulide brachiopod shells (Brand et al., 2003; Ullmann et al., 2017). This undermines the brachiopod shell MgCO_3 -temperature relationships which uses a dataset of both thecideid, rhynchonellide and terebratulide (Brand et al., 2013, 2019).

While Sr/Ca ratios, are mostly used to assess the shell preservation of fossil specimens (Brand and Veizer, 1980), this ratio also constitutes a potential marine paleothermometer to be applied to brachiopod shells, as it is for corals (McCulloch et al., 1994; Shen et al., 1996; Marshall and McCulloch, 2002; Swart et al., 2002; Ayling et al., 2006; DeLong et al., 2011). Previous studies already suggested Sr/Ca ratios as marine paleothermometers for brachiopod shells owing to their correlations with oxygen isotope ratios and seawater temperature (Lowenstam, 1961; Mii and Grossman, 1994; Pérez-Huerta et al., 2008; Butler et al., 2015; Ullmann et al., 2017). Sr/Ca ratios

of brachiopod calcite were also suggested as marine paleothermometers owing to their correlations with oxygen isotope ratios and seawater temperature (Lowenstam, 1961; Mii and Grossman, 1994; Pérez Huerta et al., 2008; Butler et al., 2015; Ullmann et al., 2017). This elemental ratio, initially used to assess the shell preservation of fossil specimens (Brand and Veizer, 1980), also constitutes a marine paleothermometer in other groups such as corals (McCulloch et al., 1994; Shen et al., 1996; Marshall and McCulloch, 2002; Swart et al., 2002; Ayling et al., 2006; DeLong et al., 2011).

Similarly, brachiopod Li/Ca ratio was also suggested as a potential temperature proxy in the seminal study of Delaney et al. (1989). Subsequent studies have also confirmed that Li/Ca may constitute a potentially accurate thermometer in brachiopods (Dellinger et al., 2018; Rollion-Bard et al., 2019; Washington et al., 2020) and in corals (Montagna et al., 2014; Marchitto et al., 2018). For this latter group, however, Li/Mg ratios have been considered as a much more precise thermometer than Li/Ca (Montagna et al., 2014; Marchitto et al., 2018), despite some identified caveats such as the scrambling role of organic matter (Cuny-Guirriec et al., 2019).

On top of element/Ca proxies, $\delta^{47}\text{C}$ lumped isotope (Δ_{47}) thermometry has been developed more recently on different carbonate materials (Ghosh et al., 2006; Zaarur et al., 2013). This parameter measures the anomaly of ^{13}C - ^{18}O bonds within the carbonate lattice relative to their stochastic abundance. Δ_{47} values are strongly correlated to crystallisation temperature and independent from $\delta^{18}\text{O}_{\text{sw}}$. This proxy has been applied to different marine calcifying organisms to estimate $\delta^{18}\text{O}_{\text{sw}}$ in deep past (Petersen et al., 2016; Bergmann et al., 2018; Henkes et al., 2018; Wierzbowski et al., 2018; Price et al., 2020; Vickers et al., 2020, 2021; de Winter et al., 2021; Meckler et al., 2022). Although promising, brachiopod shell Δ_{47} composition was only used in a couple of studies to estimate past temperatures (Henkes et al., 2013; Came et al., 2014). Indeed, the few existing modern brachiopod Δ_{47} -temperature data are in good agreement within the canonical Δ_{47} -temperature relationship established for other biogenic and abiogenic calcium carbonate minerals (Henkes et al., 2013; Came et al., 2014). Recent compilation of Δ_{47} -T calibration data from different laboratories using the new carbonate-based standardisation (Bernasconi et al., 2018, 2021) concluded to a unique Δ_{47} -T relationship for synthetic and natural carbonates (Anderson et al., 2021). The establishment of such unique calibration equation strengthens the good agreement observed between previous natural and synthetic Δ_{47} -T relationships and would suggest that this calibration is valid when applied to brachiopod shells, especially as empirical calibrations are in very good agreement with theoretical predictions (Jautzy et al., 2020). However, Δ_{47} measurements from brachiopod shells are absent from this compilation, and other studies suggest that they may be subjected to significant deviations from Δ_{47} -T relationships likely related to differences in growth rates (Bajnai et al., 2018, 2020).

Beside, a number of studies suggested that living temperature alone may not explain the variability of those different parameters within brachiopod shells. First, biogenic carbonate precipitation rate depends on environmental and biological factors (temperature and salinity tolerance, age, reproduction period, ontogeny ...) which causes biases in the paleoenvironment record extracted from the shell (Peck et al., 1997; Schöne, 2008). Brachiopod shell calcite is secreted by the outer epithelial cells of the mantle (Williams, 1968; Simonet Roda et al., 2019a, b, 2022). Calcite elements are in close relationship with organic matrices which control the mineral growth (Williams, 1968; Gaspard et al., 2008; Gaspard and Nouet, 2016; Simonet Roda et al., 2019a, b, 2022). Finally, nanoparticles are transported via ion transport through the cell membrane, highlighting a calcite precipitation as an amorphous calcium carbonate precursor stage (Griesshaber et al., 2009; Schmahl et al., 2012).

The chemistry of the calcifying environment of brachiopod shell calcite is poorly constrained (Immenhauser et al., 2016). Testing thermodynamic equilibrium at the mineralisation site is therefore, to date, out of reach. Consequently, the chemistry of brachiopod shells is usually compared to that of the surrounding water, in order to approach the processes involved in biomineralisation from a chemical point of view. Numerous studies highlighted that brachiopod shell calcite does not precipitate *sensu stricto* in isotopic equilibrium with surrounding water but as best mimic thermodynamic equilibrium (Carpenter and Lohmann, 1995; Auclair et al., 2003; Parkinson et al., 2005; Yamamoto et al., 2010a, b; Cusack et al., 2012; Takayanagi et al., 2012, 2013, 2015; Bajnai et al., 2018, 2020; Romanin et al., 2018; Rollion-Bard et al., 2019). Based on the conclusions of Bajnai et al., (2018), Brand et al., (2019) characterise their oxygen isotope fractionation equation database as representing a “brachiopod based-equilibrium”. This term is opposed to the thermodynamic equilibrium, which is best described by very slow growing calcites (Coplen, 2007; Daëron et al., 2019), thus ensuring isotopic equilibration of the dissolved inorganic carbon (DIC) species with water (Watkins et al., 2013, 2014). Trace element incorporation into calcite is also affected by kinetic effects. In synthetic carbonates, the relative abundance of elements such as Sr, Na and Li rise with higher growth rates (Lorens, 1981; Busenberg and Plummer, 1985; Tesoriero and Pankow, 1996; Gabitov et al., 2011, 2014). Such kinetic trends were evidenced at an intra-individual level among brachiopods as documented by the negative correlation of element/Ca ratios with stable isotope values ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) (Ullmann et al., 2017; Rollion-Bard et al., 2019).

In this study, we assess the potential of various geochemical proxies ($\delta^{18}\text{O}$; Δ_{47} ; Sr/Ca; Mg/Ca; Li/Ca; Na/Ca) as recorders of seawater temperature by analysing a new set of modern articulated brachiopod shells collected from various depths and latitudes during institutional oceanographic cruises, and covering a broad range of water temperatures comprised between -1.9°C and 25.5°C. We discuss the validity and robustness of our oxygen isotope fractionation equation established with measured $\delta^{18}\text{O}_c$ of calcite, and seawater temperature and $\delta^{18}\text{O}_w$ estimated from oceanographic data. The dependence of elemental ratios to seawater temperature is discussed at the light of this new available dataset of modern brachiopods. Finally, we highlight kinetic effects as non-negligible sources of isotopic and trace element variability, registered in the dataset of brachiopod shell calcite and we provide suggestions on how they may be considered for proxy data interpretation.

2. Material and Methods

2.1 Sample collection and environmental parameters

The studied material consists of 37 articulated brachiopod shells that were collected *in situ* during institutional oceanographic cruises or other scientific missions (see supplementary data for detail). Sampling location was documented with depth and geographic coordinates for most samples. Temperature and salinity, which were not measured *in situ* for most samples, were estimated independently using the NOAA World Ocean Atlas 2018 (Locarnini et al., 2018; Zweng et al., 2018) or, whenever available, using long term local records in the literature (Table 1). For both temperature and salinity, mean annual values (MAT), as well as higher and lower monthly averages, were considered. Similarly, $\delta^{18}\text{O}_{\text{sw}}$ values were usually not measured *in situ*, but calculated from the salinity data with the appropriate regional $\delta^{18}\text{O}_{\text{sw}}$ -S relationship from LeGrande and Schmidt (2006). The oceanographic parameters used in this study are listed in Table 1.

155 2.2 Sample preparation

Encrusting organisms covering the shells were first mechanically removed using stainless steel dental tools and each shell was then placed in diluted bleach (NaClO 5 %), for 5 to 10 min in an ultrasonic bath to remove organic matter and other contaminants. As preliminary tests have revealed that the more delicate shells (ex: *Macandrevia africana* Cooper, 1975) broke apart during the ultrasonic bath, immersion in diluted bleach for few hours without ultrasonic bath was subsequently preferred for few specimens. In all cases, the specimens were rinsed with deionised water and oven-dried at 50°C for a few hours.

The sampling of brachiopod shell calcite was performed following the recommendations of Romanin et al (2018). The umbo, edges and the muscle scar area were avoided because they record major kinetic or metabolic effects (Carpenter and Lohmann, 1995; Auclair et al., 2003; Parkinson et al., 2005; Yamamoto et al., 2010b; Ullmann et al., 2017; Romanin et al., 2018), and sampling was focused on the middle part of the shell. Considering the complex structure of brachiopod shells (Williams, 1968; Gaspard et al., 2018; Simonet Roda et al., 2022), relative to the precision of our sampling method (dental tools and engraving bit), we were unable to prepare pure samples for each layer of the brachiopod shell (acicular primary layer, fibrous secondary layer and columnar tertiary layer). the unambiguous distinctions between the different shell layers could not be performed during sampling. In this study, we differentiate the outer layers, i.e., a mix of the primary acicular layer and some amount of the outer secondary fibrous layer, which record significant kinetic effects (Carpenter and Lohmann, 1995; Auclair et al., 2003; Parkinson et al., 2005; Bajnai et al., 2018; Romanin et al., 2018; Rollion-Bard et al., 2019), from the inner layers, i.e., the inner secondary fibrous layer and/or the tertiary prismatic-columnar layer when present. Outer shell layers were removed manually with dental tools and resulting powders were kept to compare their elemental ratios, $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ with the inner shell layers (Carpenter and Lohmann, 1995; Auclair et al., 2003; Pérez-Huerta et al., 2008). After removal of the outer layers, the inner layers were sampled using an engraving bit fitted to Dremel Micro™ drill adjusted to the lowest possible speed. For each sample, the powder was used for all selected geochemical analyses which correspond to $\delta^{18}\text{O}$, $\delta^{13}\text{C}$, Δ_{47} , and element concentration. As clumped isotope analyses require large amounts of material (>10 mg are necessary to operate multiple replicate analyses), a large area (a few cm²) was sampled in the middle part of the shell, allowing several mg of calcitic powder to be collected. It is worthy to note that this large sampling area may correspond to several months or years of shell growth depending on the species.

This sampling protocol was applied to most of our samples, except to the smallest and most fragile shells that quickly broke apart during sampling. Consequently, in the case of the small shells of *Terebratulina latifrons*, *Frenulina sanguinolenta* and *Fallax neocaledonensis*, the outer layers were kept and only the area covering the umbo up to the muscle scar was avoided for sampling. The weakness and thinness of *M. africana* precluded a total

Table 1: Brachiopod samples with taxonomic identification, sampling location and environmental parameters. Most environmental parameters are derived from the World Ocean Atlas 2018 (Locarnini et al., 2018; Zweng et al., 2018). For a few samples, environmental parameters are derived from local literature as listed below. Almost all $\delta^{18}\text{O}_{\text{SW}}$ values are calculated using regional $\delta^{18}\text{O}_{\text{SW}}$ -Salinity relationship published by LeGrande and Schmidt (2006), unless direct measurements covering seasonal variations was available in the literature.

1. Goodwin and Cornelisen, (2012); 2. Woods et al. (2014); 3. Meredith et al. (2013); 4. Jacobson, (1983)

Specimen	Taxa	Living location	Depth (m)	MAT (°C)	Seasonal T variation (°C)	Salinity	Seasonal salinity	$\delta^{18}\text{O}_{\text{SW}}$ (‰ VSMOW)
New Caledonia								
FNEO-N4	<i>Fallax neocaledonensis</i>	22°54'S - 167°13'E	403-429	13.4	±1.2	35.11	±0.05	0.60
FNEO-M2	<i>Fallax neocaledonensis</i>	18°46'S - 163°16'E	600	7.7	±0.6	34.55	±0.03	0.45
FSAN-3	<i>Frenulina sanguinolenta</i>	20°52'S - 167°08'E	05-20	25.4	±1.8	35.18	±0.10	0.62
SCRO-1	<i>Stenosarina crosnieri</i>	22°59'S - 167°19'E	525	10.0	±0.9	34.76	±0.01	0.51
SCRO-3	<i>Stenosarina crosnieri</i>	22°59'S - 167°19'E	525	10.0	±0.9	34.76	±0.01	0.51
SGLO-S1	<i>Stenosarina globosa</i>	19°06'S - 163°30'E	215-225	19.7	±0.7	35.61	±0.01	0.73
SGLO-M1	<i>Stenosarina globosa</i>	18°59'S - 163°24'E	320	16.4	±0.7	35.40	±0.02	0.68
SGLO-M2	<i>Stenosarina globosa</i>	18°59'S - 163°24'E	320	16.4	±0.7	35.40	±0.02	0.68
Guadeloupe								
TGAL-3	<i>Tichosina cubensis</i>	15°53'N - 61°25'W	262-266	16.6	±1.0	36.22	±0.04	0.82
TGAL-4	<i>Tichosina cubensis</i>	15°53'N - 61°25'W	262-266	16.6	±1.0	36.22	±0.04	0.82
TCUB-2	<i>Tichosina cubensis</i>	16°20'N - 60°57'W	250	17.3	±1.2	36.38	±0.03	0.85
TCUB-3	<i>Tichosina cubensis</i>	16°20'N - 60°57'W	250	17.3	±1.2	36.38	±0.03	0.85
TPLI-2	<i>Tichosina cf. plicata</i>	16°21'N - 60°54'W	111-162	23.6±2	±1.0	36.8±0.1	±0.15	0.92
TPLI-5	<i>Tichosina cf. plicata</i>	16°21'N - 60°54'W	111-162	23.6±2	±1.0	36.8±0.1	±0.15	0.92
TDES-G1	<i>Tichosina sp.</i>	16°21'N - 60°54'W	111-162	23.6±2	±1.0	36.8±0.1	±0.15	0.92
TDES-G3	<i>Tichosina sp.</i>	16°21'N - 60°54'W	111-162	23.6±2	±1.0	36.8±0.1	±0.15	0.92
TDES-B2	<i>Tichosina cf. cubensis</i>	16°21'N - 60°54'W	111-162	23.6±2	±1.0	36.8±0.1	±0.15	0.92
TDES-B4	<i>Tichosina cf. cubensis</i>	16°21'N - 60°54'W	111-162	23.6±2	±1.0	36.8±0.1	±0.15	0.92
TLAT-5	<i>Terebratulina latifrons</i> ¹	16°21'N - 60°54'W	111-162	23.6±2	±1.0	36.8±0.1	±0.15	0.92

Table 1 (continued)

Specimen	Taxa	Living location	Depth (m)	MAT (°C)	Seasonal T variation (°C)	Salinity	Seasonal salinity	$\delta^{18}\text{O}_{\text{sw}}$ (‰ VSMOW)
WB5 ¹	<i>Terbratella sanguinea</i>	New Zealand	12-20	13.6	+3.3; -1.7	34.70		0.33
WB6 ¹	<i>Liothyrella neozelanica</i>	45°20.86'S - 167°02.86'E	20-30	13.2	±2.5	34.70		0.33
WB8	<i>Notosaria nigricans</i>	45°21'36"S - 170°50'24"E	20	10.4	±2.0	34.34	±0.05	0.16
WB9A ²	<i>Calloria incospicua</i>	43°34'27"S - 172°40'07"E	20	14.0	±4	33.40	±0.90	-0.26
WB4A	<i>Aerothyris kerguelenensis</i>	Crozet Islands	200	3.9	±0.6	33.97	±0.02	-0.30
WB4B	<i>Aerothyris kerguelenensis</i>	45°57'36"S - 50°03'24"E	212-230	3.7	±0.6	34.01	±0.03	-0.29
AKER-52	<i>Aerothyris kerguelenensis</i>	46°06'0"S - 50°38'18"E	355	3.1	±0.3	34.16	±0.01	-0.25
AKER-66	<i>Aerothyris kerguelenensis</i>	45°48'06"S - 49°45'45"E	325	2.8	±0.6	34.16	±0.01	-0.25
AKER-12	<i>Aerothyris kerguelenensis</i>	46°40'00"S - 51°40'30"E	290-305	3.2	±0.2	34.11	±0.01	-0.26
AKER-73	<i>Aerothyris kerguelenensis</i>	46°07'24"S - 50°46'18"E	207-215	3.3	±0.8	34.00	±0.02	-0.29
AKER-68	<i>Aerothyris kerguelenensis</i>	46°28'30"S - 51°35'00"E	200	3.4	±0.8	33.99	±0.03	-0.29
AKER-79	<i>Aerothyris kerguelenensis</i>	46°32'54"S - 51°47'00"E	140	4.2	±0.6	33.90	±0.01	-0.31
AKER-61	<i>Aerothyris kerguelenensis</i>	45°51'24"S - 50°44'00"E	105	4.1	±1.2	33.85	±0.01	-0.33
WB1A	<i>Magellania fragilis</i>	Antarctica	862-875	-1.9	+0.3	34.70	-	-0.13
MFRA-CEA	<i>Magellania fragilis</i>	66°38'S - 143°05'E	862-875	-1.9	+0.3	34.70	-	-0.13
LUA-PAL ³	<i>Liothyrella uva</i>	67°34'S - 68°08' W	10-30	-1.1	±0.4	33.30	±0.20	-0.65
MCRA-SKA ⁴	<i>Macandrevia cranium</i>	Norway	40-100	7	±1	33.5	±0.5	-0.56
MAF-5	<i>Macandrevia africana</i>	Offshore Angola	3431	2.5	-	34.90	-	0.05

removal of the outer layers, however, the umbo, muscle scars and edges were removed from the bulk shell. Fragments of these more fragile shells were ground to a fine powder in an agate mortar.

2.3 Carbon and oxygen stable isotopes $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ of modern brachiopod calcite

Stable isotope compositions of the sampled powders were determined using an auto sampler MultiprepTM coupled to a dual-inlet and a GV Isoprime® mass spectrometer. For each sample, an aliquot of about 400 µg of calcium carbonate was reacted with anhydrous oversaturated phosphoric acid at 90 °C during 20 min. Oxygen isotope ratios of calcium carbonate were computed assuming an acid fractionation factor $1000\ln\alpha(\text{CO}_2\text{--CaCO}_3)$ of 8.1 between carbon dioxide and calcite (Swart et al., 1991). All sample measurements were duplicated and adjusted to the international references NIST NBS18 ($\delta^{18}\text{O}_{\text{VPDB}} = -23.2\text{‰}$; $\delta^{13}\text{C}_{\text{VPDB}} = -5.01\text{‰}$) and NBS19 ($\delta^{18}\text{O}_{\text{VPDB}} = -2.20\text{‰}$; $\delta^{13}\text{C}_{\text{VPDB}} = +1.95\text{‰}$), and an internal standard of Carrara marble ($\delta^{18}\text{O}_{\text{VPDB}} = -1.84\text{‰}$; $\delta^{13}\text{C}_{\text{VPDB}} = +2.03\text{‰}$). Since 2019 reproducibility of the Carrara Marble in-house standard has been $\pm 0.089096\text{‰}$ for $\delta^{18}\text{O}$ (2σ , $n=6681062$) and $\pm 0.064066\text{‰}$ for $\delta^{13}\text{C}$ (2σ , $n=6681062$). These stable isotope data are completed with $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ data obtained from clumped isotope measurements for which the analytical procedures are reported in detail in section 2.5.

2.4 Elemental ratios

Elemental concentrations were obtained by dissolving 2 to 20 mg of carbonate powder in 10 ml HNO_3 (2%). For all samples, pairs of aliquots were prepared with 10 times and 100 times dilutions depending on the considered trace and major elements along with a fixed amount of Sc and In that were added to correct concentrations from instrument drift. Solutions were analysed using an Inductively Coupled Plasma-Optical Emission Spectrometer (iCAP 7000 ICP-OES) and a quadrupole ICP-Mass Spectrometer (i-CAP-Q ICP-MS), for minor and trace elements, respectively. The 100x diluted aliquots were used to calculate Ca concentration, while 10x diluted aliquots served for Mg, Sr, Na and Li concentrations. The calculated element/Ca ratios were thus calculated and are reported in mmol/mol for Mg/Ca, Sr/Ca and Na/Ca, and in µmol/mol for Li/Ca. The reproducibility of measurements was assessed through the analysis of the carbonate standard CCH1 (Roelandts and Duchesne, 1988).

2.5 Clumped isotopes

Carbonate samples were converted to CO_2 by phosphoric acid reaction at 90 °C in a common, stirred acid bath for 15 minutes. Initial phosphoric acid concentration was 103 % (1.91 g/cm^3) and each batch of acid was used for 7 days. After cryogenic removal of water, the evolved CO_2 was helium-flushed at 25 ml/min through a purification column packed with Porapak Q (50/80 mesh, 1 m length, 2.1 mm ID) and held at -20°C , then quantitatively recollected by cryogenic trapping and transferred into an Isoprime 100TM dual-inlet mass spectrometer equipped with six Faraday collectors (m/z 44–49). Each analysis took about 2.5 hours, during which analyte gas and working reference gas were allowed to flow from matching, 10 ml reservoirs into the source through deactivated fused silica capillaries (65 cm length, 110 µm ID). Every 20 minutes, gas pressures were adjusted to achieve $m/z = 44$ current of 80 nA, with differences between analyte gas and working gas generally below 0.1 nA. Pressure-

225 dependent background current corrections were measured 12 times for each analysis. All background measurements from a given session were then used to determine a mass-specific relationship linking background intensity (Zm), total $m/z = 44$ intensity (I_{44}), and time (t):

$$Zm = aI_{44} + P(t), \text{ with } P \text{ being a polynomial of degree 2 to 4.}$$

230 Background-corrected ion current ratios (δ_{45} to δ_{49}) were converted to $\delta^{13}\text{C}$, $\delta^{18}\text{O}$, and “raw” Δ_{47} values as described by Daëron et al., (2016), using the IUPAC oxygen-17 correction parameters. The isotopic composition ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) of our working reference gas was computed based on the nominal isotopic composition of carbonate standard ETH-3 (Bernasconi et al., 2018) and an oxygen-18 acid fractionation factor of 1.00813 (Kim et al., 2007). Raw Δ_{47} values were then converted to the I-CDES Δ_{47} reference frame by comparison with four “ETH” carbonate standards (ETH 1-4, Bernasconi et al., 2021) using a pooled regression approach (Daëron, 2021). Full analytical errors are derived from the external reproducibility of unknowns and standards ($N_f = 89$) and conservatively account for the uncertainties in raw Δ_{47} measurements as well as those associated with the conversion to the “absolute” Δ_{47} reference frame.

3. Results

240 All geochemical results obtained from modern brachiopods ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$, Δ_{47} , Mg/Ca , Sr/Ca , Na/Ca and Li/Ca) are reported in Table 2. Linear regression models of $\Delta^{18}\text{O}_{\text{c-w}}$, Δ_{47} , Mg/Ca , Sr/Ca , Na/Ca and Li/Ca with Mean Annual Temperature (MAT) are reported in Figure 1 and Table 3.

3.1 Carbon and oxygen stable isotopes ~~Stable isotopes~~

245 The whole dataset of $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values from modern brachiopod shells ranges from -2.24‰ to 3.472‰ and from -1.22‰ to 34.980‰ , $n=73$, respectively. There is a robust negative linear correlation ($R^2 > 0.73$, $p\text{-slope} < 0.001$) between $\Delta^{18}\text{O}_{\text{c-w}}$ ($= \delta^{18}\text{O}_{\text{c}} - \delta^{18}\text{O}_{\text{w}}$) and MAT (Figure 1, Table 3). The correlation observed in bulk shell samples is indistinguishable from that in inner layer samples (Figure 1A). On the contrary, the correlation obtained from outer layer samples shows a significant offset with both inner layers and bulk layer samples. This is in line with the differences between the outer and inner layers observed within the same specimen, which are in most cases above analytical uncertainties and range from -3.72 to -0.192‰ (mean $= -1.41 \pm 0.40\text{‰}$; 2σ ; $n=17$) and -0.788 to 0.545‰ (mean $= -0.475 \pm 0.283\text{‰}$; 2σ ; $n=17$) for $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, respectively.

Our new set of data from inner and bulk layers' samples provides the following oxygen isotope fractionation equation:

255 ~~$$T = -5.0 (\pm 0.2) (\delta^{18}\text{O}_{\text{c}} - \delta^{18}\text{O}_{\text{sw}}) + 19.4 (\pm 0.4) \quad T = -5.2 (\pm 0.4) \Delta^{18}\text{O}_{\text{c-w}} + 19.9 (\pm 0.8) \quad R^2 = 0.95 \quad (n = 53)$$~~

with T the temperature in $^{\circ}\text{C}$ and $\Delta^{18}\text{O}_{\text{c-w}}$ ($= \delta^{18}\text{O}_{\text{c}} - \delta^{18}\text{O}_{\text{w}}$) the oxygen isotope fractionation between brachiopod calcite and seawater, excluding data from the outer shell layers. As opposed to the ordinary least square regression used in Figure 1 and Table 3, this equation is established using a York regression (York et al., 2004) using seasonal

Table 2: Geochemical data of modern brachiopod samples. Stable isotope values and element/Ca ratios.

* Samples with the anterior part of the shell.

Specimen	Shell layer	$\delta^{13}\text{C}$ (‰ VPDB)	$\delta^{18}\text{O}_c$ (‰ VPDB)	Δ_{47} (‰ I-CDES)	Mg/Ca (mmol/mol)	Sr/Ca (mmol/mol)	Li/Ca ($\mu\text{mol/mol}$)
New Caledonia							
FNEO-N4	Bulk	<u>1.91-87</u>	<u>1.31-29</u>				
FNEO-N4	Bulk	<u>1.91-93</u>	<u>1.31-32</u>		13.35	1.06	30.21
FNEO-M2	Bulk*	<u>1.91-93</u>	<u>2.12-06</u>				
FNEO-M2	Bulk*	<u>2.01-97</u>	<u>2.12-05</u>		24.90	1.36	33.47
FSAN-3	Bulk*	<u>1.51-47</u>	<u>-1.2-1.20</u>				
FSAN-3	Bulk*	<u>1.41-42</u>	<u>-1.2-1.22</u>		14.65	1.08	22.45
SCRO-1	Inner	<u>2.72-72</u>	<u>2.02-04</u>	0.6387			
SCRO-3	Outer	<u>2.32-34</u>	<u>2.02-03</u>		13.21	1.07	28.90
SCRO-3	Inner	<u>3.23-17</u>	<u>2.12-13</u>		3.76	0.45	4.09
SGLO-S1	Outer	<u>2.02-04</u>	<u>0.20-20</u>		11.15	1.02	25.53
SGLO-S1	Inner	<u>3.13-06</u>	<u>0.40-44</u>		6.68	0.60	5.21
SGLO-M1	Inner	<u>2.92-87</u>	<u>1.11-10</u>		6.70	0.55	6.01
SGLO-M1	Outer	<u>2.01-98</u>	<u>0.90-91</u>		10.72	0.94	26.91
SGLO-M1	Inner	<u>3.02-99</u>	<u>1.11-12</u>				
SGLO-M2	Outer	<u>1.91-87</u>	<u>0.90-91</u>	0.6388			
SGLO-M2	Inner	<u>2.82-80</u>	<u>0.90-88</u>	0.6210			
Guadeloupe							
TGAL-3	Bulk	<u>2.01-97</u>	<u>1.21-22</u>	0.6312			
TGAL-4	Outer	<u>1.51-46</u>	<u>1.11-08</u>		15.56	1.12	30.04
TGAL-4	Inner	<u>2.72-67</u>	<u>1.31-27</u>		6.87	0.53	3.89
TGAL-4	Inner	<u>2.82-75</u>	<u>1.41-44</u>	0.6113			
TCUB-2	Bulk	<u>1.81-75</u>	<u>1.11-08</u>	0.6248			
TCUB-3	Outer	<u>1.51-45</u>	<u>1.00-98</u>		8.25	0.85	18.77
TCUB-3	Inner	<u>2.32-31</u>	<u>1.21-24</u>		4.87	0.50	3.86
TCUB-3	Inner	<u>2.32-25</u>	<u>1.31-25</u>				
TPLI-2	Inner	<u>2.82-75</u>	<u>0.10-14</u>	0.5974			
TPLI-5	Inner	<u>3.02-96</u>	<u>0.50-45</u>		9.82	0.63	5.29
TDES-G1	Inner	<u>2.72-68</u>	<u>0.20-18</u>	0.6049			
TDES-G3	Inner	<u>2.82-76</u>	<u>0.30-31</u>				
TDES-G3	Inner	<u>2.62-61</u>	<u>0.20-23</u>		3.82	0.56	3.28
TDES-B2	Inner	<u>2.62-58</u>	<u>0.30-26</u>	0.6086			
TDES-B4	Inner	<u>2.62-56</u>	<u>0.40-36</u>				
TDES-B4	Inner	<u>2.72-71</u>	<u>0.50-47</u>		6.96	0.58	4.64
TLAT-5	Bulk*	<u>1.11-07</u>	<u>0.20-19</u>				
TLAT-5	Bulk*	<u>1.01-04</u>	<u>0.20-21</u>		25.30	1.22	28.60
New Zealand							
WB5	Outer	<u>-1.5-1.54</u>	<u>-0.6-0.63</u>		6.69	1.46	40.84
WB5	Inner	<u>0.50-52</u>	<u>0.50-47</u>	0.6639	4.28	1.13	38.39
WB6	Outer	<u>1.91-93</u>	<u>1.11-05</u>		12.50	1.43	35.07
WB6	Inner	<u>2.92-91</u>	<u>1.61-55</u>	0.6295	11.14	0.82	15.92
WB8	Inner	<u>2.12-10</u>	<u>1.31-25</u>	0.6500	11.48	1.24	46.68

265	WB9A	Outer	<u>0.40</u> 43	-0.5 <u>-0.53</u>			
	WB9A	Inner	<u>2.42</u> 35	0.90 <u>85</u>	5.73	1.04	34.38


Table 2 (continued)

Specimen	Shell layer	$\delta^{13}\text{C}$ (‰ VPDB)	$\delta^{18}\text{O}_\text{c}$ (‰ VPDB)	Δ_{47} (‰ I-CDES)	Mg/Ca (mmol/mol)	Sr/Ca (mmol/mol)	Li/Ca ($\mu\text{mol/mol}$)
Crozet Islands							
WB4A	Outer	1.11-1.11	2.02-0.04	0.6664	5.59	1.18	49.33
WB4A	Inner	2.52-5.0	3.02-9.6		5.13	1.00	41.10
WB4B	Outer	1.11-1.13	2.22-2.22		5.57	1.20	51.41
WB4B	Inner	1.91-8.6	2.82-7.9	0.6752	5.08	1.01	40.80
AKER-52	Outer	1.41-3.9	2.82-7.8	0.6670	8.46	1.18	48.79
AKER-52	Inner	2.02-0.3	3.43-3.7				
AKER-52	Inner	2.02-0.4	3.23-2.3		7.47	0.94	44.70
AKER-66	Outer	2.32-2.7	3.23-1.8		9.94	1.37	48.01
AKER-66	Inner	2.52-4.6	2.72-7.3		3.57	0.60	16.27
AKER-12	Outer	0.90-8.8	2.62-5.7		7.46	1.25	51.65
AKER-12	Inner	1.91-9.2	2.52-5.1		3.92	0.89	38.68
AKER-73	Outer	0.60-5.8	2.01-9.6		9.75	1.26	46.63
AKER-73	Inner	2.01-9.7	2.42-3.5		6.80	1.08	46.41
AKER-68	Outer	-1.1-1.06	1.31-3.4	0.6811	5.53	1.37	51.73
AKER-68	Inner	1.61-5.7	2.62-5.7				
AKER-68	Inner	1.21-2.3	2.02-0.2		4.92	1.06	46.71
AKER-79	Outer	-1.1-1.08	1.21-1.5		6.57	1.33	49.67
AKER-79	Inner	0.50-5.4	1.31-3.3				
AKER-61	Outer	-2.2-2.24	0.30-3.1		6.09	1.55	58.47
AKER-61	Inner	1.51-4.8	2.12-0.9	0.6869	4.94	1.06	48.80
AKER-61	Inner	1.41-3.6	2.32-2.6				
Antarctica							
MFRA-CEA	Outer	-0.5-0.46	2.72-7.3	0.6921	5.39	1.37	56.75
MFRA-CEA	Inner	1.01-0.3	3.73-6.8		6.84	1.13	55.35
MFRA-CEA	Inner	1.01-0.1	4.03-9.8				
WB1A	Bulk	0.80-8.4	3.63-5.7	0.6941	5.70	1.21	36.69
LUVA-PAL	Inner	0.50-5.3	2.92-9.1		13.47	1.56	54.82
LUVA-PAL	Inner	0.60-6.3	3.33-3.3				
LUVA-PAL	Outer	-1.7-1.69	1.91-8.7		20.17	2.05	56.02
Norway							
MCRA-SKA	Outer	-1.3-1.32	0.50-5.4		7.28	1.24	53.14
MCRA-SKA	Inner	0.20-2.4	1.31-2.8				
Offshore Angola							
MAF-5	Bulk	1.71-6.7	3.63-5.6		12.54	1.01	54.74
MAF-5	Bulk	1.61-5.7	3.53-4.8				

3.2 Trace element/Ca ratios

The Mg/Ca, Sr/Ca, Li/Ca and Na/Ca ratios of modern brachiopod shells range from 3.57 to 25.30 mmol/mol, 0.45 to 2.05 mmol/mol, 3.28 to 58.47 $\mu\text{mol/mol}$, and 1.86 to 18.28 mmol/mol, respectively (N=47). Sr/Ca show significant but weak negative correlations ($0.28 < R^2 < 0.44$; p-slope < 0.01) with temperature. Li/Ca show significant and good negative correlations ($0.61 < R^2 < 0.86$; p-slope < 0.001) with temperature, the strongest correlation ($R^2=0.86$) being reached in outer layers samples. Na/Ca show significant negative correlations ($0.38 < R^2 < 0.60$; p-slope < 0.001) with temperature. Concentration of these three elements in brachiopod shells (Sr, Li

275 and Na) are strongly positively correlated with each other ($0.66 < R^2 < 0.93$ for the whole dataset) especially while considering only



280 **Figure 1:** Geochemical data and linear regression model for each geochemical parameter tested for its relationship with brachiopod living temperature. Parameters associated with each regression model are listed in table 3.

inner layers samples ($0.84 < R^2 < 0.96$). No significant correlation ($p\text{-slope} > 0.2$) of Mg/Ca with temperature is observed in the dataset (Figure 1; Table 3). Measured element/Ca ratios are systematically higher in the outer layers than in the inner layers of brachiopod shells (Student-t-test: $p < 0.01$; Figure 1C–E).

285

Table 3: Linear regression model parameters for the geochemical parameter-temperature relationship tested and displayed in Figure 1.

Regression model	Sample/subsample	Nb of points	Slope	Intercept	σ_{slope}	$\sigma_{\text{intercept}}$	σ_{residual}	P-slope	R^2
MAT (°C) vs $\Delta^{18}\text{O}$ (‰)	All data	74	-5.1	18.7	0.2	0.5	3.0	<0.001	0.87
	Outer layers	21	-4.4	15.2	0.6	1.3	3.7	<0.001	0.71
	Inner layers	40	-5.4	20.2	0.2	0.5	2.1	<0.001	0.94
	Bulk sample	13	-4.9	19.3	0.2	0.5	1.6	<0.001	0.97
	Bulk + inner layers	53	-5.2	19.9	0.2	0.4	2.0	<0.001	0.95
Δ_{47} (‰ I-CDES) vs MAT (°C)	All data	20	-0.0036	0.688	0.0003	0.005	0.011	<0.001	0.86
	Inner layers	17	-0.0036	0.688	0.0004	0.005	0.013	<0.001	0.86
Mg/Ca (mmol/mol) vs MAT (°C)	All data	47	0.13	7.7	0.09	1.1	5.0	0.16	0.04
	Outer layers	18	0.14	8.3	0.14	1.5	4.0	0.36	0.05
	Inner layers	23	-0.0025	6.6	0.07	0.9	2.7	0.94	<0.01
Sr/Ca (mmol/mol) vs MAT (°C)	All data	47	-0.021	1.27	0.005	0.06	0.3	<0.001	0.28
	Outer layers	18	-0.022	1.45	0.007	0.08	0.2	0.008	0.36
	Inner layers	23	-0.024	1.13	0.006	0.08	0.2	<0.001	0.44
Li/Ca ($\mu\text{mol/mol}$) vs MAT (°C)	All data	47	-1.7	51.1	0.2	2.5	11	<0.001	0.61
	Outer layers	18	-1.7	55.6	0.2	1.7	5	<0.001	0.86
	Inner layers	23	-2.1	49.5	0.3	4.1	12	<0.001	0.67
Na/Ca (mmol/mol) vs MAT (°C)	All data	47	-0.32	12.7	0.06	0.7	3	<0.001	0.38
	Outer layers	18	-0.31	14.4	0.06	0.6	2	<0.001	0.60
	Inner layers	23	-0.38	11.2	0.08	1.0	3	<0.001	0.53

3.3 Clumped isotopes

290 Δ_{47} values obtained from the selected brachiopod shells range from 0.59 ± 0.01 to 0.69 ± 0.01 ‰ I-CDES and are strongly correlated with ambient temperature (Table 3, Figure 1B). Temperatures inferred from clumped isotope data were calculated using the INTERCARB calibration determined by Anderson et al., (2021) who reprocessed the data coming from various laboratories using the same carbonate standards for data correction. Clumped isotope temperatures are significantly lower than estimated MAT with a mean deviation from environmental temperatures of $-2.4 \pm 1.3^\circ\text{C}$ (95% CI; $n=19$).

295

4. Interpretation and discussionDiscussion

4.1 Brachiopod shell paleothermometers.

4.1.1 Validity and robustness of our $^{18}\text{O}/^{16}\text{O}$ -based fractionation equation

The oxygen isotope fractionation equation derived from our data deviates substantially from that also derived from articulated brachiopod shells determined by Brand et al. (2019) (Figure 2). The two equations follow similar trends and are only offset by $\sim 0.6\text{‰}$ at tropical temperatures ($20\text{--}30^\circ\text{C}$) (Figure 2A) with the equation from Brand et al. (2019) predicting a lower fractionation factor. At temperate and polar temperatures (20 to 0°C) our new equation highlights a higher sensibility of oxygen fractionation to temperature than that of Brand et al. (2019) (Figure 2). ~~our equation has a steeper slope than that of Brand et al. (2019).~~ The two equations overlap at around 10°C and deviate substantially around $\sim 0^\circ\text{C}$ when approaching water freezing temperatures ($\sim 0^\circ\text{C}$), with a $\Delta^{18}\text{O}_{\text{c-w}}$ difference of up to $\sim 1\text{‰}$ between the equation of Brand et al. (2019) and our equation (Figure 2 A), a pattern also observed in the geochemical datasets (Figure 2B). Examining the geochemical dataset revealed that the large offset between the two equations at low temperature is partly resulting from the use of a second-order regression model by Brand et al. (2019) as the regression curve overlaps with the highest $\Delta^{18}\text{O}_{\text{c-w}}$ data points at temperatures of $\sim 1^\circ\text{C}$. We preferred to use a linear regression more appropriate to fit data in the low temperature ($<50^\circ\text{C}$) linear domain of the hyperbolic curve ($\alpha = f(1/T^2)$) predicted by equilibrium thermodynamics. Still the main discrepancies between the equation proposed herein and that of Brand et al., (2019) ~~is~~ are mainly driven by a difference in the geochemical dataset. The case of the polar brachiopods *Liothyrella uva* from Rothera island (Antarctica) highlight the likely reasons for this discrepancy at low temperatures. Indeed, $\delta^{18}\text{O}$ values reported in both studies are the same (Brand et al., 2019), indicating that the discrepancy lies in the environmental parameters used to derive the equation. Brand et al. (2019) used published environmental parameters measured locally in late February for this site, while we approached this estimation using regional annual averages. This difference in integration time of the environmental parameters is important, because the February month records the annual maximum and minimum in seawater temperature and $\delta^{18}\text{O}_{\text{sw}}$ values respectively (Meredith et al., 2013). This particular brachiopod species has a higher growth rate during the Austral winter than during the Austral summer (Peck et al., 1997), further increasing the likely seasonal bias. These discrepancies are individually relatively small ($\pm 1\text{--}2^\circ\text{C}$; $\pm 0.5\text{‰}$ VSMOW) but put together easily explain the 1‰ offset between both equations in the low temperature range. Those differences may arise from a few reasons, 1) different approaches to determine environmental parameters 2), differences in brachiopod shell preparation and sampling or 3) major differences in the geographical and taxonomical distribution of the brachiopod dataset.




Figure 2: Plot comparison of A) regression lines for different temperature-oxygen isotope fractionation for brachiopod calcite (This study, Brand et al. 2019), laboratory precipitated calcite (Kim and O'Neil, 1997) and slow growing cave calcite (Daëron et al., 2019) and B) corresponding datasets for brachiopod calcite (modified after Brand et al., 2019). Uncertainty envelope 95% CI of the York least square regression (York et al., 2004).

We use a published independent [dataset of](#) modern brachiopod $\delta^{18}\text{O}$ [dataset values](#) (Bajnai et al., 2018) to test our equation against other previously published equations in their ability to predict environmental temperatures when $\delta^{18}\text{O}_{\text{sw}}$ is known (Figure 3). This dataset [has a MAT range of 0 to 29°C and uses](#) ~~proposes~~ two sets of $\delta^{18}\text{O}_{\text{sw}}$ for most data. One derived from direct measurements and one derived from $\delta^{18}\text{O}_{\text{sw}}$ -Salinity relationships (LeGrande and Schmidt, 2006). Plus, Mg/Ca values are available for all samples, allowing us to apply the equation of Brand et al. (2013) to the dataset. Isotopic temperatures were calculated then normalised as the temperature deviation from environmental temperature (ΔT in °C) with full propagation of uncertainties. Of all the equations tested, the equations derived from measurements of modern brachiopods (Brand et al., 2013, 2019; this study) yield the most accurate temperature predictions. Mean ΔT are statistically indistinguishable from 0 using salinity-based $\delta^{18}\text{O}_{\text{sw}}$ estimates result for all three equations.




Figure 3: Offsets between the $\delta^{18}\text{O}$ temperatures and MAT from the modern brachiopods dataset of Bajnai et al., (2018) using different fractionation equations. Circles: temperature offset for each samples; vertical transparent bars: temperature offset 95% confidence interval for each samples; horizontal bar: temperature offset arithmetic mean and associated 95% confidence interval. A) Temperature offset for isotopic temperatures calculated with $\delta^{18}\text{O}_{\text{sw}}$ values calculated from the LeGrande and Schmidt (2006) database; N=18. B) Temperature offset for isotopic temperatures calculated with locally measured $\delta^{18}\text{O}_{\text{sw}}$ values; N=13. OLS = ordinary least square regression; YLS = York least square regression (York et al., 2004).

Interestingly, the accuracy of isotopic temperature estimates is poorer when measured $\delta^{18}\text{O}_{\text{sw}}$ values are used. Interestingly, while using locally measured $\delta^{18}\text{O}_{\text{sw}}$ values, the accuracy of isotopic temperature estimates is poorer. The equations of Brand et al. (2013; 2019) now significantly underestimate growing temperatures (Figure 3, $p_{(\text{mean } \Delta T = 0)} < 0.01$) while our equation yields mean ΔT that are statistically indistinguishable from 0 ($p_{(\text{mean } \Delta T = 0)} = 0.11$). The equation of Kim and O'Neil (1997), derived from synthetic carbonates, and that of Epstein et al. (1953), derived from molluscs, both underestimate brachiopod shells growing temperature (Figure 3, $p_{(\text{mean } \Delta T = 0)} < 0.001$) in both $\delta^{18}\text{O}_{\text{sw}}$ hypotheses. On the other end of the spectrum, the equation of Daëron et al. (2019), derived from slow growing calcite, overestimates brachiopod shells growing temperatures (Figure 3, $p_{(\text{mean } \Delta T = 0)} < 0.001$) in both $\delta^{18}\text{O}_{\text{sw}}$ hypotheses. Although it lessens the accuracy of $\delta^{18}\text{O}$ temperatures in this dataset, using the use of locally measured $\delta^{18}\text{O}_{\text{sw}}$ values over the salinity-based estimates improved their precision ($4.0 < 2\sigma < 5.3$ °C, n=13 rather than $5.4 < 2\sigma < 6.4$ °C, n=18). Those two $\delta^{18}\text{O}_{\text{sw}}$ hypotheses regarding $\delta^{18}\text{O}_{\text{sw}}$ values have their own benefits and drawbacks. $\delta^{18}\text{O}_{\text{sw}}$ estimated from salinity measurements and the regional $\delta^{18}\text{O}_{\text{sw}}$ -Salinity relationships offers $\delta^{18}\text{O}_{\text{sw}}$ estimates that can be averaged through time but this approach ignores coastal deviations

from the $\delta^{18}\text{O}_{\text{sw}}$ -salinity relationship. Locally measured $\delta^{18}\text{O}_{\text{sw}}$ correct for those effects, however those measurements are often scarce in time can be can be biased at the seasonal or inter annual scale for brachiopods living above the base of the thermocline. For example, Brand et al. (2019) used $\delta^{18}\text{O}_{\text{sw}}$ values measured in late February (Austral summer) to estimate the isotope fractionation of *Liothyrella uva* from Rothera island, while long term $\delta^{18}\text{O}_{\text{sw}}$ measurements at this station indicate minimal values during the austral summer (Meredith et al., 2013). Independent from the fractionation equation or the $\delta^{18}\text{O}_{\text{sw}}$ value used, temperature estimates are associated with large scatter of more than 4°C (2 σ) around mean ΔT . We suggest that this scatter is related primarily to the variability of brachiopod shell $\delta^{18}\text{O}$ values observed within a population (Figure 2, Brand et al., 2019) due to other isotopic fractionation mechanisms than temperature dependent fractionation, namely kinetic effects, metabolic effects and pH effects. Using the MgCO_3 -corrected equation of Brand et al. (2013) instead of Brand et al. (2019), both derived from very similar datasets, has the only effect of slightly improving the precision of the estimates decreasing the scatter by $\sim 0.6^\circ\text{C}$ around the mean deviation from environmental temperatures. This effect is about one order of magnitude below the remaining dispersal of data. For the different equations derived from modern brachiopod data, we push further the comparison of ΔT between isotopic temperature and environmental temperature, looking at their distribution against environmental temperature (Figure 4). This, highlights a significant correlation between ΔT and sweater temperature when applying both Brand et al. (2013) and Brand et al. (2019) equations to the comparative dataset of Bajnai et al. (2018) ($p_{\text{uncorrelated}} < 0.001$; Figure 4). On contrary, there is no significant correlation between ΔT and sweater temperature when applying both our new equation to the comparative dataset of Bajnai et al. (2018) ($p_{\text{uncorrelated}} > 0.3$). This comparison highlights that the application of the equations of Brand et al. (2013, 2019) do not completely account for temperature as an explicative variable for brachiopod $\Delta^{18}\text{O}_{\text{c-w}}$ in the dataset of Bajnai et al. (2018).




Figure 4: Distribution of isotopic temperature deviation from living temperature against living temperature using the following equations A) this study, B) Brand et al. (2013), C) Brand et al. (2019) to the dataset of Bajnai et al. (2018).

In the following, we further assess the impact of trace element incorporation in brachiopod calcite on the fractionation factor between calcite and water using our new data. Brachiopod calcite can be viewed as a three component molar mixture of CaCO_3 , MgCO_3 and SrCO_3 , and that the fractionation factor of this mixture ($\Delta_{(\text{brach.calcite-water})}$) of divalent metal carbonates is equivalent to the sum of the mineral-water fractionation factors weighted from the mineral molar fractions. This bulk isotopic fractionation factor is thus expressed as follows:

$$\Delta_{(\text{brach.calcite-water})} = \sum_{i=1}^n X_i \cdot \Delta_{(\text{mineral-water})}$$

With the n ($=3$) minerals noted i and X the molar fraction of each mineral constituting the brachiopod calcite mixture. As to our knowledge MgCO_3 -water oxygen isotope fractionation equation has not yet been established from synthetic carbonates, we used the theoretical fractionation factors ($\alpha_{(x-y)}$) of Chacko and Deines (2008) to compare the mixture and pure calcite fractionation factors. $\Delta_{(\text{brach.calcite-water})}$ calculated for brachiopod inner calcite layers with the same range of CaCO_3 , MgCO_3 and SrCO_3 contents ~~similar to that~~ recorded by our elemental data, do not exceed 0.18‰ compared to the calcite end-member, at environmental temperatures (Mean = 0.08‰; $N=24$). This difference is mainly explained by Mg^{2+} incorporation as 1) MgCO_3 is more abundant than SrCO_3 in brachiopod calcite and 2) $1000\ln \alpha_{(\text{MgCO}_3\text{-water})}$ is higher than $1000\ln \alpha_{(\text{CaCO}_3\text{-water})}$ by about 13‰ in this temperature range whereas $1000\ln \alpha_{(\text{SrCO}_3\text{-water})}$ is lower than $1000\ln \alpha_{(\text{CaCO}_3\text{-water})}$ by only ~1‰ (Chacko and Deines, 2008). This basic calculation illustrates that Mg^{2+} incorporation into brachiopod low Mg-calcite can affect calcite $\delta^{18}\text{O}$

by no more than 0.2‰, which is about twice the range of analytical uncertainties at 2σ. ~~In As a~~ consequence, the ~~incorporation effect~~ of Mg²⁺ has a weak effect on the δ¹⁸O of brachiopod calcite, the MgCO₃ correction proposed by Brand et al., (2013) is not really needed for paleoclimatic and paleoceanographic studies, which most likely justifies why it was later abandoned (Brand et al., 2019).

4.1.2 Brachiopod shells clumped-isotopes: an alternative paleothermometer?

Although there is a strong correlation between our new Δ₄₇ values from brachiopod shells and growing temperatures, Δ₄₇ values at a given temperature are generally higher than what is expected from the state-of-the-art equation of Anderson et al., (2021) (Figure 54). This observation is in line with the results of Bajnai et al., (2018) gathered in the CDES reference frame. To discuss modern brachiopod Δ₄₇ values on a larger dataset, the data from Bajnai et al., (2018) were adjusted to the CDES 90 reference frame to better compare them with our data set in the I-CDES reference frame (Bernasconi et al., 2021). We acknowledge that the different standardisation protocols used for the two datasets could hamper the strength of this comparison. However, this comparison is supported by the good agreement between sample 143 from Bajnai et al., (2018) (Δ₄₇ = 0.670 ± 0.008 ‰ CDES 90) and its replicate sample Mv143b (Δ₄₇ = 0.664 ± 0.006 ‰ I-CDES; Bajnai et al., 2020; Fiebig et al., 2021). ~~Both datasets are first compared to the Anderson et al. (2021) calibration (Figure 4). Many brachiopod shells record higher Δ₄₇ values than what is expected from the calibration of Anderson et al., (2021) (Figure 4A-B), in line with the underestimation Δ₄₇ temperatures relative to MAT when using this relationship (Figure 4C). The temperature Δ₄₇ relationship derived from the Bajnai et al. (2018) dataset appears significantly different from that of Anderson et al. (2021) equation, the latter being indistinguishable from the relationship derived from our new dataset (Figure 4A). We argue that the difference between the two datasets of modern brachiopod Δ₄₇ may largely results from distinct taxonomic assemblages and/or sampled locations in both datasets. Indeed, our new Δ₄₇ data from New Zealand brachiopods replicate very well with previous data from same brachiopod species and similar locations (Bajnai et al., 2018) (see supplementary). The Δ₄₇ temperature relationship derived using both brachiopod datasets, is still nearly within error of the equation of Anderson et al., (2021) (Figure 4B). The use of this latter equation to calculate brachiopod growing temperature results in a significant mean offset with environmental temperatures of $-3.9 \pm 1.2^{\circ}\text{C}$ (95% CI, Figure 4C) for the combined brachiopod dataset. In consequence, based on the data available to date, we support the use the following brachiopod equation: The Δ₄₇-temperature relationship derived from this combined brachiopod dataset yields the following equation:~~

$$\Delta_{47(I-CDES90^{\circ}\text{C})} = 0.042 \pm 0.002 \times \frac{10^6}{T^2} + 0.124 \pm 0.016$$

~~with, Δ₄₇ (I-CDES90°C) in ‰, and T the temperature in K. This equation while still nearly within error of the equation of Anderson et al. (2021) (Figure 5B), highlights systematically higher brachiopod Δ₄₇ values relative to what is expected from the canonical clumped isotope equation. We thus conclude that, at a given temperature brachiopod shell Δ₄₇ values are higher that what has been observed among foraminifera (Breitenbach et al., 2018; Peral et al., 2018; Meinicke et al., 2020), laboratory precipitated calcites (Jautzy et al., 2020; Anderson et al., 2021; Fiebig et al., 2021), slow growing cave calcites (Daëron et al., 2019; Anderson et al., 2021; Fiebig et al., 2021) and bivalve calcite (Huyghe et al., 2022). This observation is more relevant in the Bajnai et al. (2018) dataset with mean Δ₄₇~~

temperature deviation from seawater temperatures of $-5.6 \pm 1.6^{\circ}\text{C}$ (2 SE, N=18), than in our new dataset with mean Δ_{47} temperature deviation from seawater temperatures of $-2.4 \pm 1.3^{\circ}\text{C}$ (2 SE, N=19) (Figure 5A). However, we argue that the difference between the two datasets of modern brachiopod Δ_{47} may largely results from distinct taxonomic assemblages and/or sampled locations in both datasets. Indeed, our new Δ_{47} data from New Zealand brachiopods replicate very well with previous data from same brachiopod species and similar locations (Bajnai et al., 2018) (see supplement). The Bajnai et al. (2018) dataset is dominated by brachiopods from the suborder Terebratulidina (12/18), a group for which the combined dataset shows the most deviation from the Anderson et al. (2021) equation, at low (-2 to 5°C) but most importantly in the mid (8 - 15°C) temperature range (Figure 5C). This group is associated with a mean Δ_{47} temperature deviation from seawater temperatures of $-5.5 \pm 1.6^{\circ}\text{C}$ (2 SE, N=20), when applying the Anderson et al., (2021) equation, while the other most represented group in the combined dataset, the suborder Terebratulidina, better fits the canonical equation (Figure 5C), with mean Δ_{47} temperature deviation from seawater temperatures of $-1.7 \pm 1.5^{\circ}\text{C}$ (2 SE, N=12). This apparent taxonomic difference among brachiopod groups suggests that the deviation observed for brachiopod shell Δ_{47} values relative to other calcifying organisms, relate to processes other than ambient temperatures, controlling bound ordering within the calcite crystal of brachiopod shells (Bajnai et al., 2018). Such processes still need to be confidently identified (Section 4.2), but likely lie in different ways the shell is secreted under different growth rate dynamic and/or different crystal arrangement. Available brachiopod Δ_{47} values suggest that, while the equation of Anderson et al. (2021) could be confidently applied to brachiopod of the suborder Terebratulidina (Figure 5), this would not be the case for species of the suborder Terebratulidina. For the other brachiopod groups, data are still too scarce to propose an enlightened conclusion.

$$\Delta_{47(T=\text{CDTS}90^{\circ}\text{C})} = -0.042 \pm 0.002 \times \frac{10^6}{T^2} + 0.128 \pm 0.027$$

with, $\Delta_{47(T=\text{CDTS}90^{\circ}\text{C})}$ in ‰, and T the temperature in K.




Figure 45: Comparison of Δ_{47} values from modern brachiopods with the state-of-the-art Δ_{47} calibration of Anderson et al., (2021). Envelope and error bars at 2σ . A. Comparison of York least square

regressions (York et al., 2004) derived from two modern brachiopod datasets (Bajnai et al., 2018; this study) with the equation of Anderson et al., (2021). B. Comparison of York least square regression derived from the combined modern brachiopod dataset with the equation of Anderson et al., (2021). C) Comparison of modern brachiopods Δ_{47} temperatures calculated using the equation of Anderson et al., (2021) with environmental-seawater temperatures.

4.1.3 Mg/Ca thermometer for brachiopod shells

Our Mg/Ca data obtained from the inner layers of brachiopod shells do not show any significant correlation with growth temperature ($p=0.94$; Figure 1C, Table 3) ~~and odds with Brand et al. (2013, 2019) proposed that there is~~ a the strong correlation between the Mg/Ca molar ratio of brachiopod shell secondary layer and the growth temperature found by Brand et al. (2013, 2019). However, ~~they these authors~~ acknowledged that some species do not follow ~~that the~~ trend they illustrated as the GBMgL. ~~They, and~~ suggested that such deviations resulted from either lower growth rate (Brand et al., 2013) or higher ones (Brand et al., 2019) of the considered species relative to other brachiopods. This issue alone illustrates how from the largest dataset of modern brachiopods available so far (i.e. Brand's database), one cannot firmly determine whether or not Mg/Ca correlates with seawater temperatures. We emphasize that our dataset lack brachiopod shells samples from equatorial surface marine waters ($27^{\circ}\text{C} < T < 32^{\circ}\text{C}$), which record the highest MgCO_3 molar content in Brand's database (Brand et al., 2013, 2019). Interestingly, the highest MgCO_3 contents reported by these authors were measured on brachiopods belonging to the order Thecideida. This taxonomic singularity is of prime importance because: 1) Thecideide brachiopod shells are essentially made of a shell fabric unique to its group (Simonet Roda et al., 2022) primary shell fabric, and secondary shell fabric (fibrous layer, which form the majority of Rhynchonellida and Terebratulida shells) being is restricted to small areas of the shell such as the teeth and the inner sockets (Williams, 1968, 1973); 2) the primary shell layer has higher Mg/Ca ratios than the fibrous secondary layer in Rhynchonellid and Terebratulid brachiopods (Pérez Huerta et al., 2008; Gaspard et al., 2018; Rollion Bard et al., 2019); 3) Thecideide shell is formed of high Mg calcite Ullmann et al., (2017). ~~assigned the high Mg/Ca ratios of Thecideida brachiopod shells to the presence of high Mg calcite.~~ The exclusion of Thecideida data from the data base published by Brand et al. (2013), produces a substantially weaker ($R^2=0.23$) yet significant ($p < 0.01$) correlation between shell Mg/Ca and temperature. In addition, magnesium has been shown to be also more concentrated in major growth bands (Gaspard et al., 2018; Müller et al., 2022) than in other parts of the secondary shell layer in articulated brachiopods, which could explain the high Mg/Ca values observed from 'slow' growth and long lifespan brachiopods (Brand et al., 2019), especially when considering samples obtained from a large sampling area covering multiple growth lines. Given our new data and all these considerations, the brachiopod Mg/Ca should be regarded as an unreliable proxy for annual or seasonal temperature reconstructions, at least in the -2 to 25°C range (Figure 1, Brand et al., 2013, 2019).

An alternative or complementary way to explain the scattering of temperature-Mg/Ca paired data may rise in the variability of marine Mg/Ca. All the elements considered in this study, (Mg, Sr, Li, Na), have much larger residence times (10 Ma, 3.5 Ma and 1.8 Ma, respectively) than water molecules in the oceans, meaning that at first-order they are uniformly distributed in the main bodies of the oceans (Lécuyer, 2016). A recent study, however, has shown sizable variations in element/Ca may occur in the vicinity of continental margins, under the

influence of oceanic currents, at the interface with groundwaters or the atmosphere, or within water masses isolated from the global thermohaline circulation (e.g. Black Sea, Mediterranean Sea, Red Sea) (Lebrato et al., 2020). For example, despite the large residence time of Sr, nonhomogeneous marine strontium isotopic ratios ($^{87}\text{Sr}/^{86}\text{Sr}$) have been measured in coastal environments (Southern Okinawa Trough, South China Sea and Kao-ping Canyon) as the result of seawater mixing with groundwaters (Huang et al., 2011; El Meknassi et al., 2020). Given the poor constraint on local coastal water chemistry in the deep time, brachiopod samples from offshore environments should be ~~privileged-preferred~~ for trace-element-based paleotemperatures reconstructions, as it was suggested for marine biogenic apatite by (Balter and Lécuyer, 2010).

4.1.4 Potential of other trace element/Ca proxy

Our new, brachiopod Sr/Ca data show a significant ~~but~~ weak negative correlation with ambient temperature ($p_{\text{slope}} < 0.001$; $R^2 < 0.40$). This correlation is supported by a weak positive correlation between Sr/Ca and $\Delta^{18}\text{O}_{\text{c-w}}$ of the inner layers ($p_{\text{slope}} < 0.01$; $R^2 = 0.31$) as we also expect $\Delta^{18}\text{O}$ to decrease with increasing temperatures (i.e section 4.1). ~~However, it~~ This contradicts the pioneering findings of Lowenstam, (1961), which suggested a negative correlation between brachiopod shell Sr/Ca and $\Delta^{18}\text{O}_{\text{c-w}}$. The brachiopod database of Brand et al., (2013) shows no significant correlation between Sr/Ca and temperatures ($p_{\text{slope}} = 0.20$). Given the diverging conclusions from one dataset to the other, Sr/Ca values of brachiopod shells do not appear to be controlled by growing temperatures, at the first order.

Li/Ca in brachiopod shells is significantly correlated to temperature (Figure 1E; Table 3), a covariation initially reported by Delaney et al. (1989) and confirmed by more recent studies (Dellinger et al., 2018; Rollion-Bard et al., 2019; Washington et al., 2020). Most of the data presented here follow the Li/Ca-temperature trend derived from literature data (Washington et al., 2020), with ~~a few several~~ samples ($n=9$), exclusively collected from the inner layer, showing significantly lower Li/Ca. Those low Li/Ca values lie below the Li/Ca-temperature relationship derived from inorganic calcite (Marriott et al., 2004). Similarly low Li/Ca values were reported from brachiopod shells by previous studies (Delaney et al., 1989; Washington et al., 2020). Interestingly, most of those low Li/Ca values belong to samples from the genus *Tichosina* (Delaney et al., 1989; Washington et al., 2020, Table 2), so one could consider some peculiar species-dependent trace element partitioning driven by metabolic or kinetic processes. Caution is warranted, however, because very low Li/Ca values are also apparent in the ~~other tropical~~ genus ~~studied here~~, *Stenosarina* Cooper 1977, as well as one specimen of *Aerothyris kerguelensis* among the 9 specimens studied ~~there~~.

Other environmental parameters, ~~or biological processes~~ may control the element/Ca incorporation into calcite. Sr/Ca, Li/Ca and Na/Ca show strong significant negative correlations with the local salinity ($0.37 < R^2 < 0.76$; $p_{\text{slope}} < 0.01$) with similar scattering as for the temperature relationships. Mg/Ca again show no significant correlation with this parameter ($p_{\text{slope}}=0.7$). A wider look at the geochemical dataset highlights very strong correlations of Sr/Ca, Li/Ca, and Na/Ca ratios in brachiopod shells between one another ($0.68 < R^2 < 0.92$; $p_{\text{slope}} < 0.01$) over the whole dataset, and particularly from inner layer samples ($0.87 < R^2 < 0.97$; $p_{\text{slope}} < 0.01$). Correlation in our dataset is strongest between Sr/Ca and Na/Ca and weakest between Sr/Ca and Li/Ca. Similar correlation trends also appear at the intra-specimen level (Romanin et al., 2018; Rollion-Bard et al., 2019). Such

strong correlations, at both intra individual and inter specific scales, may suggest that the first order variability of those different element concentrations within brachiopod shells may be explained by similar processes, either environmental or biological. Ullmann et al. (2017) reported distinct geochemical signature between brachiopods species especially within the Mg/Ca-Sr/Ca space. Our inner shell layer dataset presents a similar distribution within the low Mg/Ca range (<10 mmol/mol), where Sr/Ca ratios of Terebratulidina shells are distinctly lower than those of Terebratellidina shells. Owing to the strong correlation between Sr/Ca, Li/Ca and Na/Ca, our dataset differentiate Terebratulidina shells with “low” element/Ca ($0.4 < \text{Sr/Ca} < 0.7$ mmol/mol; $0 < \text{Li/Ca} < 10$ $\mu\text{mol/mol}$; $1 < \text{Na/Ca} < 4$ mmol/mol), and Terebratellidina shells with “high” element/Ca ($0.9 < \text{Sr/Ca} < 1.2$ mmol/mol; $35 < \text{Li/Ca} < 60$ $\mu\text{mol/mol}$; $8 < \text{Na/Ca} < 12$ mmol/mol), with the exception of the Antarctic brachiopod *Liothyrella uva* (Broderip, 1833, Terebratulidina) which shows the highest Sr/Ca, Na/Ca and Li/Ca values (Figure 6 A-B). With this exception and that reported by Ullmann et al. (2017), a taxonomic relationship is not satisfactory, especially considering the significant yet noisy relationship between element/Ca and environmental parameters such as temperature and salinity within our new dataset. Indeed, we also observe a strong relationship between brachiopod shell element/Ca values and geographical location (Figure 6 C-D). Tropical shells (Guadeloupe and New Caledonia; only Terebratulidina) have low element/Ca values, the mid latitudes (New Zealand and Crozet Islands, dominated by Terebratellidina) have intermediate to high element/Ca values, and polar brachiopods (Antarctica and Norway; mixed assemblage) have high element/Ca values. Despite this first order distribution, it appears that within the same environment, species difference appears to dominate, as is the case in Doubtful Sound (New Zealand), where *Terebratella sanguinea* (Leach, 1814) (Terebratellidina) shell have higher element/Ca values than *Liothyrella neozelanica* Thomson 1918 (Terebratulidina) shell. Excluding Mg, element/Ca ratios plotted against living depth revealed two different trends below ~100m depth: a group of “low” element/Ca ($0.4 < \text{Sr/Ca} < 0.7$ mmol/mol; $0 < \text{Li/Ca} < 10$ $\mu\text{mol/mol}$; $1 < \text{Na/Ca} < 4$ mmol/mol) and a group of “high” element/Ca brachiopods ($0.9 < \text{Sr/Ca} < 1.2$ mmol/mol; $35 < \text{Li/Ca} < 60$ $\mu\text{mol/mol}$; $8 < \text{Na/Ca} < 12$ mmol/mol). Such important differences cannot be explained by variations in water element/Ca ratios, especially at this depth in the water column (Lebrato et al., 2020). This strong dichotomy, may result either from other environmental parameters not considered in this study (e.g., carbonate chemistry), or from a strong taxonomic control. Indeed, almost all specimens belonging to the “low” element/Ca group lived in tropical areas (Guadeloupe, New Caledonia), while all brachiopods belonging to the “high” element/Ca lived at higher latitudes (New Zealand, Crozet archipelago, Antarctica). Specimens from New Zealand display the largest variability in element/Ca ratios, probably partly because all samples are from coastal environments, where water element/Ca are expected to vary the most (Lebrato et al., 2020). All but one of the brachiopods studied here belong to the order Terebratulida which regrouped the suborders Terebratulidina and Terebratellidina. All but one Terebratulidine brachiopod shells are associated with “low” element/Ca ratios, while all but one Terebratellidine brachiopod shells are associated with “high” element/Ca ratios; in line with trends in Sr/Ca ratios reported by Ullmann et al., (2017). Whether there is an environmental or a taxonomic control on trace element incorporation in articulated brachiopods remains unclear but data suggest there may be both. Higher element/Ca ratios were measured from *Terebratella sanguinea* (Leach, 1814) (Terebratellidina) than from *Liothyrella neozelanica* Thomson 1918 (Terebratulidina), while living in the same environment. Similarly, the Antarctic brachiopod *Liothyrella uva* (Broderip, 1833) shows the highest Sr/Ca, Na/Ca and Li/Ca ratios, while belonging to the Terebratulidine. It is worthy to note that in the dataset presented here, tropical samples are dominated by Terebratulidine brachiopods, while high latitude samples are dominated by Terebratellidine

brachiopods, which preclude confident testing of both hypotheses from this dataset. In the fossil record, differences in Mg/Ca and Sr/Ca ratios between brachiopod genera have been associated with different shell microstructures (Ullmann et al., 2020).




Figure 6: Distribution of trace element content of the inner shell layers among the different brachiopod taxonomic groups (Same symbols as in Figure 5C). The data show a strong correlation between Sr/Ca and Li/Ca (A) and Sr/Ca and Na/Ca (B), as well as strong latitudinal partitioning of Li/Ca (C) and Sr/Ca (D) values.

Our new data do not support the use of the trace element/Ca ratios studied here as paleothermometers from a dataset of widely distributed and taxonomically diverse brachiopods. In other words, suggest that temperature is not the prime parameter that can describe trace element incorporation into articulated brachiopods shell calcite, from a dataset of widely distributed and taxonomically diverse brachiopods from various environments and belonging to various taxa, suggesting the major influence of other environmental or taxonomic parameters. This conclusion does not exclude the possibility of any a strong temperature control on trace element incorporation in brachiopod shell calcite, which may be relevant at the species level as proposed by Butler et al., (2015), and observed in other calcifying organisms such as bivalves (Freitas et al., 2006) and foraminifera (Anand et al., 2003). But the influence of other control on trace element incorporation in brachiopod shell calcite, (growth rate, environment fluid chemistry; Jurikova et al., 2020) is strong enough to blur the likely temperature control, so that the trace element/Ca ratios studied here are not fitted to be used as paleothermometers.

4.2 ~~Precipitation of brachiopod shell calcite out of equilibrium with seawater.~~ precipitation of brachiopod shell calcite with seawater.

Our new data confirm that brachiopod shell calcite does not precipitate in isotopic equilibrium with seawater, as highlighted by the significant deviation from thermodynamic isotopic equilibrium relationships constrained from very slow growing calcite (Daëron et al., 2019; Anderson et al., 2021, Figure 2-5). ~~The oxygen isotope fractionation equation presented here differs from other published equations including that of Kim and O'Neil (1997). However, it~~ It is worthy to note that most brachiopod shell data (Brand et al., 2019, this study) plot in-between thermodynamic equilibrium isotopic fractionation (Daëron et al., 2019) and the equation of Kim and O'Neil (1997) (Figure 32). Deviation of brachiopod shell calcite $\delta^{18}\text{O}$ from thermodynamic equilibrium ($\Delta^{18}\text{O}_{\text{eq}} = \delta^{18}\text{O}_{\text{brach. calcite}} - \delta^{18}\text{O}_{\text{equilibrium}}$) is obvious when we compare our data set to the equation of Daëron et al. (2019) in the clumped isotope-oxygen isotope fractionation space (Figure 5A7A) with most of our dataset showing higher Δ_{47} and lower $\delta^{18}\text{O}$ than predicted (Figure 5B7B), a feature previously highlighted by Bajnai et al., (2018). As a result, brachiopod calcite shows apparent Δ_{47} and $\delta^{18}\text{O}$ temperatures respectively lower and higher than those predicted by the thermodynamic equilibrium. This deviation from equilibrium values has a Δ_{47} - $\Delta^{18}\text{O}_{\text{eq}}$ slope of 0.019 ± 0.005 in our new brachiopod dataset, indistinguishable from the slope 0.017 ± 0.003 reported by Bajnai et al (2018) using $\delta^{18}\text{O}_{\text{sw}}$ values from the $\delta^{18}\text{O}_{\text{sw}}$ database (LeGrande and Schmidt, 2006).

Brachiopod shell $\delta^{18}\text{O}$ data (Brand et al., 2003, 2013, 2019; this study) also highlight-show a large geochemical variability among brachiopods sampled from similar marine environment (T,S), with a $\delta^{18}\text{O}$ range often higher than 1‰ (Figure 3, Brand et al., 2019). Isotopic fractionation equations (Brand et al., 2013, 2019; this study) usually rely on $\Delta^{18}\text{O}_{\text{c-w}}$ values plotted against mean annual temperatures. Thus, it would be tempting to attribute such scattering of data to seasonal variations, in both temperature and $\delta^{18}\text{O}_{\text{sw}}$, but ~~only in the case of this would only apply to~~ specimens that lived above or in the upper part of the thermocline. Such isotopic scattering is indeed expected if we consider datasets based on multiple sampling along the growth axis of an individual (Yamamoto et al., 2011; Takayanagi et al., 2015), such as in the case of Brand et al. (2013; 2019). However, seasonal scattering in data resulting from intra individual geochemical variability is unlikely for our measurements performed on a large sampling area that should correspond to multiple months or years of growth.

For example, we note a ~~great-large~~ isotopic variability in both $\delta^{18}\text{O}$ and Δ_{47} among the 9 specimens of *Aerothyris kerguelenensis* from Crozet islands even though the specimens were recovered from depths in the 105 to 355 m range with a narrow range of temperature (3-5°C) and salinity (33.8-34.2). These 9 samples, all collected from the shell inner layers from different individuals, records a $\delta^{18}\text{O}$ variability of 2‰, equivalent to a temperature range of 8 to 10°C if interpreted as such, and Δ_{47} variability of 0.02 ‰ equivalent to a temperature range of about 5°C. This subsample shows a significant Δ_{47} - $\Delta^{18}\text{O}_{\text{eq}}$ slope of 0.027 ± 0.007 (2σ ; $p_{\text{slope}} < 0.01$; $R^2=0.95$, $n=5$), as well as a strong positive correlation between $\Delta^{18}\text{O}_{\text{c-w}}$ and $\delta^{13}\text{C}$ ($p_{\text{slope}} < 0.01$; $R^2=0.69$, $n=9$). These covariations most likely result from kinetic effects as defined by McConnaughey (1989) here recorded at the intra-specific level.




Figure 57: Non equilibrium fractionation and kinetic effects in brachiopod shell calcite as evidenced by clumped isotopes and oxygen isotopes. A) Plot of brachiopod inner shell layers in the Δ_{47} - $1000 \ln \alpha_{(\text{calcite-water})}$ space compared with the equilibrium line (blue line) defined by very slow growing abiogenic calcite (Daëron et al., 2019; Anderson et al., 2021). Red highlight, results from *A. kerguelensis* (Crozet Islands) and their expected geochemical values on the equilibrium line. Note the negative correlation between Δ_{47} and $1000 \ln \alpha_{(\text{calcite-water})}$ evidenced in this subset. B) $\delta^{18}\text{O}_c$ and Δ_{47} offset from their expected equilibrium values calculated from environmental parameters using the equations of Daëron et al. (2019) and Anderson et al. (2021), respectively. The trends in followed by these offsets within our new dataset, is similar to what has been that reported from other brachiopod samples by Bajnai et al. (2018).

Our new data replicate the observations made by Bajnai et al. (2018) concerning kinetic fractionation in brachiopod shells. In consequence, we follow their conclusions and consider that different growth rates of the specimens are responsible for the second-order isotopic variability and the deviation from stable isotope and clumped isotope equilibrium observed in brachiopod datasets, both at an inter-specific and intra-specific level.

Minor and trace elements are neither incorporated into brachiopod shell calcite in equilibrium with seawater, relative to abiogenic calcite (Rollion-Bard et al., 2019; Simonet Roda et al., 2022) Trace element incorporation into calcite can also be affected by kinetic effects. Kinetic effects are known to influence trace element incorporation into calcite (Lorens, 1981; Busenberg and Plummer, 1985; Tesoriero and Pankow, 1996; Gabitov et al., 2011, 2014) and may be a process, among others (Jurikova et al., 2020), that explains trace element incorporation in brachiopod shell calcite. Owing to the very limited knowledge of brachiopod shell growth rates (only 5 out of 17 species in our dataset), we are unable to directly test this hypothesis. Consequently However, if kinetic effects have a strong such influence on element incorporation into brachiopod calcite, we should also observe strong negative correlations of Sr/Ca, Na/Ca and Li/Ca ratios with $\Delta^{18}\text{O}_{\text{eq}}$. We observe such significant correlations in the new dataset, in both the inner and outer layers' subsets ($0.40 < R^2 < 0.55$, p-slope < 0.01), with a significant slope change between the two subsets. The variation in element/Ca related to a similar variation in $\Delta^{18}\text{O}_{\text{eq}}$ as suggested

by linear regression models, is about twice as high in the inner layers' subset than in the outer layers' subset. This difference may be explained by the different pathways brachiopods use to form their different shell layers (acicular, columnar, fibrous ; Simonet Roda et al., 2022). However, while looking at *Aerothyris kerguelensis*, which showed the most important kinetic effects according to stable isotope data, there is no significant correlation of element/Ca ratios with $\Delta^{18}\text{O}_{\text{eq}}$. Element/Ca ratios appear to be independent from isotope kinetic effects at the species level, despite strong correlations of element/Ca ratios and $\Delta^{18}\text{O}_{\text{eq}}$ in the multi-specific dataset.

In line with the significant correlation between $\Delta^{18}\text{O}_{\text{eq}}$ and element/Ca values in brachiopod shell inner layer, the environmental and taxonomic distribution of isotopic offsets ($\Delta^{18}\text{O}_{\text{eq}}$, Δ_{47} offset) in our dataset, is similar to the distribution of element/Ca values. We observe more deviation from the equilibrium in high latitude environments than in tropical environments, and we observe generally more deviation from isotopic equilibrium in Terebratellidina than in Terebratulidina in the whole dataset, although the difference is not as striking as documented for trace elements. As the geographical distribution of the two suborders in the new dataset, is highly biased, this trend cannot be confidently interpreted to raise from different shell formation processes in the different groups.
~~The deviation from equilibrium isotopic fractionation observed in the new dataset seems to be linked to either taxonomic or environmental factors. As observed when comparing the new data to the equation of Daëron et al. (2019) (Figure 3), deviations from equilibrium oxygen isotope fractionation are greater at lower temperatures, and more generally we observe more deviation from the equilibrium in high latitude environments than in tropical environments. Around Crozet islands, the deeper the brachiopod lived the less deviation from equilibrium is recorded, both in oxygen isotopes and clumped isotopes. While we observe generally more deviation from isotopic equilibrium in Terebratellidina than in Terebratulidina in the whole dataset, the difference is not as striking as documented for trace elements. Given the highly geographically biased distribution of the two suborders in the new dataset, this trend may be coincidental.~~ However, several examples of Terebratellidina and Terebratulidina living in the same environment show similar pattern. In Doubtful Sound (New Zealand) *Terebratella sanguinea* (Terebratellidina) deviates more from equilibrium isotopic composition than *Liothyrella neozelanica* (Terebratulidina) (This study, Bajnai et al., 2018). In Friday Harbor (Washington, USA), *Terebratalia transversa* (Sowerby, 1846) (Terebratellidina) records lower and more variable $\delta^{18}\text{O}_c$ than *Terebratulina sp.* (Terebratulidina) (Ullmann et al., 2017). These authors also reported high isotopic variability associated with strong correlation of $\delta^{18}\text{O}$ with $\delta^{13}\text{C}$ in Terebratellidina but not among Terebratulidina. ~~It would seem that brachiopod shells from the suborder Terebratellidina record more important kinetic effects (both from stable isotopes and elemental contents) than brachiopod shells from the suborder Terebratulidina.~~

From the relationship between the different parameters studied here, we suggest that kinetic effects in brachiopod shells, may be a major source of deviation from calcite precipitation in equilibrium with seawater. Kinetic effects can be on first approximation controlled by shell growth rates.
~~Kinetic effects being strongly linked to shell growth rates~~ (McConnaughey, 1989; Bajnai et al., 2018). Therefore, both taxonomic-biologic and environmental parameters should control the extent of the kinetic effects. At first order, one can expect species growing larger shells to record more important kinetic effects. Moreover, shell growth rates in calcifying organisms increase when environmental conditions are close to optimal living range and decrease when the conditions strongly derive from that optimum, a pattern being dependant on taxon-related preferences (Schöne, 2008). However, the poor knowledge of brachiopod shell growth rate, relative to other marine calcifying organisms, prevents an in-depth

appreciation of its exact role. Kinetics effects are still insufficient to describe all of the deviation from equilibrium observe here, but our dataset does not allow to discuss other potential factors (Jurikova et al., 2020).

4.3. Consequences for the application of $\delta^{18}\text{O}$ and Δ_{47} paleothermometry to brachiopod fossils.

Brachiopod shells display disequilibrium isotopic fractionation in both clumped isotopes and stable isotopes that generates both a general offset in apparent environmental parameters (T , $\delta^{18}\text{O}_{\text{sw}}$) relative to measured ones, and an increase in the apparent variability of those parameters within a single environment. This is of concern for the accuracy and precision of paleotemperature and $\delta^{18}\text{O}_{\text{sw}}$ estimates derived from the fossil record.

In this section we illustrate this issue with the study of four fossil brachiopods from the “collections de Géologie de l'Université Claude Bernard Lyon 1”, measured alongside the modern brachiopods, and using the same sampling protocol. The four studied specimens were recovered from the Lower Jurassic of Normandy (W-France), in various locations around the cities of Caen and Bayeux. They span the Pliensbachian to Toarcian stages and are well-dated with ammonite biostratigraphy. Trace elements measured on those samples revealed Mn and Fe concentrations lower than 150 ppm, and 300 ppm, respectively. Those low values support the good preservation of the fossil material (Brand and Veizer, 1980; Brand et al., 2003; Voigt et al., 2003). In addition, the Lower Jurassic successions of Normandy lies unconformably on the Variscan basement of the Armorican massif and are hence very condensed (~10 m for the whole Pliensbachian-Toarcian) (Dommergues et al., 2008; Weis et al., 2018) due to the low rates of subsidence in the area since the Triassic (Brunet and Le Pichon, 1982). Hence alteration of the Δ_{47} during burial appears unlikely (Henkes et al., 2014; Stolper and Eiler, 2015; Hemingway and Henkes, 2021). Geochemical results and their paleoenvironmental interpretation are displayed in Table 4. To correct from the geochemical offset recorded in brachiopod shells relative to equilibrium precipitation of calcite, we suggest to use brachiopod-based calibration for both Δ_{47} temperatures and ^{18}O fractionation. Here we applied the equations derived in this study to this set of fossil brachiopods. Δ_{47} values recorded in our fossil samples suggest temperatures of $26 \pm 5^\circ\text{C}$ throughout the Pliensbachian and up to $31.8 \pm 5.2^\circ\text{C}$ in the late Toarcian. While the $\delta^{18}\text{O}$ values have very good analytical precision ($\pm 0.03\text{‰}$), they rely on the analysis of only one specimen. Before interpreting the $\delta^{18}\text{O}$ data from a single brachiopod specimen, we suggest to use a larger uncertainty of $\pm 1\text{‰}$ to account for the natural variability of $\delta^{18}\text{O}$ values within a brachiopod population (this study, Brand et al., 2019). Accordingly, $\delta^{18}\text{O}_{\text{sw}}$ estimates from single brachiopod specimen are associated with uncertainties about twice as large as if only analytical uncertainties were considered. Here, Pliensbachian to Toarcian $\delta^{18}\text{O}_{\text{sw}}$ values estimated for W-France range from 0.3 ± 1.7 to $1.1 \pm 2.0\text{‰}$ VSMOW. The uncertainties associated with $^{18}\text{O}_{\text{sw}}$ estimates determine the range of temporal or spatial variations in $\delta^{18}\text{O}_{\text{sw}}$ that could be confidently identified from brachiopod fossils. Given the very large uncertainties we thus suggest to analyse multiple well-preserved specimens within a single bed (Ullmann et al., 2020), in order to characterise the geochemical distribution of the studied population, and use central values with their associated uncertainties, for paleoenvironmental interpretations (at least for $\delta^{18}\text{O}_\text{e}$). In this case, the classically used positive correlation between $\delta^{18}\text{O}_\text{e}$ and $\delta^{13}\text{C}$ (McConnaughey, 1989) and a negative correlation between $\delta^{18}\text{O}_\text{e}$ and Δ_{47} (Bajnai et al., 2018, this study) would also provide solid evidence for kinetic effects. Finally, kinetic biases may be corrected through the use of dual-clumped isotope thermometry (Bajnai et al., 2020). However, given their instrumental and temporal costs, such clumped isotope methods will require

further analytical development before they can be routinely used to detect and correct such biases in the large number of samples needed for paleoclimate studies.

Table 4: Fossil brachiopods' geochemical results and paleoenvironmental interpretations with 2 σ analytical and propagated uncertainties. Both brachiopod-specific Δ_{47} -calibration and the Anderson et al. (2021) are compared on these samples. $\delta^{18}\text{O}_{\text{sw}}$ are calculated from Δ_{47} -T and $\delta^{18}\text{O}_{\text{e}}$ values using the brachiopod-specific equation derived in this study. An additional $\sim 1\text{‰}$ may be added to $\delta^{18}\text{O}$ uncertainties and then propagated, to consider the natural dispersal of brachiopod $\delta^{18}\text{O}$ values observed within modern and fossil brachiopod populations, which is one to two orders of magnitude larger than analytical uncertainties.

Sample	FSL 707721	FSL 707727	FSL 707729	FSL 707752
Species	<i>Platyrhynchia thalla</i>	<i>Tertrarrhynchia dumbletonensis</i>	<i>Quadratirhynchia quadrata</i>	<i>Homeorhynchia cynocephala</i>
Location	Subles	Feuguerolles	Tilly-sur-Seulles	Subles
Stage	Pliensbachian	Pliensbachian	Pliensbachian	Toarcian
Ammonite Biozone	Jamesoni	Margaritatus	Spinatum	Aalensis
Geochemical results				
$\delta^{13}\text{C}$ (‰ VPDB)	1.16 ± 0.02	2.40 ± 0.02	1.49 ± 0.02	1.33 ± 0.02
$\delta^{18}\text{O}$ (‰ VPDB)	-0.55 ± 0.03	-0.24 ± 0.03	-0.15 ± 0.03	-0.93 ± 0.03
Δ_{47} (‰ ICDES)	0.600 ± 0.011	0.602 ± 0.016	0.600 ± 0.016	0.583 ± 0.016
Mn (ppm)	26	42	121	50
Fe (ppm)	39	88	282	198

Paleoenvironment interpretation (this study)

$\Delta_{47}\text{-T} (^{\circ}\text{C})$	26.1 ± 3.4	25.4 ± 4.9	26.2 ± 4.9	31.8 ± 5.1
$\delta^{18}\text{O}_{\text{sw}} (\text{‰})$ VSMOW)	0.92 ± 0.68	1.09 ± 0.97	1.33 ± 0.98	1.63 ± 1.03

Paleoenvironment interpretation (Anderson et al., 2021)

$\Delta_{47}\text{-T} (^{\circ}\text{C})$	22.9 ± 3.5	21.1 ± 5.1	22.9 ± 5.1	28.9 ± 5.5
$\delta^{18}\text{O}_{\text{sw}} (\text{‰})$ VSMOW)	0.29 ± 0.71	0.46 ± 1.01	0.70 ± 1.02	1.05 ± 1.08

5. Conclusions

Our multi-proxy ($\delta^{18}\text{O}$ and $\delta^{13}\text{C}$, Δ_{47} , Mg/Ca, Sr/Ca, Na/Ca and Li/Ca) geochemical analyses of 37 modern brachiopods from different locations, spanning living temperatures from -1.9°C to 25.5°C allowed us to investigate the temperature relationship of several geochemical markers, to evaluate or reevaluate their potential as paleothermometers for brachiopod shell archives. $\delta^{18}\text{O}$ and Δ_{47} are the most reliable paleothermometers, although we propose a revised $\delta^{18}\text{O}$ paleotemperature equation for brachiopod shell calcite and Δ_{47} values highlight significant deviation from the canonical temperature relationship. The element/Ca ratios investigated here should not be used to infer environmental temperatures from brachiopod calcite, owing to either the absence (Mg/Ca) or a poor yet significant (Sr/Ca, Li/Ca, Na/Ca) correlation with living temperature. Our new dataset confirms previous evidence for non-equilibrium isotopic fractionation interpreted as significant kinetic effects recorded in the brachiopod shell inner layers. In general, the precipitation of brachiopod shells taking place out of isotopic equilibrium leads to 1) an under-estimation of living temperature from Δ_{47} and 2) an over-estimation of living temperature from oxygen isotopes, which once the proxies are combined could lead to 3) an underestimation of living water $\delta^{18}\text{O}_{\text{sw}}$, relative to what would be expected from equilibrium precipitation. Kinetic effects and by extension shell growth rates could explain the second order of isotopic variability in brachiopod shells and can be a larger source of isotopic variability than environmental parameters (Temperature, $\delta^{18}\text{O}_{\text{sw}}$) at the local scale. Thus, we advise in considering this variability as a significant source of uncertainties, when applying paleotemperature proxies. Although, we reject their use as a paleothermometer, we do not exclude any temperature control on trace element (Mg, Sr, Li, Na) incorporation into brachiopod shell calcite. Rather, our findings indicate that other processes, such as kinetic effects (growth rate) or shell microstructure and their associated mineralisation pathways, have a significant control on trace element incorporation into brachiopod shell calcite, resulting in the high scrambling observed in the element/Ca-temperature relationship. confirm that the brachiopod shell inner layers should be preferred to estimate water properties. There is a strong correlation between the living temperature and the $\Delta^{18}\text{O}$ fractionation between the brachiopod shell inner layers and water that allowed us to establish a new $\delta^{18}\text{O}$ paleotemperature equation for brachiopod shell calcite. Our new trace element data show no significant correlation between growing temperature and Mg/Ca ratios measured from the inner shell layers indicating that brachiopod shell Mg/Ca is not a suitable paleotemperature proxy. Sr/Ca, Li/Ca and Na/Ca of the brachiopod shell inner layers show a significant negative correlation with growing temperatures. However, partitioning of those

elements between the low Mg calcite brachiopod shell and seawater appears to show strong taxonomic patterns. Clumped isotope temperatures, derived from the calibration of (Anderson et al., 2021) are generally underestimated by -3°C relative to the environmental temperature. In consequence, we suggest to use a calibration especially dedicated to the brachiopods for reconstruction seawater temperatures. In addition, our new Δ_{47} data confirm previous evidence for non-equilibrium isotopic fractionation interpreted as significant kinetic effects recorded in the brachiopod shell inner layers. In general, the precipitation of brachiopod shells taking place out of isotopic equilibrium leads to 1) an under-estimation of living temperature from Δ_{47} , 2) an over-estimation of living temperature from oxygen isotopes and 3) an underestimation of living $\delta^{18}\text{O}_{\text{sw}}$ when combining the two proxies, relative to what would be expected from equilibrium precipitation. Locally, the kinetic effects produce geochemical apparent temperatures with high variability relative to the local temperature range, for both the oxygen isotope and the clumped isotope proxies. We therefore advise to consider this natural variability in the interpretation of paleoenvironmental parameters from brachiopod shell $\delta^{18}\text{O}$ and Δ_{47} data.

Author contribution

TL and CL designed the study in close collaboration with GS and DG, DG obtained the samples from various contributors, TL performed shell sampling in the lab, AVL and TL performed the stable isotope analysis and data treatment, TL and FAG performed the elemental analysis and data treatment, TL and MD performed the clumped isotope analysis and data treatment, TL performed the statistical analysis of the data and wrote the first draft paper with input from all authors.

Acknowledgments

This study was founded by the ANR-OXYMORE (grant no. ANR-18-CE31-0020)

The authors gratefully thank the leaders and teams of the N/O of the following expeditions: the Walda cruise in S.E. Atlantic Ocean (Dr. M. Segonzac, IFREMER, Brest); Sampling from Norway coasts (courtesy of the Trondheim Biological station); MD 30 “BIOMASS” around Crozet Islands (Arnaud, Marseille, France and Prof N. Ameziane, MNHN, who provided the material); Musorstom 4, SMIB 3 and 6 and Norfolk 1 around New Caledonia, Loyauté - Lifou Islands (Prof. B. Laurin, Univ. Bourgogne, Dijon and Richer de Forges); Karubenthos 2, around Guadeloupe: French Caribbean Island (Prof. Ph. Bouchet with Dr. P. Lozouet, Dr. L. Corbari and Ph. Maestrati and the help of J. Mainguy, MNHN, Paris, France) with a supplementary sampling (Prof. D. Lamy, Univ. Antilles-Guyane); supplementary material from New-Zealand (Prof. Lee, D., Univ Otago, Marine Biology Lab.; Prof. D.I. MacKinnon and Dr. A. Aldridge (Christchurch Univ.); CEAMARC off French Antarctic area (Prof. N. Ameziane and Dr. M. Eléaume, MNHN, Paris, France); sampling from Palmer Peninsula, Antarctic (courtesy of Th. Desvignes, USA).

~~We thank Emmanuel Robert for granting access to fossil specimens from the “collections de Géologie de l’Université Claude Bernard Lyon 1”.~~

Bibliography

- Anand, P., Elderfield, H., and Conte, M. H.: Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series: calibration of Mg/Ca thermometry in planktonic foraminifera, *Paleoceanography*, 18, 1050, <https://doi.org/10.1029/2002PA000846>, 2003.
- 835 Anderson, N. T., Kelson, J. R., Kele, S., Daëron, M., Bonifacie, M., Horita, J., Mackey, T. J., John, C. M., Kluge, T., Petschnig, P., Jost, A. B., Huntington, K. W., Bernasconi, S. M., and Bergmann, K. D.: A Unified Clumped Isotope Thermometer Calibration (0.5–1,100°C) Using Carbonate-Based Standardization, *Geophys. Res. Lett.*, 48, <https://doi.org/10.1029/2020GL092069>, 2021.
- 840 Auclair, A.-C., Joachimski, M. M., and Lécuyer, C.: Deciphering kinetic, metabolic and environmental controls on stable isotope fractionations between seawater and the shell of *Terebratalia transversa* (Brachiopoda), *Chem. Geol.*, 202, 59–78, [https://doi.org/10.1016/S0009-2541\(03\)00233-X](https://doi.org/10.1016/S0009-2541(03)00233-X), 2003.
- Ayling, B. F., McCulloch, M. T., Gagan, M. K., Stirling, C. H., Andersen, M. B., and Blake, S. G.: Sr/Ca and $\delta^{18}\text{O}$ seasonality in a *Porites* coral from the MIS 9 (339–303 ka) interglacial, *Earth Planet. Sci. Lett.*, 248, 462–475, <https://doi.org/10.1016/j.epsl.2006.06.009>, 2006.
- 845 Bajnai, D., Fiebig, J., Tomašových, A., Milner Garcia, S., Rollion-Bard, C., Raddatz, J., Löffler, N., Primo-Ramos, C., and Brand, U.: Assessing kinetic fractionation in brachiopod calcite using clumped isotopes, *Sci. Rep.*, 8, 533, <https://doi.org/10.1038/s41598-017-17353-7>, 2018.
- 850 Bajnai, D., Guo, W., Spötl, C., Coplen, T. B., Methner, K., Löffler, N., Krsnik, E., Gischler, E., Hansen, M., Henkel, D., Price, G. D., Raddatz, J., Scholz, D., and Fiebig, J.: Dual clumped isotope thermometry resolves kinetic biases in carbonate formation temperatures, *Nat. Commun.*, 11, 4005, <https://doi.org/10.1038/s41467-020-17501-0>, 2020.
- Balter, V. and Lécuyer, C.: Determination of Sr and Ba partition coefficients between apatite from fish (*Sparus aurata*) and seawater: The influence of temperature, *Geochim. Cosmochim. Acta*, 74, 3449–3458, <https://doi.org/10.1016/j.gca.2010.03.015>, 2010.
- 855 Bergmann, K. D., Finnegan, S., Creel, R., Eiler, J. M., Hughes, N. C., Popov, L. E., and Fischer, W. W.: A paired apatite and calcite clumped isotope thermometry approach to estimating Cambro-Ordovician seawater temperatures and isotopic composition, *Geochim. Cosmochim. Acta*, 224, 18–41, <https://doi.org/10.1016/j.gca.2017.11.015>, 2018.
- 860 Bernasconi, S. M., Müller, I. A., Bergmann, K. D., Breitenbach, S. F. M., Fernandez, A., Hodell, D. A., Jaggi, M., Meckler, A. N., Millan, I., and Ziegler, M.: Reducing Uncertainties in Carbonate Clumped Isotope Analysis Through Consistent Carbonate-Based Standardization, *Geochem. Geophys. Geosystems*, 19, 2895–2914, <https://doi.org/10.1029/2017GC007385>, 2018.
- 865 Bernasconi, S. M., Daëron, M., Bergmann, K. D., Bonifacie, M., Meckler, A. N., Affek, H. P., Anderson, N., Bajnai, D., Barkan, E., Beverly, E., Blamart, D., Burgener, L., Calmels, D., Chaduteau, C., Clog, M., Davidheiser-Kroll, B., Davies, A., Dux, F., Eiler, J., Elliott, B., Fetrow, A. C., Fiebig, J., Goldberg, S., Hermoso, M., Huntington, K. W., Hyland, E., Ingalls, M., Jaggi, M., John, C. M., Jost, A. B., Katz, S., Kelson, J., Kluge, T., Kocken, I. J., Laskar, A., Leutert, T. J., Liang, D., Lucarelli, J., Mackey, T. J., Manganot, X., Meinicke, N., Modestou, S. E., Müller, I. A., Murray, S., Neary, A., Packard, N., Passey, B. H., Pelletier, E., Petersen, S., Piasecki, A., Schauer, A., Snell, K. E., Swart, P. K., Tripathi, A., Upadhyay, D., Vennemann, T., Winkelstern, I., Yarian, D., Yoshida, N., Zhang, N., and Ziegler, M.: InterCarb: A Community Effort to Improve Interlaboratory Standardization of the Carbonate Clumped Isotope Thermometer Using Carbonate Standards, *Geochem. Geophys. Geosystems*, 22, e2020GC009588, <https://doi.org/10.1029/2020GC009588>, 2021.
- 870 Billups, K. and Schrag, D. P.: Application of benthic foraminiferal Mg/Ca ratios to questions of Cenozoic climate change, *Earth Planet. Sci. Lett.*, 209, 181–195, [https://doi.org/10.1016/S0012-821X\(03\)00067-0](https://doi.org/10.1016/S0012-821X(03)00067-0), 2003.
- 875 Brand, U. and Veizer, J.: Chemical diagenesis of a multicomponent carbonate system; 1, Trace elements, *J. Sediment. Res.*, 50, 1219–1236, <https://doi.org/10.1306/212F7BB7-2B24-11D7-8648000102C1865D>, 1980.

- Brand, U., Logan, A., Hiller, N., and Richardson, J.: Geochemistry of modern brachiopods: applications and implications for oceanography and paleoceanography, *Chem. Geol.*, 198, 305–334, [https://doi.org/10.1016/S0009-2541\(03\)00032-9](https://doi.org/10.1016/S0009-2541(03)00032-9), 2003.
- 880 Brand, U., Azmy, K., Bitner, M. A., Logan, A., Zuschin, M., Came, R., and Ruggiero, E.: Oxygen isotopes and MgCO₃ in brachiopod calcite and a new paleotemperature equation, *Chem. Geol.*, 359, 23–31, <https://doi.org/10.1016/j.chemgeo.2013.09.014>, 2013.
- Brand, U., Bitner, M. A., Logan, A., Azmy, K., Crippa, G., Angiolini, L., Colin, P., Griesshaber, E., Harper, E. M., Ruggiero, E. T., and Häussermann, V.: Brachiopods-based oxygen-isotope thermometer: update and review, *Riv. Ital. Paleontol. E Stratigr.*, 125, 775–787, <https://doi.org/10.13130/2039-4942/12226>, 2019.
- 885 Breitenbach, S. F. M., Mleneck-Vautravers, M. J., Grauel, A.-L., Lo, L., Bernasconi, S. M., Müller, I. A., Rolfe, J., Gázquez, F., Greaves, M., and Hodell, D. A.: Coupled Mg/Ca and clumped isotope analyses of foraminifera provide consistent water temperatures, *Geochim. Cosmochim. Acta*, 14, 2018.
- Busenberg, E. and Plummer, L. N.: Kinetic and thermodynamic factors controlling the distribution of SO₃²⁻ and Na⁺ in calcites and selected aragonites, *Geochim. Cosmochim. Acta*, 49, 713–725, [https://doi.org/10.1016/0016-7037\(85\)90166-8](https://doi.org/10.1016/0016-7037(85)90166-8), 1985.
- 890 Butler, S., Bailey, T. R., Lear, C. H., Curry, G. B., Cherns, L., and McDonald, I.: The Mg/Ca–temperature relationship in brachiopod shells: Calibrating a potential palaeoseasonality proxy, *Chem. Geol.*, 397, 106–117, <https://doi.org/10.1016/j.chemgeo.2015.01.009>, 2015.
- 895 Came, R. E., Brand, U., and Affek, H. P.: Clumped isotope signatures in modern brachiopod carbonate, *Chem. Geol.*, 377, 20–30, <https://doi.org/10.1016/j.chemgeo.2014.04.004>, 2014.
- Carpenter, S. J. and Lohmann, K. C.: $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of modern brachiopod shells, *Geochim. Cosmochim. Acta*, 59, 3749–3764, [https://doi.org/10.1016/0016-7037\(95\)00291-7](https://doi.org/10.1016/0016-7037(95)00291-7), 1995.
- 900 Chacko, T. and Deines, P.: Theoretical calculation of oxygen isotope fractionation factors in carbonate systems, *Geochim. Cosmochim. Acta*, 72, 3642–3660, <https://doi.org/10.1016/j.gca.2008.06.001>, 2008.
- Coplen, T. B.: Calibration of the calcite–water oxygen-isotope geothermometer at Devils Hole, Nevada, a natural laboratory, *Geochim. Cosmochim. Acta*, 71, 3948–3957, <https://doi.org/10.1016/j.gca.2007.05.028>, 2007.
- 905 Cuny-Guirriec, K., Douville, E., Reynaud, S., Allemand, D., Bordier, L., Canesi, M., Mazzoli, C., Taviani, M., Canese, S., McCulloch, M., Trotter, J., Rico-Esenaro, S. D., Sanchez-Cabeza, J.-A., Ruiz-Fernández, A. C., Carricart-Ganivet, J. P., Scott, P. M., Sadekov, A., and Montagna, P.: Coral Li/Mg thermometry: Caveats and constraints, *Chem. Geol.*, 523, 162–178, <https://doi.org/10.1016/j.chemgeo.2019.03.038>, 2019.
- Cusack, M., Pérez-Huerta, A., and Eimf: Brachiopods recording seawater temperature—A matter of class or maturation?, *Chem. Geol.*, 334, 139–143, <https://doi.org/10.1016/j.chemgeo.2012.10.021>, 2012.
- 910 Daëron, M.: Full Propagation of Analytical Uncertainties in Δ_{47} Measurements, *Geochem. Geophys. Geosystems*, 22, e2020GC009592, <https://doi.org/10.1029/2020GC009592>, 2021.
- Daëron, M., Blamart, D., Peral, M., and Affek, H. P.: Absolute isotopic abundance ratios and the accuracy of Δ_{47} measurements, *Chem. Geol.*, 442, 83–96, <https://doi.org/10.1016/j.chemgeo.2016.08.014>, 2016.
- 915 Daëron, M., Drysdale, R. N., Peral, M., Huyghe, D., Blamart, D., Coplen, T. B., Lartaud, F., and Zanchetta, G.: Most Earth-surface calcites precipitate out of isotopic equilibrium, *Nat. Commun.*, 10, 429, <https://doi.org/10.1038/s41467-019-08336-5>, 2019.
- Delaney, M. L., Popp, B. N., Lepzelter, C. G., and Anderson, T. F.: Lithium-to-calcium ratios in modern, cenozoic, and paleozoic articulate brachiopod shells, *Paleoceanography*, 4, 681–691, <https://doi.org/10.1029/PA004i006p00681>, 1989.
- 920 Dellinger, M., West, A. J., Paris, G., Adkins, J. F., Pogge von Strandmann, P. A. E., Ullmann, C. V., Eagle, R. A., Freitas, P., Bagard, M.-L., Ries, J. B., Corsetti, F. A., Perez-Huerta, A., and Kampf, A. R.: The Li isotope

composition of marine biogenic carbonates: Patterns and mechanisms, *Geochim. Cosmochim. Acta*, 236, 315–335, <https://doi.org/10.1016/j.gca.2018.03.014>, 2018.

DeLong, K. L., Flannery, J. A., Maupin, C. R., Poore, R. Z., and Quinn, T. M.: A coral Sr/Ca calibration and replication study of two massive corals from the Gulf of Mexico, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 307, 117–128, <https://doi.org/10.1016/j.palaeo.2011.05.005>, 2011.

El Meknassi, S., Dera, G., De Rafélis, M., Brahmi, C., Lartaud, F., Hodel, F., Jeandel, C., Menjot, L., Mounic, S., Henry, M., Besson, P., and Chavagnac, V.: Seawater $^{87}\text{Sr}/^{86}\text{Sr}$ ratios along continental margins: Patterns and processes in open and restricted shelf domains, *Chem. Geol.*, 558, 119874, <https://doi.org/10.1016/j.chemgeo.2020.119874>, 2020.

Epstein, S., Buchsbaum, R., Lowenstam, H. A., and Urey, H. C.: Revised carbonate-water isotopic temperature scale, *Geol. Soc. Am. Bull.*, 64, 1315, [https://doi.org/10.1130/0016-7606\(1953\)64\[1315:RCITS\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1953)64[1315:RCITS]2.0.CO;2), 1953.

Fiebig, J., Daëron, M., Bernecker, M., Guo, W., Schneider, G., Boch, R., Bernasconi, S. M., Jautzy, J., and Dietzel, M.: Calibration of the dual clumped isotope thermometer for carbonates, *Geochim. Cosmochim. Acta*, 312, 235–256, <https://doi.org/10.1016/j.gca.2021.07.012>, 2021.

Freitas, P. S., Clarke, L. J., Kennedy, H., Richardson, C. A., and Abrantes, F.: Environmental and biological controls on elemental (Mg/Ca, Sr/Ca and Mn/Ca) ratios in shells of the king scallop *Pecten maximus*, *Geochim. Cosmochim. Acta*, 70, 5119–5133, <https://doi.org/10.1016/j.gca.2006.07.029>, 2006.

Gabitov, R. I., Schmitt, A. K., Rosner, M., McKeegan, K. D., Gaetani, G. A., Cohen, A. L., Watson, E. B., and Harrison, T. M.: In situ $\delta^7\text{Li}$, Li/Ca, and Mg/Ca analyses of synthetic aragonites, *Geochem. Geophys. Geosystems*, 12, Q03001, <https://doi.org/10.1029/2010GC003322>, 2011.

Gabitov, R. I., Sadekov, A., and Leinweber, A.: Crystal growth rate effect on Mg/Ca and Sr/Ca partitioning between calcite and fluid: An in situ approach, *Chem. Geol.*, 367, 70–82, <https://doi.org/10.1016/j.chemgeo.2013.12.019>, 2014.

Gaspard, D. and Nouet, J.: Hierarchical architecture of the inner layers of selected extant rhynchonelliform brachiopods, *J. Struct. Biol.*, 196, 197–205, <https://doi.org/10.1016/j.jsb.2016.07.021>, 2016.

Gaspard, D., Marie, B., Luquet, G., and Marin, F.: Biochemical BlackwellPublishingLtd characteristics of the soluble organic matrix from the shell of three Recent terebratulid brachiopod species, 2008.

Gaspard, D., Aldridge, A. E., Boudouma, O., Fialin, M., Rividi, N., and Lécuyer, C.: Analysis of growth and form in *Aerothyris kerguelensis* (rhynchonelliform brachiopod) - Shell spiral deviations, microstructure, trace element contents and stable isotope ratios, *Chem. Geol.*, 483, 474–490, <https://doi.org/10.1016/j.chemgeo.2018.03.018>, 2018.

Ghosh, P., Adkins, J., Affek, H., Balta, B., Guo, W., Schauble, E. A., Schrag, D., and Eiler, J. M.: ^{13}C – ^{18}O bonds in carbonate minerals: A new kind of paleothermometer, *Geochim. Cosmochim. Acta*, 70, 1439–1456, <https://doi.org/10.1016/j.gca.2005.11.014>, 2006.

Griesshaber, E., Kelm, K., Sehrbrock, A., Mader, W., Mutterlose, J., Brand, U., and Schmahl, W. W.: Amorphous calcium carbonate in the shell material of the brachiopod *Megerlia truncata*, *Eur. J. Mineral.*, 21, 715–723, <https://doi.org/10.1127/0935-1221/2009/0021-1950>, 2009.

Henkes, G. A., Passey, B. H., Wanamaker, A. D., Grossman, E. L., Ambrose, W. G., and Carroll, M. L.: Carbonate clumped isotope compositions of modern marine mollusk and brachiopod shells, *Geochim. Cosmochim. Acta*, 106, 307–325, <https://doi.org/10.1016/j.gca.2012.12.020>, 2013.

Henkes, G. A., Passey, B. H., Grossman, E. L., Shenton, B. J., Yancey, T. E., and Pérez-Huerta, A.: Temperature evolution and the oxygen isotope composition of Phanerozoic oceans from carbonate clumped isotope thermometry, *Earth Planet. Sci. Lett.*, 490, 40–50, <https://doi.org/10.1016/j.epsl.2018.02.001>, 2018.

- 965 Huang, K.-F., You, C.-F., Chung, C.-H., and Lin, I.-T.: Nonhomogeneous seawater Sr isotopic composition in the coastal oceans: A novel tool for tracing water masses and submarine groundwater discharge, *Geochem. Geophys. Geosystems*, 12, Q05002, <https://doi.org/10.1029/2010GC003372>, 2011.
- Huyghe, D., Daëron, M., de Rafelis, M., Blamart, D., Sébilo, M., Paulet, Y.-M., and Lartaud, F.: Clumped isotopes in modern marine bivalves, *Geochim. Cosmochim. Acta*, 316, 41–58, <https://doi.org/10.1016/j.gca.2021.09.019>, 2022.
- 970 Jautzy, J. J., Savard, M. M., Dhillon, R. S., Bernasconi, S. M., and Smirnov, A.: Clumped isotope temperature calibration for calcite: Bridging theory and experimentation, *Geochem. Perspect. Lett.*, 36–41, <https://doi.org/10.7185/geochemlet.2021>, 2020.
- 975 Jiménez-López, C., Romanek, C. S., Huertas, F. J., Ohmoto, H., and Caballero, E.: Oxygen isotope fractionation in synthetic magnesian calcite, *Geochim. Cosmochim. Acta*, 68, 3367–3377, <https://doi.org/10.1016/j.gca.2003.11.033>, 2004.
- Jurikova, H., Ippach, M., Liebetrau, V., Gutjahr, M., Krause, S., Büsse, S., Gorb, S. N., Henkel, D., Hiebenthal, C., Schmidt, M., Leipe, T., Laudien, J., and Eisenhauer, A.: Incorporation of minor and trace elements into cultured brachiopods: Implications for proxy application with new insights from a biomineralisation model, *Geochim. Cosmochim. Acta*, 286, 418–440, <https://doi.org/10.1016/j.gca.2020.07.026>, 2020.
- 980 Kim, S.-T. and O’Neil, J. R.: Equilibrium and nonequilibrium oxygen isotope effects in synthetic carbonates, *Geochim. Cosmochim. Acta*, 61, 3461–3475, [https://doi.org/10.1016/S0016-7037\(97\)00169-5](https://doi.org/10.1016/S0016-7037(97)00169-5), 1997.
- Kim, S.-T., Mucci, A., and Taylor, B. E.: Phosphoric acid fractionation factors for calcite and aragonite between 25 and 75 °C: Revisited, *Chem. Geol.*, 246, 135–146, <https://doi.org/10.1016/j.chemgeo.2007.08.005>, 2007.
- 985 Lear, C. H.: Cenozoic Deep-Sea Temperatures and Global Ice Volumes from Mg/Ca in Benthic Foraminiferal Calcite, *Science*, 287, 269–272, <https://doi.org/10.1126/science.287.5451.269>, 2000.
- 990 Lebrato, M., Garbe-Schönberg, D., Müller, M. N., Blanco-Ameijeiras, S., Feely, R. A., Lorenzoni, L., Molinero, J.-C., Bremer, K., Jones, D. O. B., Iglesias-Rodriguez, D., Greeley, D., Lamare, M. D., Paulmier, A., Graco, M., Cartes, J., Barcelos e Ramos, J., de Lara, A., Sanchez-Leal, R., Jimenez, P., Paparazzo, F. E., Hartman, S. E., Westernströer, U., Küter, M., Benavides, R., da Silva, A. F., Bell, S., Payne, C., Olafsdottir, S., Robinson, K., Jantunen, L. M., Korablev, A., Webster, R. J., Jones, E. M., Gilg, O., Bailly du Bois, P., Beldowski, J., Ashjian, C., Yahia, N. D., Twining, B., Chen, X.-G., Tseng, L.-C., Hwang, J.-S., Dahms, H.-U., and Oschlies, A.: Global variability in seawater Mg:Ca and Sr:Ca ratios in the modern ocean, *Proc. Natl. Acad. Sci.*, 117, 22281–22292, <https://doi.org/10.1073/pnas.1918943117>, 2020.
- 995 Lécuyer, C.: Seawater residence times of some elements of geochemical interest and the salinity of the oceans, *Bull. Société Géologique Fr.*, 187, 245–260, <https://doi.org/10.2113/gssgfbull.187.6.245>, 2016.
- LeGrande, A. N. and Schmidt, G. A.: Global gridded data set of the oxygen isotopic composition in seawater, *Geophys. Res. Lett.*, 33, L12604, <https://doi.org/10.1029/2006GL026011>, 2006.
- 1000 Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Boyer, T. P., Zweng, M. M., Garcia, H. E., Reagan, J. R., Seidov, D., Weathers, K., Paver, C. R., and Smolyar, I.: World Ocean Atlas 2018, Volume 1: Temperature., Mishonov Tech. Ed NOAA Atlas NESDIS, 81, 52, 2018.
- Lorens, R. B.: Sr, Cd, Mn and Co distribution coefficients in calcite as a function of calcite precipitation rate, *Geochim. Cosmochim. Acta*, 45, 553–561, [https://doi.org/10.1016/0016-7037\(81\)90188-5](https://doi.org/10.1016/0016-7037(81)90188-5), 1981.
- 1005 Lowenstam, H. A.: Mineralogy, O18/O16 Ratios, and Strontium and Magnesium Contents of Recent and Fossil Brachiopods and Their Bearing on the History of the Oceans, *J. Geol.*, 69, 241–260, <https://doi.org/10.1086/626740>, 1961.
- Marchitto, T. M., Bryan, S. P., Doss, W., McCulloch, M. T., and Montagna, P.: A simple biomineralization model to explain Li, Mg, and Sr incorporation into aragonitic foraminifera and corals, *Earth Planet. Sci. Lett.*, 481, 20–29, <https://doi.org/10.1016/j.epsl.2017.10.022>, 2018.

- 1010 Marriott, C. S., Henderson, G. M., Belshaw, N. S., and Tudhope, A. W.: Temperature dependence of $\delta^7\text{Li}$, $\delta^{44}\text{Ca}$ and Li/Ca during growth of calcium carbonate, *Earth Planet. Sci. Lett.*, 222, 615–624, <https://doi.org/10.1016/j.epsl.2004.02.031>, 2004.
- Marshall, J. F. and McCulloch, M. T.: An assessment of the Sr/Ca ratio in shallow water hermatypic corals as a proxy for sea surface temperature, *Geochim. Cosmochim. Acta*, 66, 3263–3280, [https://doi.org/10.1016/S0016-7037\(02\)00926-2](https://doi.org/10.1016/S0016-7037(02)00926-2), 2002.
- 1015 McConnaughey, T.: ^{13}C and ^{18}O isotopic disequilibrium in biological carbonates: I. Patterns, *Geochim. Cosmochim. Acta*, 53, 151–162, [https://doi.org/10.1016/0016-7037\(89\)90282-2](https://doi.org/10.1016/0016-7037(89)90282-2), 1989.
- McCulloch, M. T., Gagan, M. K., Mortimer, G. E., Chivas, A. R., and Isdale, P. J.: A high-resolution Sr/Ca and $\delta^{18}\text{O}$ coral record from the Great Barrier Reef, Australia, and the 1982–1983 El Niño, *Geochim. Cosmochim. Acta*, 58, 2747–2754, [https://doi.org/10.1016/0016-7037\(94\)90142-2](https://doi.org/10.1016/0016-7037(94)90142-2), 1994.
- 1020 Meckler, A. N., Sexton, P. F., Piasecki, A. M., Leutert, T. J., Marquardt, J., Ziegler, M., Agterhuis, T., Lourens, L. J., Rae, J. W. B., Barnet, J., Tripathi, A., and Bernasconi, S. M.: Cenozoic evolution of deep ocean temperature from clumped isotope thermometry, *Science*, 377, 86–90, <https://doi.org/10.1126/science.abk0604>, 2022.
- 1025 Meinicke, N., Ho, S. L., Hannisdal, B., Nürnberg, D., Tripathi, A., Schiebel, R., and Meckler, A. N.: A robust calibration of the clumped isotopes to temperature relationship for foraminifers, *Geochim. Cosmochim. Acta*, 270, 160–183, <https://doi.org/10.1016/j.gca.2019.11.022>, 2020.
- Meredith, M. P., Venables, H. J., Clarke, A., Ducklow, H. W., Erickson, M., Leng, M. J., Lenaerts, J. T. M., and van den Broeke, M. R.: The Freshwater System West of the Antarctic Peninsula: Spatial and Temporal Changes, *J. Clim.*, 26, 1669–1684, <https://doi.org/10.1175/JCLI-D-12-00246.1>, 2013.
- 1030 Mii, H.-S. and Grossman, E. L.: Late Pennsylvanian seasonality reflected in the ^{18}O and elemental composition of a brachiopod shell, *Geology*, 22, 661, [https://doi.org/10.1130/0091-7613\(1994\)022<0661:LPSRIT>2.3.CO;2](https://doi.org/10.1130/0091-7613(1994)022<0661:LPSRIT>2.3.CO;2), 1994.
- Miller, K. G., Browning, J. V., Schmelz, W. J., Kopp, R. E., Mountain, G. S., and Wright, J. D.: Cenozoic sea-level and cryospheric evolution from deep-sea geochemical and continental margin records, *Sci. Adv.*, 6, eaaz1346, <https://doi.org/10.1126/sciadv.aaz1346>, 2020.
- 1035 Montagna, P., McCulloch, M., Douville, E., López Correa, M., Trotter, J., Rodolfo-Metalpa, R., Dissard, D., Ferrier-Pagès, C., Frank, N., Freiwald, A., Goldstein, S., Mazzoli, C., Reynaud, S., Rüggeberg, A., Russo, S., and Taviani, M.: Li/Mg systematics in scleractinian corals: Calibration of the thermometer, *Geochim. Cosmochim. Acta*, 132, 288–310, <https://doi.org/10.1016/j.gca.2014.02.005>, 2014.
- 1040 Müller, T., Tomašových, A., Correa, M. L., Mertz-Kraus, R., and Mikuš, T.: Mapping intrashell variation in Mg/Ca of brachiopods to external growth lines: Mg enrichment corresponds to seasonal growth slowdown, *Chem. Geol.*, 593, 120758, <https://doi.org/10.1016/j.chemgeo.2022.120758>, 2022.
- Nürnberg, D.: Magnesium in tests of *Neogloboquadrina pachyderma* sinistral from high northern and southern latitudes, *J. Foraminifer. Res.*, 25, 350–368, <https://doi.org/10.2113/gsjfr.25.4.350>, 1995.
- 1045 Parkinson, D., Curry, G. B., Cusack, M., and Fallick, A. E.: Shell structure, patterns and trends of oxygen and carbon stable isotopes in modern brachiopod shells, *Chem. Geol.*, 219, 193–235, <https://doi.org/10.1016/j.chemgeo.2005.02.002>, 2005.
- Peck, L. S., Brockington, S., and Brey, T.: Growth and metabolism in the Antarctic brachiopod *Liothyrella uva*, *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 352, 851–858, <https://doi.org/10.1098/rstb.1997.0065>, 1997.
- 1050 Peral, M., Daëron, M., Blamart, D., Bassinot, F., Dewilde, F., Smialkowski, N., Isguder, G., Bonnin, J., Jorissen, F., Kissel, C., Michel, E., Vázquez Riveiros, N., and Waelbroeck, C.: Updated calibration of the clumped isotope thermometer in planktonic and benthic foraminifera, *Geochim. Cosmochim. Acta*, 239, 1–16, <https://doi.org/10.1016/j.gca.2018.07.016>, 2018.

- 1055 Pérez-Huerta, A., Cusack, M., Jeffries, T. E., and Williams, C. T.: High resolution distribution of magnesium and strontium and the evaluation of Mg/Ca thermometry in Recent brachiopod shells, *Chem. Geol.*, 247, 229–241, <https://doi.org/10.1016/j.chemgeo.2007.10.014>, 2008.
- Petersen, S. V., Tabor, C. R., Lohmann, K. C., Poulsen, C. J., Meyer, K. W., Carpenter, S. J., Erickson, J. M., Matsunaga, K. K. S., Smith, S. Y., and Sheldon, N. D.: Temperature and salinity of the Late Cretaceous Western Interior Seaway, *Geology*, 44, 903–906, <https://doi.org/10.1130/G38311.1>, 2016.
- 1060 Price, G. D., Bajnai, D., and Fiebig, J.: Carbonate clumped isotope evidence for latitudinal seawater temperature gradients and the oxygen isotope composition of Early Cretaceous seas, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 552, 109777, <https://doi.org/10.1016/j.palaeo.2020.109777>, 2020.
- Prokoph, A., Shields, G. A., and Veizer, J.: Compilation and time-series analysis of a marine carbonate $\delta^{18}\text{O}$, $\delta^{13}\text{C}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{34}\text{S}$ database through Earth history, *Earth-Sci. Rev.*, 87, 113–133, <https://doi.org/10.1016/j.earscirev.2007.12.003>, 2008.
- 1065 Roelandts, I. and Duchesne, J. C.: AWI-1 SBO-1, PRI-1, AND DWA-1, Belgian Sedimentary Rock Reference Materials, *Geostand. Geoanalytical Res.*, 12, 13–38, <https://doi.org/10.1111/j.1751-908X.1988.tb00037.x>, 1988.
- 1070 Rollion-Bard, C., Milner Garcia, S., Burckel, P., Angiolini, L., Jurikova, H., Tomašových, A., and Henkel, D.: Assessing the biomineralization processes in the shell layers of modern brachiopods from oxygen isotopic composition and elemental ratios: Implications for their use as paleoenvironmental proxies, *Chem. Geol.*, 524, 49–66, <https://doi.org/10.1016/j.chemgeo.2019.05.031>, 2019.
- Romanin, M., Crippa, G., Ye, F., Brand, U., Bitner, M. A., Gaspard, D., Häusermann, V., and Laudien, J.: A sampling strategy for recent and fossil brachiopods : selecting the optimal shell segment for geochemical analyses, *Riv. Ital. Paleontol. E Stratigr. Res. Paleontol. Stratigr.*, 124, 343–359, <https://doi.org/10.13130/2039-4942/10193>, 2018.
- 1075 Schmahl, W. W., Griesshaber, E., Kelm, K., Goetz, A., Jordan, G., Ball, A., Xu, D., Merkel, C., and Brand, U.: Hierarchical structure of marine shell biomaterials: biomechanical functionalization of calcite by brachiopods, *Z. Für Krist. - Cryst. Mater.*, 227, 793–804, <https://doi.org/10.1524/zkri.2012.1542>, 2012.
- Schöne, B. R.: The curse of physiology—challenges and opportunities in the interpretation of geochemical data from mollusk shells, *Geo-Mar. Lett.*, 28, 269–285, <https://doi.org/10.1007/s00367-008-0114-6>, 2008.
- 1080 Shen, C.-C., Lee, T., Chen, C.-Y., Wang, C.-H., Dai, C.-F., and Li, L.-A.: The calibration of $\text{D}[\text{Sr}/\text{Ca}]$ versus sea surface temperature relationship for *Porites* corals, *Geochim. Cosmochim. Acta*, 60, 3849–3858, [https://doi.org/10.1016/0016-7037\(96\)00205-0](https://doi.org/10.1016/0016-7037(96)00205-0), 1996.
- 1085 Simonet Roda, M., Griesshaber, E., Ziegler, A., Rupp, U., Yin, X., Henkel, D., Häussermann, V., Laudien, J., Brand, U., Eisenhauer, A., Checa, A. G., and Schmahl, W. W.: Calcite fibre formation in modern brachiopod shells, *Sci. Rep.*, 9, 598, <https://doi.org/10.1038/s41598-018-36959-z>, 2019a.
- Simonet Roda, M., Ziegler, A., Griesshaber, E., Yin, X., Rupp, U., Greiner, M., Henkel, D., Häussermann, V., Eisenhauer, A., Laudien, J., and Schmahl, W. W.: Terebratulide brachiopod shell biomineralization by mantle epithelial cells, *J. Struct. Biol.*, 207, 136–157, <https://doi.org/10.1016/j.jsb.2019.05.002>, 2019b.
- 1090 Simonet Roda, M., Griesshaber, E., Angiolini, L., Rollion-Bard, C., Harper, E. M., Bitner, M. A., Milner Garcia, S., Ye, F., Henkel, D., Häussermann, V., Eisenhauer, A., Gnägi, H., Brand, U., Logan, A., and Schmahl, W. W.: The architecture of Recent brachiopod shells: diversity of biocrystal and biopolymer assemblages in rhynchonellide, terebratulide, thecideide and craniide shells, *Mar. Biol.*, 169, 4, <https://doi.org/10.1007/s00227-021-03962-4>, 2022.
- 1095 Swart, P. K., Burns, S. J., and Leder, J. J.: Fractionation of the stable isotopes of oxygen and carbon in carbon dioxide during the reaction of calcite with phosphoric acid as a function of temperature and technique, *Chem. Geol. Isot. Geosci. Sect.*, 86, 89–96, [https://doi.org/10.1016/0168-9622\(91\)90055-2](https://doi.org/10.1016/0168-9622(91)90055-2), 1991.

- Swart, P. K., Elderfield, H., and Greaves, M. J.: A high-resolution calibration of Sr/Ca thermometry using the Caribbean coral *Montastraea annularis*, *Geochem. Geophys. Geosystems*, 3, 1–11, <https://doi.org/10.1029/2002GC000306>, 2002.
- 1100 Takayanagi, H., Asami, R., Abe, O., Kitagawa, H., Miyajima, T., and Iryu, Y.: Carbon- and oxygen-isotope compositions of a modern deep-water brachiopod *Campagea japonica* collected off Aguni-jima, Central Ryukyu Islands, southwestern Japan, 46, 77–87, <https://doi.org/10.2343/geochemj.1.0153>, 2012.
- 1105 Takayanagi, H., Ryuji, A., Osamu, A., Toshihiro, M., Hiroyuki, K., Keiichi, S., and Yasufumi, I.: Intraspecific variations in carbon-isotope and oxygen-isotope compositions of a brachiopod *Basiliola lucida* collected off Okinawa-jima, southwestern Japan, *Geochim. Cosmochim. Acta*, 115, 115–136, <https://doi.org/10.1016/j.gca.2013.03.026>, 2013.
- 1110 Takayanagi, H., Asami, R., Otake, T., Abe, O., Miyajima, T., Kitagawa, H., and Iryu, Y.: Quantitative analysis of intraspecific variations in the carbon and oxygen isotope compositions of the modern cool-temperate brachiopod *Terebratulina crossei*, *Geochim. Cosmochim. Acta*, 170, 301–320, <https://doi.org/10.1016/j.gca.2015.08.006>, 2015.
- Tesoriero, A. J. and Pankow, J. F.: Solid solution partitioning of Sr²⁺, Ba²⁺, and Cd²⁺ to calcite, *Geochim. Cosmochim. Acta*, 60, 1053–1063, [https://doi.org/10.1016/0016-7037\(95\)00449-1](https://doi.org/10.1016/0016-7037(95)00449-1), 1996.
- 1115 Ullmann, C. V., Frei, R., Korte, C., and Lüter, C.: Element/Ca, C and O isotope ratios in modern brachiopods: Species-specific signals of biomineralization, *Chem. Geol.*, 460, 15–24, <https://doi.org/10.1016/j.chemgeo.2017.03.034>, 2017.
- Veizer, J. and Prokoph, A.: Temperatures and oxygen isotopic composition of Phanerozoic oceans, *Earth-Sci. Rev.*, 146, 92–104, <https://doi.org/10.1016/j.earscirev.2015.03.008>, 2015.
- 1120 Vickers, M. L., Fernandez, A., Hesselbo, S. P., Price, G. D., Bernasconi, S. M., Lode, S., Ullmann, C. V., Thibault, N., Hougard, I. W., and Korte, C.: Unravelling Middle to Late Jurassic palaeoceanographic and palaeoclimatic signals in the Hebrides Basin using belemnite clumped isotope thermometry, *Earth Planet. Sci. Lett.*, 546, 116401, <https://doi.org/10.1016/j.epsl.2020.116401>, 2020.
- Vickers, M. L., Bernasconi, S. M., Ullmann, C. V., Lode, S., Looser, N., Morales, L. G., Price, G. D., Wilby, P. R., Hougård, I. W., Hesselbo, S. P., and Korte, C.: Marine temperatures underestimated for past greenhouse climate, *Sci. Rep.*, 11, 19109, <https://doi.org/10.1038/s41598-021-98528-1>, 2021.
- 1125 Washington, K. E., West, A. J., Kalderon-Asael, B., Katchinoff, J. A. R., Stevenson, E. I., and Planavsky, N. J.: Lithium isotope composition of modern and fossilized Cenozoic brachiopods, *Geology*, 48, 1058–1061, <https://doi.org/10.1130/G47558.1>, 2020.
- 1130 Watkins, J. M., Nielsen, L. C., Ryerson, F. J., and DePaolo, D. J.: The influence of kinetics on the oxygen isotope composition of calcium carbonate, *Earth Planet. Sci. Lett.*, 375, 349–360, <https://doi.org/10.1016/j.epsl.2013.05.054>, 2013.
- Watkins, J. M., Hunt, J. D., Ryerson, F. J., and DePaolo, D. J.: The influence of temperature, pH, and growth rate on the $\delta^{18}\text{O}$ composition of inorganically precipitated calcite, *Earth Planet. Sci. Lett.*, 404, 332–343, <https://doi.org/10.1016/j.epsl.2014.07.036>, 2014.
- 1135 Wierzbowski, H., Bajnai, D., Wacker, U., Rogov, M. A., Fiebig, J., and Tesakova, E. M.: Clumped isotope record of salinity variations in the Subboreal Province at the Middle–Late Jurassic transition, *Glob. Planet. Change*, 167, 172–189, <https://doi.org/10.1016/j.gloplacha.2018.05.014>, 2018.
- Williams, A.: A history of skeletal secretion among articulate brachiopods, *Lethaia*, 1, 268–287, <https://doi.org/10.1111/j.1502-3931.1968.tb01741.x>, 1968.
- 1140 Williams, A.: The secretion and structural evolution of the shell of thecideidine brachiopods, *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 264, 439–478, <https://doi.org/10.1098/rstb.1973.0001>, 1973.

de Winter, N. J., Müller, I. A., Kocken, I. J., Thibault, N., Ullmann, C. V., Farnsworth, A., Lunt, D. J., Claeys, P., and Ziegler, M.: Absolute seasonal temperature estimates from clumped isotopes in bivalve shells suggest warm and variable greenhouse climate, *Commun. Earth Environ.*, 2, 121, <https://doi.org/10.1038/s43247-021-00193-9>, 2021.

1145 Yamamoto, K., Asami, R., and Iryu, Y.: Carbon and oxygen isotopic compositions of modern brachiopod shells from a warm-temperate shelf environment, Sagami Bay, central Japan, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 291, 348–359, <https://doi.org/10.1016/j.palaeo.2010.03.006>, 2010a.

1150 Yamamoto, K., Asami, R., and Iryu, Y.: Within-shell variations in carbon and oxygen isotope compositions of two modern brachiopods from a subtropical shelf environment off Amami-o-shima, southwestern Japan, *Geochem. Geophys. Geosystems*, 11, Q10009, <https://doi.org/10.1029/2010GC003190>, 2010b.

Yamamoto, K., Asami, R., and Iryu, Y.: Brachiopod taxa and shell portions reliably recording past ocean environments: Toward establishing a robust paleoceanographic proxy: BRACHIOPOD OXYGEN ISOTOPE RECORDS, *Geophys. Res. Lett.*, 38, L13601, <https://doi.org/10.1029/2011GL047134>, 2011.

1155 York, D., Evensen, N. M., Martínez, M. L., and De Basabe Delgado, J.: Unified equations for the slope, intercept, and standard errors of the best straight line, *Am. J. Phys.*, 72, 367–375, <https://doi.org/10.1119/1.1632486>, 2004.

Zaarur, S., Affek, H. P., and Brandon, M. T.: A revised calibration of the clumped isotope thermometer, *Earth Planet. Sci. Lett.*, 382, 47–57, <https://doi.org/10.1016/j.epsl.2013.07.026>, 2013.

1160 Zweng, M. M., Reagan, J. R., Seidov, D., Boyer, T. P., Locarnini, R. A., Garcia, H. E., Mishonov, A. V., Baranova, O. K., Weathers, K., Paver, C. R., and Smolyar, I.: World Ocean Atlas 2018, Volume 2: Salinity, Mishonov Tech. Ed NOAA Atlas NESDIS, 82, 50, 2018.