

# 1 **Meteotsunami in the United Kingdom: The hidden hazard.**

2 Clare Lewis<sup>1,2</sup>, Tim Smyth<sup>2</sup>, David Williams<sup>4</sup>, Jess Neumann<sup>1</sup>, Hannah Cloke<sup>1,3</sup>

3  
4 <sup>1</sup> Department of Geography & Environmental Science, University of Reading, Reading, UK

5 <sup>2</sup> Plymouth Marine Laboratory, Prospect Place, Plymouth, Devon, PL1 3DH, UK

6 <sup>3</sup> Department of Meteorology, University of Reading, Reading, UK

7 <sup>4</sup>WTW, 51 Lime Street, London, EC3M 7DQ, UK

8 Correspondence to: Clare Lewis (clare.lewis@pgr.reading.ac.uk)

## 9 **Abstract.**

10 This paper examined the occurrence and seasonality of meteotsunami in the United Kingdom (UK) to present a  
11 revised and updated catalogue of events occurring since 1750. Previous case studies have alluded to a summer  
12 prevalence and rarity of this hazard in the UK. We have verified and classified 98 events using a developed set of  
13 identification criteria. The results have revealed a prominent seasonal pattern of winter events which are related  
14 to mid latitude depressions with precipitating convective weather systems. A geographical pattern has also  
15 emerged, highlighting three ‘hotspot’ areas at the highest risk from meteotsunami. The evidence reviewed, and  
16 new data presented here shows that the hazard posed by meteotsunami has been underestimated in the UK.

17  
18 Keywords: meteotsunami, UK, hazard, mid latitude depressions.

## 20 **1 Introduction**

21 Meteotsunamis or meteorological tsunamis are globally occurring progressive shallow water waves with a period  
22 of between 2 to 120 minutes which results from an air-sea interaction. They tend to be initiated by sudden pressure  
23 changes and wind stress from moving atmospheric systems with sources ranging from convective clouds,  
24 cyclones, squalls, thunderstorms, atmospheric gravity waves and strong mid-tropospheric winds (Vilibic and  
25 Sepic, 2017). [The atmospheric pressure changes are typically only a few mb over a few tens of minutes which  
26 corresponds to only a few centimetres of sea level change occurring in a process known as the inverse barometer  
27 effect (for example, a 3 mb pressure jump will produce a 30 cm ocean wave). The atmospheric disturbance  
28 transfers energy into the ocean initiating and amplifying a water wave which travels at the same speed as the  
29 atmospheric wave, in a process known as Proudman resonance (Proudman, 1929). When the water wave reaches  
30 the coastline and shallower water, it becomes a multi resonant phenomena and is further amplified through coastal  
31 resonances. For example, if the wave reaches the entrance of a semi enclosed basin it can induce an oscillation in

32 the basin known as harbour resonance. However, if the wave reaches a beach type environment and the along  
33 shore component of the disturbance equals the phase speed of the edge wave this is a process known as Greenspan  
34 resonance (Monserrat et al. 2006). The resultant waves can elevate the coastal water level and can substantially  
35 increase flow velocities with the potential for rip currents (Linares et al. 2019). Due to the rapid onset and  
36 unexpected nature of meteotsunami waves, they have the potential to cause destruction, injuries and even fatalities  
37 (Sibley et al. 2016). For a global perspective and overview of meteotsunami observations we recommend Pellikka  
38 et al. (2020) for observations in Finland, Sepic et al. (2018) for the Adriatic, Belche et al. (2016) for seasonality  
39 of meteotsunami in the Great Lakes, Pattiaratchi and Wijeratne (2016) for observations in southwest Australia  
40 and Monserrat, Vilibic and Rabinovich (2006) provide a general overview of the mechanisms of meteotsunami.

41  
42 Meteotsunami research and monitoring is more advanced in the Mediterranean, the East Coast of the USA, and  
43 the Great Lakes due to the higher number of recorded events. However, events in the UK appear to be rare and  
44 are believed to be less devastating, meaning that research has been limited to date.

45 The two principal factors contributing to this belief are:

- 46 1. The current (since 1993) 15-minute sampling interval that is used at UK tide gauges is incapable  
47 of detecting waves with periods of between 2 – 120 minutes. This means that many events go  
48 unobserved, wave heights are underestimated, or meteotsunamis are mischaracterised as seiches,  
49 tsunamis or surge.
- 50 2. Until recently research has suggested that UK meteotsunamis are generated by precipitating,  
51 convective weather systems associated with hot weather. Such mesoscale convective systems may  
52 be associated with synoptic “Spanish plume” events. These synoptic events are more prevalent  
53 between May - October (Haslett et al. 2009b; Tappin et al. 2013; Sibley, 2012 and 2016; Thompson,  
54 2020), leading to the belief that meteotsunami occurrence is a summer-time phenomena. However,  
55 it is now emerging that embedded convection within winter frontal systems may also be responsible  
56 for a sizeable proportion of these waves (Williams et al. 2021).

57 Several issues have resulted from the untested assumption that meteotsunami events are 1) low frequency and 2)  
58 predominantly occur in summer, which has been combined with 3) the lack of high-resolution temporal data.  
59 Firstly, there is no central database of UK events. Secondly, there is no standardised methodology of meteotsunami  
60 identification. Thirdly, there is no Government or regional policy in place to cover impacts from a meteotsunami  
61 event. There is a misconception of the risk posed by meteotsunami especially for coastal areas that are already at  
62 risk from storm impacts associated with pluvial (extreme precipitation) and fluvial hazards (high levels of river

**Commented [CL1]: R2:** Introduction: The authors mention the coastal processes such as shoaling and refraction and their effect on wave amplification as the meteotsunami waves travel toward the coastline. However, meteotsunamis are multi-resonant phenomena and the major amplification mechanisms are due to those different resonance mechanisms which need to be definitely described in the Introduction part of such a manuscript. I also recommend including the fundamentals of meteotsunami generation, e.g. the inverse barometer law, only a few cm of waves would occur in the static condition, etc.

**R2:** Lines 27-28: “The characteristics of the atmospheric disturbance transfers energy into the ocean initiating and amplifying a water wave that travels at the same speed as the atmospheric wave in a process known as Proudman resonance (Proudman, 1929).” It is not the characteristics of the atmospheric disturbance transferring energy into the ocean. Please correct this sentence.

**Commented [CL2R1]: A:** amended as requested

**Commented [CL3]: R2:** Lines 34-36: It is better to provide an overall one-paragraph summary of those studies and then kindly refer the reader to those papers without directly saying “see”.

**Commented [CL4R3]: A:** Summary added as requested

**Commented [CL5]: R2:** Lines 38-39: In my opinion, it is better to write as “... due to the higher number of recorded events.” or “... due to more recorded events.” instead of “high frequency of recorded events”

**Commented [CL6R5]: A:** Amended as requested

**Commented [CL7]: R2:** Lines 55-56: “Thirdly, there is no Government or regional policy in place to cover future adaptation strategies in the case of sea-level rise.” I believe this is a too generic sentence and needs to be more specific by relating it with “meteotsunami” since it seems like the statement points out the issues related to the “sea level rise” rather than the “meteotsunami research.”

**Commented [CL8R7]: A:** Amended as requested

63 discharge). In the future the overall level of risk is likely to be greatly exacerbated by rising sea levels and an  
64 intensification of storm frequency and severity (Vilibic et al. 2018; Masselink et al. 2015).  
65 As stated by Sepic et al. (2015) the assessment of meteotsunami should become the standard in coastal hazard  
66 assessments, event cataloguing is a pre-requisite for any coastal hazard assessment especially in identifying the  
67 geographical areas that have experienced meteotsunami and the frequency of exposure.  
68 The aim of this paper is to compile, update and extend the existing list UK meteotsunami to include winter events,  
69 and to highlight the occurrence, frequency, and spatial distribution of events. Where seasonality was alluded to in  
70 Williams et al. (2021), their study was principally focused on meteotsunami in Northwest Europe from 2010 to  
71 2017. This paper will further their study by focussing in on UK waters only and will add new events up to the end  
72 of 2022. The methodology fulfils this aim by applying a set of developed identification criteria to the re-  
73 assessment of fragmented historical accounts and to the analysis of tide gauge and atmospheric data to identify  
74 new events. The outcome also highlights the potential element of winter compound hazard risk which may occur  
75 when meteotsunami waves arrive at the coast in short succession or concurrently with other storm associated  
76 hazards.

77 We propose the following research questions:

- 78 1. What standardised criteria should be used to identify meteotsunami?
- 79 2. Have events occurred which were ignored or misidentified?
- 80 3. In which regions of the UK and in what months do meteotsunami occur most frequently?
- 81 4. What are the atmospheric variables that can be correlated with meteotsunami events?

## 83 2 Methodology

84 This section outlines the data sources and identification criteria used to fulfil the objective of cataloguing and  
85

**Commented [CL9]: R1:** The authors need to clarify the original findings of this study which are different from the previous ones. Because this study shares similarities with Williams et al. (2021) in the methodology and results.

**Editor:** Address the originality of your findings from previously published papers.

**Commented [CL10R9]: A:** Where seasonality is alluded to in Williams et al (2021) and is principally focused on precipitating atmospheric systems linked to NW European meteotsunami up to 2017. We have added precision by extending this study to focus in on UK waters only up to 2022 and have subsequently introduced a geographical element with respect to seasonality. This has allowed for the examination of 'hotspot' areas.

In Line 72 we have stated that this study is indeed a form of update to the existing UK meteotsunami catalogue as presented by Thompson et al. However, their study does not address winter events and only goes up to 2016/7. Using our developed identification criteria for which there appears to be a lack of standardisation, we have verified, updated and extended the catalogue to the end of 2022.

**Commented [CL11]: R2:** The research question in #4 is not clear and difficult to understand, I suggest revising the sentence.

**Commented [CL12R11]: A:** Research question revised

86 characterising UK meteotsunami. We have extrapolated as much quantitative data as possible, to verify the  
87 event with the standardised criteria and to then arrange the results into tabular form to allow ease of use (Table  
88 1).

## 90 2.1 Meteotsunami identification criteria

91 As there are currently no fixed criteria for what qualifies as a meteotsunami, in this paper we bring together various  
92 aspects used by other researchers in the field, into one standardised system. Figure 1 (a – d) displays a visual  
93 representation of the commonly used criteria, which we explain in more detail in sections 2.1.1 – 2.1.2. The  
94 methodologies that have been previously used by researchers and studies have variations, with some using  
95 qualitative methods that base events on eyewitness accounts (Haslett et al, 2009a/b) and others using quantitative  
96 data from sea level and atmospheric observations (Tappin et al. 2013; Sibley, 2016). For the purpose of this paper,  
97 we have classified meteotsunami as atmospherically induced sea level oscillations meeting at least one sea level  
98 and one atmospheric characteristic. This allows for the distinguishing of meteotsunami from other types of  
99 waveforms and is applicable to either qualitative accounts or quantitative data.

100

### 101 2.1.1 Sea level criteria (Category 1)

- 102 a. Periods of sea level disturbance ranging from between 2 and 120 minutes (Figure 1a).
- 103 b. Wave heights exceeding 0.20 m. This threshold is within the peak thresholds of 0.2 m and 0.3  
104 m as used by other researchers in the field such as Williams et al. (2021); Dusek et al. (2019); Belche  
105 et al. (2016); Sepic et al. (2012) and Monserrat, Vilibic and Rabinovich (2006). A 0.3 m water  
106 elevation may not appear to be dangerous, but a meteotsunami in 2003 in New Zealand caused a  
107 fully laden oil tanker to be grounded through strong currents (Goring, 2009). Lynett et al. (2014)  
108 also states that any wave over 0.3 m will start to float vehicles regardless of flow velocity and is  
109 enough to sweep people off of their feet. These thresholds are a tried and tested set of characteristics  
110 that reflect meteotsunami especially those in UK waters. 0.2 m was opted to be used as the lower  
111 end of the threshold as this is more suitable for distinguishing a greater number of events that may  
112 have been missed at the higher end of the threshold (0.3m). Any anomaly below 0.2m would not be  
113 large enough to allow for accurate verification and for its separation from any other water  
114 disturbances. (Figure 1a illustrates the meteotsunami wave height criteria in the data as recorded on  
115 27 June 2011).

**Commented [CL13]: R2:** The sea level criteria used in the study seem the same as the ones by Williams et al. 2021. Any new approach should be highlighted.

**Commented [CL14R13]: A:** The sea level criteria used here are very similar to the ones used in many studies including Williams et al. 2021 and this is because they are a tried and tested set of characteristics that reflect meteotsunami. We have adjusted the wave amplitude threshold used in Williams et al. 2021 from 0.25m down to 0.20m as we feel this is more reflective of UK meteotsunami and will catch more events that may have previously been discarded.

116 c. A wave disturbance registering at two or more locations or tide gauge stations (Williams et al.  
117 2021; Kim et al. 2021).

118

### 119 2.1.2 Atmospheric criteria (Category 2)

120 a. The presence of a convective weather system at the time of the wave event displaying high radar  
121 reflectivity with precipitation rates exceeding  $2 \text{ mm/h}^{-1}$ , initiated over the sea. (Figure 1b represents  
122 the radar reflectivity of the various convective weather systems present during four different  
123 meteotsunami events).

124 b. An atmospheric pressure of 1005 mb or less with a rapid change of  $\pm 1 \text{ mb}$  in 30 minutes or a 3  
125 mb fall over three hours or less (Montserrat, Vilibic and Rabinovich, 2006). (Figure 1c illustrates this  
126 distinct air pressure change as recorded during the 28 October 2013 event).

127 c. Convective Available Potential Energy (CAPE) showing the unstable vertical profile of the  
128 atmosphere that leads to convective activity (Williams et al. 2019). (Figure 1d displays a radiosonde  
129 ascent showing sufficient CAPE to produce the event that occurred on 1 July 2015 at Stonehaven,  
130 East Scotland. Even though CAPE is a bulk atmospheric measurement and meteotsunami are  
131 localised, if this element is present in conjunction with the other indicators it supports the presence  
132 of convective activity which aids in the generation of meteotsunami.

133 d. A change in wind speed exceeding  $10 \text{ m/s}^{-1}$  (anything under this is too weak for a meteotsunami  
134 to generate) or/and a drop in air temperature of  $1.5^\circ\text{C}$  in 30 minutes (Figure 1c demonstrates this  
135 increase in wind speed as recorded during the 28 October 2013 event).

136

### 137 2.1.3 Geological criteria (Category 3)

138 a. The absence of any other explanation or data to imply an alternative source trigger. For example,  
139 the presence of seismic triggers within the continental shelf area which would produce a geological  
140 tsunami wave. However, there is one exception to this rule which for the purpose of this paper we  
141 include as a meteotsunami trigger. Volcanic eruptions, this was demonstrated on 28 August 1883  
142 (Krakatoa) and recently on 16 January 2021 (Tonga Ha'apai) where wave anomalies occurred and  
143 were the product of air pressure waves created by the eruptions. It may be argued that they are not  
144 to be classed as meteotsunami waves. However, for the purpose of this catalogue, we are classifying  
145 them as meteotsunami as they are sourced from air pressure disturbances which couple with water  
146 waves and have a wave period of 2 to 120 minutes. The force of the Tonga Ha'apai explosions sent

**Commented [CL15]: R2:** Lines 117-118: "...the event that occurred on 1 July 2015." Better to also indicate the location of the event.

**Commented [CL16R15]: A:** Amended as requested

**Commented [CL17]: R2:** Line 122: Atmospheric criteria d: Isn't 5 m/s also too low for a threshold wind speed for meteotsunami generation? In Figure 1c, the wind speed almost exceeds 20 mph~ 9 mps for a long duration. Could the authors explain the rationale behind this selection?

**Commented [CL18R17]: A:** Typo error, amended

147 a shockwave through the atmosphere that circled the globe three times. The resultant pressure  
148 wave travelled at close to the speed of sound and as a result coupled with ocean waves to create a  
149 meteotsunami which was detected as far away as Portugal and the UK (Burt. S, 2022).

150  
151 To ease the interpretation of results, the UK coastline has been partitioned into six coastal regions based on the  
152 National Tidal and Sea Level Facility (NTSLF) tide gauge network (Supplementary Table S1). The data are also  
153 separated into two seasons (each comprising of six months) that divide up the calendar year at the spring and  
154 autumn equinoxes (Haigh et al. 2016). April to September inclusive is referred to throughout this paper as  
155 'summer' and October to March is referred to as 'winter'. Finally, due to the nature of the data, two time series of  
156 meteotsunami are being referred to throughout this paper, one based primarily on historical eyewitness accounts  
157 due to a lack of high frequency instrumentation (the years 1750 to 2009 AD), and one based on and verified by  
158 quantitative instrumental data (the years 2010 to 2022 AD).

159

## 160 2.2 Historical record (1750 to 2009)

161 To gain a complete understanding of these events we follow Long (2015) and Haslett and Bryant (2008) who  
162 dated their historic tsunami catalogues back to approximately 1000 AD. We noted that any events preceding 1750  
163 AD were vaguely recorded, making validation problematic so we opted to date our catalogue back to 1750.  
164 References to meteotsunami like events in historical accounts tend to be based on descriptions of the state of the  
165 water at the coast with a lack of instrumental tidal data. There is a lack of or limited weather data so tracing back  
166 the atmospheric source is not as straightforward. It is only until the last few decades that meteorological data with  
167 sufficient resolution have been readily available. With tide gauge data, prior to 1993 the resolution was hourly,  
168 and it was not until 1996 that all the current tide gauge sites became fully operational. Therefore, we have used  
169 2009 as the upper limit of the historical record. The historical reports tend to be derived from newspaper articles,  
170 parish records, harbourmaster records and eyewitness accounts. Although there is reason to be sceptical of these  
171 accounts as they afford a level of biased review and sensationalism, they do still hold value in terms of a societal  
172 viewpoint and may help to fill in any gaps (Haslett and Bryant, 2009a/b).

173 There are certain characteristics that flag up in an historical account to verify whether it is a meteotsunami event  
174 or not. To illustrate this, we can highlight the historical account for the event of 23 May 1847 where we can look  
175 at a letter from Robert Blight of Penzance dated 24 May and published in the Cornwall Royal Gazette on 28 May.  
176 The full extract can be found in supplementary extract S1 of this paper and in Long (2015, p26).

**Commented [CL19]: R2:** It is better to indicate how this "maximum wave height" value is obtained, i.e., from measurement data or eyewitness observation?

**Commented [CL20R19]: A:** It is a general rule that any event within the historical record can be assumed to have been from eyewitness accounts due to the lack of high frequency instrumentation. Any event post 2009 has been verified by quantitative data. This will of course be noted a little clearer in the methodology section

177 “... The changes in the atmosphere during the day were very remarkable. In the morning, about six o’clock, we  
178 had a breeze from the southeast; by eight, it was a perfect calm; between ten o’clock and two, the mercury sunk  
179 several degrees; about three in the afternoon a breeze sprung up suddenly from the west, and the sky, as suddenly,  
180 became overcast..... It is very probable that all these changes, and even the agitation of the sea, were produced  
181 by electricity...”

182 In this particularly detailed account (supplementary extract S1) we can identify six of the nine criteria, including  
183 a drawback and sudden in rush of water, accompanied by a rumbling noise and the water being higher than  
184 expected at eight feet (criteria 1A and 1D), all indicating a tsunami like event. The key to the identification of a  
185 meteotsunami is in the atmospheric portion of the account, what started out as calm morning led to a change in  
186 wind speed and direction, veering from south easterly in the morning to westerly in the afternoon (criteria 2D).  
187 This variable wind was accompanied by a drop in temperature (criteria 2D) and finally, there was mention of the  
188 presence of a storm in terms of overcast sky, threatening rain and lightning (criteria 2A). As such, we identify this  
189 wave as a meteotsunami by applying both of our oceanographic and atmospheric criteria to the historic account.  
190

### 191 2.3 Tide gauge analysis for the 2010 to 2022 record

192 To identify meteotsunami from 1 January 2010 to 31 December 2022 we use data records that are available at  
193 higher frequencies meaning meteotsunami are more distinctly observable. The information for this portion of the  
194 catalogue is sourced from the British Oceanographic data centre (BODC) website (<https://www.bodc.ac.uk/>) and  
195 the International Oceanographic Commission (IOC) website (<https://ioc-sealevelmonitoring.org/>) where data are  
196 displayed from the ‘Class A’ network of tide gauges owned and funded by the Environment Agency (EA). We  
197 also use the postprocessed data of Williams et al. (2021) where the raw sea level tide gauge data has been high  
198 pass filtered to isolate high frequency disturbances. This removes periods of over 120 minutes and separates out  
199 the tidal components. In this way any signals in the tsunami frequency band (2 to 120 minutes) are isolated from  
200 the sea level elevations. Any remaining signals larger than the background noise are then identified and checked  
201 against our threshold criteria to verify events as potential meteotsunami. Apart from the standard processing to  
202 remove any erroneous spikes outside of the parameters, a visual quality control was carried out, where a seven-  
203 day plot of the data was evaluated to highlight any clear artificial spikes or gaps. Also, any data points that had  
204 no accompanying air pressure changes were also excluded from any further analysis.

**Commented [CL21]: R2:** 2.3 “Wave data analysis for the 2010 to 2022 record.” I would suggest changing the heading of this section to “Tide gauge data analysis.” or “Sea level data analysis.” not to be confusing.

**Commented [CL22R21]: A:** Amended as requested

**Commented [CL23]: R2:** I also recommend including the details of the sea level data processing such as how did the authors handle the gaps or spikes in the measured data?

**Commented [CL24R23]: A:** Details added as requested

208 **2.4 Atmospheric data analysis for the 2010 to 2022 record**

209 The time of the potential meteotsunami events are noted from the tide gauge data and they are then linked to  
210 specific precipitating convective atmospheric systems by using the meteorological C-band radar network, which  
211 is pre-processed by the UK Meteorological Office before download (Met Office 2003). The convective systems  
212 highlighted by the radar are classified into four distinct types (as shown in Figure 1b). These are: (1) open cells  
213 which are situated behind the cold front of cyclonic weather, usually where cold dry air passes over the warm sea  
214 creating shallow convection; (2) Quasi linear systems which tend to be multi-cellular and linearly organised with  
215 high CAPE, heavy precipitation, and strong winds (this type of weather feature are sometimes called squall lines  
216 and can occur within synoptic Spanish Plume events); (3) Isolated small short duration (<1h) thunderstorm cells  
217 and (4) Nonlinear clusters which are large circular, long lived clusters of precipitation and thunderstorm cells.


218 The atmospheric ascent soundings are obtained from the University of Wyoming website  
219 (<http://weather.uwyo.edu/upperair/sounding.html>) with the UK stations at Camborne (station number: 03808) and  
220 Lerwick (station number: 03005) being used. Soundings are available for 0000 UTC and 1200 UTC on each day  
221 and if a CAPE value of greater than 0 occurs then this shows a marginally unstable atmosphere leading to  
222 convective activity. Finally, the synoptic charts allow for verification of the storm system including the location  
223 of the pressure centres and fronts at the time of the meteotsunami wave event.

224


**Commented [CL25]: R2:** Lines 196-199: Following the URL provided, which stations are used for the analysis? It would be better to provide.

**Commented [CL26R25]: A:** Details added as requested







**Figure 1a:** Devonport (50°36N 4°18W) tide gauge for 27 June 2011 showing a distinct sea level disturbance at 0830 UTC as highlighted with a red circle. This is a representation of criteria 1b. The timing of this 0.25 m rise and fall in the sea level corresponds with the arrival of the meteosunami event at that specific location.



**Figure 1c:** The atmospheric pressure, wind speed and precipitation at Reigate (51°14N 0°11W) during the 27 to 28<sup>th</sup> October 2013 storm associated with the meteosunami. A representation of criteria 2b and 2d. The graph shows atmospheric pressure (red line) of less than 1005 mb and falling as the atmospheric disturbance moves over the area, with a corresponding rising wind speed of 20 mph (green line) and precipitation (blue bars). Reproduced with the kind permission of Simon Collins. <https://rgsweather.com/2013/10/29/st-jude-causes-and-impacts-of-the-october-storm-27-28-2013/amp/>



**Figure 1b:** The four different types of convective activity as shown on radar reflectivity identifying meteosunami events. A representation of criteria 2a. Orange and red in the images shows high precipitation rates ( $>4 \text{ mm h}^{-1}$ ). With idealised images shown on the left and actual examples taken from UK events on the right. All showing date, time, and direction of the storm as well as the location of the tide gauges (white dots). Image by David Williams, Journal of Physical Oceanography (<https://doi.org/10.1175/JPO-D-20-0175.1>), licensed under a [Creative Commons — Attribution 4.0 International — CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)



**Figure 1d:** The Nottingham radiosonde ascent at 1200 UTC on 1 July 2015 during a meteosunami event in the North Sea. A representation of criteria 2c which indicates sufficient CAPE (462.1  $\text{J Kg}^{-1}$ ) to produce high base convective activity, with the cloud base at an approximate height of 3000 m and cloud top at 11000m. <http://weather.uwyo.edu/upperair/sounding.html>

226

227

228

229

230

### 3 Results

231

232

233

234

235

236

In this section we highlight the seasonal occurrence and distribution of UK meteosunami events in both the historical record and the more recent instrumental data record. This is augmented by the identification of trigger systems associated with the events where available. It is prudent to note here that the catalogue cannot be considered as complete, and this is signified by dashed lines (i.e., -) in the columns where data or information are either unavailable or have not been located.

237 **3.1 Historical record (1750 to 2009)**

238 We identify 98 events as being meteotsunami occurring in UK waters between January 1750 and December 2022  
239 (Table 1), with 48 of these occurring within the historical record (1750 to 2009). This record shows that 67% of  
240 documented meteotsunamis occur in summer (April – September), with 44% of documented meteotsunamis  
241 occurring in July and August. The single year experiencing the most documented events was 1802 AD, numbering  
242 three, and the decade experiencing the most documented events was the 1840s, with six in total. The presence of  
243 a storm and/or characteristics of convective activity (thunder, and lightning) at the time of the wave event was  
244 noted for 42 of the 48 events (91%) in the historical record. There was also a defined southwest prevalence of  
245 meteotsunami in historical documents, with Devon, Cornwall and Somerset recording a combined total of 29  
246 events. Within the historical record we have identified four new events and reclassified four tsunamis, three storm  
247 surge and nine events of unknown origin as meteotsunami. Seven of these occurred within winter months (Table  
248 1).

249  
250 **3.2 Seasonal and locational frequency of UK meteotsunami events (2010 to 2022)**

251 Meteotsunamis have been thought to be a rare phenomenon in the UK and that when they do occur, it has been  
252 tended to be in the summer months due to the more abundant convective activity (Haslett et al, 2009b; Tappin et  
253 al, 2013; Sibley, 2016; Thompson, 2020). However, of the 98 identified meteotsunami events verified in this  
254 paper, 50 have been interpreted as occurring since 2010, 33 (66%) of those occurring during the winter months  
255 and nine of the winter events are identified as new. We find that not only are UK meteotsunami more common in  
256 occurrence than previous research indicates, but that they are a year-round phenomenon, as exhibited in Table 1  
257 and Figures 2 and 3.

258 The historical section of the catalogue shows an estimated return period of 5.4 years. This return period  
259 considerably decreases for the instrumental data section where the UK return period reduces to an estimated 0.25  
260 years. With an average of four events per year, we can see that certain years have experienced above average  
261 numbers and high proportions of winter events, with seven winter events out of eight in 2013, four out of seven  
262 in 2021 and five out of seven in 2022. Figure 3 displays the seasonal distribution of these events, with 34% of  
263 meteotsunami recorded in December and January, and no events being recorded in March or April. Following  
264 statistical analysis, the recorded maximum wave amplitude for each event resulted in a mean wave height of 0.33  
265 m for winter and 0.35 m for summer. With a t-test score of 0.30 and a P-value of 0.07, the tests indicate a similarity  
266 between the two sample sets, where the difference between seasonal wave heights is considered to be not  
267 statistically significant.

**Commented [CL27]:** Editor: clearly identify new events and winter events

**Commented [CL28R27]:** A: new events and winter events included in results, discussion and conclusion section

**Commented [CL29]:** R2: The number of investigated meteotsunami events is indicated as 95 in the abstract. How many of them are newly identified? Please indicate.  
Editor: clearly address the newly identified events.


**Commented [CL30R29]:** A: Many of the historical events were mis-identified in accounts as abnormal coastal flooding, non-tsunami, storm surge or unknown. By using the methodology, we re-identified them as meteotsunami which are now 'new' to the catalogue. In the recent record we have identified new events direct from the data. Table 1 has also been readjusted to distinguish these from the already correctly identified and verified events. Discussion L334 and Conclusions L488 also refer to the new events.

**Commented [CL31]:** R2: Lines 241-243: I recommend providing the details of the "statistical analysis" mentioned here. What I mean is that the following questions arise while reading: How did you obtain these average wave height values? Did you take the maximum observed "peak-to-trough" value of each event? How did you extract those values?

**Commented [CL32R31]:** A: We took the maximum wave amplitude value recorded for each event.

268 Summarising the results from the catalogue in its entirety, we suggest that there are three ‘hotspot’ regions where  
 269 meteotsunami events appear to be most frequent, these are 1) northwest Scotland, 2) Wales and 3) the southwest  
 270 UK. Up until 2009, Penzance in the southwest UK had experienced the most meteotsunami with eight in total.  
 271 Then from 2010, Kinlochbervie in Northwest Scotland experienced the maximum wave height of 0.51 m during  
 272 the 16 November 2016 event. This same location was exposed to 14 separate meteotsunami events in the 12 years  
 273 from 2010 to 2022. Harbour style geomorphology appears to be more susceptible to meteotsunami resonance  
 274 recording 71% of the events and beach environments with the remaining 29%.

275  
 276



**Figure 2:** Seasonal distribution of UK meteotsunami events, historical record (1750 to 2009) and current record (2010 to 2022).


277  
 278  
 279  
 280  
 281

**Commented [CL33]: R2:** Lines 247-248: “Then from 2010, Kinlochbervie in northwest Scotland has been exposed 14 times experiencing the highest maxima of wave height at 0.51 m.” Here it is also not clear that Kinlochbervie has experienced exactly 0.51 m maximum wave height 14 times OR the maximum wave heights that Kinlochbervie has experienced exceeded 0.51 m 14 times. Please clarify.

**Commented [CL34R33]: A:** Amended as requested

**Commented [CL35]: R2:** Figure 2. Both Figures have the heading “Seasonal Distribution of UK Meteotsunami 1750 to 2009!” The figures also look the same?

**Commented [CL36R35]: A:** upload oversight, figures have been adjusted



**Figure 3:** Seasonal and locational distribution of maximum wave heights from 1750 to 2022. Number of events at specific locations are represented by dot size as shown in the key. Base map © Crown copyright and database rights 2022 Ordnance Survey (100025252).

282

### 283 3.3 Relationship between meteotsunami and winter storms

284 In this section, we highlight two specific meteotsunami events that occurred in two consecutive winter seasons.

285 These two events have been picked as they are new events to the catalogue, and they represent a typical winter  
286 meteotsunami hidden in the associated storm data. The winter of 2021/22 saw seven sequential named storms

287 with five verifiable meteotsunami events, one of which was the 20th of October 2021. The winter of 2022/23 saw  
288 3 likely / numerically verifiable meteotsunami events, one of which was the 1 November 2022. Both meteotsunami  
289 events were low profile, localised in nature and hidden within larger scale heavily precipitating low-pressure  
290 systems.

291

#### 292 3.3.1 Event 1: 20 October 2021

293 Two low pressure systems developed in the Atlantic Ocean and propagated eastwards towards the southwest UK.

294 The first system which was detected as a mature echo signature on radar contained a sharp cold front (squall)  
295 which moved into Cornwall at approximately 0400 UTC (criteria 2a and Figure 4a) with a simultaneous leading  
296 air pressure rise of 1.6 mb over 4 minutes followed by a sharp 2°C air temperature drop (criteria 2b/d and Figure

**Commented [CL37]: R2:** Figure 3. A legend for dot size is necessary. How is maximum wave height represented here as mentioned in the figure caption?

**Commented [CL38R37]: A:** Dot size added to key

**Commented [CL39]: R2:** .3 Relationship between meteotsunami and winter storms: What is the reason behind selecting those specific two events “5 December 2013” and “20 October 2021?” I believe that it is important to mention.

**Editor:** Address newly identified winter events

**Commented [CL40R39]: A:** Reasoning included and event dates changed to reflect newly identified events

297 4c). A flattish ridge between this first system and the second system named Aurore by Meteo France led to a  
298 yellow rainfall warning being issued in the UK. At 1600 UTC the second system with a low-pressure centre of  
299 992 mb moved into the Isles of Scilly and propagated across Cornwall and Devon, it contained a heavily  
300 precipitating non-linear system with convective activity and strong winds (+70 mph) rapidly veering from west  
301 to south (criteria 2d). This system initiated a sharp air pressure rise of 0.5 mb over 2 minutes which coincided  
302 with a high tide. Both low pressure systems initiated a series of meteotsunami waves that tracked eastwards along  
303 the coast of Cornwall, Devon, and Dorset. Wave anomalies were recorded in Plymouth at 1645 UTC with a  
304 maximum wave height of 0.36 m, Totnes at 1700 UTC and Port Isaac, Weymouth, and the Isle of Wight at 1800  
305 UTC before dissipating (criteria 1b/c).

306


307 **Event 2: 1 November 2022**

308 A series of low-pressure systems over the Atlantic Ocean, swept into the southwest UK on 1 November, the first  
309 one with its centre over Cornwall at 0000 UTC, followed by a second low pressure system arriving along the  
310 southwest coast at approx. 0600 UTC then moving northeast up over the UK.

311 This synoptic situation was complicated by a series of associated cold fronts followed by low pressure troughs. A  
312 quasi-linear precipitation system with its associated convective cells developed in the vicinity (criteria 2a and c,  
313 Figures 4d and e). The arrival of the storm feature was detected in surface observations with a sharp 1 mb/35  
314 minutes air pressure rise (Figure 4f) which coincided with a series of unpredictable meteotsunami waves which  
315 reached a maximum wave height of 0.3 m (criteria 2b). The waves tracked along the southwest UK alongside of  
316 the movement of the cold fronts, the heavily precipitating cells and the convective activity where it was recorded  
317 at five tide gauge sites along the southwest coast at Port Isaac, St Marys, Newlyn, Plymouth and Totnes (criteria  
318 1c). The first series of wave anomalies occurred at 0900 UTC coinciding with a high tide followed by a second  
319 set of wave anomalies at 1600 UTC coinciding with a low tide.

320

321




a. Rain radar 20 October 2021 at 0300 UTC



b. Lightning 20 October 2021 at 0000 UTC



c. Air pressure 20 October 2021



d. Rain radar 1 November 2022 at 0900 UTC



e. Lightning 1 November 2022 at 0600 UTC



f. Air pressure 1 November 2022

**Figure 4:** The relationships between criteria of two winter meteotsunami events. All images are open source: Rain radar : [www.ventusky.com](http://www.ventusky.com) Lightning : [www.lightningmaps.org](http://www.lightningmaps.org) Air pressure: [www.starlingsroost.ddns.net](http://www.starlingsroost.ddns.net)

**Commented [CL41]: R2:** Lines 280-282: Are there any supportive figures for the statements given in Section 3.3.1 and Section 3.3.2 claiming the meteotsunami identification criteria are met. For example, is it possible to show this radar capture or data from barometric measurement or refer the reader to the source where this information is acquired? I recommend showing those relationships between the criteria and the mentioned examples of met criteria for the selected events.  
Editor: newly identified winter events

**Commented [CL42R41]: A:** Figures added as requested

323 **Table 1:** Descriptions and references for events that can be identified as UK meteotsunami events from 1750 to 2022. 1750 to 2009  
 324 are principally derived from historical sources and 2010 to 2022 are principally derived from instrumental data. [The threshold criteria  
 325 are outlined in the methodology section was used to verify previous (V) or identify new (N) events, including new winter events (NW).  
 326 (Wm represents maximum wave height in metres).

Date	Location	Wm	Time	Notes	Id criteria	Id status	Reference
1 Nov 1755	Ilfracombe	0.3	14:00	4 waves in 2 h, calm, NE wind, low tide, damage to sandbanks	1A, 1B, 2A, 3A	NW	Dawson et al 2000
27 Feb 1756	Ilfracombe	1.8	18:00	4 mins wave period, 30 mins duration, rumbling sea	1A, 1B, 2A, 3A	NW	Dawson et al 2000
31 May 1759	Lyme Regis	-	-	3 waves in 1 h, ebb and flow	1A, 2A, 3A	N	Dawson et al 2000
31 Mar 1761	Moums Bay	1.2	12:30	Ebb and flow 5 times in 1 h, NNE wind, cloudy, thunder	1A, 1C, 1C, 2A	NW	Long 2015
18 Sept 1763	Weymouth	3	-	3 waves, ebb, and flow	1A, 1B, 3A	N	<a href="http://www.phenomena.org.uk">www.phenomena.org.uk</a>
11 Feb 1764	Bristol	3	-	2 waves, ebb in 30 mins	1A, 2A, 3A	NW	<a href="http://www.phenomena.org.uk">www.phenomena.org.uk</a>
23 Dec 1791	Comwall	-	04:00	Rain, hail, extreme lightning, boats moved	2A, 3A	NW	Bortase 1758
17 July 1793	Plymouth	0.6	07:00	3 waves in 1 h, boats damaged	1A, 1B, 2A, 3A	N	-
18 Aug 1797	Lyme Regis	3	-	3 waves in 1 h, lightning	1A, 2A, 3A	N	Dawson et al 2000
9 Aug 1802	Devon	0.35	06:00	3 waves in 1 h, ebb and flow twice in 20 min	1A, 1C	N	Long 2015
10 Aug 1802	Teignmouth	0.6	08:00	10 min interval waves, fish on shoreline	1A, 1B	N	Long 2015
30 Aug 1802	Jersey	1.2	-	3 ebb and flows in 8 mins	1A, 1B, 2A	N	Long 2015
31 May 1811	Plymouth	2.4	03:00	4 h duration, rain, low pressure, ebb and flow, SW wind	1A, 1B, 2A, 2B	N	Dawson et al 2000
4 Mar 1818	Portsmouth	1.5	08:00	Rain, W to SW wind, high water for 3 h	1C, 2A, 2D	NW	<a href="http://www.surgewatch.org">www.surgewatch.org</a>
13 Sept 1821	Plymouth	1	14:00	Ebb and flow, boats moved	1A, 1B, 1C, 3A	N	Long 2015
13 July 1824	Plymouth	0.6	22:00	Ebb and flow, 4 m/s currents, ESE light wind, boats moved	1A, 1C, 2D, 3A	N	Archer, 2016
23 Nov 1824	Plymouth	2	01:00	3 waves in 10 min intervals, storm surge, 180 metres inland	1B, 1B, 2A, 3A	V	Haslett and Bryant 2009
5 July 1843	Plymouth	1	11:00	4 waves in 20 min, storm moved north, strong wind	1A, 1B, 1C, 2A	N	Thompson et al 2020
3 July 1845	Weymouth	0.6	10:30	Ebb and flow 5 times in 30 mins	1A, 2A, 3A	V	Long 2015
5 July 1846	Comwall	0.5	04:30	Thunder, severe storm reported	1C, 2A, 3A	V	Dawson et al 2000
1 Aug 1846	Penzance	0.3	04:00	30 min duration, calm sea, 6-minute ebb and flow	1C, 2A, 3A	V	Dawson et al 2000
23 May 1847	Penzance	0.9	05:00	20 mins, squally wind, sudden rush of water	1A, 1B, 2A, 2D	V	Long 2015
7 July 1848	Bristol	1.5	04:00	Thunder	1C, 2A, 3A	V	Edmonds 1862

**Commented [C143]: R1:** The authors need to clarify the original findings of this study which are different from the previous ones.  
**R2:** The number of investigated meteotsunami events is indicated as 95 in the abstract. How many of them are newly identified? Please indicate.

**Editor:** Address newly identified and winter events and originality of findings.

**Commented [C144R43]: A:** Table 1 has been adjusted to show new events and new winter events, new findings.

Date	Location	Wm (m)	Time (UTC)	Notes	ID criteria	Id Status	Reference
6 June 1855	Penzance	0.9	pm	Ebb & flow 2 to 3 times, rumbling sea, strong currents	IC, 2A, 3A	N	Dawson et al 2000
5 June 1858	Folkstone	0.9	08.00	Ebb & flow in 5 mins, ENE to WNW wind, hail, rain, seiche	IA, IC, 2A, 2D	V	Long 2015
25 June 1859	Comwall	0.3	Night	Abnormal sea oscillations x2, squall line, strong currents	IA, 2A, 3A	V	Dawson et al 2000
4 Oct 1859	Comwall	4.4	07.00	3 waves, 6 minute ebb, warm air temperatures, 1 mile upriver	IC, 2A, 3A	V	Dawson et al 2000
Oct 1865	Port Talbot	-	-	2 tides in 1 h	IA, 2A, 3A	NW	<a href="http://www.surgewatch.org">www.surgewatch.org</a>
23 Apr 1868	Lyme Regis	6	-	Swell, roar from the sea, no wind, low air pressure, calm sea	IC, 2A, 3A	V	Haslett & Bryant 2009
29 Sept 1869	Comwall	0.9	06.00	20 min wave period	IA, IB, IC, 2A	V	Dawson et al 2000
13 June 1881	Shetland	-	-	3 waves in 20 min, storm, boat damage	IA, 2A	V	Long 2015
28 Aug 1883	Plymouth	0.25	09.00	Gravity pressure wave from Krakatoa volcanic eruption	IB, 2B,	V	Garrett 1970
17 Oct 1883	Severn Est	1	08.00	1 dead, SW strong wind, high tide, precipitation, 1 mile in	IA, IC, 2A, 2D	V	Haslett & Bryant 2009
13 June 1886	Wick	0.45	16.30	Falling air pressure	IC, 2B, 3A	V	Long 2015
18 Aug 1892	Yealm	4	-	Quick ebb and flow, squall line, 3 waves, boat damage	IB, IC, 2A, 2B	V	Haslett & Bryant 2009
16 Dec 1910	Ilfracombe	4	06.15	Swell, bore, low pressure, 100 metre inland, bedrock erosion	IB, 2B	V	Haslett & Bryant 2009
26 Dec 1912	Isle of Wight	0.9	12.00	975 mb pressure low, SW wind, rain, cold front	IA, 2A, 2B, 2D	NW	<a href="http://www.surgewatch.org">www.surgewatch.org</a>
20 July 1929	Folkstone	6	19.30	8 waves, 180 metres inland, 5 mins period, low tide, 3 dead	IB, IC, 2A, 2D	V	Haslett & Bryant 2009
2 Aug 1932	Aberavon	9.3	-	4 dead, wave train, cloudy, rumbling sea, strong currents	IB, 2A, 3A	V	Haslett et al 2009
5 Aug 1938	Bridlington	4	08.00	Sea receded 4.5 m, boats moved, fish left on dry land	IA, IB, 2A	V	Haslett et al 2009
4 July 1939	Milford Haven	6	00.30	3 dead, rumbling sea, boats moved, mid tide	IB, 2A, 3A	V	Haslett & Bryant 2009
3 July 1946	Comwall	-	PM	Ebb and flow, squall line, rumbling sea, moorings broke	IA, 2A, 2B, 3A	V	Haslett & Bryant 2009
13 July 1949	Mevagissey	-	04.00	Easterly winds, boats smashed on rocks	IA, IC, 2A, 2D	V	Long 2015
6 July 1957	Bembridge	4	19.30	Wave train, 2 waves in 1 h, sultry, overcast, rocks moved	IA, IC, IC, 2A	V	Haslett & Bryant 2009
31 July 1966	Westward Ho	3	PM	Receding water, frontal trough, squall line	IA, IB, 2A	V	Haslett & Bryant 2009
1 July 1968	Folkstone	-	-	5 mb air pressure drop in 30 mins,	IA, 2B, 3A	V	Stevenson 1969
13 Feb 1979	Bristol	0.6	07.00	Spring tide, long unbroken waves, storm surge, 10 min period	IC, 2B	V	Haslett & Bryant 2009
28 May 2008	Peterhead	3	00.30	Ebb and flow in 10 mins, 4 to 6 waves	I A-C, 2 A-C	V	Sibley et al 2006
29 Jan 2010	Lowestoft	0.29	16.00	Open cell, S moving storm, 11 tide gauges	IA-C, 2A, 3A	V	Williams et al 2021
29 Aug 2010	Lowestoft	0.27	19.00	Open cell, S moving storm, 4 tide gauges	IA-C, 2A, 3A	V	Williams et al 2021



Date	Location	Wm (m)	Time (UTC)	Notes	Id criteria	Id Status	Reference
3 Feb 2011	Ullapool	0.3	22.00	Open cell, E moving, 7 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
27 June 2011	Devonport	0.3	08.30	Non-linear, N moving, 8 tide gauges plus European tide gauges	1A-C, 2A, 3A	V	Tappin et al 2013
22 Aug 2011	Newhaven	0.3	01.00	Quasi linear, N moving, 3 tide gauges, mid latitude depression	1A-C, 2A, 3A	V	Williams et al 2021
24 Nov 2011	Ullapool	0.26	04.30	Open cell, E moving, 8 tide gauges, mid latitude depression	1A-C, 2A, 3A	V	Williams et al 2021
3 Jan 2012	Lowestoft	0.33	17.15	Quasi linear, SE moving, 17 tide gauges, Low pressure	1A-C, 2A, 3A	V	Williams et al 2021
4 Feb 2013	Stornoway	0.32	07.00	Open cell, SE moving, 13 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
3 Aug 2013	Aberdeen	0.25	07.30	Non-linear cluster, NE moving, 9 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
28 Oct 2013	Devonport	0.27	03.15	Non-linear cluster, NE moving, 4 tide gauges, 1 mb/1 h drop, high	1A-C, 2A, 3A	V	Williams et al 2021
5 Dec 2013	Kinlochbervie	0.35	16.00	Quasi linear, 19 tide gauges, 1.7 mb/1 h drop, storm surge, spring tide	1A-C, 2A, 3A	V	Williams et al 2021
15 Dec 2013	Ullapool	0.25	18.00	Quasi linear, E moving, 6 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
18 Dec 2013	Millford Haven	0.33	19.00	Quasi linear, E moving, 24 tide gauges, 2.6 mb/1 h drop,	1A-C, 2A, 3A	V	Williams et al 2021
20 Dec 2013	Kinlochbervie	0.25	19.45	Quasi linear, NE moving, 5 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
21 Dec 2013	Ullapool	0.28	10.00	Individual cell, NE moving, 4 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
3 Jan 2014	Newlyn	0.33	12.30	Quasi linear, 8 tide gauges, 1.2 mb/1 h drop, high winds, high tide	1A-C, 2A, 3A	V	Williams et al 2021
8 Feb 2014	Weymouth	0.25	20.00	Open cell, E moving, 14 tide gauges, 1.3 mb/1 h drop at 18.30	1A-C, 2A, 3A	V	Williams et al 2021
12 Feb 2014	Weymouth	0.26	21.45	Quasi linear, E moving, 15 tide gauges, high winds, storm at 13.00	1A-C, 2A, 3A	V	Williams et al 2021
21 May 2014	Newhaven	0.26	23.00	Non-linear, N moving, 4 tide gauges, wave period 29 minutes	1A-C, 2A, 3A	V	Williams et al 2021
22 May 2014	Lerwick	0.33	06.45	Quasi linear, N moving, 3 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
1 Jan 2015	Ullapool	0.26	01.30	Open cell, E moving, 9 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
8 Jan 2015	Ullapool	0.27	01.00	Quasi linear, E moving, 10 tide gauges, wave period 15 minutes	1A-C, 2A, 3A	V	Williams et al 2021
1 July 2015	Stonehaven	0.25	09.00	Individual cell, NE moving,	1A-C, 2A, 3A	V	Sibley et al 2016
10 Dec 2015	Lerwick	0.31	23.00	Non-linear, NE moving,	1A-C, 2A, 3A	V	Williams et al 2021
27 Jan 2016	Workington	0.3	14.00	Non-linear, NE moving,	1A-C, 2A, 3A	V	Williams et al 2021
1 Feb 2016	Stornoway	0.27	16.30	Open cell, E moving, 11 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
23 June 2016	Newhaven	0.7	04.40	Non-linear, NE moving, 6 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021

Date	Location	Wm (m)	Time (UTC)	Notes	ID criteria	Event ID	Reference
26 Aug 2016	Devonport	0.3	22.45	Individual cell, NE moving, 7 tide gauges	1A-C, 2A, 3A	N	-
16 Nov 2016	Kinlochbervie	0.51	14.15	Open cell, E moving, 7 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
26 Dec 2016	Stomoway	0.34	08.30	Open cell, SE moving, 8 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
11 Jan 2017	Kinlochbervie	0.25	08.00	Open cell, SE moving	1A-C, 2A, 3A	V	Williams et al 2021
16 Oct 2017	Lerwick	0.35	16.00	Quasi linear, NE moving, 20 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
29 June 2019	Aberdeen	0.3	17.00	Non-linear, supercell moving from North Sea to Norway	1A-C, 2A-C	N	-
8 Feb 2020	Port Sloth	0.4	12.00	Line convection, ebb & flow, before storm Clara, Low pressure	1A-C, 2A, 2C	NW	-
21 Aug 2020	Perranporth	0.3	21.00	Spring tide, cold front, air pressure rise of 0.5 mb/2 min, bore	1C, 2B, 2C	N	-
5 July 2021	Westward Ho	0.6	12.40	S wind, Individual cell, mid tide, air pressure rise of 0.5 mb/1h, LP	1C, 2A-C, 3A	N	-
9 Aug 2021	Tonnes	0.25	11.30	S wind, Non-linear, mid tide, air pressure rise 0.5 mb/30 mins	1A, 1C, 2A-C	N	-
27 Sept 2021	Plymouth	0.32	03.00	S/SW wind, Quasi-linear, CAPE, low tide, air pressure rise 1.1 mb/20 mins	1A, 1C, 2A-D, 3A	NW	-
2 Oct 2021	Tomes	0.29	12.00	SSE wind, Non-linear, mid tide, air pressure fall 1.4 mb/1 h	1A, 1C, 2A, 2B	NW	-
20 Oct 2021	Plymouth	0.36	05.00	SSW, Non-linear, high tide, air pressure rise 1.5 mb/10 mins, CF	1A, 1C, 2A-C	NW	-
27 Nov 2021	Tomes	0.46	04.00	S/W, CAPE, mid tide, air pressure fall 1 mb/30 mins, storm surge,	1A, 1C, 2A-D	NW	-
30 Dec 2021	Tomes	0.6	00.00	S/W, non-linear, high tide, air pressure fall 0.5 mb/20 mins, Low pressure, wave period 20 minutes	1A, 1C, 2A-D, 3A	NW	-
16 Jan 2022	Port Isaac	0.3	01.00	Mid tide, air pressure fall of 1.5 mb, pressure wave from volcanic eruption	1A-C, 2B	NW	-
8 Feb 2022	Dunnet	0.3	13.15	Currents of 4 m/s, high tide, approaching cold front from north	1A, 1C, 2C, 3A	NW	-
18 June 2022	Newlyn	0.7	14.30	Spanish plume, 7+ locations, air pressure fall of 4 mb/10 mins	1A-C, 2B, 3A	N	-
19 July 2022	Anglesey	0.3	12.00	spring tide, air pressure fall 1 mb/35 mins, 5x ebb & flow, 9 m inland	1A-C, 2A-C, 3A	N	-
1 Nov 2022	Plymouth	0.3	09.00	Air pressure rise of 1 mb/5 mins, heavy ppt, thunder, SW wind	1A-C, 2A-D	NW	-
8 Nov 2022	Port Isaac	0.5	10.00	Air pressure rise 0.7 mb/2 mins, thunder, winds, precipitation	1A-C, 2A-D	NW	-
23 Nov 2022	Newlyn	0.7	09.00	Air pressure fall 1 mb/5 mins, cluster ppt, thunder,	1A-C, 2A-D	NW	-

330 **4 Discussion**

331 The aim of this paper was to introduce a revised, enhanced and current UK catalogue of meteotsunami events including a  
332 highlight of the seasonal occurrence, frequency, and spatial distribution of this hazard. This aim was set as there is no  
333 standardised identification criteria or up to date single catalogue of UK meteotsunami. This scenario has led to the mis  
334 conception that these events are non-hazardous, rare, and tend to occur more frequently in the summer months.

335

336 **4.1 The updated UK meteotsunami catalogue**

337 With the identification criteria we have laid out in this paper we have verified 98 events in UK waters since 1750 of which 38  
338 are new events containing seven new winter events in the historical record (1750 to 2009) and 8 new winter events in the  
339 modern record (2010 to 2022).

340 It was found that a selection of historical events were misidentified in accounts as either abnormal coastal flooding, non-  
341 tsunami, storm surge or of unknown origin. This was extended by an analysis of current data (since 2010) which allowed us  
342 to add a total of 38 new events to the catalogue, of which 15 occurred within winter months, these are highlighted in Table 1  
343 as new (N), new winter (NW) or verified (V) events.

344 The misidentified events were discovered after an attempt to highlight characteristics that match those listed in the  
345 methodology, in particular characteristics that suggested tsunami like phenomenon but with any associated storm like activity  
346 or air pressure fluctuations. If the account was found to contain a lack of evidence or information to suggest a meteotsunami  
347 it was rejected. An event occurring on 13 February 1979 was highlighted as a meteotsunami by Haslett et al. (2009a) but was  
348 contested by Thompson et al. (2020) as being a surge caused by a winter Atlantic storm due to its seasonal placement. In their  
349 2020 paper, Thompson et al appear not to class Atlantic storms systems as sources of meteotsunami. They state that from April  
350 to October, thunderstorms generate meteotsunami and from November to March, storms generate low pressure swells and  
351 surges. Our paper has matched descriptions in historical accounts with the criteria laid out in the methodology and we agree  
352 with Haslett and Bryant (2009a/b) that the 1979 winter event was a meteotsunami. This result was determined by the  
353 similarities in the pressure profile, geographical distribution and the speed of anomaly to the known meteotsunami event of 26  
354 June 2011.

355 In addition to the 1979 event, there were further events found that were previously labelled as meteotsunami to which our  
356 criteria have found them to be of alternative origin (tsunami) or to have insufficient detail or collaborative evidence to solidify  
357 a conclusion. These include the events presented in Long (2021), dated 14 October 1862 (found to be a tsunami due to an  
358 alternative source trigger), 15 August 1895 (insufficient information), 11 May 1912 (found to be a tidal bore) and another tidal  
359 bore dated 17 May 1964 presented in Haslett and Bryant, 2009a/b.

360 The event of the 31 March 1761 which was labelled as a tsunami by both Long (2015) and Thompson et al. (2020), was found  
361 to be a winter meteotsunami due to tsunami like waves being experienced not only along the southwest UK but also in Loch  
362 Ness in Scotland, with the mention of a calm sea before the arrival of thunderstorms.

363

**Commented [CL45]: R1, R2 and editor:** original findings and newly identified and winter events

**Commented [CL46R45]: A:** Included as requested

**Commented [CL47]: R2:** Lines 218-226: Here it is important to further explain the criteria used for the determination of those specific events as meteotsunami or not contrary to previous studies. How did the authors end up with these identifications for the mentioned events?

**Commented [CL48R47]: A:** Criteria and event selection determination explained, also highlights new events in the historical record

#### 364 4.2 Seasonal and geographical patterns of UK meteotsunami

365 The historical record (1750 to 2009) has been found to support previous studies such as Haslett and Bryant (2009a/b) that have  
366 alluded to the positive correlation between thunderstorms and meteotsunami waves with 71% of summer events displaying  
367 reports of convective activity. Our results have highlighted a summer prevalence of events with 48% of them peaking in July  
368 and August which reflects Thompson et al (2020). This prevalence has been based principally on the reliance on eyewitness  
369 reports and the volume of persons present at the shoreline during these months.

370 These summer events tend to be associated with heat waves and so called “Spanish plumes” as in the 27 June 2011 and the 18  
371 June 2022 events along the southwest UK. This is where warm air moves northwards from the European continent and Iberia,  
372 during which mesoscale convective weather tends to occur. In the summer, CAPE is at its highest and overland due to warm  
373 2 m air temperatures over landmasses (Holley et al. 2014). These types of weather event consist of single cell or clusters of  
374 small, short duration (< 1 hr) thunderstorms and squall lines with more than one convective cell (Sibley 2012 and Tappin et  
375 al. 2013).

376 The element of risk during the summer occurs when the meteotsunami wave can become fully disconnected from its source  
377 disturbance. This effect can be particularly apparent if the meteotsunami interacts with the continental slope where the wave  
378 can arrive hours after the original storm has dissipated or moved on (Greenspan 1956, Belche et al. 2016). This delayed arrival  
379 of wave disturbances can surprise people who are subsequently back out on or near the water’s edge, believing the storm has  
380 passed. This scenario was experienced during the 5 July 2021 event that occurred at Westward Ho (North Devon). Where just  
381 after midday a small yet powerful wave unexpectedly progressed 50 metres up the beach inundating many beach goers.

382  
383 Previous studies have suggested that winter wave anomalies such as meteotsunami are ‘less’ likely than storm waves, and  
384 surge, and winter data has not previously been interrogated for this reason. However, the present-day record (2010 to 2022)  
385 appears to contradict this with a winter prevalence of 66% of events peaking in December and January and with a tendency  
386 towards October and November in the 2021/22 winter season.

387 The results also show a geographical pattern to UK meteotsunami, with a large proportion of events occurring along the  
388 southwest UK and Northwest Scotland in the winter, aligning with the dominant weather direction of west to east from the  
389 Atlantic Ocean, and along the southern UK coast in the summer, aligning with Spanish Plumes bringing warm air poleward  
390 from the equator with southerly winds up and along the English Channel. The geographical pattern also reflects the influence  
391 of local bathymetry, with harbours (e.g., Penzance, Plymouth, Stornoway, and Port Talbot), bays (e.g., Kinlochbervie and Port  
392 Stoth) and river mouths (e.g., river Yealm and river Dart) containing conditions more favourable to meteotsunami initiation  
393 and amplification via resonance and seiching.

394 To further the concept presented in Williams et al. (2021) we selected two recent winter meteotsunami events and highlighted  
395 the meteotsunamigenic criteria. It has been indicated from the results that the combination of a mid-latitude depression, with  
396 frontal and convective weather moving across the UK may be important in the generation of this hazard. Results have shown  
397 that during these winter storms, convective elements are likely to be embedded around heavy rainfall (Figure 4a and b) and

**Commented [CL49]: R2:** One of the main findings is given in the abstract as “a prominent seasonal pattern of winter events” which is contrary to previous studies showing “a summer prevalence”. How do you explain this, especially referring to those previous studies? The only explanation for this is given by the reliance on eyewitness reports in the historical records period.

I would suggest including a more critical review of their findings, clearly highlighting the relationship to those referred (external) studies

Editor: Originality of findings from previous papers

**Commented [CL50R49]: A:** Our results vs previous papers in terms of similarities found.

**Commented [CL51]: R1:** L346 “This effect can be particularly apparent if the meteotsunami interacts with the continental slope where the wave can arrive hours after the original storm has dissipated or moved on.” I believe the authors mention Greenspan resurgence. Authors need to add references

**Commented [CL52R51]: A:** References added as requested

398 strong winds associated with the cold front leading to the potential for meteotsunami waves. This winter synoptic situation is  
399 a product of the combination of the cold maritime Arctic air being introduced to the rear side of the cold front passing over  
400 relatively warm water. The risk of flooding can be exacerbated due to surface water from precipitation as the front crosses a  
401 landmass (Masselink et al. 2015).

402 The results highlighted an average maximum wave height of 0.3 m which may not seem ‘dangerous’ but this hazard is not  
403 purely about this single factor. The key that makes meteotsunami a potential hazard is the rapid onset of the wave (sometimes  
404 referred to as a “wall of water”) and the associated strong currents.

405

#### 406 4.3 Constraints and Limitations

407 Identifying meteotsunami events in winter tends to be more difficult as the waves can be hidden and overshadowed by the  
408 wave characteristics of the trigger storms and may be missed unless looking specifically at the data. We strongly consider that  
409 this overshadowing means many of these winter meteotsunami do not get reported and this may have been the issue in previous  
410 research where certain winter events were identified as either storm waves or surges instead of meteotsunami. As we have seen  
411 there is a short observational record available for meteotsunami and there is evidence for severe under recording of such events.  
412 Even though the 2010 to 2022 record has shown significant improvements in recording completeness; the current 15-minute  
413 sampling interval is still too coarse. This was highlighted when certain events in the catalogue such as 2 October 2021, 20  
414 October 2021, 27 November 2021 and 19 July 2022 were uncovered in the 1-minute tide gauge data that were not so easy  
415 to locate in the 15-minute data. This creates an issue where many events with a wave period of under 15 minutes may be  
416 potentially missed. We recommend a reduction of the sampling interval to 1 to 5 minutes to yield more data to be able to draw  
417 a complete conclusion for this hazard.

418 Another limitation of this study linked to the sampling frequency was the treatment of wind-driven waves which can induce  
419 infra-gravity waves of a similar wave period to meteotsunami (2 to 5 minutes). We did initially consider wind and swell peak  
420 period and wave height; however, we discovered that the detection of infra-gravity waves from low frequency tide gauge data  
421 is uncertain and was deemed to be beyond the scope of this study. To perform such an analysis and to be confident in our  
422 results we would require 1 minute / 2 Hz data for a spectral analysis. However, it may be prudent to explore this aspect in  
423 future work.

424 We noted that historical accounts are not optimum for identifying and analysing meteotsunami due to their anecdotal nature  
425 and as such the number of events represented here may be dramatically underestimated. Data before 2008 is not readily  
426 available and records are spatially sparse which leads to incomplete data coverage and does not allow for a robust statistical  
427 analysis.

428 The placement of tide gauges used to provide data also affects results. The siting of UK tide gauges tend to be biased towards  
429 populated areas with harbours and river mouths for asset protection and is ideal for the capture of the resonant component of  
430 the meteotsunami wave. However, events in less populated areas may have been missed due to this placement. We suggest

**Commented [CL53]: R2:** What does this mean for the future?  
The first recommendation is a too general statement which is neither limited to meteotsunami hazard nor the UK region. It does not also represent a “new” finding. The necessity for the sea level data in the order of minute resolution for meteotsunami hazard has been emphasized in several studies such as Vilibic and Sepic (2017), Dusek et al. (2019), Williams et al. (2021), Zemunik et al. (2022). I believe that it is better to rewrite this recommendation by considering these issues.

**Commented [CL54R53]: A:** We agree to a point and although this may not be a ‘new’ finding it is extremely important in terms of UK meteotsunami, as we have found events in 1 minute tide gauge data that were not so easy to locate in the 15 minute sampling, proving that this is an issue for accurate cataloguing in the future.

**Commented [CL55]: R1:** It is unclear how the authors treated wind-driven waves. Wind-driven waves can induce infragravity (IG) waves which have periods of 2-30 minutes.  
**Editor:** Clearly address the treatment of wind driven waves or at least discuss them as a limitation of your study.

**Commented [CL56R55]: A:** We agree that this is a limitation of this study. It would be interesting to explore this aspect however we do not have the scope or the data to include it within this study. This may be an area for further research and a potential future manuscript as a follow on from ours.

431 potential tide gauge locations (based on the occurrence rate of previous events) could include beach or estuary locations around  
432 Devon and Cornwall such as Mevagissey or Perranporth and the North of Scotland such as Dunnet or Port Stoth.

433

#### 434 4.4 What does this mean for the future?

435 As the next few decades are likely to see sea level rise push mean and extreme water levels upward which will subsequently  
436 increase the level of risk by bringing the height of the storm tide closer to the flood stage (Masselink et al. 2015). At many UK  
437 locations, flood defences are at the design threshold of current storm surge levels, they are not designed or built for a sudden,  
438 prolonged water flow as seen in meteotsunami (Lazarus et al.2021). A question that has arisen from this paper is whether the  
439 winter seasons of 2013/14 and 2021/22 are outliers or whether this clustering of storms and meteotsunami will be a  
440 commonplace scenario in the future. Currently, we can detect and forecast mid latitude depressions nine to ten days in advance  
441 (Penn State, 2019), knowing this we can incorporate a warning of potential meteotsunami activity into the forecast. However,  
442 due to the localised nature of meteotsunami, risk level in each coastal areas needs to be considered on its own merits. The risks  
443 connected with a single meteotsunami event in two different bays can be quite different. One bay may suffer from inundation  
444 and flooding where another bay may be impacted by strong currents. This paper provides a valuable insight into the frequency,  
445 seasonality and spatial distribution of what was a hidden hazard in the UK. This new data will need to be incorporated and  
446 taken into consideration when coastal management strategies and defences are adjusted for the future.

447 Meteotsunami may well have some role to play in coastal storm impacts, however, the relative contribution of meteotsunami  
448 to storm surge in the aftermath of a storm and the full extent of the risk remains unknown and is beyond the scope of this work.  
449 It is also difficult to determine if the frequency and intensity of either low-pressure winter storms or winter meteotsunamis are  
450 on the increase. We invite a closer and more robust scrutiny of this hazard with a year-round perspective bearing in mind that  
451 no solid conclusions can be drawn without high frequency, long term, and continuous monitoring of this of hazard.

452

#### 453 5 Conclusions

454 Until recently it was thought that meteotsunami in the UK were rare and only occurred at certain times of the year, this  
455 misconception has led to a lack of provision in coastal management strategies and an under estimation of the frequency of this  
456 hazard. Motivated by coastal safety, this paper tests the hypothesis by presenting a new chronological catalogue dated from  
457 1750 to 2022 containing 98 UK meteotsunami with highlighted seasonal and geographical aspects. Using a standardised set of  
458 identification criteria developed for this study we have verified 60 previously listed events and presented 38 new events of  
459 which 15 were found to occur in the winter (Table 1).

460 Results demonstrate that meteotsunami are not restricted to the summer months and are more common than initially thought.

461 The modern record (2010 to 2022) is short and has far more winter meteotsunamis, whereas the relatively long historical record  
462 (1750 to 2009) means that the most meteotsunamis in our total have occurred in the summer which confirms the results of  
463 Thompson et al. (2020) and Haslett and Bryant (2009). During the summer months (April to September inclusive) there is a  
464 trend towards the southern UK with a 71% positive correlation between meteotsunami events and summer convective weather

Commented [CL57]: R1: L367 Then, it would be great if authors can suggest other locations for tide gauges.

Commented [CL58R57]: A: Potential tide gauge locations suggested with reasoning

465 systems, which can occur within synoptic Spanish Plume settings as suggested by Sibley (2012). During the winter months  
466 (October to March inclusive) our results demonstrate a clustering around the southwest UK and Northwest Scotland with a  
467 positive correlation between meteotsunami and the passage of mid latitude depressions where convective elements are  
468 embedded in the associated cold fronts and low pressure troughs. Subsequently meteotsunami impacts can become hidden by  
469 being superimposed on top of the storm's impacts. The meteotsunami waves are further exacerbated by the localised nature of  
470 resonance characteristics, in particular harbours and bays which can create highly dangerous situations. The immutable nature  
471 and rapid onset of this hazard means that even a sole meteotsunami event can create changes in water level and flow velocity  
472 that has the potential to cause injury, loss of life and damage to assets.

473 Increased knowledge of this hazard can be made more easily accessible through a central catalogue such as the one presented  
474 in this paper and the provision of higher frequency monitoring to detect future trends. What was thought to be a 'hidden' and  
475 rare event in historical records may soon become a common hazard in the future.

476

477 **Author contributions.** C. Lewis designed and executed the study and prepared the original draft. D. Williams pre-processed  
478 and provided data from 2010 to 2017, reviewed and edited the text. T. Smyth, J. Neumann, and H. Cloke supervised the project,  
479 provided advice, editing and feedback on the manuscript.

480

481 **Competing interests.** The authors declare that they have no conflict of interest.

482

483 **Data availability.** The datasets used in this study were derived from resources available in the public domain.

484

#### 485 **References.**

486 Bechle, A.J., Wu, C.H., Kristovich, D.A.R., Anderson, E.J., Schwab, D.J. and Rabinovich, A.B.: Meteotsunamis in the  
487 Laurentian Great Lakes. Scientific Reports 6, (37832). <https://doi.org/10.1007/s11069-014-1193-5>. 2016.

488

489 Borlase, W.: The natural history of Cornwall. Oxford. 53-4. <https://archive.org/details/naturalhistoryc00borl>. 1758.

490 British Oceanographic Data Centre: <https://www.bodc.ac.uk/> Last access: 19 February 2022.

491

492 Burt, S.: Multiple airwaves crossing Britain and Ireland following the eruption of Hunga Tonga-Hunga Ha'apai on 15

493 January 2022. Weather. Vol 77, No 3. <https://doi.org/10.1002/wea.4182>. 2022.

494

495 Chatfield, C.: Landmarks of world history: A Chronology of Remarkable Natural Phenomena Eighteenth Century 1761-1770.

496 <http://www.phenomena.org.uk/page29/page38/page38.html>. Last accessed 19 February 2022.

497

**Commented [CL59]: R2:** As I see, a major part of the findings of this study is discussed in Section 4.1 titled The UK meteotsunami. Therefore, it would be much better to provide a summary of the highlights of Section 4.1 in Section 5, Conclusions as well.

**Commented [CL60R59]: A:** Conclusions amended to focus on findings of section 4.1

498 Dawson, A.G., Musson, R.M.W., Foster, I.D.L., and Brunsden, D.: Abnormal historic sea-surface fluctuations, SW England  
499 marine Geology. Vol. 170, 59-68. [10.1016/S0025-3227\(00\)00065-7](https://doi.org/10.1016/S0025-3227(00)00065-7). 2020.  
500

501 Dusek, G., DiVeglio, C., Licate, L., Heilman, L., Kirk, K., Paternostro, C., & Miller, A.: A meteotsunami climatology along  
502 the U.S. East Coast. Bulletin of the American Meteorological Society, 100(7), 1329–1345. [https://doi.org/10.1175/BAMS-D-](https://doi.org/10.1175/BAMS-D-18-0206.1)  
503 [18-0206.1](https://doi.org/10.1175/BAMS-D-18-0206.1). 2019.  
504

505 Edmonds, R.: On extraordinary agitations of the sea not produced by winds or tides. Transactions of the Devonshire  
506 Association. 3. 144-152. <https://devonassoc.org.uk/publications/transactions/contents/>. 1869.  
507

508 Haigh, I., Wadey, M., Wahl, T.: Spatial and temporal analysis of extreme sea level and storm surge events around the coastline  
509 of the UK. Sci Data 3, 160107. <https://doi.org/10.1038/data.2016.107> 2016.  
510

511 Haslett, S.K. and Bryant, E.A.: Historic tsunamis in Britain since AD 1000: a review. Natural hazards and Earth Sciences. 8,  
512 587-601. <https://doi.org/10.5194/nhess-8-587-2008> 2008.  
513

514 Haslett, S.K. and Bryant, E.A.: Meteorological Tsunamis in Southern Britain: An Historical Review. Geographical Review.  
515 99, 146–163. <https://doi.org/10.1111/j.1931-0846.2009.tb00424.x> 2009a.  
516

517 Haslett, S.K., Mellor, H.E., and Bryant, E.A.: Meteo-tsunami hazard associated with summer thunderstorms in the United  
518 Kingdom. Physics and Chemistry of the Earth. 34, 1016-1022. <https://doi.org/10.1016/j.pce.2009.10.005> 2009b.  
519

520 Holley, D.M., Dorling, S.R., Steele, C.J. and Earl, N.: A climatology of convective available potential energy in Great  
521 Britain. International Journal of Climatology. 34: 14. 3811-3824. <https://doi.org/10.1002/joc.3976> 2014.  
522

523 Goring, D.: Meteotsunami resulting from the propagation of synoptic-scale weather systems. Physics and Chemistry of the  
524 Earth, Volume 34, Issue 17, p. 1009-1015. [10.1016/j.pce.2009.10.004](https://doi.org/10.1016/j.pce.2009.10.004) 2009.  
525

526 Kim, M., Woo, S., Eom, H. and You, S.: Pressure-forced meteotsunami occurrences in the eastern Yellow Sea over the past  
527 decade (2010–2019): monitoring guidelines. Natural Hazards and Earth System Sciences. [https://doi.org/10.5194/nhess-2021-](https://doi.org/10.5194/nhess-2021-126)  
528 [126](https://doi.org/10.5194/nhess-2021-126) 2021.  
529



530 Lazarus, E., Aldabet, S., Thompson, C., Hill, C., Nicholls, R., French, J., Brown, S., Tompkins, E., Haigh, I., Townend, I. and  
531 Penning-Rowsell, E.: The UK needs an open data portal dedicated to coastal flood and erosion hazard risk and resilience.  
532 *Anthropocene Coasts*. 4(1): 137-146. <https://doi.org/10.1139/anc-2020-0023> 2021.  
533  
534 Lightning maps: <https://www.lightningmaps.org/#m=oss;t=2;s=0;o=0;b=y=50.7086;x=-1.0547;z=4> last accessed 1 March  
535 2023.  
536  
537 Linares, Á., Wu, C.H., Bechle, A.J., Anderson, E.J. and Kristovich D.A.R.: Unexpected rip currents  
538 induced by a meteotsunami. *Sci Rep* 9:2105. <https://doi.org/10.1002/2016JCO11979> 2019.  
539  
540 Long, D.: A catalogue of tsunamis reported in the UK. British Geological Association 1R/15/043  
541 [https://nora.nerc.ac.uk/id/eprint/513298/1/IR\\_15\\_043%20%20BGS%20Tsunami%20catalogue%20update.pdf](https://nora.nerc.ac.uk/id/eprint/513298/1/IR_15_043%20%20BGS%20Tsunami%20catalogue%20update.pdf) 2015.  
542  
543 Long, D.: Comment on: Thompson et al 2020. UK meteotsunamis: a revision and update on events and their frequency.  
544 *Weather*. Vol 76. No4. 137-139. <https://doi.org/10.1144/sp456.10> 2021.  
545  
546 Lynett, P.J., Borrero, J., Son, S., Wilson, R. and Miller, K.: Assessment of the tsunami induced current hazard. *Geophysical*  
547 *Res Lett* 41 (6): 2048-2055. <https://doi.org/10.1002/2013GL058680> 2014.  
548  
549 Masselink, G., Scott, T., Poate, T., Russell, P., Davidson, M. and Conley, D.: The extreme 2013/2014 winter storms:  
550 hydrodynamic forcing and coastal response along the southwest coast of England. *Earth Surface Processes and Landforms*,  
551 volume 41, Issue 3 378-391. <https://doi.org/10.1002/esp.3836> 2015.  
552  
553 MET Office: 5 km Resolution UK Composite Rainfall Data from the Met Office Nimrod System.  
554 <https://catalogue.ceda.ac.uk/uuid/f91b2c5399c5bf689e29bb15ab45da8a> 2003.  
555  
556 Monserrat, S., Vilibic, I. and Rabinovich, A.B.: Meteotsunamis: atmospherically induced destructive ocean waves in the  
557 tsunami frequency band. *Natural Hazards and Earth System Science*. 6. 1035-1051. <https://doi.org/10.5194/nhess-6-1035-2006>  
558 2006.  
559  
560 National Oceanography Centre: <https://www.bodc.ac.uk/> Last accessed 1 March 2023.  
561  
562 National tide and sea level facility: Available at: <https://ntslf.org/> Last accessed 19 February 2022.  
563

564 Pattiaratchi, C.B. and Wijeratne, E.M.S.: Are meteotsunamis an underrated hazard? Philosophical Transactions of the Royal  
565 Society: Mathematical and Engineering Sciences 373. <https://doi.org/10.1007/s11069-014-1263-8> 2015.  
566

567 Pellikka, H., Laurila, T.K., Boman, H., Karjalainen, A., Björkqvist, J. and Kahma, K.K. Meteotsunami occurrence in the Gulf  
568 of Finland over the past century, Natural Hazards Earth System Sciences. 0. (9). 2535-2546. [https://doi.org/10.5194/nhess-20-  
569 2535-2020](https://doi.org/10.5194/nhess-20-2535-2020) 2020.  
570

571 Penn State.: Predictability limit: Scientists find bounds of weather forecasting. ScienceDaily. 15 April 2019.  
572 <https://sciencedaily.com/releases/2019/04/190415154722.htm> Last accessed 20 August 2022.  
573

574 Proudman, F.R.S.: The Effects on the Sea of Changes in Atmospheric Pressure. Geophysical Journal International 2 s4.  
575 <https://doi.org/10.1111/j.1365-246X.1929.tb05408.x> 1929.  
576

577 Reigate grammar weather station: The birth and impact of the St Jude day storm: October 2013. Available at:  
578 <https://rgsweather.com/2013/10/29/st-jude-causes-and-impacts-of-the-october-storm-27-28-2013/amp/> Last accessed 19  
579 February 2022.  
580

581 Sea Level Monitoring Facility: <https://ioc-sealevelmonitoring.org/> Last accessed 1 March 2023.  
582

583 Šepić, J., Vilibić, I. and Mahović, N.: Northern Adriatic meteorological tsunamis: observations, link to the atmosphere, and  
584 predictability. Journal of Geophysical Research Oceans. 117(C2). <https://doi.org/10.1029/2011JC007608> 2012.  
585

586 Šepić, J., Vilibić, I., Rabinovich, A. et al. Widespread tsunami-like waves of 23-27 June in the Mediterranean and Black  
587 Seas generated by high-altitude atmospheric forcing. Sci Rep 5, 11682. <https://doi.org/10.1038/srep11682> 2015.  
588

589 Šepić, J., Vilibić, I., Rabinovich, A. et al. Meteotsunami (“Marrobbio”) of 25–26 June 2014 on the Southwestern Coast of  
590 Sicily, Italy. Pure Appl. Geophysics. 175, 1573–1593. <https://doi.org/10.1007/s00024-018-1827-8> 2018.  
591

592 Sibley, A.: Thunderstorms from a Spanish Plume event on 28 June 2011. Weather. 67. No 6. 143-152.  
593 <https://doi:10.1002/wea.1928>. 2012.  
594

595 Sibley, A., Cox, D., Long, D., Tappin, D.R. and Horsburgh, K.J.: Meteorologically generated tsunami like waves in the North  
596 Sea on 1 July 2015 and 28 May 2008. Weather. 71. 68-74. <https://doi.org/10.1002/wea.2696> 2016.  
597

598 Starlings roost weather: <http://starlingsroost.ddns.net/weather/stats.php?type=day&field=&date=2023-03-05> Last accessed 1  
599 March 2023.  
600  
601 Stevenson, C. M.: The dust fall and severe storms of 1 July 1968. *Weather*. 66 (5): 125–127. DOI:[10.1002/WEA.780](https://doi.org/10.1002/WEA.780) 1969.  
602  
603 Surge Watch database: A database of UK coastal flood events. <https://www.surgewatch.org/> Last accessed 19 February 2022.  
604  
605 Tappin, D.R., Sibley, A., Horsburgh, K.J., Daubord, C., Cox, D. and Long, D.: The English Channel ‘tsunami’ of 27 June 2011  
606 - a probable meteorological source. *Weather*. 68. 144–152. <https://doi.org/10.1002/wea.2061> 2013.  
607  
608 Thompson, J., Renzi, E., Sibley, A. and Tappin, D.: UK meteotsunamis: a revision and update on events and their frequency.  
609 *Weather*. 75.9, 281–287. <https://doi.org/10.1002/wea.3741> 2020.  
610  
611 University of Wyoming: <http://weather.uwyo.edu/upperair/sounding.html> Last accessed 19 February 2022.  
612  
613 Ventusky weather: <https://www.ventusky.com/?p=54.4;-5.9;4&l=radar&t=20230313/1100&w=off> Last accessed 1 March  
614 2023.  
615  
616 Vilibić, I. and Šepić, J.: Global mapping of non-seismic sea level oscillations at tsunami timescales. *Scientific reports*. 7. (1)  
617 40818. <https://doi.org/10.1038/srep40818> 2017.  
618  
619 Vilibić, I., Šepić, J., Dunic, N., Sevault, F., Monserrat, S. and Jorda, G.: Proxy-based Assessment of Strength and Frequency  
620 of Meteotsunamis in Future Climate. *Geophysical Research Letters*. 45. 10501-10508.  
621 <https://doi.org/10.1029/2018GL079566> 2018.  
622  
623 Williams, D.A., Horsburgh, K.J., Schultz, D.M. and Hughes, C.W.: Examination of generation mechanisms for an English  
624 Channel Meteotsunami: Combining observations and modelling. *Journal of Physical Oceanography*. 49. 103-120.  
625 <https://doi.org/10.1175/JPO-D-18-0161.1> 2019.  
626  
627 Williams, D. A., Schultz, D. M., Horsburgh, K. J., & Hughes, C. W.: An 8-yr meteotsunami climatology  
628 across northwest Europe: 2010–2017. *Journal of physical oceanography*. 1145-1160. [https://doi.org/10.1175/JPO-D-20-](https://doi.org/10.1175/JPO-D-20-0175.1)  
629 [0175.1](https://doi.org/10.1175/JPO-D-20-0175.1) 2021.  
630