

# 1 **Meteotsunami in the United Kingdom: The hidden hazard.**

2 Clare Lewis<sup>1 2</sup>, Tim Smyth<sup>2</sup>, David Williams<sup>4</sup>, Jess Neumann<sup>1</sup>, Hannah Cloke<sup>1 3</sup>

3  
4 <sup>1</sup> Department of Geography & Environmental Science, University of Reading, Reading, UK

5 <sup>2</sup> Plymouth Marine Laboratory, Prospect Place, Plymouth, Devon, PL1 3DH, UK

6 <sup>3</sup> Department of Meteorology, University of Reading, Reading, UK

7 <sup>4</sup> WTW, 51 Lime Street, London, EC3M 7DQ, UK

8 Correspondence to: Clare Lewis (clare.lewis@pgr.reading.ac.uk)

## 9 **Abstract.**

10 This paper examined the occurrence and seasonality of meteotsunami in the United Kingdom (UK) to present a  
11 revised and updated catalogue of events occurring since 1750. Previous case studies have alluded to a summer  
12 prevalence and rarity of this hazard in the UK. We have verified and classified 98 events using a developed set of  
13 identification criteria. The results have revealed a prominent seasonal pattern of winter events which are related  
14 to mid latitude depressions with precipitating convective weather systems. A geographical pattern has also  
15 emerged, highlighting three ‘hotspot’ areas at the highest risk from meteotsunami. The evidence reviewed, and  
16 new data presented here shows that the hazard posed by meteotsunami has been underestimated in the UK.

17  
18 Keywords: meteotsunami, UK, hazard, mid latitude depressions.

## 19 20 **1 Introduction**

21 Meteotsunamis or meteorological tsunamis are globally occurring progressive shallow water waves with a period  
22 of between 2 to 120 minutes which results from an air-sea interaction. They tend to be initiated by sudden pressure  
23 changes and wind stress from moving atmospheric systems with sources ranging from convective clouds,  
24 cyclones, squalls, thunderstorms, atmospheric gravity waves and strong mid-tropospheric winds (Vilibic and  
25 Sepic, 2017). The atmospheric pressure changes are typically only a few mb over a few tens of minutes which  
26 corresponds to only a few centimetres of sea level change occurring in a process known as the inverse barometer  
27 effect (for example, a 3 mb pressure jump will produce a 30 cm ocean wave). The atmospheric disturbance  
28 transfers energy into the ocean initiating and amplifying a water wave which travels at the same speed as the  
29 atmospheric wave, in a process known as Proudman resonance (Proudman, 1929). When the water wave reaches  
30 the coastline and shallower water, it becomes a multi resonant phenomena and is further amplified through coastal  
31 resonances. For example, if the wave reaches the entrance of a semi enclosed basin it can induce an oscillation in

32 the basin known as harbour resonance. However, if the wave reaches a beach type environment and the along  
33 shore component of the disturbance equals the phase speed of the edge wave this is a process known as Greenspan  
34 resonance (Monserrat et al. 2006). The resultant waves can elevate the coastal water level and can substantially  
35 increase flow velocities with the potential for rip currents (Linares et al. 2019). Due to the rapid onset and  
36 unexpected nature of meteotsunami waves, they have the potential to cause destruction, injuries and even fatalities  
37 (Sibley et al. 2016). For a global perspective and overview of meteotsunami observations we recommend Pellikka  
38 et al. (2020) for observations in Finland, Sepic et al. (2018) for the Adriatic, Belche et al. (2016) for seasonality  
39 of meteotsunami in the Great Lakes, Pattiaratchi and Wijeratne (2016) for observations in southwest Australia  
40 and Monserrat, Vilibic and Rabinovich (2006) provide a general overview of the mechanisms of meteotsunami.

41

42 Meteotsunami research and monitoring is more advanced in the Mediterranean, the East Coast of the USA, and  
43 the Great Lakes due to the higher number of recorded events. However, events in the UK appear to be rare and  
44 are believed to be less devastating, meaning that research has been limited to date.

45 The two principal factors contributing to this belief are:

46 1. The current (since 1993) 15-minute sampling interval that is used at UK tide gauges is incapable  
47 of detecting waves with periods of between 2 – 120 minutes. This means that many events go  
48 unobserved, wave heights are underestimated, or meteotsunamis are mischaracterised as seiches,  
49 tsunamis or surge.

50 2. Until recently research has suggested that UK meteotsunamis are generated by precipitating,  
51 convective weather systems associated with hot weather. Such mesoscale convective systems may  
52 be associated with synoptic “Spanish plume” events. These synoptic events are more prevalent  
53 between May - October (Haslett et al. 2009b; Tappin et al. 2013; Sibley, 2012 and 2016; Thompson,  
54 2020), leading to the belief that meteotsunami occurrence is a summer-time phenomena. However,  
55 it is now emerging that embedded convection within winter frontal systems may also be responsible  
56 for a sizeable proportion of these waves (Williams et al. 2021).

57 Several issues have resulted from the untested assumption that meteotsunami events are 1) low frequency and 2)  
58 predominantly occur in summer, which has been combined with 3) the lack of high-resolution temporal data.  
59 Firstly, there is no central database of UK events. Secondly, there is no standardised methodology of meteotsunami  
60 identification. Thirdly, there is no Government or regional policy in place to cover impacts from a meteotsunami  
61 event. There is a misconception of the risk posed by meteotsunami especially for coastal areas that are already at  
62 risk from storm impacts associated with pluvial (extreme precipitation) and fluvial hazards (high levels of river

63 discharge). In the future the overall level of risk is likely to be greatly exacerbated by rising sea levels and an  
64 intensification of storm frequency and severity (Vilibic et al. 2018; Masselink et al. 2015).

65 As stated by Sepic et al. (2015) the assessment of meteotsunami should become the standard in coastal hazard  
66 assessments, event cataloguing is a pre-requisite for any coastal hazard assessment especially in identifying the  
67 geographical areas that have experienced meteotsunami and the frequency of exposure.

68 The aim of this paper is to compile, update and extend the existing list UK meteotsunami to include winter events,  
69 and to highlight the occurrence, frequency, and spatial distribution of events. Where seasonality was alluded to in  
70 Williams et al. (2021), their study was principally focused on meteotsunami in Northwest Europe from 2010 to  
71 2017. This paper will further their study by focussing in on UK waters only and will add new events up to the end  
72 of 2022. The methodology fulfils this aim by applying a set of developed identification criteria to the re-  
73 assessment of fragmented historical accounts and to the analysis of tide gauge and atmospheric data to identify  
74 new events. The outcome also highlights the potential element of winter compound hazard risk which may occur  
75 when meteotsunami waves arrive at the coast in short succession or concurrently with other storm associated  
76 hazards.

77 We propose the following research questions:

- 78 1. What standardised criteria should be used to identify meteotsunami?
- 79 2. Have events occurred which were ignored or misidentified?
- 80 3. In which regions of the UK and in what months do meteotsunami occur most frequently?
- 81 4. What are the atmospheric variables that can be correlated with meteotsunami events?

82

## 83 **2 Methodology**

84 This section outlines the data sources and identification criteria used to fulfil the objective of cataloguing and  
85 characterising UK meteotsunami. We have extrapolated as much quantitative data as possible, to verify the event  
86 with the standardised criteria and to then arrange the results into tabular form to allow ease of use (Table 1).

87

### 88 **2.1 Meteotsunami identification criteria**

89 As there are currently no fixed criteria for what qualifies as a meteotsunami, in this paper we bring together various  
90 aspects used by other researchers in the field, into one standardised system. Figure 1 (a – d) displays a visual  
91 representation of the commonly used criteria, which we explain in more detail in sections 2.1.1 – 2.1.2. The  
92 methodologies that have been previously used by researchers and studies have variations, with some using  
93 qualitative methods that base events on eyewitness accounts (Haslett et al, 2009a/b) and others using quantitative

94 data from sea level and atmospheric observations (Tappin et al. 2013; Sibley, 2016). For the purpose of this paper,  
95 we have classified meteotsunami as atmospherically induced sea level oscillations meeting at least one sea level  
96 and one atmospheric characteristic. This allows for the distinguishing of meteotsunami from other types of  
97 waveforms and is applicable to either qualitative accounts or quantitative data.

98

### 99 **2.1.1 Sea level criteria (Category 1)**

- 100 a. Periods of sea level disturbance ranging from between 2 and 120 minutes (Figure 1a).
- 101 b. Wave heights exceeding 0.20 m. This threshold is within the peak thresholds of 0.2 m and 0.3  
102 m as used by other researchers in the field such as Williams et al. (2021); Dusek et al. (2019); Belche  
103 et al. (2016); Sepic et al. (2012) and Monserrat, Vilibic and Rabinovich (2006). A 0.3 m water  
104 elevation may not appear to be dangerous, but a meteotsunami in 2003 in New Zealand caused a  
105 fully laden oil tanker to be grounded through strong currents (Goring, 2009). Lynett et al. (2014)  
106 also states that any wave over 0.3 m will start to float vehicles regardless of flow velocity and is  
107 enough to sweep people off of their feet. These thresholds are a tried and tested set of characteristics  
108 that reflect meteotsunami especially those in UK waters. 0.2 m was opted to be used as the lower  
109 end of the threshold as this is more suitable for distinguishing a greater number of events that may  
110 have been missed at the higher end of the threshold (0.3m). Any anomaly below 0.2m would not be  
111 large enough to allow for accurate verification and for its separation from any other water  
112 disturbances. (Figure 1a illustrates the meteotsunami wave height criteria in the data as recorded on  
113 27 June 2011).
- 114 c. A wave disturbance registering at two or more locations or tide gauge stations (Williams et al.  
115 2021; Kim et al. 2021).

116

### 117 **2.1.2 Atmospheric criteria (Category 2)**

- 118 a. The presence of a convective weather system at the time of the wave event displaying high radar  
119 reflectivity with precipitation rates exceeding  $2 \text{ mm/h}^{-1}$ , initiated over the sea. (Figure 1b represents  
120 the radar reflectivity of the various convective weather systems present during four different  
121 meteotsunami events).
- 122 b. An atmospheric pressure of 1005 mb or less with a rapid change of  $\pm 1 \text{ mb}$  in 30 minutes or a 3  
123 mb fall over three hours or less (Monserrat, Vilibic and Rabinovich, 2006). (Figure 1c illustrates this  
124 distinct air pressure change as recorded during the 28 October 2013 event).

125 c. Convective Available Potential Energy (CAPE) showing the unstable vertical profile of the  
126 atmosphere that leads to convective activity (Williams et al. 2019). (Figure 1d displays a radiosonde  
127 ascent showing sufficient CAPE to produce the event that occurred on 1 July 2015 at Stonehaven,  
128 East Scotland. Even though CAPE is a bulk atmospheric measurement and meteotsunami are  
129 localised, if this element is present in conjunction with the other indicators it supports the presence  
130 of convective activity which aids in the generation of meteotsunami.

131 d. A change in wind speed exceeding  $10 \text{ m/s}^{-1}$  (anything under this is too weak for a meteotsunami  
132 to generate) or/and a drop in air temperature of  $1.5^\circ\text{C}$  in 30 minutes (Figure 1c demonstrates this  
133 increase in wind speed as recorded during the 28 October 2013 event).

### 134 135 **2.1.3 Geological criteria (Category 3)**

136 a. The absence of any other explanation or data to imply an alternative source trigger. For example,  
137 the presence of seismic triggers within the continental shelf area which would produce a geological  
138 tsunami wave. However, there is one exception to this rule which for the purpose of this paper we  
139 include as a meteotsunami trigger. Volcanic eruptions, this was demonstrated on 28 August 1883  
140 (Krakatoa) and recently on 16 January 2021 (Tonga Ha'apai) where wave anomalies occurred and  
141 were the product of air pressure waves created by the eruptions. It may be argued that they are not  
142 to be classed as meteotsunami waves. However, for the purpose of this catalogue, we are classifying  
143 them as meteotsunami as they are sourced from air pressure disturbances which couple with water  
144 waves and have a wave period of 2 to 120 minutes. The force of the Tonga Ha'apai explosions sent  
145 a shockwave through the atmosphere that circled the globe three times. The resultant pressure  
146 wave travelled at close to the speed of sound and as a result coupled with ocean waves to create a  
147 meteotsunami which was detected as far away as Portugal and the UK (Burt. S, 2022).

148  
149 To ease the interpretation of results, the UK coastline has been partitioned into six coastal regions based on the  
150 National Tidal and Sea Level Facility (NTSLF) tide gauge network (Supplementary Table S1). The data are also  
151 separated into two seasons (each comprising of six months) that divide up the calendar year at the spring and  
152 autumn equinoxes (Haigh et al. 2016). April to September inclusive is referred to throughout this paper as  
153 'summer' and October to March is referred to as 'winter'. Finally, due to the nature of the data, two time series of  
154 meteotsunami are being referred to throughout this paper, one based primarily on historical eyewitness accounts

155 due to a lack of high frequency instrumentation (the years 1750 to 2009 AD), and one based on and verified by  
156 quantitative instrumental data (the years 2010 to 2022 AD).

157

## 158 **2.2 Historical record (1750 to 2009)**

159 To gain a complete understanding of these events we follow Long (2015) and Haslett and Bryant (2008) who  
160 dated their historic tsunami catalogues back to approximately 1000 AD. We noted that any events preceding 1750  
161 AD were vaguely recorded, making validation problematic so we opted to date our catalogue back to 1750.  
162 References to meteotsunami like events in historical accounts tend to be based on descriptions of the state of the  
163 water at the coast with a lack of instrumental tidal data. There is a lack of or limited weather data so tracing back  
164 the atmospheric source is not as straightforward. It is only until the last few decades that meteorological data with  
165 sufficient resolution have been readily available. With tide gauge data, prior to 1993 the resolution was hourly,  
166 and it was not until 1996 that all the current tide gauge sites became fully operational. Therefore, we have used  
167 2009 as the upper limit of the historical record. The historical reports tend to be derived from newspaper articles,  
168 parish records, harbourmaster records and eyewitness accounts. Although there is reason to be sceptical of these  
169 accounts as they afford a level of biased review and sensationalism, they do still hold value in terms of a societal  
170 viewpoint and may help to fill in any gaps (Haslett and Bryant, 2009a/b).

171 There are certain characteristics that flag up in an historical account to verify whether it is a meteotsunami event  
172 or not. To illustrate this, we can highlight the historical account for the event of 23 May 1847 where we can look  
173 at a letter from Robert Blight of Penzance dated 24 May and published in the Cornwall Royal Gazette on 28 May.  
174 The full extract can be found in supplementary extract S1 of this paper and in Long (2015, p26).

175 “... The changes in the atmosphere during the day were very remarkable. In the morning, about six o’clock, we  
176 had a breeze from the southeast; by eight, it was a perfect calm; between ten o’clock and two, the mercury sunk  
177 several degrees; about three in the afternoon a breeze sprung up suddenly from the west, and the sky, as suddenly,  
178 became overcast..... It is very probable that all these changes, and even the agitation of the sea, were produced  
179 by electricity...”

180 In this particularly detailed account (supplementary extract S1) we can identify six of the nine criteria, including  
181 a drawback and sudden in rush of water, accompanied by a rumbling noise and the water being higher than  
182 expected at eight feet (criteria 1A and 1D), all indicating a tsunami like event. The key to the identification of a  
183 meteotsunami is in the atmospheric portion of the account, what started out as calm morning led to a change in  
184 wind speed and direction, veering from south easterly in the morning to westerly in the afternoon (criteria 2D).  
185 This variable wind was accompanied by a drop in temperature (criteria 2D) and finally, there was mention of the

186 presence of a storm in terms of overcast sky, threatening rain and lightning (criteria 2A). As such, we identify this  
187 wave as a meteotsunami by applying both of our oceanographic and atmospheric criteria to the historic account.  
188

### 189 **2.3 Tide gauge analysis for the 2010 to 2022 record**

190 To identify meteotsunami from 1 January 2010 to 31 December 2022 we use data records that are available at  
191 higher frequencies meaning meteotsunami are more distinctly observable. The information for this portion of the  
192 catalogue is sourced from the British Oceanographic data centre (BODC) website (<https://www.bodc.ac.uk/>) and  
193 the International Oceanographic Commission (IOC) website (<https://ioc-sealevelmonitoring.org/>) where data are  
194 displayed from the ‘Class A’ network of tide gauges owned and funded by the Environment Agency (EA). We  
195 also use the postprocessed data of Williams et al. (2021) where the raw sea level tide gauge data has been high  
196 pass filtered to isolate high frequency disturbances. This removes periods of over 120 minutes and separates out  
197 the tidal components. In this way any signals in the tsunami frequency band (2 to 120 minutes) are isolated from  
198 the sea level elevations. Any remaining signals larger than the background noise are then identified and checked  
199 against our threshold criteria to verify events as potential meteotsunami. Apart from the standard processing to  
200 remove any erroneous spikes outside of the parameters, a visual quality control was carried out, where a seven-  
201 day plot of the data was evaluated to highlight any clear artificial spikes or gaps. Also, any data points that had  
202 no accompanying air pressure changes were also excluded from any further analysis.  
203


### 204 **2.4 Atmospheric data analysis for the 2010 to 2022 record**

205 The time of the potential meteotsunami events are noted from the tide gauge data and they are then linked to  
206 specific precipitating convective atmospheric systems by using the meteorological C-band radar network, which  
207 is pre-processed by the UK Meteorological Office before download (Met Office 2003). The convective systems  
208 highlighted by the radar are classified into four distinct types (as shown in Figure 1b). These are: (1) open cells  
209 which are situated behind the cold front of cyclonic weather, usually where cold dry air passes over the warm sea  
210 creating shallow convection; (2) Quasi linear systems which tend to be multi-cellular and linearly organised with  
211 high CAPE, heavy precipitation, and strong winds (this type of weather feature are sometimes called squall lines  
212 and can occur within synoptic Spanish Plume events); (3) Isolated small short duration (<1h) thunderstorm cells  
213 and (4) Nonlinear clusters which are large circular, long lived clusters of precipitation and thunderstorm cells.


214 The atmospheric ascent soundings are obtained from the University of Wyoming website  
215 (<http://weather.uwyo.edu/upperair/sounding.html>) with the UK stations at Camborne (station number: 03808) and  
216 Lerwick (station number: 03005) being used. Soundings are available for 0000 UTC and 1200 UTC on each day

217 and if a CAPE value of greater than 0 occurs then this shows a marginally unstable atmosphere leading to  
 218 convective activity. Finally, the synoptic charts allow for verification of the storm system including the location  
 219 of the pressure centres and fronts at the time of the meteotsunami wave event.


220  
 221




**Figure 1a:** Devonport (50°36N 4°18W) tide gauge for 27 June 2011 showing a distinct sea level disturbance at 0830 UTC as highlighted with a red circle. This is a representation of criteria 1b. The timing of this 0.25 m rise and fall in the sea level corresponds with the arrival of the meteotsunami event at that specific location.



**Figure 1b:** The four different types of convective activity as shown on radar reflectivity identifying meteotsunami events. A representation of criteria 2a. Orange and red in the images shows high precipitation rates ( $>4 \text{ mm h}^{-1}$ ). With idealised images shown on the left and actual examples taken from UK events on the right. All showing date, time, and direction of the storm as well as the location of the tide gauges (white dots). Image by David Williams, Journal of Physical Oceanography (<https://doi.org/10.1175/JPO-D-20-0175.1>), licensed under a [Creative Commons — Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) — CC BY 4.0



**Figure 1c:** The atmospheric pressure, wind speed and precipitation at Reigate (51°14N 0°11W) during the 27 to 28<sup>th</sup> October 2013 storm associated with the meteotsunami. A representation of criteria 2b and 2d. The graph shows atmospheric pressure (red line) of less than 1005 mb and falling as the atmospheric disturbance moves over the area, with a corresponding rising wind speed of 20 mph (green line) and precipitation (blue bars). Reproduced with the kind permission of Simon Collins. <https://rgsweather.com/2013/10/29/st-jude-causes-and-impacts-of-the-october-storm-27-28-2013/amp/>



**Figure 1d:** The Nottingham radiosonde ascent at 1200 UTC on 1 July 2015 during a meteotsunami event in the North Sea. A representation of criteria 2c which indicates sufficient CAPE (462.1 J Kg) to produce high base convective activity, with the cloud base at an approximate height of 3000 m and cloud top at 11000m. <http://weather.uwyo.edu/upperair/sounding.html>

222  
 223  
 224  
 225

### 226 3 Results

227 In this section we highlight the seasonal occurrence and distribution of UK meteotsunami events in both the  
 228 historical record and the more recent instrumental data record. This is augmented by the identification of trigger



229 systems associated with the events where available. It is prudent to note here that the catalogue cannot be  
230 considered as complete, and this is signified by dashed lines (i.e., -) in the columns where data or information are  
231 either unavailable or have not been located.

232

### 233 **3.1 Historical record (1750 to 2009)**

234 We identify 98 events as being meteotsunami occurring in UK waters between January 1750 and December 2022  
235 (Table 1), with 48 of these occurring within the historical record (1750 to 2009). This record shows that 67% of  
236 documented meteotsunamis occur in summer (April – September), with 44% of documented meteotsunamis  
237 occurring in July and August. The single year experiencing the most documented events was 1802 AD, numbering  
238 three, and the decade experiencing the most documented events was the 1840s, with six in total. The presence of  
239 a storm and/or characteristics of convective activity (thunder, and lightning) at the time of the wave event was  
240 noted for 42 of the 48 events (91%) in the historical record. There was also a defined southwest prevalence of  
241 meteotsunami in historical documents, with Devon, Cornwall and Somerset recording a combined total of 29  
242 events. Within the historical record we have identified four new events and reclassified four tsunamis, three storm  
243 surge and nine events of unknown origin as meteotsunami. Seven of these occurred within winter months (Table  
244 1).


245

### 246 **3.2 Seasonal and locational frequency of UK meteotsunami events (2010 to 2022)**

247 Meteotsunamis have been thought to be a rare phenomenon in the UK and that when they do occur, it has been  
248 tended to be in the summer months due to the more abundant convective activity (Haslett et al, 2009b; Tappin et  
249 al, 2013; Sibley, 2016; Thompson, 2020). However, of the 98 identified meteotsunami events verified in this  
250 paper, 50 have been interpreted as occurring since 2010, 33 (66%) of those occurring during the winter months  
251 and nine of the winter events are identified as new. We find that not only are UK meteotsunami more common in  
252 occurrence than previous research indicates, but that they are a year-round phenomenon, as exhibited in Table 1  
253 and Figures 2 and 3.


254 The historical section of the catalogue shows an estimated return period of 5.4 years. This return period  
255 considerably decreases for the instrumental data section where the UK return period reduces to an estimated 0.25  
256 years. With an average of four events per year, we can see that certain years have experienced above average  
257 numbers and high proportions of winter events, with seven winter events out of eight in 2013, four out of seven  
258 in 2021 and five out of seven in 2022. Figure 3 displays the seasonal distribution of these events, with 34% of  
259 meteotsunami recorded in December and January, and no events being recorded in March or April. Following

260 statistical analysis, the recorded maximum wave amplitude for each event resulted in a mean wave height of 0.33  
261 m for winter and 0.35 m for summer. With a t-test score of 0.30 and a P-value of 0.07, the tests indicate a similarity  
262 between the two sample sets, where the difference between seasonal wave heights is considered to be not  
263 statistically significant.  
264 Summarising the results from the catalogue in its entirety, we suggest that there are three ‘hotspot’ regions where  
265 meteotsunami events appear to be most frequent, these are 1) northwest Scotland, 2) Wales and 3) the southwest  
266 UK. Up until 2009, Penzance in the southwest UK had experienced the most meteotsunami with eight in total.  
267 Then from 2010, Kinlochbervie in Northwest Scotland experienced the maximum wave height of 0.51 m during  
268 the 16 November 2016 event. This same location was exposed to 14 separate meteotsunami events in the 12 years  
269 from 2010 to 2022. Harbour style geomorphology appears to be more susceptible to meteotsunami resonance  
270 recording 71% of the events and beach environments with the remaining 29%.  
271  
272



**Figure 2:** Seasonal distribution of UK meteotsunami events, historical record (1750 to 2009) and current record (2010 to 2022).

273  
274  
275  
276  
277



**Figure 3:** Seasonal and locational distribution of maximum wave heights from 1750 to 2022. Number of events at specific locations are represented by dot size as shown in the key. Base map © Crown copyright and database rights 2022 Ordnance Survey (100025252).

278

### 279 3.3 Relationship between meteotsunami and winter storms

280 In this section, we highlight two specific meteotsunami events that occurred in two consecutive winter seasons.  
 281 These two events have been picked as they are new events to the catalogue, and they represent a typical winter  
 282 meteotsunami hidden in the associated storm data. The winter of 2021/22 saw seven sequential named storms  
 283 with five verifiable meteotsunami events, one of which was the 20th of October 2021. The winter of 2022/23 saw  
 284 3 likely / numerically verifiable meteotsunami events, one of which was the 1 November 2022. Both meteotsunami  
 285 events were low profile, localised in nature and hidden within larger scale heavily precipitating low-pressure  
 286 systems.

287

#### 288 3.3.1 Event 1: 20 October 2021

289 Two low pressure systems developed in the Atlantic Ocean and propagated eastwards towards the southwest UK.  
 290 The first system which was detected as a mature echo signature on radar contained a sharp cold front (squall)  
 291 which moved into Cornwall at approximately 0400 UTC (criteria 2a and Figure 4a) with a simultaneous leading  
 292 air pressure rise of 1.6 mb over 4 minutes followed by a sharp 2°C air temperature drop (criteria 2b/d and Figure

293 4c). A flattish ridge between this first system and the second system named Aurore by Meteo France led to a  
294 yellow rainfall warning being issued in the UK. At 1600 UTC the second system with a low-pressure centre of  
295 992 mb moved into the Isles of Scilly and propagated across Cornwall and Devon, it contained a heavily  
296 precipitating non-linear system with convective activity and strong winds (+70 mph) rapidly veering from west  
297 to south (criteria 2d). This system initiated a sharp air pressure rise of 0.5 mb over 2 minutes which coincided  
298 with a high tide. Both low pressure systems initiated a series of meteotsunami waves that tracked eastwards along  
299 the coast of Cornwall, Devon, and Dorset. Wave anomalies were recorded in Plymouth at 1645 UTC with a  
300 maximum wave height of 0.36 m, Totnes at 1700 UTC and Port Isaac, Weymouth, and the Isle of Wight at 1800  
301 UTC before dissipating (criteria 1b/c).

302

### 303 **Event 2: 1 November 2022**

304 A series of low-pressure systems over the Atlantic Ocean, swept into the southwest UK on 1 November, the first  
305 one with its centre over Cornwall at 0000 UTC, followed by a second low pressure system arriving along the  
306 southwest coast at approx. 0600 UTC then moving northeast up over the UK.

307 This synoptic situation was complicated by a series of associated cold fronts followed by low pressure troughs. A  
308 quasi-linear precipitation system with its associated convective cells developed in the vicinity (criteria 2a and c,  
309 Figures 4d and e). The arrival of the storm feature was detected in surface observations with a sharp 1 mb/35  
310 minutes air pressure rise (Figure 4f) which coincided with a series of unpredictable meteotsunami waves which  
311 reached a maximum wave height of 0.3 m (criteria 2b). The waves tracked along the southwest UK alongside of  
312 the movement of the cold fronts, the heavily precipitating cells and the convective activity where it was recorded  
313 at five tide gauge sites along the southwest coast at Port Isaac, St Marys, Newlyn, Plymouth and Totnes (criteria  
314 1c). The first series of wave anomalies occurred at 0900 UTC coinciding with a high tide followed by a second  
315 set of wave anomalies at 1600 UTC coinciding with a low tide.

316


317



a. Rain radar 20 October 2021 at 0300 UTC



d. Rain radar 1 November 2022 at 0900 UTC



b. Lightning 20 October 2021 at 0000 UTC



e. Lightning 1 November 2022 at 0600 UTC



c. Air pressure 20 October 2021



f. Air pressure 1 November 2022

**Figure 4:** The relationships between criteria of two winter meteotsunami events.. All images are open source: Rain radar : [www.ventusy.com](http://www.ventusy.com) Lightning : [www.lightningmaps.org](http://www.lightningmaps.org) Air pressure: [www.starlingsroost.ddns.net](http://www.starlingsroost.ddns.net)

319 **Table 1:** Descriptions and references for events that can be identified as UK meteotsunami events from 1750 to 2022. 1750 to 2009  
320 are principally derived from historical sources and 2010 to 2022 are principally derived from instrumental data. The threshold criteria  
321 outlined in the methodology section was used to verify previous (V) or identify new (N) events, including new winter events (NW).  
322 (Wm represents maximum wave height in metres).

Date	Location	Wm	Time	Notes	Id criteria	Id status	Reference
1 Nov 1755	Ilfracombe	0.3	14.00	4 waves in 2 h, calm, NE wind, low tide, damage to sandbanks	IA, IB, 2A,	NW	Dawson et al 2000
27 Feb 1756	Ilfracombe	1.8	18.00	4 mins wave period, 30 mins duration, rumbling sea	IA, IB, 2A, 3A	NW	Dawson et al 2000
31 May 1759	Lyme Regis	-	-	3 waves in 1 h, ebb and flow	IA, 2A, 3A	N	Dawson et al 2000
31 Mar 1761	Mounts Bay	1.2	12.30	Ebb and flow 5 times in 1 h, NNE wind, cloudy, thunder	IA, IC, 1C, 2A	NW	Long 2015
18 Sept 1763	Weymouth	3	-	3 waves, ebb, and flow	IA, IB, 3A	N	<a href="http://www.phenomena.org.uk">www.phenomena.org.uk</a>
11 Feb 1764	Bristol	3	-	2 waves, ebb in 30 mins	IA, 2A, 3A	NW	<a href="http://www.phenomena.org.uk">www.phenomena.org.uk</a>
23 Dec 1791	Cornwall	-	04.00	Rain, hail, extreme lightning, boats moved	2A, 3A	NW	Borlase 1758
17 July 1793	Plymouth	0.6	07.00	3 waves in 1 h, boats damaged	IA, IB, 2A, 3A	N	-
18 Aug 1797	Lyme Regis	3	-	3 waves in 1 h, lightning	IA, 2A, 3A	N	Dawson et al 2000
9 Aug 1802	Devon	0.35	06.00	3 waves in 1 h, ebb and flow twice in 20 min	IA, IC	N	Long 2015
10 Aug 1802	Teignmouth	0.6	08.00	10 min interval waves, fish on shoreline	IA, IB	N	Long 2015
30 Aug 1802	Jersey	1.2	-	3 ebb and flows in 8 mins	IA, IB, 2A	N	Long 2015
31 May 1811	Plymouth	2.4	03.00	4 h duration, rain, low pressure, ebb and flow, SW wind	IA, IB, 2A, 2B	N	Dawson et al 2000
4 Mar 1818	Portsmouth	1.5	08.00	Rain, W to SW wind, high water for 3 h	IC, 2A, 2D	NW	<a href="http://www.surgewatch.org">www.surgewatch.org</a>
13 Sept 1821	Plymouth	1	14.00	Ebb and flow, boats moved	IA, IB, 1C, 3A	N	Long 2015
13 July 1824	Plymouth	0.6	22.00	Ebb and flow, 4 m/s currents, ESE light wind, boats moved	IA, IC, 2D, 3A	N	Archer, 2016
23 Nov 1824	Plymouth	2	01.00	3 waves in 10 min intervals, storm surge, 180 metres inland	IB, IB, 2A, 3A	V	Haslett and Bryant 2009
5 July 1843	Plymouth	1	11.00	4 waves in 20 min, storm moved north, strong wind	IA, IB, 1C, 2A	N	Thompson et al 2020
3 July 1845	Weymouth	0.6	10.30	Ebb and flow 5 times in 30 mins	IA, 2A, 3A	V	Long 2015
5 July 1846	Cornwall	0.5	04.30	Thunder, severe storm reported	IC, 2A, 3A	V	Dawson et al 2000
1 Aug 1846	Penzance	0.3	04.00	30 min duration, calm sea, 6-minute ebb and flow	IC, 2A, 3A	V	Dawson et al 2000
23 May 1847	Penzance	0.9	05.00	20 mins, squally wind, sudden rush of water	IA, IB, 2A, 2D	V	Long 2015
7 July 1848	Bristol	1.5	04.00	Thunder	IC, 2A, 3A	V	Edmonds 1862

323

Date	Location	Wm (m)	Time (UTC)	Notes	ID criteria	Id Status	Reference
6 June 1855	Penzance	0.9	pm	Ebb & flow 2 to 3 times, rumbling sea, strong currents	IC, 2A, 3A	N	Dawson et al 2000
5 June 1858	Folkstone	0.9	08.00	Ebb & flow in 5 mins, ENE to WNW wind, hail, rain, seiche	IA, 1C, 2A, 2D	V	Long 2015
25 June 1859	Cornwall	0.3	Night	Abnormal sea oscillations x2, squall line, strong currents	IA, 2A, 3A	V	Dawson et al 2000
4 Oct 1859	Cornwall	4.4	07.00	3 waves, 6 minute ebb, warm air temperatures, 1 mile upriver	IC, 2A, 3A	V	Dawson et al 2000
Oct 1865	Port Talbot	-	-	2 tides in 1 h	IA, 2A, 3A	NW	<a href="http://www.surewatch.org">www.surewatch.org</a>
23 Apr 1868	Lyme Regis	6	-	Swell, roar from the sea, no wind, low air pressure, calm sea	IC, 2A, 3A	V	Haslett & Bryant 2009
29 Sept 1869	Cornwall	0.9	06.00	20 min wave period	IA, 1B, 1C, 2A	V	Dawson et al 2000
13 June 1881	Shetland	-	-	3 waves in 20 min, storm, boat damage	IA, 2A	V	Long 2015
28 Aug 1883	Plymouth	0.25	09.00	Gravity pressure wave from Krakatoa volcanic eruption	IB, 2B,	V	Garrett 1970
17 Oct 1883	Severn Est	1	08.00	1 dead, SW strong wind, high tide, precipitation, 1 mile in	IA, 1C, 2A, 2D	V	Haslett & Bryant 2009
13 June 1886	Wick	0.45	16.30	Falling air pressure	IC, 2B, 3A	V	Long 2015
18 Aug 1892	Yealm	4	-	Quick ebb and flow, squall line, 3 waves, boat damage	IB, 1C, 2A, 2B	V	Haslett & Bryant 2009
16 Dec 1910	Ilfracombe	4	06.15	Swell, bore, low pressure, 100 metre inland, bedrock erosion	IB, 2B	V	Haslett & Bryant 2009
26 Dec 1912	Isle of Wight	0.9	12.00	975 mb pressure low, SW wind, rain, cold front	IA, 2A, 2B, 2D	NW	<a href="http://www.surewatch.org">www.surewatch.org</a>
20 July 1929	Folkstone	6	19.30	8 waves, 180 metres inland, 5 mins period, low tide, 3 dead	IB, 1C, 2A, 2D	V	Haslett & Bryant 2009
2 Aug 1932	Aberavon	9.3	-	4 dead, wave train, cloudy, rumbling sea, strong currents	IB, 2A, 3A	V	Haslett et al 2009
5 Aug 1938	Bridlington	4	08.00	Sea receded 4.5 m, boats moved, fish left on dry land	IA, 1B, 2A	V	Haslett et al 2009
4 July 1939	Milford Haven	6	00.30	3 dead, rumbling sea, boats moved, mid tide	IB, 2A, 3A	V	Haslett & Bryant 2009
3 July 1946	Cornwall	-	PM	Ebb and flow, squall line, rumbling sea, moorings broke	IA, 2A, 2B, 3A	V	Haslett & Bryant 2009
13 July 1949	Mevagissey	-	04.00	Easterly winds, boats smashed on rocks	IA, 1C, 2A, 2D	V	Long 2015
6 July 1957	Bembridge	4	19.30	Wave train, 2 waves in 1 h, sultry, overcast, rocks moved	IA, 1C, 1C, 2A	V	Haslett & Bryant 2009
31 July 1966	Westward Ho	3	PM	Receding water, frontal trough, squall line	IA, 1B, 2A	V	Haslett & Bryant 2009
1 July 1968	Folkstone	-	-	5 mb air pressure drop in 30 mins,	IA, 2B, 3A	V	Stevenson 1969
13 Feb 1979	Bristol	0.6	07.00	Spring tide, long unbroken waves, storm surge, 10 min period	IC, 2B	V	Haslett & Bryant 2009
28 May 2008	Peterhead	3	00.30	Ebb and flow in 10 mins, 4 to 6 waves	1 A-C, 2 A-C	V	Sibley et al 2006
29 Jan 2010	Lowestoft	0.29	16.00	Open cell, S moving storm, 11 tide gauges	IA-C, 2A, 3A	V	Williams et al 2021
29 Aug 2010	Lowestoft	0.27	19.00	Open cell, S moving storm, 4 tide gauges	IA-C, 2A, 3A	V	Williams et al 2021

Date	Location	Wm (m)	Time (UTC)	Notes	Id criteria	Id Status	Reference
3 Feb 2011	Ullapool	0.3	22.00	Open cell, E moving, 7 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
27 June 2011	Devonport	0.3	08.30	Non-linear, N moving, 8 tide gauges plus European tide gauges	1A-C, 2A, 3A	V	Tappin et al 2013
22 Aug 2011	Newhaven	0.3	01.00	Quasi linear, N moving, 3 tide gauges, mid latitude depression	1A-C, 2A, 3A	V	Williams et al 2021
24 Nov 2011	Ullapool	0.26	04.30	Open cell, E moving, 8 tide gauges, mid latitude depression	1A-C, 2A, 3A	V	Williams et al 2021
3 Jan 2012	Lowestoft	0.33	17.15	Quasi linear, SE moving, 17 tide gauges, Low pressure	1A-C, 2A, 3A	V	Williams et al 2021
4 Feb 2013	Stornoway	0.32	07.00	Open cell, SE moving, 13 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
3 Aug 2013	Aberdeen	0.25	07.30	Non-linear cluster, NE moving, 9 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
28 Oct 2013	Devonport	0.27	03.15	Non-linear cluster, NE moving, 4 tide gauges, 1 mb/1 h drop, high	1A-C, 2A, 3A	V	Williams et al 2021
5 Dec 2013	Kinlochbervie	0.35	16.00	Quasi linear, 19 tide gauges, 1.7 mb/1 h drop, storm surge, spring tide	1A-C, 2A, 3A	V	Williams et al 2021
15 Dec 2013	Ullapool	0.25	18.00	Quasi linear, E moving, 6 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
18 Dec 2013	Milford Haven	0.33	19.00	Quasi linear, E moving, 24 tide gauges, 2.6 mb/ 1 h drop,	1A-C, 2A, 3A	V	Williams et al 2021
20 Dec 2013	Kinlochbervie	0.25	19.45	Quasi linear, NE moving, 5 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
21 Dec 2013	Ullapool	0.28	10.00	Individual cell, NE moving, 4 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
3 Jan 2014	Newlyn	0.33	12.30	Quasi linear, 8 tide gauges, 1.2 mb/1 h drop, high winds, high tide	1A-C, 2A, 3A	V	Williams et al 2021
8 Feb 2014	Weymouth	0.25	20.00	Open cell, E moving, 14 tide gauges, 1.3 mb/1 h drop at 18.30	1A-C, 2A, 3A	V	Williams et al 2021
12 Feb 2014	Weymouth	0.26	21.45	Quasi linear, E moving, 15 tide gauges, high winds, storm at 13.00	1A-C, 2A, 3A	V	Williams et al 2021
21 May 2014	Newhaven	0.26	23.00	Non-linear, N moving, 4 tide gauges, wave period 29 minutes	1A-C, 2A, 3A	V	Williams et al 2021
22 May 2014	Lerwick	0.33	06.45	Quasi linear, N moving, 3 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
1 Jan 2015	Ullapool	0.26	01.30	Open cell, E moving, 9 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
8 Jan 2015	Ullapool	0.27	01.00	Quasi linear, E moving, 10 tide gauges, wave period 15 minutes	1A-C, 2A, 3A	V	Williams et al 2021
1 July 2015	Stonehaven	0.25	09.00	Individual cell, NE moving,	1A-C, 2A, 3A	V	Sibley et al 2016
2 July /2015	Lerwick	0.31	23.00	Non-linear, NE moving,	1A-C, 2A, 3A	V	Williams et al 2021
10 Dec 2015	Ullapool	0.25	08.30	Open cell, E moving, 4 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
27 Jan 2016	Workington	0.3	14.00	Non-linear, NE moving,	1A-C, 2A, 3A	V	Williams et al, 2021
1 Feb 2016	Stornoway	0.27	16.30	Open cell, E moving, 11 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
23 June 2016	Newhaven	0.7	04.40	Non-linear, NE moving, 6 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021



Date	Location	W/m (m)	Time (UTC)	Notes	ID criteria	Event ID	Reference
26 Aug 2016	Devonport	0.3	22.45	Individual cell, NE moving, 7 tide gauges	1A-C, 2A, 3A	N	-
16 Nov 2016	Kimlochbervie	0.51	14.15	Open cell, E moving, 7 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
26 Dec 2016	Stornoway	0.34	08.30	Open cell, SE moving, 8 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
11 Jan 2017	Kimlochbervie	0.25	08.00	Open cell, SE moving	1A-C, 2A, 3A	V	Williams et al 2021
16 Oct 2017	Lerwick	0.35	16.00	Quasi linear, NE moving, 20 tide gauges	1A-C, 2A, 3A	V	Williams et al 2021
29 June 2019	Aberdeen	0.3	17.00	Non-linear, supercell moving from North Sea to Norway	1A-C, 2A-C	N	-
8 Feb 2020	Port Stoth	0.4	12.00	Line convection, ebb & flow, before storm Ciara, Low pressure	1A-C, 2A, 2C	NW	-
21 Aug 2020	Perranporth	0.3	21.00	Spring tide, cold front, air pressure rise of 0.5 mb/2 min, bore	1C, 2B, 2C	N	-
5 July 2021	Westward Ho	0.6	12.40	S wind, Individual cell, mid tide, air pressure rise of 0.5 mb/1h, LP	1C, 2A-C, 3A	N	-
9 Aug 2021	Totnes	0.25	11.30	S wind, Non-linear, mid tide, air pressure rise 0.5 mb/30 mins	1A, 1C, 2A-C	N	-
27 Sept 2021	Plymouth	0.32	03.00	S/SW wind, Quasi-linear, CAPE, low tide, air pressure rise 1.1 mb/20 mins	1A, 1C, 2A-D, 3A	NW	-
2 Oct 2021	Totnes	0.29	12.00	SSE wind, Non-linear, mid tide, air pressure fall 1.4 mb/1 h	1A,1C,2A,2B	NW	-
20 Oct 2021	Plymouth	0.36	05.00	SSW, Non-linear, high tide, air pressure rise 1.5 mb/10 mins, CF	1A, 1C, 2A-C	NW	-
27 Nov 2021	Totnes	0.46	04.00	S/W, CAPE, mid tide, air pressure fall 1 mb/30 mins, storm surge,	1A,1C, 2A-D	NW	-
30 Dec 2021	Totnes	0.6	00.00	S/W, non-linear, high tide, air pressure fall 0.5 mb/20 mins, Low pressure, wave period 20 minutes	1A, 1C, 2A-D, 3A	NW	-
16 Jan 2022	Port Isaac	0.3	01.00	Mid tide, air pressure fall of 1.5 mb, pressure wave from volcanic eruption	1A-C, 2B	NW	-
8 Feb 2022	Dunnet	0.3	13.15	Currents of 4 m/s, high tide, approaching cold front from north	1A,1C,2C, 3A	NW	-
18 June 2022	Newlyn	0.7	14.30	Spanish plume, 7+ locations, air pressure fall of 4 mb/10 mins	1A-C, 2B, 3A	N	-
19 July 2022	Anglesey	0.3	12.00	spring tide, air pressure fall 1 mb/35 mins, 5x ebb & flow, 9 m inland	1A-C, 2A-C, 3A	N	-
1 Nov 2022	Plymouth	0.3	09.00	Air pressure rise of 1 mb/5 mins, heavy ppt, thunder, SW wind	1A-C, 2A-D	NW	-
8 Nov 2022	Port Isaac	0.5	10.00	Air pressure rise 0.7 mb/2 mins, thunder, winds, precipitation	1A-C, 2A-D	NW	-
23 Nov 2022	Newlyn	0.7	09.00	Air pressure fall 1 mb/5 mins, cluster ppt, thunder,	1A-C, 2A-D	NW	-

## 326 **4 Discussion**

327 The aim of this paper was to introduce a revised, enhanced and current UK catalogue of meteotsunami events including a  
328 highlight of the seasonal occurrence, frequency, and spatial distribution of this hazard. This aim was set as there is no  
329 standardised identification criteria or up to date single catalogue of UK meteotsunami. This scenario has led to the mis  
330 conception that these events are non-hazardous, rare, and tend to occur more frequently in the summer months.

331

### 332 **4.1 The updated UK meteotsunami catalogue**

333 With the identification criteria we have laid out in this paper we have verified 98 events in UK waters since 1750 of which 38  
334 are new events containing seven new winter events in the historical record (1750 to 2009) and 8 new winter events in the  
335 modern record (2010 to 2022).

336 It was found that a selection of historical events were misidentified in accounts as either abnormal coastal flooding, non-  
337 tsunami, storm surge or of unknown origin. This was extended by an analysis of current data (since 2010) which allowed us  
338 to add a total of 38 new events to the catalogue, of which 15 occurred within winter months, these are highlighted in Table 1  
339 as new (N), new winter (NW) or verified (V) events.

340 The misidentified events were discovered after an attempt to highlight characteristics that match those listed in the  
341 methodology, in particular characteristics that suggested tsunami like phenomenon but with any associated storm like activity  
342 or air pressure fluctuations. If the account was found to contain a lack of evidence or information to suggest a meteotsunami  
343 it was rejected. An event occurring on 13 February 1979 was highlighted as a meteotsunami by Haslett et al. (2009a) but was  
344 contested by Thompson et al. (2020) as being a surge caused by a winter Atlantic storm due to its seasonal placement. In their  
345 2020 paper, Thompson et al appear not to class Atlantic storms systems as sources of meteotsunami. They state that from April  
346 to October, thunderstorms generate meteotsunami and from November to March, storms generate low pressure swells and  
347 surges. Our paper has matched descriptions in historical accounts with the criteria laid out in the methodology and we agree  
348 with Haslett and Bryant (2009a/b) that the 1979 winter event was a meteotsunami. This result was determined by the  
349 similarities in the pressure profile, geographical distribution and the speed of anomaly to the known meteotsunami event of 26  
350 June 2011.

351 In addition to the 1979 event, there were further events found that were previously labelled as meteotsunami to which our  
352 criteria have found them to be of alternative origin (tsunami) or to have insufficient detail or collaborative evidence to solidify  
353 a conclusion. These include the events presented in Long (2021), dated 14 October 1862 (found to be a tsunami due to an  
354 alternative source trigger), 15 August 1895 (insufficient information), 11 May 1912 (found to be a tidal bore) and another tidal  
355 bore dated 17 May 1964 presented in Haslett and Bryant, 2009a/b.

356 The event of the 31 March 1761 which was labelled as a tsunami by both Long (2015) and Thompson et al. (2020), was found  
357 to be a winter meteotsunami due to tsunami like waves being experienced not only along the southwest UK but also in Loch  
358 Ness in Scotland, with the mention of a calm sea before the arrival of thunderstorms.

359

## 360 **4.2 Seasonal and geographical patterns of UK meteotsunami**

361 The historical record (1750 to 2009) has been found to support previous studies such as Haslett and Bryant (2009a/b) that have  
362 alluded to the positive correlation between thunderstorms and meteotsunami waves with 71% of summer events displaying  
363 reports of convective activity. Our results have highlighted a summer prevalence of events with 48% of them peaking in July  
364 and August which reflects Thompson et al (2020). This prevalence has been based principally on the reliance on eyewitness  
365 reports and the volume of persons present at the shoreline during these months.

366 These summer events tend to be associated with heat waves and so called “Spanish plumes” as in the 27 June 2011 and the 18  
367 June 2022 events along the southwest UK. This is where warm air moves northwards from the European continent and Iberia,  
368 during which mesoscale convective weather tends to occur. In the summer, CAPE is at its highest and overland due to warm  
369 2 m air temperatures over landmasses (Holley et al. 2014). These types of weather event consist of single cell or clusters of  
370 small, short duration (< 1 hr) thunderstorms and squall lines with more than one convective cell (Sibley 2012 and Tappin et  
371 al. 2013).

372 The element of risk during the summer occurs when the meteotsunami wave can become fully disconnected from its source  
373 disturbance. This effect can be particularly apparent if the meteotsunami interacts with the continental slope where the wave  
374 can arrive hours after the original storm has dissipated or moved on. This delayed arrival of wave disturbances can surprise  
375 people who are subsequently back out on or near the water’s edge, believing the storm has passed. This scenario was  
376 experienced during the 5 July 2021 event that occurred at Westward Ho (North Devon). Where just after midday a small yet  
377 powerful wave unexpectedly progressed 50 metres up the beach inundating many beach goers.

378

379 Previous studies have suggested that winter wave anomalies such as meteotsunami are ‘less’ likely than storm waves, and  
380 surge, and winter data has not previously been interrogated for this reason. However, the present-day record (2010 to 2022)  
381 appears to contradict this with a winter prevalence of 66% of events peaking in December and January and with a tendency  
382 towards October and November in the 2021/22 winter season.

383 The results also show a geographical pattern to UK meteotsunami, with a large proportion of events occurring along the  
384 southwest UK and Northwest Scotland in the winter, aligning with the dominant weather direction of west to east from the  
385 Atlantic Ocean, and along the southern UK coast in the summer, aligning with Spanish Plumes bringing warm air poleward  
386 from the equator with southerly winds up and along the English Channel. The geographical pattern also reflects the influence  
387 of local bathymetry, with harbours (e.g., Penzance, Plymouth, Stornoway, and Port Talbot), bays (e.g., Kinlochbervie and Port  
388 Stoth) and river mouths (e.g., river Yealm and river Dart) containing conditions more favourable to meteotsunami initiation  
389 and amplification via resonance and seiching.

390 To further the concept presented in Williams et al. (2021) we selected two recent winter meteotsunami events and highlighted  
391 the meteotsunamigenic criteria. It has been indicated from the results that the combination of a mid-latitude depression, with  
392 frontal and convective weather moving across the UK may be important in the generation of this hazard. Results have shown  
393 that during these winter storms, convective elements are likely to be embedded around heavy rainfall (Figure 4a and b) and

394 strong winds associated with the cold front leading to the potential for meteotsunami waves. This winter synoptic situation is  
395 a product of the combination of the cold maritime Arctic air being introduced to the rear side of the cold front passing over  
396 relatively warm water. The risk of flooding can be exacerbated due to surface water from precipitation as the front crosses a  
397 landmass (Masselink et al. 2015).

398 The results highlighted an average maximum wave height of 0.3 m which may not seem ‘dangerous’ but this hazard is not  
399 purely about this single factor. The key that makes meteotsunami a potential hazard is the rapid onset of the wave (sometimes  
400 referred to as a “wall of water”) and the associated strong currents.

401

### 402 **4.3 Constraints and Limitations**

403 Identifying meteotsunami events in winter tends to be more difficult as the waves can be hidden and overshadowed by the  
404 wave characteristics of the trigger storms and may be missed unless looking specifically at the data. We strongly consider that  
405 this overshadowing means many of these winter meteotsunami do not get reported and this may have been the issue in previous  
406 research where certain winter events were identified as either storm waves or surges instead of meteotsunami. As we have seen  
407 there is a short observational record available for meteotsunami and there is evidence for severe under recording of such events.  
408 Even though the 2010 to 2022 record has shown significant improvements in recording completeness; the current 15-minute  
409 sampling interval is still too coarse. This was highlighted when certain events in the catalogue such as 2 October 2021, 20  
410 October 2021, 27 November 2021 and 19 July 2022 were uncovered in the 1-minute tide gauge data that were not so easy  
411 to locate in the 15-minute data. This creates an issue where many events with a wave period of under 15 minutes may be  
412 potentially missed. We recommend a reduction of the sampling interval to 1 to 5 minutes to yield more data to be able to draw  
413 a complete conclusion for this hazard.

414 Another limitation of this study linked to the sampling frequency was the treatment of wind-driven waves which can induce  
415 infra-gravity waves of a similar wave period to meteotsunami (2 to 5 minutes). We did initially consider wind and swell peak  
416 period and wave height; however, we discovered that the detection of infra-gravity waves from low frequency tide gauge data  
417 is uncertain and was deemed to be beyond the scope of this study. To perform such an analysis and to be confident in our  
418 results we would require 1 minute / 2 Hz data for a spectral analysis. However, it may be prudent to explore this aspect in  
419 future work.

420 We noted that historical accounts are not optimum for identifying and analysing meteotsunami due to their anecdotal nature  
421 and as such the number of events represented here may be dramatically underestimated. Data before 2008 is not readily  
422 available and records are spatially sparse which leads to incomplete data coverage and does not allow for a robust statistical  
423 analysis.

424 The placement of tide gauges used to provide data also affects results. The siting of UK tide gauges tend to be biased towards  
425 populated areas with harbours and river mouths for asset protection and is ideal for the capture of the resonant component of  
426 the meteotsunami wave. However, events in less populated areas may have been missed due to this placement. We suggest

427 potential tide gauge locations (based on the occurrence rate of previous events) could include beach or estuary locations around  
428 Devon and Cornwall such as Mevagissey or Perranporth and the North of Scotland such as Dunnet or Port Stoth.

429

#### 430 **4.4 What does this mean for the future?**

431 As the next few decades are likely to see sea level rise push mean and extreme water levels upward which will subsequently  
432 increase the level of risk by bringing the height of the storm tide closer to the flood stage (Masselink et al. 2015). At many UK  
433 locations, flood defences are at the design threshold of current storm surge levels, they are not designed or built for a sudden,  
434 prolonged water flow as seen in meteotsunami (Lazarus et al.2021). A question that has arisen from this paper is whether the  
435 winter seasons of 2013/14 and 2021/22 are outliers or whether this clustering of storms and meteotsunami will be a  
436 commonplace scenario in the future. Currently, we can detect and forecast mid latitude depressions nine to ten days in advance  
437 (Penn State, 2019), knowing this we can incorporate a warning of potential meteotsunami activity into the forecast. However,  
438 due to the localised nature of meteotsunami, risk level in each coastal areas needs to be considered on its own merits. The risks  
439 connected with a single meteotsunami event in two different bays can be quite different. One bay may suffer from inundation  
440 and flooding where another bay may be impacted by strong currents. This paper provides a valuable insight into the frequency,  
441 seasonality and spatial distribution of what was a hidden hazard in the UK. This new data will need to be incorporated and  
442 taken into consideration when coastal management strategies and defences are adjusted for the future.

443 Meteotsunami may well have some role to play in coastal storm impacts, however, the relative contribution of meteotsunami  
444 to storm surge in the aftermath of a storm and the full extent of the risk remains unknown and is beyond the scope of this work.  
445 It is also difficult to determine if the frequency and intensity of either low-pressure winter storms or winter meteotsunamis are  
446 on the increase. We invite a closer and more robust scrutiny of this hazard with a year-round perspective bearing in mind that  
447 no solid conclusions can be drawn without high frequency, long term, and continuous monitoring of this of hazard.

448

#### 449 **5 Conclusions**

450 Until recently it was thought that meteotsunami in the UK were rare and only occurred at certain times of the year, this  
451 misconception has led to a lack of provision in coastal management strategies and an under estimation of the frequency of this  
452 hazard. Motivated by coastal safety, this paper tests the hypothesis by presenting a new chronological catalogue dated from  
453 1750 to 2022 containing 98 UK meteotsunami with highlighted seasonal and geographical aspects. Using a standardised set of  
454 identification criteria developed for this study we have verified 60 previously listed events and presented 38 new events of  
455 which 15 were found to occur in the winter (Table 1).

456 Results demonstrate that meteotsunami are not restricted to the summer months and are more common than initially thought.  
457 The modern record (2010 to 2022) is short and has far more winter meteotsunamis, whereas the relatively long historical record  
458 (1750 to 2009) means that the most meteotsunamis in our total have occurred in the summer which confirms the results of  
459 Thompson et al. (2020) and Haslett and Bryant (2009). During the summer months (April to September inclusive) there is a  
460 trend towards the southern UK with a 71% positive correlation between meteotsunami events and summer convective weather

461 systems, which can occur within synoptic Spanish Plume settings as suggested by Sibley (2012). During the winter months  
462 (October to March inclusive) our results demonstrate a clustering around the southwest UK and Northwest Scotland with a  
463 positive correlation between meteotsunami and the passage of mid latitude depressions where convective elements are  
464 embedded in the associated cold fronts and low pressure troughs. Subsequently meteotsunami impacts can become hidden by  
465 being superimposed on top of the storm's impacts. The meteotsunami waves are further exacerbated by the localised nature of  
466 resonance characteristics, in particular harbours and bays which can create highly dangerous situations. The immutable nature  
467 and rapid onset of this hazard means that even a sole meteotsunami event can create changes in water level and flow velocity  
468 that has the potential to cause injury, loss of life and damage to assets.

469 Increased knowledge of this hazard can be made more easily accessible through a central catalogue such as the one presented  
470 in this paper and the provision of higher frequency monitoring to detect future trends. What was thought to be a 'hidden' and  
471 rare event in historical records may soon become a common hazard in the future.

472

473 **Author contributions.** C. Lewis designed and executed the study and prepared the original draft. D. Williams pre-processed  
474 and provided data from 2010 to 2017, reviewed and edited the text. T. Smyth, J. Neumann, and H. Cloke supervised the project,  
475 provided advice, editing and feedback on the manuscript.

476

477 **Competing interests.** The authors declare that they have no conflict of interest.

478

479 **Data availability.** The datasets used in this study were derived from resources available in the public domain.

480

#### 481 **References.**

482 Bechle, A.J., Wu, C.H., Kristovich, D.A.R., Anderson, E.J., Schwab, D.J. and Rabinovich, A.B.: Meteotsunamis in the  
483 Laurentian Great Lakes. Scientific Reports 6, (37832). <https://doi.org/10.1007/s11069-014-1193-5>. 2016.

484

485 Borlase, W.: The natural history of Cornwall. Oxford. 53-4. <https://archive.org/details/naturalhistoryc00borl>. 1758.

486 British Oceanographic Data Centre: <https://www.bodc.ac.uk/> Last access: 19 February 2022.

487

488 Burt, S.: Multiple airwaves crossing Britain and Ireland following the eruption of Hunga Tonga-Hunga Ha'apai on 15  
489 January 2022. Weather. Vol 77, No 3. <https://doi.org/10.1002/wea.4182>. 2022.

490

491 Chatfield, C.: Landmarks of world history: A Chronology of Remarkable Natural Phenomena Eighteenth Century 1761-1770.  
492 <http://www.phenomena.org.uk/page29/page38/page38.html>. Last accessed 19 February 2022.

493

494 Dawson, A.G., Musson, R.M.W., Foster, I.D.L., and Brunsten, D.: Abnormal historic sea-surface fluctuations, SW England  
495 marine Geology. Vol. 170, 59-68. [10.1016/S0025-3227\(00\)00065-7](https://doi.org/10.1016/S0025-3227(00)00065-7). 2020.

496

497 Dusek, G., DiVeglio, C., Licate, L., Heilman, L., Kirk, K., Paternostro, C., & Miller, A.: A meteotsunami climatology along  
498 the U.S. East Coast. Bulletin of the American Meteorological Society, 100(7), 1329–1345. [https://doi.org/10.1175/BAMS-D-](https://doi.org/10.1175/BAMS-D-18-0206.1)  
499 [18-0206.1](https://doi.org/10.1175/BAMS-D-18-0206.1). 2019.

500

501 Edmonds, R.: On extraordinary agitations of the sea not produced by winds or tides. Transactions of the Devonshire  
502 Association. 3. 144-152. <https://devonassoc.org.uk/publications/transactions/contents/>. 1869.

503

504 Goring, D.: Meteotsunami resulting from the propagation of synoptic-scale weather systems. Physics and Chemistry of the  
505 Earth, Volume 34, Issue 17, p. 1009-1015. [10.1016/j.pce.2009.10.004](https://doi.org/10.1016/j.pce.2009.10.004) 2009.

506

507 Haigh, I., Wadey, M., Wahl, T.: Spatial and temporal analysis of extreme sea level and storm surge events around the coastline  
508 of the UK. Sci Data 3, 160107. <https://doi.org/10.1038/data.2016.107> 2016.

509

510 Haslett, S.K. and Bryant, E.A.: Historic tsunamis in Britain since AD 1000: a review. Natural hazards and Earth Sciences. 8,  
511 587-601. <https://doi.org/10.5194/nhess-8-587-2008> 2008.

512

513 Haslett, S.K. and Bryant, E.A.: Meteorological Tsunamis in Southern Britain: An Historical Review. Geographical Review.  
514 99, 146–163. <https://doi.org/10.1111/j.1931-0846.2009.tb00424.x> 2009a.

515

516 Haslett, S.K., Mellor, H.E., and Bryant, E.A.: Meteo-tsunami hazard associated with summer thunderstorms in the United  
517 Kingdom. Physics and Chemistry of the Earth. 34, 1016-1022. <https://doi.org/10.1016/j.pce.2009.10.005> 2009b.

518

519 Holley, D.M., Dorling, S.R., Steele, C.J. and Earl, N.: A climatology of convective available potential energy in Great  
520 Britain. International Journal of Climatology. 34: 14. 3811-3824. <https://doi.org/10.1002/joc.3976> 2014.

521

522 Kim, M., Woo, S., Eom, H. and You, S.: Pressure-forced meteotsunami occurrences in the eastern Yellow Sea over the past  
523 decade (2010–2019): monitoring guidelines. Natural Hazards and Earth System Sciences. [https://doi.org/10.5194/nhess-2021-](https://doi.org/10.5194/nhess-2021-126)  
524 [126](https://doi.org/10.5194/nhess-2021-126) 2021.

525

526 Lazarus, E., Aldabet, S., Thompson, C., Hill, C., Nicholls, R., French, J., Brown, S., Tompkins, E., Haigh, I., Townend, I. and  
527 Penning-Rowse, E.: The UK needs an open data portal dedicated to coastal flood and erosion hazard risk and resilience.  
528 *Anthropocene Coasts*. 4(1): 137-146. <https://doi.org/10.1139/anc-2020-0023> 2021.  
529  
530 Lightning maps: <https://www.lightningmaps.org/#m=oss:t=2;s=0;o=0;b=:y=50.7086;x=-1.0547;z=4> last accessed 1 March  
531 2023.  
532  
533 Linares, Á., Wu, C.H., Bechle, A.J., Anderson, E.J. and Kristovich D.A.R.: Unexpected rip currents  
534 induced by a meteotsunami. *Sci Rep* 9:2105. <https://doi.org/10.1002/2016JCO11979> 2019.  
535  
536 Long, D.: A catalogue of tsunamis reported in the UK. British Geological Association 1R/15/043  
537 [https://nora.nerc.ac.uk/id/eprint/513298/1/IR\\_15\\_043%20BGS%20Tsunami%20catalogue%20update.pdf](https://nora.nerc.ac.uk/id/eprint/513298/1/IR_15_043%20BGS%20Tsunami%20catalogue%20update.pdf) 2015.  
538  
539 Long, D.: Comment on: Thompson et al 2020. UK meteotsunamis: a revision and update on events and their frequency.  
540 *Weather*. Vol 76. No4. 137-139. <https://doi.org/10.1144/sp456.10> 2021.  
541  
542 Lynett, P.J., Borrero, J., Son, S., Wilson, R. and Miller, K.: Assessment of the tsunami induced current hazard. *Geophysical*  
543 *Res Lett* 41 (6): 2048-2055. <https://doi.org/10.1002/2013GL058680> 2014.  
544  
545 Masselink, G., Scott, T., Poate, T., Russell, P., Davidson, M. and Conley, D.: The extreme 2013/2014 winter storms:  
546 hydrodynamic forcing and coastal response along the southwest coast of England. *Earth Surface Processes and Landforms*,  
547 volume 41, Issue 3 378-391. <https://doi.org/10.1002/esp.3836> 2015.  
548  
549 MET Office: 5 km Resolution UK Composite Rainfall Data from the Met Office Nimrod System.  
550 <https://catalogue.ceda.ac.uk/uuid/f91b2c5399c5bf689e29bb15ab45da8a> 2003.  
551  
552 Monserrat, S., Vilibic, I. and Rabinovich, A.B.: Meteotsunamis: atmospherically induced destructive ocean waves in the  
553 tsunami frequency band. *Natural Hazards and Earth System Science*. 6. 1035-1051. <https://doi.org/10.5194/nhess-6-1035-2006>  
554 2006.  
555  
556 National Oceanography Centre: <https://www.bodc.ac.uk/> Last accessed 1 March 2023.  
557  
558 National tide and sea level facility: Available at: <https://ntslf.org/> Last accessed 19 February 2022.  
559



560 Pattiaratchi, C.B. and Wijeratne, E.M.S.: Are meteotsunamis an underrated hazard? Philosophical Transactions of the Royal  
561 Society: Mathematical and Engineering Sciences 373. <https://doi.org/10.1007/s11069-014-1263-8> 2015.

562

563 Pellikka, H., Laurila, T.K., Boman, H., Karjalainen, A., Björkqvist, J. and Kahma, K.K. Meteotsunami occurrence in the Gulf  
564 of Finland over the past century, Natural Hazards Earth System Sciences. 0. (9). 2535-2546. [https://doi.org/10.5194/nhess-20-  
565 2535-2020](https://doi.org/10.5194/nhess-20-2535-2020) 2020.

566

567 Penn State.: Predictability limit: Scientists find bounds of weather forecasting. ScienceDaily. 15 April 2019.  
568 <https://sciencedaily.com/releases/2019/04/190415154722.htm> Last accessed 20 August 2022.

569

570 Proudman, F.R.S.: The Effects on the Sea of Changes in Atmospheric Pressure. Geophysical Journal International 2 s4.  
571 <https://doi.org/10.1111/j.1365-246X.1929.tb05408.x> 1929.

572

573 Reigate grammar weather station: The birth and impact of the St Jude day storm: October 2013. Available at:  
574 <https://rgsweather.com/2013/10/29/st-jude-causes-and-impacts-of-the-october-storm-27-28-2013/amp/> Last accessed 19  
575 February 2022.

576

577 Sea Level Monitoring Facility: <https://ioc-sealevelmonitoring.org/> Last accessed 1 March 2023.

578

579 Šepić, J., Vilibić, I. and Mahović, N.: Northern Adriatic meteorological tsunamis: observations, link to the atmosphere, and  
580 predictability. Journal of Geophysical Research Oceans. 117(C2). <https://doi.org/10.1029/2011JC007608> 2012.

581

582 Šepić, J., Vilibić, I., Rabinovich, A. et al. Widespread tsunami-like waves of 23-27 June in the Mediterranean and Black  
583 Seas generated by high-altitude atmospheric forcing. Sci Rep 5, 11682. <https://doi.org/10.1038/srep11682> 2015.

584

585 Šepić, J., Vilibić, I., Rabinovich, A. et al. Meteotsunami (“Marrobbio”) of 25–26 June 2014 on the Southwestern Coast of  
586 Sicily, Italy. Pure Appl. Geophysics. 175, 1573–1593. <https://doi.org/10.1007/s00024-018-1827-8> 2018.

587

588 Sibley, A.: Thunderstorms from a Spanish Plume event on 28 June 2011. Weather. 67. No 6. 143-152.  
589 <https://doi:10.1002/wea.1928>. 2012.

590

591 Sibley, A., Cox, D., Long, D., Tappin, D.R. and Horsburgh, K.J.: Meteorologically generated tsunami like waves in the North  
592 Sea on 1 July 2015 and 28 May 2008. Weather. 71. 68-74. <https://doi.org/10.1002/wea.2696> 2016.

593

594 Starlings roost weather: <http://starlingsroost.ddns.net/weather/stats.php?type=day&field=&date=2023-03-05> Last accessed 1  
595 March 2023.  
596  
597 Stevenson, C. M.: The dust fall and severe storms of 1 July 1968. *Weather*. 66 (5): 125–127. DOI:[10.1002/WEA.780](https://doi.org/10.1002/WEA.780) 1969.  
598  
599 Surge Watch database: A database of UK coastal flood events. <https://www.surgewatch.org/> Last accessed 19 February 2022.  
600  
601 Tappin, D.R., Sibley, A., Horsburgh, K.J., Daubord, C., Cox, D. and Long, D.: The English Channel ‘tsunami’ of 27 June 2011  
602 - a probable meteorological source. *Weather*. 68. 144–152. <https://doi.org/10.1002/wea.2061> 2013.  
603  
604 Thompson, J., Renzi, E., Sibley, A. and Tappin, D.: UK meteotsunamis: a revision and update on events and their frequency.  
605 *Weather*. 75.9, 281–287. <https://doi.org/10.1002/wea.3741> 2020.  
606  
607 University of Wyoming: <http://weather.uwyo.edu/upperair/sounding.html> Last accessed 19 February 2022.  
608  
609 Ventusky weather: <https://www.ventusky.com/?p=54.4;-5.9;4&l=radar&t=20230313/1100&w=off> Last accessed 1 March  
610 2023.  
611  
612 Vilibić, I. and Šepić, J.: Global mapping of non-seismic sea level oscillations at tsunami timescales. *Scientific reports*. 7. (1)  
613 40818. <https://doi.org/10.1038/srep40818> 2017.  
614  
615 Vilibić, I., Šepić, J., Dunic, N., Sevault, F., Monserrat, S. and Jorda, G.: Proxy-based Assessment of Strength and Frequency  
616 of Meteotsunamis in Future Climate. *Geophysical Research Letters*. 45. 10501-10508.  
617 <https://doi.org/10.1029/2018GL079566> 2018.  
618  
619 Williams, D.A., Horsburgh, K.J., Schultz, D.M. and Hughes, C.W.: Examination of generation mechanisms for an English  
620 Channel Meteotsunami: Combining observations and modelling. *Journal of Physical Oceanography*. 49. 103-120.  
621 <https://doi.org/10.1175/JPO-D-18-0161.1> 2019.  
622  
623 Williams, D. A., Schultz, D. M., Horsburgh, K. J., & Hughes, C. W.: An 8-yr meteotsunami climatology  
624 across northwest Europe: 2010–2017. *Journal of physical oceanography*. 1145-1160. [https://doi.org/10.1175/JPO-D-20-](https://doi.org/10.1175/JPO-D-20-0175.1)  
625 [0175.1](https://doi.org/10.1175/JPO-D-20-0175.1) 2021.  
626