



Data fusion uncertainty-enabled methods to map street-scale hourly NO₂ in Barcelona city: a case study with CALIOPE-Urban v1.0

Alvaro Criado¹, Jan Mateu Armengol¹, Hervé Petetin¹, Daniel Rodríguez-Rey¹, Jaime Benavides^{1,3}, Marc Guevara¹, Carlos Pérez García-Pando^{1,2}, Albert Soret¹, and Oriol Jorba¹

¹Barcelona Supercomputing Center, Barcelona, Spain

²ICREA, Catalan Institution for Research and Advanced Studies, Barcelona, Spain

³Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, New York, NY 10032, USA

Correspondence: Alvaro Criado (alvaro.criado@bsc.es) & Jan Mateu Armengol (jan.mateu@bsc.es)

Abstract. Comprehensive monitoring of NO₂ exceedances is imperative for protecting human health, especially in trafficked urban areas. However, accurate spatial characterization of exceedances is challenging due to the typically low density of air quality monitoring stations and the inherent uncertainties of urban air quality models. We study how observational data from different sources and time scales can be combined with a dispersion air quality model to obtain bias-corrected NO₂ hourly maps at the street scale. We present a kriging-based data-fusion workflow that merges a dispersion model output with continuous hourly observations, and uses a machine-learning-based Land Use Regression (LUR) model constrained with past short intensive passive dosimeter campaigns observations. While the hourly observations allow to bias-adjust the temporal variability of the dispersion model, the microscale-LUR model adds information on the NO₂ spatial patterns. Our method includes uncertainty calculation based on the estimated error variance of the Universal Kriging technique, which is subsequently used to produce urban maps of probability of exceeding the 200 $\mu\text{g}/\text{m}^3$ hourly and the 40 $\mu\text{g}/\text{m}^3$ NO₂ annual average limits. We assess the statistical performance of this approach in the city of Barcelona for the year 2019. Our results show that simply merging the monitoring stations with the model output already significantly increases the correlation coefficient (r) by +29% and decreases the Root Mean Square Error ($RMSE$) by -32%. When adding the time-invariant LUR model in the data-fusion workflow, the improvement is even more remarkable: +46% and -48% for the r and $RMSE$, respectively. Our work highlights the usefulness of high-resolution spatial information in data-fusion methods to estimate exceedances at the street scale better.

1 Introduction

Air pollution is the leading environmental risk factor globally (WHO, 2021). Mortality, the decrease in life quality, and the detrimental economic effects associated with air pollution are pressing decision-makers to take action, especially in urban areas, where more than 50% of the global population lives and air quality standards are frequently exceeded. In the city of Barcelona (Spain), the high vehicle density (about 5800 vehicles km^{-2} (Rivas et al., 2014)) induces a chronic NO₂ problem, which makes Barcelona the sixth European city with the highest mortality associated with NO₂ exposure (ISGlobal, 2021;



Khomenko et al., 2021). In this context, obtaining high-resolution exposure to NO₂ is crucial for decision-making in urban air quality management.

During the last decades, several approaches have been developed to estimate NO₂ exposure at different spatio-temporal scales (Denby, 2011). A common one is the Land Use Regression (LUR) model, which relates explanatory variables of different nature (land use cover, population density, traffic, climate, and others) with air quality observations using regression models (Briggs et al., 1997; Hoek et al., 2008; Beelen et al., 2013). LUR models are generally skillful, relatively easy to implement, and not very demanding regarding computational resources. However, urban areas often present strong NO₂ spatial gradients that the official monitoring network cannot correctly characterize due to its low spatial representativeness (Vardoulakis et al., 2005; Santiago et al., 2013; Duyzer et al., 2015a). To overcome this limitation and produce accurate surface NO₂ maps, urban or microscale-LUR models rely on Low-Cost Sensors (LCS), typically restricting the temporal coverage to a few weeks. Works dealing with microscale-LUR models have used different types of LCS, including passive dosimeters, which report period-averaged concentrations (Perelló et al., 2021a; Su et al., 2009), time-dependent LCS (Munir et al., 2020; Weissert et al., 2019), or mobile LCS campaigns (Wang et al., 2021). Due to the lack of experimental campaigns monitoring at high spatial and temporal (hourly) resolutions consistently over a whole year, current microscale-LUR studies typically cannot target the hourly averaged NO₂ maximum level (200 $\mu\text{g}/\text{m}^3$) regulated by the 2008 European Ambient Air Quality Directive (AAQD) (2008/EC/50).

Physics-based urban air quality models can generate hourly pollutant concentration estimates, overcoming the temporal limitation of microscale-LUR models. Currently, these systems usually consist of the coupling between a regional chemical transport model, which accounts for the long-range transport of pollutants, and an urban scale dispersion model. The last one can be based on semi-empirical relations such as Gaussian dispersion models and mass exchange global parameterizations (e.g. Soulhac et al. (2017); Kim et al. (2018); Benavides et al. (2019); Denby et al. (2020); Hood et al. (2021), or an obstacle resolving dispersion model using Computational Fluid Dynamics (Kwak et al., 2015; Auvinen et al., 2017). Despite the recent efforts to improve urban dispersion modeling systems, they are afflicted by persistent uncertainties and biases, notably due to the difficulty of prescribing accurate boundary conditions and emissions at the street scale, and reproducing the turbulent phenomena within the urban canopy.

In order to reduce model uncertainties, data-fusion methods can be employed to post-process model outputs and obtain more reliable NO₂ exposure maps. Several works have used monitoring station data to build data-fusion methods, either relying solely on urban dispersion models to explain the spatial distribution (Tilloy et al., 2013) or adding different spatial information (e.g. traffic intensity, satellite data, or land use cover) as proxies in addition to the model output (Horálek et al., 2006; Chen et al., 2019; Zhang et al., 2021; Dimakopoulou et al., 2022). In urban areas, the usual low density of monitoring stations has motivated the development of data-fusion methods that integrate LCS campaigns to better explain the spatial distribution of NO₂ at the street-scale. For instance, the works of Schneider et al. (2017) and Mijling (2020) combine time-resolved LCS hourly data with an urban model output to improve the NO₂ characterization at a high-spatial resolution. Schneider et al. (2017) use a popular geostatistic technique, Universal Kriging, considering the time-aggregated annual mean of an urban model as a *basemap* (or *climatology*) to explain the long-term spatial gradients at the street-scale, while the time-dependent



LCS network explains the temporal behavior. However, the temporal coverage of their results is restricted to a few weeks in which measurements are available. Thus, compromising their use to systematically estimate hourly NO₂ exposure levels for extended periods, in the order of years.

60 By combining model and observational data, advanced data-fusion methods can provide typically unbiased estimates of pollutant concentrations at the street scale. However, another piece of information of crucial importance is the uncertainty of the estimated concentrations, as it can help decision-making or support the design of environmental epidemiological studies (Gryparis et al., 2009). The Universal Kriging methodology provides the error variance of its predictions, which has already been used as a measure of the uncertainty on data-fusion results of NO₂ at the street scale (Schneider et al., 2017). However, 65 the validity of the confidence intervals and the normality of error distribution in this application remains to be investigated.

Our study presents a data-fusion methodology considering a microscale-LUR model, in addition to the hourly monitoring data, to bias-correct hourly NO₂ estimates of an urban dispersion model at high spatial resolution (20m × 20m). Similarly to Schneider et al. (2017), our work also relies on the *basemap* concept. However, contrary to previous studies, we have derived it using a microscale-LUR model based on 840 samplers from recent passive dosimeters campaigns (Perelló et al., 2021a; 70 Benavides et al., 2019). Thus, the basemap accounts for the spatial patterns, whereas the temporal behavior is characterized by the hourly urban model output and hourly monitoring data. This approach can be very convenient for applying data-fusion methods in cities where period-averaged LCS campaigns are available but lack time-dependent LCS data throughout the year, which is usually the case. To assess the benefits of considering such microscale-LUR basemap, we compare two different data-fusion methods: (i) Universal Kriging combining hourly observations with the hourly outputs of a street-scale Gaussian 75 dispersion model, namely UK-DM, and (ii) Universal Kriging combining the above items and the microscale-LUR model, namely UK-DM-LUR. The data-fusion methods are applied in Barcelona city (Spain) for the entire year 2019. An original aspect of the present study is the empirical validation of the UK-based uncertainties and their translation into street-scale probabilities of exceedance of the hourly and annual regulatory thresholds.

The paper is structured as follows: the observational data is described in Sect. 2.1. The Gaussian dispersion air quality 80 model CALIOPE-Urban used to produce hourly high-resolution fields of surface NO₂ concentrations is described in Sect. 2.2, while the microscale-LUR method is explained in Sect. 2.3. A detailed description of the data-fusion methods is given in Sect. 2.4. Results of the microscale-LUR model are presented in Sect. 3.1, and Sect. 3.2 discusses the results of the data-fusion methodologies. Finally, conclusions and final remarks are provided in Sect. 4.

2 Data and methodology

85 We compare two data-fusion methods (UK-DM and UK-DM-LUR) illustrated in Fig. 1. Below we describe each process and dataset used to derive them.




Figure 1. Workflow of the two studied data-fusion methodologies. Hourly data from monitoring stations are combined with hourly dispersion model results (UK-DM) and the time-invariant microscale-LUR basemap (UK-DM-LUR). PBL stands for Planetary Boundary Layer.

2.1 Study domain and observational NO₂ data

Barcelona (Fig. 2) is the second most populated city in Spain and the tenth in Europe, with approximately 1.660.000 inhabitants and 102 km² (~ 16.300 people per km²). It is located on the northeast coast of Spain, between the Mediterranean Sea and the Collserola mountains. The city has a Mediterranean climate characterized by the dominance of sea breeze during the warm season, shallow boundary layer development, and recirculation of air pollutants (Jorba et al., 2004).

Hourly NO₂ observational data for 2019 are obtained from the Catalan Atmospheric Pollution Surveillance Network (XVPCA) measurement points in the Barcelona urban and surrounding areas. There are 12 stations available on the Barcelona agglomeration (Fig. 2), with a percentage of availability of hourly data greater than 93%. *Gràcia* and *Eixample* are urban traffic monitoring stations, *Segnier* and *Jardins* are sub-urban background stations, and the remaining 8 correspond to urban background stations.

Two different NO₂ passive dosimeter experimental campaigns (Fig. 3) are considered to derive the microscale-LUR model: the xAire citizen science campaign (Perelló et al., 2021a, b) composed of 725 samplers and deployed between February




Figure 2. Domain of study and location of the referenced monitoring stations. The map has been generated using *ggplot2* (Wickham (2016)) and *ggmap* (Kahle and Wickham (2013)) R packages (R Core Team (2013)), and data from OpenStreetMap. © OpenStreetMap contributors 2017. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.

16th and March 15th, 2018, and the 2-week measurement campaign of the *Institute of Environmental Assessment and Water Research - Spanish National Research Council (IDAEA-CSIC)*, that deployed 175 NO₂ samplers across Barcelona during
100 February and March 2017 (Benavides et al., 2019). Both campaigns used Palmes-type NO₂ diffusion tubes (Palmes et al. (1976)) to sample the NO₂ levels, which implies an estimated uncertainty of $\pm 25\%$, as reported in Kuklinska et al. (2015).

2.2 Street-scale air quality model: CALIOPE-Urban

Hourly high-resolution concentrations of surface NO₂ at street-scale over the city of Barcelona are estimated using the CALIOPE-Urban multi-scale air quality model (Benavides et al. (2019)). CALIOPE-Urban accounts for the dispersion of
105 traffic emissions at high spatial resolution using the R-LINE Gaussian dispersion model (Snyder et al. (2013); Venkatram et al. (2013)). As described in more detail in Benavides et al. (2019), R-LINE is adapted to street-canyons by taking into account road-link traffic emissions (Guevara et al. (2020)), meteorological variables (e.g., wind speed and direction, Monin–Obukhov length and planetary boundary layer height) and buildings morphology (e.g., building density and height, and street orienta-



tion). The chemical balance between NO_x and NO_2 is computed based on the generic reaction set (Valencia et al. (2018))
110 assuming clear-sky conditions and uncoupling chemistry from transport phenomena; in other words, the aging of pollutants is
solely a function of wind speed and the distance between source and receptors.

At the regional scale, CALIOPE-Urban relies on the regional air quality modeling system CALIOPE (Baldasano Recio et al.
(2011)) for predicting urban background NO_2 concentration. The regional CALIOPE accounts for long-range transport of
pollutants using three nested domains at increasing resolutions: $12 \text{ km} \times 12 \text{ km}$ for the European region, $4 \text{ km} \times 4 \text{ km}$ for the
115 Iberian Peninsula, and $1 \text{ km} \times 1 \text{ km}$ for the region of Catalonia (Baldasano Recio et al. (2011), Pay et al. (2014)). The urban
background NO_2 concentrations obtained with regional CALIOPE are combined with the R-LINE dispersion results using a
dedicated parameterization of the vertical mixing (Benavides et al. (2019)).

In this work, CALIOPE-Urban employs a non-uniform mesh refined at the edge of traffic roads and coarser in low-gradient
regions of NO_2 . This type of mesh accelerates the calculations and reduces memory demand. The refined grid zones have a
120 resolution of $25 \text{ m} \times 25 \text{ m}$, progressively degrading to $500 \text{ m} \times 500 \text{ m}$ in the regions of low NO_2 gradients. To facilitate
their visualization, these NO_2 concentrations are finally interpolated over a uniform mesh with a resolution of $20 \text{ m} \times 20 \text{ m}$.
CALIOPE-Urban has been evaluated and successfully used in the framework of several impact studies, including the works of
Benavides et al. (2021) and Rodriguez-Rey et al. (2022).

2.3 Microscale-LUR model using Gradient Boosting Machine (GBM)

125 A non-linear microscale-LUR model based on passive dosimeter campaigns is used to produce an observation-based clima-
tological view of the NO_2 concentrations at a high spatial resolution over Barcelona city. While the monitoring stations and
the urban dispersion model provide information on the pollutants' temporal behavior, the microscale-LUR basemap (long-
term mean) remains constant in time. Its main goal is to provide reliable long-term spatial variability patterns of NO_2 at high
resolution using observational data and other urban information.

130 The target variable of the microscale-LUR model is the time-averaged concentrations of the two different NO_2 experimental
campaigns described in Sect. 2.1 and represented in Fig. 3. We have discarded the xAire samplers related to playgrounds
and classrooms, so we are using the remaining 669. In order to combine the xAire and IDAEA-CSIC campaigns, we have
annualized both following the procedure described in Perelló et al. (2021b): for each station, an adjustment factor is computed
as the ratio between the observed 2017 annual mean and the average over the period of the experimental campaign. Then, the
135 average of this factor over all stations is used to scale all passive samplers to the 2017 annual mean. This scaling assumes that
the ratio does not depend on location and can be applied to all samplers. Despite adding some noise to the experimental results,
it corrects the environmental conditions influence and allows combining both campaigns, producing a dataset of 844 samplers
on which the microscale-LUR model relies.

The potential predictors of the microscale-LUR model are shown in Table 1. The geometric variables are calculated from
140 the *Institut Cartogràfic i Geològic de Catalunya* (ICGC) and *Plan Nacional de Ortografía Aérea* (PNOA (2020)). Traffic-
related predictors consist of traffic density (t) for different circular buffer sizes. Being a the radius of the buffer, t_a is computed
following Eq. (1) expressed in [$vehicles \cdot m / s$]:




Figure 3. Sampler locations of the two different NO₂ experimental campaigns used to train the microscale-LUR model. The left panel shows the NO₂ values and the locations of the combined campaigns. The top- and bottom-right panels show the CSIC and xAire campaign locations, respectively. The colour scale refers to the 2017 annualized NO₂ values, in $\mu\text{g}/\text{m}^3$. The map has been generated using *ggplot2* (Wickham (2016)) and *ggmap* (Kahle and Wickham (2013)) R packages (R Core Team (2013)), and data from OpenStreetMap. © OpenStreetMap contributors 2017. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.

$$t_a = \sum_{i=1}^n AADT_i \cdot l_{a,i}, \quad (1)$$

where i represents the street segment, n is the number of street segments over the circular area of πa^2 , l_i is the length of the street segment i within the buffer, and $AADT_s$ is the average daily traffic of the street segment s expressed in vehicles per second. The t_a predictors associated with the smaller buffers (5, 10 and 15 m) have highly skewed distributions, given that most values across the map are null. To avoid training the microscale-LUR with skewed predictors, we introduce here the *traffic scaled* variable, s , which combines all buffers as follows:



Type	Num.	Variable	Resolution
Urban geometric	1	Average building density	Squared buffer of 250m × 250m
	2	Average building height	
	3	Maximum building height	
	4	Standard deviation building height	
Traffic-related	5-16	Simulated vehicular traffic densities	Circular buffers of 5, 10, 15, 25, 50, 100, 300, 500, 1000, 2000, 3000 and 4000 m of radius
	17	Traffic scaled	Linear combination of the buffers above
Output from the regional modeling system CALIOPE (lowest layer)	18	NO ₂	Uniform mesh of 1km × 1km
	19	Planetary boundary layer height	
	20	Wind speed	
Output from the CALIOPE-Urban model	21	NO ₂	Non uniform mesh (25m × 25m to 500m × 500m)

Table 1. The final microscale-LUR model contemplates the use of these 21 potential predictors.

$$s = \frac{1}{N} \sum_a \frac{t_a}{\pi a^2}, \quad (2)$$

150 where N is the number of buffers (12 in our case). Traffic data are extracted from the road-link traffic network of the HER-MESv3 bottom-up emission model (Guevara et al. (2020)). We also considered NO₂, Planetary Boundary Layer height and wind speed annual means from the regional air quality modeling system CALIOPE as potential predictors, together with the NO₂ annual mean from the air quality model CALIOPE-Urban.

A recursive feature elimination method has been applied to remove highly correlated or uninformative features. We have 155 used the simple backward selection algorithm implemented in the R package (Kuhn, 2008), which starts with the full-featured model and gradually removes the least important feature while monitoring the *RMSE* in Cross-Validation (CV). The goal is to obtain the simplest model with the lowest *RMSE* to gain generalization and interpretability. The final microscale-LUR model includes the following eight predictors: average building density; traffic buffers of 25, 100 m, and the traffic scaled variable; all the annually averaged data from the regional CALIOPE modeling system; and the annual average NO₂ from CALIOPE-Urban.

160 To account for non-linear relations among the predictors and the target variable, we used the Gradient Boosting Machine (GBM) algorithm implemented in the R package *gbm* (Ridgeway, 2004). GBM is a popular machine learning algorithm (Natekin and Knoll, 2013) that has shown excellent results in terms of accuracy and generalization when compared to other learning algorithms (Caruana and Niculescu-Mizil (2006)). The GBM hyper-parameters (shrinkage rate, interaction depth, minimum observation per node, and bag fraction) are optimized based on the minimum mean cross-validated error and a grid 165 search algorithm. Additionally, following the work of Chen et al. (2019), we exploit the potential spatial correlation of the



GBM residuals by interpolating and adding them to the predicted values. The interpolation is done with an Ordinary Kriging (OK) (Wackernagel, 2003).

One could think that skipping over the LUR computation by directly using all its time-invariant information (passive dosimeter campaigns, urban geometry, traffic-related data, and annual-averaged model results) as covariates in the Universal Kriging would simplify the workflow. However, there are two main drawbacks to doing so. On the one hand, in contrast to the GBM, the Universal Kriging assumes linear relations between covariates and the observed NO₂, which is not necessarily true for this case. On the other hand, when considering a large number of covariates with only 12 monitoring stations, strong spurious correlations lacking physical meaning are prone to happen, wrongly driving the final solution (Hengl et al. (2007)). Thus, gathering all static information in the single LUR covariate offers more robust results while permitting the addition of predictors using non-linear regression models.

2.4 Universal Kriging as a data fusion methodology for spatial bias correction

Microscale-LUR model and the hourly CALIOPE-Urban outputs are combined with observational NO₂ data from the monitoring stations using the geostatistical technique Universal Kriging, which is commonly used for spatial interpolation. This methodology predicts a random variable Z , in a target point \mathbf{x} , based on a combination between a multi-linear regression analysis with external variables f , referred as *covariates*, and a pure spatial interpolation considering the auto-correlation structure of the regression residuals. In our case, the variable Z corresponds to the monitoring data while the covariates are CALIOPE-Urban and our microscale-LUR model. A simple (multi)linear regression model is convenient here, given the low number (12) of available monitoring stations within the computational domain. Universal Kriging assumes the following relation (Cressie, 1993):

$$Z(\mathbf{x}) = \sum_{l=0}^L a_l f_l(\mathbf{x}) + R(\mathbf{x}), \quad (3)$$

where L equals 1 in the UK-DM approach and 2 in the UK-DM-LUR; a_l are the non-zero coefficients from the multi-linear regression between the observations and the covariates f_l , with $f_0(\mathbf{x}) = 1$ by convention; and $R(\mathbf{x})$ is the residual random field. The deterministic part of the variable Z is explained by a linear combination of the covariates, while the residual random field is considered to have zero mean and to be spatially auto-correlated. The main advantage of this method is that depending on the strength of the correlation between covariates and observations, Universal Kriging gives more weight either to the multi-linear regression or to the spatial interpolation of the residuals (Hengl (2009)). Thus, providing a robust data-fusion method that adapts to the quality of the model output.

As a Gaussian process, Universal Kriging estimates the variance of its predictions (σ^2) coming from both, the multi-linear regression (σ_{MLR}^2), and the spatial interpolation (σ_{SI}^2) steps, as



$$195 \quad \sigma^2(\mathbf{x}) = \underbrace{\sum_{\alpha=1}^m w_{\alpha} \cdot \gamma_R(\mathbf{x}_{\alpha} - \mathbf{x})}_{\sigma_{SI}^2} + \underbrace{\sum_{l=0}^L \lambda_l f_l(\mathbf{x})}_{\sigma_{MLR}^2}, \quad (4)$$

where m is the number of monitoring stations; w_{α} are the spatial interpolation weights associated with each measurement point; λ_l are the $L+1$ Lagrange multipliers used to minimize the variance error; and $\gamma_R(\mathbf{x}_{\alpha} - \mathbf{x})$ stands for the variogram, which characterizes the spatial structure of the residuals (Chiles and Delfiner (1999)). Thus, the variance of a prediction reflects how far the unmeasured location is from the observation points, as well as from the feature space in which the regression model has
 200 been calibrated, i.e., the extrapolation effect (Hengl (2009)). Our Universal Kriging implementation relies on the *R* package *gstat* (Pebesma (2004), Gräler et al. (2016)).

To normalize the distribution of the NO_2 data and to ensure positive predicted values, we have applied the Universal Kriging described above after transforming NO_2 data into the log-space. However, results need to be back-transformed to the original scale. Following the work of Cressie (1993), the back-transformed is performed as

$$205 \quad \hat{Z}(\mathbf{x}) = \exp(Z_l(\mathbf{x}) + \sigma_l^2(\mathbf{x})/2), \quad (5)$$

$$\hat{\sigma}^2(\mathbf{x}) = (\exp(\sigma_l^2(\mathbf{x})) - 1) \cdot \exp(2 \cdot Z_l(\mathbf{x}) + \sigma_l^2(\mathbf{x})), \quad (6)$$

where $\hat{Z}(\mathbf{x})$ and $\hat{\sigma}^2(\mathbf{x})$ respectively represent back-transformed prediction and variance at the target point, while $Z_l(\mathbf{x})$ and $\sigma_l^2(\mathbf{x})$ are prediction and variance in log-space, respectively.

210 Assuming a normal distribution of the error, the probability of exceedance (\mathcal{P}) a certain limit value (\mathcal{L}) can be computed as

$$\mathcal{P}(\mathbf{x}) = 1 - F\left(\frac{\mathcal{L} - \hat{Z}(\mathbf{x})}{\hat{\sigma}(\mathbf{x})}\right), \quad (7)$$

where F is the normal cumulative distribution function.

2.4.1 Statistical metrics to evaluation data-fusion skills

Statistical performance is assessed by Leave-One-Out-Cross-Validation (LOOCV), which consists of performing the data-
 215 fusion considering all monitoring stations except one kept to cross-validate the results. For each LOOCV we present the *Coefficient of Efficiency (COE)*, the *Root Mean Square Error (RMSE)*, the *Mean Bias (MB)*, and the *Correlation Coefficient (r)* defined as:

$$COE = 1 - \frac{\sum_{i=1}^k |M_i - O_i|}{\sum_{i=1}^k |O_i - O|} \quad (8)$$



$$MB = \frac{1}{k} \sum_{i=1}^k M_i - O_i \quad (9)$$

$$220 \quad r = \frac{1}{k-1} \sum_{i=1}^k \left(\frac{M_i - M}{\sigma_M} \right) \left(\frac{O_i - O}{\sigma_O} \right) \quad (10)$$

$$RMSE = \sqrt{\frac{1}{k} \sum_{i=1}^k (M_i - O_i)^2} \quad (11)$$

where k is the total number of observations; O_i and M_i are the observed and modelled i values, respectively; O and M are their respective means; and σ_O and σ_M refer to their standard deviation.

2.4.2 Spatial auto-correlation structure of NO₂ levels

225 In the Universal Kriging context, the variogram describes the spatial auto-correlation structure of the residual random field. In our case, the limited number of monitoring stations makes extracting a meaningful spatial structure challenging. For this reason, we estimate the residual variogram based on the dosimeters campaigns. This decision, however, entails a substantial limitation due to the assumption of a static variogram. We rely only on the IDAEA-CSIC campaign (discarding the xAire campaign for the variogram derivation) to avoid extra assumptions on combining campaigns. Additionally, we considered an isotropic variogram. All these assumptions impact the variance error estimated by the Universal Kriging (Brus and Heuvelink
 230 (2007)). To assess the impact of such assumptions, an analysis of the estimated variance in LOOCV is carried out in Sect. 3.2.

The variogram is fitted using the Matérn model with the Stein's parametrization implemented in the *automap* package (Hiemstra et al., 2008) setting the smoothing parameter $\kappa = 0.2$. The resulting variogram model is characterized by a $5 \cdot 10^{-2}$ *partial sill*, $3 \cdot 10^{-5}$ *nugget* and a *range* of 620 m. Following the work of Denby et al. (2007), we have optimized the *range* value
 235 to minimize the RMSE of the Universal Kriging. The *range* estimates the distance at which the data are no longer correlated. To optimize it, we performed a CV at all monitoring stations varying the range from 1 to 10 km every 1 km, keeping all other model parameters constant. We obtained the best results for the *range* of 5 km, improving the r by 4 %, the *COE* by 14 %, and the *RMSE* by -9 % with respect to the Universal Kriging using the original range value of 620 m.

2.4.3 Statistical quality assurance of the (multi)linear regression

240 The correlation coefficient and the slope of the regression model between covariates (CALIOPE-Urban and the microscale-LUR model) and observations are checked before including covariates in the Universal Kriging workflow as indicated in Fig. 1. If a covariate shows a low correlation (p -value > 0.05) with the observations at a specific hour, it is not considered in the regression model, as in the works of Zhang et al. (2021) and Oh et al. (2021). Additionally, if none of the covariates show a significant correlation, we use all covariates to build the regression model. However, to avoid nonphysical hourly maps, the




Figure 4. Scheme of the outer 10-fold CV and the inner 4-fold CV applied for the GBM training.

245 covariates are used only if their slope is positive, as suggested by Denby et al. (2007). In case all slopes are negative, or there are less than four observations available in a specific hour, the Universal Kriging is not performed and the results of the data-fusion method are directly the raw dispersion-model output. Following the above criteria, the dispersion model is not corrected 14 % and 2 % of the hours in 2019 for the UK-DM and UK-DM-LUR approaches, respectively.

3 Results and Discussion

250 Results are organized into two sections. Firstly, in Sect. 3.1, we estimate the microscale-LUR model performance and present the obtained NO₂ basemap. Secondly, in Sect. 3.2, the data fusion methodologies are discussed in terms of statistical performance, uncertainty quantification, and exceedance probability maps.

3.1 Microscale-LUR model

3.1.1 Performance assessment

255 The GBM-based microscale-LUR model is evaluated using two nested K-fold CV, the inner one for tuning the model (*Training-validation set*) and the outer one for testing the model on different parts of the dataset (*Test set*). Such a procedure aims at giving a reliable estimate of the expected performance. We use an outer 10-fold CV and an inner 4-fold CV as illustrated in Fig. 4. The tuning of the model is performed through a grid search over the following hyperparameters: shrinkage rate (with values ranging from 0.001 to 0.05 every 0.001), the interaction depth (from 1 to 4 every 1), the minimum observation in a node (from 260 5 to 15 every 1), and the bag fraction (0.5 and 0.65).

Results are given in Table 2, together with the performance reference of the annual mean NO₂ concentration obtained directly from CALIOPE-Urban. As explained in Sect. 2.3, we exploit the spatial auto-correlation of the LUR residuals to improve its estimation. To do so, the LUR residuals at the training locations are interpolated at the test locations by applying an OK. Then, they are added to the predictions to obtain the corrected results (*Test set adding the residuals* in Table 2).



Model		n	COE	MB ($\mu\text{g}/\text{m}^3$)	r	RMSE ($\mu\text{g}/\text{m}^3$)
	Training-validation set	7600	0.30	0.15	0.69	11.38
Microscale-LUR	Test set without adding the residuals	840	0.24	0.22	0.62	12.17
	Test set adding the residuals	840	0.27	-0.27	0.64	11.87
Raw CALIOPE-Urban	Annual mean	840	0.13	-0.81	0.54	13.68

Table 2. Statistical results of the microscale-LUR model in nested CV. The 2017 annual mean concentration of NO_2 of the raw dispersion model (CALIOPE-Urban) is also shown. The parameter n stands for the number of data points used to compute the statistics.




Figure 5. The left panel shows the raw annual mean of CALIOPE-Urban NO_2 concentrations versus the annualized passive dosimeters campaigns. The right and center panels correspond to the microscale-LUR results with and without the interpolated residuals, respectively. These figures use the test sets in which the performance of the microscale-LUR model has been assessed. The red dashed lines report passive dosimeter uncertainty ($\pm 25\%$) and the identity line is represented in blue. The statistical results are shown in Table 2.

265 Table 2 shows that the LUR model significantly improves the CALIOPE-Urban results. Also, the addition of the residuals slightly increases the statistical performance. The *Training-validation set* results are not perfectly fitted and are only slightly better than the *Test set* results, indicating that the LUR is not overfitted and has good capabilities to predict unseen data.

We show in Fig. 5 the scatter plots of the *annual mean CALIOPE-Urban* and the *Test dataset* results with and without adding the residuals, along with the observational uncertainty ranges indicated by the dashed red lines ($\pm 25\%$ according to Kuklinska et al. (2015)). Although a large portion of the predicted values for the LUR model with the residual correction lies within the uncertainty range, difficulty in predicting values of NO_2 higher than $80 \mu\text{g}/\text{m}^3$ can be observed. We attribute this behaviour to the limited number of points in this range, which can weaken the model training, particularly in the nested CV context, but also to the already poor predictive skills of CALIOPE-Urban in this concentration range.

275 Comparing these results with previous works, the resulting correlation coefficient (r) is lower than a LUR model fitted with the xAire campaign data ($r = 0.74$ in LOOCV) reported in Perelló et al. (2021a). However, Perelló et al. (2021a) used only 370 outdoor sampling sites out of the 669 available. They excluded samplers close to traffic and street intersections, achieving a



skilled urban LUR model. Even if the correlation coefficient is slightly lowered, we have considered all outdoor sampling sites (along with the IDAEA-CSIC campaign) to capture as much as possible the NO_2 spatial trends. On the other hand, the work of Munir et al. (2020) reported a microscale-LUR model based on 40 time-dependent LCS with slightly lower performance in CV than the present one ($r = 0.56$). Moreover, Munir et al. (2020) also reported a r value of 0.53 when the non-linear LUR model is based on the combination of 188 period-averaged and 40 time-dependent LCS. A key aspect of the present data-fusion methodology is that the microscale-LUR model results better explain the annualized passive dosimeters campaigns compared to the reference CALIOPE-Urban annual mean. Therefore, they are subsequently considered as a covariate in the Universal Kriging methodology, as further explained in the next Sect. 3.1.2.

285 3.1.2 Microscale-LUR basemap




Figure 6. (a) Resulting microscale-LUR basemap using all available sampling sites and adding the interpolated residuals, and (b) 2019 annual mean concentration of NO_2 of the raw dispersion model CALIOPE-Urban.

We proceed to train the microscale-LUR model with the residual correction using all available sampling sites. Figure 6 compares the long-term NO_2 patterns from the microscale-LUR basemap (Fig. 6a) with the NO_2 2019 annual mean of CALIOPE-Urban (Fig. 6b). Notice that the goal of the basemap is to correct the long-term spatial variability of NO_2 ; thus, Figure 6 highlights the differences in spatial patterns rather than differences in absolute NO_2 values. The resulting basemap shows a



290 qualitatively consistent NO₂ distribution: the major trafficked roads of the city and the port area are the most polluted locations, while the Collserola Mountains and the sea bordering the city have moderate NO₂ levels. Although both figures show similar NO₂ patterns, local differences from experimental information can be observed in Fig. 6a. As further shown in the statistical results, considering extensive passive dosimeters information through the microscale-LUR model avoids relying only on the urban model to describe the NO₂ gradients and significantly improves the data-fusion methodology.

295 The influence of each predictor in the final microscale-LUR model has been computed based on the methodology proposed by Friedman (2001) and implemented in the *R* package *gbm* (Ridgeway (2004)), in which the relative importance of each variable is associated to the reduction of the GBM cost function. Given the chosen set of predictors, the most influential variable is the NO₂ CALIOPE-Urban annual mean with a relative importance of 25.1 %, followed by 17.7 % for the traffic scaled variable and 15.7 % for the average building density. The other predictors exhibited a relative influence under the 15 %, 300 with the NO₂ CALIOPE regional mean as the lowest one with 4.3 %.

3.2 Data-fusion methodologies

3.2.1 Statistical evaluation

In order to quantify the added value of including the microscale-LUR basemap in the data-fusion methodology, two different post-processes (see Fig.1) have been carried out. First, the output of the urban dispersion model CALIOPE-Urban is merged 305 with the monitoring data using Universal Kriging, named UK-DM. Second, the microscale-LUR basemap is added as a covariate in the Universal Kriging workflow, named UK-DM-LUR.

Hourly statistical results for the raw CALIOPE-Urban, UK-DM and UK-DM-LUR are shown in Fig. 7 for each monitoring station using all available data of 2019. UK-DM and UK-DM-LUR results have been computed in LOOCV as explained in Sect. 2.4. *Gràcia* and *Eixample* are the urban traffic monitoring stations, and the last row in Fig. 7 corresponds to the average 310 results over all stations. This figure shows that the hourly scale post-processes consistently improve all studied statistical metrics at all monitoring stations, regardless of the monitoring station type. Moreover, adding the LUR basemap as a covariate (UK-DM-LUR) further improves the spatial correction at all stations and for all statistical metrics, except for the *MB* which does not have a clear trend. A negative value of the *COE* reflects a poor predictive capacity, so we highlight that both data-fusion methods achieve a positive *COE* at all considered stations. Almost all stations show a positive *MB* for CALIOPE-Urban 315 indicating a general overestimation of the model, while UK-DM and UK-DM-LUR present almost null *MB* averaged over all stations. The overestimation of CALIOPE-Urban in the monitoring stations may seem contradictory with the negative bias presented in Table 2 for the passive dosimeters campaigns. However, this could indicate that the highest NO₂ values in Barcelona city are not routinely monitored, as already pointed out in the work of Duyzer et al. (2015b). Regarding the *RMSE*, the averaged reduction between CALIOPE-Urban and UK-DM is about 32 %, and 24 % between UK-DM and UK-DM-LUR. 320 For *r*, an averaged improvement of 29 % between CALIOPE-Urban and UK-DM is observed, while the improvement is of 13 % between UK-DM and UK-DM-LUR.




Figure 7. Statistical results for each station after applying UK-DM and UK-DM-LUR to 2019 hourly data in LOOCV. In addition, we show the statistical results for the CALIOPE-Urban estimates at each station. The *All stations* row refers to the average over all stations.

3.2.2 Uncertainty quantification

The uncertainty of the Universal Kriging predictions is estimated from the (multi)linear regression and the spatial interpolation variances, as formulated in Sect. 2.4. The spatial interpolation is based on the variogram, which has been modelled from a period-averaged passive dosimeter campaign; thus, assuming a static behaviour as pointed out in Sect. 2.4.2. Additionally, the variogram is considered isotropic for simplicity, while we know that in the urban scale, the NO₂ autocorrelation structure may significantly vary depending on the direction with respect to traffic road-links. These assumptions directly impact the error variance estimated by the Universal Kriging, $\hat{\sigma}^2$. Considering that the interpolation error is normally distributed (N_{ref}), the observation at a specific monitoring station when performing a LOOCV should be within $\pm\hat{\sigma}$ of the predicted value 68% of the times, while 95% and 99,7% respectively for $\pm 2\hat{\sigma}$ and $\pm 3\hat{\sigma}$, as indicated in Table 3. To assess the normality of these distributions, Table 3 reports an empirical validation of the percentage of observations falling within the corresponding error range, again computed in LOOCV. These percentages show that uncertainty is underpredicted for both methods, being UK-DM-LUR overconfident results slightly better than the UK-DM ones.

To better understand the behaviour of uncertainty estimates, we show in Fig. 8 the probability density functions (PDFs) of the hourly bias, normalized by the error standard deviation $\hat{\sigma}$, for UK-DM and UK-DM-LUR using all studied monitoring stations in LOOCV over all available hours in 2019. The error PDFs have normal trends with a slightly negative skew and are



	$\pm 1\hat{\sigma}$	$\pm 2\hat{\sigma}$	$\pm 3\hat{\sigma}$
N_{ref}	68 %	95 %	99,7 %
UK-DM	47.9 %	78.0 %	91.3 %
UK-DM-LUR	51.2 %	81.3 %	92.9 %

Table 3. Percentages of observations falling in the $\pm\hat{\sigma}$, $\pm 2\hat{\sigma}$, $\pm 3\hat{\sigma}$ confidence intervals using all stations in LOOCV during 2019. Confidence intervals are computed based on the hourly predicted values and their standard deviation.

overconfident in accordance with Table 3. The negative skew and the overall null mean bias (shown in Fig. 7) indicate that samples with positive ($M_h > O_h$) bias are more frequent and have lower biases.




Figure 8. PDF of the hourly bias, normalized by the Universal Kriging standard deviation, for all monitoring stations in LOOCV during 2019. The PDFs correspond to the reference Normal distribution (N_{ref}), UK-DM and UK-DM-LUR hourly results.

In addition, in Fig. 9, the PDFs are computed by splitting the observed NO_2 concentration levels in three different ranges: 340 lesser than 40, greater than 100, and between 40 and 100 $\mu\text{g}/\text{m}^3$. These PDFs allow us to study the behavior of the error distribution for different NO_2 values. This figure shows that larger concentration levels tend to be underestimated, while the smaller ones are overestimated. In all ranges and for both methodologies, the normal trends of the error PDFs are conserved, being the intermediate ranges the closest to the theoretical normal distribution.

3.2.3 Street-scale maps

345 We first analyze the annual mean concentration levels of NO_2 . Table 4 presents the evaluation in LOOCV of the results post-processed by the UK-DM and the UK-DM-LUR methodologies applied directly to the 2019 annual mean. The presented statistics are computed using a single NO_2 averaged value for each station. The annual-based statistics are similar to the hourly




Figure 9. PDF by observed NO_2 ranges of the hourly bias, normalized by the standard deviation error, for all monitoring stations in LOOCV during 2019. The left panel corresponds to UK-DM, and the right panel to UK-DM-LUR.

results shown in Fig. 7; however, there is a substantial drop in the *RMSE* associated with the bias compensation when averaging the hourly data.

	COE	MB ($\mu\text{g}/\text{m}^3$)	r	RMSE ($\mu\text{g}/\text{m}^3$)
UK-DM	0.25	0.20	0.74	3.93
UK-DM-LUR	0.41	0.76	0.84	3.27

Table 4. Statistical results using the 12 monitoring stations after applying UK-DM and UK-DM-LUR directly to the annual mean in LOOCV.

350 Fig. 10 presents the NO_2 annual mean, their associated relative uncertainty, and probability maps of exceeding the $40 \mu\text{g}/\text{m}^3$ NO_2 annual limit value (AAQD 2008/EC/50) for both, UK-DM and UK-DM-LUR methodologies. The annual mean levels combining the raw model with the monitoring stations data (UK-DM) (Fig. 10a) have similar trends to the raw CALIOPE-Urban (Fig. 6b); however, pollution levels are significantly reduced. Adding the passive dosimeters information through the microscale-LUR basemap (UK-DM-LUR), in Fig. 10d, slightly increases NO_2 concentrations, particularly in the city center and secondary roads, where the microscale-LUR basemap (Fig. 6a) exhibits steeper NO_2 gradients than CALIOPE-Urban (Fig. 355 6b).

As expected, the areas surrounding the monitoring stations (presented in Fig.2) show lower relative uncertainty, as can be seen in Figs. 10b and 10e. The higher uncertainty regions, on the other hand, correspond to areas far from monitoring sites and with extreme concentration levels which causes an extrapolation effect in the regression model. When comparing the two uncertainty maps (Figs. 10b and 10e), UK-DM-LUR has regions with higher relative uncertainty than the UK-DM. This behaviour is due to the addition of the microscale-LUR covariate, which increases the standard deviation associated with the regression model. In addition, some localized regions of high uncertainty can be observed in Fig. 10e. They are associated 360




Figure 10. (a) NO₂ 2019 annual map resulting from applying UK-DM with the annual values, (b) Relative uncertainty associated with the predictions in (a), (c) Annual probability map of exceeding the $40 \mu\text{g}/\text{m}^3$ NO₂ limit value using the values in (a) and (b), (d) NO₂ 2019 annual map resulting from applying UK-DM-LUR with the annual values, (e) Relative uncertainty associated to the predictions in (d), and (f) Annual probability map of exceeding the $40 \mu\text{g}/\text{m}^3$ NO₂ limit value using the values in (d) and (e).




Figure 11. (a) Hourly NO₂ concentration map resulting of applying the UK-DM-LUR methodology at 9h UTC on 28/02/2019. (b) Relative uncertainty associated to the predictions in (a), (c) Hourly probability map of exceeding the 200 µg/m³ NO₂ hourly averaged limit value using the UK-DM-LUR method at 9h UTC on 28/02/2019.

with passive dosimeters' locations and trafficked roads where the microscale-LUR covariate has caused an increase in NO₂ concentrations, rising the level of extrapolation in the regression model.

365 Regardless of the data-fusion method, the most polluted regions correspond to probabilities exceeding the annual limit above 0.7, as shown in Figs. 10c and 10f. When considering the UK-DM-LUR method, 13 % of the Barcelona municipality area has 0.7 or higher probabilities of exceeding the annual limit; and this percentage rises to 30 % when considering probabilities equal to or higher than 0.5. The *Eixample* district, which is the most polluted while being the most populous and densely populated (approximately 270000 inhabitants and 36000 inhabitants per square kilometer (Ajuntament de Barcelona, 2019)),
370 has 95 % of its area exceeding the annual limit with a probability equal to or higher to 0.5, and 69 % in the case of 0.7. Thus, significant evidence indicates that the annual legal limit was broadly exceeded in Barcelona in 2019. Stronger evidence could be obtained by reducing the uncertainty associated with the results, either by a better correlated urban model or by increasing the monitoring system's coverage. To test a more restrictive threshold, we have analyzed the exceedance probability annual maps using the recommended WHO 2021 annual limit of 10 µg/m³ (not shown here), obtaining probabilities above 0.9 over
375 all the domain for both methodologies.



Figure 11 presents the NO₂ prediction at a specific hour, its associated relative uncertainty, and the exceedance probability map based on the 200 $\mu\text{g}/\text{m}^3$ NO₂ hourly threshold (AAQD 2008/EC/50) for the UK-DM-LUR methodology. The goal is to illustrate that, apart from studying the long-term NO₂ mean values, the present methodology can also be used to correct short NO₂ exposure episodes such as the ones observed during traffic rush hours. Figure 11 corresponds to the peak traffic hour at 9 UTC on February 28, 2019, which was a particularly polluted hour reporting 138 and 201 $\mu\text{g}/\text{m}^3$ at the traffic monitoring stations of *Eixample* and *Gràcia*, respectively. Similar to Fig. 10, low uncertainty regions are obtained around the monitoring stations' locations. Likewise, high relative uncertainty regions are associated with pollution hot-spots due to the extrapolation effect in the regression step. Concerning the exceedance probability maps shown in Fig. 11c, the city center and its major trafficked streets have the highest values (> 0.7). For the *Eixample* district, 19 % of the area exceeds the hourly limit with a probability equal to or higher than 0.5, and 6 % in the case of 0.7.

4 Conclusions

The present work assesses the added value of including a microscale-LUR basemap into a data-fusion method to obtain spatially bias-corrected urban maps of NO₂ at the hourly scale. To do so, we have compared two different data-fusion methods: (i) merging an urban dispersion model with the observational data of 12 monitoring stations using Universal Kriging (UK-DM), and (ii) adding to UK-DM a non-linear LUR model as a covariate in the Kriging workflow based on the GBM algorithm (UK-DM-LUR). The comparison is based on the statistical performance in LOOCV at each monitoring station, the resulting NO₂ maps, and their associated uncertainty.

The statistical performance of the microscale-LUR model has been assessed using a comprehensive nested cross-validation. The obtained microscale-LUR basemap ($r = 0.64$, $RMSE = 11.87 \mu\text{g}/\text{m}^3$) outperformed the raw annual-averaged dispersion model results ($r = 0.54$, $RMSE = 13.68 \mu\text{g}/\text{m}^3$), highlighting the convenience of using passive dosimeters campaigns to explain the spatial distribution of NO₂. Moreover, a novel traffic density variable based on the combination of different traffic buffer sizes has been shown to have a significant influence (17.7 %) in the microscale-LUR basemap, suggesting its relevance in future microscale-LUR models.

Adding the microscale-LUR time-invariant spatial information (UK-DM-LUR) has proven to significantly improve the skills of the more straightforward data-fusion UK-DM method at the hourly scale, increasing the correlation coefficient (r) by 13 % and reducing the $RMSE$ by -24 % in average over all monitoring stations during 2019. Thus, our results suggest that data-fusion methods applied at the street-scale benefit from high-spatial resolution data such as passive dosimeters campaigns, urban morphology, or traffic intensity estimates. When only using monitoring stations in the data-fusion approach, the spatial patterns of NO₂ rely mainly on the urban model patterns. Generally, the better the temporal and spatial coverage of observational data, the better statistical performance can be achieved.

To check the consistency of the estimated uncertainty, we have empirically validated the UK-based uncertainties through a LOOCV. Despite the Universal Kriging's predicted variance is slightly overconfident and tends to degrade for extreme concentration values, we found that it is a meaningful estimate of the uncertainty. The PDFs of the error are close to the



normal distribution, especially for the UK-DM-LUR approach. The spatial characterization of the uncertainty adds value to
410 the NO₂ concentration maps, making data-fusion results more comprehensive for regulatory purposes, decision-makers, and
health impact assessment. For instance, uncertainty maps can be used to allocate new observational stations or to plan future
LCS campaigns. In this regard, our results show that pollution hot-spots are areas of high uncertainty underrepresented by the
current monitoring system. We thus stress the need to monitor the vicinity of heavily trafficked roads better to increase the
performance of data-fusion methods in predicting hourly and annual exceedances.

415 *Code and data availability.* The R source code, input data, and results are available via Zenodo on Criado et al. (2022).

Author contributions. AC implemented the data-fusion code and generated the figures. AC and JMA conducted the study and drafted the
manuscript. JMA and JB processed the CALIOPE-Urban data. HP supported the validation of the microscale-LUR model. All authors
contributed to the analysis and objectives of the document, and internally reviewed and supported the text.

Competing interests. The authors declare that they have no conflict of interest.

420 *Acknowledgements.* We acknowledge support from the Ministerio de Ciencia, Innovación y Universidades (MICINN) as part of the BROWN-
ING project RTI2018-099894-BI00, from the VITALISE project (PID2019-108086RA-I00) funded by the MCIN/AEI/10.13039/501100011033,
the MITIGATE project (PID2020-116324RA695 I00 / AEI /10.13039/501100011033) from the Agencia Estatal de Investigacion (AEI), and
the AXA Research Fund. This project has also received funding from the European Union's Horizon 2020 research and innovation pro-
gramme under the Marie Skłodowska-Curie grant agreement H2020-MSCA-COFUND-2016-754433. BSC researchers thankfully acknowl-
425 edge the computer resources at Marenstrum and the technical support provided by Barcelona Supercomputing Center (RES-AECT-2021-1-
0027, RES-AECT-2021-2-0001).



References

- Ajuntament de Barcelona: Open Data BCN, <https://opendata-ajuntament.barcelona.cat/es>, under license Creative Commons by 4.0, 2019.
- Auvinen, M., Järvi, L., Hellsten, A., Rannik, Ü., and Vesala, T.: Numerical framework for the computation of urban flux footprints employing large-eddy simulation and Lagrangian stochastic modeling, *Geoscientific Model Development*, 10, 4187–4205, 2017.
- 430 Baldasano Recio, J. M., Pay Pérez, M. T., Jorba, O., Gassó, S., and Jiménez-Guerrero, P.: An annual assessment of air quality with the CALIOPE modeling system over Spain, *Science of the Total Environment*, 2011, vol. 409, num. 11, p. 2163-2178, 2011.
- Beelen, R., Hoek, G., Vienneau, D., Eeftens, M., Dimakopoulou, K., Pedeli, X., Tsai, M.-Y., Künzli, N., Schikowski, T., Marcon, A., Eriksen, K. T., Raaschou-Nielsen, O., Stephanou, E., Patelarou, E., Lanki, T., Yli-Tuomi, T., Declercq, C., Falq, G., Stempfelet, M., Birk, M., Cyrus, J., von Klot, S., Nádor, G., Varró, M. J., Dèdelè, A., Gražulevičienė, R., Mölter, A., Lindley, S., Madsen, C., Cesaroni, G., Ranzi, A., Badaloni, C., Hoffmann, B., Nonnemacher, M., Krämer, U., Kuhlbusch, T., Cirach, M., de Nazelle, A., Nieuwenhuijsen, M., Bellander, T., Korek, M., Olsson, D., Strömgen, M., Dons, E., Jerrett, M., Fischer, P., Wang, M., Brunekreef, B., and de Hoogh, K.: Development of NO₂ and NO_x land use regression models for estimating air pollution exposure in 36 study areas in Europe – The ESCAPE project, *Atmospheric Environment*, 72, 10–23, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2013.02.037>, 2013.
- 440 Benavides, J., Snyder, M., Guevara, M., Soret, A., Pérez García-Pando, C., Amato, F., Querol, X., and Jorba, O.: CALIOPE-Urban v1.0: coupling R-LINE with a mesoscale air quality modelling system for urban air quality forecasts over Barcelona city (Spain), *Geoscientific Model Development*, 12, 2811–2835, 2019.
- Benavides, J., Guevara, M., Snyder, M. G., Rodríguez-Rey, D., Soret, A., García-Pando, C. P., and Jorba, O.: On the impact of excess diesel NO_x emissions upon NO₂ pollution in a compact city, *Environmental Research Letters*, 16, 024024, 2021.
- 445 Briggs, D. J., Collins, S., Elliot, P., FISCHER, P., KINGHAM, S., LEBRET, E., PRYL, K., REEUWIJK, H. V., SMALLBONE, K., and VEEN, A. V. D.: Mapping urban air pollution using GIS: a regression-based approach, *International Journal of Geographical Information Science*, 11, 699–718, <https://doi.org/10.1080/136588197242158>, 1997.
- Brus, D. J. and Heuvelink, G. B.: Optimization of sample patterns for universal kriging of environmental variables, *Geoderma*, 138, 86–95, <https://doi.org/https://doi.org/10.1016/j.geoderma.2006.10.016>, 2007.
- 450 Caruana, R. and Niculescu-Mizil, A.: An empirical comparison of supervised learning algorithms, in: *Proceedings of the 23rd international conference on Machine learning*, pp. 161–168, 2006.
- Chen, J., de Hoogh, K., Gulliver, J., Hoffmann, B., Hertel, O., Ketzel, M., Bauwelinck, M., van Donkelaar, A., Hvidtfeldt, U. A., Katsouyanni, K., Janssen, N. A., Martin, R. V., Samoli, E., Schwartz, P. E., Stafoggia, M., Bellander, T., Strak, M., Wolf, K., Vienneau, D., Vermeulen, R., Brunekreef, B., and Hoek, G.: A comparison of linear regression, regularization, and machine learning algorithms to develop Europe-wide spatial models of fine particles and nitrogen dioxide, *Environment International*, 130, 104934, <https://doi.org/https://doi.org/10.1016/j.envint.2019.104934>, 2019.
- Chiles, J.-P. and Delfiner, P.: *Geostatistics: modeling spatial uncertainty*, Wiley, New York, 1999.
- Cressie: *Statistics for Spatial Data*, chap. 1, pp. 1–26, John Wiley & Sons, Ltd, <https://doi.org/https://doi.org/10.1002/9781119115151.ch1>, 1993.
- 460 Criado, A., Mateu Armengol, J., Petetin, H., Rodríguez-Rey, D., Benavides, J., Guevara, M., Pérez García-Pando, C., Soret, A., and Jorba, O.: Code and data set from data fusion uncertainty-enabled methods to map street-scale hourly NO₂ in Barcelona city: a case study with CALIOPE-Urban v1.0, <https://doi.org/10.5281/zenodo.7185913>, 2022.



- Denby, B.: Guide on modelling Nitrogen Dioxide (NO₂) for air quality assessment and planning relevant to the European Air Quality Directive, ETC/ACM Technical Paper 2011/15, European Topic Centre on Air Pollution and Climate Change Mitigation, 2011.
- 465 Denby, B., Horálek, J., de Smet, P., de Leeuw, F., and Kurfürst, P.: European scale exceedance mapping for PM₁₀ and ozone based on daily interpolation fields, ETC/ACC Technical paper, 8, 2007.
- Denby, B. R., Gauss, M., Wind, P., Mu, Q., Grøtting Wærsted, E., Fagerli, H., Valdebenito, A., and Klein, H.: Description of the uEMEP_v5 downscaling approach for the EMEP MSC-W chemistry transport model, *Geoscientific Model Development*, 13, 6303–6323, 2020.
- Dimakopoulou, K., Samoli, E., Analitis, A., Schwartz, J., Beevers, S., Kitwiroon, N., Beddows, A., Barratt, B., Rodopoulou, S., Zafeiratou, S., et al.: Development and Evaluation of Spatio-Temporal Air Pollution Exposure Models and Their Combinations in the Greater London Area, UK, *International Journal of Environmental Research and Public Health*, 19, 5401, 2022.
- 470 S., et al.: Development and Evaluation of Spatio-Temporal Air Pollution Exposure Models and Their Combinations in the Greater London Area, UK, *International Journal of Environmental Research and Public Health*, 19, 5401, 2022.
- Duyzer, J., van den Hout, D., Zandveld, P., and van Ratingen, S.: Representativeness of air quality monitoring networks, *Atmospheric Environment*, 104, 88–101, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2014.12.067>, 2015a.
- Duyzer, J., van den Hout, D., Zandveld, P., and van Ratingen, S.: Representativeness of air quality monitoring networks, *Atmospheric Environment*, 104, 88–101, 2015b.
- 475 Environment, 104, 88–101, 2015b.
- Friedman, J. H.: Greedy function approximation: a gradient boosting machine, *Annals of statistics*, pp. 1189–1232, 2001.
- Gryparis, A., Paciorek, C. J., Zeka, A., Schwartz, J., and Coull, B. A.: Measurement error caused by spatial misalignment in environmental epidemiology, *Biostatistics*, 10, 258–274, 2009.
- Gräler, B., Pebesma, E., and Heuvelink, G.: Spatio-Temporal Interpolation using gstat, *The R Journal*, 8, 204–218, <https://journal.r-project.org/archive/2016/RJ-2016-014/index.html>, 2016.
- 480 org/archive/2016/RJ-2016-014/index.html, 2016.
- Guevara, M., Tena, C., Porquet, M., Jorba, O., and Pérez García-Pando, C.: HERMESv3, a stand-alone multi-scale atmospheric emission modelling framework–Part 2: The bottom–up module, *Geoscientific Model Development*, 13, 873–903, 2020.
- Hengl, T.: A practical guide to geostatistical mapping, Hengl Amsterdam, 2009.
- Hengl, T., Heuvelink, G. B., and Rossiter, D. G.: About regression-kriging: From equations to case studies, *Computers & Geosciences*, 33, 1301–1315, <https://doi.org/https://doi.org/10.1016/j.cageo.2007.05.001>, *spatial Analysis*, 2007.
- 485 1301–1315, <https://doi.org/https://doi.org/10.1016/j.cageo.2007.05.001>, *spatial Analysis*, 2007.
- Hiemstra, P., Pebesma, E., Twenh"ofel, C., and Heuvelink, G.: Real-time automatic interpolation of ambient gamma dose rates from the Dutch Radioactivity Monitoring Network, *Computers & Geosciences*, DOI: <http://dx.doi.org/10.1016/j.cageo.2008.10.011>, 2008.
- Hoek, G., Beelen, R., de Hoogh, K., Vienneau, D., Gulliver, J., Fischer, P., and Briggs, D.: A review of land-use regression models to assess spatial variation of outdoor air pollution, *Atmospheric Environment*, 42, 7561–7578, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2008.05.057>, 2008.
- 490 https://doi.org/https://doi.org/10.1016/j.atmosenv.2008.05.057, 2008.
- Hood, C., Stocker, J., Seaton, M., Johnson, K., O'Neill, J., Thorne, L., and Carruthers, D.: Comprehensive evaluation of an advanced street canyon air pollution model, *Journal of the Air & Waste Management Association*, 71, 247–267, 2021.
- Horálek, J., Denby, B., de Smet, P., de Leeuw, F., Kurfürst, P., Swart, R., and van Noije, T.: Spatial mapping of air quality for European scale assessment, Tech. rep., ETC/ACC, 2006.
- 495 ICGC: Orthopoto of Catalunya, Generalitat de Catalunya, Institut Cartogràfic i Geològic de Catalunya (ICGC), <http://www.icc.cat/appdownloads/?c=dlftopo5m>, under license Creative Commons by 4.0.
- ISGlobal: ISGlobal ranking of cities, <https://isglobalranking.org/>, 2021.
- Jorba, O., Pérez, C., Rocadenbosch, F., and Baldasano, J.: Cluster analysis of 4-day back trajectories arriving in the Barcelona area, Spain, from 1997 to 2002, *Journal of Applied Meteorology*, 43, 887–901, 2004.



- 500 Kahle, D. and Wickham, H.: ggmap: Spatial Visualization with ggplot2, *The R Journal*, 5, 144–161, <https://journal.r-project.org/archive/2013-1/kahle-wickham.pdf>, 2013.
- Khomenko, S., Cirach, M., Pereira-Barboza, E., Mueller, N., Barrera-Gómez, J., Rojas-Rueda, D., de Hoogh, K., Hoek, G., and Nieuwenhuijsen, M.: Premature mortality due to air pollution in European cities: a health impact assessment, *The Lancet Planetary Health*, 5, e121–e134, [https://doi.org/https://doi.org/10.1016/S2542-5196\(20\)30272-2](https://doi.org/https://doi.org/10.1016/S2542-5196(20)30272-2), 2021.
- 505 Kim, Y., Wu, Y., Seigneur, C., and Roustan, Y.: Multi-scale modeling of urban air pollution: development and application of a Street-in-Grid model (v1. 0) by coupling MUNICH (v1. 0) and Polair3D (v1. 8.1), *Geoscientific Model Development*, 11, 611–629, 2018.
- Kuhn, M.: Building Predictive Models in R Using the caret Package, *Journal of Statistical Software, Articles*, 28, 1–26, <https://doi.org/10.18637/jss.v028.i05>, 2008.
- Kuklinska, K., Wolska, L., and Namiesnik, J.: Air quality policy in the US and the EU—a review, *Atmospheric Pollution Research*, 6, 129–137, 510 2015.
- Kwak, K.-H., Baik, J.-J., Ryu, Y.-H., and Lee, S.-H.: Urban air quality simulation in a high-rise building area using a CFD model coupled with mesoscale meteorological and chemistry-transport models, *Atmospheric Environment*, 100, 167–177, 2015.
- Mijling, B.: High-resolution mapping of urban air quality with heterogeneous observations: a new methodology and its application to Amsterdam, *Atmospheric Measurement Techniques*, 13, 4601–4617, <https://doi.org/10.5194/amt-13-4601-2020>, 2020.
- 515 Munir, S., Mayfield, M., Coca, D., and Mihaylova, L. S.: A nonlinear land use regression approach for modelling NO₂ concentrations in urban areas—Using data from low-cost sensors and diffusion tubes, *Atmosphere*, 11, 736, 2020.
- Natekin, A. and Knoll, A.: Gradient boosting machines, a tutorial, *Frontiers in Neurorobotics*, 7, <https://doi.org/10.3389/fnbot.2013.00021>, 2013.
- Oh, I., Hwang, M.-K., Bang, J.-H., Yang, W., Kim, S., Lee, K., Seo, S., Lee, J., and Kim, Y.: Comparison of dif- 520 ferent hybrid modeling methods to estimate intraurban NO₂ concentrations, *Atmospheric Environment*, 244, 117907, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2020.117907>, 2021.
- Palmes, E., Gunnison, A., DiMattio, J., and Tomczyk, C.: Personal sampler for nitrogen dioxide, *American Industrial Hygiene Association Journal*, 37, 570–577, 1976.
- Pay, M. T., Martínez, F., Guevara, M., and Baldasano, J. M.: Air quality forecasts on a kilometer-scale grid over complex Spanish terrains, 525 *Geoscientific Model Development*, 7, 1979–1999, <https://doi.org/10.5194/gmd-7-1979-2014>, 2014.
- Pebesma, E. J.: Multivariable geostatistics in S: the gstat package, *Computers & Geosciences*, 30, 683–691, 2004.
- Perelló, J., Cigarini, A., Vicens, J., Bonhoure, I., Rojas-Rueda, D., Nieuwenhuijsen, M. J., Cirach, M., Daher, C., Targa, J., and Ripoll, A.: Large-scale citizen science provides high-resolution nitrogen dioxide values and health impact while enhancing community knowledge and collective action, *Science of The Total Environment*, 789, 147750, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2021.147750>, 530 2021a.
- Perelló, J., Cigarini, A., Vicens, J., Bonhoure, I., Rojas-Rueda, D., Nieuwenhuijsen, M. J., Cirach, M., Daher, C., Targa, J., and Ripoll, A.: Data set from large-scale citizen science provides high-resolution nitrogen dioxide values for enhancing community knowledge and collective action to related health issues, *Data in Brief*, 37, 107269, <https://doi.org/https://doi.org/10.1016/j.dib.2021.107269>, 2021b.
- PNOA: Ministerio de transportes, movilidad y agenda urbana: LIDAR, https://pnoa.ign.es/productos_lidar, under license Creative Commons 535 by 4.0 scne.es, 2020.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <http://www.R-project.org/>, 2013.



- Ridgeway, G.: Gbm: Generalized Boosted Regression Models. R Package, 1.5, R package version, 1, 2004.
- Rivas, I., Viana, M., Moreno, T., Pandolfi, M., Amato, F., Reche, C., Bouso, L., Álvarez-Pedrerol, M., Alastuey, A., Sunyer, J., et al.: Child
540 exposure to indoor and outdoor air pollutants in schools in Barcelona, Spain, *Environment international*, 69, 200–212, 2014.
- Rodriguez-Rey, D., Guevara, M., Linares, M. P., Casanovas, J., Armengol, J. M., Benavides, J., Soret, A., Jorba, O., Tena, C., and García-
Pando, C. P.: To what extent the traffic restriction policies applied in Barcelona city can improve its air quality?, *Science of the Total
Environment*, 807, 150 743, 2022.
- Santiago, J. L., Martín, F., and Martilli, A.: A computational fluid dynamic modelling approach to assess the representativeness of urban
545 monitoring stations, *Science of The Total Environment*, 454-455, 61–72, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2013.02.068>,
2013.
- Schneider, P., Castell, N., Vogt, M., Dauge, F. R., Lahoz, W. A., and Bartonova, A.: Mapping urban air quality in
near real-time using observations from low-cost sensors and model information, *Environment International*, 106, 234–247,
<https://doi.org/https://doi.org/10.1016/j.envint.2017.05.005>, 2017.
- 550 Snyder, M. G., Venkatram, A., Heist, D. K., Perry, S. G., Petersen, W. B., and Isakov, V.: RLINE: A line source dispersion model for
near-surface releases, *Atmospheric environment*, 77, 748–756, 2013.
- Soulhac, L., Nguyen, C. V., Volta, P., and Salizzoni, P.: The model SIRANE for atmospheric urban pollutant dispersion. PART III: Validation
against NO₂ yearly concentration measurements in a large urban agglomeration, *Atmospheric environment*, 167, 377–388, 2017.
- Su, J. G., Jerrett, M., Beckerman, B., Wilhelm, M., Ghosh, J. K., and Ritz, B.: Predicting traffic-related air pol-
555 lution in Los Angeles using a distance decay regression selection strategy, *Environmental Research*, 109, 657–670,
<https://doi.org/https://doi.org/10.1016/j.envres.2009.06.001>, 2009.
- Tilloy, A., Mallet, V., Poulet, D., Pesin, C., and Brocheton, F.: BLUE-based NO₂ data assimilation at urban scale, *Journal of Geophysical
Research*, 118, 2031–2040, <https://doi.org/10.1002/jgrd.50233>, 2013.
- Valencia, A., Venkatram, A., Heist, D., Carruthers, D., and Arunachalam, S.: Development and evaluation of the R-LINE model algorithms
560 to account for chemical transformation in the near-road environment, *Transportation Research Part D: Transport and Environment*, 59,
464–477, <https://doi.org/https://doi.org/10.1016/j.trd.2018.01.028>, 2018.
- Vardoulakis, S., Gonzalez-Flesca, N., Fisher, B. E., and Pericleous, K.: Spatial variability of air pollution in the vicinity of a permanent mon-
itoring station in central Paris, *Atmospheric Environment*, 39, 2725–2736, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2004.05.067>,
fourth International Conference on Urban Air Quality: Measurement, Modelling and Management, 25-28 March 2003, 2005.
- 565 Venkatram, A., Snyder, M. G., Heist, D. K., Perry, S. G., Petersen, W. B., and Isakov, V.: Re-formulation of plume spread for near-surface
dispersion, *Atmospheric environment*, 77, 846–855, 2013.
- Wackernagel, H.: Ordinary kriging, in: *Multivariate geostatistics*, pp. 79–88, Springer, 2003.
- Wang, S., Ma, Y., Wang, Z., Wang, L., Chi, X., Ding, A., Yao, M., Li, Y., Li, Q., Wu, M., Zhang, L., Xiao, Y., and Zhang, Y.: Mobile
monitoring of urban air quality at high spatial resolution by low-cost sensors: impacts of COVID-19 pandemic lockdown, *Atmospheric
570 Chemistry and Physics*, 21, 7199–7215, <https://doi.org/10.5194/acp-21-7199-2021>, 2021.
- Weissert, L., Alberti, K., Miskell, G., Pattinson, W., Salmond, J., Henshaw, G., and Williams, D. E.: Low-cost sensors and microscale land
use regression: Data fusion to resolve air quality variations with high spatial and temporal resolution, *Atmospheric environment*, 213,
285–295, 2019.
- WHO: WHO (World Health Organization) global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide,
575 sulfur dioxide and carbon monoxide, World Health Organization, 2021.



Wickham, H.: ggplot2: Elegant Graphics for Data Analysis, Springer-Verlag New York, <https://ggplot2.tidyverse.org>, 2016.

Zhang, X., Just, A. C., Hsu, H.-H. L., Kloog, I., Woody, M., Mi, Z., Rush, J., Georgopoulos, P., Wright, R. O., and Stroustrup, A.: A hybrid approach to predict daily NO₂ concentrations at city block scale, *Science of The Total Environment*, 761, 143 279, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.143279>, 2021.