

Uncertainty analysis of single- and multiple-size-class frazil ice models

Fabien Souillé¹, Cédric Goeury¹, and Rem-Sophia Mouradi^{2,3}

¹EDF R&D, National Laboratory for Hydraulics and Environment (LNHE), 6 Quai Watier, 78400 Chatou, France

²EDF R&D, Fluid Dynamics, Energy and Environment Department (MFEE), 6 Quai Watier, 78400 Chatou, France

³CEREA (Centre d'Enseignement et de Recherche en Environnement Atmosphérique), Joint Laboratory École des Ponts ParisTech and EDF R&D, Université Paris-Est, 77455 Marne-la-Vallée, France

Correspondence: Fabien Souillé (fabien.souille@edf.fr)

Abstract. The formation of frazil ice in supercooled waters has been extensively studied, both experimentally and numerically, in recent years. Numerical models, with varying degrees of complexity, have been proposed; these are often based on many parameters, the values of which are uncertain and difficult to estimate. In this paper, an uncertainty analysis of two mathematical models that simulate supercooling and frazil ice formation is carried out within a probabilistic framework. The two main goals are (i) to provide quantitative insight into the relative importance of contributing uncertain parameters, to help identify parameters for optimal calibration, and (ii) to compare the output scatter of frazil ice models with single and multiple crystal size classes. The derivation of single- and multi-class models is presented in light of recent work, their numerical resolution is discussed, and a list of the main uncertain parameters is proposed. An uncertainty analysis is then carried out in three steps. Parameter uncertainty is first quantified, based on recent field, laboratory and numerical studies. Uncertainties are then propagated through the models using Monte Carlo simulations. Finally, the relative influence of uncertain parameters on the output time series - i.e. the total frazil volume fraction and water temperature - is assessed by means of Sobol indices. The influence of input parameters on the long-term asymptotic as well as short-term transient evolution of the systems is discussed, depending on whether gravitational removal is included or not in the models.

1 Introduction

Formation of frazil ice in water bodies has been widely investigated because of its impact on submerged structures (Daly, 1991, 2006; Richard and Morse, 2008) and because it often precedes formation of ice cover in rivers and oceans (Daly, 1994; Smedsrud and Jenkins, 2004). Frazil ice also plays an important role in geophysical flows such as plumes of ice shelf water under floating ice sheets (Bombosch and Jenkins, 1995; Rees Jones and Wells, 2018; Frazer et al., 2020). For these reasons, the study of frazil formation processes is an active area of research, with a large variety of applications in river and coastal engineering.

The main drivers for the formation of frazil are the water cooling rate resulting from heat exchanges with the atmosphere, the initial seeding of frazil nuclei and the turbulent mixing. Then, the thermal growth process deriving from the heat exchange between water and primitive crystals, allows a fine description of the balance between growth frazil ice and water supercooling.

Previous mathematical models describing the evolution of frazil ice and water temperature, were based on ideas pioneered by
25 Daly (1984, 1994). Omstedt (1985) developed a model based on a turbulent channel-flow boundary-layer theory, in which a
mean particle is considered of constant geometric properties and constant Nusselt number is considered. Mean particle-size
models have been incorporated in numerical hydraulic tools such as CRISSP2D (Shen and Wasantha Lal, 1993; Shen et al.,
1995; Shen, 2010). More complex models are based on the ice-number continuity equation introduced by Daly (1984), taking
30 into account the complexity of crystal distribution. In contrast with single-size-class models, the crystal number continuity
equations allow introduction of secondary nucleation and flocculation processes. When combined with thermal growth, these
complement the modeling of frazil crystal size evolution. Svensson and Omstedt (1994) proposed a numerical model for
solving the equations introduced by Daly (1984) by considering discrete radius intervals. Conceptual models for secondary
nucleation and flocculation were introduced and calibrated to fit the experimental data of Michel (1963) and Carstens (1966).
Hammar and Shen (1991) derived a variable particle-size model in two dimensions, and later improved it with secondary
35 nucleation and flocculation (Hammar and Shen, 1995). Variable particle-size models has also been integrated in TELEMAC-
MASCARET (Souillé et al., 2020). Numerous simplified implementations have been made that consider a well-mixed water
body. For example, Wang and Doering (2005) worked with the same model as Svensson and Omstedt (1994) and further
discussed calibration of influential parameters in initial seeding, secondary nucleation, and flocculation by comparing them
with the experimental data of Clark and Doering (2004). Implementation of multiple size class frazil dynamics in the context
40 of sea ice and ice shelf water plumes has also been proposed by Smedsrud (2002), Smedsrud and Jenkins (2004) and Holland
and Feltham (2005). In the same context, Rees Jones and Wells (2018) have recently shed new light on multiple-size-class
models and discussed crystal growth rate, and the occurrence of frazil explosion. They also identified and characterized steady
state crystal size distribution. Henceforth, single- and multiple-size-class models will be referred to as (SSC)¹ and (MSC)²
models respectively.

45 Common to all numerical model of frazil dynamics is their use of a large number of parameters, making calibration difficult.
The modeling studies mentioned above show that frazil ice models can be fitted to reproduce evolution of temperature and frazil
volume fraction. However, given the uncertainty of fitting parameters, it is questionable whether these models are predictive.
As Rees Jones and Wells (2018) point out, the consistency between experiments and models does not necessarily means
parameterization is correct. This is particularly true when different processes compete and have similar impact on the crystal
50 size distribution. Besides, introducing new processes in the models increases the number of parameters that need calibration.
This comes at the cost of introducing new uncertain parameters and raises the question of models trustworthiness.

This trade-off between uncertainties and model complexity, is often discussed in geophysical and environmental modeling
(MacGillivray, 2021; Van Zelm and Huijbregts, 2013). On this matter Saltelli (2019) advises use of statistics to aid mathemat-
ical modeling via “*a systemic appraisal of model uncertainties and parametric sensitivities*”. There is, moreover, a growing
55 appreciation of how sensitivity analysis enhance understanding of the intricate physical processes involved in ice formation
(Sheikholeslami et al., 2017). The sensitivity of multiple-size-class frazil models to initial seeding, secondary nucleation and

¹SSC: Single Size Class

²MSC: Multiple Size Class

flocculation parameters has been investigated by Wang and Doering (2005) and Smedsrud and Jenkins (2004), by simple modifications of specific parameters. Although probabilistic methods are sometimes used in processing observational data (Frazer et al., 2020), to the author’s knowledge, probabilistic sensitivity analysis of frazil ice models has never been performed. In addition, comparison of (SSC) and (MSC) models in terms of uncertainty of output has never been tackled, despite that the prohibitive computational cost of the (MSC) model, making the (SSC) approach still relevant to many applications.

In this paper, we intend to bridge that gap by proposing an uncertainty and sensitivity analysis of single and multiple-class frazil ice models, using a of full description of parameters uncertainty (by probability density functions) and variance decomposition of model outputs. First, we introduce the models, and main hypotheses and discuss their implementation in section 2. The uncertainty assessment methodology is presented in section 3. We then rely on numerous field and experimental measurements to quantify the uncertainties of input parameters of the two frazil numerical models in section 4. Using Monte Carlo experiments, we next study propagation of these uncertainties on model outputs in section 5: in other words, the evolution of the frazil ice volume fraction and the water temperature. The evolution of output scatter is discussed and compared to the asymptotes of the dynamic systems. Finally, sensitivity analysis based on Sobol indices (Sobol, 2001) and aggregated Sobol indices enables us to propose a selection of the most influential parameters in both models. We conclude about this study’s perspectives in section 6.

2 Frazil ice dynamics

This section introduces the continuum equations used to describe evolution of water temperature and frazil volume fraction, as well as their discrete counterparts and their numerical resolution under the assumption of well-mixed water bodies. The parameters of the models, that can be considered uncertain, are also introduced.

2.1 Mathematical description

Frazil crystals are assumed to be discs of radius r and thickness λ . The ratio between diameter and thickness, denoted $R = 2r/\lambda$ is supposed to be constant as crystals grow. Let us define n as the crystal number density, which corresponds to the number of crystals per unit volume per unit length in radial space. The total number of particles per unit volume is then defined as $N = \int_0^\infty n(r)dr$. We also introduce the frazil volume fraction density as $c = nV$ in which $V = \pi r^2 \lambda$ is the crystal volume. The volume proportion of the water-ice mixture occupied by frazil ice, i.e., the total volume fraction of frazil, is then defined as $C = \int_0^\infty c(r)dr$. An incompressible water-ice mixture of velocity \mathbf{u} and depth h is considered. Following Daly (1984) the number density balance equation can be defined as

$$\frac{\partial n}{\partial t} + \mathbf{u} \cdot \nabla n - \nabla \cdot (\nu_c \nabla n) = - \underbrace{\frac{\partial}{\partial r}(Gn)}_{(1)} + \underbrace{(\dot{N}_T + \dot{N}_I) \delta(r - r_c)}_{(2)} - \underbrace{\frac{1}{V} \frac{\partial}{\partial r}(FVn)}_{(3)} - \underbrace{w_r \frac{\partial n}{\partial z}}_{(4)}, \quad (1)$$

where ν_c is the frazil particles diffusivity, w_r is the buoyancy velocity, and remaining right hand side terms are source terms described hereafter:

- The source term (1) represents the crystal size evolution due to thermal growth, resulting from the heat exchange between the supercooled water and ice particles. The crystals can be considered to grow mainly on their peripheral area, defined by $a = 2\pi r\lambda$ (Daly, 1994). Frazil evolution is then driven by the radial growth rate G , equal to

$$\rho_i L_i G = \frac{N_u k_w}{\delta_T} \Delta T, \quad (2)$$

in which ρ_i is the frazil ice density, L_i is the latent heat of ice fusion and the right hand side is the heat exchange between the crystal of temperature T_i and the surrounding water of temperature T . The latter is modeled as a function of the temperature delta $\Delta T = T_i - T$, the thermal conductivity of water k_w , the Nusselt number N_u , which represents the ratio between turbulent heat transfer and conduction heat transfer, and a thermal boundary layer length scale, denoted δ_T (which is either chosen as the radius or thickness in literature). The crystal temperature T_i is assumed to be equal to the freezing temperature denoted T_f . The Nusselt number can be described as a function of ratio $m^* = r/\eta$, where η is the Kolmogorov length scale, which can be defined as a function of the turbulent dissipation rate ε as $\eta = (\nu^3/\varepsilon)^{1/4}$, in which ν is the molecular viscosity of water (Daly, 1984). The Nusselt number formulation described by Holland et al. (2007) is used. For small particles, i.e. $m^* \leq 1$, heat transfer is governed by diffusion and convection, and the Nusselt number can therefore be written as :

$$N_u = \begin{cases} 1 + 0.17m^*P_r^{1/2} & \text{if } m^* \leq P_r^{-1/2} \\ 1 + 0.55m^{*2/3}P_r^{1/3} & \text{if } P_r^{-1/2} < m^* \leq 1, \end{cases} \quad (3)$$

where P_r is the Prandtl number, defined as the ratio between molecular and thermal diffusivity. For larger particles ($m^* > 1$), heat transfer is governed by turbulent mixing of the boundary layer around the crystal, and the Nusselt number is defined by

$$N_u = \begin{cases} 1.1 + 0.77\alpha_T^{0.035}m^{*2/3}P_r^{1/3} & \text{if } \alpha_T m^{*4/3} \leq 1000 \\ 1.1 + 0.77\alpha_T^{0.25}m^*P_r^{1/3} & \text{otherwise,} \end{cases} \quad (4)$$

where α_T is the turbulent intensity.

- Heterogeneous nucleation (growth from foreign nuclei) and secondary nucleation (birth of new nuclei due to breakage of parent crystals) are invoked to explain the continuous feed of frazil nuclei during the rapid initial frazil growth (Daly, 1984, 1994). Heterogeneous nucleation is mainly caused by impurities that come either from the water body itself (biological elements, suspended sediments, etc.) or by penetration of new nuclei from the atmosphere (meteorological conditions or artificial seeding during experiments). The source term (2) of Equation (1) models the introduction of new nuclei, as well as the secondary nucleation, where \dot{N}_I is a seeding rate, and \dot{N}_T the secondary nucleation rate function of collisions between crystals. The function $\delta(r - r_c)$ is the Dirac delta function and r_c is the critical radius (radius of new nuclei). Note that seeding and secondary nucleation are assumed to introduce particles of the same size. Collisions are supposed to cause small nuclei to break off from parent crystals with a frequency of collision proportional to the crystal velocity relative to the fluid U_r and the total number of particles N that are contained in the volume swept by the crystal

(Svensson and Omstedt, 1994). The secondary nucleation rate \dot{N}_T is then defined as

$$\dot{N}_T = \tilde{n} \int_0^\infty \pi r^2 U_r(r) n(r) dr, \quad (5)$$

where $U_r = \sqrt{\frac{4\varepsilon r^2}{15\nu} + w_r^2}$ is the geometric mean between turbulent velocity and buoyant rise velocity. The number $\tilde{n} = \max(N, n_{max})$ is the average number of particles per unit volume that take part in the collisions, and n_{max} is a fitting parameter controlling the efficiency of the collision breeding. This parametrization was also followed by Smedsrud (2002), Smedsrud and Jenkins (2004), Wang and Doering (2005) and Holland and Feltham (2005).

- The source term (3) of Equation (1) is a flocculation source term supposed to represent the net effect of both flocculation and breakup, as introduced by Svensson and Omstedt (1994), who chose $F = F_0 r^2$, where F_0 is a constant. Their choice was motivated by the intuition that larger crystals are more prone to flocculate. However, as explained by Rees Jones and Wells (2018), this yields a linear flocculation in the number of crystals. However one could argue that flocculation should be quadratic in the number of crystals like secondary nucleation since it is physically based on the same collision mechanisms. Moreover, it also depends on hydrodynamic conditions, of which the effects on flocculation are still poorly characterized in literature. To be consistent with models in the literature, the formulation proposed by Svensson and Omstedt (1994) is chosen for this paper.
- The source term (4) of Equation (1) represents the buoyancy of frazil ice crystals, where w_r is the rise velocity. In the present paper, it is set to $w_r = 32.8(2r)^{1.2}$, following Svensson and Omstedt (1994), who simplified the model proposed by Daly (1984). For consistency with previous modeling studies, we retain this simple formulation, even if many other formulations have been proposed as summarized by Morse and Richard (2009) and McFarlane et al. (2014).

Finally, the thermal balance of the water-ice mixture complements Equation (1). Assuming $C \ll 0$, the water fraction temperature equation is obtained :

$$\frac{\partial T}{\partial t} + \mathbf{u} \cdot \nabla T - \nabla \cdot (\nu_t \nabla T) = \frac{\phi}{\rho c_p} + \frac{\rho_i L_i}{\rho c_p} \int_0^\infty G(r) a(r) n(r) dr, \quad (6)$$

where T is the temperature of the water fraction of the water-ice mixture, ν_t is the thermal diffusivity, ρ the density of water, c_p the specific heat of water and ϕ the net heat source resulting from heat exchanges with the atmosphere in W.m^{-3} .

In the following sections, a well-mixed water body is considered. Equations (1) and (6) are written in terms of space-averaged quantities. This assumption allows us to neglect the convection and diffusion operators and to focus on solving the average temperature and average frazil volume fraction. Therefore, the partial differential equations of frazil and temperature are simplified to a coupled set of Ordinary Differential Equations (ODEs) including only the source terms.

2.2 Radial space discretization

In this section we introduce a multiple-size-class (MSC) frazil ice model, which is a discrete version of Equations (1) and (6) in radial space. It consists of considering m classes of constant radius chosen between a minimum and a maximum radius

(Svensson and Omstedt, 1994). For each class i , the radius and the thickness are supposed to be equal to r_i and $\lambda_i = 2r_i/R$ ($1 \leq i \leq m$). The peripheral area as well as the surface and volume of frazil crystals are defined as $a_i = a(r_i) = 2\pi r_i \lambda_i$, $s_i = s(r_i) = 2\pi(r_i \lambda_i + r_i^2)$ and $V_i = V(r_i) = \pi r_i^2 \lambda_i$, respectively. A log-Uniform discretization of the radial space is chosen
150 as in previous studies (Svensson and Omstedt, 1994; Wang and Doering, 2005; Rees Jones and Wells, 2018). The number of crystals of class i is noted n_i , and similarly, the volume fraction of crystals of class i is noted c_i . The total volume fraction of frazil, and total number of particles, are then defined as $C = \sum_{i=1}^m c_i$ and $N = \sum_{i=1}^m n_i$ with $c_i = n_i V_i$, respectively. The discrete version of Equation (1) written in terms of volume fraction balance reads:

$$\frac{dc_i}{dt} = \underbrace{V_i (\Gamma_{i-1} c_{i-1} + (\Lambda_i - \Gamma_i) c_i - \Lambda_{i+1} c_{i+1})}_{(1)} + \underbrace{\tau_i}_{(2)} + \underbrace{\beta_{i-1} c_{i-1} - \beta_i c_i}_{(3)} - \underbrace{\gamma_i c_i}_{(4)} \quad (1 \leq i \leq m), \quad (7)$$

155 with the boundary conditions $V_0 = V_{m+1} = \Gamma_0 = \Gamma_m = \Lambda_{m+1} = \beta_0 = \beta_m = 0$. The discrete versions of the source terms introduced in Equation 1 are defined following previous work (Svensson and Omstedt, 1994; Wang and Doering, 2005; Holland and Feltham, 2005):

- The thermal growth source term (1) is composed of thermal growth and melt functions Γ_i and Λ_i , which are defined as

$$\Gamma_i = H \frac{G_i a_i}{V_i \Delta V_i}, \quad (8)$$

160 $\Lambda_i = (1 - H) \frac{G_i s_i}{V_i \Delta V_{i-1}}, \quad (9)$

where $G_i = G(r_i)$, $\Delta V_i = V_{i+1} - V_i$ and $H = H_e(T_f - T)$, with H_e the Heaviside function allowing to a switch between melting or freezing. We assume that frazil crystals grow from their peripheral area a_i but melt from their surface s_i (Holland and Feltham, 2005).

- The crystal birth source term (2), composed of the secondary nucleation and seeding, reads

165 $\tau_{i \neq 1} = -\zeta \alpha_i c_i, \quad (10)$

$$\tau_{i=1} = \tau_s V_1 / h + \sum_{j=2}^m \alpha_j c_j, \quad (11)$$

where $\zeta = V_1 / V_i$ is a coefficient to conserve crystal volume, $\alpha_i = \pi \tilde{n} U_r(r_i) r_i^2$, $\tilde{n} = \max(N, n_{max})$, with n_{max} a fitting parameter for collision breeding and τ_s is a constant seeding rate in $\text{m}^{-2} \cdot \text{s}^{-1}$.

- The flocculation source term (3) is defined as $\beta_i = a_f r_i / r_1$, where a_f is a flocculation coefficient in s^{-1} .

170 – The buoyancy of crystals is simplified into a gravitational removal sink term (4) defined as $\gamma_i = -w_r(r_i) a_d / h$, in which h is the water depth. We introduce a coefficient a_d , to account for the uncertainty of the rise velocity and gravitational removal process.

Finally, writing the thermal growth source term as $S_i = V_i (\Gamma_{i-1} c_{i-1} + (\Lambda_i - \Gamma_i) c_i - \Lambda_{i+1} c_{i+1})$, one can write the discrete version of Equation (6) as

175 $\frac{dT}{dt} = \frac{\phi}{\rho c_p} + \frac{\rho_i L_i}{\rho c_p} \sum_{i=1}^m S_i. \quad (12)$

A general discrete frazil ice model is implemented in the present work (Equations 7 and 12) so that one can easily retrieve the formulations developed in previous papers. For example, supposing $H = 1$, $a_d = 1$, $\tau_s = 0$ and writing $c_i = n_i V_i$, Equation (7) is equivalent to Equation (4) in Wang and Doering (2005) and Equation (1) in Svensson and Omstedt (1994) with $\zeta = 1$. By also neglecting flocculation ($a_f = 0$), it is equivalent to Equation (10) in Rees Jones and Wells (2018).

180 2.3 Single-size-class simplification

A simplified approach is to take a single-size-class (SSC) composed only of particles of radius \bar{r} representative of the whole crystal distribution. A simplified set of equations for the frazil volume fraction and water temperature can then be written as

$$\frac{dC}{dt} = GaN + \frac{1}{h} (V\tau_s - w_r a_d C), \quad (13)$$

$$\frac{dT}{dt} = \frac{\phi}{\rho c_p} + \frac{\rho_i L_i}{\rho c_p} GaN, \quad (14)$$

185 where $a = a(\bar{r})$, $V = V(\bar{r})$, $w_r = w_r(\bar{r})$ and $G = G(\bar{r})$, with $G(r)$ being defined by Equation (2) and the Nusselt number by Equations (3) and (4). In the previous equations, τ_s is the seeding rate, a_d is the buoyancy coefficient, h is the water depth and w_r is the buoyancy velocity. The (SSC) system can either be expressed in terms of total number of frazil crystals N or total frazil volume fraction C using the expression $N = C/V$.

2.4 Numerical methods to solve governing equations

190 A semi-implicit theta scheme is implemented for the MSC model, with a constant time step Δt and a fully implicit method is used for the SSC model. Details on the numerical resolution are provided in Appendix A. Note that the semi-implicit time scheme is subject to a stability condition function of the smallest radius. In the present study, we found that, for the range of radius tested in the sensitivity analysis, decreasing the time step below $\Delta t = 0.25$ s did not impact the results, so a value of 0.25 s was retained for all simulations.

195 An important aspect of the numerical frazil ice models presented in this paper is the need to provide a non-zero initial condition for the frazil volume fraction (in absence of seeding: i.e. $\tau_s = 0$). In the case of MSC models, Svensson and Omstedt (1994) assumed a uniform initial number of particles n_0 in each radius class i.e. $n_i(t_0) = n_0$ ($1 \leq i \leq m$). We followed the same principle to initialize the system, but fixed the number of initial particles at zero for classes with a radius exceeding a threshold r_0 : i.e. $n_{r_i \leq r_0}(t_0) = n_0$ like proposed in other works (Holland and Feltham, 2005; Rees Jones and Wells, 2018). To
200 be able to compare SSC and MSC models, we worked in terms of initial volume fraction of frazil $C(t_0) = C_0$. The system was then initialized with $n_0 = C_0 / \sum_{r_i \leq r_0} V_i$ and $c_i(t_0) = n_i(t_0) V_i$ ($1 \leq i \leq m$). As the initial setup has a significant influence on the results (Holland and Feltham, 2005), C_0 and r_0 are considered in the following sections as uncertain parameters.

To test convergence in terms of the number of radius classes, we make it vary from $m = 10$ to $m = 200$. The results, presented in Figure 1, are consistent with observations by Rees Jones and Wells (2018) that the system requires $m \gtrsim 100$ to converge.
205 Note that Svensson and Omstedt (1994) and Wang and Doering (2005) took $m = 20$ and $m = 40$, respectively. It should also be noted that convergence in the number of classes depends on the initialization method of the system, as highlighted by Holland




Figure 1. MSC model convergence in the number of classes with $C_0 = 4.5 \times 10^{-8}$, $r_0 = 0.2 \text{ mm}$, $r_{min} = 10^{-5} \text{ m}$, $r_{max} = 10^{-2} \text{ m}$, $dt = 0.1 \text{ s}$, $\phi = -1400 \text{ W.m}^{-3}$, $\varepsilon = 1.2 \times 10^{-3} \text{ m}^2.\text{s}^{-3}$, $\alpha_T = 0.0876$, $n_{max} = 8 \times 10^6 \text{ m}^{-1}$, $a_f = 10^{-4} \text{ s}^{-1}$, $\tau_s = 0 \text{ m}^{-2}.\text{s}^{-1}$ and $a_d = 0$. Number of crystals per class (a) at initial time (black) and at time $t = 300 \text{ s}$ (color). Water temperature and frazil volume fraction vs time (b).

and Feltham (2005). To perform all the numerical simulations required for a complete sensitivity analysis, $m = 100$ was chosen as a trade-off between numerical convergence and computational cost.

2.5 Study cases and model parameters

In the present study, we focus on the evolution of water temperature and total frazil volume fraction in a supercooled, well mixed water body of depth $h = 1 \text{ m}$. To focus on the frazil modeling rather than heat budget, uncertainties deriving from exchanges with the atmosphere are neglected and the cooling rate ϕ is considered deterministic and constant over time in all experiments. Nonetheless, different values of ϕ were tested, ranging from -50 to -1000 W.m^{-3} , to test the variability of the results in different cooling rate situations. As described in previous sections, frazil ice models are driven by many parameters, some of which are subject to a significant degree of uncertainty. The list of parameters considered in the present study for conducting the uncertainty analysis is shown in Table (1). Some physical properties are considered constant and are taken at $T = 0^\circ \text{C}$ such that $\nu = 1.792 \times 10^{-6} \text{ m}^2.\text{s}^{-1}$, $\rho = 999.82 \text{ kg.m}^{-3}$, $\rho_i = 916.8 \text{ kg.m}^{-3}$, $L_i = 3.35 \times 10^5 \text{ J.kg}^{-1}$, $c_p = 4.1855 \times 10^3 \text{ J.kg}^{-1}.\text{K}^{-1}$, and $k_w = 0.561 \text{ W.m}^{-1}.\text{K}^{-1}$.

Taking the SSC model as a reference, one can simply visualize the main feature of frazil dynamic systems. When time is close to zero, the initial temperature decrease rate is defined by $\lim_{t \rightarrow 0} \dot{T} = \phi / \rho c_p$. In the absence of seeding and gravitational removal (i.e. $\tau_s = 0$ and $a_d = 0$), the temperature decreases to the maximum supercooling point, characterized by a maximal temperature depression, denoted $\theta = \max(T_f - T)$, and a critical time, denoted t_c . After the maximum supercooling, the frazil production rate releases more heat compared to that released in the atmosphere, leading to an increase in water temperature which tends toward freezing point. Finally, when time tends to infinity, there is a balance between the frazil growth source term and the cooling rate. This leads to a linear increase in frazil volume fraction at a constant rate i.e. $\lim_{t \rightarrow +\infty} \dot{C} = -\phi / \rho_i L_i$. This typical evolution of water temperature and frazil volume fraction is described in Figure (2) in which frazil volume fraction

Table 1. Description of uncertain parameters of the frazil ice models.

Parameter	Unit	Description	Category	Model
C_0	-	Initial frazil volume fraction	Initial condition	Both
r_0	m	Initial maximum radius	Initial condition	MSC
r_{min}	m	Minimum radius	Discretization	MSC
r_{max}	m	Maximum radius	Discretization	MSC
\bar{r}	m	Mean radius	Discretization	SSC
R	-	Diameter to thickness ratio	Source term ①	Both
δ_T	m	Thermal growth length scale	Source term ①	Both
ε	$\text{m}^2.\text{s}^{-3}$	Turbulent dissipation rate	Source terms ①, ②	Both
α_T	-	Turbulent intensity	Source term ①	Both
n_{max}	m^{-3}	Secondary nucleation efficiency cap	Source term ②	MSC
τ_s	$\text{m}^{-2}.\text{s}^{-1}$	Seeding rate	Source term ②	Both
a_f	s^{-1}	Flocculation coefficient	Source term ③	MSC
a_d	-	Buoyancy coefficient	Source term ④	Both




Figure 2. Typical evolution of water temperature and frazil volume fraction vs time (a) and reproduction of Carstens (1966) case I experiments (b) with $dt = 0.1\text{s}$, $\phi = -1400 \text{ W.m}^{-3}$, $\varepsilon = 1.2 \times 10^{-3} \text{ m}^2.\text{s}^{-3}$, $\alpha_T = 0.0876$, $n_{max} = 8 \times 10^6 \text{ m}^{-1}$, $a_f = 10^{-4} \text{ s}^{-1}$, $\tau_s = 0 \text{ m}^{-2}.\text{s}^{-1}$, $a_d = 0$, $\bar{r} = 3.8 \times 10^{-4} \text{ m}$, $R = 10$, $C_0 = 5.1 \times 10^{-6}$ (SSC) and $C_0 = 4.5 \times 10^{-8}$ (MSC), $r_0 = 0.2\text{mm}$, $r_{min} = 10^{-5} \text{ m}$, $r_{max} = 10^{-3} \text{ m}$, $m = 100$, and $\delta_T = \lambda$.

asymptote, denoted C^∞ , is defined as

$$C^\infty = C_0 - \phi t / \rho_i L_i. \quad (15)$$

By introducing seeding and gravitational removal, i.e. $\tau_s \neq 0$ and $a_d \neq 0$, the frazil ice long term asymptotic is modified, and we have $\lim_{t \rightarrow +\infty} \dot{C} = -\phi / \rho_i L_i + (V\tau_s - w_r a_d C) / h$. Considering ϕ and τ_s as constant over time, this yields a convergence

of the frazil volume fraction towards a finite limit, defined by the ratio between the buoyancy removal and the production rate due to thermal growth and seeding, which reads

$$C^\infty = \frac{h}{a_d w_r} \left(-\frac{\phi}{\rho_i L_i} + \frac{V \tau_s}{h} \right). \quad (16)$$

It should be observed that the steady states are not affected by the crystal growth rate. Similar observations were discussed by
 235 Rees Jones and Wells (2018) for the MSC model.

In the following the recovery time is defined as the time when temperature has recovered 90% of its supercooling depression. We refer to the recovery phase to describe the long-term evolution of the system i.e. all times past recovery time. Similarly, we will refer to the transient phase as times between initial time and recovery time.

In the present study, we analysed the uncertainty of input parameters in the main transient and steady-state features of the
 240 frazil ice models with and without gravitational removal. We started the simulation with a water temperature equal to the freezing temperature (which is supposed to be equal to zero in the absence of salinity : i.e. $T_f = 0^\circ C$). We checked that all the simulations ran beyond the critical time. For the range of parameters considered there, it was found that choosing a final time of $t_f \simeq 1\text{h}$ was enough to capture the transient features of the ODE systems.

Note that both the SSC and MSC models are able to reproduce the experiments of Michel (1963), Carstens (1966) and
 245 (Clark and Doering, 2004) as shown by Svensson and Omstedt (1994) and Wang and Doering (2005). In Figure (2) we give an example of the results of both SSC and MSC models compared to Carstens (1966) experimental results (Case I).

3 Probabilistic framework for uncertainty analysis

The objective of this section is to describe the mathematical tools necessary to study the uncertainties of SSC and MSC frazil ice models. An uncertainty study of a numerical model can be performed in three steps (Sudret, 2007). The first step consists
 250 in identifying the uncertain parameters, and characterizing the probabilities of occurrence for their values through Probability Density Functions (PDFs), which is referred to as *quantification of uncertainty sources*. The second step concerns the *propagation of uncertainties* through the interest models, generally using sampling techniques (e.g., Monte Carlo) to obtain possible values of the target outputs (here frazil ice concentration, water temperature) and compute their statistical estimates (mean, standard deviation, percentiles, etc.). The third step, called *sensitivity analysis*, focuses on ranking the uncertain parameters
 255 in terms of influence on the target output. In the following, formulations for the statistical estimates and sensitivity analysis indices are given, with a summary of elements essential to understand this paper. Theoretical details can be found in Soize (2017) and Sudret (2007). All uncertainty analysis computations were performed with the OpenTURNS library (version 1.18) developed by a partnership between Airbus Group, EDF R&D, IMACS, ONERA and Phimeca Engineering (Baudin et al., 2016b).

Let $\mathbf{X} = (X^1, \dots, X^{n_X})$ be the vector of uncertain parameters of the frazil ice model, where n_X is the number of uncertain parameters. Let g be the interest frazil model, either SSC or MSC models presented in Equations (14) and (12) for temperature and (13) and (7) for frazil volume fraction. The output of the interest models are the frazil volume fraction and water temperature discrete time series i.e. $T(t_k)$ and $C(t_k)$ ($1 \leq k \leq n_t$), which we will generally denote $\mathbf{Y} = (Y^1, \dots, Y^{n_t})$ below for the sake of simplicity. The random vectors \mathbf{X} and \mathbf{Y} are linked through frazil models g such that $\mathbf{Y} = g(\mathbf{X}, \mathbf{d})$, where \mathbf{d} is a deterministic vector, i.e. fixed parameters in contrast with the uncertain set of inputs \mathbf{X} .

To undertake the uncertainty studies, both the inputs \mathbf{X} and the interest outputs \mathbf{Y} are considered to be random vectors. We assume that \mathbf{X} has a density $p_{\mathbf{X}}$, so that $\mathbb{P}(\mathbf{X} \in E \subseteq D_{\mathbf{X}}) = \int_E p_{\mathbf{X}}(\mathbf{x}) d\mathbf{x}$, where E is a subset from the space of all possible values $D_{\mathbf{X}}$. Each element X^i and Y^k of the input and output vectors is hence characterized by a PDF.

The results of the uncertainty analysis are directly linked to the UQ (Uncertainty Quantification) study specification and consequently to the description of the uncertain input parameters \mathbf{X} . Thus, special attention is paid to propose a relevant quantification of uncertainty sources. This particular point is addressed in section (4), in which frazil literature is explored to provide adequate bounds and PDFs for each parameter identified in Table (1).

3.2 Uncertainty propagation of random variables

In the following, we suppose that PDFs of inputs \mathbf{X} are known, in contrast to those of \mathbf{Y} . The objective is therefore to characterize the latter by estimating statistical indicators as the PDF's mean and standard deviation. To achieve this, a Monte Carlo sampling method can be used.

In the Monte-Carlo method, we generate a sample of independent observations of the random vector \mathbf{X} using the joint PDF of the input random vector. For each observation of the input \mathbf{X} , we evaluate the corresponding output \mathbf{Y} . The resulting experimental design of the input random vector \mathbf{X} is an array of size $N \times n_X$ where each row, denoted $\mathbf{x}_i = \{x^1, \dots, x^{n_X}\}$, represents a possible configuration of the frazil model. The output realizations i.e. $\mathbf{y}_i = \{y_i^1, \dots, y_i^{n_t}\}$, are generated by N deterministic simulations with corresponding inputs. For a sample of size N , the output of the model is a $N \times n_t$ array: $\mathbf{Y} = [y_i^k]_{i,k} = [g^k(\mathbf{x}_i, \mathbf{d})]_{i,k} \in \mathbb{R}^{N \times n_t}$. Each row of the matrix represents one output time series for a given set of input parameters, while each column represents the output realizations at a given time t_k .

Statistical estimators of the response can then be computed. The Monte-Carlo estimation of the mean, denoted $\hat{\mu}_{Y^k}$ and standard deviation, denoted $\hat{\sigma}_{Y^k}$ at time t_k reads:

$$\hat{\mu}_{Y^k} = \frac{1}{N} \sum_{j=1}^N y_j^k \quad \text{and} \quad \hat{\sigma}_{Y^k} = \sqrt{\frac{1}{N} \sum_{j=1}^N (y_j^k - \hat{m}_{Y^k})^2}. \quad (17)$$

These estimations converge to the true values following the law of large numbers (conditioned by the existence of corresponding PDF's first and second moment, i.e. expectation and variance (Soize, 2017)). The convergence order of the Monte-Carlo method is given by the central limit theorem leading to decrease of the confidence intervals' sizes proportional to $1/\sqrt{N}$. In the results of this paper, confidence intervals of the estimated mean and standard deviation are computed using a bootstrap method.

3.3 Sensitivity analysis using Sobol indices

Sensitivity analysis is essential in understanding numerical models (Razavi et al., 2021). It aims at quantifying the impact of inputs variables imprecision on the accuracy of the model output variables. Conventional approaches to Global Sensitivity Analysis (GSA) imply the stochastic estimation of statistical moments and indices classically achieved with the Monte-Carlo technique. In the present work, the relative influence of uncertain parameters on the output time series is assessed by means of Sobol indices resulting from the ANOVA (ANalysis Of VAriance) variance decomposition (Sobol, 2001). Sobol indices are computed via a modified version of the method proposed by Saltelli (2002) which is described in Appendix B. For each time step t_k ($1 \leq k \leq n_t$) and $i \in \{1, \dots, n_X\}$, first order and total Sobol indices, denoted S_i^k and ST_i^k respectively, are computed. By definition, at any time t_k , the total Sobol sensitivity index (ST_i^k) measures the contribution to the output variance of the uncertain variable X^i , including all variance caused by its interaction, of any order, with any other input variables. Thus, the variance part explained by variable interactions of the input factor X^i with the other uncertain parameters is determined by subtracting the total Sobol sensitivity index (ST_i^k) and the first order Sobol index (S_i^k) characterizing the influence of the variable X^i alone. Finally, the discrete time first order and total Sobol indices are aggregated, denoted AS_i and AST_i respectively, as proposed by Gamboa et al. (2014).

4 Uncertainty source quantification

Many laboratory experiments have been carried out to better understand frazil ice dynamics as summed up by Barrette (2020, 2021). These experiments, as well as field measurements, help us define parameter variability. In this section, we consider the uncertain parameters listed in Table (1), and deduce adequate variability and PDFs for uncertainty propagation and sensitivity analysis using the aforementioned data. The main geometrical properties of crystals and radial space discretization are first discussed, as well as initial concentration. We then review all the parameters involved in frazil source terms in the same chronology as presented in section (2).

4.1 Crystals' geometrical properties

Description of the frazil crystals' shape has been the subject of several field and laboratory measurements (Arakawa, 1954; Daly, 1984, 1994). The crystals have been described as thin discs that grow mainly from their peripheral area. More recently, photos have brought valuable confirmation of the disc shape of frazil crystals, but also highlighted the complexity in larger flocs' geometry (Clark and Doering, 2006; Ghobrial et al., 2012; McFarlane et al., 2014, 2015; Schneck et al., 2019). Let us recall that, in both SSC and MSC models, discretization is done in radial space. Therefore, either the ratio R or the thickness λ must be considered constant to fully describe disc-shaped particles. In the present study, frazil crystals are assumed to have the same aspect ratios, which is consistent with previous studies hypothesis (Svensson and Omstedt, 1994; Holland et al., 2007). Note that Rees Jones and Wells (2018) considered constant thickness instead, but highlighted a weak dependency of the

thermal growth to aspect ratio. Let us discuss the uncertainties associated with the choice of the radius discretization as well as the diameter to thickness ratio.




Figure 3. Particle size ranges (thick grey line), mean range (thin dark line) and log-normal distributions' mean radius (red vertical ticks) reported in field or laboratory experiments (Schaefer, 1950; Arakawa, 1954; Carstens, 1966; Gosink and Osterkamp, 1983; Osterkamp and Gosink, 1983b; Ettema et al., 1984; Wuebben, 1984; Kempema et al., 1986; Daly and Colbeck, 1986; Doering and Morris, 2003; Ye et al., 2004; Ye and Doering, 2004; Clark and Doering, 2004, 2006, 2009; Marko and Jasek, 2010; Ghobrial et al., 2012; McFarlane et al., 2012; Ghobrial et al., 2013; McFarlane et al., 2015; Kempema and Ettema, 2016; McFarlane et al., 2016, 2017; Schneck et al., 2019)

Either a mean radius or a radius space discretization must be specified for SSC and MSC models, respectively. In both cases, observed radius distribution and particle size, reported in a many studies, are taken into consideration. For the sake of synthesis, we analyzed the corresponding publications from 1950 to 2019, and report the available observations in Figure 3. This figure is complementary to Table 7 of McFarlane et al. (2017), who summarize particle sizes from both laboratory and field measurements. Daly and Colbeck (1986) and Clark and Doering (2006) reported log-normal distributions of particles in laboratory experiments, with a mean radius ranging from 0.12 to 0.25 mm and 0.49 to 1.4 mm, respectively. This was later confirmed by Ghobrial et al. (2012) and McFarlane et al. (2015) in the University of Alberta Cold Room Facility, with mean size ranging from 0.66 to 0.94 mm. Schneck et al. (2019) recently published similar results under different salinities, with mean radius ranging from 0.45 to 0.52 mm. Log-normal distributions were also observed in field studies, with the mean ranging from 0.59 to 0.94 mm in a report by McFarlane et al. (2017). Gathering together all reported mean radius ranges, an uncertainty interval of 1.2×10^{-4} to 2.1×10^{-3} m was chosen for the mean radius in the SSC model. When their full variation range is considered, frazil crystal sizes are known to follow a log-normal distribution as previously argued, but there is no evidence or reason for the mean radius being log-normal as well. The number of field and laboratory observations is still too small to fit

an empirical PDF on the mean radius with reasonable confidence. Hence, a log-uniform PDF is chosen with the previously described bounds. Using Log-uniform approximate distributions allows us to explore by means of evenly distributed values, parameters that vary over several orders of magnitude.

340 For the MSC model, both a minimum and a maximum radius are to be determined for the bounds of the radial discretization. The minimum radius can be determined from Lal et al. (1969) survival theory and is about $4\text{ }\mu\text{m}$ (Mercier, 1984). The maximum size of frazil particles is about 1 mm to 5 mm according to Clark and Doering (2004), but frazil flocs can be significantly larger as shown by Schneck et al. (2019), and can range up to several cm (McFarlane et al., 2017). Svensson and Omstedt (1994) worked with $[4 \times 10^{-6}, 3 \times 10^{-3}]$ m while Wang and Doering (2005) and Rees Jones and Wells (2018) chose $[7 \times 10^{-5}, 5 \times 10^{-3}]$ m and $[5 \times 10^{-5}, 5 \times 10^{-2}]$ m, respectively. In literature, r_{min} is taken to be between 10^{-6} and 10^{-4} , so we chose an intermediate order of magnitude of $r_{min} = 10^{-5}$ m. The same choice was made for the maximum radius, leading to a value of $r_{max} = 10^{-3}$ m. Both parameters were then considered as uncertain parameters with log-uniform PDFs and were taken to be within $[10^{-6}, 10^{-4}]$ for r_{min} and within $[10^{-3}, 10^{-1}]$ for r_{max} .

350 The aspect ratio ranged from 6 to 13 in the study by Daly and Colbeck (1986), while there were greater values, from 12.2 to 16.33, in Clark and Doering (2004, 2006). Arakawa (1954) reported an even wider aspect ratio range, from 5 to 100. More recently, McFarlane et al. (2014) reported aspect ratios ranging from 11 to 71 with a mean of 37 and a standard deviation of 11. Considering all these observations, an uncertainty range of 5 to 100 was chosen for the diameter-to-thickness ratio. As not enough data to properly fit a PDF on aspect ratios was provided, a uniform PDF is chosen for the sensibility analysis (Maximum Entropy Principle Soize (2017)).

355 Weak dependency of the thickness to the radius was reported by McFarlane et al. (2014), who suggest assuming a increasing aspect ratio as discs grow instead of a constant aspect ratio. Note also that, as frazil ice forms larger flocs, the disc shape hypothesis commonly accepted in models does not hold anymore. This could lead to erroneous estimations of the thermal growth rate of larger flocs. However, neither the variability of the aspect ratio in the crystals' distribution, nor that of the shape, are considered in the present study and we make the assumption that r and R are independent.

360 4.2 Initial concentration

As previously mentioned, either a non-zero initial volume fraction or a non-zero seeding rate is required to trigger thermal growth in the model. In experimental facilities, frazil ice nuclei are sometimes artificially introduced in water to initiate frazil ice growth (Muller, 1978; Tsang and Hanley, 1985) but the initial concentration is rarely addressed and depends on the method of initial seeding. As such, a large uncertainty domain is considered in this study for the initial volume fraction of frazil, ranging from 10^{-8} to 10^{-4} , to account for the lack of data on this parameter. Additionally, a log-uniform PDF is retained since the parameter range variation vary over several orders of magnitude.

For the MSC model, it is necessary to choose how the initial volume fraction is distributed over size classes. One possibility is to initialize the system with log-uniform distributions, similar to the one observed in laboratory experiments. However, observations of nuclei close to r_c in size are limited. Consequently, authors have preferred simpler initialization methods, whereby a constant number of crystals is distributed equally over all classes (Svensson and Omstedt, 1994; Wang and Doering,

370

2005; Rees Jones and Wells, 2018). Holland et al. (2007) argued that distributing the initial volume fraction over a range of radii, thus changing the initial number of particles per class, significantly impacts results. They found that filling only the first size class, as in Hammar and Shen (1995) has less impact on results and should therefore be the preferred initialization method. Given poor evidence of initial predominance of each class in nature, we decided to test both initialization methods i.e. distributing the initial volume fraction over a range of small radii as presented in section (2.4) then feeding only the first class as suggested by Holland et al. (2007). For the first method, ice nuclei are supposed to be initially spread between the minimum radius r_{min} and a threshold radius r_0 , which we suppose can only be smaller than the mean radius (cf. Figure 3). Therefore, the threshold r_0 is considered uncertain within the bounds $[1.2 \times 10^{-4}, 2.1 \times 10^{-3}]$. For the second method, we set $r_0 = r_{min}$, so that only the first receives the initial concentration.

380 4.3 Thermal growth and turbulent parameters

As shown in Equation (2), thermal growth (1) is mainly affected by two uncertain parameters, which are the Nusselt number and the thermal boundary layer thickness. In this section, we discuss the uncertainty of both parameters. Since the Nusselt number is being modeled via turbulent parameters, namely the turbulent dissipation rate and the turbulent intensity, their uncertainty is also addressed.

385 As discussed by Rees Jones and Wells (2018), the thermal boundary layer thickness δ_T is not constant around a crystal and in recent studies there has been an inconsistent scaling of thermal growth. Svensson and Omstedt (1994) chose the length scale as $\delta_T = \lambda$ while others have chose $\delta_T = r$ (Smedsrud and Jenkins, 2004; Holland et al., 2007), leading to a serious underestimation of the growth rate, since $\lambda < r$ for frazil discs. Rees Jones and Wells (2015) have shown that there is a logarithmic dependency of the thermal growth on the aspect ratio that favors the $\delta_T = \lambda$ scaling. To account for the variability of the thermal boundary layer thickness, δ_T should be taken as an uncertain parameter. However, it should be mentioned that, with the variation range being $\delta_T \in [\lambda, r]$, the ANOVA methodology (section 3.3) cannot be applied, since the set of uncertain inputs, which contains δ_T and λ/r , cannot be considered independent anymore. To overcome this difficulty, the sensitivity analysis is conducted with the most appropriate scaling: i.e., $\delta_T = \lambda$, for both the SSC and MSC models. Nevertheless, we propose to investigate the impact of scaling by considering $\delta_T \in [\min(\lambda), \max(r)]$ to avoid modeling dependency. A log-uniform PDF is used and $\min(\lambda)$ and $\max(r)$ are estimated from their full variation ranges. A third experiment was also carried out with the $\delta_T = r$ scaling. The three results are then compared for the SSC model.

Two main turbulence parameters are considered: turbulent dissipation rate ε and turbulent intensity α_T , both impacting the Nusselt number (cf. Equations 3 and 4). For rivers, the dissipation rate can be estimated using friction velocity $u_* = \sqrt{gR_hS}$ such that

$$400 \quad \varepsilon = \frac{u_*^3}{\kappa R_h} \left[\ln \left(\frac{u_* R_h}{\nu} \right) - 1 \right] \quad (18)$$

where g is gravity, R_h the hydraulic radius, S the slope of the river, κ the von Karman constant (generally taken as equal to 0.4), and ν the viscosity of water. Daly (1994) summarized the dissipation rate for several experiments (Michel, 1963; Carstens, 1966; Tsang and Hanley, 1985; Muller, 1978) using Equation (18) and found values ranging from 7×10^{-5} to $0.4667 \text{ m}^2.\text{s}^{-3}$.

A similar method was described by McFarlane et al. (2015) to estimate the dissipation rate in rivers leading to values ranging from 4.2×10^{-4} to $1.4968 \text{ m}^2.\text{s}^{-3}$. Schneck et al. (2019) summarized dissipation rates observed in oceans, ranging from 10^{-9} to $10^{-2} \text{ m}^2.\text{s}^{-3}$ and from 10^{-10} to $10^{-3} \text{ m}^2.\text{s}^{-3}$ in polar regions. In the present study, we consider dissipation rates varying between 10^{-9} and $1.5 \text{ m}^2.\text{s}^{-3}$, which cover most flows encountered in rivers and oceans. Similarly, we consider a wide range of flows from low turbulence to high turbulence intensity which include most of the work presented in Figure (3), leading to a turbulent intensity ranging from 1% to 20%. Log-uniform PDFs are considered for both the dissipation rate and the turbulent intensity. Note that turbulence influences not only the rate of growth of frazil crystals but also the rate of secondary nucleation, thus impacting the frazil size distribution.

4.4 Other source terms

Following the same order as in Equation (1), let us discuss uncertain parameters involved in frazil source terms other than thermal growth: secondary nucleation (2), flocculation (3) and gravitational removal (4). It should be mentioned that almost no direct observations of these processes have been reported in the literature. Therefore, the definition of the uncertainty intervals is mainly based on expert knowledge and past numerical experiments in which parameters were determined by comparison to observed radius distributions.

- The uncertainty interval of the seeding rate is set after Daly (1994) to $[3 \times 10^{-1}, 10^{-4}] \text{ m}^{-2}.\text{s}^{-1}$, and to simplify the uncertainty analysis, is considered constant over time. For the secondary nucleation, Svensson and Omstedt (1994) proposed setting a common value for n_{max} that would allow focus on calibration of the initial seeding. They found that a value of $n_{max} = 4 \times 10^6$, along with initial seeding of the order of magnitude of $n_0 = 10^4$, gave satisfactory results compared to the experimental results of Michel (1963) and Carstens (1966). Wang and Doering (2005) found that fitting a single n_{max} value for all experiments was unsatisfactory. Instead, they proposed fitting both the initial seeding and n_{max} , leading to values ranging from 2×10^4 to 2×10^5 for Clark and Doering (2004) experiments, and from 2×10^4 to 2×10^5 for Carstens (1966) experiments. Smedsrud (2002) proposed a different value of $n_{max} = 10^3$, which was also used by Smedsrud and Jenkins (2004) and Holland and Feltham (2005).
- Similarly, the flocculation parameter a_f was calibrated to a value of 10^{-4} s^{-1} by Svensson and Omstedt (1994), who compared the results of their simulation to the expected size distribution spectra (other parameters in their model, like the number of class, the heat sink, the turbulent dissipation rate and initial condition were set to $m = 20$, $\phi = -1000 \text{ W.m}^{-3}$, $\varepsilon = 10^{-3} \text{ m}^2.\text{s}^{-3}$ and $n_i(t_0) = 10^4$ ($1 \leq i \leq m$)). From all previous calibration studies, one could choose n_{max} from 10^3 to 10^7 and $a_f = 10^{-4}$. But given the fact that secondary nucleation and flocculation are still very poorly understood and modeled, we choose large uncertainty intervals for these parameters. We added an order of magnitude for the bounds leading to n_{max} ranging from 10^2 to 10^8 and we suppose that flocculation can, depending on the conditions, be negligible, so that a_f was varied from 10^{-8} to 10^{-3} . The parameters n_{max} and a_f are considered independent from hydrodynamic parameters, and Log-uniform PDFs were chosen for both.

– The gravitational removal is affected by both buoyant rise velocity of frazil particles, and deposition process once particles reach the surface of the water column. Several attempts to measure the frazil rise velocity were done (Osterkamp and Gosink, 1983a; Wueben, 1984; McFarlane et al., 2014) and many formulations were proposed as summarized by McFarlane et al. (2014). Significant scatter can be observed in the data as shown on Figure (10) of McFarlane et al. (2014), with velocities ranging from approximately 0.7 to 16 mm/s for a radius of 1 mm. Models exhibit significant differences as well (cf. figure 4). To take into account uncertainties inherent to the choice of the rise velocity model, a buoyancy parameter a_d is introduced. A rise velocity envelope, combining the results of all models, can be defined by upper and lower bounds denoted $w_{max}(r, R)$ and $w_{min}(r, R)$ (cf. figure 4). The interval of a_d is defined from the mean of the gap between the simplified formulation used the present study for w_r (Svensson and Omstedt, 1994) and upper and lower bounds of the rise velocity envelope. To avoid the modeling of dependencies, a constant value is taken for a_d , even if the envelope depends on the radius and on the diameter to thickness ratio. Finally, we propose $a_d \in [a^-, a^+]$ in which $a^+ = \text{mean}(w_{max}(r, R)/w_r(r))$ and $a^- = \text{mean}(w_{min}(r, R)/w_r(r))$. By considering $r \in [10^{-5}, 10^{-2}]$ m and $R \in [5, 100]$, the following interval is obtained: $a_d \in [0.086, 1.51]$. Finally, it should be noted that this uncertainty quantification is rather imprecise since the rise velocity dispersion depends on the shape of the particles and flow conditions. Therefore, our analysis could be refined by taking into account dependencies. Also note that low values of a_d could also be justified from the uncertainty of the deposition process, which is not modelled in the present study.




Figure 4. Comparison of frazil rise velocity models with $R = 10$ (a) and rise velocity envelope chosen for the uncertainty analysis (b) (Zacharov et al., 1972; Gosink and Osterkamp, 1983; Ashton, 1983; Wueben, 1984; Daly, 1984; Svensson and Omstedt, 1994; Matoušek, 1992; Shen and Wang, 1995; Morse and Richard, 2009)

4.5 Summary of uncertain parameters

To conclude the uncertainty quantification, all the uncertain parameters, their bounds, and PDFs are summarized in Table (2).

Table 2. Uncertain parameters of the frazil ice models and their PDFs

Parameter	Unit	Description	Uncertainty interval	PDF
C_0	-	Initial frazil volume fraction	$[10^{-8}, 10^{-4}]$	Log-Uniform
r_0	m	Initial maximum radius	$[1.2 \times 10^{-4}, 2.1 \times 10^{-3}]$	Log-Uniform
r_{min}	m	Minimum radius	$[10^{-6}, 10^{-4}]$	Log-Uniform
r_{max}	m	Maximum radius	$[10^{-3}, 10^{-1}]$	Log-Uniform
\bar{r}	m	Mean radius	$[1.2 \times 10^{-4}, 2.1 \times 10^{-3}]$	Log-Uniform
R	-	Diameter to thickness ratio	$[5, 100]$	Uniform
δ_T	m	Thermal growth length scale	$[7.34 \times 10^{-6}, 2.1 \times 10^{-3}]$	Log-Uniform
ε	$\text{m}^2.\text{s}^{-3}$	Turbulent dissipation rate	$[10^{-9}, 1.5]$	Log-Uniform
α_T	-	Turbulent intensity	$[0.01, 0.2]$	Log-Uniform
n_{max}	m^{-3}	Secondary nucleation efficiency cap	$[10^2, 10^8]$	Log-Uniform
τ_s	$\text{m}^{-2}.\text{s}^{-1}$	Seeding rate	$[3 \times 10^{-1}, 10^4]$	Log-Uniform
a_f	s^{-1}	Flocculation coefficient	$[10^{-8}, 10^{-3}]$	Log-Uniform
a_d	-	Buoyancy coefficient	$[0.086, 1.51]$	Uniform

5 Uncertainty propagation and sensitivity analysis

455 In this section, we present the different Monte Carlo simulation cases considered for the uncertainty analysis, as synthesized in Table (3). Then we discuss the results of the uncertainty propagation and the sensitivity analysis obtained for both SSC and MSC models.

5.1 Propagation cases

Monte Carlo simulations are first carried out without seeding and gravitational removal ($\tau_s = 0$ and $a_d = 0$) for both (SSC) and (MSC) models, which correspond to cases (1) and (2), respectively. Seeding and gravitational removal processes are subsequently considered in cases (3) and (4). For each case, the set of uncertain parameters is described in Table (2). When not in X , parameters are considered deterministic with the following default values: $a_d = 0$, $\tau_s = 0 \text{ m}^2.\text{s}^{-1}$, $\delta_T = e$, $r_{min} = 10^{-5} \text{ m}$ and $r_{max} = 10^{-3} \text{ m}$.

465 Additional Monte Carlo simulations were carried out to examine the influence of specific parameters. For example, although $\delta_T = e$ is the preferred scaling and is chosen by default in all experiments, we also tested $\delta_T = r$ and $\delta_T \in [e, r]$ in experiments (1b) and (1c) to investigate the impact on results. In experiments (2b) and (2c), modifications of the MSC model uncertain parameters are also taken into account, in order to investigate the impact of the radius bounds r_{min} and r_{max} , as well as the alternate methods of initialization discussed in (4.2).

Statistical estimators are evaluated every 10s of physical time, leading to $n_t = 400$. To cope with the computational cost of multi-class experiments we used clusters to run all simulations in parallel. The computation time of 4.5×10^5 multi-class

Table 3. List of Monte Carlo simulations and their associated input random vectors X , sampling size, and number of function calls. Seeding rate and buoyancy velocity are set to $\tau_s = 0$, $a_d = 0$ in cases 1 and 2 .

Case	Model	Uncertain parameters	Case specificity	Sample size	No. of calls
1	SSC	$X = (C_0, \bar{r}, R, \varepsilon, \alpha_T)$	-	5×10^4	3.5×10^5
1b	SSC	$X = (C_0, \bar{r}, R, \varepsilon, \alpha_T)$	$\delta_T = r$	5×10^4	3.5×10^5
1c	SSC	$X = (C_0, \bar{r}, R, \delta_T, \varepsilon, \alpha_T)$	$\delta_T \in X$	5×10^4	4.8×10^5
2	MSC	$X = (C_0, r_0, R, \varepsilon, \alpha_T, n_{max}, a_f)$	-	5×10^4	4.5×10^5
2b	MSC	$X = (C_0, r_0, r_{min}, r_{max}, R, \varepsilon, \alpha_T, n_{max}, a_f)$	$r_{min}, r_{max} \in X$	6×10^4	6.6×10^5
2c	MSC	$X = (C_0, R, \varepsilon, \alpha_T, n_{max}, a_f)$	$r_0 = r_{min}$	5×10^4	4×10^5
3	SSC	$X = (C_0, \bar{r}, R, \varepsilon, \alpha_T, \tau_s, a_d)$	$\tau_s, a_d \in X$	7×10^4	6.3×10^5
4	MSC	$X = (C_0, r_0, R, \varepsilon, \alpha_T, n_{max}, \tau_s, a_f, a_d)$	$\tau_s, a_d \in X$	6×10^4	6.6×10^5

simulations with $m = 100$ is ~ 24 hours with 960 processes. A total of 4 million simulations were carried out for uncertainty propagation.

5.2 Results without gravitational removal

By neglecting the seeding rate and gravitational removal source terms, the steady state corresponds to the constant frazil
475 production rate that only depends on the heat sink ϕ , as shown in section (2.5). As expected, the statistical estimator time series for temperature and frazil volume fraction, presented in Figure (5) for cases (1) and (2), show a very narrow scatter of the output PDF at the start of simulation and at steady state. The results converge towards the two asymptotes (T^0 and C^∞ respectively) of the mono-class ODE system: i.e., the constant cooling rate when $t \rightarrow 0$ (initial supercooling phase) and the constant frazil production rate when $t \rightarrow \infty$ (recovery phase). However, for both SSC and MSC models, the transition between
480 the two asymptotes of the ODE system is spread out, and there is a significant difference in maximum supercooling between the 5^{th} , 25^{th} , 75^{th} and 95^{th} percentiles. For the median, a maximum supercooling of $T(t_c) \simeq -0.018^\circ C$ is reached at $t_c = 180s$ for the SSC model. The gap between the 5^{th} and 95^{th} percentiles maximum supercooling is $\Delta t_c = 890$ s and $\Delta \theta = 0.097^\circ C$. Note that the envelope obtained with the standard deviation gives a poor description of the output since its PDF is not normal, as it can be seen from the $\hat{\mu} \pm \hat{\sigma}$ lines in the figure 5.

485 Similar results were obtained with the MSC model, which recovers the same asymptote at steady state, however, there is a slight residual scatter at recovery. For case (2), we observed slightly less scatter at the maximum supercooling than with the SSC model, and the gap between the 5^{th} and 95^{th} percentiles maximum supercooling was $\Delta t_c = 820$ s and $\Delta \theta = 0.08^\circ C$.

Time series of the first-order Sobol indices are presented in Figure (6) (see Appendix E and F for details). In Appendix C, we also present the first- and total-order Sobol indices at times t_c , $2 \times t_c$ and t_f with 95% confidence intervals, as well as
490 the aggregated Sobol indices, computed via Equation (B7). The time evolution of Sobol indices of temperature and frazil were similar for both the SSC and MSC with the exception of the initial concentration, which, as expected had more impact on frazil volume fraction at the start of simulation.




Figure 5. Uncertainty propagation results for SSC (case 1) for temperature (a) and frazil (c) and MSC (case 2) for temperature (b) and frazil (d): mean, standard deviation, median, 5^{th} , 25^{th} , 75^{th} and 95^{th} percentiles are computed for t_k ($0 \leq k \leq n_t$).

For the SSC model, the initial concentration plays an important role at the start of the simulation. However, its influence quickly decreases, and the most influential parameter becomes the radius, with a peak influence reached at the median value of the time of maximum supercooling. Thus, using the average radius as the calibration parameter of the SSC model seems to be the most relevant choice. The most influential parameters after radius are diameter-to-thickness ratio and dissipation rate of turbulent kinetic energy. In experiments, turbulent dissipation rate is often better known than the initial concentration or crystals shape. In natural water bodies, turbulent dissipation rate measurements are rare but turbulence models can be used to estimate these parameters. Therefore, one could then choose initial frazil volume fraction or diameter-to-thickness ratio as a secondary calibration parameters.

For the MSC model, the most influential parameters prior to maximum supercooling are initial concentration (C_0) and maximum initial radius (r_0), that both control the initial distribution (and consequently the initial condition of the system). Using these initial distribution parameters to calibrate the transient phase until maximum supercooling seems to be the right approach. However, specifying more accurate initial distribution (not necessarily uniform) by comparison with what can be observed in nature would be a welcome improvement, although this requires further research.

At the recovery, the parameters of secondary nucleation and flocculation processes (n_{max} and a_f), both impacting steady-state crystal distribution, become more influential. However, the hierarchy of parameter is less obvious than for the (SSC) model. In addition, we observed strong interactions between parameters when the model reaches steady state (cf. blank space on Figure 6 and Total Sobol indices in Appendix F), which might lead to difficulties in the calibration process. Aggregated




Figure 6. Time series of first-order Sobol indices (S_i) for SSC (case 1) for temperature (a) and frazil (b) and MSC (case 2) for temperature (c) and frazil (d).

Sobol indices summarized in Figure (7), confirm the relative influence of each parameter over the whole duration of the simulation, taking into account both the transient phase and steady state.

While the approach adopted by Svensson and Omstedt (1994): i.e., tweaking the values of n_{max} and a_f , was adequate to calibrate the frazil distribution in (MSC) models, the present sensitivity analysis shows that it might not be the best option to calibrate water temperature and frazil total volume fraction. Results suggests that more focus should be on the initial condition to calibrate supercooling, by modifying initial seeding like it was done by Wang and Doering (2005) or by modifying the initial distribution itself. We therefore suggest calibrating the supercooling curve with the help of the initial distribution, along with secondary nucleation and flocculation parameters to calibrate the evolution of size distribution over time. Hopefully, recent observations of the transient evolution of frazil size distribution (McFarlane et al., 2015; Schneck et al., 2019), will provide the necessary data to carry out an optimal calibration of the identified parameters.

5.3 Influence of gravitational removal and seeding

Long-term evolution that does not take account of gravitational removal leads to infinite increase in frazil concentration as long as the cooling rate remains constant (cf. Equation 15). This asymptotic behavior of the models has never been observed in experiment nor in nature. Clark and Doering (2006) observed a peak in the number of particles per image they recorded, located shortly after maximum supercooling, after which there was a small decrease in the number of particles and a stagnation at residual supercooling. Similar observations were also reported by McFarlane et al. (2015) and Schneck et al. (2019). The




Figure 7. Aggregated first order Sobol indices (dots) and aggregated total Sobol indices (bars) for temperature (a) and total frazil volume fraction (b) for cases (1), (2), (3) and (4).

models in which only thermal growth is considered do not incorporate the required physics to properly reproduce what is observed. However by introducing gravitational removal, as shown in section (2.5), the models converge towards a constant frazil volume fraction (cf. Equation 16). In this section we analyze the results of the uncertainty propagation and sensitivity analysis for cases (3) and (4) which include the seeding rate and gravitational removal source terms.




Figure 8. Uncertainty propagation results for SSC (case 3) for temperature (a) and frazil (c) and MSC (case 4) for temperature (b) and frazil (d): mean, standard deviation, median, 5^{th} , 25^{th} , 75^{th} and 95^{th} percentiles computed for t_k ($0 \leq k \leq n_t$).

530 At steady state, the introduction of gravitational removal leads to wide scatter of both temperature and frazil as shown by the comparison of Figure (8) and (5) (see Appendix G and H for details). This is caused by the variation of the steady-state




Figure 9. Time series of first-order Sobol indices (S_i) for SSC (case 3) for temperature (a) and frazil (c) and MSC (case 4) for temperature (b) and frazil (d).

frazil volume fraction, which depends on the heat flux ϕ and model parameters inherent in secondary nucleation, flocculation and rise velocity (n_{max} , a_f and a_d). Significant supercooling values are observed as the water temperature (5th percentile) drops below -0.2°C which is really low compared to what can be observed in rivers. This is due to the highest values of the buoyancy velocity combined with low thermal growth rate. High gravitational removal withdraw a large amount of the frazil volume fraction from the water and thus limits the amount of latent heat release to water that would increase its temperature. No significant difference is observed in the output scatter between SSC and MSC models similarly to cases (1) and (2). However, minimum supercooling is not reached on the 5th percentile with the MSC as opposed to the SSC model. This is due to the dependency of the buoyancy velocity to the radius, which is not constant in the MSC model as opposed to the SSC model. This also cause the median frazil volume fraction at steady state to be slightly different in the two models.

With the SSC model at recovery, the first-order Sobol indices on frazil volume fraction ($S_i - C$) in Figure (9), show a major influence of the radius and the buoyancy coefficient ($S_{\bar{r}} = 0.381$ and $S_{a_d} = 0.382$, respectively at $t = t_f$), the main parameters influencing the gravitational removal (cf. Equation 15). For the MSC model, the most influential parameters at recovery are n_{max} and a_d ($S_{n_{max}} = 0.37$ and $S_{a_d} = 0.26$, respectively at $t = t_f$), which is consistent with Equation (27) and (29) of Rees Jones and Wells (2018). The hierarchy of the most influential parameters is similar to cases (1) and (2) prior to maximum supercooling. However accurate modeling of the buoyancy velocity of frazil crystals is essential, as it has a very important influence on the long-term evolution of the system, and therefore merits particular attention.

5.4 Maximum supercooling scatter

The results discussed above were obtained with a cooling rate of -500 W.m^{-3} . Several cooling rates, ranging from -50 to -1000 W.m^{-3} , were tested with case (1) to assess variations in maximum supercooling predictions. We found that the higher the cooling rate, the greater the scatter of the predicted maximum supercooling temperature, as presented in Figures (10) and (11). This is the opposite for the time until maximum supercooling peak, where the higher the cooling rate, the lower the scatter in supercooling time. These results are conform to observations of Carstens (1966) and Ye et al. (2004) that the degree of supercooling increases with the heat sink rate while the time to supercooling decreases. Note that the gap between 5^{th} and 95^{th} percentile maximum supercooling predictions is as much as $\Delta t_c = 2540 \text{ s}$ with $\phi = -50 \text{ W.m}^{-3}$, and $\Delta\theta = 0.142^\circ\text{C}$ with $\phi = -1000 \text{ W.m}^{-3}$ as shown in Figure (11). The dispersion both in time to supercooling and degree of supercooling is significant. This shows the importance of having a good good quantification of input parameters to be able to predict the maximum supercooling point.




Figure 10. Maximum supercooling point scatter computed from median, 5^{th} , 25^{th} , 75^{th} and 95^{th} percentile time series at different cooling rates ($\phi = -50, -100, -250, -500$, and -1000 W.m^{-3}) for case (1) and with $\phi = -500 \text{ W.m}^{-3}$ for case (2). Global view (a) and focus on initial time (b).

Results obtained from different scalings of the thermal boundary layer are also shown in Figure (11). A significant increase in scatter is observed for the $\delta_T = r$ compared to $\delta_T = \lambda$ scaling, consistent with the fact that thermal growth is clearly underestimated (Rees Jones and Wells, 2018). With δ_T taken as an uncertain parameter with a log-uniform PDF within the bounds $[\min(\lambda), \min(r)]$, the result is also more widely scattered by the same order of magnitude as with $\delta_T = r$. The results highlight the significant influence of the choice of scaling for the boundary layer. Choice of scaling also explains inconsistencies in calibrated parameters in literature.

Finally, let us discuss the results obtained with the MSC model, which need to be viewed from the standpoint of the way it is initialized. By taking the minimum and maximum radius as uncertain parameters (case 2b), we observe an increased gap between 5^{th} and 95^{th} percentiles and we have $\Delta t_c = 930 \text{ s}$ and $\Delta\theta = 0.09^\circ\text{C}$ (cf. Figure 11). The constant feed of first-




Figure 11. Maximum supercooling point scatter computed from median, 5^{th} , 25^{th} , 75^{th} and 95^{th} percentile time series at different cooling rates ($-50, -100, -250, -500$, and -1000 W.m^{-3}) for case (1), comparison between the choice of the length scale δ_T (cases 1b and 1c) and comparison between different initial conditions (cases 2b and 2c) with $\phi = -500 \text{ W.m}^{-3}$. Maximum degree of supercooling (a) and maximum supercooling time (b). White line is the median, wide box is delimited by 25^{th} and 75^{th} percentiles, thin line is delimited by 5^{th} and 95^{th} percentiles and round dots corresponds to minimum and maximum values.

class nuclei due to secondary nucleation accentuate the influence of the minimum radius parameter. This explains why the minimum radius has more influence on the results than the maximum radius as shown in Appendix D. Additionally the volume growth rate being higher for small classes, makes the initial distribution of concentration a determining choice. The more initial concentration is attributed to the smallest classes, the quicker the model reaches steady state, and the less scatter is observed in maximum supercooling time. The extreme case is when the initial concentration is only applied to the first class (case 2c), when an astonishingly narrow scatter of results is observed. In fact, the transient evolution is so quick that the model almost instantaneously converges towards its steady state. The median maximum supercooling is only -0.001°C and is reached in only 20 seconds. This confirms the sensitivity test carried out by Holland and Feltham (2005), who suggested distributing initial concentration on one class. However, the results show that this type of initialization might not be the best option, as it almost totally does away with transient evolution of the model.

5.5 Limits and perspectives

The importance of the availability of quantitative field and laboratory data, to support the uncertainty quantification of model parameters, cannot be stressed enough. As we have shown, many model parameter uncertainties could not be characterized by means of direct data. We had to use expert knowledge as well as past numerical implementation (Daly, 1984, 1994; Svensson and Omstedt, 1994; Smedsrud, 2002; Smedsrud and Jenkins, 2004; Wang and Doering, 2005; Holland and Feltham, 2005; Rees Jones and Wells, 2018). In part, this is because most physical processes, such as secondary nucleation or flocculation, are not directly measured, but instead are inferred from frazil distribution observations. Moreover, some parameters, such as the

585 initial condition, are dependent on the discretization method. Consequently, several parameter uncertainty bounds and PDFs should be refined. The modeling of dependency should also be taken into account in futur work as many parameters, such as δ_T and a_d , depend on other parameters (e.g., r and R). This would add a degree of complexity to the sensitivity analysis but innovative methods to tackle parameter dependency could bring valuable new insight on the models.

In this paper, we considered a well-mixed water body and a simplified gravitational removal sink term. Spatial variations for
590 temperature and frazil may result in different conclusions. A poorly mixed water body, where the cooling rate at higher layers is more severe than at the bottom, would cause an heterogeneity in the initial formation of frazil. Additionally, the meteorological seeding of frazil nuclei occurs mainly at the free surface, increasing the heterogeneity. Furthermore, as the rise velocity is higher for larger particles, the rise of frazil crystals and flocs yet again increases the heterogeneity of frazil on the vertical axis, altering the steady state distributions, as it was shown in the study by Hammar and Shen (1991). All these complex processes
595 make the conclusions for a well-mixed body difficult to extrapolate to multidimensional cases. Further research therefore may be warranted into uncertainty quantification and sensitivity analyses of frazil ice models in non-well mixed conditions with multidimensional models. The emergence of efficient APIs (Application Programming Interface) in numerical tools such as TELEMAC-MASCARET (Goeury et al., 2022), together with meta-modeling techniques that synthesize the essence of the multi-dimensional fields (Mouradi et al., 2021), could greatly assist such an undertaking.

600 Viewed from a different perspective, one may use the same probabilistic framework to compare modeling approaches for each process. This could be done by focusing on the volume fraction of each class, to obtain a good picture of how frazil is distributed over radius. In the same way that we considered time series of total frazil volume fraction as output, one could easily transpose the analysis to multi-dimensional class volume fraction as output. Uncertainty propagation could then allow for the characterization of PDFs associated with each class, and sensitivity analysis would shed light on the properties of each
605 model and how they affect frazil distributions. This could be a valuable tool for inferring new laws of secondary nucleation or flocculation by comparison to the observed evolution of frazil distributions.

6 Summary and conclusions

In this paper, two mathematical models for predicting of the evolution of frazil ice volume fraction and temperature have been studied. We developed a multiple-size-class MSC model, relying on the radial space discretization (Svensson and Omstedt,
610 1994) of frazil ice dynamic equations introduced by Daly (1984), which includes processes such as thermal growth, secondary nucleation, flocculation, seeding and gravitational removal. A simplified single-size-class SSC model, including only thermal growth, seeding and gravitational removal, was also developed for comparison. Properties of the two models, such as their steady states, were highlighted, and details provided for their numerical resolution. We found that, for proper resolution of the transient phase with the MSC model, a class number of about 100 is necessary for convergence of the results, which corroborates the observations by Rees Jones and Wells (2018). Uncertainties in both models were then studied within a probabilistic
615 framework. Aided by recent experimental and field studies of frazil ice, and also by numerical studies carried out in the recent years, the uncertainty of the main parameters of the frazil ice models were quantified. Various Monte Carlo experiments were

considered to propagate the uncertainties. Time dependent statistical estimates of the models' outputs (temperature and frazil volume fraction) were then analysed and a sensitivity analysis was carried out by means of a variance decomposition method.

620 Given the uncertainty bounds defined in the present study, SSC and MSC models yield very similar results for the prediction of water temperature and total frazil volume fraction. In the absence of gravitational removal, we have shown that the uncertainties have a great impact on the maximum supercooling and recovery time, but scarcely any impact on the steady state, which is governed only by cooling rate. The more detailed physics of the multi-class model, although providing valuable new information on size distribution of the crystals, does not make it possible to obtain a more reliable estimate of water temperature and total frazil volume fraction in the transient phase. The development of MSC models raises the possibility that uncertainty may be removed from choosing a mean radius. We have shown, however, that scatter is similar somehow in both models, and derives from new uncertain parameters inherent in radial space discretization. Note that, in many numerical tools, modeling frazil distribution requires the resolution of multiple advection-diffusion equations. Given the number of classes required for a model convergence, one can easily grasp the high numerical cost of using the MSC model for large scale, multi-dimensional applications. This makes the SSC model a suitable candidate for multi-dimensional frazil ice modeling, and the present study shows that it is still a very good compromise between uncertainty and model complexity. However, it should be noted that such uncertainty in the MSC model could be overcome in future laboratory experiments by a better estimation of the initial crystal size distribution.

635 The sensitivity analysis, allowed us to address with confidence the choice of calibration parameters. Relying on first- and total-order Sobol indices, we quantified the relative influence of each uncertain parameter on the output distribution, for both SSC and MSC models, and proposed a selection of parameters to be used for calibration. For the SSC model, the most influential factor for both temperature and frazil is the mean radius. Initial concentration played a secondary role although it was initially identified as a predominant factor. We therefore suggest using the average radius as the main calibration parameter. The turbulent dissipation rate also plays a major role and as such should be specified with care. As it can be estimated via turbulence models, we suggest using initial concentration and diameter-to-thickness ratio as secondary calibration parameters. With the MSC model, we showed that the dispersion is somehow similar to what we observed for the SSC model but originates from new uncertain parameters. Thus, the most influential parameters on the transient phase are the parameters specific to the initial condition. However, once the steady state was reached, we observed an increasing influence of the secondary nucleation and flocculation parameters. The long-term evolution of the system also showed increasing interactions between parameters. This could be explained by the balance in the physical processes involved in class interactions, and further investigation using class volume fraction as model outputs would help. When gravitational removal was introduced in the models, the stationary state was modified and the concentration converged towards a finite limit instead of diverging. In the case of the SSC model, the asymptotic limit is a function of the ratio between the gravitational removal term and the heat flux while in the case of the MSC model, the stationary state is also a function of the steady state radius distribution (which depends on the balance between secondary nucleation, flocculation and gravitational removal). Our study, confirming previous asymptotic analyses, showed that both secondary nucleation and gravitational removal parameters are the most influential on total frazil volume fraction. The buoyant rise velocity, the uncertainty of which was rarely taken into account in previous modeling studies, should

therefore be one of the main foci of future efforts to calibrate frazil ice models. Contrary to frazil, water temperature was mostly influenced by the initial condition, even at steady state. This should impel us to use both water temperature and frazil
655 volume fraction measurements to calibrate the models. Fortunately, recent laboratory and field studies (Schneck et al., 2019; McFarlane et al., 2015, 2017), particularly on the evolution of frazil distributions over time, offer precious data that can assist in developing appropriate calibration of the models. In this regard, using optimal calibration techniques, that allow consideration of both modeling and data uncertainties, would be a natural and complementary extension of the present study.

Appendix A: Semi-implicit theta scheme matrices

660 In this section, the semi-implicit time discretization of Equations 7 and 12 is described. Let us denote $t_k = t_0 + k\Delta t$ the time at iteration k , t_0 the initial time and $c^k = c(t_k)$. The semi-implicit time scheme consists in posing $c_i = \theta c_i^{k+1} + (1 - \theta)c_i^k$ in the right-hand side of Equation 7. Choosing $\theta = 0$ leads to a fully explicit time scheme while choosing $\theta = 1$ is equivalent to an implicit scheme on c . The non linear terms are treated semi-implicitly i.e. $G_i = G_i(t_k)$ in order to retrieve a linear system of the form $A[c_1^{k+1}, \dots, c_m^{k+1}]^T = B$ in which A and B are two matrices defined below. The temperature balance equation is then
665 solved with a forward Euler time scheme.

The matrix system reads: $A[c_1^{k+1}, \dots, c_m^{k+1}]^T = B \Leftrightarrow$

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1m} \\ a_{21} & a_{22} & a_{23} & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \ddots & & \\ & & a_{ii-1} & a_{ii} & a_{ii+1} & \vdots \\ \vdots & & & \ddots & \ddots & 0 \\ & & & & a_{m-1m} & \\ 0 & \dots & 0 & a_{mm-1} & a_{mm} \end{pmatrix} \begin{pmatrix} c_1^{k+1} \\ \vdots \\ c_i^{k+1} \\ \vdots \\ c_m^{k+1} \end{pmatrix} = \begin{pmatrix} b_1 \\ \vdots \\ b_i \\ \vdots \\ b_N \end{pmatrix}, \quad (\text{A1})$$

in which the diagonal terms are defined as:

$$\begin{aligned} a_{11} &= 1 - \theta\Delta t (V_1(\Lambda_1 - \Gamma_1) - \beta_1 - \gamma_1), \\ 670 \quad a_{ii} &= 1 - \theta\Delta t (V_i(\Lambda_i - \Gamma_i) - \beta_i - \gamma_i - \zeta\alpha_i) \quad \text{for } 2 \leq i \leq m-1, \\ a_{mm} &= 1 - \theta\Delta t (V_m\Lambda_m - \gamma_m - \zeta\alpha_m), \end{aligned}$$

the lower off-diagonal terms are defined as:

$$a_{ii-1} = -\theta\Delta t (V_i\Gamma_{i-1} + \beta_{i-1}) \quad \text{for } 2 \leq i \leq m,$$

the upper off-diagonal terms are defined as:

$$\begin{aligned} 675 \quad a_{1i} &= -\theta\Delta t\alpha_i \quad \text{for } 2 \leq i \leq m, \\ a_{ii+1} &= \theta\Delta t V_i\Lambda_{i+1} \quad \text{for } 2 \leq i \leq m-1, \end{aligned}$$

and the matrix B is defined as:

$$\begin{aligned} b_1 &= c_1^k + \Delta t\tau_s V_1/h + (1 - \theta)\Delta t [V_1((\Lambda_1 - \Gamma_1)c_1^k - \Lambda_2 c_2^k) - \beta_1 c_1^k - \gamma_1 c_1^k] + (1 - \theta)\Delta t \sum_{j=2}^m \alpha_j c_j^k, \\ b_i &= c_i^k + (1 - \theta)\Delta t [V_i(\Gamma_{i-1}c_{i-1}^k + (\Lambda_i - \Gamma_i)c_i^k - \Lambda_{i+1}c_{i+1}^k) + \beta_{i-1}c_{i-1}^k - \beta_i c_i^k - \gamma_i c_i^k - \zeta\alpha_i c_i^k] \\ 680 \quad &\text{for } 2 \leq i \leq m-1, \\ b_m &= c_m^k + (1 - \theta)\Delta t [V_m(\Gamma_{m-1}c_{m-1}^k + \Lambda_m c_m^k) + \beta_{m-1}c_{m-1}^k - \gamma_m c_m^k - \zeta\alpha_m c_m^k], \end{aligned}$$

Appendix B: Sensitivity analysis using Sobol indices

For a given set of independent input parameters, the ANOVA (ANalyses Of VAriance) decomposition allows us to compute the variance of output $\mathbf{Y} = g(\mathbf{X}, \mathbf{d})$, for each time step t_k ($1 \leq k \leq n_t$) as:

$$685 \quad \text{Var}[Y^k] = \sum_{i=1}^{n_X} V_i(Y^k) + \sum_{i < j} V_{ij}(Y^k) + \dots + V_{1\dots n_X}(Y^k), \quad (\text{B1})$$

where :

$$\begin{aligned} V_i(Y^k) &= \text{Var}[\mathbb{E}[Y^k | X^i]] \\ V_{ij}(Y^k) &= \text{Var}[\mathbb{E}[Y^k | X^i, X^j] - \mathbb{E}[Y^k | X^i] - \mathbb{E}[Y^k | X^j]] \\ &= \text{Var}[\mathbb{E}[Y^k | X^i, X^j]] - V_i(Y^k) - V_j(Y^k) \end{aligned}$$

690 in which $\mathbb{E}[Y^k | X^i]$ represents the conditional expectation of Y^k with the condition that X^i remains constant.

To evaluate the influence of each input parameter, the so-called Sobol indices are used (Sobol, 2001). The first and second order Sobol indices are defined as follows, for $k \in \{1, \dots, n_t\}$ and $i \in \{1, \dots, n_X\}$:

$$\begin{aligned} S_i^k &= \frac{V_i(Y^k)}{\text{Var}[Y^k]} \\ S_{ij}^k &= \frac{V_{ij}(Y^k)}{\text{Var}[Y^k]} \end{aligned} \quad (\text{B2})$$

The first-order Sobol index S_i^k indicates the part of output variance explained by a single parameter X_i without interactions, whereas second-order indices S_{ij}^k quantify the part of variance of the output explained by the interaction between two inputs X_i and X_j . The number of second order indices is given by $\binom{n_X}{2} = n_X(n_X - 1)/2$. When the number of input parameters is too large, it may be difficult to estimate second-order indices. In that case, we only estimate first-order and total indices. Total indices quantify the part of variance of the output explained by an input and its interactions with all the other inputs parameters. Total Sobol indices are defined as follows, for $k \in \{1, \dots, n_t\}$ and $i \in \{1, \dots, n_X\}$:

$$700 \quad ST_i^k = S_i^k + \sum_{i \neq j} S_{ij}^k + \sum_{i \neq j, l \neq i, j \leq l} S_{ijl}^k + \dots = \frac{VT_i(Y^k)}{\text{Var}[Y^k]} = 1 - \frac{V_{-i}(Y^k)}{\text{Var}[Y^k]} \quad (\text{B3})$$

where $V_{-i}(Y^k) = \text{Var}[\mathbb{E}[Y^k | X^1, \dots, X^{i-1}, X^{i+1}, \dots, X^{n_X}]]$.

To compute Sobol indices, a modified version of the method proposed by Saltelli (2002) is used (Baudin et al., 2016a), in which two independent samples of size N , denoted \mathbf{A} and \mathbf{B} are generated. Both can be written as matrices (cf. Equation B4) in which each line is a realization of the random vector \mathbf{X} . A third matrix, denoted \mathbf{C}^i is then created by replacing only the column i of the matrix \mathbf{A} by the column i of the matrix \mathbf{B} (cf. Equation B4).

$$705 \quad \mathbf{A} = \begin{pmatrix} x_1^{A,1} & x_2^{A,1} & \dots & x_{n_X}^{A,1} \\ x_1^{A,2} & x_2^{A,2} & \dots & x_{n_X}^{A,2} \\ \vdots & \vdots & \ddots & \vdots \\ x_1^{A,N} & x_2^{A,N} & \dots & x_{n_X}^{A,N} \end{pmatrix}, \mathbf{C}^i = \begin{pmatrix} x_1^{A,1} & \dots & x_i^{B,1} & \dots & x_{n_X}^{A,1} \\ x_1^{A,2} & \dots & x_i^{B,2} & \dots & x_{n_X}^{A,2} \\ \vdots & & \vdots & \ddots & \vdots \\ x_1^{A,N} & \dots & x_i^{B,N} & \dots & x_{n_X}^{A,N} \end{pmatrix}. \quad (\text{B4})$$

First order and total Sobol indices are computed using estimations of $V_i(Y^k)$ and $V_{-i}(Y^k)$ computed using samples \mathbf{A} , \mathbf{B} and \mathbf{C}^i and noted $\hat{V}_i(Y^k)$ and $\hat{V}_{-i}(Y^k)$ respectively. These estimations are defined as follows:

$$\begin{aligned}\hat{V}_i(Y^k) &= \frac{1}{N-1} \sum_{j=1}^N \tilde{g}(\mathbf{B}_j) \tilde{g}(\mathbf{C}_j^i) - \left(\frac{1}{N} \sum_{j=1}^N \tilde{g}(\mathbf{A}_j) \right) \left(\frac{1}{N} \sum_{j=1}^N \tilde{g}(\mathbf{B}_j) \right), \\ \hat{V}_{-i}(Y^k) &= \frac{1}{N-1} \sum_{j=1}^N \tilde{g}(\mathbf{A}_j) \tilde{g}(\mathbf{C}_j^i) - \left(\frac{1}{N} \sum_{j=1}^N \tilde{g}(\mathbf{A}_j) \right) \left(\frac{1}{N} \sum_{j=1}^N \tilde{g}(\mathbf{B}_j) \right),\end{aligned}\tag{B5}$$

715 where \tilde{g} is the centered model defined by $\tilde{g} = g - \bar{g}$ in which \bar{g} is the sample mean of the combined output samples $g(\mathbf{A})$ and $g(\mathbf{B})$. To compute the second-order Sobol indices, an additional matrix is used, noted \mathbf{C}'^j which is created by replacing only the column i of the matrix \mathbf{B} by the column i of the matrix \mathbf{A} . Then the estimation $\hat{V}_{ij}(Y^k)$ is computed as:

$$\hat{V}_{ij}(Y^k) = \frac{1}{N-1} \sum_{m=1}^N \tilde{g}(\mathbf{C}_m^i) \tilde{g}(\mathbf{C}_m'^j) - \frac{1}{N} \sum_{m=1}^N \tilde{g}(\mathbf{A}_m) \tilde{g}(\mathbf{B}_m) - \hat{V}_i(Y^k) - \hat{V}_j(Y^k).\tag{B6}$$

For a sample size N , estimation of the first-order and total Sobol indices requires $(n_X + 2) \times N$ simulations.

715 For multivariate outputs, the indices can be aggregated as proposed by Gamboa et al. (2014). The aggregated first order and total Sobol indices are defined as:

$$\begin{aligned}AS_i &= \frac{\sum_{k=1}^{n_t} V_i(Y^k)}{\sum_{k=1}^{n_t} \text{Var}[Y^k]} \\ AST_i &= \frac{\sum_{k=1}^{n_t} VT_i(Y^k)}{\sum_{k=1}^{n_t} \text{Var}[Y^k]}\end{aligned}\tag{B7}$$

This means that Sobol indices S_i^k and ST_i^k quantify the influence of X_i on the variance of Y at time t_k , while the aggregated indices AS_i and AST_i quantify the influence of X_i over the whole time series of Y .

Convergence of statistical estimators is addressed by running several Monte Carlo simulations with increasing sampling size. An example of the convergence of mean and standard deviation at different times is given in Figure C.1 for case (1). Similarly, testing of Sobol indices convergence is shown in Figure C.2 for case (1). Note that the 5th and 95th confidence intervals are systematically computed by a bootstrap method, and plotted in all Sobol index Figures.




Figure C.1. Mean and standard deviation convergence for case (1) for temperature (a) and frazil (b).




Figure C.2. First order Sobol indices convergence for case (1) for temperature (a) and frazil (b).




Figure D.1. First order and total order Sobol indices at times t_{min} , $2t_{min}$ and t_f and aggregated Sobol indices for temperature (left) and frazil (right) the case 1 (a), 2 (b), 3 (c) and 4 (d) from top to bottom.

Appendix E: Aggregated Sobol indices




Figure E.1. Aggregated first order Sobol indices (dots) and aggregated total Sobol indices (bars) for temperature (a) and total frazil volume fraction (b) for the case 1, 1b, 1c, 2, 2b, 2c, 3 and 4.

Appendix F: Results of case (1)




Figure F.1. Results of case (1). From top to bottom: uncertainty propagation result (a), first-order Sobol indices (b), first order Sobol indices with 95% confidence intervals (c), total order Sobol indices with 95% confidence intervals (d), for temperature (left) and frazil (right).

Appendix G: Results of case (2)




Figure G.1. Results of case (2). From top to bottom: uncertainty propagation result (a), first-order Sobol indices (b), first order Sobol indices with 95% confidence intervals (c), total order Sobol indices with 95% confidence intervals (d), for temperature (left) and frazil (right).

Appendix H: Results of case (3)




Figure H.1. Results of case (3). From top to bottom: uncertainty propagation result (a), first-order Sobol indices (b), first order Sobol indices with 95% confidence intervals (c), total order Sobol indices with 95% confidence intervals (d), for temperature (left) and frazil (right).




Figure I.1. Results of case (4). From top to bottom: uncertainty propagation result (a), first-order Sobol indices (b), first order Sobol indices with 95% confidence intervals (c), total order Sobol indices with 95% confidence intervals (d), for temperature (left) and frazil (right).

Code availability. Frazil ice models presented in this paper have been implemented in the TELEMAC-MASCARET open source suite of solvers: <http://www.opentelemac.org>. The uncertainty analysis was carried out using OpenTURNS (Open source initiative for the Treatment of Uncertainties, Risks'N Statistics) which can be found here: <http://openturns.github.io/openturns/latest/index.html>.

735 *Author contributions.* FS, CG, and RSM conceptualized the study. FS conducted the analyses, interpreted the results, and wrote the manuscript with input from CG and RSM.

Competing interests. The contact author has declared that none of the authors has any competing interests.

740 *Acknowledgements.* This research was funded by EDF. The authors would like to thank Mark Loewen and David W. Rees Jones whose comments and suggestions helped improve the scientific quality of this paper. The authors would also like to address special thanks to Michael Baudin for his reading and discussion that enriched this paper. The authors gratefully acknowledge contributions from the OpenTURNS open-source community.

References

- Arakawa, K.: Studies on the Freezing of Water (II) Formation of disc crystals, *Journal of the Faculty of Science, Hokkaido University*, 4, 311–339, 1954.
- Ashton, G.: Frazil ice, in: *Theory of dispersed multiphase flow*, pp. 271–289, Elsevier, 1983.
- 745 Barrette, P. D.: A tabulated review of 83 laboratory studies on frazil ice, in: *IAHR International Symposium on Ice*, 2020.
- Barrette, P. D.: Understanding frazil ice: The contribution of laboratory studies, *Cold Regions Science and Technology*, 189, 103 334, <https://doi.org/10.1016/j.coldregions.2021.103334>, 2021.
- Baudin, M., Boumhaout, K., Delage, T., Iooss, B., and Martinez, J.-M.: Numerical stability of Sobol’ indices estimation formula, in: *Proceedings of the 8th International Conference on Sensitivity Analysis of Model Output (SAMO 2016)*. Le Tampon, Réunion Island, France, 2016a.
- 750 Baudin, M., Dutfoy, A., Iooss, B., and Popelin, A.-L.: OpenTURNS: An Industrial Software for Uncertainty Quantification in Simulation, pp. 1–38, Springer International Publishing, https://doi.org/10.1007/978-3-319-11259-6_64-1, 2016b.
- Bombosch, A. and Jenkins, A.: Modeling the formation and deposition of frazil ice beneath Filchner-Ronne Ice Shelf, *Journal of Geophysical Research: Oceans*, 100, 6983–6992, 1995.
- 755 Carstens, T.: Experiments with supercooling and ice formation in flowing water, *Geofys. Publ. Norway*, 26(9-18), 1966.
- Clark, S. and Doering, J.: Laboratory Experiments on Frazil-Size Characteristics in a Counterrotating Flume, *Journal of Hydraulic Engineering*, 132, 94–101, [https://doi.org/10.1061/\(ASCE\)0733-9429\(2006\)132:1\(94\)](https://doi.org/10.1061/(ASCE)0733-9429(2006)132:1(94)), 2006.
- Clark, S. and Doering, J.: Frazil flocculation and secondary nucleation in a counter-rotating flume, *Cold Regions Science and Technology*, 55, 221–229, <https://doi.org/10.1016/j.coldregions.2008.04.002>, 2009.
- 760 Clark, S. and Doering, J. C.: A laboratory study of frazil ice size distributions, *Proc.*, 17th Int.Symp.on Ice, pp. 291–297, 2004.
- Daly, S. F.: Frazil Ice Dynamics, Technical Report 84-1. U.S. Army Cold Regions Research and Engineering Laboratory, Hanover, New Hampshire., 1984.
- Daly, S. F.: Frazil Ice Blockage of Intake Trash Racks, *Cold Regions Technical Digest No. 91-1*, Tech. rep., US Army Corps of Engineers, 1991.
- 765 Daly, S. F.: Report on Frazil Ice, Tech. rep., International Association for Hydraulic Research, Working Group on Thermal Regimes, Special Report 94-23, 1994.
- Daly, S. F.: Frazil Ice Blockage of Water Intakes in the Great Lakes, *Journal of Hydraulic Engineering*, 2006.
- Daly, S. F. and Colbeck, S. C.: Frazil ice measurements in CRREL’s flume facility., in: *IAHR Symposium on Ice 1986 - Proceedings.*, vol. 1, pp. 427–438, 1986.
- 770 Doering, J. C. and Morris, M. P.: River Ice Engineering / Ingénierie des glaces fluviales A digital image processing system to characterize frazil ice, *Canadian Journal of Civil Engineering*, 30, 1–10, <https://doi.org/10.1139/102-028>, 2003.
- Ettema, R., Karim, M., and Kennedy, J.: Laboratory experiments on frazil ice growth in supercooled water, *Cold Regions Science and Technology*, 10, 43–58, [https://doi.org/10.1016/0165-232X\(84\)90032-6](https://doi.org/10.1016/0165-232X(84)90032-6), 1984.
- Frazer, E. K., Langhorne, P. J., Leonard, G. H., Robinson, N. J., and Schumayer, D.: Observations of the size distribution of frazil ice in an Ice Shelf Water plume, *Geophysical Research Letters*, 47, e2020GL090498, 2020.
- 775 Gamboa, F., Janon, A., Klein, T., and Lagnoux, A.: Sensitivity analysis for multidimensional and functional outputs, *Electronic Journal of Statistics*, 8, 575–603, 2014.

- Ghobrial, T. R., Loewen, M. R., and Hicks, F.: Laboratory calibration of upward looking sonars for suspended frazil ice concentration, *Cold Regions Science and Technology*, 70, 19–31, <https://doi.org/10.1016/j.coldregions.2011.08.010>, 2012.
- 780 Ghobrial, T. R., Loewen, M. R., and Hicks, F. E.: Characterizing suspended frazil ice in rivers using upward looking sonars, *Cold Regions Science and Technology*, 86, 113–126, <https://doi.org/10.1016/j.coldregions.2012.10.002>, 2013.
- Goeury, C., Audouin, Y., and Zaoui, F.: Interoperability and computational framework for simulating open channel hydraulics: Application to sensitivity analysis and calibration of Gironde Estuary model, *Environmental Modelling & Software*, 148, 105 243, 2022.
- Gosink, J. P. and Osterkamp, T. E.: Measurements and Analyses of Velocity Profiles and Frazil Ice-Crystal Rise Velocities During Periods of Frazil-Ice Formation in Rivers, *Annals of Glaciology*, 4, 79–84, <https://doi.org/10.3189/S0260305500005279>, 1983.
- 785 Hammar, L. and Shen, H. T.: A mathematical model for frazil ice evolution and transport in channels, in: *Proc., 6th Workshop on the Hydraulics of River Ice*, pp. 201–206, 1991.
- Hammar, L. and Shen, H. T.: Frazil evolution in channels, *Journal of Hydraulic Research*, 33, 291–306, <https://doi.org/10.1080/00221689509498572>, 1995.
- 790 Holland, P. R. and Feltham, D. L.: Frazil dynamics and precipitation in a water column with depth-dependent supercooling, *Journal of Fluid Mechanics*, 530, 101–124., 2005.
- Holland, P. R., Feltham, D. L., and Daly, S. F.: On the Nusselt number for frazil ice growth—a correction to “Frazil evolution in channels” by Lars Hammar and Hung-Tao Shen, *Journal of Hydraulic Research*, 45, 421–424, <https://doi.org/10.1080/00221686.2007.9521775>, 2007.
- Kempema, E., Reimnitz, E., and Hunter, R. E.: Flume studies and field observations of the interaction of frazil ice and anchor ice with sediments, US Department of the Interior, Geological Survey, 1986.
- 795 Kempema, E. W. and Ettema, R.: Fish, Ice, and Wedge-Wire Screen Water Intakes, *Journal of Cold Regions Engineering*, 30, 04015 004, [https://doi.org/10.1061/\(ASCE\)CR.1943-5495.0000097](https://doi.org/10.1061/(ASCE)CR.1943-5495.0000097), 2016.
- Lal, D., Mason, R., and Strickland-Constable, R.: Collision breeding of crystal nuclei, *Journal of crystal growth*, 5, 1–8, 1969.
- MacGillivray, B. H.: Handling uncertainty in models of seismic and postseismic hazards: toward robust methods and resilient societies, *Risk analysis*, 41, 1499–1512, 2021.
- 800 Marko, J. and Jasek, M.: Sonar detection and measurement of ice in a freezing river II: Observations and results on frazil ice, *Cold Regions Science and Technology*, 63, 135–153, <https://doi.org/10.1016/j.coldregions.2010.05.003>, 2010.
- Matoušek, V.: Frazil and skim ice formation in rivers, in: *Proceedings of the IAHR Ice Symposium*, 1992.
- McFarlane, V., Loewen, M., and Hicks, F.: Laboratory experiments to determine frazil properties, *Proceedings of the Annual General Conference of the Canadian Society for Civil Engineering*, Edmonton, Alberta, Canada, p. 10, 2012.
- 805 McFarlane, V., Loewen, M., and Hicks, F.: Laboratory measurements of the rise velocity of frazil ice particles, *Cold regions science and technology*, 106, 120–130, 2014.
- McFarlane, V., Loewen, M., and Hicks, F.: Measurements of the evolution of frazil ice particle size distributions, *Cold Regions Science and Technology*, 120, 45–55, <https://doi.org/10.1016/j.coldregions.2015.09.001>, 2015.
- 810 McFarlane, V., Loewen, M., and Hicks, F.: Field observations of the growth rate of anchor ice crystals, in: *Proceedings of the 23rd IAHR International Symposium on Ice*, Ann Arbor, MI, USA, 2016.
- McFarlane, V., Loewen, M., and Hicks, F.: Measurements of the size distribution of frazil ice particles in three Alberta rivers, *Cold Regions Science and Technology*, 142, 100–117, <https://doi.org/10.1016/j.coldregions.2017.08.001>, 2017.
- Mercier, R. S.: The reactive transport of suspended particles: Mechanics and modeling, in: *PhD dissertation*, Joint Committee on Oceanographic Engineering, 1984.
- 815

- Michel, B.: Theory of formation and deposit of frazil ice, Eastern Snow Conference, Proc. Annual Meeting, Quebec., 1963.
- Morse, B. and Richard, M.: A field study of suspended frazil ice particles, *Cold Regions Science and Technology*, 55, 86–102, <https://doi.org/10.1016/j.coldregions.2008.03.004>, 2009.
- 820 Mouradi, R.-S., Goeury, C., Thual, O., Zaoui, F., and Tassi, P.: Physically interpretable machine learning algorithm on multidimensional non-linear fields, *Journal of Computational Physics*, 428, 110 074, 2021.
- Muller, A.: Frazil ice formation in turburemt flow, *Proc. Int. Assoc. of Hydr. Res., Sympo. on Ice*, Lulea, Sweden, 1978, 1978.
- Omstedt, A.: On Supercooling and Ice Formation in Turbulent Sea-Water, *Journal of Glaciology*, 31, 1985.
- Osterkamp, T. and Gosink, J.: Frazil ice formation and ice cover development in interior Alaska streams, *Cold Regions Science and Technology*, 8, 43–56, 1983a.
- 825 Osterkamp, T. and Gosink, J.: Frazil ice formation and ice cover development in interior Alaska streams, *Cold Regions Science and Technology*, 8, 43–56, [https://doi.org/10.1016/0165-232X\(83\)90016-2](https://doi.org/10.1016/0165-232X(83)90016-2), 1983b.
- Razavi, S., Jakeman, A., Saltelli, A., Prieur, C., Iooss, B., Borgonovo, E., Plischke, E., Lo Piano, S., Iwanaga, T., Becker, W., Tarantola, S., Guillaume, J. H., Jakeman, J., Gupta, H., Melillo, N., Rabitti, G., Chabridon, V., Duan, Q., Sun, X., Smith, S., Sheikholeslami, R., Hosseini, N., Asadzadeh, M., Puy, A., Kucherenko, S., and Maier, H. R.: The Future of Sensitivity Analysis: An essential discipline for
- 830 systems modeling and policy support, *Environmental Modelling & Software*, 137, 104 954, 2021.
- Rees Jones, D. W. and Wells, A. J.: Solidification of a disk-shaped crystal from a weakly supercooled binary melt, *Physical Review E*, 92, 022 406, 2015.
- Rees Jones, D. W. and Wells, A. J.: Frazil-ice growth rate and dynamics in mixed layers and sub-ice-shelf plumes, *The Cryosphere*, 12, 25–38, <https://doi.org/10.5194/tc-12-25-2018>, 2018.
- 835 Richard, M. and Morse, B.: Multiple frazil ice blockages at a water intake in the St. Lawrence River, *Cold Regions Science and Technology*, 53, 131–149, 2008.
- Saltelli, A.: Making best use of model evaluations to compute sensitivity indices, *Computer physics communications*, 145, 280–297, 2002.
- Saltelli, A.: A short comment on statistical versus mathematical modelling, *Nature communications*, 10, 1–3, 2019.
- Schaefer, V. J.: The formation of frazil and anchor ice in cold water, *Eos, Transactions American Geophysical Union*, 31, 885–893, <https://doi.org/10.1029/TR031i006p00885>, 1950.
- 840 Schneck, C. C., Ghobrial, T. R., and Loewen, M. R.: Laboratory study of the properties of frazil ice particles and flocs in water of different salinities, *The Cryosphere*, 2019.
- Sheikholeslami, R., Yassin, F., Lindenschmidt, K.-E., and Razavi, S.: Improved understanding of river ice processes using global sensitivity analysis approaches, *Journal of Hydrologic Engineering*, 22, 04017 048, 2017.
- 845 Shen, H. T.: Mathematical modeling of river ice processes, *Cold Regions Science and Technology*, 62, 3–13, 2010.
- Shen, H. T. and Wang, D. S.: Under cover transport and accumulation of frazil granules, *Journal of Hydraulic Engineering*, 121, 184–195, 1995.
- Shen, H. T. and Wasantha Lal, A. M.: A Mathematical Model for River Ice Processes, Tech. rep., US Army Corps of Engineers Cold Regions Research and Engineering Laboratory, 1993.
- 850 Shen, H. T., Wang, D. S., and Lal, A. W.: Numerical simulation of river ice processes, *Journal of Cold Regions Engineering*, 9, 107–118, 1995.
- Smedsrud, L. H.: A model for entrainment of sediment into sea ice by aggregation between frazil-ice crystals and sediment grains, *Journal of Glaciology*, 48, 51–61, 2002.

- Smedsrud, L. H. and Jenkins, A.: Frazil ice formation in an ice shelf water plume, *Journal of Geophysical Research: Oceans*, 109, 2004.
- 855 Sobol, I. M.: Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates, *Mathematics and computers in simulation*, 55, 271–280, 2001.
- Soize, C.: *Uncertainty quantification*, Springer, 2017.
- Souill  , F., Taccone, F., and El Mertahi, C.: A Multi-class Frazil Ice Model for Shallow Water Flows, in: *TELEMAC-MASCARET User Conference October 2020*, pp. 122–129, 2020.
- 860 Sudret, B.: Uncertainty propagation and sensitivity analysis in mechanical models—Contributions to structural reliability and stochastic spectral methods, *Habilitationa diriger des recherches*, Universit   Blaise Pascal, Clermont-Ferrand, France, 147, 53, 2007.
- Svensson, U. and Omstedt, A.: Simulation of supercooling and size distribution in frazil ice dynamics, *Cold regions science and technology*, 1994.
- Tsang, G. and Hanley, T. O.: Frazil Formation in Water of Different Salinities and Supercoolings, *Journal of Glaciology*, 31, 1985.
- 865 Van Zelm, R. and Huijbregts, M. A.: Quantifying the trade-off between parameter and model structure uncertainty in life cycle impact assessment, *Environmental science & technology*, 47, 9274–9280, 2013.
- Wang, S. M. and Doering, J. C.: Numerical Simulation of Supercooling Process and Frazil Ice Evolution, *Journal of Hydraulic Engineering*, 2005.
- Wuebben, J. L.: The rise pattern and velocity of frazil ice, in: *the proceedings of the Third Workshop on the Hydraulics of Ice Covered*
- 870 *Rivers*, Session F, pp. 297–316, 1984.
- Wueben, J.: The rise pattern and velocity of frazil ice, *CRIPE*, in: *Proceedings of the 3rd Workshop on the Hydraulics of Ice Covered Rivers*, Fredericton, NB, 1984.
- Ye, S. Q. and Doering, J.: Simulation of the supercooling process and frazil evolution in turbulent flows, *Canadian Journal of Civil Engineering*, 31, 915–926, <https://doi.org/10.1139/I04-055>, 2004.
- 875 Ye, S. Q., Doering, J., and Shen, H. T.: A laboratory study of frazil evolution in a counter-rotating flume, *Canadian Journal of Civil Engineering*, 31, 899–914, <https://doi.org/10.1139/I04-056>, 2004.
- Zacharov, V., Bejlinson, M., and   atalina, I.: Features of ice conditions in rivers and reservoirs of central Asia, in: *Proceedings of IAHR Symposium on Ice and its action on Hydraulic Structures*, Leningrad, Russia, pp. 224–228, 1972.