



Review article: large fluctuations in non-equilibrium physics

Giovanni Jona-Lasinio

Dipartimento di Fisica and INFN, Università di Roma “La Sapienza”
Piazzale A. Moro 2, Roma 00185, Italy

Correspondence: Gianni Jona Lasinio (gianni.jona@roma1.infn.it)

Abstract. Non-equilibrium is dominant in geophysical and climate phenomena. However the study of non-equilibrium is much more difficult than equilibrium and the relevance of probabilistic simplified models has been emphasized. Large deviation rates have been used recently in climate science. In this paper after recalling progress during the last decades in understanding the role of large deviations in a class of non-equilibrium systems we point out differences between equilibrium and non-equilibrium. For example in non-equilibrium: a) large deviation rates may be extensive but not simply additive. b) In non-equilibrium there are generically long range space correlations so large deviation rates are non-local. c) Singularities in large deviation rates denote the existence of phase transitions often not possible in equilibrium. To exemplify we shall refer to lattice gas models like the symmetric simple exclusion process and other models which are playing an important role in the understanding of non-equilibrium physics. Reasons why all this may be of interest in climate physics will be briefly indicated.

10 **Keywords:**

1 Introduction

This paper, is an enlarged version of a seminar in the series "*Perspectives in Climate Sciences 2021*" and provides an introduction to the main ideas of the so called macroscopic fluctuation theory (MFT). This theory developed mainly for non-equilibrium diffusive systems is very well supported by mathematical models and provides a methodological approach that can be followed for other non-equilibrium systems.

The theory is based on variational principles determining for each model the large deviation rates (LDR) of thermodynamic variables like a density or a time average like the time averaged current flowing through the system in stationary states. Models in climate science are considerably more complex than those for which MFT has been proved to hold. However MFT may provide a guide for more complex problems and it shows that there are considerable differences in fluctuations going from an equilibrium to a non-equilibrium state.

In the last decades the theory of large deviations has become a main tool in statistical mechanics especially in the study of non-equilibrium. MFT has been formulated and verified in probabilistic models of lattice gases which macroscopically lead to hydrodynamic diffusion equations which represent laws of large numbers. Therefore large deviations from hydrodynamic behavior have been studied. Here we shall follow the formulation given in (Bertini et al., 2001, 2002) and the reviews (Bertini et al., 2007, 2015). The reader is advised to refer also to (Derrida, 2007, 2011). Although the theory has been developed especially for diffusions, several conclusions seem to hold more generally, e.g. for **reaction-diffusion systems**.



Non-equilibrium is dominant in geophysical and climate phenomena and we refer to the following papers for the relevance of large deviation estimates in this domain (Galfi et al., 2019; Galfi & Lucarini, 2021; Galfi et al., 2021). In (Galfi et al., 2019) a Simplified Model of the General Circulation of the Atmosphere is adopted and assumed to be a chaotic dynamical system so that probabilistic concepts and methods can be applied. Stochastic climate models were introduced by Hasselmann (Hasselmann, 1976).

Non-equilibrium includes an enormous variety of phenomena so we cannot hope to formulate a unique theory having a generality comparable to classical thermodynamics. We have to restrict to subclasses of problems. One difficulty is to define suitable thermodynamic functionals in far from equilibrium situations. Large fluctuations have offered a way out as large deviation rates provide genuine thermodynamic functionals in non-equilibrium stationary states.

Studying irreversible processes is much more difficult than understanding equilibrium phenomena. In equilibrium statistical mechanics we do not have to solve any equation of motion and the Gibbs distribution provides the basis for the calculation of macroscopic quantities and their fluctuations. In non equilibrium we cannot bypass the dynamics even in the study of stationary states (SNS) which we may consider as the simplest beyond equilibrium. Examples are the heat flow in an iron rod whose endpoints are thermostated at different temperatures or the stationary flow of electrical current in a given potential difference.

For such states the fluctuations exhibit novel and rich features with respect to the equilibrium situation. As experimentally observed, the space correlations of the thermodynamic variables generically extend to macroscopic distances, which means that the fluctuations of the density in different points of the system are not independent. This is reflected in the structure of large deviation rates which in non-equilibrium stationary states are in general non local in space.

In this paper we consider a class of systems for which it is possible at the macroscopic scale to define local thermodynamic variables and behave hydrodynamically as diffusions. The systems considered locally deviate only slightly from equilibrium so that small gradients and linear response to external fields are reasonable approximations. Microscopically, this implies that the system reaches a local equilibrium in a time which is short compared to the times typical of macroscopic evolution. So what characterizes situations in which this description applies is a separation of scales both in space and time. Far from equilibrium states are those which exhibit differences over the size of the whole system. In other words far from equilibrium is obtained from building up small local differences. **Local equilibrium is the first assumption we make.**

This restriction allows us to understand some typical phenomena induced by non-equilibrium in somewhat ideal cases which however give a hint of what may happen in more realistic cases.

The content of the paper: in the next section we recall the use by Einstein of the Boltzmann relation between entropy and probability to estimate the probability of a thermodynamic fluctuation in equilibrium. In section 3 we describe the essentials of the macroscopic fluctuation theory in the version reviewed in (Bertini et al., 2015). Section 4 is devoted to the illustration of the differences between equilibrium and non-equilibrium through explicit calculations for the symmetric simple exclusion process (SSEP). Final comments are in section 5. Our exposition is not very detailed and if the reader wants to understand the calculations behind we suggest to consult the reviews (Bertini et al., 2015; Derrida, 2007, 2011; Mallick, 2015) and references therein.



2 Macroscopic fluctuations in equilibrium

The first explicit large deviation theory in equilibrium states is presumably Einstein theory of opalescence (Einstein, 1910), see also (Landau & Lifshitz, 1980). He uses the Boltzmann-Planck formula connecting entropy and probability to estimate probabilities of fluctuations of thermodynamic variables, e.g. of densities. Therefore entropy appears as a large deviation rate. The small parameter representing the intensity of the noise in this case is the Boltzmann constant (Bertini et al., 2015), $\kappa = R/N$, R the gas constant, N Avogadro's number, so that its inverse is proportional to the typical number of the microscopic degrees of freedom of the system. Einstein paper, which is entirely correct far from the critical point, unfortunately contains an improper application to the critical point where long range correlations are present. This paper was criticized later by Ornstein and Zernike (Onsager & Zernike, 1914) but provided an important source of inspiration for non-equilibrium.

Starting from what he calls the Boltzmann principle,

$$S = \kappa \ln W + \text{const}, \quad (1)$$

it is interesting to quote from (Einstein, 1910)

W is commonly equated with the number of different possible ways (complexions) in which the state considered - which is incompletely defined in the sense of a molecular theory, by parameters of a system - can conceivably be realized. To be able to calculate W, one needs a complete theory of the system under consideration. If considered from a phenomenological point of view equation (1) appears devoid of content.

However, Boltzmann's principle does acquire some content independent of any elementary theory if one assumes and generalizes from molecular kinetics the proposition that the irreversibility of physical processes is only apparent."

It follows from (1) that

$$W = \text{const} \cdot e^{\frac{N}{R}S}. \quad (2)$$

The order of magnitude of the constant is determined by taking into account that for the state of maximum entropy (entropy S_0) W is of the order of magnitude one, so that we then have, with order of magnitude accuracy,

$$W = e^{\frac{N}{R}(S-S_0)}. \quad (3)$$

From this we can conclude that the probability dW that the quantities $\lambda_1 \dots \lambda_k$ lie between λ_1 and $\lambda_1 + d\lambda_1 \dots \lambda_k$ and $\lambda_k + d\lambda_k$ is given, in order of magnitude, by the equation

$$dW = e^{\frac{N}{R}(S-S_0)} d\lambda_1 \dots d\lambda_k, \quad (4)$$


in the case when the system is determined only incompletely by $\lambda_1 \dots \lambda_k$.

In 1931 Onsager (Onsager, 1931), in the same vein as (Einstein, 1910) that he quoted, made use of Boltzmann formula in the study of fluctuations in non-equilibrium phenomena under the condition of small global deviations from equilibrium. The



theory was developed further by Onsager and Machlup in (Onsager & Machlup, 1953) where fluctuations of time trajectories of thermodynamic variables were considered under the same hypotheses. In the next section we discuss how it is possible to give a meaning to a formula like (3) in the more general situation of stationary states non necessarily close to equilibrium.

95 Typically we think of systems in contact with thermostats at different temperatures and/or reservoirs characterized by different chemical potentials and under the action of external fields. The MFT represents a step forward with respect to Onsager and Onsager–Machlup theory.

To study the fluctuations in states far from equilibrium let us analyze the meaning of the difference $S - S_0$ in (3). In an isolated system energy is conserved so that, if the volume remains constant, $S - S_0 = -\frac{F - F_0}{T}$ where F is the Helmholtz free energy. The quantity $F_0 - F$ represents the minimal work to bring the equilibrium state to the state corresponding to F at constant temperature and volume (Landau & Lifshitz, 1980). 

The concept of minimal work is meaningful also in non–equilibrium and can be taken as a generalization of the free energy. This is essentially the starting point of the macroscopic fluctuation theory. In the following section we shall present the basic ideas of the MFT stating explicitly the main assumptions, following mainly (Bertini et al., 2015).

105 3 Macroscopic Fluctuation Theory (MFT)

The MFT, as we mentioned, was inspired by non-equilibrium microscopic probabilistic models, the so called lattice gases, in particular the symmetric simple exclusion process (SSEP) for which the macroscopic dynamics is diffusive and can be proved rigorously. Also the probabilities of large deviations can be obtained and the rates computed.

3.1 Macroscopic fluctuations in stationary states

110 The macroscopic dynamics of diffusive systems is described by hydrodynamic equations often provided by conservation laws and constitutive equations, that is equations expressing the current in terms of the thermodynamic variables. More precisely on the basis of a local equilibrium assumption, at the macroscopic level the system is completely described by a local density $\rho(t, x)$ which may have several components and the local current $j(t, x)$. The evolution equations are of the form

$$\begin{cases} \partial_t \rho(t) + \nabla \cdot j(t) = 0, \\ j(t) = J(t, \rho(t)), \end{cases} \quad (5)$$

115 the space variable $x \in \Lambda \in R^d$.

For diffusive systems the constitutive equations take the form

$$J(\rho) = -D(\rho)\nabla\rho + \chi(\rho) E, \quad (6)$$

where the *diffusion coefficient* $D(\rho)$ and the *mobility* $\chi(\rho)$ are $d \times d$ symmetric and positive definite matrices, E is an external field acting on the **bulk**.



120 These equations have to be supplemented by appropriate boundary conditions on $\partial\Lambda$, the boundary of Λ , due to the interac-
 tion with the external reservoirs. If $\lambda(t, x)$, $x \in \partial\Lambda$, is the chemical potential of the external reservoirs, the boundary conditions
 read

$$f'(\rho(t, x)) = \lambda(t, x), \quad x \in \partial\Lambda. \quad (7)$$

125 where $f(\rho)$ is the **equilibrium free energy density**. Much of what we shall say applies to equations which are not in divergence
 form.

We assume that the microscopic evolution is given by a Markov process X_t which represents the configuration of the system
 at time t . The stationary non equilibrium state (SNS) is described by a stationary, i.e. invariant with respect to time shifts,
 probability distribution P_{st} over the trajectories of X_t .

130 The macroscopic equations are supposed to derive from an underlying microscopic dynamics through an appropriate scaling
 limit where the microscopic time is divided by a factor $\approx N^2$ and space is divided by $\approx N$ for N tending to ∞ where N is the
 number of degrees of freedom of the system.

The hydrodynamic equations represent laws of large numbers with respect the probability measure P_{st} conditioned on an
 initial state X_0 . The initial conditions are determined by X_0 . Of course many microscopic configurations give rise to the same
 value of $\rho_0(x)$. In general $\rho_t(x)$ is an appropriate limit as the number of degrees of freedom diverges.

135 Classically we should start from molecules interacting with realistic forces and evolving with Newtonian dynamics. This is
 beyond the reach of present day mathematical theory and much simpler models have to be adopted in the reasonable hope that
 some essential features are adequately captured.

140 We further assume that the stationary measure P_{st} admits a principle of large deviations describing the fluctuations from the
 typical hydrodynamic behavior. This means that the probability that the macroscopic variable ρ_t deviates from the solutions of
 the hydrodynamic equations and is close to some trajectory $\hat{\rho}_t$, is exponentially small and of the form

$$\begin{aligned} P_{st}(\rho(X_t) \approx \hat{\rho}_t(u), t \in [t_1, t_2]) &\approx e^{-\frac{1}{\epsilon^d} [V(\hat{\rho}_{t_1}) + \mathcal{J}(\hat{\rho})]} \\ &= e^{-\frac{1}{\epsilon^d} I(\hat{\rho})} \end{aligned} \quad (8)$$

145 where $\mathcal{J}(\hat{\rho})$ is a positive functional which vanishes if $\hat{\rho}_t = \bar{\rho}$ is a stationary solution of (5) and $V(\hat{\rho}_{t_1})$ is the cost to produce
 the initial value $\hat{\rho}_{t_1}$. The parameter ϵ is a scaling factor of the order of the ratio between the microscopic length scale (typical
 intermolecular distance) and the macroscopic one. The factor ϵ^{-d} is of the order of the number of particles in a macroscopic
 volume. The role of Avogadro's number in (3) is played here by ϵ^{-d} . We normalize $V(\hat{\rho})$ so that $V(\hat{\rho}) = 0$ when $\hat{\rho} = \bar{\rho}$ is the
 stationary solution. Therefore $\mathcal{J}(\hat{\rho})$ represents the extra cost necessary in order that the system follow the trajectory $\hat{\rho}_t$. Finally
 $\rho(X_t) \approx \hat{\rho}_t$ means closeness in some metric. V is the generalization to the infinite dimensional case of the Freidlin-Wentzell
quasi-potential (Freidlin & Wentzell, 2012) and is the large deviation rate of the stationary probability measure.



150 Let us denote by θ the time inversion operator defined by $\theta X_t = X_{-t} = X_t^*$. The probability measure P_{st}^* describing the evolution of the time reversed process X_t^* is given by the composition of P_{st} and θ^{-1} that is

$$P_{st}^*(X_t^* = \phi_t, t \in [t_1, t_2]) = P_{st}(X_t = \phi_{-t}, t \in [-t_2, -t_1]) \quad (9)$$

We assume that the time reversed process also admits a hydrodynamic description. This hypothesis is physically very reasonable: by acting on a system from the outside we can invert the currents flowing through the system and for example have heat
 155 passing from lower temperature to higher temperature.

At the level of large deviations (9) implies

$$V(\hat{\rho}_{t_1}) + \mathcal{J}_{[t_1, t_2]}^*(\hat{\rho}) = V(\hat{\rho}_{t_2}) + \mathcal{J}_{[-t_2, -t_1]}(\theta \hat{\rho}) \quad (10)$$

where $\hat{\rho}_{t_1}, \hat{\rho}_{t_2}$ are the initial and final points of the trajectory and $V(\hat{\rho}_{t_i})$ the free energies associated with the creation of the fluctuations $\hat{\rho}_{t_i}$ starting from the stationary non-equilibrium state (SNS). The functional \mathcal{J}^* vanishes on the solutions of the
 160 hydrodynamics associated to the time reversed process.

The physical situation we are considering is the following. The system is in the stationary state $\bar{\rho}$ at $t = -\infty$ but at $t = 0$ we find it in the state $\hat{\rho}_0$. We want to determine the most probable trajectory followed in the spontaneous creation of this fluctuation. According to (8) this trajectory is the one that minimizes \mathcal{J} among all trajectories connecting $\bar{\rho}$ to $\hat{\rho}_0$ in the time interval $[-\infty, 0]$. From (10) we have

$$165 \mathcal{J}_{[-\infty, 0]}(\hat{\rho}) = \mathcal{J}_{[0, \infty]}^*(\theta \hat{\rho}) + V(\hat{\rho}_0) \quad (11)$$

The right hand side is minimal if $\mathcal{J}_{[0, \infty]}^*(\theta \hat{\rho}) = 0$ that is if $\theta \hat{\rho}$ is a solution of the time reversed hydrodynamics. The existence of such a relaxation solution is due to the fact that the stationary solution $\bar{\rho}$ is attractive also for the time reversed hydrodynamics. We have therefore the following generalization of Onsager-Machlup to SNS

170 *"In a SNS the spontaneous emergence of a macroscopic fluctuation takes place most likely following a trajectory which is the time reversal of the relaxation path according to the time reversed hydrodynamics"*

From (10) or (11) we have that the free energy is related to J by

$$V(\rho) = \inf_{\hat{\rho}} \mathcal{J}_{[-\infty, 0]}(\hat{\rho}) \quad (12)$$

where the infimum is taken over all trajectories connecting $\bar{\rho}$ to ρ .

3.2 Density fluctuations

175 The functional \mathcal{J} for diffusive systems has the Freidlin-Wentzell form (Freidlin & Wentzell, 2012) generalized to the infinite dimensional situation.

$$\mathcal{J} = \frac{1}{2} \int dt \int dx (\partial_t \rho + \nabla J(\rho)) K(\rho)^{-1} (\partial_t \rho + \nabla J(\rho)) \quad (13)$$



where the kernel $K(\rho)$ is the elliptic operator defined on functions $\pi : \Lambda \rightarrow \mathbf{R}$ vanishing at the boundary $\partial\Lambda$ by

$$K(\rho)\pi = -\nabla \cdot (\chi(\rho)\nabla\pi). \quad (14)$$

180 This operator is the generalization of Onsager matrix L . Interpreting $\mathcal{L}(\rho, \partial_t \rho)$ as a Lagrangian there corresponds by Legendre duality the Hamiltonian

$$\mathcal{H} = \frac{1}{2} \langle \nabla H, \chi(\rho)\nabla H \rangle + \frac{1}{2} \langle H, \nabla \cdot (D(\rho)\nabla\rho) \rangle \quad (15)$$

where H is the momentum conjugate to ρ , that is $H = \frac{\delta \mathcal{L}}{\delta(\partial_t \rho)}$. The scalar product \langle, \rangle means integration over x .

The associated Hamilton equations are

$$185 \begin{cases} \partial_t \rho = \frac{\delta \mathcal{H}}{\delta H} = \frac{1}{2} \nabla \cdot (D(\rho)\nabla\rho) - \nabla \cdot (\chi(\rho)\nabla H) \\ \partial_t H = -\frac{\delta \mathcal{H}}{\delta \rho} = -\frac{1}{2} \sum_{1 \leq i, j \leq d} [\chi'_{i,j}(\rho) \partial_{x_i} H \partial_{x_j} H + D_{i,j}(\rho) \partial_{x_i} \partial_{x_j} H]. \end{cases} \quad (16)$$

These equations with appropriate boundary conditions are the variational equations to calculate the optimal trajectory creating ρ . They characterize the MFT. They are difficult to solve for generic dependence on ρ of the transport coefficients $D(\rho), \chi(\rho)$. The variational problem has been solved for constant D and quadratic χ and for the special model zero-range where V turns out to be local.

190 The quasi-potential or non-equilibrium free energy satisfies the associated Hamilton-Jacobi equation (HJ)

$$\langle \nabla \frac{\delta V}{\delta \rho} \cdot \chi(\rho)\nabla \frac{\delta V}{\delta \rho} \rangle - \langle \frac{\delta V}{\delta \rho} \nabla \cdot J(\rho) \rangle = 0. \quad (17)$$

As we shall see the expression of $V(\rho)$ for SSEP can be obtained by solving (17).

3.3 Current fluctuations

For lattice gases the following expression has been derived (Bertini et al., 2005, 2006) for the joint fluctuations of density and
 195 current

$$\mathcal{P}(\rho, j) \asymp \exp \left\{ -\frac{1}{\epsilon^d} \frac{1}{4} \int dt \int dx (j - J(\rho)) \cdot \chi(\rho)^{-1} (j - J(\rho)) \right\} \quad (18)$$

$$= \exp \left\{ -\frac{1}{\epsilon^d} \mathcal{I}_{[t_0, t_1]}(\rho, j) \right\},$$

where

$$\mathcal{I}_{[T_0, T_1]}(\rho, j) = \frac{1}{4} \int_{T_0}^{T_1} dt \int_{\Lambda} dx [j - J(\rho)] \cdot \chi(\rho)^{-1} [j - J(\rho)]. \quad (19)$$



200 and j is the actual value of the current, which is connected to ρ by the continuity equation $\partial_t \rho + \nabla \cdot j = 0$, while $J(\rho)$ is the hydrodynamic current for the given value of ρ . For a simple interpretation of the exponent think of an electric circuit. In this case χ^{-1} is the resistance and the double integral is the **energy dissipated by** $j(t) - J(\rho(t))$.

By minimizing first with respect to the current j it is possible to show that

$$V(\rho) = \inf_{\substack{\rho(t), j(t): \\ \nabla \cdot j = -\partial_t \rho \\ \rho(-\infty) = \bar{\rho}, \rho(0) = \rho}} \mathcal{I}_{[-\infty, 0]}(\rho, j), \quad (20)$$

205 where $\bar{\rho}$ is the stationary solution.

In a stationary state it is natural to consider the fluctuations of the local time averaged current

$$\tilde{j}(x) = \frac{1}{\tau} \int_0^\tau dt j(x, t) \quad (21)$$

For fluctuations of \tilde{j} the following large deviation principle has been derived

$$P(\tilde{j}) \asymp \exp -\epsilon^{-d} \tau \Phi(\tilde{j}) \quad (22)$$

210 where

$$\Phi(\tilde{j}) = \lim_{\tau \rightarrow \infty} \inf_{j \in \mathcal{A}_{\tau, j}} \frac{1}{\tau} \mathcal{I}_{[0, \tau]}(\rho, \tilde{j}) \quad (23)$$

The set $\mathcal{A}_{\tau, j}$ is the set of all currents j such that $\tau^{-1} \int_0^\tau dt j(t, x) = \tilde{j}(x)$.

This is a more general form of a large fluctuation principle proposed by Bodineau and Derrida (Bodineau & Derrida, 2004) and called *additivity principle*. Suppose we split a one-dimensional system in two segments of different length L_1 and L_2 .

215 In this case we must specify the boundary condition in the intermediate point, that is a value ρ of the density. The *additivity principle* takes the following form

$$P_{L_1+L_2}(j, \rho_a, \rho_b) \approx \max_{\rho} [P_{L_1}(j, \rho_a, \rho) P_{L_2}(j, \rho, \rho_b)] \quad (24)$$

where P_L is the probability corresponding to the length L and ρ_a, ρ_b are the boundary values of the density. This principle is correct for various models and equivalent to (23). However in this approach there is no time dependence and it is assumed
 220 that in the variational calculation is enough to consider minimizers that are independent of time while in (23) we admit time dependence. This is a non trivial difference because as it has been clarified a phase transition may be involved. In such a case the method of (Bodineau & Derrida, 2004) would underestimate the probability of a current fluctuation. Such a transition has been proved to exist in the model of Kipnis-Marchioro-Presutti (Kipnis et al., 1982) in equilibrium (Bertini et al., 2006; Bodineau & Derrida, 2005) and found numerically in (Hurtado & Garrido, 2011).



225 3.4 Phase transitions

In general the appearance of singularities in the large deviation rates denotes the presence of a non-equilibrium phase **transition**. Actually the variational principle of Bodineau and Derrida may provide several time independent solutions which in fact have



been found in models discussed in (Baek et al., 2018) representing different phases. There is another type of phase transition whose appearance is signaled by the non differentiability of the quasi-potential $V(\rho)$. This type of transition has been found in
 230 the weakly asymmetric exclusion process (WASEP), that is in presence of an external field, for sufficiently high values of the field (Bertini et al., 2010). The existence of non-equilibrium phase transitions, often impossible in equilibrium, is a generic feature which has to be taken in account when analysing e phenomenon.

3.5 Long range correlations

We are concerned with *macroscopic correlations* which are a generic feature of nonequilibrium non-linear models. Microscopic
 235 correlations which decay as a **summable power law** disappear at the macroscopic level.

We introduce the *pressure* functional as the Legendre transform of the quasi-potential V

$$G(h) = \sup_{\rho} \{ \langle h\rho \rangle - V(\rho) \}$$

By Legendre duality we have the change of variable formulae $h = \frac{\delta V}{\delta \rho}$, $\rho = \frac{\delta G}{\delta h}$, so that the Hamilton-Jacobi **equation** can then be rewritten in terms of G as

$$240 \quad \left\langle \nabla h \cdot \chi \left(\frac{\delta G}{\delta h} \right) \nabla h \right\rangle - \left\langle \nabla h \cdot D \left(\frac{\delta G}{\delta h} \right) \nabla \frac{\delta G}{\delta h} - \chi \left(\frac{\delta G}{\delta h} \right) E \right\rangle = 0 \quad (25)$$

where h vanishes at the boundary of Λ . As for equilibrium systems, G is the **generating functional of the correlation functions**.

We define

$$C_n(x_1, \dots, x_n) = \frac{\delta^n G}{\delta h(x_1) \dots \delta h(x_n)} \Big|_{h=0} \quad (26)$$

By expanding (25) around the stationary state one obtains, after non trivial manipulations and combinatorics, recursive equa-
 245 tions for the $C_n(x_1, \dots, x_n)$, see (Bertini et al., 2009). We discuss the pair correlation function by splitting it into the local equilibrium part and a possibly non-local term

$$C(x, y) = C_{\text{eq}}(x) \delta(x - y) + B(x, y) \quad (27)$$

where

$$C_{\text{eq}}(x) = D^{-1}(\bar{\rho}(x)) \chi(\bar{\rho}(x)) \quad (28)$$

250 and $\bar{\rho}$ is the stationary solution. We then obtain from the general equations the following equation for B

$$\mathcal{L}^\dagger B(x, y) = \alpha(x) \delta(x - y) \quad (29)$$

where \mathcal{L}^\dagger is the formal adjoint of the elliptic operator $\mathcal{L} = L_x + L_y$ given by, using the usual convention that repeated indices are summed,

$$L_x = D_{ij}(\bar{\rho}(x)) \partial_{x_i} \partial_{x_j} + \chi'_{ij}(\bar{\rho}(x)) E_j(x) \partial_{x_i} \quad (30)$$




255 and

$$\alpha(x) = \partial_{x_i} [\chi'_{ij}(\bar{\rho}(x)) D_{jk}^{-1}(\bar{\rho}(x)) \bar{J}_k(x)] \quad (31)$$

where $\bar{J} = J(\bar{\rho}) = -D(\bar{\rho}(x))\nabla\bar{\rho}(x) + \chi(\bar{\rho}(x))E(x)$ is the macroscopic current in the stationary state. In order that (29) may have a non trivial solution α must be different from zero which is generically the case when χ and D have a non-linear dependence on ρ . For non-trivial α long range correlations appear. In particular for the SSEP where $\chi(\rho) = \rho(1 - \rho)$

$$260 \quad B(x, y) = -\frac{1}{2}(\rho_1 - \rho_2)^2 \Delta^{-1}(x, y) \quad (32)$$

where Δ^{-1} is the Green function of the Dirichlet laplacian. 

In (Basile & Jona-Lasinio., 2004) it was shown that long range correlations may appear also in equilibrium in a reaction–diffusion dynamics when microscopic time reversal invariance is strongly violated.

4 The simple exclusion process (SSEP)

265 The simple exclusion process SSEP is the most studied system in far from equilibrium situations and is having a role similar to the Ising model in the study of phase transitions. The SSEP in one dimensional lattice is a process in which particles perform symmetric random walks subject to hard core exclusion. In non-equilibrium the boundary conditions at the two ends of the lattice are different, an external field may act on the system, or both, so that a current is flowing through the system.

Stationary states and their large deviations functions have been calculated exactly both microscopically and macroscopically
 270 via the MFT obtaining the same results. This happens also in the case of other models like the zero range process or the Kipnis-Marchioro-Presutti model.


4.1 Non-locality of the quasi-potential V in non-equilibrium stationary states

We consider the variational problem defining $V(\rho)$ for the one-dimensional simple exclusion process characterized by $D = 1$ and $\chi = \rho(1 - \rho)$. By performing the change of variable and inserting it in the HJ equation

$$275 \quad \frac{\delta V}{\delta \rho(x)} = \log \frac{\rho(x)}{1 - \rho(x)} - \phi(x; \rho) \quad (33)$$

for some functional $\phi(x; \rho)$ to be determined satisfying the boundary conditions $\phi(\pm 1) = \log \rho(\pm 1)/[1 - \rho(\pm 1)]$. We obtain a solution of the variational problem if we solve the following ordinary differential equation which relates the functional $\phi(x) = \phi(x; \rho)$ to ρ

$$\begin{cases} \frac{\Delta \phi(x)}{[\nabla \phi(x)]^2} + \frac{1}{1 + e^{\phi(x)}} = \rho(x) & x \in (-1, 1), \\ \phi(\pm 1) = \log \rho(\pm 1)/[1 - \rho(\pm 1)]. \end{cases} \quad (34)$$



The step initial condition with a density ρ_a at the left of the origin and ρ_b at the right of the origin

Figure 1. Step initial condition

280 This equation admits a unique monotone solution which is the relevant one for the quasi-potential. A computation shows that

$$V(\rho) = F(\rho) + \int_{\Lambda} dx \left\{ (1 - \rho)\phi + \log \left[\frac{\nabla \phi}{\nabla \bar{\rho}(1 + e^{\phi})} \right] \right\} \quad (35)$$

where $F(\rho)$ is the equilibrium free energy

$$F(\rho) = \int dx f(\rho(x)) = \int dx \{ \rho \log \rho + (1 - \rho) \log(1 - \rho) \} \quad (36)$$

This expression was first obtained by Derrida, Lebowitz and Speer solving the microscopic model (Derrida et al., 2002). They
 285 also proved that if one splits the system in two parts the rates obey an additivity rule more complicated than a simple sum and
 similar to the *additivity principle* of (Bodineau & Derrida, 2004). The above macroscopic calculation via the Hamilton-Jacobi
 equation was done in (Bertini et al., 2002).

If an external field is present the large deviation rate has been calculated by Derrida and Enaud (Enaud & Derrida, 2004).

4.2 Non-stationary states

290 The time evolution depends on the initial condition. Basic work on non stationary problems for SSEP is due to Derrida and
 Gerschenfeld (Derrida & Gerschenfeld, 2009a, b) with interesting developments in recent work by Mallick-Moriya-Sasamoto
 (Mallick et al., 2022). They studied the evolution of a step initial condition like in Fig.1

by considering the time integral of the local current through the origin $Q_t = \int_0^t j(0, t') dt'$ where t is the microscopic time.

We expect in a diffusive one-dimensional regime a law of large numbers for the quantity Q_t/\sqrt{t} for large t and a **large deviation**

295 **principle**

$$P\left(\frac{Q_t}{\sqrt{t}} = q\right) \asymp e^{-\sqrt{t}\phi(q)} \quad (37)$$

Define the cumulant generating function $\langle e^{\lambda Q} \rangle \simeq e^{\sqrt{t}\mu(\lambda)}$ where λ is a real parameter and \langle, \rangle now stands for stochastic
 average. The functions $\Phi(q)$ and $\mu(\lambda)$ are Legendre transform of each other. This problem was studied by Derrida and
 Gerschenfeld in 2009.



300 The variational equations are

$$\begin{aligned}\partial_t \rho &= \partial_x (\partial_x \rho - \chi(\rho) \partial_x H) \\ \partial_t H &= -\partial_x^2 H - \frac{1}{2} \chi'(\rho) \partial_x^2 H\end{aligned}\quad (38)$$

where $\chi(\rho)$ is the mobility. Notice that the equation for H is backward in time, the boundary conditions being $H(t, x) = \lambda \theta(x)$ and $H(0, x) = \lambda \theta(x) + f'(\rho(0, x)) - f'(\bar{\rho})$, $f(\rho)$ is the equilibrium free energy density.

In (Derrida & Gerschenfeld, 2009b) the authors calculated the moment generating function with quenched and fluctuating
 305 initial conditions, microscopically with the Bethe ansatz and macroscopically with the MFT. The variational equations could
 be solved only in special cases. Recently Mallick-Moriya-Sasamoto (Mallick et al., 2022) have discovered that the following
 non-local transformations, which generalize the Cole–Hopf transformation, allow to map the variational equations for the SSEP
 to the Ablowitz-Kaup-Newell-Segur (AKNS) equations

$$u(t, x) = \frac{1}{\chi'(\rho)} \partial_x \chi(\rho) \exp\left[-\int_{-\infty}^x dy \frac{\chi'(\rho)}{2} \partial_y H\right]\quad (39)$$

310

$$v(t, x) = -\frac{2}{\chi'(\rho)} \partial_x \exp\left[\int_{-\infty}^x dy \frac{\chi'(\rho)}{2} \partial_y H\right]\quad (40)$$

The AKNS equations

$$\partial_t u(t, x) = \partial_{xx} u(t, x) - 2u^2(t, x)v(x, t)\quad (41)$$

$$\partial_t v(t, x) = \partial_{xx} v(t, x) - 2v^2(t, x)u(x, t)\quad (42)$$

315 which were solved for the SSEP, that is $\chi = \rho(1 - \rho)$, with the inverse scattering method. These transformations are valid for
 quadratic χ . The boundary conditions for the step initial condition

$$\begin{aligned}u(0, x) &= \omega \delta(x) \\ v(t, x) &= \delta(x)\end{aligned}\quad (43)$$

From their solution they obtained the moment generating function

$$\mu(\lambda) = \frac{1}{\sqrt{\pi}} \sum_1^{\infty} \frac{(-1)^{n-1} \omega^n}{n^{3/2}}\quad (44)$$

320 where

$$\omega = (e^\lambda - 1)\rho_a(1 - \rho_b) + (e^{-\lambda} - 1)\rho_b(1 - \rho_a)\quad (45)$$

For integrability results in the Kipnis–Marchioro–Presutti model of heat transfer see the recent work by Bettelheim–Smith–
 Meerson (Bettelheim et al., 2022) and references therein.




5 Final remarks

325 The MFT shows that once macroscopic evolution equations like hydrodynamics are available and a separation of scales holds, a self-consistent macroscopic description of non-equilibrium phenomena can be obtained through a study of rare fluctuations. The origin of the probabilistic behaviour may be due to the influence of a smaller scale on a larger one or to chaotic properties of the underlying dynamical system.

The discovery that a purely macroscopic theory could reproduce in the case of SSEP the large deviation function for non-
330 equilibrium stationary states obtained from a microscopic calculation has been a source of inspiration for the MFT. The work of Derrida and Gerschenfeld extended the theory to time dependent evolutions and its correctness received an important mathematical support by the recent work of Mallick-Moriya-Sasamoto.

We have illustrated the theory in an idealized case: we have considered i) purely diffusive systems and ii) simplified stochastic models. However as the Ising model allowed us to understand a lot about phase transitions and the critical point, we believe
335 that the SSEP and other solvable models are providing a guide to what may happen out of equilibrium. Furthermore the MFT applies to all variants of SSEP or of other diffusive models that macroscopically have the same transport coefficients D, χ .

The general approach of MFT has been extended to systems with more than one conservation law (Bernardin., 2008) and to some reaction diffusion process like the Glauber-Kawasaki dynamics (Bodineau & Lagouge., 2010). It is reasonable to expect that it will be applicable to more general cases which may lead to a different structure of the large deviation rates. The
340 experience so far indicates that the phenomenon of long range space correlations is not limited to purely diffusive systems and is rather general in non-equilibrium. The same can be expected for non-equilibrium phase transitions.

In climate science the models are comparatively more complicated. The approach of the MFT can be presumably applied to circulation models whose equations have the structure of a reaction-diffusion system in which one can separate the conservation laws from the non-conservative part. A preliminary step should be the study of a simpler system exhibiting some essential
345 mathematical features. Quasi-geostrophic models may be a possibility. 

Appendix A: Fluctuating Hydrodynamics

Formally the equations of the MFT can be derived also from assuming an extension to non-equilibrium of the so called Fluctuating Hydrodynamics (FH). The idea of fluctuating hydrodynamics goes back to Landau (Landau & Lifshitz, 1980) who considered only linear hydrodynamic equations near equilibrium while the most interesting phenomena are generated by non
350 linearities far from equilibrium. It consists in adding to the macroscopic equations a noise term. In the case of hydrodynamic equations in divergence form we add to the current a fluctuating term

$$j = J(\rho) + \xi, \tag{A1}$$

where, conditionally on ρ , ξ is a Gaussian random term with variance

$$\langle \xi_i(t, x), \xi_j(t', x') \rangle = \epsilon^d \chi_{ij}(\rho) \delta(t - t') \delta(x - x') \tag{A2}$$



355 The hydrodynamic equation takes the form

$$\partial_t \rho + \nabla \cdot (J(\rho) + \xi) = 0 \quad (\text{A3})$$

However we have to emphasize that such equations, in the case of non-linear hydrodynamics, are very singular and need renormalization and so far there is no mathematical theory applicable even in one dimension.

From the previous equations we obtain

$$360 \quad P(\rho, j) \asymp \exp \left\{ -\frac{1}{\epsilon^d} \frac{1}{4} \int dt \int dx (j - J(\rho)) \cdot \chi(\rho)^{-1} (j - J(\rho)) \right\}$$

$$= \exp \left\{ -\frac{1}{\epsilon^d} \mathcal{I}_{[t_0, t_1]}(\rho, j) \right\},$$

the same as formula (18).

It would be interesting to provide a rigorous foundation to FH however this requires as a first step to give a clear mathematical meaning to the stochastic partial differential equations on which it is founded.

365 *Competing interests.* The author declares that he has no known conflict of interests

Acknowledgements. I wish to acknowledge my long standing collaboration on the topics of this paper with L. Bertini, A. De Sole, D. Gabrielli, C. Landim. I thank Vera Melinda Galfi for a critical reading of a preliminary version and for very useful comments.



References

- Bertini, L., A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, 2001, *Fluctuations in stationary nonequilibrium states of irreversible processes*, Phys. Rev. Lett. **87**, 040601.
- 370 Bertini, L., A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, 2002, *Macroscopic fluctuation theory for stationary non-equilibrium states*, J. Statist. Phys. **107**, 635–675.
- Bertini, L., A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, 2007, *Stochastic interacting particle systems out of equilibrium*, J. Stat. Mech. Theory Exp. P07014.
- 375 Derrida B., 2007, *Non-equilibrium steady states: fluctuations and large deviations of the density and of the current*, J. Stat. Mech. Theory Exp. P07023.
- Derrida B., 2011, *Microscopic versus macroscopic approaches to non-equilibrium systems*, J. Stat. Mech. Theory Exp. P01030.
- Bertini, L., A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, 2015, *Macroscopic Fluctuation Theory* Rev. Mod. Phys. **87**, 593–636.
- V. M. Galfi, V. Lucarini, J. Wouters, 2019 *A Large Deviation Theory-based Analysis of Heat Waves and Cold Spells in a Simplified Model of the General Circulation of the Atmosphere*, J. Stat. Mech. 033404, arXiv:1807.08261.
- 380 V. M. Galfi, V. Lucarini, 2021 *Fingerprinting heatwaves and cold spells and assessing their response to climate change using large deviation theory* Phys. Rev. Lett. bf 127, 058701.
- V.M. Galfi, V. Lucarini, F. Ragone, J. Wouters, 2021, *Applications of large deviation theory in geophysical fluid dynamics and climate science*, Riv. Nuovo Cim. **44**, 291363, arXiv:2106.13546.
- 385 Hasselmann K., 1976, *Stochastic climate models Part I. Theory*, Tellus, 28:6, 473–485.
- Mallick K., (2015) *The Exclusion Process: A paradigm for non-equilibrium behaviour*, Physica A **418** 17–48.
- Einstein A., 1910 *Theorie der Opaleszenz von homogenen Flüssigkeiten und Flüssigkeitsgemischen in der Nähe des kritischen Zustandes*. Annalen der Physik, **33**, 1275–1298, English translation in *The collected papers of Albert Einstein*, vol.3 p.231–249, Princeton University Press, 1993.
- 390 Landau L. D., E. M. Lifshitz, *Statistical Physics* third edition 1980, Ch. XII.
- Onsager, L., 1931, *Reciprocal relations in irreversible processes. I, II.*, Phys. Rev. **37**, 405–426; **38**, 2265–2279.
- Onsager L., Machlup, 1953 *Fluctuations and irreversible processes*, Phys. Rev. **91**, 1505–1512.
- Ornstein L. S., F. Zernike, 1914 *Accidental deviations of density and opalescence at the critical point of a single substance*, Proc. Acad. Sci. (Amsterdam) **17** 793–806.
- 395 Freidlin M., A. Wentzell, 2012 *Random Perturbations of Dynamical Systems* third edition, Springer.
- Bertini, L., A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, 2005, *Current fluctuations in stochastic lattice gases*, Phys. Rev. Lett. **94**, 030601.
- Bertini, L., A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, 2006, *Non-equilibrium current fluctuations in stochastic lattice gases*, J. Stat. Phys. **123**, 237–276.
- 400 Bodineau T., B. Derrida, 2004, *Current fluctuations in non-equilibrium diffusive systems: an additivity principle*, Phys. Rev. Lett. **92**, 180601.
- Bodineau T., B. Derrida, 2005, *Distribution of current in non-equilibrium diffusive systems and phase transitions*, Phys. Rev. E **72**, 066110.
- Kipnis C., C. Marchioro, E. Presutti, 1982, *Heat flow in an exactly solvable model*, J. Stat. Phys. **27**, 65.
- Hurtado P. I., P. L. Garrido, 2011, *Spontaneous symmetry breaking at the fluctuating level*, Phys. Rev. Lett. **107**, 180601.



- 405 Baek Y., Y. Kafri, V. Lecomte, 2018, *Dynamical phase transitions in the current distribution of driven diffusive channels*, J. Phys. A: Math. Theor. **51** 105001.
- Bertini, L., A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, 2010, *Lagrangian phase transitions in non-equilibrium thermodynamic systems*, J. Stat. Mech. Theory Exp., L11001.
- Bertini, L., A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, 2009, *Towards a non-equilibrium thermodynamics: a selfcontained macroscopic description of driven diffusive systems*, J. Stat. Phys. **135**, 857–872.
- 410 Basile G., G. Jona-Lasinio, 2004, *Equilibrium states with macroscopic correlations*, Int. J. Mod. Phys. B **18**, 479.
- Derrida B., J. L. Lebowitz, E. R. Speer, *Large deviation of the density profile in the steady state of the open symmetric exclusion process* 2002, J. Stat. Phys. **107**, 599-634.
- Enaud C., B. Derrida, (2004) *Large Deviation Functional of the Weakly Asymmetric Exclusion Process*, J. Stat. Phys. **114**, 537–562.
- Derrida B., A. Gerschenfeld, 2009a, *Current fluctuations in the one-dimensional symmetric simple exclusion process with step initial condi-*
415 *tion*, J. Stat. Phys. **136**, 1.
- Derrida B., A. Gerschenfeld, 2009b, *Current fluctuations in the one-dimensional diffusive systems with step initial density profile*, J. Stat. Phys. **137**, 978.
- Mallick K., H. Moriya, T. Sasamoto, 2022, *Exact solution of the macroscopic fluctuation theory for the symmetric exclusion process*, arXiv:2202.05213.
- 420 Bettelheim E., N. R. Smith, B. Meerson, 2022, *Full statistics of nonstationary heat transfer in the Kipnis–Marchioro–Presutti model*, arXiv:2204.06278.
- Bernardin C., 2008, *Stationary non-equilibrium properties for a heat conduction model*, Phys. Rev. E **78**, 021134.
- Bodineau T., M. Lagouge, 2010, *Current large deviations in a driven dissipative model*, J. Stat. Phys. **139**, 201.