

# Description and Evaluation of the JULES-ES setup for ISIMIP2b

Camilla Mathison<sup>1,2</sup>, Eleanor Burke<sup>1</sup>, Andrew Hartley<sup>1</sup>, Douglas I Kelley<sup>4</sup>, Chantelle Burton<sup>1</sup>, Eddy Robertson<sup>1</sup>, Nicola Gedney<sup>1</sup>, Karina Williams<sup>1,3</sup>, Andy Wiltshire<sup>1,3</sup>, Richard J. Ellis<sup>4</sup>, Alistair A. Sellar<sup>1</sup>, Chris D. Jones<sup>1</sup>

5 <sup>1</sup> Met Office Hadley Centre, FitzRoy Road, Exeter, UK

<sup>2</sup> School of Earth and Environment, Institute for Climate and Atmospheric Science, University of Leeds, Leeds, UK

<sup>3</sup> Global Systems Institute, University of Exeter, Laver Building, North Park Road, Exeter, UK

<sup>4</sup> UK Centre for Ecology and Hydrology, Wallingford, Oxfordshire, OX10 8BB, UK

*Correspondence to:* Andrew Hartley ([andrew.hartley@metoffice.gov.uk](mailto:andrew.hartley@metoffice.gov.uk))

10 **Abstract.** Global studies of climate change impacts that use future climate model projections also require projections of land surface changes. Simulated land surface performance in Earth System models is often affected by the atmospheric models' climate biases, leading to errors in land surface projections. Here we run the [Joint UK Land Environment Simulator Earth System configuration](#) (JULES-ES) land surface model with [Inter Sectoral Impacts Model Intercomparison project 2<sup>nd</sup> phase future projections](#) (ISIMIP2b) bias-corrected climate model data from 4 global climate models (GCMs). The bias correction  
15 reduces the impact of the climate biases present in individual models. We evaluate JULES-ES performance against present-day observations to demonstrate its usefulness for providing required information for impacts such as fire and river flow. ~~We simulate a historical and two future scenarios; a mitigation scenario RCP2.6 and RCP6.0, which has very little mitigation.~~ We include a standard JULES-ES configuration without fire as a contribution to ISIMIP2b and JULES-ES with fire as a potential future development. Simulations for gross primary productivity (GPP), evapotranspiration (ET) and albedo compare well  
20 against observations. Including fire improves the simulations, especially for ET and albedo and vegetation distribution, with some degradation in shrub cover and river flow. This configuration represents some of the most current earth system science for land surface modelling. The suite associated with this configuration provides a basis for past and future phases of ISIMIP, providing a simulation setup, postprocessing and initial evaluation using [the International Land Model Benchmarking \(ILAMB\) project](#). This suite ensures that it is as straightforward, reproducible and transparent as possible to follow the  
25 protocols and participate fully in ISIMIP using JULES.

## 1 Introduction

The Joint UK Land Environment Simulator (JULES) ([Clark et al., 2011; Best et al., 2011](#))(Clark et al., 2011; Best et al., 2011) is a community-supported and developed land surface model used by land, hydrological, weather and climate communities. JULES is a configurable code base supporting weather, climate and earth system science applications. Here, we describe and  
30 evaluate the JULES Earth System (JULES-ES) configuration and experimental setup used in the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP; [Frieler et al., 2017](#))Frieler et al., 2017). JULES-ES builds on the JULES-GL7 configuration

described in (Wiltshire et al., 2020) Wiltshire et al. (2020) by including ~~additional processes necessary for representing impacts and terrestrial~~ biogeochemical ~~fluxes governing carbon and nitrogen cycles that influence earth system processes; these are considered more relevant to ecosystems and people, which could be affected by climate change processes.~~ Whilst we run  
35 JULES-ES in offline mode, it is also coupled to the atmosphere within the earth system model UKESM (Sellar et al., 2019)(Sellar et al., 2019). Climate change impacts are already a feature of everyday life for much of the world, and quantifying these allows us to understand future benefits and trade-offs of climate mitigation and adaptation policies. ISIMIP provides a consistent framework for assessing impacts using a large ensemble of impacts models across various sectors (Warszawski et al., 2013, 2014)(Warszawski et al., 2013, 2014). ISIMIP has recently completed its second phase, having more than 60  
40 modelling groups contributing simulations to ISIMIP2a (reanalysis-driven hindcasts) and ISIMIP2b (bias-corrected GCM-driven historical and future scenarios). We present the JULES-ES configuration and experimental setup that has contributed to ISIMIP2b and will be the basis for further development in subsequent ISIMIP phases. An advantage of using JULES-ES as an offline impacts model is that it ~~uses a prescribed atmosphere, without these feedbacks JULES-ES is more~~is computationally efficient ~~than compared to~~ the closely aligned land surface scheme ~~coupled to the atmosphere used~~ in UKESM1 (Sellar et al., 2019)(Sellar et al., 2019), ~~and, while~~ using multi-model climate ensembles sample scientific uncertainty in ~~our understanding of the climate system and land surface forcing~~ that would not be possible within a single climate model framework.

This paper briefly describes the changes to JULES-GL7 (Wiltshire et al., 2020)(Wiltshire et al., 2020) that form the JULES-ES configuration, the ISIMIP setup and an evaluation of the arising simulations. JULES-ES has been widely evaluated and applied for global biogeochemical modelling (Sellar et al., 2019; Slevin et al., 2017)(Sellar et al., 2019; Slevin et al., 2017),  
50 including in the Global Carbon Budget (Friedlingstein et al., 2020)(Friedlingstein et al., 2020). Here we focus on using JULES for impacts applications. Alongside this manuscript, we provide a suite to run JULES-ES following the ISIMIP2b modelling protocol (Frieler et al., 2017)(Frieler et al., 2017) for tailored impact projections that are consistent across sectors such as water and biomes. The suite includes the code to post-process output into ISIMIP formatted netcdf output and run the International Land Model Benchmarking (ILAMB) system to allow for quick evaluation (see Text S1). Data from the JULES-ES ISIMIP2b  
55 suite have been submitted to the biomes and water ISIMIP2b sectors and are available via the ISIMIP model archive (<https://data.isimip.org/>) and provide simulations of the historical and future land surface in the (United Nations Environment Programme, 2022)United Nations Environment Programme (2022) wildfire report. The JULES ISIMIP2b simulations with fire provide the basis for the contribution of JULES to the next FireMIP, which will use the ISIMIP3 set-up. The historical simulations and their evaluation are shown in section 3, with discussion and conclusions in Sections 4 and 5 respectively.

## 2 Materials and Methods

### 2.1 JULES-ES Configuration

To better represent variation in plant traits and managed land, we extend the standard representation of 5 Plant Functional Types (PFTs) [to 13](#) in JULES-ES ~~to 13~~, building on [\(Harper et al., 2016\)](#) Harper et al. (2016), with 4 managed and 9 natural PFTs. Natural PFTs are extended by splitting trees into deciduous and evergreen types and then distinguishing between temperate and tropical broadleaf evergreen trees. These additional PFTs represent a wider range of leaf life spans and metabolic capacities. Evergreen trees typically have less access to nutrients, higher leaf mass per unit area, longer lifespans and ~~lower~~ carbon assimilation and respiration rates, whereas a deciduous PFT typically has leaves with a higher nutrient concentration, shorter lifespan and lower leaf mass per unit area. Tropical broadleaf evergreen trees have lower maximum carbon assimilation rates than temperate trees. The 9 natural PFTs used are: tropical broadleaf evergreen trees (BET-Tr), temperate broadleaf evergreen trees (BET-Te), broadleaf deciduous trees (BDT), needle-leaf evergreen trees (NET), needle-leaf deciduous trees (NDT), ~~C3C4~~ grasses [\(C3G\)](#), C4 grasses [\(C4G\)](#), evergreen shrubs (ESh), and deciduous shrubs (DSH). [\(Harper et al., 2016\)](#) Harper et al. (2016) also updated several parameters required for calculating photosynthesis and respiration using the TRY database [\(Kattge et al., 2011\)](#) (Kattge et al., 2011). They also reduced the bias in model simulations by tuning parameters relating to leaf dark respiration, canopy radiation, canopy nitrogen, stomatal conductance, root depth, and temperature sensitivities of the maximum carboxylation rate of Rubisco ( $V_{cmax}$ ) based on available observations. The 4 managed PFTs are C3 and C4 crop and pasture [\(C3Cr, C4Cr, C3Pa, C4Pa\)](#) PFTs and are functionally similar to natural grasses but, in the case of crops are assumed not to be nitrogen limited and litter carbon is removed as a simple representation of crop harvest [\(Sellar et al., 2019; Robertson, 2019\)](#). [This simple fertilization of crops through lack of nitrogen limitation does not distinction between mineral and manure fertilizer.](#) Both crop and pasture surface types undergo land-use change according to externally forced time-varying land use. Within the respective crop/pasture fraction, only the C3 and C4 crop/pasture PFTs are allowed to grow, with the area of each determined by the TRIFFID dynamic vegetation module [\(Burton et al., 2019\)](#) (Burton et al., 2019).

Outside of the managed land area, ~~only~~ the nine natural PFTs (including natural ~~C3GC3~~ and ~~C4GC4~~ grasses) can grow in the remainder of the grid box once the non-vegetated surfaces have been accounted for (urban, ice, lakes, [represented by corresponding non-vegetation surface types, and ocean which is not simulated](#)). ~~As the net~~. As the prescribed crop or pasture fraction increases with land-use change, natural vegetation is removed from the portion of the grid box into which agriculture has expanded, representing anthropogenic land clearance. Conversely, when crop and pasture areas are reduced, the natural PFTs are allowed to recolonize the vacated grid box fraction. [The vacated gridbox fraction is initially bare soil and the existing natural PFTs gradually expand their coverage into this area, the rate of expansion is determined by the TRIFFID vegetation dynamics scheme and will follow a succession of faster growing grass PFTs, followed by shrubs and then trees.](#) Bare soil occupies any remaining space once the vegetation dynamics have been simulated. Simple representations of fertilisation and harvesting are applied to the crop PFTs, but otherwise these are physiologically identical to the natural grasses. After

accounting for land-use, the fractional coverage and biomass of each PFT within a grid box is determined by the TRIFFID dynamic vegetation model. Inter-PFT competition is based on vegetation height, with the taller vegetation shading and therefore dominating other PFTs ([Harper et al., 2018](#))(Harper et al., 2018).

The other major change introduced in JULES-ES is a representation of nitrogen and nutrient limitation effects on ecosystem carbon assimilation. The nitrogen component of JULES is described in ([Wiltshire et al., 2021](#))Wiltshire et al. (2021). In brief, JULES-ES represents all the key terrestrial N processes. Inputs to the land surface are via biological fixation, fertilization and nitrogen deposition, with losses from the land surface occurring via leaching and gas loss, with Nitrogen deposition being externally provided to the model. JULES simulates a nitrogen-limited ecosystem by reducing the ~~carbon use efficiency~~~~net primary productivity~~ if there is insufficient available N to satisfy plant N demand. ~~This results in a reduced carbon assimilation under nutrient support. Any excess carbon is added to the plant respiration.~~ The soil biogeochemistry is based on the representation of the four-pool RothC soil carbon ([Clark et al., 2011](#))~~model~~(Clark et al., 2011) consisting of decomposable plant material (DPM), resistant plant material (RPM), microbial biomass (BIO), and humus (HUM). For each soil carbon pool there is an equivalent soil nitrogen pool ([Wiltshire et al., 2021](#))(Wiltshire et al., 2021). Nitrogen transfers between the organic and inorganic nitrogen pools depend on decomposition rates and the C to N ratio of the organic pool.

Another important change is the inclusion of a fire module. Fire is simulated in JULES by the fire model INFERNO (INteractive Fire and Emission algoRithm for Natural enviroNments; ([Mangeon et al., 2016](#))Mangeon et al. (2016)). Burned area is calculated from flammability and ignitions. Temperature, saturation vapour pressure, relative humidity, precipitation, together with soil moisture and fuel load from JULES give flammability by PFT, prescribed population density ([Goldewijk et al., 2017](#))~~from HYDE3.2~~(Goldewijk et al., 2017) gives human ignitions, and prescribed climatological lightning ~~from LIS/OTD version 2.3.2015~~(Cecil, 2006) gives natural ignitions. Previous evaluation of JULES-INFERNO, which compared model output against observational constraints, showed that burnt area is relatively insensitive to changes in lightning ([Mangeon et al., 2016; Burton et al., 2022](#)). Here we use INFERNO coupled to the dynamic vegetation model TRIFFID ([Burton et al., 2019](#))(Burton et al., 2019), enabling carbon cycle feedbacks from fire onto the land surface via vegetation mortality, regrowth, and burnt litter fluxes. Recent updates to INFERNO allow fire mortality to vary by PFT, and updates to the representation of land-use and PFTs in JULES allows for reduced burning in ~~C3CrC3~~ and ~~C4CrC4 crop~~ PFTs ([Burton et al., 2020](#))(Burton et al., 2020), based on global trends of agricultural fire suppression ([Bistinas et al., 2014; Andela et al., 2017](#)). Background mortality rates were reduced compared to the model without fire to account for the extra vegetation mortality from fire, as per Burton et al. (2019). We use fire and background mortality in (Burton et al., 2020). C3Pa and C4Pa burning remain as per Burton et al. (2019), which reflects observational constraints that show an increase in burnt area in pasture areas ([Kelley et al., 2019; Bistinas et al., 2014](#))(Bistinas et al., 2014; Andela et al., 2017).

## 2.2 Modifications for ISIMIP2b

In the ISIMIP JULES-ES configuration the TRIFFID period, [this is a number \(in days\) that defines the frequency that the TRIFFID dynamic vegetation code is called](#), has been reduced from a 10-day to a 1-day period in order to allow a shorter restart periods necessary to meet the diagnostic requirements of ISIMIP (a large number of variables on short temporal scales). A daily TRIFFID period also allows the vegetation dynamics to respond more realistically to variations in plant productivity on shorter timescales, although the effect of this change is minimal in test historical runs.

Another key difference to the standard setup of JULES is the use of daily meteorological driving data. JULES needs a model timestep of no more than 1 hour to accurately simulate the diurnal cycle and exchange of heat, water and momentum and avoid numerical instabilities. In the ISIMIP experimental setup, we use the internal disaggregator [\(Williams and Clark, 2014\)](#) [\(Williams and Clark, 2014\)](#) to calculate driving data values at the model timestep of 1 hour, ~~based on the method used by the IMOGEN model~~ [disaggregator \(Huntingford et al., 2010\)](#) [\(Huntingford et al., 2010\)](#) ~~to initially disaggregates to 3-hourly, which the model linearly interpolates to 1 hour.~~ The diurnal cycle of downward shortwave radiation is calculated from the position of the sun in the sky. Temperature is calculated from a sinusoidal function with a maximum 0.15 of a daylength after local noon, normalised by the diurnal temperature range. Downward longwave radiation is a linear function of temperature, and specific humidity is kept below saturation at each timestep. Precipitation is considered to occur in a single event, with a globally specified ‘duration’ parameter (6h for convective rainfall, 1h for large-scale rainfall, convective snowfall and large-scale snowfall). [Convective occurs when temperature exceeds 288.15°K. The model assumes convective rainfall is more intense and so leads to more runoff and less infiltration into the soil. Given that precipitation events do](#) ~~Given that this event does~~ not, by construction, overlap with midnight GMT, on average, this produces a spurious trapezoidal diurnal cycle, which is zero at midnight GMT [\(Williams and Clark, 2014\)](#) [\(Williams and Clark, 2014\)](#). Precipitation above 350 mm/day is redistributed. Note that convective precipitation occurs only on a fraction of the grid box [\(Best et al., 2011\)](#) [\(Best et al., 2011\)](#), set to 30% in the ISMIP2b runs, and within this fraction is modelled as a negative exponential distribution [\(Johannes Dolman and Gregory, 1992\)](#) [\(Johannes Dolman and Gregory, 1992\)](#). Therefore, the grid box average intensity is not the same as the effective intensity at a point. Given the strong effect of intensity on canopy interception and runoff, the water cycle in the model is sensitive to the duration parameter choices [\(Williams and Clark, 2014\)](#) [\(Williams and Clark, 2014\)](#). See Figure S1 for plots showing that using the disaggregator has little effect on vegetation that we would expect to be influenced by rainfall.

## 2.3 ISIMIP2b protocol

The ISIMIP2b experiments focus on understanding different levels of mitigation. They ~~use two RCP scenarios to explore~~ [are consistent with the international commitment](#) made under the Paris Agreement to stabilise global warming at well below 2°C, relative to pre-industrial mean temperatures. ISIMIP2b uses simulations from the Coupled Model Inter-

comparison Project 5 (CMIP5), using an historical scenario (1860-~~2005~~~~2006~~) and the RCP2.6 and RCP6.0 ~~future~~ concentration pathways (~~post 2005~~~~2006~~ ~~2099~~) to represent a higher ambition, lower temperature outcome and a low ambition pathway respectively ([Riahi et al., 2017](#))(Riahi et al., 2017). Land-use data and population density are based on the Shared Socioeconomic Pathway (SSP2) scenario and applied to RCP2.6 and RCP6.0 simulations. [Lightning ignitions for fire are from the LIS/OTD version 2.3.2015 climatology \(Cecil, 2006\), and applied for the historical and future as per Rabin et al. \(2017\) and Hantson et al. \(2020\).](#) To capture a range of climate sensitivities, four CMIP5 GCM driving models are chosen: GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR and ~~MIOCMiROC5~~. The GCM driving data is global bias-corrected daily data at 0.5° resolution ([Hempel et al., 2013; Frieler et al., 2017; Lange, 2018](#))(Hempel et al. (2013),Lange (2018)~~and~~ Frieler et al. (2017). [Bias-correction means we can compare the observed and simulated impacts during the historical reference period with a smooth transition into the future period.](#) The bias-correction methodology adjusts multi-year monthly [mean distribution throughout the historical and RCP periods based on comparison of GCM output against EWEMBI reanalysis data distributions \(Dee et al., 2011\).](#) ~~mean distribution~~ for the period 1979 to 2013, using CMIP5 RCP8.5 post-~~2005~~~~2006~~ (the end of CMIP5 “historic” period), such that trends and inter-annual variability are preserved in absolute and relative terms for temperature and non-negative variables respectively ([Lange, 2018](#))Lange (2018). [EWEMBI combines data from multiple reanalysis source to cover all required variables. The variables bias-corrected for ISIMIP2b are listed in Table 1 reproduced from Frieler et al. \(2017\).](#) ~~Transfer functions are used to adjust the distributions of daily anomalies from monthly mean values.~~ The ISIMIP2b bias correction includes humidity as well as shortwave and longwave radiation using quantile mapping. [Transfer functions are used to adjust the distributions of daily anomalies from monthly mean values.](#) ISIMIP2b bias-correction methods adjust distributions independently for each variable, grid cell and month, preserving the statistical dependencies between variables, in space and changes over time. The bias correction approach preserves the trends (and therefore sensitivities) from different GCMs but removes absolute biases [found](#) over the reference period [from the historical and RCP periods](#). Each GCM therefore has a different variability and simulated climate outside of the reference period, [but with a smooth transiting going from historical into the reference period, or from the reference period into the future.](#) Some small biases remain after bias-correction, particularly in precipitation (Figure S2), where biases are a small fraction of total local rainfall but can affect precipitation particularly in the South America (Figure S3). As part of the setup provided here, we include code for preparing JULES data for submitting to ISIMIP and ensuring it conforms to the strict protocols (see Text S1) and the ILAMB system for rapid evaluation of the simulations (see Text S2).

[Table Bias corrected variables for ISIMIP2b simulations, reproduced from Table 1 of Frieler et al. \(2017\)](#)

## 2.4 Model Evaluation

We evaluate the model for key impacts sectors, and use the International Land Model Benchmarking (ILAMB) tool ([Collier et al., 2018](#))(Collier et al., 2018) to assess model performance for GPP, ET, runoff and albedo. ILAMB evaluates performance

against observations from remote sensing, reanalysis data and fluxnet site measurements and produces graphical and statistical scores of model results. ~~We compare model GPP against the upscaled fluxnet product from Jung et al. (2011), ET against GLEAM (Miralles et al., 2011) and MODIS (Mu et al., 2011) estimates, runoff against Dai and Trenberth (2002), albedo against the GEWEX SRB radiation observations (Stackhouse et al., 2011) and burnt area against Global Fire Emission Database version 4 with small fires (GFED4s; Van Der Werf et al. (2017))~~ The model values and the observation datasets we use for the evaluation are given in Table 1. As ILAMB doesn't include vegetation cover evaluation, we also include the Manhattan Metric (Kelley et al., 2013)(Kelley et al., 2013) comparison ~~with~~against from ESA CCI Land Cover tree, shrub, wood and grass cover (Harper et al., 2022)(Harper et al., 2022). Results and further details of the ILAMB and vegetation cover analysis are provided in Text S2. We evaluate the historical simulations separately for each GCM because the bias correction preserves inter-annual differences between GCMs. We conduct evaluation over common time periods between observations and simulation using the ~~historical~~historie and, for observational periods beyond the end of 2006, RCP6.0.

## Table 2 the observations used to evaluate the JULES-ES fields

Click or tap here to enter text.

## 2.5 The experimental suite

To run JULES we collect all the tasks and input files that are needed together into what is known as a suite, this allow runs to be reproduced using the exact same applications, options and commands as previously run possibly by another user, scheduling them to run based on the dependencies between the tasks. Full instructions on how to run JULES in a Rose suite are provided on github (). We provide a full setup for running the ISIMIP2b simulations using JULES-ES in the form of the suite u-cc669 available via the Met Office Science Repository Service (MOSRS - <https://code.metoffice.gov.uk/trac/roses-u> see data availability section for information). The bias corrected driving data is available from ISIMIP at <https://www.isimip.org/gettingstarted/input-data-bias-adjustment/>. We also use the following datasets from ISIMIP. Where preprocessing of these is required to use this data within JULES, this preprocessing code is also part of the suite.

- CO<sub>2</sub> concentration
- Future land-use patterns
- Nitrogen deposition
- Land-sea mask

~~Total Runoff Integrating Pathways (TRIP) is used for~~ river routing model allows JULES to collect and route water through river channels essentially converting runoff to river discharge or river flow. ~~A in JULES, therefore a~~ TRIP 0.5° river routing ancillary is also required ~~for these runs, which is~~and available from <http://hydro.iis.u-tokyo.ac.jp/~taikan/TRIPDATA/>.




3 Model evaluation

3.1 Water

225 We evaluated the water cycle using runoff derived from [\(Dai, 2021\)](#)Dai and Trenberth (2002) for the 50 largest river catchments. Estimated observed mean basin runoff combines river flow measurements at downstream gauge stations with a river flow model to estimate the flow at the river mouth. By assuming there are no losses from the river, we calculated the long-term mean, basin averaged runoff by dividing the river flow at the river mouth by the basin area.

230 **Particularly in temperate regions north and south of the equator, simulations using all 4 sets of driving data show similar biases. However, differences between the driving datasets are greatest in tropical or sub-tropical river catchments (**



**Figure 1**Figure 1). This is particularly evident in the Amazon basin, where mean runoff biases range from approximately 0 in the simulation driven by HADGEM2-ES, to more than -0.6mm day<sup>-1</sup> in the simulation driven by IPSL-CM5A-LR. Strong variations between the simulations are also seen in the Brahmaputra basin [with some smaller in the Chang Jiang](#) (Figure 2). The general underestimate of runoff in the higher latitudes may be due to the treatment of moisture infiltration into partially frozen soils (see below) but could also be caused by biases in precipitation estimates due to the gauge “under-catch” of snow [\(Adam and Lettenmaier, 2003\)](#)(Adam and Lettenmaier, 2003). In arid and semi-arid basins river flow and runoff tends to be over-estimated, which could be due to [missing processes such as](#) anthropogenic water extraction, [primarily irrigation](#) (Richey



240 [et al., 2015](#)) and river channel evaporation and transmission losses ([Haddeland et al., 2011](#); [Döll and Siebert, 2002](#))—which disproportionately affects groundwater depletion ([Richey et al., 2015](#)).

Figure 2 compares the long-term monthly mean river flow (1980-2014 inclusive where observations are available) over the six largest rivers to those of downstream observations in ([Dai, 2021](#)) Dai (2021). All the simulations reproduce the overall seasonal river flow over the Amazon. After the wet season, the modelled river flows decline earlier than observed and the simulated river flows at "low water" are too low. This could be due to too much evaporation in the drier months (see [ET for JJA](#) in Figure S4) and/or the simulated speed of flow through the soil and/or river channel being too fast. Though GFDL-ESM2M and IPSL-CM5A-LR driving precipitation data also has a dry bias during the dry season (Figure S3). All simulations over-estimate river flow over the Congo, mainly due to over-estimates in the rainy months. This could be driven by too little evaporation from the vegetation canopy or from flooded areas ([See ET in Figure 3 and Figure S4](#)). The simulations reproduce the seasonal river flow over the Orinoco well. The timings of peak river flow for the Chiangjiang and Brahmaputra are well simulated although the amplitudes are too low. Dams, which we do not model, are likely to affect the observed seasonal cycle. The simulated river flow over the Mississippi lags observations by several months. This lag is also evident over many high latitude basins (not shown). The observed river flow peak is mainly driven by Spring melt, whereas the simulated river flow peak is in line with that of precipitation. This is due to this configuration allowing significant soil infiltration of snow melt when the soil surface is mainly frozen, rather than resulting in surface runoff. This also may result in the underestimate of annual river flow because once the water has infiltrated the soil it may then be evaporated.

### 3.2 Surface fluxes

260 Global Gross Primary Productivity (GPP) is 134-137 PgC/yr, depending on driving data, which is above the estimate [of IPCC AR6](#) ([Canadell et al., 2021](#)) (Canadell et al., 2021) of 113 PgC yr<sup>-1</sup>, but agrees well with the estimates of  $146 \pm 21$  PgC/yr of ([Cheng et al., 2017](#)) Cheng et al. (2017; Table S1). Net Biome Productivity (NBP) is 0.94-1.46 PgC/yr between 2011-2020, within the 1.0-2.8 PgC/yr range estimated by the Global Carbon Budget ([Friedlingstein et al., 2022](#)) (Friedlingstein et al., 2022). All simulations show positive GPP biases in similar regions, such as central and southern Africa, south of the Himalaya and east towards Bangladesh and Myanmar compared to ([Jung et al., 2011](#)) Jung et al. (2011) observations (Figure 3). South America is a more complicated picture with Brazil broadly split between negative GPP to the northeast and positive GPP to the country's southeast. For Brazil, the ET bias has a more northwest – southeast split, with the northwest having a slightly negative bias and the southeast more positive. The northwest bias in ET and the bias in GPP in South America is more prominent and wide-spread in early wet season (September-November) when driven by climate data from GFDL-ESM2M and IPSL-CM5A-LR (see Figure [S5S4](#)) and is due to a longer dry season in both sets of driving data (Figure S3), with rains starting in October rather than September. The far north of Columbia, Bolivia and Argentina also have a negative bias in both GPP

and ET across the simulations. Australia also has a north-south split with a slight positive ET bias to the north and the inverse to the south.

275 Albedo (Figure 3, right column) generally shows a positive bias across most regions and simulations. However, there are small regions with a negative bias, for example, south of the Sahel and small regions at higher northern latitudes. Eastern Siberia has a positive bias in all simulations (see Discussion).

### 3.3 Biomes

280 All simulations show similar vegetation cover patterns that largely follow observations, capturing high tree cover fractions in boreal and tropical forests, grass cover in tropical, temperate and boreal grasslands and bare ground in arid regions (Figure 4). There are, however, some biases common to all simulations. Tree cover fraction is too high globally, with a simulated range of 4.97-5.31 Mkm<sup>2</sup>, which is higher than observations and depends~~depending~~ on driving data (Table S2). ~~In the observations,~~ shrub and grasses dominate eastern Siberian Taiga in the model instead of observed high tree cover (Figure 4). ~~There is also~~ 285 slightly too much shrub cover in tropical forests at the expense of tree cover, contributing to a global bias in shrub cover of 1.22-1.31 Mkm<sup>2</sup>. Conversely, simulated tropical tree cover is too high in savanna regions, giving the impression of more continuous and less fragmented forests across the tropics (Figure 4). Boundaries between temperate and warm temperate woodlands/forests and tropical forests are too sharp, suggesting JULES-ES does not capture processes in temperate woodland transition. Savanna and grasslands tend to be too narrow, with more bare soil in the models in semi-arid regions such as 290 southern Africa, the Mojave desert and the Sahel.

### 3.4 Fire

In addition to the simulations without fire submitted to the archive, we performed additional simulations with fire and fire feedbacks switched on. These fire simulations provide burnt area and alter vegetation cover, carbon, fluxes, albedo and runoff 295 (Burton et al., 2019). Burnt area is similar across all four simulations (Figure 5). ~~The model simulates present-day burnt area well compared to satellite observations, with the global total burnt area average for 2000-2020 observed by MODIS CCI v5.1 as 4.55 Mkm<sup>2</sup>, and the model simulating between 3.94-4.43 Mkm<sup>2</sup> depending on driving data (Table S3). The model captures the high burnt area in southern hemisphere Africa - a common area of low bias in global fire models (Hantson et al., 2020)(Hantson et al., 2020). The model also performs better than other FireMIP models at simulating the high burnt areas in~~ 300 northern hemisphere Africa (Hantson et al., 2020), though fire is still lower than in observations. This is partly due to very low simulated burnt areas in Nigeria's Guinean savanna, which (Kelley et al., 2019)Kelley et al. (2019) show is a consequence of show is likely inevitable in global parameterisations of population density and agricultural drivers of burnt area. Other regional biases include too high burnt area in South America and too low in Eurasia and northern Australia (Figure 5).

305 [We also show to low burnt area in Australia in a common problem across fire models \(Hantson et al., 2020\), even in model optimized to burnt area observations \(Kelley et al., 2019; Bistinas et al., 2014\). This may be due to the unique fire ecology in Northern Australia \(Kelley, 2014\) and to high uncertainty in observations of burnt area \(Giglio et al., 2010\). High burning in South America occurs in areas where cropland fragmentation reduces burnt area beyond the extent of agricultural areas \(Kelley et al., 2019; Andela et al., 2017\), which is hard to reproduce in tile and PFT-based models \(Hantson et al., 2016\). We also](#)  
310 [simulate too low burnt area in Eurasia. Some of this observed burnt area at these high latitudes comes from peatland fires, which, like most other fire models \(Rabin et al., 2017\), are not simulated in INFERNO. Observational burnt area products tend to underestimate levels of burning in forests \(Randerson et al., 2012\), which may explain the slight bias towards to high simulated burnt area in tropical forests.](#)

315 Fire is simulated well in savanna bands (15 degrees north and south), which improves the representation of tree cover by reducing the positive bias compared to observations (Figure 6; Table S2). The inclusion of interactive fire has the overall effect of decreasing simulated global tree cover (from 38.66-39, depending on driving data to 34.32-35.65 Mkm<sup>2</sup>) to more in line with observations (34.86 Mkm<sup>2</sup>) and increasing grasses and bare soil. In the tropics, this tends to bring the modelled “forest” (areas dominated by trees) more in line with the observations, with high simulated tree cover restricted to observed  
320 [forested areas. However, including fire does reduce tree cover in Savanna areas \(Figure ~~S8S5~~\), particularly in Africa, which means the spatial distribution of trees in this region compared with observations is not as good as without fire \(Table S2\). There is an increase and improvement in tree cover in some high latitude North America due to changes in background mortality in the with fire simulation.](#)~~diminishes performance in spatial distribution (Table S2).~~ Fire reduces shrub cover to well below observations (Table S2), though given the well documented issues in distinguishing tall and short woody vegetation  
325 [\(Gerard et al., 2017; Adzhar et al., 2022\)](#)(Gerard et al., 2017), it is probably more meaningful to assess total woody cover. Here, including fire reduces model bias (from 6.19-6.59 to -3.60- -2.25 Mkm<sup>2</sup>) and improves the spatial pattern (Table S2). Including fire reduces the global bias of high GPP [when compared against estimated by the Global Carbon Budget \(Friedlingstein et al., 2020\).](#), bringing global total down by ~2 PgC/yr across all simulations (Table S1). However, fire does slightly degrade ~~GPP’sGPPs~~ spatial pattern (Table S1). Including fire also reduced global NBP (Table S1) by 0.12-0.38 PgC/yr,  
330 depending on driving data.

Fire alleviates some of the high bias in ET, improving the ~~model’smodels~~<sup>2</sup> overall performance (Table S4). In semi-arid areas we already over-estimate river flow (in part probably due to human extraction). The addition of fire lowers ET, thereby increasing river flow bias in semi-arid regions, which slightly degrades overall runoff performance (Figure ~~S9S6~~). Fire also  
335 improves spatial pattern of albedo (Table S5), though seasonal performance decreases. This is in line with well-documented biases in fire seasonal cycles across all global fire models, [which tend to have longer than observed fire seasons in tropical savannas, and, in human dominated fire regimes, season timing shifted and often late compared to observations \(Hantson et](#)

al., 2016, 2020). Previous evaluations of JULES configurations incorporating INFERNO also show these biases (Burton et al., 2019, 2022, submitted; Hantson et al., 2020) (Hantson et al., 2020, 2016), ~~including in previous JULES configurations incorporating INFERNO~~ (Burton et al., 2019, 2022).

## 4 Discussion

We have presented simulations of the JULES-ES land surface model, run according to the ISIMIP2b protocols using bias-corrected climate model data from 4 GCMs for the historical period. The configuration will be used to perform simulations under 2 future scenarios (RCP2.6 and RCP6.0). The JULES-ES ISIMIP2b configuration simulates the surface fluxes (GPP, ET, albedo) reasonably well. Including fire improves the ET and albedo, but not the GPP which is biased high. Including fire in the simulations currently degrades runoff (Table S6).

The configurations of JULES can capture the ~~seasonal~~ annual cycle of many of the largest rivers, although high latitude rivers and managed rivers, ~~although high latitude rivers and managed rivers~~ are generally not captured as well. Including irrigation and structural hydrological developments, such as dams and reservoirs, would likely improve the simulations of managed rivers. Previously, (Falloon et al., 2011) Falloon et al. (2011) found that GCM precipitation biases contribute to errors in TRIP river flows for some basins in both HadGEM1 and HadCM3. In this study, we use bias-corrected data which reduces these errors, meaning that differences in JULES-ES results between the driving models are due to differences in inter-seasonal or inter-annual variability between driving models (e.g. Figure S3). However, errors in evapotranspiration, runoff generation or other missing processes e.g., snow accumulation and snow melt processes, could also contribute. Uncertainty in precipitation due to the sparsity of observation networks and the under-catch of solid precipitation for high latitude (Falloon et al., 2011) (Falloon et al., 2011) and altitude rivers e.g., in the Himalaya (Mathison et al., 2015) (Mathison et al., 2015), means that it is difficult to interpret model performance in these basins.

Some basins show the same bias direction in runoff (




Figure 1 (Figure 1) and ET (Figure 3), notably that the Amazon is too dry for both variables and the Nile too wet, whereas we would expect opposing biases if they were from land surface simulation. In these basins, there is a dry and wet bias (respectively) in the driving precipitation data (Figure S2). HadGEM2-ES and IPSL-CM5A-LR have particularly dry driving data in the Amazon, and [this](#) results in the driest runoff and ET in the simulation. This translates to biases in GPP (Figure 3) and vegetation cover (Figure 4). So, while ISIMIP bias-correction reduces climate model biases compared to those in an Earth System Model (see evaluation in [\(Sellar et al., 2019\)](#) Sellar et al. (2019)) or when run with a [non-bias](#)-bias corrected climate [\(Burton et al., 2022\)](#) (Burton et al., 2022), they are not eliminated.

Land cover is an important factor for the surface fluxes. Grassy regions for example, correlate with the regions of positive ET bias. The annual mean global GPP biases are small, but this is not the case for GPP on a seasonal timescale and over a smaller region such as South America. The albedo and land cover area bias are also closely related. For example, [if JULES simulates a larger number of trees than observed, this may lead to a higher or lower albedo than observed or if JULES simulates a lower number of trees this may lead to a higher albedo than observed. Conversely, higher grass cover just North of the observed Sahel corresponds to simulated low-albedo biases. Vegetation impacts on albedo are. This is](#) particularly important at high latitudes where there is snow cover, for example the positive albedo bias in Eastern Siberia is because JULES simulates too few trees and too much grass there. [Grasses are more readily buried below snow than trees making these areas more reflective](#)

(Sellar et al., 2019) ~~This positive bias in grass cover, affects the snow cover in these regions,~~ which in turn affects the albedo.

380 JULES represents the bending and partial burying of vegetation by snow (Ménard et al., 2014) (Ménard et al., 2014), however the settings controlling this interaction described in (Sellar et al., 2019) Sellar et al. (2019) have been tuned for the coupled UKESM1 model rather than the standalone JULES model. Eastern Siberia is a vulnerable region which has experienced increased climate-related impacts, including heatwaves (Ciavarella et al., 2021) (Ciavarella et al., 2021) and fires (Kelley et al., 2019) (Kelley et al., 2019); it is very likely that climate change will exacerbate these feedbacks by the end of the 21st century  
385 (United Nations Environment Programme, 2022) (United Nations Environment Programme, 2022). Developments by Mercado et al. (2018) (Mercado et al., 2018) which improve the representation of plant acclimation to thermal stress may improve spatial variations across different vegetation types in JULES.

In general, the simulations with fire improve the vegetation and productivity distribution, particularly the distribution of  
390 tropical forests, boundary between forests and savannas, and North America. Developing JULES to include ~~fire~~ these processes will improve simulations for these areas and properly capture the climate impacts on vegetation cover and carbon fluxes. ~~there.~~ The results show that there are too few trees compared to observations for western parts of Brazil. The simulations with fire on improve tree cover in savanna, which is consistent with the findings of (Staver et al., 2011; Lasslop et al., 2016) Lasslop et al. (2016) and Staver et al. (2011), however there is still ongoing discussion around how much impact fire really has on tree  
395 cover in the savanna compared to other dry disturbances such as wind throw, heat stress, rainfall distribution (Veenendaal et al., 2018; Brovkin et al., 2009) (Veenendaal et al., 2018; Brovkin et al., 2009).

## 5 Conclusions

We have presented a configuration of JULES-ES set up to run and generate output following the ISIMIP protocols. We provide  
400 a suite for running the simulations that includes driving data, ancillaries, postprocessing and first look evaluation (ILAMB) for any phase of ISIMIP. Outputs using this set up were submitted to the biomes and water ISIMIP2b sectors, and our evaluation helps inform any difference between JULES-ES and other models participating in ISIMIP2b. The suite also provides a starting point for further JULES-ES developments. We ~~evaluate~~ include a ~~fire~~ set up ~~for running~~ with ~~INFERNO to~~  
~~include~~ representation of fire using the INFERNO fire model in anticipation of ISIMIP3 which will include a Fire Sector. We  
405 show that including fire has an impact on model results and is important to include in simulations of climate impacts. While fire mostly improves model performance, it does degrade certain vegetation distributions, including lack of larch forest, and runoff. However, fire has a substantial impact on both ecosystem composition and hydrological processes and should therefore still be included when studying impacts under changing climate and environmental conditions. Therefore, while documentation of the configuration without fire will be useful for anyone using previously submitted results, we recommend using the  
410 configuration with fire in future JULES-ES development. Future work using this configuration and new phases of ISIMIP will



focus on using the full benefit and extent of the ISIMIP ensemble to enable more in-depth exploration of climate impacts together with the quantification of earth system uncertainties, co-benefits of mitigation and adaptation to climate change.

### Data and code availability

415 The JULES-ES for ISIMIP configuration (based on JULES version 5.5) is preserved at  
<https://code.metoffice.gov.uk/trac/roses-u/browser/b/k/8/8/6> (fire off) and <https://code.metoffice.gov.uk/trac/roses-u/browser/c/f/1/3/7> (fire on). JULES and associated configurations are freely available for non-commercial research use as set out in the JULES user terms and conditions ([http://jules-lsm.github.io/access\\_req/JULES\\_Licence.pdf](http://jules-lsm.github.io/access_req/JULES_Licence.pdf)). For a comprehensive guide on how to access, install and run the configurations used in this research, we direct the reader to Appendix A of (Wiltshire  
420 [et al., 2020](https://gmd.copernicus.org/articles/13/483/2020/#section6)) Wiltshire et al. (2020) available at <https://gmd.copernicus.org/articles/13/483/2020/#section6>  
The data and code used for the evaluation of the JULES-ES outputs with iLAMB in the study are available at  
<https://www.ilamb.org/datasets.html> and <https://github.com/rubisco-sfa/iLAMB> with a BSD 3-clause “New” license  
(<https://github.com/rubisco-sfa/iLAMB/blob/master/LICENSE.rst>)  
The JULES model data output used in the model evaluation in the study are available at <https://data.isimip.org/>, using the  
425 search tag ‘jules-es-55’ <https://data.isimip.org/search/query/jules-es-55/> with a Creative Commons Attribution 4.0  
International license (<https://creativecommons.org/licenses/by/4.0/>)

© British Crown Copyright 2022, the Met Office. All rights reserved. The software is provided by the Met Office to the Topical  
Editor at Geoscientific Model Development under the Software Licence for Peer Review (Use, duplication or disclosure of  
430 [this code is subject to the restrictions as set forth in the aforementioned Software Licence for Peer Review](#)). The software is  
provided to facilitate the peer review of submitted manuscript ‘Description and Evaluation of the JULES-ES setup for  
ISIMIP2b’ by Mathison et al. and should be used and distributed to Authorised Persons for this purpose only. The software is  
extracted from the UM and JULES trunks at the revisions of the MOSRS repositories corresponding to the stated version  
[having passed both science and code reviews according to the UM and JULES working practices.](#)

435

### References

[Adam, J. C. and Lettenmaier, D. P.: Adjustment of global gridded precipitation for systematic bias, Journal of Geophysical Research: Atmospheres, 108, 4257, <https://doi.org/10.1029/2002JD002499>, 2003.](#)  
[Adzhar, R., Kelley, D. I., Dong, N., George, C., Raventos, M. T., Veenendaal, E., Feldpausch, T. R., Phillips, O. L., Lewis, S.](#)  
440 [L., Sonké, B., Taedoumg, H., Marimon, B. S., Domingues, T., Arroyo, L., Djangbletey, G., Saiz, G., and Gerard, F.: MODIS](#)  
[Vegetation Continuous Fields tree cover needs calibrating in tropical savannas, 2022.](#)

- Andela, N., Morton, D. C., Giglio, L., Chen, Y., van der Werf, G. R., Kasibhatla, P. S., DeFries, R. S., Collatz G J and Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., and Yue C and Randerson, J. T.: A human-driven decline in global burned area, *Science* (1979), 356, 1356–1362, 2017.
- 445 Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. . L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes, *Geosci Model Dev*, 4, 677–699, <https://doi.org/10.5194/gmd-4-677-2011>, 2011.
- 450 Bistinas, I., Harrison, S. P., Prentice, I. C., and Pereira, J. M. C.: Causal relationships versus emergent patterns in the global controls of fire frequency, *Biogeosciences*, 11, 5087–5101, 2014.
- Brovkin, V., Raddatz, T., Reick, C. H., Claussen, M., and Gayler, V.: Global biogeophysical interactions between forest and climate, *Geophys Res Lett*, 36, <https://doi.org/10.1029/2009GL037543>, 2009.
- Burton, C., Betts, R., Cardoso, M., Feldpausch, R. T., Harper, A., Jones, C. D., Kelley, D. I., Robertson, E., and Wiltshire, A.: Representation of fire, land-use change and vegetation dynamics in the Joint UK Land Environment Simulator vn4.9 (JULES), *Geosci Model Dev*, 12, 179–193, <https://doi.org/10.5194/GMD-12-179-2019>, 2019a.
- 455 Burton, C., Betts, R., Cardoso, M., Feldpausch, T. R., Harper, A., Jones, C. D., Kelley, D. I., Robertson, E., and Wiltshire, A.: Representation of fire, land-use change and vegetation dynamics in the Joint UK Land Environment Simulator vn4.9 (JULES), *Geosci Model Dev*, 12, 179–193, 2019b.
- Burton, C., Betts, R. A., Jones, C. D., Feldpausch, T. R., Cardoso, M., and Anderson, L. O.: El Niño Driven Changes in Global Fire 2015/16, *Front Earth Sci (Lausanne)*, 8, 199, <https://doi.org/10.3389/FEART.2020.00199/BIBTEX>, 2020.
- 460 Burton, C., Kelley, D. I., Jones, C. D., Betts, R. A., Cardoso, M., and Anderson, L.: South American fires and their impacts on ecosystems increase with continued emissions, *Climate Resilience and Sustainability*, 1, e8, <https://doi.org/https://doi.org/10.1002/cli2.8>, 2022.
- Burton, C., Kelley, D. I., Burke, E., Mathison, C., Jones, C. D., Betts, R., Robertson, E., Teixeira, J. C. M., Cardoso, M., and 465 Anderson, L. O.: Fire weakens land carbon sinks before 1.5°C, n.d.
- Canadell, J. G., Monteiro, P. M. S., Costa, M. H., Cotrim da Cunha, L., Cox, P. M., Eliseev, A. V., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P. K., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global Carbon and other Biogeochemical Cycles and Feedbacks, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: V., M.-D., 470 Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 673–816, <https://doi.org/https://doi.org/10.1017/9781009157896.007>, 2021.
- Cecil, D. J.: LIS/OTD 0.5 Degree High Resolution Monthly Climatology (HRMC) (No. [1995-2014]), 2006.

475 [Cheng, L., Zhang, L., Wang, Y. P., Canadell, J. G., Chiew, F. H. S., Beringer, J., Li, L., Miralles, D. G., Piao, S., and Zhang, Y.: Recent increases in terrestrial carbon uptake at little cost to the water cycle, \*Nature Communications\* 2017 8:1, 8, 1–10, <https://doi.org/10.1038/s41467-017-00114-5>, 2017.](#)

[Chuvieco, E., Lizundia-Loiola, J., Lucrecia Pettinari, M., Ramo, R., Padilla, M., Tansey, K., Mouillot, F., Laurent, P., Storm, T., Heil, A., and Plummer, S.: Generation and analysis of a new global burned area product based on MODIS 250 m reflectance](#)

480 [bands and thermal anomalies, \*Earth Syst Sci Data\*, 10, 2015–2031, <https://doi.org/10.5194/essd-10-2015-2018>, 2018.](#)

[Ciavarella, A., Cotterill, D., Stott, P., Kew, S., Philip, S., van Oldenborgh, G. J., Skålevåg, A., Lorenz, P., Robin, Y., Otto, F., Hauser, M., Seneviratne, S. I., Lehner, F., and Zolina, O.: Prolonged Siberian heat of 2020 almost impossible without human influence, \*Clim Change\*, 166, 1–18, <https://doi.org/10.1007/S10584-021-03052-W/TABLES/3>, 2021.](#)

[Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H.,](#)

485 [Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator \(JULES\), model description – Part 2: Carbon fluxes and vegetation dynamics, \*Geosci Model Dev\*, 4, 701–722, <https://doi.org/10.5194/gmd-4-701-2011>, 2011.](#)

[Collier, N., Hoffman, F. M., Lawrence, D. M., Keppel-Aleks, G., Koven, C. D., Riley, W. J., Mu, M., and Randerson, J. T.: The International Land Model Benchmarking \(ILAMB\) System: Design, Theory, and Implementation, \*J Adv Model Earth\*](#)

490 [Syst, 10, 2731–2754, <https://doi.org/10.1029/2018MS001354>, 2018.](#)

[Dai, A.: Hydroclimatic trends during 1950–2018 over global land, \*Clim Dyn\*, 56, 4027–4049, <https://doi.org/10.1007/S00382-021-05684-1/FIGURES/14>, 2021.](#)

[Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, d P., and others: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, \*Quarterly\*](#)

495 [Journal of the royal meteorological society, 137, 553–597, 2011.](#)

[Döll, P. and Siebert, S.: Global modeling of irrigation water requirements, \*Water Resour Res\*, 38, 1–8, 2002.](#)

[Falloon, P., Betts, R., Wiltshire, A., Dankers, R., Mathison, C., Mcneall, D., Bates, P., and Trigg, M.: Validation of River Flows in HadGEM1 and HadCM3 with the TRIP River Flow Model, \*J Hydrometeorol\*, 12, 1157–1180, <https://doi.org/10.1175/2011JHM1388.1>, 2011.](#)

500 [Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzier, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I.,](#)

505 [Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S. I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Van Der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan,](#)

W., Yue, X., and Zaehle, S.: Global Carbon Budget 2020, *Earth Syst Sci Data*, 12, 3269–3340, <https://doi.org/10.5194/ESSD-12-3269-2020>, 2020.

Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Goldewijk, K. K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S. I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., Van Der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global Carbon Budget 2021, *Earth Syst Sci Data*, 14, 1917–2005, <https://doi.org/10.5194/ESSD-14-1917-2022>, 2022.

Frieler, K., Lange, S., Piontek, F., Reyer, C. P. O., Schewe, J., Warszawski, L., Zhao, F., Chini, L., Denvil, S., Emanuel Kerry and Geiger, T., Halladay, K., Hurtt, G., Mengel, M., Murakami, D., Ostberg, S., Popp, A., Riva, R., Stevanovic, M., Suzuki, T., Volkholz, J., Burke, E., Ciais, P., Ebi, K., Eddy, T. D., Elliott Joshua and Galbraith, E., Gosling, S. N., Hattermann, F., Hickler, T., Hinkel, J., Hof, C., Huber, V., Jägermeyr, J., Krysanova, V., Marcé, R., Schmied, H. M., Mouratiadou, I., Pierson, D., Tittensor, D. P., Vautard Robert and van Vliet, M., Biber, M. F., Betts Richard A and Bodirsky, B. L., Deryng, D., Froliking, S., Jones, C. D., Lotze, H. K., Lotze-Campen, H., Sahajpal, R., Thonicke, K., Tian, H., and Yamagata, Y.: Assessing the impacts of 1.5 °C global warming – simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b), 2017.

Gerard, F., Hooftman, D., van Langevelde, F., Veenendaal, E., White, S. M., and Lloyd, J.: MODIS VCF should not be used to detect discontinuities in tree cover due to binning bias. A comment on Hanan et al. (2014) and Staver and Hansen (2015). *Global Ecology and Biogeography*, 26, 854–859, <https://doi.org/10.1111/GEB.12592>, 2017.

Giglio, L., Randerson, J. T., der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries, R. S.: Assessing variability and long-term trends in burned area by merging multiple satellite fire products, *Biogeosciences*, 7, 1171–1186, 2010.

Goldewijk, K. K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates for the Holocene - HYDE 3.2, *Earth Syst Sci Data*, 9, 927–953, <https://doi.org/10.5194/ESSD-9-927-2017>, 2017.

Haddeland, I., Clark, D. B., Franssen, W., Ludwig, F., Voß, F., Arnell, N. W., Bertrand, N., Best, M., Folwell, S., Gerten, D., and others: Multimodel estimate of the global terrestrial water balance: setup and first results, *J Hydrometeorol*, 12, 869–884, 2011.

Hantson, S., Arneth, A., Harrison, S. P., Kelley, D. I., Colin Prentice, I., Rabin, S. S., Archibald, S., Mouillot, F., Arnold, S. R., Artaxo, P., Bachelet, D., Ciais, P., Forrest, M., Friedlingstein, P., Hickler, T., Kaplan, J. O., Kloster, S., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Meyn, A., Sitch, S., Spessa, A., Van Der Werf, G. R., Voulgarakis, A., and Yue, C.: The status and challenge of global fire modelling, *Biogeosciences*, 13, 3359–3375, <https://doi.org/10.5194/BG-13-3359-2016>, 2016.

Hantson, S., Kelley, D. I., Arneth, A., Harrison, S. P., Archibald, S., Bachelet, D., Forrest, M., Hickler, T., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Nieradzik Lars and Rabin, S. S., Colin Prentice, I., Sheehan, T., Sitch, S., Teckentrup, L., Voulgarakis, A., and Yue, C.: Quantitative assessment of fire and vegetation properties in historical simulations with fire-enabled vegetation models from the Fire Model Intercomparison Project, *Geoscientific Model Development Discussions*, 2020.

Harper, A. B., Cox, P. M., Friedlingstein, P., Wiltshire, A. J., Jones, C. D., Sitch, S., Mercado, L. M., Groenendijk, M., Robertson, E., Kattge, J., Bönisch, G., Atkin, O. K., Bahn, M., Cornelissen, J., Niinemets, Ü., Onipchenko, V., Peñuelas, J., Poorter, L., Reich, P. B., Soudzilovskaia, N. A., and Bodegom, P. van: Improved representation of plant functional types and physiology in the Joint UK Land Environment Simulator (JULES v4.2) using plant trait information, *Geosci Model Dev*, 9, 2415–2440, <https://doi.org/10.5194/gmd-9-2415-2016>, 2016.

Harper, A. B., Wiltshire, A. J., Cox, P. M., Friedlingstein, P., Jones, C. D., Mercado, L. M., Sitch, S., Williams, K., and Duran-Rojas, C.: Vegetation distribution and terrestrial carbon cycle in a carbon cycle configuration of JULES4.6 with new plant functional types, *Geosci Model Dev*, 11, 2857–2873, <https://doi.org/10.5194/GMD-11-2857-2018>, 2018.

Harper, K. L., Lamarche, C., Hartley, A., Peylin, P., Ottlé, C., Bastrikov, V., San Martín, R., Bohnenstengel, S. I., Kirches, G., Boettcher, M., Shevchuk, R., Brockmann, C., and Defourny, P.: A 29-year time series of annual 300-metre resolution plant functional type maps for climate models, *Earth System Science Data Discussions*, Submitted, 2022.

Hempel, S., Frieler, K., Warszawski, L., Schewe, J., and Piontek, F.: A trend-preserving bias correction: The ISI-MIP approach, *Earth System Dynamics*, 4, 219–236, <https://doi.org/10.5194/ESD-4-219-2013>, 2013.

Huntingford, C., Booth, B. B. B., Sitch, S., Gedney, N., Lowe, J. A., Liddicoat, S. K., Mercado, L. M., Best, M. J., Weedon, G. P., Fisher, R. A., Lomas, M. R., Good, P., Zelazowski, P., Everitt, A. C., Spessa, A. C., and Jones, C. D.: IMOGEN: An intermediate complexity model to evaluate terrestrial impacts of a changing climate, *Geosci Model Dev*, 3, 679–687, <https://doi.org/10.5194/GMD-3-679-2010>, 2010.

Johannes Dolman, A. and Gregory, D.: The Parametrization of Rainfall Interception In GCMs, *Quarterly Journal of the Royal Meteorological Society*, 118, 455–467, <https://doi.org/10.1002/QJ.49711850504>, 1992.

Jung, M., Reichstein, M., Margolis, H. a., Cescatti, A., Richardson, A. D., Arain, M. A., Arneth, A., Bernhofer, C., Bonal, D., Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari, F., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations, *J Geophys Res*, 116, G00J07, <https://doi.org/10.1029/2010JG001566>, 2011.

Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Bönisch, G., Garnier, E., Westoby, M., Reich, P. B., Wright, I. J.,  
 Cornelissen, J. H. C., Violle, C., Harrison, S. P., Van BODEGOM, P. M., Reichstein, M., Enquist, B. J., Soudzilovskaia, N.  
 a., Ackerly, D. D., Anand, M., Atkin, O., Bahn, M., Baker, T. R., Baldocchi, D., Bekker, R., Blanco, C. C., Blonder, B., Bond,  
 W. J., Bradstock, R., Bunker, D. E., Casanoves, F., Cavender-Bares, J., Chambers, J. Q., Chapin Iii, F. S., Chave, J., Coomes,  
 580 D., Cornwell, W. K., Craine, J. M., Dobrin, B. H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M., Fagan, W. F., Fang,  
 J., Fernández-Méndez, F., Fidelis, a., Finegan, B., Flores, O., Ford, H., Frank, D., Freschet, G. T., Fyllas, N. M., Gallagher, R.  
 V., Green, W. a., Gutierrez, a. G., Hickler, T., Higgins, S. I., Hodgson, J. G., Jalili, a., Jansen, S., Joly, C. a., Kerkhoff, a. J.,  
 Kirkup, D., Kitajima, K., Kleyer, M., Klotz, S., Knops, J. M. H., Kramer, K., Kühn, I., Kurokawa, H., Laughlin, D., Lee, T.  
 D., Leishman, M., Lens, F., Lenz, T., Lewis, S. L., Lloyd, J., Llusià, J., Louault, F., Ma, S., Mahecha, M. D., Manning, P.,  
 585 Massad, T., Medlyn, B. E., Messier, J., Moles, a. T., Müller, S. C., Nadrowski, K., Naeem, S., Niinemets, Ü., Nöller, S.,  
 Nüske, a., Ogaya, R., Oleksyn, J., Onipchenko, V. G., Onoda, Y., Ordoñez, J., Overbeck, G., et al.: TRY - a global database  
 of plant traits, *Glob Chang Biol*, 17, 2905–2935, <https://doi.org/10.1111/j.1365-2486.2011.02451.x>, 2011.  
 Kelley, D. I.: *Modelling Australian fire regimes*, Macquarie University, 2014.  
 Kelley, D. I., Prentice, I. C., Harrison, S. P., Wang, H., Simard, M., Fisher, J. B., and Willis, K. O.: A comprehensive  
 590 benchmarking system for evaluating global vegetation models, *Biogeosciences*, 10, 3313–3340, <https://doi.org/10.5194/BG-10-3313-2013>, 2013.  
 Kelley, D. I., Bistinas, I., Whitley, R., Burton, C., Marthews, T. R., and Dong, N.: How contemporary bioclimatic and human  
 controls change global fire regimes, *Nature Climate Change* 2019 9:9, 9, 690–696, <https://doi.org/10.1038/s41558-019-0540-7>, 2019.  
 595 Lange, S.: Bias correction of surface downwelling longwave and shortwave radiation for the EWEMBI dataset, *Earth System  
 Dynamics*, 9, 627–645, 2018.  
 Lasslop, G., Brovkin, V., Reick, C. H., Bathiany, S., and Kloster, S.: Multiple stable states of tree cover in a global land surface  
 model due to a fire-vegetation feedback, *Geophys Res Lett*, 43, 6324–6331, <https://doi.org/10.1002/2016GL069365>, 2016.  
 Mangeon, S., Voulgarakis, A., Gilham, R., Harper, A., Sitch, S., and Folberth, G.: INFERNO: A fire and emissions scheme  
 600 for the UK Met Office’s Unified Model, *Geosci Model Dev*, 9, 2685–2700, <https://doi.org/10.5194/GMD-9-2685-2016>, 2016.  
 Mathison, C., Wiltshire, A. J., Falloon, P., and Challinor, A. J.: South Asia river-flow projections and their implications for  
 water resources, *Hydrol Earth Syst Sci*, 19, 4783–4810, <https://doi.org/10.5194/HESS-19-4783-2015>, 2015.  
 Ménard, C. B., Essery, R., Pomeroy, J., Marsh, P., and Clark, D. B.: A shrub bending model to calculate the albedo of shrub-  
 tundra, *Hydrol Process*, 28, 341–351, <https://doi.org/10.1002/HYP.9582>, 2014.  
 605 Mercado, L. M., Medlyn, B. E., Huntingford, C., Oliver, R. J., Clark, D. B., Sitch, S., Zelazowski, P., Kattge, J., Harper, A.  
 B., and Cox, P. M.: Large sensitivity in land carbon storage due to geographical and temporal variation in the thermal response  
 of photosynthetic capacity, *New Phytologist*, 218, 1462–1477, <https://doi.org/10.1111/NPH.15100>, 2018.



610 Miralles, D. G., De Jeu, R. A. M., Gash, J. H., Holmes, T. R. H., and Dolman, A. J.: Magnitude and variability of land evaporation and its components at the global scale, *Hydrol Earth Syst Sci*, 15, 967–981, [https://doi.org/10.5194/HESS-15-967-](https://doi.org/10.5194/HESS-15-967-2011) 2011, 2011.

Mu, Q., Zhao, M., and Running, S. W.: Improvements to a MODIS global terrestrial evapotranspiration algorithm, *Remote Sens Environ*, 115, 1781–1800, <https://doi.org/10.1016/j.rse.2011.02.019>, 2011.

Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Li Fang and Mangeon, S., Yue, C., Arora, V. K., and Others: The Fire Modeling Intercomparison Project (FireMIP), phase 1: Experimental and analytical protocols, *Geosci Model Dev*, 20, 1175–1197, 2017.

615 Randerson, J. T., Chen, Y., Van Der Werf, G. R., Rogers, B. M., and Morton, D. C.: Global burned area and biomass burning emissions from small fires, *J Geophys Res Biogeosci*, 117, 2012.

Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environmental Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.

620 Richey, A. S., Thomas, B. F., Lo, M. H., Reager, J. T., Famiglietti, J. S., Voss, K., Swenson, S., and Rodell, M.: Quantifying renewable groundwater stress with GRACE, *Water Resour Res*, 51, 5217–5238, <https://doi.org/10.1002/2015WR017349>, 2015.

Robertson, E.: The Local Biophysical Response to Land-Use Change in HadGEM2-ES, *J. Clim.*, 32, 7611–7627, 2019.

630 Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahayan, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *J Adv Model Earth Syst*, 11, 4513–4558, <https://doi.org/10.1029/2019MS001739>, 2019.

635 Slevin, D., Tett, S. F. B., Exbrayat, J.-F., Bloom, A. A., and Williams, M.: Global evaluation of gross primary productivity in the JULES land surface model v3.4.1, *Geosci Model Dev*, 10, 2651–2670, <https://doi.org/10.5194/gmd-10-2651-2017>, 2017.

Stackhouse, P. W. Jr., Gupta, S. K., Cox, S. J., Mikovitz, J. C., Zhang, T., and Hinkelman, L. M.: The NASA/GEWEX Surface Radiation Budget Release 3.0: 24.5-Year Dataset, *GEWEX News*, 21, 2011.

- 640 [Staver, A. C., Archibald, S., and Levin, S. A.: The global extent and determinants of savanna and forest as alternative biome states, \*Science\* \(1979\), 334, 230–232, \[https://doi.org/10.1126/SCIENCE.1210465/SUPPL\\\_FILE/STAVERSOM.REVISION.1.PDF\]\(https://doi.org/10.1126/SCIENCE.1210465/SUPPL\_FILE/STAVERSOM.REVISION.1.PDF\), 2011.](#)  
[United Nations Environment Programme: Spreading like wildfire - The rising threat of extraordinary landscape fires, Nairobi, 1–126 pp., 2022.](#)
- 645 [Veenendaal, E. M., Torello-Raventos, M., Miranda, H. S., Sato, N. M., Oliveras, I., van Langevelde, F., Asner, G. P., and Lloyd, J.: On the relationship between fire regime and vegetation structure in the tropics, \*New Phytologist\*, 218, 153–166, <https://doi.org/10.1111/NPH.14940>, 2018.](#)  
[Warszawski, L., Friend, A., Ostberg, S., Frieler, K., Lucht, W., Schaphoff, S., Beerling, D., Cadule, P., Ciais, P., Clark, D. B., Kahana, R., Ito, A., Keribin, R., Kleidon, A., Lomas, M., Nishina, K., Pavlick, R., Rademacher, T. T., Buechner, M., Piontek,](#)  
650 [F., Schewe, J., Serdeczny, O., and Schellnhuber, H. J.: A multi-model analysis of risk of ecosystem shifts under climate change, \*Environmental Research Letters\*, 8, 044018, <https://doi.org/10.1088/1748-9326/8/4/044018>, 2013.](#)  
[Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., and Schewe, J.: The Inter-Sectoral Impact Model Intercomparison Project \(ISI-MIP\): Project framework, \*Proceedings of the National Academy of Sciences\*, 111, 3228–3232, <https://doi.org/10.1073/PNAS.1312330110>, 2014.](#)
- 655 [Van Der Werf, G. R., Randerson, J. T., Giglio, L., Van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., Van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, \*Earth Syst Sci Data\*, 9, 697–720, <https://doi.org/10.5194/ESSD-9-697-2017>, 2017.](#)  
[Williams, K. and Clark, D. B.: Disaggregation of daily data in JULES, Hadley Centre Technical Note, 96, 2014.](#)  
[Wiltshire, A. J., Duran Rojas, M. C., Edwards, J. M., Gedney, N., Harper, A. B., Hartley, A. J., Hendry, M. A., Robertson, E.,](#)  
660 [and Smout-Day, K.: JULES-GL7: the Global Land configuration of the Joint UK Land Environment Simulator version 7.0 and 7.2, \*Geosci Model Dev\*, 13, 483–505, <https://doi.org/10.5194/gmd-13-483-2020>, 2020.](#)  
[Wiltshire, A. J., Burke, E. J., Chadburn, S. E., Jones, C. D., Cox, P. M., Davies-Barnard, T., Friedlingstein, P., Harper, A. B., Liddicoat, S., Sitch, S., and Zaehle, S.: Jules-cn: A coupled terrestrial carbon-nitrogen scheme \(jules vn5.1\), \*Geosci Model Dev\*, 14, 2161–2186, <https://doi.org/10.5194/GMD-14-2161-2021>, 2021.](#)
- 665 -
- [Adam, J. C. and Lettenmaier, D. P.: Adjustment of global gridded precipitation for systematic bias, \*J. Geophys. Res. Atmos.\*, 108, 4257, <https://doi.org/10.1029/2002JD002499>, 2003.](#)
- [Andela, N., Morton, D. C., Giglio, L., Chen, Y., Werf, G. R. van der, Kasibhatla, P. S., DeFries, R. S., Collatz, G. J., Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Yue, C., and Randerson, J. T.: A](#)  
670 [human-driven decline in global burned area, \*Science\* \(80-. \), <https://doi.org/10.1126/SCIENCE.AAL4108>, 2017.](#)
- [Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. . L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator \(JULES\), model description – Part 1: Energy and water fluxes, \*Geosci. Model Dev.\*,](#)

- 4, 677–699, <https://doi.org/10.5194/gmd-4-677-2011>, 2011.
- 675 Bistinas, I., Harrison, S. P., Prentice, I. C., and Pereira, J. M. C.: Causal relationships versus emergent patterns in the global controls of fire frequency, *Biogeosciences*, 11, 5087–5101, <https://doi.org/10.5194/BG-11-5087-2014>, 2014.
- Brovkin, V., Raddatz, T., Reick, C. H., Claussen, M., and Gayler, V.: Global biogeophysical interactions between forest and climate, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2009GL037543>, 2009.
- Burton, C., Betts, R., Cardoso, M., Feldpausch, R. T., Harper, A., Jones, C. D., Kelley, D. I., Robertson, E., and Wiltshire, A.: Representation of fire, land-use change and vegetation dynamics in the Joint UK Land Environment Simulator vn4.9 (JULES),  
680 *Geosci. Model Dev.*, 12, 179–193, <https://doi.org/10.5194/GMD-12-179-2019>, 2019.
- Burton, C., Betts, R. A., Jones, C. D., Feldpausch, T. R., Cardoso, M., and Anderson, L. O.: El Niño Driven Changes in Global Fire 2015/16, *Front. Earth Sci.*, 8, 199, <https://doi.org/10.3389/FEART.2020.00199>/BIBTEX, 2020.
- Burton, C., Kelley, D. I., Jones, C. D., Betts, R. A., Cardoso, M., and Anderson, L.: South American fires and their impacts on  
685 ecosystems increase with continued emissions, *Clim. Resil. Sustain.*, 1, e8, <https://doi.org/10.1002/CLI2.8>, 2022.
- Canadell, J. G., Monteiro, P. M. S., Costa, M. H., Cotrim da Cunha, L., Cox, P. M., Eliseev, A. V., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P. K., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global Carbon and other Biogeochemical Cycles and Feedbacks, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: V., M.-D.,  
690 Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 673–816, <https://doi.org/https://doi.org/10.1017/9781009157896.007>, 2021.
- Cecil, D. J.: LIS/OTD 0.5 Degree High Resolution Monthly Climatology (HRMC) (No. [1995-2014]),  
695 <http://dx.doi.org/10.5067/LIS/LIS-OTD/DATA303>, 2006.
- Cheng, L., Zhang, L., Wang, Y. P., Canadell, J. G., Chiew, F. H. S., Beringer, J., Li, L., Miralles, D. G., Piao, S., and Zhang, Y.: Recent increases in terrestrial carbon uptake at little cost to the water cycle, *Nat. Commun.* 2017 81, 8, 1–10, <https://doi.org/10.1038/s41467-017-00114-5>, 2017.
- Chuvieco, E., Lizundia-Loiola, J., Lucrecia Pettinari, M., Ramo, R., Padilla, M., Tansey, K., Mouillot, F., Laurent, P., Storm,  
700 T., Heil, A., and Plummer, S.: Generation and analysis of a new global burned area product based on MODIS 250 m reflectance bands and thermal anomalies, *Earth Syst. Sci. Data*, 10, 2015–2031, <https://doi.org/10.5194/essd-10-2015-2018>, 2018.
- Ciavarella, A., Cotterill, D., Stott, P., Kew, S., Philip, S., van Oldenborgh, G. J., Skålevåg, A., Lorenz, P., Robin, Y., Otto, F., Hauser, M., Seneviratne, S. I., Lehner, F., and Zolina, O.: Prolonged Siberian heat of 2020 almost impossible without human influence, *Clim. Change*, 166, 1–18, <https://doi.org/10.1007/S10584-021-03052-W/TABLES/3>, 2021.
- 705 Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics, *Geosci. Model Dev.*, 4, 701–722,

<https://doi.org/10.5194/gmd-4-701-2011>, 2011.

- Collier, N., Hoffman, F. M., Lawrence, D. M., Keppel-Aleks, G., Koven, C. D., Riley, W. J., Mu, M., and Randerson, J. T.:  
710 The International Land Model Benchmarking (ILAMB) System: Design, Theory, and Implementation, *J. Adv. Model. Earth Syst.*, 10, 2731–2754, <https://doi.org/10.1029/2018MS001354>, 2018.
- Dai, A.: Hydroclimatic trends during 1950–2018 over global land, *Clim. Dyn.*, 56, 4027–4049,  
<https://doi.org/10.1007/S00382-021-05684-1/FIGURES/14>, 2021.
- Dai, A. and Trenberth, K. E.: Estimates of Freshwater Discharge from Continents: Latitudinal and Seasonal Variations, *J.*  
715 *Hydrometeorol.*, 3, 660–687, 2002.
- Falloon, P., Betts, R., Wiltshire, A., Dankers, R., Mathison, C., Mcneall, D., Bates, P., and Trigg, M.: Validation of River  
Flows in HadGEM1 and HadCM3 with the TRIP River Flow Model, *J. Hydrometeorol.*, 12, 1157–1180,  
<https://doi.org/10.1175/2011JHM1388.1>, 2011.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J.,  
720 Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneeth, A., Arora, V., Bates,  
N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W.,  
Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung,  
K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I.,  
Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D. R., Nabel,  
725 J. E. M. S., Nakaoka, S. I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E.,  
Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H.,  
Tilbrook, B., Van Der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan,  
W., Yue, X., and Zaehle, S.: Global Carbon Budget 2020, *Earth Syst. Sci. Data*, 12, 3269–3340, [https://doi.org/10.5194/ESSD-](https://doi.org/10.5194/ESSD-12-3269-2020)  
12-3269-2020, 2020.
- Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., Peters, G. P.,  
Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M.,  
Bellouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djéutchouang, L.  
M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N.,  
Gürses, Ö., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy,  
735 D., Goldewijk, K. K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S.,  
Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S. I., Niwa, Y., Ono, T., Pierrot,  
D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Schwingshackl, C.,  
Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., Van Der Werf, G. R.,  
Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S.,  
740 and Zeng, J.: Global Carbon Budget 2021, *Earth Syst. Sci. Data*, 14, 1917–2005, <https://doi.org/10.5194/ESSD-14-1917-2022>,  
2022.

- Frieler, K., Lange, S., Piontek, F., Reyer, C. P. O., Schewe, J., Warszawski, L., Zhao, F., Chini, L., Denvil, S., Emanuel, K., Geiger, T., Halladay, K., Hurtt, G., Mengel, M., Murakami, D., Ostberg, S., Popp, A., Riva, R., Stevanovic, M., SuzGBRi, T., Volkholz, J., Burke, E., Ciais, P., Ebi, K., Eddy, T. D., Elliott, J., Galbraith, E., Gosling, S. N., Hattermann, F., Hickler, T.,
- 745 Hinkel, J., Hof, C., Huber, V., Jägermeyr, J., Krysanova, V., Marcé, R., Müller Schmied, H., Mouratiadou, I., Pierson, D., Tittensor, D. P., Vautard, R., Van Vliet, M., Biber, M. F., Betts, R. A., Leon Bodirsky, B., Deryng, D., Frolking, S., Jones, C. D., Lotze, H. K., Lotze-Campen, H., Sahajpal, R., Thonicke, K., Tian, H., and Yamagata, Y.: Assessing the impacts of 1.5°C global warming - Simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b), *Geosci. Model Dev.*, 10, 4321–4345, <https://doi.org/10.5194/GMD-10-4321-2017>, 2017.
- 750 Gerard, F., Hooftman, D., van Langevelde, F., Veenendaal, E., White, S. M., and Lloyd, J.: MODIS VCF should not be used to detect discontinuities in tree cover due to binning bias. A comment on Hanan et al. (2014) and Staver and Hansen (2015), *Glob. Ecol. Biogeogr.*, 26, 854–859, <https://doi.org/10.1111/GEB.12592>, 2017.
- Goldewijk, K. K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates for the Holocene - HYDE 3.2, *Earth Syst. Sci. Data*, 9, 927–953, <https://doi.org/10.5194/ESSD-9-927-2017>, 2017.
- 755 Hantson, S., Arneth, A., Harrison, S. P., Kelley, D. I., Colin Prentice, I., Rabin, S. S., Archibald, S., Mouillot, F., Arnold, S. R., Artaxo, P., Bachelet, D., Ciais, P., Forrest, M., Friedlingstein, P., Hickler, T., Kaplan, J. O., Kloster, S., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Meyn, A., Sitch, S., Spessa, A., Van Der Werf, G. R., Voulgarakis, A., and Yue, C.: The status and challenge of global fire modelling, *Biogeosciences*, 13, 3359–3375, <https://doi.org/10.5194/BG-13-3359-2016>, 2016.
- 760 Hantson, S., Kelley, D. I., Arneth, A., Harrison, S. P., Archibald, S., Bachelet, D., Forrest, M., Hickler, T., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Nieradzik, L., Rabin, S. S., Colin Prentice, I., Sheehan, T., Sitch, S., Teckentrup, L., Voulgarakis, A., and Yue, C.: Quantitative assessment of fire and vegetation properties in simulations with fire-enabled vegetation models from the Fire Model Intercomparison Project, *Geosci. Model Dev.*, 13, 3299–3318, <https://doi.org/10.5194/GMD-13-3299-2020>, 2020.
- 765 Harper, A. B., Cox, P. M., Friedlingstein, P., Wiltshire, A. J., Jones, C. D., Sitch, S., Mercado, L. M., Groenendijk, M., Robertson, E., Kattge, J., Bönisch, G., Atkin, O. K., Bahn, M., Cornelissen, J., Niinemets, Ü., Onipchenko, V., Peñuelas, J., Poorter, L., Reich, P. B., Soudzilovskaia, N. A., and Bodegom, P. van: Improved representation of plant functional types and physiology in the Joint UK Land Environment Simulator (JULES v4.2) using plant trait information, *Geosci. Model Dev.*, 9, 2415–2440, <https://doi.org/10.5194/gmd-9-2415-2016>, 2016.
- 770 Harper, A. B., Wiltshire, A. J., Cox, P. M., Friedlingstein, P., Jones, C. D., Mercado, L. M., Sitch, S., Williams, K., and Duran-Rojas, C.: Vegetation distribution and terrestrial carbon cycle in a carbon cycle configuration of JULES4.6 with new plant functional types, *Geosci. Model Dev.*, 11, 2857–2873, <https://doi.org/10.5194/GMD-11-2857-2018>, 2018.
- Harper, K. L., Lamarche, C., Hartley, A., Peylin, P., Ottlé, C., Bastrikov, V., San Martín, R., Bohnenstengel, S. I., Kirches, G., Boettcher, M., Shevchuk, R., Brockmann, C., and Defourny, P.: A 29-year time series of annual 300-metre resolution plant
- 775 functional type maps for climate models, *Earth Syst. Sci. Data Discuss.*, Submitted, 2022.

- Hempel, S., Frieler, K., Warszawski, L., Schewe, J., and Piontek, F.: A trend-preserving bias correction: The ISI-MIP approach, *Earth Syst. Dyn.*, 4, 219–236, <https://doi.org/10.5194/ESD-4-219-2013>, 2013.
- Huntingford, C., Booth, B. B. B., Sitch, S., Gedney, N., Lowe, J. A., Liddicoat, S. K., Mercado, L. M., Best, M. J., Weedon, G. P., Fisher, R. A., Lomas, M. R., Good, P., Zelazowski, P., Everitt, A. C., Spessa, A. C., and Jones, C. D.: IMOGEN: An  
780 intermediate complexity model to evaluate terrestrial impacts of a changing climate, *Geosci. Model Dev.*, 3, 679–687, <https://doi.org/10.5194/GMD-3-679-2010>, 2010.
- Johannes Dolman, A. and Gregory, D.: The Parametrization of Rainfall Interception In GCMs, *Q. J. R. Meteorol. Soc.*, 118, 455–467, <https://doi.org/10.1002/QJ.49711850504>, 1992.
- Jung, M., Reichstein, M., Margolis, H. a., Cescatti, A., Richardson, A. D., Arain, M. A., Arneth, A., Bernhofer, C., Bonal, D.,  
785 Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari, F., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations, *J. Geophys. Res.*, 116, G00J07, <https://doi.org/10.1029/2010JG001566>, 2011.
- Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Bönsch, G., Garnier, E., Westoby, M., Reich, P. B., Wright, I. J.,  
790 Cornelissen, J. H. C., Violle, C., Harrison, S. P., Van BODEGOM, P. M., Reichstein, M., Enquist, B. J., Soudzilovskaia, N. a., Ackerly, D. D., Anand, M., Atkin, O., Bahn, M., Baker, T. R., Baldocchi, D., Bekker, R., Blanco, C. C., Blonder, B., Bond, W. J., Bradstock, R., Bunker, D. E., Casanoves, F., Cavender-Bares, J., Chambers, J. Q., Chapin Iii, F. S., Chave, J., Coomes, D., Cornwell, W. K., Craine, J. M., Dobrin, B. H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M., Fagan, W. F., Fang, J., Fernández-Méndez, F., Fidelis, a., Finegan, B., Flores, O., Ford, H., Frank, D., Freschet, G. T., Fyllas, N. M., Gallagher,  
795 R. V., Green, W. a., Gutierrez, a. G., Hickler, T., Higgins, S. I., Hodgson, J. G., Jalili, a., Jansen, S., Joly, C. a., Kerkhoff, a. J., Kirkup, D., Kitajima, K., Kleyer, M., Klotz, S., Knops, J. M. H., Kramer, K., Kühn, I., Kurokawa, H., Laughlin, D., Lee, T. D., Leishman, M., Lens, F., Lenz, T., Lewis, S. L., Lloyd, J., Llusià, J., Louault, F., Ma, S., Mahecha, M. D., Manning, P., Massad, T., Medlyn, B. E., Messier, J., Moles, a. T., Müller, S. C., Nadrowski, K., Naeem, S., Niinemets, Ü., Nöller, S., Nüske, a., Ogaya, R., Oleksyn, J., Onipchenko, V. G., Onoda, Y., Ordoñez, J., Overbeck, G., et al.: TRY - a global database  
800 of plant traits, *Glob. Chang. Biol.*, 17, 2905–2935, <https://doi.org/10.1111/j.1365-2486.2011.02451.x>, 2011.
- Kelley, D. I., Prentice, I. C., Harrison, S. P., Wang, H., Simard, M., Fisher, J. B., and Willis, K. O.: A comprehensive benchmarking system for evaluating global vegetation models, *Biogeosciences*, 10, 3313–3340, <https://doi.org/10.5194/BG-10-3313-2013>, 2013.
- Kelley, D. I., Bistinas, I., Whitley, R., Burton, C., Marthews, T. R., and Dong, N.: How contemporary bioclimatic and human  
805 controls change global fire regimes, *Nat. Clim. Chang.* 2019 99, 9, 690–696, <https://doi.org/10.1038/s41558-019-0540-7>, 2019.
- Lange, S.: Bias correction of surface downwelling longwave and shortwave radiation for the EWEMBI dataset, *Earth Syst. Dyn.*, 9, 627–645, <https://doi.org/10.5194/ESD-9-627-2018>, 2018.
- Lasslop, G., Brovkin, V., Reick, C. H., Bathiany, S., and Kloster, S.: Multiple stable states of tree cover in a global land surface



- 810 model due to a fire-vegetation feedback, *Geophys. Res. Lett.*, 43, 6324–6331, <https://doi.org/10.1002/2016GL069365>, 2016.
- Mangeon, S., Voulgarakis, A., Gilham, R., Harper, A., Sitch, S., and Folberth, G.: INFERNO: A fire and emissions scheme for the UK Met Office’s Unified Model, *Geosci. Model Dev.*, 9, 2685–2700, <https://doi.org/10.5194/GMD-9-2685-2016>, 2016.
- Mathison, C., Wiltshire, A. J., Falloon, P., and Challinor, A. J.: South Asia river-flow projections and their implications for water resources, *Hydrol. Earth Syst. Sci.*, 19, 4783–4810, <https://doi.org/10.5194/HESS-19-4783-2015>, 2015.
- 815 Ménard, C. B., Essery, R., Pomeroy, J., Marsh, P., and Clark, D. B.: A shrub bending model to calculate the albedo of shrub-tundra, *Hydrol. Process.*, 28, 341–351, <https://doi.org/10.1002/HYP.9582>, 2014.
- Mercado, L. M., Medlyn, B. E., Huntingford, C., Oliver, R. J., Clark, D. B., Sitch, S., Zelazowski, P., Kattge, J., Harper, A. B., and Cox, P. M.: Large sensitivity in land carbon storage due to geographical and temporal variation in the thermal response of photosynthetic capacity, *New Phytol.*, 218, 1462–1477, <https://doi.org/10.1111/NPH.15100>, 2018.
- 820 Miralles, D. G., De Jeu, R. A. M., Gash, J. H., Holmes, T. R. H., and Dolman, A. J.: Magnitude and variability of land evaporation and its components at the global scale, *Hydrol. Earth Syst. Sci.*, 15, 967–981, <https://doi.org/10.5194/HESS-15-967-2011>, 2011.
- Mu, Q., Zhao, M., and Running, S. W.: Improvements to a MODIS global terrestrial evapotranspiration algorithm, *Remote Sens. Environ.*, 115, 1781–1800, <https://doi.org/10.1016/j.rse.2011.02.019>, 2011.
- 825 Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic
- 830 Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Glob. Environ. Chang.*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- Richey, A. S., Thomas, B. F., Lo, M. H., Reager, J. T., Famiglietti, J. S., Voss, K., Swenson, S., and Rodell, M.: Quantifying renewable groundwater stress with GRACE, *Water Resour. Res.*, 51, 5217–5238, <https://doi.org/10.1002/2015WR017349>, 2015.
- 835 Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O’Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahann, A., Smith, R. S., Swaminathan,
- 840 R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *J. Adv. Model. Earth Syst.*, 11, 4513–4558, <https://doi.org/10.1029/2019MS001739>, 2019.
- Slevin, D., Tett, S. F. B., Exbrayat, J.-F., Bloom, A. A., and Williams, M.: Global evaluation of gross primary productivity in the JULES land surface model v3.4.1, *Geosci. Model Dev.*, 10, 2651–2670, <https://doi.org/10.5194/gmd-10-2651-2017>, 2017.

Stackhouse, P. W. J., Gupta, S. K., Cox, S. J., Mikovitz, J. C., Zhang, T., and Hinkelman, L. M.: The NASA/GEWEX Surface  
845 Radiation Budget Release 3.0: 24.5-Year Dataset, *GEWEX News*, 21, 2011.

Staver, A. C., Archibald, S., and Levin, S. A.: The global extent and determinants of savanna and forest as alternative biome  
states, *Science* (80-. ),, 334, 230–232,  
[https://doi.org/10.1126/SCIENCE.1210465/SUPPL\\_FILE/STAVERSOM.REVISION.1.PDF](https://doi.org/10.1126/SCIENCE.1210465/SUPPL_FILE/STAVERSOM.REVISION.1.PDF), 2011.

United Nations Environment Programme: Spreading like wildfire - The rising threat of extraordinary landscape fires, Nairobi,  
850 1–126 pp., 2022.

Veenendaal, E. M., Torello-Raventos, M., Miranda, H. S., Sato, N. M., Oliveras, I., van Langevelde, F., Asner, G. P., and  
Lloyd, J.: On the relationship between fire regime and vegetation structure in the tropics, *New Phytol.*, 218, 153–166,  
<https://doi.org/10.1111/NPH.14940>, 2018.

Warszawski, L., Friend, A., Ostberg, S., Frieler, K., Lucht, W., Schaphoff, S., Beerling, D., Cadule, P., Ciais, P., Clark, D. B.,  
855 Kahana, R., Ito, A., Keribin, R., Kleidon, A., Lomas, M., Nishina, K., Pavlick, R., Rademacher, T. T., Buechner, M., Piontek,  
F., Schewe, J., Serdeczny, O., and Schellnhuber, H. J.: A multi-model analysis of risk of ecosystem shifts under climate change,  
*Environ. Res. Lett.*, 8, 044018, <https://doi.org/10.1088/1748-9326/8/4/044018>, 2013.

Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., and Schewe, J.: The Inter-Sectoral Impact Model  
Intercomparison Project (ISI-MIP): Project framework, *Proc. Natl. Acad. Sci.*, 111, 3228–3232,  
860 <https://doi.org/10.1073/PNAS.1312330110>, 2014.

Van Der Werf, G. R., Randerson, J. T., Giglio, L., Van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., Van Marle, M. J.  
E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997-2016,  
*Earth Syst. Sci. Data*, 9, 697–720, <https://doi.org/10.5194/ESSD-9-697-2017>, 2017.


Williams, K. and Clark, D. B.: Disaggregation of daily data in JULES, Hadley Cent. Tech. Note, 96, 2014.

865 Wiltshire, A. J., Duran Rojas, M. C., Edwards, J. M., Gedney, N., Harper, A. B., Hartley, A. J., Hendry, M. A., Robertson, E.,  
and Smout-Day, K.: JULES-GL7: the Global Land configuration of the Joint UK Land Environment Simulator version 7.0  
and 7.2, *Geosci. Model Dev.*, 13, 483–505, <https://doi.org/10.5194/gmd-13-483-2020>, 2020.


Wiltshire, A. J., Burke, E. J., Chadburn, S. E., Jones, C. D., Cox, P. M., Davies-Barnard, T., Friedlingstein, P., Harper, A. B.,  
Liddicoat, S., Sitch, S., and Zaehle, S.: Jules-cn: A coupled terrestrial carbon-nitrogen scheme (jules vn5.1), *Geosci. Model*  
870 *Dev.*, 14, 2161–2186, <https://doi.org/10.5194/GMD-14-2161-2021>, 2021.

## Figures


GFDL-ESM2M




HADGEM2-ES




IPSL-CM5A-LR




MIROC5






875 **Figure 1: Multi-year mean bias of catchment scale runoff simulated by JULES driven by 4 sets of climate driving data compared to runoff derived from (Dai, 2021) (Dai, 2021). Number of years of observations contributing to the multi-year mean varies depending on catchment, however, observations within the period 1980-2006 were accepted. ISIMIP2b forcing data derived from 4 CMIP5 GCMs: GFDL-ESM2M; HadGEM2-ES; IPSL-CM5A-LR; MIROC5.**




**Figure 2: Comparison of the simulated long term monthly mean river flow with observations (Dai, 2021) for the six largest rivers: a) Amazon (Obidos, -1.95N,-55.51E); b) Congo (Kinshasa, -4.3N, 15.3E); c) Orinoco (Pte Angostu , 8.15N,-63.6E); d) Changjiang (Datong, 30.77N, 117.62E); e) Brahmaputra (Bahadurabad, 25.18N, 89.67E); f) Mississippi (Vicksburg 32.31N, -90.91E). The observations are over the time period 1980 to 2010.**

# ISIMIP: Multi-year Annual Mean






# ISIMIP: Multi-year Annual Mean





890 **Figure 3: Multi-year annual mean for GPP (column 1), Evapotranspiration (Column 2) and Albedo (column 3) for observations**  
 (row 1) and subsequent rows show the anomaly compared with observations for each set of ISIMIP2b forcing data derived from:  
 GFDL-ESM2M (row 2), HADGEM2-ES (row 3), IPSL-CM5A-LR (row 4) and MIROC5 (row 5). Observations are downloaded  
 from iLAMB ([https://www.ilamb.org/doc/ilamb\\_fetch.html](https://www.ilamb.org/doc/ilamb_fetch.html)) and the ~~datasets~~ shown are GBAF for GPP (Jung et al.,  
 2011)(Jung et al., 2011), GLEAM for ET (Miralles et al., 2011)(Miralles et al., 2011) and GEWEX.SRB for Albedo (Stackhouse et  
 895 al., 2011)(Stackhouse et al., 2011).




**Figure 4: Observed vegetation fractional cover, derived from ESACCI Land Cover (v2.0.7) for 2010 (K. L. Harper et al., 2022), for (left to right) tree, shrub, grass and unvegetated (bare) fraction. Subsequent rows show the difference between model (without fire) and ES CCI observations for each set of driving data.**




a) Fire CCI




b) GFDL-ESM2M





c) HADGEM2-ES



d) IPSL-CM5A-LR



e) MIROC5



Burnt Area Fraction





Figure 5: Present day [percentage](#) burned area [fraction](#) (2000-2020) from and Fire CCI observations (a) and modelled by JULES driven by [GFDL-ESM2M](#) (b), [HadGEM2-ES](#) (c), [IPSL-CM5A-LR](#) (d), [MIROC5](#) (e)



**Figure 6: Modelled vegetation cover without fire (left) and with fire (middle) compared to observations from ESA CCI Land Cover and Fire (right). Dark green, light green, light brown, dark brown indicates tree, shrub, grass and unvegetated fraction of the latitude band. Shaded transition between colours indicates ensemble range, which is quite narrow indicating agreement across ensemble members. Black hashing indicates burnt area, with observations taken from MODIS CCI v5.1 (Chuvieco et al., 2018)(Chuvieco et al., 2018). In “Model with fire”, burnt area from the 4 driving models is shown by hatching at 4 different angles.**

## Author contribution

Camilla Mathison: Conceptualisation; Data curation; Formal analysis; Investigation; Methodology; Project administration; Software; Writing – original draft; Writing – review & editing

920 Eleanor Burke: Conceptualisation; Data curation; Investigation; Methodology; Software; Writing – original draft; Writing – review & editing

Andrew Hartley: Data curation; Formal analysis; Investigation; Methodology; Project administration; Validation; Visualisation; Writing – original draft; Writing – review & editing

Douglas I Kelley: Formal analysis; Investigation; Methodology; Validation; Visualisation; Writing – original draft; Writing – review & editing

925 Chantelle Burton: Data curation; Methodology; Formal analysis; Investigation; Validation; Visualisation; Writing – original draft; Writing – review & editing

Eddy Robertson: Formal analysis; Software; Validation; Writing – review & editing

Nicola Gedney: Validation; Visualisation; Writing – original draft; Writing – review & editing

Karina Williams: Methodology; Writing – review & editing

930 Andy Wiltshire: Writing – review & editing

Richard J Ellis: Software; [Writing – review & editing](#)

Alistair A Sellar: Software

Chris D. Jones: Validation; Writing – review & editing

## 935 Acknowledgments

Work and its contributors were supported by the Newton Fund through the Met Office Climate Science for Service Partnership Brazil (CSSP Brazil) (CB, CM, AH, AW [CJ and NG](#)), the Met Office Climate Science for Service Partnership South Africa (CSSP South Africa) (AH and CM). CM, KW, AW, AH, AS, EB, ER, ~~CJ~~ and ~~CJ NG~~ were supported by the Met Office Hadley Centre Climate Programme funded by BEIS and Defra. The contribution by DIK and RJE was supported by the UK Natural Environment Research Council through The UK Earth System Modelling Project (UKESM, grant no. NE/N017951/1). DIK was [additionally](#) supported by UK Natural Environment Research Council as part of the NC-International programme (NE/X006247/1). ~~EB~~ and [CJ were supported by](#) the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 101003536 (ESM2025 - Earth System Models for the Future)

940