

Differential impact of isolated topographic bumps on ice sheet flow and subglacial processes

Marion A. McKenzie¹, Lauren E. Miller¹, Jacob S. Slawson^{1*}, Emma J. MacKie², and Shujie Wang³

¹Department of Environmental Sciences, University of Virginia, 291 McCormick Rd., Charlottesville, VA, USA 22904

²Department of Geological Sciences, University of Florida, 241 Williamson Hall, Gainesville, FL, USA 32611-2120

³Department of Geography, Pennsylvania State University, 302 N Burrowes St., University Park, PA, USA 16802

*Current affiliation: Department of Geology and Geological Engineering, Colorado School of Mines, 1516 Illinois St., Golden, CO, USA 80401

10 Correspondence to: Marion A. McKenzie (mm8dt@virginia.edu)

Abstract. Topographic highs (“bumps”) across glaciated landscapes have the potential to temporarily slow ice sheet flow or, conversely, accelerate ice flow through subglacial strain heating and meltwater production. Isolated bumps of variable size across the deglaciated landscape of the Cordilleran Ice Sheet (CIS) of Washington state present an opportunity to study the influence of topographic highs on ice-bed interactions and ice flow organization. This work utilizes semi-automatic mapping techniques of subglacial bedforms to characterize the morphology of streamlined subglacial bedforms including elongation, surface relief, and orientation, all of which provide insight into subglacial processes during post-Last Glacial Maximum deglaciation. We identify a bump-size threshold of several cubic kilometers, around 4.5 km³, in which bumps larger than this size will consistently and significantly disrupt both ice-flow organization and subglacial sedimentary processes which are fundamental to the genesis of streamlined subglacial bedforms. Additionally, sedimentary processes are persistent and well-developed downstream of bumps as reflected by enhanced bedform elongation and reduced surface relief, likely due to increased availability and production of subglacial sediment and meltwater. While isolated topography plays a role in disrupting ice flow, larger bumps have a greater disruption to ice flow organization, while bumps below the identified threshold seem to have little effect on ice and subglacial processes. The variable influence of isolated topographic bumps on ice flow of the CIS has significant implications for outlet glaciers of the Greenland Ice Sheet (GrIS) due to similarities in regional topography where local bumps are largely unresolved.

1 Introduction

Isolated topographic highs in the terrain beneath ice sheets can contribute to increased basal drag and decreased ice flow velocity and, for marine-based margins, offer pinning-points to halt or slow down margin retreat (Durand et al., 2011; Favre

Deleted: glacial

Formatted: Top: 0.39", Bottom: 0.93", Footer distance from edge: 0.51"

Commented [MM1]: Author (profile number 415616) has changed her name.

Deleted: . Simkins

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: P

Deleted: g

Deleted: glacial

Formatted: Font color: Text 1

Deleted: increase

Deleted: subglacial

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: assess

Formatted: Font color: Text 1

Deleted: --

Deleted: of the landscape

Deleted: ~ 4.5

Deleted: km³

Deleted: --

Formatted: Font color: Text 1

Deleted: most mature

Deleted: is found to

Formatted: Font color: Text 1

Deleted: not all bumps have the same degree of impact.

Formatted: Font color: Text 1

Deleted: in this system

Deleted: significance

Deleted: general topographic

Formatted: Font color: Text 1

Formatted: Font color: Text 1

et al., 2016; Alley et al., 2021; Robel et al., 2022). Conversely, ice flow over topographic highs can increase strain heating and basal meltwater production, elevating basal meltwater pressure and reducing basal friction in the downstream environment (Payne and Dongelmans, 1997; Cuffey and Paterson, 2010). However, identifying which forms and scales of “bumps” across a glaciated landscape may increase, decrease, or not affect ice-flow velocity, basal water pressure, and basal friction is not well understood outside of simple geophysical modeling (Alley et al., 2021). Additionally, topography at the base of the ice sheet, and even for most glacier catchments, is poorly resolved (e.g., MacKie et al., 2020; Morlighem et al., 2020). Therefore, we turn to a formerly glaciated landscape in Washington state where geomorphological indicators of ice-flow conditions in the form of streamlined subglacial bedforms (e.g., glacial lineations, whalebacks, and drumlins) can be used to better understand the sensitivity of ice sheets to isolated bumps in the subglacial environment. Due to the controls of basal shear, meltwater and sediment availability, and sediment processes including transport, deposition, and erosion on bedform synthesis, morphometrics of streamlined subglacial bedforms offer information on ice-bed interactions (Schoof and Clark, 2008; Shaw et al., 2008; King et al., 2009). Quantitatively assessing streamlined subglacial bedform morphometrics, provides an opportunity to assess characteristics of paleo-ice flow organization and relative speeds across a landscape (e.g., Clark, 1997, 1999; King et al., 2009; Clark et al., 2003, 2009; Spagnolo et al., 2012, 2014; Principato et al., 2016). Assessment of streamlined subglacial bedforms and their implications for ice flow are applicable to modern ice sheets (MacKie et al., 2021), where observations and theory of subglacial conditions are spatially and temporally limited.

1.1 Site Characteristics

The Puget Lowland of Washington state was glaciated by the southwestern Cordilleran Ice Sheet (CIS) during the Last Glacial Maximum (LGM), when the region was largely depressed below sea level due to glacial isostatic adjustment (GIA; Booth and Hallet, 1993; Dethier et al., 1995; Kovanen and Slaymaker, 2004; Eyles et al., 2018); therefore, the southwestern CIS was predominantly marine based. The Puget Lowland was glaciated at least six times by the CIS during the Quaternary (Clague & James, 2002; Booth et al., 2003) indicating the highly active growth and decay of the marine-terminating southern lobe. The drivers of CIS behavior are not well understood, as the growth and decay of this formidable-size ice sheet has not been found to coincide with global perturbations like its eastern neighbor, the Laurentide Ice Sheet (Broecker, 1994; Cheng et al., 2016; Blunier & Brook, 2001; Walczak et al., 2020). Active tectonics and volcanic activity across the Puget Lowland led to exposed crystalline and volcanic bedrock of Eocene age interrupting sedimentary bedrock across the region (Khazaradze et al., 1999; Booth et al., 2004). On a larger scale, the Puget Lowland is a basin surrounded by mountainous terrain near the coast of the Pacific Ocean with these isolated topographic highs, similar to the terrain beneath the margins of the Greenland Ice Sheet (Bamber et al. 2013; Eyles et al., 2018). The variability in the hard to soft-bed conditions in the Puget Lowland also bears similarity to the mixed-bed conditions seen beneath Thwaites Glacier (Schroeder et al., 2014; Holschuh et al., 2020; Alley et al., 2021). Based on simple geophysical subglacial models, focused on Thwaites Glacier (Alley et al., 2021), it is likely, high relief regions influence marine-based ice-bed interactions through basal drag and sediment reorganization. As such, the

- Deleted: basal
- Formatted: Font color: Text 1
- Deleted: -
- Deleted: -
- Deleted: , such as
- Deleted: ,
- Formatted: Font color: Text 1
- Deleted: M
- Deleted: and
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: empirical
- Deleted: (
- Deleted: y)
- Formatted: Font color: Text 1

- Deleted: have
- Formatted: Font: Not Bold
- Deleted: ing
- Deleted: highly possible
- Deleted: the
- Deleted: er
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1

95 crystalline and volcanic bedrock exposures or “bumps” across the Puget Lowland likely record bump influence in the
 geomorphic record, but this concept has yet to be empirically tested across the region. This work aims to determine the role of
 topographic bumps on glacial ice flow and sedimentary processes via streamlined subglacial bedform morphology and
 distribution due to its relevance for constraining the influence of subglacial topography beneath contemporary glacial ice. By
 investigating ice flow behavior within a single glacial system, the effects of isolated crystalline bedrock highs on ice flow will
 100 neither be confounded by geographically variable conditions such as local climate and ocean forcings, nor the vastly different
 timescales of deglaciation and post-glacial landscape evolution.

2. Methodology

2.1 Topographic “Bump” Classification

Digital elevation models (DEMs) with horizontal resolution of 1.83 x 1.83 meters and vertical resolution of 2 meters from
 105 across the Puget Lowland (Clallam County, Olympic Department of Natural Resources, WA, 2008; Quantum Spatial Inc.,
 2017, 2019; OCM Partners, 2019a, 2019b) and ambient occlusion hillshading techniques (c.f., McKenzie et al., 2022) were
 used to analyze nine crystalline and volcanic bedrock bumps for streamlined subglacial bedforms across the Puget Lowland.
 The nine identified bumps span a wide range in peak elevation, bump surface area, and bump volume (Fig. 1). While some
 bumps have more coarse topography than others, the lack of ice streaming seen in-between singular bump topography, as
 110 determined by the presence of streamlined subglacial bedforms, allowed us to treat all bumps as “aggregate” features. The
 outermost 100-foot closed contour across each aggregate bump was expanded to three times the surface area to classify the
 region of interest, following the modeled influence of bump perturbations on basal hydrologic potential (Alley et al., 2021).
 While present-day elevations of these deglaciated sites differ from elevations during glaciation due to GIA, tectonics, and post-
 glacial landscape evolution, the scale of bump relief in relation to streamlined bedforms is well preserved. Fractures, faults,
 115 and joints from tectonic activity and brittle deformation of the crust across bumps are below the scale of analysis for this work
 and are therefore not considered here.

2.2 Streamlined subglacial bedform identification

Streamlined subglacial bedforms were identified across the nine bump sites using a combination of Topographic Position Index
 120 (TPI) analysis (McKenzie et al., 2022), contour-tree mapping (Wang et al., 2017), and manual identification. TPI utilizes DEM
 slope variations across defined cell-neighborhood sizes to semi-automatically identify positive relief features (McKenzie et
 al., 2022). Morphometric threshold limits were applied to TPI-mapped bedform length, width, and area to increase the accuracy
 of the tool. In the larger Puget Lowland region, the TPI tool generally maps 80% of the bedforms correctly (McKenzie et al.,
 2022). Additionally, localized contour-tree mapping was utilized on the DEM data to isolate closed contours within a defined
 125 elevation (Wang et al., 2017). Both tool outputs were validated and corrected through manual removal of incorrectly identified

- Deleted: influenced marine and terrestrial-based ice-bed ... [2]
- Formatted ... [1]
- Deleted: The Puget Lowland is a basin surrounded by ... [4]
- Formatted ... [3]
- Formatted ... [5]
- Formatted ... [6]
- Deleted: assessing
- Formatted ... [7]
- Formatted ... [8]
- Deleted: not
- Deleted: and
- Formatted ... [9]
- Formatted ... [10]
- Formatted ... [11]
- Deleted: coarsest
- Formatted ... [12]
- Deleted: utilized
- Deleted: to
- Formatted ... [13]
- Formatted ... [14]
- Deleted: assess
- Deleted: with
- Deleted: , and topographic setting
- Formatted ... [15]
- Formatted ... [16]
- Formatted ... [17]
- Formatted ... [18]
- Formatted ... [19]
- Deleted: bump
- Deleted: by
- Formatted ... [20]
- Formatted ... [21]
- Deleted: .
- Deleted: relative relief and influence of these
- Deleted: s
- Deleted: on the presence of
- Formatted ... [22]
- Formatted ... [23]
- Deleted: Localized contour-tree mapping utilizes DEM da ... [24]
- Deleted: if needed by

155 features and manual addition of bedforms missed by the automated tools, resulting in a highly accurate dataset of bedforms across the region (McKenzie et al., 2022).

160 All bedforms in the final dataset (n=3,273) have an associated long-axis length, cardinal orientation, width orthogonal to long-axis length, and range in elevation across the long axis. These metrics were calculated by the ArcGIS Pro “Minimum Bounding Geometry” and “Add Z Information” tools (McKenzie et al., 2023). Long axis cardinal direction, or orientation, of streamlined bedforms is used to infer direction of ice flow (Clark, 1997; Kleman et al., 2006; Kleman and Borgström, 1996). Bedform elongation ratio, calculated by dividing a bedform’s length by its width is used to infer relative speed of ice flow velocity (Clark 1997, 1999; Clark et al., 2003) and relative duration of ice presence in a region (Benediktsson et al., 2016). Bedform surface relief, the difference between the highest and lowest elevation along the bedform long axis, is used to infer persistence of ice flow and sedimentary processes in the subglacial environment (McKenzie et al., 2022). Smaller surface relief values indicate more persistent, warm-based ice flow across the bed with well-developed sedimentary processes in the subglacial environment than larger values of bedform surface relief (McKenzie et al., 2022). The rate of GIA in the region (centimeters per year; Dethier et al., 1995), is much larger than rates of regional tectonics (millimeters per year; Sherrod et al., 2008). Therefore, along the long axis of a single bedform, impacts of post-glacial erosion and regional-scale tectonics cannot be isolated from larger-scale dynamics such as GIA and glacial sedimentary processes. Due to these timescale discrepancies, a high resolution assessment of where bedform surface relief occurs across an individual long axis was not conducted in this work.

170 For each site, bedforms were categorized into groups “upstream”, “on top of”, and “downstream” as determined by bedform location with respect to the outermost 100-foot closed contour of the topographic high. There is a potential for bumps to develop lateral shear margins as a result of increasing ice speed along the edge of the bump (Alley, 1993; Stokes et al., 2007). Therefore, bedforms mapped alongside the bump were classified as “downstream” of the bump, considering the overlying ice in the lateral direction would have contacted the bump. While bedforms themselves cause shear stress on ice (Damsgaard et al., 2020), the scale of the bumps to bedforms makes the influence of bedforms on ice flow negligible in this case study. Additionally, the large timescales of uniform glaciation in this region contributes to a consideration of these bedforms as a final “snap shot” of bedform development.

180 Bedforms in the upstream and downstream environments are likely to be sedimentary forms due to large sediment availability, while bedforms on top of bumps are interpreted to be erosional features developed in the sediment-starved environment of the crystalline bedrock highs. While the difference in generation is loosely referred to throughout this work, the specific sedimentary composition of the bedform was not explicitly considered as we analyze bedform metrics collectively for all erosional and depositional features. Defining bedforms solely on shape rather than composition is a strength of this work as it considers topographic influence on ice flow and sediment processes without commenting on specific landform-generating processes (McKenzie et al., 2022).

185 To quantify bump influence, we performed analysis of variance (ANOVA) and non-parametric Kruskal-Wallis tests on groups of streamlined subglacial bedforms across bumps to compare the statistical significance of the means and

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: Bedforms identified “downstream” of bumps include bump-lateral features

Formatted: Font color: Text 1

Formatted: Font: Not Bold, Not Italic, Font color: Text 1

Formatted: Font: Not Bold, Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Indent: First line: 0.5"

Formatted: Font color: Text 1

Formatted: Font: Not Bold, Not Italic, Font color: Text 1

Formatted: Font: Not Bold, Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Bold, Italic, Font color: Text 1

Deleted: Long axis cardinal direction, or orientation, of streamlined bedforms is used to infer direction of ice flow (Clark, 1997; Kleman et al., 2006; Kleman and Borgström, 1996). Bedform elongation ratio, calculated by dividing a bedform’s length by its width is used to infer relative speed of ice flow velocity (Clark 1997, 1999; Clark et al., 2003) and relative duration of ice presence in a region (Benediktsson et al., 2016). Bedform surface relief, the difference between the highest and lowest elevation along the bedform long axis, is used to infer maturity of ice flow and sedimentary processes in the subglacial environment, where smaller surface relief values indicate more mature sedimentary processes in the subglacial environment than larger values (McKenzie et al., 2022). ...

Deleted: W

Formatted: Font color: Text 1

Formatted: Font color: Text 1

distributions between populations, respectively, in “R”. Results of statistical analyses were used to determine significance ($p < 0.05$) of bedform characteristics at each site (i.e., upstream, on top of, and downstream of bumps) as well as significance of bedform morphometrics across sites, where significance is defined as a p value less than 0.05 between groups.

3. Results

The number of streamlined bedforms per site is positively correlated with bump surface area and volume (Fig. 1). When considering morphologies of streamlined bedform location across the bumps, on top of all bump sites, bedform elongation for the full dataset ($n = 3,273$) is lowest and bedform surface relief is highest (Fig. 2). Differences between upstream and downstream subglacial streamlined bedform morphometrics are not notably different across many of the sites (Fig. 3), however, one difference of note is the increased number of bedforms downstream of bumps than upstream. At seven of the nine sites, surface relief along bedform crests increases significantly between populations upstream and on top of bumps (average increase of 21 meters; Fig. 3B). The greatest number and most elongate bedforms, as well as the greatest proportion of bedforms with low surface relief, occur downstream of bumps (Fig. 1A; Fig. 2; Fig. 4A, 4B). Notably, there is a statistically significant decrease in bedform elongation between upstream and on top of the two largest bumps (average decrease of 1.5; Fig. 3A), Blue Hills and Devils Mountain.

While bedform surface relief and elongation ranges overlap across all site populations, bedforms associated with smaller bumps tend to have outliers below the 1σ (68%) confidence level for all populations (e.g., San Juan Island, Fidalgo Island, and Black Hills) while those associated with larger bumps have outliers above the 1σ confidence level (e.g., Blue Hills and Cougar Mountain; Fig. 2). The greatest proportion of bedforms with low surface relief (Fig. 4B) are located at the smallest bump sites ($< 0.3 \text{ km}^3$). Many sites showcase an increase in disorganization of bedform orientation on top of the bump (Fig. 4C), but only at the two largest bumps ($> 4.5 \text{ km}^3$) does downstream bedform orientation recover to patterns present in the upstream bedform populations (average orientation within 0.5 degrees similarity; Fig. 4C). The remaining sites have bedform orientations that either remain unchanged or develop less agreement in bedform orientation downstream (Fig. 4C).

4. Discussion and Interpretation

Positive correlation between site area and number of bedforms indicates spatial continuity in the bedform distribution across the Puget Lowland. Where a reduction in bedform elongation and surface relief occurs on top of bumps, this relationship suggests a slow in ice flow speed or persistence at the same time as a reduction in spatial homogeneity of sedimentary processes in developing streamlined subglacial bedforms. As such, we interpret that bumps in the subglacial environment of the CIS generally led to ice-flow deceleration and a reduction of spatial homogeneity (i.e., efficiency) of sedimentary processes including bedrock erosion and sediment transport and deposition – all of which are important for bedform genesis (Schoof and Clark, 2008; Shaw et al., 2008; King et al., 2009).

The effect of bumps on subglacial processes and ice flow are seen on top of almost all site bumps (Fig. 3). The nearly ubiquitous observation of bedform surface relief decrease on top of bumps is most likely due to a transition in lithology from

Deleted: and Discussion

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Indent: First line: 0.5"

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ¶

The number of streamlined bedforms per site is positively correlated with site surface area and volume (Fig. 1B),

Deleted: ing

Deleted: On top of all bump sites, bedform elongation for the full dataset ($n = 3,273$) is lowest and bedform surface relief is highest (Fig. 2).

Deleted: We, thus,

Deleted: Cordilleran Ice Sheet

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: efficiency or

Formatted: Font color: Text 1

Deleted: ¶

Deleted: While bedform surface relief and elongation ranges overlap across all site populations, bedforms associated with smaller bumps tend to have outliers below the 1σ (68%) confidence level for all populations (e.g., San Juan Island, Fidalgo Island, and Black Hills) while those associated with larger bumps have outliers above the 1σ confidence level (e.g., Blue Hills and Cougar Mountain; Fig. 2). This trend in outliers demonstrates a linkage between bump size and possible bedform morphometrics in a relatively systemic manner across the Puget Lowland. Notably, there is a significant decrease in bedform elongation between upstream and on top of the two largest bumps (Fig. 3A), Blue Hills and Devils Mountain, suggesting bump volume larger than 4.5 km^3 significantly slows or causes disorganization in ice flow (Fig. 1B; Clark 1997, 1999; Clark et al., 2003).

At seven of the nine sites, surface relief along bedform crests increases significantly between populations upstream and on top of bumps (Fig. 3B).

Formatted: Font color: Text 1

Deleted: subglacial

270 sedimentary to crystalline or volcanic bedrock, ~~which disrupts along-flow~~ sedimentary processes as ice ~~begins developing~~
~~subglacial streamlined bedforms on~~ more-erosion-resistant ~~and sediment-starved~~ bed compositions. The two exceptions to this
trend are Big Skidder Hill and Lopez Island, where there is no appreciable change in streamlined subglacial bedform surface
relief across the bumps (Fig. 3B). ~~This suggests that the conditions at these two sites were able to overcome direct lithologic~~
impact on bedform relief. Due to the more-erosion-resistant lithologies of the bumps, combined with increased pressure and
basal drag in the subglacial environment, ~~we propose~~ decreased efficiency in which the ice is able to facilitate streamlined
subglacial bedform formation through bedrock erosion (Eyles and Doughty, 2016; Krabbendam et al., 2016), leading to
truncated bedforms with high surface relief (McKenzie et al., 2022; Fig. 2). ~~Despite the influence of bumps on ice flow and~~
275 ~~sedimentary processes at almost all sites, the similarities between upstream and downstream bedform metrics are notably~~
~~similar. However, there are far more streamlined subglacial bedforms identified downstream of bumps.~~ Increased sediment
availability and basal meltwater that results from the strain heating on top of the bump (Payne and Dongelmanns, 1997) increases
downstream sediment transport efficiency (McIntyre, 1985; Pohjola and Hedfors, 2003; Winsborrow et al., 2010b) ~~which~~
~~could contribute to development of a greater number of bedforms and recovery of the ice flow and sedimentary systems to~~
280 ~~reflect bedform metrics seen upstream, prior to bump influence.~~

Bumps with surface relief generally below a few square kilometers and volume smaller than around 0.3 cubic
kilometers ~~have predominantly low-elongation, low-surface relief bedform outliers while large sites have highly elongate, high~~
~~surface relief bedform outliers demonstrates a linkage between bump size and possible bedform morphometrics in a relatively~~
285 ~~systemic manner across the Puget Lowland (Fig. 2). We postulate that bump size – through its impact on ice flow and subglacial~~
~~processes – is a control on downstream bedform metrics. From observations of bedform orientation recovery only at the largest~~
~~two sites, we infer a bump volume of over several cubic kilometers (>4.5 km³) will influence reorganization of downstream~~
ice-flow orientation and subglacial sedimentary processes, while bumps below this threshold cannot regain the same
organization seen upstream of bumps. This analysis found ~~little~~ evidence of channelized meltwater in the subglacial
environment, ~~suggesting that~~ meltwater development across these bumps was distributed and saturated ~~in the porous upstream~~
290 ~~and downstream environments with water films supported atop the crystalline bedrock highs. An increase in subglacial~~
~~meltwater downstream of bumps is supported by the presence of homogeneity in bedform metrics, which would support~~
~~homogenous deformation of sediments into streamlined bedforms (Shaw et al., 2008; King et al., 2009).~~

An additional consideration for this work is its relevance to modern subglacial systems with similar topographic
conditions. While local climatic factors render the specific thresholds for bump influence on ice and sedimentary process
295 behavior relevant only in this glacial system, the process-based understanding derived here can be applied to systems where
isolated topographic highs are identified. Topographic similarities between this region and outlet glaciers in Greenland (Eyles
et al., 2018), modern Thwaites (Holschuh et al., 2020; Alley et al., 2021), and other contemporary glaciers with isolated
topographic highs (Greenwood et al., 2021) highlight the need for closer study of the effect of subglacial bumps on ice flow
in these systems with their own unique set of local climatic influences.

300

Deleted: disrupting

Deleted: contacted

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ; therefore, suggesting

Formatted: Font color: Text 1

Deleted: there is

Deleted:

Formatted: Font color: Text 1

Deleted: We postulate that bump size – through its impact on ice flow and subglacial processes – is a major control on bedform surface relief, where the greatest proportion of bedforms with low surface relief (Fig. 4B) are located at the smallest bump sites (< 0.2 km³).

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ,

Deleted: , resulting in the greatest number and most elongate bedforms, as well as the greatest proportion of bedforms with low surface relief, downstream of bumps (Fig. 1A; Fig. 2; Fig. 4A, 4B).

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ¶

While many sites showcase an increase in disorganization of bedform orientation on top of the bump, only at the two largest bumps (> 4.5 km³) does downstream bedform orientation recover to patterns present in the upstream bedform populations (Fig. 4C). The rest of the sites have bedform orientations that either remain unchanged or become more disorganized downstream (Fig. 4C).

Deleted: these findings

Formatted: Font color: Text 1

Deleted: ~ 4.5 km³

Formatted: Font color: Text 1

Deleted: no

Deleted: potentially

Formatted: Font color: Text 1

Deleted: , which would explain

Deleted: formation

Deleted: observed in bedforms downstream of bumps

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: Bold, Italic, Font color: Text 1

Formatted: Font color: Text 1

4. Conclusions

330 Overall, there is general ice flow deceleration and reduction of bedrock erosion efficiency on top of bumps, which results from a subglacial lithology transition. Sedimentary processes, essential to streamlined bedform genesis, are organized and efficient downstream of bumps despite disruption from crystalline bedrock highs - likely as a result of increased sediment availability and subglacial meltwater sourced from strain heating on top of the bump. The two largest bumps in both volume and surface area considered in this work notably disturb ice-flow orientation and speed on top of the bump with only bumps 335 larger than several cubic kilometers indicating recovery of ice flow orientation and speed downstream of bumps. Findings from these paleo-subglacial bumps may be used as an analog for ice flow in contemporary ice sheets and support process-based understanding of subglacial terrain influence on overlying ice-sheet behavior in similar systems such as those beneath Thwaites Glacier and outlet glaciers within the Greenland Ice Sheet.

Deleted: most

Deleted: sites

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ~4.5 km³

Formatted: Font color: Text 1

340 5. Data availability

All bedform data produced from this work is publicly available through PANGAEA and are available by request from the corresponding author.

6. Author contribution

345 Project conceptualization, data curation, methodology, formal analysis, initial draft writing, and editing were conducted by M. McKenzie. Conceptualization, funding acquisition, formal analysis, editing, and supervision were conducted by L. Miller. Conceptualization, preliminary research, and editing were conducted by J. Slawson. Partial conceptualization and editing were conducted by E. MacKie. Data curation support and editing were conducted by S. Wang.

Deleted: Simkins

Deleted: Support with data

350 7. Competing interests

The authors declare that they have no conflict of interest.

8. Acknowledgments

355 We acknowledge the Washington Department of Natural Resources for their accessible LiDAR data that made this project possible. The sites analyzed for this work are located on land historically cultivated and inhabited by the Skokomish, Suquamish, Squaxin, Stl'pulmsh, Steilacoom, Puyallup, Muckleshoot, and Duwamish peoples, while much of the data analysis and interpretation were conducted on land cultivated and inhabited by the Monacan Nation. The peoples of these Nations were custodians of the land for time immemorial before forced removal and genocide during colonization. The authors acknowledge their ongoing stewardship of the lands. This work was funded by the Chamberlain Endowment and the H.G. Goodell 360 Endowment at the University of Virginia.

References

- Alley, R.B.: In search of ice-stream sticky-spots, *Journal of Glaciology*, 39(133), 447-454, doi: 10.3189/S0022143000016336, 1993.
- Alley, R.B., Holschuh, N., MacAyeal, D.R., Parizek, B.R., Zoet, L., Riverman, K., Muto, A., Christianson, K., Clyne, E., Anandakrishnan, S., Stevens, N., GHOST Collaboration: Bedforms of Thwaites Glacier, West Antarctica: Character and Origin, *Journal of Geophysical Research: Earth Surface*, 126:12, <https://doi.org/10.1029/2021JF006339>, 2021.
- 370 Bamber, J.L., Griggs, J.A., Hurkmans, R.T.W.L., Dowdeswell, J.A., Gogineni, S.P., Howat, I., Mougnot, J., Paden, J., Palmer, S., Rignot, E., and Steinhage, D.: A new bed elevation dataset for Greenland, *The Cryosphere*, 7, 499-510, <https://doi.org/10.5194/tc-7-499-2013>, 2013.
- 375 Benediktsson, Í., Aradóttir, N., Ingólfsson, Ó. and Brynjólfsson, S.: Cross-cutting palaeo-ice streams in NE-Iceland reveal shifting Iceland Ice Sheet dynamics, *Geomorphology*, 396, 1-16. <https://doi.org/10.1016/j.geomorph.2021.108009>, 2022.
- Blunier, T., & Brook, E. J.: Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial period. *Science*, 291(5501), 109-112, <https://doi.org/10.1126/science.291.5501.109>, 2001.
- 380 Booth, D.B., and Hallet, B.: Channel networks carved by subglacial water: Observations and reconstruction in the eastern Puget Lowland of Washington, *Geological Society of America Bulletin*, 105, 671-683, 1993.
- Booth, D.B., Troost, K.G., and Hagstrum, J.T.: Deformation of quaternary strata and its relationship to crustal folds and faults, south-central Puget Lowland, Washington State, *Geology*, 32:6, 505-508, <https://doi.org/10.1130/G20355>, 2004.
- Booth, D.B., Troost, K.G., Clague, J.J., & Waitt, R.B.: The cordilleran ice sheet. *Development in Quaternary Science*, 1, 17-43, 2003.
- 385 Broecker, W. S.: Massive iceberg discharges as triggers for global climate change, *Nature*, 372(6505), 421-424, <https://doi.org/10.1038/372421a0>, 1994.
- Cheng, H., Edwards, R. L., Sinha, A., Spötl, C., Yi, L., Chen, S., ... & Zhang, H.: The Asian monsoon over the past 640,000 years and ice age terminations, *Nature*, 534(7609), 640-646, <https://doi.org/10.1038/nature18591>, 2016
- 390 Clague, J.J., & James, T.S. (2002). History and isostatic effects of the last ice sheet in southern British Columbia. *Quaternary Science Review*, 21(1-3), 71-87.
- Clallam County, Olympic Department of Natural Resources, Washington Department of Transportation: Puget Lowlands 2005 [datafile]. Retrieved from <https://lidarportal.dnr.wa.gov/#47.85003:-122.92053:7>, 2008.
- Clark, C.D.: Reconstructing the evolutionary dynamics of former ice sheets using multi-temporal evidence, remote sensing and GIS, *Quaternary Science Reviews*, 16:9, 1067-1092, [https://doi.org/10.1016/S0277-3791\(97\)00037-1](https://doi.org/10.1016/S0277-3791(97)00037-1), 1997.
- 395 Clark, C.D.: Glaciodynamic context of subglacial bedform generation and preservation, *Annals of Glaciology*, 28, 23-32, <https://doi.org/10.3189/172756499781821832>, 1999.

Formatted: Indent: Left: 0", Hanging: 0.5"

Formatted: Font: (Default) Times New Roman, Font color: Text 1

Deleted: ¶

Formatted: Font: (Default) Times New Roman, Font color: Text 1

Formatted: Font: (Default) Times New Roman, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Text 1

Formatted: Font: (Default) Times New Roman, Font color: Text 1

Formatted: Font: (Default) Times New Roman, Font color: Text 1

Formatted: Font: (Default) Times New Roman, Font color: Text 1

Formatted: Font: (Default) Times New Roman, Font color: Text 1

Formatted: Font: (Default) Times New Roman, Font color: Text 1

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Text 1

Deleted: ¶

Formatted: Font color: Text 1

- 400 Clark, C.D., Evans, D.J.A., and Piotrowski, J.A.: Palaeo-ice streams: an introduction, *Boreas*, 32:1, 1-3, <https://doi.org/10.1080/03009480310001182>, 2003.
- Clark, C.D., Hughes, A.L.C., Greenwood, S.L., Spagnolo, M., and Ng, F.S.L.: Size and shape characteristics of drumlins, derived from a large sample, and associated scaling laws, *Quaternary Science Reviews*, 28:7-8, 677-696, <https://doi.org/10.1016/j.quascirev.2008.08.035>, 2009.
- 405 Cuffey, K.M., and Paterson, W.S.B.: *The Physics of Glaciers: Fourth Edition*, Academic Press, Burlington, Massachusetts, ISBN: 9780123694614, 2010.
- Damsgaard, A., Goren, L., & Suckale, J.: Water pressure fluctuations control variability in sediment flux and slip dynamics beneath glaciers and ice streams. *Communications Earth & Environment*, v. 1, no. 1, p. 66. <https://doi.org/10.1038/s43247-020-00074-7>, 2020.
- 410 Dethier, D.P., Pessl, F., Keuler, R.F., Balzarini, M.A., and Pevear, D.R.: Late Wisconsinan glaciomarine deposition and isostatic rebound, northern Puget Lowland, Washington, *Geological Society of America Bulletin*, 107:11, [https://doi.org/10.1130/0016-7606\(1995\)1072.3.CO;2](https://doi.org/10.1130/0016-7606(1995)1072.3.CO;2), 1995.
- Durand, G., Gagliardini, O., Favier, L., Zwinger, T., and le Meur, E.: Impact of bedrock description on modeling ice sheet dynamics, *Geophysical Research Letters*, 38:20, <https://doi.org/10.1029/2011GL048892>, 2011.
- 415 Ehlers, J., Gibbard, P.L., and Hughes, P.D.: *Quaternary Glaciations – Extent and Chronology*, Elsevier, <http://booksite.elsevier.com/9780444534477/index>, 2010.
- Eyles, N., Arbelaez Moreno, L., and Sookhan, S.: Ice streams of the Late Wisconsin Cordilleran Ice Sheet in western North America, *Quaternary Science Reviews*, 179, 87-122, <https://doi.org/10.1016/j.quascirev.2017.10.027>, 2018.
- Eyles, N., and Doughty, M.: Glacially-streamlined hard and soft beds of the paleo-Ontario ice stream in Southern Ontario and New York state, *Sedimentary Geology*, 338, 51-71, <https://doi.org/10.1016/j.sedgeo.2016.01.019>, 2016.
- 420 Favier, L., Pattyn, F., Berger, S., and Drews, R.: Dynamic influence of pinning points on marine ice-sheet stability: a numerical study in Dronning Maud Land: East Antarctica, *European Geosciences Union*, 10:6, 2623-2635, <https://doi.org/10.5194/tc-10-2623-2016>, 2016.
- Greenwood, S.L., Simkins, L.M., Winsborrow, M.C.M., & Bjarnadóttir, L.R.: Exceptions to bed-controlled ice sheet flow and retreat from glaciated continental margins worldwide. *Science Advances*, 7(3), <https://www.science.org/doi/full/10.1126/sciadv.abb6291>, 2021.
- 425 Holschuh, N., Christianson, K., Paden, J., Alley, R. B. & Anandakrishnan, S.: Linking postglacial landscapes to glacier dynamics using swath radar at Thwaites Glacier. *Antarctica. Geology*, p. 1–5. doi:10.1130/g46772.1., 2020.
- Khazaradze, G., Qamar, A., and Dragert, H.: Tectonic deformation in western Washington from continuous GPS measurements, *Geophysical Research Letters*, 26:20, 3153-3156, <https://doi.org/10.1029/1999GL010458>, 1999.
- 430 King, E., Hindmarsh, R., and Stokes, C.: Formation of mega-scale glacial lineations observed beneath a West Antarctic ice stream, *Nature Geoscience*, 2 : 8, 585-588, <https://doi.org/10.1038/ngeo581>, 2009.

Deleted: ¶

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ¶

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

- 435 Kleman, J., and Borgström, I.: Reconstruction of palaeo-ice sheets: The use of geomorphological data, *Earth Surface Processes and Landforms*, 21:10, 893-909, [https://doi.org/10.1002/\(SICI\)1096-9837\(199610\)21:103.0.CO;2-U](https://doi.org/10.1002/(SICI)1096-9837(199610)21:103.0.CO;2-U), 1996.
- Kleman, J., Hättestrand, C., Stroeven, A.P., Jansson, K.N., Angelis, H.D. and Borgström, I.: Reconstruction of Palaeo-Ice Sheets- Inversion of their Glacial Geomorphological Record, *Glacier Science and Environmental Change*, 192-198, <https://doi.org/10.1002/9780470750636.ch38>, 2006.
- 440 Kovanen, D.J., and Slaymaker, O.: Relict Shorelines and Ice Flow Patterns of the Northern Puget Lowland from Lidar Data and Digital Terrain Modelling, *Geografiska Annaler, Series A, Physical Geography*, 86:4, 385-400, [jstor.org/stable/3566155](https://doi.org/10.1002/9780470750636.ch38), 2004.
- Krabbendam, M., Eyles, N., Putkinen, N., Bradwell, T., and Arbelaez-Moreno, L.: Streamlined hard beds formed by palaeo ice stream: a review, *Sedimentary Geology*, 338, 24-50, <https://doi.org/10.1016/j.sedgeo.2015.12.007>, 2016.
- 445 MacKie, E. J., Schroeder, D. M., Caers, J., Siegfried, M. R., and Scheidt, C.: Antarctic topographic realizations and geostatistical modeling used to map subglacial lakes, *Journal of Geophysical Research: Earth Surface*, 125:3, <https://doi.org/10.1029/2019JF005420>, 2020.
- MacKie, E.J., Schroeder, D.M., Zuo, C., Yin, Z., and Caers, J.: Stochastic modeling of subglacial topography exposes uncertainty in water routing at Jakobshavn Glacier, *Journal of Glaciology*, 67:261, 75-83, <https://doi.org/10.1017/jog.2020.84>, 2021.
- 450 McIntyre, N.F.: The Dynamics of Ice-Sheet Outlets: *Journal of Glaciology*, 31:108, 99-107. <https://doi.org/10.1017/S0022143000006328>, 1985.
- McKenzie, M.A., Simkins, L.M., Principato, S., and Munevar-Garcia, S.: Subglacial bedform sensitivity to bed characteristics across the deglaciated Northern Hemisphere, *Earth Surface Processes and Landforms*, 47:9, 2341-2356, <https://doi.org/10.1002/esp.5382>, 2022.
- 455 McKenzie, M.A., Simkins, L.M., Slawson, J.S., and Wang, S.: Streamlined subglacial bedforms across isolated topographic highs in the Puget Lowland, Washington state, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.954529>, 2023.
- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., et al.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13:2, 132– 137, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- OCM Partners: 2017 USGS Lidar: Olympic Peninsula, WA from 2010-06-15 to 2010-08-15, NOAA National Centers for Environmental Information [datafile], <https://www.fisheries.noaa.gov/inport/item/59232>, 2019a.
- OCM Partners: 2019 WA DNR Lidar: San Juan County, WA from 2010-06-15 to 2010-08-15, NOAA National Centers for Environmental Information [datafile], <https://www.fisheries.noaa.gov/inport/item/67199>, 2019b.
- 465 Payne, A., and Dongelmans, P.: Self-organization in the thermomechanical flow of ice sheets, *Journal of Geophysical Research B: Solid Earth*, 102:6, 12219-12233, <https://doi.org/10.1029/97JB00513>, 1997.

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Pohjola, V.A., and Hedfors, J.: Studying the effects of strain heating on glacial flow within outlet glaciers from the Heimefrontfjella Range, Dronning Maud Land, Antarctica, *Annals of Glaciology*, 37, 134-142, <https://doi.org/10.3189/172756403781815843>, 2003.

Principato, S., Moyer, A., Hampsch, A. and Ipsen, H.: Using GIS and streamlined landforms to interpret palaeo-ice flow in northern Iceland, *Boreas*, 45:3, 470– 482, <https://doi.org/10.1111/bor.12164>, 2016.

Quantum Spatial, Inc.: Western Washington 3DEP LiDAR, United States Geological Survey [datafile], https://gismaps.snoco.org/metadata/topography/Western_Washington_3DEP_Technical_Data_Report.pdf, 2017.

Quantum Spatial, Inc.: Olympic Peninsula , Washington 3DEP LiDAR, United States Geological Survey [datafile], https://coast.noaa.gov/htdata/lidar3_z/geoid18/data/9072/supplemental/Olympic_Peninsula_3DEP_Area_1_LiDAR_Technical_Data_Report_revised_110419.pdf, 2019.

Robel, A. A., Pegler, S. S., Catania, G., Felikson, D., and Simkins, L. M.: Ambiguous stability of glaciers at bed peaks, *Journal of Glaciology*, 1-8, <https://doi.org/10.1017/jog.2022.31>, 2022.

Schoof, C., and Clarke, G.: A model for spiral flows in basal ice and the formation of subglacial flutes based on a Reiner-Rivlin rheology for glacial ice: *Journal of Geophysical Research: Solid Earth*, v. 113 no. 5, p. 1-12. <https://doi.org/10.1029/2007JB004957>, 2008.

Shaw, J., Pugin, A., and Young, R.R.: A meltwater origin for Antarctic shelf bedforms with special attention to megalineations, *Geomorphology*, 102:3-4, 364-375, [https://doi.org/10.1016/0037-0738\(89\)90114-0](https://doi.org/10.1016/0037-0738(89)90114-0), 2008.

Sherrod, B. L., Blakely, R.J., Weaver, C.S., Kelsey, H.M., Barnett, E., Liberty, L., Meagher, K.L., and Pape, K.: Finding concealed active faults: Extending the southern Whidbey Island fault across the Puget Lowland, Washington. *Journal of Geophysical Research*, 113, B05313, 2008.

Schroeder, D. M., Blankenship, D. D., Young, D. A., Witus, A. E., & Anderson, J. B. Airborne radar sounding evidence for deformable sediments and outcropping bedrock beneath Thwaites Glacier, West Antarctica. *Geophysical Research Letters*, v. 41, no. 20, p. 7200–7208. <https://doi.org/10.1002/2014GL061645>, 2014.

Spagnolo, M., Clark, C.D., Ely, J.C., Stokes, C.R., Anderson, J.B., Andreassen, K., ... and King, E.C.: Size, shape and spatial arrangement of mega-scale glacial lineations from a large and diverse dataset, *Earth Surface Processes and Landforms*, 39:11, 1432-1448, <https://doi.org/10.1002/esp.3532>, 2014.

Spagnolo, M., Clark, C.D., and Hughes, A.L.C.: Drumlin relief, *Geomorphology*, 153-154, 179-191, <https://doi.org/10.1016/j.geomorph.2012.02.023>, 2012.

Stokes, C.R., Clark, C.D., Lian, O.B., and Tulaczyk, S.: Ice stream sticky spots: A review of their identification and influence beneath contemporary and palaeo-ice streams. *Earth-Science Reviews*, 81, 3-4, 217-249. <https://doi.org/10.1016/j.earscirev.2007.01.002>, 2007.

Walczak MH, Mix AC, Cowan EA, Fallon S, Fifield LK, Alder JR, Du J, Haley B, Hobern T, Padman J, Praetorius SK, Schmittner A, Stoner JS, Zellers SD.: Phasing of millennial-scale climate variability in the Pacific and Atlantic Oceans, *Science*, 370(6517), 716-720. doi: 10.1126/science.aba7096, 2020.

Deleted: 20

Deleted: 08,

Deleted:

Formatted: Font color: Text 1

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Text 1

Deleted: ¶

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Indent: Left: 0", Hanging: 0.5"

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Deleted: ¶

Formatted: Font color: Text 1

Wang, S., Wu, Q., and Ward, D.: Automated delineation and characterization of drumlins using a contour tree approach, *Int J Appl Earth Obs Geoinformation*, 62, 144-156, <https://doi.org/10.1016/j.jag.2017.06.006>, 2017.

Winsborrow, M.C.M., Clark, C.D., and Stokes, C.R.: What controls the location of ice streams?, *Earth-Science Reviews*, 103, 45-59, <https://doi.org/10.1016/j.earscirev.2010.07.003>, 2010b.

510

515

Deleted: ¶

Formatted: Font color: Text 1





Figure 1: (A) Site overview across the Puget Lowland with inset reference of CIS glaciation (14 kya; Ehlers et al., 2010). Site maps for i) the Blue Hills, ii) Devils Mountain, iii) Cougar Mountain, iv) Dow Mountain, v) Big Skidder Mountain, vi) Fidalgo Island, vii) San Juan Island, viii) the Black Hills, and ix) Lopez Island. Streamlined subglacial bedforms are mapped upstream (gray polygons), on top of (black polygons), and downstream (blue polygons) of bed bumps (small white outlines) within larger site regions (large white outlines). (B) Relative volume and surface area of bed bump sites.

Formatted: Font color: Text 1

Formatted: Font color: Text 1

520 **Figure 1.** A) Site overview across the Puget Lowland with inset reference of CIS glaciation (14kya; Ehlers et al., 2010). Site maps for i) the Blue Hills, ii) Devils Mountain, iii) Cougar Mountain, iv) Dow Mountain, v) Big Skidder Mountain, vi) Fidalgo Island, vii) San Juan Island, viii) the Black Hills, and ix) Lopez Island. Streamlined subglacial bedforms are mapped upstream (gray polygons), on top of (black polygons), and downstream (blue polygons) of bed bumps (small white outlines) within larger site regions (large white outlines). B) Relative volume and surface area of bed bump sites.




Figure 2: Scatterplots of the log of bedform elongation ratio and surface relief in meters. The ellipses are 1σ (68%) confidence levels for multivariate t-distributions for all bedforms ($n=3,273$). Sites are listed in the legend from largest surface area ((i) Blue Hills) to smallest surface area ((ix) Lopez Island).

525

Figure 2. Scatterplots of the log of bedform elongation ratio and surface relief in meters. The ellipses are 1σ (68%) confidence levels for multivariate t-distributions for all bedforms ($n=3,273$). Sites are listed in the legend from largest surface area ((i) Blue Hills) to smallest surface area ((ix) Lopez Island).

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1
 Formatted: Font color: Text 1




Figure 3: Box plots of A) bedform elongation data at each site with populations characterized upstream of, on top of, and downstream of bumps and B) bedform surface relief data at each site with populations characterized upstream of, on top of, and downstream of bumps. Statistically significant differences between groups are indicated by asterisks. Multiple asterisks indicate a separate population with significant differences, independent of other groups of statistical significance.

530 **Figure 3.** Box plots of A) bedform elongation data at each site with populations characterized upstream of, on top of, and downstream of bumps and B) surface relief data at each site with populations characterized upstream of, on top of, and downstream of bumps. Statistically significant differences between groups are indicated by asterisks. Multiple asterisks indicate a separate population with significant differences, independent of other groups of statistical significance.





Figure 4: (A) Density curves of bedform elongation upstream, on top of, and downstream of nine bumps. Density curves with color are less than 4 square kilometers, all other sites are shown in gray. (B) Density curves of bedform surface relief in meters upstream, on top of, and downstream of nine bumps. Density curves with color are less than 4 square kilometers in surface area, all other sites are shown in gray. (C) Cardinal orientations of streamlined bedforms upstream, on top of, and downstream of nine bumps.

535 **Figure 4. A) Density curves of bedform elongation upstream, on top of, and downstream of nine bumps. Density curves with color are less than 4 square kilometers, all other sites are shown in gray. B) Density curves of bedform surface relief in meters upstream, on top of, and downstream of nine bumps. Density curves with color are less than 4 square kilometers in surface area, all other sites are shown in gray. C) Cardinal orientations of streamlined bedforms upstream, on top of, and downstream of nine bumps.**

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Page 3: [1] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [2] Deleted Marion A. McKenzie 5/4/23 11:46:00 AM

Page 3: [3] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [4] Deleted Marion A. McKenzie 5/4/23 11:55:00 AM

Page 3: [5] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [6] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [7] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [8] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [9] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [10] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [11] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [12] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [13] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [14] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Page 3: [15] Formatted Marion A. McKenzie 4/27/23 9:21:00 AM

Font color: Text 1

Font color: Text 1

▲
Page 3: [17] Formatted **Marion A. McKenzie** **4/27/23 9:21:00 AM**

Font color: Text 1

▲
Page 3: [18] Formatted **Marion A. McKenzie** **4/27/23 9:21:00 AM**

Font color: Text 1

▲
Page 3: [19] Formatted **Marion A. McKenzie** **4/27/23 9:21:00 AM**

Font color: Text 1

▲
Page 3: [20] Formatted **Marion A. McKenzie** **4/27/23 9:21:00 AM**

Font color: Text 1

▲
Page 3: [21] Formatted **Marion A. McKenzie** **4/27/23 9:21:00 AM**

Font color: Text 1

▲
Page 3: [22] Formatted **Marion A. McKenzie** **4/27/23 9:21:00 AM**

Font color: Text 1

▲
Page 3: [23] Formatted **Marion A. McKenzie** **4/27/23 9:21:00 AM**

Font color: Text 1

▲
Page 3: [24] Deleted **Marion A. McKenzie** **4/24/23 10:46:00 AM**

✖

▲