

Global evaluation of Doppler velocity errors of EarthCARE Cloud Profiling Radar using global storm-resolving simulation

Yuichiro Hagihara¹, Yuichi Ohno¹, Hiroaki Horie¹, Woosub Roh², Masaki Satoh², and Takuji Kubota³

¹Radio Research Institute, National Institute of Information and Communications Technology, Koganei, Tokyo 184-8795, Japan

²Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Chiba 277-8564, Japan

³Earth Observation Research Center, Japan Aerospace Exploration Agency, Tsukuba, Ibaraki 305-8505, Japan

Correspondence: Yuichiro Hagihara (hagihara@nict.go.jp)

Abstract. The Cloud Profiling Radar (CPR) on the Earth Clouds, Aerosol, and Radiation Explorer (EarthCARE) satellite is the first satellite-borne Doppler radar (EC-CPR). In our previous study, we examined the effects of horizontal (along-track) integration and simple unfolding methods on the reduction of Doppler errors in the EC-CPR observations, and those effects were evaluated using two limited scenes in limited latitude and low pulse repetition frequency (PRF) settings. In this study, the amount of data used was significantly increased, and the area of the data used was extended globally. Not only low PRF but also high PRF settings were examined. We calculated the EC-CPR-observed Doppler velocity from pulse-pair covariances using the radar reflectivity factor and Doppler velocity obtained from a satellite data simulator and a global storm-resolving simulation. The global data were divided into five latitudinal zones, and mean Doppler errors for 5 dBZ_e after 10 km integration were calculated. In the case of low PRF setting, the error without unfolding correction for the tropics reached a maximum of 2.2 m s⁻¹ and then decreased toward the poles (0.43 m s⁻¹). The error with unfolding correction for the tropics became much smaller at 0.63 m s⁻¹. In the case of high PRF setting, the error without unfolding correction for the tropics reached a maximum of 0.78 m s⁻¹ and then decreased toward the poles (0.19 m s⁻¹). The error with unfolding correction for the tropics was 0.29 m s⁻¹, less than half the value without the correction. The results of the analyses of the simulated data indicated that the zonal mean frequency of precipitation echoes was highest in the tropics and decreased toward the poles. Considering a limitation of the unfolding correction for discrimination between large upward velocity and large precipitation falling velocity, the latitudinal variation of the Doppler error can be explained by the precipitation echo distribution.

1 Introduction

The Earth Clouds, Aerosol and Radiation Explorer (EarthCARE; hereafter EC) is a joint satellite mission by the Japan Aerospace Exploration Agency (JAXA) and European Space Agency (ESA) that will carry a Cloud Profiling Radar (CPR), an ATmospheric LIDar (ATLID), a Multi Spectral Imager (MSI), and a Broad Band Radiometer (BBR). From the derived 3D cloud and aerosol scene profiles, heating rates and radiation flux profiles are systematically determined with a resolution of 100 km² (Illingworth et al., 2015). Active sensors of EC will be regarded as an evolutionary successor of the 94-GHz CloudSat

削除: Copyright statement. TEXT

CPR (Stephens et al., 2008) and the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO; Winker et al., 2009) lidar (Stephens et al., 2018).

35 Because of EC's low orbit (~400 km) and the EC-CPR's large antenna (2.5 m), it has a better sensitivity (~36 dBZ_e at the top-of-atmosphere (TOA)) than the CloudSat CPR (~30 dBZ_e) and can observe 98 % of radiatively significant ice clouds and 40 % of all stratocumulus clouds (Stephens et al., 2002; Hagihara et al., 2010). Moreover, the EC-CPR has the vertical Doppler measurement capability that the CloudSat CPR does not have. It will reveal, for the first time, the vertical motion of cloud particles globally. Such an entirely new dataset would improve the discrimination between clouds and precipitation (Ceccaldi et al., 2013; Kikuchi et al., 2017), as well as the retrieval of cloud microphysical parameters (Heymsfield et al., 2008).
40 Consequently, it should improve various parameterization schemes used in atmospheric models and the understanding of the processes related to cloud and precipitation (Roh and Satoh, 2014; Roh et al., 2017; Roh and Satoh, 2018; Hagihara et al., 2014; Mülmenstädt et al., 2020; Takahashi et al., 2021).

Vertical Doppler velocity estimation from space suffers from Doppler broadening and velocity folding or aliasing (e.g., Kobayashi et al., 2002; Sy et al., 2014). The EC-CPR measures Doppler velocities using the pulse-pair method. It measures phase shift of echoes from two successive transmitted pulses. Since the EC-CPR is a finite beamwidth on fast moving spaceborne platform, targets have a broad Doppler width, which causes a worsening of the correlation of the phase. Then, large Doppler errors are introduced. Hagihara et al. (2022; hereafter, H22) examined the effect of horizontal (along-track) integration and unfolding methods on the reduction of Doppler velocity measurement errors, in order to improve Doppler data processing in the JAXA standard algorithm. They obtained EC-CPR data simulated by a satellite data simulator, the Joint-Simulator (Hashino et al., 2016; Satoh et al., 2016; Roh et al., 2020) using a global storm-resolving simulation data with the Nonhydrostatic Icosahedral Atmospheric Model (NICAM; Tomita and Satoh, 2004; Satoh et al., 2008; Satoh et al., 2014). They evaluated the Doppler errors for each Z_e for two cases (cirrus clouds and precipitation). They found that the error was reduced by horizontal integration alone in the case of cirrus clouds, whereas the error became large without unfolding correction in addition to the horizontal integration in the case of precipitation.
55

In H22, the evaluation was limited to two scenes in the mid-latitudes of the Northern Hemisphere and a low pulse repetition frequency (PRF) setting. In this study, we used more data than in H22 and performed the evaluation on a global scale. We also adopted different PRF settings. In Sect. 2, the simulation methods for EC-CPR data, the horizontal integration and unfolding correction of Doppler velocity, and the CloudSat-observed data are described. In Sect. 3, we investigate the Doppler errors on a global scale. To examine the characteristics of each latitude, we separated the data into five latitudinal zones. Two PRF
60 modes were also included. The summary and conclusions are given in Sect. 4.

2 Data and Method

We utilized the global storm-resolving simulation data simulated by the NICAM with a 3.5 km horizontal resolution. Moreover, we obtained the simulated EC-CPR data using the data and the Joint-Simulator following H22. Note that attenuations of the gas and particle are considered in the calculation of the radar reflectivity factor, whereas Doppler velocity is the total velocity.

削除: of the hydrometeor echo, including reflectivity-weighted particle fall speed and vertical air motion. Those




Figure 1. Zonal mean frequency of hydrometeors obtained by (a) NICAM/J-Sim and (b) CloudSat observations for 19 June 2008.

70 of the hydrometeor echo, including reflectivity-weighted particle fall speed and vertical air motion. Our forward model is based
 on the single scattering assumption. There are some studies on multiple scattering using Monte Carlo methods (e.g., Matrosov
 et al. 2008; Battaglia and Tanelli, 2011). Especially the effect of multiple scattering to the Doppler velocity is discussed in
 Battaglia and Tanelli (2011). In this study, we focus on Doppler errors caused by Doppler broadening and folding, so we do
 not consider multiple scattering for simplicity. This issue will be the subject of future research. The simulated data were then
 75 calculated along an EC orbit and interpolated into the EC-CPR sampling interval (100 m in vertical and 500 m in horizontal).
 The radar reflectivity factor ($Z_{e, jsim}$) and Doppler velocity (V_{jsim}) curtain data were obtained (hereinafter referred to as
 “NICAM/J-Sim data”). In H22, only two scenes extracted from two orbits of data were used, but in this study, the amount of
 data used was significantly increased to 16 orbits of data, which is equivalent to one day of satellite tracks.

80 We note that there may be fast updrafts on the km or sub km scale. However, such events are rare globally and would be
 negligible in statistics such as latitudinal zonal means. This study focuses on global statistical results and therefore we use the
 NICAM. When higher horizontal resolution NICAM data becomes available, we would like to study similar evaluation with
 it.

In using the NICAM/J-Sim data, we first performed the following statistical analyses. We examined the zonal mean
 frequencies of hydrometeors obtained from the NICAM/J-Sim data and the CloudSat observations for 19 June 2008 (Fig. 1).
 85 We used the CloudSat Z_e (the standard geometrical profile of cloud product, 2B-GEOPROF) (Stephens et al., 2008) for
 comparison with $Z_{e, jsim}$. For the observed data, we defined the hydrometeor bin as where the cloud mask value is greater or
 equal than 20 from the CPR Level 2B-GEOPROF product, which means a weak, good, or strong echo detection (Marchand et
 al., 2008). These are estimated to gives an estimated false detection rate smaller than 5%. This value is adopted in many other
 CloudSat-based hydrometeor studies (e.g., Sassen & Wang 2008). The frequency of cloud occurrence at a given altitude was

書式変更: インデント: 最初の行: 1.42 字

書式変更: インデント: 最初の行: 1.42 字, 間隔 段落前: 6 pt, 段落後: 6 pt

削除: For the observed data, we used the CPR Level 2B-GEOPROF cloud mask product to extract bins with threshold ≥ 20 that are less affected by surface clutter and other factors. These are estimated to have a false-positive probability of 5% (Marchand et al., 2008).

95 defined as the number of cloud echo bins ($Z_e > -24$ dBZ_e) divided by the total number of observations at that level. The bin size was 240 m in vertical and 2.0° latitude in horizontal. The overall frequencies of the NICAM/J-Sim simulated cloud field are comparable to the results of the CloudSat observations.

We simulated the measured vertical Doppler velocity (V_m) as

$$V_m = V_{jsim} + V_{random}, \quad (1)$$

100 where V_{random} is the random error caused by the spread of Doppler velocities within the beam width. This is a Gaussian error distribution, and its SD of random error (SD_{random}) is determined by perturbation approximation (Doviak and Zrnich, 1993) as

$$SD_{random} = C \sqrt{\frac{\lambda^2}{32\pi^2 M \rho^2 \left(\frac{1}{PRF}\right)^2} \left[\left(1 + \frac{N}{S}\right)^2 - \rho^2 \right]}, \quad (2)$$

105 and C is a correction factor. We set $C = 1.3$ following H22. The wavelength is λ ($\lambda = 3.2$ mm for EC-CPR), M is the number of pulse pairs within an integration length, ρ is the correlation function, and S/N is the SNR. In nominal operation, the EC-CPR will change the observation window, that is, low mode (-1 to 16 km altitude) at latitudes of 60 to 90° and high mode (-1 to 20 km) at latitudes of 0 to 60°. The PRF is determined on the basis of the satellite altitude and changes in the range of 6100 to 7500 Hz with the latitude and observation window, as illustrated in Fig. 2. The high mode has a lower PRF and worse




Figure 2. Satellite altitude and PRF as functions of latitude and observation mode.

110 Doppler accuracy, as discussed in H22, although cloud echoes up to an altitude of 20 km can be observed. On the other hand, the low mode has a higher PRF and better Doppler accuracy, but cloud echoes higher than 16 km cannot be observed. M is 357 to 420 for 500 m integration depending on the PRF. The SNR is determined by the received echo power calculated from

削除: [1]

書式変更: 間隔 段落前: 0 pt, 段落後: 0 pt

下へ移動 [1]: [1]
<オブジェクト>Figure 2. Satellite altitude and PRF as functions of latitude and observation mode.

削除: [1]

書式変更: 英語 (米国)

移動 (挿入) [1]

削除:

120 the radar equation and estimated EC-CPR noise level. In the case of EC-CPR, the SNR is 0 dB, which is a signal equivalent to -21.2 dBZ_e echo intensity. If $Z_{e,j\text{sim}}$ is less than -24 dBZ_e, we assume the Doppler velocity of its echo to be random noise in this study. The correlation function ρ is defined as

$$\rho = \exp \left\{ -8 \left(\frac{\pi \sigma_v}{\lambda \cdot \text{PRF}} \right)^2 \right\}, \quad (3)$$

where σ_v is the total Doppler velocity spectrum width.

125 The width σ_v can be considered as a sum of contributions by each. That is,

$$\sigma_v^2 = \sigma_{sm}^2 + \sigma_{\sigma}^2 + \sigma_{psd}^2 \quad (4)$$

130 where σ_{sm} is the spread due to satellite motion, given by $\sigma_{sm} \sim 0.3 V_{\text{sat}} \theta_{\text{dB}}$, V_{sat} is the satellite velocity, and θ_{dB} is the beam width (Sloss and Atlas 1968). When V_{sat} is 7738 m/s and θ_{dB} is 0.00166 rad (0.095°), σ_{sm} becomes 3.85 m/s. The spread σ is due to turbulence and σ_{psd} to the distributions of hydrometeor falling velocities, respectively, which are assumed to be $\sigma = 1.0$ m/s (Amayenc et al., 1993), and $\sigma_{psd} = 0.5$ m/s (Gossard et al., 1997). As for the latter term, it is reported to spread to 1.0 m/s for rain (Lhermitte 1963). In this study, we assumed the $\sigma_{psd} = 0.5$ m/s so that σ_v becomes 4.01 m/s.

The EC-CPR measures the phase change of the echo between two successive pulses by pulse-pair processing to estimate the Doppler velocities. The real and imaginary parts of pulse-pair covariances R_τ integrated onboard corresponding to a 500 m along-track are simulated in this study as

135 $Re(R_\tau) = Z_{e,j\text{sim}} \cdot \cos \left(\frac{4\pi V_m}{\lambda \cdot \text{PRF}} \right), \quad (5)$

$Im(R_\tau) = Z_{e,j\text{sim}} \cdot \sin \left(\frac{4\pi V_m}{\lambda \cdot \text{PRF}} \right). \quad (6)$

V_{500m} is calculated using the arctangent of the real and imaginary parts of the 500-m-integrated R_τ simulated by Eqs. (5) and (6). The sign of Doppler velocity is defined as being those of radial Doppler velocity (i.e., downward motion is positive) following the EC-CPR data processing. To reduce random error, V_{1km} and V_{10km} are also calculated using 1 and 10 km horizontally integrated R_τ respectively, that are calculated from the 500 m-integrated R_τ .

140 Velocity folding or aliasing is inherent to Doppler radar. V_{max} can be measured by the pulse-pair method and is defined by PRF ($V_{\text{max}} = \lambda \cdot \text{PRF}/4$). In the PRF of the high mode (lower PRF), V_{max} ranges from 4.9 to 5.2 m s⁻¹, whereas in the PRF of the low mode (higher PRF), it ranges from 5.7 to 6.0 m s⁻¹.

145 The simulated EC-CPR Doppler velocities are required for unfolding correction. To correct the velocity folding in spaceborne radar, it is difficult to use the conventional unfolding method generally used by ground-based Doppler weather radar (e.g., Bargaen and Brown, 1980). From the ground-based vertically pointing cloud radar observations (Horie et al., 2000),

- 書式を変更: フォント: 10 pt
- 書式を変更: インデント: 最初の行: 1.42 字
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 書式を変更: フォント: 10 pt
- 削除: In this study, we assumed $\sigma = 4.01$ m s⁻¹.
- 削除: (4)
- 削除: (5)
- 削除: (4)
- 削除: (5)
- 削除: high-mode PRF
- 削除: low-mode PRF




Figure 3. Standard deviation of random error of simulated Doppler velocities for PRF of the high mode (lower PRF) as a function of Z_e for (a) 1 km integration and (b) 10 km integration. The solid lines denote the results with unfolding correction. The black lines indicate the precipitation case in Hagihara et al. (2022).

upward motion above 3 m s^{-1} was rarely observed. On the basis of this, we thus assumed that the echoes with velocities higher than 3 m s^{-1} are upward folded precipitation echoes. We used the simple unfolding method as follows:

$$V_{\text{unfolded}} = \begin{cases} V_{\text{folded}} + 2 \cdot V_{\text{max}} & \text{for } V_{1\text{km},10\text{km}} < -3\text{m/s} \\ V_{\text{folded}} & \text{otherwise.} \end{cases} \quad (7)$$

3 Results

We first evaluated the global mean Doppler errors in the PRF of the high mode (lower PRF) as well as PRF of the low mode (higher PRF). Then, we separated the NICAM/J-Sim data into five latitudinal zones (Arctic, Northern midlatitude, tropics, Southern midlatitude, and Antarctic). The Doppler errors for each latitudinal zone are investigated in both PRF modes.

Figure 3 shows the global mean Doppler errors in the PRF of the high mode (lower PRF). The vertical axis indicates the SD of Doppler error that is calculated from the difference between the simulated velocity (i.e., $V_{1\text{km}}$, $V_{10\text{km}}$) and V_{sim} (hereafter, SD_{diff}). The horizontal axis indicates Z_e of the NICAM/J-Sim data. The red dashed lines show SD_{diff} and the solid lines indicate SD_{diff} with unfolding correction using Eq. (7). Figure 3a shows SD_{diff} of $V_{1\text{km}}$ and SD_{diff} of $V_{1\text{km}}$ with unfolding correction. SD_{diff}

書式変更: インデント: 最初の行: 0 字

削除: (6)

削除: high-mode PRF


削除: low-mode PRF

削除: high-mode PRF

書式変更: インデント: 最初の行: 1.42 字

削除: (6)

of $V_{1\text{km}}$ decreases for Z_e below -10 dBZ_e. This is attributed to the reduction of random error owing to the increase in S/N and decrease in SD_{random} in Eq. (2) as Z_e increases. SD_{diff} of $V_{1\text{km}}$ increases for Z_e above -10 dBZ_e. This is due to the increase in the occurrence of velocity folding. That is, an increase in Z_e results in an increase in the intensity of precipitation echoes and an increase in mean fall velocity. When the unfolding method is applied, SD_{diff} of $V_{1\text{km}}$ is noticeably reduced because the folded negative velocities are corrected and the occurrence of the velocity folding is reduced. In Fig. 3b, SD_{diff} of $V_{10\text{km}}$ decreases for Z_e below -7 dBZ_e and increases for Z_e above -7 dBZ_e. SD_{diff} of $V_{10\text{km}}$ is much smaller than that of $V_{1\text{km}}$, reaching 0.8 m s⁻¹ for -9 dBZ_e. This is because of the increase in M and the decrease in SD_{random} in Eq. (2). If the unfolding method is applied, SD_{diff} of $V_{10\text{km}}$ becomes smaller since the effect of folding Doppler errors of precipitation echoes is reduced, as shown in Fig. 3a. For instance, SD_{diff} of $V_{10\text{km}}$ is less than 0.5 m s⁻¹ above -5 dBZ_e.



185 **Figure 4.** Standard deviation of random error of simulated Doppler velocities for PRF of the low mode (higher PRF) as a function of Z_e for (a) 1 km integration and (b) 10 km integration. The solid lines denote the results with unfolding correction.

What has been described so far is consistent with what was shown in the analysis of the precipitation case in H22. Note that PRF varied from 6106 to 6464 Hz in the high mode illustrated in Fig. 2 but was a single value of 6279 Hz in the

削除:
 <オブジェクト> Figure 3. Standard deviation of random error of simulated Doppler velocities for high-mode PRF as a function of Z_e for (a) 1 km integration and (b) 10 km integration. The solid lines denote the results with unfolding correction. The black lines indicate the precipitation case in Hagihara et al. (2022).⁴¹

削除: low-mode PRF

precipitation case in H22. Note the black dashed and solid lines in Fig. 3. In both Figs. 3a and 3b, the results in H22 are in good agreement with those of this study.

削除: Incidentally, we added black dashed and solid lines to Fig. 3 to show the results.

Figure 4 illustrates the global mean Doppler errors in the low-mode PRF. The dashed lines show SD_{diff} without unfolding correction and the solid lines indicate SD_{diff} with unfolding correction using Eq. (6). The PRF varies from 7156 to 7500 Hz, with a corresponding SD_{random} of 0.8 to 1.5, for 0 to -19 dBZ_e (see Fig. 2 in H22). On the other hand, in the high mode, the PRF varies from 6106 to 6464 Hz, with a corresponding SD_{random} of 1.5 to 3.4, for 0 to -19 dBZ_e. Similarly, V_{max} takes values between 5.7 and 6.0 m s⁻¹, whereas in the high mode, it is between 4.9 and 5.2 m s⁻¹. Comparison of Figs. 3 and 4 clearly shows that the Doppler error is much smaller in the latter because of SD_{random} described above. Furthermore, SD_{diff} without unfolding correction is smaller than that in the PRF of the high mode (lower PRF) because V_{max} is larger in addition to the effect of SD_{random} .

削除: 1.5 to 3.4

削除: 0.8 to 1.5

削除: high-mode PRF

Since the frequencies of cloud and precipitation echoes differ in latitude and the PRF varies with latitude, as shown in Fig. 2, we investigated the change in SD_{diff} with latitude. We defined five latitudinal zones, namely, Arctic (>60°), Northern midlatitude (60° to 30°), tropics (30° to -30°), Southern midlatitude (-30° to -60°), and Antarctic (<-60°). In the following analysis, we focused on SD_{diff} of V_{10km} . Figs. 5a-5e show the Doppler error for the five latitudinal zones in the PRF of the high mode (lower PRF). The dashed lines show SD_{diff} without unfolding correction and the solid lines indicate SD_{diff} with unfolding correction using Eq. (7). SD_{diff} of V_{10km} without unfolding correction decreases up to a certain value of Z_e and increases after that value. SD_{diff} with unfolding correction decreases as Z_e increases. These tendencies observed in the five latitudinal zones are similar to those of the global mean SD_{diff} of V_{10km} shown in Fig. 3b, although their magnitudes are not the same. We compared SD_{diff} without unfolding correction. SD_{diff} for the tropics, shown in Fig. 5c, has the largest value and is larger than the global mean result. The SD_{diff} values for both midlatitudes (Figs. 5b and 5d) are smaller than that for the tropics but slightly larger than or comparable to the global mean result. The SD_{diff} values for both polar regions (Figs. 5a and 5e) are even smaller than those for both midlatitudes and smaller than the global mean result. SD_{diff} for the Antarctic in Fig. 5e shows the smallest value. The tendency of the magnitude relation of SD_{diff} for each latitudinal zone was the similar between without and with unfolding correction. From the PRF variation shown in Fig. 2, in the PRF of the high mode (lower PRF), the Doppler accuracy should be higher in the tropics and lower toward the poles. However, the results we have seen so far are opposite. On the other hand, the frequency of precipitation echoes is considered to be the highest in the tropics, and the folding Doppler error may have resulted in the largest SD_{diff} in the tropics. This does not mean, however, that the mean Doppler velocity of the precipitation echo exceeds V_{max} . In H22, Fig. 9(a) shows a 2D-histogram of V_{sim} without the random error as a function of the Z_e for the precipitation case. Large fall velocities are not seen due to Mie scattering of the larger drops at 94-GHz. As shown in Fig. 9(b-d) in H22, considering the random error due to the Doppler broadening, the velocity folding occurs.

削除: high-mode PRF

削除: (6)

削除: resulting

削除: being

書式を変更: 英語 (米国)

削除: low-mode PRF

書式変更: インデント: 最初の行: 1.42 字

削除: (6)

Figures 5f-5j demonstrate the Doppler error for the five latitudinal zones in the PRF of the low mode (higher PRF). The dashed lines show SD_{diff} without unfolding correction and the solid lines indicate SD_{diff} with unfolding correction using Eq. (7). Similarly to Figs. 3 and 4, comparison of Figs. 5a-5e and 5f-j shows that SD_{diff} is much smaller in the latter. There is a difference between with and without unfolding correction only for SD_{diff} for the tropics shown in Fig. 5h, but not for the others.




Figure 5. Standard deviation of random error of simulated Doppler velocities for (a-e) PRF of the high mode (lower PRF) and (f-i) PRF of the low mode (higher PRF) as a function of Z_e after 10 km integration for (a, f) Arctic, (b, g) Northern midlatitude, (c, h) tropics, (d, i) Southern midlatitude, and (e, j) Antarctic zones. The solid lines denote the results with unfolding correction.

This may be related to the frequency of precipitation echoes, as also explained in Figs. 5a–5e. In the low-mode PRF, V_{\max} is larger and SD_{random} is smaller owing to the higher PRF.

To summarize what has been discussed so far, the SD_{diff} values for the five latitudinal zones for 5 dBZ_e were extracted and shown in Fig. 6. The red crosses indicate SD_{diff} without unfolding correction of the high-mode PRF, and the red circles denote SD_{diff} with unfolding correction using Eq. (6). The red dashed line is SD_{diff} for 5 dBZ_e without unfolding correction, and the red solid line is that with unfolding correction shown in Fig. 3b. SD_{diff} without unfolding correction (red crosses) for the tropics is the largest at 2.2 m s⁻¹ and decreases in both polar directions, with the smallest value at 0.43 m s⁻¹ in the Antarctic. The SD_{diff} values for the Northern midlatitude and Arctic are slightly larger than those for the Southern midlatitude and Antarctic. In comparison with the global mean SD_{diff} without unfolding correction, the values for the tropics and




Figure 6. Standard deviation of random error of Doppler velocities with and without unfolding correction for 5 dBZ_e after 10 km integration as a function of latitude.

Northern midlatitude are larger, but the other values are smaller. SD_{diff} with unfolding correction (red circles) for the tropics is 0.63 m s⁻¹, which is above the global mean result of 0.54 m s⁻¹ in Fig. 3b. The SD_{diff} values with unfolding correction for the Southern midlatitude, Northern midlatitude, and Arctic are comparable to the global mean result, but the value for the Antarctic is smaller than the global mean result. Next, we examine the PRF of the low mode (higher PRF) results. The blue crosses indicate SD_{diff} without unfolding correction of the PRF of the low mode (higher PRF), and the blue circles denote SD_{diff} with unfolding correction using Eq. (7). The blue dashed line is SD_{diff} for 5 dBZ_e without unfolding correction, and blue solid line is the value with unfolding correction illustrated in Fig. 4b. SD_{diff} without unfolding correction (blue crosses) for the tropics is

書式変更: インデント: 最初の行: 0 字

削除: <オブジェクト>Figure 5. Standard deviation of random error of simulated Doppler velocities for (a–e) high-mode and (f–i) low-mode PRF as a function of Z_e after 10 km integration for (a, f) Arctic, (b, g) Northern midlatitude, (c, h) tropics, (d, i) Southern midlatitude, and (e, j) Antarctic zones. The solid lines denote the results with unfolding correction.

書式変更: インデント: 左: 0 mm, 最初の行: 0 字, 行間: 1.5 行, 句読点のぶら下げを行わない

削除: low-mode PRF

削除: low-mode PRF

削除: (6)

275 the largest at 0.78 m s^{-1} and decreases toward the poles, with the smallest value being 0.19 m s^{-1} at the Antarctic. SD_{diff} with
 unfolding correction (blue circles) for the tropics is 0.29 m s^{-1} , which is above the global mean of 0.22 m s^{-1} in Fig. 4b. The
 SD_{diff} values with unfolding correction for the other zones are comparable to the global mean result. As already explained in
 Figs. 5, the latitudinal variation of SD_{diff} without unfolding correction may be due to the frequency of precipitation echoes. If
 the unfolding correction were perfect, there would be no relationship between the latitudinal variation of SD_{diff} with unfolding
 correction and the frequency of precipitation echoes. However, there is actually a relationship between the two, which indicates
 280 a limitation of the unfolding correction.




Figure 7. Zonal mean frequency of precipitation echoes obtained by NICAM/J-Sim for 19 June 2008.

We examined the zonal mean frequencies of precipitation echoes obtained from the NICAM/J-Sim data for 19 June 2008.
 285 First, to obtain precipitation echoes, we used the same method as in Fig. 1a but added a Doppler velocity condition ($V_{\text{sim}} > 3$
 m s^{-1} , downward motion). Then, using the same bin size as in Fig. 1a, we obtained Fig. 7. The extracted precipitation echoes
 show that the frequency decreases at higher altitudes compared with that shown in Fig. 1a. The frequency is high in the tropics
 and decreases toward the poles. The frequencies at altitudes of less than 5 km were averaged by latitudinal zone and found to
 be as follows: 27.8 % in the Arctic, 60.3 % in the Northern midlatitude, 68.5 % in the tropics, 36.7 % in the Southern
 290 midlatitude, and 2.6 % in the Antarctic. This is because it was summer in the Northern Hemisphere in the simulation. The
 latitudinal variation of SD_{diff} described so far can be explained on the basis of the precipitation echo distribution.

削除: ↵

4 Conclusions

295 We examined the vertical Doppler velocity error due to Doppler broadening and velocity folding in the EarthCARE CPR (EC-CPR) observations throughout the globe. We used simulated observation data (NICAM/J-Sim Z_e , v_{sim} and V_{sim}) for 16 satellite orbits with the same sampling interval as the EC-CPR, obtained using the NICAM and a satellite data simulator, the Joint-Simulator. The EC-CPR observed 500 m horizontally integrated pulse-pair covariances and Doppler velocity. The 1 and 10 km horizontally integrated Doppler velocities were calculated from them. We evaluated the Doppler error, i.e., the standard deviation of random error (SD_{diff}), and investigated the effectiveness of error reduction by horizontal integration. We also
300 evaluated the Doppler folding error by comparing the corrected Doppler velocities using our simple unfolding method.

We first evaluated the global mean Doppler error in the PRF of the high mode (lower PRF) as well as the PRF of the low mode (higher PRF) and compared the results with those of our previous study. In the PRF of the high mode (lower PRF), SD_{diff} without unfolding correction for 1 km integration decreases up to a certain value of Z_e and increases after that value. This
305 decreasing feature is due to the decrease in the SD of random error as the SNR increases, and the increasing feature is the result of an increase in the frequency of the folded Doppler error of precipitation echoes. SD_{diff} without unfolding correction is much smaller for 10 km integration than for 1 km integration, because of the increased number of pulse pairs. When the unfolding correction is applied, SD_{diff} becomes considerably smaller regardless of the integration length and the PRF mode. The results of PRF of the low mode (higher PRF) (higher PRF) show very small Doppler error both without and with unfolding
310 correction.

To investigate the latitudinal variation of Doppler error, we separated the data into five latitudinal zones, namely, Arctic ($>60^\circ$), Northern midlatitude ($60^\circ - 30^\circ$), tropics (30° to -30°), Southern midlatitude (-30° to -60°), and Antarctic ($<-60^\circ$). In the present work, we focused on SD_{diff} for 10 km integration. In the PRF of the high mode (lower PRF), SD_{diff} for the tropics without unfolding correction is the largest and is larger than the global mean result. SD_{diff} without unfolding correction
315 decreases toward the poles with the smallest value for the Antarctic, which is smaller than the global mean. The tendency of the magnitude relation of SD_{diff} for each latitudinal zone was similar between without and with unfolding correction. The frequency of precipitation echoes is expected to be highest in the tropics, and the folding Doppler error is also likely to be the largest. Therefore, SD_{diff} for the tropics without unfolding correction is considered to be the largest. SD_{diff} is much smaller in the PRF of the low mode (higher PRF) than in the PRF of the high mode (lower PRF), as shown by the global mean results
320 described earlier.

In summary, SD_{diff} for the five latitudinal zones for 5 dBZ_e is described as follows. In the PRF of the high mode (lower PRF), SD_{diff} without unfolding correction for the tropics reached a maximum of 2.2 m s⁻¹ and then decreased toward the poles. SD_{diff} with unfolding correction for the tropics was much smaller at 0.63 m s⁻¹. In the PRF of the low mode (higher PRF), SD_{diff} without unfolding correction for the tropics reached a maximum of 0.78 m s⁻¹ and then decreased toward the poles. SD_{diff} with
325 unfolding correction for the tropics was 0.29 m s⁻¹, which is less than half the value without correction. As explained previously, the latitudinal variation of SD_{diff} can be attributed to the frequency of precipitation echoes. The zonal mean frequency of

削除: high-mode PRF

削除: low-mode PRF

削除: high-mode PRF

削除: low-mode PRF

削除: high-mode PRF

削除: low-mode PRF

削除: high-mode PRF

削除: high-mode PRF

削除: low-mode PRF

precipitation echoes obtained from the NICAM/J-Sim data was higher in the tropics and decreased toward the poles. Therefore, the latitudinal variation of SD_{diff} can be explained on the basis of the precipitation echo distribution.

We found that the Doppler error was higher in the tropics than in the other latitudes. In the tropics, the unfolding correction reduced the large Doppler errors more efficiently. However, there is also a limitation of the unfolding correction for discrimination between large upward velocity and large precipitation falling velocity. Comparison of the results of the low-mode and PRF of the high mode (lower PRF) settings showed that the Doppler error for the PRF of the low mode (higher PRF) setting was significantly reduced, although cloud echoes for altitudes higher than 16 km cannot be observed.

削除: high-mode PRF

削除: low-mode PRF

Data availability

CloudSat CPR data are available from the CloudSat data processing center (<http://www.cloudsat.cira.colostate.edu/>, last access: 12 November 2022). We can share the NICAM/J-Sim data. Please send the email (ws-roh@aori.u-tokyo.ac.jp) if interested.

Author contributions

Y.H. performed the data analysis and produced the final manuscript draft. Y.O. provided feedback on the analysis methods as well as on the manuscript draft. H.H. developed and maintained the algorithm code and provided feedback on the manuscript draft. W.R. performed the Joint-Simulator simulations and provided feedback on the manuscript draft. M.S. led the NICAM development. T.K. led the Joint-Simulator development and provided feedback on the manuscript draft.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

The authors would like to thank the members of the JAXA EarthCARE Science Team and CPR project. The computational resources were partly provided by the National Institute for Environmental Studies.

Financial support

This work has been supported by the National Institute of Information and Communications Technology. This research is part of the EarthCARE satellite study commissioned by the Japan Aerospace Exploration Agency, “Enhancement of the Joint Simulator for Satellite Sensors”. MS and WR were supported by the Grant-in-Aid for Scientific Research B (20H01967) and Program for Promoting Technological Development of Transportation of the Ministry of Land, Infrastructure, Transport and Tourism (MLIT).

References

- Amayenc, P., Testud, J., and Marzoug, M.: Proposal for a Spaceborne Dual-Beam Rain Radar with Doppler Capability, *J. Atmos. Ocean. Technol.*, **10**, 262–276, [https://doi.org/10.1175/1520-0426\(1993\)010<0262:PFASDB>2.0.CO;2](https://doi.org/10.1175/1520-0426(1993)010<0262:PFASDB>2.0.CO;2), 2002.
- Bargen, D. W. and Brown, R. C.: Interactive radar velocity unfolding, 19th Conference on Radar Meteorology, 278–285, 1980.
- 370 Battaglia, A. and Tanelli, S.: DOMUS: DOppler MULTiple-Scattering simulator, *IEEE Trans. Geosci. Remote Sens.*, **49**, 442–450, <https://doi.org/10.1109/TGRS.2010.2052818>, 2011.
- Ceccaldi, M., Delanoë, J., Hogan, R. J., Pounder, N. L., Protat, A., and Pelon, J.: From CloudSat-CALIPSO to EarthCare: Evolution of the DARDAR cloud classification and its comparison to airborne radar-lidar observations, *J. Geophys. Res. Atmos.*, **118**, 7962–7981, <https://doi.org/10.1002/jgrd.50579>, 2013.
- 375 Doviak, R. J. and Zrnic, D. S.: Doppler Radar and Weather Observations, Academic Press, 562 pp., 1993.
- Gossard, E. E., Snider, J. B., Clothiaux, E. E., Martner, B., Gibson, J. S., Kropfli, R. A., and Frisch, A. S.: The potential of 8-mm radars for remotely sensing cloud drop size distributions, *J. Atmos. Ocean. Technol.*, **14**, 76–87, [https://doi.org/10.1175/1520-0426\(1997\)014<0076:TPOMRF>2.0.CO;2](https://doi.org/10.1175/1520-0426(1997)014<0076:TPOMRF>2.0.CO;2), 1997.
- Hagihara, Y., Ohno, Y., Horie, H., Roh, W., Satoh, M., Kubota, T., and Oki, R.: Assessments of Doppler velocity errors of EarthCARE cloud profiling radar using global cloud system resolving simulations: Effects of Doppler broadening and folding, *IEEE Trans. Geosci. Remote Sens.*, **60**, 1–9, <https://doi.org/10.1109/TGRS.2021.3060828>, 2022.
- 380 Hagihara, Y., Okamoto, H., and Luo, Z. J.: Joint analysis of cloud-top heights from CloudSat and CALIPSO: New insights into cloud-top microphysics, *J. Geophys. Res. Atmos.*, **119**, 4087–4106, <https://doi.org/10.1002/2013JD020919>, 2014.
- Hagihara, Y., Okamoto, H., and Yoshida, R.: Development of a combined CloudSat-CALIPSO cloud mask to show global cloud distribution, *J. Geophys. Res. Atmos.*, **115**, D00H33, <https://doi.org/10.1029/2009JD012344>, 2010.
- 385 Hashino, T., Satoh, M., Hagihara, Y., Kubota, T., Matsui, T., Nasuno, T., and Okamoto, H.: Evaluating cloud microphysics from NICAM against CloudSat and CALIPSO, *J. Geophys. Res. Atmos.*, **118**, 7273–7292, <https://doi.org/10.1002/jgrd.50564>, 2013.
- Hashino, T., Satoh, M., Hagihara, Y., Kato, S., Kubota, T., Matsui, T., Nasuno, T., Okamoto, H., and Sekiguchi, M.: Evaluating arctic cloud radiative effects simulated by NICAM with A-train, *J. Geophys. Res.*, **121**, 7041–7063, <https://doi.org/10.1002/2016JD024775>, 2016.
- 390 Heymsfield, A. J., Protat, A., Austin, R. T., Bouniol, D., Hogan, R. J., Delanoë, J., Okamoto, H., Sato, K., van Zadelhoff, G. J., Donovan, D. P., and Wang, Z.: Testing IWC retrieval methods using radar and ancillary measurements with in situ data, *J. Appl. Meteorol. Climatol.*, **47**, 135–163, <https://doi.org/10.1175/2007JAMC1606.1>, 2008.
- 395 Horie, H., Iguchi, T., Hanado, H., Kuroiwa, H., Okamoto, H., and Kumagai, H.: Development of a 95-GHz airborne cloud profiling radar (SPIDER) - Technical aspects, *IEICE Trans. Commun.*, E83-B, 2010–2019, 2000.
- Illingworth, A. J., Barker, H. W., Beljaars, A., Ceccaldi, M., Chepfer, H., Clerbaux, N., Cole, J., Delanoë, J., Domenech, C., Donovan, D. P., Fukuda, S., Hirakata, M., Hogan, R. J., Huenerbein, A., Kollias, P., Kubota, T., Nakajima, T., Nakajima,

- 400 T. Y., Nishizawa, T., Ohno, Y., Okamoto, H., Oki, R., Sato, K., Satoh, M., Shephard, M. W., Velázquez-Blázquez, A.,
Wandinger, U., Wehr, T., and Van Zadelhoff, G. J.: The EarthCARE satellite: The next step forward in global
measurements of clouds, aerosols, precipitation, and radiation, *Bull. Am. Meteorol. Soc.*, 96, 1311–1332,
<https://doi.org/10.1175/BAMS-D-12-00227.1>, 2015.
- Kikuchi, M., Okamoto, H., Sato, K., Suzuki, K., Cesana, G., Hagihara, Y., Takahashi, N., Hayasaka, T., and Oki, R.:
405 Development of algorithm for discriminating hydrometeor particle types with a synergistic use of CloudSat and CALIPSO,
J. Geophys. Res. Atmos., 122, 11,022–11,044, <https://doi.org/10.1002/2017JD027113>, 2017.
- Kobayashi, S., Kumagai, H., and Kuroiwa, H.: A proposal of pulse-pair Doppler operation on a spaceborne cloud-profiling
radar in the W band, *J. Atmos. Ocean. Technol.*, 19, 1294–1306, [https://doi.org/10.1175/1520-0426\(2002\)019<1294:AOPPPD>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<1294:AOPPPD>2.0.CO;2), 2002.
- Lhermitte, R. M.: Motions of scatterers and the variance of the mean intensity of weather radar signals, Rep. SRRC-RR-63-57, Sperry Rand Research Center, 43 pp., 1963.
- 410 Marchand, R., Mace, G. G., Ackerman, T., and Stephens, G.: Hydrometeor detection using Cloudsat - An earth-orbiting 94-GHz cloud radar, *J. Atmos. Oceanic Technol.*, 25, 519–533, <https://doi.org/10.1175/2007JTECHA1006.1>, 2008.
- Matrosov, S. Y., Battaglia, A., and Rodriguez, P.: Effects of multiple scattering on attenuation-based retrievals of stratiform rainfall from CloudSat, *J. Atmos. Ocean. Technol.*, 25, 2199–2208, <https://doi.org/https://doi.org/10.1175/2008JTECHA1095.1>, 2008.
- 415 Mülmenstädt, J., Nam, C., Salzmann, M., Kretzschmar, J., L'Ecuyer, T. S., Lohmann, U., Ma, P. L., Myhre, G., Neubauer, D., Stier, P., Suzuki, K., Wang, M., and Quaas, J.: Reducing the aerosol forcing uncertainty using observational constraints on warm rain processes, *Sci. Adv.*, 6, 1–8, <https://doi.org/10.1126/sciadv.aaz6433>, 2020.
- Roh, W. and Satoh, M.: Evaluation of precipitating hydrometeor parameterizations in a single-moment bulk microphysics
420 scheme for deep convective systems over the tropical central Pacific, *J. Atmos. Sci.*, 71, 2654–2673,
<https://doi.org/10.1175/JAS-D-13-0252.1>, 2014.
- Roh, W. and Satoh, M.: Extension of a multisensor satellite radiance-based evaluation for cloud system resolving models, *J. Meteor. Soc. Japan*, 96, 55–63, <https://doi.org/10.2151/jmsj.2018-002>, 2018.
- Roh, W., Satoh, M., Hashino, T., Okamoto, H., and Seiki, T.: Evaluations of the thermodynamic phases of clouds in a cloud-
425 system-resolving model using CALIPSO and a satellite simulator over the Southern Ocean, *J. Atmos. Sci.*, 77, 3781–3801,
<https://doi.org/10.1175/JAS-D-19-0273.1>, 2020.
- Roh, W., Satoh, M., and Nasuno, T.: Improvement of a cloud microphysics scheme for a global nonhydrostatic model using TRMM and a satellite simulator, *J. Atmos. Sci.*, 74, 167–184, <https://doi.org/10.1175/JAS-D-16-0027.1>, 2017.
- Sassen, K. and Wang, Z.: Classifying clouds around the globe with the CloudSat radar: 1-year of results, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2007GL032591>, 2008.
- 430 35, <https://doi.org/10.1029/2007GL032591>, 2008.

書式を変更: 英語 (米国)

- Satoh, M., Matsuno, T., Tomita, H., Miura, H., Nasuno, T., and Iga, S.: Nonhydrostatic icosahedral atmospheric model (NICAM) for global cloud resolving simulations, *J. Comput. Phys.*, 227, 3486–3514, <https://doi.org/10.1016/j.jcp.2007.02.006>, 2008.
- 435 Satoh, M., Roh, W., and Hashino, T.: Evaluations of clouds and precipitations in NICAM using the Joint Simulator for Satellite Sensors, CGER's Supercomput. Monogr. Rep., 22, 110, <https://doi.org/CGER-I127-2016>, 2016.
- Satoh, M., Tomita, H., Yashiro, H., Miura, H., Kodama, C., Seiki, T., Noda, A. T., Yamada, Y., Goto, D., Sawada, M., Miyoshi, T., Niwa, Y., Hara, M., Ohno, T., Iga, S., Arakawa, T., Inoue, T., and Kubokawa, H.: The Non-hydrostatic Icosahedral Atmospheric Model: description and development, *Prog. Earth Planet. Sci.*, 1, 18, doi:10.1186/s40645-014-0018-1, 2014.
- 440 [Sloss, P. W. and Atlas, D.: Wind shear and reflectivity gradient effects on Doppler radar spectra, *J. Atmos. Sci.*, 25, 1080–1089, 1968.](#)
- Stephens, G. L., Vane, D. G., Boain, R. J., Mace, G. G., Sassen, K., Wang, Z., Illingworth, A. J., O'Connor, E. J., Rossow, W. B., Durden, S. L., Miller, S. D., Austin, R. T., Benedetti, A., and Mitrescu, C.: The CloudSat mission and the A-Train: A new dimension of space-based observations of clouds and precipitation, *Bull. Am. Meteorol. Soc.*, 83, 1771–1790, <https://doi.org/10.1175/bams-83-12-1771>, 2002.
- 445 Stephens, G. L., Vane, D. G., Tanelli, S., Im, E., Durden, S., Rokey, M., Reinke, D., Partain, P., Mace, G. G., Austin, R., L'Ecuyer, T., Haynes, J., Lebsock, M., Suzuki, K., Waliser, D., Wu, D., Kay, J., Gettelman, A., Wang, Z., and Marchand, R.: CloudSat mission: Performance and early science after the first year of operation, *J. Geophys. Res. Atmos.*, 114, 1–18, <https://doi.org/10.1029/2008JD009982>, 2008.
- Stephens, G., Winker, D., Pelon, J., Trepte, C., Vane, D., Yuhas, C., L'Ecuyer, T., and Lebsock, M.: CloudSat and CALIPSO within the A-Train: Ten years of actively observing the Earth system, *Bull. Am. Meteorol. Soc.*, 99, 569–581, <https://doi.org/10.1175/BAMS-D-16-0324.1>, 2018.
- Sy, O. O., Tanelli, S., Takahashi, N., Ohno, Y., Horie, H., and Kollias, P.: Simulation of EarthCARE spaceborne Doppler radar products using ground-based and airborne data: Effects of aliasing and nonuniform beam-filling, *IEEE Trans. Geosci. Remote Sens.*, 52, 1463–1479, <https://doi.org/10.1109/TGRS.2013.2251639>, 2014.
- 445 Takahashi, H., Luo, Z. J., and Stephens, G.: Revisiting the entrainment relationship of convective plumes: A perspective from global observations, *Geophys. Res. Lett.*, 48, 1–7, <https://doi.org/10.1029/2020GL092349>, 2021.
- Tomita, H. and Satoh, M.: A new dynamical framework of nonhydrostatic global model using the icosahedral grid, *Fluid Dyn. Res.*, 34, 357–400, <https://doi.org/https://doi.org/10.1016/j.fluiddyn.2004.03.003>, 2004.
- Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the 460 CALIPSO mission and CALIOP data processing algorithms, *J. Atmos. Ocean. Technol.*, 26, 2310–2323, <https://doi.org/10.1175/2009JTECHA1281.1>, 2009. Smith, A. A., Carter, C., and Miller, B. B.: More test articles, *J. Adv. Res.*, 35, 13–28, doi:10.2345/67890, 2014.