

# 1 Improving Statistical Projections of Ocean Dynamic Sea-level Change

## 2 Using Pattern Recognition Techniques.

3 Víctor Malagón-Santos<sup>1</sup>, Aimée B.A. Slangen<sup>1</sup>, Tim H.J. Hermans<sup>1,2</sup>, Sönke Dangendorf<sup>3</sup>, Marta  
4 Marcos<sup>4</sup>, Nicola Maher<sup>5,6,7</sup>.

5 <sup>1</sup>NIOZ Royal Netherlands Institute for Sea Research, Department of Estuarine & Delta Systems, P.O. Box 140, 4400 AC  
6 Yerseke, the Netherlands.

7 <sup>2</sup>University of Utrecht, Institute for Marine and Atmospheric research Utrecht (IMAU), Utrecht, The Netherlands

8 <sup>3</sup>Department of River-Coastal Science and Engineering, Tulane University, New Orleans, USA.

9 <sup>4</sup>Mediterranean Institute for Advanced Studies (IMEDEA), Spanish National Research Council-University of Balearic Islands  
10 (CSIC-UIB), Esporles, Spain.

11 <sup>5</sup>Cooperative Institute for Research in Environmental Science, University of Colorado, Boulder, CO, USA.

12 <sup>6</sup>Department of Atmospheric and Oceanic Sciences, University of Colorado, Boulder, CO, USA.

13 <sup>7</sup>Max Planck Institute for Meteorology, Hamburg, Germany.

14  
15  
16 *Correspondence to:* Víctor Malagón-Santos (victor.malagon.santos@nioz.nl)

17 **Abstract.** Regional emulation tools based on statistical relationships, such as pattern scaling, provide a computationally  
18 inexpensive way of projecting ocean dynamic sea-level change for a broad range of climate change scenarios. Such approaches  
19 usually require a careful selection of one or more predictor variables of climate change so that the statistical model is properly  
20 optimized. Even when appropriate predictors have been selected, spatiotemporal oscillations driven by internal climate  
21 variability can be a large source of statistical model error. Using pattern recognition techniques that exploit spatial covariance  
22 information can effectively reduce internal variability in simulations of ocean dynamic sea level, significantly reducing random  
23 errors in regional emulation tools. Here, we test two pattern recognition methods based on Empirical Orthogonal Functions  
24 (EOF), namely signal-to-noise maximising EOF pattern filtering and low-frequency component analysis, for their ability to  
25 reduce errors in pattern scaling of ocean dynamic sea-level change. We use the Max Planck Institute Grand Ensemble (MPI-  
26 GE) as a testbed for both methods, as it is a type of initial-condition large ensemble designed for an optimal characterization  
27 of the externally forced response. We show that the two methods tested here more efficiently reduce errors than conventional  
28 approaches such as a simple ensemble average. For instance, filtering only two realizations by characterising their common  
29 response to external forcing reduces the random error by almost 60%, a reduction that is only achieved by averaging at least  
30 12 realizations. We further investigate the applicability of both methods to single realization modelling experiments, including  
31 four CMIP5 simulations for comparison with previous regional emulation analyses. Pattern filtering leads to a varying degree  
32 of error reduction depending on the model and scenario, ranging from more than 20% to about 70% reduction in global-mean

33 root-mean-squared error compared with unfiltered simulations. Our results highlight the relevance of pattern recognition  
34 methods as a tool to reduce errors in regional emulation tools of ocean dynamic sea-level change, especially when one or only  
35 a few realizations are available. Removing internal variability prior to tuning regional emulation tools can optimize the  
36 performance of the statistical model, leading to substantial differences in emulated dynamic sea level compared to unfiltered  
37 simulations.

## 38 **1 Introduction**

39 Sea levels are closely linked to the state of the climate. Understanding how increased radiative forcing in the atmosphere will  
40 affect sea-level rise is of utmost importance given the devastating impacts to coastal systems. Global-mean sea level has been  
41 increasing over the 20<sup>th</sup> century (Fox-Kemper et al, 2021), and its rate has been accelerating over the past decades both globally  
42 (e.g., Dangendorf et al., 2019; Fox-Kemper, 2021; Frederikse et al., 2020; Nerem et al., 2006) and regionally (e.g., Steffelbauer  
43 et al., 2022). This acceleration is expected to continue over the next century for all greenhouse gas (GHG) emissions scenarios  
44 (Fox-Kemper et al., 2021) with the potential to further increase widespread impacts in coastal areas (Cooley et al., 2022).  
45 Increased sea levels will change coastal flood risk through expanding areas under permanent inundation, increasing frequencies  
46 of extreme coastal flooding events (Vitousek et al., 2017; Wahl et al., 2017), and modifying tides (Haigh et al., 2020) and thus  
47 potentially increasing the frequency of tidal-induced flooding (Moftakhari et al., 2015). These processes will not only impact  
48 coastal infrastructure and assets (Hinkel et al., 2014) but also alter coastal ecosystems and the services they provide, from  
49 ecosystem value to natural flood risk protection (Cooley et al., 2022). Understanding how global and regional sea levels evolve  
50 under different scenarios will help to better adapt to changing risks and mitigate their potential impacts in coastal zones  
51 (Haasnoot et al., 2019, 2021).

52 Global-mean sea-level change is driven by a combination of processes. The melting of the Greenland and Antarctica ice sheets  
53 and glaciers and ice caps, changes in land-water storage, and thermal expansion of the ocean are the processes driving global  
54 mean sea-level rise (e.g., Gregory et al., 2019; Fox-Kemper, 2021). Analogously to global warming, sea-level rise is a global  
55 concern but it is not spatially uniform (e.g., Slangen et al., 2017). There are several processes that determine regional sea-level  
56 change. First, the redistribution of mass on the Earth's surface, as a result of melting land ice and changes in land-water storage,  
57 causes a regionally variable sea-level change due to gravitational, rotational, and deformational effects (Farrell and Clark,  
58 1976; Mitrovica et al., 2001). Second, vertical land motion also causes relative sea-level changes. The viscoelastic relaxation  
59 of the Earth induced by deglaciation following the last glacial maximum, defined as glacial isostatic adjustment (GIA; e.g.,  
60 Peltier, 1999, 2001) and more local processes driving subsidence (e.g., Nicholls et al., 2021), are the main processes driving  
61 changes in land elevation. Third, ocean circulation, and heat and freshwater fluxes over the ocean, also known as ocean  
62 dynamics (Gregory et al., 2019), change local densities and move water mass around the ocean. Fourth, changes in sea-level  
63 pressure over the oceans, also known as inverted barometer (IB) effects, may lead to regionally varying rates of sea-level  
64 change (Stammer and Hüttemann, 2008). These regional drivers of sea-level change act on a wide range of spatial and temporal

65 scales, which makes their local assessment essential for impact studies, planning, and adaptation needs. For instance, while  
66 ocean dynamics have a typical temporal scale ranging from days to decades, vertical land movements presents a much wider  
67 range (Durand et al., 2022), as the latter is governed by processes affecting land elevation on significantly different timescales  
68 from earthquakes (on the order of seconds) to GIA (on the order of millennia). This study focuses on ocean dynamic sea-level  
69 (DSL) change, which is governed by changes in ocean circulation and density. DSL features large spatiotemporal variations  
70 across the oceans, which makes it a crucial component to predict regional sea-level changes accurately, yet also one that  
71 provides significant uncertainty (Couldrey et al., 2021). Spatial and temporal variability in DSL is driven by internal climate  
72 variability (ICV), which are defined as naturally occurring climatic variations controlled by interactions between different  
73 components of the Earth system (Hasselmann, 1976; Schwarzwald and Lenssen, 2022), and by a forced response associated  
74 with increased radiative forcing in the climate system. DSL is typically projected with Global Climate Models (or related  
75 models, hereinafter GCMs), which are state-of-the-art comprehensive climate models that solves a range of environmental  
76 variables controlling the Earth's system, including its climate. GCMs require vast computational resources, and therefore  
77 climate modelling experiments have been designed for a limited range of GHG concentration scenarios (O'Neill et al., 2017;  
78 Riahi et al., 2017; van Vuuren et al., 2011) within the climate model intercomparison (CMIP) framework (Eyring et al., 2016),  
79 so that model differences are somewhat comparable.

80 To reduce the computational demand, complementary approaches based on parameterizing process-based models are  
81 commonly used. This method, also known as emulation, aims to mimic the output of complex models at a reduced  
82 computational cost and has been widely used in recent literature to model different aspects of the climate system (e.g., Thomas  
83 and Lin, 2018; Edwards et al., 2021; Schwarber et al., 2019). Regional emulation follows the same principle and aims to  
84 estimate a spatiotemporal varying variable by mimicking GCMs behavior. One of the most commonly used emulation  
85 approaches for projecting changes in a regional variable is pattern scaling (Mitchell, 2003; Perrette et al., 2013; Santer et al.,  
86 1990), which consists of relating a local, grid-point variable (predictand) to one or a few global-mean change variables  
87 (predictors) via regression. Based on that statistical relationship, a change in a regional variable can be emulated by projecting  
88 the global-mean variables via simpler climate models (Goodwin et al., 2018; Meinshausen et al., 2011; Millar et al., 2017;  
89 Smith et al., 2018)

90 Here, we build on the approach proposed by Bilbao et al. (2015), who applied a linear pattern scaling approach to assess the  
91 ensemble mean DSL computed from five CMIP5 models and their simulations of several variables describing global changes,  
92 including Global Surface Air Temperature (GSAT), Global-Mean Thermosteric Sea-Level Rise (GMTSLR), and ocean-  
93 volume mean temperature. While GSAT turned out to be the best predictor of 21<sup>st</sup>-century DSL change in a high emissions  
94 scenario (Representative Concentration Pathway (RCP) 8.5), ocean-volume mean temperature and GMTSLR outperformed  
95 the rest of variables considered in lower emissions scenarios (RCP 2.6 and 4.5). As the surface ocean layer responds quicker  
96 to air temperature changes than the deeper ocean layer, they speculated that surface warming had a more important role relative  
97 to deep warming in a high emissions scenario. Based on Bilbao et al. (2015)'s findings, Yuan and Kopp (2021) used the same  
98 set of CMIP5 models to develop a bivariate pattern scaling approach, accounting for the surface and deep ocean layers

99 separately. Their goal was to capture the different delayed response of those two layers by using GSAT and global-mean deep  
100 ocean temperature changes as predictors. By employing a bivariate pattern scaling approach, Yuan and Kopp (2021) reported  
101 a reduction of the predicted DSL error for the period 2271-2290 of 36%, 24%, and 34% for RCP 2.6, 4.5, and 8.5, respectively,  
102 compared to a univariate approach based on only GSAT.

103 The aforementioned studies highlight the importance of selecting appropriate predictors to attain an optimized regional  
104 emulator of DSL, and how accounting for different processes driving DSL change (in different layers of the ocean) can help  
105 further improve emulator performance. While designing a regional emulator based on performance metrics may provide  
106 insights into the global processes driving DSL changes, this process can be obscured by other drivers of emulator error. In  
107 particular, random errors contained in the regression forming the pattern scaling approach, are assumed to be mostly caused  
108 by ICV (Bilbao et al., 2015) and may be a source of large uncertainty. Thus, if random errors are not minimized prior to  
109 emulator training with GCM simulations, their presence could impair a proper selection of global predictors, such that it would  
110 be uncertain whether an increase in model performance is due to an appropriate selection of predictors or an artifact of ICV  
111 causing a biased selection. In previous studies, this effect has been minimized by computing 30-year means, assuming this  
112 cancels out ICV. This step, however, entails a substantial loss of data and does not guarantee ICV is optimally subtracted, and  
113 residual ICV, for instance caused by long-memory processes (e.g., Becker et al., 2014; Dangendorf et al., 2014), can remain.  
114 We therefore propose to take a different approach to separate ICV from the response driven by external radiative forcing in  
115 the Earth, by employing state-of-the-art modelling experiments specifically designed to do so. These are known as Single-  
116 Model Initial Condition Large Ensembles (SMILES) and consist of a set of simulations with the same forcing but with the  
117 variability evolving in a different phase (Deser et al., 2020). These realizations can be combined through different methods  
118 (e.g., Frankcombe et al., 2015) so that ICV cancels out. However, conventional approaches such as computing the ensemble  
119 mean or linear trends are not the most efficient tools to do so and tend to lead to the loss of much of the information gained  
120 from running large ensembles (Wills et al., 2020). Other methods based on pattern recognition via Empirical Orthogonal  
121 Functions (EOFs) exploit spatial covariance information to remove ICV more efficiently (Wills et al., 2020) and have  
122 demonstrated to provide a superior agreement between observations and simulations than an ensemble average (Marcos and  
123 Amores, 2014). These types of efficient methods for removing ICV hold potential to benefit emulation experiments of DSL  
124 for which the number of simulations is limited.

125 The aim of this study is to characterise the importance of ICV as a driver of random errors in statistically based (pattern-scaled)  
126 projections of DSL change. To achieve this aim, we will compare different pattern recognition techniques, including Signal-  
127 to-Noise Maximising (S/N M) EOF pattern filtering (Wills et al., 2020) and Low Frequency Component Analysis (LFCA,  
128 Wills et al., 2018, 2020). We will use these techniques to truncate ICV in DSL simulations from the Max Planck Institute  
129 Grand Ensemble (MPI-GE) SMILE (Maher et al., 2019), and explore their applicability to single realization modelling  
130 experiments, including a set of CMIP5 simulations used in previous pattern scaling studies. In this paper, we particularly aim  
131 to attain the following objectives:

132

- 1) Use a large ensemble (MPI-GE) to determine the forced pattern and examine to which extent pattern recognition techniques isolate the forced response in DSL change more efficiently than conventional methods (Section 4.1)
- 2) Determine the error reduction in pattern scaling of DSL provided by pattern recognition methods relative to more conventional methods (Section 4.2).
- 3) Test whether filtering improves pattern scaling in single-realization modelling experiments of DSL (Section 4.3).

## 2 Climate model data and pre-processing

Separating ICV from the forced response is key for detection and attribution studies in climate change (Labe and Barnes, 2021) and to understand its effects on the climate system (Deser et al., 2020; Mankin et al., 2020). However, the combination of distinct GCMs to analyse ICV should be performed with caution, as this may conflate ICV with model biases (Maher et al., 2021b). In recent literature, this has motivated the development and use of SMILES, which branch each realization at a different model stage in the pre-industrial control simulation (Danabasoglu et al., 2020; Deser et al., 2020; Fasullo et al., 2020; Kay et al., 2015; Maher et al., 2019, 2021a; Mankin et al., 2020). This results in simulations with the same forced response but with variability evolving in a different phase, enabling a separation of the variability from the forced response.

There are two main procedures for creating SMILES: 1) inducing small round-off level differences in their atmospheric initial conditions (micro-initialization); 2) branching simulations at different times in the control simulation (macro-initialization). Both micro and macro initialization are useful to characterize unpredictable ICV within a model. Macro-initialization, however, provides larger differences in the initial states in both the atmosphere and ocean. Macro-initialized ensembles are therefore better suited than ‘micro’ ensembles to sample uncertainty in an initialized framework (Hawkins et al., 2016; Stainforth et al., 2007), facilitating an assessment of ICV in different aspects of the climate system.

Since we are assessing ocean processes, a macro-initialized ensemble is most suitable for the purpose of this study. From the available macro-initialized SMILES (Deser et al., 2020; Maher et al., 2021a), we decided to use the Max-Planck Institute Grand Ensemble (MPI-GE; Maher et al., 2019) because it contains the largest number of ensemble members available (100) in a SMILE for different RCP scenarios (RCP 2.6, 4.5, and 8.5) up to 2100. MPI-GE simulations assume a stationary and volcano free 1850 climate, and are macro-initialized on the first of January in different years of the control simulation (Table 1 in Maher et al., 2019). The branching separation between realizations varies along the pre-industrial control, ranging from 6 to 24 years and with a median of 16 years. MPI-GE has a relatively lower resolution than other GCMs, representing the atmosphere at an approximate horizontal resolution of 200 km (1.875 degrees) with 47 layers (up to 0.01 hPa ~ 80 km in height). The horizontal resolution of the ocean (including biogeochemistry) varies from 12 to 150 km at 40 layers, whereas the land biosphere has the same horizontal resolution as the atmosphere. Despite its relatively low resolution, Suarez-Gutierrez et al. (2021) show that MPI-GE samples observed ocean variability well in all regions except for the Southern Ocean.

Additionally, we use four CMIP5 models that were used in previous studies of DSL pattern scaling (Bilbao et al., 2015; Yuan and Kopp, 2021), including GISS-E2-R, HadGEM2-ES, IPSL-CM5A-LR, and MPI-ESM-LR. These four GCMs were selected

165 in the afore-mentioned studies because they were used to calibrate the parameters of the simple climate model used by Geoffroy  
166 et al. (2013a, b), which facilitated the design of their emulation tool. Also, these models provide multi-century data (up to  
167 2300) in three emissions scenarios, granting an assessment of the suitability of pattern scaling for long-term projections. We  
168 use them here for comparison purposes.

169 The focus of this study is on DSL change, which is defined at each location and time as the change in local sea-surface height  
170 relative to the geoid, with the IB correction applied (Gregory et al., 2019). DSL varies locally due to ocean circulation and  
171 horizontal gradients, and its global mean is zero at every time step (Eq. 15 in Gregory et al. 2019), i.e., GMTSLR is excluded.  
172 In CMIP models, DSL is diagnosed as ‘*zos*’ (Griffies et al., 2016) and often expressed as differences in relation to a control  
173 state. DSL simulations from GCMs, however, do not include the effect of sea-level pressure on sea level (IB effect) and such  
174 effect is not subject of study in our analysis, hence it is not considered here.

175 Since we are interested in assessing the forced response in DSL for historical and future GHG emissions we will use *zos* from  
176 a range of GCMs for historical and future radiative forcing scenarios, including RCP 2.6, 4.5, and 8.5 (Meinshausen et al.,  
177 2011). Once the forced DSL has been characterized, we will proceed to pattern scale each model and scenario using a variable  
178 from their respective GCM simulation representing a global change in the state of the climate system. Among other potential  
179 global predictors, we chose GMTSLR (diagnosed as ‘*zostoga*’ in CMIP models), defined as the part of global-mean sea-level  
180 rise due to thermal expansion. We deemed GMTSLR as a suitable predictor candidate because it is closely related to DSL and  
181 has been successfully used in previous pattern scaling analysis of DSL (e.g., Bilbao et al., 2015; Thomas and Lin, 2018). We  
182 refrain from testing other global variables as predictors to ease comparing models and scenarios, and determining to which  
183 extent pattern filtering reduces statistical error via reducing ICV.

184 In this study, we are particularly interested in removing interannual variability, thus we compute annual mean *zostoga* and *zos*  
185 time series from the raw monthly mean GCM data. In addition, since GCMs are run for a few centuries and the deep ocean  
186 usually takes millennia to reach an equilibrium, both *zos* and *zostoga* are subject to model drift (Sen Gupta et al., 2013). Model  
187 drift in the historical and scenario simulations can be corrected for by subtracting the smoothed long-term change of the pre-  
188 industrial control run. To avoid contaminating the drift correction with ICV, ideally the full length of the control run is used  
189 to determine the drift (Sen Gupta et al., 2013). Therefore, to dedrift the historical and scenario simulations of *zostoga* and *zos*  
190 (the latter on a grid cell by grid cell basis) we first fit a quadratic polynomial to the full pre-industrial control simulations of  
191 these variables. Then, we evaluate and subtract the polynomial fit over the time period in which the pre-industrial control run  
192 and historical and scenario runs overlap, as identified by the branch times of the different simulation realizations and their  
193 length, from the historical and scenario runs. Similar to what was found by Hermans et al., (2020) and Hobbs et al. (2016),  
194 fitting a linear or quadratic polynomial to the pre-industrial control simulations yields little difference for the drift-correction  
195 of the *zostoga* simulations of GISS-E2-R, HadGEM2-ES, IPSL-CM5A-LR, and MPI-ESM-LR. However, in the pre-industrial  
196 simulation of MPI-GE, the increase of *zostoga* behaves non-linearly and levels off toward the branching time of ensemble  
197 member 40, so we only dedrift ensemble members 1 to 39. For *zos*, some differences are found between linear and quadratic  
198 drift correction depending on the model, variant, and location. We assume linear dedrift is suitable for our purpose, since

199 we verified that the dedrifting does not substantially affect the pattern scaling performance and it is tedious to assess the best  
200 fit on a grid-point basis. After dedrifting, the area-weighted mean of  $zos$  is removed at each timestep, and the resulting fields  
201 are bilinearly regridded to a common 1 by 1 degree grid.

## 202 **3 Methods**

### 203 **3.1 Pattern filtering techniques**


204 Both S/N M EOF pattern filtering and LFCA aim to identify those spatial patterns in the data than explain most of the forced  
205 climate change signal, by decomposing the data into EOFs. Effectively, this allows to distinguish the forced signal from  
206 noise caused by ICV. The difference between S/N M EOF and LFCA lies in their definition of what type of variance (or  
207 patterns of variance) in the data belongs to the signal and the noise. Here, only the basics of both methods will be explained.  
208 Interested readers can find an extensive methodological explanation about S/N ME EOF pattern filtering applied to an  
209 ensemble and LFCA in Wills et al. (2020) and Wills et al (2018), respectively.

210 S/N M EOF pattern filtering diagnoses the variance that is forced by either assessing a simulation of forced climate change  
211 relative to a preindustrial control simulations (DelSole et al., 2011; Marcos and Amores, 2014), or by using an ensemble mean  
212 of realizations with the same forcing (Wills et al., 2020). The former is advantageous in single realization GCM experiments,  
213 as it only requires one forced realization and one preindustrial control run. However, this could neglect the forced response  
214 when external forcing only affects the phase of an ICV mode (Wills et al., 2020). The latter allows to effectively reduce ICV  
215 while avoiding phase neglect issues but requires the availability of two or more ensemble members. Since one of our  
216 objectives is to determine how efficient pattern filtering methods are compared to an ensemble mean of realizations to reduce  
217 ICV in DSL, here we focus on the latter approach.

218 Essentially, S/N M EOF pattern filtering exploits a SMILE to find patterns where different ensemble members agree on the  
219 temporal evolution (forced response), whereas those patterns in which members disagree are considered ICV. S/N M EOF  
220 pattern filtering finds spatial patterns (r.h.s. of Fig. 3, for example) associated with the time series  $t_k$  of each pattern  $k$  (l.h.s. of  
221 Fig. 3, for example) that maximize the ratio of (ensemble mean) signal to total variance  $s_k$ :

$$222 \quad s_k = \frac{\langle t_k \rangle^T \langle t_k \rangle}{t_k^T t_k}, \quad (1)$$

224  
225 where angle brackets represent an ensemble average. The leading S/N patterns (i.e., anomaly patterns with high signal fraction  
226  $s_k$ ) can be combined to isolate the forced response from the ICV (Fig. 1).



227

228 **Figure 1.** Main steps involved in isolating the forced response, including variability decomposition (EOF analysis), finding  
 229 leading anomaly patterns, and combining leading patterns above a significant statistical level.

230

231 To apply S/N M EOF pattern filtering, we must determine two parameters: 1) the number of EOFs retained (N), and 2) the  
 232 number of S/N patterns used to compose the forced response (M). Following the approach by Wills et al. (2020), we choose  
 233 N to retain between 75% and 95% of the total variance. We use a block bootstrapping approach to determine M, which consists  
 234 of taking block samples with replacement from the ensemble members to construct a randomized ensemble where the forced  
 235 response timing of their realizations should not agree with one another. Here, we choose 30-yr blocks to distinguish forced  
 236 patterns from ICV, so that most of the ICV in DSL is excluded. S/N EOF pattern filtering is then applied to randomized  
 237 ensembles and the  $s_k$  value of the pattern with the highest S/N ratio is taken as a threshold. This allows us to obtain a distribution  
 238 of  $s_k$  values (one for each randomized ensemble produced) from which a desired confidence level can be estimated. S/N M  
 239 EOF patterns with a higher  $s_k$  value than the threshold can be considered as part of the forced response with the chosen  
 240 confidence level (Fig. 1). As there is no sufficient statistical evidence to include patterns with a lower  $s_k$  value in the forced  
 241 response, those are considered noise (ICV).



242 In contrast to S/N M EOF, LFCA identifies the signal that makes it through a low-pass filter. The advantage of LFCA is that  
 243 it can analyse the forced response in a single ensemble member without relying on the preindustrial control run (Schneider and  
 244 Held, 2001; Wills et al., 2018). LFCA is similar to S/NP M EOF pattern filtering but, instead of using an ensemble mean, it  
 245 detects anomaly patterns associated with time series  $t_k$  (Eq. 2) that maximize the ratio of low-frequency signal to total variance.  
 246 The failure to detect some forced variations such as those driven by volcanic activity in surface air temperature and some  
 247 changes in the seasonal cycle is the main disadvantage of this method being documented in the literature (Wills et al., 2020).

$$249 \quad r_k = \frac{\widetilde{t}_k^T \widetilde{t}_k}{t_k^T t_k}, \quad (2)$$

250  
 251 Variations that make it through a low-pass filter (denoted by a tilde), constitute the low-frequency signal (forced response).  
 252 Here, we apply a linear Lanczos filter (Duchon, 1979) with a 30-yr lowpass filter, so only variability at larger timescales is  
 253 included. Following the same process as in S/N M EOF, a forced response can be constructed by linearly combining leading  
 254 anomaly patterns, as illustrated in Fig. 1.

### 255 **3.2 Pattern scaling**

256 Pattern scaling is usually based on grid-point regression against a global variable, and it assumes that a regional change in DSL  
 257 can be explained by global changes of the predictor(s) of choice. Previous studies have shown such relationships can be a  
 258 reasonable approximation for different variables of the climate system. For instance, local surface air temperature change  
 259 (Collins et al., 2013; Hawkins and Sutton, 2012) and local precipitation (Osborn et al., 2016) have successfully been linked to  
 260 GSAT change. Regional emulation based on pattern scaling assumes that patterns of local response to external forcing remains  
 261 constant (Tebaldi and Arblaster, 2014), an assumption that can lead to errors (Wells et al., 2022). However, its simplicity and  
 262 transferability to many regional variables have made it a popular approach for exploring regional changes in climate change  
 263 studies (Bilbao et al., 2015; Fox-Kemper, 2021; Herger et al., 2015; Mitchell, 2003; Osborn et al., 2016; Perrette et al., 2013;  
 264 Tebaldi and Arblaster, 2014; Thomas and Lin, 2018; Wells et al., 2022; Wu et al., 2021; Yuan and Kopp, 2021).

265 Once we have identified the forced DSL within an ensemble of realizations or a single simulation (as outlined in Section 3.1),  
 266 we will use this forced response as a predictand in our statistical model for projecting regional DSL. There are different forms  
 267 of pattern scaling, mostly differing in the number of predictors included in the analysis (e.g., univariate, Bilbao et al., 2015;  
 268 bivariate, Yuan & Kopp, 2021). Here, for simplicity and to ease comparison between raw (de-drifted) DSL and its pattern-  
 269 filtered equivalent, we only test pattern scaling based on GMTSLR (or *zostoga*) as a predictor. The univariate case of pattern  
 270 scaling for relating DSL with GMTSLR can be described by the following linear regression relationship:

271

$$\zeta(t, x, y) = \alpha(x, y) \bar{\eta}(t) + b(x, y) + \varepsilon(t, x, y) \quad (3)$$


273

274 Where  $\zeta$  and  $\bar{\eta}$  denote DSL and GMTSLR, respectively. Longitude and latitude are represented by  $x$  and  $y$ , whereas  $t$  denotes  
 275 time.  $\alpha$  is a spatial pattern that captures the scaling relationship between DSL and GMTSLR, and  $b$  is an intercept term, both  
 276 being only a function of location.  $\varepsilon$  is a residual term regarded as random noise and often assumed to be driven by internally  
 277 generated variability (Bilbao et al., 2015).

## 278 4 Results & Discussion

### 279 4.1 Forced response in MPI-GE and efficiency of pattern filtering.

280 In this section, we focus on determining the forced response in DSL within a SMILE (MPI-GE) using S/N M EOF pattern  
 281 filtering and show the efficiency of the latter to remove ICV compared to the more conventional approach of ensemble  
 282 averaging. To construct the forced response based on S/N patterns, we follow the block-bootstrapping approach described in  
 283 Section 3.1 we define blocks in terms of thirty years, so most ICV in DSL is excluded. 30-yr block samples are taken from the  
 284 100 historical realizations of the MPI-GE to construct 20 randomized ensembles. A value of 20 is chosen because increasing  
 285 it further does not lead to substantial changes in the estimation of the 95<sup>th</sup> percentile of  $S_k$ . The estimated ratio  $S_k$  (Eq. 1) for a  
 286 95 % confidence level is 0.08, leading to a total of eight patterns that can be considered as part of the forced response at such  
 287 a confidence level (Figure 1).



288


289

290 **Figure 2:** Signal fraction of the leading S/N M EOF patterns along with their respective explained forced response variance  
291 (%). The significance level (95%) computed using 30-year block-bootstrapping is represented as a dashed line. Patterns are  
292 sorted based on the magnitude of their signal fraction, as illustrated in Figure 1.

293

294 Even though patterns constructed based on EOFs are created from mathematical constraints, known physical processes can be  
295 identified in some patterns. For instance, the S/N M EOF pattern with the highest  $S_k$  value pattern 1, (Fig. 3) explains 62% of  
296 the forced response variance (Fig. 2) and is similar to the main forced pattern of DSL change field driven by increased radiative  
297 forcing due to increased GHG emissions. There is a zonal dipole in the Southern Ocean, with decreased and increased sea level  
298 relative to the mean below and above  $50^\circ\text{S}$ , respectively (e.g., Frankcombe et al., 2013). Another dipole structure is found in  
299 the North Atlantic with a decreased DSL in the north compared to an increased DSL in the southern section, a feature which  
300 appears to disagree with some models (e.g., Bouttes et al., 2014). Nonetheless, the North Atlantic Ocean is an area of large  
301 model spread in both CMIP5 and CMIP6 models (Lyu et al., 2020), which suggests the representation of such zonal dipole  
302 may be model dependent. Other relevant features include a large DSL rise in the Beaufort Sea and an increased DSL in the  
303 North-West Pacific Ocean. Most of these features agree with those documented among CMIP6 and earlier models (Church et  
304 al., 2013; Ferrero et al., 2021; Landerer et al., 2007; Lowe and Gregory, 2006; Lyu et al., 2020; Slangen et al., 2014). Patterns  
305 are similar between RCP scenarios, mainly differing on their intensity.

306 The three following resulting patterns (patterns 2, 3 and 4, Fig. S1, S2 and S3) represent between 4-1% (Fig. 2) of the forced  
307 response variance and, although with a much lower importance than pattern 1, when combined together represent non-linear  
308 processes that start to have an effect in DSL after 2050. Patterns 5, 6, 7 and 8 (Fig. S4, S5, S6, and S7) explain between 1-  
309 0.7% of the forced response variance (Fig. 2) and show a rather stable temporal evolution except for perturbations that coincide  
310 with historical volcanic eruptions from Krakatoa, Agung, El Chinchón, and Pinatubo. Volcano-induced perturbations were  
311 also observed in the analysis by Wills et al. (2020), as aerosol changes in the atmosphere can affect global and regional  
312 temperatures, subsequently affecting DSL. Patterns number 9 and beyond explain a variance of less than 0,6% and since their  
313  $S_k$  value is not statistically significant at the 95% level they could be caused by chance.



314

315

316

317

318

319

**Figure 3:** Time evolution of DSL standard deviation (a, c, and e) and associated S/N M EOF pattern number 1 for RCP 2.6, 4.5, and 8.5 (b, d, and f respectively). Light coloured lines in a, c, and d represent standard deviation anomalies from ensemble members, whereas dark coloured lines depict ensemble mean evolution of the pattern. In the historical + RCP scenarios DSL is calculated relative to the mean of 1993–2012.

320

321

322

323

324

325

326

327

328

329

330


331

332

We first compare the efficiency of pattern filtering techniques to that of conventional methods, in particular an ensemble mean, to isolate the forced response in DSL. We follow the approach used by Wills et al. (2020) based on the number of ensemble members needed to constrain a certain level of variance of the forced response using the coefficient of determination  $r^2$ , which indicates the proportion of variance shared between two datasets. As we need two datasets for such a comparison, the 100-member MPI-GE ensemble is divided into two sub-ensembles: one is used for testing (estimate ensemble) and the other is left for reference (reference ensemble). This leaves us with two 50-member sub-ensembles, where all 50 members in the reference sub-ensemble are used to estimate the forced response by either using ensemble averaging or S/N M EOF pattern filtering and this reference sub-ensemble is considered as ground truth. The other (estimate) 50-member ensemble is also used to estimate the forced response, but instead of using all sub-ensemble members we estimate the forced response in an iterative process by increasing the number of members included in the analysis from 2 to 50. As an illustration of the procedure, we start with only 2 members which are used to characterize the forced response in the estimate sub-ensemble and compare the result with the forced response from the 50-member reference sub-ensemble. This comparison is performed via the coefficient of determination between two estimated forced responses on a grid-point basis, identifying where the 80% level is exceeded.

333 Grid points where the threshold is not reached are used for subsequent analysis where an additional member (3 in total) is  
334 included in the estimate sub-ensemble, repeating the same process until the latter reaches 50 members. This procedure enables  
335 an evaluation of the number of ensemble members needed in the estimate sub-ensemble to characterize the forced response  
336 based on explained variance (i.e.,  $r^2$ ) in the reference sub-ensemble. To consider sampling uncertainty, this process is repeated  
337 ten times for random choices of realizations, taking the median value of all iterations.

338 When simple averaging is used, we find that 50 members are not sufficient to constrain at least 80% of the forced response  
339 variance of the reference ensemble over most of the ocean surface (Fig. 4a). In contrast, S/N M EOF pattern filtering  
340 characterises the forced response more efficiently than simply averaging, as it requires a much smaller number of realizations  
341 to remove ICV (Fig. 4b). While the grid-point median value of the number of ensemble members required is 50 or more when  
342 using simple averaging, the median estimate for the filtering method is reduced to eight. Large areas of the ocean benefit from  
343 filtering and there are significant reductions, especially the Indian Ocean, South and Northwest Atlantic Ocean, as well as  
344 large areas in the Pacific Ocean (Fig. 4b). Other areas, however, remain over the 50-member threshold to explain forced  
345 response variance after filtering. Those areas are mostly found where strong western boundary currents exist (Imawaki et al.,  
346 2013), as well as in areas influenced by the Antarctic Circumpolar Current (Rintoul et al., 2001). In those locations, variability  
347 is higher, and a larger number of realizations is needed to characterize it. Yet, there clearly is an advantage in using S/N M  
348 EOF over simple averaging methods, as less realizations are required to explain a significant part of the forced response in  
349 DSL, which means that the forced response can also be determined in models with smaller ensembles.



350  
351  
352 **Figure 4.** The number of ensemble members (realizations) needed to form an MPI-GE sub-ensemble that shares at least 80%  
353 of the variance of the forced response with a reference 50-member MPI-GE sub-ensemble using an ensemble average (a) and


354 using S/N M EOF pattern filtering (b) for RCP 2.6. The reference dataset is an average (a) or S/M EOF-filtered sub-ensemble  
355 (b) of 50 members which does not share realizations with the sub-ensemble used for estimation. Values represent the median  
356 of ten random choices of realizations sampling for both estimate and reference sub-ensembles. Note that bright yellow indicates  
357 more than 50 ensemble members required.

## 358 **4.2 Improved Pattern Scaling Using SMILES**

359 In this section, we demonstrate how S/N M EOF pattern filtering can increase the capabilities of statistical approaches for  
360 explaining DSL based in GMTSLR by reducing ICV within SMILES. For comparison, we first show pattern scaling  
361 performance when using single realizations and how conventional methods (ensemble mean) reduces RMSE when using a  
362 couple of realizations instead. Second, we examine S/N M EOF as a method for reducing RMSE more efficiently. We compare  
363 regional RSME from both ensemble mean and pattern filtering on only two realizations to allow an assessment of the areas  
364 that benefit the most from filtering when a few simulations are available. Lastly, we contrast how both ensemble mean and  
365 S/N M EOF pattern filtering reduce global mean RMSE as the number of realizations included in the analysis is increased.

366 As pattern scaling is performed on a grid-point basis, regression performances can be location dependent (Fig. 4a). Despite  
367 such regional variations, we found no substantial differences between GHG scenarios for both the regional and global mean  
368 RMSE estimates when pattern scaling DSL simulations extending up to 2100. Thus, results shown and discussed here are  
369 pertinent to the historical+RCP2.6 scenario for illustrative purposes, unless otherwise stated. When applying pattern scaling  
370 on a single realization of DSL from MPI-GE, the area-weighted, ensemble average RMSE is 3.78 cm, a value which is similar  
371 to previous estimates from studies performed on some of the CMIP5 models (Bilbao et al., 2015; Yuan and Kopp, 2021).  
372 However, pattern scaling performance shows a large spatial variability, ranging from 1.13 to 14.95 cm regionally (Fig. 5a).  
373 High RMSE values (i.e., lower regression performance) can be found in places subject to non-linear mesoscale processes  
374 driven by strong currents, coinciding with the places where the S/N M EOF technique requires many realizations to explain at  
375 least 80% of the forced response variance (Fig. 4b). These are the Antarctic Circumpolar Current (Southern Ocean) or western  
376 boundary currents, including the Gulf Stream (West North Atlantic), and Agulhas Current (South Africa), the Kuroshio Current  
377 (West North Pacific), and at the Brazil-Malvinas Confluence (West South Atlantic). Low RMSE values are found in the more  
378 stable eastern boundary currents, such as the Humboldt (Peru) Current, and in equatorial locations where DSL is relatively less  
379 influenced by large modes of climate variability (e.g., Equatorial Atlantic and Indian Ocean).

380 Despite its inefficiency, using an ensemble average cancels out some of the ICV that varies in a different phase between  
381 realizations. When using a 2-member ensemble mean, RMSE reduction is observed both globally and regionally: The area-  
382 weighted average RMSE estimate is reduced from 3.78 to 2.77 cm (27% reduction) when two ensembles are used, with regional  
383 values ranging from 0.87 to 11.00 cm (Fig. 5b). This translates to increased statistical model capabilities within the entire  
384 model domain. While grid-point RMSE reduction ranges from 10 to 30%, the majority of the ocean benefits from a decrease  
385 of more than 25% due to the removal of some of the ICV (Fig. 5c). Locations experiencing a lower improvement in regression  
386 performance include those that already performed relatively well prior averaging and those with a high ICV.



387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

**Figure 5.** Regional pattern scaling performance based on regression RMSE when one realization (a) and a two-member ensemble average (b) are used in the univariate regression. Sampling uncertainty is accounted for in (a) by averaging RMSE from pattern scaling performed individually to the 100 realizations, whereas in (b) random pairs (without replacement) are taken for the two-member ensemble average. The difference in regression performance between (a) and (b) is shown in (c) in terms of percentage. Results are shown for RCP 2.6 as an example.


To compare how S/N M EOF pattern filtering improves pattern scaling as opposed to averaging, we take two ensemble members from the MPI-GE historical+RCP2.6 experiment and proceed to remove their ICV by pattern filtering. The 2-member pattern-filtered DSL (Fig. 6a) shows an improved RMSE with similar regional structures compared to its averaged counterpart (Fig. 5b), featuring higher values in western boundary currents and Southern Ocean. Nonetheless, the overall improvement is apparent in all areas: the global estimated RMSE from the regression decreases almost 60% from an average value of 2.77 to 1.12 cm (Fig. 6c and d). Regionally, RMSE ranges from 0.39 to 6.05 cm when filtering is applied on two ensemble members (Fig. 6a and c). The differences between averaged and filtered approaches are substantial and location dependent, with filtering yielding a decrease in RMSE ranging from 12% to about 80% (Fig. 6b). The tropical Indian and

402 Eastern Pacific Ocean are among the locations benefiting the most from the largest performance improvement, which  
403 highlights the skill of pattern filtering to remove variability associated with large climate modes (e.g., ENSO has a large  
404 influence on sea level in the Eastern Pacific Ocean). Similar to previous findings when using averaging (Fig. 4c), pattern  
405 filtering offers a reduced improvement in areas where regression already performed relatively well or where the presence of  
406 meso-scale processes is significant. Regardless of improvement magnitude, pattern filtering provides an overall increase in  
407 regression performance that is observable in the entire ocean domain. While averaging also offers an enhancement of pattern  
408 scaling skill, filtered 2-member pairs produce a distribution of RMSE that is significantly superior (Fig. 6c).

409 We further investigate how pattern filtering enhances regression compared to averaging by increasing the number of  
410 members included in the analysis (Fig. 6d). Increasing the number of realizations grants ensemble averaging a considerable  
411 decrease in RMSE. Yet, performance improvement asymptotically reaches a plateau around 20 members after which further  
412 reductions in RMSE are modest. Regression based on pattern-filtered DSL also shows an improvement as the number of  
413 realizations increases. Such improvement is very limited compared to the one undergone by averaging, although filtering  
414 always provides a superior performance regardless of the number of members incorporated in the analysis. Importantly, area-  
415 weighted RMSE values differ significantly between the considered approaches when only a small number of realizations are  
416 available and become more similar for a larger number. This highlights the role of pattern filtering techniques when only a  
417 few ensemble members are available. Based on the analysis performed on the DSL simulations from the MPI-GE, filtering  
418 two members provides a regression performance that would only be achieved by averaging at least 12 members.

419





420


421 **Figure 6.** Regional pattern scaling performance based on regression RMSE when two ensemble members are used to estimate  
 422 the forced response via S/N M EOF pattern filtering (a). Panel (b): the difference in regression performance between the 2-  
 423 member average pattern scaling (Fig. 5b) and the S/N M EOF-filtered equivalent (a). Panel (c): Violin plots of RMSE  
 424 distributions from the 1-member, 2-member average, and 2-member S/N M EOF-filtered approaches. Panel (d): The area  
 425 weighted average RMSE obtained in the regression as a function of the number ensemble members included when using an  
 426 ensemble mean (yellow) and filtering (blue). The difference in performances in terms of percentage is shown in green. Analysis  
 427 for the RCP 2.6 scenario (we observed no discernible differences between scenarios).

428

### 429 4.3 Improved Pattern Scaling Using Single Realizations

430 Most models in CMIP prior to CMIP6 (and some in CMIP6) provided only one realization of historical and scenario  
 431 simulations. Therefore, we now test whether pattern filtering could improve regional emulation of single-realization models.  
 432 To do so, we apply LFCA which uses a similar approach to S/N M EOF (as explained in Section 3.1). In this section, we first  
 433 examine how LFCA improves the regression RMSE by truncating ICV in a single simulation from the MPI-GE. We then apply  
 434 LFCA to a range of CMIP5 models that were used in previous patterns scaling analyses of DSL, focusing on the differences  
 435 between models and RCP scenarios in longer simulations.

436 LFCA filtering uses the same linear algebra machinery as S/N M EOF, providing a similar regional improvement in pattern  
 437 scaling (compare Fig. 6a and 6a). Slightly higher RMSE values are observed in LFCA-based regression, for instance, in the  
 438 equatorial Pacific. This is expected because only one simulation is used, compared to two simulations in S/N M EOF filtering,  
 439 which enables the latter to identify a larger proportion of ICV. LFCA provides a substantial reduction in RMSE, as compared  
 440 to using a single simulation in pattern scaling (Fig. 7b-c). Regionally, it shows a similar qualitative pattern of improvement as  
 441 the other methods shown here (Fig. 7b vs 4c and 5b; averaging and S/N M EOF filtering, respectively). Quantitatively,  
 442 however, LFCA provides a larger RMSE reduction on a single realization than S/N M EOF performed on two. LFCA provides  
 443 a reduction of the area weighted average RMSE of 68% for all radiative forcing scenarios (Fig. 7c), while S/N M EOF yields  
 444 67% when using two realizations relative to unfiltered 1-member pattern scaling. While both estimates are quite similar, it is  
 445 worth noting that S/N M EOF requires two ensemble members to provide such reduction, while LFCA leads to a similar  
 446 performance just using one simulation. Similar to S/N M EOF pattern filtering, no substantial differences are found in pattern  
 447 scaling RMSE between RCP scenarios up to 2100 (Fig., 6c). This implies that ICV is analogous for different RCP scenarios  
 448 which, since a reduction in RMSE is due to the removal of ICV, leads to a similar improvement in performance for all RCPs  
 449 both globally (Fig. 7c) and regionally (not shown).  
 450




451

452 **Figure 7.** Regional pattern scaling performance based on regression RMSE when one (RCP 2.6) ensemble member is filtered  
453 via LFCA (a). Filtering is performed individually for each ensemble member to compute 100 scaling patterns whose results  
454 are averaged to diminish sampling issues. Differences in regression performance between Fig. 5a (unfiltered 1-member pattern  
455 scaling) and (a) are shown in (b) in terms of percentage. The area-weighted average RMSE is shown in (c) for RCPs 2.6, 4.5,  
456 and 8.5 and depending on whether the ensemble member is (blue) or not (yellow) filtered. Green indicates RMSE reduction  
457 between approaches in terms of percentage, whereas values on top of the bars are the absolute differences in cm.  
458

459 Since the aim of this study is to explore differences in emulated DSL when ICV is reduced, we also assess potential differences  
460 between unfiltered and filtered simulations (Fig. 8) when predicting DSL at 2100 using GMTSLR as a predictor. Emulated  
461 DSL differences caused by filtering may differ depending on the realization used, as each realization features an ICV evolving  
462 in a different phase. Thus, we focus on the maximum emulated DSL differences that filtering causes out of all 100 MPI-GE  
463 simulations. Exploring the maximal potential difference in statistically projected DSL is an added benefit of using SMILES,  
464 as such analysis can only be done with a large set of realizations with out-of-phase variability.

465 The difference in emulated DSL varies geographically (Fig. 8), with a spatial variability resembling the RMSE when ICV is  
466 reduced (e.g., Fig. 6a and 7a). Areas characterized by high temporal variability, which pattern filtering does not completely  
467 remove, experience greater difference in DSL projections (Fig. 8). Unlike RMSE (e.g., Fig. 7a), the difference between  
468 emulated DSL differs between RCP scenarios, increasing in magnitude with radiative forcing (Fig. 8). RMSE measures the  
469 error throughout the entire regression without accounting for the predictor, so only the effect of reduced ICV is captured. On  
470 the other hand, an increasing difference in predicted DSL with stronger RCP is expected since the magnitude of the predictor  
471 (GMTSLR) is larger for higher emissions scenarios. However, we observe the opposite behavior when assessing the difference  
472 in emulated DSL in relative terms, i.e., when the difference is divided by the emulated unfiltered DSL or by GMTSLR in 2100  
473 (not shown). Despite contrast between RCPs either in total difference (slightly increasing with forcing) or relative terms  
474 (decreasing with increasing forcing), RMSE being similar between RCPs highlights pattern filtering may be relevant for all  
475 scenarios.

476 The effect of pattern filtering on differences in slope  $\alpha$ , a key parameter in pattern scaling, shows again a similar spatial  
477 variability to RMSE (Fig. 7 vs Fig. S8). Changes in slopes are substantial in places with high variability, sometimes even  
478 showing a sign change (e.g., Fig. S13). Contrary to the total difference in emulated DSL and similar to the relative one, slope  
479 differences tend to decrease with higher emissions scenarios (Fig. S8). Since lower radiative forcing means lower signal-to-  
480 noise ratio, noise (ICV) can drive large differences in slopes between filtered and unfiltered results, and vice versa. Apart from  
481 reducing RMSE and leading to narrower confidence intervals (e.g., Fig., S10-14), pattern filtering finds slopes that are  
482 significantly different than the one obtained from applying a moving mean (e.g., Fig., S12 and 14), as the latter does not remove  
483 ICV as efficiently and requires neglecting data points for its computation (Fig., S10b-14b). It is worth highlighting that these  
484 differences in emulated DSL and slopes showcase an example for a GCM and may not hold as ground truth for other GCMs,  
485 scenarios, or predictors used.




486

487 **Figure 8.** Maximum difference between DSL change in 2100 obtained by pattern scaling with coefficients fitted to unfiltered  
 488 and LFCA-filtered realizations, considering all 100 MPI-GE members, for RCP 2.6, RC P4.5, and RCP 8.5 (a, b, and c,  
 489 respectively).

490 We further explore the performance of LFCA by comparing the pattern scaling results when isolating the forced response for  
 491 other GCMs. We identify the forced DSL in four CMIP5 models, being GISS-E2-R, HadGEM2-ES, IPSL-CM5A-LR, and  
 492 MPI-ESM-LR (Fig. 9a-d, respectively), which all provide scenario simulations up to 2300. To ease comparison with results  
 493 from the MPI-GE, however, we first examine results up to 2100 (Fig. 9a-d, small r.h.s. insets). RMSE from unfiltered  
 494 simulations up to 2100 vary between models, and so does RMSE reduction provided by LFCA. Nonetheless, error reduction  
 495 within a model and between scenarios is very similar, as previously observed for the MPI-GE. This implies that, for all models  
 496 considered here, there are no significant changing behaviours in the relationship between DSL and GMTLSR between RCP  
 497 scenarios up to 2100.

498 When considering results up to 2300, pattern scaling of unfiltered DSL against GMTSLR yields similar results as previous  
 499 studies (Bilbao et al., 2015), showing a global area-weighted mean RMSE between 2 and 4 cm. RMSE in both unfiltered and  
 500 filtered simulations of DSL increases with radiative forcing for all models considered. As simulations run up to 2300, a  
 501 decrease in pattern scaling performance for higher RCPs may indicate a more important role of the deeper ocean layer driving

502 non-linear processes (Bilbao et al., 2015; Yuan and Kopp, 2021). This tendency is also reflected in the error reduction after  
 503 filtering, which decreases as radiative forcing increases both over time and because of the higher emissions scenario, but the  
 504 latter is more apparent. Although LFCA filtering improves the performance of pattern scaling for all four CMIP5 models,  
 505 considerable differences in error reductions are observed. For instance, HadGEM2-ES benefits the most from pattern filtering  
 506 between all the models, with a ~70% decrease in error for RCP 2.6. Conversely, GISS-E2-R undergoes the lowest reduction  
 507 after pattern filtering, with about a 50% increase in performance for the same RCP scenario. Differences in model performance  
 508 pre- and post-filtering do not only highlight differences in how ICV is represented in distinct models but may also reflect  
 509 model differences in terms of physics representation and modelled forced response.  
 510



511  
 512 **Figure 9.** Area-weighted average RMSE for RCP 2.6, 4.5, and 8.5, indicating whether the ensemble member is (blue) or is not  
 513 (yellow) filtered via LFCA. Green indicates relative RMSE reduction between approaches (%), whereas values on top of the  
 514 bars are the absolute differences in cm. Different panels represent different CMIP5 models. The main panel includes simulation  
 515 data up to 2300, whereas the small inset on the right-hand top corner shows RMSE results up to 2100. Small insets share the  
 516 same axes as main panels.

## 517 **5 Conclusions**

518 Regional emulation tools for DSL change are complementary approaches to GCMs that allow for computationally cheap  
519 statistical projections. Most DSL regional emulators are based on pattern scaling, a statistical model usually based on a grid-  
520 point regression against a global variable representing change in the climate system driven by external forcing. While choosing  
521 suitable global predictors is essential for appropriate tuning of the statistical model, random errors can remain leading to high  
522 uncertainties in statistically based projections. A portion of these random errors are driven by ICV in DSL and can be  
523 characterised using macro-initialized initial condition large ensembles (SMILES), which are designed to facilitate a separation  
524 between ICV and external forcings within a model. Here, we applied pattern recognition techniques to a SMILE with the aim  
525 to efficiently truncate ICV, demonstrating how these approaches could significantly reduce random errors in regional emulators  
526 of DSL and provide substantially different emulated results in areas with high ICV.

527 Although ICV can be also reduced by using more conventional methods, such as computing an ensemble mean or linear trends,  
528 this requires a relatively large number of realizations to do it effectively. This is a significant constraint particularly for  
529 modelling experiments featuring a limited number of realizations. A more efficient alternative consists of employing methods  
530 that exploit spatial covariance information, such as S/N M EOF pattern filtering and LFCA. We have demonstrated that S/N  
531 M EOF applied to two realizations attains the same level of error reduction as averaging 12 realizations. The largest  
532 improvement relative to unfiltered simulations was observed when only a few simulations were available, whereas both S/N-  
533 filtered and ensemble average model performance tended to converge for a large number of ensemble members. By identifying  
534 spatiotemporal coherent structures, the S/N M EOF filtering was particularly skilful at removing ICV due to large modes of  
535 climate variability, such as the ENSO influence on sea level in the Eastern Pacific.

536 S/N M EOF pattern filtering can identify the common response within at least two realizations. This motivated us to also test  
537 LFCA, which can remove variability in single realization modelling experiments by applying a lowpass filter. Apart from  
538 being computationally more efficient, LFCA outperforms S/N M EOF in improving the performance of DSL pattern scaling  
539 when using one or two realizations. Moreover, LFCA applied to individual SMILE realizations allows exploring the maximal  
540 potential difference between statistically projected unfiltered and filtered DSL. We found substantial differences in emulated  
541 DSL and regression slopes in places with high variability, highlighting the relevance of pattern filtering methods in areas  
542 subject to non-mesoscale processes. Despite LFCA versatility and performances results, previous studies have emphasized  
543 that S/N M EOF pattern filtering provides a range of benefits compared to LFCA, including: 1) a better isolation of the forced  
544 response when the number of ensemble members is large, and 2) the detection of relatively less important forced patterns, such  
545 as those driven by volcanism.

546 We have also investigated LFCA by applying it to longer (up to 2300) CMIP5 simulations. We found that pattern scaling  
547 performance is independent of the GHG emission scenario up to 2100 and decreases with radiative forcing beyond 2100. Since  
548 we used a linear model, this implies that non-linear processes have different effects on DSL depending on the GHG scenario  
549 and this is reflected in a decrease in model performance depending on the emissions. We also found substantial differences

550 between CMIP5 models, due to variability being represented differently as well as distinct model physics. Nonetheless, the  
551 performance improvement of pattern scaling when applying LFCA filtering is considerable for all models and scenarios,  
552 ranging from 20% to more than 70% reduction relative to the unfiltered results.

553 Here, we have demonstrated that reducing ICV increases the capabilities of statistical approaches to project DSL. Pattern  
554 recognition techniques are especially advantageous for such a task, as they do not require numerous realizations to significantly  
555 reduce uncertainties in statistical projections and no data is lost (as in 30-year means) when reducing ICV. Previous studies  
556 have not considered removing ICV, which could significantly reduce uncertainties in statistically projected DSL and lead to  
557 substantial differences in emulated DSL. Although the difference in emulated DSL and regression slope varies depending on  
558 scenario, and results shown here are an example and may differ depending on GCM, RCPs, and predictor used, we show that  
559 pattern filtering is a useful approach to consider as a means of enhancing emulated DSL simulations.

### 560 **Code availability**

561 The methods used to perform this study are an adaptation from the ones used by Wills et al. (2020). The code is available at  
562 <https://github.com/rcjwills/forced-patterns> and <https://github.com/rcjwills/lfca>.

### 563 **Data availability**

564 Simulations from the MPI-GE can be obtained at <https://esgf-data.dkrz.de/projects/mip-ge/>, whereas CMIP5 data can be found  
565 at <https://esgf-node.llnl.gov/search/cmip5/>.

### 566 **Author contribution**

567 VMS devised, designed, and performed the analysis, and wrote the manuscript. ABAS supervised the study and contributed  
568 to writing. THJH contributed to data pre-processing and manuscript writing. SD and MM provided valuable feedback on  
569 methods and contributed to writing. NM provided MPI-GE data, information on MPI-GE methods, and contributed to writing.

### 570 **Competing interests**

571 The authors declare that they have no conflict of interest.

### 572 **Acknowledgements**

573 VMS, ABAS, THJH were supported by PROTECT. This project has received funding from the European Union's Horizon  
574 2020 research and innovation programme under grant agreement No 869304, PROTECT contribution number XX. SD

575 acknowledges David and Jane Flowerree for their support. We acknowledge the World Climate Research Programme's  
576 Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modelling groups for  
577 producing and making available their model output. For CMIP the U.S. Department of Energy's Program for Climate Model  
578 Diagnosis and Intercomparison provides coordinating support and led development of software infrastructure in partnership  
579 with the Global Organization for Earth System Science Portals.

## 580 **References**

581 Becker, M., Karpytchev, M., and Lennartz-Sassinek, S.: Long-term sea level trends: Natural or anthropogenic?, *Geophysical*  
582 *Research Letters*, 41, 5571–5580, <https://doi.org/10.1002/2014GL061027>, 2014.

583 Bilbao, R. A. F., Gregory, J. M., and Bouttes, N.: Analysis of the regional pattern of sea level change due to ocean dynamics  
584 and density change for 1993–2099 in observations and CMIP5 AOGCMs, *Clim Dyn*, 45, 2647–2666,  
585 <https://doi.org/10.1007/s00382-015-2499-z>, 2015.

586 Bouttes, N., Gregory, J. M., Kuhlbrodt, T., and Smith, R. S.: The drivers of projected North Atlantic sea level change, *Clim*  
587 *Dyn*, 43, 1531–1544, <https://doi.org/10.1007/s00382-013-1973-8>, 2014.

588 Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A.,  
589 Nerem, R. S., and Nunn, P. D.: *Sea level change*, PM Cambridge University Press, 2013.

590 Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W. J., Johns, T.,  
591 Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A. J., Wehner, M. F., Allen, M. R., Andrews, T., Beyerle, U., Bitz, C. M.,  
592 Bony, S., and Booth, B. B. B.: Long-term Climate Change: Projections, Commitments and Irreversibility, *Climate Change*  
593 *2013 - The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental*  
594 *Panel on Climate Change*, 1029–1136, 2013.

595 Cooley, S., Schoeman, D., Bopp, L., Boyd, P., Donner, S., Ito, S., Kiessling, W., Martinetto, P., Ojea, E., and Racault, M.-F.:  
596 *Oceans and Coastal Ecosystems and their Services*, in: *IPCC AR6 WGII*, Cambridge University Press, 2022.

597 Couldrey, M. P., Gregory, J. M., Boeira Dias, F., Dobrohotoff, P., Domingues, C. M., Garuba, O., Griffies, S. M., Haak, H.,  
598 Hu, A., Ishii, M., Jungclaus, J., Köhl, A., Marsland, S. J., Ojha, S., Saenko, O. A., Savita, A., Shao, A., Stammer, D., Suzuki,  
599 T., Todd, A., and Zanna, L.: What causes the spread of model projections of ocean dynamic sea-level change in response to  
600 greenhouse gas forcing?, *Clim Dyn*, 56, 155–187, <https://doi.org/10.1007/s00382-020-05471-4>, 2021.

601 Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J.,  
602 Garcia, R., and Gettelman, A.: The community earth system model version 2 (CESM2), *Journal of Advances in Modeling*  
603 *Earth Systems*, 12, e2019MS001916, 2020.

604 Dangendorf, S., Rybski, D., Mudersbach, C., Müller, A., Kaufmann, E., Zorita, E., and Jensen, J.: Evidence for long-term  
605 memory in sea level, *Geophysical Research Letters*, 41, 5530–5537, <https://doi.org/10.1002/2014GL060538>, 2014.

606 Dangendorf, S., Hay, C., Calafat, F. M., Marcos, M., Piecuch, C. G., Berk, K., and Jensen, J.: Persistent acceleration in global  
607 sea-level rise since the 1960s, *Nat. Clim. Chang.*, 9, 705–710, <https://doi.org/10.1038/s41558-019-0531-8>, 2019.



- 608 DelSole, T., Tippett, M. K., and Shukla, J.: A significant component of unforced multidecadal variability in the recent  
609 acceleration of global warming, *Journal of Climate*, 24, 909–926, 2011.
- 610 Deser, C., Lehner, F., Rodgers, K. B., Ault, T., Delworth, T. L., DiNezio, P. N., Fiore, A., Frankignoul, C., Fyfe, J. C., Horton,  
611 D. E., Kay, J. E., Knutti, R., Lovenduski, N. S., Marotzke, J., McKinnon, K. A., Minobe, S., Randerson, J., Screen, J. A.,  
612 Simpson, I. R., and Ting, M.: Insights from Earth system model initial-condition large ensembles and future prospects, *Nat.*  
613 *Clim. Chang.*, 10, 277–286, <https://doi.org/10.1038/s41558-020-0731-2>, 2020.
- 614 Duchon, C. E.: Lanczos Filtering in One and Two Dimensions, *Journal of Applied Meteorology and Climatology*, 18, 1016–  
615 1022, [https://doi.org/10.1175/1520-0450\(1979\)018<1016:LFIOAT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1979)018<1016:LFIOAT>2.0.CO;2), 1979.
- 616 Durand, G., van den Broeke, M. R., Le Cozannet, G., Edwards, T. L., Holland, P. R., Jourdain, N. C., Marzeion, B., Mottram,  
617 R., Nicholls, R. J., Pattyn, F., Paul, F., Slangen, A. B. A., Winkelmann, R., Burgard, C., van Calcar, C. J., Barré, J.-B., Bataille,  
618 A., and Chapuis, A.: Sea-Level Rise: From Global Perspectives to Local Services, *Frontiers in Marine Science*, 8, 2022.
- 619 Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E.,  
620 Smith, C. J., McKenna, C. M., Simon, E., Abe-Ouchi, A., Gregory, J. M., Larour, E., Lipscomb, W. H., Payne, A. J., Shepherd,  
621 A., Agosta, C., Alexander, P., Albrecht, T., Anderson, B., Asay-Davis, X., Aschwanden, A., Barthel, A., Bliss, A., Calov, R.,  
622 Chambers, C., Champollion, N., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Felikson, D., Fettweis, X., Fujita, K., Galton-  
623 Fenzi, B. K., Gladstone, R., Golledge, N. R., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huss, M., Huybrechts,  
624 P., Immerzeel, W., Kleiner, T., Kraaijenbrink, P., Le Clec'h, S., Lee, V., Leguy, G. R., Little, C. M., Lowry, D. P., Malles, J.-  
625 H., Martin, D. F., Maussion, F., Morlighem, M., O'Neill, J. F., Nias, I., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Radić,  
626 V., Reese, R., Rounce, D. R., Rückamp, M., Sakai, A., Shafer, C., Schlegel, N.-J., Shannon, S., Smith, R. S., Straneo, F., Sun,  
627 S., Tarasov, L., Trusel, L. D., Van Breedam, J., van de Wal, R., van den Broeke, M., Winkelmann, R., Zekollari, H., Zhao, C.,  
628 Zhang, T., and Zwinger, T.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74–82,  
629 <https://doi.org/10.1038/s41586-021-03302-y>, 2021.
- 630 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled  
631 Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*,  
632 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 633 Farrell, W. E. and Clark, J. A.: On postglacial sea level, *Geophysical Journal International*, 46, 647–667, 1976.
- 634 Fasullo, J. T., Gent, P. R., and Nerem, R. S.: Forced Patterns of Sea Level Rise in the Community Earth System Model Large  
635 Ensemble From 1920 to 2100, *Journal of Geophysical Research: Oceans*, 125, e2019JC016030,  
636 <https://doi.org/10.1029/2019JC016030>, 2020.
- 637 Ferrero, B., Tonelli, M., Marcello, F., and Wainer, I.: Long-term Regional Dynamic Sea Level Changes from CMIP6  
638 Projections, *Adv. Atmos. Sci.*, 38, 157–167, <https://doi.org/10.1007/s00376-020-0178-4>, 2021.
- 639 **Fox-Kemper, B., Hewitt, H. T., Xiao, C., Aalgeirsdottir, G., Drijfhout, S. S., Edwards, T. L., ... & Yu, Y. (2021). Ocean,**  
640 **Cryosphere and Sea Level Change. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to**  
641 **the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.**
- 642 Frankcombe, L. M., Spence, P., Hogg, A. M., England, M. H., and Griffies, S. M.: Sea level changes forced by Southern Ocean  
643 winds, *Geophysical Research Letters*, 40, 5710–5715, 2013.
- 644 Frankcombe, L. M., England, M. H., Mann, M. E., and Steinman, B. A.: Separating Internal Variability from the Externally  
645 Forced Climate Response, *Journal of Climate*, 28, 8184–8202, <https://doi.org/10.1175/JCLI-D-15-0069.1>, 2015.

- 646 Frederikse, T., Landerer, F., Caron, L., Adhikari, S., Parkes, D., Humphrey, V. W., Dangendorf, S., Hogarth, P., Zanna, L.,  
647 and Cheng, L.: The causes of sea-level rise since 1900, *Nature*, 584, 393–397, 2020.
- 648 Geoffroy, O., Saint-Martin, D., Olivié, D. J., Voldoire, A., Bellon, G., and Tytéca, S.: Transient climate response in a two-  
649 layer energy-balance model. Part I: Analytical solution and parameter calibration using CMIP5 AOGCM experiments, *Journal*  
650 *of Climate*, 26, 1841–1857, 2013a.
- 651 Geoffroy, O., Saint-Martin, D., Bellon, G., Voldoire, A., Olivié, D. J. L., and Tytéca, S.: Transient climate response in a two-  
652 layer energy-balance model. Part II: Representation of the efficacy of deep-ocean heat uptake and validation for CMIP5  
653 AOGCMs, *Journal of Climate*, 26, 1859–1876, 2013b.
- 654 Goodwin, P., Katavouta, A., Roussenov, V. M., Foster, G. L., Rohling, E. J., and Williams, R. G.: Pathways to 1.5 C and 2 C  
655 warming based on observational and geological constraints, *Nature Geoscience*, 11, 102–107, 2018.
- 656 Gregory, J. M., Griffies, S. M., Hughes, C. W., Lowe, J. A., Church, J. A., Fukimori, I., Gomez, N., Kopp, R. E., Landerer, F.,  
657 Cozannet, G. L., Ponte, R. M., Stammer, D., Tamisiea, M. E., and van de Wal, R. S. W.: Concepts and Terminology for Sea  
658 Level: Mean, Variability and Change, Both Local and Global, *Surv Geophys*, 40, 1251–1289, [https://doi.org/10.1007/s10712-](https://doi.org/10.1007/s10712-019-09525-z)  
659 [019-09525-z](https://doi.org/10.1007/s10712-019-09525-z), 2019.
- 660 Griffies, S. M., Danabasoglu, G., Durack, P. J., Adcroft, A. J., Balaji, V., Böning, C. W., Chassignet, E. P., Curchitser, E.,  
661 Deshayes, J., Drange, H., Fox-Kemper, B., Gleckler, P. J., Gregory, J. M., Haak, H., Hallberg, R. W., Heimbach, P., Hewitt,  
662 H. T., Holland, D. M., Ilyina, T., Jungclaus, J. H., Komuro, Y., Krasting, J. P., Large, W. G., Marsland, S. J., Masina, S.,  
663 McDougall, T. J., Nurser, A. J. G., Orr, J. C., Pirani, A., Qiao, F., Stouffer, R. J., Taylor, K. E., Treguier, A. M., Tsujino, H.,  
664 Uotila, P., Valdivieso, M., Wang, Q., Winton, M., and Yeager, S. G.: OMIP contribution to CMIP6: experimental and  
665 diagnostic protocol for the physical component of the Ocean Model Intercomparison Project, *Geoscientific Model*  
666 *Development*, 9, 3231–3296, <https://doi.org/10.5194/gmd-9-3231-2016>, 2016.
- 667 Gupta, A. S., Jourdain, N. C., Brown, J. N., and Monselesan, D.: Climate drift in the CMIP5 models, *Journal of Climate*, 26,  
668 8597–8615, 2013.
- 669 Haasnoot, M., Brown, S., Scussolini, P., Jimenez, J. A., Vafeidis, A. T., and Nicholls, R. J.: Generic adaptation pathways for  
670 coastal archetypes under uncertain sea-level rise, *Environmental Research Communications*, 1, 071006, 2019.
- 671 Haasnoot, M., Winter, G., Brown, S., Dawson, R. J., Ward, P. J., and Eilander, D.: Long-term sea-level rise necessitates a  
672 commitment to adaptation: A first order assessment, *Climate Risk Management*, 34, 100355,  
673 <https://doi.org/10.1016/j.crm.2021.100355>, 2021.
- 674 Haigh, I. D., Pickering, M. D., Green, J. A. M., Arbic, B. K., Arns, A., Dangendorf, S., Hill, D. F., Horsburgh, K., Howard,  
675 T., Idier, D., Jay, D. A., Jänicke, L., Lee, S. B., Müller, M., Schindelegger, M., Talke, S. A., Wilmes, S.-B., and Woodworth,  
676 P. L.: The Tides They Are A-Changin’: A Comprehensive Review of Past and Future Nonastronomical Changes in Tides,  
677 Their Driving Mechanisms, and Future Implications, *Reviews of Geophysics*, 58, e2018RG000636,  
678 <https://doi.org/10.1029/2018RG000636>, 2020.
- 679 Hasselmann, K.: Stochastic climate models part I. Theory, *tellus*, 28, 473–485, 1976.
- 680 Hawkins, E. and Sutton, R.: Time of emergence of climate signals, *Geophysical Research Letters*, 39,  
681 <https://doi.org/10.1029/2011GL050087>, 2012.
- 682 Hawkins, E., Smith, R. S., Gregory, J. M., and Stainforth, D. A.: Irreducible uncertainty in near-term climate projections, *Clim*  
683 *Dyn*, 46, 3807–3819, <https://doi.org/10.1007/s00382-015-2806-8>, 2016.

- 684 Herger, N., Sanderson, B. M., and Knutti, R.: Improved pattern scaling approaches for the use in climate impact studies,  
685 *Geophysical Research Letters*, 42, 3486–3494, <https://doi.org/10.1002/2015GL063569>, 2015.
- 686 Hermans, T. H. J., Tinker, J., Palmer, M. D., Katsman, C. A., Vermeersen, B. L. A., and Slangen, A. B. A.: Improving sea-  
687 level projections on the Northwestern European shelf using dynamical downscaling, *Clim Dyn*, 54, 1987–2011,  
688 <https://doi.org/10.1007/s00382-019-05104-5>, 2020.
- 689 Hinkel, J., Lincke, D., Vafeidis, A. T., Perrette, M., Nicholls, R. J., Tol, R. S., Marzeion, B., Fettweis, X., Ionescu, C., and  
690 Levermann, A.: Coastal flood damage and adaptation costs under 21st century sea-level rise, *Proceedings of the National  
691 Academy of Sciences*, 111, 3292–3297, 2014.
- 692 Hobbs, W., Palmer, M. D., and Monselesan, D.: An energy conservation analysis of ocean drift in the CMIP5 global coupled  
693 models, *Journal of Climate*, 29, 1639–1653, 2016.
- 694 Imawaki, S., Bower, A. S., Beal, L., and Qiu, B.: Chapter 13 - Western Boundary Currents, in: *International Geophysics*, vol.  
695 103, edited by: Siedler, G., Griffies, S. M., Gould, J., and Church, J. A., Academic Press, 305–338,  
696 <https://doi.org/10.1016/B978-0-12-391851-2.00013-1>, 2013.
- 697 Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S. C., Danabasoglu, G., and Edwards,  
698 J.: The Community Earth System Model (CESM) large ensemble project: A community resource for studying climate change  
699 in the presence of internal climate variability, *Bulletin of the American Meteorological Society*, 96, 1333–1349, 2015.
- 700 Labe, Z. M. and Barnes, E. A.: Detecting Climate Signals Using Explainable AI With Single-Forcing Large Ensembles, *Journal  
701 of Advances in Modeling Earth Systems*, 13, e2021MS002464, <https://doi.org/10.1029/2021MS002464>, 2021.
- 702 Landerer, F. W., Jungclaus, J. H., and Marotzke, J.: Regional dynamic and steric sea level change in response to the IPCC-  
703 A1B scenario, *Journal of Physical Oceanography*, 37, 296–312, 2007.
- 704 Lowe, J. A. and Gregory, J. M.: Understanding projections of sea level rise in a Hadley Centre coupled climate model, *Journal  
705 of Geophysical Research: Oceans*, 111, 2006.
- 706 Lyu, K., Zhang, X., and Church, J. A.: Regional Dynamic Sea Level Simulated in the CMIP5 and CMIP6 Models: Mean  
707 Biases, Future Projections, and Their Linkages, *Journal of Climate*, 33, 6377–6398, <https://doi.org/10.1175/JCLI-D-19-1029.1>,  
708 2020.
- 709 Maher, N., Milinski, S., Suarez-Gutierrez, L., Botzet, M., Dobrynin, M., Kornblueh, L., Kröger, J., Takano, Y., Ghosh, R.,  
710 Hedemann, C., Li, C., Li, H., Manzini, E., Notz, D., Putrasahan, D., Boysen, L., Claussen, M., Ilyina, T., Olonscheck, D.,  
711 Raddatz, T., Stevens, B., and Marotzke, J.: The Max Planck Institute Grand Ensemble: Enabling the Exploration of Climate  
712 System Variability, *Journal of Advances in Modeling Earth Systems*, 11, 2050–2069, <https://doi.org/10.1029/2019MS001639>,  
713 2019.
- 714 Maher, N., Milinski, S., and Ludwig, R.: Large ensemble climate model simulations: introduction, overview, and future  
715 prospects for utilising multiple types of large ensemble, *Earth System Dynamics*, 12, 401–418, [https://doi.org/10.5194/esd-  
716 12-401-2021](https://doi.org/10.5194/esd-12-401-2021), 2021a.
- 717 Maher, N., Power, S., and Marotzke, J.: More accurate quantification of model-to-model agreement in externally forced  
718 climatic responses over the coming century, *Nature Communications*, 12, 788, <https://doi.org/10.1038/s41467-020-20635-w>,  
719 2021b.

- 720 Mankin, J. S., Lehner, F., Coats, S., and McKinnon, K. A.: The Value of Initial Condition Large Ensembles to Robust  
721 Adaptation Decision-Making, *Earth's Future*, 8, e2012EF001610, <https://doi.org/10.1029/2020EF001610>, 2020.
- 722 Marcos, M. and Amores, A.: Quantifying anthropogenic and natural contributions to thermosteric sea level rise, *Geophysical  
723 Research Letters*, 41, 2502–2507, <https://doi.org/10.1002/2014GL059766>, 2014.
- 724 Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L.: Emulating coupled atmosphere-ocean and carbon cycle models with  
725 a simpler model, *MAGICC6 – Part 1: Model description and calibration*, *Atmospheric Chemistry and Physics*, 11, 1417–1456,  
726 <https://doi.org/10.5194/acp-11-1417-2011>, 2011.
- 727 Millar, R. J., Nicholls, Z. R., Friedlingstein, P., and Allen, M. R.: A modified impulse-response representation of the global  
728 near-surface air temperature and atmospheric concentration response to carbon dioxide emissions, *Atmospheric Chemistry and  
729 Physics*, 17, 7213–7228, 2017.
- 730 Mitchell, T. D.: Pattern Scaling: An Examination of the Accuracy of the Technique for Describing Future Climates, *Climatic  
731 Change*, 60, 217–242, <https://doi.org/10.1023/A:1026035305597>, 2003.
- 732 Mitrovica, J. X., Tamisiea, M. E., Davis, J. L., and Milne, G. A.: Recent mass balance of polar ice sheets inferred from patterns  
733 of global sea-level change, *Nature*, 409, 1026–1029, 2001.
- 734 Moftakhari, H. R., AghaKouchak, A., Sanders, B. F., Feldman, D. L., Sweet, W., Matthew, R. A., and Luke, A.: Increased  
735 nuisance flooding along the coasts of the United States due to sea level rise: Past and future, *Geophysical Research Letters*,  
736 42, 9846–9852, 2015.
- 737 Nerem, R. S., Leuliette, É., and Cazenave, A.: Present-day sea-level change: A review, *Comptes Rendus Geoscience*, 338,  
738 1077–1083, <https://doi.org/10.1016/j.crte.2006.09.001>, 2006.
- 739 Nicholls, R. J., Lincke, D., Hinkel, J., Brown, S., Vafeidis, A. T., Meyssignac, B., Hanson, S. E., Merken, J.-L., and Fang, J.:  
740 A global analysis of subsidence, relative sea-level change and coastal flood exposure, *Nature Climate Change*, 11, 338–342,  
741 <https://doi.org/10.1038/s41558-021-00993-z>, 2021.
- 742 O'Neill, B. C., Kriegler, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B. J., van Vuuren, D. P.,  
743 Birkmann, J., Kok, K., Levy, M., and Solecki, W.: The roads ahead: Narratives for shared socioeconomic pathways describing  
744 world futures in the 21st century, *Global Environmental Change*, 42, 169–180,  
745 <https://doi.org/10.1016/j.gloenvcha.2015.01.004>, 2017.
- 746 Osborn, T. J., Wallace, C. J., Harris, I. C., and Melvin, T. M.: Pattern scaling using ClimGen: monthly-resolution future climate  
747 scenarios including changes in the variability of precipitation, *Climatic Change*, 134, 353–369, [https://doi.org/10.1007/s10584-  
748 015-1509-9](https://doi.org/10.1007/s10584-015-1509-9), 2016.
- 749 Peltier, W. R.: Global sea level rise and glacial isostatic adjustment, *Global and Planetary Change*, 20, 93–123, 1999.
- 750 Peltier, W. R.: Global glacial isostatic adjustment and modern instrumental records of relative sea level history, in: *International  
751 geophysics*, vol. 75, Elsevier, 65–95, 2001.
- 752 Perrette, M., Landerer, F., Riva, R., Frieler, K., and Meinshausen, M.: A scaling approach to project regional sea level rise and  
753 its uncertainties, *Earth System Dynamics*, 4, 11–29, 2013.

- 754 R. Rintoul, S., W. Hughes, C., and Olbers, D.: Chapter 4.6 The antarctic circumpolar current system, in: *International*  
755 *Geophysics*, vol. 77, edited by: Siedler, G., Church, J., and Gould, J., Academic Press, 271–XXXVI,  
756 [https://doi.org/10.1016/S0074-6142\(01\)80124-8](https://doi.org/10.1016/S0074-6142(01)80124-8), 2001.
- 757 Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko,  
758 O., Lutz, W., Popp, A., Cuaresma, J. C., Kc, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa,  
759 T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj,  
760 J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M.,  
761 Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic  
762 Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environmental Change*,  
763 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- 764 Santer, B. D., Wigley, T. M., Schlesinger, M. E., and Mitchell, J. F.: Developing climate scenarios from equilibrium GCM  
765 results, 1990.
- 766 Schneider, T. and Held, I. M.: Discriminants of twentieth-century changes in Earth surface temperatures, *Journal of Climate*,  
767 14, 249–254, 2001.
- 768 Schwarber, A. K., Smith, S. J., Hartin, C. A., Vega-Westhoff, B. A., and Sriver, R.: Evaluating climate emulation: fundamental  
769 impulse testing of simple climate models, *Earth System Dynamics Discussions*, 10, 729–739, [https://doi.org/10.5194/esd-10-](https://doi.org/10.5194/esd-10-729-2019)  
770 729-2019, 2019.
- 771 Schwarzwald, K. and Lenssen, N.: The importance of internal climate variability in climate impact projections, *Proceedings*  
772 *of the National Academy of Sciences*, 119, e2208095119, 2022.
- 773 Slangen, A. B. A., Carson, M., Katsman, C. A., Van de Wal, R. S. W., Köhl, A., Vermeersen, L. L. A., and Stammer, D.:  
774 Projecting twenty-first century regional sea-level changes, *Climatic Change*, 124, 317–332, 2014.
- 775 Slangen, A. B. A., Adloff, F., Jevrejeva, S., Leclercq, P. W., Marzeion, B., Wada, Y., and Winkelmann, R.: A Review of  
776 Recent Updates of Sea-Level Projections at Global and Regional Scales, *Surv Geophys*, 38, 385–406,  
777 <https://doi.org/10.1007/s10712-016-9374-2>, 2017.
- 778 Smith, C. J., Forster, P. M., Allen, M., Leach, N., Millar, R. J., Passerello, G. A., and Regayre, L. A.: FAIR v1. 3: a simple  
779 emissions-based impulse response and carbon cycle model, *Geoscientific Model Development*, 11, 2273–2297, 2018.
- 780 Stainforth, D. a, Allen, M. r, Tredger, E. r, and Smith, L. a: Confidence, uncertainty and decision-support relevance in climate  
781 predictions, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 365, 2145–  
782 2161, <https://doi.org/10.1098/rsta.2007.2074>, 2007.
- 783 Stammer, D. and Hüttemann, S.: Response of regional sea level to atmospheric pressure loading in a climate change scenario,  
784 *Journal of Climate*, 21, 2093–2101, 2008.
- 785 Steffelbauer, D. B., Riva, R. E. M., Timmermans, J. S., Kwakkel, J. H., and Bakker, M.: Evidence of regional sea-level rise  
786 acceleration for the North Sea, *Environ. Res. Lett.*, 17, 074002, <https://doi.org/10.1088/1748-9326/ac753a>, 2022.
- 787 Suarez-Gutierrez, L., Milinski, S., and Maher, N.: Exploiting large ensembles for a better yet simpler climate model evaluation,  
788 *Clim Dyn*, 57, 2557–2580, <https://doi.org/10.1007/s00382-021-05821-w>, 2021.
- 789 Tebaldi, C. and Arblaster, J. M.: Pattern scaling: Its strengths and limitations, and an update on the latest model simulations,  
790 *Climatic Change*, 122, 459–471, <https://doi.org/10.1007/s10584-013-1032-9>, 2014.

- 791 Thomas, M. A. and Lin, T.: A dual model for emulation of thermosteric and dynamic sea-level change, *Climatic Change*, 148,  
792 311–324, 2018.
- 793 Vitousek, S., Barnard, P. L., Fletcher, C. H., Frazer, N., Erikson, L., and Storlazzi, C. D.: Doubling of coastal flooding  
794 frequency within decades due to sea-level rise, *Scientific reports*, 7, 1–9, 2017.
- 795 van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V.,  
796 Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration  
797 pathways: an overview, *Climatic Change*, 109, 5, <https://doi.org/10.1007/s10584-011-0148-z>, 2011.
- 798 Wahl, T., Haigh, I. D., Nicholls, R. J., Arns, A., Dangendorf, S., Hinkel, J., and Slangen, A. B. A.: Understanding extreme sea  
799 levels for broad-scale coastal impact and adaptation analysis, *Nat Commun*, 8, 16075, <https://doi.org/10.1038/ncomms16075>,  
800 2017.
- 801 Wells, C. D., Jackson, L. S., Maycock, A. C., and Forster, P. M.: Understanding pattern scaling errors across a range of  
802 emissions pathways, 30, 2022.
- 803 Wills, R. C., Schneider, T., Wallace, J. M., Battisti, D. S., and Hartmann, D. L.: Disentangling Global Warming, Multidecadal  
804 Variability, and El Niño in Pacific Temperatures, *Geophysical Research Letters*, 45, 2487–2496,  
805 <https://doi.org/10.1002/2017GL076327>, 2018.
- 806 Wills, R. C. J., Battisti, D. S., Armour, K. C., Schneider, T., and Deser, C.: Pattern Recognition Methods to Separate Forced  
807 Responses from Internal Variability in Climate Model Ensembles and Observations, *Journal of Climate*, 33, 8693–8719,  
808 <https://doi.org/10.1175/JCLI-D-19-0855.1>, 2020.
- 809 Wu, Q., Zhang, X., Church, J. A., Hu, J., and Gregory, J. M.: Evolving patterns of steric sea-level rise under mitigation  
810 scenarios and insights from linear system theory, *Clim Dyn*, 57, 635–656, <https://doi.org/10.1007/s00382-021-05727-7>, 2021.
- 811 Yuan, J. and Kopp, R. E.: Emulating Ocean Dynamic Sea Level by Two-Layer Pattern Scaling, *Journal of Advances in*  
812 *Modeling Earth Systems*, 13, e2020MS002323, <https://doi.org/10.1029/2020MS002323>, 2021.

813