

Supplementary Material: ‘Improving Statistical Projections of Ocean Dynamic Sea-level Change Using Pattern Recognition Techniques.’

Víctor Malagón-Santos¹, Aimée B.A. Slangen¹, Tim H.J. Hermans^{1,2}, Sönke Dangendorf³, Marta Marcos⁴, Nicola Maher^{5,6,7}.

¹NIOZ Royal Netherlands Institute for Sea Research, Department of Estuarine & Delta Systems, P.O. Box 140, 4400 AC Yerseke, the Netherlands.

²University of Utrecht, Institute for Marine and Atmospheric research Utrecht (IMAU), Utrecht, The Netherlands

³Department of River-Coastal Science and Engineering, Tulane University, New Orleans, USA.

⁴Mediterranean Institute for Advanced Studies (IMEDEA), Spanish National Research Council-University of Balearic Islands (CSIC-UIB), Esporles, Spain.

⁵Cooperative Institute for Research in Environmental Science, University of Colorado, Boulder, CO, USA.

⁶Department of Atmospheric and Oceanic Sciences, University of Colorado, Boulder, CO, USA.

⁷Max Planck Institute for Meteorology, Hamburg, Germany.

Correspondence to: Víctor Malagón-Santos (victor.malagon.santos@nioz.nl)




Figure S1: Time evolution in terms of standard deviations (a, c, and e, respectively) and associated S/N M EOF pattern number 2 for RCP 2.6, 4.5, and 8.5 (b, d, and f respectively). Light coloured lines in a, c, and d represent standard deviation anomalies from ensemble members, whereas dark coloured lines depict ensemble mean evolution of the pattern.




Figure S2: Time evolution in terms of standard deviations (a, c, and e, respectively) and associated S/N M EOF pattern number 3 for RCP 2.6, 4.5, and 8.5 (b, d, and f respectively). Light coloured lines in a, c, and d represent standard deviation anomalies from ensemble members, whereas dark coloured lines depict ensemble mean evolution of the pattern.




Figure S3: Time evolution in terms of standard deviations (a, c, and e, respectively) and associated S/N M EOF pattern number 4 for RCP 2.6, 4.5, and 8.5 (b, d, and f respectively). Light coloured lines in a, c, and d represent standard deviation anomalies from ensemble members, whereas dark coloured lines depict ensemble mean evolution of the pattern.




Figure S4: Time evolution in terms of standard deviations (a, c, and e, respectively) and associated S/N M EOF pattern number 5 for RCP 2.6, 4.5, and 8.5 (b, d, and f respectively). Light coloured lines in a, c, and d represent standard deviation anomalies from ensemble members, whereas dark coloured lines depict ensemble mean evolution of the pattern.




Figure S5: Time evolution in terms of standard deviations (a, c, and e, respectively) and associated S/N M EOF pattern number 6 for RCP 2.6, 4.5, and 8.5 (b, d, and f respectively). Light coloured lines in a, c, and d represent standard deviation anomalies from ensemble members, whereas dark coloured lines depict ensemble mean evolution of the pattern.




Figure S6: Time evolution in terms of standard deviations (a, c, and e, respectively) and associated S/N M EOF pattern number 7 for RCP 2.6, 4.5, and 8.5 (b, d, and f respectively). Light coloured lines in a, c, and d represent standard deviation anomalies from ensemble members, whereas dark coloured lines depict ensemble mean evolution of the pattern.




Figure S7: Time evolution in terms of standard deviations (a, c, and e, respectively) and associated S/N M EOF pattern number 8 for RCP 2.6, 4.5, and 8.5 (b, d, and f respectively). Light coloured lines in a, c, and d represent standard deviation anomalies from ensemble members, whereas dark coloured lines depict ensemble mean evolution of the pattern.