

# Identification and ranking of subaerial volcanic tsunami hazard sources in Southeast Asia

Edgar U. Zorn<sup>1</sup>, Aiyem Orynbaiyzy<sup>2</sup>, Simon Plank<sup>2</sup>, Andrey Babeyko<sup>1</sup>, Herlan Darmawan<sup>3</sup>, Ismail Fata Robbany<sup>2</sup>, Thomas R. Walter<sup>1</sup>

<sup>1</sup>German Research Centre for Geosciences GFZ, Telegrafenberg, 14473 Potsdam, Germany

<sup>2</sup>German Aerospace Center DLR, Münchenerstr. 20, 82234 Weßling, Germany

<sup>3</sup>Geophysics Study Program, Department of Physics, Faculty of Mathematics and Natural Sciences, Universitas Gadjah Mada, Sekip Utara, Bulaksumur, Yogyakarta, Indonesia

*Correspondence to:* Edgar U. Zorn (zorn@gfz-potsdam.de)

## Abstract.

Tsunamis caused by large volcanic eruptions and flanks collapsing into the sea are major hazards for nearby coastal regions.

They often occur with little precursory activity, and are thus challenging to detect in a timely manner. This makes the pre-emptive identification of volcanoes prone to causing tsunamis particularly important, as it allows for better hazard assessment and denser monitoring in these areas. Here, we present a catalogue of potentially tsunamigenic volcanoes in Southeast Asia and rank these volcanoes by their tsunami hazard. The ranking is based on a Multicriteria Decision Analysis (MCDA) composed of five individually weighted factors impacting flank stability and tsunami hazard. The data is sourced from geological databases, remote sensing data, historical volcano induced tsunami records and our topographic analyses, mainly considering the eruptive and tsunami history, elevation relative to the distance from the sea, flank steepness, hydrothermal alteration as well as vegetation coverage. Out of 131 analysed volcanoes, we found 19 with particularly high tsunamigenic hazard potential in Indonesia (Anak Krakatau, Batu Tara, Iliwerung, Gamalama, Sangeang Api, Karangetang, Sirung, Wetar, Nila, Ruang, Serua) and Papua New Guinea (Kadovar, Ritter Island, Rabaul, Manam, Langila, Ulawun, Bam), but also in the Philippines (Didicas). While some of these volcanoes, such as Anak Krakatau, are well-known for their deadly tsunamis, many others on this list are lesser known and monitored. We further performed tsunami travel time modelling on these high-hazard volcanoes, which indicates that future events could affect large coastal areas in a short time. This highlights the importance of individual tsunami hazard assessment for these volcanoes, dedicated volcanological monitoring, and the need for increased preparedness on the potentially affected coasts.

## 30 1 Introduction

Tsunamis are among the deadliest hazards affecting coastal regions around the world. While most tsunamis are caused by tectonic earthquakes, tsunamis induced by volcanic sources account for ~6% of global tsunamis (NGDC, 2021). As a result, volcanic tsunamis are far less researched, but have resulted in many deadly events that were often unexpected and hit shores without warning. This is because volcanic tsunamis are low-probability, high-impact and hardly predictable black swan events ~~and 26% of all fatalities recorded at volcanoes since 1800 have been due to such tsunamis, causing some 26% of all volcano-induced fatalities since 1800~~ (Brown et al., 2017). Due to the very high density of active volcanoes near coastlines, Southeast (SE) Asia is one of the most prominent regions in the world for volcano induced tsunami events. The most well-known example is Krakatau volcano, Indonesia, where a tsunami caused by a major eruption in 1883 had an estimated death toll of ~36,000 people (Hamzah et al., 2000; Brown et al., 2017). In 2018, another tsunami by the same volcano killed 437 people due to the instability of its regrown volcano flank, which caused a sector collapse into the sea (Walter et al., 2019; Darmawan et al., 2020; Omira and Ramalho, 2020). In 1888, Ritter Island in Papua New Guinea experienced a similar catastrophic collapse, resulting in a tsunami with a death toll exceeding several hundred people (Ward and Day, 2003). Even without flank or edifice instability, volcanic eruptions can still cause deadly tsunamis by the expulsion of pyroclastic density currents (PDCs) into the sea. Such events repeatedly occurred at Awu volcano, Indonesia, in 1856, 1892 and 1913 with a cumulative ~4,500 fatalities (Hidayat et al., 2020). The recent explosive eruption of the Hunga Tonga-Hunga Ha'apai volcano near Tonga caused a tsunami affecting the entire Pacific Ocean, which travelled faster towards the coasts than was expected. While the mechanisms of this tsunami are not fully understood yet, reports suggest the interaction of acoustic and water gravity waves as a possible explanation (Somerville et al., 2022). This may be similar to airwaves attributed to tsunamis produced by Taal volcano, Philippines, in 1911 and 1965 (Paris et al., 2014 and references therein). These and many further examples emphasise that such potentially catastrophic tsunamis occur frequently and may pose a severe threat for coastal regions even hundreds of kilometres away from the source.

As historical databases reveal, the details of the tsunami triggering source are often a subject of debate. Even for the largest and deadliest volcanogenic tsunami, at Krakatau in 1883, discussed processes include caldera collapse, pyroclastic flows into the sea, sector collapse, explosion, or combinations thereof (e.g., Yokoyama, 1981; Francis, 1985; Nomanbhoy and Satake, 1995; Maeno et al., 2006). This highlights that there are multiple ways in which a volcano can cause a tsunami, which are considered a secondary volcanic hazard and result from eruptions or sector collapses of volcanic edifices (McGuire, 2006; Paris, 2015). Specifically, eruptions are known to generate tsunamis through large PDCs resulting from column or lava dome collapses entering the sea (Carey et al., 2000; Watts and Waythomas, 2003), through phreatic explosions when lava interacts with seawater (Smith and Shepherd, 1993; Belousov et al., 2000), and through the collapse of a lava delta when large amounts of lava construct unstable new land in the sea (Poland and Orr, 2014; Di Traglia et al., 2018). The formation of a caldera during particularly large eruptions is also known to generate tsunamis as large parts of the volcanic edifice are moving or collapsing (Maeno et al., 2006). On the other hand, a landslide or sector collapse (or lateral collapse) and resulting

debris avalanche from a volcanic edifice may also occur in association with volcanic activity, i.e., flank instability triggered through the intrusion of cryptodomes or dykes, earthquakes, explosions or loading of the flank with eruptive products (Lipman and Mullineaux, 1981; Siebert, 1984; Murray and Voight, 1996; McGuire, 2006; Romero et al., 2021). Flank instability can also be gravitationally driven without current volcanic unrest or eruptive activity through a deep-seated and slow-moving décollement (van Wyk De Vries and Borgia, 1996) and may be seen as a precursory stage of a catastrophic flank collapse. Hydrothermal alteration may further facilitate gravitational instability by altering the chemical and structural composition of the volcanic flanks or basement (van Wyk De Vries and Borgia, 1996; Heap et al., 2013; Heap et al., 2021). However, the interrelation between gravitational and magmatic flank instability are still poorly understood and may strongly depend on the local geologic setting and structural architecture of the volcano (Poland et al., 2017).

The inherent problem of volcanogenic tsunamis is the lack of warning time and quick response options because these tsunamis are not recognized by the early warning systems designed to detect tectonic earthquake events via seismic monitoring (Hanka et al., 2010; Lauterjung et al., 2010) and scenario-based modelling (Harig et al., 2020). This shortcoming also affected the 2018 tsunami originating from Anak Krakatau, where a warning system was in operation, but not designed to correctly interpret the moderate-sized seismic energy as a tsunamigenic event (Annunziato et al., 2019). ~~This may then allow for a targeted implementation of disaster mitigation strategies and warning systems at critical sites, e.g. by placing additional tide gauges as proposed for Krakatau (Annunziato et al., 2019) or improved volcano monitoring. To compensate for this blind spot and improve local preparedness, a comprehensive understanding of the tsunami hazards posed by volcanic sources is required to identify the most likely individual volcanoes to produce such an event.~~ Here, we present an in-depth analysis of the tsunami hazards by volcanoes in the Southeast Asian seas, including Indonesia, Papua New Guinea, the Philippines, and India. ~~While we incorporate some elements of review studies, which have been done extensively for the volcanic tsunamis in Southeast Asia (see e.g. Paris et al., 2014; Mutaqin et al., 2019), we expand on this by attempting to place the potential source volcanoes in a hierarchical order and identify the most likely volcanoes to cause tsunamis in the future. For this purpose, we~~We create a comprehensive catalogue of potentially tsunamigenic volcanoes and further use this data to create a point-based hierarchical ranking and identify the most likely candidates for sourcing potentially catastrophic tsunamis in the future.


## 2 Methods

### 2.1 Morphological evaluation and catalogue

For creating the catalogue we considered all active volcanoes in the SE-Asian region (Fig. 1), here including India, Indonesia, Papua New Guinea, and the Philippines, that were listed in the Global Volcanism Program (GVP) database (Global Volcanism Program, 2013) with a maximum distance of 20 km to the sea (with one exception: Peuet Sague, which has an uncertain historical tsunami associated with it). Volcanoes further inland were not considered as mass movements from eruptions or flank/sector collapses are unlikely to exceed such a distance, although in some circumstances this may still

be possible. Deposits of debris avalanches from sector failures of stratovolcanoes, e.g. the Gotemba deposit from Fuji volcano, Japan, were found as far as 24 km from the summit (Yoshida et al., 2012), or 35 km at Shiveluch, Kamtchatka (Belousov et al., 1999). Pyroclastic flows at Mt St. Helens, USA, also reached more than 25 km from the vent (Kieffer, 1981). However, these distances are exceptional and likely limited to very large volume collapses or highly energetic lateral eruptions.

(Kieffer, 1981; Yoshida et al., 2012).



**Figure 1:** The overview map of all active volcanoes located in the SE-Asian region. Considered for our catalogue (in colour) are all subaerial volcanoes in Indonesia, India, Papua New Guinea and the Philippines at least less than 20 km away from the nearest coastline. The colour corresponds to the country where a volcano is located. In total, there are 214 active volcanoes between the four countries and 131 of them are considered in our catalogue. The rectangles indicate the extent of the scoring maps in figure 3. Base map data source: © OpenStreetMap contributors 2022. Distributed under the Open Data Commons Open Database License-Licence (ODbL) v1.0.

The catalogue contains a total of 131 volcanoes, colour-coded according to the country in Fig. 1. The information on the individual volcanoes was collected from databases, including the GVP (Global Volcanism Program, 2013) and the Global

Historical Tsunami Database (NGDC, 2021), scientific literature, optical remote sensing data from Sentinel-2 satellites accessed via Sentinel Playground (ESA, 2022), as well as our own measurements using the Copernicus digital elevation model (DEM) (Airbus, 2020). [The Copernicus DEM is based on the previously available DEMs such as ASTER, SRTM90, SRTM30, SRTM30plus, GMTED2010, TerraSAR-X Radargrammetric DEM, and ALOS World 3D-30m.](#)

115 We started by extracting morphometric data from the Copernicus DEMs using the NETVOLC (Euillades et al., 2013) and  
MORVOLC (Grosse et al., 2009; Grosse et al., 2012) codes and automatically delineated the edifice boundary of the  
volcanoes. NETVOLC initiates with a starting location (usually the volcano summit) and a pixel range to draw the edifice  
boundary by iteratively looking for the minimum profile convexity. This is based on the assumption that volcanic edifices  
are bound by concave breaks in their slopes. Then, MORVOLC allowed us to use that boundary (and an optional crater  
120 outline, manually added) to collect data such as maximum elevation, slope steepness, or edifice volume for further  
evaluation. [A full example for the NETVOLC and MORVOLC output and an illustration of the edifice boundary definition  
is provided in the supplementary material A.](#) For most volcanoes in our catalogue this approach worked very well. For some  
volcanoes (28/131 cases or ~21%), however, we had to delineate the edifice boundary manually as the automatic NETVOLC  
approach would produce no or visibly wrong boundaries. This is due to the more complex morphology of some volcanoes,  
125 which does not allow for a clear delineation of concave slope breaks.

The final catalogue (see supplementary material [B](#)) includes (1) the coordinates of the volcano, (2) the volcano type, (3) its  
activity status as the last eruption year, (4) how many tsunamis it has historically sourced, (5) its height and distance from  
the sea (as a ratio) as well as (6) the maximum average slope angle, (7) the possible collapse azimuth range (flanks facing the  
sea) and (8) the most likely azimuth for tsunamis (manually selected base on slope steepness or location relative to the sea),  
130 (9) the edifice volume, and (10) a summary of other hazardous features (see below).

## 2.2 The volcanic tsunami-hazard ranking

One key challenge in creating a *meaningful* ranking of the tsunami hazard posed by many individual volcanoes is the  
application of consistent criteria to allow for strong comparability. Since volcanoes have a large variety of shapes,  
135 morphologies and styles of activity, this is not a trivial task. We therefore only considered the data representing all volcanoes  
under the same conditions and we used as many objectively measurable criteria as possible to minimise human subjectivity  
and inconsistency. Here, we decided on a Multicriteria Decision Analysis (MCDA), which is a method frequently used to aid  
in prioritising and decision-making for complex and multifaceted problems across many scientific disciplines. For example,  
in medical sciences evaluating the impact of drugs (Nutt et al., 2010), in Earth sciences assisting in the management of  
140 nuclear waste disposal sites (Morton et al., 2009), or in hazard evaluations of flood-prone sites (Fernández and Lutz, 2010;  
Rahmati et al., 2016; Toosi et al., 2019). Our MCDA system uses weighted point scores based on five major factors to

influence our final ranking. These factors were assigned (i) factor points based on their likelihood to contribute to the volcano's tsunami hazard, and (ii) factor weights based on how much the factor contributes to the volcano's tsunami hazard. By adding these weighed factor points to a final score we could rank the tsunami-hazard from the individual volcanoes:

145 
$$Score = F_1 \cdot W_1 + F_2 \cdot W_2 + \dots + F_n \cdot W_n \quad (1)$$

where  $F$  represents the individual factor points from 0-100 (100 representing the highest tsunami-hazard) and  $W$  represents the factor weight as a percentage (the total of all factor weights adding up to 100%). Due to the specific nature of some factors, the full 100 points could not always be reached, but the point systems were designed to scale well between 0 and 100. This results in a final score that also has points between 0-100 and makes our ranking easier to comprehend and visualise. While there are numerous factors that can be considered to reflect the tsunami hazard from a volcano, most of them do not have a known empirical relation to the hazard. For example, it is reasonable that a steep volcano close to the sea is more likely to produce a tsunami than a gently sloped one far inland, but exactly how much more likely this makes a tsunami is not known. With our ranking, we can therefore only aim to compare these factors by assuming that certain higher values equal a higher hazard.

150

We consider the following five factors and point systems for the ranking. Each represents a set of data that can be recorded or quantified objectively, which is then assigned a subjective but consistent point scale in order to create a comparable hierarchy:

155

i) **H/D-Ratio:** This is the height  $H$  of the volcanic edifice (i.e., the maximum elevation of the peak) as a fraction of the distance from that point to the sea  $D$ . To measure the height of volcano edifice, we used the 30 m Copernicus DEM GLO-30 (Airbus, 2020), which is considered the most reliable choice among the freely available and global DEMs (Guth and Geoffroy, 2021). Only for Anak Krakatau no post-collapse DEM was available, so we used a downsampled photogrammetric DEM previously published by Darmawan et al. (2020). We selected the maximum elevation of the edifice (extracted using the MORVOLC-code) and from that location measured the minimum distance to the shoreline using OpenStreetMap land polygons (OpenStreetMap, 2022). This parameter is highly relevant to the tsunami hazard of a volcano. Firstly, because a volcano with higher elevation can produce larger collapses and allow for more potential energy in mass movements, both from flank/sector/dome collapses and PDCs directed towards the sea. Secondly, if the mass movement source is closer to the sea it is more likely to actually reach it and produce a tsunami. The resulting values ranged from 0.02 to 0.89, so in order to convert the factor to the 0-100 point scale, the values were multiplied by 100. The only exception is Ritter Island, Papua New Guinea, which achieved a ratio of 1.77 due to the remnant collapse scar, which drops steeply towards the sea. We simplified this by assigning a maximum of 100 points.

160

165

ii) **Volcanic activity:** Frequent eruptive activity can exert strain on the flanks of a volcano as the volcano deforms, inflates and deflates as a result of pressurisation or the movement of intrusions. Additionally, erupted lava or

170

175 tephra can quickly pile up and over steepen flanks, while constant high seismic activity can act as a trigger to  
mass movements. This means a volcano with a high activity level is far more likely to experience a  
sector/flank/dome collapse or produce pyroclastic flows compared to a quiescent one. While the exact  
mechanisms and timescales are generally not well understood, it is known that many volcanic systems have  
extended periods or cycles of quiescence or low eruptive frequency, followed by more frequent eruptions (e.g.,  
180 Crisci et al., 1991; Gertisser and Keller, 2003; Turner, 2008). This suggests that a volcano that recently erupted  
is more likely to become more active or erupt again in the near-future compared to ones that have been quiet  
for a long time. Considering all above, we based our factor points on the time since the last known eruption of  
the volcano, with the data sourced from the GVP (Global Volcanism Program, 2013). However, due to the  
different timescales of eruptive cycles and the decreasing certainty of historic records, we decided to apply a  
non-linear point scale. Any volcano that erupted since the beginning of 2020 received a full 100 points. Every  
185 year before 2020 resulted in one point deduction and from the year 2000 only every decade is worth one point  
less, but to a minimum of 20 points as long as the volcano erupted within the Holocene. Similarly, if these  
volcanoes experienced historical unrest episodes but no eruption, they were assigned 20 points. For volcanoes  
where Holocene eruptions are presumed or considered likely, we assigned 10 points and 0 points were assigned  
for volcanoes that are presumed to be extinct.

190 iii) **Tsunamigenic history:** Some volcanoes have an increased tendency to produce tsunamis, either through  
frequent large eruptions or inherently unstable flanks. Here, we counted the number of tsunamis known from  
historic records sourced from the Global Historical Tsunami Database (NGDC, 2021) and relevant review  
papers (Hamzah et al., 2000; Paris et al., 2014; Mutaqin et al., 2019; Hidayat et al., 2020). Our compilation is  
shown in table 1. We further considered signs of previous edifice instability that include collapse scars  
195 (amphitheatres) and known submarine debris avalanche deposits. We assigned 10 points for every known  
historical tsunami, 10 for a collapse scar or submarine deposit and 20 if the volcano has multiple scars or  
submarine deposits.

200 iv) **Slope angle:** The steeper an edifice is the less stable it becomes. For volcanoes, steeper slope angles are  
correlated with higher frequencies of edifice failures (Siebert, 1984). Typically, the angle of repose for natural  
volcanic rocks lies between 30–40° and edifices exceeding this angle may experience gravitational instability.  
Here, we measured the maximum average slope angle that is part of the MORVOLC output, meaning the  
steepest part of the edifice calculated as the average slope value of the 50 m elevation interval of the edifice  
with the highest mean average slope, see Grosse et al. (2009). However, we limited the output to only those  
flanks facing towards the sea and excluded slopes facing inland. For the factor points, we doubled the steepness  
205 value in degrees, so a 50° slope would equal the maximum of 100 points. We also note that this factor is a  
major distinction from the H/D-ratio since it considers real measured local steepness. This may differ strongly  
from the H/D-Ratio, e.g., if the volcano is very steep, but the summit is located far from the sea.


v) **Hazardous Features:** This factor is deliberately designed to encompass a broad collection of features impacting the likelihood of a tsunami-generating event at the volcanoes. These features do not really fit a typical quantitative scheme cannot be easily quantified and have to be arbitrarily evaluated by a human and are easily missed or misinterpreted. Thus, in order to minimise user bias and avoid unrealistic points, all listed features below were combined into one factor. The data was collected by visually examining the Copernicus DEM GLO-30 (Airbus, 2020) as well as optical satellite imagery from Sentinel-2 satellites accessed via Sentinel Playground (ESA, 2022). The considered features include:

- Underwater extent: Whether a volcano and its flanks are submerged in the sea plays a very important role when assessing tsunami hazards as an edifice failure or an erupting vent may be located partially or fully underwater. We assigned 10 points if the edifice is partially submerged (so at least part of the edifice outline used in the NETVOLC and MORVOLC codes reaches into the sea) and 20 points if the edifice is fully in the water and all flanks reach into the sea
- Morphological features: Breached craters highlight a volcano's tendency to experience partial instability and provide an easy path for eruption products to reach the sea. We assigned 15 points to volcanoes with breached craters. Similarly, calderas signal that one or many large eruptions have occurred at this volcano and were also given 15 points. Finally, a larger volcanic edifice may have multiple secondary edifices and vents, adding further potential source locations to volcanogenic tsunamis. We thus added 15 points if one or more secondary peaks/vents were found.
- Vegetation: Dense vegetation and plant roots can significantly enhance a flank's stability and make a flank/sector collapse less likely (Gonzalez-Ollauri and Mickovski, 2017). Contrary, a flank's stability may significantly decrease due to the vegetation loss after major eruptions (Korup et al., 2019). Here, we visually inspected the volcano flanks using Sentinel-2 data (10 m spatial resolution). We gave no points if the volcano flanks were densely vegetated, 5 points if the volcano had vegetation free portions (most commonly, this is near the crater as a result of recent eruptions), and 15 points if at least one flank between summit and sea was free of vegetation.
- Hydrothermal alteration: Fumarole systems and weathering may significantly alter the rocks of a volcano flank, changing their appearance, composition and strength. Most commonly, hydrothermal alteration can weaken the rocks and promote failure or close permeable fluid paths and induce phreatic explosions (Heap and Violay, 2021). Alteration through geothermal systems is sometimes listed as the main reason for flank destabilisation, as with the 1979 landslide causing a tsunami at Iliwerung volcano, Indonesia (Yudhicara et al., 2015). Thus, we attempted to identify areas of localised alteration, by visually inspecting the volcano flanks using Sentinel-2 data. We specifically looked for characteristic bright spots indicative of alteration minerals and localised vegetation loss (outside the main crater). If these were identified, we added 10 points.




- 245
- Topography between an edifice and the sea: Within 20 km of the shore, the volcano's flank may not directly face the sea. While past events at St. Helens, USA, 1980 (Fisher, 1990) and Merapi, Indonesia, 2010 (Cronin et al., 2013) have shown that sector collapses or PDCs can overcome significant topography, it is less likely to reach the sea in such circumstances. To account for that, we added 30 points if we found no major topographic obstacle between the summit of the edifice towards the sea, thus adding more points to volcanoes close to the coast.

250 For the factor weights, we have to choose values based on the importance of the factor data. A higher weight of a factor will result in a larger impact of this factor on the final score and thus make it more important. Here too, these choices are largely subjective, but allow reducing the impact or importance of e.g. less reliable factor data and in-turn raise the impact of more reliable factors. We ~~For the chosen factor weights we~~ decided to favour morphometry and eruptive activity over the others factors. Morphometry, here meaning H/D-ratio and slope angle, measure both the feasibility of gravitational mass movements (flank collapses or PDCs) reaching the sea, as well as quantify oversteepening of individual flanks. This data  
255 also represents the most reliable quantitative data in our ranking as it can be precisely measured. Eruptive activity is also favoured as tsunamis do not only occur by flank collapse but also through PDCs or explosions. Additionally, unrest or eruptions may also act as a trigger for gravitational failures. In turn, we decided to weigh the Hazardous Features less since these are not quantitatively determined and more prone to human subjectivity and misjudgement. Thus, they are less reliable.  
260 Consequently, the final factor weights used were the H/D-ratio (20%) and the slope angle (20%) as morphometry factors, then volcanic activity (30%), tsunamigenic history (20%), and hazardous features (10%). An example of how the score was calculated is provided in Fig. 2.



c	H/D-Ratio	Volcanic Activity	Tsunamigenic History	Slope Angle	Hazardous Features	Total Score
Factor Value/ Feature	771/1560 = 0.50	Last erupted: 1968	None are known	Max: 34°	- Caldera (15) - Secondary vent (15) - Mostly vegetation free slope (5) - Hydrothermal Alteration (10) - Submerged edifice (20) - Unobstructed path to sea (30)	<b>56</b>
Factor Points (Weight)	50 (20%)	77 (30%)	0 (20%)	67 (20%)	95 (10%)	



265 **Figure 2:** Exemplary calculation of the ranking score using Nila volcano, Indonesia. a) is a false colour image from Sentinel-2 from bands 8 (NIR), 4 (red), and 3 (green). This highlights vegetated areas as red, but leaves barren areas grey. Here, Nila Island is densely vegetated, except for the southeast slope, which appears to be unvegetated due to ongoing hydrothermal alteration. b) is a hillshade DEM from Copernicus GLO30 and c) is a summary of our MCDA score calculation using data marked in a) and b) as well as the GVP and Global Historical Tsunami databases.

270

We further tested how robust our ranking is with respect to used factor weights. This is done to confirm that the highest scoring volcanoes still retain their high score even when the weighing is significantly different, which can confirm that these volcanoes really pose the highest tsunami hazard despite possible human error or misjudgement. The test was carried out by

275 changing the five factor weights, increasing one factor to 60% and all others are set to 10%. The procedure was repeated for  
 all five factor weights, so that every single factor was once set as the strongest influence. We also added one instance of all  
 weights being considered equal (i.e., all five factors being weighed at 20%). This then enabled us to calculate both an  
 average score and standard deviations within this variability of weights. The results could then be used to judge whether our  
 ranking can generally identify the highest scoring and most hazardous volcanoes well, despite the subjective weight choices.  
 280 We could further determine which volcanoes were more sensitive to the influence of single factors, as this would result in  
 higher deviations.  
~~We then counted how many times the volcanoes would be placed within the top 5, 10, 15, and 20 spots on the ranking. For  
 example, if a volcano is always in the top 5, no matter how the factor weights change, then it is 5 times in the top 5, 10, 15,  
 and 20 each, giving it the maximum count of 20. However, if the volcano is once in the top 10, twice in the top 15 and four  
 285 times in the top 20 it receives a count of 7. The resulting compilation was used to judge whether our ranking can generally  
 identify the highest scoring and most hazardous volcanoes well.~~

Volcano	Country	Year	Deaths	Cause	References
Agung	Indonesia	1963			3, 5
Awu	Indonesia	1856	3000	Pyroclastic flows	1, 2, 3, 4, 5
Awu	Indonesia	1892	1532	Pyroclastic surges?	1, 2, 3, 4, 5
Awu	Indonesia	1913		Pyroclastic surges?	4
Banua Wuhu	Indonesia	1889		Underwater explosion?	2, 3, 4
Banua Wuhu	Indonesia	1918		Underwater explosion	2, 3, 4
Banua Wuhu	Indonesia	1919		Underwater explosion	2, 3, 4
Gamalama	Indonesia	1608	<50		2, 3, 4, 5
Gamalama	Indonesia	1771			3
Gamalama	Indonesia	1772	35		4
Gamalama	Indonesia	1840			2, 3, 4
Gamalama	Indonesia	1871			1

Gamkonora	Indonesia	1673		Earthquake/ Landslide	5
Gamkonora	Indonesia	1673			2, 3, 5
Iliwerung	Indonesia	1973		Underwater explosions?	2
Iliwerung	Indonesia	1979	>539	Landslide	2, 3, 4
Iliwerung	Indonesia	1983		Underwater explosion	2, 4, 5
Krakatau	Indonesia	416	<1000		3, 4, 5
Krakatau	Indonesia	1883	34417	Pyroclastic flows	1, 2, 3, 5
Krakatau	Indonesia	1883		Landslide?	2, 5
Krakatau	Indonesia	1884		Underwater explosion?	2, 3, 4
Krakatau	Indonesia	1928		Underwater explosion	1, 2, 3, 4, 5
Krakatau	Indonesia	1930		Underwater explosion	2, 3, 4, 5
Krakatau	Indonesia	1981		Landslide?	2, 3
Krakatau	Indonesia	2018	437	Landslide	4
Makian	Indonesia	1550			2
Peuet Sague	Indonesia	1837			3
Rokatenda/Paluweh	Indonesia	1927	226	Underwater explosion?	1, 4
Rokatenda/Paluweh	Indonesia	1928	98	Landslide	1, 2, 3, 4
Ruang	Indonesia	1871	>400	Lava dome collapse	2, 3, 5
Ruang	Indonesia	1889			1
Soputan	Indonesia	1845	118	Earthquake?	2, 3
Tambora	Indonesia	1815	<1000	Pyroclastic flows	1, 2, 3, 4, 5

Teon/Serawerna	Indonesia	1659		Pyroclastic flows?	2, 3, 4, 5
Unknown Volcano	Indonesia	1773			5
Unknown Volcano	Indonesia	1878			5
Unknown Volcano	Indonesia	1883		Pyroclastic flows?	3
Unknown Volcano	Indonesia	1892			5
Unknown Volcano	Indonesia	1918			5
Unknown Volcano	Indonesia	1919			5
Kadovar	Papua New Guinea	2018		Lava dome collapse	5
Long Island	Papua New Guinea	1660		Pyroclastic flows?	2, 5
Rabaul	Papua New Guinea	1878		Earthquake	2
Rabaul	Papua New Guinea	1937	<50	Pyroclastic flows/ Explosions	2, 5
Rabaul	Papua New Guinea	1994		Pyroclastic flows	2, 5
Ritter Island	Papua New Guinea	1888	<3000	Landslide	2
Ritter Island	Papua New Guinea	1972		Underwater explosions?	2, 5
Ritter Island	Papua New Guinea	1974		Landslide?	2, 5
Ritter Island	Papua New Guinea	2007		Landslides?	2
Unknown Volcano	Papua New Guinea	1857		Earthquake	2
Unknown Volcano	Papua New Guinea	1953			5
Bulusan?	Philippines	1933	9		2, 5
Camiguin	Philippines	1871		Pyroclastic flows?	2, 5
Didicas	Philippines	1969	3	Underwater explosions?	2, 5

Taal	Philippines	1716		Underwater explosions	2, 5
Taal	Philippines	1749		Pyroclastic flows?	2, 5
Taal	Philippines	1754	12	Pyroclastic flows	2, 5
Taal	Philippines	1911	>50	Pyroclastic surges/ Air waves?	2, 5
Taal	Philippines	1965	355	Air waves?	2, 5

**Table 1:** Compiled list of historic Tsunami events in the Southeast Asia region. References are 1: Hamzah et al. (2000); 2: Paris et al. (2014); 3: Mutaqin et al. (2019); 4: Hidayat et al. (2020); 5: NGDC (2021). A cause with question mark indicates that the actual cause is unclear and/or debated.

### 2.3 Ranking assumptions and limitations

There are many factors influencing the tsunami hazard posed by volcanoes and not all of them can be quantitatively evaluated. Furthermore, even for the ones which can be evaluated, certain assumptions have to be made in order to create a comparable baseline to judge all volcanoes on an equal basis. Here we address some of the issues with the data used in our ranking approach:

It is important to emphasise that the point system used in our MCDA is based on arbitrary and thus subjective point scales, assigned to best cover the range of values used to build the ranking score. Any evaluation of this type will inherently involve arbitrarily chosen scores that have to consider and weigh the many multifaceted factors that contribute to volcano flank instability and the hazards of generating a tsunami. In studies previously ranking volcanic hazards and risks, these could be done by a simple count of “yes” or “no” features adding 1 or 0 points, respectively (Yokoyama et al., 1984), or similar variations (Ewert, 2007; Ewert et al., 2018), or via the creation of index values and adding them up to create a score (Scandone et al., 2015). Then categories (e.g. high, medium and low hazard) are defined to best cover the range of scores (e.g. Ewert et al., 2018), which is also what we do in our ranking. MCDAs in other fields often have more quantitative scales such as 0-9 points (Fernández and Lutz, 2010; Rahmati et al., 2016) or 0-100 (Nutt et al., 2010), but the score systems are still assigned arbitrarily. Thus, all these approaches and our ranking presented here use some degree of subjective judgement, as not all factors directly translate into an empirical hazard or risk value. Without this, no meaningful comparison between volcanoes could be made. However, as the rules with which points are given are kept strictly the same for all volcanoes, the comparability of scores is retained, allowing for a meaningful hierarchical order or scores. For our ranking, this means that the hazard score by itself should be seen as a rough hierarchy estimation rather than a strict empirical value as it has little

315 meaning in terms of hard data, such as expected tsunami event frequency, possible wave heights, or impacts on shorelines and population. Similarly, we can thus not adequately assess the risk to shores and population in the traditional sense. Instead, we identify which volcanoes are the most likely to cause a tsunami in the future as these are expected to produce the highest hazard score. This also means that the factor points and weights are based on our subjective judgement by how important we think these factors are. Naturally, this leaves a lot of room for arguments on how the points and weights could be assigned differently, which may significantly change the final score. However, as the rules with which points are given are kept strictly the same for all volcanoes, we retain the comparability of our results, allowing a meaningful identification of the most hazardous volcanoes for tsunami generation. This is despite the individual score by itself having little meaning in terms of hard data, such as expected tsunami event frequency, possible wave heights, or impacts on shorelines and population. Similarly, we can thus not adequately assess the risk to shores and population in the traditional sense.

320 Another important consideration is that we cannot consider all factors that are known or suspected to impact the tsunamigenic potential of a volcano. The most noteworthy ones are (1) ongoing flank deformations (e.g., through dyke intrusions or slow décollement movement), which can destabilise parts of the edifice, and (2) absence of bathymetric data covering the underwater geometry of the volcanoes. Because the lack of high quality and accessible data for all volcanoes makes meaningful comparison of the impact of these factors on the tsunami hazard impossible, we could only consider these factors by proxy. For instance, eruptive activity of a volcano increases the likelihood of ongoing deformation and thus it is indirectly tied to the activity factor points. However, not all erupting volcanoes are likely to experience significant deformation and, on the other hand, there may be significant deformation without an eruption (e.g., due to the intrusion of magma underground). For the underwater geometry, our analysis is simplified to whether the volcano is partially or fully in the water, thus counting towards the hazardous features factor points. But it neglects potential steep underwater flanks, which also means that some potentially hazardous submarine volcanoes had to be excluded from our catalogue as they had no subaerial edifice to examine. Here, these are Banua Wuhu, Indonesia, and Hankow Reef, Papua New Guinea as well as two unnamed seamounts. Future studies could take these missing factors into account properly and warrant their own point scores, provided that data quality, regularity, and availability of surveys improve.

335 Conducting a comparative ranking can be more challenging if there are major dependencies between the used factors. As an example for our case, it would be reasonable to assume that recent eruptive activity would more likely cause hydrothermal alteration, thus making the eruptive history and hazardous features factors interdependent. However, in our catalogue, only few volcanoes are recorded to have extensive hydrothermal alteration on their flanks and for many of these, no eruption occurred for decades to centuries (e.g. Manuk, Teon, Serua). Hence, we think that these issues are unlikely to significantly affect our results. The only exception is a direct dependence between the H/D-ratio and the slope angle as it is essentially the same value if the volcano is close to the coast, however the separation does allow for a more distinct look at volcanoes that may be far from the coast, but still have steep slopes on a local level.



Data based on historic observations is often flawed as they were not always systematically recorded. This includes historical tsunamis, flank collapses, and eruptions. The problem is facilitated the further the events lie back in time, and many events may be missed or wrongly interpreted because they were either not understood properly or simply not noticed or remembered. For instance, some volcanoes have unknown eruption dates and thus received a relatively low rating, even when it is quite likely that its last eruption happened only decades ago. One example is Balbi volcano, Papua New Guinea, which likely erupted around the early 1800s, but this is unconfirmed (Global Volcanism Program, 2013). This creates a slight bias towards volcanoes that are well-known and researched, whereas volcanoes with less scientific attention may receive lower ratings. Thus, our analyses may be prone to missing or incorrectly recorded events. This is also the reason we prefer to use just the last eruption year known from a volcano rather than creating a score based on eruption frequency as these frequencies can simply not be reliably established for many volcanoes in our catalogue. For tsunamis, we tried to minimise this bias by incorporating data from multiple databases and studies (Hamzah et al., 2000; Paris et al., 2014; Mutaqin et al., 2019; Hidayat et al., 2020; NGDC, 2021). Surprisingly, no study had the exact same tsunami events listed and some were only found in certain reviews. Our compilation thus likely marks the most comprehensive list of historic tsunamis in SE-Asia to date, however, there are likely some further events that were missed or not recorded.

For evidence of edifice instability the problem is similar. Many collapse scars related to such instability or lateral blasts can be obscured by the regrowth of the volcano and are thus easy to miss. On the other hand, gradual erosion and destabilisation due to hydrothermally weakened rocks (Darmawan et al., 2022) may produce scars that are very similar to collapse scars and consequently very challenging to distinguish using satellite data. Finally, submarine debris avalanche deposits which are evidence for past edifice failures that reached the sea are poorly studied since the required bathymetry data is rarely acquired. Here, Silver et al. (2009) was specifically investigating volcanic debris avalanches for most volcanoes in Papua New Guinea, so this region can be considered reasonably well covered. However, no similar studies exist for Indonesia or the Philippines, making some oversights likely.


When analysing the vegetation cover of the volcanoes using the Sentinel-2 satellite images we found that the vegetation is subject to seasonal changes. Mainly this encompasses vegetation colour changes due to dry or rainy periods. Since our analysis is kept rather simple and qualitative rather than quantitative, this effect is unlikely to impact our results.


While we consider gravitational instability of volcanic edifices by measuring the slope steepness, additional factors can play an important role that could not be considered here as they are unknown for most volcanoes. These are mainly the lithological properties of the flank and the rock mass strength (Watters et al., 2000). Furthermore, the nature of the volcanic flanks deposits and their state of consolidation is likely to play a role on how resistant they are to oversteepening and destabilisation. For instance, one factor that is speculated to have contributed to the 2018 Anak Krakatau flank collapse is that it consisted of loose unconsolidated pyroclastics covered with lavas (Grilli et al., 2019), which likely contributed to the instability.

## 375 3 Results

### 3.1 Volcano catalogue and ranking

Using the factor points and weights described in section 2.2, we ranked the volcanoes in our catalogue by their tsunami hazard. An overview is presented in Fig. 3 and a list of the 40 highest-scoring volcanoes is shown in table 2. A complete, more detailed, and interactive version of this list with individual entries relating to how the points were counted can be found in ~~the~~ supplementary material [B](#). The points range from 76 (Batu Tara, Indonesia), representing the highest tsunami hazard, to 13 (Baluan, Papua New Guinea), representing the lowest tsunami hazard. Using these results, we grouped the volcanoes into high, medium, and low tsunami hazard categories based on their relative score. High hazard is assigned for volcanoes having more than 55 points (~14% of the volcanoes in the catalogue), medium hazard is assigned to volcanoes with scores between 40 and 55 points (~36% of the volcanoes in the catalogue) and low hazard category is assigned to volcanoes with less than 40 points (~49% of volcanoes in the catalogue). Volcanoes like Anak Krakatau, Indonesia, and Ritter Island, Papua New Guinea are among the highest tsunami hazard volcanoes in our ranking, which is ~~little-surprise~~ [unsurprising](#) considering both their history of powerful eruptions and catastrophic tsunamis (Paris et al., 2014). However, we also identify ~~high~~ high-hazard volcanoes that are not as ~~prominently-considered~~ [well known](#) for their tsunamigenic potential, but received similarly high scores. In Indonesia these include Batu Tara, Gamalama, Iliwerung, Karangetang, Nila, Sangeang Api, Wetar, Sirung, Serua and Ruang. For Papua New Guinea we identify Kadovar, Rabaul, Ritter Island, Manam, Bam, Langila and Ulawun as volcanoes with high tsunami hazard. For the Philippines, only Didicas is classified as a high tsunami hazard volcano.





395 **Figure 3:** Maps of the volcanoes in the catalogue with their corresponding ranking score and the resulting hazard category [in](#) [\(a\) Indonesia](#), [\(b\) Philippines](#), [\(c\) Papua New Guinea](#) and [\(d\) India](#). For a detailed overview of tsunami hazard scores for all volcanoes in the catalogue, see table 2. Base map data source: © OpenStreetMap contributors 2022. Distributed under the Open Data Commons Open Database [License-Licence](#) (ODbL) v1.0.

<b>Volcano</b>	<b>Country</b>	<b>Points: H/D-Ratio</b>	<b>Points: Volcanic Activity</b>	<b>Points: Tsunamigenic History</b>	<b>Points: Slope Angle</b>	<b>Points: Hazardous Features</b>	<b>Total weighted Score</b>
Batu Tara	Indonesia	83	95	20	100	65	<b>76</b>
Anak Krakatau (pre-2018)	Indonesia	52	100	80	60	80	<b>76</b>
Kadovar	Papua New Guinea	79	100	20	74	80	<b>72</b>
Ritter Island	Papua New Guinea	100	87	60	39	65	<b>72</b>
Iliwering	Indonesia	45	100	40	70	70	<b>68</b>
Anak Krakatau	Indonesia	21	100	90	36	80	<b>67</b>
Gamalama	Indonesia	40	98	50	63	70	<b>67</b>
Rabaul	Papua New Guinea	62	94	30	67	70	<b>67</b>
Manam	Papua New Guinea	41	100	20	76	90	<b>66</b>
Didicas	Philippines	89	78	10	73	65	<b>64</b>
Sangeang Api	Indonesia	36	100	20	67	70	<b>61</b>
Karangetang	Indonesia	49	100	0	72	70	<b>61</b>
Langila	Papua New Guinea	29	100	20	65	75	<b>60</b>
Sirung	Indonesia	47	95	0	62	85	<b>59</b>
Ulawun	Papua New Guinea	25	100	20	68	60	<b>59</b>
Wetar	Indonesia	86	50	10	86	65	<b>58</b>
Nila	Indonesia	50	77	0	67	95	<b>56</b>
Ruang	Indonesia	45	82	20	64	55	<b>56</b>
Bam	Papua New Guinea	50	76	20	68	55	<b>56</b>
Serua/ Legatala	Indonesia	70	73	0	67	65	<b>56</b>
Lewotolok	Indonesia	40	100	0	56	55	<b>55</b>
Agung	Indonesia	25	99	10	65	50	<b>55</b>


Paluweh/ Rokatenda	Indonesia	32	93	20	49	65	<b>55</b>
Barren Island	India	33	100	0	48	80	<b>54</b>
Teon/ Serawerna	Indonesia	51	71	20	63	60	<b>54</b>
Rinjani	Indonesia	20	96	0	67	75	<b>54</b>
Iya	Indonesia	49	77	10	64	60	<b>54</b>
Banda Api	Indonesia	52	79	0	62	70	<b>54</b>
Gamkonora	Indonesia	33	87	20	53	60	<b>53</b>
Karkar/ Uluman	Papua New Guinea	26	94	10	49	80	<b>53</b>
Awu	Indonesia	24	84	40	48	55	<b>53</b>
Taal	Philippines	2	100	50	21	75	<b>52</b>
Soputan	Indonesia	10	100	20	53	50	<b>52</b>
Mayon	Philippines	25	99	0	64	35	<b>51</b>
Makian	Indonesia	36	79	10	57	65	<b>51</b>
Biliran	Philippines	32	74	20	70	40	<b>51</b>
Camiguin	Philippines	30	76	10	63	65	<b>50</b>
Bagana	Papua New Guinea	12	100	0	70	35	<b>50</b>
Babuyan Claro	Philippines	32	73	20	43	90	<b>50</b>

**Table 2:** List of the 40 highest scoring volcanoes in our catalogue and their respective MCDA ranking of the relative tsunami hazard, showing individual factor points and the final score. The factor weights for the total score are H/D-ratio (20%), volcanic activity (30%), tsunamigenic history (20%), slope angle (20%), and hazardous features (10%). For the full catalogue and ranking table see [the-supplementary material B](#).


The results of the ranking robustness testing are summarised in Fig. 4, showing that the higher scoring volcanoes in our ranking [with our chosen weights generally also score the highest average points independent of the factor weight. This adds confidence to our results and confirms that we are generally able to identify the higher hazard volcanoes well, even if their](#)

410 individual rank may vary slightly (Fig. 4a), generally still score high and are independent of the factor weight, which adds  
confidence to our results. The only notable exceptions here are at the transitions between low and medium hazard categories  
(e.g., Hiri, Indonesia, or Tolokiwa, Papua New Guinea) in Fig 4a and 4b, which is indicating that these volcanoes are likely  
more sensitive to individual weights and, thus, less robustly ranked. The lower the volcanoes are ranked, the less robust the  
415 ranking order becomes, which is likely due to a higher number of volcanoes with similar scores (Fig 4b, c). Here, this means  
that our ranking is most meaningful for the higher scoring volcanoes and puts them in a more reliable order. Standard  
deviations vary between 7 and 15 points for each individual volcano, being higher when the scoring strongly depends on  
single factors. For instance, the pre-2018 tsunami Anak Krakatau achieved high points in all factors and is thus not very  
variably in its total score, while Batu Tara has low points in tsunamigenic history and may thus have a greater variance in its  
total score for different weights (Fig. 4a).

420








**Figure 4:** Robustness test of the factor weights used in the ranking. This was done by calculating an average score and standard deviations from repeat scoring while systematically changing the factor weights. It shows that the volcanoes we classed as high hazard volcanoes are generally well distinguished, with the highest values being independent of factor weights. This demonstrates that This counts the number of times a volcano made it into the top 5, 10, 15 and 20 places in the ranking with different individual weights. This is displayed per countries a) Indonesia, b) Papua New Guinea and c) Philippines. It generally shows that the volcanoes we classed as high, medium and low hazard are well generally sorted in the order we classed them in, despite different factor weights and with only few exceptions. This shows that changing the factor weights may slightly change the order in which the volcanoes are ranked, but our analysis is generally classifying

higher hazard volcanoes correctly, confirming the robustness of our ranking. However, for the medium and low hazard volcanoes the ranking is less robust, due to a high number of volcanoes with similar scores, which can significantly change the hierarchical order depending on the chosen factor weights.

435 **3.2 Volcano distribution and tsunami causes**


Most of the high and medium tsunami hazard volcanoes are located in Indonesia, which by itself is not surprising since Indonesia also has the most volcanoes in our catalogue (~46%, Fig 5a). However, the relative amount of these volcanoes in those categories is significantly higher (Fig. 5), suggesting that Indonesia has an over-proportionally high number of hazardous volcanoes. This is further evident in the low tsunami hazard category, which are dominantly from Papua New Guinea (Fig. 5c, d). Volcanoes of the Philippines are only underrepresented in the high tsunami hazard category, but this may be due to the lower number of overall volcanoes.



445 **Figure 5:** Tsunami hazard from individual volcanoes by country. Shown are (a) all considered volcanoes by country and (c) their hazard categories according to our ranking. These are then split in their and their hazard categories according to

our ranking, then their respective distribution between the countries for (b) high hazard, (ed) medium hazard, and (de) low hazard volcanoes. We find a disproportionately large number of high and medium hazard category volcanoes located in Indonesia (~64% and ~59%, respectively), which is much larger than expected since Indonesian volcanoes make up only ~46% of all volcanoes in our catalogue. Contrary, the Philippines have a lower share of high-hazard volcanoes (~4%, only one volcano), but make up 18% of all volcanoes in our catalogues. Papua New Guinean volcanoes make up ~35% of the catalogue, so they can be considered underrepresented in the medium hazard category and overrepresented in the low hazard category.

Our review of historical tsunamis in Southeast Asia contains 61 distinct events (Table 1) and shows that the majority of the historical volcanogenic tsunamis still have an unknown or uncertain cause (Fig. 6). However, we can still extract that the most known or suspected causes were explosions (21%), followed by mass movements like pyroclastic flows (19%) and landslides (14%). Volcanic earthquakes rarely cause tsunamis by themselves since volcanic quakes are usually much weaker compared to tectonic events and lava dome growth is a rather specific eruption style and thus less frequent in producing tsunamis (here only 2 cases). Some cascading events also occurred involving multiple causes (here 3 cases), which were once an earthquake and a landslide, and twice an explosion and pyroclastic flow. It is noteworthy that most known cases of volcanogenic tsunamis are produced by pyroclastic flows and explosions, both of which are commonly associated with strong eruptive activity. Gravitational failures such as landslides or lava dome collapses occur less frequently, but do not necessarily require ongoing eruptions.



**Figure 6:** The historical causes of volcanogenic tsunamis in SE-Asia sourced from table 1, uncertain causes are marked transparent. While there are still many unknown tsunami sources, about 66% including both unknown and uncertain events, most known cases are A third of all cases is caused by mass movements (pyroclastic flows and landslides) and about a fifth is attributed to explosions, whereas earthquakes and lava domes rarely produce tsunamis. For the purpose of this figure, air waves were grouped with explosions as they also require a strong explosion.

## 4 Discussion

### 4.1 Evaluating scores and ranking through recent tsunami events

In ~~recent years~~ 2018, both Anak Krakatau, Indonesia, and Kadovar, Papua New Guinea, produced tsunamis, with good-quality topographic data available and their circumstances being well known (Plank et al., 2019; Walter et al., 2019). The changes resulting from the events are displayed in figure 7. One important aspect when judging the usefulness of our ranking is its ability to correctly identify volcanoes that are most likely to produce tsunamis in the future. We tested this by comparing the scores of both volcanoes as it is now compared to how it was before the ~~event~~ tsunami events. The morphology of Anak Krakatau has changed quite significantly following the 2018 flank collapse (Darmawan et al., 2020). The island is reduced in size and became lower in elevation, which consequently reduced the H/D-ratio factor points (Fig. 7a-d). However, it now has a crater that is open to the sea and it still has a history of many tsunamis and regular recent eruptions, thus only changing the score slightly from 76 points before the collapse to 67 points after (Table 2). Consequently, Anak Krakatau was the highest hazard volcano before its collapse and tsunami occurred, confirming that our approach would have correctly identified the volcano as a threat. Now, after the collapse, it is still among the highest scoring volcanoes, meaning the recent 2018 tsunami and the event changed little in its overall status. However, the 2018 flank collapse indeed lowered the score, which is reasonable since the main volcanic edifice has yet to rebuild after the collapse. Further tsunamis through further collapses, explosions or pyroclastic flows are still potential tsunami causes that may occur at Anak Krakatau in the near future.


For Kadovar, the changes were different. While the 2018 eruption produced a new vent and formed a littoral lava dome (Plank et al., 2019), the overall topography of the island did not change significantly (Fig. 7e-g). Here, the tsunami was not generated as a result of a major flank collapse but rather due to a collapse of the littoral dome and smaller parts of the southern flank (Plank et al., 2019). However, before the eruption, Kadovar had no known historic eruptions, although it is possible that one occurred in 1700, but this is unconfirmed (Llanes et al., 2009; Global Volcanism Program, 2013). In 1976 the island residents were briefly evacuated due to strong fumarolic activity and fears of an eruption, which ultimately did not occur (Llanes et al., 2009). Thus, before its 2018 eruption, the island could only be considered to have historic unrest, which lowered the score significantly. The score was 45 points before the 2018 tsunami, meaning Kadovar would still have been identified as a volcano with an elevated hazard (medium category) before the eruption. However, given its high eruptive

activity now, the score has increased strongly. If the unconfirmed eruption in 1700 is included, the hazard score would be significantly higher at 54 points, just outside the high hazard category. This highlights that a better constrained volcanic history on some volcanoes can significantly improve the meaningfulness of this ranking. Now, after the 2018 tsunami, Kadovar scores 72 points, making it the third highest tsunami hazard in SE-Asia based on our ranking. Both the steep south-facing flank and the littoral lava dome are still in place and may pose a large tsunami hazard in the future, especially if the eruptive activity continues.

From both these cases we can conclude that our ranking system is able to identify hazardous volcanoes reasonably well before a tsunami occurs. While individual volcano scores may change significantly over time due to eruptions or morphological changes, the applied multi-categorized approach has worked well for both Anak Krakatau and Kadovar. However, these cases also emphasise that particular attention should be devoted to coastal volcanoes with unclear eruptive histories, especially when they become active after decades or centuries of no activity.

We further investigate the score of the currently highest ranking volcano with our method, Batu Tara, Indonesia, with 76 points. This lone and small volcanic island is located north of the Lesser Sunda Islands and positioned centrally between the Flores- and Banda Sea. The island's high eruptive activity is well documented and appears to be mostly strombolian to vulcanian (Spina et al., 2021), but also pyroclastic flows and rockfalls were reported to reach the sea (Global Volcanism Program, 2013). The volcano is also exceptionally steep with the eastern flank measuring an incline of about 50 degrees. Despite its relatively small size, the Island has an elevation of 753 m, making it 2.5 times higher than Anak Krakatau before its collapse. Despite its activity and morphology, no recorded instances of historical tsunamis are known from this volcano, although smaller past events could easily have been missed as the island is uninhabited. However, there are multiple scars on different sides of the island, indicating major past collapse events (Fig. 7h, i). All this results in very high scores in all factors except its tsunamigenic history.

Batu Tara is often compared to Stromboli volcano, Italy, which has a similar style of eruptive activity and morphological features, including multiple sector collapse scars (Laiolo et al., 2018; Spina et al., 2021). The morphology of Batu Tara's eastern flank is strikingly reminiscent of the Sciarra del Fuoco, both of which channel recent eruptive material towards the sea, which is clearly distinguishable in the loss of vegetation on these flanks (Fig 7h). Contrary to Batu Tara, Stromboli is inhabited and a well-known tsunami hazards evident both through its sector collapses in its geological history (Tibaldi, 2001; Tinti et al., 2003) and more recent tsunamis such as in 2002 caused by partial flank failure and landslide (Bonaccorso et al., 2003; Fornaciai et al., 2019). For this reason, the high score of Batu Tara is very likely justified and its consideration of a high-hazard tsunamigenic volcano reasonable. Since this volcano is a lot less known compared to islands like Anak Krakatau, future monitoring efforts and tsunami risk modelling should make Batu Tara a high priority target.



**Figure 7:** Remote sensing data for Anak Krakatau (a-d) and Kadovar (e-g) and Batu Tara (h, i) volcanoes. Anak Krakatau and Kadovar are shown before and after their 2018 tsunamis. Sentinel-2 images in false colour bands 8 (NIR), 4 (red), and 3 (green) show a drastic reduction of vegetation (seen as red) after both events due to the volcanic activity. For the scoring, this mainly affects Kadovar, which appeared fully vegetated before the event (e). The DEMs also illustrate the massive morphological changes on Anak Krakatau due to the flank collapse (b, d), which strongly affected the H/D-ratio and the slope angle in our scoring. Both islands also show amphitheatre collapse scars (dashed lines) visible on the DEMs. Notably, the new failure scar from Anak Krakatau after the collapse (d) is nearly identical to the scar at this volcano before the collapse (b). On the other hand, the old scar at Kadovar (g) did not produce another failure, but the tsunami was triggered by a partial collapse at the newly grown littoral lava dome. For Batu Tara we highlight the development of a steep slope

reminiscent of the Sciara del Fuoco at Stromboli volcano, Italy, where scoria, lava flows and PDCs are directed towards the sea. The island also features multiple collapse scars, suggesting past flank failures.


540

#### **4.2 Future tsunami hazards in SE-Asia**


To allow for a more detailed look at future tsunami hazards in SE-Asia we summarised in which locations a high concentration of hazardous volcanoes is located. This was done by performing a weighted point density calculation, highlighting areas where many tsunamigenic volcanoes are located closely together, with their impact multiplied by the hazards score (Fig. 78). The result shows that the area with the highest volcanogenic tsunami hazard is located around the Indonesian Lesser Sunda Islands, particularly between East Nusa Tenggara and the Alor archipelago, at the Molucca Sea coast between northern Sulawesi and Halmahera, and at the southern Bismarck Sea in Papua New Guinea. Further elevated hazard areas can be found within the Indonesian Banda Sea, the Philippine Luzon Strait, the central Philippine Islands, and along the southern Solomon Sea coast of Papua New Guinea. These areas can thus be considered to be the most likely areas to source volcanic tsunamis and are therefore important to prioritise for tsunami monitoring, modelling and forecasting. A brief feature of individual high-hazard volcanoes can be found in the supplementary material C.

545

550







555 **Figure 78:** Travel-distance plots of tsunamis originating at a high-hazard volcano after 60 minutes in a) Indonesia and b)  
 Papua New Guinea. How much coastline is affected by each volcano is summarised in table 3. The additional heat map  
 shows the most likely source locations based on a weighted point density calculation using our hazard score. A higher  
 density means the area is closer to many tsunamigenic volcanoes. This highlights that particularly the Lesser Sunda Islands  
 and the northern Molucca Sea in Indonesia as well as the southern Bismarck Arc in Papua New Guinea are likely hot-spots  
 560 for future volcanogenic tsunamis and events from high hazard volcanoes identified by our ranking may affect large amounts  
 of coastline in a short time. Base map data source: © OpenStreetMap contributors 2022. Distributed under the Open Data

Commons Open Database Licence (ODbL) v1.0. Heat map resulting from weighted point density calculation using the

565 In order to assess the risks and impacts of volcanogenic tsunamis, numerical simulations are commonly used, both for  
distinct future scenarios and in retrospect for past events. For Southeast Asia, a large number of such studies had been  
conducted. Most models were done for Anak Krakatau looking specifically at the 2018 flank collapse with some using the  
known event to calibrate and confirm the quality of current simulation methods (Grilli et al., 2019; Borrero et al., 2020;  
Mulia et al., 2020; Omira and Ramalho, 2020; Paris et al., 2020; Zengaffinen et al., 2020), some using the known tsunami  
data (e.g. from tide gauges) to identify source parameters (Heidarzadeh et al., 2020; Ren et al., 2020; Grilli et al., 2021) and  
570 some testing variations in the source parameters to characterise potential future events (Dogan et al., 2021). In general, the  
consensus is that a landslide between 0.1 and 0.3 km<sup>3</sup> volume that occurred both with a subaerial and a submarine  
component is mostly consistent with the observed and modelled runup heights at the adjacent shores. Similar models also  
exist for the 1883 tsunami at Krakatau, with the main purpose being the identification of its generation mechanism (Maeno  
and Imamura, 2011) and how such a tsunami propagates in the far-field (Choi et al., 2003). Predictive studies only  
575 considering possible future events are not as abundant, but have been done for Anak Krakatau before the 2018 tsunami  
(Giachetti et al., 2012; Badriana et al., 2017), with Giachetti et al. (2012) making a remarkably close prediction to the later  
event. Other volcanoes in Southeast Asia are not as commonly considered. Pranantyo et al. (2021) test the tsunami  
propagation from Ruang volcano, Indonesia, using and comparing both historical observations and data from the 2018 Anak  
Krakatau event and reproducing a 25 m runup in the near-field. In Papua New Guinea numerical tsunami models have  
580 almost exclusively been considered for the Ritter Island tsunami in 1888 and the reconstruction of its generation (Ward and  
Day, 2003; Karstens et al., 2020). Similarly, numerical tsunami models in the Philippines are mostly limited to Taal volcano,  
where models are based both on a past tsunami in 1716 (Pakoksung et al., 2021) and a predictive study considering scenarios  
with different explosion sites and energies (Paris and Ulvrova, 2019). Considering these works, it is clear that tsunamis  
sourced by volcanoes can be well explained with numerical models, but the considered volcanoes remain limited to a few  
585 select sites and scenarios. These models are also typically restricted to one particular volcano and one specific mechanism of  
tsunami generation as a retrospectively investigation.

Consequently, predictive studies remain rare (Giachetti et al., 2012; Paris and Ulvrova, 2019) and are only possible because  
the specific local circumstances leading to the tsunami are very well understood, which is knowledge that is lacking for most  
coastal volcanoes. Here, we provide multiple predictive models for the volcanoes we classified as posing a high  
590 tsunamigenic hazard. As volcanogenic tsunamis are caused by a large variety of mechanisms (Fig. 6) we contribute to this  
aspect by providing a simplified and broader view at the travel times of potential future tsunamis that are unspecific to the  
mechanism of tsunami generation and their magnitude (with the possible exception of meteotsunamis as seen at Hunga  
Tonga Ha'apai in 2022, which appear to have different wave propagation properties). We mainly account for the potential  
spatial impact of volcanogenic tsunamis and To account for the potential spatial impact of volcanogenic tsunamis, we extend

595 our tsunami hazard evaluation by assessing the total length of a coastline affected within one and two hours of tsunami  
propagation for the volcanoes categorised as high hazard in our ranking (except Didicas). For that, we compute tsunami  
travel times (TTT) from point sources centred at each volcano position using the SRM30+ bathymetry (Becker et al., 2009)  
resampled to 1 arc minute resolution and the numerical algorithm as proposed by Marchuk (2008). This means that we can  
simulate the travel and arrival times of specific volcanoes independent of how the tsunami was generated (as long as it is a  
point source), but we also cannot consider specific wave heights or runup as these depend strongly on the specific source  
mechanism and magnitude of the event and require additional and much more specific modelling data for individual sites.  
600 Fig. 8 shows our calculated tsunami propagation fronts after 1 hour of wave propagation. Table 3, in turn, lists the total  
lengths of the coastlines affected after 1 and 2 hours of propagation, respectively. Detailed tsunami travel time plots for the  
individual volcanoes can be found in the supplementary figures 1, 2 and 3 material D. It is important to note that this type of  
simulation only includes the travel time of the tsunami, but not other important characteristics such as wave amplitude.  
605

<b>Volcano</b>	<b>Coastline 1h in km</b>	<b>Coastline 2h in km</b>
Anak Krakatau	1621	3969
Batu Tara	7322	25511
Kadovar	3078	10195
Ritter Island	3933	13370
Gamalama	6069	27900
Rabaul	5007	16380
Manam	2720	9061
Iliwerung	5974	21244
Sangeang Api	5451	18112
Karangetang	5819	24147
Langila	3955	13174
Sirung	5866	22289
Ulawun	2521	11527
Wetar	8512	27519
Nila	6117	23948

Ruang	6019	24998
Bam	2841	9012
Serua	6498	23998

**Table 3:** The affected coastline of a tsunami originating from high hazard volcanoes extracted using the tsunami travel time modelling. We note that Anak Krakatau, where the most prominent event in recent years occurred in 2018, affects the lowest length of coastline. This highlights that similar events at other locations can have much more widespread impacts.

610

The results show the potentially wide reach of volcanogenic tsunamis in both Indonesia and Papua New Guinea (Fig. 8), but it also reveals surprising differences. Anak Krakatau is only projected to affect ~1600 km of coastline within one hour (Table 1) and the area is restricted mostly within the Sunda Strait coast and parts of southern Java and Sumatra. This is nearly consistent with the 2018 tsunami, where the impacts were only within the Sunda Strait (e.g., Paris et al., 2020). All other volcanoes may produce tsunamis affecting a lot more coasts (Table 3). Batu Tara, which is now the highest scoring volcano in our catalogue, is projected to affect a coastline of ~7300 km, almost 5 times as much as Anak Krakatau within the same time. The longest total affected coastline of ~8500 km (after 1 hour of propagation) is calculated for Wetar, covering almost the entire Banda Sea. Tsunamis from the northern volcanoes in Indonesia are shown to potentially reach up to the Philippines as well as cover large parts of northern Sulawesi and Halmahera and almost all volcanoes in Papua New Guinea (with the exception of Rabaul) reach through most of the Bismarck Sea.

615

620

While our models are limited to the travel time, they can be used to estimate the warning time for shores in case a tsunami occurs at one of the considered volcanoes. Despite demonstrating the potential to affect large coastal areas, our TTT modelling does not inform how severe a tsunami from a volcanic source could actually turn out. In fact, tsunami travel time modelling neither accounts for source magnitude, nor for energy transfer. Full source process simulation coupled to modelling of the full wave propagation is needed to assess the magnitude of the coastal impact. In particular, the models are expected to strongly depend on the mechanism triggering the tsunami, i.e., a landslide, PDC or explosion, as well as the magnitude of the event. For example, in landslide or sector collapse events, the largest runups and the most severe impacts are expected to be largest in the near-field of the volcano, but may still be significant in the far-field (Harris et al., 2012; Grilli et al., 2021). Furthermore, the recent Hunga Tonga-Hunga Ha'apai eruption has shown that large eruptions are capable of generating meteotsunamis travelling long distances without significant loss of amplitude. This particular tsunami also travelled faster than initially expected (Somerville et al., 2022). To compensate for these knowledge gaps, full physics-based source and propagation modelling, especially if they are coupled to the population density information along the coasts, may become the most useful tool to improve our understanding of the risk posed by volcanogenic tsunamis. So far, published models almost exclusively consider Anak Krakatau (e.g., Grilli et al., 2019; Heidarzadeh et al., 2020; Mulia et al., 2020;

625

630

635 Omira and Ramalho, 2020; Paris et al., 2020), whereas the risk posed by the other high-hazard volcanoes in our catalogue is largely unknown, emphasising the need for future investigations and modelling efforts. For future hazard and risk assessments, we thus recommend supplementing the knowledge from our TTT-models with specific detailed scenario calculations using established numerical modelling approaches, particularly for those high-hazard volcanoes where no such models exist (e.g. Batu Tara, Iliwerung, Nila). This coupled with the prioritisation of specific volcanoes provided by our ranking can provide a well-founded basis for future disaster mitigation strategies. While the detection of volcanic processes triggering a tsunami will remain challenging to detect due to the multiple possible generation mechanisms, other steps can be done to improve warning times. These include the addition of strategically placed tide gauges as suggested by Annunziato et al. (2019), or improved real-time volcano monitoring through seismometers, radar, cameras or infrasound sensors. Regular use of satellite data (e.g. InSAR) can also help to preemptively identify volcanic unrest or destabilising flanks.


645

Based on our MCDM analysis considering 131 volcanoes in SE-Asia we identify 19 that pose a high tsunami hazard and another 48 with moderate tsunami hazard. We find our ranking system to be robust for the higher scoring volcanoes, meaning that we can reliably identify the most likely volcanoes to produce a tsunami in the future. For volcanoes with moderate to low scores the ranking is less robust and more susceptible to subjective judgement. The main limitations remaining are (1) a lack of knowledge how much individual factors contribute to the tsunami hazard of a volcano, instead requiring subjective assumptions, (2) erroneous, incomplete or insufficient data availability for many volcanoes (e.g. bathymetry or historical data), and (3) the multitude of different mechanisms which may cause a volcanic tsunami (i.e. PDCs, landslides, explosions) making a clear scenario assessment challenging.

650

655 Our results show that the Indonesian Lesser Sunda Islands and northern Molucca Sea as well as the southern Bismarck Sea in Papua New Guinea are areas with a high number of hazardous volcanoes and may thus be particularly prone to tsunami sources sourced by volcanoes. Many of these volcanoes such as Batu Tara, Indonesia, are not commonly considered for this type of hazard. We therefore emphasise the need to reconsider the current state of monitoring and risk assessment in these areas. Since tsunami warning systems are largely unable to detect volcanogenic tsunamis, our results highlight the

660 importance of a reassessment of the current network and additional suitable equipment on the ground and through earth observation satellites. Due to the inherently short warning times of these events, we also recommended increased preemptive measures on a local level, such as increased public education programs for coastal communities and the marking evacuation routes along populated coasts.



## 6 Acknowledgements

665 The authors acknowledge the financial support by the Federal Ministry of Education and Research of Germany in the framework of the TSUNAMI\_RISK project (project numbers 03G0906A and 03G0906B), which is a part of the funding initiative CLIENT-II. We further acknowledge the past contributions by GITEWS, on which this projects builds upon. [We also thank Raphaël Paris for friendly and helpful feedback on the manuscript.](#)

## 7 Author contributions

670 EZ conceptualised and wrote the manuscript. Figures were prepared by EZ and AO. EZ also performed the analyses and evaluations of the individual volcanoes. EZ, AO, SP, TW and IR jointly developed the point score system. Tsunami-travel-time modelling was performed by AB and photogrammetric data for Anak Krakatau was provided by HD. All authors contributed to the writing and editing of the manuscript.

## 8 Competing interests

675 The authors declare no competing interests.

## 9 Code/Data availability

[The data used to create the MCDA ranking and more detailed TTT model plots can be freely accessed via the supplementary files. NETVOLC and MORVOLC data for individual volcanoes are available from the corresponding author upon reasonable request.](#)

680

## 109 References

- Airbus, 2020. Copernicus DEM Validation Report (v2.1), [https://spacedata.copernicus.eu/documents/20126/0/GEO1988-CopernicusDEM-SPE-002\\_ProductHandbook\\_I1.00.pdf](https://spacedata.copernicus.eu/documents/20126/0/GEO1988-CopernicusDEM-SPE-002_ProductHandbook_I1.00.pdf).
- 685 Annunziato, A., Prasetya, G. and Husrin, S., 2019. Anak Krakatau volcano emergency tsunami early warning system. *Science of Tsunami Hazards*, 38(2).
- Badriana, M.R., Bachtiar, H., Adytia, D., Sembiring, L., Andonowati and Groesen, E.v., 2017. Wave run-up of a possible Anak-Krakatau tsunami on planned and optimized Jakarta Sea Dike. *AIP Conference Proceedings*, 1857(1): 090004. DOI: 10.1063/1.4987103
- 690 Becker, J.J., Sandwell, D.T., Smith, W.H.F., Braud, J., Binder, B., Depner, J., Fabre, D., Factor, J., Ingalls, S., Kim, S.H., Ladner, R., Marks, K., Nelson, S., Pharaoh, A., Trimmer, R., Von Rosenberg, J., Wallace, G. and Weatherall, P., 2009. Global Bathymetry and Elevation Data at 30 Arc Seconds Resolution: SRTM30\_PLUS. *Marine Geodesy*, 32(4): 355-371. DOI: 10.1080/01490410903297766

Feldfunktion geändert

- 695 Belousov, A., Belousova, M. and Voight, B., 1999. Multiple edifice failures, debris avalanches and associated eruptions in the Holocene history of Shiveluch volcano, Kamchatka, Russia. *Bulletin of Volcanology*, 61(5): 324-342. DOI: 10.1007/s004450050300
- Belousov, A., Voight, B., Belousova, M. and Muravyev, Y., 2000. Tsunamis Generated by Subaquatic Volcanic Explosions: Unique Data from 1996 Eruption in Karymskoye Lake, Kamchatka, Russia. *pure and applied geophysics*, 157(6): 1135-1143. DOI: 10.1007/s000240050021
- 700 Bonaccorso, A., Calvari, S., Garfi, G., Lodato, L. and Patanè, D., 2003. Dynamics of the December 2002 flank failure and tsunami at Stromboli volcano inferred by volcanological and geophysical observations. *Geophysical Research Letters*, 30(18). DOI: 10.1029/2003GL017702
- Borrero, J.C., Solihuddin, T., Fritz, H.M., Lynett, P.J., Prasetya, G.S., Skanavis, V., Husrin, S., Kushendratno, Kongko, W., Istiyanto, D.C., Daulat, A., Purbani, D., Salim, H.L., Hidayat, R., Asvaliantina, V., Usman, M., Kodijat, A., Son, S. and Synolakis, C.E., 2020. Field Survey and Numerical Modelling of the December 22, 2018 Anak Krakatau Tsunami. *Pure and Applied Geophysics*, 177(6): 2457-2475. DOI: 10.1007/s00024-020-02515-y
- 705 Brown, S.K., Jenkins, S.F., Sparks, R.S.J., Odbert, H. and Auker, M.R., 2017. Volcanic fatalities database: analysis of volcanic threat with distance and victim classification. *Journal of Applied Volcanology*, 6(1): 15. DOI: 10.1186/s13617-017-0067-4
- Carey, S., Sigurdsson, H., Mandeville, C. and Bronto, S., 2000. Volcanic hazards from pyroclastic flow discharge into the sea: Examples from the 1883 eruption of Krakatau, Indonesia. In: F.W. McCoy and G. Heiken (Editors), *Volcanic Hazards and Disasters in Human Antiquity*. Geological Society of America, pp. 0. DOI: 10.1130/0-8137-2345-0.1
- 710 Choi, B.H., Pelinovsky, E., Kim, K.O. and Lee, J.S., 2003. Simulation of the trans-oceanic tsunami propagation due to the 1883 Krakatau volcanic eruption. *Nat. Hazards Earth Syst. Sci.*, 3(5): 321-332. DOI: 10.5194/nhess-3-321-2003
- 715 Crisci, G.M., De Rosa, R., Esperança, S., Mazzuoli, R. and Sonnino, M., 1991. Temporal evolution of a three component system: the island of Lipari (Aeolian Arc, southern Italy). *Bulletin of Volcanology*, 53(3): 207-221. DOI: 10.1007/BF00301231
- Cronin, S.J., Lube, G., Dayudi, D.S., Sumarti, S., Subrandiyo, S. and Surono, 2013. Insights into the October–November 2010 Gunung Merapi eruption (Central Java, Indonesia) from the stratigraphy, volume and characteristics of its pyroclastic deposits. *Journal of Volcanology and Geothermal Research*, 261: 244-259. DOI: 10.1016/j.jvolgeores.2013.01.005
- 720 Darmawan, H., Mutaqin, B.W., Harijoko, A., Wibowo, H.E., Haerani, N., Surmayadi, M., Jati, R. and Asriningrum, W., 2020. Topography and structural changes of Anak Krakatau due to the December 2018 catastrophic events. *The Indonesian Journal of Geography*, 52(3): 402-410. DOI: 10.22146/ijg.53740
- 725 Darmawan, H., Troll, V.R., Walter, T.R., Deegan, F.M., Geiger, H., Heap, M.J., Seraphine, N., Harris, C., Humaida, H. and Müller, D., 2022. Hidden mechanical weaknesses within lava domes provided by buried high-porosity hydrothermal alteration zones. *Scientific Reports*, 12(1): 3202. DOI: 10.1038/s41598-022-06765-9
- Di Traglia, F., Nolesini, T., Solari, L., Ciampalini, A., Frodella, W., Steri, D., Allotta, B., Rindi, A., Marini, L., Monni, N., Galardi, E. and Casagli, N., 2018. Lava delta deformation as a proxy for submarine slope instability. *Earth and Planetary Science Letters*, 488: 46-58. DOI: 10.1016/j.epsl.2018.01.038
- 730 Dogan, G.G., Annunziato, A., Hidayat, R., Husrin, S., Prasetya, G., Kongko, W., Zaytsev, A., Pelinovsky, E., Imamura, F. and Yalciner, A.C., 2021. Numerical Simulations of December 22, 2018 Anak Krakatau Tsunami and Examination of Possible Submarine Landslide Scenarios. *Pure and Applied Geophysics*, 178(1): 1-20. DOI: 10.1007/s00024-020-02641-7
- 735 ESA, 2022. Copernicus Sentinel-2 data. Retrieved from Sentinel Playground accessed on 21st September 2021, <https://apps.sentinel-hub.com/sentinel-playground>.
- Euillades, L.D., Grosse, P. and Euillades, P.A., 2013. NETVOLC: An algorithm for automatic delimitation of volcano edifice boundaries using DEMs. *Computers & Geosciences*, 56: 151-160. DOI: 10.1016/j.cageo.2013.03.011
- Ewert, J.W., 2007. System for Ranking Relative Threats of U.S. Volcanoes. *Natural Hazards Review*, 8(4): 112-124. DOI: doi:10.1061/(ASCE)1527-6988(2007)8:4(112)
- 740 Ewert, J.W., Diefenbach, A.K. and Ramsey, D.W., 2018. 2018 update to the U.S. Geological Survey national volcanic threat assessment. 2018-5140, Reston, VA. DOI: 10.3133/sir20185140

Feldfunktion geändert

- Fernández, D.S. and Lutz, M.A., 2010. Urban flood hazard zoning in Tucumán Province, Argentina, using GIS and multicriteria decision analysis. *Engineering Geology*, 111(1): 90-98. DOI: 10.1016/j.enggeo.2009.12.006
- 745 Fisher, R.V., 1990. Transport and deposition of a pyroclastic surge across an area of high relief: The 18 May 1980 eruption of Mount St. Helens, Washington. *GSA Bulletin*, 102(8): 1038-1054. DOI: 10.1130/0016-7606(1990)102<1038:TADOAP>2.3.CO;2
- Fornaciai, A., Favalli, M. and Nannipieri, L., 2019. Numerical simulation of the tsunamis generated by the Sciara del Fuoco landslides (Stromboli Island, Italy). *Scientific Reports*, 9(1): 18542. DOI: 10.1038/s41598-019-54949-7
- 750 Francis, P.W., 1985. The origin of the 1883 Krakatau tsunamis. *Journal of Volcanology and Geothermal Research*, 25(3): 349-363. DOI: 10.1016/0377-0273(85)90021-6
- Gertisser, R. and Keller, J., 2003. Temporal variations in magma composition at Merapi Volcano (Central Java, Indonesia): magmatic cycles during the past 2000 years of explosive activity. *Journal of Volcanology and Geothermal Research*, 123(1): 1-23. DOI: 10.1016/S0377-0273(03)00025-8
- 755 Giachetti, T., Paris, R., Kelfoun, K. and Ontowirjo, B., 2012. Tsunami hazard related to a flank collapse of Anak Krakatau Volcano, Sunda Strait, Indonesia. *Geological Society, London, Special Publications*, 361(1): 79. DOI: 10.1144/SP361.7
- Global Volcanism Program, 2013. *Volcanoes of the World*, v. 4.10.6 (24 Mar 2020). Venzke, E. Smithsonian Institution. DOI: 10.5479/si.GVP.VOTW4-2013
- 760 Gonzalez-Ollauri, A. and Mickovski, S.B., 2017. Hydrological effect of vegetation against rainfall-induced landslides. *Journal of Hydrology*, 549: 374-387. DOI: 10.1016/j.jhydrol.2017.04.014
- Grilli, S.T., Tappin, D.R., Carey, S., Watt, S.F.L., Ward, S.N., Grilli, A.R., Engwell, S.L., Zhang, C., Kirby, J.T., Schambach, L. and Muin, M., 2019. Modelling of the tsunamis from the December 22, 2018 lateral collapse of Anak Krakatau volcano in the Sunda Straits, Indonesia. *Scientific Reports*, 9(1): 11946. DOI: 10.1038/s41598-019-48327-6
- 765 Grilli, S.T., Zhang, C., Kirby, J.T., Grilli, A.R., Tappin, D.R., Watt, S.F.L., Hunt, J.E., Novellino, A., Engwell, S., Nurshal, M.E.M., Abdurrachman, M., Cassidy, M., Madden-Nadeau, A.L. and Day, S., 2021. Modeling of the Dec. 22nd 2018 Anak Krakatau volcano lateral collapse and tsunami based on recent field surveys: Comparison with observed tsunami impact. *Marine Geology*, 440: 106566. DOI: 10.1016/j.margeo.2021.106566
- 770 Grosse, P., van Wyk de Vries, B., Petrinovic, I.A., Euillades, P.A. and Alvarado, G.E., 2009. Morphometry and evolution of arc volcanoes. *Geology*, 37(7): 651-654. DOI: 10.1130/G25734A.1
- Grosse, P., van Wyk de Vries, B., Euillades, P.A., Kervyn, M. and Petrinovic, I.A., 2012. Systematic morphometric characterization of volcanic edifices using digital elevation models. *Geomorphology*, 136(1): 114-131. DOI: 10.1016/j.geomorph.2011.06.001
- 775 Guth, P.L. and Geoffroy, T.M., 2021. LiDAR point cloud and ICESat-2 evaluation of 1 second global digital elevation models: Copernicus wins. *Transactions in GIS*, 25(5): 2245-2261. DOI: 10.1111/tgis.12825
- Hamzah, L., Puspito, N.T. and Imamura, F., 2000. Tsunami Catalog and Zones in Indonesia. *Journal of Natural Disaster Science*, 22(1): 25-43. DOI: 10.2328/jnds.22.25
- Hanka, W., Saul, J., Weber, B., Becker, J., Harjadi, P., Fauzi and Group, G.S., 2010. Real-time earthquake monitoring for tsunami warning in the Indian Ocean and beyond. *Nat. Hazards Earth Syst. Sci.*, 10(12): 2611-2622. DOI: 10.5194/nhess-10-2611-2010
- 780 Harig, S., Immerz, A., Weniza, Griffin, J., Weber, B., Babeyko, A., Rakowsky, N., Hartanto, D., Nurokhim, A., Handayani, T. and Weber, R., 2020. The Tsunami Scenario Database of the Indonesia Tsunami Early Warning System (InaTEWS): Evolution of the Coverage and the Involved Modeling Approaches. *Pure and Applied Geophysics*, 177(3): 1379-1401. DOI: 10.1007/s00024-019-02305-1
- 785 Harris, J.C., Grilli, S.T., Abadie, S. and Bakhsh, T.T., 2012. Near- And Far-field Tsunami Hazard From the Potential Flank Collapse of the Cumbre Vieja Volcano, The Twenty-second International Offshore and Polar Engineering Conference, pp. ISOPE-I-12-515.
- 790 Heap, M.J., Mollo, S., Vinciguerra, S., Lavallée, Y., Hess, K.U., Dingwell, D.B., Baud, P. and Iezzi, G., 2013. Thermal weakening of the carbonate basement under Mt. Etna volcano (Italy): Implications for volcano instability. *Journal of Volcanology and Geothermal Research*, 250: 42-60. DOI: 10.1016/j.jvolgeores.2012.10.004



- Heap, M.J., Baumann, T.S., Rosas-Carbajal, M., Komorowski, J.C., Gilg, H.A., Villeneuve, M., Moretti, R., Baud, P., Carbillet, L., Harnett, C. and Reuschlé, T., 2021. Alteration-Induced Volcano Instability at La Soufrière de Guadeloupe (Eastern Caribbean). *Journal of Geophysical Research: Solid Earth*, 126(8): e2021JB022514. DOI: 10.1029/2021JB022514
- 795 Heap, M.J. and Violay, M.E.S., 2021. The mechanical behaviour and failure modes of volcanic rocks: a review. *Bulletin of Volcanology*, 83(5): 33. DOI: 10.1007/s00445-021-01447-2
- Heidarzadeh, M., Ishibe, T., Sandanbata, O., Muhari, A. and Wijanarto, A.B., 2020. Numerical modeling of the subaerial landslide source of the 22 December 2018 Anak Krakatoa volcanic tsunami, Indonesia. *Ocean Engineering*, 195: 106733. DOI: 10.1016/j.oceaneng.2019.106733
- 800 Hidayat, A., Marfai, M.A. and Hadmoko, D.S., 2020. Eruption on Indonesia's volcanic islands: a review of potential hazards, fatalities, and management. *IOP Conference Series: Earth and Environmental Science*, 485(1): 012061. DOI: 10.1088/1755-1315/485/1/012061
- Karstens, J., Kelfoun, K., Watt, S.F.L. and Berndt, C., 2020. Combining 3D seismics, eyewitness accounts and numerical simulations to reconstruct the 1888 Ritter Island sector collapse and tsunamis. *International Journal of Earth Sciences*, 109(8): 2659-2677. DOI: 10.1007/s00531-020-01854-4
- 805 Kieffer, S.W., 1981. Blast dynamics at Mount St Helens on 18 May 1980. *Nature*, 291(5816): 568-570. DOI: 10.1038/291568a0
- Korup, O., Seidemann, J. and Mohr, C.H., 2019. Increased landslide activity on forested hillslopes following two recent volcanic eruptions in Chile. *Nature Geoscience*, 12(4): 284-289. DOI: 10.1038/s41561-019-0315-9
- 810 Laiolo, M., Massimetti, F., Cigolini, C., Ripepe, M. and Coppola, D., 2018. Long-term eruptive trends from space-based thermal and SO<sub>2</sub> emissions: a comparative analysis of Stromboli, Batu Tara and Tinakula volcanoes. *Bulletin of Volcanology*, 80(9): 68. DOI: 10.1007/s00445-018-1242-0
- Lauterjung, J., Münch, U. and Rudloff, A., 2010. The challenge of installing a tsunami early warning system in the vicinity of the Sunda Arc, Indonesia. *Nat. Hazards Earth Syst. Sci.*, 10(4): 641-646. DOI: 10.5194/nhess-10-641-2010
- 815 Lipman, P.W. and Mullineaux, D.R., 1981. The 1980 eruptions of Mount St. Helens, Washington. 1250. DOI: 10.3133/pp1250
- Llanes, P., Silver, E., Day, S. and Hoffman, G., 2009. Interactions between a transform fault and arc volcanism in the Bismarck Sea, Papua New Guinea. *Geochemistry, Geophysics, Geosystems*, 10(6). DOI: 10.1029/2009GC002430
- Maeno, F., Imamura, F. and Taniguchi, H., 2006. Numerical simulation of tsunamis generated by caldera collapse during the 7.3 ka Kikai eruption, Kyushu, Japan. *Earth, Planets and Space*, 58(8): 1013-1024. DOI: 10.1186/BF03352606
- 820 Maeno, F. and Imamura, F., 2011. Tsunami generation by a rapid entrance of pyroclastic flow into the sea during the 1883 Krakatau eruption, Indonesia. *Journal of Geophysical Research: Solid Earth*, 116(B9). DOI: 10.1029/2011JB008253
- Marchuk, A.G., 2008. Minimizing computational errors of tsunami wave-ray and travel time. *Science of Tsunami Hazards*, 27(4): 12-24.
- 825 McGuire, W.J., 2006. Lateral collapse and tsunamigenic potential of marine volcanoes. Geological Society, London, Special Publications, 269(1): 121. DOI: 10.1144/GSL.SP.2006.269.01.08
- Morton, A., Airoldi, M. and Phillips, L.D., 2009. Nuclear Risk Management on Stage: A Decision Analysis Perspective on the UK's Committee on Radioactive Waste Management. *Risk Analysis*, 29(5): 764-779. DOI: 10.1111/j.1539-6924.2008.01192.x
- 830 Mulia, I.E., Watada, S., Ho, T.C., Satake, K., Wang, Y. and Aditiya, A., 2020. Simulation of the 2018 Tsunami Due to the Flank Failure of Anak Krakatau Volcano and Implication for Future Observing Systems. *Geophysical Research Letters*, 47(14): e2020GL087334. DOI: 10.1029/2020GL087334
- Murray, J.B. and Voight, B., 1996. Slope stability and eruption prediction on the eastern flank of Mount Etna. Geological Society, London, Special Publications, 110(1): 111. DOI: 10.1144/GSL.SP.1996.110.01.08
- 835 Mutaqin, B.W., Lavigne, F., Hadmoko, D.S. and Ngalawani, M.N., 2019. Volcanic Eruption-Induced Tsunami in Indonesia: A Review. *IOP Conference Series: Earth and Environmental Science*, 256: 012023. DOI: 10.1088/1755-1315/256/1/012023
- NGDC, 2021. National Geophysical Data Center / World Data Service: NCEI/WDS Global Historical Tsunami Database, NOAA National Centers for Environmental Information. DOI: 10.7289/V5PN93H7
- 840

- Nomanbhoj, N. and Satake, K., 1995. Generation mechanism of tsunamis from the 1883 Krakatau Eruption. *Geophysical Research Letters*, 22(4): 509-512. DOI: 10.1029/94GL03219
- Nutt, D.J., King, L.A. and Phillips, L.D., 2010. Drug harms in the UK: a multicriteria decision analysis. *The Lancet*, 376(9752): 1558-1565. DOI: 10.1016/S0140-6736(10)61462-6
- 845 Omira, R. and Ramalho, I., 2020. Evidence-Calibrated Numerical Model of December 22, 2018, Anak Krakatau Flank Collapse and Tsunami. *Pure and Applied Geophysics*, 177(7): 3059-3071. DOI: 10.1007/s00024-020-02532-x
- OpenStreetMap, 2022. OSM Land polygons, <https://www.openstreetmap.org/>.
- Pakoksung, K., Suppasri, A. and Imamura, F., 2021. Probabilistic Tsunami Hazard Analysis of Inundated Buildings Following a Subaqueous Volcanic Explosion Based on the 1716 Tsunami Scenario in Taal Lake, Philippines. *Geosciences*, 11(2): 92. DOI: 10.3390/geosciences11020092
- 850 Paris, A., Heinrich, P., Paris, R. and Abadie, S., 2020. The December 22, 2018 Anak Krakatau, Indonesia, Landslide and Tsunami: Preliminary Modeling Results. *Pure and Applied Geophysics*, 177(2): 571-590. DOI: 10.1007/s00024-019-02394-y
- Paris, R., Switzer, A.D., Belousova, M., Belousov, A., Ontowirjo, B., Whelley, P.L. and Ulvrova, M., 2014. Volcanic tsunamis: a review of source mechanisms, past events and hazards in Southeast Asia (Indonesia, Philippines, Papua New Guinea). *Natural Hazards*, 70(1): 447-470. DOI: 10.1007/s11069-013-0822-8
- 855 Paris, R., 2015. Source mechanisms of volcanic tsunamis. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 373(2053): 20140380. DOI: 10.1098/rsta.2014.0380
- Paris, R. and Ulvrova, M., 2019. Tsunamis generated by subaqueous volcanic explosions in Taal Caldera Lake, Philippines. *Bulletin of Volcanology*, 81(3): 14. DOI: 10.1007/s00445-019-1272-2
- 860 Plank, S., Walter, T.R., Martinis, S. and Cesca, S., 2019. Growth and collapse of a littoral lava dome during the 2018/19 eruption of Kadovar Volcano, Papua New Guinea, analyzed by multi-sensor satellite imagery. *Journal of Volcanology and Geothermal Research*, 388: 106704. DOI: 10.1016/j.jvolgeores.2019.106704
- Poland, M.P. and Orr, T.R., 2014. Identifying hazards associated with lava deltas. *Bulletin of Volcanology*, 76(12): 880. DOI: 10.1007/s00445-014-0880-0
- 865 Poland, M.P., Peltier, A., Bonforte, A. and Puglisi, G., 2017. The spectrum of persistent volcanic flank instability: A review and proposed framework based on Kīlauea, Piton de la Fournaise, and Etna. *Journal of Volcanology and Geothermal Research*, 339: 63-80. DOI: 10.1016/j.jvolgeores.2017.05.004
- Pranantyo, I.R., Heidarzadeh, M. and Cummins, P.R., 2021. Complex tsunami hazards in eastern Indonesia from seismic and non-seismic sources: Deterministic modelling based on historical and modern data. *Geoscience Letters*, 8(1): 20. DOI: 10.1186/s40562-021-00190-y
- 870 Rahmati, O., Zeinivand, H. and Besharat, M., 2016. Flood hazard zoning in Yasooj region, Iran, using GIS and multi-criteria decision analysis. *Geomatics, Natural Hazards and Risk*, 7(3): 1000-1017. DOI: 10.1080/19475705.2015.1045043
- Ren, Z., Wang, Y., Wang, P., Hou, J., Gao, Y. and Zhao, L., 2020. Numerical study of the triggering mechanism of the 2018 Anak Krakatau tsunami: eruption or collapsed landslide? *Natural Hazards*, 102(1): 1-13. DOI: 10.1007/s11069-020-03907-y
- 875 Romero, J.E., Polacci, M., Watt, S., Kitamura, S., Tormey, D., Sielfeld, G., Arzilli, F., La Spina, G., Franco, L., Burton, M. and Polanco, E., 2021. Volcanic Lateral Collapse Processes in Mafic Arc Edifices: A Review of Their Driving Processes, Types and Consequences. *Frontiers in Earth Science*, 9. DOI: 10.3389/feart.2021.639825
- 880 Scandone, R., Bartolini, S. and Martí, J., 2015. A scale for ranking volcanoes by risk. *Bulletin of Volcanology*, 78(1): 2. DOI: 10.1007/s00445-015-0995-y
- Siebert, L., 1984. Large volcanic debris avalanches: Characteristics of source areas, deposits, and associated eruptions. *Journal of Volcanology and Geothermal Research*, 22(3): 163-197. DOI: 10.1016/0377-0273(84)90002-7
- 885 Silver, E., Day, S., Ward, S., Hoffmann, G., Llanes, P., Driscoll, N., Appelgate, B. and Saunders, S., 2009. Volcano collapse and tsunami generation in the Bismarck Volcanic Arc, Papua New Guinea. *Journal of Volcanology and Geothermal Research*, 186(3): 210-222. DOI: 10.1016/j.jvolgeores.2009.06.013
- Smith, M.S. and Shepherd, J.B., 1993. Preliminary investigations of the tsunami hazard of Kick'em Jenny submarine volcano. *Natural Hazards*, 7(3): 257-277. DOI: 10.1007/BF00662650
- 890 Somerville, P., Blong, R. and Gissing, A., 2022. Hunga Tonga-Hunga Ha'apai Eruption 15th of January 2022, Risk Frontiers Briefing Note 460.

Feldfunktion geändert

- Spina, L., Del Bello, E., Ricci, T., Taddeucci, J. and Scarlato, P., 2021. Multi-parametric characterization of explosive activity at Batu Tara Volcano (Flores Sea, Indonesia). *Journal of Volcanology and Geothermal Research*, 413: 107199. DOI: 10.1016/j.jvolgeores.2021.107199
- 895 Tibaldi, A., 2001. Multiple sector collapses at Stromboli volcano, Italy: how they work. *Bulletin of Volcanology*, 63(2): 112-125. DOI: 10.1007/s004450100129
- Tinti, S., Pagnoni, G., Zaniboni, F. and Bortolucci, E., 2003. Tsunami generation in Stromboli island and impact on the south-east Tyrrhenian coasts. *Nat. Hazards Earth Syst. Sci.*, 3(5): 299-309. DOI: 10.5194/nhess-3-299-2003
- Toosi, A.S., Calbimonte, G.H., Nouri, H. and Alaghmand, S., 2019. River basin-scale flood hazard assessment using a modified multi-criteria decision analysis approach: A case study. *Journal of Hydrology*, 574: 660-671. DOI: 900 10.1016/j.jhydrol.2019.04.072
- Turner, M.B., 2008. Eruption cycles and magmatic processes at a reawakening volcano, Mt. Taranaki, New Zealand : a thesis presented in partial fulfilment of the requirements for the degree of Doctor of Philosophy in Earth Science at Massey University, Palmerston North, New Zealand. Doctoral Thesis, Massey University.
- van Wyk De Vries, B. and Borgia, A., 1996. The role of basement in volcano deformation. *Geological Society, London, Special Publications*, 110(1): 95. DOI: 10.1144/GSL.SP.1996.110.01.07
- 905 Walter, T.R., Haghshenas Haghghi, M., Schneider, F.M., Coppola, D., Motagh, M., Saul, J., Babeyko, A., Dahm, T., Troll, V.R., Tilmann, F., Heimann, S., Valade, S., Triyono, R., Khomarudin, R., Kartadinata, N., Laiolo, M., Massimetti, F. and Gaebler, P., 2019. Complex hazard cascade culminating in the Anak Krakatau sector collapse. *Nature Communications*, 10(1): 4339. DOI: 10.1038/s41467-019-12284-5
- 910 Ward, S.N. and Day, S., 2003. Ritter Island Volcano—lateral collapse and the tsunami of 1888. *Geophysical Journal International*, 154(3): 891-902. DOI: 10.1046/j.1365-246X.2003.02016.x
- Watters, R.J., Zimbelman, D.R., Bowman, S.D. and Crowley, J.K., 2000. Rock Mass Strength Assessment and Significance to Edifice Stability, Mount Rainier and Mount Hood, Cascade Range Volcanoes. *pure and applied geophysics*, 157(6): 957-976. DOI: 10.1007/s000240050012
- 915 Watts, P. and Waythomas, C.F., 2003. Theoretical analysis of tsunami generation by pyroclastic flows. *Journal of Geophysical Research: Solid Earth*, 108(B12). DOI: 10.1029/2002JB002265
- Yokoyama, I., 1981. A geophysical interpretation of the 1883 Krakatau eruption. *Journal of Volcanology and Geothermal Research*, 9(4): 359-378. DOI: 10.1016/0377-0273(81)90044-5
- Yokoyama, I., Tilling, R.I. and Scarpa, R., 1984. International Mobile Early-Warning System (s) for Volcanic Eruptions and 920 Related Seismic Activities: Report of an Unesco/UNEP Sponsored Preparatory Study in 1982-84. Unesco.
- Yoshida, H., Sugai, T. and Ohmori, H., 2012. Size–distance relationships for hummocks on volcanic rockslide-debris avalanche deposits in Japan. *Geomorphology*, 136(1): 76-87. DOI: 10.1016/j.geomorph.2011.04.044
- Yudhicara, Y., Bani, P. and Darmawan, A., 2015. Geothermal System as the Cause of the 1979 Landslide Tsunami in Lembata Island, Indonesia. *Indonesian Journal on Geoscience*, 2(2). DOI: 10.17014/ijog.2.2.91-99
- 925 Zengaffinen, T., Løvholt, F., Pedersen, G.K. and Muhari, A., 2020. Modelling 2018 Anak Krakatoa Flank Collapse and Tsunami: Effect of Landslide Failure Mechanism and Dynamics on Tsunami Generation. *Pure and Applied Geophysics*, 177(6): 2493-2516. DOI: 10.1007/s00024-020-02489-x