

Improving numerical snowpack simulations by assimilating land surface temperature By Esteban Alonso-González, Simon Gascoin, Sara Arioli, Ghislain Picard

This manuscript describes a study about the point-scale assimilation of land surface temperature data in an energy balance snowpack model. This is a synthetic experiment, given that there are not currently many data sets for observed land surface temperature data. The authors explain that future satellite missions should provide more data in the coming years and that it is important to be prepared to use that data. Consequently, their study aims at assessing the potential of LST to improve the representation of SWE in snowpack models. They have designed different scenarios for their synthetic data, in order to simulate the presence of clouds and also to simulate different revisiting time for the satellites taking the measures. They also compare two methods for data assimilation: the particle filter and a particle batch smoother.

In my opinion the manuscript is interesting, well-written and relevant for the readers of The Cryosphere. I have only a few minor comments that I would like the authors to address.

Specific minor comments:

- Line 80: I am curious about OSSE, as I had never heard or read this specific appellation before. Is it a just a fancy name to refer to any form of synthetic experiment, or is it a specific *type* of synthetic experiment, guided by a series of principles/rules? Later in this paragraph you just refer to « synthetic experiments ». Are they interchangeable terms? Maybe uniformize?
- Table 1: There are multiple polices in this table. Please uniformize.
- Lines 113-114: There are certain methodological choices for the construction of the synthetic data sets for which I would have liked to have more detail. First, I understand that you selected the specific value for the multiplicative perturbation of precip after Beck et al. (2019), but I still would have liked to have more detail about that, and why you did not test more than one value for this multiplicative factor.
- Similarly, on page 5, I would have liked to have more detail about:
 - Line 125-129: The trial and error process that led to the selection of mu and sigma. Just a little bit more detail. Also, why 300 particles?
 - Line 139: what are the « new perturbation parameters »? Do you mean « new particles »? Or is it a more sophisticated way of preventing the collapsing of the filter?
- On page 6, the authors switch back and forth between the present tense and the past. I think it would be better to use the present all the time.
- Figure 2: If I understand correctly, in your synthetic experiment, the model can only underestimate SWE (the open loop is consistently inferior to the « synthetic truth »). Is this on purpose? If so, why? I have seem many cases for which the model overestimate SWE, for instance because it overestimated the density on some occasions. I think it would be important to explain why FSM2 can only underestimate SWE.
- Figure 3: it would be good to add a legend.
- Figure 5: Maybe it is obvious but I don't understand why the perturbation of precip on this graph is 2 while on page 4 it is 0.5. Is it because it is inverted (2 vs 1/2)? I guess it is something like that, but it is not completely clear to me the way it is written. It would be nice if it was the same number on the figure and in the text.
- Figure 6: the caption announces a « dashed line » but there is no dashed line.
- Page 8: Your entire experiment if at point scale. What would be the challenges of applying this at the basin scale? It would be interesting to add some discussion about that, given that basin

scale is very important for hydrology and hydrologists are also potential readers of the Cryosphere

- I have noted a few typos, misplaced comas, etc in the manuscript. I include my annotated version, in which they are all encircled.



Improving numerical snowpack simulations by assimilating land surface temperature.

Esteban Alonso-González¹, Simon Gascoin¹, Sara Arioli², Ghislain Picard²

¹Centre d'Etudes Spatiales de la Biosphère, Université de Toulouse, CNRS/CNES/IRD/INRA/UPS, Toulouse, France

5 ²Univ. Grenoble Alpes, CNRS, Institut des Géosciences de l'Environnement (IGE), Grenoble, France

Correspondence: Alonso-Gonzalez E.: esteban.alonso-gonzalez@univ-tlse3.fr

Abstract. The assimilation of data from Earth observation satellites into numerical models is considered as the path forward to estimate SWE distribution in mountain catchments. The land surface temperature (LST) can be observed from space, but its potential to improve SWE simulations remains underexplored. This is likely due to the insufficient temporal or spatial
10 resolution offered by the current thermal infrared (TIR) missions. However, three planned missions will provide global-scale TIR data at much higher spatio-temporal resolution in the coming years.

To investigate the value of TIR data to improve SWE estimation, we developed a **synthetic data assimilation experiment** at **five snow-dominated sites** covering a latitudinal gradient in the northern hemisphere. We generated synthetic true LST and SWE series by forcing an energy-balance snowpack model with the ERA5-Land reanalysis. We used this synthetic true LST
15 to recover the synthetic true SWE from a degraded version of ERA5-Land. **We defined different observation scenarios to emulate the revisiting times of Landsat 8 (16 days) and the Thermal infraRed Imaging Satellite for High-resolution Natural resource Assessment (TRISHNA) (3 days), while accounting for cloud cover.** We replicated the experiments 100 times at each experimental site to assess the robustness of the assimilation process. We performed the assimilation using two different approaches: a sequential scheme (particle filter) and a smoother (particle batch smoother).

20 The results show that LST data assimilation using the smoother reduced the normalized Root Mean Square Error (nRMSE) of the simulations from 57% (open loop) to 7% and 3% for 16 day revisit and 3 day revisit respectively, in the absence of clouds. We found similar but higher nRMSE values by removing observations due to cloud cover but with a substantial increase of the standard deviation of the nRMSE of the replicates, highlighting the importance of revisiting times in the stability of the assimilation output. **The smoother largely outperformed the particle filter** algorithm, **suggesting that the**
25 **capability of a smoother to propagate the information along the season is key** to exploit LST information for snow modeling. These results suggest that the LST data assimilation has an underappreciated potential to improve snowpack simulations and highlight the value of upcoming TIR missions to advance snow hydrology.



1 Introduction

The seasonal snowpack plays a key role in many ecological and hydrological processes worldwide (Barnett et al., 2005).

30 Due to its high albedo and insulating capabilities, the extensive snow-covered area of the northern hemisphere influences the Earth climate system (Henderson et al., 2018). In mountain regions, the seasonal snowpack is also an important source of runoff during the summer when the water demand peaks. Hence, an accurate knowledge of the snowpack conditions has an **important economic value** (Sturm et al., 2017). The snow cover is also a source of natural hazards such as floods caused by rain-on-snow events or snow avalanches, events that are expected to increase due to the impacts of climate warming
35 (Ballesteros-Cánovas et al., 2018; Musselman et al., 2018).

Despite its importance, monitoring the snowpack remains challenging for both scientists and water management agencies. The variable nature of the snowpack makes it difficult to deploy and maintain ground based snow monitoring networks (Kinar & Pomeroy, 2015). Therefore, snow hydrologists have developed methods to take advantage of satellite remote sensing since the beginning of the Landsat programme (Rango & Martinec, 1979). Yet, **the application of remote sensing in**
40 **snow hydrology remains hindered by the lack of direct observations of the snow water equivalent (SWE) in mountain regions** (Dozier et al., 2016).

Numerical models allow simulating many snowpack state variables, including the SWE. However, their accuracy is constrained by the **large uncertainty of the meteorological forcing** (Raleigh et al., 2015). Recent studies suggest that data assimilation (DA) of remotely sensed products is the path forward to estimate the spatial distribution of relevant snowpack
45 characteristics (Aalstad et al., 2018; Charrois et al., 2016; Cortés & Margulis, 2017; Margulis et al., 2016; Smyth et al., 2020; Stigter et al., 2017). Using DA techniques it is possible to fuse model simulations and multiple remote sensing datasets to improve the snowpack simulations. In particular, the **snow cover area** was assimilated in many case studies due to its **widespread availability** (e.g. Baba et al., 2018, Alonso-González et al., 2021). Yet, **the extent of the snow cover provides no direct information on the internal state of the snowpack and is blind to snowpack changes when the pixel is fully snow**
50 **covered.**

The ice-surface temperature (IST), is a key state variable for simulating the snowpack evolution. Physically-based snowpack models solve the energy balance equation iteratively along the time dimension, estimating the IST at each timestep of the model (Essery, 2015; Liston & Elder, 2006). Also it is a key parameter to estimate the emitted energy as outgoing longwave radiation. The estimation of the IST **allows forecasting the occurrence of the melting events, as when the IST reaches 0°C all**
55 **the added energy is converted to melt.** Thus, assimilating IST may provide key information about the timing of the melting events. Also, the assimilation of the land surface temperature (LST) (i.e. the temperature of the earth surface independently if it is snow covered) may improve the snowpack simulations by different mechanisms. The IST is physically bound to the melting point temperature, **while** ^{since?} ^{about} once the snow melts, the LST can exceed 0°C. Thus the assimilation of the LST may indirectly provide information of the snow cover area too. **Also, it should be possible to improve the snow simulations by**
60 **retrieving thermal information when there is no sunlight, like during the night time or during the polar night at high latitudes.**



A previous study showed that IST DA improved the surface ice mass balance simulations of the Greenland ice sheet (Navari et al., 2016), fusing IST stimations with the CROCUS snow model. On the other hand, previous research suggested little improvements in the mass simulations after assimilating LST simulations retrieved from the Meteosat Second Generation (MSG) (~6km spatial resolution) in the Alps. However, the coarse resolution of the LST products of MSG prevents ^{its} use in
65 complex terrain, and more research is needed to assess the potential of high resolution LST DA.

The thermal imagery already available only offers coarse resolution for the snow applications over complex terrain (MODIS, Sentinel-3) or long revisiting times (Landsat). This has probably prevented the study of the impact of LST DA, although recent research suggests that LST can provide useful information to retrieve internal snowpack properties (Colombo et al., 2019), a capability that can be exploited from satellites (Colombo et al., 2023). The availability of high spatiotemporal
70 resolution LST products will be improved in the short term with the appearance of new satellites, such as the French-Indian mission Thermal infraRed Imaging Satellite for High-resolution Natural resource Assessment (TRISHNA) (Lagouarde et al., 2018). TRISHNA is expected to provide surface temperature measurements at 60 m spatial resolutions every 3 days. Also, given the agenda of the space agencies high resolution thermal infrared retrievals will be readily accessible in the near future. For instance the LSTM (Copernicus Land Surface Temperature Monitoring) (Koetz et al., 2018) which will offer similar
75 observations as TRISHNA but with improved spectral, spatial and temporal resolutions and the SBG satellite (Surface Biology and Geology)(Cawse-Nicholson et al., 2021) from NASA which will provide similar high resolution TIR images of the surface of the earth. The combination of these three missions may eventually provide close to bi-daily (day and night time) high resolution thermal infrared observations of the earth surface. In this context the objectives of this work are; i) to test the potential of the LST to improve the snowpack simulations and ii) explore the effect of increasing the temporal
80 observations.

A convenient approach to emulate future remote sensing observations is to use an **Observing System Simulation Experiment (OSSE)**. For example, Navari et al. (2016) assessed the feasibility to integrate ice surface temperatures in a regional climate model estimate of Greenland ice sheet surface mass balance through an OSSE. Synthetic experiments were also used to explore the potential of data assimilation techniques in improving the snowpack simulations (Clark et al., 2006; Revuelto et al., 2021; Smyth et al., 2019, Deschamps-Berger et al., 2022).
85

Here, we designed an OSSE to evaluate the benefit of future remote sensing LST to simulate seasonal SWE. In this experiment, synthetic LST and SWE data were generated in several climatic regions. Synthetic LST data were assimilated into a snowpack model under different cloud cover scenarios and satellite revisit times. The benefit of assimilating LST was studied by comparing the posterior SWE to the synthetic SWE.

90 2 Data and methods

We selected five sites in snow-dominated regions of Europe, spanning 40° of latitude from the Pyrenees mountains to the Svalbard archipelago. The sites were chosen every 10° of latitude approximately to sample different climatic influences .



The southern sites (Gerlachovský štít, Bigorre) are located in high mountain regions (Pyrenees, Tatra), Finse is located on a high elevation plateau (Hardangervidda), whereas Tromsø and Ny-Ålesund are near sea level. Gerlachovský štít is in the Eastern Europe and its climate is influenced by its continental characteristics, Tromsø and Ny-Ålesund exhibit obvious polar climates and Bigorre shows a montane climate with Mediterranean influences.

Table 1: Geographical coordinates and elevation (m a.s.l.) of the ERA5-Land centroids used in the study

Centroid	Longitude	Latitude	Elevation
Ny-Ålesund	12.0°	78.9°	124
Tromsø	19.8°	69.5°	647
Finse	07.5°	60.6°	147
Gerlachovský štít	20.2°	49.2°	1479
Bigorre	00.1°	42.9°	1843

We used ERA5-land surface reanalysis data (Muñoz-Sabater et al., 2021) to force the Flexible Snow Model (FSM2) (Essery, 2015) over four consecutive hydrological years from 01/Sep/2017 to 31/Aug/2021. From this simulation, we retrieved the SWE and LST time series, which were considered as the synthetic truth. The LST was exported at 13:00 local time, corresponding to foreseen TRISHNA overpass time. To mimic instrumental noise, we added to the LST time series a Gaussian noise with a zero mean and a standard deviation of 1.5 K. This standard deviation was chosen as an intermediate value between the reported RMSE of errors obtained by the comparison of Landsat 8 with in situ measurements of the snow surface temperature (RMSE = 2.0 °K) (Robledano et al., 2022) and the expected performance of LST products delivered by the TRISHNA mission (Lagouarde et al., 2018).

The synthetic LST time series were downsampled with a period of 16 and 3 days, to emulate revisit times of Landsat 8 and TRISHNA respectively. We simulated the impact of cloud cover by further removing values in the synthetic LST time series at random dates selected from an uniform distribution. We defined four different cloud cover scenarios with probabilities of 0%, 25%, 50% and 75%.

For each site, we created a new, degraded meteorological forcing to run FSM2 by averaging ERA5-Land data from the nearest 9 cells (i.e. resampling the spatial resolution from 10 km to 30 km approximately). We further perturbed the precipitation field after aggregation using a multiplicative factor of 0.5. This strong perturbation was chosen to emulate precipitation biases that are typically found in global reanalyses and large scale precipitation products (Beck et al., 2019) potentially leading to large underestimation of SWE in mountain regions (Wrzesien et al., 2019). A similar value was already used in previous synthetic snow data assimilation experiment (Deschamps-Berger et al., 2022).



Intéressant! (à la fois le modèle F2M2 et MuSA)

The degraded meteorological forcing and synthetic LST were used to feed the **Multiple Snow data Assimilation system (MuSA)**. MuSA is an open source ensemble based data assimilation toolbox **built around the FSM2 model** (Alonso-González et al., 2022). We used the same initial conditions to run FSM2 within MuSA (soil temperature profile, initial LST and absence of snow), therefore we did not perform a spin-up. The assimilation experiments were done using the **Particle Batch Smoother (PBS)** (Margulis et al., 2015). **Smoother algorithms are typically used to develop reanalyses** as the whole time series of information are available, whereas **filtering is rather used for operational forecasting** where future observations respective to the analysis step are not available (Largeron et al., 2020). A description of these algorithms and the MuSA toolbox can be found in Alonso-González et al., (2022).

The prior ensemble of FSM2 simulations was composed of **300 particles** that were generated by perturbing the air temperature (additive perturbation) and the precipitation (multiplicative perturbation). The perturbations were time-invariant and randomly drawn from a **normal distribution of mean $\mu = 0$ and standard deviation $\sigma = 2$ (temperature) and a log-normal distribution with mean $\mu = 0.45$ and standard deviation $\sigma = 0.8$** of the underlying normal distribution (precipitation). These parameters were chosen to cover the expected differences between the “truth” forcing and the degraded forcing and were obtained by **preliminary trial and error tests**. **Plus de détails?**

While the PF performs the analysis sequentially, i.e. each time an observation occurs, the PBS is a smoother, hence it **assimilates all the available observations in a single time window, propagating information from the observations forward and backward in time**. Here the assimilation time **window** was defined as **a hydrological year** (i.e. one snow season). In both the PF and PBS, prior weights of ensemble members (particles) are updated based on the likelihood, i.e. a measure of the distance between the predictions of each particle and the observations. The posterior weights are then used to estimate posterior statistics from the ensemble, typically its weighted mean and weighted standard deviation. In the case of the PF, we used the **bootstrap** resampling algorithm to eliminate particles with low weights and replicate particles with high weights **by sampling with replacement randomly from the probability distribution of the updated weights**. To prevent the filter from collapsing (all the weight is shared by few and eventually just one particle), **new perturbation parameters** were drawn from a normal approximation of the posterior from the previous analysis step at each new analysis step, instead of resampling both the states of the model and the parameters. A rigorous description of the algorithms, the underlying theory and implementation details can be found in Alonso-González et al., (2022).

For each site and each cloud cover scenario, we ran MuSA and generated **a posterior SWE**. However, the output of MuSA is **stochastic** due to the random generation of the forcing perturbation parameters. Also the position of the gaps in the different cloud cover scenarios and the gaussian noise are random. Therefore, to increase the statistical robustness of our results, we repeated each assimilation experiment 100 times, drawing new gaps and gaussian noise for each replicate of the experiment. **J'aimerais bien avoir un schéma, en particulier pour illustrer tous les endroits où il y a de l'échantillonnage**. This created an **ensemble of posterior SWE** which was compared to the synthetic true SWE.

In total, for a given site, MuSA was run **1600 times** (**100 replicates x 4 cloud cover scenarios x 2 revisit times x 2 DA algorithms**). This corresponds to **480,000 FSM2 runs** (**300 particles by MuSA run**), summing up to **2,400,000 FSM2 runs** considering the 5 study areas.

Temps de calcul?



3 Results

As expected, the degradation of the ERA5-Land meteorological forcing had a large impact on the **open loop SWE simulations** (Fig. 2). In comparison with the true SWE, the average normalized RMSE (nRMSE) was 57% (after removing the summer time months of July and August). The aggregation of the data from neighboring ERA5-Land cells and the scaling of the precipitation caused an **overall reduction in the simulated SWE**, leading to a **shorter snow seasons** at all sites (Fig. 2). However, the LST assimilation with the PBS substantially improved the SWE simulations (Fig. 2). This improvement was evident at both revisit times, although the 3 day revisit scenario (nRMSE = 3%) outperformed the 16 day revisit scenario (nRMSE = 7%).

155 The posterior SWE series in Figure 2 were averaged from an ensemble of 100 replicates (see Sect. 2. Data and methods). **Figure 3** shows every posterior SWE realization in the case of the Tromsø site under different cloud cover scenarios when assimilating LST at 3-day resolution. This Figure shows that the spread of the posterior ensemble increased with the cloud cover probability. The standard deviation of the obtained nRMSE values over each cloud cover scenario ranged from 1% to 7% in this particular case. **rester au présent**

160 Figure 4 summarizes the results of the PBS from all experiments under every cloud cover and revisit scenarios. In all cases, the data assimilation significantly reduced the nRMSE in comparison with the open loop simulations. The nRMSE was always higher for the 16-day revisit compared to the 3-day revisit, but the difference was more pronounced under the 50% and 75% cloud cover scenarios. In addition, the standard deviation of the nRMSE of the 100 replicates was higher for the 16-day compared to the 3-day revisit scenarios. Both the averaged nRMSE and standard deviations increased with the cloud cover with an average nRMSE for all the sites ranging between 3% and 8% in the case of the 3-day revisit experiments and 7% and 22% for the 16-day revisit experiments.

170 Figure 5 shows the distribution of the mean of the posterior precipitation perturbation parameters obtained from the 100 data assimilation runs using the PBS. **présent** It demonstrates that the assimilation of LST reduced the error on the precipitation forcing, since the posterior parameter distributions intersect the actual perturbation factor of 2 that was used to degrade the input precipitation. However, the difference between the true precipitation and the degraded precipitation **présent** may not exactly equal to the scaling factor of 2 since the precipitation forcing was also perturbed by aggregating precipitation from the surrounding ERA5-Land cells (Sect. 2). As observed above in Figure 4, the standard deviation of the posterior perturbation parameters of the replicates increased when comparing the 3-day with the 16-day revisit scenarios, as well as with the cloud cover probability.

180 Whereas the above results **présent** show that the PBS algorithm clearly improved the SWE simulation, it was not the case with the PF. Figure 6 summarizes the results of the same experiments shown in Figure 4, but using the PF instead of the PBS. In this case the improvement in the average nRMSE of the posterior simulations was not as obvious with respect to the open loop as



in the PBS experiments. Although, in most cases the nRMSE showed a moderate improvement compared with the open loop on average, several runs among the 100 replicates had a higher nRMSE than the open loop run. The revisit or cloud cover scenarios had no clear effect on the nRMSE. As in the PBS case, there was no significant difference between the different locations, except for Ny-Ålesund which yielded a higher nRMSE standard deviation than for the other locations. This is a consequence of the very cold conditions in this polar region, as some particles became “glaciers” due to the perturbed forcing (non zero SWE at the end of the hydrological year).

4 Discussion

Navari et al. (2016) showed the potential of IST data assimilation to improve the surface mass balance of the Greenland ice sheet in a regional climate simulation with an Ensemble batch smoother. Our study also suggests that the assimilation of LST can improve seasonal snow simulations in sites with different climate contexts. With the PBS, the improvement was substantial independently of the site, i.e. the climatic context did not exhibit an obvious influence on the results. However, our results with the PF also supports the conclusions of Piazzì et al. (2019), who did not obtain obvious improvements in the posterior SWE simulations after assimilating LST using an Ensemble Kalman Filter. Therefore, our study provide an explanation of the contrasting performances found by Navari et al. (2016) and Piazzì et al. (2019). While Navari et al. (2016) used smoothers, Piazzì et al. (2019) used a filter. A filter updates the simulations sequentially while smoothers update the whole season in batch. This in-batch assimilation allows the propagation of the information of the observations backwards in the simulation. The saw tooth pattern reported by Piazzì et al. (2019) is reminiscent of dynamical inconsistencies associated with filters (Dunne & Entekhabi, 2005). Also, the performance of the LST data assimilation reported by Piazzì et al. (2019) was probably hampered by the coarse resolution of the MSG LST products that were used to update snowpack simulations at the point scale. In the specific case of the LST, considering the observations of the whole snow season in a batch may be key to have a positive impact on the posterior SWE. The trajectory of the LST in seasonal snow dominated regions exhibits a characteristic pattern as the physical bounds of the IST are different from the LST. Once the snow melts, the LST can rise above the water melting point and therefore the trajectory of the LST may be a good indicator of the length of the snow season. However, this should not be the only reason, as Navari et al. (2016) experiments were developed over the Greenland ice sheet where there is a permanent ice cover. During the melting season the IST is fixed to the melting point temperature, providing information on the duration of the melting period. Also the occurrence of winter time melt events should be visible in the TIR domain. The information of the whole seasonal trajectory of the LST is propagated to the posterior by using a smoother, but not by using a filter. This is highlighted at the Ny-Ålesund site, where the polar conditions made snowmelt impossible at the end of the hydrological year for some of the replicates leading to very high nRMSE using the PF. These results suggest that the LST may be less beneficial to snowmelt forecasting applications where the use of filters is more extended to update the model as new observations arise, but it should be valuable information to improve snow reanalyses which aim to reconstruct snow cover climatologies.



215 Our results also suggest that even the currently available thermal infrared estimations of the LST from Landsat missions
have the potential to significantly improve SWE simulations despite a revisit time of 16 days. The emulated revisiting times
of both Landsat and TRISHNA are the expected values at the equator, and can be lower in other latitudes. Here we did not
study the effect of the spatial resolution but hypothesized that high resolution (i.e. Landsat-like) is needed for snow cover
simulations in the studied regions. Landsat TIR images have a 100 m resolution which makes them suitable to sample the
220 slope scale in mountain terrain, hence homogeneous conditions in the energy balance budget (Baba et al., 2019). Despite the
low revisiting times of the Landsat mission, Landsat TIR imagery may be useful to improve SWE simulations using a
smoother data assimilation algorithm, an approach that to our knowledge has not been explored yet. **More research should be
carried out on this topic, especially in the context of joint assimilation experiments where more than one variable is
assimilated.**

225 **Comment leurs résultats changeraient-ils à l'échelle du bassin versant? Application pour le débit, par exemple**
Nevertheless, the change in revisit from 3 to 16 days in our experiments translated into an approximate doubling of the
posterior nRMSE. Therefore, we expect significant progress with TRISHNA observations, not to mention the enhanced
spatial resolution (approximately 60 m). The benefit of the 3 day revisit was particularly evident under 50% to 75% cloud
cover. This should be considered, as previous global estimates of the cloud cover suggest values closer to our highest cloud
230 cover scenario (Wylie et al., 2005). For instance, cloud cover probability in MODIS products reached 60% in the Alps and
50% in the Pyrenees (Gascoin et al., 2015; Parajka & Blöschl, 2008).

In any case, under both revisit scenarios, the cloud cover decreased the precision of the replicates of the posterior SWE, i.e.
the variability between repeated experiments, but the average was only marginally affected. In other words, the cloud cover
reduced the robustness of the data assimilation, but even regions with a persistent cloud cover could benefit from LST
235 assimilation. The different replicates of each experiment exhibited different results, with a variance that increased with the
number of gaps introduced in the synthetic LST observations, suggesting that not all the combinations of observations are
equally informative. This was also obvious regarding the posterior precipitation perturbation parameters, as the standard
deviation of the different replicates increased with the percentage of cloud cover.

Despite the promising potential of the LST to improve SWE simulation, some **limitations** of the current study inherent to the
240 synthetic nature of the OSSE should be taken into consideration. **The use of the same model to generate the real synthetic
SWE and for the assimilation processes avoids the uncertainty as a consequence of the model internal parameters and
structural uncertainty.** This simplifies the interpretation of the results of our work, but needs to be taken into account in real-
world scenarios. The simulation of the cloud cover scenarios was generated by selecting random dates from an uniform
distribution. **However, in some regions the cloud cover exhibits marked seasonal patterns** (Sudmanns et al., 2020), that may
245 challenge to update the snowpack simulations **even with smoothers** if cloud cover is more frequent during key periods in the
snow season.



5 Conclusions

The motivation of this study on LST data assimilation is the upcoming launch of high resolution thermal infrared spatial missions with improved revisit time in the next few years. We implemented a synthetic data assimilation experiment to study the potential of the LST in improving SWE simulations along a latitudinal gradient in the northern hemisphere. The methodology was based on the generation of synthetic LST estimations and SWE true estimates, and a prior ensemble of SWE simulations generated by forcing the FSM2 model with degraded meteorological fields. The MuSA snow data assimilation software was used to generate SWE posterior time series using the particle batch smoother and particle filter algorithms to be compared with synthetic true SWE.

The results suggest that the assimilation of LST has a great potential to improve seasonal snowpack simulations across all the tested sites. Gap-free LST series, improved the average nRMSE of the open loop simulations from 57% to 7% and 3% for the 16 days and 3 day revisiting times respectively. However, a lower revisit frequency caused an increase in the variance of the nRMSE when the runs are replicated 100 times, meaning that the assimilation becomes less robust. This conclusion was more evident with high cloud cover scenarios, highlighting the importance of the revisit time in thermal infrared remote sensing to reduce the uncertainty on the updated SWE.

The type of data assimilation was also key to explain the role of LST in improving SWE simulations. The particle batch smoother strongly improved the simulations, whereas the particle filter was much less performant and could even cause a degradation of the simulations at a cold polar site.

Overall, our results encourage a more systematic use of the current LST products within snow data assimilation studies, especially if the objective is to perform a snow reanalysis which can benefit from observations acquired over an entire snow season.

Acknowledgements

This work was supported by the Centre National d'Etudes Spatiales (CNES) through the postdoctoral grant of Esteban Alonso-González, the PhD grant of Sara Arioli and the TOSCA program.

Code and data availability.

The MuSA code is available at github.com/ealonsogzl/MuSA. The original FSM2 code is found at github.com/RichardEssery/FSM2, and in the MuSA repository with slight modifications from the original version. ERA5-Land data is available for download from the Copernicus Climate Data Store.

275



Author contributions. Conceptualization: EAG, SG. Data curation: EAG. Formal analysis: EAG, SG, SA, GP. Funding acquisition: EAG, SG. Investigation: EAG, SG. Methodology: EAG. Project administration: EAG, SG. Resources: EAG, SG. Software: EAG. Supervision: SG, GP. Validation: EAG. Visualization: EAG. Writing - original draft preparation: EAG. Writing review & editing: EAG, SG, SA, GP.

280

Competing interests. The authors declare that they have no conflict of interest.

References

- Aalstad, K., Westermann, S., Schuler, T. V., Boike, J., & Bertino, L. (2018). Ensemble-based assimilation of fractional snow-covered area satellite retrievals to estimate the snow distribution at Arctic sites. *Cryosphere*, 12(1), 247–270.
285 <https://doi.org/10.5194/tc-12-247-2018>
- Alonso-González, E., Gutmann, E., Aalstad, K., Fayad, A., Bouchet, M., & Gascoin, S. (2021). Snowpack dynamics in the Lebanese mountains from quasi-dynamically downscaled ERA5 reanalysis updated by assimilating remotely sensed fractional snow-covered area. *Hydrology and Earth System Sciences*, 25(8), 4455–4471. <https://doi.org/10.5194/hess-25-4455-2021>
- 290 Alonso-González, Esteban, Aalstad, K., Baba, M. W., Revuelto, J., López-Moreno, J. I., Fiddes, J., et al. (2022). MuSA: The Multiscale Snow Data Assimilation System (v1.0). *Geoscientific Model Development Discussions*, 1–43. <https://doi.org/10.5194/gmd-2022-137>
- Baba, M. W., Gascoin, S., & Hanich, L. (2018). Assimilation of Sentinel-2 data into a snowpack model in the High Atlas of Morocco. *Remote Sensing*, 10(12), 1982. <https://doi.org/10.3390/rs10121982>
- 295 Baba, M. W., Gascoin, S., Kinnard, C., Marchane, A., & Hanich, L. (2019). Effect of Digital Elevation Model Resolution on the Simulation of the Snow Cover Evolution in the High Atlas. *Water Resources Research*, 55(7), 5360–5378. <https://doi.org/10.1029/2018WR023789>
- Ballesteros-Cánovas, J. A., Trappmann, D., Madrigal-González, J., Eckert, N., & Stoffel, M. (2018). Climate warming enhances snow avalanche risk in the Western Himalayas. *Proceedings of the National Academy of Sciences of the United States of America*, 115(13), 3410–3415. <https://doi.org/10.1073/pnas.1716913115>
- 300 Barnett, T. P., Adam, J. C., & Lettenmaier, D. P. (2005). Potential impacts of a warming climate on water availability in snow-dominated regions. *Nature*, 438(7066), 303–309. <https://doi.org/10.1038/nature04141>
- Beck, H. E., Pan, M., Roy, T., Weedon, G. P., Pappenberger, F., van Dijk, A. I. J. M., et al. (2019). Daily evaluation of 26 precipitation datasets using Stage-IV gauge-radar data for the CONUS. *Hydrology and Earth System Sciences*, 23(1), 207–
305 224. <https://doi.org/10.5194/hess-23-207-2019>



- Cawse-Nicholson, K., Townsend, P. A., Schimel, D., Assiri, A. M., Blake, P. L., Buongiorno, M. F., et al. (2021). NASA's surface biology and geology designated observable: A perspective on surface imaging algorithms. *Remote Sensing of Environment*, 257, 112349. <https://doi.org/10.1016/j.rse.2021.112349>
- Charrois, L., Cosme, E., Dumont, M., Lafaysse, M., Morin, S., Libois, Q., & Picard, G. (2016). On the assimilation of optical reflectances and snow depth observations into a detailed snowpack model. *Cryosphere*, 10(3), 1021–1038. <https://doi.org/10.5194/tc-10-1021-2016>
- 310 Clark, M. P., Slater, A. G., Barrett, A. P., Hay, L. E., McCabe, G. J., Rajagopalan, B., & Leavesley, G. H. (2006). Assimilation of snow covered area information into hydrologic and land-surface models. *Advances in Water Resources*, 29(8), 1209–1221. <https://doi.org/10.1016/j.advwatres.2005.10.001>
- 315 Colombo, R., Garzonio, R., Di Mauro, B., Dumont, M., Tuzet, F., Cogliati, S., et al. (2019). Introducing Thermal Inertia for Monitoring Snowmelt Processes With Remote Sensing. *Geophysical Research Letters*, 46(8), 4308–4319. <https://doi.org/10.1029/2019GL082193>
- Colombo, Roberto, Pennati, G., Pozzi, G., Garzonio, R., Di Mauro, B., Giardino, C., et al. (2023). Mapping snow density through thermal inertia observations. *Remote Sensing of Environment*, 284, 113323. <https://doi.org/10.1016/j.rse.2022.113323>
- 320 Cortés, G., & Margulis, S. (2017). Impacts of El Niño and La Niña on interannual snow accumulation in the Andes: Results from a high-resolution 31 year reanalysis: El Niño Effects on Andes Snow. *Geophysical Research Letters*, 44(13), 6859–6867. <https://doi.org/10.1002/2017GL073826>
- Deschamps-Berger, C., Cluzet, B., Dumont, M., Lafaysse, M., Berthier, E., Fanise, P., & Gascoin, S. (2022). Improving the Spatial Distribution of Snow Cover Simulations by Assimilation of Satellite Stereoscopic Imagery. *Water Resources Research*, 58(3), e2021WR030271. <https://doi.org/10.1029/2021WR030271>
- 325 Dozier, J., Bair, E. H., & Davis, R. E. (2016). Estimating the spatial distribution of snow water equivalent in the world's mountains. *Wiley Interdisciplinary Reviews: Water*, n/a-n/a. <https://doi.org/10.1002/wat2.1140>
- Dunne, S., & Entekhabi, D. (2005). An ensemble-based reanalysis approach to land data assimilation. *Water Resources Research*, 41(2). <https://doi.org/10.1029/2004WR003449>
- 330 Essery, R. (2015). A factorial snowpack model (FSM 1.0). *Geoscientific Model Development*, 8(12), 3867–3876. <https://doi.org/10.5194/gmd-8-3867-2015>
- Gascoin, S., Hagolle, O., Huc, M., Jarlan, L., Dejoux, J. F., Szczypta, C., et al. (2015). A snow cover climatology for the Pyrenees from MODIS snow products. *Hydrology and Earth System Sciences*, 19(5), 2337–2351. <https://doi.org/10.5194/hess-19-2337-2015>
- 335 Henderson, G. R., Peings, Y., Furtado, J. C., & Kushner, P. J. (2018). Snow–atmosphere coupling in the Northern Hemisphere. *Nature Climate Change*, 8(11), 954–963. <https://doi.org/10.1038/s41558-018-0295-6>
- Kinar, N. J., & Pomeroy, J. W. (2015). Measurement of the physical properties of the snowpack. *Reviews of Geophysics*, 53(2), 481–544. <https://doi.org/10.1002/2015RG000481>



- 340 Koetz, B., Bastiaanssen, W., Berger, M., Defournay, P., Del Bello, U., Drusch, M., et al. (2018). High Spatio- Temporal Resolution Land Surface Temperature Mission - a Copernicus Candidate Mission in Support of Agricultural Monitoring. In IGARSS 2018 - 2018 IEEE International Geoscience and Remote Sensing Symposium (pp. 8160–8162). <https://doi.org/10.1109/IGARSS.2018.8517433>
- Lagouarde, J., Bhattacharya, B. K., Crébassol, P., Gamet, P., Babu, S. S., Boulet, G., et al. (2018). The Indian-French
345 Trishna Mission: Earth Observation in the Thermal Infrared with High Spatio-Temporal Resolution (pp. 4078–4081). Presented at the IGARSS 2018 - 2018 IEEE International Geoscience and Remote Sensing Symposium. <https://doi.org/10.1109/IGARSS.2018.8518720>
- Largeron, C., Dumont, M., Morin, S., Boone, A., Lafayesse, M., Metref, S., et al. (2020). Toward Snow Cover Estimation in Mountainous Areas Using Modern Data Assimilation Methods: A Review. *Frontiers in Earth Science*, 8. Retrieved from
350 <https://www.frontiersin.org/article/10.3389/feart.2020.00325>
- Liston, G. E., & Elder, K. (2006). A distributed snow-evolution modeling system (snowmodel). *Journal of Hydrometeorology*, 7(6), 1259–1276. <https://doi.org/10.1175/JHM548.1>
- Margulis, S. A., Giroto, M., Cortés, G., & Durand, M. (2015). A particle batch smoother approach to snow water equivalent estimation. *Journal of Hydrometeorology*, 16(4), 1752–1772. <https://doi.org/10.1175/JHM-D-14-0177.1>
- 355 Margulis, S. A., Cortés, G., Giroto, M., & Durand, M. (2016). A landsat-era Sierra Nevada snow reanalysis (1985-2015). *Journal of Hydrometeorology*, 17(4), 1203–1221. <https://doi.org/10.1175/JHM-D-15-0177.1>
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., et al. (2021). ERA5-Land: A state-of-the-art global reanalysis dataset for land applications. *Earth System Science Data*, 13(9), 4349–4383. <https://doi.org/10.5194/essd-13-4349-2021>
- 360 Musselman, K. N., Lehner, F., Ikeda, K., Clark, M. P., Prein, A. F., Liu, C., et al. (2018). Projected increases and shifts in rain-on-snow flood risk over western North America. *Nature Climate Change*, 8(9), 808–812. <https://doi.org/10.1038/s41558-018-0236-4>
- Navari, M., Margulis, S. A., Bateni, S. M., Tedesco, M., Alexander, P., & Fettweis, X. (2016). Feasibility of improving a priori regional climate model estimates of Greenland ice sheet surface mass loss through assimilation of measured ice
365 surface temperatures. *The Cryosphere*, 10(1), 103–120. <https://doi.org/10.5194/tc-10-103-2016>
- Parajka, J., & Blöschl, G. (2008). Spatio-temporal combination of MODIS images - Potential for snow cover mapping. *Water Resources Research*, 44(3). <https://doi.org/10.1029/2007WR006204>
- Piazzzi, G., Campo, L., Gabellani, S., Castelli, F., Cremonese, E., Cella, U. M. di, et al. (2019). An Enkf-Based Scheme for Snow Multivariable Data Assimilation at an Alpine Site. *Journal of Hydrology and Hydromechanics*, 67(1), 4–19.
370 <https://doi.org/10.2478/johh-2018-0013>
- Raleigh, M. S., Lundquist, J. D., & Clark, M. P. (2015). Exploring the impact of forcing error characteristics on physically based snow simulations within a global sensitivity analysis framework. *Hydrology and Earth System Sciences*, 19(7), 3153–3179. <https://doi.org/10.5194/hess-19-3153-2015>



- 375 Rango, A., & Martinec, J. (1979). Application of a Snowmelt-Runoff Model Using Landsat Data. *Nordic Hydrology*, 10(4), 225–238. <https://doi.org/10.2166/nh.1979.0006>
- Revuelto, J., Cluzet, B., Duran, N., Fructus, M., Lafaysse, M., Cosme, E., & Dumont, M. (2021). Assimilation of surface reflectance in snow simulations: Impact on bulk snow variables. *Journal of Hydrology*, 603, 126966. <https://doi.org/10.1016/j.jhydrol.2021.126966>
- 380 Robledano, A., Picard, G., Arnaud, L., Larue, F., & Ollivier, I. (2022). Modelling surface temperature and radiation budget of snow-covered complex terrain. *The Cryosphere*, 16(2), 559–579. <https://doi.org/10.5194/tc-16-559-2022>
- Smyth, E. J., Raleigh, M. S., & Small, E. E. (2019). Particle Filter Data Assimilation of Monthly Snow Depth Observations Improves Estimation of Snow Density and SWE. *Water Resources Research*, 55(2), 1296–1311. <https://doi.org/10.1029/2018WR023400>
- 385 Smyth, E. J., Raleigh, M. S., & Small, E. E. (2020). Improving SWE Estimation With Data Assimilation: The Influence of Snow Depth Observation Timing and Uncertainty. *Water Resources Research*, 56(5). <https://doi.org/10.1029/2019WR026853>
- Stigter, E. E., Wanders, N., Saloranta, T. M., Shea, J. M., Bierkens, M. F. P., & Immerzeel, W. W. (2017). Assimilation of snow cover and snow depth into a snow model to estimate snow water equivalent and snowmelt runoff in a Himalayan catchment. *The Cryosphere*, 11(4), 1647–1664. <https://doi.org/10.5194/tc-11-1647-2017>
- 390 **Sturm, M., Goldstein, M. A., & Parr, C. (2017). *Water and life from snow: A trillion dollar science question.***
- Sudmanns, M., Tiede, D., Augustin, H., & Lang, S. (2020). Assessing global Sentinel-2 coverage dynamics and data availability for operational Earth observation (EO) applications using the EO-Compass. *International Journal of Digital Earth*, 13(7), 768–784. <https://doi.org/10.1080/17538947.2019.1572799>
- 395 **Wrzesien, M. L., Pavelsky, T. M., Durand, M. T., Dozier, J., & Lundquist, J. D. (2019). *Characterizing Biases in Mountain Snow Accumulation From Global Data Sets.* *Water Resources Research*, 55(11), 9873–9891. <https://doi.org/10.1029/2019WR025350>**
- Wylie, D., Jackson, D. L., Menzel, W. P., & Bates, J. J. (2005). Trends in Global Cloud Cover in Two Decades of HIRS Observations. *Journal of Climate*, 18(15), 3021–3031. <https://doi.org/10.1175/JCLI3461.1>

400

405





Figure 1: Main workflow of the OSSE.



Une mauvaise modélisation de la densité pourrait-elle mener à une surestimation de l'ÉÉN?



Ne serait-il pas possible de surestimer le couvert de neige? Il me semble qu'en réalité ça peut arriver

Figure 2: Comparison of **hourly** time series of synthetic true SWE with the open loop simulations (degraded forcing) and the posterior SWE after assimilating LST with a revisit time of 16 days or 3 days (**0% cloud cover scenario**) using the PBS. Here, **the posterior SWE is the average of the 100 replicates.**

Why not use the entire ensemble and compute the CRPS? Or make use of the ensemble in any way?



415

L'ajout d'une légende serait bien




Figure 3: Comparison of hourly time series of synthetic true SWE (green) with the open loop simulations (degraded forcing, in black), and the 100 replicates of posterior SWE after assimilating LST with a revisit time of 3 days and varying cloud cover scenarios and the average of the experiments (blue and dark blue respectively).




Figure 4: Normalized root mean square error (nRMSE) of the posterior SWE from the PBS compared with the synthetic true SWE for each experiment. The bars indicate the mean nRMSE while the error bars indicate the standard deviation in the 100 replicates.

Ce serait bien d'avoir aussi des barres d'incertitude pour le open loop



420




Figure 5: Boxplots showing the distribution of the posterior precipitation perturbation parameter for each experiment estimated by the PBS. **The dashed line indicates the true perturbation that was applied to the forcing.**

Je ne comprends pas pourquoi la perturbation (ligne pointillée) ici est de 2 alors qu'à la page 4 ligne 110 on indique que c'est 0.5 Est-ce que c'est pris dans l'autre sens? 0.5 vs x 2?




Figure 6: Boxplots showing the distribution of the *posterior precipitation perturbation parameter* for each experiment estimated by the PF. The *dashed line* indicates the true perturbation that was applied to the forcing.

Je ne vois pas la dashed line