

Why is ozone in South Korea and the Seoul Metropolitan Area so high and increasing?

5 Nadia K. Colombi^{1*}, Daniel J. Jacob¹, Laura Hyesung Yang², Shixian Zhai², Viral Shah^{3,4}, Stuart K. Grange⁵, Robert M. Yantosca², Soontae Kim⁶, and Hong Liao⁷

¹Harvard University, Department of Earth and Planetary Science, Cambridge, MA 02138, USA

²Harvard University, John A. Paulson School of Engineering and Applied Sciences, Cambridge, MA 02138, USA

³NASA Global Modeling and Assimilation Office, Goddard Space Flight Center, Greenbelt, MD 20771, USA

10 ⁴Science Systems and Applications, Inc., Lanham, MD 20706, USA

⁵Empa, Swiss Federal Laboratories for Materials Science and Technology, Überlandstrasse 129 8600 Dübendorf, Switzerland

⁶Ajou University, Department of Environmental and Safety Engineering, Suwon, Gyeonggi 16499, Republic of Korea

15 ⁷Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control, Jiangsu Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing 210044, China

*Correspondence to: Nadia Colombi (ncolombi@g.harvard.edu)

20

Abstract. Surface ozone pollution in South Korea has increased over the past two decades, despite efforts to decrease emissions, and is pervasively in exceedance of the maximum daily 8-hr average (MDA8) standard of 60 ppb. Here, we investigate the 2015-2019 trends in surface ozone and NO₂ concentrations over South Korea and the Seoul Metropolitan Area (SMA), focusing on the 90th percentile MDA8 ozone as an air quality metric. We use a random forest algorithm to remove the effect of meteorological variability on the 2015-2019 trends and find an ozone increase of up to 1.5 ppb a⁻¹ in April-May while NO₂ decreases by 22%. GEOS-Chem model simulations including recent chemical updates can successfully simulate surface ozone over South Korea and China as well as the very high free tropospheric ozone observed above 2 km altitude (mean 75 ppb in May-June), and can reproduce the observed 2015-2019 emission-driven ozone trend over the SMA including its seasonality. Further investigation of the model trend for May, when meteorology-corrected ozone and its increase are the highest, reveals that a decrease in South Korea NO_x emissions is the main driver for the SMA ozone increase. Although this result implies that decreasing volatile organic compound (VOC) emissions is necessary to decrease ozone, we find that ozone would still remain above 80 ppb even if all anthropogenic emissions in South Korea were shut off. China contributes only 8 ppb to this elevated South Korea background and ship emissions contribute only a few ppb. Zeroing out all anthropogenic emissions in East Asia in the model indicates a remarkably high external background

25

30

35

Deleted: emission-driven

Deleted: SMA

of 56 ppb, consistent with the high concentrations observed in the free troposphere, implying that the air quality standard in South Korea is not practically achievable unless this background external to East Asia can be decreased.

1 Introduction

40

Surface ozone is a severe air quality problem in South Korea and has become steadily worse over the past two decades (Gaudel et al., 2018; Yeo and Kim, 2021; Kim et al., 2021). Ozone often exceeds 90 ppb in the Seoul Metropolitan Area (SMA) where 50% of South Korea's population is located (Miyazaki et al., 2018). In 2015, Phase 2 of the Seoul Metropolitan Air Quality Control Master Plan established a national standard of 60 ppb for the maximum daily 8-hour average (MDA8) ozone concentration (MOE, 2016). However, no monitoring sites have been compliant with this standard in recent years and ozone has continued to increase (NIER, 2020). Improved understanding of the causes of elevated ozone in South Korea is crucial for developing effective emission control strategies.

Formatted: Indent: First line: 0", Right: 0.08", Space Before: 5.75 pt, Tab stops: Not at 0.68"

Deleted:

Deleted: ¶
45→

50

Ozone is produced in the troposphere by photochemical oxidation of volatile organic compounds (VOCs) in the presence of nitrogen oxides ($\text{NO}_x \equiv \text{NO} + \text{NO}_2$). Both VOCs and NO_x have large anthropogenic sources from combustion, and VOCs also have fugitive industrial and residential, as well as biogenic sources. Effectively reducing ozone concentrations requires knowledge of whether ozone production is NO_x - or VOC-limited. In the NO_x -limited regime, decreasing NO_x emissions decreases ozone while decreasing VOC emissions has little effect. In the VOC-limited regime, when NO_x concentrations are very high, decreasing NO_x emissions drives an increase in ozone while decreasing VOC emissions decreases ozone (Sillman et al., 1990). The Clean Air Policy Support System (CAPSS) bottom-up emission inventory in South Korea reports emission declines of 26% for NO_x and 25% for VOCs in Seoul over the 2000-2019 period (<https://www.air.go.kr/eng/capss/emission/sido.do?menuId=100>). Using satellite and surface observations of NO_2 , Seo et al. (2021) found that NO_x emissions declined in Seoul by 30% during the 2015-2019 period, and Bae et al. (2021) found an 18% decrease for the 2015-2018 period. On the other hand, Bauwens et al. (2022) found an increase in satellite-observed HCHO columns over South Korea by 1-2% a^{-1} for the 2005-2019 period, which does not support a decrease in VOC emissions.

Deleted: for

60

Ozone concentrations over South Korea depend not only on domestic emissions but also on the background from external sources. The KORUS-AQ aircraft campaign in May-June 2016 found free tropospheric concentrations above 2 km altitude frequently exceeding 80 ppb (Miyazaki et al., 2018; Sullivan et al., 2019; Gaubert et al., 2020; Crawford et al., 2021), which would affect surface ozone through subsidence. An obvious source of background ozone is China, where ozone is very high and increasing (K. Li et al., 2019, 2021), and would be transported to South Korea by westerly winds (Cuesta et al., 2018). But other background sources could also contribute. Lam and Cheung (2022) found that strong transport from the stratosphere can enhance springtime surface ozone by up to 8 ppb in East Asia. J. Li et al. (2016) estimated from a global model that long-range transport from outside East Asia could contribute 50-80% to annual surface ozone in the Korean peninsula. Wang et al. (2022) found an increase of

Deleted: Gaudel et al., 2018,

70

free tropospheric ozone over East Asia of 3.8 to 6.7 ppb per decade over the 1995-2017 period.

Deleted: from aircraft and ozonesonde data

75 The high and increasing ozone over South Korea could thus reflect a combination of decreasing NO_x emissions and/or increasing VOC emissions under VOC-limited conditions for ozone production (Jung et al, 2018), as well as high and increasing background ozone. Here we aim to better understand the factors controlling ozone and its increase in the SMA and more broadly over South Korea during the 2015-2019 period. We focus our analysis on the 90th percentile MDA8 ozone as a robust metric for polluted conditions. (Fiore et al., 1998; Sun et al., 2017; Wells et al., 2021). We use a random forest (RF) method (Grange et al., 2018) to correct for the role of meteorology in driving the 2015-2019 ozone trend in the SMA, and show that meteorology-corrected ozone is highest in May-June and increases the fastest in April-May, while showing no significant trend in July-August. We find that the GEOS-Chem chemical transport model can successfully capture the magnitude and trends of ozone concentrations, including their seasonality, and we use the model to quantify the importance of domestic and different background contributions in driving elevated ozone and its increase over South Korea.

Deleted: metric, since the upper tail of ozone concentrations are most harmful to human and ecosystem health. The 90th percentile is

Deleted: common

Deleted: in

Deleted: ozone pollution statistics

Formatted: Superscript

Deleted:

2 AirKorea data and trends, 2015-2019

90 We use ozone and NO₂ concentrations measured hourly by the AirKorea national air quality network of the South Korea Ministry of Environment (<http://www.airkorea.or.kr/web>). There are 255 sites in South Korea covering the 2015 to 2019 period including 79 sites in the SMA defined here as the rectilinear domain (126.7°E–127.3°E, 37.2°N–37.8°N). Figure 1 shows the maximum monthly 90th percentile MDA8 ozone at the ensemble of AirKorea sites for each year from 2015 to 2019. Ozone rises steadily over that period except for a dip in 2018, reaching 96 ppb in 2019 averaged across all AirKorea sites. High values are spread throughout South Korea and no site meets the 60 ppb air quality standard. Also shown is the monthly timeseries of ozone for the SMA. Ozone levels are similar to the rest of South Korea though do not show the 2018 dip. The seasonal maximum is in May-August depending on the year.

100 Figure 2 shows the annual mean 24-hr average NO₂ concentration at the ensemble of AirKorea sites. Concentrations of NO₂ in the SMA are generally 10 ppb larger than averaged across South Korea. NO₂ concentrations peak in winter and are minimum in summer, as observed elsewhere in East Asia and mostly driven by longer NO_x lifetime and reduced vertical mixing in winter (Lamsal et al., 2010; Shah et al., 2019; Lin et al., 2019; Kim et al., 2020). There is a decreasing trend over the 2015-2019 period as previously reported by Seo et al. (2021), more so in summer than in winter.

105

3 Meteorological Correction of 2015-2019 trends in the SMA

The 2015-2019 trends in ozone and NO₂ concentrations from Figures 1 and 2 could reflect emission trends but also meteorological variability. Here we use a random forest (RF) non-parametric statistical model (Breiman, 2001; Tong

Commented [JDJ1]: Not sure what you're doing here with section numbering?

Commented [CN2R1]: Issue with word autocorrect. Fixed.

Deleted: ¶

Formatted: Heading 1, Tab stops: 0.33", Left

et al., 2003) to isolate and remove the effect of meteorological variability for the 79 AirKorea sites in the SMA (Figure 3). RF is a supervised ensemble machine learning method, where many individual uncorrelated decision trees are fit to the training data to predict an output value, with the average value taken as best estimate (Breiman, 2001). The RF model was constructed using R "normalweatherr" packages (<https://github.com/skgrange/normalweatherr>; Grange et al., 2018). Hourly meteorological data are from two sites operated by the Korean Meteorological Administration (KMA) within the SMA (<https://data.kma.go.kr/data/grnd>). The RF model is trained to predict the hourly ozone and NO₂ concentrations averaged across the 79 AirKorea sites using the meteorological data averaged for the 2 KMA sites as well as time of day, day of year, and a long-term linear trend (Unix time stamp). Explanatory variables for the RF algorithm are listed in Table 1. Training of the RF model was conducted on 70% of the input data and the other 30% were withheld as testing data. The number of variables used to grow a tree was set to three, the minimum node-size was five, and the number of trees within a forest was set to 300. Once trained, the RF model is then used to predict ozone and NO₂ concentrations based on randomly sampled meteorological data and predictions are aggregated as described below.

Formatted: Font: 10 pt, Not Italic

Formatted: Font: 10 pt, Not Italic

Formatted: Font: 10 pt, Not Italic

Formatted: Font: 10 pt, Not Italic

Figure 4 compares observed and predicted hourly concentrations of ozone and NO₂ for the data withheld from training. The RF model shows a strong predictive ability ($R = 0.93$ for ozone, 0.90 for NO₂) with negligible mean bias (0.23 ppb for ozone, 0.01 ppb for NO₂), and root-mean-square errors (RMSE) of 5.9 ppb for ozone and 6.2 ppb for NO₂. The model has difficulty in capturing the tails of the distribution, which is a well-recognized problem in RF algorithms (Zhang and Lu, 2012; Pendergrass et al, 2021).

Formatted: Font: Italic

Deleted: NO₂

The top predictors in the RF fit for ozone are temperature, day of year, relative humidity, hour of day, and wind speed, in that order, consistent with previous studies for urban areas (Sillman and Samson, 1995; Jacob and Winner, 2009; K. Li et al., 2020). The top predictors for NO₂ are wind speed, day of year, temperature, hour of day, and surface pressure, again consistent with previous studies (Liu et al., 2020; Richmond-Bryant et al., 2018).

We use the RF model to remove the effect of meteorological variability in driving the 2015-2019 ozone and NO₂ trends by following the technique outlined in Vu et al. (2019). Meteorological variables for a specific hour and date in the input dataset are replaced by randomly selecting weather data over the entire study period (2015-2019) at that hour of day but for different day of year within a 4-week period (2 weeks before to 2 weeks after the selected date). This process is repeated 1000 times, and the resulting 1000 RF predictions of ozone and NO₂ for that hour and date are then averaged to produce meteorology-corrected concentrations from which we recalculate MDA8 ozone and 24-h averaged NO₂ to infer 2015-2019 emission-driven trends.

4 Emissions-driven trends in ozone and NO₂ concentrations in the SMA, 2015-2019

Figure 5 shows the observed and meteorology-corrected trends of monthly 90th percentile MDA8 ozone concentrations in the SMA from 2015 to 2019. The observations show peak increase in May but highly variable trend from month to month driven in part by interannual meteorological variability. The meteorology-corrected data show a much smoother behavior with a broad springtime (March-May) maximum in the increasing ozone trend, and a decreasing trend in August. Meteorology-corrected ozone is highest in May-June for all years. The 2015-2019

trend in meteorology-corrected 90th percentile ozone is 0.7, 1.4, and 0.4 ppb a⁻¹ for winter, spring, and autumn. The overall trend for summer is not statistically significant, but the trend for June alone is 0.9 ppb a⁻¹. This seasonality of ozone trends in the SMA from 2015 to 2019 is consistent with the 2000-2014 results of Jung et al. (2018), who reported a maximum springtime ozone increase in South Korea and an advancement of the ozone season by 2.1 days per year. Similar seasonality in the ozone trend has been reported for the North China Plain (K. Li et al., 2021), showing a twofold increase in May ozone exceedances above the 75 ppb standard from 2014 to 2019. Ozone production is most likely to be NO_x-limited in summer ~~when solar radiation and biogenic emissions are highest, and~~ and VOC-limited in spring and fall (Jacob et al., 1995), thus the seasonality of the trend is consistent with VOC-limited conditions.

Deleted: due to higher

Figure 6 is same as Figure 5 but for 24-h average NO₂ concentrations. The observations show a >2% a⁻¹ decrease in all months except November-February. The meteorology-corrected data show a consistent 5.6% a⁻¹ decrease in March-October, or 22% over the four years, and a consistent but weaker 1.5% a⁻¹ decrease in November-February. This is consistent with findings from Bae et al. (2020), who reported a 4.4% a⁻¹ decline of annual mean NO₂ in the SMA for 2015-2018 using surface and satellite observations. Declining NO₂ in the SMA can be attributed to policies to decrease vehicular NO_x emissions (Kim and Lee, 2018). The weaker decline in winter is consistent with findings from Seo et al. (2021), who found that surface NO₂ concentrations in the SMA in 2015-2019 declined by 5.3% a⁻¹ during morning commute time for the ozone season, but only by 2.6% a⁻¹ for the non-ozone season. The weaker response of NO₂ to reduced NO_x emissions in winter could be due to ozone titration by emitted NO, which would take place most systematically at night but also extend to daytime if the ozone supply is weak. We find that in November-February the decline of NO₂ during midday (11:00-15:00 LT) is 2.4% a⁻¹, greater than twice that at night (23:00-03:00 LT), consistent with ozone titration. For the GEOS-Chem simulations in the following Sections we will assume a 22% decrease of NO_x emissions from 2015 to 2019.

Deleted: r

5 GEOS-Chem simulation

We use the GEOS-Chem chemical transport model version 13.3.4 (<http://geos-chem.org>) to interpret the observed ozone and its 2015-2019 trend in the SMA and more broadly in South Korea, including influences from China and the global background. GEOS-Chem has been applied previously in South Korea to investigate ozone production efficiency (Oak et al., 2019), the factors determining ozone seasonality (Lee and Park, 2022), and the photochemical environment for ozone production (Yang et al., 2022). Park et al. (2021) previously found that GEOS-Chem version 12.7.2 underestimated free tropospheric ozone over South Korea by 20-30 ppb. Addition of detailed aromatic chemistry in version 13.3.4 (Bates et al., 2021) was subsequently found to increase net ozone production over South Korea by 37% (Oak et al., 2019). Here we also add particulate nitrate photolysis and suppression of sea salt aerosol debromination to the model following Shah et al. (2022), and as we will see this largely corrects the remaining model ozone bias over East Asia.

185 We use a nested-grid version of GEOS-Chem driven by MERRA-2 assimilated meteorological data with a horizontal resolution of $0.5 \times 0.625^\circ$ over East Asia (25° - 50° N, 105° - 140° E; domain of Figure 7). Chemical boundary conditions at the edges of the nested domain are updated every 3 hours from a global simulation with $4^\circ \times 5^\circ$ resolution. We conduct a full-year simulation for 2016 with six months of initialization. Global anthropogenic emissions are from the Community Emissions Data System (CEDS) global inventory (Hoesly et al., 2018) and are superseded with regional emission inventories for South Korea (KORUSv5, <http://aisl.konkuk.ac.kr>) and China (Multi-resolution Emission Inventory, Zheng et al., 2018). Natural emissions include NO_x from lightning (Murray et al., 2012) and soil (Hudman et al., 2012), MEGANv2 biogenic volatile organic compounds (VOCs) (Guenther et al., 2012), dust (Meng et al., 2021), and sea salt (Jaeglé et al., 2011). Open-fire emissions are from the Global Fire Emissions Database version 4 (GFED4; van der Werf et al., 2017).

195 Ships are a relatively large source of NO_x in East Asia. The standard GEOS-Chem model includes pre-processing of ship emissions with the PARAMeterization of emitted NO_x (PARANOX) algorithm (Vinken et al., 2011) to account for the non-linear chemistry occurring during the dispersion of ship exhaust plumes. PARANOX is a plume-in-grid formulation where ship emissions are aged chemically for 5 hours before being released into the model grid. This greatly reduces the ozone yield from ship NO_x emissions, which would otherwise be diluted by the model in a relatively clean environment where the ozone production efficiency is very high. PARANOX was intended for global model simulations with grid resolution of hundreds of km (Holmes et al., 2014) and its application to higher-resolution simulations is questionable, particularly over East Asia where the maritime environment is highly polluted (Cuesta et al., 2018; Peterson et al., 2019; Jung et al., 2022). Here we disable PARANOX for the nested simulation and find that this increases ozone over the Yellow Sea in May by 1 ppb on average.

205 We evaluate our GEOS-Chem simulation for 2016 with MDA8 ozone observations from the AirKorea network in South Korea and the Ministry of Energy and Environment (MEE) monitoring network in China (<http://data.epmap.org/page/index>). Observations of seasonal mean 90th percentile MDA8 ozone overlaid on our GEOS-Chem simulation are shown in Figure 7. There is good agreement between GEOS-Chem and observations in all seasons, with a spatial correlation coefficient $R > 0.7$ and a mean bias < 4 ppb.

210 We evaluated GEOS-Chem's ability to reproduce the seasonal cycle of ozone in the three megacity clusters of Seoul Metropolitan Area (SMA), Beijing-Tianjin-Hebei (BTH), and Yangtze River Delta (YRD). Figure 8 shows the monthly 90th percentile MDA8 ozone for 2016 averaged over all network sites in each cluster. The simulated seasonal cycle is consistent with observations ($R > 0.95$ and mean bias < 6.0 ppb).

215 May is of particular interest in the SMA because this is when ozone and its increasing trend are highest in the meteorology-corrected data (Figure 5). Previous model comparisons to extensive vertical profiles taken during the KORUS-AQ aircraft campaign over South Korea in May-June 2016 showed large underestimates, with GEOS-Chem version 12.7.2 being too low by 20-30 ppb (Park et al., 2021). The model updates described above largely

Deleted: against

220 correct this underestimate (Yang et al., 2022). Figure 9 compares our simulated GEOS-Chem ozone profile to the
mean of 15 ozonesonde observations over Olympic Park in Seoul taken during the KORUS-AQ campaign on DC-8
flight observation days (15 profiles in total). Our simulation has a low bias of only 5.4 ppb in the free troposphere.

225 To investigate and diagnose the ability of GEOS-Chem to reproduce the observed 2015-2019 ozone trend in the
SMA, we performed simulations with 2016 meteorology (January 2016- December 2016) and perturbed emissions
in China and South Korea for 2015 and 2019 to simulate the 2015-2019 trend. The sensitivity simulations used 6
months of initialization. China emissions in 2015 are from MEIC (Zheng et al., 2018) but MEIC does not extend
beyond 2017. Following K. Li et al. (2021), we scaled 2017 MEIC emissions to 2019 based on observed MEE
network trends. Overall, emissions in China declined from 2015 to 2019 by 16% for NO_x, 50% for SO₂, 23%, for
CO, and 32% for primary PM_{2.5}, with flat VOC emissions (K. Li et al., 2021). Anthropogenic emissions for South
230 Korea in 2015 are taken from the KORUSv5 inventory (<http://aisl.konkuk.ac.kr>). For 2019 we decrease NO_x
emissions in South Korea by 22% (Section 4) and apply no other changes to South Korea emissions, including
VOCs for which emission trends are not clear as mentioned in the Introduction. We also do not apply trends to ship
emissions.

235 Figure 10 shows the emission-driven trends of 90th percentile MDA8 ozone from 2015 to 2019 in the SMA for both
meteorology-corrected observations (data from Figure 5) and GEOS-Chem in individual months. The model trend is
obtained by subtraction of results from simulations with 2015 and 2019 emissions, both using the same 2016
meteorology. GEOS-Chem reproduces the general magnitude and seasonality of the observed trend. It reproduces in
particular the April-May maximum in the trend.

240

6 Attribution of ozone and its 2015-2019 trend over South Korea

We exploit the success of GEOS-Chem in simulating ozone over East Asia and its trend over the SMA to investigate
the causes. We focus on May, where both ozone concentrations and its increasing trend in the meteorology-
245 corrected data for the SMA are the highest. In addition to the baseline simulation described in Section 5, we also
conduct sensitivity simulations for both emission years to isolate the effects of anthropogenic emissions from South
Korea, China, ships, and East Asia as a whole by zeroing the corresponding emissions including NO_x, VOCs, CO,
and PM_{2.5}. The same global boundary conditions described above are used for each of these cases, with 6 months of
initialization.

250

Figure 11 shows the distribution of simulated 90th percentile MDA8 ozone for May using 2015 emissions, the
difference when using 2019 emissions, and the contributions from South Korea and China as determined from the
sensitivity simulations with the corresponding emissions shut off. 2015 values in the baseline simulation average
85.8 ppb in the SMA and 90.1 ppb for all of South Korea, and the 2019-2015 difference averages +6.2 ppb for the
255 SMA while southern parts of the country show decreases. Zeroing out South Korea emissions has remarkably little

effect on SMA concentrations, which remain at 84.1 ppb for 2015, though the 2015-2019 trend is now near zero. Zeroing out China emissions decreases SMA ozone concentrations to 79.8 ppb but the 2015-2019 increase remains at 5.6 ppb. We conclude that the 2015-2019 ozone increase in the SMA can be attributed to the decrease of domestic NO_x emissions under VOC-limited conditions. When emissions from China are zeroed out, we find a 6 ppb ozone decrease in the SMA and a 8 ppb decrease in South Korea as a whole ~~compared to the baseline simulation~~. The 2015-2019 ozone trend over the SMA is affected by less than 1 ppb, confirming that this trend is mainly driven by domestic emission changes.

Deleted: , representing a significant but relatively modest improvement

A notable result is that ozone levels over South Korea remain very high at about 80 ppb even when emissions from either South Korea or China are totally shut off. Lee and Park (2022) previously found with GEOS-Chem that surface ozone over South Korea in April hardly changes when domestic emissions are shut off, and here we find that zeroing China emissions also has only a modest effect over South Korea. This resilience is indicative of a major contribution to ozone pollution from the northern mid-latitudes background external to East Asia.

Figure 12 further explores the role of this East Asia background in a simulation with anthropogenic emissions shut off throughout the nested model domain. The 90th percentile MDA8 ozone drops to 55 ppb in South Korea, meeting the 60 ppb standard but still extremely high, and indicating that even small anthropogenic emissions would cause ozone to rise above the standard. This high East Asia background affects northern China even more. Lam and Cheung (2022) previously found with GEOS-Chem that the mean MDA8 background ozone over China in April is 53 ppb, and we find here that the 90th percentile over northern China reaches 70 ppb. This East Asia background ozone is much higher than the corresponding North American background of 20-40 ppb previously reported in studies of US ozone pollution (Fiore et al., 2003; Zhang et al., 2011; Emery et al., 2012; Jaffe et al. 2018). Such a high East Asia background is reflected in the observation of 75 ppb ozone in the free troposphere (Figure 9) while comparable ozonesonde observations over the western US in spring show mean values of 60 ppb (Zhang et al., 2014). Satellite observations of free tropospheric ozone also show particularly high values over East Asia (Hu et al., 2017; Gaudel et al., 2018). High free tropospheric ozone over East Asia in spring could reflect regional downwelling from the stratosphere associated with cyclogenesis (Hwang et al., 2007). It could also reflect the observed rise of free tropospheric ozone at northern mid-latitudes and particularly over East Asia in recent decades (Gaudel et al., 2018; Lee et al., 2021; Wang et al., 2022) which could possibly be due to increasing emissions in India and the Middle East (Anwar et al., 2021; Ding et al., 2022; Anenberg et al., 2022). We find from analysis of sonde observations ~~at Pohang, Hong Kong, and Tateno (https://woudc.org/data/dataset_info.php?id=ozonesond https://woudc.org/data/dataset_info.php?id=ozonesonde) no~~ significant trend in free tropospheric ozone over South Korea during 2015-2019, meaning that the background is not responsible for the observed increase in surface ozone over that period. Domestic emissions are likely responsible, as discussed above.

Deleted: at Pohang, Hong Kong, and Tano

Additional panels in Figure 12 show the enhancement of ozone above the East Asian background due to emissions from the Yellow Sea (ships), South Korea, and China. Emissions from ships in the Yellow Sea enhance 90th percentile MDA8 ozone over South Korea by only a few ppb, although they can drive ozone concentrations over the

ocean in excess of 90 ppb. Despite ship traffic in the Yellow Sea being intense, the NO_x emissions are still small relative to continental emissions. Emissions from South Korea alone push ozone to almost 80 ppb over South Korea, with even larger increases over the surrounding oceans reflecting VOC-limited conditions over land. In this way, emissions in South Korea push ozone in the Shandong peninsula in China to over 80 ppb. Emissions in China have a comparable effect on ozone over South Korea as domestic emissions.

7 Conclusions

We examined the factors controlling the high and increasing surface ozone concentrations over South Korea and particularly in the Seoul Metropolitan Area (SMA). Ozone in South Korea has risen steadily over the past two decades and is everywhere far in excess of the 60 ppb air quality standard set by the South Korean government in 2015. Improved understanding of the causes of elevated ozone in South Korea is critical for developing effective emission control strategies.

~~We find a continuation of the multidecadal increase of surface ozone in South Korea.~~ Analysis of 2015-2019 data from the AirKorea network of air quality monitoring sites shows elevated ozone throughout South Korea, with 90th percentile ozone averaged across all sites exceeding 75 ppb every year and increasing over the period. NO₂ concentrations also measured at AirKorea sites are typically >10 ppb higher in the SMA than elsewhere, with maximum concentrations in winter and a decrease over the 2015-2019 period.

We used a random forest (RF) non-parametric statistical model to isolate and remove the effect of meteorological variability on 2015-2019 ozone and NO₂ trends in the SMA. Meteorology-corrected ozone is highest in May-June for all years and increases at the fastest rate of 1.5 ppb a⁻¹ in April-May. Meteorology-corrected NO₂ is highest during November-March and lowest in July-August. During the ozone season of March-October, NO₂ shows a consistent decline of 5.6% a⁻¹ over the 2015-2019 period, whereas in winter the decline is lower at 1.3% a⁻¹. The March-October trend in NO₂ concentrations suggests that NO_x emissions declined by 22% from 2015 to 2019.

We used the GEOS-Chem chemical transport model to interpret the elevated ozone and its 2015-2019 trend in the SMA and more broadly in South Korea, including influences from China and the global background. We improved on previous versions of the model, which substantially underestimated tropospheric ozone over South Korea, through the addition of detailed aromatic chemistry in version 13.3.4 (Bates et al., 2021), the removal of sea salt aerosol debromination, and the addition of particulate nitrate photolysis (Shah et al., 2022). The resulting model can reproduce the seasonality and spatial distribution of surface ozone in South Korea and China without significant bias. It reproduces the high free tropospheric ozone concentrations observed over Seoul during the KORUS-AQ campaign in May-June 2016 (75 ± 7 ppb) with only a 5 ppb low bias. Implementing in the model the 2015-2019 emission decreases in Korea and China reproduces the observed seasonality and magnitude of the meteorology-corrected ozone trend over the SMA.

Formatted: Body Text, Indent: Left: 0.22", Right: 0.1"

Deleted: ¶

Deleted: Here w

330 ~~We used~~ GEOS-Chem sensitivity simulations for emission years 2015 and 2019 to better understand the factors contributing to elevated ozone in the SMA and South Korea, focusing on May when meteorology-corrected ozone and its increase are the highest. We find that the 2015-2019 ozone increase in the SMA can be explained by the 22% decrease of ~~surface NO_x concentrations, which act as a proxy for declining NO_x emissions, reflecting the VOC-~~ limited conditions for ozone production. We also find that emissions in China and South Korea contribute equally to elevated ozone over South Korea, while ships only contribute a small amount. VOC emission reductions would be expected to decrease ozone in South Korea, but we find that concentrations remain over 80 ppb even if all anthropogenic emissions from South Korea or from China are zeroed out. The East Asia background, defined by zeroing out all anthropogenic emissions over East Asia, is very high at 55 ppb, implying that the 60 ppb air quality standard in South Korea is not achievable without addressing the origin of this elevated background.

- Deleted: We
- Deleted: went on
- Deleted: to use
- Deleted: domestic
- Deleted: x
- Deleted: emissions in South Korea
- Formatted: Subscript

340 **Code Availability**

The code used in this work is available upon request.

Data Availability

345 Ground-based measurements from the AirKorea national air quality network of the South Korea Ministry of Environment are available at <http://www.airkorea.or.kr/web>. Ozonesonde data from the KORUS-AQ data archive are available at <https://www-air.larc.nasa.gov> (KORUS-AQ Science Team, 2019). Meteorological data from the Korean Meteorological Administration (KMA) are found at <https://data.kma.go.kr/data/grnd>.

350 **Author Contribution**

The original draft preparation was done by NKC, with review and editing by DJJ, LHY, SZ, VS, SKG, RMY, SK, and HL. DJJ contributed to project conceptualization. Modeling was done by NKC, with additional support from LHY, SZ, VS, and RMY. The formal analysis was conducted by NKC with additional support from DJJ, LHY, SZ, VS, SKG, and SK.

355 **Acknowledgments**

This work was funded by the Samsung Advanced Institute of Technology and the Harvard–NUIST Joint Laboratory for Air Quality and Climate (JLAQC). We thank Zongbo Shi and Tuan Vu for their helpful insight on removing the effect of meteorology on pollutant trends.

360 **References**

365 Anenberg, Susan C, Arash Mohegh, Daniel L Goldberg, Gaige H Kerr, Michael Brauer, Katrin Burkart, Perry Hystad, Andrew Larkin, Sarah Wozniak, and Lok Lamsal. 2022. "Long-Term Trends in Urban NO₂ Concentrations and Associated Paediatric Asthma Incidence: Estimates from Global Datasets." *The Lancet Planetary Health* 6 (1): e49–58. [https://doi.org/10.1016/S2542-5196\(21\)00255-2](https://doi.org/10.1016/S2542-5196(21)00255-2).

Anwar, Muhammad Naveed, Muneeba Shabbir, Eza Tahir, Mahnoor Iftikhar, Hira Saif, Ajwa Tahir, Malik Ashir Murtaza, et al. 2021. "Emerging Challenges of Air Pollution and Particulate Matter in China, India, and Pakistan and

- Mitigating Solutions.” *Journal of Hazardous Materials* 416 (August): 125851.
 https://doi.org/10.1016/j.jhazmat.2021.125851.
- 370 Bae, Minah, Byeong-Uk Kim, Hyun Cheol Kim, Jhoon Kim, and Soontae Kim. 2021. “Role of Emissions and Meteorology in the Recent PM_{2.5} Changes in China and South Korea from 2015 to 2018.” *Environmental Pollution* 270 (February): 116233. https://doi.org/10.1016/j.envpol.2020.116233.
- 375 Bates, Kelvin H., Daniel J. Jacob, Ke Li, Peter D. Ivatt, Mat J. Evans, Yingying Yan, and Jintai Lin. 2021. “Development and Evaluation of a New Compact Mechanism for Aromatic Oxidation in Atmospheric Models.” *Atmospheric Chemistry and Physics* 21 (24): 18351–74.
- Bauwens, M. *et al.*, 2022. “Spaceborne Evidence for Significant Anthropogenic VOC Trends in Asian Cities over 2005– 2019.” *Environ. Res. Lett.* 17 015008 https://doi.org/10.5194/acp-21-18351-2021.
- Breiman, Leo. 2001. “Random Forests.” *Machine Learning* 45 (1): 5–32. https://doi.org/10.1023/A:1010933404324.
- 380 Carslaw, David C., Karl Ropkins, and Margaret C. Bell. 2006. “Change-Point Detection of Gaseous and Particulate Traffic-Related Pollutants at a Roadside Location.” *Environmental Science & Technology* 40 (22): 6912–18. https://doi.org/10.1021/es060543u.
- Crawford, James H., Joon-Young Ahn, Jassim Al-Saadi, Limseok Chang, Louisa K. Emmons, Jhoon Kim, Gangwoong Lee, *et al.* 2021. “The Korea–United States Air Quality (KORUS-AQ) Field Study.” *Elementa: Science of the Anthropocene* 9 (1): 00163. https://doi.org/10.1525/elementa.2020.00163.
- 385 Cuesta, Juan, Yugo Kanaya, Masayuki Takigawa, Gaëlle Dufour, Maxim Eremanenko, Gilles Foret, Kazuyuki Miyazaki, and Matthias Beekmann. 2018. “Transboundary Ozone Pollution across East Asia: Daily Evolution and Photochemical Production Analysed by IASI + GOME2 Multispectral Satellite Observations and Models.” *Atmospheric Chemistry and Physics* 18 (13): 9499–9525. https://doi.org/10.5194/acp-18-9499-2018.
- 390 Ding, Jieying, Ronald van der A, Bas Mijling, Jos de Laat, Henk Eskes, and K. Folkert Boersma. 2022. “NO_x Emissions in India Derived from OMI Satellite Observations.” *Atmospheric Environment: X* 14 (April): 100174. https://doi.org/10.1016/j.aeaoa.2022.100174.
- Emery, Christopher, Jaegun Jung, Nicole Downey, Jeremiah Johnson, Michele Jimenez, Greg Yarwood, and Ralph Morris. 2012. “Regional and Global Modeling Estimates of Policy Relevant Background Ozone over the United States.” *Atmospheric Environment* 47 (February): 206–17. https://doi.org/10.1016/j.atmosenv.2011.11.012.
- 395 Fiore, A., Daniel J. Jacob, Jennifer A. Logan, and Jeffrey H. Yin. 1998. “Long-Term Trends in Ground Level Ozone over the Contiguous United States, 1980–1995.” *Journal of Geophysical Research: Atmospheres* 103 (D1): 1471–80. https://doi.org/10.1029/97JD03036.
- 400 Fiore, A., D. J. Jacob, H. Liu, R. M. Yantosca, T. D. Fairlie, and Q. Li. 2003. “Variability in Surface Ozone Background over the United States: Implications for Air Quality Policy.” *Journal of Geophysical Research: Atmospheres* 108 (D24). https://doi.org/10.1029/2003JD003855.
- Gaubert, Benjamin, Louisa K. Emmons, Kevin Raeder, Simone Tilmes, Kazuyuki Miyazaki, Avelino F. Arellano Jr., Nellie Elguindi, *et al.* 2020. “Correcting Model Biases of CO in East Asia: Impact on Oxidant Distributions during KORUS-AQ.” *Atmospheric Chemistry and Physics* 20 (23): 14617–47. https://doi.org/10.5194/acp-20-14617-2020.
- 405 Gaudel, A., O. R. Cooper, G. Ancellet, B. Barret, A. Boynard, J. P. Burrows, C. Clerbaux, *et al.* 2018. “Tropospheric Ozone Assessment Report: Present-Day Distribution and Trends of Tropospheric Ozone Relevant to Climate and Global Atmospheric Chemistry Model Evaluation.” Edited by Detlev Helmig and Alastair Lewis. *Elementa: Science of the Anthropocene* 6 (May): 39. https://doi.org/10.1525/elementa.291.
- 410 Grange, Stuart K., David C. Carslaw, Alastair C. Lewis, Eirini Boleti, and Christoph Hueglin. 2018. “Random Forest Meteorological Normalisation Models for Swiss PM₁₀ Trend Analysis.” *Atmospheric Chemistry and Physics* 18 (9): 6223–39. https://doi.org/10.5194/acp-18-6223-2018.
- Guenther, A. B., X. Jiang, C. L. Heald, T. Sakulyanontvittaya, T. Duhl, L. K. Emmons, and X. Wang. 2012. “The Model of Emissions of Gases and Aerosols from Nature Version 2.1 (MEGAN2.1): An Extended and Updated Framework for Modeling Biogenic Emissions.” *Geoscientific Model Development* 5 (6): 1471–92. https://doi.org/10.5194/gmd-5-1471-2012.
- Han, Kyung M. 2019. “Temporal Analysis of OMI-Observed Tropospheric NO₂ Columns over East Asia during 2006– 2015.” *Atmosphere* 10 (11): 658. https://doi.org/10.3390/atmos10110658.
- 415 He, Jianjun, Sunling Gong, Ye Yu, Lijuan Yu, Lin Wu, Hongjun Mao, Congbo Song, *et al.* 2017. “Air Pollution Characteristics and Their Relation to Meteorological Conditions during 2014–2015 in Major Chinese Cities.” *Environmental Pollution* 223 (April): 484–96. https://doi.org/10.1016/j.envpol.2017.01.050.
- Hoesly, Rachel M., Steven J. Smith, Leyang Feng, Zbigniew Klimont, Greet Janssens-Maenhout, Tyler Pitkanen, Jonathan J. Seibert, *et al.* 2018. “Historical (1750–2014) Anthropogenic Emissions of Reactive Gases and Aerosols from the Community Emissions Data System (CEDS).” *Geoscientific Model Development* 11 (1): 369–408. https://doi.org/10.5194/gmd-11-369-2018.
- 420 Holmes, C. D., M. J. Prather, and G. C. M. Vinken. 2014. “The Climate Impact of Ship NO_x Emissions: An Improved Estimate Accounting for Plume Chemistry.” *Atmospheric Chemistry and Physics* 14 (13): 6801–12. https://doi.org/10.5194/acp-14-6801-2014.

Field Code Changed

- 425 Hu, Lu, Daniel Jacob, Xiong Liu, Yi Zhang, Lin Zhang, Patrick Kim, Melissa Sulprizio, and Robert Yantosca. 2017. "Global Budget of Tropospheric Ozone: Evaluating Recent Model Advances with Satellite (OMI), Aircraft (IAGOS), and Ozone-sonde Observations." *Atmospheric Environment* 167 (August). <https://doi.org/10.1016/j.atmosenv.2017.08.036>.
- 430 Hudman, R. C., N. E. Moore, A. K. Mebust, R. V. Martin, A. R. Russell, L. C. Valin, and R. C. Cohen. 2012. "Steps towards a Mechanistic Model of Global Soil Nitric Oxide Emissions: Implementation and Space Based-Constraints." *Atmospheric Chemistry and Physics* 12 (16): 7779–95. <https://doi.org/10.5194/acp-12-7779-2012>.
- Hwang, Seung-Hyun, Jhoon Kim, and Gwang-Rae Cho. 2007. "Observation of Secondary Ozone Peaks near the Tropopause over the Korean Peninsula Associated with Stratosphere-Troposphere Exchange." *Journal of Geophysical Research: Atmospheres* 112 (D16). <https://doi.org/10.1029/2006JD007978>.
- 435 Jacob, Daniel J., Larry W. Horowitz, J. William Munger, Brian G. Heikes, Russell R. Dickerson, Richard S. Artz, and William C. Keene. 1995. "Seasonal Transition from NO_x- to Hydrocarbon-Limited Conditions for Ozone Production over the Eastern United States in September." *Journal of Geophysical Research: Atmospheres* 100 (D5): 9315–24. <https://doi.org/10.1029/94JD03125>.
- Jacob, Daniel J., and Darrel A. Winner. 2009. "Effect of Climate Change on Air Quality." *Atmospheric Environment*. <https://doi.org/10.1016/j.atmosenv.2008.09.051>.
- 440 Jaeglé, L., P. K. Quinn, T. S. Bates, B. Alexander, and J.-T. Lin. 2011. "Global Distribution of Sea Salt Aerosols: New Constraints from in Situ and Remote Sensing Observations." *Atmospheric Chemistry and Physics* 11 (7): 3137–57. <https://doi.org/10.5194/acp-11-3137-2011>.
- 445 Jaffe, Daniel A., Owen R. Cooper, Arlene M. Fiore, Barron H. Henderson, Gail S. Tonnesen, Armistead G. Russell, Daven K. Henze, Andrew O. Langford, Meiyun Lin, and Tom Moore. 2018. "Scientific Assessment of Background Ozone over the U.S.: Implications for Air Quality Management." Edited by Detlev Helmig and Allen Goldstein. *Elementa: Science of the Anthropocene* 6 (July): 56. <https://doi.org/10.1525/elementa.309>.
- Jung, Hyun-Chae, Byung-Kwon Moon, and Jieun Wie. 2018. "Seasonal Changes in Surface Ozone over South Korea." *Heliyon* 4 (1). <https://doi.org/10.1016/j.heliyon.2018.e00515>.
- 450 Jung, Jia, Yunsoo Choi, Amir H. Souri, Seyedali Mousavinezhad, Alqamah Sayeed, and Kyunghwa Lee. 2022. "The Impact of Springtime-Transported Air Pollutants on Local Air Quality With Satellite-Constrained NO_x Emission Adjustments Over East Asia." *Journal of Geophysical Research: Atmospheres* 127 (5): e2021JD035251. <https://doi.org/10.1029/2021JD035251>.
- Kang, Yoojin, Hyunyoung Choi, Jungho Im, Seohui Park, Minso Shin, Chang-Keun Song, and Sangmin Kim. 2021. "Estimation of Surface-Level NO₂ and O₃ Concentrations Using TROPOMI Data and Machine Learning over East Asia." *Environmental Pollution* 288 (November): 117711. <https://doi.org/10.1016/j.envpol.2021.117711>.
- Kim, Heejeong, Junsu Gil, Meehye Lee, Jinsang Jung, Andrew Whitehill, James Szykman, Gangwoong Lee, et al. 2020. "Factors Controlling Surface Ozone in the Seoul Metropolitan Area during the KORUS-AQ Campaign." *Elementa (Washington, D.C.)* 8 (46): 10.1525/elementa.444. <https://doi.org/10.1525/elementa.444>.
- 460 Kim, Hyun Cheol, Soontae Kim, Sang-Hyun Lee, Byeong-Uk Kim, and Pius Lee. 2020. "Fine-Scale Columnar and Surface NO_x Concentrations over South Korea: Comparison of Surface Monitors, TROPOMI, CMAQ and CAPSS Inventory." *Atmosphere* 11 (1): 101. <https://doi.org/10.3390/atmos11010101>.
- Kim, Jeonghwan, Jimin Lee, Jinseok Han, Jinsoo Choi, Dai-Gon Kim, Jinsoo Park, and Gangwoong Lee. 2021. "Long-Term Assessment of Ozone Nonattainment Changes in South Korea Compared to US, and EU Ozone Guidelines." *Asian Journal of Atmospheric Environment* 15 (December): 20–32. <https://doi.org/10.5572/ajae.2021.098>.
- 465 Kim, Yong Pyo, and Gangwoong Lee. 2018. "Trend of Air Quality in Seoul: Policy and Science." *Aerosol and Air Quality Research* 18 (9): 2141–56. <https://doi.org/10.4209/aaqr.2018.03.0081>.
- Kuhn, Max. n.d. "Conventions in R Data Splitting and Estimating Performance Data Pre-Processing Over-Fitting and Resampling Training and Tuning Tree Models Training and Tuning A Support Vector Machine Comparing Models Parallel Processing," 63.
- 470 Lam, Yun Fat, and Hung Ming Cheung. 2022. "Investigation of Policy Relevant Background (PRB) Ozone in East Asia." *Atmosphere* 13 (5): 723. <https://doi.org/10.3390/atmos13050723>.
- Lamsal, L. N., R. V. Martin, A. van Donkelaar, E. A. Celarier, E. J. Bucsela, K. F. Boersma, R. Dirksen, C. Luo, and Y. Wang. 2010. "Indirect Validation of Tropospheric Nitrogen Dioxide Retrieved from the OMI Satellite Instrument: Insight into the Seasonal Variation of Nitrogen Oxides at Northern Midlatitudes." *Journal of Geophysical Research: Atmospheres* 115 (D5). <https://doi.org/10.1029/2009JD013351>.
- 475 Lee, Hyo-Jung, Lim-Seok Chang, Daniel A. Jaffe, Juseon Bak, Xiong Liu, Gonzalo González Abad, Hyun-Young Jo, Yu-Jin Jo, Jae-Bum Lee, and Cheol-Hee Kim. 2021. "Ozone Continues to Increase in East Asia Despite Decreasing NO₂: Causes and Abatements." *Remote Sensing* 13 (11): 2177. <https://doi.org/10.3390/rs13112177>.

- 480 Lee, Hyung-Min, and Rokjin J. Park. 2022. "Factors Determining the Seasonal Variation of Ozone Air Quality in South Korea: Regional Background versus Domestic Emission Contributions." *Environmental Pollution* 308 (September): 119645. <https://doi.org/10.1016/j.envpol.2022.119645>.
- Li, Jie, Wenyi Yang, Zifa Wang, Huansheng Chen, Bo Hu, Jianjun. Li, Yele. Sun, Pingqing Fu, and Yuqia Zhang. 2016. "Modeling Study of Surface Ozone Source-Receptor Relationships in East Asia." *Atmospheric Research* 167 (January): 77–88. <https://doi.org/10.1016/j.atmosres.2015.07.010>.
- 485 Li, Ke, Daniel J. Jacob, Hong Liao, Lu Shen, Qiang Zhang, and Kelvin H. Bates. 2019. "Anthropogenic Drivers of 2013–2017 Trends in Summer Surface Ozone in China." *Proceedings of the National Academy of Sciences* 116 (2): 422–27. <https://doi.org/10.1073/pnas.1812168116>.
- 490 Li, Ke, Daniel J. Jacob, Lu Shen, Xiao Lu, Isabelle De Smedt, and Hong Liao. 2020. "Increases in Surface Ozone Pollution in China from 2013 to 2019: Anthropogenic and Meteorological Influences." *Atmospheric Chemistry and Physics* 20 (19): 11423–33. <https://doi.org/10.5194/acp-20-11423-2020>.
- Li, Ke, Daniel J. Jacob, Hong Liao, Yulu Qiu, Lu Shen, Shixian Zhai, Kelvin H. Bates, et al. 2021. "Ozone Pollution in the North China Plain Spreading into the Late-Winter Haze Season." *Proceedings of the National Academy of Sciences* 118 (10): e2015797118. <https://doi.org/10.1073/pnas.2015797118>.
- 495 Lin, Chin-An, Yi-Chun Chen, Chian-Yi Liu, Wei-Ting Chen, John H. Seinfeld, and Charles C.-K. Chou. 2019. "Satellite-Derived Correlation of SO₂, NO₂, and Aerosol Optical Depth with Meteorological Conditions over East Asia from 2005 to 2015." *Remote Sensing* 11 (15): 1738. <https://doi.org/10.3390/rs11151738>.
- Liu, Yansui, Yang Zhou, and Jiaxin Lu. 2020. "Exploring the Relationship between Air Pollution and Meteorological Conditions in China under Environmental Governance." *Scientific Reports* 10 (1): 14518. <https://doi.org/10.1038/s41598-020-71338-7>.
- 500 Meng, Jun, Randall V. Martin, Paul Ginoux, Melanie Hammer, Melissa P. Sulprizio, David A. Ridley, and Aaron van Donkelaar. 2021. "Grid-Independent High-Resolution Dust Emissions (v1.0) for Chemical Transport Models: Application to GEOS-Chem (12.5.0)." *Geoscientific Model Development* 14 (7): 4249–60. <https://doi.org/10.5194/gmd-14-4249-2021>.
- 505 Miyazaki, K., T. Sekiya, D. Fu, K. W. Bowman, S. S. Kulawik, K. Sudo, T. Walker, et al. 2019. "Balance of Emission and Dynamical Controls on Ozone During the Korea-United States Air Quality Campaign From Multiconstituent Satellite Data Assimilation." *Journal of Geophysical Research: Atmospheres* 124 (1): 387–413. <https://doi.org/10.1029/2018JD028912>.
- MOE, Ministry of Environment, Korea (2016) *White Paper of the Environment*, Seoul.
- 510 Murray, Lee T., Daniel J. Jacob, Jennifer A. Logan, Rynda C. Hudman, and William J. Koshak. 2012. "Optimized Regional and Interannual Variability of Lightning in a Global Chemical Transport Model Constrained by LIS/OTD Satellite Data." *Journal of Geophysical Research: Atmospheres* 117 (D20). <https://doi.org/10.1029/2012JD017934>.
- NIER, National Institute of Environmental Research, Korea (2020). *2020 Annual Report*.
- Oak, Yujin J., Rokjin J. Park, Jason R. Schroeder, James H. Crawford, Donald R. Blake, Andrew J. Weinheimer, Jung-Hun Woo, et al. 2019. "Evaluation of Simulated O₃ Production Efficiency during the KORUS-AQ Campaign: Implications for Anthropogenic NO_x Emissions in Korea." Edited by Detlev Helmig and Jochen Stutz. *Elementa: Science of the Anthropocene* 7 (December): 56. <https://doi.org/10.1525/elementa.394>.
- Oh, In-Bo, Yoo-Keun Kim, Mi-Kyung Hwang, Cheol-Hee Kim, and Soon Kim. 2010. "Elevated Ozone Layers over the Seoul Metropolitan Region in Korea: Evidence for Long-Range Ozone Transport from Eastern China and Its Contribution to Surface Concentrations." *Journal of Applied Meteorology and Climatology* 49 (February): 203–20. <https://doi.org/10.1175/2009JAMC2213.1>.
- 520 Park, Rokjin J., Yujin J. Oak, Louisa K. Emmons, Cheol-Hee Kim, Gabriele G. Pfister, Gregory R. Carmichael, Pablo E. Saide, et al. 2021. "Multi-Model Intercomparisons of Air Quality Simulations for the KORUS-AQ Campaign." *Elementa: Science of the Anthropocene* 9 (1): 00139. <https://doi.org/10.1525/elementa.2021.00139>.
- 525 Pendergrass, Drew C., Shixian Zhai, Jhoon Kim, Ja-Ho Koo, Seoyoung Lee, Minah Bae, Soontae Kim, Hong Liao, and Daniel J. Jacob. 2022. "Continuous Mapping of Fine Particulate Matter (PM_{2.5}) Air Quality in East Asia at Daily 6x6 km² Resolution by Application of a Random Forest Algorithm to 2011–2019 GOCI Geostationary Satellite Data." *Atmospheric Measurement Techniques* 15 (4): 1075–91. <https://doi.org/10.5194/amt-15-1075-2022>.
- Peterson, David A., Edward J. Hyer, Sang-Ok Han, James H. Crawford, Rokjin J. Park, Robert Holz, Ralph E. Kuehn, et al. 2019. "Meteorology Influencing Springtime Air Quality, Pollution Transport, and Visibility in Korea." Edited by Detlev Helmig and Md Firoz Khan. *Elementa: Science of the Anthropocene* 7 (December): 57. <https://doi.org/10.1525/elementa.395>.
- 530 Richmond-Bryant, J., M.G. Snyder, R.C. Owen, and S. Kimbrough. 2017. "Factors Associated with NO₂ and NO_x Concentration Gradients near a Highway." *Atmospheric Environment (Oxford, England: 1994)* 174 (November): 214–26. <https://doi.org/10.1016/j.atmosenv.2017.11.026>.
- 535

- Seo, Seunghwan, Si-Wan Kim, Kyoung-Min Kim, Lok N. Lamsal, and Hyungah Jin. 2021. "Reductions in NO₂ Concentrations in Seoul, South Korea Detected from Space and Ground-Based Monitors Prior to and during the COVID-19 Pandemic." *Environmental Research Communications* 3 (5): 051005. <https://doi.org/10.1088/2515-7620/abcd92>.
- 540 Shah, Viral, Daniel J. Jacob, Ke Li, Rachel F. Silvern, Shixian Zhai, Mengyao Liu, Jintai Lin, and Qiang Zhang. 2020. "Effect of Changing NO_x Lifetime on the Seasonality and Long-Term Trends of Satellite-Observed Tropospheric NO₂ Columns over China." *Atmospheric Chemistry and Physics* 20 (3): 1483–95. <https://doi.org/10.5194/acp-20-1483-2020>.
- Sillman, Sanford, Jennifer A. Logan, and Steven C. Wofsy. 1990. "The Sensitivity of Ozone to Nitrogen Oxides and Hydrocarbons in Regional Ozone Episodes." *Journal of Geophysical Research: Atmospheres* 95 (D2): 1837–51. <https://doi.org/10.1029/JD095iD02p01837>.
- Sillman, Sanford, and Perry J. Samson. 1995. "Impact of Temperature on Oxidant Photochemistry in Urban, Polluted Rural and Remote Environments." *Journal of Geophysical Research: Atmospheres* 100 (D6): 11497–508. <https://doi.org/10.1029/94JD02146>.
- Sullivan, John T., Thomas J. McGee, Ryan M. Stauffer, Anne M. Thompson, Andrew Weinheimer, Christoph Knote, Scott Janz, et al. 2019. "Taehwa Research Forest: A Receptor Site for Severe Domestic Pollution Events in Korea during 2016." *Atmospheric Chemistry and Physics* 19 (7): 5051–67. <https://doi.org/10.5194/acp-19-5051-2019>.
- Sun, Wenxiu, Peter Hess, and Chengji Liu. 2017. "The Impact of Meteorological Persistence on the Distribution and Extremes of Ozone." *Geophysical Research Letters* 44 (3): 1545–53. <https://doi.org/10.1002/2016GL071731>.
- 550 KOSTAT (Statistics Korea) (2021). Population, Land Area of the Seoul Metropolitan Area. <https://www.index.go.kr/main.do>. Last Access: 3 November 2022.
- Tong, Weida, Huixiao Hong, Hong Fang, Qian Xie, and Roger Perkins. 2003. "Decision Forest: Combining the Predictions of Multiple Independent Decision Tree Models." *Journal of Chemical Information and Computer Sciences* 43 (2): 525–31. <https://doi.org/10.1021/ci020058s>.
- 555 Vinken, G. C. M., K. F. Boersma, D. J. Jacob, and E. W. Meijer. 2011. "Accounting for Non-Linear Chemistry of Ship Plumes in the GEOS-Chem Global Chemistry Transport Model." *Atmospheric Chemistry and Physics* 11 (22): 11707–22. <https://doi.org/10.5194/acp-11-11707-2011>.
- Vu, Tuan V., Zongbo Shi, Jing Cheng, Qiang Zhang, Kebin He, Shuxiao Wang, and Roy M. Harrison. 2019. "Assessing the Impact of Clean Air Action on Air Quality Trends in Beijing Using a Machine Learning Technique." *Atmospheric Chemistry and Physics* 19 (17): 11303–14. <https://doi.org/10.5194/acp-19-11303-2019>.
- 560 Wang, Haolin, Xiao Lu, Daniel J. Jacob, Owen R. Cooper, Kai-Lan Chang, Ke Li, Meng Gao, et al. n.d. "Global Tropospheric Ozone Trends, Attributions, and Radiative Impacts in 1995–2017: An Integrated Analysis Using Aircraft (IAGOS) Observations, Ozone-sonde, and Multi-Decadal Chemical Model Simulations," 55.
- Wells, Benjamin, Pat Dolwick, Brian Eder, Mark Evangelista, Kristen Foley, Elizabeth Mannshardt, Chris Misenis, and Anthony Weishampel. 2021. "Improved Estimation of Trends in U.S. Ozone Concentrations Adjusted for Interannual Variability in Meteorological Conditions." *Atmospheric Environment* 248 (March): 118234. <https://doi.org/10.1016/j.atmosenv.2021.118234>.
- Werf, Guido R. van der, James T. Randerson, Louis Giglio, Thijs T. van Leeuwen, Yang Chen, Brendan M. Rogers, Mingquan Mu, et al. 2017. "Global Fire Emissions Estimates during 1997–2016." *Earth System Science Data* 9 (2): 697–720. <https://doi.org/10.5194/essd-9-697-2017>.
- 570 Yang, Laura, Daniel J. Jacob, Nadia K. Colombi, Shixian Zhai, Kelvin H. Bates, Viral Shah, Ellie Beaudry, Robert M. Yantosca, Haipeng Lin, Jared F. Brewer, Heesung Chong, Katherine R. Travis, James H. Crawford, Lok Lamsal, Ja-Ho Koo, and Jhoon Kim. 2022. NO₂ vertical profiles over South Korea and their relation to oxidant chemistry: Implications for geostationary satellite retrievals and the observation of NO₂ diurnal variation from space. Manuscript submitted for publication.
- 575 Yeo, Min Ju, and Yong Pyo Kim. 2021. "Long-Term Trends of Surface Ozone in Korea." *Journal of Cleaner Production* 294 (April): 125352. <https://doi.org/10.1016/j.jclepro.2020.125352>.
- Yun, Sug-yeong, and Changhyun Yoo. 2019. "The Effects of Spring and Winter Blocking on PM₁₀ Concentration in Korea." *Atmosphere* 10 (7): 410. <https://doi.org/10.3390/atmos10070410>.
- 580 Zhang, Guoyi, and Yan Lu. 2012. "Bias-Corrected Random Forests in Regression." *Journal of Applied Statistics* 39 (1): 151–60. <https://doi.org/10.1080/02664763.2011.578621>.
- Zhang, Lin, Daniel J. Jacob, Nicole V. Downey, Dana A. Wood, Doug Blewitt, Claire C. Carouge, Aaron van Donkelaar, Dylan B. A. Jones, Lee T. Murray, and Yuxuan Wang. 2011. "Improved Estimate of the Policy-Relevant Background Ozone in the United States Using the GEOS-Chem Global Model with 1/2° × 2/3° Horizontal Resolution over North America." *Atmospheric Environment* 45 (37): 6769–76. <https://doi.org/10.1016/j.atmosenv.2011.07.054>.
- 585 Zhang, Bo, Dan Tong, Meng Li, Fei Liu, Chaopeng Hong, Guannan Geng, Haiyan Li, et al. 2018. "Trends in China's Anthropogenic Emissions since 2010 as the Consequence of Clean Air Actions." *Atmospheric Chemistry and Physics* 18 (19): 14095–111. <https://doi.org/10.5194/acp-18-14095-2018>.

Formatted

Field Code Changed

Deleted: ¶

Maximum monthly 90th percentile MDA8 ozone concentration, ppb




Figure 1: 90th percentile maximum daily 8-h average (MDA8) ozone concentrations in South Korea for 2015-2019. The top row shows the maximum monthly 90th percentile ozone at individual AirKorea sites. The mean of this statistic across the ensemble of sites is shown inset. The bottom row shows 90th percentile MDA8 ozone averaged for individual months over sites within the Seoul Metropolitan Area (SMA, 126.7°E–127.3°E, 37.2°N–37.8°N), as well as for all AirKorea sites. Tick marks are for June and dashed lines are for December. Only sites with over 90% of observational coverage for the 2015 to 2019 period are included in this analysis.

590

24-h average NO₂ concentration, ppb





Figure 2: Same as Figure 1 but for 24-h average annual mean NO₂ concentrations.


595

Maximum monthly 90th percentile MDA



Deleted:

Deleted: <object>



600 **Figure 3:** AirKorea monitoring sites in the Seoul Meteorological Area (SMA) with hourly ozone and NO₂ concentration data for 2015-2019. Red diamonds show the two meteorological sites in the SMA operated by the Korean Meteorological Administration (<https://data.kma.go.kr/data/grnd>).

Deleted: ¶

605

Table 1. Random forest predictor variables for hourly ozone and NO₂ concentrations^a

Meteorology ^b	
	Wind speed
	Wind direction
	Temperature
	Surface pressure
	Relative humidity
Time	
	Day of Year ^c
	Unix time ^d
	Hour of day


^a Hourly explanatory variables in the random forest (RF) model fitted to hourly ozone and NO₂ concentrations averaged across 79 AirKorea sites in the Seoul Metropolitan Area (SMA) for 2015-2019.

^b Meteorological data are from the two SMA Synoptic Meteorological Observation stations (<https://data.kma.go.kr/data/grnd>) located at Gwanaksan (126.975°E, 37.345°N) and Seoul (126.980°E, 37.585°N). Data are averaged across the two stations for input to the RF model.


^c Day of year, used as a seasonal term

^d Used as a linear trend term


610



615 **Figure 4:** Performance of the random forest (RF) model in fitting 2015-2019 hourly ozone and NO₂ concentrations
 in the Seoul Metropolitan Area (SMA). The RF model is trained on hourly concentrations averaged across 79
 AirKorea monitoring sites in the SMA (Figure 3). The Figure compares predicted and observed values for the 30%
 of data withheld from training. Comparison statistics are shown inset including root-mean-square error (RMSE),
 620 correlation coefficient (R), and mean bias (MB). Also shown are the 1:1 lines. Count refers to the number of data
 points within a given (ozone, NO₂) data bin (individual symbol).




625 **Figure 5:** 2015-2019 trends in monthly 90th percentile MDA8 ozone averaged across the 79 AirKorea sites in the
 Seoul Metropolitan Area (SMA). The left panels show the observed trends for individual months and the right
 panels show meteorology-corrected trends. The bottom panels show the 2015-2019 slopes for individual months
 obtained by ordinary least square regressions of the data in the top panels.





630 **Figure 6:** Same as Figure 5 but for 24-h average NO₂ concentrations. Trends are shown in % a⁻¹ relative to the 2015-2019 mean.

GEOS-Chem and observed MDA8 ozone, ppb




635 **Figure 7:** Monthly 90th percentile MDA8 ozone over South Korea and China for different seasons in 2016. GEOS-Chem model results for each season (background contours) are compared to AirKorea and MEE network observations (symbols). 50% of network sites have been culled randomly for visualization purposes. GEOS-Chem correlation coefficient (R) and mean bias (MB) relative to observations are shown inset.



640 **Figure 8:** Seasonal variation of monthly 90th percentile MDA8 ozone in three megacity clusters in 2016. The
 645 clusters are the Seoul Metropolitan area (SMA; 126.7°E–127.3°E, 37.2°N–37.8°N), Yangtze River Delta (YRD;
 30°–33°N, 118°–122°E), and Beijing-Tianjin-Hebei (BTH; 37°–41°N, 114°–118°E). GEOS-Chem results are
 compared to observations and the corresponding root mean square error (RMSE) and mean bias (MB) are shown
 inset. The 90th percentiles are computed from the time series of spatial mean concentrations for each cluster, with
 GEOS-Chem sampled at the network sites.

Deleted: correlation coefficient (R)



Deleted: <object>


Figure 9: Mean vertical ozone profile over Olympic Park (37.522°N, 127.124°E) and Taehwa Research Forest
 (37.312°N, 127.310°E) in Seoul during KORUS-AQ (May- June 2016). Observations from Olympic Park are from 15
 650 ozonesondes and from Taehwa 42 ozonesondes. Ozonesondes were launched between 13:00 and 14:00 local time on
 KORUS-AQ flight days. GEOS-Chem model results are sampled at the observation times. The circles show the surface
 ozone concentrations from GEOS-Chem and at the AirKorea site in closest proximity.

Deleted: ,

Formatted: Indent: Left: 0"


Deleted: at

Deleted: at 13:00 local time in



655 **Figure 10:** Emission-driven trends in 90th percentile MDA8 ozone from 2015 to 2019 in the Seoul Metropolitan Area (SMA) for individual months. The observed meteorology-corrected trend is as shown in Figure 5. The modeled trend is obtained by subtraction of results from simulations with 2015 and 2019 emissions, both using the same 2016 meteorology.

Deleted: <object>




660 **Figure 11:** Emission-driven ozone changes over East Asia from 2015 to 2019 in GEOS-Chem. Results show the 90th percentile MDA8 ozone for May simulated by GEOS-Chem using 2015 emissions, and the difference using 2019 emissions, both for the same meteorological year. The top row shows the baseline simulation described and evaluated with observations in Section 5. The middle and bottom rows show sensitivity simulations with zero

665

anthropogenic emissions in South Korea and China respectively. Spatially averaged values for the Seoul Metropolitan Area (SMA) and South Korea are given inset.

GEOS-Chem 90th percentile MDA8 ozone in May



Deleted: <object>

670

Figure 12: East Asia background ozone and individual enhancements due to anthropogenic emissions from ships in the Yellow Sea (north of 30.5° N), South Korea, and China. Results show the monthly mean 90th percentile MDA8 ozone for May simulated by GEOS-Chem for meteorological year 2016.