

# Environmental Controls on Observed Spatial Variability of Soil Pore Water Geochemistry in Small Headwater Catchments Underlain with Permafrost

Nathan Alec Conroy,<sup>1\*</sup> Jeffrey M. Heikoop,<sup>1</sup> Emma Lathrop,<sup>1,2</sup> Dea Musa,<sup>1</sup> Brent D. Newman,<sup>1</sup> Chonggang Xu,<sup>1</sup> Rachael E. McCaully,<sup>3</sup> Carli A. Arendt,<sup>3</sup> Verity G. Salmon,<sup>4</sup> Amy Breen,<sup>5</sup> Vladimir Romanovsky,<sup>6</sup> Katrina E. Bennett,<sup>1</sup> Cathy J. Wilson,<sup>1</sup> and Stan D. Wulfschleger<sup>4</sup>

<sup>1</sup>Earth and Environmental Sciences Division, Los Alamos National Laboratory, Bikini Atoll Road, Los Alamos, New Mexico, 87545, USA

<sup>2</sup>Center for Ecosystem Science and Society, Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ, 86011, USA

<sup>3</sup>Department of Marine Earth and Atmospheric Sciences, North Carolina State University, Raleigh, North Carolina, 27695, USA

<sup>4</sup>Biological and Environmental Systems Science Division and Climate Change Science Institute, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 37831, USA

<sup>5</sup>International Arctic Research Center, P.O. Box 757340, University of Alaska, Fairbanks, Alaska 99775-7340, USA

<sup>6</sup>Geophysical Institute, University of Alaska Fairbanks, Fairbanks, Alaska, 99775, USA

*Correspondence to:* Nathan Alec Conroy (nconroy@lanl.gov)

**Abstract.** Soil pore water (SPW) chemistry can vary substantially across multiple scales in Arctic permafrost landscapes. The magnitude of these variations and their relationship to scale are critical considerations for understanding current controls on geochemical cycling and for predicting future changes. These aspects are especially important for Arctic change modelling where accurate representation of sub-grid variability may be necessary to predict watershed scale behaviours. Our research goal was to characterize intra- and inter-watershed soil water geochemical variations at two contrasting locations in the Seward Peninsula of Alaska, USA. We then attempt to establish which environmental factors were important for controlling concentrations of important pore water solutes in these systems. The SPW geochemistry of 18 locations spanning two small Arctic catchments were examined for spatial variability and its dominant environmental controls. The primary environmental controls considered were vegetation, soil moisture/redox condition, water/soil interactions and hydrologic transport, and mineral solubility. The sampling locations varied in terms of vegetation type and canopy height, presence or absence of near-surface permafrost, soil moisture, and hillslope position. Vegetation was found to have a significant impact on SPW  $\text{NO}_3^-$  concentrations, associated with the localized presence of nitrogen-fixing alders and mineralization and nitrification of leaf litter from tall willow shrubs. The elevated  $\text{NO}_3^-$  concentrations were however, frequently equivoiced by increased microbial denitrification in regions with sufficient moisture to support it. Vegetation also had an observable impact on soil moisture sensitive constituents, but the effect was less significant. The redox conditions in both catchments were generally limited by Fe reduction, seemingly well-buffered by a cache of amorphous Fe hydroxides, with the most reducing conditions found at sampling locations with the highest soil moisture content. Non-redox-sensitive cations were affected by a wide variety of water-soil interactions that affect mineral solubility and transport. Identification of the dominant controls on current SPW hydrogeochemistry allows for qualitative prediction of future geochemical trends in small Arctic catchments that are likely to experience warming and permafrost thaw. As source

39 areas for geochemical fluxes to the broader Arctic hydrologic system, geochemical processes occurring in these  
40 environments are particularly important to understand and predict with regards to such environmental changes.

## 41 **1. Introduction**

42 Permafrost thaw in the Arctic is causing significant changes to landscape structure (Kokelj and Jorgenson, 2013;  
43 Rowland et al., 2010), hydrology (Hiyama et al., 2021; Kurylyk et al., 2021; Liljedahl et al., 2016; Vonk, Tank, and  
44 Walvoord, 2019; Walvoord and Kurylyk, 2016), vegetation (Lara et al., 2018; Myers-Smith et al., 2011; Sturm,  
45 Racine, and Tape, 2001; K. D. Tape et al., 2012; Tape, Sturm, and Racine, 2006;), and biogeochemistry (O'Donnell  
46 et al., 2021; Frey and McClelland, 2009; Salmon et al., 2019; Vonk, Tank, and Walvoord, 2019). The integrated  
47 hydrogeochemical effects of these environmental changes are already apparent in the chemistry of the large Arctic  
48 rivers, where fluxes of carbon and nutrients are increasing, leading to enhanced nutrient loadings, with strong  
49 implications for the global carbon cycle (Bring et al., 2016; Fuchs et al., 2020; McClelland et al., 2016). While the  
50 watershed areas of large Arctic rivers are vast, recent studies suggest that solute concentrations in these large rivers  
51 are likely controlled by solute generation processes occurring at much smaller scales (Harms and Ludwig, 2016; Koch  
52 et al., 2013; Shogren et al., 2019; Vonk et al., 2015).

53 While there is a rapidly growing body of literature focused on observing and understanding environmental changes  
54 over time with further Arctic warming, relatively few studies directly address the existing spatial variability, within  
55 catchments or across catchments, and we are not aware of any studies that have combined field observations with  
56 thermodynamic modelling in an effort to understand the causes of the existing spatial variability. Therefore, we have  
57 a limited understanding of the key environmental controls on the spatial distribution of soil pore water solute  
58 concentrations. In this study, we quantitatively evaluate the spatial variability of soil pore water (SPW) geochemistry  
59 within and between two distinct catchments underlain with permafrost, and then seek to identify the source of the  
60 observed spatial variability.

61 This study takes advantage of a scientifically diverse array of observations and datasets made available by the Next  
62 Generation Ecosystem Experiment (NGEE) Arctic project, sponsored by the US Department of Energy Office of  
63 Science. Most of the locations studied herein were selected by the NGEE Arctic project to provide co-located  
64 measurements in a wide range of vegetation types, nested within representative hillslopes and catchments. Although  
65 selected largely to represent a range of vegetation structure, such as shrub abundance and canopy height, these  
66 locations also have considerable variability in other environmental parameters including, but not limited to: soil  
67 moisture and temperature, presence or absence of near-surface permafrost, and maximum observed thaw depth (**Table**  
68 **1** and **Table 2**). The vegetation-delineated sampling approach provides an opportunity to not only quantify the  
69 biogeochemical variability of SPW in Arctic environments, but also to investigate the root causes of that observed  
70 variability. Data from additional sampling locations, available from a co-located study, were also utilized when  
71 possible.

72 Our overarching hypothesis is that vegetation-type and hillslope position are the dominant controls on spatial  
73 variability of SPW geochemistry at the NGEE Arctic field sites located on the Seward Peninsula. Vegetation-type  
74 seems likely to have a significant effect on SPW geochemistry both directly and indirectly. Indirect effects would

75 include vegetation canopy impacts on soil moisture (through evapotranspiration and snow trapping). Direct effects of  
76 vegetation would include nutrient cycle changes resulting from the annual deposition of plant litter. Such a direct  
77 effect can be augmented at sites populated by alder shrubs due to this genus of deciduous shrubs ability to form a  
78 symbiotic relationship with nitrogen-fixing *Frankia*, which they host in underground root nodules. Nitrogen fixation  
79 associated with alders has previously been shown to accelerate local nitrogen cycling (Binkley et al., 1992; Clein and  
80 Schimel, 1995; Bühlmann et al., 2014). Soil moisture will also affect SPW geochemistry, particularly of redox  
81 sensitive species, by limiting oxygen diffusion and thus controlling which regions develop anoxic/reducing  
82 geochemical conditions. Soil moisture impacts will likely be correlated with vegetation-type as well as hillslope  
83 position, and the presence or absence of perching layers, including permafrost, all of which impact the vertical and  
84 horizontal drainage characteristics of a watershed. Chemical species that are not redox-sensitive or controlled by  
85 biogeochemical reactions are likely to be affected by transport, solubility, and water/sediment/organic matter  
86 interactions, and therefore largely controlled by hillslope position as well as soil moisture.

87 Identifying the dominant controls on solute concentration variability within each catchment and across catchments  
88 will facilitate better projections of future soil pore hydrogeochemistry in permafrost landscapes, and how these  
89 signatures are related to changing soil moisture and increasing tundra shrub abundance in a changing Arctic (Bring et  
90 al., 2016; Myers-Smith et al., 2011; Prowse et al., 2015; Salmon et al., 2019; Sturm et al., 2001; Tape et al., 2012;  
91 Tape et al., 2006; Wrona et al., 2016, 2016). Arctic warming and associated permafrost thaw will increase hydrological  
92 connectedness between terrestrial and aquatic environments through deepening of the active layer and the formation  
93 of deeper, more coherent groundwater flow paths (Bring et al., 2016; Harms and Jones, 2012; Prowse et al., 2015a;  
94 Prowse et al., 2015b). Meanwhile, changes in hydrogeochemical signatures in larger Arctic rivers are likely to  
95 originate in smaller catchments (McClelland et al., 2016; Prowse et al., 2015; Shogren et al., 2019; Spence et al.,  
96 2015). In this sense, changes in hydrogeochemistry in small Arctic catchments not only impact hydrogeochemistry at  
97 much larger scales, but also prognosticate the future hydrogeochemistry of larger Arctic rivers.

## 98 **2. Methods**

### 99 **2.1 Site Descriptions**

100 This study focuses on two sites with permafrost on the Seward Peninsula of western Alaska, the Teller-27 Catchment  
101 and the Kougarok-64 Hillslope (**Figure 1**). The Teller-27 Catchment, henceforth “Teller,” is a small (~2.25 km<sup>2</sup>)  
102 headwater catchment located west of mile marker 27 along the Nome-Teller Highway northwest of Nome, Alaska.  
103 The Kougarok-64 Hillslope, henceforth “Kougarok,” is a hillslope (~2.0 km<sup>2</sup>) located west of mile marker 64 along  
104 the Nome-Taylor Highway northeast of Nome, Alaska. We utilized data from “intensive stations” at both Teller and  
105 Kougarok where concentrated, multi-year, co-located observations of soil water chemistry, vegetation characteristics,  
106 soil moisture and temperature, and other measurements have been collected as part of the NGEE Arctic Research  
107 Project. These are identified as TL# (Teller Station #) or KG# (Kougarok Station #) in **Figure 2** and **Figure 3**,  
108 respectively. It should be noted that Teller and Kougarok are not “paired watersheds” in the classical sense, differing  
109 in only one major characteristic, which provides the basis for comparison. Instead, Teller and Kougarok differ in many

110 respects and are both representative of the broad range of hillslope conditions common on the Seward Peninsula.  
111 Detailed descriptions of Teller and Kougarok have been published previously (Jafarov et al., 2018; Léger et al., 2019;  
112 Philben et al., 2019, 2020; Salmon et al., 2019; Yang et al., 2020), therefore, only the catchment characteristics that  
113 are probable sources of variability in SPW chemistry will be highlighted here.

114 Teller is a discrete catchment with a well-defined central drainage, a vertical declivity of approximately 200 m, and a  
115 catchment area of approximately 2.25 km<sup>2</sup>. Temperature probes, soil pits, coring activities, and geophysical  
116 interpretations at Teller have confirmed the catchment is underlain with discontinuous permafrost (Léger et al., 2019).  
117 The upper shoulder of Teller (near Station 5, **Figure 2** and **Figure 5**) is underlain with near-surface permafrost and  
118 appears to be a degraded peat plateau. The resultant microtopography of the degraded peat and the shallow perching  
119 horizon caused by the permafrost creates a landscape of unsaturated peat mounds surrounded by ponds and saturated  
120 soils. Downslope of the peat plateau, the Teller hillslope has highly variable soil moisture and vegetation (**Table 1**).  
121 The microtopography within the lower footslope looks similar to the upper shoulder, but the peat appears more  
122 severely degraded and the cause of the perched water table is less clear. Léger et al. (2019) suggest the presence of  
123 permafrost at a depth of 1 – 2 m at Teller Station 9 (**Figure 2**), but the perching could also be caused by a layer of silt,  
124 at a depth of approximately 30 cm (Graham et al., 2018). The full extent of permafrost and silt in this region of the  
125 catchment remains unknown, but the thaw depth in July 2018 was greater than 1 m and maintained a perched water  
126 table (Philben et al., 2020), suggesting perching could be the result of silt rather than permafrost. Vegetation type,  
127 moisture content, permafrost extent, and hillslope position for all Teller Stations are summarized in **Table 1**.

128 Kougarok differs in many ways from Teller, although both have characteristics that are typical of hillslopes on the  
129 Seward Peninsula. Kougarok is a convex hillslope, with a vertical declivity of approximately 70 meters. The study  
130 area at Kougarok is approximately 2.0 km<sup>2</sup>. Soil temperature measurements at Kougarok suggest that the vast majority  
131 of the site is underlain by shallow continuous permafrost (Romanovsky, Cable, and Dolgikh, 2020a); Kougarok  
132 Station 5 is an exception, where the permafrost is deeper (Romanovsky et al., 2020a). The upper shoulder of Kougarok  
133 is a well-drained rocky outcrop composed of metagranitic rock (Hopkins et al., 1955; Till, Dumoulin, Werdon, and  
134 Bleick, 2011). Saturated soils are not prevalent until the footslope and the lower backslope, where Kougarok Stations  
135 2, 11, 10, 1, and 6 are situated (**Figure 3**). The lower backslope is characterized by persistent saturation between  
136 ubiquitous tussocks, formed by the tussock cotton grass *Eriophorum vaginatum*. The tussock-lichen tundra at  
137 Kougarok introduces microtopography and spatially variable saturation; in this sense, the Kougarok tussocks are  
138 analogous to the peat mounds and hummocks at Teller, but on different spatial scales and formed by different  
139 processes. Kougarok has numerous patches of alder shrubland in an altitudinal band within the upper backslope; it  
140 should be emphasized that Teller lacks tussock-lichen tundra and alder (*Alnus viridis* ssp. *fruticosa*) shrubs that are a  
141 feature of Kougarok. While continuous permafrost largely remains, the Kougarok site appears to be undergoing  
142 environmental changes as evidenced by an increase in alder coverage over the past decades (Salmon et al., 2019). Soil  
143 profiles underneath the alder patches are rocky with shallow bedrock and warmer permafrost (**Table 2**). Shrub tundra  
144 (alder savanna in tussock tundra and willow-birch tundra) dominates the lower backslope, where the annual active  
145 layer thickness is typically less than 100 cm. Vegetation type, moisture content, permafrost extent, and hillslope  
146 position at all Kougarok stations are summarized in **Table 2**.

## 147 2.2 Sampling & Analytical Approach

148 SPWs were sampled using two complimentary techniques. Fiberglass wicks (Frisbee et al., 2010) were deployed in  
149 the upper 30 cm of soils at stations where shallow soils were unsaturated. These wicks were left in place from year-  
150 to-year and only replaced if damage was observed or suspected. The sample reservoirs from the wicks were collected  
151 whenever possible, usually a few times each summer. MacroRhizons (Rhizosphere Research Products; Netherlands)  
152 were used at stations that were more saturated, also targeting the upper 30 cm of soils. Both techniques were used at  
153 stations of intermediate saturation, where both could be deployed effectively. MacroRhizons represent a relatively  
154 discrete temporal sampling event (minutes to hours), whereas wicks represent a cumulative water collected over longer  
155 periods (weeks to months). It is in this sense, that the two techniques are complimentary. Unfortunately, due to  
156 saturation variability both techniques could not be used at all stations and conditions at some Kougarok stations were  
157 sometimes too dry to collect meaningful volumes of SPW using either method. Additional SPW data from Kougarok  
158 were supplemented from a separate study focused on alder-related nutrient dynamics (McCaully et al., 2022). These  
159 data were collected by MacroRhizons and are captured as Kougarok Stations 10 – 13, which were not part of the  
160 original stations established by the NGEE Arctic Program. A total of 309 SPW samples from Kougarok were collected  
161 and analysed, whereas a total of 89 SPW samples from Teller were collected and analysed.

162 After collection, SPW cation concentrations were measured in triplicate by inductively coupled plasma optical  
163 emission spectroscopy (Optima 2100 DV; PerkinElmer, USA) following US EPA Method 200.7. Inorganic anion  
164 concentrations were measured by ion chromatography (DX-600; Dionex, USA) following US EPA Method 300.0. B,  
165 F, K, Na, and Si concentrations collected by wicks were excluded from the dataset due to known issues with these  
166 ions leeching from fiberglass wick samplers (Perdrial et al., 2014; Wallenberger and Bingham, 2009). This effect is  
167 illustrated in **Supplementary Figure 1** and the lack of such an effect for divalent cations is shown in **Supplementary**  
168 **Figure 2**. Comparison of data from wicks and MacroRhizons, along with the observations from (Perdrial et al., 2014),  
169 demonstrates that remaining constituents discussed herein were not affected by collection with fiberglass wicks.  
170 Alkalinity, pH, and  $E_H$  are all critical geochemical parameters that are susceptible to change during storage (Petroni  
171 et al., 2007); because of the large amount of data from wicks these parameters were not considered further, except in  
172 the context of thermodynamic modelling.

173 Observations related to vegetation, soil moisture, and permafrost extend were compiled from datasets made available  
174 by the NGEE Arctic project and are given for Teller in **Table 1** and for Kougarok in **Table 2**. The reported soil  
175 moisture contents were derived from an average of gravimetric measurements (2017 and 2018) and time domain  
176 reflectometry measurements (2017 and 2019), and from remotely-sensed P-band Synthetic Aperture Radar (2017).  
177 End-of-winter snow depths were measured in March and April of 2016, 2017, and 2018. The annual average ground  
178 temperature was measured using in-situ temperature sensors (HOBO U30 DataLogger) at a depth of 1.5 meters below  
179 the ground surface (Romanovsky et al., 2020a; Romanovsky, Cable, and Dolgikh, 2020b) and the active layer  
180 thicknesses were determined by frost probe in September 2019 at the end of the growing season. Vegetation data were  
181 collected at the peak of the growing season in mid to late July 2016 and 2017 at the NGEE Arctic Kougarok and Teller  
182 field sites, respectively. The distribution of plant communities in the Arctic is primarily controlled by landscape,  
183 topography, soil chemistry, soil moisture, and the plants that historically colonized an area (Raynolds et al., 2019).

184 Soil available rooting depth, which can be limited by shallow depths to bedrock, permafrost, or the water table, can  
185 also restrict plant growth and survival of certain species by reducing access to water and nutrients. We surveyed the  
186 dominant plant communities along each hillslope, which varied in their shrub abundance, canopy height, and structure,  
187 to characterize the vegetation composition at the sites following the recommended protocol of Walker et al. (2016).  
188 Extensive field site details and vegetation sampling methods are more thoroughly described in previous studies  
189 (Salmon et al., 2019; Langford et al., 2019; Yang et al., 2020; Sulman et al., 2021; Yang et al. 2021).  
190 For this study, we provide summary statistics for vegetation plots associated with intensive stations. Vegetation  
191 composition plots within each intensive station were chosen subjectively in areas of homogeneous and representative  
192 vegetation varying in size from 1 to 25 m<sup>2</sup> depending on canopy structure and height. The surveyed plot area was 1 ×  
193 1 m for all plant communities except for the taller stature willow-birch tundra, mesic willow shrubland (2.5 × 2.5 m),  
194 and alder shrubland (5 × 5 m). For each plot, all plant species (vascular plants, lichens, and bryophytes) were recorded  
195 along with visual estimates of their percent cover. For plots with multiple canopies, field cover estimates were recorded  
196 as absolute cover, meaning that the total cover per plot can be >100%. We calculated relative cover values (adding to  
197 100%) from the field data and use these for all subsequent analyses.  
198 Plant species were further aggregated into nine plant functional types (PFTs), groupings of plant species that share  
199 similar growth forms and roles in ecosystem function (Wullschleger et al., 2014), based on growth patterns and plant  
200 traits. PFTs in this study include: (1) nonvascular mosses and lichens, (2) deciduous and evergreen shrubs of various  
201 height classes, including an alder PFT, (3) graminoids, and (4) forbs. Photos of representative PFTs from both sites  
202 are given in **Supplementary Figures 9-17**. Canopy height was estimated within each plot for each PFT as the average  
203 of 4 measurements, including a maximum canopy height. Active layer depth was measured at the end of the growing  
204 season for all plots in September 2018 using a frost probe. A temperature probe was used to determine if the resistive  
205 layer was permafrost ( $\leq 0$  °C) or rock ( $> 2$  °C). Thaw depth is an average of 4 measurements from the vegetation plot  
206 corners.

### 207 **2.3 Statistical Analysis**

208 Principal Components Analysis (PCA) and the Mann-Whitney U-Test (MWUT) were both used to investigate  
209 dominant environmental controls on solute concentrations in SPWs at Teller and Kougarak. PCA is an exploratory  
210 data analysis tool that reduces the dimensionality of large complex datasets and considers how components (i.e. solute  
211 concentrations) vary together. Because PCA was predominately used as a screening tool to reveal geochemical  
212 correlations that may not have been evident by traditional geochemical causations or inference, a detailed discussion  
213 of the PCA results is reserved to the Supplementary Materials. The MWUT was used to test for significant differences  
214 in solute concentrations between Teller and Kougarak (inter-site variability) and between stations at each site (intra-  
215 site variability). The MWUT is a non-parametric method of challenging a null hypothesis, which in this case is the  
216 assumption that the concentrations of a given solute are not systematically greater at either site nor at any particular  
217 station. Water chemistry data are typically not normally distributed and thus, non-parametric difference tests such as  
218 the MWUT are preferred. The MWUT challenges the distribution of values, not the means. In this work, the level of  
219 significance associated with the null hypothesis was operationally defined as 0.05, which equates to a 95 % chance

220 that an observed statistical difference is real and not coincidental. This error rate is operationally defined per contrast  
221 (i.e. a 95 % chance that the observed statistical difference in nitrate concentrations between Teller Station 9 and Teller  
222 Station 7 is real or that the observed statistical difference in sulphate concentrations between Teller and Kougarok is  
223 real) as opposed to familywise (i.e. a 95 % chance that all of the observed/reported statistical differences are real and  
224 not coincidental). MWUTs were completed using the methods described in Corder and Foreman (2009) and PCA was  
225 completed using packages available in R statistical software, version 3.3.6 (Corder and Foreman, 2009; R Core Team,  
226 2020). For all analyses, concentrations below the method detection limit were operationally defined as half the  
227 detection limit, in agreeance with (Helsel, 2005, p. 43). While the emphasis of this study was on site/station (i.e.  
228 spatial) variability, it should be recognized that seasonal and inter-annual variability could also be significant. To  
229 minimize seasonal forcing on the variability observed, all SPW geochemical data presented were collected during the  
230 thaw season between June and September.

## 231 **2.4 Thermodynamic Modelling**

232 To investigate thermodynamic controls on solute behaviour, particularly solubility limitations, thermodynamic  
233 modelling exercises were undertaken using PHREEQC, a thermodynamic geochemical modelling code, and  
234 PhreePlot, which facilitates repetitive PHREEQC calculations through looping (Kinniburgh and Cooper, 2011;  
235 Parkhurst and Appelo, 2013). Because this study was focused on elucidating the primary geochemical controls on  
236 solute concentrations in SPWs and not on developing a rigorous transport model, representative concentrations were  
237 used instead of station specific concentrations. Representative “low”, “median”, and “high” concentration conditions  
238 were proxied from the 25th, 50th, and 100th concentration percentiles, respectively, taken from both Teller and  
239 Kougarok (Supplementary **Table 4**). Meanwhile, representative pH and  $E_H$  ranges were determined either through  
240 direct measurement (pH), or indirectly by correlating dissolved  $Fe^{2+}$  concentrations and pH with a redox condition  
241 through geochemical models and the Nernst equation. Modelling exercises were performed utilizing the phreeqc.dat  
242 database, with the only modification being the suppression of methane production by inorganic carbonate reduction.  
243 Modelling exercises were performed at the default PHREEQC modelling temperature (25 °C), as the selection of an  
244 alternative defensible temperature was non-trivial; temperatures on the Seward Peninsula span a very wide range and  
245 its unclear what temperature would be most suitable for mineral solubility limitation modelling. Ultimately, because  
246 the thermodynamic models were used as a tool understand what could be controlling soil pore water solute  
247 concentrations and were not intended to model the system or to predict future concentrations, the default temperature  
248 was decided to be the most suitable. While there is some temperature dependence of mineral solubility, the differences  
249 in predicted solubility between 4 °C and 25 °C did not impact the interpretation of our results (**Supplementary Figure**  
250 **8**). Methane production was “turned-off” to maintain carbonate availability under reducing conditions to help identify  
251 any possible carbonate minerals that could be precipitating. Because alkalinity was only measured in a small number  
252 of samples, carbonate concentration percentiles were estimated from charge imbalances. Alkalinity and charge  
253 imbalance were very well correlated in samples where alkalinity was measured (**Supplementary Figure 3**). Although  
254 not a particularly rigorous modelling exercise, this approach was sufficient to identify mineral phases that could be  
255 controlling solute generation processes through solubility limitations.

256 **3. Results**

257 **3.1 Physical Characteristics of Stations (Co-Located Studies)**

258 Controls on the observed spatial variability of SWP solute concentrations at Teller and Kougarok stations were  
259 deduced, in part, from differences in physical features and conditions of each station. Quantitative measures of many  
260 of these physical characteristics were available from the interdisciplinary studies co-located at the Teller and Kougarok  
261 stations. The extent of permafrost, ground temperature, active layer depth, soil moisture content, snow depth,  
262 vegetation type, vegetation canopy height, dominant plant functional type, and hillslope position were all compiled  
263 from these co-located studies. Using these measures, the physical characteristics of each station are summarized in  
264 **Table 1** and **Table 2**, grouped by vegetation type.

265 **3.2 Inter-site Variability: Teller versus Kougarok**

266 Mann-Whitney U-Testing revealed that the concentrations of 14 of the 23 constituents analysed were significantly  
267 different ( $1.96 < |z|$ ) between Teller and Kougarok (**Table 3**). The effect size, a measure of how significantly different  
268 the concentrations were, were large for Na and F; medium-large for K and Si; medium for Al, Oxalate, B, Zn,  $\text{SO}_4^{2-}$ ,  
269 Fe, Ba, Ti, and  $\text{NO}_2^-$ ; and small-medium for Li. The terminology and thresholds for these semi-quantitative differences  
270 in correlation were taken from Corder and Foreman (2009). Mann-Whitney U-testing revealed that SPW  
271 concentrations of many constituents were significantly different between Teller and Kougarok (**Table 3**). When  
272 concentrations were significantly different between the sites, Kougarok generally exhibited the higher concentrations  
273 of the two. SPW concentrations of Na, F, K, Si, Al, oxalate, B, Zn, Fe, Ba, Ti,  $\text{NO}_2$ , and Li were all significantly  
274 greater at Kougarok than Teller, while only  $\text{SO}_4^{2-}$  concentrations were significantly greater at Teller. Meanwhile, the  
275 concentrations of Br,  $\text{NO}_3^-$ , Sr,  $\text{PO}_4$ , Mg, Cr, Mn, Cl, and Ca were not significantly different between Teller and  
276 Kougarok. A summary of the inter-site MWUT results are given in **Table 3** with the constituents that exhibited  
277 significant differences between the sites displayed over a darkened background.

278 **3.3 Intra-site Variability: Teller and Kougarok Stations**

279 Mann-Whitney U-Testing was also used to test for intra-site differences between stations at both Teller and Kougarok.  
280 Boxplots and compact letter displays are used to visualize the within-site variability of a select group of constituents  
281 of interest (COIs), which are given in **Figure 4**. Tables of the results of the intra-site MWUTs for all constituents that  
282 were monitored, including those that did not demonstrate some systematic inter-station variability or were not  
283 otherwise of interest, are given in the Supplementary Materials. Our interpretation of the major environmental controls  
284 on the observed spatial variability of SPW solute concentrations between stations are shown in **Table 4**. Each of these  
285 controls, including vegetation effects, soil moisture and redox effects, weathering, water/soil interactions and  
286 hydrological transport effects, and mineral solubility effects, is considered in detail in the following sections.

287 **3.3.1 Vegetation Effects**

288 Vegetation can influence hydrogeochemical variability directly via vegetation-induced changes to elemental cycling  
289 and soil moisture contents, or indirectly via the secondary impacts changes in soil moisture can have on mineral



290 solubility or on the soil redox condition. The geochemical consequences of solubility and redox conditions are the  
291 focus of sections to follow, thus, this section will focus on direct vegetation effects via influences on elemental cycling  
292 and soil moisture via evapotranspiration and preferential trapping of snow.

293  $\text{NO}_3^-$  was the only COI that showed a distinct effect from vegetation via elemental cycling. Elevated  $\text{NO}_3^-$   
294 concentrations were associated with the presence of alder shrubs and, in some cases, willow shrubs.  $\text{NO}_3^-$   
295 concentrations at both sites were generally low, with the exception of Kougarak Stations 3, 5, and 12, and Teller  
296 Station 7 (**Figure 4**). Low to tall alder shrubs are the dominant vegetation type at Kougarak Stations 3 and 12 (**Table**  
297 **2**). Meanwhile, alders are present at Kougarak Station 5 despite the dominant vegetation type being low willow and  
298 birch shrubs. Alders increase soil nitrogen through a symbiotic relationship with nitrogen-fixing bacteria that reside  
299 in their root nodules, thus, an association between  $\text{NO}_3^-$  concentrations and alder vegetation is expected (Salmon et  
300 al., 2019).

301 Perhaps more noteworthy was the lack of elevated  $\text{NO}_3^-$  concentrations at Kougarak Stations 1, 2, 6, 10, and 11. The  
302 vegetation type at Kougarak Stations 1, 2, 6, 10, and 11 is alder savanna in tussock tundra, which is a mixed graminoid-  
303 shrub tundra with shorter stature and lower density of alder shrubs, yet nonetheless nitrogen input via alder derived  
304 nitrogen-fixation is anticipated to occur. The lack of elevated  $\text{NO}_3^-$  suggests either that 1) nitrogen-fixation in alder  
305 savanna in tussock tundra is insufficient to result in an increase in  $\text{NO}_3^-$  concentrations, 2) that the Kougarak footslope  
306 and lower backslope is very nitrogen-limited, and thus, that  $\text{NO}_3^-$  is largely consumed by vegetation as it is fixed, or  
307 3) that microbes in the Kougarak footslope and lower backslope rapidly denitrify the available  $\text{NO}_3^-$  as a substitute  
308 for oxygen in their metabolisms. The smaller shrub size and density in the alder savanna in tussock tundra certainly  
309 results in less accumulated leaf litter relative to the denser and larger alder shrubland intensive stations, as such, it  
310 seems reasonable that less nitrogen would be available at stations in alder savanna in tussock tundra. Meanwhile,  
311 isotopic measurements of nitrogen downslope of alder patches at Kougarak Stations 12 and 3 also support the  
312 occurrence of denitrification (McCaully et al., 2022). Therefore, we believe the lack of elevated  $\text{NO}_3^-$  concentrations  
313 at Kougarak Stations 1, 2, 6, 10, and 11 is a combination of less alder leaf litter and greater denitrification, than at  
314 Kougarak Stations 3, 5, or 12.

315 At Teller, only Station 7 exhibited elevated  $\text{NO}_3^-$  concentrations relative to the rest of the catchment (**Figure 4**). Teller  
316 Station 7 is dominated by tall willow shrubs and is relatively dry. Mineralization and nitrification of willow leaf litter  
317 coupled with limited microbial denitrification is the presumed cause of elevated  $\text{NO}_3^-$  concentrations at Teller Station  
318 7. Teller Station 2 also has tall willow shrubs but did not exhibit elevated  $\text{NO}_3^-$  concentrations. From the limited scope  
319 of this study, it is unclear why Teller Station 2 did not exhibit elevated  $\text{NO}_3^-$  while Station 7 did, but we suspect that  
320 higher seasonal moisture content and greater microbial denitrification at Teller Station 2 likely played a role. Also of  
321 note was that despite significant intra-site  $\text{NO}_3^-$  concentration differences, inter-site differences were not significant  
322 ( $|z| = 1.59$ ) and that relatively few Kougarak stations showed elevated  $\text{NO}_3^-$  concentrations, despite a widespread alder  
323 presence. Increased microbial denitrification is suspected to balance increased nitrogen-fixation at these stations. This  
324 is consistent with previous studies that have noted higher nitrogen mineralization rates in acidic tundra than non-acidic  
325 tundra (Weiss et al., 2005); Kougarak is predominantly acidic tundra and Teller is non-acidic tundra.

326 The effect of vegetation on spatial variability of soil moisture was not readily observed in the volumetric moisture  
327 content of soil (**Table 1** and **Table 2**) but was somewhat apparent in the spatial variability of moisture sensitive  
328 constituents, such as Cl (**Figure 4**). The lack of a clear correlation between vegetation and soil moisture by TDR or  
329 P-band SAR observations is perhaps do to the coarseness of the P-band SAR observations and the strong seasonality  
330 associated with both methods. Moisture sensitive constituents, such as Cl, may provide a more seasonally averaged  
331 tracer of soil moisture content at the stations. An increase in Cl concentrations with vegetation canopy height was  
332 apparent at Teller stations suggesting an evapotranspiration effect. This trend was also apparent at Kougarok, but the  
333 differences were rarely significant. Overall, the spatial variability of soil moisture sensitive constituents, like Cl, was  
334 far less correlated with vegetation-type than expected; perhaps due to preferential trapping of snow, which may offset  
335 the increased evapotranspiration of tall shrubs more than previously realized. Overall, Cl concentrations at Kougarok  
336 appeared to be more correlated with hillslope position than with vegetation canopy height (**Figure 4**).

### 337 **3.3.2 Soil Moisture and Redox Effects**

338 Soil moisture content can have a profound effect on redox sensitive elements. Saturation limits oxygen diffusion into  
339 soil, and thus, forces microorganisms to utilize less energetic electron acceptors to metabolize organic matter. In an  
340 ideal system, soil microorganisms will use the strongest electron acceptor available, until it is exhausted. Although  
341 natural environments are not ideal systems, redox conditions in soils follow a more or less stepwise progression.  
342 Therefore, by evaluating the dissolved concentrations of  $\text{NO}_3^-$ , Mn, Fe, and  $\text{SO}_4^{2-}$  in SPWs, it is possible to  
343 qualitatively assess soil redox conditions and their impact on hydrogeochemical variability.

344 The redox conditions at both Teller and Kougarok are generally limited by Fe reduction, with the most reducing  
345 conditions found at stations with the highest soil moisture content. As such,  $\text{NO}_3^-$  concentrations are generally low  
346 (**Table 3**),  $\text{SO}_4^{2-}$  concentrations are relatively consistent (**Figure 4**), and Mn and Fe concentrations increase with  
347 increasing soil moisture (**Figure 4**).  $\text{NO}_3^-$  concentrations were generally low, except for drier stations in the proximity  
348 of tall alders or willows. While  $\text{NO}_3^-$  inputs are discussed in the vegetation effects section, the lack of high  $\text{NO}_3^-$   
349 concentrations at wetter stations that contain alders suggests that soil moisture coupled with microbial denitrification  
350 bares a strong control on SPW  $\text{NO}_3^-$  concentrations. Meanwhile,  $\text{SO}_4^{2-}$  concentrations at both sites are relatively  
351 constant across clear moisture and redox gradients (**Figure 4**). This suggests that  $\text{SO}_4^{2-}$  reduction is not pervasive at  
352 either site. Dissolved Fe concentrations were higher at stations with higher soil moisture content, consistent with Fe  
353 reduction. Similarly, Mn concentrations were slightly elevated at wetter stations. The concentrations of Mn, however,  
354 rarely rose above  $0.05 \text{ mg}\cdot\text{L}^{-1}$ , suggesting either Mn solubility limitations or a lack of a significant Mn weathering  
355 source. Low Mn concentrations at Teller Station 5, a wetter station on the upper shoulder of the Teller watershed  
356 (**Table 1; Figure 2**) seems to support the latter conclusion, as do geochemical modelling exercises (Section 4.5).  
357 Together, these results suggest that the most reducing condition at both sites is typically limited to Fe reduction and  
358 that this only occurs at stations with the highest soil moisture contents.

### 3.3.3 Weathering, Water/Soil Interaction, and Hydrological Transport Effects

A combination of weathering, water/soil interactions, and hydrological transport were identified as probable drivers of hydrogeochemical variability for some solutes. As noted by Philben et al. (2020), soil derived solutes tend to accumulate in low-lying areas within watersheds. This is observed at Teller, where the concentrations of Ca, Sr, and Mg all increase dramatically at the transition from lower backslope to footslope (**Figure 5**). Both Teller and Kougarok are underlain by carbonate-rich metamorphic facies, and Ca, Sr, and Mg are probable carbonate counter-cations. Weathering of Ca, Sr, and Mg carbonates and subsequent transport of these cations downslope explains this pattern of spatial variability. At Kougarok, concentrations of Ca, Mg, and Sr similarly increase from upper backslope to footslope, but concentrations of Ca and Sr decrease further down the lower backslope (Stations 10 and 1), while Mg concentrations continue to increase. A possible explanation for this behaviour is the greater affinity of cation exchange surfaces for Ca and Sr compared to Mg, thus, Ca and Sr are preferentially retained in the footslope whilst Mg is transported further down the lower backslope (Sparks, 2003, p. 189).

### 3.3.4 Mineral Solubility Effects

Although redox reactions are rarely at equilibrium in natural environments, comparison of field data with equilibrium models provides valuable semi-quantitative insight into the redox condition of natural environments. Because Fe appeared to be limiting the development of more reducing conditions (Section 4.3), select samples from both sites were measured for soluble  $\text{Fe}^{2+}$  following methods presented in Viollier et al. (2000). These concentrations of aqueous  $\text{Fe}^{2+}$  were then compared with model-predicted concentrations of  $\text{Fe}^{2+}$ , when coupled with an infinite  $\text{Fe}(\text{OH})_{3(\text{am})}$  phase, across a range of pH values (2 – 10) and fixed  $E_{\text{H}}$  values of 400 mV, 200 mV, 0 mV, and -200 mV; activity coefficients were assumed to be equal to 1. The measured and modelled  $\text{Fe}^{2+}$  concentrations are compared in **Figure 6**, where concentrations that were below the method detection limit (0.05 mg·L<sup>-1</sup>) are set equal to 0.025 mg·L<sup>-1</sup> (half the detection limit). Comparison of model predicted  $\text{Fe}^{2+}$  concentrations with field data suggests that while Teller exhibits a narrower range of pH conditions than Kougarok, it exhibits a broader range of redox conditions (**Figure 6**). Although several  $\text{Fe}^{2+}$  measurements were below the detection limit, suggesting oxidizing conditions, high  $\text{Fe}^{2+}$  concentrations in some samples suggested  $E_{\text{H}}$  values below 0 mV. Therefore, Fe redox conditions at Teller ranged from mildly reducing to oxic and Fe redox conditions at Kougarok ranged from mildly oxic to oxic. Oxidation-reduction potentials (ORPs), calculated from pH,  $\text{Fe}^{2+}$  concentrations, and the Nernst equation suggest that ORPs at Teller were as low as -69 mV, while the lowest ORP at Kougarok was +134 mV (**Figure 6**). Maximum ORP values could not be determined quantitatively as some  $\text{Fe}^{2+}$  concentrations were below  $\text{Fe}^{2+}$  detection limits, at both sites.  $E_{\text{H}}$ /pH predominance diagrams were created from the 25th, 50th, and 100th concentration percentiles and are shown in **Figure 7** for the COIs where precipitation of mineral phases were predicted under some conditions. The concentrations for these diagrams were taken from filtered aqueous concentration data, thus, predicted mineral precipitation is an indication of nearly saturated or over-saturated conditions. The range of  $E_{\text{H}}$  and pH conditions observed at Teller and Kougarok are overlaid as solid yellow and solid blue lines, respectively. Only the predominance diagrams that indicated possible mineral formation under the  $E_{\text{H}}$ /pH conditions present at either site are shown in **Figure 7**. These phases included  $\text{Fe}(\text{OH})_{3(\text{am})}$  (Fe), siderite (Fe),  $\text{Al}(\text{OH})_{3(\text{am})}$  (Al), chalcedony (Si), barite (Ba and

395 SO<sub>4</sub>), calcite (Ca), dolomite (Ca and Mg), and rhodochrosite (Mn). Predominance diagrams for the remaining key  
396 COIs that were not predicted to form any mineral phases under any site conditions are given in **Supplementary Figure**  
397 **4**.

398 To further examine which mineral phases could be controlling SPW solute concentrations, saturated conditions for  
399 the mineral phases identified in **Figure 7** were modelled using sweeps of pH values from 2 – 10 at various fixed E<sub>H</sub>  
400 values (400mV, 200mV, 0mV, and -200mV). Predicted solute concentrations under the modelled saturated conditions  
401 were then compared with field data to find common trends. In general, if solute concentrations were frequently  
402 measured near the saturation of a mineral, or were identified to have similar dependence on pH or E<sub>H</sub>, it was inferred  
403 that the mineral phase could be controlling the generation of that solute. The mineral phases that were identified to  
404 possibly be controlling solute concentrations were Al(OH)<sub>3(am)</sub>, Fe(OH)<sub>3(am)</sub>, chalcidony, and barite. This does not  
405 preclude the presence of significant concentrations of other mineral phases, it only identifies these as possibly  
406 controlling the dissolved concentrations of Al, Fe, Si, and Ba, respectively. Although it does not provide mineralogical  
407 information, X-ray fluorescence (XRF) data reported by another study at Teller confirmed high concentrations of Al,  
408 Fe, Si, and Ba in the organic and mineral soil layers at that site (Graham et al., 2018). We are unaware of any similar  
409 studies at Kougarok, nor are we aware of any studies that provide would provide confirmatory mineralogical  
410 information, for example by X-ray diffraction (XRD).

411 Aluminium concentrations in SPWs at both Teller and Kougarok appear to be controlled by the  
412 dissolution/precipitation of amorphous Al hydroxide (Al(OH)<sub>3(am)</sub>) (**Figure 8**). The solubility limit of Al(OH)<sub>3(am)</sub> has  
413 no redox dependence, but is highly pH dependent. Aluminium concentrations were generally clustered near the  
414 solubility limit of Al(OH)<sub>3(am)</sub>;  $\text{Al(OH)}_{3(\text{am})} + 3\text{H}^+ \leftrightarrow \text{Al}^{3+} + 3\text{H}_2\text{O}$ ;  $\log k = 10.8$ . This suggests that Al SPW  
415 concentrations at both sites are controlled by wetting/drying (dissolution/precipitation) processes. It also suggests that  
416 there could be a significant amount of Al(OH)<sub>3(am)</sub> in the soils at both sites. While organic matter may also sorb to  
417 alumina surfaces, the adherence to the solubility of Al(OH)<sub>3(am)</sub> suggests that significant concentrations of Al are not  
418 complexed with dissolved organic matter. The predominance diagrams highlight 1) the strong pH dependence on the  
419 stability of Al(OH)<sub>3(am)</sub>, 2) the influence of dissolved F can have on Al speciation when Al concentrations are low,  
420 and 3) that Al is a cation at low pH and an anion at high pH (**Figure 7**). Despite being a weathering product, Al  
421 concentrations show a dissimilar downslope trend to other weathering products, especially at Teller (**Supplementary**  
422 **Figure 5**). While the concentrations of weathering products generally increase with distance downslope, Al  
423 concentrations decrease. We suspect this can be attributed to increasing pH with distance downslope. Philben et al.  
424 (2020) reported a 1 pH unit increase in pH in organic soils along the Teller transect (**Figure 2**), increasing from 5.6 at  
425 Station 5 to 6.7 at Station 9. Such an increase would decrease the solubility of Al(OH)<sub>3(am)</sub>, and thus, decrease the  
426 concentration of dissolved Al (**Figure 8**).

427 Similar to Al, Fe concentrations in SPWs at both Teller and Kougarok appear to be controlled by the  
428 dissolution/precipitation of amorphous Fe hydroxide (Fe(OH)<sub>3(am)</sub>). Fe concentrations were generally clustered near  
429 the solubility limit of Fe(OH)<sub>3(am)</sub> (**Figure 8**). Unlike Al(OH)<sub>3(am)</sub> however, Fe(OH)<sub>3(am)</sub> solubility is dependent on the  
430 redox condition in addition to the pH;  $\text{Fe(OH)}_{3(\text{am})} + 3\text{H}^+ + \text{e}^- \leftrightarrow \text{Fe}^{2+} + 3\text{H}_2\text{O}$ ;  $\log k = 16.0$  (**Figure 8**). Fe(III) is only  
431 sparingly soluble in aqueous solutions and reduction to Fe(II) significantly increases the solubility of Fe, thus, at a

432 given pH value higher aqueous concentrations are predicted and observed under more reducing conditions (**Figure 8**).  
433 Iron concentrations in SPWs at both sites generally follow the pH dependence of  $\text{Fe}(\text{OH})_{3(\text{am})}$  solubility (**Figure 8**).  
434 This suggests that SPW concentrations of Fe at both sites are controlled by wetting/drying (dissolution/precipitation)  
435 processes, coupled with the redox condition.  
436 Si concentrations are frequently limited by the solubility of chalcedony, a very finely grained form of  $\text{SiO}_2$ , which is  
437 much more soluble than quartz;  $\text{SiO}_2 + 2\text{H}_2\text{O} \leftrightarrow \text{H}_4\text{SiO}_4$ ;  $\log k = -3.55$ . Particularly at Kougarok, the dissolved Si  
438 concentrations, coupled with a lack of a strong pH or  $E_{\text{H}}$  dependence, suggest a controlling influence of chalcedony.  
439 Ba concentrations also appear to be controlled by solubility, but rather than by the solubility of an oxide or a hydroxide  
440 phase, by the solubility of barite [ $\text{Ba}^{2+} + \text{SO}_4^{2-} \leftrightarrow \text{BaSO}_4(\text{s})$ ;  $\log k = 9.97$ ]. Unlike Al hydroxide or Fe hydroxide, barite  
441 solubility lacks a strong pH dependence and instead is dependent solely on the activities of  $\text{Ba}^{2+}$  and  $\text{SO}_4^{2-}$ . Unlike Ba,  
442  $\text{SO}_4^{2-}$  concentrations are not limited by the solubility limit of barite and are generally higher and not well correlated  
443 with Ba concentrations. Together, these suggest that  $\text{SO}_4^{2-}$  from another source (likely, atmospheric deposition or  
444 sulfidic mineral oxidation), is suppressing barite dissolution, and thus, is reducing dissolved Ba concentrations. Barite  
445 solubility can exhibit a redox dependence if conditions are sufficiently reducing to reduce  $\text{SO}_4^{2-}$  to sulphide. This shifts  
446 the equilibrium to greater dissolution of barite, and therefore higher conditions of Ba. The lack of  $E_{\text{H}}$  dependence in  
447 observational data further suggests that neither site exhibits significant  $\text{SO}_4^{2-}$  reduction.

#### 448 **4. Discussion**

449 The 18 stations examined herein (8 at Teller and 10 at Kougarok) represent a wide range of vegetation types, soil  
450 moisture contents, permafrost extents, and hillslope positions. Coupling the spatial variability of these landscape  
451 characteristics with the spatial variability of SPW solute concentrations provides valuable insight into the dominant  
452 environmental controls on observed spatial variability of SPW geochemistry. It is our hope that correlating SPW  
453 geochemistry with readily observable and scalable landscape features will inform earth system modelling efforts in  
454 permafrost regions and provide fast and easy methods to determine if earth system models are working properly (i.e.  
455 predicting the correct trends). The inferred dominant environmental controls on the observed inter-site and intra-site  
456 variability of SPW solute concentrations are discussed in the following sections.

#### 457 **4.1 The Dominant Environmental Controls on Inter-site Variability of SPW Solute Concentrations**

458 Overall, the more frequent instance of significantly greater constituent concentrations at Kougarok suggests a  
459 systematic cause. The extensive low-gradient toeslope (**Figure 2**) and lack of a well-defined drainage channel at  
460 Kougarok, are likely causes of the systematically higher SPW solute concentrations at Kougarok. Water perching, the  
461 result of near-surface permafrost in the lower-backslope and toeslope, increases evapotranspiration and, thus, SPW  
462 solute concentrations. Meanwhile, the lack of a drainage channel at Kougarok suggests that runoff (and therefore  
463 solute exports) is more limited than at Teller. Without a relatively rapid export mechanism such as a stream channel,  
464 solute transport is likely limited to interflow within the Kougarok hillslope over much of the thaw season, allowing  
465 weathering products to increase to significantly greater concentrations than those observed at Teller, where a well-

466 defined drainage/export mechanism does exist. Field observations from pits at Kougarok confirm observable interflow  
467 at the site. Overall, our study suggests that evaporative concentration could be a significant control on SPW solute  
468 concentrations in permafrost catchments, especially in those with limited drainage and therefore a perched near-  
469 surface water table. This effect has been reported previously (Raudina et al., 2017), but does not appear to be widely  
470 considered, perhaps due to the generally few studies of SPW solutes in permafrost regions. We suggest future efforts  
471 to predict future SPW solute and nutrient dynamics directly address the impacts of evaporative concentration on  
472 permafrost catchments, especially with future permafrost thaw.

473 The exception to the general observation of elevated concentrations at Kougarok versus Teller was  $\text{SO}_4^{2-}$ . Although  
474 the cause of consistently higher  $\text{SO}_4^{2-}$  concentrations at Teller is unclear from the limited scope of this study, it seems  
475 likely to be due to a greater abundance of sulfidic bedrock material. The presence of sulfidic bedrock in the vicinity  
476 of Teller has been reported by mineral prospecting efforts (Brobst, Pinckney, and Sainsbury, 1971; Herreid, 1966;  
477 Mulligan, 1965); we are unaware of any such reports near Kougarok.

#### 478 **4.2 The Dominant Environmental Controls on Intra-site Spatial Variability of SPW Solute Concentrations**

479 Vegetation influences on elemental cycles were only readily apparent for nitrogen and although vegetation induced  
480 changes to soil moisture content were discernible, they were far less significant than anticipated.  $\text{NO}_3^-$  was the only  
481 COI that exhibited a clear vegetation effect; elevated concentrations were associated with the presence of alder shrubs  
482 and, in some cases, tall willow shrubs. These increases in  $\text{NO}_3^-$  concentrations associated with alder nitrogen-fixation  
483 and the mineralization and nitrification of willow leaf litter were frequently equivoiced by increased microbial  
484 denitrification in regions sufficiently moist to support it, this is perhaps one of the most significant finding of this  
485 work. Although both Kougarok and Teller exhibited some indications of increased Cl concentrations in the presence  
486 of tall shrubs, the net vegetation effect on soil moisture was far less than hypothesized. Redox sensitivity was also less  
487 than hypothesized and most stations seemed well-buffered at Fe redox conditions. The result of this buffering was  
488 generally low  $\text{NO}_3^-$  concentrations (except where vegetation effects dominated), consistent  $\text{SO}_4^{2-}$  concentrations across  
489 clear redox gradients, and variable Mn and Fe concentrations. Mn concentrations were generally low, likely due to a  
490 limited source. Fe concentrations were higher at stations with higher soil moisture content, consistent with Fe  
491 reduction. Similar Fe redox cycling between soluble Fe(II) species and precipitated Fe oxyhydroxides in permafrost  
492 catchments has been reported recently (Patzner et al., 2022), which suggests that Fe redox buffering in permafrost  
493 landscapes is widespread. Weathering, water/soil interactions, and hydrological transport were probable drivers of  
494 variability for Ca, Sr, and Mg. Ca, Sr, and Mg all tended to accumulate in low-lying areas, although Ca and Sr  
495 demonstrated greater accumulation potential than Mg, likely via greater affinity of cation exchange surfaces for Ca  
496 and Sr compared to Mg. Mineral solubility limitations were the primary controls on Al ( $\text{Al}(\text{OH})_{3(\text{am})}$ ), Fe ( $\text{Fe}(\text{OH})_{3(\text{am})}$ ),  
497 Ba (barite), and Si (chalcedony) concentrations. This suggests that the SPW concentrations of these constituents will  
498 remain stable until those mineral phases are exhausted or soil pore hydrochemistry changes sufficiently to alter the  
499 solubility of those mineral phases. Supersaturation of Al with respect to gibbsite (crystalline  $\text{Al}(\text{OH})_3$ ) and Si with  
500 respect to chalcedony in a permafrost wetland has been reported previously (Jesson et al., 2014). The solubility curves  
501 for gibbsite and  $\text{Al}(\text{OH})_{3(\text{am})}$  are similar, with  $\text{Al}(\text{OH})_{3(\text{am})}$  being slightly more soluble at all pH values do to the

502 increased thermodynamic stability of the crystalline Al hydroxide mineral, gibbsite. Meanwhile, seasonal precipitation  
503 of Fe oxyhydroxides in permafrost peatlands and their effect of carbon cycling was the subject of an excellent paper  
504 by Patzner et al. (2022). Our study is the first observation we are aware of the saturation controls of barite on Ba in  
505 permafrost SPWs, although that could be because few relatively few studies consider barium concentrations; it is  
506 worthwhile noting the Ba was not supersaturated with respect to barite but approached a saturated condition. Future  
507 studies should also note that changes in redox condition would significantly alter  $\text{Fe}(\text{OH})_{3(\text{am})}$  solubility, whereas  
508 changes in pH conditions would significantly alter  $\text{Al}(\text{OH})_{3(\text{am})}$  and  $\text{Fe}(\text{OH})_{3(\text{am})}$  solubility.  
509 Although discerning the environmental controls on spatial variability of SPW solute concentrations provides some  
510 high-level insight into the effects changes in landscape character may have on soil pore hydrochemistry, our scope  
511 was limited and leveraged on previously available datasets. The significance of SPW in small Arctic headwater  
512 catchments as a key initial component in the freshwater hydrologic continuum is under recognized, and such  
513 catchments warrant more detailed and systematic investigations.

## 514 **5. Acknowledgements**

515 We would like to thank the Sitnasuak Native Corporation and the Mary's Igloo Native Corporation for their guidance  
516 and for allowing us to conduct this research on the traditional homelands of the Inupiat people. Funding was provided  
517 by the Next-Generation Ecosystem Experiments (NGEE Arctic) project, supported by the Office of Biological and  
518 Environmental Research in the U.S. DOE Office of Science. We wish to thank Lauren Charsley-Groffman and Nathan  
519 Wales for their assistance with fieldwork, as well as, George Perkins, Oana Marina, Rose Harris, and Emily Kluk for  
520 their assistance with laboratory analyses.

## 521 **6. Data availability statement**

522 The data that support the findings of this study are made openly available in the NGEE-Arctic data repository at (DOI:  
523 10.5440/1735757).

## 524 **7. References**

- 525 Anderson, M. D., Ruess, R. W., Myrold, D. D., and Taylor, D. L. (2009). Host species and habitat affect nodulation  
526 by specific *Frankia* genotypes in two species of *Alnus* in interior Alaska. *Oecologia* 160, 619–630. doi:  
527 10.1007/s00442-009-1330-0
- 528 Anderson, M. D., Ruess, R. W., Uliassi, D. D., and Mitchell, J. S. (2004). Estimating N<sub>2</sub> fixation in two species of  
529 *Alnus* in interior Alaska using acetylene reduction and <sup>15</sup>N<sub>2</sub> uptake. *Ecoscience* 11, 102–112. doi: 10.1080/  
530 11956860.2004.11682814
- 531 Binkley, D., Sollins, P., Bell, R., Sachs, D., and Myrold, D.: Biogeochemistry of adjacent conifer and alder-conifer  
532 stands, *Ecology*, 73, 2022–2033, 1992.

533 Breen, A., Iversen, C., Salmon, V., VanderStel, H., Busey, B., and Wulschleger, S. 2020a.: NGEE Arctic Plant Traits:  
534 Plant Community Composition, Kougarok Road Mile Marker 64, Seward Peninsula, Alaska, 2016 [Data set], doi:  
535 <https://doi.org/10.5440/1465967>.

536 Breen, A., Iversen, C., Salmon, V., VanderStel, H., Busey, B., and Wulschleger, S. 2020b.: NGEE Arctic Plant Traits:  
537 Plant Community Composition, Kougarok Road Mile Marker 64, Seward Peninsula, Alaska, 2016 [Data set], doi:  
538 <https://doi.org/10.5440/1465967>.

539 Bring, A., Fedorova, I., Dibike, Y., Hinzman, L., Mård, J., Mernild, S. H., ... Woo, M.-K. 2016: Arctic terrestrial  
540 hydrology: A synthesis of processes, regional effects, and research challenges. *Journal of Geophysical Research:*  
541 *Biogeosciences*, 121: 621–649, doi: <https://doi.org/10.1002/2015JG003131>.

542 Brobst, D. A., Pinckney, D. M., and Sainsbury, C. L. 1971: *Geology and Geochemistry of the Sinuk River Barite*  
543 *Deposit, Seward Peninsula, Alaska (No. 463 (?)). United States Department of the Interior Geological Survey.*

544 Bühlmann, T., Hiltbrunner, E., and Körner, C.: *Alnus viridis* expansion contributes to excess reactive nitrogen release,  
545 reduces biodiversity and constrains forest succession in the Alps, *Alpine Botany*, 124, 187–191,  
546 <https://doi.org/10.1007/s00035-014-0134-y>, 2014.

547 Clein, J. S. and Schimel, J. P.: Nitrogen turnover and availability during succession from alder to poplar in Alaskan  
548 taiga forests, *Soil Biology and Biochemistry*, 27, 743–752, [https://doi.org/10.1016/0038-0717\(94\)00232-P](https://doi.org/10.1016/0038-0717(94)00232-P), 1995.

549 Conroy, N., Heikoop, J., Newman, B., Wilson, C., Arendt, C., Perkins, G., and Wulschleger, S. 2021: *Soil Water*  
550 *Chemistry and Water and Nitrogen Isotopes, Teller Road Site and Kougarok Hillslope, Seward Peninsula, Alaska,*  
551 *2016 - 2019 [Data set], doi: <https://doi.org/10.5440/1735757>.*

552 Corder, G. W., and Foreman, D. I. 2009: *Nonparametric statistics for non-statisticians: a step-by-step approach.*  
553 *Hoboken, N.J.: Wiley, 247 pp.*

554 Frey, K. E., and McClelland, J. W. 2009: Impacts of permafrost degradation on arctic river biogeochemistry.  
555 *Hydrological Processes*, 23: 169–182, doi: <https://doi.org/10.1002/hyp.7196>.

556 Frisbee, M. D., Phillips, F. M., Campbell, A. R., and Hendrickx, J. M. H. 2010: Modified passive capillary samplers  
557 for collecting samples of snowmelt infiltration for stable isotope analysis in remote, seasonally inaccessible  
558 watersheds 1: laboratory evaluation. *Hydrological Processes*, 24: 825–833, doi: <https://doi.org/10.1002/hyp.7523>.

559 Fuchs, M., Nitze, I., Strauss, J., Günther, F., Wetterich, S., Kizyakov, A., ... Grosse, G. 2020: Rapid Fluvio-Thermal  
560 Erosion of a Yedoma Permafrost Cliff in the Lena River Delta. *Frontiers in Earth Science*, 8: 336, doi:  
561 <https://doi.org/10.3389/feart.2020.00336>.

562 Graham, D. E., Kholodov, A., Wilson, C. J., Moon, J.-W., Romanovsky, V. E., and Busey, B. 2018: *Soil Physical,*  
563 *Chemical, and Thermal Characterization, Teller Road Site, Seward Peninsula, Alaska, 2016., doi:*  
564 <https://doi.org/10.5440/1342956>.

565 Harms, T. K., and Jones, J. B. 2012: Thaw depth determines reaction and transport of inorganic nitrogen in valley  
566 bottom permafrost soils. *Global Change Biology*, 18: 2958–2968, doi:  
567 <https://doi.org/10.1111/j.13652486.2012.02731.x>.

568 Harms, T. K., and Ludwig, S. M. 2016: Retention and removal of nitrogen and phosphorus in saturated soils of arctic  
569 hillslopes. *Biogeochemistry*, 127: 291–304, doi: <https://doi.org/10.1007/s10533-016-0181-0>.



570 Helsel, D. R. 2005: Nondetects and data analysis: statistics for censored environmental data. Hoboken, NJ.; Wiley-  
571 Interscience, 250 pp.

572 Herreid, G. 1966: Preliminary geology and geochemistry of the Sinuk River area. Seward Peninsula, Alaska: Alaska  
573 Division of Mines and Minerals Geologic Report, 24: 19.

574 Hiyama, T., Yang, D. and Kane, D.L., 2021. Permafrost Hydrology: Linkages and Feedbacks. In Arctic Hydrology,  
575 Permafrost and Ecosystems (pp. 471-491). Springer, Cham.

576 Hollingsworth, T. N., Lloyd, A. H., Nosssov, D. R., Ruess, R. W., Charlton, B. A., and Kielland, K. (2010). Twenty-  
577 five years of vegetation change along a putative successional chronosequence on the Tanana River, Alaska. *Can. J.*  
578 *For. Res.* 40, 1273–1287. doi: 10.1139/X10-094

579 Hopkins, D. M., Karlstrom, T. N. V., Black, R. F., Williams, J. R., Pewe, T. L., Fernald, A. T., and Muller, E. H.  
580 1955: Permafrost and ground water in Alaska. U.S. Geological Survey Professional Paper 264-F.

581 Jafarov, E. E., Coon, E. T., Harp, D. R., Wilson, C. J., Painter, S. L., Atchley, A. L., and Romanovsky, V. E. 2018:  
582 Modelling the role of preferential snow accumulation in through talik development and hillslope groundwater flow in  
583 a transitional permafrost landscape. *Environmental Research Letters*, 13: 105006, doi: [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/aadd30)  
584 [9326/aadd30](https://doi.org/10.1088/1748-9326/aadd30).

585 Jessen, Søren, Hanne D. Holmslykke, Kristine Rasmussen, Niels Richardt, and Peter E. Holm. 2014. “Hydrology and  
586 Pore Water Chemistry in a Permafrost Wetland, Ilulissat, Greenland.” *Water Resources Research* 50 (6): 4760–74.  
587 <https://doi.org/10.1002/2013WR014376>.

588 Kinniburgh, D., and Cooper, D. 2011: PhreePlot: Creating Graphical Output with Phreeqc.

589 Koch, J. C., Runkel, R. L., Striegl, R., and McKnight, D. M. 2013: Hydrologic controls on the transport and cycling  
590 of carbon and nitrogen in a boreal catchment underlain by continuous permafrost. *Journal of Geophysical Research:*  
591 *Biogeosciences*, 118: 698–712, doi: <https://doi.org/10.1002/jgrg.20058>.

592 Kokelj, S. V., and Jorgenson, M. T. 2013: Advances in Thermokarst Research. *Permafrost and Periglacial Processes*,  
593 24: 108–119, doi: <https://doi.org/10.1002/ppp.1779>.

594 Kurylyk and Walvoord, 2021 Kurylyk, B.L. and Walvoord, M.A., 2021. Permafrost Hydrogeology. In Arctic  
595 Hydrology, Permafrost and Ecosystems (pp. 493-523). Springer, Cham.

596 Langford, Z. L., Kumar, J., Hoffman, F. M., Breen, A. L., & Iversen, C. M. (2019). Arctic vegetation mapping using  
597 unsupervised training datasets and convolutional neural networks. *Remote Sensing*, 11(1), 1–23.  
598 <https://doi.org/10.3390/rs11010069>

599 Lara, M. J., Nitze, I., Grosse, G., and McGuire, A. D. 2018: Tundra landform and vegetation productivity trend maps  
600 for the Arctic Coastal Plain of northern Alaska. *Scientific Data*, 5: 1–10, doi: <https://doi.org/10.1038/sdata.2018.58>.

601 Léger, E., Dafflon, B., Robert, Y., Ulrich, C., Peterson, J. E., Biraud, S. C., ... Hubbard, S. S. 2019: A distributed  
602 temperature profiling method for assessing spatial variability in ground temperatures in a discontinuous permafrost  
603 region of Alaska. *The Cryosphere*, 13: 2853–2867, doi: <https://doi.org/10.5194/tc-13-2853-2019>.

604 Liljedahl, A. K., Boike, J., Daanen, R. P., Fedorov, A. N., Frost, G. V., Grosse, G., ... Zona, D. 2016: Pan-Arctic ice-  
605 wedge degradation in warming permafrost and its influence on tundra hydrology. *Nature Geoscience*, 9: 312–318,  
606 doi: <https://doi.org/10.1038/ngeo2674>.

607 McCaully, R. E., Arendt, C. A., Newman, B. D., Salmon, V. G., Heikoop, J. M., Wilson, C. J., Sevanto, S., Wales, N.  
608 A., Perkins, G. B., Marina, O. C., and Wulfschleger, S. D.: High nitrate variability on an Alaskan permafrost hillslope  
609 dominated by alder shrubs, *The Cryosphere*, 16, 1889–1901, <https://doi.org/10.5194/tc-16-1889-2022>, 2022.

610 McClelland, J. W., Holmes, R. M., Peterson, B. J., Raymond, P. A., Striegl, R. G., Zhulidov, A. V., ... Griffin, C. G.  
611 2016: Particulate organic carbon and nitrogen export from major Arctic rivers. *Global Biogeochemical Cycles*, 30:  
612 629–643, doi: <https://doi.org/10.1002/2015GB005351>.

613 Mitchell, J. S., and Ruess, R. W. (2009). N<sub>2</sub> fixing alder (*Alnus viridis* spp. *fruticosa*) effects on soil properties across  
614 a secondary successional chronosequence in interior Alaska. *Biogeochemistry* 95, 215–229. doi: 10.1007/s10533-009-  
615 9332-x

616 Mulligan, J. J. 1965: Examination of the Sinuk Iron Deposits Seward Peninsula, Alaska. United States Department of  
617 the Interior, 37.

618 Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., ... Hik, D. S. 2011: Shrub  
619 expansion in tundra ecosystems: dynamics, impacts and research priorities. *Environmental Research Letters*, 6:  
620 045509, doi: <https://doi.org/10.1088/1748-9326/6/4/045509>.

621 Nossov, D. R., Hollingsworth, T. N., Ruess, R. W., and Kielland, K. (2011). Development of *Alnus tenuifolia* stands  
622 on an Alaskan floodplain: patterns of recruitment, disease and succession. *J. Ecol.* 99, 621–633. doi: 10.1111/j.1365-  
623 2745.2010.01792.x

624 O'Donnell, J., Douglas, T., Barker, A. and Guo, L., 2021. Changing Biogeochemical Cycles of Organic Carbon,  
625 Nitrogen, Phosphorus, and Trace Elements in Arctic Rivers. In *Arctic Hydrology, Permafrost and Ecosystems* (pp.  
626 315-348). Springer, Cham.

627 Patzner, M.S., Kainz, N., Lundin, E., Barczok, M., Smith, C., Herndon, E., Kinsman-Costello, L., Fischer, S., Straub,  
628 D., Kleindienst, S., Kappler, A., Bryce, C., 2022. Seasonal Fluctuations in Iron Cycling in Thawing Permafrost  
629 Peatlands. *Environ. Sci. Technol.* 56, 4620–4631. <https://doi.org/10.1021/acs.est.1c06937>

630 Parkhurst, D., and Appelo, C. A. J. 2013: Description of input and examples for PHREEQC version 3: a computer  
631 program for speciation, batch-reaction, one-dimensional transport, and inverse geochemical calculations (USGS  
632 Numbered Series No. 6-A43). Reston, VA: U.S. Geological Survey.

633 Perdrial, J. N., Perdrial, N., Vazquez-Ortega, A., Porter, C., Leedy, J., and Chorover, J. 2014: Experimental  
634 Assessment of Passive Capillary Wick Sampler Suitability for Inorganic Soil Solution Constituents. *Soil Science*  
635 *Society of America Journal*, 78: 486–495, doi: <https://doi.org/10.2136/sssaj2013.07.0279>.

636 Petrone, K. C., Hinzman, L. D., Shibata, H., Jones, J. B., and Boone, R. D. 2007: The influence of fire and permafrost  
637 on sub-arctic stream chemistry during storms. *Hydrological Processes*, 21: 423–434, doi:  
638 <https://doi.org/10.1002/hyp.6247>.

639 Philben, M., Taş, N., Chen, H., Wulfschleger, S. D., Kholodov, A., Graham, D. E., and Gu, B. 2020: Influences of  
640 hillslope biogeochemistry on anaerobic soil organic matter decomposition in a tundra watershed. *Journal of*  
641 *Geophysical Research: Biogeosciences*, n/a: e2019JG005512, doi: <https://doi.org/10.1029/2019JG005512>.

642 Philben, M., Zheng, J., Bill, M., Heikoop, J. M., Perkins, G., Yang, Z., ... Gu, B. 2019: Stimulation of anaerobic  
643 organic matter decomposition by subsurface organic N addition in tundra soils. *Soil Biology and Biochemistry*, 130:  
644 195–204, doi: <https://doi.org/10.1016/j.soilbio.2018.12.009>.

645 Prowse, T., Bring, A., Mård, J., and Carmack, E. 2015: Arctic Freshwater Synthesis: Introduction. *Journal of*  
646 *Geophysical Research: Biogeosciences*, 120: 2121–2131, doi: <https://doi.org/10.1002/2015JG003127>.

647 Prowse, T., Bring, A., Mård, J., Carmack, E., Holland, M., Instanes, A., ... Wrona, F. J. 2015: Arctic Freshwater  
648 Synthesis: Summary of key emerging issues. *Journal of Geophysical Research: Biogeosciences*, 120: 1887–1893, doi:  
649 <https://doi.org/10.1002/2015JG003128>.

650 R Core Team. 2020: R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for  
651 Statistical Computing. Retrieved from <https://www.R-project.org/>.

652 Raudina, Tatiana V., Sergey V. Loiko, Artyom G. Lim, Ivan V. Krickov, Liudmila S. Shirokova, Georgy I. Istigechev,  
653 Daria M. Kuzmina, Sergey P. Kulizhsky, Sergey N. Vorobyev, and Oleg S. Pokrovsky. 2017. “Dissolved Organic  
654 Carbon and Major and Trace Elements in Peat Porewater of Sporadic, Discontinuous, and Continuous Permafrost  
655 Zones of Western Siberia.” *Biogeosciences* 14 (14): 3561–84. <https://doi.org/10.5194/bg-14-3561-2017>.

656 Reynolds, M.K.; Walker, D.A.; Maier, H.A. Plant community-level mapping of arctic Alaska based on the  
657 Circumpolar Arctic Vegetation Map. *Phytocoenologia* 2005, 35, 821–848.

658 Reynolds, M. K., Walker, D. A., Balser, A., Bay, C., Campbell, M., Cherosov, M. M., et al. (2019). A raster version  
659 of the Circumpolar Arctic Vegetation Map (CAVM). *Remote Sensing of Environment*, 232.  
660 <https://doi.org/10.1016/j.rse.2019.111297>

661 Romanovsky, V., Cable, W., and Dolgikh, K. 2020a:. Soil Temperature and Moisture, Kougarak Road Mile Marker  
662 64, Seward Peninsula, Alaska, beginning 2016 [Data set], doi: <https://doi.org/10.5440/1581586>.

663 Romanovsky, V., Cable, W., and Dolgikh, K. 2020b:. Soil Temperature and Moisture, Kougarak Road Mile Marker  
664 64, Seward Peninsula, Alaska, beginning 2016 [Data set], doi: <https://doi.org/10.5440/1581586>.

665 Romanovsky, V., Cable, W., and Dolgikh, K. 2020c:. Soil Temperature and Moisture, Teller Road Mile Marker 27,  
666 Seward Peninsula, Alaska, beginning 2016 [Data set], doi: <https://doi.org/10.5440/1581437>.

667 Romanovsky, V., Cable, W., and Dolgikh, K. 2020d:. Soil Temperature and Moisture, Teller Road Mile Marker 27,  
668 Seward Peninsula, Alaska, beginning 2016 [Data set], doi: <https://doi.org/10.5440/1581437>.

669 Rowland, J. C., Jones, C. E., Altmann, G., Bryan, R., Crosby, B. T., Hinzman, L. D., ... Geernaert, G. L. 2010: Arctic  
670 Landscapes in Transition: Responses to Thawing Permafrost. *Eos, Transactions American Geophysical Union*, 91:  
671 229–230, doi: <https://doi.org/10.1029/2010EO260001>.

672 Ruess, R. W., Anderson, M. D., Mcfarland, J. M., Kielland, K., Olson, K., and Taylor, D. L. (2013). Ecosystem-level  
673 consequences of symbiont partnerships in an N-fixing shrub from interior Alaskan floodplains. *Ecol. Monogr.* 83,  
674 177– 194. doi: 10.1890/12-0782.1

675 Salmon, V. G., Breen, A. L., Kumar, J., Lara, M. J., Thornton, P. E., Wulschleger, S. D., and Iversen, C. M. 2019:  
676 Alder Distribution and Expansion Across a Tundra Hillslope: Implications for Local N Cycling. *Frontiers in Plant*  
677 *Science*, 10, doi: <https://doi.org/10.3389/fpls.2019.01099>.

678 Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., ... Vonk, J. E. 2015: Climate  
679 change and the permafrost carbon feedback. *Nature*, 520: 171–179, doi: <https://doi.org/10.1038/nature14338>.

680 Shaver, G. R., Billings, W. D., Chapin, F. S., Giblin, A. E., Nadelhoffer, K. J., Oechel, W. C., and Rastetter, E. B.  
681 1992: Global Change and the Carbon Balance of Arctic Ecosystems. *BioScience*, 42: 433–441, doi:  
682 <https://doi.org/10.2307/1311862>.

683 Shogren, A. J., Zarnetske, J. P., Abbott, B. W., Iannucci, F., Frei, R. J., Griffin, N. A., and Bowden, W. B. 2019:  
684 Revealing biogeochemical signatures of Arctic landscapes with river chemistry. *Scientific Reports*, 9: 1–11, doi:  
685 <https://doi.org/10.1038/s41598-019-49296-6>.

686 Smith, L. C., Sheng, Y., MacDonald, G. M., and Hinzman, L. D. 2005: Disappearing Arctic Lakes. *Science*, 308:  
687 1429–1429, doi: <https://doi.org/10.1126/science.1108142>.

688 Sparks, D. L. 2003: *Environmental soil chemistry* (2nd ed). Amsterdam ; Boston:, Academic Press, 352 pp.

689 Spence, C., Kokelj, S., McCluskie, M., and Hedstrom, N. 2015: Impacts of Hydrological and Biogeochemical Process  
690 Synchrony Transcend Scale. In *AGU Fall Meeting Abstracts* (Vol. 2015).

691 Sturm, M., Racine, C., and Tape, K. 2001: Increasing shrub abundance in the Arctic. *Nature*, 411: 546–547, doi:  
692 <https://doi.org/10.1038/35079180>.

693 Sulman, B. N., Salmon, V. G., Iversen, C. M., Breen, A. L., Yuan, F., & Thornton, P. E. (2021). Integrating arctic  
694 plant functional types in a land surface model using above- and belowground field observations. *Journal of Advances*  
695 *in Modeling Earth Systems*, 13, e2020MS002396. <https://doi.org/10.1029/2020MS002396>

696 Tape, K. D., Hallinger, M., Welker, J. M., and Ruess, R. W. 2012: Landscape Heterogeneity of Shrub Expansion in  
697 Arctic Alaska. *Ecosystems*, 15: 711–724, doi: <https://doi.org/10.1007/s10021-012-9540-4>.

698 Tape, K., Sturm, M., and Racine, C. 2006: The evidence for shrub expansion in Northern Alaska and the Pan-Arctic.  
699 *Global Change Biology*, 12: 686–702, doi: <https://doi.org/10.1111/j.1365-2486.2006.01128.x>.

700 Till, A. B., Dumoulin, J. A., Weldon, M. B., and Bleick, H. A. 2011: Bedrock geologic map of the Seward Peninsula,  
701 Alaska, and accompanying conodont data. US Department of the Interior, US Geological Survey.

702 Uren, N. C. 2018: Calcium oxalate in soils, its origins and fate – a review. *Soil Research*, 56: 443, doi:  
703 <https://doi.org/10.1071/SR17244>.

704 Vonk, J. E., Tank, S. E., Bowden, W. B., Laurion, I., Vincent, W. F., Alekseychik, P., ... Wickland, K. P. 2015:  
705 Reviews and syntheses: Effects of permafrost thaw on Arctic aquatic ecosystems. *Biogeosciences*, 12: 7129–7167,  
706 doi: <https://doi.org/10.5194/bg-12-7129-2015>.

707 Vonk, J. E., Tank, S. E., and Walvoord, M. A. 2019: Integrating hydrology and biogeochemistry across frozen  
708 landscapes. *Nature Communications*, 10: 1–4, doi: <https://doi.org/10.1038/s41467-019-13361-5>.

709 Walker, D. A., Breen, A. L., Druckenmiller, L. A., Wirth, L. W., Fisher, W., Reynolds, M. K., Sibik, J., Walker, M.  
710 D., Hennekens, S., Boggs, K., Boucher, T., Buchhorn, M., Bultmann, H., Cooper, D. J., Daniels, F. J. A., Davidson,  
711 S. J., Ebersole, J. J., Elmendorf, S. C., Epstein, H. E., Gould, W. A., Hollister, R. D., Iversen, C. M., Jorgenson, M.  
712 T., Kade, A., Lee, M. T., MacKenzie, W. H., Peet, R. K., Peirce, J. L., Schickhoff, U., Sloan, V. L., Talbot, S. S.,  
713 Tweedie, C. E., Villarreal, S., Webber, P. J., and Zona, D.: The Alaska Arctic Vegetation Archive (AVA-AK),  
714 *Phytocoenologia*, 46, 221–229, <https://doi.org/10.1127/phyto/2016/0128>, 2016.

715 Wallenberger, F. T., and Bingham, P. A. 2009: *Fiberglass and Glass Technology: Energy-Friendly Compositions and*  
716 *Applications*. Springer Science & Business Media, 479 pp.

717 Walvoord, M. A., and Kurylyk, B. L. 2016: Hydrologic Impacts of Thawing Permafrost—A Review. *Vadose Zone*  
718 *Journal*, 15, doi: <https://doi.org/10.2136/vzj2016.01.0010>.

719 Weiss, M., Hobbie, S. E., & Gettel, G. M. (2005). Contrasting Responses of Nitrogen-Fixation in Arctic Lichens to  
720 Experimental and Ambient Nitrogen and Phosphorus Availability. *Arctic, Antarctic, and Alpine Research*, 37(3), 396–  
721 401. [https://doi.org/10.1657/1523-0430\(2005\)037\[0396:CRONIA\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0396:CRONIA]2.0.CO;2)

722 Wilson, C., Bolton, R., Busey, R., Lathrop, E., and Dann, J. 2019: End-of-Winter Snow Depth, Temperature, Density  
723 and SWE Measurements at Kougarok Road Site, Seward Peninsula, Alaska, 2018 [Data set], doi:  
724 <https://doi.org/10.5440/1593874>.

725 Wilson, C., Bolton, R., Busey, R., Lathrop, E., Dann, J., and Charsley-Groffman, L. 2019: End-of-Winter Snow Depth,  
726 Temperature, Density and SWE Measurements at Teller Road Site, Seward Peninsula, Alaska, 2016-2018 [Data set],  
727 doi: <https://doi.org/10.5440/1592103>.

728 Wilson, C., Dann, J., Bolton, R., Charsley-Groffman, L., Jafarov, E., Musa, D., and Wullschleger, S. 2021: In Situ  
729 Soil Moisture and Thaw Depth Measurements Coincident with Airborne SAR Data Collections, Barrow and Seward  
730 Peninsulas, Alaska, 2017 [Data set], doi: <https://doi.org/10.5440/1423892>.


731 Wrona, F. J., Johansson, M., Culp, J. M., Jenkins, A., Mård, J., Myers-Smith, I. H., ... Wookey, P. A. 2016: Transitions  
732 in Arctic ecosystems: Ecological implications of a changing hydrological regime. *Journal of Geophysical Research:*  
733 *Biogeosciences*, 121: 650–674, doi: <https://doi.org/10.1002/2015JG003133>.

734 Wullschleger, S. D., Epstein, H. E., Box, E. O., Euskirchen, E. S., Goswami, S., Iversen, C. M., et al. (2014). Plant  
735 functional types in earth system models: past experiences and future directions for application of dynamic vegetation  
736 models in high-latitude ecosystems. *Ann. Bot.* 114, 1–16. doi: 10.1093/aob/mcu077

737 Yang, D., Meng, R., Morrison, B. D., McMahon, A., Hantson, W., Hayes, D. J., ... Serbin, S. P. 2020: A Multi-Sensor  
738 Unoccupied Aerial System Improves Characterization of Vegetation Composition and Canopy Properties in the Arctic  
739 Tundra. *Remote Sensing*, 12: 2638, doi: <https://doi.org/10.3390/rs12162638>.

740 Yang, D., et al. "Landscape-Scale Characterization of Arctic Tundra Vegetation Composition, Structure, and Function  
741 with a Multi-Sensor Unoccupied Aerial System." *Environmental Research Letters* 16 (8), 085005 (2021).  
742 <https://doi.org/10.1088/1748-9326/ac1291>.


743



745  
746 **Figure 1. Location of the Teller and Kougarak field sites with respect to the municipalities of Teller, Nome, and Council.**  
747 **All are located on the Seward Peninsula in northwestern Alaska. RGB composite imagery from the 8-band WorldView-2**  
748 **imagery obtained on July 14, 2017 at 1.5 m resolution downloaded from the DigitalGlobe website**  
749 **(<https://www.digitalglobe.com/>).**

750






751

752 Figure 2. Topographic map of Teller. Station areas are shown as red polygons and the topographic station transect is given  
 753 as a solid blue line. The hillslope transect elevation profile is given below the map in green, with stations along the transect  
 754 in blue and hillslope positions noted with red arrows and text. RGB composite imagery from the 8-band WorldView-2  
 755 imagery obtained on July 27, 2011 at 1.5 m resolution downloaded from the DigitalGlobe website  
 756 (<https://www.digitalglobe.com/>).





758

759


760

761

762

Figure 3. Topographic map of Kougarok. Station areas are shown as red polygons and the station transect is given as a solid blue line. The transect elevation profile is given below the map in green, with stations along the transect in yellow and hillslope positions noted with red arrows and text. RGB composite imagery from the 8-band WorldView-2 imagery obtained on July 14, 2017 at 1.5 m resolution downloaded from the DigitalGlobe website (<https://www.digitalglobe.com/>).





764

765



766

767

768


769

**Figure 4. Mean COI concentrations at Teller (blue) and Kougarak (yellow) stations. Stations are arranged (left to right) by soil moisture content determined by P-Band SAR (top right). Boxplots show the first, second, and third data quartiles, with box whiskers representing either 150% of the inner quartile range (IQR), or the maximum or minimum value, when that value was less than 1.5×IQR. Red circles represent data points outside of the 1.5×IQR whiskers (i.e. outliers). Note that the concentration scales on the Teller and Kougarak plots often differ.**



772 Figure 5. Median (50<sup>th</sup> percentile) concentrations (grey diamonds with dashed black lines) of Ca, Sr, and Mg, with distance  
 773 downslope at Teller (blue) and Kougarak (yellow) along topographic transects; areas of stations are indicated by blue and  
 774 yellow colouring, respectively. The elevation profiles of the hillslopes are plotted in green, on separate y-axes (right axes).  
 775 Topographic regions of both catchments are indicated by red arrows along the elevation gradient.


776



777

778 Figure 6. Left: Model-predicted Fe<sup>2+</sup> concentrations in saturated solutions of Fe(OH)<sub>3(am)</sub> at fixed E<sub>H</sub> conditions of 400 mV  
779 (green), 200 mV (orange), 0 mV (grey), and -200 mV (red), compared with field concentrations of Fe<sup>2+</sup> at Teller (red circles)  
780 and Kougarok (yellow circles). Right: Fe predominance diagram, showing the dominant specie of Fe under a range of  
781 E<sub>H</sub>/pH conditions. E<sub>H</sub>/pH regions relevant to Teller and Kougarok are outlined in blue and yellow, respectively. Samples  
782 with Fe<sup>2+</sup> concentrations below the detection limit are given as colour coordinated open circles set at 0.025 mg·L<sup>-1</sup> (half the  
783 detection limit) in both sides of the figure.

784



785

786

787


788

789

790

Figure 7.  $E_H/pH$  diagrams for key species that indicated possible mineral formation under the  $E_H/pH$  conditions present at either Teller or Kougarok. The  $E_H$  and pH conditions observed at Teller and Kougarok are overlaid as blue and yellow lines, respectively. Mineral species (solids) are shown in grey, cations are shown in blue, anions are shown in purple, neutral species are shown in white. Predominance diagrams were created in PhreePlot using the phreeqc.dat database, with inorganic carbonate reduction to methane “turned off.”

791



792

793

794

**Figure 8. Modelled solute concentrations in solutions saturated with  $\text{Al(OH)}_{3(\text{am})}$ ,  $\text{Fe(OH)}_{3(\text{am})}$ , chalcedony, and barite, with respect to pH (x-axis) and  $E_H$  (model lines), overlaid with observed solute concentrations.**

795

797 Table 1. Teller Station Physical Characteristics

	Hillslope Position	Vegetation					Relative wetness			Permafrost		
		Vegetation type	Average (maximum) canopy height (cm)	Dominant PFT		Low to tall shrub cover	Average TDR soil moisture (VMC)	Average P-band SAR (VMC)	Average snow depth (cm)	Average Ground Temperature (°C)	Permafrost Extent	Average (maximum) thaw depth (cm)
TL9	Lower Footslope	Wetland complex	28 (41)	Bryophyte	44 %	10 %	NA	0.46	68.4	0	Marginal	101 (>120)
TL5	Upper Shoulder		12 (45)	Graminoid	45 %	7 %	0.55	0.37	103.3	-0.45	Near-surface	97 <sup>r</sup> (>114 <sup>r</sup> )
TL8	Upper Footslope		7 (34)	Bryophyte	33 %	20 %	0.55	0.36	77.7	-0.6		69 <sup>r</sup> (>120)
TL3	Upper Backslope	Cassiope dwarf shrub tundra	9 (23)	Evergreen dwarf shrub	47 %	12 %	NA	0.25	62.1	2.2	None/ deep	72 <sup>r</sup> (82 <sup>r</sup> )
TL4	Upper Footslope		8 (14)		58 %	4 %	0.35	0.39	89.5	0.5	Marginal	40 <sup>r</sup> (70 <sup>r</sup> )
TL2	Upper Backslope	Mesic willow shrubland	84 (141)	Deciduous low to tall shrub (willow)	44 %	44 %	0.4	0.34	124	2.4	None/ deep	75 <sup>r</sup> (>120)
TL7	Lower Shoulder		151 (189)		37 %	37 %	0.46	0.26	128.8	2.4		51 <sup>r</sup> (66 <sup>r</sup> )
TL6	Upper Backslope	Willow-birch tundra	64 (115)	Forb	23 %	32 %	0.38	0.34	86.4	1.2	None/ deep	67 <sup>r</sup> (102 <sup>r</sup> )

798

799 PFT – plant functional type; dwarf shrub (height <40 cm), low shrub (height 40-200 cm), low to tall shrub (height 40 to >200 cm tall).  
800 Deciduous shrub PFT classes identify the dominant species in the plant community as either willow or willow and birch. There is no alder at  
801 the Teller site. Low to tall shrub cover represents the sum of deciduous low shrubs, deciduous low to potentially tall willow and birch, and  
802 deciduous low to tall alder.

803 <sup>1</sup>Single point soil moisture measurements. Data are more accurate than P-band SAR but represent a much smaller spatial scale.

804 <sup>2</sup>P-band SAR has 30m resolution.

805 <sup>r</sup>Resistive layer was rock; all others are permafrost. A temperature probe was used to determine if the resistive layer was permafrost ( $\leq 0$  °C)  
806 or rock ( $> 2$  °C). Thaw depth is an average of 4 measurements from the vegetation plot corners within the IS and was measured at the end of  
807 the growing season.

808

809 **Table 2. Kougarok Station Physical Characteristics**

	<i>Hillslope Position</i>	<i>Vegetation</i>					<i>Relative wetness</i>			<i>Permafrost</i>		
		Vegetation type	Average (maximum) canopy height (cm)	Dominant PFT		Low to tall shrub cover	Average TDR soil moisture (VMC)	Average P-band SAR (VMC)	Average snow depth (cm)	Average Ground Temperature (°C)	Permafrost Extent	Average (maximum) thaw depth (cm)
<i>KG3</i>	Upper Backslope	Alder shrubland	204 (265)	Deciduous low to tall shrub (alder)	30 %	53 %	0.19	0.39	131.3	-0.01	Near-surface	48 <sup>†</sup> (53 <sup>r</sup> )
<i>KG12</i>	Footslope		NA	NA	NA	NA	0.30*	0.39	NA	NA		NA
<i>KG1</i>	Lower Backslope	Alder savanna in tussock tundra	60 (90)	Deciduous low shrub	31 %	44 %	NA	0.51	83.4	-2.5	Near-surface	61 (68)
<i>KG2</i>	Footslope		48 (73)	Graminoid	30 %	42 %	0.63	0.52	102.3	-1.2		75 (89)
<i>KG6</i>	Lower Backslope		24 (61)	Graminoid	46 %	17 %	0.36	0.48	66.2	-2.2		58 (62)
<i>KG10</i>	Lower Backslope		NA	NA	NA	NA	NA*	0.44	71.4	NA		NA
<i>KG11</i>	Footslope		NA	NA	NA	NA	0.59*	0.42	NA	NA		NA
<i>KG7</i>	Upper Backslope	Tussock-lichen tundra	20 (22)	Graminoid	34 %	14 %	0.51	0.45	54.7	-2.1	Near-surface	76 (100)
<i>KG4</i>	Shoulder	Dryas-lichen shrub tundra	6 (12)	Evergreen dwarf shrub	62 %	1 %	NA	0.37	NA	-1.9	Near-surface	0 <sup>†</sup> (0 <sup>†</sup> )
<i>KG13</i>	Upper Backslope		NA	NA	NA	NA	0.41*	0.39	92.1	NA		NA
<i>KG5</i>	Upper Backslope	Willow-birch tundra	62 (137)	Deciduous low shrub	60 %	62 %	NA	0.4	178.4	> 0	Deep	88 (96)
<i>KG8</i>	Upper Backslope		45 (120)	Evergreen dwarf shrub	52 %	42 %	0.23	0.24	85.5	-0.04	Near-surface	44 <sup>†</sup> (55 <sup>†</sup> )

810  
 811 Note: PFT – plant functional type. Deciduous shrub PFT classes identify the dominant species in the community as either willow, alder,  
 812 willow and birch, or alder, willow, and birch. Low to tall shrub cover represents the sum of deciduous low shrubs, deciduous low to potentially  
 813 tall willow and birch, and deciduous low to tall alder.  
 814 <sup>1</sup>Single point soil moisture measurements. Data are more accurate than P-band SAR but represent a much smaller spatial scale.  
 815 <sup>2</sup>P-band SAR has 30m resolution.  
 816 \*Average gravimetric water content measurements, corrected to VMC by bulk density.  
 817 <sup>†</sup>Resistive layer was rock; all others are permafrost. A temperature probe was used to determine if the resistive layer was permafrost ( $\leq 0$  °C)  
 818 or rock ( $>2$  °C). Thaw depth is an average of 4 measurements from the vegetation plot corners within the IS and was measured at the end of  
 819 the growing season.

Table 3. Inter-Site Mann-Whitney U-Test Results

	<u>Teller</u>			<u>Kougarok</u>			z	Site with Higher Median	Effect Size	Difference in Correlation
	n	$\sum R_i$	$U_i$	n	$\sum R_i$	$U_i$				
Na	59	3184	14811.5	275	52761.5	1413.5	<b>9.95</b>	Kougarok	0.54	large
F	59	3502	14375.5	273	51776.5	1731.5	<b>9.46</b>	Kougarok	0.52	large
K	59	3882	14113	275	52063	2112	<b>8.92</b>	Kougarok	0.49	medium-large
Si	59	4119	13876.5	275	51826.5	2348.5	<b>8.56</b>	Kougarok	0.47	medium-large
Al	58	4952	12709	275	50659	3241	<b>7.11</b>	Kougarok	0.39	medium
Oxalate	57	4996	12161.5	272	49289.5	3342.5	<b>6.75</b>	Kougarok	0.37	medium
B	59	5429	12566.5	275	50516.5	3658.5	<b>6.62</b>	Kougarok	0.36	medium
Zn	58	5605	12056	275	50006	3894	<b>6.12</b>	Kougarok	0.34	medium
SO <sub>4</sub>	58	13653	3892.5	273	41293.5	11941.5	<b>6.08</b>	Teller	0.33	medium
Fe	58	5958	11703	275	49653	4247	<b>5.60</b>	Kougarok	0.31	medium
Ba	58	6256	11405.5	275	49355.5	4544.5	<b>5.15</b>	Kougarok	0.28	medium
Ti	58	6266	11395.5	275	49345.5	4554.5	<b>5.13</b>	Kougarok	0.28	medium
NO <sub>2</sub>	54	5588	10585.5	272	47713.5	4102.5	<b>5.12</b>	Kougarok	0.28	medium
Li	58	7778	9883	275	47833	6067	<b>2.86</b>	Kougarok	0.16	small-medium
Br	58	8485	9060.5	273	46461.5	6773.5	1.73	Equal	0.09	small
NO <sub>3</sub>	58	8576	8969	273	46370	6865	1.59	Kougarok	0.09	small
Sr	58	8683	8978	275	46928	6972	1.51	Kougarok	0.08	small
PO <sub>4</sub>	54	9659	6460.5	271	43316.5	8173.5	1.36	Equal	0.08	small
Mg	58	10495	7166	275	45116	8784	1.21	Teller	0.07	small
Cr	58	8884	8777	275	46727	7173	1.20	Kougarok	0.07	small
Mn	58	9164	8497	275	46447	7453	0.78	Teller	0.04	small
Cl	58	9221	8266.5	272	45394.5	7509.5	0.57	Kougarok	0.03	small
Ca	58	10016	7645	275	45595	8305	0.50	Teller	0.03	small

Table 4. Dominant Environmental Controls on SPW Geochemistry at Teller and Kougarok

<b>Environmental Control</b>	<b>Analytes Affected</b>
Vegetation	NO <sub>3</sub> <sup>-</sup>
Soil Moisture/Redox	NO <sub>3</sub> <sup>-</sup> , Mn, Fe, SO <sub>4</sub> (occasionally)
Water/Soil Interactions & Hydrologic Transport	Ca, Mg, Sr
Mineral Solubility	Al, Ba, Si, Fe