



## Revealing the chemical characteristics of Arctic low-level cloud residuals – in situ observations from a mountain site

Yvette Gramlich<sup>1,2</sup>, Karolina Siegel<sup>1,2,3</sup>, Sophie L. Haslett<sup>1,2</sup>, Gabriel Freitas<sup>1,2</sup>, Radovan Krejci<sup>1,2</sup>, Paul Zieger<sup>1,2</sup>, and Claudia Mohr<sup>1,2</sup>

<sup>1</sup>Department of Environmental Science, Stockholm University, Stockholm, Sweden

<sup>2</sup>Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

<sup>3</sup>Department of Meteorology, Stockholm University, Stockholm, Sweden

Correspondence to: Claudia Mohr (claudia.mohr@aces.su.se)

10

**Abstract.** The role aerosol chemical composition plays in Arctic low-level cloud formation is still poorly understood. In this study we address this issue by combining in situ observations of the chemical characteristics of cloud residuals (dried liquid cloud droplets or ice crystals) and aerosol particles from the Zeppelin Observatory in Ny-Ålesund, Svalbard (approx. 480 m a. s. l.). These measurements were part of the one-year long Ny-Ålesund Aerosol and Cloud Experiment 2019-2020 (NASCENT). To obtain the chemical composition of cloud residuals at molecular level, we deployed a Filter Inlet for Gases and AEROsols coupled to a Chemical Ionization Mass Spectrometer (FIGAERO-CIMS) with iodide as the reagent ion behind a Ground-based Counterflow Virtual Impactor (GCVI). The station was enshrouded in clouds roughly 15 % of the time during NASCENT, out of which we analyzed 14 cloud events between December 2019 and December 2020. During the entire year, the composition of the cloud residuals shows contributions from oxygenated organic compounds, including organonitrates, and traces of the biomass burning tracer levoglucosan. In summer, methanesulfonic acid (MSA), an oxidation product of dimethyl sulfide (DMS), shows large contributions to the sampled mass, indicating marine natural sources of cloud condensation nuclei (CCN) and ice nucleating particles (INP) mass during the sunlit part of the year. In addition, we also find contributions of the inorganic acids nitric and sulfuric acid, with outstanding high absolute signals of sulfuric acid in one cloud residual sample in spring and one in late summer (May 21 and September 12, 2020), probably caused by high anthropogenic sulfur emissions near the Barents and Kara Sea. During one particular cloud event, on May 18, 2020, the air mass origin did not change from before to during and after the cloud. We therefore chose it as a case study to investigate cloud impact on aerosol physicochemical properties. We show that the overall chemical composition of the organic aerosol particles is similar before, during, and after the cloud, indicating that the particles have already undergone one or several cycles of cloud processing before being measured as residuals at Zeppelin, and/or that on the timescales of the observed cloud event, cloud processing of the organic fraction can be neglected. Meanwhile there are on average fewer particles, but relatively more in the accumulation mode after the cloud. Comparing the signal of sulfur-containing compounds of cloud residuals with aerosols during cloud-free conditions, we find that sulfuric acid has a higher relative contribution to the cloud residuals compared to aerosols during cloud-free conditions, but we did not observe an increase in particulate MSA due to the cloud. Overall, the chemical composition, especially of the organic fraction of the Arctic cloud residuals, reflects the overall composition of the general aerosol population well. Our results thus suggest that most aerosols can serve as seeds for low-level clouds in the Arctic.

30

35

### 1 Introduction

Aerosol particles interact with solar radiation either directly by light scattering and/or absorption, or indirectly by acting as cloud seeds (Haywood and Boucher, 2000). On an annual scale, Arctic clouds have an overall warming effect on the surface

40



(Shupe and Intrieri, 2004). The formation of a cloud particle requires the availability of sufficient water vapor and updraft to create supersaturated conditions, and aerosol particles that provide a surface for the water vapor to condense onto. This subfraction of aerosol particles is termed cloud condensation nuclei (CCN) or ice nucleating particles (INP). The physical and chemical properties of aerosols play an important role for describing aerosol-cloud interactions (Köhler, 1936; Seinfeld and Pandis, 2006). Better knowledge on their composition, especially of the subfraction activating as cloud droplets, helps to better constrain their effect in the warming Arctic, in addition to an improved understanding of their sources.

While in most of the regions on Earth water vapor is the limiting factor, in the Arctic the availability of aerosol particles able to act as CCN or INP can be limited (Mauritsen et al., 2011). Due to this limit in aerosol number concentrations, aerosol particles in the Aitken mode (diameter < 80 nm), even as small as 20 nm in diameter, are able to serve as CCN in the Arctic (Korhonen et al., 2008; Leaitch et al., 2016; Bulatovic et al., 2021; Pöhlker et al., 2021; Karlsson et al., 2021, 2022; Siegel et al., 2022).

Overall, the Arctic aerosol particle composition follows a distinct annual cycle, where long-range atmospheric transport dominates in winter and spring, and local sources increase in importance during the summer. One specific phenomenon in the annual aerosol cycle is called “Arctic Haze”, occurring in late winter and spring, when enhanced mass concentrations of aerosol particles mainly composed of long-range transported sulfate, particulate organic matter, and heavy metals, but also black carbon, nitrate, ammonium and dust (Quinn et al., 2007) are observed. Among the natural sources, the ocean plays an important role. Aerosol particles emitted by sea spray can comprise sea salt and organic material from the sea surface microlayer (Cavalli et al., 2004; Kirpes et al., 2019). Sea salt is the largest contributor to particulate matter by mass across the Arctic. Its relative contribution to total particulate matter has been shown to be higher in the dark period compared to the bright season (Moschos et al., 2022a). In addition to open ocean, the origin of sea salt has been attributed to blowing snow, especially during the dark season at high wind speeds over sea ice, e.g. in the central Arctic (Huang and Jaeglé, 2017). Model simulations from a decade ago predicted the sea salt emissions to increase with less sea ice cover (Struthers et al., 2011). An increasing trend in sea spray emissions was found recently at the Arctic monitoring station Ny-Ålesund on Svalbard, (Heslin-Rees et al., 2020); however, this trend was linked to changes in the circulation pattern rather than decreasing sea ice cover.

Also, the organic fraction of Arctic aerosol is dominated by anthropogenic sources in winter, and natural emissions in summer. The primary source region for wintertime anthropogenic aerosol is Eurasia (Moschos et al., 2022b). In summer, the growth of aerosol particles has been associated with the presence of methanesulfonic acid (MSA), produced from the oxidation of dimethylsulfide (DMS) released by marine phytoplankton, and trimethylamine ( $N(CH_3)_3$ ) and other organic compounds (Willis et al., 2016; Beck et al., 2021). The highest particulate MSA concentrations in Ny-Ålesund were found to occur in May or June, when phytoplankton biomass is active in the surrounding Greenland and Barents Sea (Jang et al., 2021).

The presence of various particulate organic and sulfate-containing compounds was recently reported for the summertime Arctic Ocean using offline filter analysis (Siegel et al., 2021). These observations were among the first molecular-level measurements of semi-volatile aerosols from the high Arctic. Organic molecules with carbon chains of up to 18 carbon atoms were identified. The largest signal was observed for compounds with 5 to 10 carbon atoms and 3 to 4 oxygen atoms. Semi-volatile organics were found to be involved in aerosol particle growth during summer in the marine Arctic environment (Burkart et al., 2017). Although new information about the chemical composition of Arctic aerosol is emerging, very little is known yet about the composition of Arctic cloud residuals (dried liquid cloud droplets or ice crystals), from which information on the chemical composition of CCN and INP, aqueous-phase processing, or condensation of gaseous compounds (co-condensation, (Topping and McFiggans, 2012; Topping et al., 2013)) can be derived. Direct observations are scarce (Sect. S1, Fig. S1) and if available limited to intensive campaigns over just a couple of weeks; hence, there is no information on how the properties of cloud seeds change throughout the year (McFarquhar et al., 2011; Wendisch et al., 2019). Aircraft observations in spring 2008 near Barrow (Alaska, ISDAC campaign) showed that compared to ambient aerosols, cloud residuals contained relatively less organics and more sea salt and black carbon (Hiranuma et al., 2013). However, no details on the organic molecular composition was



reported. In addition, the cloud residuals were generally larger than the ambient aerosol particles (Zelenyuk et al., 2010;  
85 Hiranuma et al., 2013). In a case study during ISDAC, looking at the change in chemical composition of single particles at  
different altitudes in a cloud, enriched sulfate was observed in the cloud residuals compared to the aerosol population below  
the cloud (Zelenyuk et al., 2010). Wendisch et al. (2019) measured cloud residuals by aircraft over Svalbard during the  
transition from spring to summer (end of May until beginning of June in 2017). They observed trimethylamine and sulfate in  
the cloud residuals, and to a lesser extent metals, organic carbon, and levoglucosan, a tracer for biomass burning. Higher levels  
90 of trimethylamine were found in cloud residuals sampled over sea ice, compared to over open ocean and drift ice. One very  
recent study investigated the composition of cloud residuals in Ny-Ålesund based on 4 years of observations (Adachi et al.,  
2022). They focused on the fraction of sea salt, mineral dust, sulfate, K-bearing and carbonaceous material-containing particles  
and found that the cloud residuals have the same composition as ambient aerosol particles at positive temperatures, while at  
negative temperatures the residuals contained more mineral dust and sea salt compared to the ambient aerosol.  
95 To the best of our knowledge, this study presents the first molecular-level observations of the chemical composition of Arctic  
cloud residuals. We identify organic and inorganic compounds in cloud residuals in the Arctic region using in situ  
measurements from a Chemical Ionization high-resolution time-of-flight Mass Spectrometer (CIMS) coupled to a Filter Inlet  
for Gases and AEROsols (FIGAERO), referred to as the FIGAERO-CIMS, set up behind a Ground-based Counterflow Virtual  
Impactor (GCVI) for a full year at the Zeppelin Observatory, Ny-Ålesund, Svalbard (Platt et al., 2022). We investigate the  
100 changes in chemical composition of cloud residuals during different parts of the year, and of the aerosol population before,  
during and after one particular cloud event.

## 2 Methods

This section gives an overview of the instrumentation deployed during NASCENT for this study, and the data processing  
procedures used to determine the chemical characteristics of cloud residuals. Further information on the campaign and  
105 additional instrumentation used during NASCENT can be found in Pasquier et al. (2022).

### 2.1 NASCENT campaign

The data presented here were acquired during the one-year long Ny-Ålesund Aerosol and Cloud Experiment 2019-2020  
(NASCENT). In brief, the aim of the campaign was to determine the physical and chemical properties of trace gases, aerosol  
110 and cloud particles in high detail with state-of-the-art instrumentation over the course of a full year. During the campaign,  
measurements were conducted at different locations and altitudes near Ny-Ålesund, including the Zeppelin Observatory. In  
this study we only focus on the measurements conducted at the Zeppelin Observatory.

### 2.2 Zeppelin Observatory

The Zeppelin Observatory near Ny-Ålesund, Svalbard (78°54'N 11°53'E), is one of a few permanent measurement stations in  
115 the Arctic. The observatory itself is located about 2 km south of the small research settlement in Ny-Ålesund, on Mt. Zeppelin  
at an altitude of 474 m a.s.l., and is equipped with instrumentation to continuously measure atmospheric trace gases, particles  
and other atmospheric properties. The measurement station was established in 1989 and is now part of several monitoring  
programs as a global background station, e.g. the Global Atmosphere Watch Programme (GAW) (Platt et al., 2022). The  
remote location of the observatory is characterized by the 26 km long Kongsfjorden located towards the north and east, and  
120 mountainous landscape with glaciers in the south and west. During the NASCENT year, low-level mixed phase clouds were  
present in all seasons, covering between 20 % to around 40 % of the monthly cloud cases (Pasquier et al., 2022). The Zeppelin  
Observatory itself is frequently covered in clouds (between October 2019 and December 2020 about 15 % of the time (visibility




< 1000 m for at least 5 min)), providing an ideal location to study the physicochemical properties of Arctic clouds and aerosol-cloud interactions.

### 125 2.3 Instrumental setup during NASCENT including GCVI

The instruments used in this study at the Zeppelin Observatory were connected through a three-way switching valve (Fig. 1) to a whole-air inlet during cloud-free conditions and to a GCVI (Brechtel Manufacturing Inc., USA, Model 1205) inlet during cloudy conditions. The inlet height was approx. 480 m.a.s.l.. With the whole-air inlet, both interstitial (non-activated) aerosol particles and cloud droplets  $\sim < 40 \mu\text{m}$  were collected, whereas the GCVI inlet only sampled cloud droplets and ice crystals > 6 to 7  $\mu\text{m}$  (aerodynamic diameter). In brief, the working principle of the GCVI is as follows: within a wind tunnel, the cloud particles are accelerated onto the GCVI tip, where an opposing air flow is generated such that only those particles with high enough inertia make it into the sampling flow, thereby removing the interstitial aerosol and particles  $< 6\text{-}7 \mu\text{m}$  (Ogren et al., 1985; Noone et al., 1988). A more detailed description and evaluation of the inlet system, the GCVI and additional instrumentation used to characterize microphysical aerosol and cloud residual properties is available in a previous study (Karlsson et al., 2021).

During the NASCENT campaign, the GCVI was partly operated in manual mode and partly in automatic mode. Manual mode means that the GCVI was manually turned on in the presence of a cloud. As an indication of cloudy conditions, a visibility sensor was used. Similar to Karlsson et al. (2021) we used a threshold of 1000 m, where visibility below this threshold meant presence of a cloud. In automatic mode the GCVI was automatically turned on whenever the visibility at the observatory was equal or less than 1000 m.



145 **Figure 1:** Setup of the FIGAERO-CIMS and the DMPS at the Zeppelin Observatory. A three-way-valve (3-way-v) defines the type of particles sampled. With the whole-air inlet, all particles (referred to as total particles, i.e. both aerosol particles and cloud particles) are sampled, whereas the GCVI inlet only samples cloud particles. Upstream of the FIGAERO-CIMS particle-phase inlet, a second filter (b-filter) was in place to perform regular particle-phase blanks. DMPS2 consists of two separate DMPS systems measuring partly overlapping size ranges. These two size ranges have been harmonized to one dataset in this study. Modified version of Fig. 1 from Karlsson et al. (2021), where DMPS2 consists of DMPS 2a and DMPS 2b.



## 150 2.4 FIGAERO-CIMS

The chemical composition of cloud residuals at molecular level was obtained from a FIGAERO-CIMS (Aerodyne Research Inc., USA) (Lopez-Hilfiker et al., 2014; Thornton et al., 2020) using iodide (I<sup>-</sup>) as reagent ion. The FIGAERO-CIMS was installed in October 2019, and data is available almost continuously from November 2019 until December 2020, with a larger gap in July and August 2020.

155 The FIGAERO-CIMS was installed inside one of the rooms at the Zeppelin Observatory and connected to two separate inlet lines (Fig. 1). The particle-phase inlet was connected via ½ inch stainless steel tubing to the three-way valve of the main inlet switching between the whole-air inlet and the GCVI inlet. The gas-phase inlet was connected via ¼ PTFE tubing to ambient air directly through a hole in the wall.

The FIGAERO automatically cycled between its two modes: gas-phase analysis with simultaneous particle deposition, and 160 thermal desorption and analysis of the deposited particulate matter. These two modes are hereafter referred to as the sampling and heating periods, respectively. The sampling period with simultaneous particle deposition was 2.5 h. The particles were sampled with a flow of 4 LPM. During the gas-phase measurements, zero air was introduced every 40 min for 15 min, to obtain the gas-phase background signal.

After the sampling period, the particle-loaden filter was automatically moved to the desorption position. A 2 LPM flow of 165 heated ultrapure nitrogen (from a Nitrogen-generator, Peak Scientific NG5000) was passed through the filter to thermally desorb the particles. This heating period was divided in three phases of 20 min each: ramping, soaking, and cooling. During ramping, the temperature of the nitrogen flow passing through the filter was gradually increased from room temperature to approx. 200°C. During the soaking phase, the heated nitrogen flow was held at this temperature to ensure complete evaporation of the deposited material. During the cooling phase, the nitrogen flow was cooled back down to room temperature. The signal 170 obtained during the heating procedure (start of the ramping until end of the soaking) are hereafter referred to as thermograms. Particle-phase background samples, also called blanks, were collected using a filter upstream of the sampling filter (Fig. 1). Automatically switching valves allowed sample air to either bypass or pass through the zeroing filter (blank). Every third sampling period was a blank.

The FIGAERO-CIMS data was acquired at 1 s time resolution until February 2020 and at 2 s until the end of the campaign, 175 and pre-averaged to 30 s for analysis. To account for the instrument's sensitivity on the water vapor pressure in the ion-molecule reaction region, the data was normalized to the sum of iodide (I<sup>-</sup>; mass-to-charge ratio (m/z): 126.905) and the iodide-water-cluster (H<sub>2</sub>OI<sup>-</sup>; m/z: 144.916) (Lee et al., 2014). Since dry nitrogen (N<sub>2</sub>) is used to vaporize the particulate compounds during the heating period, accounting for the iodide-water-cluster signal is not relevant for the particle-phase data. Using iodide as reagent ion allows for mainly the detection of polar and oxygenated compounds (e.g. carboxylic acids), of which we can 180 obtain the molecular composition up to an m/z of around 600 Th (Lee et al., 2014).

Since the cloud residual samples and blanks of this study were distributed over the entire year with long periods of cloud-free conditions and/or instrumental breaks in between, we treated the signals as if they were offline filter samples following the procedures detailed by Cai et al., 2022. In order to take varying instrument backgrounds into account, for each compound, the signal of the blank thermogram was scaled by the signal ratio of blank and corresponding sample thermogram at the end of 185 the soaking period (last 3-6 datapoints/last ~1.5 to 3 min, Method 2b in Cai et al., 2022, Sect. S2, Fig. S2). After this step, the thermograms of the sample and the scaled blank were integrated from the beginning of the ramping period until the end of the soaking period. To obtain the total signal, the integrated thermogram signal from the scaled blank was subtracted from the integrated thermogram signal of the sample. The integration of the thermograms resulted in the total number of detected ions, which is also the unit of the data reported here.

190 To obtain the ambient aerosol composition (sampled via the whole-air inlet) of the two samples before and after the cloud for the case study of May 18, 2020, we used the neighboring ambient blank and subtracted the integrated thermogram blank from



the integrated thermogram sample. We did not use the scaling approach here, as the instrument background of neighboring heatings usually does not differ substantially.

## 2.5 DMPS

195 The particle number size distribution at the Zeppelin Observatory is continuously measured with a Differential Mobility Particle Sizer (DMPS) since 2000 (Ström et al., 2003; Tunved et al., 2013). In total there are three DMPS systems installed at the observatory, one, similar to the FIGAERO-CIMS, behind a three-way valve switching between the whole-air and GCVI inlets (DMPS1, installed in November 2015), and two behind the whole-air inlet. The two DMPS systems behind the whole-air inlet measure partly overlapping size ranges, which is why the data of these two were harmonized and are used hereafter  
200 as one size range (DMPS2, Fig. 1). DMPS1 measured cloud residuals during cloudy conditions (when the GCVI was on) and ambient aerosol during cloud-free conditions (when the GCVI was off) in the size range of 10 to 945 nm. DMPS2 measured all ambient particles in the size range of 5 to 708 nm. More technical specifications of both DMPS systems can be found in Karlsson et al. (2021).

The number of cloud particles sampled with the GCVI is higher compared to the actual ambient cloud particle concentration.  
205 This enrichment of particles can be corrected for by calculating an enrichment factor (EF), which is determined by the settings of the GCVI (sampling flow and airspeed in the wind tunnel) and its geometry (Shingler et al., 2012). We used the median EF ( $EF_{med}$ ) of the respective sample to correct the DMPS1 data during times when we sampled cloud residuals. For the entire dataset discussed here, the  $EF_{med}$  was in the range of 6.8 and 19.5 (mean 11.1, median 10.5, Table 1).

In addition, a sampling efficiency ( $eff$ ) needs to be determined for particle number concentrations sampled behind the GCVI.  
210  $eff$  depends on the cloud particle number size distribution and has been experimentally calculated for our site to be 0.46 (Karlsson et al., 2021). All the cloud residual number size distributions in this study were corrected by  $k$  (Eq. 1). No other particle loss calculations were applied. The data for the number size distributions of some samples was filtered for potential droplet splashing according to the procedure described in the supplementary (Sect. S3).

For the conversion from number to mass concentrations, we used a density of  $1.3 \text{ g cm}^{-3}$ , representing secondary organic  
215 aerosol (e.g. Alfarra et al., 2006; Malloy et al., 2009). The mass-based measurements (e.g. FIGAERO-CIMS) were not corrected, as we only show relative changes when comparing in cloud with out of cloud conditions, and no ambient concentrations when comparing the cloud residual samples. All given sizes in this study refer to the diameter ( $D$ ).

$$k = \frac{1}{eff * EF_{med}} \quad (\text{Eq. 1})$$

220

## 2.6 Back trajectories, meteorological data and cloud target classification

Air mass back trajectories were obtained using HYSPLIT (Stein et al., 2015), starting at 474 m a.s.l. at the Zeppelin Observatory (3-hourly archive data from the Global Data Assimilation System (GDAS), operated by the National Centers for Environmental Prediction (NCEP), 1 degree horizontal grid resolution). The back trajectories were calculated for 10 days, out  
225 of which the most recent 5 days were used here. Data within and above the boundary layer were used. The hourly temperature ( $T$ ) and relative humidity (RH) data was downloaded from EBAS (<https://ebas-data.nilu.no>). The cloud target classification for Ny-Ålesund (Nomokonova et al., 2019) was taken from Cloudnet (<https://cloudnet.fmi.fi/>).

230



**Table 1: Cloud residual samples and the respective sampling start and end date and time of the FIGAERO-CIMS collection period, and corresponding median visibility, mean temperature (T) and median enrichment factor ( $EF_{med}$ ) during the sampling time.**

Cloud residual name	Sampling start	Sampling end	Median visibility (m)	Mean T (°C)	$EF_{med}$
Dec 25, 2019	2019-12-25 16:42:29	2019-12-25 19:12:50	233.9	-13.7	6.8
May 18, 2020	2020-05-18 07:32:54	2020-05-18 10:03:02	48.8	-3.1	9.7
May 21, 2020	2020-05-21 12:47:44	2020-05-21 15:17:46	244.2	-2.4	10.2
Jun 2, 2020	2020-06-02 14:08:20	2020-06-02 16:38:22	135.3	-0.5	10.6
Jun 12-1, 2020	2020-06-12 18:20:20	2020-06-12 20:50:22	76.5	-0.8	10.2
Jun 12-2, 2020	2020-06-12 21:50:23	2020-06-13 00:20:25	265.3	-1.9	10.6
Jun 14, 2020	2020-06-14 01:50:04	2020-06-14 04:20:06	162.9	-1.9	10.3
Jun 25, 2020	2020-06-25 16:59:09	2020-06-25 19:29:11	87.7	4.2	9.6
Jun 26, 2020	2020-06-26 17:29:11	2020-06-26 19:58:44	41.4	-0.3	10.7
Jun 27-1, 2020	2020-06-27 00:28:53	2020-06-27 02:58:55	101.1	-0.5	12.0
Jun 27-2, 2020	2020-06-27 03:58:56	2020-06-27 06:28:58	105.0	-1.1	12.3
Sep 12, 2020	2020-09-12 02:31:29	2020-09-12 05:01:39	168.9	1.2	9.3
Oct 28, 2020	2020-10-28 20:38:37	2020-10-28 23:09:01	407.6	-9.8	19.5
Dec 9, 2020	2020-12-09 10:18:37	2020-12-09 12:49:04	432.6	-4.2	14.1

235

## 2.7 Definition of cloud residuals and selection of samples

In analogy to the previous study by Karlsson et al., 2021, we here define cloud residuals operationally as the particles remaining after drying the cloud droplets and ice crystals collected with the GCVI. This is to clarify that with our instrumental setup, we do not measure CCN or INP directly. Even though cloud residuals are often termed as CCN or INP in the literature, this is not entirely correct, as coalescence or chemical reactions may occur inside the cloud particles and thereby modify the original cloud nuclei (Twohy and Anderson, 2008).

As mentioned above (Sect. 2.3), the GCVI was not operated automatically throughout the entire duration of the campaign. As a consequence, there are times when the visibility at the observatory was above 1000 m, but the GCVI was still on.



Following from the above we selected the FIGAERO-CIMS cloud residual samples according to the following criteria:

- 245
- The inlet was set to GCVI during the entire 2.5 h sampling time of the FIGAERO-CIMS, and GCVI flows were fully operational for at least 95 % percent of the sampling time.
  - The median visibility during the 2.5 h sampling time was below 1000 m.

Following these criteria, we are left with in total 14 cloud residual samples and 10 cloud residual blanks from the NASCENT campaign. An overview of the cloud residual samples and the respective ambient conditions during the sampling times are  
250 listed in Table 1.

### 3 Results and Discussion


#### 3.1 Seasonal distribution of cloud residuals characterized by FIGAERO-CIMS

Figure 2 shows the monthly number of cloud residual samples and related blanks measured with the FIGAERO-CIMS during NASCENT, as well as the corresponding monthly total number of hours with visibility below 1000 m, ambient temperature  
255 and RH. The monthly average temperature during the year-long campaign was lowest in March (-19°C) and highest in July (7°C) (Fig. 2a). Based on the temperature, most of our cloud residual samples are probably originating from mixed phase clouds. Compared to the meteorological average (1994-2018), the winter months of NASCENT were around 6 K colder, and the summer slightly warmer (around 2 K, Pasquier et al., 2022). Monthly average RH was lowest in December 2019 and January 2020 (67 %), and highest in June and July (91 %, Fig. 2b).

260 Overall, the cloud occurrence at the measurement station was higher from February to September than during the winter months, as observed previously (Chernokulsky et al., 2017; Dekhtyareva et al., 2018; Gierens et al., 2020). March and June were the two months with the most hours of cloudy conditions at the observatory (Fig. 2c). Unfortunately, during that period, the GCVI was run mostly in manual mode, and therefore not always following the visibility threshold criterion. The sampling time with the GCVI inlet was highest in June, resulting in total eight cloud residual samples and six blanks. In May we collected  
265 two samples and two blanks, and in September only one cloud residual sample and one blank. During October 2020 and December 2019, 2020, we could only measure one cloud residual sample in each month. Although the haze season (March-May) showed cloudy conditions (19 % of time), we did not measure cloud residual samples in this period due to problems with the instrumentation (in only 15 % of the cloudy conditions the GCVI was on).

In the following we will focus first on the number size distribution of ambient and cloud particles during NASCENT before  
270 we discuss their chemical composition in more detail.





275 **Figure 2: Atmospheric conditions during the entire year of NASCENT measurement, and number of cloud residual samples. (a) Monthly mean and median temperature, (b) monthly mean and median relative humidity. The grey shaded area represents the standard deviation. (c) Monthly total hours of visibility below 1000 m for at least 5 min (solid line), and corresponding total hours of sampling with the GCVI inlet (GCVI ON, dashed line), as well as the number of FIGAERO-CIMS cloud residual samples (sample) and blanks (blank) collected in the individual months.**

### 3.2 Number size distributions of cloud residuals during NASCENT

280 Karlsson et al. (2021) analyzed 2 years of cloud residual and corresponding total particle (cloud residuals and non-activated particles) sizes and number concentrations at the Zeppelin Observatory. They found that cloud residuals and aerosol particles have a similar annual cycle with respect to number, with highest concentrations in spring and summer, and a minimum in fall and winter. In the winter months (Dec-Feb) they observed a dominating fraction of Aitken mode particles ( $D < 80$  nm) in the cloud residuals, while during Arctic haze until summer (Apr-Jun) the cloud residuals were most frequently found in the accumulation mode. Cloud residual sizes down to 20-30 nm were observed throughout all seasons. For liquid clouds, smaller cloud residual sizes occurred at lower total particle numbers.

285

To illustrate how both the number and the size of the cloud residuals and the total particles behave for our cloud cases, we show the number size distributions of cloud residuals and total particles for a cloud case of winter, spring, summer and late fall (Fig. 3). In agreement with Karlsson et al. (2021), we observe more aerosol particles and cloud residuals in the spring and




290 summer, and fewer in winter and fall. Also, the size distributions largely follow the findings from their long-term study: the  
ambient aerosol size distribution is dominated by accumulation mode in winter and fall, and shifts to being more Aitken mode-  
dominated in summer. For the cloud residuals, we observe the largest sizes in spring (144 nm on May 18, 2020), and smaller  
sizes (56 nm on Jun 12-1, 2020) with a more pronounced Aitken mode in summer. In winter the cloud residuals show the  
smallest sizes (18 nm on Dec 25, 2020). The cloud residual sizes from late fall show a plateau in the average number size  
295 distribution from around 66 to 144 nm.

As we focus on a few cloudy events only throughout an entire year, deviations from the seasonal pattern of particle number  
and size are expected. This can be illustrated in the case of the cloud event on June 27, 2020 (samples June 27-1 and June 27-  
2, Sect. S3, Fig. S3), where the ambient average particle number concentrations ( $46 \text{ cm}^{-3}$ ) are much lower compared to typical  
conditions during this time of the year (Ström et al., 2003; Tunved et al., 2013; Karlsson et al., 2021). The visibility was below  
300 1000 m for several hours before we sampled the two cloud residual samples, and the low aerosol number concentrations are  
most likely a result of cloud scavenging.

For most of the cloud samples, the average number of cloud residuals per size bin is smaller than the total number of particles  
measured at the same time in the same size bin (Sect. S3, Fig. S3). This is largely expected, since the number of cloud residuals  
is just a fraction of the number of total particles. However, for three cloud residual samples (June 27-1, June 25, and September  
305 12, 2020), we observe on average more Aitken mode particles (14 nm - 16 nm) in the cloud residuals than the average total  
particles. It is possible that this behavior is a sampling artifact from the GCVI. Karlsson et al., (2021) have discussed potential  
artifacts in the cloud residuals due to shattering of ice crystals when hitting the tip of the GCVI, mainly during the winter  
months when the fraction of ice crystals exceeds the number of liquid droplets. Our sample from December 25, 2019 ( $T: -$   
13.7°C) can be grouped in this category. In the June 27-1, June 25, and September 12, 2020 samples, the temperature was  
310 between  $-0.5^\circ\text{C}$  and  $4.2^\circ\text{C}$  and the sampled clouds composed of mostly liquid droplets, but contained some ice as well (Sect.  
S3, Fig. S4). In analogy to ice crystal shattering, it can be possible that the enhanced number of cloud residuals with a peak  
below 20 nm arises from splashing drizzle droplets within the funnel of the GCVI wind tunnel. Another possible reasons could  
be shrinking of the cloud residuals in the GCVI when the water evaporates, the capture of particles due to the wake effect  
(Pekour and Cziczo, 2011), or entrainment of drier air (Targino et al., 2007). Based on the target classification on all the three  
315 days, it is likely that there was drizzle present (Sect. S3, Fig. S4).

These potential artifacts are of negligible relevance for the discussion of the chemical composition of the cloud residuals the  
following below, as their contribution to the bulk mass, and hence bulk chemical composition measured by the FIGAERO-  
CIMS is small (Sect. S3, Fig. S5). Moreover, in the case of droplet splashing, only the size of the cloud residuals is affected  
and not the chemical composition, as the material of the splashing droplets would remain in the cloud residuals when the cloud  
320 droplets evaporate.



**Figure 3:** Average number size distributions of cloud residuals and total particles during the corresponding 2.5 h FIGAERO-CIMS sampling time of (a) December 25, 2019 (winter), (b) May 18, 2020 (spring), (c) June 12-1, 2020 (summer), (d) October 28, 2020 (late fall). The insert in (d) shows the number size distributions of cloud residuals and total particles with the y-axis adjusted such that the shapes of the distributions are visible. The solid and the dashed lines represent the mean and the median, respectively. The shaded areas are the standard deviations. The orange and blue numbers show the mean particle size of the total particles and the cloud residuals, respectively.

325

### 330 3.4 Chemical characteristics of cloud residuals

In the following we discuss the chemical composition of the cloud residuals of the 14 cloud events measured during the year-long NASCENT campaign with the FIGAERO-CIMS behind the GCVI as a function of season, and in relationship to the composition of the total particle population. With the FIGAERO-CIMS and using iodide as reagent ion, we were able to identify the molecular composition of in total 1558 different compounds. These include inorganic compounds, and organic compounds following the formula  $I_hC_iH_jO_kY_l^-$  ( $Y = \{N, S, Cl, Br, Si\}$ ), where  $h$  is the number of iodide atoms,  $i$  the number of carbon atoms,  $j$  the number of hydrogen atoms,  $k$  the number of oxygen atoms, and  $l$  the number of either nitrogen, sulfate, chloride, bromide or silicon atoms. In total 1094 of the detected compounds were clustered with iodide ( $h = 1-3$ ).

335

The observed compounds can be grouped as following based on their composition: ICHO ( $h = 1, i > 0, j > 0, k > 0$ ), CHO ( $h = 0, i > 0, j > 0, k > 0$ ), ICHON ( $h = 1, i > 0, j > 0, k > 0, Y = N, l = 1$ ), ICHOS ( $h = 1, i > 0, j > 0, k > 0, Y = S, l = 1$ ), I<sub>2</sub>CHO ( $h = 2, i > 0, j > 0, k > 0$ ), I<sub>3</sub>CHO ( $h = 3, i > 0, j > 0, k > 0$ ), and other. A subgroup of ICHO and CHO are fatty acids (FA), with  $h < 2, 3 < i < 29, k = 2, 0 < j < 2 * i$ . Levoglucosan (IC<sub>6</sub>H<sub>10</sub>O<sub>5</sub>-), a tracer for biomass burning, is part of the ICHO group. We exclude formic acid (ICH<sub>2</sub>O<sub>2</sub>-) from the following analyses due to interference from the gas phase (Sect. S4, Fig. S7). From the ICHON group we exclude IC<sub>6</sub>H<sub>15</sub>O<sub>7</sub>N-, as it has a very high signal in the cloud residual blanks. For the ICHOS group, our focus hereafter lies on MSA (detected as ICH<sub>4</sub>O<sub>3</sub>S-, with a clear signal above background, Sect. S2, Fig. S2). In the “others” group we include the inorganic compounds sulfuric acid (SA, detected as IH<sub>2</sub>SO<sub>4</sub>-) and nitric acid (NA, detected as IHNO<sub>3</sub>-), as well as organic compounds clustered with more than one iodide ion, I<sub>2</sub>CHO and I<sub>3</sub>CHO, the identification of the molecular formula of which was not straightforward.

345


To illustrate the difference in the molecular composition of the cloud residuals observed during the different seasons of NASCENT we present a mass spectrum from winter (Dec 25, 2019), spring (May 18, 2020), summer (Jun 12-1, 2020) and



350 late fall (Oct 28, 2020) each (Fig. 4). For all seasons except winter, we find a clear contribution of organic compounds across  
the entire mass spectrum, and higher signals of a few individual compounds such as MSA, SA, and NA. For most of the cloud  
residuals in spring, summer and fall, the ICHO show a similar pattern to that in mass spectra from offline filter samples from  
the summertime high Arctic measured by FIGAERO-CIMS (Siegel et al., 2021), with several molecules in the mass range of  
about 220 until 360 Th. In contrast to spring, summer and fall, the winter cloud residuals show signal above background for  
355 only very few compounds, mainly NA and  $\text{IC}_3\text{H}_6\text{O}_3^-$ . In the dark months, the contribution of organic aerosol to the total  
particulate matter is expected to be lower than during the sunlit part of the year (Moschos et al., 2022b). The absence of more  
oxygenated organics in the wintertime cloud residual samples seems to reflect this. The average signal-weighted O:C ratio of  
the ICHO group is slightly larger than the ratio of the ICHON group (0.55 - 0.81, and 0.59 - 0.73, respectively), which is in  
agreement with offline FIGAERO-CIMS measurements conducted in the central Arctic (Siegel et al., 2021).

360 The cloud residuals observed in spring, summer and fall (May 18, 2020 until Sep 12, 2020) show a clear MSA signal, whereas  
the cloud residuals observed in winter (Dec 25, 2019; Oct 28 and Dec 9, 2020) do not (Sect. S5, Fig. S8). The large signal of  
MSA suggests a marine contribution to CCN in the summertime. In a recent study from the Southern Ocean it was suggested  
that MSA is formed in the aqueous phase and thereby contributes to the growth of aerosol particles (Baccarini et al., 2021). In  
Sect. 2.7 we investigate this further.

365



**Figure 4: Mass spectra of cloud residuals. (a) December 25, 2019, (b) May 18, 2020, (c) June 12-1, 2020, (d) October 28, 2020. The detected compounds are presented as I-clusters, as detected by the FIGAERO-CIMS.**

370

### 3.5 Seasonal patterns of cloud residual chemical components

In general, the ambient organic aerosol mass is much lower in the dark period of the year than during the bright season. The  
particle composition in Ny-Ålesund during the dark period is dominated by anthropogenic emissions reaching the station due  
to atmospheric long-range transport, whereas in the sunlit period of the year, natural emissions can account for almost half of  
375 the organic submicron aerosol burden (Moschos et al., 2022a, b). The transition period from late winter to spring is known as  
the Arctic haze, characterized by enhanced aerosol mass concentrations (e.g. Tunved et al., 2013). In the following, we will



investigate how this general pattern is reflected in the cloud residuals. We note that we have less cloud residual data during the winter months due to less cloudy conditions and lower concentrations close to the detection limit, and in the haze month because of the mentioned problems with the instrumentation (Sect. 3.1); hence, our observations are skewed towards the summer months.

In Fig. 5 we present the cloud residual signal of various compounds and groups of compounds measured by FIGAERO-CIMS behind the GCVI as absolute signal (Fig. 5a) and as relative signal (Fig. 5b). We include compounds related to natural (MSA, SA, levoglucosan, FA) and anthropogenic (ICHON, SA, NA) emissions and show that they are present in the cloud residuals throughout the year.

The largest fraction of compounds in all cloud residuals is made up by the organic CHO+ICHO groups. The relative contribution of these compounds to the total measured signal is highest in the two cloud residuals on Jun 12, whereas the absolute signal is highest in the cloud residual sample of Jun 25. Except for the sample of Sep 12, 2020, the observed absolute signal contribution follows the expected ambient organic aerosol seasonality (Moschos et al., 2022b). It is interesting to note that from June 25 until September 12, the signal of CHO+ICHO and MSA both decrease in absolute numbers, but MSA has a more pronounced decrease in relative terms. This could be linked to the different source regions: While the MSA precursor DMS peaks in late spring until early summer (May-June), other organic compounds are present all year round and have both natural and anthropogenic continental sources (Behrenfeldt et al., 2008; Jang et al., 2021). The elevated levels of CHO+ICHO in the late summer (Sep 12) in combination with the elevated levels of NA and levoglucosan at the same time could indicate an anthropogenic influence or a combination of different sources the air masses encountered before arriving at the station.

The absolute signal of organonitrates (ICHON) follows the pattern of the CHO+ICHO group, and their relative contribution to the total signal observed with the FIGAERO-CIMS is similar for the individual cloud residual samples. Compounds in the ICHON group might have been formed via ICHO and NO<sub>x</sub> emissions from year-round ship traffic in the surrounding seas and the Arctic Ocean, or from the nearby power plant (Eckhardt et al., 2013).

The seasonal contribution of compounds with molecular formulae corresponding to fatty acids with 4 to 28 carbon atoms follows the contribution of the CHO+ICHO. FA-like compounds were also found in ambient aerosol samples from the high Arctic with offline FIGAERO-CIMS (Siegel et al., 2021). Their presence indicates organic enriched sea spray as a natural source of CCN (Mashayekhy Rad et al., 2018).


The seasonal pattern of the absolute signal of NA in the cloud residual samples is similar to that of the CHO+ICHO group for the majority of the samples. However, while CHO+ICHO has the highest absolute signal on June 25, NA shows the largest absolute and relative signal on May 18. Given that NA is related to anthropogenic emissions, and back trajectories (Sect. S6, Fig. S9) indicate the northern coast of Russia as the source region, it is possible that there was a prominent continental contribution to the aerosol population on May 18.

Levoglucosan (C<sub>6</sub>H<sub>10</sub>O<sub>5</sub>) is a known tracer for biomass burning (Simoneit et al., 1999). Of all the compounds and compound groups we present in Fig. 5, levoglucosan has the lowest contribution to the observed signal in each of the cloud residual samples (below 1 % in all samples in Fig. 5b). We observe elevated absolute signals of levoglucosan in the cloud residuals on May 21, from mid-June until September 12 and also on December 9, 2020. These results show that long-range transport of biomass burning aerosol (Stohl et al., 2007; Zangrando et al., 2013; Moroni et al., 2020) is contributing to the CCN fraction during large parts of the year.

The measured total contribution of MSA to the cloud residual mass is overall low in the beginning of the year, increases towards June, and decreases again from September onwards. We observe the highest contributions to the cloud residual mass at the end of June 2020. Elevated MSA absolute signal can also be found in the cloud residuals in May, in the beginning of June and in mid-September 2020. The increase of MSA in the cloud residuals towards June follows the overall pattern of ambient particulate MSA levels measured with the FIGAERO-CIMS in 2020 (Siegel et al., in prep.). Also in previous years, the ambient concentration of particulate MSA in Ny-Ålesund was found to follow a seasonal pattern with the highest



420 concentrations in May or June (Jang et al., 2021; Park et al., 2021; Moschos et al., 2022b). Since MSA is an oxidation product  
of DMS, the presence of MSA in the cloud residuals during the sunlit time of the year indicates a marine contribution to the  
aerosol particle population able to act as CCN. This confirms previous ambient aerosol observations at the same measurement  
location suggesting that DMS oxidation products are relevant to grow aerosol particles to CCN-active sizes (Park et al., 2021).  
Another oxidation product of DMS is SA, which was observed in the cloud residuals as well. Since SA is formed via oxidation  
425 (in both gas and aqueous phase) of sulfur dioxide (SO<sub>2</sub>), which has both anthropogenic (e.g. emissions from coal burning) and  
natural origins (e.g. oxidation product of DMS), source regions of the air mass need to be considered to estimate its origin.  
The absolute signal of SA in general shows a similar pattern as the absolute signal of MSA, except for the cloud residuals on  
May 21 and September 12, when both the absolute and relative SA signal is highest of all cloud residual samples. For May 21,  
HYSPLIT trajectories show that the air mass originated from the south-west and spent some time over the Greenland Sea, and  
430 the Barents and Kara Sea further east of Svalbard (Sect. S6, Fig. S9). On Sep 12, the trajectories show air arriving from the  
Barents Sea and the northern coast of Norway. The Greenland Sea south-west of Svalbard provides a source for DMS emissions  
and air masses from this area have been previously observed to contain high levels of SA (Lee et al., 2020). A large  
anthropogenic source region of SO<sub>2</sub> emissions, the Kola Peninsula, is located near the coastal region of the Barents Sea. SO<sub>2</sub>  
emissions from this region have been identified to form SA, which drives new particle formation far away from the source  
435 region (Sipilä et al., 2021). Given that these high SA concentrations do not correlate with the observed MSA, the high SA  
signal in the cloud residuals on May 21 and September 12 can probably be attributed to be mainly of anthropogenic origin. It  
is interesting to note that on Jun 25, 2020, the HYSPLIT trajectories indicate air coming from near the continental border  
between Norway and Russia, similar to September 12, but with less exposure over the Barents Sea. However, the cloud residual  
sample on Jun 25 contains much less SA and high MSA. The reason for this could be linked to the seasonality of MSA, with  
440 a large source in June (Jang et al., 2021). The different signals of the two DMS oxidation products suggests that the aerosol  
particles acting as CCN in the Arctic can have a strong seasonality, and natural sulfur sources might contribute to their mass  
to a large extent during the phytoplankton bloom season.



445 **Figure 5:** (a) Absolute signal of different compound groups (CHO+ICHO, ICHON,  $I_{0-1}C_{i=4-28}H_{2i}O_2^-$  (fatty acids),  $IHNO_3^-$  (NA  $IC_6H_{10}O_5^-$  (levoglucosan),  $IH_2SO_4^-$  (SA),  $ICH_4SO_3^-$  (MSA)) in the different cloud residual samples, and the respective  $PM_{1, \text{mass}}$ . (b) Relative signal of different compound groups in the different cloud residual samples. Note: in the absolute signal view in (a) the CHO+ICHO group contains also the signal of  $IC_6H_{10}O_5^-$ , and  $I_{0-1}C_{i=4-28}H_{2i}O_2^-$ , whereas for the relative signal in (b) the signal from these two groups have been subtracted from CHO+ICHO.

450

### 3.6 Cloud residual composition and size

To investigate if there is a link between the size and the sources of the cloud residuals, we investigated the mass fraction of Aitken mode particles ( $< 80$  nm) to  $PM_{1, \text{mass}}$ , and the number fraction of Aitken mode particles to the sum of Aitken and accumulation mode particles (Fig. 6a), and the ratio of MSA-to-SA (Fig. 6b). Since MSA is only produced from natural sulfur emissions and SA can be produced from both natural and anthropogenic emissions, their ratio provides a relative estimate of the dominating source region. We note again that this is not a direct comparison of size and chemistry, as the overall contribution of the Aitken mode to the chemical composition is much smaller than that of the accumulation mode.


455

The number fraction of Aitken mode particles in the cloud residuals shows a minimum in spring and late fall (May 18 and Oct 28, 2020) and a maximum in the summer (Jun 27-1 and June 27-2, 2020) and in the winter (Dec 25, 2019 and Dec 9, 2020,



460 Fig. 6a). The contribution of Aitken mode particles to the  $PM_{10}$  mass of the cloud residuals is clearly increasing from spring towards summer (May 18 until Jun 27-2, 2020), and is low during the rest of the year. The dominating number of Aitken mode particles in the winter is most likely linked to the presence of ice particles that create artifacts in the GCVI (Karlsson et al., 2021).

In general, the summer period with a high contribution of Aitken-mode particles in the cloud residuals coincides with high  
 465 MSA signals (Fig. 6b). The highest MSA/SA ratio is found in the cloud residual sample on June 2, 2020, suggesting that during this cloud case the contribution of natural sulfate was highest of all the cloud residual samples. In absolute terms, the largest signal of MSA is found in the cloud residual sample on Jun 25, 2020. Aitken-mode aerosol particles measured in Ny-Ålesund have previously been associated with new particle formation (NPF) events during the summer time when local sources dominate the aerosol population (Tunved et al., 2013; Lee et al., 2020). Ambient particulate MSA concentrations have been  
 470 reported to be highest during NPF events, and based on gas-phase measurements, this DMS oxidation product has been shown to be an important contributor to the growth of newly formed particles in Ny-Ålesund (Dall'Osto et al., 2017; Beck et al., 2021; Park et al., 2021). Aitken-mode particles have previously been shown to play a role as CCN in the summertime Arctic (Kecorius et al., 2019; Bulatovic et al., 2021; Karlsson et al., 2022). The combination of the observed dominating Aitken mode and elevated levels of MSA in the cloud residual samples at the end of June 2020 provides further evidence that during this  
 475 time of the year, MSA clearly contributes to the growth of newly formed particles into CCN sizes.



480 **Figure 6: (a) Mass fraction of Aitken mode particles to total  $PM_{10}$ , and number fraction of Aitken mode particles to the total number of Aitken and accumulation (acc) mode particles. As Aitken mode particles we consider particles < 80 nm. (b) Ratio of MSA to SA and their absolute signal for all cloud residuals.**

### 3.7 Cloud case study – May 18, 2020

In Fig. 7 we present different ambient properties during a cloud event on May 18, 2020. On this day we observed a cloud event  
 485 before, during, and after which the air masses were originating from the same region (Fig. 7a). This situation allowed us to study potential cloud processing of aerosol particles in detail. To do so, we identified FIGAERO-CIMS samples before, during, and after the cloud occurrence based on the observed visibility (Fig. 7b). *Before* defines the 2.5 h directly before the cloud was observed at the Zeppelin Observatory and the visibility was around 60 km. *During* is equal to the sampling time of the FIGAERO-CIMS cloud residual sample of May 18, 2020, measured in the beginning of the cloud when the median visibility  
 490 was 49 m. For this time period we measured the size distribution of both the cloud residuals and the interstitial aerosol particles





(together termed total particles), and the chemical composition of the cloud residuals only. *After* refers to the sampling time after the cloud had dissipated and the visibility was around 75 km. The average temperature during the cloud event was between -1.7 and -3.5°C, indicating most likely a low-level mixed-phase cloud, which is also identified as such in the target classification (Fig. 7c). The air mass reached the Zeppelin Observatory from the east, where it spent time over the northern coast of Russia. Throughout the cloud event the back trajectories show very similar source regions, with a slight shift to more marine influence over time. The air mass spent most of the time above the boundary layer (Sect. S7, Fig. S10).


### 3.7.1 Changes in aerosol number size distribution

*Before*, the aerosol number size distribution was composed of a dominating Aitken mode with a peak at around 45 nm and a smaller contribution of accumulation mode particles with a peak at around 125 nm (Fig. 7d). The total aerosol population also included particles in the nucleation mode (8-17 nm).

*During*, the accumulation mode particles with a peak at around 144 nm dominated the number concentration in the cloud residuals (Fig. 7d). A few Aitken mode particles down to around 56 nm were observed in the cloud residuals as well. Overall, the cloud residuals span a size range of around 60 to 300 nm. Due to the larger size of the accumulation mode particles, they are expected to activate into cloud droplets first. The peak in number occurring at 144 nm for the cloud residuals and lower numbers towards the 56 nm sized cloud residuals confirm this expected behavior. The total particle population (activated and non-activated aerosol particles) exhibited fewer particles in the nucleation and Aitken mode, and slightly more particles in the accumulation mode compared to *before*. Additionally, another peak in the size distribution occurred at around 79 nm in the total particle population, and a minimum at around 56 nm. The reduced number of nucleation and Aitken mode particles in the total particle population was probably a result of their coagulation with the cloud droplets. The second peak in the Aitken mode could indicate the advection of a different air mass.

*After*, the number size distribution showed a bimodal distribution of Aitken and accumulation mode particles (Fig. 7d). Compared to *during*, the nucleation mode particles were completely removed and the number of Aitken mode particles had decreased further. Additionally, compared to *before*, the peak in the accumulation mode had shifted to a larger size at around 141 nm, similar to *during*. The average particle number concentration had decreased by about half, from 559 cm<sup>-3</sup> *before* to 344 cm<sup>-3</sup> *after*. *During*, there were on average 456 cm<sup>-3</sup> total particles and 105 cm<sup>-3</sup> cloud residuals. This decrease in total number can be mainly attributed to the decrease in Aitken mode particles. In combination with the shift to larger particles *after*, this could indicate growth of the accumulation mode particles due to coalescence with the Aitken mode particles *during*, which resulted in larger aerosol particles *after* when the cloud droplets had evaporated. It is also possible that the accumulation mode particles grew due to in-cloud processing. We investigate this in the next section.

The *after* bimodal size distribution with a minimum between Aitken and accumulation mode at 63 nm (known as the “Hoppel minimum” (Hoppel et al., 1986)) indicates a non-precipitating cloud, and agrees well with observations from the central Arctic (64 to 71 nm, Karlsson et al., 2022). According to Hoppel et al. (1986), this minimum is related to the activation diameter of the aerosol particles. The presence of a Hoppel minimum in the aerosol size distribution at the Zeppelin Observatory has frequently been observed (Tunved et al., 2013).



530 **Figure 7:** Cloud case on May 18, 2020. (a) 5 days back trajectories of air masses arriving at the Zeppelin Observatory before, during and after the cloud event color-coded by time. (b) Time evolution of the visibility and sampling times of the FIGAERO-CIMS samples before, during and after the cloud event (grey shaded area). (c) Target classification on May 18, 2020 ((Cloudnet (2021), last access: 2022-11-20, 20:17 UTC) (d) Number size distributions before, during and after the cloud event. Before and after the cloud event the two DMPS both measure behind the whole-air inlet; hence, they both measure the same particle population during these times. For before and after we only show the DMPS2 data. During the cloud event, DMPS1 measured behind the GCVI and DMPS2 measured behind the whole-air inlet; hence, DMPS1 measured the cloud residuals and DMPS2 the total particle population. Average total particle number concentrations: 552 cm<sup>3</sup> before, 379 cm<sup>3</sup> during – total particles, 96 cm<sup>3</sup> during – cloud residuals, 285 cm<sup>3</sup> after.

535

### 3.7.2 Changes in aerosol chemical composition

Overall, we observed only small changes in the chemical composition throughout the cloud event (Fig. 8a, pie charts). The “other” group, containing largely inorganic sulfur- and chloride-containing species, contributes the same fraction throughout the cloud event. The largest change from *before* to *after* is the increased fraction of CHO+ICHO (also in absolute terms, not shown here), whereas MSA and NA show a slight decrease. Also, the absolute signal of NA is decreasing from *before* to *after* (not shown here). MSA and NA show a decreasing trend also from *before* to *during*. However, the changes in chemical composition are not that pronounced, which is similar to a study conducted at a mountain site in Sweden using a Quadrupole Aerosol Mass Spectrometer behind a GCVI observing no significant change in aerosol composition between before and after a cloud event (Drewnick et al., 2006).

545

As the CHO+ICHO contribute the largest fraction of cloud residual signal for the cloud case on May 18, 2020, and also of most of the cloud residuals measured with the FIGAERO-CIMS during the year (see Sect. 3.5), we investigate this group in more detail and look at their molecular composition. Overall, the molecular composition of the CHO+ICHO does not change



much during to the cloud event. The majority of the CHO+ICHO are composed of 2 to 11 carbon atoms and 3 to 6 oxygen atoms (Fig. 8a). Among these, the compounds with 4 oxygen atoms, likely dicarboxylic acids, dominate the signal *before*, *during* and *after*. Dicarboxylic acids have been shown to be part of the water-soluble fraction of organic aerosols (Saxena and Hildemann, 1996). *During*, the number of oxygen atoms shifts to smaller numbers. Most of the CHO+ICHO signal comes from compounds with 5 and 7 carbon atoms. The largest change is observed for compounds with 3 carbon atoms. Their relative contribution is much lower *before* than *during* and *after*. Compounds that show a smaller change in relative contribution are those with 4 and 10 carbon atoms. Their contribution is much higher *before* and *after* than *during* (Sect. S7, Fig. S11). The increase in absolute and relative signal *after* compared to *before*, is possibly linked to aqueous phase oxidation in the cloud droplets.

In the following we identify individual compounds that showed an increasing (79 compounds, hereafter termed as *increasing*) or decreasing (46 compounds, hereafter termed as *decreasing*) trend from *before* to *after* the cloud event, and compounds that are highest *during* in the cloud residuals (Fig. 8b, including all CHO+CHOI, CHION, NA, MSA, SA groups). In Fig. 8b we show the trends in form of a mass defect, to show how the compounds are spread over the entire  $m/z$  range. The *highest during* category comprises 428 compounds. Of all the compounds with a change in relative signal, no specific chemical families could be identified. We therefore focus on the three compounds showing the most pronounced relative change in signal in each of the three groups (*increasing*, *decreasing*, *highest during*).

The three compounds with the strongest relative increase were  $\text{IC}_4\text{H}_6\text{O}_4^-$ ,  $\text{IC}_3\text{H}_4\text{O}_4^-$ ,  $\text{IC}_5\text{H}_8\text{O}_4^-$ , which are the chemical formulae of succinic acid, malonic acid and glutaric acid, respectively. Succinic and malonic acid have previously been found in cloud water in continental Europe (Löflund et al., 2002). We also observed levoglucosan in this group. Except for succinic acid, these acids have been found to be able to serve as CCN in the laboratory (Cruz and Pandis, 1997; Giebl et al., 2002; Rosenørn et al., 2006). It is likely that the compounds in the *increasing* category are chemical tracers of good CCN, or produced in the aqueous phase through oxidation of water-soluble or co-condensing gases, and due to their higher oxidation state are more likely to stay in the particulate phase after the cloud droplets evaporate.

The three compounds that showed the strongest relative decrease were NA,  $\text{IC}_2\text{H}_4\text{O}_3^-$  (might be glycolic acid), and  $\text{IC}_5\text{H}_6\text{O}_4^-$  (might be citraconic acid or glutaconic acid). Their decrease could be due to either aqueous-phase reactions or evaporation.

The three compounds that showed the highest signal *during* were  $\text{IC}_3\text{H}_6\text{O}_3^-$ ,  $\text{IC}_7\text{H}_8\text{O}_4^-$  and  $\text{IC}_8\text{H}_{10}\text{O}_4^-$ .  $\text{IC}_3\text{H}_6\text{O}_3^-$  can likely be attributed to lactic acid, whereas the other two compounds could not be unambiguously identified. The compounds in the *highest during* category may be particularly efficient as CCN or INP (ice nucleating particle), or be formed in the aqueous phase. However, lactic acid is also a tracer of human activity, and the high lactic acid signal could therefore indicate contamination from handling of the GCVI during maintenance.

The normalized signal of SA showed that it belongs to the category *highest during*. The elevated level of SA in the cloud residuals indicates aqueous-phase reaction of dissolved  $\text{SO}_2$  in the cloud droplets (Hoppel et al., 1994). Elevated contributions of sulfate in the cloud residuals compared to the ambient aerosol particles were also found in an aircraft study during the haze season near Alaska (Zelenyuk et al., 2010). In contrast to our study, they measured the difference in ambient particle and cloud residual composition vertically, where ambient particles were measured below the cloud. It is interesting that the SA levels *after* drop down to a similar level as *before* (Fig. 8c). If SA is produced inside the cloud droplets, we would expect the evaporated cloud droplets *after* to be enriched with SA. SA is rather non-volatile and partitioning into the gas phase therefore unlikely. From the changes in the total particle number size distribution we saw an additional Aitken mode at around 79 nm *during*, which we attributed to be linked to advection of a different air mass. Hence, a possible explanation for the lack of enrichment in SA after the cloud could be that we were not measuring the same air mass throughout the cloud event, or different air mass got entrained from above throughout the cloud event.


MSA does not show a clear trend from *before* towards *after* (Fig. 8b). It showed the highest normalized signal *before* and the lowest *during* (Fig. 8c). Hence, our data does not support previous studies stating that MSA is mainly produced in the aqueous



phase (Hoffmann et al., 2016; Chen et al., 2018; Baccarini et al., 2021; Xavier et al., 2022). The ratio of MSA to SA *before* and *after* is very similar, whereas *during* the ratio is much smaller (Fig. 8c). This decrease in the cloud residuals is almost entirely driven by changes in SA.

595 Overall, the chemical composition of the aerosol particles is similar throughout the cloud event. This indicates that the aerosol particles were subject of cloud processing already before we observed this cloud case, or cloud processing occurs on longer time scales than regarded in this study.

600



605

610

**Figure 8:** Cloud case on May 18, 2020, chemical composition of total particles before, and after, of cloud residuals during the cloud event. (a) Pie chart showing the relative contribution of groups and individual compounds to the measured FIGAERO-CIMS signal. Green: CHO+ICHO. Purple: fatty acids, blue: ICHON, yellow: ICH<sub>4</sub>SO<sub>3</sub><sup>-</sup> (MSA), red: IH<sub>2</sub>O<sub>4</sub>S<sup>-</sup> (SA), grey: IHNO<sub>3</sub><sup>-</sup> (NA). Bar chart with absolute signal of CHO+ICHO compounds before, during and after the cloud event grouped by the number of carbon (1-19) and oxygen (O1-O10) atoms. (b) Mass defect highlighting individual compounds that show an increasing or decreasing trend during the cloud event (all: all compounds detected with the FIGAERO-CIMS; increasing: compounds of which the relative signal is increasing from before, via during until after the cloud; decreasing: compounds of which the relative signal is decreasing from before, via during until after the cloud; highest during: compounds of which the relative signal is highest during the cloud compared to before and after). (c) Ratio of MSA to sulfuric acid (SA) and their normalized signal before, during and after the cloud event.



#### 4 Conclusions

615 In this study we provide the first year-long in situ chemical characterization of cloud residuals at molecular level from the Arctic. We investigated the chemical composition of cloud residuals measured with a FIGAERO-CIMS behind a GCVI inlet at the Zeppelin Observatory, Svalbard, during the NASCENT campaign. From December 2019 until December 2020, we analyzed in total 14 cloud events, two in December (one in 2019 and one in 2020), two in May 2020, eight in June 2020, one in September 2020, and one in October 2020. No cloud residual sample was collected during the haze period.

620 The number concentrations and size distributions of the cloud residuals agreed roughly with the seasonal pattern observed in the long-term study by Karlsson et al. (2021). We found that the overall chemical composition of the cloud residuals follows the expected annual cycle of aerosol chemical composition in the Arctic, with a large contribution of naturally derived secondary aerosol in the form of MSA during spring and summer. Organic aerosol was present in the cloud residuals during the entire year, with higher absolute signals in the summer compared to winter. Inorganic acids, namely nitric and sulfuric acid, had their largest contribution, both absolute and relative, in spring and late summer, indicating anthropogenic influence.

625 The biomass burning tracer levoglucosan was observed in the summer cloud residuals as well, although its relative contribution to the total measured signal was below 1 % during the entire year. Our results indicate that most of the large enough aerosol particles serve as cloud seeds in the Arctic. In this aerosol-limited environment the chemical properties of the aerosol particles are not as relevant as the physical properties (size and number).

630 We observed a general relation between the amount of Aitken mode particles and the amount of MSA in the cloud residuals, where higher contributions of Aitken mode particles were present at elevated levels of MSA in the summer. This indicates that during this time of the year, MSA clearly contributes to the growth of newly formed particles into CCN sizes.

In a cloud case study from May 18, 2020 we investigated the change in the aerosol chemical composition due to cloud processing. To do so, we identified distinct periods before, during and after the cloud event, during which the air was coming

635 from the same region east of Svalbard. This cloud case revealed that after the cloud event, the aerosol population contained a larger relative and absolute signal of oxygenated hydrocarbons (CHO+ICHO), possibly linked to aqueous phase oxidation in the cloud droplets. Nitric acid showed decreasing relative and absolute signal, possibly linked to evaporation. Among individual compounds that showed a relative increase throughout the cloud event are succinic, malonic and glutaric acid, as well as levoglucosan. No clear increase of MSA in the cloud residuals was observed, thus not supporting results from previous

640 studies that indicated MSA to be formed in cloud droplets. However, we note that the observed cloud event was rather short (9 hours). Overall, the presence of the cloud did not seem to change the chemical composition of the aerosol particles much. This indicates, at least for the timescales of a few hours the cloud could be observed, that cloud processing mainly has an impact on the mass of the aerosol particles, and to a lower extent on the relative contribution of individual compounds.

Whereas little change was observed for the chemical composition of the aerosol particles in comparison to the cloud residuals,

645 the presence of the cloud reduced the number of nucleation and Aitken mode particles. This can be attributed to coagulation of particles and cloud droplets. Overall, the cloud event shifted the average submicron particle population to fewer but slightly larger accumulation mode particles.

The method used in this study allowed us to only investigate the organic fraction and some inorganic acids in the cloud residuals. As the Arctic aerosol also contains other species such as sea salt, heavy metals or dust, our results only reflect part

650 of the cloud residual composition. Nevertheless, our study highlights the importance of natural marine sources as CCN in the Arctic region and further suggests that the entire organic aerosol fraction can serve as CCN in the Arctic.

For future experiments it would be crucial to identify the size-segregated cloud residual chemical composition. Especially of the Aitken mode particles, which play an important role in acting as cloud seeds during a large part of the year (Lawler et al., 2021). In addition, the chemical composition of the total particle composition (cloud residuals and interstitial aerosol particles)

655 along with the cloud residual composition should be investigated – in analogy to the data available for the size distributions. This additional information would shed light on the chemical composition of the interstitial aerosol particles, and allow us to



better determine the properties of CCN. Offline and size segregated filter sampling for later FIGAERO-CIMS analysis could be a suitable option.

In the sunlit part of the year, the contribution of natural, local aerosol particle sources to the Arctic aerosol are higher than in  
660 the dark period. Our results show that these sources provide relevant seeds for cloud formation in the Arctic. Since the highest  
cloud cover in Ny-Ålesund is usually expected in summer, the locally produced aerosol particles could be more important for  
cloud formation than the long-range transported aerosol particles in the dark period. Especially in the context of a warming  
Arctic, where more open ocean provides a larger source for phytoplankton emissions and thereby also MSA formation, the  
marine environment could provide an important source of atmospheric aerosol particles able to activate as cloud droplets.

665

#### Data availability

The data of this study will be available on the Bolin Centre Database (<https://bolin.su.se/data/>). The meteorological data is available on the EBAS data base (<https://ebas-data.nilu.no>). The Cloudnet data used in this study are generated by the Aerosol, Clouds and Trace Gases Research Infrastructure (ACTRIS) and are available from the ACTRIS Data Centre using the  
670 following links: <https://hdl.handle.net/21.12132/1.6ad98e2e244346e8>, <https://hdl.handle.net/21.12132/1.015016f142ae45b7>,  
<https://hdl.handle.net/21.12132/1.2068d2269e014b15>, <https://hdl.handle.net/21.12132/1.61025d8bee914c66>.

#### Author contributions

CM, PZ and RK were responsible for funding and conceptualization. YG, KS, SH and CM performed the FIGAERO-CIMS measurements. YG, KS and SH wrote code for FIGAERO-CIMS analysis. RK, PZ and GF operated the GCVI and provided  
675 the GCVI data. GF provided the DMPS data. YG analyzed the FIGAERO-CIMS, and visualized the GCVI, DMPS and FIGAERO-CIMS data. KS visualized the trajectories. YG wrote the manuscript with input from CM. CM was responsible for supervision. All co-authors read and commented on the manuscript.

#### Competing interests

Some of the authors are co-editors at ACP.

#### 680 Acknowledgements

We gratefully acknowledge financial support from the Knut and Alice Wallenberg (KAW) foundation (WAF project CLOUDFORM, grant no. 2017.0165 and ACAS project # 2016.0024). The work was also financially supported by Swedish Environmental Protection Agency (Naturvårdsverket) and by the funding agency FORMAS (IWCAA project # 2016-01427). This work was supported by the Swedish Research Council (Vetenskapsrådet starting grant, project number 2018-05045 and  
685 project number 2016-05100). This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 821205 (FORCeS). We owe great thanks to Peter Tunved from ACES for harmonizing the DMPS2 data. We are grateful to Kerstin Ebell for help with the interpretation of the Cloudnet classification data. The authors thank NILU, Norway, especially Wenche Aas, Anne Hjøllbrekke, and Ove Hermansen for providing the meteorological data (T, RH). We are very thankful for all the support from Helge Tore Markussen and Vera Sklet from the  
690 Norwegian Polar Institute in Ny-Ålesund, and from the technicians from the Norwegian Polar Institute in Ny-Ålesund and from ACES, especially, Christer Søren, Christelle Guesnon, Håkon Jonsson Ruud, Svein-Torgar Oland Paulsen, Filip Heitmann, Ondrej Tesar and Tabea Hennig. We owe great thanks to NILU for providing zero air. We acknowledge ACTRIS



and Finnish Meteorological Institute for providing the data set which is available for download from <https://cloudnet.fmi.fi/>.  
The cloud radar data for Ny Ålesund was provided by the University of Cologne, the ceilometer and microwave radiometer  
695 data by the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research. We gratefully acknowledge the funding  
by the Deutsche Forschungsgemeinschaft DFG (German Research Foundation) - project number 268020496 - TRR 172, within  
the "Transregional Collaborative Research Center 'Arctic Amplification: Climate Relevant Atmospheric and Surface  
Processes, and Feedback Mechanisms (AC)3'".

## References

- 700 Adachi, K., Tobo, Y., Koike, M., Freitas, G., Zieger, P., and Krejci, R.: Composition and mixing state of Arctic aerosol and  
cloud residual particles from long-term single-particle observations at Zeppelin Observatory, Svalbard, *Atmospheric Chem.*  
*Phys.*, 22, 14421–14439, <https://doi.org/10.5194/acp-22-14421-2022>, 2022.
- Alfarra, M. R., Paulsen, D., Gysel, M., Garforth, A. A., Dommen, J., Prévôt, A. S. H., Worsnop, D. R., Baltensperger, U., and  
Coe, H.: A mass spectrometric study of secondary organic aerosols formed from the photooxidation of anthropogenic and  
705 biogenic precursors in a reaction chamber, *Atmospheric Chem. Phys.*, 6, 5279–5293, <https://doi.org/10.5194/acp-6-5279-2006>,  
2006.
- Baccarini, A., Dommen, J., Lehtipalo, K., Henning, S., Modini, R. L., Gysel-Beer, M., Baltensperger, U., and Schmale, J.:  
Low-Volatility Vapors and New Particle Formation Over the Southern Ocean During the Antarctic Circumnavigation  
Expedition, *J. Geophys. Res. Atmospheres*, 126, <https://doi.org/10.1029/2021JD035126>, 2021.
- 710 Beck, L. J., Sarnela, N., Junninen, H., Hoppe, C. J. M., Garmash, O., Bianchi, F., Riva, M., Rose, C., Peräkylä, O., Wimmer,  
D., Kausiala, O., Jokinen, T., Ahonen, L., Mikkilä, J., Hakala, J., He, X., Kontkanen, J., Wolf, K. K. E., Cappelletti, D.,  
Mazzola, M., Traversi, R., Petroselli, C., Viola, A. P., Vitale, V., Lange, R., Massling, A., Nøjgaard, J. K., Krejci, R., Karlsson,  
L., Zieger, P., Jang, S., Lee, K., Vakkari, V., Lampilahti, J., Thakur, R. C., Leino, K., Kangasluoma, J., Duplissy, E., Siivola,  
E., Marbouti, M., Tham, Y. J., Saiz-Lopez, A., Petäjä, T., Ehn, M., Worsnop, D. R., Skov, H., Kulmala, M., Kerminen, V.,  
715 and Sipilä, M.: Differing Mechanisms of New Particle Formation at Two Arctic Sites, *Geophys. Res. Lett.*, 48,  
<https://doi.org/10.1029/2020GL091334>, 2021.
- Behrenfeldt, U., Krejci, R., Ström, J., and Stohl, A.: Chemical properties of Arctic aerosol particles collected at the Zeppelin  
station during the aerosol transition period in May and June of 2004, *Tellus B Chem. Phys. Meteorol.*, 60, 405–415,  
<https://doi.org/10.1111/j.1600-0889.2008.00349.x>, 2008.
- 720 Bulatovic, I., Igel, A. L., Leck, C., Heintzenberg, J., Riipinen, I., and Ekman, A. M. L.: The importance of Aitken mode aerosol  
particles for cloud sustenance in the summertime high Arctic – a simulation study supported by observational data,  
*Atmospheric Chem. Phys.*, 21, 3871–3897, <https://doi.org/10.5194/acp-21-3871-2021>, 2021.
- Burkart, J., Hodshire, A. L., Mungall, E. L., Pierce, J. R., Collins, D. B., Ladino, L. A., Lee, A. K. Y., Irish, V., Wentzell, J. J.  
B., Liggio, J., Papakyriakou, T., Murphy, J., and Abbatt, J.: Organic Condensation and Particle Growth to CCN Sizes in the  
725 Summertime Marine Arctic Is Driven by Materials More Semivolatile Than at Continental Sites: Organic Condensation and  
Particle Growth, *Geophys. Res. Lett.*, 44, 7255–7274, <https://doi.org/10.1002/2017GL075671>, 2017.
- Cai, J., Daellenbach, K., Wu, C., Zheng, Y., Zheng, F., Du, W., Haslett, S., Chen, Q., Kulmala, M., and Mohr, C.:  
Characterization of offline analysis of particulate matter with FIGAERO-CIMS, *Aerosols/Laboratory Measurement/Data*  
*Processing and Information Retrieval*, <https://doi.org/10.5194/amt-2022-248>, 2022.
- 730 Cavalli, F., Facchini, M. C., Decesari, S., Mircea, M., Emblico, L., Fuzzi, S., Ceburnis, D., Yoon, Y. J., O'Dowd, C. D.,  
Putaud, J.-P., and Dell'Acqua, A.: Advances in characterization of size-resolved organic matter in marine aerosol over the  
North Atlantic, *J. Geophys. Res.*, 109, D24215, <https://doi.org/10.1029/2004JD005137>, 2004.



- Chen, Q., Sherwen, T., Evans, M., and Alexander, B.: DMS oxidation and sulfur aerosol formation in the marine troposphere: a focus on reactive halogen and multiphase chemistry, *Atmospheric Chem. Phys.*, 18, 13617–13637, <https://doi.org/10.5194/acp-18-13617-2018>, 2018.
- 735 Chernokulsky, A. V., Esau, I., Bulygina, O. N., Davy, R., Mokhov, I. I., Outten, S., and Semenov, V. A.: Climatology and Interannual Variability of Cloudiness in the Atlantic Arctic from Surface Observations since the Late Nineteenth Century, *J. Clim.*, 30, 2103–2120, <https://doi.org/10.1175/JCLI-D-16-0329.1>, 2017.
- Cloudnet: Classification data from Ny-Ålesund on 18 May 2020, <https://hdl.handle.net/21.12132/1.6ad98e2e244346e8>, 2021.
- 740 Cruz, C. N. and Pandis, S. N.: A study of the ability of pure secondary organic aerosol to act as cloud condensation nuclei, *Atmos. Environ.*, 31, 2205–2214, [https://doi.org/10.1016/S1352-2310\(97\)00054-X](https://doi.org/10.1016/S1352-2310(97)00054-X), 1997.
- Dall’Osto, M., Beddows, D. C. S., Tunved, P., Krejci, R., Ström, J., Hansson, H.-C., Yoon, Y. J., Park, K.-T., Becagli, S., Udisti, R., Onasch, T., O’Dowd, C. D., Simó, R., and Harrison, R. M.: Arctic sea ice melt leads to atmospheric new particle formation, *Sci. Rep.*, 7, 3318, <https://doi.org/10.1038/s41598-017-03328-1>, 2017.
- 745 Dekhtyareva, A., Holmén, K., Maturilli, M., Hermansen, O., and Graversen, R.: Effect of seasonal mesoscale and microscale meteorological conditions in Ny-Ålesund on results of monitoring of long-range transported pollution, *Polar Res.*, 37, 1508196, <https://doi.org/10.1080/17518369.2018.1508196>, 2018.
- Drewnick, F., Schneider, J., Hings, S. S., Hock, N., Noone, K., Targino, A., Weimer, S., and Borrmann, S.: Measurement of Ambient, Interstitial, and Residual Aerosol Particles on a Mountaintop Site in Central Sweden using an Aerosol Mass Spectrometer and a CVI, *J. Atmospheric Chem.*, 56, 1–20, <https://doi.org/10.1007/s10874-006-9036-8>, 2006.
- 750 Eckhardt, S., Hermansen, O., Grythe, H., Fiebig, M., Stebel, K., Cassiani, M., Baecklund, A., and Stohl, A.: The influence of cruise ship emissions on air pollution in Svalbard – a harbinger of a more polluted Arctic?, *Atmospheric Chem. Phys.*, 13, 8401–8409, <https://doi.org/10.5194/acp-13-8401-2013>, 2013.
- Giebl, H., Berner, A., Reischl, G., Puxbaum, H., Kasper-Giebl, A., and Hitzenberger, R.: CCN activation of oxalic and malonic acid test aerosols with the University of Vienna cloud condensation nuclei counter, *J. Aerosol Sci.*, 33, 1623–1634, [https://doi.org/10.1016/S0021-8502\(02\)00115-5](https://doi.org/10.1016/S0021-8502(02)00115-5), 2002.
- 755 Gierens, R., Kneifel, S., Shupe, M. D., Ebell, K., Maturilli, M., and Löhnert, U.: Low-level mixed-phase clouds in a complex Arctic environment, *Atmospheric Chem. Phys.*, 20, 3459–3481, <https://doi.org/10.5194/acp-20-3459-2020>, 2020.
- Haywood, J. and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: A review, *760 Rev. Geophys.*, 38, 513–543, <https://doi.org/10.1029/1999RG000078>, 2000.
- Heslin-Rees, D., Burgos, M., Hansson, H.-C., Krejci, R., Ström, J., Tunved, P., and Zieger, P.: From a polar to a marine environment: has the changing Arctic led to a shift in aerosol light scattering properties?, *Atmospheric Chem. Phys.*, 20, 13671–13686, <https://doi.org/10.5194/acp-20-13671-2020>, 2020.
- Hiranuma, N., Brooks, S. D., Moffet, R. C., Glen, A., Laskin, A., Gilles, M. K., Liu, P., Macdonald, A. M., Strapp, J. W., and 765 McFarquhar, G. M.: Chemical characterization of individual particles and residuals of cloud droplets and ice crystals collected on board research aircraft in the ISDAC 2008 study: ISDAC CLOUD PARTICLE RESIDUALS, *J. Geophys. Res. Atmospheres*, 118, 6564–6579, <https://doi.org/10.1002/jgrd.50484>, 2013.
- Hoffmann, E. H., Tilgner, A., Schrödner, R., Bräuer, P., Wolke, R., and Herrmann, H.: An advanced modeling study on the impacts and atmospheric implications of multiphase dimethyl sulfide chemistry, *Proc. Natl. Acad. Sci.*, 113, 11776–11781, 770 <https://doi.org/10.1073/pnas.1606320113>, 2016.
- Hoppel, W. A., Frick, G. M., and Larson, R. E.: Effect of nonprecipitating clouds on the aerosol size distribution in the marine boundary layer, *Geophys. Res. Lett.*, 13, 125–128, <https://doi.org/10.1029/GL013i002p00125>, 1986.
- Hoppel, W. A., Frick, G. M., Fitzgerald, J. W., and Wattle, B. J.: A Cloud Chamber Study of the Effect That Nonprecipitating Water Clouds Have on the Aerosol Size Distribution, *Aerosol Sci. Technol.*, 20, 1–30, 775 <https://doi.org/10.1080/02786829408959660>, 1994.





- Huang, J. and Jaeglé, L.: Wintertime enhancements of sea salt aerosol in polar regions consistent with a sea ice source from blowing snow, *Atmospheric Chem. Phys.*, 17, 3699–3712, <https://doi.org/10.5194/acp-17-3699-2017>, 2017.
- Jang, S., Park, K.-T., Lee, K., Yoon, Y. J., Kim, K., Chung, H. Y., Jang, E., Becagli, S., Lee, B. Y., Traversi, R., Eleftheriadis, K., Krejci, R., and Hermansen, O.: Large seasonal and interannual variations of biogenic sulfur compounds in the Arctic atmosphere (Svalbard; 78.9° N, 11.9° E), *Atmospheric Chem. Phys.*, 21, 9761–9777, <https://doi.org/10.5194/acp-21-9761-2021>, 2021.
- 780 Karlsson, L., Krejci, R., Koike, M., Ebell, K., and Zieger, P.: A long-term study of cloud residuals from low-level Arctic clouds, *Atmospheric Chem. Phys.*, 21, 8933–8959, <https://doi.org/10.5194/acp-21-8933-2021>, 2021.
- Karlsson, L., Baccarini, A., Duplessis, P., Baumgardner, D., Brooks, I. M., Chang, R. Y. -W., Dada, L., Dällenbach, K. R., Heikkinen, L., Krejci, R., Leaitch, W. R., Leck, C., Partridge, D. G., Salter, M. E., Wernli, H., Wheeler, M. J., Schmale, J., and Zieger, P.: Physical and Chemical Properties of Cloud Droplet Residuals and Aerosol Particles During the Arctic Ocean 2018 Expedition, *J. Geophys. Res. Atmospheres*, 127, <https://doi.org/10.1029/2021JD036383>, 2022.
- 805 Kecorius, S., Vogl, T., Paasonen, P., Lampilahti, J., Rothenberg, D., Wex, H., Zeppenfeld, S., van Pinxteren, M., Hartmann, M., Henning, S., Gong, X., Welti, A., Kulmala, M., Stratmann, F., Herrmann, H., and Wiedensohler, A.: New particle formation and its effect on cloud condensation nuclei abundance in the summer Arctic: a case study in the Fram Strait and Barents Sea, *Atmospheric Chem. Phys.*, 19, 14339–14364, <https://doi.org/10.5194/acp-19-14339-2019>, 2019.
- 790 Kirpes, R. M., Bonanno, D., May, N. W., Fraund, M., Barget, A. J., Moffet, R. C., Ault, A. P., and Pratt, K. A.: Wintertime Arctic Sea Spray Aerosol Composition Controlled by Sea Ice Lead Microbiology, *ACS Cent. Sci.*, 5, 1760–1767, <https://doi.org/10.1021/acscentsci.9b00541>, 2019.
- 795 Köhler, H.: The nucleus in and the growth of hygroscopic droplets, *Trans Faraday Soc.*, 32, 1152–1161, <https://doi.org/10.1039/TF9363201152>, 1936.
- Korhonen, H., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., and Ström, J.: A global model study of processes controlling aerosol size distributions in the Arctic spring and summer, *J. Geophys. Res.*, 113, D08211, <https://doi.org/10.1029/2007JD009114>, 2008.
- 800 Lawler, M. J., Saltzman, E. S., Karlsson, L., Zieger, P., Salter, M., Baccarini, A., Schmale, J., and Leck, C.: New Insights Into the Composition and Origins of Ultrafine Aerosol in the Summertime High Arctic, *Geophys. Res. Lett.*, 48, <https://doi.org/10.1029/2021GL094395>, 2021.
- Leaitch, W. R., Korolev, A., Aliabadi, A. A., Burkart, J., Willis, M. D., Abbatt, J. P. D., Bozem, H., Hoor, P., Köllner, F., Schneider, J., Herber, A., Konrad, C., and Brauner, R.: Effects of 20–100 nm particles on liquid clouds in the clean summertime Arctic, *Atmospheric Chem. Phys.*, 16, 11107–11124, <https://doi.org/10.5194/acp-16-11107-2016>, 2016.
- 805 Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Kurtén, T., Worsnop, D. R., and Thornton, J. A.: An Iodide-Adduct High-Resolution Time-of-Flight Chemical-Ionization Mass Spectrometer: Application to Atmospheric Inorganic and Organic Compounds, *Environ. Sci. Technol.*, 48, 6309–6317, <https://doi.org/10.1021/es500362a>, 2014.
- Lee, H., Lee, K., Lunder, C. R., Krejci, R., Aas, W., Park, J., Park, K.-T., Lee, B. Y., Yoon, Y. J., and Park, K.: Atmospheric new particle formation characteristics in the Arctic as measured at Mount Zeppelin, Svalbard, from 2016 to 2018, *Atmospheric Chem. Phys.*, 20, 13425–13441, <https://doi.org/10.5194/acp-20-13425-2020>, 2020.
- 810 Löflund, M., Kasper-Giebl, A., Schuster, B., Giebl, H., Hitznerberger, R., and Puxbaum, H.: Formic, acetic, oxalic, malonic and succinic acid concentrations and their contribution to organic carbon in cloud water, *Atmos. Environ.*, 36, 1553–1558, [https://doi.org/10.1016/S1352-2310\(01\)00573-8](https://doi.org/10.1016/S1352-2310(01)00573-8), 2002.
- 815 Lopez-Hilfiker, F. D., Mohr, C., Ehn, M., Rubach, F., Kleist, E., Wildt, J., Mentel, Th. F., Lutz, A., Hallquist, M., Worsnop, D., and Thornton, J. A.: A novel method for online analysis of gas and particle composition: description and evaluation of a Filter Inlet for Gases and AEROsols (FIGAERO), *Atmospheric Meas. Tech.*, 7, 983–1001, <https://doi.org/10.5194/amt-7-983-2014>, 2014.



- Malloy, Q. G. J., Nakao, S., Qi, L., Austin, R., Stothers, C., Hagino, H., and Cocker, D. R.: Real-Time Aerosol Density  
820 Determination Utilizing a Modified Scanning Mobility Particle Sizer—Aerosol Particle Mass Analyzer System, *Aerosol Sci. Technol.*, 43, 673–678, <https://doi.org/10.1080/02786820902832960>, 2009.
- Mashayekhy Rad, F., Leck, C., Ilag, L. L., and Nilsson, U.: Investigation of ultrahigh-performance liquid chromatography/travelling-wave ion mobility/time-of-flight mass spectrometry for fast profiling of fatty acids in the high Arctic sea surface microlayer, *Rapid Commun. Mass Spectrom.*, 32, 942–950, <https://doi.org/10.1002/rcm.8109>, 2018.
- 825 Mauritsen, T., Sedlar, J., Tjernström, M., Leck, C., Martin, M., Shupe, M., Sjogren, S., Sierau, B., Persson, P. O. G., Brooks, I. M., and Swietlicki, E.: An Arctic CCN-limited cloud-aerosol regime, *Atmospheric Chem. Phys.*, 11, 165–173, <https://doi.org/10.5194/acp-11-165-2011>, 2011.
- McFarquhar, G. M., Ghan, S., Verlinde, J., Korolev, A., Strapp, J. W., Schmid, B., Tomlinson, J. M., Wolde, M., Brooks, S. D., Cziczo, D., Dubey, M. K., Fan, J., Flynn, C., Gultepe, I., Hubbe, J., Gilles, M. K., Laskin, A., Lawson, P., Leaitch, W. R.,  
830 Liu, P., Liu, X., Lubin, D., Mazzoleni, C., Macdonald, A.-M., Moffet, R. C., Morrison, H., Ovchinnikov, M., Shupe, M. D., Turner, D. D., Xie, S., Zelenyuk, A., Bae, K., Freer, M., and Glen, A.: Indirect and Semi-direct Aerosol Campaign: The Impact of Arctic Aerosols on Clouds, *Bull. Am. Meteorol. Soc.*, 92, 183–201, <https://doi.org/10.1175/2010BAMS2935.1>, 2011.
- Moroni, B., Ritter, C., Crocchianti, S., Markowicz, K., Mazzola, M., Becagli, S., Traversi, R., Krejci, R., Tunved, P., and Cappelletti, D.: Individual Particle Characteristics, Optical Properties and Evolution of an Extreme Long-Range Transported  
835 Biomass Burning Event in the European Arctic (Ny-Ålesund, Svalbard Islands), *J. Geophys. Res. Atmospheres*, 125, <https://doi.org/10.1029/2019JD031535>, 2020.
- Moschos, V., Schmale, J., Aas, W., Becagli, S., Calzolari, G., Eleftheriadis, K., Moffett, C. E., Schnelle-Kreis, J., Severi, M., Sharma, S., Skov, H., Vestenius, M., Zhang, W., Hakola, H., Hellén, H., Huang, L., Jaffrezo, J.-L., Massling, A., Nøjgaard, J. K., Petäjä, T., Popovicheva, O., Sheesley, R. J., Traversi, R., Yttri, K. E., Prévôt, A. S. H., Baltensperger, U., and El Haddad,  
840 I.: Elucidating the present-day chemical composition, seasonality and source regions of climate-relevant aerosols across the Arctic land surface, *Environ. Res. Lett.*, 17, 034032, <https://doi.org/10.1088/1748-9326/ac444b>, 2022a.
- Moschos, V., Dzepina, K., Bhattu, D., Lamkaddam, H., Casotto, R., Daellenbach, K. R., Canonaco, F., Rai, P., Aas, W., Becagli, S., Calzolari, G., Eleftheriadis, K., Moffett, C. E., Schnelle-Kreis, J., Severi, M., Sharma, S., Skov, H., Vestenius, M., Zhang, W., Hakola, H., Hellén, H., Huang, L., Jaffrezo, J.-L., Massling, A., Nøjgaard, J. K., Petäjä, T., Popovicheva, O.,  
845 Sheesley, R. J., Traversi, R., Yttri, K. E., Schmale, J., Prévôt, A. S. H., Baltensperger, U., and El Haddad, I.: Equal abundance of summertime natural and wintertime anthropogenic Arctic organic aerosols, *Nat. Geosci.*, <https://doi.org/10.1038/s41561-021-00891-1>, 2022b.
- Nomokonova, T., Ebell, K., Löhnert, U., Maturilli, M., and Ritter, C.: The influence of anomalous atmospheric conditions at Ny-Ålesund on clouds and their radiative effect, *Clouds and Precipitation/Remote Sensing/Troposphere/Physics*  
850 (physical properties and processes), <https://doi.org/10.5194/acp-2019-985>, 2019.
- Noone, K. J., Ogren, J. A., Heintzenberg, J., Charlson, R. J., and Covert, D. S.: Design and Calibration of a Counterflow Virtual Impactor for Sampling of Atmospheric Fog and Cloud Droplets, *Aerosol Sci. Technol.*, 8, 235–244, <https://doi.org/10.1080/02786828808959186>, 1988.
- Ogren, J. A., Heintzenberg, J., and Charlson, R. J.: In-situ sampling of clouds with a droplet to aerosol converter, *Geophys. Res. Lett.*, 12, 121–124, <https://doi.org/10.1029/GL012i003p00121>, 1985.
- Park, K., Yoon, Y. J., Lee, K., Tunved, P., Krejci, R., Ström, J., Jang, E., Kang, H. J., Jang, S., Park, J., Lee, B. Y., Traversi, R., Becagli, S., and Hermansen, O.: Dimethyl Sulfide-Induced Increase in Cloud Condensation Nuclei in the Arctic Atmosphere, *Glob. Biogeochem. Cycles*, 35, <https://doi.org/10.1029/2021GB006969>, 2021.
- Pasquier, J. T., David, R. O., Freitas, G., Gierens, R., Gramlich, Y., Haslett, S., Li, G., Schäfer, B., Siegel, K., Wieder, J.,  
860 Adachi, K., Belosi, F., Carlsen, T., Decesari, S., Ebell, K., Gilardoni, S., Gysel-Beer, M., Henneberger, J., Inoue, J., Kanji, Z. A., Koike, M., Kondo, Y., Krejci, R., Lohmann, U., Maturilli, M., Mazzolla, M., Modini, R., Mohr, C., Motos, G., Nenes, A.,



- Nicosia, A., Ohata, S., Paglione, M., Park, S., Pileci, R. E., Ramelli, F., Rinaldi, M., Ritter, C., Sato, K., Storelvmo, T., Tobo, Y., Traversi, R., Viola, A., and Zieger, P.: The Ny-Ålesund Aerosol Cloud Experiment (NASCENT): Overview and First Results, *Bull. Am. Meteorol. Soc.*, <https://doi.org/10.1175/BAMS-D-21-0034.1>, 2022.
- 865 Pekour, M. S. and Cziczko, D. J.: Wake Capture, Particle Breakup, and Other Artifacts Associated with Counterflow Virtual Impaction, *Aerosol Sci. Technol.*, 45, 758–764, <https://doi.org/10.1080/02786826.2011.558942>, 2011.
- Platt, S. M., Hov, Ø., Berg, T., Breivik, K., Eckhardt, S., Eleftheriadis, K., Evangeliou, N., Fiebig, M., Fisher, R., Hansen, G., Hansson, H.-C., Heintzenberg, J., Hermansen, O., Heslin-Rees, D., Holmén, K., Hudson, S., Kallenborn, R., Krejci, R., Krognes, T., Larssen, S., Lowry, D., Lund Myhre, C., Lunder, C., Nisbet, E., Nizzetto, P. B., Park, K.-T., Pedersen, C. A.,  
870 Aspö Pfaffhuber, K., Röckmann, T., Schmidbauer, N., Solberg, S., Stohl, A., Ström, J., Svendby, T., Tunved, P., Tørnkvist, K., van der Veen, C., Vratolis, S., Yoon, Y. J., Yttri, K. E., Zieger, P., Aas, W., and Tørseth, K.: Atmospheric composition in the European Arctic and 30 years of the Zeppelin Observatory, Ny-Ålesund, *Atmospheric Chem. Phys.*, 22, 3321–3369, <https://doi.org/10.5194/acp-22-3321-2022>, 2022.
- Pöhlker, M. L., Zhang, M., Campos Braga, R., Krüger, O. O., Pöschl, U., and Ervens, B.: Aitken mode particles as CCN in  
875 aerosol- and updraft-sensitive regimes of cloud droplet formation, *Atmospheric Chem. Phys.*, 21, 11723–11740, <https://doi.org/10.5194/acp-21-11723-2021>, 2021.
- Quinn, P. K., Shaw, G., Andrews, E., Dutton, E. G., Ruoho-Airola, T., and Gong, S. L.: Arctic haze: current trends and knowledge gaps, *Tellus B Chem. Phys. Meteorol.*, 59, 99–114, <https://doi.org/10.1111/j.1600-0889.2006.00236.x>, 2007.
- Rosenørn, T., Kiss, G., and Bilde, M.: Cloud droplet activation of saccharides and levoglucosan particles, *Atmos. Environ.*,  
880 40, 1794–1802, <https://doi.org/10.1016/j.atmosenv.2005.11.024>, 2006.
- Saxena, P. and Hildemann, L. M.: Water-soluble organics in atmospheric particles: A critical review of the literature and application of thermodynamics to identify candidate compounds, *J. Atmospheric Chem.*, 24, 57–109, <https://doi.org/10.1007/BF00053823>, 1996.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, 2nd ed., J. Wiley,  
885 Hoboken, N.J., 1203 pp., 2006.
- Shingler, T., Dey, S., Sorooshian, A., Brechtel, F. J., Wang, Z., Metcalf, A., Coggon, M., Mülmenstädt, J., Russell, L. M., Jonsson, H. H., and Seinfeld, J. H.: Characterisation and airborne deployment of a new counterflow virtual impactor inlet, *Atmospheric Meas. Tech.*, 5, 1259–1269, <https://doi.org/10.5194/amt-5-1259-2012>, 2012.
- Shupe, M. D. and Intrieri, J. M.: Cloud Radiative Forcing of the Arctic Surface: The Influence of Cloud Properties, Surface  
890 Albedo, and Solar Zenith Angle, *J. Clim.*, 17, 616–628, [https://doi.org/10.1175/1520-0442\(2004\)017<0616:CRFOTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0616:CRFOTA>2.0.CO;2), 2004.
- Siegel, K., Karlsson, L., Zieger, P., Baccarini, A., Schmale, J., Lawler, M., Salter, M., Leck, C., Ekman, A. M. L., Riipinen, I., and Mohr, C.: Insights into the molecular composition of semi-volatile aerosols in the summertime central Arctic Ocean using FIGAERO-CIMS, *Environ. Sci. Atmospheres*, 1, 161–175, <https://doi.org/10.1039/D0EA00023J>, 2021.
- 895 Siegel, K., Neuberger, A., Karlsson, L., Zieger, P., Mattsson, F., Duplessis, P., Dada, L., Daellenbach, K., Schmale, J., Baccarini, A., Krejci, R., Svenningsson, B., Chang, R., Ekman, A. M. L., Riipinen, I., and Mohr, C.: Using Novel Molecular-Level Chemical Composition Observations of High Arctic Organic Aerosol for Predictions of Cloud Condensation Nuclei, *Environ. Sci. Technol.*, [acs.est.2c02162](https://doi.org/10.1021/acs.est.2c02162), <https://doi.org/10.1021/acs.est.2c02162>, 2022.
- Siegel, K., Gramlich, Y., Haslett, S., Freitas, G., Tunved, P., Krejci, R., Zieger, P., and Mohr, C.: Measurement report: Year-  
900 long observation of Hydroperoxymethyl Thioformate (HPMTF) at the Zeppelin Observatory, Svalbard – seasonal behaviour and relationship to other oxidation products from dimethyl sulfide, in prep.
- Simoneit, B. R. T., Schauer, J. J., Nolte, C. G., Oros, D. R., Elias, V. O., Fraser, M. P., Rogge, W. F., and Cass, G. R.: Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles, *Atmos. Environ.*, 33, 173–182, [https://doi.org/10.1016/S1352-2310\(98\)00145-9](https://doi.org/10.1016/S1352-2310(98)00145-9), 1999.



- 905 Sipilä, M., Sarnela, N., Neitola, K., Laitinen, T., Kemppainen, D., Beck, L., Duplissy, E.-M., Kuittinen, S., Lehmusjärvi, T., Lampilahti, J., Kerminen, V.-M., Lehtipalo, K., Aalto, P. P., Keronen, P., Siivola, E., Rantala, P. A., Worsnop, D. R., Kulmala, M., Jokinen, T., and Petäjä, T.: Wintertime subarctic new particle formation from Kola Peninsula sulfur emissions, *Atmospheric Chem. Phys.*, 21, 17559–17576, <https://doi.org/10.5194/acp-21-17559-2021>, 2021.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bull. Am. Meteorol. Soc.*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- 910 Stohl, A., Berg, T., Burkhardt, J. F., Fjérraa, A. M., Forster, C., Herber, A., Hov, Ø., Lunder, C., McMillan, W. W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Tørseth, K., Treffeisen, R., Virkkunen, K., and Yttri, K. E.: Arctic smoke – record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006, *Atmospheric Chem. Phys.*, 7, 511–534, <https://doi.org/10.5194/acp-7-511-2007>, 2007.
- Ström, J., Umegård, J., Tørseth, K., Tunved, P., Hansson, H.-C., Holmén, K., Wismann, V., Herber, A., and König-Langlo, G.: One year of particle size distribution and aerosol chemical composition measurements at the Zeppelin Station, Svalbard, March 2000–March 2001, *Phys. Chem. Earth Parts ABC*, 28, 1181–1190, <https://doi.org/10.1016/j.pce.2003.08.058>, 2003.
- 915 Struthers, H., Ekman, A. M. L., Glantz, P., Iversen, T., Kirkevåg, A., Mårtensson, E. M., Seland, Ø., and Nilsson, E. D.: The effect of sea ice loss on sea salt aerosol concentrations and the radiative balance in the Arctic, *Atmospheric Chem. Phys.*, 11, 3459–3477, <https://doi.org/10.5194/acp-11-3459-2011>, 2011.
- Targino, A. C., Noone, K. J., Drewnick, F., Schneider, J., Krejci, R., Olivares, G., Hings, S., and Borrmann, S.: Microphysical and chemical characteristics of cloud droplet residuals and interstitial particles in continental stratocumulus clouds, *Atmospheric Res.*, 86, 225–240, <https://doi.org/10.1016/j.atmosres.2007.05.001>, 2007.
- 925 Thornton, J. A., Mohr, C., Schobesberger, S., D'Ambro, E. L., Lee, B. H., and Lopez-Hilfiker, F. D.: Evaluating Organic Aerosol Sources and Evolution with a Combined Molecular Composition and Volatility Framework Using the Filter Inlet for Gases and Aerosols (FIGAERO), *Acc. Chem. Res.*, 53, 1415–1426, <https://doi.org/10.1021/acs.accounts.0c00259>, 2020.
- Topping, D., Connolly, P., and McFiggans, G.: Cloud droplet number enhanced by co-condensation of organic vapours, *Nat. Geosci.*, 6, 443–446, <https://doi.org/10.1038/ngeo1809>, 2013.
- 930 Topping, D. O. and McFiggans, G.: Tight coupling of particle size, number and composition in atmospheric cloud droplet activation, *Atmospheric Chem. Phys.*, 12, 3253–3260, <https://doi.org/10.5194/acp-12-3253-2012>, 2012.
- Tunved, P., Ström, J., and Krejci, R.: Arctic aerosol life cycle: linking aerosol size distributions observed between 2000 and 2010 with air mass transport and precipitation at Zeppelin station, Ny-Ålesund, Svalbard, *Atmospheric Chem. Phys.*, 13, 3643–3660, <https://doi.org/10.5194/acp-13-3643-2013>, 2013.
- 935 Twohy, C. H. and Anderson, J. R.: Droplet nuclei in non-precipitating clouds: composition and size matter, *Environ. Res. Lett.*, 3, 045002, <https://doi.org/10.1088/1748-9326/3/4/045002>, 2008.
- Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Dethloff, K., Velasco, C. B., Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Engler, C., Eppers, O., Gehrmann, M., Gong, X., Gottschalk, M., Goubeyre, C., Griesche, H., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L., Leroy, D., Maturilli, M., Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T., Notholt, J., Palm, M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder, K., Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P., Shupe, M. D., Siebert, H., Spreen, G., Stapf, J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M., and Zeppenfeld, S.: The Arctic Cloud Puzzle: Using 945 ALOUD/PASCAL Multiplatform Observations to Unravel the Role of Clouds and Aerosol Particles in Arctic Amplification, *Bull. Am. Meteorol. Soc.*, 100, 841–871, <https://doi.org/10.1175/BAMS-D-18-0072.1>, 2019.



- Willis, M. D., Burkart, J., Thomas, J. L., Köllner, F., Schneider, J., Bozem, H., Hoor, P. M., Aliabadi, A. A., Schulz, H., Herber, A. B., Leitch, W. R., and Abbatt, J. P. D.: Growth of nucleation mode particles in the summertime Arctic: a case study, *Atmospheric Chem. Phys.*, 16, 7663–7679, <https://doi.org/10.5194/acp-16-7663-2016>, 2016.
- 950 Xavier, C., Baykara, M., Wollesen de Jonge, R., Altstädter, B., Clusius, P., Vakkari, V., Thakur, R., Beck, L., Becagli, S., Severi, M., Traversi, R., Krejci, R., Tunved, P., Mazzola, M., Wehner, B., Sipilä, M., Kulmala, M., Boy, M., and Roldin, P.: Secondary aerosol formation in marine Arctic environments: a model measurement comparison at Ny-Ålesund, *Atmospheric Chem. Phys.*, 22, 10023–10043, <https://doi.org/10.5194/acp-22-10023-2022>, 2022.
- Zangrando, R., Barbaro, E., Zennaro, P., Rossi, S., Kehrwald, N. M., Gabrieli, J., Barbante, C., and Gambaro, A.: Molecular  
955 Markers of Biomass Burning in Arctic Aerosols, *Environ. Sci. Technol.*, 130716103911002, <https://doi.org/10.1021/es400125r>, 2013.
- Zelenyuk, A., Imre, D., Earle, M., Easter, R., Korolev, A., Leitch, R., Liu, P., Macdonald, A. M., Ovchinnikov, M., and Strapp, W.: In Situ Characterization of Cloud Condensation Nuclei, Interstitial, and Background Particles Using the Single Particle Mass Spectrometer, SPLAT II, *Anal. Chem.*, 82, 7943–7951, <https://doi.org/10.1021/ac1013892>, 2010.

960