

Effects of denitrification on the distributions of trace gas abundances in the polar regions: a comparison of WACCM with observations

Michael Weimer¹, Douglas E. Kinnison², Catherine Wilka³, and Susan Solomon¹

¹Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA

²Atmospheric Chemistry Observations & Modeling Laboratory, National Center for Atmospheric Research, Boulder, CO, USA

³Department of Earth System Science, Stanford University, Stanford, CA, USA

Correspondence: Michael Weimer (mweimer@mit.edu)

Abstract. Polar stratospheric clouds (PSCs) play a key role in the polar chemistry of the stratosphere. Nitric acid trihydrate (NAT) particles have been shown to lead to denitrification of the lower stratosphere. While the existence of large NAT particles (NAT "rocks") has been verified by many measurements especially in the Northern Hemisphere (NH), most current chemistry-climate models use simplified parameterizations, often based on evaluations in the Southern Hemisphere where the polar vortex is stable enough that accounting for NAT rocks is not as important as in the NH. Here, we evaluate the probability density functions of various gaseous species in the polar vortex using one such model, the Whole Atmosphere Community Climate Model (WACCM), and compare these with measurements by the Michelson Interferometer for Passive Atmospheric Sounding onboard the Environmental Satellite (MIPAS/Envisat) and two ozonesonde stations for a range of years and in both hemispheres. Using the maximum difference between the distributions of MIPAS and WACCM as a measure of coherence, we find better agreement for HNO_3 when reducing the NAT number density from the standard value of 10^{-2} used in this model to $5 \times 10^{-4} \text{ cm}^{-3}$ for almost all spring seasons during the MIPAS period in both hemispheres. The distributions of ClONO_2 and O_3 are not greatly affected by the NAT density. The average difference of WACCM to ozonesondes supports the need to reduce the NAT number density in the model. Therefore, this study suggests to use a NAT number density of $5 \times 10^{-4} \text{ cm}^{-3}$ for future simulations with WACCM.

1 Introduction

Polar stratospheric clouds (PSCs) have been known for decades to play a key role in explaining the stratospheric ozone hole (Solomon et al., 1986; Solomon, 1999; Tritscher et al., 2021). Reactions on their surfaces lead to activation of chlorine reservoirs, known as chlorine activation. In addition, sedimentation of PSCs result in dehydration and denitrification of the lower stratosphere. Three types of PSCs with different composition and roles have been found to be important: liquid supercooled ternary solution droplets (STS, Carslaw et al., 1994) are major contributors to chlorine activation (e.g., Peter, 1997). Ice PSCs lead to dehydration of the stratosphere (e.g., Kelly et al., 1989). Nitric acid trihydrate particles (NAT) dominate the irreversible removal of nitric acid (HNO_3) from the lower stratosphere via sedimentation, known as denitrification, thus potentially reducing the reformation of chlorine reservoir species which can affect ozone loss (e.g., Waibel et al., 1999).

The existence of NAT particles in the stratosphere has been verified by a variety of airborne measurements (Fahey et al., 1989; Voigt et al., 2000; Fahey, 2001; Molleker et al., 2014; Woiwode et al., 2016). Since the polar vortex in the Southern Hemisphere (SH) generally is more stable than in the Northern Hemisphere (NH) (e.g., Schoeberl et al., 1992), the time period when denitrification can occur is much longer in the SH polar vortex than the NH. On the other hand, denitrification in the NH has been found to occur locally and the role of low-number density large-size NAT particles, so-called NAT rocks, has been discussed and verified by measurements (Fahey, 2001; Fueglistaler et al., 2002; Drdla and Müller, 2012; Adriani et al., 2004; Woiwode et al., 2014, 2016). These large particles can lead to significant sedimentation even on the shorter time scales needed to explain the occurrence of low HNO_3 volume mixing ratios (VMR) in the NH spring.

Satellite instruments are able to provide measurements of PSCs and gaseous HNO_3 with daily near-global coverage (Höpfner et al., 2018; Spang et al., 2018; Pitts et al., 2018; Santee et al., 2007; Wespes et al., 2022). Höpfner et al. (2006) found the first evidence of a NAT belt around the Antarctic Continent using the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) onboard the Environmental Satellite. Therefore, satellite measurements provide a unique opportunity for comparisons of HNO_3 and denitrification with current chemistry-climate models in both hemispheres. They also allow study of the range of the probability distributions in the data for different years and conditions.

Accounting for denitrification is an important process in chemistry-climate and in chemistry-transport models, which is why many parameterizations with different levels of detail have been developed to account for the microphysics and sedimentation of NAT particles in these models (e.g., Considine et al., 2000; Carslaw et al., 2002; Grooß et al., 2005; Wohltmann et al., 2010; Wegner et al., 2013; Zhu et al., 2015; Kirner et al., 2011; Weimer et al., 2021). Some models account for comprehensive microphysics of NAT particles (e.g., Zhu et al., 2015) to determine radii and concentrations which then redistribute gaseous HNO_3 . Others use an intermediate approach accounting not for the full microphysics but constraining the amount of NAT to a measured NAT distribution (e.g., Kirner et al., 2011; Weimer et al., 2021). In addition, there are models with diagnostic approaches, as suggested e.g. by Considine et al. (2000), where NAT is formed based on the available gaseous HNO_3 , sedimentation is computed and the HNO_3 contained in NAT particles is released back to the gaseous phase within the same model time step.

The standard version of the Whole Atmosphere component of the Community Earth System Model includes a diagnostic parameterization of NAT with a NAT number density set to a global value of 10^{-2} cm^{-3} (Wegner et al., 2013). However, Wilka et al. (2021) found that this value leads to an overestimation of gaseous HNO_3 in the NH for the extraordinarily cold NH winter 2019/2020 in comparison to measurements of the Microwave Limb Sounder. They found a better agreement with the measurements using an adopted NAT number density of $N_{\text{NAT}} = 10^{-5} \text{ cm}^{-3}$. Here, we investigate this further by applying an adapted version of the method by Zambri et al. (2021) to compare the distributions of various gaseous species within the polar vortex with MIPAS, which provided measurements during 2002 to 2012. MIPAS data is well suited to this task, because it provides high signal to noise local measurements over multiple years in both hemispheres, with good dynamic range; other datasets may also be appropriate but here we use MIPAS for this initial test. Further, using MIPAS and varying the NAT number density of the model, we can also investigate its associated impact on HNO_3 , ClONO_2 and O_3 for many spring seasons in both hemispheres.

Table 1. Simulations in this study. In the “noHetAll” simulation, the heterogeneous reactions are removed apart from the reaction $\text{N}_2\text{O}_5 + \text{H}_2\text{O}$.

| Simulation | with heterogeneous chemistry? | $N_{\text{NAT}} (\text{cm}^{-3})$ |
|---------------|-------------------------------|-----------------------------------|
| HetAll.1e-2 | Yes | 10^{-2} |
| noHetAll.5e-4 | No | 5×10^{-4} |
| HetAll.5e-4 | Yes | 5×10^{-4} |
| HetAll.1e-5 | Yes | 10^{-5} |

Details about the model and the observation as well as the methods to compare them can be found in Sect. 2. Section 3 shows the comparison between the datasets. Finally, some concluding remarks are given in Sect. 4.

2 Datasets

2.1 WACCM

In this study, we use the Whole Atmosphere component (WACCM6) of the Community Earth System Model (CESM2.1) in Specified Dynamics (SD) mode (Gettelman et al., 2019; Danabasoglu et al., 2020) to compare distributions of the trace gases with the satellite measurements. The model is relaxed towards the Modern-Era Retrospective Analysis for Research and Applications version 2 (MERRA2, Gelaro et al., 2017). In this study, we use a horizontal resolution of 1.9° latitude \times 2.5° longitude and 88 vertical levels up to about 140 km. In the component set “FWmaSD”, a comprehensive chemistry for the middle atmosphere is included, as also used e.g. by Zambri et al. (2021).

Polar stratospheric clouds in WACCM are calculated using a diagnostic parameterization described by Considine et al. (2000), Kinnison et al. (2007) and Wegner et al. (2013). NAT particles are formed in thermodynamical equilibrium with the gaseous H_2O and HNO_3 below the NAT threshold temperature (Hanson and Mauersberger, 1988). Coexistence of NAT and STS is accounted for by allowing 20% of HNO_3 to form NAT whereas the rest is available for STS (Wegner et al., 2013; Solomon et al., 2015). Sedimentation of the NAT particles, i.e. the vertical redistribution of gaseous HNO_3 , is calculated using a simple upwind scheme (Considine et al., 2000). The radius of the particles is determined in this scheme by using the amount of condensed HNO_3 and the NAT number density, which is a global parameter in this scheme. Larger NAT density leads to smaller particles and vice versa, assuming the mass of HNO_3 condensed in NAT is constant in the grid box. Therefore, the NAT density is a parameter in this scheme that can be used to tune the denitrification in the model. Examination of observations of NAT particle number densities (Pitts et al., 2009, 2011) and an emphasis on HNO_3 data for the SH in 2005 led Wegner et al. (2013) to adopt a NAT value of 0.01 cm^{-3} . Fahey (2001) measured NAT number densities between 2.3×10^{-4} and $2 \times 10^{-3} \text{ cm}^{-3}$. Pitts et al. (2009) and Pitts et al. (2011) used a wide range of N_{NAT} between 10^{-4} and 10^{-1} cm^{-3} based on in-situ measurements. In mountain waves, high NAT densities larger than 10^{-2} cm^{-3} have been measured (e.g., Carslaw et al.,

1998). Voigt et al. (2005) observed a NAT PSC with 10^{-4} cm^{-3} . Here we have had the benefit of the unusual NH year 2020 and have placed more emphasis on both hemispheres to derive a parameterization that better represents the remaining gas phase HNO_3 in the maximum number of years and for both hemispheres. However, it is important to emphasize that the model's
85 NAT parameterization is subject to multiple simplifications of complex microphysics. For instance, NAT particles in a grid box with a horizontal extent of about $100 \times 100 \text{ km}^2$ do not necessarily have the same radius but usually follow a multi-modal size distribution (e.g., Fahey, 2001). In addition, NAT particles are not allowed to change their size over time or to be transported while interacting with the atmosphere by nucleation and (re-)sublimation and not all of the supersaturated amount of gaseous HNO_3 will become NAT. Therefore, observed NAT particle abundances may not be the best guide for this parameter choice.

90 We performed sensitivity simulations varying the NAT number density within the observed range from 10^{-2} to 10^{-5} cm^{-3} in the current version 6 of WACCM. We also performed a simulation excluding stratospheric heterogeneous reactions, apart from the reaction $\text{N}_2\text{O}_5 + \text{H}_2\text{O}$, to evaluate the impact of heterogeneous processes on the gas-phase species. The $\text{N}_2\text{O}_5 + \text{H}_2\text{O}$ reaction rate is nearly independent of temperature, happens also on the background aerosols, does not directly affect the halogen chemistry and is important for the partitioning of reactive nitrogen in the atmosphere which is why we kept this reaction in that
95 simulation. Without this reaction, the chemistry in the model would be completely unrealistic. All simulations are summarized in Table 1. They cover the satellite period starting from 1979 to present.

2.2 MIPAS

The Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) operated in limb geometry on board the Environmental Satellite (Envisat) between July 2002 and April 2012 (Fischer et al., 2008). Envisat was placed in a sun-synchronous
100 polar orbit at an altitude of around 800 km with more than 14 orbits per day. MIPAS measured a variety of trace gases including HNO_3 , ClONO_2 and O_3 using a Fourier transform spectrometer in the infrared spectral range between 4.15 and $14.6 \mu\text{m}$ at tangent altitudes from 7 to 72 km (Fischer et al., 2008). The spatial resolution was approximately 3 km in the vertical and 30 km in the horizontal. The spectral resolution was 0.05 cm^{-1} between 2002 and March 2004. Due to technical issues with the satellite and a corresponding gap of measurements, MIPAS started measuring again in January 2005 with a reduced spectral
105 resolution of 0.12 cm^{-1} .

For this study, we use the V8 level 2 MIPAS retrievals from the IMKIAA processor (Kiefer et al., 2021, 2022; Von Clarmann et al., 2009) of HNO_3 , ClONO_2 and O_3 to compare their distributions with the WACCM simulations. The reported precision in the region between 30 and 150 hPa is 5 – 15 % (around 100 pptv) for HNO_3 (Sheese et al., 2016, 2017), 10 % (around 150 ppbv) for O_3 (Sheese et al., 2017; Kiefer et al., 2022) and 7 – 32 % (24 – 89 pptv) for ClONO_2 (Sheese et al., 2016;
110 Höpfner et al., 2007).

The method used to compare the MIPAS data with WACCM is based on the approach by Zambri et al. (2021). It takes advantage of WACCM's ability to provide output at the locations and times closest to the MIPAS profiles. In combination with interpolation of the MIPAS vertical levels to the WACCM altitudes, this enables a direct comparison of the two datasets. As a result, probability density functions (PDFs) can be evaluated in the spatiotemporal range of interest and compared to the
115 observations.

Since NAT formation and denitrification are strongly temperature-dependent and most efficient at the lowest temperatures, we compute PDFs for profiles inside the polar vortex, determined by MERRA2 using the Nash criterion (Nash et al., 1996). As the sedimentation of the NAT particles takes several weeks (Tabazadeh et al., 2001), the largest effects of denitrification can be expected to be most amplified after the local winter, which is why we restrict our analysis to the early local spring, i.e. 120 1 February to 15 March in the Northern Hemisphere and 1 September to 15 October in the Southern Hemisphere, and to the pressure range 30 to 150 hPa. Sensitivity studies by changing these two ranges did not affect the main message of the results shown in the next section. In order to make the datasets comparable, we remove the profiles from both WACCM and MIPAS data where negative values occur in the measurements. This does not have a significant effect on the results. The number of profiles depends on the area of the polar vortex and differs from year to year. Generally, the number of profiles is larger in the 125 SH due to the larger size of the SH polar vortex compared to the NH.

Zambri et al. (2021) used the Kolmogorov-Smirnov test to evaluate the difference in the PDFs between the model and observations, which uses the number of data points and the maximum difference in the cumulative density functions (CDFs) to check whether the distributions are distinguishable in a statistical sense related to a significance level α . Here, we use the maximum difference in the CDFs F of MIPAS and WACCM “max(d)” of the respective WACCM simulation to MIPAS directly 130 to evaluate the difference in the distributions (Zambri et al., 2021):

$$\max(d) = \max_x |F_{\text{MIPAS}}(x) - F_{\text{WACCM}}(x)| \quad (1)$$

with x being the binned trace gas VMR. A more detailed discussion can be found in Zambri et al. (2021).

2.3 Ozonesondes

We also compare the WACCM simulations to balloon-borne in-situ measurements of ozonesondes, made available by the 135 World Ozone and Ultraviolet Data Centre. They use an electrochemical concentration cell to measure ozone profiles with a precision of 3 to 5 % and an uncertainty of about $\pm 10\%$ in the pressure range of interest here (Smit et al., 2007). We use ozonesondes of two stations: Eureka in the NH (80.04 °N, 86.18 °W) and Syowa in the SH (69.00 °S, 39.58 °E).

The method to compare the WACCM simulations with the ozonesonde measurements is similar to Wilka et al. (2021): The data is compared to the daily averaged WACCM ozone concentration at the grid point closest to the respective station for 140 profiles within the polar vortex. Averages and spread of the differences between WACCM and the observations are evaluated.

3 Impact on the distribution of gas-phase species

In this section, we will analyze the influence of changing denitrification in WACCM on the trace gases HNO_3 , ClONO_2 and O_3 during early spring in both hemispheres. Generally, these species are expected to be influenced by a changed NAT number density because it leads to a redistribution of HNO_3 (lower HNO_3 at high altitudes and higher HNO_3 at lower altitudes). 145 During early spring, HNO_3 is photolyzed and forms NO_2 which combines with ClO to form ClONO_2 . ClO is responsible for part of the catalytic ozone depletion and deactivated by the reaction into ClONO_2 .




Figure 1. Daily vortex-averaged HNO_3 VMR profiles in the analyzed pressure range between 30 and 150 hPa during local spring for MIPAS (first row) and the WACCM simulations (other rows) in the NH (left two columns) and SH (right two columns) and for the years 2003 and 2011 (first and second column in the hemispheres). The number of profiles used to average each day depends on the availability of MIPAS data and the size of the polar vortex.

As a starting point, Fig. 1 shows vortex-mean profiles of the HNO_3 VMR during the local spring months for both hemispheres and the years 2003 and 2011 for MIPAS and all WACCM simulations of this study. Analysis of the other years of the MIPAS period showed similar results, which is why we focus on the two example years 2003 and 2011 here. The signature of denitrification is indicated in the MIPAS measurements (first row) by HNO_3 VMR decreasing over time at around 30 to 60 hPa, with corresponding increases due to reinitiation at the end of the shown timeseries at lower levels, like in 2011 in the NH (panel f). In the simulation with the current operational NAT number density of 10^{-2} cm^{-3} , HetAll.1e-2 in the second row of Fig. 1, the NAT particles do not get large enough for the simulated HNO_3 to compare well with the measured denitrification of the shown years. This means that larger HNO_3 VMRs than observed remain at higher altitudes at the end of the timeseries, whereas the VMR is underestimated at lower altitudes. In contrast to this, with the smallest tested number density of 10^{-5} cm^{-3} , HetAll.1e-5 in the last row, the HNO_3 VMRs become too low, which indicates that the NAT particles become too large. By using a number density of $5 \times 10^{-4} \text{ cm}^{-3}$ (HetAll.5e-4 in the third row), the differences compared to MIPAS HNO_3 are smallest and the altitude regions of denitrification and reinitiation are comparable to the measurements.

In order to generalize this comparison, Fig. 2 shows correlation plots of the HetAll simulations using different NAT number densities compared to the MIPAS measurements for the whole pressure range (30 – 150 hPa) and all MIPAS years in the NH and SH. The simulation HetAll.5e-4 (green) has a relatively compact correlation with MIPAS in both hemispheres, whereas the other simulations show a larger scatter. The simulation HetAll.1e-2 shows HNO_3 VMRs that are too large compared to the observations because of NAT particles that are too small. As already suggested by Fig. 1, the HNO_3 VMRs in HetAll.1e-5 are




Figure 2. Scatter plots of MIPAS and WACCM HNO_3 in (a) the NH and (b) the SH for the HetAll simulation using the three NAT number densities described in the text. The dots include all measurement and model data between 30 and 150 hPa. The dashed line corresponds to the one-by-one line.

too small compared to MIPAS as a result of too large NAT particles which lead to sedimentation below the shown pressure
 165 range. As this figure includes the data of all MIPAS years and the whole pressure range, we will investigate the PDFs of this
 pressure range in the following, using the approach based on Zambri et al. (2021).

Probability density functions of HNO_3 for the whole MIPAS period 2002 to 2012 and for the years 2003 and 2011 are shown
 in Fig. 3 within the polar vortex in both hemispheres. All individual years can be found in the supplement of this study (Fig.
 S1). The blue line corresponds to the simulation with the largest NAT density, the red line shows the simulation with smallest
 170 NAT density and the green and yellow lines show the simulations using $N_{\text{NAT}} = 5 \times 10^{-4} \text{ cm}^{-3}$, with the latter probing
 turning off all heterogeneous chemistry apart from $\text{N}_2\text{O}_5 + \text{H}_2\text{O}$. The black line corresponds to the MIPAS HNO_3 PDF.

Figure 3 demonstrates that the NAT density has a large impact on the PDFs in the model between 30 and 150 hPa. In
 HetAll.1e-2, larger HNO_3 values are more frequent compared to the other simulations in all the panels. In contrast to this,
 HetAll.1e-5 underestimates HNO_3 as manifested by a higher frequency at smaller values. The simulation with peak values
 175 closest to the MIPAS observations in all panels of Fig. 3 is HetAll.5e-4, indicating that heterogeneous chemistry is relevant and
 that $N_{\text{NAT}} = 5 \times 10^{-4} \text{ cm}^{-3}$ leads to PDFs comparable to MIPAS for all years, see also Fig. S1 in the supplement. However,
 the NAT parameterization in WACCM is not able to capture the minimum and maximum values, especially in the SH where
 VMRs as high as 12.5 ppbv have been measured by MIPAS whereas the largest values in HetAll.5e-4 are around 10 ppbv, e.g.
 in panel d of Fig. 3.




Figure 3. HNO_3 probability density functions within the polar vortex for MIPAS and the sensitivity simulations of WACCM for the NH (left column) and SH (right column) spring months in (a,d) the MIPAS period 2002 to 2012, (b,e) 2003 and (c,f) 2011. Vertical dashed lines show the average VMR of the respective simulations. The number of profiles used to derive the PDFs in each panel is denoted by “nprof”. The maximum differences between the CDFs of MIPAS and the respective WACCM simulation are denoted by “max(d)” in the panels.

180 The overall agreement is also shown by the mean values illustrated by the dashed vertical lines in the figure. The mean HNO_3 value for HetAll.5e-4 is closest to the MIPAS mean value in all the panels, too. In addition, the maximum difference “max(d)” of the 5e-4 simulations is smallest in the shown years compared to the other simulations.

In almost all years, heterogeneous chemistry leads to about 0.5 ppbv smaller HNO_3 mixing ratios in the NH, see also Fig. S1 in the supplement. In the SH, this effect is not as large as in the NH. In total, the effect of halogen heterogeneous chemistry
185 on HNO_3 is small. Mean and shape of the distributions are similar for noHetAll.5e-4 and HetAll.5e-4.

Since the number of profiles differs from year to year depending on the development of the polar vortex, more weight in the distributions of the whole period 2002 to 2012 (panels a and d in Fig. 3) is given to years with a larger polar vortex. This is why the distributions of all individual years are shown in the supplement (Figs. S1 to S3). As a summary, the max(d) values of the individual years are shown in Tables 2 and 3 for NH and SH, respectively. The simulation with the minimum difference
190 is highlighted for each year. Apart from some exceptions in the NH, the minimum difference occurs for the simulations using $N_{\text{NAT}} = 5 \times 10^{-4} \text{ cm}^{-3}$. This is another indication that this value improves denitrification and the distribution of HNO_3 in the polar vortex in both hemispheres.

Exceptions are the years 2004 and 2012 in the NH, where the simulation with the smallest NAT density (HetAll.1e-5) shows the minimum differences compared to the MIPAS distribution, see Table 2. In 2004, relatively warm temperatures in

Table 2. Maximum differences of WACCM to MIPAS HNO₃ CDFs (“max(d)” in the figures) for all years, 1 February to 15 March in the NH. Minimum values of each row are highlighted.

| Years | HetAll.1e-2 | HetAll.5e-4 | noHetAll.5e-4 | HetAll.1e-5 |
|-------------|-------------|--------------|---------------|--------------|
| 2002 – 2012 | 0.278 | 0.086 | 0.153 | 0.172 |
| 2003 | 0.379 | 0.105 | 0.306 | 0.246 |
| 2004 | 0.187 | 0.167 | 0.214 | 0.136 |
| 2005 | 0.283 | 0.119 | 0.190 | 0.296 |
| 2006 | 0.155 | 0.045 | 0.166 | 0.325 |
| 2007 | 0.297 | 0.194 | 0.302 | 0.220 |
| 2008 | 0.443 | 0.177 | 0.309 | 0.242 |
| 2009 | 0.148 | 0.095 | 0.193 | 0.141 |
| 2010 | 0.319 | 0.080 | 0.169 | 0.195 |
| 2011 | 0.385 | 0.087 | 0.128 | 0.225 |
| 2012 | 0.226 | 0.150 | 0.205 | 0.139 |

Table 3. Same as Table 2, but for the SH spring (1 September to 15 October).

| Years | HetAll.1e-2 | HetAll.5e-4 | noHetAll.5e-4 | HetAll.1e-5 |
|-------------|-------------|--------------|---------------|-------------|
| 2002 – 2012 | 0.453 | 0.099 | 0.078 | 0.410 |
| 2002 | 0.558 | 0.173 | 0.141 | 0.642 |
| 2003 | 0.477 | 0.099 | 0.105 | 0.452 |
| 2005 | 0.598 | 0.095 | 0.124 | 0.559 |
| 2006 | 0.553 | 0.141 | 0.210 | 0.366 |
| 2007 | 0.467 | 0.124 | 0.110 | 0.404 |
| 2008 | 0.495 | 0.112 | 0.089 | 0.432 |
| 2009 | 0.409 | 0.078 | 0.072 | 0.290 |
| 2010 | 0.419 | 0.156 | 0.129 | 0.425 |
| 2011 | 0.429 | 0.107 | 0.082 | 0.360 |

195 the NH stratosphere lead to small denitrification (Manney et al., 2005). As shown in Fig. 4a, this is reflected by the WACCM HNO₃ distributions which do not show large differences when changing the NAT density during that year. There seems to be a systematic difference between WACCM and MIPAS in this year, though, maybe due to a lower accuracy of the instrument or for instance due to a mismatch in the location of the polar vortex in the reanalysis data used for the model and the temperature history during that winter. The Arctic winter 2011/2012 was characterized by an early breakdown of the polar vortex at the end
200 of December (Roy and Kuttippurath, 2022), but there is some evidence of large NAT particles during the cold period during




Figure 4. Same as Fig. 3, but for the years (a) 2004 and (b) 2012 in the NH.

December 2011 (Woiwode et al., 2019). This is why the HNO_3 distributions change in WACCM and probably large NAT particles play a larger role in determining the distributions for that year, see Fig. 4b.

Moderate changes due to the different NAT densities can also be found in other species. Figure 5 shows the distributions of chlorine nitrate (ClONO_2) for the same years and simulations as in Fig. 3. Both HetAll.1e-2 and HetAll.5e-4 have similar distributions and max(d) values. Depending on the year, minimum max(d) values occur in both simulations. HetAll.1e-5 seems to slightly underestimate the ClONO_2 mixing ratios in both hemispheres, illustrated by higher frequencies of low mixing ratios.

As a result of larger denitrification, impacts on the maximum values in the ClONO_2 distributions can be seen in both hemispheres. While the maximum ClONO_2 mixing ratios in the NH are closer to the MIPAS distributions using $N_{\text{NAT}} = 10^{-5} \text{ cm}^{-3}$, the HetAll.5e-4 simulation captures the maximum values in the SH distributions.

There is also a clear signature of heterogeneous chemistry in the ClONO_2 distributions. The deactivation reaction forming ClONO_2



is usually faster than the deactivation reaction forming HCl



as long as NO_2 is available and the ozone VMRs are large enough (e.g., Kawa et al., 1992; Prather and Jaffe, 1990). At the edge regions of the polar vortex, some of the chlorine originally in HCl at the start of the winter shows up in ClONO_2 at




Figure 5. Same as Fig. 3, but for ClONO₂.

the end of the winter, known as the ClONO₂ “collar” (e.g., Toon et al., 1989; Douglass et al., 1995; Solomon et al., 2016). Therefore, while the simulation without heterogeneous chemistry noHetAll.5e-4 generally shows ClONO₂ VMRs that are too low compared to MIPAS, it is notable that the maximum values are increased by about 1 ppbv and then are comparable to MIPAS in all simulations with heterogeneous chemistry.

Similar impacts can be seen in the distributions of ozone (O₃), shown in Fig. 6. Ozone VMRs are about 1 ppmv too large in the SH for noHetAll.5e-4 compared to MIPAS. In the NH, the difference is smaller but ozone is still overestimated in the simulation without heterogeneous chemistry.

Wilka et al. (2021) showed that the agreement of WACCM model calculations of Arctic spring ozone losses in 2020 with ozonesonde data was heavily dependent on the denitrification parameterization used in the model, providing an important check. Figure 7 displays comparisons between WACCM and ozonesonde data at Syowa and Eureka for their respective spring seasons for all years between 2002-2012. A figure showing the comparison to Syowa without noHetAll.5e-4 can be found in the supplement (Fig. S4). Similar comparisons are provided for the South Pole in supplemental Fig. S5. Because local Antarctic ozone abundances can drop to values lower than 1 ppmv between about 15-20 km in years with large ozone losses (see e.g. Fig. 7a), care must be taken in interpreting percentage differences; we also provide absolute differences for that reason. The Syowa sondes agree considerably better with the model when the NAT number density of $5 \times 10^{-4} \text{ cm}^{-3}$ or less is used, compared to larger values of this parameter. At Eureka, results are far less sensitive due to smaller ozone losses, except in very cold years (like 2020 shown in Wilka et al., 2021).




Figure 6. Same as Fig. 3, but for O₃.

235 In summary, the results in this section showed that the diagnostic parameterization of NAT particles in WACCM is able to reflect the main features in the three key gas-phase species HNO₃, ClONO₂ and ozone. When using a reduced NAT particle number density of $N_{\text{NAT}} = 5 \times 10^{-4} \text{ cm}^{-3}$ compared to the standard WACCM setup, the difference in the cumulative density functions is minimized for almost all spring months of the MIPAS period 2002 to 2012 and the bias between 30 and 150 hPa in comparison to the ozonesondes is reduced. This indicates that $N_{\text{NAT}} = 5 \times 10^{-4} \text{ cm}^{-3}$ improves the distribution of these

240 species in the polar regions and that it should be used in future WACCM simulations.

4 Discussion and Conclusions

In this study, we investigated the impact of the NAT number density (N_{NAT}) on the distributions of key gaseous species in the WACCM model. We compared probability density functions within the polar vortex with the MIPAS satellite instrument which operated between 2002 and 2012. As a measure of the difference between the distributions of WACCM and MIPAS, we used

245 the maximum difference, max(d), between the cumulative density functions. To identify the impact of NAT on the distributions of the species, we performed sensitivity simulations varying N_{NAT} between 10^{-2} and 10^{-5} cm^{-3} . In the method applied here to compare the satellite measurements with the model, negative values in the satellite retrievals are cut off, which is why it cannot be used for noisy data (such as N₂O₅ and ClO in MIPAS) and we restricted the analysis to the gaseous species HNO₃, ClONO₂ and O₃.




Figure 7. Mean and interquartile (0.25 and 0.75) range of (a,d) the volume mixing ratio and (b,e) relative differences and (c,f) the absolute differences between the respective WACCM simulations and two ozonesonde stations for profiles within the polar vortex using the same definition as in the previous analysis. The first row shows Syowa Station in the Antarctic and the second row shows Eureka Station in the Arctic. For WACCM, the grid point closest to the station is chosen. Average and distributions are taken for all profiles within the MIPAS period and the same months as for the satellite comparison. The left and right columns show the relative and absolute difference to the measurements, respectively.

250 The diagnostic parameterization of NAT in the WACCM model was shown to be able to reproduce the general shape of the MIPAS distributions for almost all years of the MIPAS period in the polar vortices of both hemispheres. There was a general overestimation of gaseous HNO₃, though, when using the standard N_{NAT} of 10^{-2} cm^{-3} . Reducing this number concentration to $5 \times 10^{-4} \text{ cm}^{-3}$ was shown to also reduce the differences in the HNO₃ distributions between MIPAS and WACCM for almost all spring seasons in both hemispheres during the MIPAS period. Changes in the distributions of ClONO₂ and O₃
 255 due to the new value of N_{NAT} were either negligible or the differences were reduced. Mean differences between ozonesonde profiles and the WACCM simulations at the grid point closest to the sonde stations Syowa and Eureka were also shown to be decreased when reducing N_{NAT} . Therefore, this study suggests the use of $N_{\text{NAT}} = 5 \times 10^{-4} \text{ cm}^{-3}$ for future simulations. This new value is within the range of measured NAT number densities.

A simplified parameterization as applied in WACCM cannot capture all features of the distributions of all years. This was demonstrated by the years 2004 and 2012 where differences between the MIPAS and WACCM distributions were increased. Further, there are many uncertainties in the detailed chemistry from one year to another that may also be important for denitrification and ozone loss, including for example volcanic particle inputs, gravity and planetary wave amplitudes, and changes in circulation, to name only a few. It can also be seen that the smallest number density of 10^{-5} cm^{-3} is comparable to measurements during the exceptionally cold northern winter 2019/2020 (Wilka et al., 2021). In addition, although overall differences in the PDFs are reduced with $N_{\text{NAT}} = 5 \times 10^{-4} \text{ cm}^{-3}$, the Kolmogorov-Smirnov test, used e.g. by Zambri et al. (2021) with a similar approach as in this study, will fail if not using a significance level α that is set to a very small value ($O(10^{-70})$). Therefore, although the differences are decreased, the distributions of MIPAS and WACCM are still different in a statistical sense, probably due to the simplifications in the NAT parameterization of WACCM.

Future studies could apply this methodology to other long-term satellite measurements to evaluate the usage of this new NAT number density for further years, although their precision has to be low enough to allow for comparisons with individual profiles. Nevertheless, as a follow-up of the publication by Wilka et al. (2021), we recommend using $N_{\text{NAT}} = 5 \times 10^{-4} \text{ cm}^{-3}$ for future simulations with WACCM, indicating that NAT rocks play an important role, especially in the NH where we saw the largest impact on changing the NAT number density.

Code and data availability. Instructions for the access to MIPAS data can be found at <https://www.imk-asf.kit.edu/english/308.php> (last access on 18 November 2022). Instructions how to download CESM version 2 can be found in Danabasoglu et al. (2020). Ozonezone station data are publicly available through the World Ozone and Ultraviolet Data Center (WOUDC; WMO/GAW Ozone Monitoring Community, 2022). The scripts and data needed to create the figures of this paper can be found at https://g-27eb33.7a577b.6fbd.data.globus.org/ACP_Weimer_2022/ACP_Weimer_2022.tar.gz (last access on 7 December 2022).

Author contributions. MW added the comparison with MIPAS and wrote the first draft of the manuscript. DEK performed the simulations with WACCM used for the comparisons. CW added the comparison to ozonesondes. All authors contributed to prepare the manuscript.

Competing interests. The authors declare that they have no competing interests.

Acknowledgements. Michael Weimer and Douglas E. Kinnison were funded in part by NASA grant 80NSSC19K0952. Susan Solomon was funded in part by grant no. 1906719 of the U.S. National Science Foundation (NSF). We would like to thank Brian Zambri for initiating this work and helping with the comparison to MIPAS. This material is based upon work supported by National Center for Atmospheric Research (NCAR), which is a major facility sponsored by NSF under the Cooperative Agreement 1852977. The CESM project is supported primarily by NSF. We would like to acknowledge support from the Svante cluster at MIT used for the comparison of model and the data,

see <https://svante.mit.edu/intro.html> (last access 18 November 2022) for more information. We would like to acknowledge high-performance computing support from Cheyenne and Casper provided by NCAR's Computational and Information Systems Laboratory (2019), sponsored by NSF.

290 References

- Adriani, A., Massoli, P., Di Donfrancesco, G., Cairo, F., Moriconi, M. L., and Snels, M.: Climatology of polar stratospheric clouds based on lidar observations from 1993 to 2001 over McMurdo Station, Antarctica, *J. Geophys. Res.: Atmospheres*, 109, 1–17, <https://doi.org/10.1029/2004JD004800>, 2004.
- 295 Carslaw, K. S., Luo, B. P., Clegg, S. L., Peter, T., Brimblecombe, P., and Crutzen, P. J.: Stratospheric aerosol growth and HNO₃ gas phase depletion from coupled HNO₃ and water uptake by liquid particles, *Geophys. Res. Lett.*, 21, 2479–2482, <https://doi.org/10.1029/94GL02799>, 1994.
- Carslaw, K. S., Wirth, M., Tsias, A., Luo, B. P., Dörnbrack, A., Leutbecher, M., Volkert, H., Renger, W., Bacmeister, J. T., and Peter, T.: Particle microphysics and chemistry in remotely observed mountain polar stratospheric clouds, *J. of Geophys. Res.: Atmospheres*, 103, 5785–5796, <https://doi.org/https://doi.org/10.1029/97JD03626>, 1998.
- 300 Carslaw, K. S., Kettleborough, J. A., Northway, M. J., Davies, S., Gao, R.-S., Fahey, D. W., Baumgardner, D. G., Chipperfield, M. P., and Kleinböhl, A.: A vortex-scale simulation of the growth and sedimentation of large nitric acid hydrate particles, *J. Geophys. Res.: Atmospheres*, 107, SOL 43–1–SOL 43–16, <https://doi.org/https://doi.org/10.1029/2001JD000467>, 2002.
- Computational and Information Systems Laboratory: Cheyenne: HPE/SGI ICE XA System (University Community Computing), Tech. rep., National Center for Atmospheric Research, Boulder, CO, <https://doi.org/10.5065/D6RX99HX>, 2019.
- 305 Considine, D. B., Douglass, A. R., Connell, P. S., Kinnison, D. E., and Rotman, D. A.: A polar stratospheric cloud parameterization for the global modeling initiative three-dimensional model and its response to stratospheric aircraft, *J. Geophys. Res.: Atmospheres*, 105, 3955–3973, <https://doi.org/10.1029/1999JD900932>, 2000.
- Danabasoglu, G., Lamarque, J. F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *J. Adv. Model. Earth Syst.*, 12, e2019MS001 916, <https://doi.org/10.1029/2019MS001916>, 2020.
- 310 Douglass, A. R., Schoeberl, M. R., Stolarski, R. S., Waters, J. W., Russell III, J. M., Roche, A. E., and Massie, S. T.: Interhemispheric differences in springtime production of HCl and ClONO₂ in the polar vortices, *J. Geophys. Res.: Atmospheres*, 100, 13 967–13 978, <https://doi.org/https://doi.org/10.1029/95JD00698>, 1995.
- Drdla, K. and Müller, R.: Temperature thresholds for chlorine activation and ozone loss in the polar stratosphere, *Ann. Geophys.*, 30, 1055–1073, <https://doi.org/10.5194/ANGE0-30-1055-2012>, 2012.
- Fahey, D. W.: The Detection of Large HNO₃-Containing Particles in the Winter Arctic Stratosphere, *Science*, 291, 1026–1031, <https://doi.org/10.1126/science.1057265>, 2001.
- 320 Fahey, D. W., Kelly, K. K., Ferry, G. V., Poole, L. R., Wilson, J. C., Murphy, D. M., Loewenstein, M., and Chan, K. R.: In situ measurements of total reactive nitrogen, total water, and aerosol in a polar stratospheric cloud in the Antarctic, *J. Geophys. Res.*, 94, 11 299, <https://doi.org/10.1029/JD094iD09p11299>, 1989.
- Fischer, H., Birk, M., Blom, C., Carli, B., Carlotti, M., von Clarmann, T., Delbouille, L., Dudhia, A., Ehhalt, D., Endemann, M., Flaud, J. M., Gessner, R., Kleinert, A., Koopman, R., Langen, J., López-Puertas, M., Mosner, P., Nett, H., Oelhaf, H., Perron, G., Remedios, J.,
- 325

- Ridolfi, M., Stiller, G., and Zander, R.: MIPAS: an instrument for atmospheric and climate research, *Atmos. Chem. Phys.*, 8, 2151–2188, <https://doi.org/10.5194/acp-8-2151-2008>, 2008.
- Fueglistaler, S., Luo, B. P., Voigt, C., Carslaw, K. S., and Peter, T.: NAT-rock formation by mother clouds: A microphysical model study, *Atmos. Chem. Phys.*, 2, 99–101, <https://doi.org/10.5194/ACP-2-93-2002>, 2002.
- 330 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Clim.*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 335 Gettelman, A., Mills, M. J., Kinnison, D. E., Garcia, R. R., Smith, A. K., Marsh, D. R., Tilmes, S., Vitt, F., Bardeen, C. G., McInerny, J., Liu, H. L., Solomon, S. C., Polvani, L. M., Emmons, L. K., Lamarque, J. F., Richter, J. H., Glanville, A. S., Bacmeister, J. T., Phillips, A. S., Neale, R. B., Simpson, I. R., DuVivier, A. K., Hodzic, A., and Randel, W. J.: The Whole Atmosphere Community Climate Model Version 6 (WACCM6), *J. Geophys. Res.: Atmospheres*, 124, 12 380–12 403, <https://doi.org/10.1029/2019JD030943>, 2019.
- Groß, J.-U., Günther, G., Müller, R., Konopka, P., Bausch, S., Schlager, H., Voigt, C., Volk, C. M., and Toon, G. C.: Simulation of denitrification and ozone loss for the Arctic winter 2002/2003, *Atmos. Chem. Phys.*, 5, 1437–1448, <https://doi.org/10.5194/acp-5-1437-2005>, 2005.
- Hanson, D. and Mauersberger, K.: Laboratory studies of the nitric acid trihydrate: Implications for the south polar stratosphere, *Geophysical Research Letters*, 15, 855–858, <https://doi.org/https://doi.org/10.1029/GL015i008p00855>, 1988.
- Höpfner, M., Larsen, N., Spang, R., Luo, B. P., Ma, J., Svendsen, S. H., Eckermann, S. D., Knudsen, B., Massoli, P., Cairo, F., Stiller, G., 345 v. Clarmann, T., and Fischer, H.: MIPAS detects Antarctic stratospheric belt of NAT PSCs caused by mountain waves, *Atmos. Chem. Phys.*, 6, 1221–1230, <https://doi.org/10.5194/acp-6-1221-2006>, 2006.
- Höpfner, M., Von Clarmann, T., Fischer, H., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Milz, M., Steck, T., Stiller, G. P., Bernath, P., Blom, C. E., Blumenstock, T., Boone, C., Chance, K., Coffey, M. T., Friedl-Vallon, F., Griffith, D., Hannigan, J. W., Hase, F., Jones, N., Jucks, K. W., Keim, C., Kleinert, A., Kouker, W., Liu, G. Y., Mahieu, E., Mellqvist, J., Mikuteit, S., Notholt, 350 J., Oelhaf, H., Piesch, C., Reddman, T., Ruhnke, R., Schneider, M., Strandberg, A., Toon, G., Walker, K. A., Warneke, T., Wetzell, G., Wood, S., and Zander, R.: Validation of MIPAS ClONO₂ measurements, *Atmos. Chem. Phys.*, 7, 257–281, <https://doi.org/10.5194/ACP-7-257-2007>, 2007.
- Höpfner, M., Deshler, T., Pitts, M., Poole, L., Spang, R., Stiller, G., and von Clarmann, T.: The MIPAS/Envisat climatology (2002–2012) of polar stratospheric cloud volume density profiles, *Atmos. Meas. Tech.*, 11, 5901–5923, <https://doi.org/10.5194/amt-11-5901-2018>, 2018.
- 355 Kawa, S. R., Fahey, D. W., Heidt, L. E., Pollock, W. H., Solomon, S., Anderson, D. E., Loewenstein, M., Proffitt, M. H., Margitan, J. J., and Chan, K. R.: Photochemical partitioning of the reactive nitrogen and chlorine reservoirs in the high-latitude stratosphere, *J. Geophys. Res.: Atmospheres*, 97, 7905–7923, <https://doi.org/https://doi.org/10.1029/91JD02399>, 1992.
- Kelly, K. K., Tuck, A. F., Murphy, D. M., Proffitt, M. H., Fahey, D. W., Jones, R. L., McKenna, D. S., Loewenstein, M., Podolske, J. R., Strahan, S. E., Ferry, G. V., Chan, K. R., Vedder, J. F., Gregory, G. L., Hypes, W. D., McCormick, M. P., Browell, E. V., and 360 Heidt, L. E.: Dehydration in the lower Antarctic stratosphere during late winter and early spring, 1987, *J. Geophys. Res.*, 94, 11 317, <https://doi.org/10.1029/JD094iD09p11317>, 1989.

- Kiefer, M., von Clarmann, T., Funke, B., García-Comas, M., Glatthor, N., Grabowski, U., Kellmann, S., Kleinert, A., Laeng, A., Linden, A., López-Puertas, M., Marsh, D. R., and Stiller, G. P.: IMK/IAA MIPAS temperature retrieval version 8: nominal measurements, *Atmos. Meas. Tech.*, 14, 4111–4138, <https://doi.org/10.5194/amt-14-4111-2021>, 2021.
- 365 Kiefer, M., von Clarmann, T., Funke, B., Garcia-Comas, M., Glatthor, N., Grabowski, U., Höpfner, M., Kellmann, S., Laeng, A., Linden, A., Lopez-Puertas, M., and Stiller, G.: Version 8 IMK/IAA MIPAS ozone profiles: nominal observation mode, *Atmos. Meas. Tech. Discuss.*, 2022, 1–30, <https://doi.org/10.5194/amt-2022-257>, under review, 2022.
- Kinnison, D. E., Brasseur, G. P., Walters, S., Garcia, R. R., Marsh, D. R., Sassi, F., Harvey, V. L., Randall, C. E., Emmons, L., Lamarque, J. F., Hess, P., Orlando, J. J., Tie, X. X., Randel, W., Pan, L. L., Gettelman, A., Granier, C., Diehl, T., Niemeier, U., and Simmons, A. J.:
 370 Sensitivity of chemical tracers to meteorological parameters in the MOZART-3 chemical transport model, *J. Geophys. Res.: Atmospheres*, 112, 20302, <https://doi.org/10.1029/2006JD007879>, 2007.
- Kirner, O., Ruhnke, R., Buchholz-Dietsch, J., Jöckel, P., Brühl, C., and Steil, B.: Simulation of polar stratospheric clouds in the chemistry-climate-model EMAC via the submodel PSC, *Geosci. Model Dev.*, 4, 169–182, <https://doi.org/10.5194/gmd-4-169-2011>, 2011.
- Manney, G. L., Krüger, K., Sabutis, J. L., Sena, S. A., and Pawson, S.: The remarkable 2003–2004 winter and other recent warm winters in
 375 the Arctic stratosphere since the late 1990s, *J. Geophys. Res.: Atmospheres*, 110, 1–14, <https://doi.org/10.1029/2004JD005367>, 2005.
- Molleker, S., Borrmann, S., Schlager, H., Luo, B., Frey, W., Klingebiel, M., Weigel, R., Ebert, M., Mitev, V., Matthey, R., Woiwode, W., Oelhaf, H., Dörnbrack, A., Stratmann, G., Groöß, J.-U., Günther, G., Vogel, B., Müller, R., Krämer, M., Meyer, J., and Cairo, F.: Microphysical properties of synoptic-scale polar stratospheric clouds: in situ measurements of unexpectedly large HNO₃-containing particles in the Arctic vortex, *Atmos. Chem. Phys.*, 14, 10785–10801, <https://doi.org/10.5194/acp-14-10785-2014>, 2014.
- 380 Nash, E. R., Newman, P. A., Rosenfield, J. E., and Schoeberl, M. R.: An objective determination of the polar vortex using Ertel’s potential vorticity, *J. Geophys. Res.: Atmospheres*, 101, 9471–9478, <https://doi.org/10.1029/96JD00066>, 1996.
- Peter, T.: Microphysics and heterogeneous chemistry of polar stratospheric clouds, *Annu. Rev. Phys. Chem.*, 48, 785–822, <https://doi.org/10.1146/annurev.physchem.48.1.785>, 1997.
- Pitts, M. C., Poole, L. R., and Thomason, L. W.: CALIPSO polar stratospheric cloud observations: second-generation detection algorithm
 385 and composition discrimination, *Atmos. Chem. Phys.*, 9, 7577–7589, <https://doi.org/10.5194/acp-9-7577-2009>, 2009.
- Pitts, M. C., Poole, L. R., Dörnbrack, A., and Thomason, L. W.: The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective, *Atmos. Chem. Phys.*, 11, 2161–2177, <https://doi.org/10.5194/acp-11-2161-2011>, 2011.
- Pitts, M. C., Poole, L. R., and Gonzalez, R.: Polar stratospheric cloud climatology based on CALIPSO spaceborne lidar measurements from 2006 to 2017, *Atmos. Chem. Phys.*, 18, 10881–10913, <https://doi.org/10.5194/acp-18-10881-2018>, 2018.
- 390 Prather, M. and Jaffe, A. H.: Global impact of the Antarctic ozone hole: Chemical propagation, *J. Geophys. Res.: Atmospheres*, 95, 3473–3492, <https://doi.org/10.1029/JD095iD04p03473>, 1990.
- Roy, R. and Kuttippurath, J.: The dynamical evolution of Sudden Stratospheric Warmings of the Arctic winters in the past decade 2011–2021, *SN Applied Sciences*, 4, 1–12, <https://doi.org/10.1007/S42452-022-04983-4/TABLES/1>, 2022.
- Santee, M. L., Lambert, A., Read, W. G., Livesey, N. J., Cofield, R. E., Cuddy, D. T., Daffer, W. H., Drouin, B. J., Froidevaux, L., Fuller, R. A., Jarnot, R. F., Knosp, B. W., Manney, G. L., Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Waters, J. W.,
 395 Muscari, G., de Zafra, R. L., Dibb, J. E., Fahey, D. W., Popp, P. J., Marcy, T. P., Jucks, K. W., Toon, G. C., Stachnik, R. A., Bernath, P. F., Boone, C. D., Walker, K. A., Urban, J., and Murtagh, D.: Validation of the Aura Microwave Limb Sounder HNO₃ measurements, *J. Geophys. Res.: Atmospheres*, 112, 24–40, <https://doi.org/10.1029/2007JD008721>, 2007.

- Schoeberl, M. R., Lait, L. R., Newman, P. A., and Rosenfield, J. E.: The structure of the polar vortex, *J. Geophys. Res.: Atmospheres*, 97, 400 7859–7882, <https://doi.org/10.1029/91JD02168>, 1992.
- Sheese, P. E., Walker, K. A., Boone, C. D., McLinden, C. A., Bernath, P. F., Bourassa, A. E., Burrows, J. P., Degenstein, D. A., Funke, B., Fussen, D., Manney, G. L., Thomas McElroy, C., Murtagh, D., Randall, C. E., Raspollini, P., Rozanov, A., Russell, J. M., Suzuki, M., Shiotani, M., Urban, J., Von Clarmann, T., and Zawodny, J. M.: Validation of ACE-FTS version 3.5 NO_y species profiles using correlative satellite measurements, *Atmos. Meas. Tech.*, 9, 5781–5810, <https://doi.org/10.5194/AMT-9-5781-2016>, 2016.
- 405 Sheese, P. E., Walker, K. A., Boone, C. D., Bernath, P. F., Froidevaux, L., Funke, B., Raspollini, P., and von Clarmann, T.: ACE-FTS ozone, water vapour, nitrous oxide, nitric acid, and carbon monoxide profile comparisons with MIPAS and MLS, *J. Quant. Spectrosc. Radiat. Transf.*, 186, 63–80, <https://doi.org/10.1016/J.JQSRT.2016.06.026>, 2017.
- Smit, H. G. J., Straeter, W., Johnson, B. J., Oltmans, S. J., Davies, J., Tarasick, D. W., Hoegger, B., Stubi, R., Schmidlin, F. J., Northam, T., Thompson, A. M., Witte, J. C., Boyd, I., and Posny, F.: Assessment of the performance of ECC-ozone sondes under quasi-flight conditions
410 in the environmental simulation chamber: Insights from the Juelich Ozone Sonde Intercomparison Experiment (JOSIE), *J. Geophys. Res.: Atmospheres*, 112, <https://doi.org/https://doi.org/10.1029/2006JD007308>, 2007.
- Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37, 275–316, <https://doi.org/10.1029/1999RG900008>, 1999.
- Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone, *Nature*, 321, 755–758,
415 <https://doi.org/10.1038/321755a0>, 1986.
- Solomon, S., Kinnison, D., Bandoro, J., and Garcia, R.: Simulation of polar ozone depletion: An update, *J. Geophys. Res.: Atmospheres*, 120, 7958–7974, <https://doi.org/10.1002/2015JD023365>, 2015.
- Solomon, S., Kinnison, D., Garcia, R. R., Bandoro, J., Mills, M., Wilka, C., Neely, R. R., Schmidt, A., Barnes, J. E., Vernier, J. P., and Höpfner, M.: Monsoon circulations and tropical heterogeneous chlorine chemistry in the stratosphere, *Geophys. Res. Lett.*, 43, 624–12,
420 <https://doi.org/10.1002/2016GL071778>, 2016.
- Spang, R., Hoffmann, L., Müller, R., Groöß, J.-U., Tritscher, I., Höpfner, M., Pitts, M., Orr, A., and Riese, M.: A climatology of polar stratospheric cloud composition between 2002 and 2012 based on MIPAS/Envisat observations, *Atmos. Chem. Phys.*, 18, 5089–5113, <https://doi.org/10.5194/acp-18-5089-2018>, 2018.
- Tabazadeh, A., Jensen, E. J., Toon, O. B., Drdla, K., and Schoeberl, M. R.: Role of the Stratospheric Polar Freezing Belt in Denitrification,
425 *Science*, 291, 2591–2594, <https://doi.org/10.1126/science.1057228>, 2001.
- Toon, G. C., Farmer, C. B., Lowes, L. L., Schaper, P. W., Blavier, J.-F., and Norton, R. H.: Infrared aircraft measurements of stratospheric composition over Antarctica during September 1987, *J. Geophys. Res.: Atmospheres*, 94, 16 571–16 596, <https://doi.org/https://doi.org/10.1029/JD094iD14p16571>, 1989.
- Tritscher, I., Pitts, M. C., Poole, L. R., Alexander, S. P., Cairo, F., Chipperfield, M. P., Groöß, J., Höpfner, M., Lambert, A., Luo, B., Mollerker,
430 S., Orr, A., Salawitch, R., Snels, M., Spang, R., Woiwode, W., and Peter, T.: Polar Stratospheric Clouds: Satellite Observations, Processes, and Role in Ozone Depletion, *Rev. Geophys.*, 59, e2020RG000 702, <https://doi.org/10.1029/2020RG000702>, 2021.
- Voigt, C., Schreiner, J., Kohlmann, A., Zink, P., Mauersberger, K., Larsen, N., Deshler, T., Kroger, C., Rosen, J., Adriani, A., Cairo, F., Di Donfrancesco, G., Viterbin, M., Ovarlez, J., Ovarlez, H., David, C., and Dornbrack, A.: Nitric Acid Trihydrate (NAT) in Polar Stratospheric Clouds, *Science*, 290, 1756–1758, <https://doi.org/10.1126/SCIENCE.290.5497.1756>, 2000.

- 435 Voigt, C., Schlager, H., Luo, B. P., Dörnbrack, A., Roiger, A., Stock, P., Curtius, J., Vössing, H., Borrmann, S., Davies, S., Konopka, P., Schiller, C., Shur, G., and Peter, T.: Nitric Acid Trihydrate (NAT) formation at low NAT supersaturation in Polar Stratospheric Clouds (PSCs), *Atmos. Chem. Phys.*, 5, 1371–1380, <https://doi.org/10.5194/acp-5-1371-2005>, 2005.
- Von Clarmann, T., Höpfner, M., Kellmann, S., Linden, A., Chauhan, S., Funke, B., Grabowski, U., Glatthor, N., Kiefer, M., Schieferdecker, T., Stiller, G. P., and Versick, S.: Retrieval of temperature, H₂O, O₃, HNO₃, CH₄, N₂O, ClONO₂ and ClO from MIPAS reduced resolution
440 nominal mode limb emission measurements, *Atmos. Meas. Tech.*, 2, 159–175, <https://doi.org/10.5194/AMT-2-159-2009>, 2009.
- Waibel, A. E., Peter, T., Carslaw, K. S., Oelhaf, H., Wetzel, G., Crutzen, P. J., Pöschl, U., Tsias, A., Reimer, E., and Fischer, H.: Arctic Ozone Loss Due to Denitrification, *Science*, 283, 2064–2069, <https://doi.org/10.1126/science.283.5410.2064>, 1999.
- Wegner, T., Kinnison, D. E., Garcia, R. R., and Solomon, S.: Simulation of polar stratospheric clouds in the specified dynamics version of the whole atmosphere community climate model, *J. Geophys. Res.: Atmospheres*, 118, 4991–5002, <https://doi.org/10.1002/jgrd.50415>,
445 2013.
- Weimer, M., Buchmüller, J., Hoffmann, L., Kirner, O., Luo, B., Ruhnke, R., Steiner, M., Tritscher, I., and Braesicke, P.: Mountain-wave-induced polar stratospheric clouds and their representation in the global chemistry model ICON-ART, *Atmos. Chem. Phys.*, 21, 9515–9543, <https://doi.org/10.5194/acp-21-9515-2021>, 2021.
- Wespes, C., Ronsmans, G., Clarisse, L., Solomon, S., Hurtmans, D., Clerbaux, C., and Coheur, P.-F.: Polar stratospheric nitric acid depletion
450 surveyed from a decadal dataset of IASI total columns, *Atmos. Chem. Phys.*, 22, 10993–11007, <https://doi.org/10.5194/ACP-22-10993-2022>, 2022.
- Wilka, C., Solomon, S., Kinnison, D., and Tarasick, D.: An Arctic ozone hole in 2020 if not for the Montreal Protocol, *Atmos. Chem. Phys.*, 21, 15771–15781, <https://doi.org/10.5194/ACP-21-15771-2021>, 2021.
- WMO/GAW Ozone Monitoring Community: World Meteorological Organization-Global Atmosphere Watch Program (WMO-GAW):
455 Ozone, World Ozone and Ultraviolet Radiation Data Centre (WOUDC) [data set], <https://doi.org/10.14287/10000001>, 2022.
- Wohlmann, I., Lehmann, R., and Rex, M.: The Lagrangian chemistry and transport model ATLAS: simulation and validation of stratospheric chemistry and ozone loss in the winter 1999/2000, *Geosci. Model Dev.*, 3, 585–601, <https://doi.org/10.5194/gmd-3-585-2010>, 2010.
- Woiwode, W., Grooß, J.-U., Oelhaf, H., Molleker, S., Borrmann, S., Ebersoldt, A., Frey, W., Gulde, T., Khaykin, S., Maucher, G., Piesch, C., and Orphal, J.: Denitrification by large NAT particles: the impact of reduced settling velocities and hints on particle characteristics,
460 *Atmos. Chem. Phys.*, 14, 11525–11544, <https://doi.org/10.5194/acp-14-11525-2014>, 2014.
- Woiwode, W., Höpfner, M., Bi, L., Pitts, M. C., Poole, L. R., Oelhaf, H., Molleker, S., Borrmann, S., Klingebiel, M., Belyaev, G., Ebersoldt, A., Griessbach, S., Grooß, J.-U., Gulde, T., Krämer, M., Maucher, G., Piesch, C., Rolf, C., Sartorius, C., Spang, R., and Orphal, J.: Spectroscopic evidence of large aspherical β -NAT particles involved in denitrification in the December 2011 Arctic stratosphere, *Atmos. Chem. Phys.*, 16, 9505–9532, <https://doi.org/10.5194/acp-16-9505-2016>, 2016.
- 465 Woiwode, W., Höpfner, M., Bi, L., Khosrawi, F., and Santee, M. L.: Vortex-Wide Detection of Large Aspherical NAT Particles in the Arctic Winter 2011/12 Stratosphere, *Geophys. Res. Lett.*, 46, 13420–13429, <https://doi.org/10.1029/2019GL084145>, 2019.
- Zambri, B., Kinnison, D. E., and Solomon, S.: Subpolar Activation of Halogen Heterogeneous Chemistry in Austral Spring, *Geophys. Res. Lett.*, 48, e2020GL090036, <https://doi.org/10.1029/2020GL090036>, 2021.
- Zhu, Y., Toon, O. B., Lambert, A., Kinnison, D. E., Brakebusch, M., Bardeen, C. G., Mills, M. J., and English, J. M.: Development of a Polar
470 Stratospheric Cloud Model within the Community Earth System Model using constraints on Type I PSCs from the 2010-2011 Arctic winter, *J. Adv. Model. Earth Syst.*, 7, 551–585, <https://doi.org/10.1002/2015MS000427>, 2015.