# Elucidating ozone and PM<sub>2.5</sub> pollution in Fenwei Plain reveals the cobenefits of controlling precursor gas emissions in winter haze

Chunshui Lin<sup>1</sup>, Ru-Jin Huang<sup>1,2,3,4\*</sup>, Haobin Zhong<sup>1,5</sup>, Jing Duan<sup>1</sup>, Zixi Wang<sup>3,6</sup>, Wei Huang<sup>1</sup>, and Wei Xu<sup>1</sup>

<sup>5</sup>School of Advanced Materials Engineering, Jiaxing Nanhu University, Jiaxing 314001, China. <sup>6</sup>State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

Correspondence to: Ru-Jin Huang (rujin.huang@ieecas.cn)

- 15 Abstract. Fenwei Plain, home to 50 million people in central China, is one of the most polluted regions in China. In 2018, Fenwei Plain is designated as one of the three key regions for the "Blue Sky Protection Campaign", along with the Beijing-Tianjin-Hebei (BTH) and Yangtze River Delta (YRD) regions. However, compared to BTH and YRD, our understanding of the current status of air pollution in the Fenwei Plain is limited partly due to a lack of detailed analysis of the transformation from precursor gases to secondary products including secondary organic aerosol (SOA) and ozone. Through the analysis of 7
- 20 years (2015-2021) of surface monitoring of the air pollutants in Xi'an, the largest city in the Fenwei Plain, we show that roughly 2/3 of the days exceeded either the PM<sub>2.5</sub> or the O<sub>3</sub> level-1 air quality standard, highlighting the severity of air pollution. Moreover, an increase in O<sub>3</sub> pollution in the winter haze was also revealed, due to the constantly elevated reactive oxygenated volatile organic compounds (OVOCs), and in particular formaldehyde with ozone formation potential of over 50 µg m<sup>-3</sup> in combination with the reduced NO<sub>2</sub>. The abrupt decrease of NO<sub>2</sub>, as observed during the lockdown in 2020, provided real-
- 25 world evidence of the control measures, targeting only NOx (70% decrease on average), were insufficient to reduce ozone pollution because reactive OVOCs remained constantly high in a VOC-limited regime. Model simulation results showed that with NO<sub>2</sub> reduction from 20-70%, the self-reaction rate between peroxy radicals, a pathway for SOA formation, was intensified by up to 75%, while the self-reaction rate was only reduced with a further reduction of VOCs of >50%. Therefore, a synergic reduction in PM<sub>2.5</sub> and O<sub>3</sub> pollution can only be achieved through a more aggressive reduction of their precursor gases. This
- 30 study elucidates the status of ozone and PM<sub>2.5</sub> pollution in one of the most polluted regions in China, revealing a general trend of increasing secondary pollution i.e., ozone and SOA in winter haze. Controlling precursor gas emissions is anticipated to curb both ozone and SOA formation which will benefit not just the Fenwei Plain but also other regions in China.

#### 1 Introduction

	The Fenwei Plain (about 760 km in length and 40-100 km in width) is the largest plain in the middle reaches of the Yellow
35	River. It is home to over 50 million people in central China, surrounded by the Chinese Loess Plateau to the northwest and
	Qinling Mountains to the south. The rapid growth and urbanization of Fenwei Plain are accompanied by air pollution that is
	characterized by high concentrations of fine particulate matter $(PM_{2.5})$ in the heating season and high concentrations of ground-
	level ozone in the warm season (Elser et al., 2016; Lin et al., 2021; Song et al., 2021; Lin et al., 2022; Lin et al., 2022). Recently,
	air pollution in Fenwei Plain has been found to be more severe than in the Beijing-Tianjin-Heibei (BTH) region, Yangtze River

40 Delta (YRD) region, and Pearl River Delta (PRD) region, making Fenwei Plain one of the most polluted regions in China (Cao and Cui, 2021). In addition to being the emission hotspot of air pollutants, the unique topography of the Fenwei Plain is

• <b>设置了格式:</b> 字体: 10 磅	
<b>设置了格式:</b> 字体: 10 磅	
<b>设置了格式:</b> 字体: 10 磅	

<sup>&</sup>lt;sup>5</sup> <sup>1</sup>State Key Laboratory of Loess and Quaternary Geology (SKLLQG), Center for Excellence in Quaternary Science and Global Change, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710061, China <sup>2</sup>Institute of Global Environmental Change, Xi'an Jiaotong University, Xi'an 710049, China <sup>3</sup>University of Chinese Academy of Sciences, Beijing 100049, China <sup>4</sup>Laoshan Laboratory, Oingdao 266061, China

favorable for accumulating the air pollutants inside the basin (Cao and Cui, 2021), rapidly building up high levels of air pollutants e.g., under typical cold-haze conditions in calm weather with a wind speed of less than 2 m s<sup>-1</sup>. In 2018, the Fenwei Plain was designated as one of the three key regions for the "Blue Sky Protection Campaign" (the other two are the BTH and

45 YRD regions). To evaluate the effectiveness of the clean air policies and to further develop cost-effective mitigation policies in the Fenwei Plain, a better understanding of the trend in pollution patterns, sources, and formation mechanism of key pollutants i.e., PM<sub>2.5</sub> and ozone, is required.

As the largest city in the Fenwei Plain, Xi'an city is home to over 12 million people, with severe winter haze pollution events being recorded frequently, featuring high PM2.5 concentration levels rivaling that in Beijing (Elser et al., 2016; Lin et al., 2021; Wang et al., 2022). However, ozone is generally viewed as a summertime problem, with very low concentrations in winter (Li

et al., 2019). Most ozone studies in the Fenwei Plain were conducted in summer, with few studies performed in winter (Song et al., 2021; Yan et al., 2021; Li et al., 2022). In a recent study, Li et al. (2021) show a tendency of increasing winter-spring ozone with winter haze events in the North China Plain due to the reduced NO<sub>x</sub> emission, while the emission of reactive volatile organic compounds (VOCs) remained constantly high. In particular, the increased association of high ozone with

- 55 winter haze events is highlighted during the strict lockdown period in January and February 2020, whereas the formation of secondary organic aerosol (SOA), a major fraction of PM<sub>2.5</sub> which is produced from the oxidation of VOCs, was found to be largely increased (He et al., 2020; Zhao et al., 2020; Duan et al., 2021; Li et al., 2021; Zhong et al., 2021). The city to nationwide lockdown provides a real-world experiment to study the impact of emission control, which mostly targeted NO<sub>x</sub> (Li et al., 2021), on the haze and ozone pollution, which, however, does not appear to be abating in the Fenwei Plain (Duan et al., 2021; Zhong et al., 2021).
  - Reactive VOCs and oxygenated VOCs (OVOCs) are key precursor gases of ozone and SOA (Li et al., 2022; Wang et al., 2022), the formation of which involves a series of photochemical reactions of VOC/OVOCs and NOx. Briefly, upon solar radiation at wavelength <424 nm, NO<sub>2</sub> is photolyzed in the atmosphere (Li et al., 2019; Li et al., 2022). Following the photolysis of NO<sub>2</sub>, the resulting O atom quickly becomes ozone, while NO<sub>2</sub> is formed from the oxidation of NO by peroxy
- 65 radicals, including the peroxy radical (RO<sub>2</sub>) and hydroperoxyl radical (HO<sub>2</sub>), which are initiated by the oxidation of VOCs/OVOCs by the hydroxyl radical (OH) (Wang et al., 2009; Wang et al., 2019; Li et al., 2022). High levels of NOx, as currently experienced in most Chinese cities, can contribute to radical termination (e.g., OH+NO<sub>2</sub>) (Li et al., 2022; Wang et al., 2022), while, in NOx-lean environments, RO<sub>2</sub> mainly reacts with other peroxy radicals (i.e., self-reactions), some of which can lead to SOA formation albeit not the only way (Zhao et al., 2018; Lyu et al., 2022). Previous studies in the Fenwei Plain
- 70 mostly focus on hydrocarbon-like VOCs, while the OVOCs were rarely studied partly due to the limitation of sampling instrumentation (Song et al., 2021; Li et al., 2022). However, OVOCs were found to be the dominant species in other urban areas in China, showing high ozone and SOA formation potential (Li et al., 2019; Luo et al., 2020; Li et al., 2021). With the development of the state-of-the-art instrument e.g., Vocus proton-transfer-reaction time-of-flight mass spectrometer (PTR-TOF) (Krechmer et al., 2018), a better characterization of OVOCs is crucial for the co-benefits of ozone and PM<sub>2.5</sub> reduction
- 75 in the Fenwei Plain. This is particularly important as the synergic control of ozone and PM<sub>2.5</sub> pollution is one of the key policies to improve air quality during the 14th 5-year Plan period (2021-2025). In this study, a trend analysis of PM<sub>2.5</sub> and ozone from 2015 to 2021 were performed to understand the current status of air pollution. To disentangle the meteorological impacts on the measured trend of PM<sub>2.5</sub> and ozone, a machine learning (Grange
- et al., 2018; Dai et al., 2021; Shi et al., 2021) based meteorological normalization technique was applied, revealing the trend
   of de-weathered PM<sub>2.5</sub> and ozone. Under the background of decreasing anthropogenic emissions, changes in the chemical composition of PM<sub>2.5</sub> and the organic aerosol (OA) factors were investigated based on the available studies using online aerosol mass spectrometry (Elser et al., 2016; Zhong et al., 2020; Duan et al., 2021; Duan et al., 2022). As a case study, the ozone formation potential of OVOCs was analyzed based on its reactivity (Carter, 2010), while the SOA formation potential was analyzed using a 0-D chemical box modeling (Wolfe et al., 2016) in different NO<sub>x</sub> and VOC reduction scenarios. Based on the

85 ground surface and satellite observation, in combination with the modeling results, the reduction of reactive VOCs/OVOCs for the co-benefits of ozone and PM<sub>2.5</sub> reduction are discussed.

## 2. Method

## 2.1 Surface measurement data

- Hourly O<sub>3</sub>, PM<sub>2.5</sub>, NO<sub>2</sub>, SO<sub>2</sub>, and CO were continuously monitored at 13 sites within the city center of Xi'an from 2015 to
   2021 (Fig. S1). Measurements of these pollutants were routinely managed by the China National Environmental Monitoring
   Centre. The distance between the sampling sites is up to 40 km. Despite the distance, the time series of PM<sub>2.5</sub> and O<sub>3</sub> monitored at different sites were well correlated with similar magnitude in concentrations (Fig. S2). The consistency of the measurements between different sampling sites confirmed the data quality. The data from 13 sites were averaged to get the city-wide mean concentration of each pollutant and were used to perform the trend analysis.
- 95 VOC and OVOCs were measured using a Vocus-PTR which was equipped with a Long Time-of-Flight (LToF) mass analyzer, from 24 January to 6 February 2021 along with a high-resolution Long Time-of-Flight Aerosol Mass Spectrometry (AMS). The equipped LToF mass analyzer of Vocus-PTR had a mass-resolving power of approximately 12,000. Vocus-PTR measured organic vapors with a wide range of volatilities using a low-pressure reagent ion source and focusing ion-molecule reactor (FIMR). FIMR focused ions to the central axis efficiently with the quadrupole radio frequency (RF) field inside, greatly
- 00 improving the detection efficiency of product ions. Moreover, no humidity dependence for sensitivity was found for Vocus-PTR since a high-water mixing ratio was applied in the FIMR. To protect the microchannel plate detector from degrading too quickly, the big segmented quadrupole (BSQ) was set up as a high-pass band filter because the focusing effect of RF fields caused the low mass signal to be very high. More instrument details are available in Krechmer et al. (2018). In this study, Vocus-PTR was calibrated with VOC/OVOC standard mixture (Table S1). The sensitivity of Vocus-PTR towards uncalibrated
- 05 VOC compounds was calculated from the kinetic rate constant following Krechmer et al. (2018). Uncertainties for the calibrated VOC/OVOC species were 15%, while the uncertainty was 30% for the uncalibrated VOC compounds (Table S2) with known reaction rate constant (k<sub>FTR</sub>) (https://kb.tofwerk.com/tofware/; last access: 1 March 2023). Due to the low transmission efficiency for low-mass molecules that caused high uncertainties in quantification, formaldehyde (HCHO) was scaled to the mean concentration of the surface HCHO based on satellite observation using the empirical relationship (Zhang
- 10 et al., 2012). Vocus-PTR data was only available in 2021 and was used as a case study to perform an analysis of ozone formation potential and 0-D box modeling of self-reaction rate between peroxy radicals in different scenarios. The chemical composition of non-refractory particulate matter (NR-PM) was measured using an AMS or an aerosol chemical speciation monitor (ACSM) at the old campus of the Institute of Earth Environment, Chinese Academy of Sciences (Table S3) in the winter of 2012-2014 (Zhong et al., 2020) (Elser et al., 2016), and more recently in 2018-2019 (Duan et al., 2022), 2020
- 15 (Duan et al., 2021), and 2021 (this study; Fig. S3). For AMS measurement, Elser et al. (2016) first deployed an AMS with a <u>PM<sub>2.5</sub> inlet and comprehensively evaluated its performance during the winter campaign in Xi'an. Based on the same design, a soot particle AMS (SP-AMS) equipped with a PM<sub>2.5</sub> inlet was recently deployed at the same sampling site, quantifying PM<sub>2.5</sub> chemical composition with high mass and temporal resolution (Lin et al., 2021; Duan et al., 2022). OA factors were apportioned using the Positive Matrix Factorization (PMF) with the Multilinear Engine (ME-2) (Elser et al., 2016; Zhong et al., 2016; Zhong et al., 2017).</u>
- 20 al., 2020; Duan et al., 2021; Duan et al., 2022), Similarly, OA source apportionment was performed for the 2021 dataset using the Igor Pro (WaveMetrics Inc.) - based interface of SoFi (version 8.2.1) (Canonaco et al., 2013). The apportioned factors explained the input OA matrix well (Fig. S4-S6).

2.2 Satellite and meteorological data

**设置了格式:** 下标

(	删除了:2.5
(	删除了:3
1	删除了: and 2014
Ì	删除了:
Ì	刪除了: (Zhong et al., 2020)
N	<b>设置了格式:</b> 下标
$\overline{\langle}$	<b>设置了格式:</b> 下标
Y	<b>设置了格式:</b> 下标

删除了: (Grange and Carslaw, 2019)

Vertical column densities (molecules cm<sup>-2</sup>) of tropospheric formaldehyde and NO<sub>2</sub> were obtained from Sentinel-5P Level-3 Near Real-Time dataset based on the observation of the TROPOspheric Monitoring Instrument (TROPOMI) using the Google Earth Engine (Gorelick et al., 2017). This dataset was used to study the spatiotemporal variation of reactive OVOCs as a response to strict lockdown measures implemented in 2020. As a comparison, satellite images were also obtained for the same period in 2019 and 2021.

Boundary layer height (blh) and downward ultraviolet (UV) radiation at the surface was accessed from European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis v5 (ERA5). Wind speed, wind direction, relative humidity, and air temperature were obtained from the Integrated Surface Database (ISD).

## 40 2.3 Meteorological normalization

35

A machine-learning based meteorological normalization using the random forest (<u>RF</u>) algorithm (Grange et al., 2018) was used to decouple the meteorological impacts on the observed O<sub>3</sub> and PM<sub>2.5</sub>. Firstly, the random forest was grown using the meteorological and time variables as input. The meteorological variables included the data obtained from ERA5 and ISD, while time variables included date unix, day julian, weekday, hour of the day, and day of the lunar year. 80% of the dataset was randomly selected to train the model, while 20% was used to validate the model. The "rmweather" R packages was applied

- during the random forest modeling (Grange et al., 2018). As shown in Fig. <u>\$7</u> and <u>\$8</u>, the predicted and measured O<sub>3</sub> and PM<sub>2.5</sub> were well correlated (R<sup>2</sup>>0.89), suggesting the developed model rebuilt the measurements very well. <u>RF</u>-based meteorological normalization was first introduced by Grange and Carslaw (2019). However, it is not straightforward
- to investigate the seasonal variation in the trends of de-weathered air pollutants using the proposed normalization method
   (Grange and Carslaw, 2019), Vu et al. (2019) enhanced the meteorological normalization procedure by repeatedly resampling
   the meteorological variables for a particular time point within a four-week period and the resampled variables were fed to the
   developed RF model. The meteorological normalization method proposed by Vu et al. (2019), is widely applied in different
   sampling sites (Dai et al., 2021); Shi et al., 2021). In this study, we resampled the meteorological variables by averaging the
- meteorological variables for the same time point across 7 years. Our proposed method was more straightforward and subject
   to less variation given that meteorological variables varied a lot over a four-week period (e.g., see the comparison between the resampled and observed temperatures in Fig. S19). Also, our method is less time-consuming. Therefore, the meteorological participation to a propose the period of the period
- normalization technology assumed the meteorological variables were invariant across the years. In this scenario, the predicted values represent the changes in air pollutant concentrations that were not affected by meteorological variables (i.e., deweathered pollutants).

## 60 2.4 0-D box modelling

65

Master Chemical Mechanism (MCM) v3.3.1 (Jenkin et al., 2003; Jenkin et al., 2019) was incorporated in a chemical box model using the framework for 0-D Atmospheric Modeling (F0AM) version 4.2.2 (Wolfe et al., 2016). Surface observational data were averaged for the overlapping period with OVOC measurements to get the diurnal profiles of each pollutant. The averaged air pollutants, including NO<sub>2</sub>, O<sub>3</sub>(not included when it was simulated, Fig. S10), CO, and the measured VOC/OVOCs (Table S2) were used to constrain the model at hourly resolution (base run). The box model reproduced the measured ozone reasonably well (Fig. S10) after optimizing the correction of the simulated photolysis rate. However, due to the uncertainties

of the box model which did not consider factors like regional transport and heterogenous chemistry, as well as the parameterized ozone deposition, the modelling results were only used to understand the atmospheric chemistry involving OVOCs by comparing different scenarios. Note that HCHQ, was not included when it was being simulated. Using the averaged diurnal profiles reduced modeling uncertainty caused by occasional data flaws and data gaps due to instrument maintenance. The photolysis frequencies were calculated based on TUV v5.2 solar spectra. For each scenario simulation, the box model was set to spin up for 48 hours. 删除了: The reduction in NO<sub>2</sub> as observed from the satellite image was used as a scenario input for subsequent 0-D box modeling analysis....

## 删除了: S3 删除了: S4

删除了: Based on the developed random forest model, meteorological normalization was performed by resampling the meteorological variables and predicting the values at a specific time point on a rolling basis. The resampled meteorological variables at a specific time point represented the averaged meteorological condition across the sampling years. The mean meteorological variables at a specific time were fed to the random forest model to predict the dependent variables. The predicted variables represent the deweathered pollutants, i.e., not affected by a specific meteorological condition over the sampling years, and, therefore, revealing the trend of air pollutants that were mainly affected by emission and/or chemistry (Dai et al., 2021; Grange et al., 2021; Shi et al., 2021).

<b>设置了格式:</b> 字体颜色: 文字 1
<b>设置了格式:</b> 字体颜色: 文字 1
( <b>设置了格式:</b> 英语(美国)
(删除了:,
(删除了: except for
(删除了: (which
(删除了: i
(删除了: the
删除了: target) were used to constrain the model at hourly resolution (base run)

## 3. Results and Discussion

## 3.1 Ozone pollution spreading into the late-winter haze season

Figure 1 shows the observed exceedance frequency (in days month<sup>-1</sup>) of China's National Ambient Air Quality Standard
 (NAAQS level-1) for 24-h PM<sub>2.5</sub> (35 μg m<sup>-3</sup>) and for maximum daily 8-hour (MDA8) ozone (100 μg m<sup>-3</sup>), averaged for 2015-2018 and 2019-2021 in the biggest city (i.e., Xi'an) in the Fenwei Plain in central China (Fig. S1). For the observed PM<sub>2.5</sub>, most exceedances were observed in the heating season from November to March, with over 2/3 of the days in individual months exceeding the 24-h PM<sub>2.5</sub> standard (Fig. 1a). In contrast, for surface ozone, most exceedances were observed in the warm season from May to August, with approximately 2/3 of the days in individual month exceeding the MDA8 ozone standard (Fig. 1b). Combined, over 2/3 of the days throughout the year (except October) exceeded either the ozone or the

PM<sub>2.5</sub> standard, reflecting the severe air pollution in Fenwei Plain (Fig. 1c), despite the decreasing trend found for PM<sub>2.5</sub> (Fig. S11). A similar trend was observed for the deweathered ozone and PM<sub>2.5</sub>, suggesting meteorological conditions were not affecting the trend significantly (Fig. S12). Moreover, in terms of exceedance of the NAAQS level-2 standard (i.e., 75 µg m<sup>-3</sup> for 24-h PM<sub>2.5</sub> and 160 µg m<sup>-3</sup> for MDA8 ozone), similar pollution patterns were observed, although the exceedance frequency was less due to the relatively loose standard (Fig. S13).

Compared to 2015-2018, the observed exceedances of PM<sub>2.5</sub> standard during the warm season in 2019-2021 decreased largely e.g., by over 10 days in August (Fig. 1a), while the differences in the observed exceedances of PM<sub>2.5</sub> standard were only marginal during the heating season in 2019-2021, i.e., less than 3 days for an individual month in winter. In particular, the exceedance for January in 2019-2021 even increased by approximately 1 day (Fig. 1a). Compared to PM<sub>2.5</sub>, the difference

in the observed exceedances of the ozone standard between 2015-2018 and 2019-2021 was small, with values mostly below 5 days month<sup>-1</sup> (Fig. 1b). In contrast to the slightly decreasing trend of PM<sub>2.5</sub>, the observed exceedances of ozone standard even increased by 1 4 days month<sup>-1</sup> for January and February in 2019-2021 when compared to 2015-2018 (Fig. 1b and Fig. S14). Therefore, while the winter particulate pollution does not appear abating in recent years in terms of frequency of exceedance of PM<sub>2.5</sub> standard, the spread of ozone pollution into the late winter (i.e., January and February) may aggravate the air pollution issue in the Fenwei Plain.

Figure 2 shows the observed and deweathered (using a machine learning-based deweathering technique; see Method section) diurnal cycle of PM<sub>2.5</sub> and O<sub>3</sub> in late winter (January-February) averaged over 2015-2018 and 2019-2021. For the observed PM<sub>2.5</sub>, elevated concentrations (>100 µg m<sup>-3</sup>) were observed from the evening to the next noon (20:00 - 12:00; Fig. 2a). The elevated concentration in the evening and morning was due to the primary emission from various sources and secondary

- 25 particulate formation (discussed in Sect. 3.3), coupled with a shallow boundary layer (Fig. S15) and the topography of the Fenwei Plain (Cao and Cui, 2021), favoring the build-up of air pollutants. However, PM<sub>2.5</sub> started to decrease at 12:00, reaching the lowest concentrations (<100 µg m<sup>-3</sup>) at 16:00-18: 00. The decrease in the afternoon was likely due to the reduced primary emission and increased boundary layer height (BLH). The increasing BLH in the afternoon diluted the air pollutants, which was confirmed by the decreased CO at the same time (Fig. S16). The ΔCO corrected PM<sub>2.5</sub> first showed an increase at noon
- 30 and decreased afterward (Fig. S16) probably because the PM<sub>2.5</sub> emissions and/or secondary formation were insufficient to counter the diluting effects in the afternoon. Compared to 2015-2018, the observed PM<sub>2.5</sub> in 2019-2021 showed a larger decrease (10 µg m<sup>-3</sup>) in the late evening and morning, while the decrease in the afternoon was less significant (~5 µg m<sup>-3</sup>; Fig. 2). A similar trend was also observed for the deweathered PM<sub>2.5</sub>, suggesting the reduction in PM<sub>2.5</sub> in recent years was mainly caused by emission reduction.
- Different from PM<sub>2.5</sub>, both the observed (Fig. 2b) and deweathered O<sub>3</sub> (Fig. 2d) showed small differences between 2019-2021 and 2015-2018 during the morning hours (00:00-12:00; sunrise at 8:00) when PM<sub>2.5</sub> showed the largest reduction during the same period. In contrast, the increase in O<sub>3</sub> concentration in 2019-2021 was most prominent in the afternoon when PM<sub>2.5</sub> reduction was less prominent. Given that the observed NO<sub>2</sub> also showed the largest reduction (20 µg m<sup>-3</sup>, Fig. <u>\$17</u>) in the

e	
<u>.</u>	<b>设置了格式:</b> 下标
ot	
-3	删除了: S5
у	
	删除了: S6
d	
e	
r,	
e	
5	
n	删除了: For January and February,
).	<b>设置了格式:</b> 下标
e	删除了:2
n	
d	
e	
у	
e	删除了:7

(	删除了:, creating a dispersion condition	
(	<b>设置了格式:</b> 下标	2
(	<b>设置了格式:</b> 下标	-

afternoon, the increase in O<sub>3</sub> in the afternoon for 2019-2021 was likely caused by the reduced titration effects, coupled with the reduced  $PM_{2.5}$  that scavenged less  $RO_x$  (OH+HO<sub>2</sub>+RO<sub>2</sub>) radicals. This is consistent with the VOC-limited regime for the O<sub>3</sub> formation in the Fenwei Plain for 2019-2021 (discussed in Section 3.4). In particular, for a VOC-limited regime, a moderate decrease in NO<sub>x</sub> emission is accompanied by an increase in O<sub>3</sub>.

## 150 3.2 OVOCs and ozone formation potential in the winter: a case study

To understand the formation processes of ozone in winter, precursor gases of VOCs and OVOCs were measured using a Vocus-PTR in Xi'an from 24 January 2021 to 6 February 2021 (Sect. 2.1). Table 1 shows the top 10 VOC/OVOC species that were quantified, along with the ozone formation potential (OFP) that was estimated using the maximum incremental reactivity (MIR) (Carter, 2010). Formaldehyde (HCHO) was the most dominant OVOC, with a mean concentration of 4.16 ppb. Its

- is ozone formation potential (OFP) was also the highest (54.4 μg m<sup>-3</sup>), reflecting its high reactivity. The important role of formaldehyde in ozone formation is consistent with the findings in the North China Plain (Li et al., 2021). Note that formaldehyde can be directly emitted from incomplete combustion of e.g., biomass and fossil fuel, while it can also be produced from the oxidation of reactive VOCs (Su et al., 2019).
- By constraining ozone and other VOCs/OVOCs (except formaldehyde) in a 0-D box model, the secondary formation of formaldehyde, fueled by photochemical reaction, was simulated (Fig. 3). Compared to the mixing ratio of the measured formaldehyde, the model reproduced the measured formaldehyde at a similar level of around 4.2 ppb, However, compared to the simulated formaldehyde, the elevated concentration of the observed formaldehyde in the evening and morning (22:00 – 8:00; Fig. 3) was likely associated with the primary emission from biomass and fossil fuel combustion, coupled with a shallow boundary layer (Fig. \$15). Note that the primary emission of formaldehyde as well as the regional transport was not considered
- in the box model. The modelled formaldehyde showed an increased mixing ratio at noon (Fig. 3a), corresponding to the increased local formation as indicated by the increased solar radiation and OH radical (Fig. 3b). This is because the box model only considered photochemical formation, not taking into account the scavenging effects of RO<sub>x</sub> (OH+HO<sub>2</sub>+RO<sub>2</sub>). High levels of PM<sub>2.5</sub> (> 80 µg m<sup>-3</sup>) were observed in the morning (Fig. 2), which were likely playing an important role in scavenging RO<sub>x</sub> radicals (Shao et al., 2021). Therefore, the modelled spike in formaldehyde at noon was not observed. Similarly, the highest concentration of ozone was coincident with the lowest level of PM<sub>2.5</sub> in the late afternoon (15:00-17:00) likely due to the
- reduced scavenging effects which were associated with reduced PM2.5 concentration. As a comparison, the highest level of downward solar radiation or the estimated OH radical was in midday (12:00-14:00), corresponding to the highest ozone production rate simulated in the box model (Fig. S18).
- Acetone was the second most abundant OVOC, with a mean mixing ratio of 3.3 ppb. However, due to the low reactivity of
   acetone, its OFP was less significant (3.2 μg m<sup>-3</sup>). Acetone was also found to be one of the most abundant OVOC in a summer
   campaign conducted in Xi'an, although the acetone mixing ratio in this study was at the lower end of the concentration ranges
   (2.1-6.6 ppb) sampled in summer 2019 (Song et al., 2021). Nitrogen-containing VOCs of acetonitrile (mean: 3.0 ppb) and
   formamide (1.1 ppb) were also abundant. However, these nitrogen-containing VOCs were generally considered minor
   contributors to ozone formation (Carter, 2010), and their OFPs were not evaluated here. Other important ozone precursors of
   include butanedione (1.6 ppb), acetaldehyde (1.2 ppb), and methyl furans (0.65 ppb).

Aromatics including benzene (1.2 ppb), phenol (0.85 ppb), and toluene (0.76 ppb) were also the top VOCs or OVOCs, contributing to OFP of 3.0 μg m<sup>-3</sup>, 9.9 μg m<sup>-3</sup>, 12.6 μg m<sup>-3</sup>, respectively (Table 1). Toluene (0.76 ppb) was largely reduced when compared to the values (1.9-10.5 ppb) reported previously e.g., in autumn to winter 2017 (Li et al., 2022), although the concentrations were not directly comparable given the different sampling techniques and time. The large reduction was partly due to the reduction in emission intensities. It is important to note that, without the high mass-resolution Vocus-PTR, toluene

due to the reduction in emission intensities. It is important to note that, without the high mass-resolution Vocus-PTR, toluene can be largely overestimated due to the dominant ion  $(C_3H_8O_3H^+)$  at the same nominal mass to charge (m/z) ratio as toluene 删除了:, suggesting secondary formation were the major source of formaldehyde...

(删除了: which

删除了:9

i.e., at *m/z* 93 (Fig. <u>\$19</u>). In this study, aromatics were estimated to contribute to 17% of total OFP, slightly higher than its mass fraction (16%) in VOCs.

· 删除了: S10

## 3.3 An increased role of secondary particulate formation in winter haze

- 95 To understand the changes in PM<sub>2.5</sub> and organic aerosol (OA) sources in the winter season, the chemical composition of non-refractory particulate matter (NR-PM) and the corresponding OA factors were compiled (Fig. 4 and Table <u>\$3</u>) based on the available studies conducted at the same sampling site in urban Xi'an using an on-line AMS or ACSM (<u>Table \$4</u>). Compared to the winter <u>mean</u> NR-PM concentration (261.2 µg m<sup>-3</sup>) in 2012<sub>4</sub> 2014, a reduction of over 60% in the NR-PM concentration (100.6 µg m<sup>-3</sup>) in 2018<sub>4</sub> 2021 was observed (Fig. <u>4</u>). Such a large reduction was mainly due to the reduction in primary particulate emissions including primary OA and chloride, as well as the reduction in the precursor gases of e.g., SO<sub>2</sub> that can form secondary species of sulfate, starting from the implementation of CO, a surrogate of primary emission due to the incomplete combustion of fossil and biomass burning, showed a similar level of (60%) reduction from 2015 to 2021 (Fig. S2Q). For the same period, an approximately 60% reduction in city-wide SO<sub>2</sub> was also seen where SO<sub>2</sub> is mainly associated with fossil fuel combustion (Fig. <u>\$20</u>). Despite the large reduction in NR-PM concentrations, exceedances of NAQQS standards
- were still frequently observed (see Sect. 3.1). In terms of chemical composition, OA fraction in NR-PM remained similar, accounting for approximately half of the NR-PM, for both periods (Fig. 4a), although its absolute concentration decreased by 62% from 117.1 µg m<sup>-3</sup> in <u>2012</u>-2014 to 44.6 µg
- m<sup>-3</sup> in <u>2018</u>-2021. The reduction in OA concentration was mainly due to the large reduction in primary OA while oxygenated
   OA remains at a similar level (Fig. 4b). Specifically, the primary fossil fuel OA factor was reduced by 77% (28.9 μg m<sup>-3</sup>), while biomass burning OA was reduced by 72% (24.2 μg m<sup>-3</sup>). The reduction in fossil fuel and biomass burning was consistent with the reduction in CO and SO<sub>2</sub> as discussed above. Similarly, the primary cooking OA factor was also largely reduced by 84% (16.4 μg m<sup>-3</sup>). The reduction in cooking emissions could be partly due to upgraded kitchen facilities (Liu et al., 2022), although we note that the sampling time and duration of the compiled studies were not the same (Table S3), causing
- 15 uncertainties in the evaluations of changes in OA factors years. Future studies with long-term continuous measurements e.g., using ACSM (Chen et al., 2022), will improve the understanding of trends in OA factors in this region. Nevertheless, compared to POA factors, OOA was marginally reduced by 11% (3.0 μg m<sup>-3</sup>). Due to the larger reduction in primary OA factors, the OOA fraction accounted for over half (52%) of the total OA in 2018-2021, twice higher in terms of fractional contribution compared to the value (22% of OA) in 2012-2014. Biomass burning OA (21% of OA) and fossil fuel OA (20% of OA) had
- 20 similar contributions to the total OA (Fig. 4b) in 2018-2021. The increased fractional contribution of OOA in recent years was likely due to the emission of its precursor VOCs that were not decreasing in the Fenwei Plain and the increased oxidizing capacity as indicated by the increased ozone concentration. The increase in secondary OA was consistently observed in other areas in China (Li et al., 2022); Nie et al., 2022), which may change of volatility and viscosity of OA (Feng et al., 2023). Among the inorganic components of NR-PM, chloride showed the largest reduction (84% or 22.9 µg m<sup>-3</sup>) in the 2018-2021
- winter when compared to that in 2012-2014 (Fig. 4a). Chloride was mainly associated with the primary emission from coal and biomass burning. The large reduction in chloride was consistent with the reduction in the primary fossil fuel and biomass burning OA factors (Fig. 4b). Sulfate showed a reduction of 61% (32.3 μg m<sup>-3</sup>), while nitrate showed a reduction of 50% (17.3 μg m<sup>-3</sup>). Ammonium showed a reduction of 51% (10.8 μg m<sup>-3</sup>). The reduction in sulfate was consistent with the reduction in its precursor SO<sub>2</sub> (Fig. S2Q). Because nitrate reduction was less significant, nitrate increased its fractional contribution (17%)
- of NR-PM<sub>2.5</sub>) in <u>2018-2021</u> from 13% of NR-PM<sub>2.5</sub> in <u>2012-2014</u>. Compared to SO<sub>2</sub>, the reduction in NO<sub>2</sub> was only observed in recent years from 2018 to 2021, while NO<sub>2</sub> showed a slight increase from 2015 to 2018 (Fig. <u>\$20</u>). Therefore, nitrate is likely to increase its fractional contribution to the total PM<sub>2.5</sub> mass because NO<sub>2</sub> was not decreasing as significantly as SO<sub>2</sub>. Also, the increased oxidizing capacity leads to enhanced nitrogen oxidation ratio and thus enhanced nitrate yield.

(	删除了: 2.5
~(	删除了: S1
(	删除了: See method section
(	删除了: 2.5
$\geq$	删除了:3
X	删除了: 2.5
$\langle \rangle$	删除了:9
) (	删除了:3
(	删除了:11

(	删除了: S1
~(	删除了:1
7	删除了: 2.5
^(	删除了: as discussed in
(	删除了: 2.5
X	删除了: 2.5
X	删除了: 2013
Y(	删除了: 2019
(	删除了:T

	刪除了:
	miller J. La contract
	mijror J . in contrast
(	删除了:2019
(	删除了: 2013
(	删除了: 2019

删除了: 2.5	
删除了: 2019	
删除了:2013	
删除了: 3b	

删除了:11	
删除了:2019	
删除了: 2013	
删除了: S1	
删除了:1	

## 3.4 Impact of control policies on the ozone and secondary aerosol formation: observation-based modelling

As a response to the coronavirus pandemic (i.e., COVID-19), the lockdown measures, which were implemented in the Fenwei Plain from 24 January to 28 February 2020, provided a unique opportunity to investigate the impact of strict emission control on the ozone and SOA formation potential in comparison to that before (2019) and after the pandemic (2021). Figure 5 shows

- the vertical column density of NO<sub>2</sub>, a surrogate of anthropogenic emission, and formaldehyde, a surrogate of reactive OVOCs, over the same period (24 January 28 February) for three consecutive years from 2019 to 2021. Compared to NO<sub>2</sub> (18-25 ×10<sup>15</sup> molecules cm<sup>-2</sup>) before the pandemic in 2019, the mean vertical column density of NO<sub>2</sub> during the lockdown (6-10 ×10<sup>15</sup> molecules cm<sup>-2</sup>) in Xi an 2020 reduced by up to over 70%. However, without the lockdown measures in 2021, the NO<sub>2</sub> column density bounced back quickly, but with a slightly lower density in the city center of Xi'an (16-20 ×10<sup>15</sup> molecules cm<sup>-2</sup>). In
- particular, the NO<sub>2</sub> in other cities in the Fenwei Plain was only 70-80% (or  $14-20 \times 10^{15}$  molecules cm<sup>-2</sup>) of that in 2019 (Fig. 5).

The large reduction in NO<sub>2</sub> during the lockdown period in 2020 implies a sharp reduction of anthropogenic emissions. However, different from NO<sub>2</sub>, the vertical column concentration for formaldehyde remained at the same level for the lockdown year in 2020 when compared to before the lockdown in 2019 with densities in the range of 8-10 ×10<sup>15</sup> molecules cm<sup>-2</sup> in urban Xi'an

- 2020 when compared to before the lockdown in 2019 with densities in the range of 8-10 ×10<sup>15</sup> molecules cm<sup>-2</sup> in urban Xi'an (Fig. 5), suggesting a weaker reduction of the reactive OVOC despite the significant reduction in anthropogenic activities including traffic and some industries that were not operative during the pandemic in 2020. In 2021, the formaldehyde densities even increased a little bit when compared to 2019 (Fig. 5). Due to the large ozone formation potential from formaldehyde (as discussed in Sect. 3.3) and the reduced NO<sub>x</sub> emissions (titration effects), surface ozone concentrations increased its NAAQS exceedance frequency in recent winters as demonstrated in Fig. 1. This is also consistent with the fact that ozone formation
- pathways in the city centers of the Fenwei Plain are in the VOC-limited regime as corroborated by the low TROPOMI HCHO/NO<sub>2</sub> ratios with values below 1 (Fig. S21). The low HCHO/NO<sub>2</sub> values in the Fenwei Plain were consistent with that observed in most Chinese cities where O<sub>3</sub> was in the VOC-limited region (Li et al., 2022; Tan and Wang, 2022).
- To examine the response of in situ photochemistry to up to 70% reduction in NO<sub>2</sub> as the scenario of the strict emission control during the lockdown, simulations of the changes in the mean rate of self-reaction between peroxy radicals were performed using a box model. Figure 6 shows that the mean self-reaction rate between peroxy radicals increase by 12-75% in response to the reduction of 20-70% of NO<sub>2</sub> while the VOCs were constant. Self-reaction between peroxy radicals includes the autooxidations of RO<sub>2</sub> (RO<sub>2</sub> + RO<sub>2</sub>) and HO<sub>2</sub> (HO<sub>2</sub> + HO<sub>2</sub>) and cross reactions (RO<sub>2</sub> + HO<sub>2</sub>), some of which lead to SOA formation (Zhao et al., 2018). Given that self-reaction between peroxy radicals is an important pathway for SOA formation
- 95 potential, the increase in self-reaction rate can partly explain the increase in SOA formation potential in recent winters (Fig. 3). In addition to the self-reaction, some of the reaction between NO<sub>g</sub>/NO<sub>g</sub> and peroxy radicals can also produce organic nitrate that could partition to the particle phase (Lin et al., 2021). However, such formation pathways were likely playing a less important role when NOx were largely reduced. In a scenario of a further 20% reduction in VOCs, self-reaction rate was reduced as compared to the reduction of NO<sub>2</sub> alone. This is because reducing VOCs reduces the availability of peroxy radicals,
- .00 while the competition between peroxy radicals and NOx was also reduced at the same time. However, the rate is still 14% higher than the base case. A large reduction (50-80%) in the self-reaction rate can be only achieved in response to a further 50-70% reduction of VOCs as compared to the base case. The modeling results imply the co-benefits of controlling VOCs in reducing ozone and SOA pollution.

## 3.5 Atmospheric implication

70

.05 The increase of ozone pollution in winter, concurrent with haze pollution in the Fenwei Plain is concerning since ozone pollution is generally considered a summertime problem (Li et al., 2019; Li et al., 2022). To make things worse, the increase in ozone pollution would stimulate the formation of secondary PM<sub>2.5</sub>, including SOA, nitrate and sulfate, and, therefore, partly negating the efforts to reduce haze pollution in winter. This is demonstrated in this study as the secondary pollutants, and in

· 删除了: good surrogate

**设置了格式:** 下标 **设置了格式:** 下标 particular, SOA and nitrate, are becoming increasingly important in recent winter haze. Due to the unique topology of the Fenwei Plain (Cao and Cui, 2021), the air pollutants, both locally emitted/produced and regionally transported, are favorably trapped inside the basin on winter night when the boundary layer height is less than 100 m, accumulating air pollutants to health-threatening level. As the development of boundary layer after sunrise, ozone formation is increased in the late afternoon.

- As a result of the increased O<sub>3</sub>, secondary particle pollutants are further increased and subsequently accumulated at night. Such a feedback loop makes it more challenging to reduce ozone and PM<sub>2.5</sub> in the Fenwei Plain than in other regions e.g., North China Plain. If inappropriate emission control measures are implemented, one may expect more frequent ozone-PM<sub>2.5</sub> pollution in winter.
- The increase of ozone in winter haze is explained by the in-situ photochemical reaction caused by the reduction of NO<sub>x</sub>, while reactive OVOCs remain constantly elevated. The abrupt decrease of NO<sub>x</sub>, as observed during the lockdown in 2020, provides real-world evidence of the side effects of implementing inappropriate emission-control measures, targeting NO<sub>x</sub> but letting loose VOCs. Here, we also show that NO<sub>2</sub> levels without the lockdown in 2021, were still 10-20% reduced when compared to before the pandemic in 2019, while the reactive HCHO was even higher. Such a trend needs to be reversed in order to achieve the synergic reduction of O<sub>3</sub> and PM<sub>2.5</sub>.
- The complexity of the O<sub>3</sub>-PM<sub>2.5</sub> interaction also needs to be considered because PM<sub>2.5</sub> scavenges HO<sub>2</sub> radicals that would otherwise produce O<sub>3</sub>. Such an interaction is more important when the starting PM<sub>2.5</sub> concentration is high (>40 µg m<sup>-3</sup>) (Shao et al., 2021). Here, we show that, although the city-wide mean PM<sub>2.5</sub> concentration decreased in recent years, the PM<sub>2.5</sub> concentration is still above 80 µg m<sup>-3</sup> for both observed and de-weathered PM<sub>2.5</sub>, i.e., still in the range of PM<sub>2.5</sub> concentration where the reduction of PM<sub>2.5</sub> can increase O<sub>3</sub> due to the reduced scavenging effect. Since the reduction in PM<sub>2.5</sub> in Xi'an is slower (-3 3% yr<sup>-1</sup>) (Wang et al. 2022) than in other areas e.g. Beijing (-7 8% yr<sup>-1</sup>) (Vu et al. 2019) and Shanghai (-6 4% yr.
- 30 slower (-3.3% yr<sup>-1</sup>) (Wang et al., 2022) than in other areas, e.g., Beijing (-7.8% yr<sup>-1</sup>) (Vu et al., 2019) and Shanghai (-6.4% yr<sup>-1</sup>) (Wang et al., 2022), a large increase in O<sub>3</sub> in winter is expected as PM<sub>2.5</sub> is only reduced at a similarly slow rate in the near future. Only with a more aggressive reduction of the precursor gases can we achieve the goal of synergic reduction O<sub>3</sub> and PM<sub>2.5</sub> pollution in Fenwei Plain.

## 4. Conclusion

- In this study, an increase in O<sub>3</sub> pollution in the winter haze was revealed through the analysis of city-wide mean concentrations of air pollutants for 7 years (2015-2021). The increased O<sub>3</sub> was not due to the changes in meteorological variations, as confirmed by the trend of deweathered O<sub>3</sub>, but due to the constantly elevated reactive OVOCs, and in particular formaldehyde in combination with the reduced NO<sub>2</sub>. An increase in O<sub>3</sub> stimulated the formation of secondary aerosols, including SOA and nitrate, which were shown to increase their fractional contributions to the total PM<sub>2.5</sub>. A further extension of ozone pollution in winter haze is expected if NO<sub>x</sub> and PM<sub>2.5</sub> were reducing at the current rate, while reactive OVOC were not reducing. Through
- scenario analysis in a chemical box model, we show the co-benefits of reducing NO<sub>x</sub> and VOCs simultaneously in reducing ozone and SOA. The complex interaction between O<sub>3</sub> and PM<sub>2.5</sub>, coupled with the topology of the Fenwei Plain and the evolution of the boundary layer height, highlight the challenges in further reducing particulate pollution in winter despite years of efforts to reduce emissions. A synergic reduction in PM<sub>2.5</sub> and O<sub>3</sub> pollution can only be achieved through a more aggressive reduction of their precursor gases.

## ASSOCIATE CONTENT

**Data availability.** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplement. Also, all data used in the study are available from the corresponding author upon request. Supplementary Figures (Fig. S1-S12) and Table (Table S1) AUTHOR INFORMATION **Corresponding author** 

Ru-Jin Huang (rujin.huang@ieecas.cn)

#### .55 Author Contributions

CL and RJH designed the study. CL, JD, and HBZ conducted measurements, data analysis, and source apportionment. CL prepared the manuscript with contributions from all co-authors

Competing interests. The authors declare that they have no conflicting interests.

## ACKNOWLEDGMENT

-60 This work was supported by National Natural Science Foundation of China (NSFC) under Grant No. 42277092, 42107126, 42207137, and 41925015, the Strategic Priority Research Program of Chinese Academy of Sciences (No. XDB40000000), the Chinese Academy of Sciences (no. ZDBS-LY-DQC001), Institute of Earth Environment (E051QB2837), and the Cross Innovative Team fund from the State Key Laboratory of Loess and Quaternary Geology (No. SKLLQGTD1801).

## **References:**

Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and Prévôt, A. S. H.: SoFi, an IGOR-based interface for the -65 efficient use of the generalized multilinear engine (ME-2) for the source apportionment: ME-2 application to aerosol mass spectrometer data, Atmos. Meas. Tech., 6, 3649-3661, doi:10.5194/amt-6-3649-2013, 2013. Cao, J. J. and Cui, L.: Current Status, Characteristics and Causes of Particulate Air Pollution in the Fenwei Plain, China: A Review, Journal of Geophysical Research: Atmospheres, 126, e2020JD034472, https://doi.org/10.1029/2020JD034472, .70 2021.

Carter, W. P. L.: Development of the SAPRC-07 chemical mechanism, Atmospheric Environment, 44, 5324-5335, https://doi.org/10.1016/j.atmosenv.2010.01.026, 2010. Chen, G., Canonaco, F., Tobler, A., Aas, W., Alastuey, A., Allan, J., Atabakhsh, S., Aurela, M., Baltensperger, U.

- Bougiatioti, A., De Brito, J. F., Ceburnis, D., Chazeau, B., Chebaicheb, H., Daellenbach, K. R., Ehn, M., El Haddad, I., .75 Eleftheriadis, K., Favez, O., Flentje, H., Font, A., Fossum, K., Freney, E., Gini, M., Green, D. C., Heikkinen, L., Herrmann, H., Kalogridis, A.-C., Keernik, H., Lhotka, R., Lin, C., Lunder, C., Maasikmets, M., Manousakas, M. I., Marchand, N., Marin, C., Marmureanu, L., Mihalopoulos, N., Močnik, G., Necki, J., O'Dowd, C., Ovadnevaite, J., Peter, T., Petit, J.-E.,
- Pikridas, M., Matthew Platt, S., Pokorná, P., Poulain, L., Priestman, M., Riffault, V., Rinaldi, M., Różański, K., Schwarz, J., Sciare, J., Simon, L., Skiba, A., Slowik, J. G., Sosedova, Y., Stavroulas, I., Styszko, K., Teinemaa, E., Timonen, H., Tremper, A., Vasilescu, J., Via, M., Vodička, P., Wiedensohler, A., Zografou, O., Cruz Minguillón, M., and Prévôt, A. S. H.: -80 European Aerosol Phenomenology - 8: Harmonised Source Apportionment of Organic Aerosol using 22 Year-long
- ACSM/AMS Datasets, Environment International, 107325, https://doi.org/10.1016/j.envint.2022.107325, 2022. Dai, Q., Hou, L., Liu, B., Zhang, Y., Song, C., Shi, Z., Hopke, P. K., and Feng, Y.: Spring Festival and COVID-19 Lockdown: Disentangling PM Sources in Major Chinese Cities, Geophys. Res. Lett., 48, e2021GL093403, .85 https://doi.org/10.1029/2021GL093403, 2021.
- Duan, J., Huang, R.-J., Chang, Y., Zhong, H., Gu, Y., Lin, C., Hoffmann, T., and O'Dowd, C.: Measurement report of the change of PM2.5 composition during the COVID-19 lockdown in urban Xi'an: enhanced secondary formation and oxidation, Science of The Total Environment, 148126, https://doi.org/10.1016/j.scitotenv.2021.148126, 2021. Duan, J., Huang, R. J., Gu, Y., Lin, C., Zhong, H., Xu, W., Liu, Q., You, Y., Ovadnevaite, J., Ceburnis, D., Hoffmann, T.,
- .90 and O'Dowd, C.: Measurement report: Large contribution of biomass burning and aqueous-phase processes to the wintertime secondary organic aerosol formation in Xi'an, Northwest China, Atmos. Chem. Phys., 22, 10139-10153, 10.5194/acp-22-10139-2022, 2022.
- Elser, M., Huang, R. J., Wolf, R., Slowik, J. G., Wang, Q., Canonaco, F., Li, G., Bozzetti, C., Daellenbach, K. R., Huang, Y., Zhang, R., Li, Z., Cao, J., Baltensperger, U., El-Haddad, I., and André, P.: New insights into PM2.5 chemical composition .95 and sources in two major cities in China during extreme haze events using aerosol mass spectrometry, Atmospheric Chemistry and Physics, 16, 3207-3225, 10.5194/acp-16-3207-2016, 2016.
- Freng, T., Wang, Y., Hu, W., Zhu, M., Song, W., Chen, W., Sang, Y., Fang, Z., Deng, W., Fang, H., Yu, X., Wu, C., Yuan, B., Huang, S., Shao, M., Huang, X., He, L., Lee, Y. R., Huey, L. G., Canonaco, F., Prevot, A. S. H., and Wang, X.: Impact of aging on the sources, volatility, and viscosity of organic aerosols in Chinese outflows, Atmos. Chem. Phys., 23, 611-636, 00
- 10.5194/acp-23-611-2023, 2023.

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone, Remote Sensing of Environment, 202, 18-27, <u>https://doi.org/10.1016/j.rse.2017.06.031</u>, 2017.

Grange, S. K. and Carslaw, D. C.: Using meteorological normalisation to detect interventions in air quality time series,
 Science of The Total Environment, 653, 578-588, <u>https://doi.org/10.1016/j.scitotenv.2018.10.344</u>, 2019.
 Grange, S. K., Carslaw, D. C., Lewis, A. C., Boleti, E., and Hueglin, C.: Random forest meteorological normalisation

- Grange, S. K., Carslaw, D. C., Lewis, A. C., Boleti, E., and Hueglin, C.: Random forest meteorological normalisation models for Swiss PM10 trend analysis, Atmos. Chem. Phys., 18, 6223-6239, 10.5194/acp-18-6223-2018, 2018. He, G., Pan, Y., and Tanaka, T.: The short-term impacts of COVID-19 lockdown on urban air pollution in China, Nat. Sustain, 3, 1005-1011, 10.1038/s41893-020-0581-y, 2020.
- Jonkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, Atmos. Chem. Phys., 3, 181-193, 10.5194/acp-3-181-2003, 2003.

Jenkin, M. E., Valorso, R., Aumont, B., and Rickard, A. R.: Estimation of rate coefficients and branching ratios for reactions of organic peroxy radicals for use in automated mechanism construction, Atmos. Chem. Phys., 19, 7691-7717, 10.5194/acp-19-7691-2019, 2019.

- Krechmer, J., Lopez-Hilfiker, F., Koss, A., Hutterli, M., Stoermer, C., Deming, B., Kimmel, J., Warneke, C., Holzinger, R., Jayne, J., Worsnop, D., Fuhrer, K., Gonin, M., and de Gouw, J.: Evaluation of a New Reagent-Ion Source and Focusing Ion– Molecule Reactor for Use in Proton-Transfer-Reaction Mass Spectrometry, Analytical Chemistry, 90, 12011-12018, 10.1021/acs.analchem.8b02641, 2018.
- 20 Li, C., Zhu, Q., Jin, X., and Cohen, R. C.: Elucidating Contributions of Anthropogenic Volatile Organic Compounds and Particulate Matter to Ozone Trends over China, Environmental Science & Technology, 56, 12906-12916, 10.1021/acs.est.2c03315, 2022.

 Li, J., Deng, S., Tohti, A., Li, G., Yi, X., Lu, Z., Liu, J., and Zhang, S.: Spatial characteristics of VOCs and their ozone and secondary organic aerosol formation potentials in autumn and winter in the Guanzhong Plain, China, Environmental
 Research 211 113036 https://doi.org/10.1016/j.envres.2022.113036.2022

25 Research, 211, 113036, <u>https://doi.org/10.1016/j.envres.2022.113036</u>, 2022. Li, J., Han, Z., Wu, J., Tao, J., Li, J., Sun, Y., Liang, L., Liang, M., and Wang, Q. g.: Secondary organic aerosol formation and source contributions over east China in summertime, Environmental Pollution, 306, 119383, <u>https://doi.org/10.1016/j.envpol.2022.119383</u>, 2022.

- Li, K., Jacob, D. J., Liao, H., Zhu, J., Shah, V., Shen, L., Bates, K. H., Zhang, Q., and Zhai, S.: A two-pollutant strategy for improving ozone and particulate air quality in China, Nature Geoscience, 12, 906-910, 10.1038/s41561-019-0464-x, 2019.
  Li, K., Jacob, D. J., Liao, H., Qiu, Y., Shen, L., Zhai, S., Bates, K. H., Sulprizio, M. P., Song, S., Lu, X., Zhang, Q., Zheng, B., Zhang, Y., Zhang, J., Lee, H. C., and Kuk, S. K.: Ozone pollution in the North China Plain spreading into the late-winter haze season, Proceedings of the National Academy of Sciences, 118, e2015797118, 10.1073/pnas.2015797118, 2021.
  Lin, C., Huang, R.-J., Duan, J., Zhong, H., and Xu, W.: Primary and Secondary Organic Nitrate in Northwest China: A Case
- 35 Study, Environmental Science & Technology Letters, 8, 947-953, 10.1021/acs.estlett.1c00692, 2021. Lin, C., Huang, R.-J., Duan, J., Zhong, H., and Xu, W.: Polycyclic aromatic hydrocarbons from cooking emissions, Science of The Total Environment, 818, 151700, <u>https://doi.org/10.1016/j.scitotenv.2021.151700</u>, 2022. Lin, C., Huang, R.-J., Duan, J., Zhong, H., Xu, W., Wu, Y., and Zhang, R.: Large contribution from worship activities to the

atmospheric soot particles in northwest China, Environmental Pollution, 299, 118907,
https://doi.org/10.1016/j.envpol.2022.118907, 2022.
Liu, Q., Son, Y. J., Li, L., Wood, N., Senerat, A. M., and Pantelic, J.: Healthy home interventions: Distribution of PM2.5 emitted during cooking in residential settings, Build Environ, 207, 108448, https://doi.org/10.1016/j.buildenv.2021.108448, 2022.

Luo, H., Li, G., Chen, J., Lin, Q., Ma, S., Wang, Y., and An, T.: Spatial and temporal distribution characteristics and ozone formation potentials of volatile organic compounds from three typical functional areas in China, Environmental Research, 183, 109141, <u>https://doi.org/10.1016/j.envres.2020.109141</u>, 2020.

Lyu, X., Guo, H., Zou, Q., Li, K., Xiong, E., Zhou, B., Guo, P., Jiang, F., and Tian, X.: Evidence for Reducing Volatile Organic Compounds to Improve Air Quality from Concurrent Observations and In Situ Simulations at 10 Stations in Eastern China, Environmental Science & Technology, 10.1021/acs.est.2c04340, 2022.

- 50 Nie, W., Yan, C., Huang, D. D., Wang, Z., Liu, Y., Qiao, X., Guo, Y., Tian, L., Zheng, P., Xu, Z., Li, Y., Xu, Z., Qi, X., Sun, P., Wang, J., Zheng, F., Li, X., Yin, R., Dallenbach, K. R., Bianchi, F., Petäjä, T., Zhang, Y., Wang, M., Schervish, M., Wang, S., Qiao, L., Wang, Q., Zhou, M., Wang, H., Yu, C., Yao, D., Guo, H., Ye, P., Lee, S., Li, Y. J., Liu, Y., Chi, X., Kerminen, V.-M., Ehn, M., Donahue, N. M., Wang, T., Huang, C., Kulmala, M., Worsnop, D., Jiang, J., and Ding, A.: Secondary organic aerosol formed by condensing anthropogenic vapours over China's megacities, Nature Geoscience, 15,
- 155 255-261, 10.1038/s41561-022-00922-5, 2022. Shao, M., Wang, W., Yuan, B., Parrish, D. D., Li, X., Lu, K., Wu, L., Wang, X., Mo, Z., Yang, S., Peng, Y., Kuang, Y., Chen, W., Hu, M., Zeng, L., Su, H., Cheng, Y., Zheng, J., and Zhang, Y.: Quantifying the role of PM2.5 dropping in variations of ground-level ozone: Inter-comparison between Beijing and Los Angeles, Science of The Total Environment, 788, 147712, <u>https://doi.org/10.1016/j.scitotenv.2021.147712</u>, 2021.
- 60 Shi, Z., Song, C., Liu, B., Lu, G., Xu, J., Vu, T. V., Elliott, R. J. R., Li, W., Bloss, W. J., and Harrison, R. M.: Abrupt but smaller than expected changes in surface air quality attributable to COVID-19 lockdowns, Science Advances, 7, eabd6696, doi:10.1126/sciadv.abd6696, 2021.

Song, M., Li, X., Yang, S., Yu, X., Zhou, S., Yang, Y., Chen, S., Dong, H., Liao, K., Chen, Q., Lu, K., Zhang, N., Cao, J., Zeng, L., and Zhang, Y.: Spatiotemporal variation, sources, and secondary transformation potential of volatile organic

- 65 compounds in Xi'an, China, Atmospheric Chemistry and Physics, 21, 4939-4958, 10.5194/acp-21-4939-2021, 2021. Su, W., Liu, C., Hu, Q., Zhao, S., Sun, Y., Wang, W., Zhu, Y., Liu, J., and Kim, J.: Primary and secondary sources of ambient formaldehyde in the Yangtze River Delta based on Ozone Mapping and Profiler Suite (OMPS) observations, Atmos. Chem. Phys., 19, 6717-6736, 10.5194/acp-19-6717-2019, 2019.
- Tan, Y. and Wang, T.: What caused ozone pollution during the 2022 Shanghai lockdown? Insights from ground and satellite observations, Atmos. Chem. Phys., 22, 14455-14466, 10,5194/acp-22-14455-2022, 2022.
- Vu, T. V., Shi, Z., Cheng, J., Zhang, Q., He, K., Wang, S., and Harrison, R. M.: Assessing the impact of clean air action on air quality trends in Beijing using a machine learning technique, Atmos. Chem. Phys., 19, 11303-11314, 10.5194/acp-19-11303-2019, 2019.
- Wang, M., Duan, Y., Zhang, Z., Huo, J., Huang, Y., Fu, Q., Wang, T., Cao, J., and Lee, S.-c.: Increased contribution to
   PM2.5 from traffic-influenced road dust in Shanghai over recent years and predictable future, Environmental Pollution, 120119, https://doi.org/10.1016/j.envpol.2022.120119, 2022.
- Wang, M., Zhang, Z., Yuan, Q., Li, X., Han, S., Lam, Y., Cui, L., Huang, Y., Cao, J., and Lee, S.-c.: Slower than expected reduction in annual PM2.5 in Xi'an revealed by machine learning-based meteorological normalization, Science of The Total Environment, 156740, https://doi.org/10.1016/j.scitotenv.2022.156740, 2022.
- Wang, R., Wang, X., Cheng, S., Zhu, J., Zhang, X., Cheng, L., and Wang, K.: Determining an optimal control strategy for anthropogenic VOC emissions in China based on source emissions and reactivity, Journal of Environmental Sciences, https://doi.org/10.1016/j.jes.2022.10.034, 2022.

 Wang, T., Dai, J., Lam, K. S., Nan Poon, C., and Brasseur, G. P.: Twenty-Five Years of Lower Tropospheric Ozone Observations in Tropical East Asia: The Influence of Emissions and Weather Patterns, Geophysical Research Letters, 46, 11463-11470, https://doi.org/10.1029/2019GL084459, 2019.

Wang, T., Xue, L., Feng, Z., Dai, J., Zhang, Y., and Tan, Y.: Ground-level ozone pollution in China: a synthesis of recent findings on influencing factors and impacts, Environmental Research Letters, 2022.
Wang, T., Wei, X. L., Ding, A. J., Poon, C. N., Lam, K. S., Li, Y. S., Chan, L. Y., and Anson, M.: Increasing surface ozone

wang, T., wei, A. E., Ding, A. J., Pool, C. N., Lani, K. S., Li, T. S., Chan, L. T., and Anson, M.: increasing surface ozone concentrations in the background atmosphere of Southern China, 1994–2007, Atmos. Chem. Phys., 9, 6217-6227, 10.5194/acp-9-6217-2009, 2009.

Wolfe, G. M., Marvin, M. R., Roberts, S. J., Travis, K. R., and Liao, J.: The Framework for 0-D Atmospheric Modeling (F0AM) v3.1, Geosci. Model Dev., 9, 3309-3319, 10.5194/gmd-9-3309-2016, 2016.

- Yan, Y., Wang, X., Qu, K., Li, X., Shi, W., Peng, Z., and Zhang, Y.: Impacts of synoptic circulations on summertime ozone pollution in Guanzhong Basin, northwestern China, Atmospheric Environment, 262, 118660, https://doi.org/10.1016/j.atmosenv.2021.118660, 2021.
- <sup>595</sup> https://doi.org/10.1016/j.atmosenv.2021.18600, 2021.
   Zhang, Q., Shao, M., Li, Y., Lu, S. H., Yuan, B., and Chen, W. T.: Increase of ambient formaldehyde in Beijing and its implication for VOC reactivity, Chinese Chemical Letters, 23, 1059-1062, <u>https://doi.org/10.1016/j.cclet.2012.06.015</u>, 2012.
   Zhao, Y., Thornton, J. A., and Pye, H. O. T.: Quantitative constraints on autoxidation and dimer formation from direct probing of monoterpene-derived peroxy radical chemistry, Proceedings of the National Academy of Sciences, 115, 12142-12147, doi:10.1073/pnas.1812147115, 2018.

Zhao, Y., Zhang, K., Xu, X., Shen, H., Zhu, X., Zhang, Y., Hu, Y., and Shen, G.: Substantial changes in nitrate oxide and ozone after excluding meteorological impacts during the COVID-19 outbreak in mainland China, Environmental Science & Technology Letters, XXXX, 10.1021/acs.estlett.0c00304, 2020.

Zhong, H., Huang, R.-J., Chang, Y., Duan, J., Lin, C., and Chen, Y.: Enhanced formation of secondary organic aerosol from photochemical oxidation during the COVID-19 lockdown in a background site in Northwest China, Science of The Total Environment, 778, 144947, <u>https://doi.org/10.1016/j.scitotenv.2021.144947</u>, 2021.
 Zhong, H., Huang, R.-J., Duan, J., Lin, C., Gu, Y., Wang, Y., Li, Y., Zheng, Y., Chen, Q., Chen, Y., Dai, W., Ni, H., Chang,

Zhong, H., Huang, R.-J., Duan, J., Lin, C., Gu, Y., Wang, Y., Li, Y., Zheng, Y., Chen, Q., Chen, Y., Dai, W., Ni, H., Chang, Y., Worsnop, D. R., Xu, W., Ovadnevaite, J., Ceburnis, D., and O'Dowd, C. D.: Seasonal variations in the sources of organic aerosol in Xi'an, Northwest China: The importance of biomass burning and secondary formation, Science of The Total
 Environment, 139666, https://doi.org/10.1016/j.scitotenv.2020.139666, 2020.

	Formula	Exact Mass	Assignment	Mean (ppb)	OFP (µg m <sup>-3</sup> )
1	НСНОН	31.017841	Formaldehyde <sup>a</sup>	4.16	54.4
2	C3H6OH	59.049141	acetone	3.39	3.2
3	C2H3NH	42.033825	acetonitrile	3.03	-
4	C4H6O2H	87.044055	butanedione	1.63	27.7
5	C2H4OH	45.03349	acetaldehyde	1.20	15.8
6	C6H6H	79.054226	benzene	1.19	3.0
7	CH3NOH	46.028740	formamide	1.10	-
8	C6H6OH	95.049141	phenol	0.85	9.9
9	C7H8H	93.069876	toluene	0.76	12.6
10	C5H6OH	83.04914127	methylfurans	0.65	20.0
Total			TVOCs	18.0	146.8

## 615 Table 1. Top 10 volatile organic compounds (VOCs)/oxygenated VOCs (OVOCs) and the ozone formation potential (OFP).

<sup>a</sup>Formaldehyde concentration was scaled using the vertical column concentration.





Figure 1. Observed exceedance frequency (in days month<sup>-1</sup>) of  $PM_{2.5}(a)$ ; Ozone (b); Ozone or  $PM_{2.5}(c)$  standard (NAAQS level-1) in the biggest city (i.e., Xi'an) in Fenwei Plain, averaged over 2015-2018 and 2019-2021. Ozone and  $PM_{2.5}$  were averaged from 13 monitoring sites in Xi'an. Error bar represents one standard deviation.

620

· 删除了: Ozone or





Figure 2. Observed and deweathered diurnal patterns for PM<sub>2.5</sub> (a,c) and O<sub>3</sub> (b, d) for January-February in 2015-2018 and 2019-2021

630



Figure 3. (a) Observed and simulated formaldehyde (HCHO) and (b) the EAR5 UV radiation flux and the simulated OH radical in the box model.





Figure 5. Tropospheric NO<sub>2</sub> column densities, a surrogate of anthropogenic emissions, and tropospheric HCHO column densities, a surrogate of OVOCs, measured by the TROPOMI satellite instrument in January-February in 2019, 2020, and 2021.



