

Thermal regime of the Grigoriev ice cap and the Sary-Tor glacier in the Inner Tien Shan, Kyrgyzstan

Lander VAN TRICHT¹, Philippe HUYBRECHTS¹

5 ¹Earth System Science and Departement Geografie, Vrije Universiteit Brussel, Brussels, Belgium

Correspondence to: Lander Van Tricht (lander.van.tricht@vub.be)

Abstract. The thermal regime of glaciers and ice caps represents the internal distribution of ice temperatures. An accurate knowledge of the thermal regime of glaciers and ice caps is important to understand their dynamics and response to climate change, and to model their evolution. Although the assumption is that most ice masses in the Tien Shan are polythermal, this has not been examined in appropriate detail so far. In this research, we investigate the thermal regime of the Grigoriev ice cap and the Sary-Tor glacier, both located in the Inner Tien Shan in Kyrgyzstan, using a 3D higher-order thermomechanical ice flow model. Input data and boundary conditions are inferred from a surface energy mass balance model, a historical air temperature and precipitation series, ice thickness measurements and reconstructions, and digital elevation models. Calibration and validation of the englacial temperatures is performed using historical borehole measurements on the Grigoriev ice cap and radar measurements for the Sary-Tor glacier. The results of this study reveal a polythermal structure of the Sary-Tor glacier and a cold structure of the Grigoriev ice cap. The difference is related to the larger amount of snow (insulation) and refreezing meltwater (release of latent heat) for the Sary-Tor glacier resulting in higher surface layer temperature, especially in the accumulation area, which are subsequently advected downstream. Further, ice velocities are much lower for the Grigoriev ice cap with consequent lower horizontal advection rates. A detailed analysis concerning the influence of temperature and precipitation changes at the surface reveals that the thermal structure of both ice bodies is not constant over time, with recent climate change causing increasing ice temperatures in higher areas. The selected ice masses are representative examples of the (Inner) Tien Shan glaciers and ice caps. Therefore, our findings and the calibrated parameters can be generalised allowing to improve the understanding of the dynamics and future evolution of other glaciers and ice caps in the region.

25 1 Introduction

Global warming is causing an unprecedented loss of glaciers that is expected to continue and have far-reaching consequences for worldwide water availability, even without further temperature increase (Zemp et al., 2019). High Mountain Asia (HMA), comprising several glaciated mountain ranges such as the Himalayas, Karakoram, Pamir and Tien Shan, contains the highest concentration of glaciers globally (Bhattacharya et al., 2021). However, glaciers in HMA also involve one of the largest uncertainties in current and future ice volume assessments (Farinotti et al., 2019). An important characteristic influencing the dynamics and behaviour of glacier ice are the englacial temperatures (Gilbert et al., 2020). The thermal regime of an ice mass,

heeft verwijderd: of ice masses

heeft verwijderd: superimposed ice

heeft verwijderd: ice

heeft verwijderd: s

heeft verwijderd: a

heeft verwijderd: more temperate ice

heeft verplaatst (invoeging) [1]

heeft verwijderd: is

heeft verwijderd: Englacial temperatures represent the internal distribution of the ice temperatures of ice bodies.

heeft opmaak toegepast: Lettertype: 10 pt, Tekstkleur: Auto, Engels (Groot-Brittannië)

describing the englacial temperature structure, is universally categorised into three types: cold (containing cold ice, possibly with a thin layer of temperate ice at the base), temperate (containing only temperate ice, i.e. the ice temperature reaches the pressure melting point everywhere), and polythermal (containing cold and temperate ice) (Blatter and Hutter, 1991). Accurate knowledge of the ice temperatures is important because ice deformation depends on the ice temperature, meltwater routing is affected by ice temperature and the basal temperature determines whether the ice mass can slide over its base (Blatter and Hutter, 1991; Hambrey and Glasser, 2012; Gilbert et al., 2020). Hence, precise knowledge of the thermal regime of ice bodies is needed to understand and accurately model their response to climate change (Gilbert et al., 2020).

The focus of this study is on the Tien Shan, where ice masses are assumed to be mostly cold and polythermal (Vilesov, 1961; 50 Maohuan et al., 1982; Echelmeyer and Zhongxiang, 1987; Vasilenko et al., 1988; Petrakov et al., 2014; Nosenko et al., 2016). Previous studies demonstrated a polythermal structure using borehole measurements for the Ürümqi glacier (Cai et al., 1988) and the Davydov glacier (Vasilenko et al., 1988). Using Ground Penetrating Radar (GPR) measurements, other studies revealed the presence of cold and temperate layers on the Sary-Tor glacier (Petrakov et al., 2014), the Tuyuksu glacier (Nosenko et al., 2016), and the Abramov glacier, located in the nearby Pamir Alay (Kislov et al., 1977). According to measurements on the 55 Abramov glacier, the Ürümqi glacier and the Tuyuksu glacier, temperate ice was present near the surface of the lower part of the accumulation area (Kronenberg et al., 2020), and over a large area at the base (Nosenko et al., 2016). The latter was corroborated by larger summer velocities compared to winter velocities indicating that basal sliding was occurring on the Ürümqi glacier in the Chinese Tien Shan (Maohuan et al., 1989, 1992). Part of the Inner Tien Shan glaciers have also been categorized as surge-type glaciers (Mukherjee et al., 2018; Bhattacharya et al., 2021). The surge behavior has been interpreted 60 by the polythermal structure of the ice masses with a cold mantle at the front restraining ice movement. After a long period with ice thickening and steepening in the upper reaches, the cold mantle at the front is supposed to be destructed, followed by an advance over a short period of time (Nosenko et al., 2016; Mukherjee et al., 2017). Few (11) of such glaciers were identified in the Ak-Shyirak massive, including the Northern Bordu glacier and the Davydov glacier, located in the valleys next to the Sary-Tor glacier (Bondarev, 1961; Mukherjee et al., 2017).

65 The temperate ice in the lowest part of the accumulation area has typically been explained by an infiltration congelation/recrystallisation zone, characteristic for the continental climate in which the ice masses are located (Shumskiy, 1955; Vilesov, 1961; Maohuan et al., 1982, 1990; Osmonov et al., 2013; Petrakov et al., 2014). In the lowest part of the 70 accumulation area, the ice and snow surface are warmed significantly by the infiltration and refreezing of meltwater releasing latent heat (Shumskiy, 1955; Lüthi et al., 2015). Such a temperate area in the infiltration zone has also been observed at other sub-polar glaciers such as Storglaciären in Sweden (Hooke et al., 1983). But, in addition to the influence of refreezing meltwater, other processes play a role in the warming and cooling of surface layers, such as a seasonal snow cover that shields the underlying ice from the extremely cold air temperatures during winter (Hooke et al., 1983) or the ablation of ice which removes the warm or cold ice from above (Blatter et al., 1987; Wohlleben et al., 2009).

heeft opmaak toegepast: Lettertype: 10 pt, Tekstkleur: Auto, Engels (Groot-Brittannië)

heeft verwijderd: over

heeft opmaak toegepast: Lettertype: 10 pt, Tekstkleur: Auto, Engels (Groot-Brittannië)

heeft opmaak toegepast: Lettertype: 10 pt, Engels (Groot-Brittannië)

heeft verwijderd: Englacial temperatures can be measured directly using boreholes. However, the distinction between cold and temperate ice can also be recognised on radargrams (Delf et al., 2022).¹

heeft verwijderd: Accurate

heeft opmaak toegepast: Lettertype: 10 pt, Engels (Groot-Brittannië)

heeft opmaak toegepast: Lettertype: (Standaard +Hoofdtekst) (Times New Roman)

heeft verwijderd: therefore

heeft verwijderd: of ice bodies

heeft opmaak toegepast: Lettertype: (Standaard +Hoofdtekst) (Times New Roman)

heeft opmaak toegepast: Lettertype: (Standaard +Hoofdtekst) (Times New Roman)

heeft opmaak toegepast: Lettertype: (Standaard +Hoofdtekst) (Times New Roman)

heeft verwijderd: ¹

heeft verwijderd: The latter means that the ice mass consists of both warm/temperate ice, i.e. the ice temperature reaches the pressure melting point, and cold ice.

heeft verwijderd: warm

heeft verwijderd: on the Abramov glacier (Kronenberg et al., 2020), ...

heeft verwijderd: and

heeft verwijderd: 0

heeft verwijderd: an

heeft verwijderd: reconnection

heeft verwijderd: Vilesov, 1961;

heeft verwijderd: ; Kronenberg et al., 2020

heeft verwijderd: In the infiltration zone, which is the

heeft verwijderd: is

heeft verwijderd: thawed

heeft verwijderd: which can heat up the layers to zero-degree

heeft verwijderd: warm

heeft verwijderd: ¹

Here, we investigate the thermal regime of the Grigoriev ice cap and the Sary-Tor glacier, both located in the Inner Tien Shan, Kyrgyzstan (Central-Asia), using a 3-dimensional (3D) higher-order (HO) thermomechanical ice flow model coupled to a simplified 2D surface energy mass balance model. The selected ice flow model has been successfully applied at different scales, ranging from a medium sized mountain glacier in the Alps (Zekollari et al., 2013; Zekollari and Huybrechts, 2015), a medium-sized ice cap in Greenland (Zekollari et al., 2017) to the entire Greenland ice sheet (Fürst et al., 2013, 2015). To date, most of the 3D glacier and ice cap model studies have been performed assuming an isothermal state of the ice mass (e.g. Jouvet et al., 2011; Zekollari et al., 2014). Only a few studies have been performed applying a 3D thermomechanical ice flow model including variations in englacial temperatures for glaciers (Zwinger et al., 2007; Zwinger and Moore, 2009; Zhao et al., 2014; Rowan et al., 2015; Li et al., 2017; Gilbert et al., 2017, 2020), and ice caps (Flowers et al., 2007; Schäfer et al., 2015; Zekollari et al., 2017).

heeft verwijderd: 1

heeft verplaatst (invoeging) [2]

heeft verwijderd: small

heeft verwijderd: s

heeft verwijderd:

heeft verwijderd: s

In this study, we perform different experiments under constant 1960-1990 average climatic conditions. Afterwards, multiple sensitivity experiments are carried out to determine the influence of parameter choice, to assess the uncertainty of the results, and to determine the impact of changing climatic conditions at the surface on the thermal structure of both ice bodies. Several datasets are used for calibration and validation. For the Grigoriev ice cap, ice thickness measurements were performed in August 2021 with a GPR system and used for a glacier wide ice thickness reconstruction. Ice temperatures derived from ice cores made in 1962 (Dikikh, 1962), 1990 (Thompson et al., 1993), 2001 (Arkhipov et al., 2004), and 2007 (Takeuchi et al., 2014), distributed over the ice cap, are used as calibration and validation for the thermal structure. Historical surface mass balance measurements from different periods are used to calibrate the mass balance model (Dyurgerov, 2002; Arkhipov et al., 2004; Fujita et al., 2011). Concerning the Sary-Tor glacier, observed and reconstructed ice thicknesses from 2013 are adopted (Petrakov et al., 2014), recent mass balance measurements are used, and radargrams displaying cold and temperate ice are considered for the validation of the thermal structure.

heeft verplaatst (invoeging) [5]

heeft verwijderd: 1

heeft verwijderd: ground penetrating radar (

heeft verwijderd:)

125 2 Study area

2.1 The Grigoriev ice cap and the Sary-Tor glacier

The Grigoriev ice cap ($\sim 7 \text{ km}^2$) (also called flat-top glacier) is located at the southern slopes of the Terskey Ala-Too mountain range in the Inner Tien Shan in Kyrgyzstan (Central Asia) (Fig. 1). The ice cap is located between the valleys of the Chontor glacier (to the west) and the Popova glacier (to the east). The top of the Grigoriev ice cap is a flat snow field at an elevation between 4500 and 4600 m a.s.l. In contrast to a typical valley glacier, the ice of the Grigoriev ice cap flows in multiple directions, ending at steeper glacier fronts and at multiple smaller outlet glaciers (Fig. 1). The northern boundary of the Grigoriev ice cap is also characterised by ice cliffs where mass is lost due to dry calving. The Grigoriev ice cap has been subject to numerous glaciological investigations such as mass balance measurements (Mikhailenko, 1989; Fujita et al., 2011)

heeft omhoog verplaatst [5]: under constant 1960-1990 average climatic conditions. Afterwards, multiple sensitivity experiments are carried out to determine the influence of parameter choice, to assess the uncertainty of the results, and to determine the impact of changing climatic conditions at the surface on the thermal structure of both ice bodies.

heeft verwijderd: used

heeft verwijderd: Different experiments are performed

heeft verwijderd: 1

heeft verwijderd: and data

heeft verwijderd: is a small ice cap



and ice core drilling (Dikikh, 1965; Thompson et al., 1993; Arkhipov et al., 2004; Takeuchi et al., 2014; Takeuchi et al., 2019).

In August 2021, we measured the ice thickness of the Grigoriev ice cap with a GPR system, following the approach of Van

155 Tricht et al. (2021a).

The Sary-Tor glacier is a small valley glacier ($\sim 2 \text{ km}^2$), located in the north-western part of the Ak-Shyirak massif, roughly 30 km to the southeast of the Grigoriev ice cap (Fig. 1). Its ice thickness was measured in 2013 with a GPR system showing a maximum ice thickness of almost 160 m (Petrakov et al., 2014). Annual mass balance measurements on this glacier are 160 performed since 2014 and these are deposited in the World Glacier Monitoring Surface (WGMS) database. Because debris cover is almost entirely absent on the Sary-Tor glacier, the glacier changes are assumed to directly reflect the changing climate in the Inner Tien Shan (Petrakov et al., 2014). The Sary-Tor glacier has therefore typically been used as a reference glacier for the surrounding area, especially for the Ak-Shyirak massive (Mikhailenko, 1993). Van Tricht et al. (2021b) calibrated a surface energy mass balance model for the Sary-Tor glacier and reconstructed the historical mass balance between 1750 and 2020, 165 showing a significant decrease in the mean specific mass balance, especially since the seventies. Recently, the concession of the Kumtor gold mine, which is located north of the glacier (Fig. 1), was extended towards the valley of the Sary-Tor glacier, removing the terminal moraines of the Little Ice Age (LIA).

heeft verwijderd: on the other hand



170 heeft verwijderd:

175 heeft verwijderd: a light blue

Figure 1: The Grigoriev ice cap and the Sary-Tor glacier located in their respective mountain range in the Inner Tien Shan. All the glaciers of the Randolph Glacier Inventory version 6 (from 2002) are indicated in [white colour on the upper map](#). The black outline of the glaciers [in the lower maps](#) represents the extent of the ice mass in August 2021. Glacier elevation contours are added for every 50 metres. The location of the Kumtor-Tien Shan meteorological station is added with a red dot. The background topography is from TanDEM-X, the satellite images are from Sentinel-2 on the 26th of July 2021.

2.2 Historical englacial temperature measurements

- 180 Over the past 60 years, various temperature measurements have been carried out in the study area (Dikikh, 1965; Vasilenko et al., 1988; Thompson et al., 1993; Arkhipov et al., 2004; Takeuchi et al., 2014). In 1962, a first drilling campaign was carried out on the Grigoriev ice cap between 4170 and 4420 m a.s.l. (Dikikh, 1965). The measurements revealed the presence of cold ice. In July 1985, the USSR Academy of Sciences performed a thermal drilling in the central part of the Davydov glacier, located in the valley next to the Sary-Tor glacier, at about 3950 m a.s.l. down to a depth of 102 m (Vasilenko et al., 1988).
- 185 Temperatures were measured down to a depth of 30 m. The results showed that ice temperatures decreased sharply to almost -6°C at a depth of 6 m. Further down, ice temperatures increased quickly, reaching -0.2°C at a depth of 30 m. As such, the results strongly suggested a two-layer structure with a temperate ice core over the bed.

In 1990, a drilling of a shallow ice core (16.5 and 20 m) was carried out at the summit of the Grigoriev ice cap, covering the last 50 years. (Thompson et al., 1993). Thompson et al. (1993) measured an ice temperature of -2°C at a depth of 20 m, substantially higher than the 1962 measured temperatures (Dikikh, 1965). One of the conclusions of the study of Thompson et al (1993) was that the 10 to 20 m temperatures generally exceeded the mean annual air temperature because of the refreezing of meltwater, and that ice temperatures increased strongly between 1960 and 1990. Thompson et al (1993) also observed a significant amount of melting and runoff during the 1990 summer season, as well as a few large pools of standing water on flat areas of the ice cap below 4500 m elevation. Such supraglacial water suggested cold, impermeable ice.

In September 2001, Arkhipov et al. (2004) made a drilling of 21.5 meter on the summit of the Grigoriev ice cap. During their survey, they noticed the presence of a thick water saturated firm layer by the end of the ablation period leading to a significant warming of the upper layers. Besides, Arkhipov et al. (2004) measured an abrupt ice temperature increase of several degrees over the transition from the ablation region to the accumulation region (at the EL, around 4300 m), which they explained by the heat released from refreezing meltwater.

Finally, in September 2007, a deep drilling was performed near the summit of the Grigoriev ice cap (4563 m a.s.l.) down to the bedrock (87.46 m) (Takeuchi et al., 2014). Temperature measurements of the drilled ice revealed that the ice was -2.7°C at 10 m depth and down to -3.9 °C at the bottom. Hence, the basal ice was not at the pressure melting point (Takeuchi et al., 2014). The lower temperatures at the bottom of the ice suggested a recent warming of the top layers that had not yet reached the bottom.

On the Sary-Tor glacier, no direct ice temperature measurements were made. However, the radar measurements of Petrakov et al. (2014) clearly showed a distinction between temperate ice over the bed and cold ice above at a surface elevation of 4150 m.▼

heeft verwijderd: E

heeft omhoog verplaatst [1]: Englacial temperatures represent the internal distribution of the ice temperatures of ice bodies. The thermal regime is universally categorised into three types: cold (containing cold ice, possibly with a thin layer of temperate ice over the base), temperate (containing only temperate ice), and polythermal (containing cold and temperate ice) (Blatter and Hutter, 1991). Accurate knowledge of the ice temperatures is important because ice deformation depends on the ice temperature, meltwater routing is affected by ice temperature and the basal temperature determines whether the ice mass can slide over its base. Englacial temperatures can be measured directly using boreholes. However, the distinction between cold and temperate ice can also be recognised on radargrams (Delf et al., 2022).

heeft verwijderd: ↗

heeft verwijderd: up

heeft verwijderd: up

heeft verwijderd: warm

heeft verwijderd: new

heeft verwijderd:

heeft verwijderd: last

heeft verwijderd: close to the top

heeft verwijderd: up

heeft verwijderd: warm

heeft verwijderd: This demonstrated a polythermal structure of this glacier....

3 Methods, data, and calibration

240 3.1 Higher-Order glacier model

A 3-dimensional glacier model resolves the spatially distributed [x,y,z]-velocity fields in an ice mass. In this study, we apply a Blatter-Pattyn type of a 3D higher-order thermomechanical ice flow model to derive the horizontal velocities (Fürst et al., 2011). The HO model includes longitudinal and transverse stress gradients, which are neglected in the SIA approximation (Zekollari et al., 2013). Compared to a Full Stokes solution, the HO model assumes cryostatic equilibrium in the vertical, 245 neglecting the vertical resistive stresses (bridging effects). [In the following, the model is briefly described. Please refer to Fürst et al. \(2011\) and references therein for a more detailed description. Model parameters and constants are summarised in Table 1.](#) In the applied HO model, the problem of determining velocities in 3D is reduced to determining only the horizontal velocities, because the vertical velocities are obtained through incompressibility. Nye's generalisation of Glen's flow law, which links deviatoric stresses (τ_{ij}) to strain rates (ε_{ij}) (deformation and velocity gradients) (eq. 1), is used to quantify the 250 deformation (and the corresponding flow velocity due to internal deformation) of the ice based on the stresses that act on it.

$$\tau_{ij} = 2\eta\varepsilon_{ij} \quad (1)$$

$$\eta = \frac{1}{2}A(T_{pmp})^{-\frac{1}{n}}(\varepsilon_e + \varepsilon_0)^{\frac{1}{n}-1} \quad (2)$$

$$\varepsilon^2_e = \frac{1}{2}\varepsilon_{ij}\varepsilon_{ij} \quad (3)$$

$$255 \quad \varepsilon_{ij} = \frac{1}{2}(\frac{\partial_i u}{v} + \frac{\partial_j v}{u}) \quad (4)$$

The symbols 'i' and 'j' denote the space derivative with respect to the ith and jth spatial components. n is the power-law exponent, which is set to 3, its most common value used in previous studies (Zekollari et al., 2013). η is the effective viscosity of the ice (resistance to deformation) (eq. 2), which is defined via the second invariant of the strain rate tensor (eq. 3). ε₀ is a small offset (10⁻³⁰) that ensures finite viscosity (Fürst et al., 2011). [u and v are the horizontal components of the velocity vector.](#) The flow rate factor A(T_{pmp}) is a function of the ice temperature corrected for pressure melting (see section 3.2). The strain rate tensor is defined in terms of velocity gradients (eq. 4). [In the model, a grid with a spatial resolution of 25 m is used, and 21 layers in the vertical are considered. This resembles to an average layer thickness of 3-5 m i.e.](#) The 3D HO model has a temporal resolution of approximately one week (0.02 year). [In the model, a Weertman-type sliding law is implemented in which the basal sliding velocity \(u_b\) is proportional to the basal drag \(τ_b\) to the third power. This is a common approach in glacier modelling \(Jouvet et al., 2011; Zekollari et al., 2013\).](#)

$$u_b = -A_{sl}\tau_b^3 \quad (5)$$

heeft verwijderd: (3D)

heeft verwijderd: (HO)

heeft omhoog verplaatst [2]: The selected ice flow model has been successfully applied at different scales, ranging from small mountain glaciers (Zekollari et al., 2013; Zekollari and Huybrechts, 2015), a medium-sized ice cap (Zekollari et al., 2017) to entire ice sheets (Fürst et al., 2013, 2015).

heeft verwijderd: 1

heeft verwijderd: T

heeft verwijderd: E

Met opmaak: Standaard, Links, Regelafstand: enkel, Lettertype-uitlijning: Automatisch, Patroon: Doorzichtig

heeft verwijderd: exponent

heeft verwijderd: .

heeft verwijderd: with u the 3D components of the velocity vector...

heeft verwijderd: We use

heeft verwijderd: 1

The basal drag is calculated following the HO approximation and corresponds to the sum of all basal resistive stresses (Fürst et al., 2011). α_s is the sliding parameter which is defined as a constant in the model (Table 1). When the basal temperature is below the pressure melting point, the ice is frozen to the bedrock and there is no basal sliding. Knowing all the velocities in 3D, the continuity equation (eq. 6) is used to link the ice dynamics and the surface mass balance (m_s) (see section 3.4) to calculate the evolution of the ice thickness (H) over time (t). \bar{v} is the vertically averaged horizontal velocity vector.

$$\frac{\partial H}{\partial t} = -\nabla(\bar{v}H) + m_s \quad (6)$$

heeft verwijderd: [1](#)
 heeft verwijderd: [\(u\)](#)
 heeft verwijderd: [E](#)
 heeft verwijderd: [.](#)

3.2 3D temperature calculation and influence on flow rate factor

A full 3D calculation of the ice temperatures is performed simultaneously with the velocity calculations. This is crucial because ice temperature regulates the ice stiffness (viscosity) (see eq. 2 and eq. 7) and determines whether basal sliding occurs. The flow rate factor, $A(T_{pmp})$, affects the speed at which ice deforms under a given stress. Higher values comprise lower viscosities and hence faster deformation. Hence, the deformation rate increases with increasing temperatures because dislocations in the ice become more mobile. The temperature dependence is described using an Arrhenius relationship. At each time step, the ice temperature, which is corrected for pressure melting (eq. 8), is used to obtain the local flow rate factor using (eq. 7).

$$A(T_{pmp}) = m_a \exp(-\frac{Q}{RT_{pmp}}) \quad (7)$$

heeft verwijderd: [1](#)
 Variable
 ... [1]

heeft verwijderd: [E](#)
 heeft verwijderd: [E](#)
 heeft verwijderd: [6](#)

T_{pmp} is the ice temperature corrected for pressure melting, which is also called the homologous temperature.

$$T_{pmp} = T - \gamma H \quad (8)$$

With γ being 8.7×10^{-4} , H the ice thickness and T the ice temperature. Hence, for increasing pressure, the ice potentially starts melting at sub-zero temperatures. As such, the ice's melting point decreases with increasing pressure. E.g. ice starts to melt at -0.1°C below an ice column of 115 metres. Q is the activation energy for creep, a is the flow law coefficient, and R is the gas constant ($8.31 \text{ J mol}^{-1} \text{ K}^{-1}$) (see Table 1). For temperatures lower than -10°C , $a = 1.14 \times 10^5 \text{ Pa}^{\frac{1}{2}} \text{ yr}^{-1}$ and $Q = 60 \text{ kJ mol}^{-1}$ while for temperatures larger than -10°C , $a = 5.47 \times 10^{10} \text{ Pa}^{\frac{1}{2}} \text{ yr}^{-1}$ and $Q = 139 \text{ kJ mol}^{-1}$. m is an enhancement factor which is typically used for tuning, and which is generally comprised between 1 and 10 (Huybrechts et al., 1991; Zekollari et al., 2013). The applied HO model is a cold ice model which implies that it is not energy conserving when temperate ice is present, as it does not account for the part of internal energy coming from the latent heat of liquid water (Aschwanden et al., 2012, 2017). Hence, it does not take into account the presence of water in the ice. Nevertheless, the tuning of the enhancement factor, (m) is an implicit way to include softening effects due to factors such as water content and impurity content (Huybrechts et al., 1991).

heeft verplaatst (invoeging) [4]
 heeft verwijderd:
 heeft verwijderd: [1](#)
 heeft verwijderd: ["](#)
 heeft verwijderd: [a](#)
 heeft verwijderd: ["](#)
 heeft verwijderd: [a](#)
 heeft verwijderd: tuning
 heeft verwijderd: [T](#)
 heeft verwijderd:
 heeft omhoog verplaatst [4]: Q is the activation energy for creep and R is the gas constant ($8.31 \text{ J mol}^{-1} \text{ K}^{-1}$).
 heeft verwijderd: the
 heeft opmaak toegepast: Tekstkleur: Tekst 1

The ice temperature is determined from vertical diffusion, three-dimensional advection and the heating due to deformation and basal sliding (eq. 9). Diffusion depends on the specific heat capacity and the thermal conductivity of the ice, which depends

heeft verwijderd: E

350 on local ice temperature. Advection (horizontal and vertical) depends on the 3D velocity fields (\vec{V} is the 3D velocity vector).

The internal heat production (P) is calculated from strain heating (Huybrechts, 1996).

heeft verwijderd: I

heeft verwijderd: used by internal deformation and basal sliding

$$\frac{\partial T}{\partial t} = \frac{k}{\rho c_p} \nabla^2 T - \vec{V} \cdot \nabla T + P \quad (9)$$

355 k is the thermal conductivity; c_p is the specific heat capacity. They both are functions of the temperature (eq. 10 and eq. 11). ρ is the ice density (910 kg m^{-3}).

heeft verwijderd: E

heeft verwijderd: E

$$c_p = 146.3 + 7.253 T(K) \quad [\text{in J kg}^{-1} \text{ K}^{-1}] \quad (10)$$

$$k = 9.828 * e^{-0.0057T(K)} \quad [\text{in W m}^{-1} \text{ K}^{-1}] \quad (11)$$

360

The vertical velocity, needed for the vertical advection, is obtained through vertical integration of the incompressibility condition from the base of the glacier to a height z (eq. 12), using the kinematic boundary conditions at the bed (eq. 13). u_b and v_b are the horizontal velocity components at the bedrock, which correspond to the basal sliding components.

heeft verwijderd:

heeft verwijderd: E

heeft verwijderd: E

$$365 w_z = w_{z=h-H} - \int_{z=h-H}^{z=h} \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) dz \quad (12)$$

$$w_{z=h-H} = u_b \frac{\partial(h-H)}{\partial x} + v_b \frac{\partial(h-H)}{\partial y} - m_b \quad (13)$$

The bedrock elevation is assumed to be constant in time. h is the surface elevation (m a.s.l.), m_b is the basal melt rate, which is positive in case of melting.

370

375

Table 1. Variables, constants and their symbols, values, and units. ST refers to the Sary-Tor glacier while GR stands for the Grigoriev ice cap.

Variable	Symbol	Unit
Basal drag	τ_b	Pa
Basal melt rate	m_b	m w.e.
Basal sliding velocity components	u_b, v_b	m yr ⁻¹
Deviatoric stresses	τ_{ij}	Pa
Effective ice viscosity	η	Pa s
Enhancement factor	m	/
Flow rate factor	$\Delta(T_{\text{pmp}})$	Pa ⁻³ yr ⁻¹
Horizontal velocity components	u, v	m yr ⁻¹
Ice temperature	T	°C
Ice thickness	H	m i.e.
Ice temperature corrected for pressure melting	T _{temp}	°C
Local geothermal heat flux	g _{hf,ij}	mW m ⁻²
Local geothermal heat flux anomaly	$\Delta g_{hf,ij}$	/
	g _{hf,ij}	
Maximum winter snow depth	s _{max}	m w.e.
Mean annual air temperature	T _{ma}	°C
Mean elevation averaged within a 10x10 km moving window centred over position ij	h _{ij}	m a.s.l.
Refrozen meltwater remaining	f _{rem}	m w.e.
Second invariant of the strain rate tensor	ϵ_e^2	/
Specific heat capacity	c _p	J kg ⁻¹ K ⁻¹
Strain rate	$\dot{\epsilon}_{ij}$	/
Surface elevation	h	m a.s.l.
Surface mass balance	m _b	m w.e.
Temperature of the surface layer	T _s	°C
Thermal conductivity	k	W m ⁻¹ K ⁻¹
Time	t	yr
Vertically averaged horizontal velocity	v	m yr ⁻¹
3D velocity vector	v	m yr ⁻¹

Constant	Symbol	Value	Unit
Activation energy for creep (T < -10°C)	Q ₁	60	kJ mol ⁻¹
Activation energy for creep (T > -10°C)	Q ₂	139	kJ mol ⁻¹
Average geothermal heat flux	g _{hf}	50	mW m ⁻²
Basal sliding parameter	A _d	5 x 10 ⁻¹⁴	m ⁸ N ³ yr ⁻¹
Characteristic height	a	950	m
Dependence of the melting point with ice depth	γ	8.7 x 10 ⁻⁴	°C m ⁻¹
Exponent in Glen's flow law	n	3	/
Firn warming parameter	w _f	41	°C (m w.e.) ⁻¹
Flow law coefficient (T < -10°C)	a ₁	1.14 x 10 ⁻⁵	Pa ³ yr ⁻¹
Flow law coefficient (T > -10°C)	a ₂	5.47 x 10 ¹⁰	Pa ³ yr ⁻¹
Ice density	ρ	910	kg m ⁻³
Maximum fraction of refrozen water	P _{max}	0.6	/
Small number to make viscosity a finite number	ε ₀	10 ⁻³⁰	/
Snow insulation effect parameter	b	21	°C (m w.e.) ⁻¹
Tuning parameter of the mass balance model	c ₁	24	W m ⁻² °C ⁻¹
Tuning parameter of the mass balance model	c ₀	-199 (ST) -197 (GR)	W m ⁻²
Universal gas constant	R	8.31	J mol ⁻¹ K ⁻¹

Tabel met opmaak

3.3 Boundary conditions

395 The calculation of the ice temperature distribution requires the specification of two boundary conditions: the ice temperature of the surface layer and the temperature gradient at the bottom of the ice mass~~es~~.

3.3.1 Basal boundary conditions

At the bottom, heat is produced by geothermal heating (energy received by the ice at its base) and friction induced by basal sliding and ice deformation. Heat flow studies at regional scales in the Kyrgyz Tien Shan indicate the presence of a “cold

400 Cenozoic orogeny” with no additional heating associated to tectonic deformation and basin development. The area is characterised by an average geothermal heat flux (g_{hf}) of approximately 50 mW m⁻² (Duchkov et al., 2001; Vermeesch et al.,

heeft verwijderd: herefore, t

heeft verwijderd: relatively low

heeft verwijderd: /

heeft verwijderd: c

2015). It is important to take this effect into account which raises the temperatures, relaxing associated viscosity and hence increasing local flow velocities (Colgan et al., 2015). To consider the effect of meltwater refreezing and its warming effect on the surface layer, a parametrisation is applied that warms the surface layer for the locations where meltwater refreezes and remains at the end of a mass balance year (Reeh, 1991; Huybrechts et al., 1991; Zekollari et al., 2017). Below the equilibrium

455 line, this latent heat is lost together with meltwater runoff over the impermeable ice surface.

heeft verwijderd: warming

heeft verwijderd: is retained

However, besides a warming effect due to refreezing meltwater, other processes play a role in determining the ice temperature of the surface layer. The presence or absence of snow has been proven to be crucial for the surface layer temperature (Hooke et al., 1983; Zhang et al., 2005; Meierbach et al., 2015). The most pronounced effect of snow on the surface layer temperature 460 of ice masses occurs in the winter. During the winter months, because of the lower thermal conductivity of snow compared to ice, a significant snow cover insulates the ice below as a blanket from the strongest winter cold penetrating in the ice (Hooke et al., 1983; Zhang et al., 2005; Kronenberg et al., 2021). This ensures that average temperatures below the snow cover are usually much higher compared to the average air temperatures above the snow (Hooke et al., 1983).

heeft verwijderd: at

heeft verwijderd: s

heeft verwijderd: the

heeft verwijderd: 0

heeft verwijderd: For instance, Zhang (2005) reported an increase of the mean annual surface temperature of permafrost of $0.1^{\circ}\text{C cm}^{-1}$ of maximum snow depth.

465 In addition to the effect of the insulating snow layer, in the ablation area, the vertical velocity is positive (upward) implying that ice from deeper layers is advected upwards. Further, the upper ice layers are removed due to melting. Both these processes ensure that the ice temperature of the surface layer in the ablation area is strongly influenced by the ice temperature of the 470 deeper layers (upward vertical advection). For this reason, Wohlböck et al. (2009) and Blatter et al. (1987) proposed to use the ice temperature gradient to prescribe the ice temperature of the surface layer as boundary condition. However, this still requires a prescription of the air temperature that influences the ice surface temperature from above.

heeft verwijderd: at

heeft verwijderd: at

A last effect that can influence the surface layer temperature is the percolation of water through crevasses and moulin. This 475 water can provide heat of friction or it can refreeze, contributing to ice temperature warming (Philips et al., 2010; Gilbert et al., 2020). These processes are not considered in this study because neither for the Grigoriev nor for the Sary-Tor glacier significantly large areas with crevasses or moulins were detected. Observations show that meltwater on both ice masses is mainly discharged supraglacially.

heeft verwijderd: ice

heeft verwijderd: s

heeft verwijderd: flow

heeft verwijderd: in supraglacial melt channels and deep-water percolation

heeft verwijderd: which

heeft verwijderd:

heeft verwijderd: in the ablation area

heeft verwijderd:

heeft verwijderd: for all the effects

$$T_s = \begin{cases} \min(0^\circ\text{C}, T_{ma} + w_f r_{rem} + i_s s_{max}) & H > \text{ELA} \\ \min(0^\circ\text{C}, T_{ma} + i_s s_{max} + m_s (\text{in m i.e.}) \frac{T_{20m} - T_{10m}}{10 (\text{m i.e.})}) & H < \text{ELA} \end{cases}$$

H > ELA

(18)

H < ELA

505

T_s is the ice temperature of the surface layer which is assumed not to be influenced by the seasonal cycle. Hence, it corresponds typically to the ice temperature at 10-15 m depth. T_s is used as the upper boundary condition in the model. T_{ma} is the mean annual air temperature, which is calculated by taking the average values of all hours in one year. For the lapse rate, a sinusoidal function is used with a limit of -0.0065°C m⁻¹ in June and -0.0023°C m⁻¹ in January, based on data from Aizen et al. (1995),

510 to ensure a smooth transition of the lapse rates during the year. For the calculation of T_{ma}, as for the mass balance model (see section 3.4.), temperature data from the Kumtor-Tien Shan meteorological station (Fig. 1) is used (see Van Tricht et al. (2021b) for more information about this meteorological station). w_f corresponds to the firn warming parameter (in °C (m w.e.)⁻¹) which determines the warming for every metre of refrozen meltwater remaining (r_{rem}) at the end of each mass balance year. The i_s

515 parameter (in °C (m w.e.)⁻¹) corresponds to the snow insulation effect based on the maximum snow depth at the end of spring, (s_{max}), when snow thickness on the ice masses is typically maximum. When the ice temperature of the surface layer reaches the melting temperature, the ice temperature is kept constant at 0°C. The temperature gradient between 10-20 m ice depth ((T_{20m} - T_{10m})/10) and the surface mass balance (m_s) are used to correct for the effect of surface layer removal in the ablation area. Both the w_f and i_s parameters are calibrated based on historical and spatially distributed ice temperature measurement at

520 10-15 m depth (assumed to reflect the average annual temperature) on the Grigoriev ice cap (Dikikh et al., 1965; Thompson et al., 1993; Arkhipov et al., 2004; Takeuchi et al., 2014) (Table 2 and section 2.2). The calibrated mass balance model (see section 3.4) is used to estimate the average amount of r_{rem} at the end of the year, and s_{max}. The 15 years prior to each of the individual measurements in Table 2 are used for the average values of T_{ma}, r_{rem} and s_{max}. This period is chosen because it is large enough to filter out interannual variability and at the same time considers sufficient fluctuations in the conditions at the surface. An overview of the temperature measurements used for calibration in this study (see section 3.3) is given in Table 2.

525 As can be seen in Table 2, the ice temperatures at 10-15 m depth increase with altitude, while the mean annual air temperatures decrease with altitude. This difference can be explained by the effect of an increasing warming of the surface layer due to refreezing meltwater and maximum snow thickness for insulation.

530

heeft verwijderd: ↗

heeft verwijderd: 6

heeft verwijderd: at

heeft verwijderd: not

heeft verwijderd: winter

heeft verwijderd: at

heeft verwijderd: ↗

heeft verwijderd: .

heeft verwijderd: at the end of winter which is typically in May in the area

545 **Table 2. Measured ice temperatures of the Grigoriev ice cap at 10-15 m depth from 1962 (Dikikh et al., 1965), 1990 (Thompson et al., 1990), 2001 (Arkhipov et al., 2004) and 2007 (Takeuchi et al., 2014). T_{ma} represents the mean annual air temperature at the considered surface elevation of the 15 years prior to each individual measurement.**

Year	Height (m a.s.l.)	$T_{10-15\text{ m}} (\text{°C})$	$T_{ma} (\text{°C})$
1962	4250	-6.7	-9.9
1962	4293	-6.2	-10.0
1962	4305	-6.2	-10.1
1962	4420	-5.1	-10.4
1990	4600	-4.0	-10.9
2001	4600	-2.7	-10.8
2007	4600	-2.6	-10.7

550 3.4 Mass balance model

To obtain m_s , s_{max} , and r_{rem} , we apply the surface energy mass balance model that has been optimised, calibrated and used in Van Tricht et al. (2021b). This simple model is based on incoming solar radiation, temperature and precipitation and contains two additional tuning parameters (c_0 and c_1) representing the sum of the net longwave radiation balance and the turbulent sensible heat exchange. The temporal resolution of the mass balance model is 1 hour, while the coupling with the ice flow model occurs yearly. The input data of the mass balance model are limited to a digital elevation model (DEM), air temperatures and solid precipitation at hourly intervals. For meteorological data, the reconstructed data series of the Kumtor-Tien Shan meteorological station is used (Van Tricht et al. 2021b). The DEM is used to calculate the hourly (solar) insolation with respect to the slope and aspect of every grid point, and the effect of shadow, needed for the energy balance, and to estimate the temperature and precipitation based on the absolute elevation of the grid point. For the Sary-Tor glacier, we directly use the calibrated parameters (c_0 and c_1) from Van Tricht et al. (2021b). However, incorporating the mass balance measurements from 2020/21 in the calibration procedure reveals an optimal c_0 -value of -199 which is slightly lower than the obtained value of -193 based on calibration data from 2014-2020. Concerning the Grigoriev ice cap, initially, the same parameters are used as for the Sary-Tor glacier. The c_1 parameter is fixed at 24 as this was found to be optimal for the other glaciers in the area (Van Tricht et al., 2021b). However, the c_0 parameter as well as the precipitation gradient are then tuned by minimising the root mean square error (RMSE) between modelled and measured mass balances from Dyurgerov (2002), Arkhipov et al. (2004) and Fujijita et al. (2011) (Table 3). For instance, Arkhipov et al. (2004) determined the average accumulation at the summit of the Grigoriev ice cap based on the detection of annual layers to be 0.26 m w.e. yr^{-1} between 1990 and 2000 which appeared to be slightly less than the 0.32 m w.e. yr^{-1} between 1963 and 1989.

560
570

heeft verwijderd:

heeft opmaak toegepast: Letertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië)

heeft opmaak toegepast: Letertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië)

heeft opmaak toegepast: Letertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië)

heeft verwijderd: 1

heeft verwijderd: 2

575 Table 3 Data used for the calibration of the mass balance model for the Grigoriev ice cap. m_s represent the surface mass balance.

Year	Mean specific m_s	m_s at the summit	Average m_s (<4200 m a.s.l.)	Reference
Average 1963/64-1988/89		0.32 m w.e. yr ⁻¹		(Arkhipov et al., 2004)
1986/87	-0.22 m w.e. yr ⁻¹		-0.50 m w.e. yr ⁻¹	(Dyurgerov, 2002)
1987/88	-0.30 m w.e. yr ⁻¹		-0.90 m w.e. yr ⁻¹	(Dyurgerov, 2002)
Average 1990/91-1999/00		0.26 m w.e. yr ⁻¹		(Arkhipov et al., 2004)
2005/06		0.16 m w.e. yr ⁻¹		(Fujita et al., 2011)
2006/07	-0.36 m w.e. yr ⁻¹	0.33 m w.e. yr ⁻¹	-1.80 m w.e. yr ⁻¹	(Fujita et al., 2011)

The calibration of the mass balance model for the Grigoriev ice cap reveals optimal values for a precipitation gradient of +150 mm km⁻¹ yr⁻¹ and a c_0 value of -197 W m⁻². This rather low value for the precipitation gradient (compared to the Sary-Tor glacier) appears to be necessary to limit the accumulation on the summit of the ice cap, as was measured and reconstructed by Arkhipov et al. (2004) and Fujita et al. (2011). Such a low gradient can be explained by the location of the Grigoriev ice cap on the top of an isolated mountain, lower than the surrounding mountains, reducing the effect of orographic precipitation enhancement and making it more exposed to wind erosion. On the other hand, the Sary-Tor glacier is located at the north-western side of the Ak-Shyirak massive which could be more exposed to precipitation. The calibrated c_0 value is of equal magnitude compared to the values found for the nearby located Bordu and Sary-Tor glaciers (Van Tricht et al., 2021b).

A crucial output of the mass balance model for the determination of the thermal regime is the amount of meltwater that refreezes in the snowpack (see section 3.3). Following Van Tricht et al. (2021b), we assume all snowmelt to percolate into the snow cover and to be retained and to refreeze until a maximum amount of P_{max} (taken as 60% of the snow depth) is reached (Reeh, 1991; Huybrechts et al., 1991). At every timestep, the snow depth and the amount of refrozen water are updated. Finally, at the end of the mass balance year, it is assumed that the remaining refrozen meltwater forms r_{rem} . Refreezing of meltwater below the last summer horizon can occur if not all pore spaces of the existing snow cover are occupied (< 60%), as it was shown to be an important process for glaciers and ice caps in the area (Dyurgerov and Mikhaleko, 1995; Takeuchi et al., 2014; Kronenberg et al., 2016). The maximum snow depth s_{max} is determined by considering the maximum snow depth before the end of May when average snow depths are typically maximal on glaciers in the area (Van Tricht et al., 2021c). During the summer months, there can still occur regular snow events during convective showers, but melt then compensates for the accumulation, so that the snow thicknesses on the ice masses usually no longer increase, and snow insulation is focused on winter and spring.

heeft verwijderd: 2... Data used for the calibration of the mass balance model for the Grigoriev ice cap. m_s represent the surface mass balance. All values are in m w.e. yr⁻¹. ... [2]

heeft opmaak toegepast: Lettertype: 5 pt

heeft verwijderd: at the front

heeft opmaak toegepast: Lettertype: Vet

heeft opmaak toegepast: Lettertype: Vet, Niet Cursief

heeft opmaak toegepast: Lettertype: Vet

heeft opmaak toegepast: Lettertype: Vet, Engels (V.S.)

heeft opmaak toegepast: Lettertype: Vet

heeft opmaak toegepast: Lettertype: Vet, Niet Cursief

Tabel met opmaak

heeft verwijderd: (Arkhipov et al., 2004)

heeft opmaak toegepast: Lettertype: 9 pt

heeft verwijderd: -19...7 (Dyurgerov, 2002) ... [3]

heeft opmaak toegepast: Lettertype: 9 pt

Tabel met opmaak

heeft verwijderd: -19...8 (Dyurgerov, 2002) ... [4]

heeft opmaak toegepast: Lettertype: 9 pt

heeft verwijderd: 20... (Arkhipov et al., 2004) ... [5]

heeft opmaak toegepast: Lettertype: 9 pt

Tabel met opmaak

heeft verwijderd: -20...6 (Fujita et al., 2011) ... [6]

heeft opmaak toegepast: Lettertype: 9 pt

heeft verwijderd: -20...7 (Fujita et al., 2011) ... [7]

heeft opmaak toegepast: Lettertype: 9 pt

heeft opmaak toegepast: Lettertype: 9 pt

Tabel met opmaak

heeft verwijderd: remains ...refreezes in the snowpack and refreezes at the end of the mass balance year ...see section....3.3. Following Van Tricht et al. (2021b), we assume all snowmelt to be retained...percolate into the snow cover and to be retained and to refreeze ...in the snowpack ...until a maximum amount of P_{max} (taken as 60% of the snow depth) is reached (Reeh, 1991; Huybrechts et al., 1991). At every timestep, the snow depth and the amount of retained refrozen water are updated. Finally, at the end of the mass balance year, it is assumed that the remaining retained water refreezes...refrozen meltwater and ...forms r_{rem} . Refreezing of meltwater below the last summer horizon can occur if not all pore spaces of the existing snow cover are occupied (< 60%), as it was shown to be an important process for glaciers and ice caps in the area (Dyurgerov and Mikhaleko, 1995; Takeuchi et al., 2014; Kronenberg et al., 2016). The maximum snow depth s_{max} is determined by considering the maximum snow depth at the ... [8]

665 **3.5 Geometric data**

Several geometric datasets are needed to apply the ice dynamic model and the surface mass balance model. The experiments in this study are performed under average 1960-1990 climatic conditions. Therefore, the geometry of the Grigoriev ice cap and the Sary-Tor glacier is reconstructed for 1990 [as a starting point for all simulations](#).

670 **3.5.1 Outlines**

The outlines of both ice masses are initially taken from the Randolph Glacier Inventory v6 (RGI, 2017). Subsequently, optical satellite images from Landsat 5, valid for August 1990, are used to update the outline using manual delineation based on visual inspection of the ice – bedrock boundary. [Furthermore, a LIA ice mask is created from visual detection of unique LIA moraines, clearly present on satellite images \(Landsat and Sentinel-2\) for both the Grigoriev ice cap and the Sary-Tor glacier. This ice mask serves as the maximum area of both ice masses which is allowed to be occupied by ice. When the ice were to expand beyond the mask, the ice is removed. However, this never applies to the geometry for average 1960-1990 climatic conditions as this remains entirely within the LIA mask. Concerning the northern boundary of the Grigoriev ice cap, when ice flows beyond the predefined calving front, it is automatically removed.](#)

heeft verplaatst (invoeging) [3]


heeft verwijderd: A

heeft verwijderd: ↗

675 **3.5.2 Bedrock DEM**

Concerning the Sary-Tor glacier, the ice thickness measurements were performed with a VIRL-6 GPR radar at 20 MHz central frequency (Petrakov et al. 2014). A maximum part of the accessible area of the glacier was covered with a grid of longitudinal and transverse profiles. Then, an ice thickness field was reconstructed using interpolation with an estimated accuracy of 15-20 m and a volume of $0.126 \pm 0.001 \text{ km}^3$ was obtained (Petrakov et al., 2014). A bedrock DEM is obtained by subtracting the ice thickness of Sary-Tor glacier in 2013 from the surface elevation in 2013 (obtained from the TanDEM-X DEM). [Corrected for ice melt between 2013 and 2021, a maximum ice thickness of \$157 \pm 15 \text{ m}\$ in 2021 is obtained, and a total ice volume \$0.118 \pm 0.001 \text{ km}^3\$. Concerning the ice thickness of the Grigoriev ice cap, we performed ~ 500 GPR-measurements over the entire ice cap in the summer of 2021 using a Narod Radio Echo Sounding system with a lower frequency of 5 MHz, reducing the necessity for interpolation over larger distances between the measurement transects \(Fig. 2\). A maximum ice thickness of \$114 \pm 10 \text{ m}\$ was measured.](#) Subsequently, following Van Tricht et al. (2021a), a glacier wide ice thickness field is reconstructed using a constant yield stress model for the unmeasured areas [resulting in a total ice volume of \$0.391 \pm 0.07 \text{ km}^3\$](#) . During the field campaign, UAV images were collected using a DJI Phantom 4 RTK from about 200 m above the ice cap. These images were subsequently used in Pix4D to infer an accurate and centimetre resolution digital surface model (DSM), as in Van Tricht et al. (2021c). A bedrock DEM is determined by subtracting the reconstructed ice thickness distribution from the DSM.

heeft verwijderd: ↗



Met opmaak: Centrum

Figure 2: Ice thickness distribution of the Grigoriev ice cap and the Sary-Tor glacier in 2021. The locations of the ice thickness measurements of 2021 on the Grigoriev ice cap are added in red. The background is a Sentinel-2 image from 26th of July 2021. The location of the ice core made in 2007 (Takeuchi et al., 2014) is added with a black triangle.

heeft opmaak toegepast: Lettertype: 9 pt, Vet

heeft opmaak toegepast: Lettertype: 9 pt, Vet

Met opmaak: Regelafstand: Meerdere 1.15 rg

3.5.3 DEM of 1990

For consistency with the 1960-1990 climatic conditions, a DEM is created for 1990 as initial state to be used in the glacier model. Using the calibrated mass balance model (see section 3.4), the cumulative mass balance is calculated between 1990 and the year in which the ice thickness was measured (2021 for the Grigoriev ice cap and 2013 for the Sary-Tor glacier). Under the assumption that $\Delta\text{SMB} = \Delta H$ (hence neglecting the role of ice dynamics for these periods), the surface elevation of 1990 is then inferred by adding ΔSMB to the DEM of 2021 and 2013 for the glacial area (Fig. 3). Outside the glacial area of 2013 and 2021, the surface elevation is reconstructed by setting the elevation equal to the bedrock elevation at the 1990 front, attaching to the DEM of 2013/2021 at the observed glacier front, an interpolating in between (Fig. 3).

heeft verwijderd: .

heeft verwijderd: 2

heeft verwijderd: 2




Figure 3. Reconstruction of the 1990 geometry from a combination of modelling the cumulative surface mass balance (Δm_s) and interpolation.

720 Reconstructed for the 1990 geometry, we obtain an area of the Grigoriev ice cap of $8.746 \pm 0.42 \text{ km}^2$ (~15% larger than the present-day area), with a maximum ice thickness of $115 \pm 10 \text{ m}$, and a total ice volume of $0.513 \pm 0.11 \text{ km}^3$ (~30% more than the present-day area). The corresponding values for the Sary-Tor glacier are $2.789 \pm 0.22 \text{ km}^2$ (~21% larger than the present-day area), a maximum ice thickness of $158 \pm 15 \text{ m}$, and a total volume of $0.158 \pm 0.002 \text{ km}^3$ (~34% more than the present-day volume). Hence, both ice bodies lost a substantial fraction of their area and volume over the last 31 years, similar to observations in other studies (Aizen et al., 2006; Kutuzov and Shahgedanova, 2009; Sorg et al., 2012). The created 1990 geometry of both ice masses is used to initiate the experiments below. The eastern tributary of the Sary-Tor glacier (visible at the right panel of Figure 2) is not included for the 1960-1990 runs, as it was already disconnected from the Sary-Tor main glacier at the end of this time frame.

730 3.6 Experiments and sensitivity analysis

First, multiple experiments are performed using the coupled surface mass balance and the ice flow models with standard settings for the basal sliding parameter A_s ($5.10^{-14} \text{ m}^8 \text{ N}^{-3} \text{ yr}^{-1}$, see Table 1). The velocity and temperature fields are run under constant 1960-1990 climatic conditions. During the runs, the geometry is freely evolving with a maximum limit of the LIA extent, until a steady state is reached. When a steady state is reached, modelled ice thickness is compared with the 1990 reconstructed thickness, and temperature fields are compared with measured temperatures as well as with detected boundaries between cold and temperate ice (Petrakov et al., 2014). A constant mass balance correction over the entire ice mass is implemented to reach the extent of the 1990 geometry.

heeft verwijderd: 2

Met opmaak: Bijchrift

heeft omhoog verplaatst [3]: A LIA ice mask is created from visual detection of unique LIA moraines, clearly present on satellite images (Landsat and Sentinel-2) for both the Grigoriev ice cap and the Sary-Tor glacier. This ice mask serves as the maximum area of both ice masses which is allowed to be occupied by ice. When the ice were to expand beyond the mask, the ice is removed. However, this never applies to the geometry for average 1960-1990 climatic conditions as this remains entirely within the LIA mask.

heeft verwijderd: 3.5.4 Little Ice Age mask

heeft verwijderd: 3.6

heeft opmaak toegepast: Lettertype: (Standaard) Times New Roman, Tekstkleur: Auto

heeft verwijderd:

Met opmaak: Standaard

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), Tekstkleur: Tekst 1

heeft opmaak toegepast: Tekstkleur: Auto

heeft verwijderd: SL

755 Then, a sensitivity analysis is performed with regards to the different parameters affecting the temperature distribution in the ice masses. This concerns specifically the i_s and w_f parameters, which are altered within $\pm 10\%$. In each run, one parameter is changed while the other remains fixed at its optimal value. The sensitivity with regard to the geothermal heat flux is also assessed by using 50% and 200% of the average geothermal heat flux. In every experiment, first, the geometry is freely evolving until a steady state is reached using the optimal parameter values. Then, the parameter under consideration is altered, 760 and the geometry is kept fixed to avoid an effect of a thinner or thicker ice mass caused by a change in the ice viscosity and to simplify the interpretation of the results.

heeft verwijderd: initial

heeft verwijderd: g_{hf} -value

Finally, a sensitivity to changes in boundary conditions is investigated by altering the precipitation {-40% to 40%} and the surface temperatures {-2°C to +3°C}. These ranges were selected based on observed (Van Tricht et al., 2021b) and projected 765 changes for the area. For this analysis, the geometry can evolve freely until a new steady state is reached in equilibrium with the altered surface boundary conditions. This analysis can give an indication about the evolution of the thermal regime in the past and into the future, which is part of the discussion section.

heeft verwijderd: 4.1

4.1 Geometric input data and characteristics

During the 2021 field campaign, ~500 well-spread ice thickness measurements were performed on the Grigoriev ice cap, reducing the necessity for interpolation over larger distances between the measurement transects (Fig. 3). In 2021, the Grigoriev ice cap is characterised by an area of 7.545 km², a maximum ice thickness of 114 m, and a total ice volume of 0.513 km³ (~30% more). Regarding the Sary-Tor glacier in 2021, the glacier area amounts to 2.250 km² with a maximum ice thickness of 157 m and a total ice volume 0.118 km³, based on measurements of Petrakov et al. (2014) which we corrected for melt between 2013 and 2021. 4

Reconstructed for the 1990 geometry, we obtain an area of the Grigoriev ice cap of 8.746 km² (~15% larger), with a maximum ice thickness of 115 m, and a total ice volume of 0.513 km³ (~30% more). The corresponding values for the Sary-Tor glacier are 2.789 km² (~21% larger), a maximum ice thickness of 158 m, and a total volume of 0.158 km³ (~34% more). Hence, both ice bodies lost a substantial fraction of their area and volume over the last 31 years, similar to observations in other studies (Aizen et al., 2006; Kutuzov and Shahgedanova, 2009; Sorg et al., 2012). The created 1990 geometry of both ice masses is used to initiate the experiments below. The eastern tributary of the Sary-Tor glacier is not included for the 1960–1990 runs, as it was already disconnected from the Sary-Tor main glacier at the end of this time frame. 4

[9]

heeft opmaak toegepast: Tekstkleur: Auto

Met opmaak: Kop 1, Regelafstand: enkel

heeft verwijderd: 4.2 Calibrated parameters and boundary conditions


heeft verwijderd: 2

heeft verwijderd: .1 Surface mass balance

The calibration of the surface mass balance model for the Grigoriev ice cap reveals optimal values for a precipitation gradient of +150 mm km⁻¹ yr⁻¹ and a c_0 value of -197 W m⁻². This rather low value for the precipitation gradient (compared ... [10]

heeft verwijderd: input data

heeft verwijderd: .




870

Figure 4: Average surface mass balance (m_s) in m w.e. yr^{-1} of the Grigoriev ice cap and the Sary-Tor glacier for the mean 1960-1990 climatic conditions. The background is a Sentinel-2 image from 26th of July 2021.

4.2 Boundary conditions

4.2.1 Firn warming and snow insulation

875 Using the output from the surface mass balance model, r_{rem} and s_{max} , and the temperature data, the optimal combination of the w_f and the i_s parameters is 41 and 22 (in $^{\circ}C$ (m w.e.) $^{-1}$) (Fig. 5a). The corresponding RMSE between the measured (see Table 2) and modelled temperatures at 10-15 m depth is equal to $0.28^{\circ}C$ (Fig. 5a,b), which is a close correspondence.



heeft verwijderd:

heeft verwijderd: 1

heeft verwijderd: 2.2

heeft verwijderd: 1





Figure 5: (a) Calibration of the w_f and the i_s parameters based on ice temperature measurements of the Grigoriev ice cap. (b) Modelled and measured temperatures for different elevations and times of the Grigoriev ice cap.

885

The obtained firm warming parameter (w_f) value ($41^{\circ}\text{C (m w.e.)}^{-1}$) is larger than values found in previous studies for e.g. the Greenland ice sheet ($26.6^{\circ}\text{C (m w.e.)}^{-1}$; Reeh, 1991; Huybrechts et al., 1991), and Hans Tausen ice cap ($22^{\circ}\text{C (m w.e.)}^{-1}$; Zekollari et al., 2017). The parameter to consider for the insulation of snow (optimal $i_s = 22^{\circ}\text{C (m w.e.)}^{-1}$) is slightly larger than the values found in Hooke et al. (1983) (about 4°C for 1 metre of snow ($\sim 12\text{--}20^{\circ}\text{C (m w.e.)}^{-1} \text{yr}^{-1}$) on the Storglaciären glacier, or about 2.5°C warming for 0.6 metres of snow ($\sim 13\text{--}21^{\circ}\text{C (m w.e.)}^{-1} \text{yr}^{-1}$) for the Barnes ice cap). However, Zhang (2005) and Yershov (1998) reported a mean annual temperature increase of about $0.1^{\circ}\text{C cm}^{-1}$ of snow, which is equivalent to $10^{\circ}\text{C m}^{-1} \text{yr}^{-1}$ of snow ($\sim 30\text{--}50^{\circ}\text{C (m w.e.)}^{-1} \text{yr}^{-1}$). In the applied model, the calibrated values of w_f and i_s are kept constant in space and time.

890

895

4.2.2 Surface warming and temperatures

The input of latent heat from percolating and refreezing meltwater is the strongest in the middle part of the accumulation zones (Fig. 6) (infiltration congelation/recrystallisation zone). At the highest elevations, the amount of refrozen water is low, caused by the absence of substantial melt (too low temperatures). This area corresponds to the dry snow zone or recrystallisation zone (Shumskiy, 1955). The effect related to snow insulation is more homogeneously distributed over the ice bodies (Fig. 6) because almost all precipitation in winter and spring is snow, independent from the elevation on the glacier, which ensures that differences in maximum snow depth at the end of spring are only related to precipitation enhancement with elevation. Combining the effect of firm warming and snow insulation, an entirely cold surface condition is obtained for the Grigoriev ice cap while a polythermal surface condition is obtained for the Sary-Tor glacier (Fig. 6).

900

heeft verwijderd: Closer to the equilibrium line, the snowpack present at the end of the year is too thin for large amounts of refrozen meltwater. ...

heeft verwijderd: Further, values of equal magnitude were also obtained in Liang and Zhou (1993) and Zhou et al. (2000), which increases the confidence in the value derived in this study.

heeft verwijderd: 2.3

heeft verwijderd: 1

heeft verwijderd: Further, values of equal magnitude were also obtained in Liang and Zhou (1993) and Zhou et al. (2000), which increases the confidence in the value derived in this study.

heeft verwijderd: 2.3

heeft verwijderd: Further, values of equal magnitude were also obtained in Liang and Zhou (1993) and Zhou et al. (2000), which increases the confidence in the value derived in this study.

For both ice masses, the coldest ice temperatures at the surface can be found in the ablation areas, where the formation of superimposed ice at the end of summer is absent because the meltwater is evacuated, and where a thinner snow cover in winter allows for a larger surface cooling (Fig. 6). The temperatures are the highest in the accumulation areas. Concerning the Sary-

930 Tor glacier, there is a temperate band between ~4200-4500 m with temperatures equal to zero degree. The hypothesis is that these higher ice temperatures are dissipated downwards through conduction and vertical movement towards lower elevations being the reason for the expected polythermal regime of the valley glaciers (Petrakov et al., 2014), such as also observed on the Tuyuksu glacier in the Northern Tien Shan (Nosenko et al., 2016). At the highest elevations of the Sary-Tor glacier, the
935 ice temperature at the surface are again lower because of the absence of meltwater available for refreezing.





Figure 6: Warming due to refreezing of meltwater (firn warming) (panels a and d), due to snow insulation (panels b and e), and resulting mean annual surface layer temperature (T_s) (panels c and f) for both ice masses according to constant 1960-1990 average
940 climatic conditions and masked for the 1990 outline (black lines). The background is a Sentinel-2 image from 26th of July 2021. The x and y-axes on the left panels stand for the other panels as well.

heeft verwijderd: and not retained in the snow

heeft verwijderd: ensures a larger cooling of the surface



heeft verwijderd:

Met opmaak: Centrum, Regelafstand: 1.5 regel

heeft verwijderd: and

heeft verwijderd: 10 m

heeft verwijderd: ice

950 4.3 Modelled steady state velocities and thickness

Under constant 1960-1990 climatic conditions and with the applied surface mass balance adjustments, the obtained steady states match the reconstructed 1990 geometry (extent and ice thickness) relatively well (Fig. 7). The RMSE between the reconstructed and modelled ice thickness is about 22.01 m for the Sary-Tor glacier and 15.20 m for the Grigoriev ice cap. The enhancement factor is set at 3 for the Grigoriev ice cap and at 4 for the Sary-Tor glacier, which appears to be the optimal value to match with the reconstructed ice thickness of 1990. For the Sary-Tor glacier, in general, modelled and reconstructed ice thickness match well (Fig. 7), and discrepancies are mainly occurring in areas where no GPR measurements were performed (Petrakov et al., 2014). Regarding the Grigoriev ice cap, the modelled ice thickness is slightly thicker and extends generally further at the edges of the ice cap compared to the 1990 reconstruction. At the top, on the other hand, the modelled ice thickness is thinner (15-25 m). The latter might be associated to the complexity of modelling ice thickness in the vicinity of ice divides (Pattyn, 2003), or to the measurements of the ice thickness itself. It is unlikely that the underestimated modelled ice thickness at the top of the Grigoriev ice cap is related to an excessively high ice temperature, as to correctly model the ice thickness here would require an enhancement factor of 1, which corresponds to a lower temperature of -5°C (Huybrechts and Oerlemans, 1988). This is far outside the limit of the values of the observed ice temperatures (Thompson et al., 1993; Arkhipov et al., 2004; Takeuchi et al., 2014).

965




Figure 7: (a,c) Ice thickness difference between the modelled steady state (~1960-1990 average climatic conditions) and the reconstructed geometry of 1990. The used flowline is added on the map with a black line. (b,d) Reconstructed (initial glacier surface) and modelled (glacier surface) ice thickness along the central flowlines. The upper panels are for the Sary-Tor glacier, the lower for the Grigoriev ice cap. The background of the left maps is a Sentinel-2 image from 26th of July 2021. The outlines are from 1990.

970


heeft verwijderd: 1

Even though there is generally a good agreement between modelled and reconstructed ice thickness, a few larger discrepancies exist (Fig. 7). For the highest parts of the Sary-Tor glacier, the modelled ice thickness is smaller than the reconstructed ice thickness. However, in this area, no GPR measurements were performed, demonstrating that this area is associated with the largest uncertainty in ice thickness

heeft verwijderd: In addition, the modelled ice thickness is specifically larger than the reconstructed ice thickness on the lateral margins of the Sary-Tor glacier in the ablation area. This is likely related to the reconstructed geometry of 1990, or it might be related to the reconstructed geometry not being in equilibrium with the 1960-1990 climatic conditions (Fig. 7). Furthermore, the modelled ice thickness is thicker for the eastern tributary, but here, as well, no direct ice thickness measurements were performed (Petrakov et al., 2014). 1

heeft verwijderd: 1

heeft verwijderd: degrees



heeft verwijderd:

heeft verwijderd: 1

Regarding the horizontal (surface) ice velocities, the Grigoriev ice cap is generally characterised by a very low velocity between 0 and 5 m yr⁻¹ (Fig. 8c). Such low values were also observed by Fujijita et al. (2011) and modelled by Nagornov et al. 995 (2006). The very low ice velocities imply that the ice is only advected very slowly downstream, keeping ice fluxes low. Only for the outlet glacier in the eastern part, which flows towards the Popova glacier, velocities increase to > 20 m yr⁻¹ due to the steeper slope in this area. The ice velocities are mostly significantly higher for the Sary-Tor glacier with a maximum slightly above 30 m yr⁻¹ (Fig. 8b). This implies that the horizontal advection of ice for Sary-Tor glacier is much more pronounced.

Comparison between modelled velocities and velocities derived from mass balance stake displacements at the Sary-Tor glacier shows a close correspondence (RMSE = 1.5 m yr⁻¹).





Figure 8: (a,c) Horizontal surface ice velocities of the Grigoriev ice cap and the Sary-Tor glacier. The flowline is added on the map with a black line. (b,d) Horizontal ice velocities along the flowlines of both ice masses. The upper panels are for the Sary-Tor glacier, the lower for the Grigoriev ice cap. The background of the left maps is a Sentinel-2 image from 26th of July 2021. The outlines are from 1990.

heeft verwijderd: .

heeft opmaak toegepast: Superscript



heeft verwijderd:

heeft opmaak toegepast: Tekstkleur: Auto

Met opmaak: Standaard, Regelaafstand: enkel

1010 4.4 Thermal regime under the 1960-1990 climatic conditions

With the imposed boundary conditions and the modelled thickness and velocity structure, the steady state englacial ice temperatures are determined. They range between -7.70°C and 0°C for the Sary-Tor glacier and between -6.61°C and -0.03°C for the Grigoriev ice cap. A polythermal regime is obtained for the Sary-Tor glacier (Fig. 9). Temperate ice is present over a large part of the glacier base, originating in the accumulation area which is subsequently advected downstream with the ice flow. At the snout of the Sary-Tor glacier, and in the largest part of the ablation area, the ice is cold. Over the entire glacier volume, about 38% of the ice of the Sary-Tor glacier is temperate, classifying this glacier reliably as a polythermal glacier. Concerning the Grigoriev ice cap, the model results show cold ice except for a very thin layer at the bed in the central part, which is characterised by temperate ice. This concerns less than 1% of the total volume. Hence, the Grigoriev ice cap can be classified as a cold ice cap. The thin layer of temperate ice in the central area at the bed was also observed on GPR profiles measured by I. Lavrentiev in 2018 (personal communication), increasing the confidence in the obtained results.





Figure 9: Modelled ice temperatures of the Sary-Tor glacier and the Grigoriev ice cap along the central flowline using a free evolving geometry and corrected for the pressure melting point. The vertical line in the plot of the Sary-Tor glacier corresponds to the cross-section used in Fig. 10.

The difference between the thermal regime of the two ice masses can be attributed to a combination of several factors. These are related to the surface conditions, the geothermal heat flux corrected for topographic relief, the ice thickness, and the ice dynamics. First, the accumulation area of the Grigoriev is located at higher altitudes and has therefore a lower mean annual air temperature (T_{ma}). Next to that, the Grigoriev ice cap is characterised by a thinner maximum snow thickness (S_{max}) and remaining refrozen water (r_{ref}) reducing the effect of insulating snow and firm warming. Further, the ice thickness of the Grigoriev ice cap is thinner, resulting in an improved transport of the geothermal heat. Finally, the faster ice flow of the Sary-

heeft verwijderd: (personal communication)

heeft verwijderd: .

T_{pmp} , Sary-Tor glac



heeft verwijderd:

heeft verwijderd: indicated on

heeft verwijderd: .

heeft verwijderd: smaller amount of

heeft verwijderd: :

heeft verwijderd: :

heeft verwijderd: r

heeft verwijderd: the firm warming effect and the

heeft verwijderd: flux

Tor glacier leads to significantly more efficient advection that can transport the heat produced in the lower accumulation area downstream, which is absent for the Grigoriev ice cap. The difference between the thermal regime of the Grigoriev ice cap and the Sary-Tor glacier found in this study shows that both ice masses are characterised by different dynamics and characteristics. Because of the lower average ice temperature, the ice viscosity of the Grigoriev ice cap is larger than that of the Sary-Tor glacier. This causes the ice to deform more slowly. Further, the presence of temperate basal ice for the Sary-Tor glacier implies that this ice mass can slide over its base, while this is not the case for the Grigoriev ice cap. Using a typical basal sliding parameter (see section 3.6 and Table 1), a maximum basal sliding speed of 13 m yr^{-1} is found for the Sary-Tor glacier, for which 55% of the bed is at the pressure melting point. Such basal sliding was also observed for the Ürümqi glacier in the Chinese Tien Shan (Maohuan et al., 1989, 1992). In addition, the presence of temperate ice at the surface causes the Sary-Tor glacier to have zones around the elevation of the EL where water can infiltrate in the glacier (Fig. 9), as temperate ice contains networks of intragranular veins where the flow of water is possible if it is not blocked by air bubbles (Nye and Frank, 1973; Liboutry, 1996; Ryser et al., 2013). Cold ice in contrast, is fracture free (water refreezes within hours in smaller cracks), making the ice impermeable to water which facilitates the formation of persistent and deeply incised water streams and supraglacial lakes (Boon and Sharp, 2003; Ryser et al., 2013). In the ablation area of the Sary-Tor glacier, all the water drains away superficially, which can most likely be explained by the impermeable cold ice (Fig. 9). This is the case for the largest part of the surface of the Grigoriev ice cap. Only in the upper part of the ice cap, ice lenses were observed when boreholes were made (Thompson et al., 1997; Takeuchi et al., 2014). As a consequence of the ice temperature differences, it might be more difficult to reproduce measured flow velocities correctly with ice flow models employing isothermal (cold or temperate) ice (Riesen et al., 2010; Ryser et al., 2013).

heeft verwijderd: 1

heeft verwijderd: ensures

heeft verwijderd: ensures a greater stiffness which causes the

heeft verwijderd: intrinsically

heeft verwijderd: slower to

heeft verwijderd: .

heeft verwijderd: .

heeft verwijderd: .

heeft verwijderd: entire

4.4.1 Validation of the thermal regime of the Sary-Tor glacier

While the ice temperature measurements of the Grigoriev ice cap are used to calibrate the firn warming and snow insulation parameters (see section 4.2.2), a validation of the thermal regime of the Sary-Tor glacier is performed by comparing modelled ice temperatures with GPR profiles of Petrakov et al. (2014) showing the boundary between temperate and cold ice. For this, a transverse profile in the middle part of the glacier is considered (Fig. 9). There is clearly a similarity regarding the presence of temperate ice over the bedrock, which is about 1/3 of the total ice thickness (black line) (Fig. 10).

heeft verwijderd: .




Figure 10: Validation of the thermal regime with a GPR profile showing the boundary between cold and temperate ice. The purple line is the modelled boundary between cold and temperate ice while the yellow line corresponds to the observed boundary between cold and temperate ice. Panel b is a radargram reproduced from Petrakov et al. (2014) with permission from the publisher. The figure clearly shows two sections of cold (upper part) and temperate (lower part) ice. The latter is characterised by the presence of small hyperbolic diffraction features due to the presence of liquid water in the ice, typical for temperate (shown here as warm) ice.

In addition, a point-by-point comparison with measurements of Petrakov et al. (2014) is performed which reveals a mean difference of -2 m i.e., and a root mean square error of 28 m i.e. between modelled and measured thickness of the temperate ice layer. Further, for all locations where temperate ice was effectively observed on the GPR profiles, the model output also shows the presence of temperate ice. Given the uncertainty in both the measurements and the model output, this is a strong agreement, and it increases the confidence in the obtained results.

4.4.2 Ice temperature profile at the summit of the Grigoriev ice cap

For the Grigoriev ice cap, the modelled temperature profile for average 1960-1990 climatic conditions is compared with the measured ice temperatures of Takeuchi et al. (2014) from 2007 near the summit of the ice cap (Fig. 2 and Fig. 11). A large mismatch is found for depths below 20 metres which increases with depth. Since the temperatures of the measured profile decrease with depth, while there cannot be advection of cold ice from upper areas close to the summit, this must be related to a recent warming at the surface which has not yet reached the lower ice layers (Vincent et al., 2007; Cuffey and Paterson, 2010). In other words, the measured ice temperature profile of 2007 is not in equilibrium with the imposed constant 1960-1990 climatic conditions.

For a bottom temperature of -4°C (as was measured by Takeuchi et al. (2014)), the surface temperatures should have been at least 2-3 degrees colder. Such a temperature increase was noticed by Thompson et al. (1993) by comparing borehole temperatures at 10-15 m depth of the 1960's (Dikikh, 1965) with 1990 measurements on the Grigoriev ice cap. However, as no such temperature increase is recorded at the nearby Kumtor-Tien Shan weather station nor a significant increase in

085

090

100

1105

095

105

115

120

125

130

135

140

145

150

155

160

165

170

175

180

185

190

195

200

205

210

215

220

225

230

235

240

245

250

255

260

265

270

275

280

285

290

295

300

305

310

315

320

325

330

335

340

345

350

355

360

365

370

375

380

385

390

395

400

405

410

415

420

425

430

435

440

445

450

455

460

465

470

475

480

485

490

495

500

505

510

515

520

525

530

535

540

545

550

555

560

565

570

575

580

585

590

595

600

605

610

615

620

625

630

635

640

645

650

655

660

665

670

675

680

685

690

695

700

705

710

715

720

725

730

735

740

745

750

755

760

765

770

775

780

785

790

795

800

805

810

815

820

825

830

835

840

845

850

855

860

865

870

875

880

885

890

895

900

905

910

915

920

925

930

935

940

945

950

955

960

965

970

975

980

985

990

995

1000

1005

1010

1015

1020

1025

1030

1035

1040

1045

1050

1055

1060

1065

1070

1075

1080

1085

1090

1095

1100

1105

1110

1115

1120

1125

1130

1135

1140

1145

1150

1155

1160

1165

1170

1175

1180

1185

1190

1195

1200

1205

1210

1215

1220

1225

1230

1235

1240

1245

1250

1255

1260

1265

1270

1275

1280

1285

1290

1295

1300

1305

1310

1315

1320

1325

1330

1335

1340

1345

1350

1355

1360

1365

1370

1375

1380

1385

1390

1395

1400

1405

1410

1415

1420

1425

1430

1435

1440

1445

1450

1455

1460

1465

1470

1475

1480

1485

1490

1495

1500

1505

1510

1515

1520

1525

1530

1535

1540

1545

1550

1555


1560

precipitation (Van Tricht et al., 2021b), this rapid increase in ice temperature is likely related due to an increase in the formation of r_{rem} or because of a lowering of the surface.

To assess the time varying ice temperature for the location of the borehole, the surface layer temperature is calculated for the 1130 1920-1950 period and the 1980-2010. The latter is about 4 degrees lower for the 1920-1950 mean climatic conditions. The difference is caused by a T_{ma} difference of 1.1°C (recorded at the Kumtor meteorological station), a difference of 2.5°C caused by an increase in r_{rem} , and a difference of 0.4°C related to a change in snow insulation. By gradually increasing the surface temperatures between 1920-1950 and 1980-2010 ($0.06^{\circ}\text{C yr}^{-1}$) for a time-dependent temperature profile, the mismatch between modelled and measured temperatures is strongly reduced (yellow curve in Fig. 11). The above analysis shows clearly that the 135 largest contributor to the ice temperature increase is the strong increase in r_{rem} . Figure 11 also demonstrates the influence of the uncertainty of the average geothermal heat flux on the obtained vertical temperature profile. It is clear that the use of the initial average geothermal heat flux of 50 mW m^{-2} results in modelled ice temperatures matching most closely to the measured temperatures.


heeft verwijderd: ice

heeft verwijderd: also demonstrates



140
Figure 11: Measured (2007) and modelled (T_{mod}) temperature profile at the summit of the Grigoriev ice cap for a fixed geometry. The steady state temperature profile is modelled for the 1920-1950, 1960-1990, and the 1980-2010 mean climatic conditions. A time-dependent temperature profile is obtained for 1980-2010 by increasing the surface temperature from 1920-1950 with $0.06^{\circ}\text{C yr}^{-1}$. The purple profile is a time-dependent temperature profile obtained by using 50% of the average geothermal while the green dashed line corresponds to the time-dependent temperature profile obtained by using 200% of the average geothermal heat flux.

heeft verwijderd:
Met opmaak: Centrum



5 Sensitivity study

1150 5.1 Uncertainty and sensitivity to parameters

The sensitivity experiments bring to light that the fractional amount of temperate ice varies between 35–42% for the Sary-Tor glacier, and between 0 and 2% for the Grigoriev ice cap depending on the choice of the parameters associated with the firm warming, the insulation of snow, and the magnitude of the average geothermal heat flux (Table 4). Hence, the fractional amount of temperate ice can vary significantly, but in every case, there exists a substantial amount of temperate ice for the Sary-Tor glacier and very little to none for the Grigoriev ice cap. Further, and in general, we find a lower sensitivity of the results for the Grigoriev ice cap, which can mainly be attributed to the lower amounts of s_{max} and t_{ref} for the ice cap and its lower average temperature. For both ice bodies, the average ice temperature changes only slightly when the parameters and the average geothermal heat flux are altered within the predefined ranges (Table 4). Hence, based on the sensitivity analysis, it can be concluded that for the selected ice masses, parameter uncertainty and the magnitude of the geothermal heat flux do not have a major influence on the obtained thermal regime, making the results of this study robust.

Other assumptions and parameters that influence the results of this study concern the air temperature lapse rates (based on Aizen et al. (1995)), the precipitation gradients, the P_{max} value, and the timing of the s_{max} . The impact of the selected lapse rates is likely small because its uncertainty is implicitly taken into account in the calibration of the w_f and i_s parameter (through obtaining T_{ms}). Concerning the precipitation gradient, a lower gradient (compared to the Sary-Tor glacier, this differs by 100 mm at 4600 m altitude) proved to be necessary to obtain a sufficiently low accumulation on the summit of the Grigoriev ice cap. In the case of a larger precipitation gradient, a similar one as for the Sary-Tor glacier, s_{max} and t_{ref} would increase. Using these higher values would result in a slightly lower value of w_f and i_s . Nevertheless, even with the use of a lower w_f and i_s value, a polythermal regime is obtained for the Sary-Tor glacier and a cold regime for the Grigoriev ice cap (Table 4). As such, the precipitation gradient also has a small influence on the results achieved. Concerning the refreezing of meltwater, which appeared to be crucial for the determination of the surface layer temperature, it must be emphasised that in this study, a simple parametrisation is used with a constant P_{max} value (Reeh, 1991; Wright et al., 2007), to model its warming effect on the surface layer (Huybrechts et al., 1991; Zekollari et al., 2017). Other, more complex, approaches and models exist (Wright et al., 2007; Reijmer et al., 2012). Nevertheless, in this study, direct and multiple temperature observations in boreholes were used to calibrate the parameters of the applied parameterisation, which means that the effects of the uncertainties of the parametrisation on the results are expected to be limited. Regarding s_{max} and the snow insulation method, the largest uncertainty arises from the date of the maximum snow depth. In some years, the maximum snow depth is only reached after the end of May, for example, when substantial snow accumulates during the beginning of the summer season, or when the ablation season initiates later. In addition, significant accumulation of snow in autumn/winter is sometimes delayed, causing the snow to have a smaller insulating effect in this period (Hooke, 1983).

heeft verwijderd: 4

heeft verwijderd: 7

heeft verwijderd: 3

heeft verwijderd: 3

heeft verwijderd: 1

F

heeft verwijderd: is most sensitive to changes of the geothermal heat flux, while the fractional amount of temperate ice changes only slightly when the geothermal heat flux is reduced to 50% or increased to 200% ...

heeft verwijderd: 3

Met opmaak: mb0, Regelafstand: 1.5 regel, Patroon: Doorzichtig (Wit)

heeft verwijderd: ,

heeft verwijderd:) and

heeft verwijderd: .

heeft opmaak toegepast: Subscript

heeft verwijderd: .

heeft verwijderd: 3

heeft verwijderd: 1

1200 Other uncertainties that contribute to the results found in this study concern uncertainties in the mass balance measurements used for the calibration of c_0 , and the measured ice temperatures used for the calibration of w_f and i_s . However, the calibration of these parameters has been optimised by selecting all available independent borehole ice temperature and mass balance measurements. Due to the large amount of data available in time and space (see Table 2 and Table 3), the impact of inaccurate measurements is minimised, and the values found are reliable. Finally, all experiments in this study are performed under constant mean 1960-1990 climatic conditions. However, in reality, warmer and colder years, drier and wetter years, alternate, and the equilibrium line moves up and down. The values of s_{max} and r_{rem} vary accordingly. These interannual variations ensure that the transitions in surface layer ice temperature are in reality more gradual than found in this study.

210 **Table 4: Change in percentage of temperate ice and average temperature of both ice masses based on parameter changes of $\pm 10\%$ for the firn warming (w_f) and the snow insulation (i_s), and 50% vs 200% of the average geothermal heat flux (g_{hf}). The values for the initial parameter values are also given.**

Parameter	New value	Grigoriev ice cap	Sary-Tor glacier
w_f	26.9 °C (m w.e.) ⁻¹	0% (-3.4°C)	35% (-1.8°C)
	45.1 °C (m w.e.) ⁻¹	2% (-2.8°C)	42% (-1.6°C)
i_s	19.8 °C (m w.e.) ⁻¹	0% (-3.4°C)	35% (-1.8°C)
	24.2 °C (m w.e.) ⁻¹	2% (-2.7°C)	41% (-1.5°C)
g_{hf}	25 mW m ⁻²	0% (-3.4°C)	37% (-1.7°C)
	100 mW m ⁻²	2% (-2.8°C)	38% (-1.6°C)

5.2 Thermal regime since the LIA and into the future

1215 A glacier/ice cap thermal regime responds to climate change because the surface boundary condition strongly depends on air temperature and precipitation and because the ice mass geometry adjusts itself. To assess the evolution of the thermal regime of the Grigoriev ice cap and the Sary-Tor glacier, an analysis is performed to examine the effect of changing climatic conditions on the thermal regime of both ice masses by altering the air temperatures (between -2 and +3°C) and precipitation (between -40 and +40%). This range was selected based on historical temperature and precipitation records at the Kumtor – Tien Shan 1220 weather station (Van Tricht et al., 2021b), and expected changes in the future. The retreat or expansion of both ice masses is considered by letting the geometry freely evolve until a new steady state is reached, in equilibrium with the adjusted climate.

For a temperature increase of 2-3°C, the Sary-Tor glacier retreats to high elevations with a remaining volume which becomes significantly smaller. When the amount of precipitation remains constant or increases only slightly, which is suggested by 1225 most climate models (Sorg et al., 2012; Shahgedanova et al., 2020), the fractional amount of temperate ice becomes smaller (Fig. 12a). However, when the temperature increase is accompanied by substantial increase in precipitation (+40%), slightly

- heeft verwijderd: for the Grigoriev ice cap... and the m ... [11]
- heeft verwijderd: 3... Change in average temperature ... [12]
- heeft opmaak toegepast ... [13]
- Met opmaak ... [14]
- heeft verwijderd: ¶ ... [15]
- heeft opmaak toegepast ... [16]
- heeft opmaak toegepast ... [17]
- heeft opmaak toegepast ... [18]
- heeft verwijderd: Reduction
- heeft verwijderd: ¶
- heeft verwijderd: Reduction
- heeft opmaak toegepast ... [19]
- heeft opmaak toegepast ... [20]
- heeft opmaak toegepast ... [24]
- heeft verwijderd: ¶
- heeft opmaak toegepast ... [21]
- heeft opmaak toegepast ... [28]
- heeft opmaak toegepast ... [29]
- heeft opmaak toegepast ... [30]
- heeft opmaak toegepast ... [31]
- heeft opmaak toegepast ... [32]
- heeft opmaak toegepast ... [33]
- heeft verwijderd: 1
- heeft opmaak toegepast ... [25]
- heeft opmaak toegepast ... [34]
- heeft opmaak toegepast ... [35]
- heeft opmaak toegepast ... [36]
- Met opmaak ... [22]
- heeft verwijderd: ¶
- heeft verwijderd: 7...C¶ ... [27]
- heeft opmaak toegepast ... [23]
- Met opmaak ... [26]
- heeft opmaak toegepast ... [37]
- heeft opmaak toegepast ... [38]
- heeft opmaak toegepast ... [39]
- heeft opmaak toegepast ... [40]
- heeft opmaak toegepast ... [41]
- heeft opmaak toegepast ... [42]
- heeft opmaak toegepast ... [43]

more ice remains, and the ratio becomes larger again. This can be attributed to increased snow insulation and a thicker snow layer in which meltwater can refreeze. For colder climates, the Sary-Tor glacier would most likely have been polythermal as well (Fig. 12a). Only for a very strong decrease in temperature (-2°C) and a strong decrease in precipitation (-40%) does the Sary-Tor glacier become significantly colder with a fractional amount of temperate ice smaller than 5%. However, this climate configuration is rather unlikely as was shown by the climate reconstruction in Van Tricht et al. (2021b). Therefore, it can be concluded that the Sary-Tor glacier can always be considered a polythermal glacier since the LIA and into the future, until it disappears as soon as temperatures rise too much.

heeft verwijderd: be retained

Concerning the Grigoriev ice cap, the fractional amount of temperate ice remains low in all scenarios except for a strong increase in precipitation (Fig. 12b). For instance, increasing the amount of precipitation with 40% while keeping the temperatures fixed at the 1960-1990 average results in a transition from a cold to a polythermal ice cap. This can be explained by a larger insulation by snow and a larger amount of f_{refr} . For increasing temperatures, the Grigoriev ice cap retreats strongly (it disappears when temperatures rise between 2-3°C for equal precipitation) and it remains cold. For lower temperatures, the Grigoriev ice cap expands, but it remains cold as well. Hence, the Grigoriev ice cap can be regarded as a cold ice cap over time.▼

heeft verwijderd: formation

heeft verwijderd: ↗




Figure 12: Fractional amount of temperate ice for different combinations of temperature and precipitation changes **for the Sary-Tor glacier (panel a)** and the Grigoriev ice cap (panel b). For combinations in the red area, the ice mass disappears completely.

As demonstrated, ice temperatures can vary substantially over time. Ice temperatures at higher altitudes, where melting is currently limited by the colder climate, rise stronger than air temperatures, because of the enhanced formation of refrozen water (and the associated release of latent heat) (see section 4.4.2). Such a firm warming has also been observed and modelled at other glaciers (Hoelzle et al., 2011; Vincent et al., 2020; Mattea et al., 2021). Further, most climate models indicate a slight increase in winter precipitation (Sorg et al., 2012), which would imply an increase of s_{max} and thus a strengthening of the snow insulation effect. Evidence of a precipitation increase has been observed on Abramov glacier in the Pamir Alay (Kronenberg

heeft verwijderd:

Met opmaak: Uitvullen

heeft verwijderd: the thermal regime is not a constant and

heeft verwijderd: of superimposed ice

heeft verwijderd:

heeft opmaak toegepast: Lettertype: 10 pt

|310 **et al., 2021**) As a result, englacial temperatures generally rise, and part of the glaciers or ice caps in the Inner Tien Shan can undergo a transition from a cold to a polythermal or from a polythermal to a temperate regime (Arkhipov et al., 2004; Gusmeroli et al., 2012). Such a transition enhances ice flow and can fundamentally alter the ice dynamics (Marshall, 2021).

heeft verwijderd: This has already been observed at the Abramov glacier in the southwestern Tien Shan (Kronenberg et al., 2020)

6 Conclusion

In this study, we investigated the thermal regime of the Grigoriev ice cap and the Sary-Tor glacier, both located in the Inner
1315 Tien Shan in Kyrgyzstan (Central Asia), using a time-dependent 2-dimensional surface energy mass balance model coupled to a 3-dimensional higher-order thermomechanical ice flow model. Both models were calibrated with mass balance observations and temperature measurements for different periods and elevations. After calibration with the time-dependent output from the mass balance model, a firm warming parameter of $41^{\circ}\text{C} (\text{m w.e.})^{-1} \text{yr}^{-1}$ and a snow insulation parameter of $22^{\circ}\text{C} (\text{m w.e.})^{-1} \text{yr}^{-1}$ was found. Using these parameters, the ice temperature at 10-15 m depth of the Grigoriev ice cap could
1320 be modelled with a root mean square error of 0.28°C at locations with observations. The calibrated parameters were then used to model the thermal regime of both ice masses for constant 1960-1990 climatic conditions.

The simulations revealed a polythermal structure of the Sary-Tor glacier and a cold structure of the Grigoriev ice cap, apart from a thin layer of temperate ice near the bed in the central part. The difference can be attributed to a combination of a smaller
|325 amount of maximum snow depth at the end of winter, **reduced refreezing of meltwater**, as well as lower ice flow velocities and thinner ice of the Grigoriev ice cap. As a result, the Sary-Tor glacier is characterised by basal sliding and temperate ice in the upper layers where **in water can infiltrate through intragranular veins**, while this is not the case for the Grigoriev ice cap. An extensive sensitivity analysis was performed to analyse the thermal structure for combinations of historical and future air temperature and precipitation variations. This demonstrated that the Sary-Tor glacier can always be considered a polythermal
1330 glacier since the LIA and into the future, while the Grigoriev ice cap has always been and will always be characterised by a cold structure, despite the ice being **warmed in the higher parts of the ice cap when air temperatures increase**. The reason for this is that at these higher temperatures, the ice cap will disappear faster than it warms up to a temperate state.

heeft verwijderd: reduced formation of refrozen meltwater at the end of summer

heeft opmaak toegepast: Lettertype: 10 pt

heeft verwijderd: e

heeft verwijderd: is able to

heeft verwijderd: significantly

Because the selected ice masses are typical examples of ice bodies in the Inner Tien Shan, and because a detailed sensitivity
1335 analysis revealed robust results to changes in the parameters and the magnitude of the geothermal heat flux, it is reasonable to conclude that the results of this study can be generalised for similar type of glaciers and ice caps in the study area. Valley glaciers in the Inner Tien Shan are probably almost all polythermal, while the high-altitude ice caps are expected to be mainly cold. Glaciers in the outer ranges, which are associated to larger amounts of precipitation might be temperate as well. These findings are important as the dynamics of the ice masses can only be understood and modelled precisely if ice temperature is
1340 considered correctly in ice flow models. The calibrated parameters of this study can be used in applications with ice flow models for individual ice masses as well as to optimise more general models for large-scale regional simulations.

1350 Data availability

Research data is provided through an online public repository, accessible via <https://www.doi.org/10.5281/zenodo.6556313>.
(Van Tricht et al., 2022). Information and specific details model code will be specified on request by Lander Van Tricht.

Gewijzigde veldcode

Author contributions

LVT developed the method, performed the experiments and wrote the manuscript. PH provided the model code, and gave
1355 guidance in implementing the research and interpreting the results, assisting during the entire process.

Competing interests

The authors have declared that they do not have any competing interests.

Disclaimer

/

1360 Acknowledgements

The authors would like to thank everyone who contributed to the fieldwork to carry out the ice thickness measurements. We would also like to specifically thank Benjamin Vanbiervliet for processing the RES data and Oleg Rybak and Rysbek Satylkanov for assisting and organising the fieldwork. We also thank Ivan Lavrentiev for the GPR data from the Sary-Tor glacier that was used to distinguish between temperate and cold ice. Finally, we thank Victor Popovnin for providing the mass
1365 balance data of the Sary-Tor glacier.

Financial support

Lander Van Tricht holds a PhD fellowship of the Research Foundation – Flanders (FWO-Vlaanderen) and is affiliated with the Vrije Universiteit Brussel (VUB). [Local logistics were mainly organised and funded by the Tien Shan High Mountain Research Center and the Kumtor mining company.](#)

1370 References

Aschwanden, A., Bueler, E., Khroulev, C. and Blatter, H.: An enthalpy formulation for glaciers and ice sheets. *Journal of Glaciology*, 58(209), 441–457, <https://www.doi.org/10.3189/2012JoG11J088>, 2012

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië)

Aschwanden, A., Bueler, E., Khroulev, C. and Blatter, H.: An enthalpy formulation for glaciers and ice sheets. *Journal of Glaciology*, 58(209), 441-457, <https://www.doi.org/10.3189/2012JoG11J088>, 2012

1375

Aizen, V.B., Aizen, E.M., and Melack, J.M.: Climate, Snow Cover, Glaciers, and Runoff in the Tien Shan, Central Asia. *Journal of the American Water Resources Association*, 31(6), 1113–1129, <https://www.doi.org/10.1111/j.1752-1688.1995.tb03426.x>, 1995

1380

Aizen, V.B., Kuzmichenok, V., Surazakov, A. and Aizen, E.M.: Glacier changes in the central and northern Tien Shan during the last 140 years based on surface and remote-sensing data. *Annals of Glaciology*, 43(1), 202-213. <https://www.doi.org/10.3189/172756406781812465>, 2006

1385

Arkhipov, S.M., Mikhaleko, V.N., Kunakhovich, M.G., Dikikh, A.N., and Nagornov, O.V.: Termicheskii rezhim, usloviia l'doobrazovaniia i akkumulyatsii na ladnike Grigor'eva (Tyan'-Shan') v 1962–2001 gg. [Thermal regime, ice types and accumulation in Grigoriev Glacier, Tien Shan, 1962–2001]. Mater. Glyatsiol. Issled. 96, 77–83, 2004 (in Russian with English summary)

1390

Bagdassarov, N., Batalev, V. and Egorova, V.: State of lithosphere beneath Tien Shan from petrology and electrical conductivity of xenoliths. *Journal of Geophysical Research: Solid Earth*, 116(B1), <https://www.doi.org/10.1029/2009JB007125>, 2011

1395

Barandun, M., Huss, M., Sold, L., Farinotti, D., Azisov, E., Salzmann, N., Usualiev, R., Merkushkin, A. and Hoelzle, M.: Re-analysis of seasonal mass balance at Abramov glacier 1968–2014. *Journal of Glaciology*, 61(230), 1103–1117, <https://www.doi.org/10.3189/2015JoG14J239>, 2015

1400

Bhattacharya, A., Bolch, T., Mukherjee, K., King O., Menounos B., Kapitsa V., Neckel, N., Yang W. and Yao T.: High Mountain Asian glacier response to climate revealed by multi-temporal satellite observations since the 1960s. *Nature Communications*, 12, 4133, <https://www.doi.org/10.1038/s41467-021-24180-y>, 2021

Blatter, H.: On the Thermal Regime of an Arctic Valley Glacier: A Study of White Glacier, Axel Heiberg Island, N.W.T., Canada. *Journal of Glaciology*, 33(114), 200-211, <https://www.doi.org/10.3189/S002214300008704>, 1987

Gewijzigde veldcode

1405

Blatter, H. and Hutter K.: Polythermal conditions in Arctic glaciers. *Journal of Glaciology*, 37(126), 261–269, <https://www.doi.org/10.3189/S002214300007279>, 1991

Bondarev, L.G.: Evolution of some Tien Shan glaciers during the last quarter of the century. IAHS Publication , 54: 412–419, 1961

410

Boon, S. and Sharp, M.: The role of hydrologically-driven ice fracture in drainage system evolution on an arctic glacier, Geophysical Research Letters, 30(18), 1916, <https://www.doi.org/10.1029/2003GL018034>, 2003

heeft opmaak toegepast: Standaardlinea-lettertype, Lettertype: (Standaard) Times New Roman, Spelling en grammatica controleren

415 Braithwaite, R.J., Laternser, M. and Pfeffer, W.T.: Variations of near-surface firm density in the lower accumulation area of the Greenland ice sheet, Pakitsoq, West Greenland, Journal of Glaciology, 40 (136), 477–485, 1994

Cai, B., Huang M. and Zichu X.: A preliminary research on the temperature in deep boreholes of Glacier No. 1, Ürümqi headwaters. Kexue Tongbao (Science Bulletin), 33(24), 2054–2056, 1988 (in Chinese)

420 Colgan, W., Sommers, A., Rajaram, H., Abdalati, W. and Frahm, J.: Considering thermal-viscous collapse of the Greenland ice sheet. Earth's Future, 3(7), 252–267, <https://www.doi.org/10.1002/2015EF000301>, 2015

Colgan, W., MacGregor, J.A., Mankoff, K.D., Haagenson, R., Rajaram, H., Martos, Y.M., Morlighem, M., Fahnestock, M.A. and Kjeldsen, K.K.: Topographic correction of geothermal heat flux in Greenland and Antarctica, Journal of Geophysical Research: Earth Surface, 126, <https://www.doi.org/10.1029/2020JF005598>, 2021

Met opmaak: Links

Gewijzigde veldcode

heeft verwijderd: ↪

Delf, R., Bingham, R.G., Curtis, A., Singh, S., Giannopoulos, A., Schwarz, B. and Borstad, C.P.: Reanalysis of polythermal glacier thermal structure using radar diffraction focussing. Journal of Geophysical Research: Earth Surface, 127, 2, <https://doi.org/10.1029/2021JF006382>, 2022

430 Delvaux, D., Cloetingh, S., Beekman, F., Sokoutis, D., Burov, E., Buslov, M.M. and Abdurakhmatov, K.E.: Basin evolution in a folding lithosphere: Altai-Sayan and Tien Shan belts in Central Asia. Tectonophysics, 602, 194–222, <https://www.doi.org/10.1016/j.tecto.2013.01.010>, 2013

Dikikh, A.N.: Temperature regime of flat-top glaciers (using Grigoriev as an Example). - Glyatsiol. Issledovaniya na Tyan-Shane, Frunze , N.11, p.32-35, 1965 (in Russian)

heeft verwijderd:

Duchkov, A.D., Shvartsman Yu.G. and Sokolova L.S.: Deep heat flow in Tien Shan: advances and drawbacks, Geologiya I Geofizika (Russian Geology and Geophysics), 42, 10, 1516–1531(1436–1452), 2001 (in Russian)

Dyurgerov, M.B. and Mikhalenko, V.N.: *Glaciation of Tien Shan* (M.B. Dyurgerov and V.N. Mikhalenko, Eds.), VINITI, 1995 (in Russian)

- 450 Dyurgerov, M.B.: *Glacier Mass Balance and Regime: Data of Measurements and Analysis*. Occasional Paper 55, Boulder, CO: Institute of Arctic and Alpine Research, University of Colorado, 2002

heeft verwijderd: 

- Echelmeyer, K. and Zhongxiang, W.: Direct Observation of Basal Sliding and Deformation of Basal Drift at Sub-Freezing Temperatures. *Journal of Glaciology*, 33(113), 83–98,
1455 <https://www.doi.org/10.3189/S0022143000005396>, 1987

Gewijzigde veldcode

- Farinotti, D., Huss, M., Fürst, J.J., Landmann, J., Machguth, H., Maussion, F. and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth. *Nature Geoscience*, 12, 168–173, <https://www.doi.org/10.1038/s41561-019-0300-3>, 2019

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië)

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië)

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië)

heeft verwijderd: 

- Flowers, G.E., Björnsson, H., Geirsdóttir, Á., Miller, G.H. and Clarke, G.K.C.: *Glacier fluctuation and inferred climatology of Langjökull ice cap through the Little Ice Age*. *Quaternary Science Reviews*, 26(19–21), 2337–2353, <https://www.doi.org/10.1016/j.quascirev.2007.07.016>, 2007
- Fujita, K., Takeuchi, N., Nikitin, S.A., Surazakov, A.B., Okamoto, S., Aizen, V.B. and Kubota, J.: Favorable climatic regime for maintaining the present-day geometry of the Grigoriev Glacier, Inner Tien Shan, The Cryosphere, 5, 539–549, <https://www.doi.org/10.5194/tc-5-539-2011>, 2011

- 1470 Fürst, J.J., Rybak, O., Goelzer, H., De Smedt, B., de Groen, P. and Huybrechts, P.: Improved convergence and stability properties in a three-dimensional higher-order ice sheet model. *Geoscientific Model Development*, 4(4), 1133–1149, <https://www.doi.org/10.5194/gmd-4-1133-2011>, 2011

heeft verwijderd: 

Gewijzigde veldcode

- Fürst, J.J., Goelzer, H. and Huybrechts, P.: Effect of higher-order stress gradients on the centennial mass evolution of the Greenland ice sheet. *The Cryosphere*, 7, 183–199, <https://www.doi.org/10.5194/tc-7-183-2013>, 2013

heeft verwijderd: 

Gewijzigde veldcode

- 1475 Fürst, J.J., Goelzer, H. and Huybrechts, P.: Ice-dynamic projections of the Greenland ice sheet in response to atmospheric and oceanic warming. *The Cryosphere*, 9, 1039–1062, <https://www.doi.org/10.5194/tc-9-1039-2015>, 2015

- 485 Gilbert, A., Flowers, G. E., Miller, G. H., Refsnider, K., Young, N. E. and Radic, V.: The projected demise of Barnes Ice Cap: Evidence of an unusually warm 21st century Arctic. *Geophysical Research Letters*, 44(6), 2810–2816. <https://www.doi.org/10.1002/2016GL072394>, 2017.
- 490 Gilbert, A., Sinisalo, A., Gurung, T.R., Fujita, K., Maharjan, S.B., Sherpa, T.C. and Fukuda, T.: The influence of water percolation through crevasses on the thermal regime of a Himalayan mountain glacier. *The Cryosphere*, 14, 1273–1288, <https://www.doi.org/10.5194/tc-14-1273-2020>, 2020
- 495 Gusmeroli, A., Jansson P., Pettersson R. and Murray T.: Twenty years of cold surface layer thinning at Storglaciären, sub-Arctic Sweden, 1989–2009. *Journal of Glaciology*, 58(207), 3–10, <https://www.doi.org/10.3189/2012JoG11J018>, 2012
- 500 Hambrey, M.J. and Glasser, N.F.: Discriminating glacier thermal and dynamic regimes in the sedimentary record. *Sedimentary Geology*, 251–252(0): p.1-33, 2012
- 505 Hoelzle, M., Darms, G., Lüthi, M. P. and Suter, S.: Evidence of accelerated englacial warming in the Monte Rosa area, Switzerland/Italy. *The Cryosphere*, 5, 231–243, <https://www.doi.org/10.5194/tc-5-231-2011>, 2011
- 510 Hooke, R. L.: Pleistocene ice at the base of the Barnes Ice Cap, Baffin Island, N.W.T., Canada. *Journal of Glaciology*, Vol. 17, No. 75, p.49–59, <https://www.doi.org/10.3189/S0022143000030719>, 1976.
- 515 Hooke, R. LeB., Gould, J.E. and Brzozowski, J.: Near-surface temperatures near and below the equilibrium line on polar and subpolar glaciers. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 19(1), 1–25, 1983
- 520 Huybrechts, P. and Oerlemans, J.: Evolution of the East Antarctic Ice Sheet: A Numerical Study of Thermo-Mechanical Response Patterns With Changing Climate. *Annals of Glaciology*, 11, 52-59. <https://www.doi.org/10.3189/S026030550006327>, 1988
- 525 Huybrechts, P., Letreguilly, A. and Reeh, N.: The Greenland ice sheet and greenhouse warming. *Global And Planetary Change*, 3(4), 399–412, [https://www.doi.org/10.1016/0921-8181\(91\)90119-H](https://www.doi.org/10.1016/0921-8181(91)90119-H), 1991
- 530 Huybrechts, P.: Basal temperature conditions of the Greenland ice sheet during the glacial cycles. *Annals of Glaciology* 23, 226-236, <https://www.doi.org/10.3189/S0260305500013483>, 1996.
- 535 Jouvet, G., Huss, M., Funk, M. and Blatter, H.: Modelling the retreat of Grosser Aletschgletscher,
- heeft opmaak toegepast:** Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren
- heeft opmaak toegepast:** Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren
- heeft opmaak toegepast:** Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren
- Met opmaak:** Regelafstand: 1.5 regel
- heeft opmaak toegepast:** Hyperlink, Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren
- heeft opmaak toegepast:** Hyperlink, Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren
- heeft opmaak toegepast:** Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren
- Gewijzigde veldcode**
- heeft verwijderd:** 1
- Gewijzigde veldcode**
- heeft opmaak toegepast:** Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren, Rand: : (Geen rand), Patroon: Doorzichtig
- heeft opmaak toegepast:** Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren, Patroon: Doorzichtig
- Met opmaak:** Regelafstand: 1.5 regel
- heeft opmaak toegepast:** Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren, Patroon: Doorzichtig
- heeft opmaak toegepast:** Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren
- heeft verwijderd:** 1
- heeft verwijderd:** 1
- Met opmaak:** Inspriking: Links: 0 cm, Verkeerd-om: 0.85 cm, Zwenende begin- en eindregels niet voorkomen, Spatiëring tussen Aziatische en Latijnse tekst niet aanpassen, Spatiëring tussen Aziatische tekst en nummers niet aanpassen

Switzerland, in a changing climate. *Journal of Glaciology*, 57(206), 1033–1045,
<https://www.doi.org/10.3189/00221431179843359>, 2011

- 525 Kislov, B.v., Nozdrukhin, V.K. and Pertziger, F.I.: Temperature regime of the active layer of Abramov Glacier. Materialy
Glaciologicheskikh Issledovanii (Data of Glaciological Studies), 30, 199–204, 1977 (in Russian)

heeft opmaak toegepast: Engels (V.S.)

- Kronenberg, M., Barandun, M., Hoelzle, M., Huss, M., Farinotti, D., Azisov, E., Usualiev, R., Gafurov, A., Petrakov, D. and
Kääb, A.: Mass-balance reconstruction for Glacier No. 354, Tien Shan, from 2003 to 2014. *Annals of Glaciology*, 57(71), 92–
530 102, <https://www.doi.org/10.3189/2016AoG71A032>, 2016

heeft opmaak toegepast: Engels (V.S.)

Gewijzigde veldcode

heeft verwijderd: 1

heeft verwijderd: 0

- Kronenberg, M., Machguth, H., Eichler, A., Schwikowski, M. and Hoelzle, M.: Comparison of historical and
recent accumulation rates on Abramov Glacier, Pamir Alay. *Journal of Glaciology*, 67(262), 253–268,
<https://www.doi.org/10.1017/jog.2020.103>, 2021

heeft verwijderd: .

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

Met opmaak: Regelstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

Gewijzigde veldcode

heeft verwijderd: 1

Liang, L. and Y. Zhou: The characteristics of the distribution of snow
cover and its warm effect on the temperature of permafrost, Amur
Region, Da Hinggan Ling. In Proceedings of the 6th International
Conference on Permafrost, Beijing, China, July 5–9, 1993, vol. 1,
pp. 393–396, S. China Univ. of Technol. Press, Guangzhou, China,
1993

- 1535 Kutuzov, S. and Shahgedanova, M.: Glacier retreat and climatic variability in the eastern Terskey-Alatau, inner Tien Shan
between the middle of the 19th century and beginning of the 21st century. *Global and Planetary Change*, 69, 59–70,
<https://www.doi.org/10.1016/j.gloplacha.2009.07.001>, 2009

- 540 Libloutry, L.: Temperate ice permeability, stability of water veins and percolation of internal meltwater. *Journal of
Glaciology*, 42(141), 201–211, <https://www.doi.org/10.3189/S0022143000004068>, 1996

heeft verwijderd: .

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

Met opmaak: Regelstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

Gewijzigde veldcode

heeft verwijderd: 1

Li, Y., Tian, L., Yi, Y., Moore, J. C., Sun, S. and Zhao, L.: Simulating the evolution of Qiangtang No. 1 glacier in the central
Tibetan Plateau to 2050. *Arctic Antarctic and Alpine Research*, 49(1), 1–12, <https://www.doi.org/10.1657/aaar0016-008>,
2017

- 545 Lüthi, M.P., Ryser, C., Andrews, L.C., Catania, G.A., Funk, M., Hawley, R.L., Hoffman, M.J. and Neumann, T.A.:
Heat sources within the Greenland Ice Sheet: dissipation, temperate paleo-firn and cryo-hydrologic warming.
The Cryosphere, 9, 245–253, <https://www.doi.org/10.5194/tc-9-245-2015>, 2015

heeft verwijderd: .

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

Met opmaak: Regelstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Engels
(Groot-Brittannië), Geen spelling- of grammaticacontrole
uitvoeren

Gewijzigde veldcode

heeft verwijderd: 1

Maohtuan, H., Zhongxiang, W. and Jiawen, R.: On the Temperature Regime of Continental-Type Glaciers in
1555 China. *Journal of Glaciology*, 28(98), 117–128, <https://www.doi.org/10.3189/S0022143000011837>, 1982

Maohuan, H., Zhongxiang, W., Baolin, C. and Jiankang, H.: Some Dynamics Studies on Urumqi Glacier No. 1, Tianshan Glaciological Station, China. Annals of Glaciology, 12, 70-73, <https://www.doi.org/10.3189/S0260305500006972>, 1989

1570

Maohuan, H.: On the Temperature Distribution of Glaciers in China. Journal of Glaciology, 36(123), 210–216, <https://www.doi.org/10.3189/S002214300000945X>, 1990

Maohuan, H.: The movement mechanisms of Ürümqi Glacier No. 1, Tien Shan mountains, China. Annals of Glaciology, 16(1), 39–44, <https://www.doi.org/10.3189/1992AoG16-1-39-44>, 1992

1575

Mattea, E., Machguth, H., Kronenberg, M., van Pelt, W., Bassi, M. and Hoelzle, M.: Firn changes at Colle Gnifetti revealed with a high-resolution process-based physical model approach. The Cryosphere, 15, 3181–3205, <https://www.doi.org/10.5194/tc-15-3181-2021>, 2021

1580

Marshall, S.J.: Regime Shifts in Glacier and Ice Sheet Response to Climate Change: Examples From the Northern Hemisphere. Frontiers in Climate, 3, <https://www.doi.org/10.3389/fclim.2021.702585>, 2021

Meierbachtol, T.W., Harper J. T., Johnson J. V., Humphrey N. F. and Brinkerhoff D. J.: Thermal boundary conditions on western Greenland: Observational constraints and impacts on the modeled thermomechanical state. Journal of Geophysical Research Earth Surface, 120(3), 623-636, <https://www.doi.org/10.1002/2014JF003375>, 2015

Mikhailenko, V.N.: Osobennosti massoobmena lednikov ploskikh vershin vnutrennego Tyan'-Shanya [Peculiarities of the mass exchange of flat summit glaciers of interior Tyan'-Shan']. Materialy Glyatsiologicheskikh

1590

Issledovaniy. 65, 86–92, 1989.(in Russian)

Mikhailenko, V.N.: Calculation and modeling of the mass balance of the Aj-Shyirak massif changes in the Tien Shan.

Materialy Glyatsiologicheskikh Issledovaniy, 76, 102–107, 1993.(in Russian)

1595 Mukherjee K., Bolch T., Goerlich G., Kutuzov S., Osmonov A., Pieczonka T. and Shesterova I.: Surge-Type Glaciers in the Tien Shan (Central Asia). Arctic, Antarctic, and Alpine Research, 49(1), 147-171, <https://www.doi.org/10.1657/AAAR0016-021>, 2017

Nagornov, O., Konovalov, Y. and Mikhailenko, V.: Prediction of thermodynamic state of the

Met opmaak: Zwevende begin- en eindregels voorkomen, Spatiëring tussen Aziaatse en Latijnse tekst aanpassen, Spatiëring tussen Aziaatse tekst en nummers aanpassen

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), Spelling en grammatica controleren

Gewijzigde veldcode

heeft verwijderd: ,

heeft verwijderd: ,

Grigoriev ice cap, Tien Shan, central Asia, in the future. Annals of Glaciology, 43, 307-312,
<https://www.doi.org/10.3189/172756406781812221>, 2006

- 1605 Nosenko, G.A., Lavrentiev, I.I., Glazovskii, A.F., Kazatkin, N.E. and Kokarev, A.L.: The polythermal structure of Central
Tuyuksu glacier. Криосфера Земли 20(4), 105–115, [https://www.doi.org/10.21782/KZ1560-7496-2016-4\(105-115\)](https://www.doi.org/10.21782/KZ1560-7496-2016-4(105-115)), 2016

Nye, J. F. and Frank, F.C.: Hydrology of the intergranular veins in a temperate glacier, in Symposium on the Hydrology of
Glaciers, IAHS Publ. 95., 157–161, 1973

- 1610 Osmonov, A., Bolch, T., Xi, C., Kurban, A. and Guo, W.: Glacier characteristics and changes in the Sary-Jaz River
Basin (Central Tien Shan, Kyrgyzstan) – 1990–2010, Remote Sensing Letters, 4(8), 725–734,
<https://www.doi.org/10.1080/2150704X.2013.789146>, 2013.

- 615 Pattyn, F.: A new three-dimensional higher-order thermomechanical ice sheet model: Basic sensitivity, ice stream
development, and iceflow across subglacial lakes. Journal of Geophysical Research, 108(B8), B82382,
<https://www.doi.org/10.1029/2002JB002329>, 2003

- Petrakov, D.A., Lavrentiev, I.I., Kovalenko, N.V. and Usualiev, R.A.: Ice thickness, volume and modern change of the
1620 Sary-Tor Glacier area (Ak-Shiyirak Massif, Inner Tian Shan). Earth's Cryosphere, 18(3), 91–100, 2014

Phillips, T., Rajaram, H. and Steffen, K.: Cryo-hydrologic warming: A potential mechanism for rapid thermal response of ice
sheets, Geophysical Research Letters, 37(20), <https://www.doi.org/10.1029/2010GL044397>, 2010

- 1625 Reeh, N.: Parameterization of Melt Rate and Surface Temperature in the Greenland Ice Sheet, Polarforschung, Bremerhaven,
Alfred Wegener Institute for Polar and Marine Research & German Society of Polar Research, 59 (3), pp. 113-128, 1991

Reijmer, C. H., van den Broeke, M. R., Fettweis, X., Ettema, J. and Stap, L.B.: Refreezing on the Greenland ice sheet: a
comparison of parameterizations, The Cryosphere, 6, 743–762, <https://www.doi.org/10.5194/tc-6-743-2012>, 2012.

- 630 RGI Consortium: Randolph Glacier Inventory – A Dataset of Global Glacier Outlines: Version 6.0: Technical Report, Global
Land Ice Measurements from Space. Colorado, USA: Digital Media, <https://www.doi.org/10.7265/N5-RGI-60>, 2017

heeft opmaak toegepast: Letertype: (Standaard) +Koppen
(Times New Roman), 10 pt, Tekstkleur: Tekst 1, Patroon:
Doorzichtig (Wit)

heeft opmaak toegepast: Letertype: (Standaard) +Koppen
(Times New Roman), 10 pt, Tekstkleur: Tekst 1, Patroon:
Doorzichtig (Wit)

heeft opmaak toegepast: Letertype: (Standaard) +Koppen
(Times New Roman), 10 pt, Niet Vet, Tekstkleur: Blauw,
Patroon: Doorzichtig (Wit)

heeft opmaak toegepast: Letertype: (Standaard) +Koppen
(Times New Roman), 10 pt, Niet Vet, Tekstkleur: Blauw,
Patroon: Doorzichtig (Wit)

Gewijzigde veldcode

heeft verwijderd: 1

heeft opmaak toegepast: Letertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1,
Engels (Groot-Brittannië)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Letertype: (Standaard)
+Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1,
Engels (Groot-Brittannië)

heeft verwijderd: 1

Riesen, P., Sugiyama, S. and Funk, M.: The influence of the presence and drainage of an ice-marginal lake on the ice flow of Gornergletscher, Switzerland. *Journal of Glaciology*, 56(196), 278–286, <https://www.doi.org/10.3189/002214310791968575>, 2010

640 Rowan, A.V., Egholm, D.L., Quincey, D.J. and Glasser, N.F.: Modelling the feedbacks between mass balance, ice flow and debris transport to predict the response to climate change of debris-covered glaciers in the Himalaya. *Earth and Planetary Science Letters*, 430, 427–438, <https://www.doi.org/10.1016/j.epsl.2015.09.004>, 2015

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Patroon: Doorzichtig (Wit)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (Groot-Brittannië), Patroon: Doorzichtig (Wit)

heeft verwijderd: ↵

Ryser, C., Lüthi, M., Blindow, N., Suckro, S., Funk, M. and Bauder, A.: Cold ice in the ablation zone: its relation to glacier hydrology and ice water content. *Journal of Geophysical Research*, 118(2), 693–705, <https://www.doi.org/10.1029/2012JF002526>, 2013.

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

heeft verwijderd: ↵

645 Schäfer, M., Möller, M., Zwinger, T. & Moore, J.C.: Dynamic modelling of future glacier changes: Mass-balance/elevation feedback in projections for the Vestfonna ice cap, Nordaustlandet, Svalbard. *Journal of Glaciology*, 61(230), 1121–1136, <https://www.doi.org/10.3189/2015JoG14J184>, 2015

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

heeft verwijderd: ↵

650 Shahgedanova, M., Afzal, M., Hagg, W., Kapitsa, V., Kasatkin, N., Mayr, E., Rybak, O., Saidaliyeva, Z., Severskiy, I., Usmanova, Z., Wade, A., Yaitskaya, N. and Zhumbabayev, D.: Emptying Water Towers? Impacts of Future Climate and Glacier Change on River Discharge in the Northern Tien Shan, Central Asia. *Water*, 12(3), 627, <https://www.doi.org/10.3390/w12030627>, 2020

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

heeft verwijderd: ↵

655 Shumskiy, P.A.: Osnovy strukturnogo ledovedeniya. Petrografiya presnogo l'da kak metod glyatsiologicheskogo issledovaniya. Moscow, Izdatel 'stvo Akademii Nauk SSSR. [English translation: Principles of structural glaciology: the petrography of fresh-water ice as a method of glaciological investigation Translated from the Russian by David Kraus. New York, Dover Publications, 1964]

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

heeft verwijderd: ↵

Sorg, A., Bolch, T., Stoffel, M., Solomina, O. and Beniston, M.: Climate change impacts on glaciers and runoff in Tien Shan (Central Asia). *Nature Climate Change* 2, 725–731, <https://www.doi.org/10.1038/nclimate1592>, 2012

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Tekstkleur: Tekst 1, Engels (V.S.), Patroon: Doorzichtig (Wit)

heeft verwijderd: ↵

1665 Takeuchi, N., Fujita, K., Aizen, V.B., Narama, C., Yokoyama, Y., Okamoto, S., Naoki, K. and Kubota, J.: The disappearance of glaciers in the Tien Shan Mountains in Central Asia at the end of Pleistocene. *Quaternary Science Revision*, 103, 26–33, <https://www.doi.org/10.1016/j.quascirev.2014.09.006>, 2014

heeft verwijderd:

Gewijzigde veldcode

Takeuchi, N., Sera, S., Fujita, K., Aizen, V.B., and Kubota, J.: Annual layer counting using pollen grains of the Grigoriev ice core from the Tien Shan Mountains, central Asia, Arctic Antarctic and Alpine Research, 51, 299–312, <https://www.doi.org/10.1080/15230430.2019.1638202>, 2019.

1675 Thompson, L.G., Mosley-Thompson, E., Davis, M., Lin, P.N., Yao, T., Dyurgerov, M. and Dai, J.: Recent warming: ice core evidence from tropical ice cores with emphasis on Central Asia. Global Planetary Change, 7(1–3), 145–156, [https://www.doi.org/10.1016/0921-8181\(93\)90046-Q](https://www.doi.org/10.1016/0921-8181(93)90046-Q), 1993

1680 van Pelt, W. J., Pohjola, V. A. and Reijmer, C. H.: The changing impact of snow conditions and refreezing on the mass balance of an idealized Svalbard glacier, Frontiers in Earth Science, 4, 102, <https://www.doi.org/10.3389/feart.2016.00102>, 2016

1685 Van Tricht, L., Huybrechts, P., Van Breedam, J., Fürst, J., Rybak, O., Satylkanov, R., Ermenbaiev B., Popovnin V., Neyns, R., Paice C.M. and Malz, P.: Measuring and inferring the ice thickness distribution of four glaciers in the Tien Shan, Kyrgyzstan. Journal of Glaciology, 67(262), 269–286. <https://www.doi.org/10.1017/jog.2020.104>, 2021a

Van Tricht L., Paice C.M., Rybak O., Satylkanov R., Popovnin V., Solomina O. and Huybrechts P.: Reconstruction of the Historical (1750–2020) Mass Balance of Bordu, Kara-Batkak and Sary-Tor Glaciers in the Inner Tien Shan, Kyrgyzstan. 1690 Frontiers in Earth Science, 9, <https://www.doi.org/10.3389/feart.2021.734802>, 2021b

Van Tricht, L., Huybrechts, P., Van Breedam, J., Vanhulle, A., Van Oost, K., and Zekollari, H.: Estimating surface mass balance patterns from unoccupied aerial vehicle measurements in the ablation area of the Morteratsch–Pers glacier complex (Switzerland), The Cryosphere, 15, 4445–4464, <https://www.doi.org/10.5194/tc-15-4445-2021>, 2021c.

Gewijzigde veldcode

1695 Vasilenko, E.V., Gromyko, A.N., Dmitriev, D.N. and Macheret, Ya Yu.: Stroenie lednika Davydova po dannym radiozonirovaniya i termobureniyu [Structure of the Davydov Glacier according with radio sounding and thermal drilling data]. Akademiya nauk SSSR. Institut geografii. Materialy gliatsiologicheskikh issledovanii [Academy of Sciences of the USSR. Institute of Geography. Data of Glaciological Studies]. Vol. 62, p. 208–215, 1988 (in Russian with English summary.)

700 Vermeesch, P., Poort, J., Duchkov, A., Klerkx, J. and De Batist, M.: Lake Issyk-Kul (Tien Shan): Unusually low heat flow in an active intermontane basin. GEOLOGIYA I GEOFIZIKA, 45(5), 616–625, 2004

Vilesov, E.: Temperature of ice in the lower parts of the Tuyuksu glaciers. Union Géodésique et Géophysique Internationale. Association Internationale d'Hydrologie Scientifique. Assemblée générale de Helsinki, 25-7-6-8 1960. Commission des Neiges et Glaces, p. 313–24. (Publication No. 54 de l'Association Internationale d'Hydrologie Scientifique.), 1961

- 710 Vincent, C., Le Meur, E., Six, D., Possenti, P., Lefebvre, E. and Funk, M.: Climate warming revealed by englacial temperatures at Col du Dôme (4250 m, Mont-Blanc area), *Geophysical Research Letters*, 34, L16502,

[https://www.doi.org/10.1029/2007GL029933, 2007](https://www.doi.org/10.1029/2007GL029933)

- 715 Vincent, C., Gilbert, A., Jourdain, B., Piard, L., Ginot, P., Mikhalenko, V., Possenti, P., le Meur, E., Laarman, O. and Six, D.: Strong changes in englacial temperatures despite insignificant changes in ice thickness at Dôme du Goûter glacier (Mont Blanc area). *The Cryosphere*, 14, 925–934, [https://www.doi.org/10.5194/tc-14-925-2020, 2020](https://www.doi.org/10.5194/tc-14-925-2020)

Wohlleben, T., Sharp, M. and Bush, A.: Factors influencing the basal temperatures of a High Arctic polythermal glacier. *Annals of Glaciology*, 50(52), 9–16, [https://www.doi.org/10.3189/172756409789624210, 2009](https://www.doi.org/10.3189/172756409789624210)

- 720 Wright, A.P., Wadhams, J.L., Siegert, M.J., Luckman, A., Kohler, J. and Nuttall, A.M.: Modeling the refreezing of meltwater as superimposed ice on a high Arctic glacier: A comparison of approaches. *Journal of Geophysical Research*, 112, F04016, [https://www.doi.org/10.1029/2007JF000818, 2007](https://www.doi.org/10.1029/2007JF000818)

- 725 Wu, Z., Zhang, H., Liu, S., Ren, D., Bai, X., Xun, Z. and Ma, Z.: Fluctuation analysis in the dynamic characteristics of continental glacier based on Full-Stokes model. *Scientific Reports*, 9(1), 20245, [https://www.doi.org/10.1038/s41598-019-56864-3, 2019](https://www.doi.org/10.1038/s41598-019-56864-3)

Yershov, E.D.: General Geocryology, Stud. Polar Res., 580 pp., Cambridge Univ. Press, New York, 1998

- 1730 Zekollari H, Huybrechts P, Fürst J.J., Rybak O. and Eisen O.: Calibration of a higher-order 3-D ice-flow model of the Morteratsch glacier complex, Engadin, Switzerland. *Annals of Glaciology*, 54(63), 343–351, [https://www.doi.org/10.3189/2013AoG63A434, 2013](https://www.doi.org/10.3189/2013AoG63A434)

- 1735 Zekollari, H. and Huybrechts, P.: On the climate–geometry imbalance, response time and volume–area scaling of an alpine glacier: insights from a 3-D flow model applied to Vadret da Morteratsch, Switzerland. *Annals of Glaciology*, 56(70), 51–62, [https://www.doi.org/10.3189/2015AoG70A921, 2015](https://www.doi.org/10.3189/2015AoG70A921)

Zekollari, H., Huybrechts, P., Noël, B., van de Berg, W.J. and van den Broeke, M.R.: Sensitivity, stability

heeft verwijderd: 

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Koppen (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

Gewijzigde veldcode

heeft verwijderd: 

Gewijzigde veldcode

and future evolution of the world's northernmost ice cap, Hans Tausen Iskappe (Greenland). The Cryosphere, 11(2), 805–825, <https://www.doi.org/10.5194/tc-11-805-2017>, 2017

1745

Zemp, M., Huss, M., Thibert, E., McNabb R., Huber, J., Barandun M., Machguth H., Nussbaumer S.U., Gärtnner-Roer I., Thomson L., Paul F., Maussion F., Kutuzov S. and Cogley J.G.: Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016. Nature, 568, 382–386, <https://www.doi.org/10.1038/s41586-019-1071-0>, 2019

1750 Zhang, T: Influence of the seasonal snow cover on the ground thermal regime: An overview, Rev. Geophys., 43, RG4002, <https://www.doi.org/10.1029/2004RG000157>, 2005

755

Zhao, L., Tian, L., Zwinger, T., Ding, R., Zong, J., Ye, Q. and Moore, J. C.: Numerical simulations of Gurenhekou Glacier on the Tibetan Plateau. Journal of Glaciology, 60(219), 71–82, <https://www.doi.org/10.3189/2014JoG13J126>, 2014.

760

Zhou, Y., Guo, D., Qiu, G., Cheng, G. and Li, S.: Geocryology in China, 450 pp., Sci. Press, Beijing, 2000 (in Chinese)

Zwinger, T., Greve, R., Gagliardini, O., Shiraiwa, T. and Lylly, M.: A full Stokes-flow thermo-mechanical model for firm and ice applied to the Gorshkov crater glacier, Kamchatka. Annals of Glaciology, 45, 29–37,

<https://www.doi.org/10.3189/172756407782282543>, 2007

Zwinger, T. and Moore, J. C.: Diagnostic and prognostic simulations with a full Stokes model accounting for superimposed ice of Midtre Lovénbreen, Svalbard. The Cryosphere, 3(2), 217–229, <https://www.doi.org/10.5194/tc-3-217-2009>, 2009

Gewijzigde veldcode

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (V.S.), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Engels (V.S.)

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (V.S.), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Engels (V.S.)

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Hyperlink, Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Hyperlink, Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

Gewijzigde veldcode

heeft verwijderd: :

heeft verwijderd: ,

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast: Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

heeft opmaak toegepast ... [44]

Met opmaak: Regelafstand: 1.5 regel

heeft opmaak toegepast ... [45]

heeft opmaak toegepast ... [46]

heeft opmaak toegepast ... [47]

heeft opmaak toegepast ... [48]

Gewijzigde veldcode

Pagina 8: [1] heeft verwijderd	Lander Van Tricht	18-08-2022 08:20:00
Pagina 15: [2] heeft verwijderd	Lander Van Tricht	16-08-2022 17:22:00
▼		
Pagina 15: [2] heeft verwijderd	Lander Van Tricht	16-08-2022 17:22:00
▼		
Pagina 15: [3] heeft verwijderd	Lander Van Tricht	18-08-2022 20:12:00
▼		
Pagina 15: [3] heeft verwijderd	Lander Van Tricht	18-08-2022 20:12:00
▼		
Pagina 15: [4] heeft verwijderd	Lander Van Tricht	18-08-2022 20:12:00
▼		
Pagina 15: [4] heeft verwijderd	Lander Van Tricht	18-08-2022 20:12:00
▼		
Pagina 15: [5] heeft verwijderd	Lander Van Tricht	18-08-2022 20:13:00
▼		
Pagina 15: [5] heeft verwijderd	Lander Van Tricht	18-08-2022 20:13:00
▼		
Pagina 15: [6] heeft verwijderd	Lander Van Tricht	18-08-2022 20:12:00
▼		
Pagina 15: [6] heeft verwijderd	Lander Van Tricht	18-08-2022 20:12:00
▼		
Pagina 15: [7] heeft verwijderd	Lander Van Tricht	18-08-2022 20:12:00
▼		
Pagina 15: [7] heeft verwijderd	Lander Van Tricht	18-08-2022 20:12:00
▼		
Pagina 15: [8] heeft verwijderd	Lander Van Tricht	25-08-2022 15:39:00
▼		
Pagina 15: [8] heeft verwijderd	Lander Van Tricht	25-08-2022 15:39:00
▼		
Pagina 15: [8] heeft verwijderd	Lander Van Tricht	25-08-2022 15:39:00
▼		
Pagina 15: [8] heeft verwijderd	Lander Van Tricht	25-08-2022 15:39:00
▼		

▼
▲ **Pagina 15: [8] heeft verwijderd** Lander Van Tricht **25-08-2022 15:39:00**

▼
▲ **Pagina 15: [8] heeft verwijderd** Lander Van Tricht **25-08-2022 15:39:00**

▼
▲ **Pagina 15: [8] heeft verwijderd** Lander Van Tricht **25-08-2022 15:39:00**

▼
▲ **Pagina 19: [9] heeft verwijderd** Lander Van Tricht **19-08-2022 16:51:00**

▼
▲ **Pagina 19: [10] heeft verwijderd** Lander Van Tricht **19-08-2022 16:50:00**

▼
▲ **Pagina 30: [11] heeft verwijderd** Lander Van Tricht **26-08-2022 13:44:00**

▼
▲ **Pagina 30: [11] heeft verwijderd** Lander Van Tricht **26-08-2022 13:44:00**

▼
▲ **Pagina 30: [11] heeft verwijderd** Lander Van Tricht **26-08-2022 13:44:00**

▼
▲ **Pagina 30: [12] heeft verwijderd** Lander Van Tricht **16-08-2022 17:23:00**

▼
▲ **Pagina 30: [12] heeft verwijderd** Lander Van Tricht **16-08-2022 17:23:00**

▼
▲ **Pagina 30: [12] heeft verwijderd** Lander Van Tricht **16-08-2022 17:23:00**

▼
▲ **Pagina 30: [12] heeft verwijderd** Lander Van Tricht **16-08-2022 17:23:00**

▼
▲ **Pagina 30: [13] heeft opmaak toegepast** Lander Van Tricht **19-08-2022 17:53:00**

Lettertype: Vet

▲ **Pagina 30: [14] Met opmaak** Lander Van Tricht **25-08-2022 09:34:00**

Centrum

▲ **Pagina 30: [15] heeft verwijderd** Lander Van Tricht **18-08-2022 13:43:00**

▲ **Pagina 30: [16] heeft opmaak toegepast** Lander Van Tricht **02-09-2022 10:41:00**

Lettertype: Vet, Tekstkleur: Tekst 1

▲ **Pagina 30: [17] heeft opmaak toegepast** Lander Van Tricht **02-09-2022 10:41:00**

Lettertype: 9 pt, Vet, Tekstkleur: Tekst 1

Pagina 30: [18] heeft opmaak toegepast Lander Van Tricht 28-08-2022 09:20:00

Lettertype: 9 pt, Vet, Tekstkleur: Tekst 1

Pagina 30: [19] heeft opmaak toegepast Lander Van Tricht 02-09-2022 10:41:00

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [19] heeft opmaak toegepast Lander Van Tricht 02-09-2022 10:41:00

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [20] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [21] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt

Pagina 30: [22] Met opmaak Lander Van Tricht 25-08-2022 09:34:00

Centrum

Pagina 30: [23] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [24] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt

Pagina 30: [24] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt

Pagina 30: [25] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [26] Met opmaak Lander Van Tricht 25-08-2022 09:34:00

Centrum

Pagina 30: [27] heeft verwijderd Lander Van Tricht 19-08-2022 17:46:00

x

x

Pagina 30: [27] heeft verwijderd Lander Van Tricht 19-08-2022 17:46:00

x

Pagina 30: [28] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [28] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [29] heeft opmaak toegepast Lander Van Tricht 26-08-2022 13:46:00

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [30] heeft opmaak toegepast Lander Van Tricht **02-09-2022 10:41:00**

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [31] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [31] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [32] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [32] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [33] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [33] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [33] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [33] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [34] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [34] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 30: [35] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [35] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [36] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [36] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Lettertype: 9 pt

Pagina 30: [39] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [39] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [40] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [40] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [40] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [40] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [41] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [41] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [41] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [42] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [42] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [42] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [42] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt

Pagina 30: [43] heeft opmaak toegepast Lander Van Tricht **26-08-2022 13:46:00**

Lettertype: 9 pt, Tekstkleur: Tekst 1

Pagina 44: [44] heeft opmaak toegepast Lander Van Tricht **18-08-2022 15:40:00**

Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

| Hyperlink, Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

| **Pagina 44: [47] heeft opmaak toegepast Lander Van Tricht 18-08-2022 15:42:00**

| Hyperlink, Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren

| **Pagina 44: [48] heeft opmaak toegepast Lander Van Tricht 18-08-2022 15:40:00**

| Lettertype: (Standaard) +Hoofdtekst (Times New Roman), 10 pt, Engels (Groot-Brittannië), Geen spelling- of grammaticacontrole uitvoeren