

The ozone–climate penalty over South America and Africa by 2100

Flossie Brown¹, Gerd A. Folberth², Stephen Sitch³, Susanne Bauer^{4,5}, Marijn Bauters⁶, Pascal Boeckx⁶, Alexander W. Cheesman^{3,7}, Makoto Deushi⁸, Inês Dos Santos⁶, Corinne Galy-Lacaux⁹, James Haywood^{1,2}, James Keeble^{10,11}, Lina M. Mercado^{3,12}, Fiona M. O'Connor², Naga Oshima⁸, Kostas Tsigaridis^{4,5}, Hans Verbeeck⁶

¹ College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK

² UK Met Office Hadley Centre, Exeter, UK

³ College of Life and Environmental Sciences, University of Exeter, Exeter, UK

10 ⁴ Center for Climate Systems Research, Columbia University, New York, NY, USA

⁵ NASA Goddard Institute for Space Studies, New York, NY, USA

⁶ Department of Environment, Ghent University, Ghent, Belgium

⁷ Centre for Tropical Environmental and Sustainability Science, James Cook University, Australia

⁸ Meteorological Research Institute, Japan Meteorological Agency, Tsukuba, Ibaraki, Japan

15 ⁹ Laboratoire d'Aerologie, Université Toulouse III Paul Sabatier, CNRS, Toulouse, France

¹⁰ Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge, UK

¹¹ National Centre for Atmospheric Science (NCAS), University of Cambridge, UK

¹² UK Centre for Ecology and Hydrology, Wallingford, UK

20 *Correspondence to:* Flossie Brown (fb428@exeter.ac.uk)

Abstract

Climate change has the potential to increase surface ozone (O₃) concentrations, known as the ‘ozone–climate penalty’, through changes to atmospheric chemistry, transport and dry deposition. In the tropics, the response of surface O₃ to changing climate is relatively understudied, but has important consequences for air pollution, human and ecosystem health. In this study, we

25 evaluate the change in surface O₃ due to climate change over South America and Africa using three state-of-the-art Earth system models that follow the Shared Socioeconomic Pathway 3-7.0 emissions scenario from CMIP6. In order to quantify changes due to climate change alone, we evaluate the difference between simulations including climate change and simulations with a fixed present-day climate. We find that by 2100, models predict an ozone–climate penalty in areas where O₃ is already

30 predicted to be high due to the impacts of precursor emissions, namely urban and biomass burning areas, although on average, models predict a decrease in surface O₃ due to climate change. We identify a small but robust positive trend in annual mean surface O₃ over polluted areas. Additionally, during biomass burning seasons, seasonal mean O₃ concentrations increase by 15 ppb (model range 12 to 18 ppb) in areas with substantial biomass burning such as the arc of deforestation in the Amazon. The ozone–climate penalty in polluted areas is shown to be driven by an increased rate of O₃ chemical production, which is strongly influenced by NO_x concentrations and is therefore specific to the emissions pathway chosen. Multiple linear regression finds

35 the change in NO_x concentration to be a strong predictor of the change in O₃ production whereas increased isoprene emission

rate is positively correlated with increased O₃ destruction, suggesting NO_x-limited conditions over the majority of tropical Africa and South America. However, models disagree on the role of climate change in remote, low-NO_x regions, partly because of significant differences in NO_x concentrations produced by each model. We also find that the magnitude and location of the ozone–climate penalty in the Congo basin has greater inter-model variation than in the Amazon, so further model development and validation is needed to constrain the response in central Africa. We conclude that if the climate were to change according to the emissions scenario used here, models predict that forested areas in biomass burning locations and urban populations will be at increasing risk of high O₃ exposure, irrespective of any direct impacts on O₃ via the prescribed emissions scenario.

1. Introduction

Climate change threatens to bring new pressures to the tropical forests, grasslands and agricultural lands of Africa and South America. As a result of shifts in emissions, atmospheric chemistry, and meteorology, as well as changes in vegetation behaviour such as transpiration rate, surface O₃ concentrations are likely to change (e.g. Turnock et al., 2019; Griffiths et al., 2021; Zanis et al., 2022), which may impair or benefit human and vegetation health (Agathokleous et al., 2019; Emberson, 2020) depending on the direction of change. O₃ is a highly oxidising compound, formed in the atmosphere through reaction of volatile organic compounds (VOC) or carbon monoxide (CO) with hydroxyl radicals (OH), and nitrate radicals (NO₃) in the presence of nitrogen oxides (NO_x). However, it can also be removed from the atmosphere through reactions with many of the same chemical species (NO_x, VOC, OH) depending on their relative concentrations. In addition to chemical pathways, O₃ can be removed from the lower atmosphere by dry deposition, which includes stomatal uptake by plants (Silva & Heald, 2018). Stomatal uptake of O₃ and subsequent ozone–plant damage, can lead to reduced carbon drawdown from the atmosphere (Sitch et al., 2007; Yue & Unger, 2018; Franz & Zaehle, 2020), and changes to biosphere–climate interactions (Sadiq et al., 2017). Sitch et al. (2007) showed that the tropics may be especially sensitive to high O₃ concentrations and therefore susceptible to large productivity losses if surface O₃ were to increase. Additionally, O₃ is a near-term climate forcer with impacts on the radiative balance leading to a positive radiative forcing of climate through absorption of longwave radiation (Myhre et al., 2017). As the tropical forests are vital as sinks for atmospheric CO₂, and tropical ecosystems play a vital role in regional and global climate (Lewis, 2006; Bonan, 2008), an understanding of the impact of climate change on surface O₃ concentrations in the tropics is critical.

Whilst there have been no studies specifically assessing changes in surface O₃ due to climate change in the tropics, global studies have suggested that chemical and biological changes in temperature-dependent chemistry, natural emissions of precursors and land surface properties; as well as dynamical changes including circulation changes and transport from the stratosphere may lead to an ‘ozone–climate penalty’ over some continental areas (Jacob & Winner, 2009; Doherty et al., 2013;

Zanis et al., 2022). The ‘ozone–climate penalty’ is defined as an increase in surface O₃ concentrations due to climate change alone. It is influenced by many complex chemical and biological processes, which are not all well-understood or represented in current climate models, although there has been substantial recent research to reduce uncertainty in temperature-sensitive chemistry, meteorology and land–atmosphere feedbacks (Oswald et al., 2015; Coates et al., 2016; Sadiq et al., 2017; Romer et al., 2018; Archibald et al., 2020b). There has been relatively little research focusing on O₃ in tropical environments. Unlike the more commonly studied extra-tropical Northern hemisphere, the tropics and subtropics have relatively low (natural) NO_x emissions and high biogenic VOC emissions, high actinic flux and strong atmospheric convection (Bond et al., 2002; Pugh et al., 2010; Paulot et al., 2012). This paper focuses on South America and Africa. We exclude equatorial Asia because the atmospheric chemistry in this region is more uncertain due to difficulties in detecting and accounting for peat fire emissions (Prosperi et al., 2020). Equatorial Asia also has a greater marine influence with most model grid boxes containing ocean as well as land, so they may follow a different chemical regime.

Many areas in Africa and South America are considered remote (defined in this paper by low emissions of NO_x), although increasing anthropogenic activity such as urbanisation and biomass burning causes moderate NO_x emissions in some regions (e.g. Kuhn et al., 2010; Pacifico et al., 2015; Shi et al., 2020). The sensitivity of O₃ production to NO_x depends on the relative concentrations of NO_x and VOCs. Isoprene is the most abundant biogenic VOC in remote Africa and South America and must be oxidised in the atmosphere before it can form O₃ (Liu et al., 2016). NO_x is produced from both natural and anthropogenic sources including soils, lightning, transport and biomass burning. Regions are defined as NO_x-limited when increasing VOCs or OH acts to reduce O₃ concentrations through oxidation and formation of organic peroxides (Pacifico et al., 2012). In this NO_x-limited case, increasing NO_x will lead to greater O₃ formation. Conversely, in VOC-limited regions with sufficient NO_x present, increasing NO_x concentrations may reduce O₃ concentrations by removal of the key O₃-forming radicals OH (reaction: OH + NO₂ → HNO₃).

Earlier studies have found that South America and Africa are generally NO_x-limited (Ziemke et al., 2009; Bela et al., 2015), and that increases in NO_x concentration associated with climate change will be a key driver of O₃ increases over South America and Africa. Doherty et al. (2013) attribute the NO_x increase predominantly to enhanced decomposition of the NO_x reservoir species, peroxyacetyl nitrate (PAN). A fraction of emitted NO_x is locked up as PAN, which decomposes back into NO_x in warmer temperatures, sometimes after having travelled long distances from the NO_x source. As PAN is unstable at high temperatures, climate change will result in a smaller fraction of NO_x being stored as PAN in NO_x source regions and may also decrease the amount of NO_x that is transported into remote regions (Schultz et al., 1999; Finney et al., 2018). Lightning NO_x is known to contribute to O₃ formation, however studies project both increases and decreases in future lightning frequency (Clark et al., 2017; Finney et al., 2018) leading to low confidence in how O₃ will be affected by climate-driven changes in lightning (Fu & Tian, 2019; Murray, 2016).

The role of isoprene in the ozone–climate penalty is debated as there is uncertainty about how isoprene emissions will change in the future in response to temperature, CO₂ and land-use change (Fu & Liao, 2016; Fu & Tian, 2019) and how to best represent isoprene chemistry in climate models (Weber et al., 2021). Biogenic isoprene emissions increase strongly with temperature and vegetation stress (e.g. Guenther et al., 1993; Niinemets et al., 1999; Unger et al., 2013; Morfopoulos et al., 2021), but very high temperatures or moisture stress may cause ‘die-back’ of vegetated areas, which would decrease isoprene emissions overall (Sanderson et al., 2003; Cox et al., 2004; Malhi et al., 2009). On the other hand, elevated CO₂ concentrations directly inhibit isoprene emission but can indirectly increase emission if this CO₂ fertilisation effect results in increased plant productivity (Pacifico et al., 2011; Squire et al., 2014; Hantson et al., 2017). Isoprene, NO_x and OH concentrations are influenced by isoprene chemistry. Formation of isoprene nitrates partially recycles NO_x, and oxidation of isoprene partially recycles HO_x (Bates and Jacob, 2019). Difference between models in their calculation of the yields and recycling rates of these species is likely to affect O₃ concentrations.

Besides chemical processes, dry deposition of O₃ to vegetation is a major O₃ sink, accounting for 20% of O₃ removal (Wedow et al., 2021). Most O₃ deposition occurs via plant stomata, which respond to changes in the climate. (Silva & Heald, 2018; Clifton et al., 2020). Increased CO₂, temperature and vapour pressure deficit (VPD) will decrease stomatal conductance and therefore decrease O₃ deposition rate. This could lead to increased concentrations of O₃ in the atmosphere, although it would have a protective effect for plants (Emberson et al., 2013; Lin et al., 2020).

Studies agree that over the ocean, average surface O₃ concentrations will decrease under the influence of climate change (Zeng et al., 2008; Doherty et al., 2013; Zanis et al., 2022). The warmer air can hold more water vapour, a major species contributing to O₃ loss (reaction: O(1D) + H₂O → 2OH leading to O₃ loss via OH₂ + O₃ → OH + 2O₂). There may also be O₃ reductions in remote regions over land due to this process although natural emissions can change the atmospheric chemistry over the continents. Other factors contributing to the O₃ concentration over the continents are numerous and complex, including changes to oxidising capacity, stratospheric transport and land-use (Archibald et al., 2010; Squire et al., 2015).

This paper quantifies the effect of climate change on surface O₃ concentrations in the future, and aims to understand its uncertainty and the relative contributions of the underlying processes. We focus on areas with robust O₃ changes to identify areas in South America and Africa at greatest risk by 2100. Sect. 1 has introduced the key chemical species involved in O₃ chemistry and the most important changes that may result from climate change. In Sect. 2, we provide the model details, data used for evaluation, and description of analysis of model output. Sect. 3 evaluates model predictions of surface O₃ in the present-day, evaluates model predictions for surface O₃ changes in 2100 and examines the importance of chemical and deposition changes in controlling the ozone–climate penalty in models. Finally, Sect. 4 discusses the key trends predicted by the models, limitations of the study and crucial uncertainties in the models.

2. Methods

We analyse surface O₃ for simulations that follow a medium-high emissions pathway created for CMIP6 (Pascoe et al., 2019). The simulations were carried out as part of an ensemble of Earth system model experiments designed to quantify the climate and air quality impacts of aerosols and trace gases in climate models (Collins et al., 2017), named Aerosol-Chemistry Model
140 Intercomparison Project (AerChemMIP). For this study three Earth system models were used: UKESM1-0-LL (abbr. UKESM1), GISS-E2-1-G (abbr. GISS), MRI-ESM2-0 (abbr. MRI). These were selected because of their detailed tropospheric chemistry schemes and availability of output for O₃ and O₃ precursor concentrations on the Earth System Grid Federation (ESGF).

145 The simulations follow the Shared Socioeconomic Pathway 3-7.0 (SSP3-7.0) emissions scenario, a scenario assuming low international cooperation to protect the environment. This includes high emissions of non-methane near-term climate forcers and substantial land-use change (O'Neill et al., 2016; Gidden et al., 2019). The prescribed emissions include anthropogenic and biomass burning emissions of NO, CO₂ and CO (Rao et al. 2017; Riahi et al., 2017) and the future pathway for CH₄ is calculated as an atmospheric concentration (Meinshausen et al., 2019). Emissions due to growing populations and poor
150 international cooperation results in significant temperature increases by 2100 (Turnock et al., 2019) and societies that are highly vulnerable to climate change. This emissions pathway was chosen in order to understand changes in end-of-century O₃ concentration if there is no international cooperation to reduce precursor emissions.

2.1 Model descriptions

Here we provide a brief description of the Earth system models and their tropospheric chemistry schemes that are relevant to
155 this study.

UKESM1-0-LL (abbr. UKESM1)

UKESM1-0-LL is a combination of HadGEM3 (Williams et al., 2018) with additional land and atmospheric chemistry components (Sellar et al., 2019). The UK Chemistry and Aerosol scheme (UKCA) contains stratospheric and tropospheric chemistry (Archibald et al., 2020a) combined with the GLOMAP-mode aerosol microphysics scheme (Mulcahy et al., 2018,
160 Mulcahy et al., 2020).

Interactive emissions include isoprene, monoterpenes, lightning NO_x and soil NO_x. Isoprene and monoterpene emissions respond to light and temperature and the isoprene scheme also includes CO₂ inhibition (Archibald et al., 2020a; Mulcahy et al., 2018) following the emission model of Pacifico et al. (2011). Lightning NO_x is calculated using the parameterisation of
165 Price and Rind (1992). Secondary organic aerosols (SOA) are calculated as a fixed yield of 26% from gas-phase oxidation reactions involving monoterpene sources.

The terrestrial biogeochemistry is provided by JULES (Wiltshire et al., 2019; Wiltshire et al., 2021). Stomatal conductance in JULES is similar to the Ball-Berry-Leuning model (Leuning, 1995) and responds to the ratio of internal to external CO₂ concentrations and leaf humidity deficit (Jacobs, 1994).

The model has a horizontal resolution of 1.25° latitude by 1.875° longitude with 85 vertical levels in a hybrid height coordinate.

MRI-ESM2-0 (abbr. MRI)

MRI-ESM2-0 (Yukimoto et al., 2019; Kawai et al., 2019; Oshima et al., 2020) contains an atmospheric and land-surface model (MRI-AGCM3.5), an ocean and sea-ice model (MRI.COMv4), an aerosol model (MASINGAR mk-2r4c) and an atmospheric chemistry model (MRI-CCM2.1). The chemistry model includes tropospheric, stratospheric and mesospheric chemistry with 90 chemical species and 259 chemical reactions (Deushi & Shibata, 2011).

Lightning NO_x is interactive (Price & Rind, 1992) but natural emissions from land and ocean are prescribed as monthly climatologies, including isoprene and soil NO_x (Deushi & Shibata, 2011). 15% of natural terpene emissions at the surface form SOA and SOA have identical properties to POA.

Each component employs different horizontal resolutions but the outputs used in this paper are from the chemistry component which uses a horizontal resolution of 2.8125° latitude by 2.8125° longitude with 80 vertical levels in a hybrid sigma pressure coordinate.

GISS-E2-1-G (abbr. GISS)

GISS-E2-1-G contains a coupled troposphere and stratosphere chemistry scheme using the G-PUCCINI chemistry mechanism (Shindell et al., 2013; Kelley et al., 2020) combined with the One-Moment Aerosol (OMA) scheme for aerosols (Bauer et al., 2020).

Lightning NO_x is interactive as described by Kelley et al. (2020). Natural emissions include soil NO_x and isoprene, which respond to light and temperature (Shindell et al., 2006) following the algorithm defined by Guenther et al., (1995). Monoterpenes are prescribed. SOA are calculated using the CBM4 chemical mechanism to describe the gas phase tropospheric chemistry together with all main aerosol components including SOA formation and nitrate, and is calculated using four tracers in the model. Isoprene (VOCs) contribute to the formation of SOA (Tsigaridis et al., 2018).

GISS has a horizontal resolution of 2.00° latitude by 2.25° longitude with 40 vertical levels output on hybrid sigma pressure coordinate.

2.2 Data analysis methods

200 **Model evaluation**

In situ observations from 65 sites across South America and tropical Africa, covering key biomes and land-use types, are used for grid level model evaluation. Monthly mean O₃ concentrations from individual sites have been aggregated into seven regions by grouping together sites within latitude and longitude bounds (see Table S1). To compare to models, the coordinates of the in situ measurement sites are matched to the nearest model grid cell coordinates. To create an average seasonal cycle for each
205 region, sites with the same nearest grid cell are averaged together to create a grid cell seasonal cycle. Then, grid cell seasonal cycles in the same region are averaged together. Monthly mean data from 1990 up to 2021 were used, although most sites only provide a few years of data. Data were excluded if there was an unequal distribution of data points over the monthly mean diel cycle.

210 To produce a model seasonal cycle, the monthly mean O₃ concentration was calculated using CMIP6 historical simulations for the period 1990–2014. This was done for each grid cell that contained an observation site. The standard deviation in monthly mean O₃ concentrations between the grid cells used was calculated for each region (i.e. it represents variation in O₃ geographically between the sites rather than inter-annual variation).

215 For model evaluation against Tropospheric Emission Spectrometer (TES) data from the Aura satellite, which retrieves O₃ in ppb over 67 pressure levels, we use O₃ concentrations at the lowest level available from TES (825 hPa). Monthly mean gridded outputs are used for the period 2004–2011, the period for which complete monthly mean data is available. As with the model output, satellite grid cell coordinates closest to the in situ site coordinates were selected. The monthly mean and standard deviation for each region was calculated, only using data from grid cells containing in situ sites.

220 **CMIP6 model output**

We isolate the effect of climate change on surface O₃ concentrations using the difference between two simulations which consider the same trajectory of anthropogenic emissions changes but differ in climate. The simulations are global model runs driven with prescribed sea surface temperatures (SSTs) over the period 2015–2100. We use a simulation driven with changing SSTs from the coupled simulation ssp370 so that the climate changes in accordance with the emissions changes, this simulation
225 is named ssp370SST. We use a second simulation with prescribed SSTs and sea ice concentrations taken from a present-day climatology (2005–2014) in historical simulations, named ssp370pdSST. Importantly, although anthropogenic emissions are identical in both simulations, ssp370pdSST does not include the resulting climate change.

To isolate the effect of climate change on tropospheric O₃, we subtract ssp370pdSST (present-day constant climate + future emissions) from ssp370SST (future climate + future emissions) following Zanis et al. (2022) and Szopa et al. (2021). Biomass burning and land-use change are considered anthropogenic and are prescribed for both models but natural emissions are allowed to change depending on the model set-up. In this way, the background atmospheric composition is based on the future emissions scenario used, since the response of atmospheric chemistry to climate change may depend on the background concentrations of precursors. In UKESM1, CO₂ is also fixed to present-day concentrations in ssp370pdSST so that the effect of climate change includes the effect of CO₂ inhibition.

Model output is taken as monthly means during the period 2090–2100 between 40° S and 40° N from experiments ssp370SST and ssp370pdSST. All variables used are outlined in Table S2. The change due to climate change refers to subtracting ssp370pdSST from ssp370SST, where positive values are considered an O₃–climate penalty. When evaluating the change due to climate change regionally, this study distinguishes between ‘high-NOx’ and ‘remote’ areas (Fig. 1). High-NOx areas are defined as areas where the annual mean NOx emissions are above the 95th percentile for the tropics and subtropics (40° S–40° N).

O₃ concentrations are taken from the lowest model level (~ 20 m above orography). Where multi-model means are shown, data has been re-gridded to the 2.8125° by 2.8125° grid used by MRI. We evaluate the ozone–climate penalty as a yearly average and seasonally. To identify seasonal patterns we aggregate the data by burning season. The Western African burning season is defined as Dec–Feb, the Southern African burning season is June and July and the Southern Amazon burning season is Aug–Oct.

To attribute the ozone–climate penalty to precursor variables, we also use NOx, isoprene emission rate, OH and surface temperature variables. When presenting these variables, the ocean has been masked so that only land surface changes are presented. ‘Surface concentrations’ shown in this study refer to chemical mixing ratios in the lowest model grid cells and ‘background concentrations’ refer to chemical mixing ratios in the absence of climate change (using data from ssp370pdSST).

To evaluate the O₃ budget, we also use O₃ chemical production, O₃ chemical loss and dry deposition variables. O₃ chemical production is defined as O₃ produced from the reaction NO+RO₂ / HO₂, O₃ chemical loss is the sum of (i) O(1D)+H₂O; (ii) O₃+HO₂; (iii) O₃+OH; (iv) O₃+alkenes (e.g. isoprene). To compare these three variables on the same scale, we convert the units to Tg year⁻¹ and sum production and loss over the lowest 1 km, the approximate boundary layer height. We choose 1 km to establish an approximate region that can contribute to surface O₃ concentrations.

In Sect. 3.4, we present a sensitivity study relating changes in NO_x concentration and isoprene emissions to changes in O₃ chemical production. We use monthly mean data in South America and Africa within 30° S and 30° N to calculate a monthly climatology for 2090–2100, masking the ocean and the non-vegetated region of Saharan Africa. To identify the limiting O₃ precursor in the tropics and subtropics, the percentage change in O₃ production rate (n>500) is modelled with an ordinary least squares linear regression using the percentage change in NO_x and isoprene as predictor variables (see S4: The relationship between NO_x and O₃ production). To interpret the ability of the model we rely on the central limit theorem to assume that the normalised sum of the residuals can be approximated by a normal distribution. Unique months and grid cells are treated as separate data points. To present the results graphically, we highlight values using star markers if they are above a threshold of the 95th percentile for NO_x concentrations using the monthly climatology for each model individually, with aims to identify biomass burning areas and cities. The atmospheric chemistry in these areas may be different due to the elevated NO_x concentrations. Therefore, some grid cells will be above the 95th percentile for specific months only (biomass burning seasons).

3. Results

3.1. Evaluation of model skill for present-day surface O₃ concentrations

Results show that climate models are able to capture the observed seasonal cycle in most regions except for West Africa and DR Congo (Fig. 1). However, the models overpredict monthly mean surface O₃ concentrations by up to 50 ppb, with the largest bias present in remote forest locations such as the Congo area (Fig. 1e). GISS overall has the smallest positive bias out of all the models, and MRI has the largest. UKESM1 shows the smallest seasonal variation in O₃ concentration, which is often closer to the observed seasonal pattern.




Figure 1: (a)–(g) compare monthly mean O₃ (black dashed line) from site measurements averaged over 7 regions to model predictions for the period 1990–2014 and satellite products from the TES satellite at 825 hPa (navy dash-dot line). Model means are shown for UKESM1 (purple solid line), GISS (blue solid line) and MRI (red solid line) and 2 standard deviations from the mean are shaded. Grey crosses indicate monthly mean O₃ measurements from individual sites and years.

The central panel marks the locations where O₃ measurement sites are located (purple crosses) and how the sites have been grouped (coloured shading). Observations (a)–(c) and (f)–(g) use data from the Tropospheric O₃ Assessment Report (TOAR I, Schröder, 2021; Schultz et al., 2017), (d) and (e) from INDAAF (<http://www.indAAF.obs-mip.fr>) and (e) from CONGOFLUX in Yangambi, DR Congo. Grey grid cells cover areas where NO_x emissions are above the 95th percentile for the region 40° S–40° N in 2100 (prescribed by (Rao et al., 2017; Riahi et al., 2017), indicative of biomass burning areas and cities. Un-shaded areas are referred to as ‘remote’ throughout.

Grey shading on Fig. 1 highlights the areas in South America and Africa with the highest NO_x emissions. The shaded areas represent areas with high biomass burning emissions and urban areas and are referred to as ‘high-NO_x’ areas in this paper. In the Southern Amazon (Fig. 1b), the biomass burning months are August and September and both models and observations predict the highest O₃ concentrations in this season. However, the observed monthly mean O₃ concentrations range between 9 to 20 ppb whereas models predict values up to 40 ppb, with GISS displaying the smallest positive bias. In Africa (Figs 1d–1f), the biomass burning months are December–February (North / West Africa) moving to June–July (Southern Africa). Whilst models predict concentrations of up to 80 ppb in the Congo during these months due to transport of precursors from biomass burning, observations show substantially lower surface O₃ concentrations of less than 20 ppb at the remote locations sampled, although the highest O₃ concentrations occur during December–February (Adon et al., 2010; 2013; Ossouhou et al., 2019). In

fact, seasonal variation is low at the Congo sites whereas models predict strong seasonal patterns (Fig. 1e). In months without substantial burning, GISS captures the low O₃ values well and UKESM1 and MRI overestimate by 10 to 15 ppb.

305 The enhanced O₃ concentrations predicted by models in burning seasons over the Congo and West Africa are also captured in satellite retrievals at 825 hPa. Satellite O₃ concentrations in the DR Congo are 10 ppb higher in December–February compared to months without burning (Fig. 1e). At 825 hPa, these satellite capture O₃ concentrations above the altitude of in situ observation sites and the lowest model level. Model predictions for O₃ at 825 hPa are ~10 ppb higher than the lowest model level and therefore compare well to satellite retrievals (Fig. S1), especially in Amazonia (Fig. S1b) and West Africa (Fig. S1d).

310 Similar to biases over the remote continents, measurements from Réunion Island (Fig. 1f), which capture oceanic air masses, are overestimated by 5 ppb in GISS and by 12 ppb in UKESM1 and MRI, although the seasonal cycle is reproduced. On the other hand, over the urban sites in South Africa, Johannesburg, UKESM1 replicates the observed mean with moderate accuracy, whereas GISS and MRI overestimate by 15 ppb (Fig. 1g).

3.2. Average changes to atmospheric composition over Africa and South America at the end of 2100

315 The overall change in O₃ concentration due to climate change over the African and South American land surface is shown in Fig. 2a for each model compared to the simulation with a fixed present-day climate. All models predict that on average O₃ concentrations will decrease due to climate change over the tropical land surface in this study. The magnitude of the predicted O₃ change for each model will depend on background concentrations of O₃ chemical precursors, their change due to climate change and individual model mechanistic details. There is significant diversity in background atmospheric composition

320 between models (Fig. 2, blue marker) and the direction of change in surface NO_x and OH concentration due to climate change (Figs 2b, 2c, arrows).

Different temperature sensitivities of the models (differing by up to 2.7 K) and different biogenic VOC schemes will contribute to the inter-model variation (Fig. S2). UKESM1 has the greatest temperature sensitivity with a 6.5 K increase in temperature

325 over the tropical land surface due to climate change (Fig. 2e). The temperature change due to climate change varies seasonally and regionally, which may affect concentration of O₃ precursors locally, with dry seasons temperatures predicted to rise by 1–1.5 K more than wet season temperatures (Fig. S3).

The change in NO_x concentration between models is determined by the balance of changes in isoprene nitrate formation, OH concentrations, PAN decomposition and lightning in the models. A decrease in NO_x concentrations could be related to changes

330 in OH concentration and precipitation (and thus NO_x removal via reaction $\text{NO}_2 + \text{OH} \rightarrow \text{HNO}_3$) and isoprene (and thus NO_x removal via isoprene nitrate formation), whereas NO_x concentration increases may be due to increased PAN decomposition or lightning.




Figure 2: The change in surface concentration of (a) O₃, (b) NO_x, (c) OH, and the change in (d) isoprene emission rate and (e) surface temperature from experiment ssp370pdSST with no climate change (blue line) to experiment ssp370SST with climate change (red line) for the three climate models in this study. Variables have been averaged over the African and South American continents between 12° N–30° S for the period 2090–2100. The change due to climate change is significant at the 5% level for all variables and models except isoprene in MRI (which is prescribed so does not change).

NO_x emissions, including biomass burning emissions, are prescribed based on the SSP3-7.0 scenario but lightning NO_x and soil NO_x differ between the models based on the chosen parameterisation of individual models. Compared to the present-day, NO_x emissions in biomass burning areas decrease in Africa to follow projected trends, but do not change in South America. NO_x emissions increase in cities and Nigeria especially has major growth in urban areas. Compared to the scenario without climate change, total lightning NO_x emissions increase in all models, and the increases occur during the wet season (Fig. S4). MRI predicts much larger increases than GISS and UKESM1, and UKESM1 shows a decrease in lightning NO_x over the Amazon basin in December–February (Fig. S4a) although the net effect over all seasons is positive. Peroxyacetyl nitrate (PAN) decreases in all models (–94 ppt, –61 ppt, –30 ppt for UKESM1, GISS and MRI respectively) due to increased thermal

350 decomposition. In GISS and UKESM1, the increase in isoprene emissions can increase removal of NO_x via formation of isoprene nitrates. Soil NO_x does not change in response to climate change in any model.


Hydroxyl radical (OH) concentration determines the oxidising capacity of the atmosphere and affects rates of reaction such as VOC oxidation and ozone destruction. Increased temperatures will increase atmospheric water vapour and OH production, 355 however OH concentrations decrease in the GISS model when climate change is included. A portion of this decrease can be attributed to an increase in isoprene emissions, which is much larger in GISS than UKESM1 (Fig. S2).

The increase in isoprene emission rate due to climate change depends on the isoprene emission scheme used, or in MRI, isoprene emissions are prescribed as a climatology. The greatest increase in isoprene emissions rate occurs in the GISS model, 360 which increases from $2.5 \times 10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$ to $3.3 \times 10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$ when climate change is considered, whereas UKESM1, which accounts for CO₂ inhibition, increases more modestly from $2.9 \text{ kg m}^{-2} \text{ s}^{-1}$ to $3.3 \times 10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$. Isoprene emissions are presented throughout rather than isoprene concentrations (see S2: Isoprene representation in this paper).



3.3 Changes in surface O₃ concentration due to climate change over remote regions compared to high-NO_x areas

This study focuses on the change in surface O₃ concentration over land in 2100, although we note there are significant decreases 365 in O₃ concentration over the oceans and non-vegetated areas such as Saharan Africa (Fig. 3). Over land, the multi-model mean shows increases of up to 4 ppb over urban areas and the biomass burning areas of South America and Africa (Fig. 3a) whereas ocean-influenced locations such as Northeast Brazil are expected to benefit by a 4 to 5 ppb decrease in surface O₃. However, the direction of change in surface O₃ concentration over central Africa and the remote Amazon (North West) is not robust between models (Fig. 3c, 3d, 3e).


(a) O₃ due to climate change




(b) O₃ inter-model SD



(c) UKESM1 O₃



(d) GISS O₃



(e) MRI O₃





Figure 3: The average change in surface O₃ concentration due to climate change for the period 2090–2100 for (a) the multimodel mean, (c) UKESM1 only, (d) GISS only, (e) MRI only. (b) shows the inter-model standard deviation in the same units. Grey hatching in (a) covers areas where the inter-model standard deviation is greater than 20% of the multimodel mean value. Grey dots in panels (c)–(e) cover areas that are not significant at the 5% level from a student's t-test. Black dotted lines outline areas where background O₃ is higher than 40 ppb.

UKESM1 and MRI predict increases in surface O₃ concentration of up to 5 ppb in the Amazon and central Africa, with decreases over coastal regions due to climate change (Figs 3c, 3e). In the remote Amazon, MRI predicts an increase and UKESM1 a decrease in O₃ concentration, but neither change is significant. On the other hand, GISS predicts significant O₃ decreases across remote regions of up to 4 ppb (Fig. 3d), including central Africa, which experiences O₃ increases in the other simulations (Figs 3c, 3e).

Changes in surface O₃ concentration due to climate change in 2100 are shown in Fig. 4, grouped by regional biomass burning season, with dotted contours where background O₃ is 40 ppb (a number assumed associated with thresholds for plant O₃ damage) and 70 ppb. High background O₃ is associated with biomass burning and pollution in and around cities due to their higher NO_x emissions. These high O₃ areas also show the greatest increase in O₃ due to climate change (Fig. 4).




Figure 4: The multimodel mean change in surface O_3 concentration due to climate change for the period 2090–2100 for (a) the Western African burning season (Dec–Feb), (b) the Southern African burning season (June, July), (c) the Southern Amazon burning season (Aug–Oct), and (d) the remaining months with limited burning (March–May, Nov). Grey hatching covers areas where models disagree on the sign of the change due to climate change. Black dotted lines outline areas where background O_3 is 40 ppb and 70 ppb.

During December–February, the biomass burning area in Western Africa coincides with O_3 increases of 9 ppb (5 ppb for UKESM1, 7 ppb for MRI and 15 ppb for GISS; Fig. 4a) and similar O_3 penalties are seen for the Southern African biomass burning season during June–July (Fig. 4b). During the Amazon biomass burning season, there are even larger increases of up to 12 ppb in the Southern Amazon (Fig. 4c). In months without biomass burning, these areas have minor increases of 2 ppb for UKESM1 and MRI and a decrease of 3 ppb for GISS.

GISS is the only model to show significant decreases in monthly mean surface O_3 concentration over land, which consistently occur in areas and seasons with low background O_3 (not shown). This includes biomass burning areas but in seasons without burning, which are followed by large increases in the biomass burning season. The result is that seasonal changes in surface O_3 concentration due to climate change in GISS are much larger than UKESM1 and MRI, and models do not agree on the direction of change in remote areas, although the yearly average increase is similar between models (Fig. 3). This results in

405 uncertainty in the response to climate change from regions and seasons with low background O₃, but likely increases in areas and seasons with high background O₃ from anthropogenic NO_x emissions (Fig. 4, black dashed lines).

Grid cells which include highly populated regions and megacities are often associated with an increase in O₃ concentration in all months, and an average ozone–climate penalty of 3 ppb in the yearly average. In particular, there is an ozone–climate
410 penalty of 3 ppb that shows limited seasonal variation in grid cell containing the megacities in Nigeria (Lagos), Brazil (São Paulo, Rio de Janeiro) and Colombia (Bogotá, Medellín). This penalty is robust over Southeast Brazil in all seasons (Fig. 4).

3.4 Changes in chemical production and deposition of O₃ due to climate change

Attribution of changes in surface O₃ to changes in chemical production, chemical loss and dry deposition at the surface are
415 shown in Fig. 5. The increase in O₃ production due to climate change is the largest out of these terms (Fig. 5c) and increases the most (over 0.25 Tg year⁻¹) in high-NO_x areas where surface O₃ increases (Fig. 5a). Therefore, the increase in the rate of O₃ production is likely to be the main cause of the ozone–climate penalty in high-NO_x areas (high-NO_x defined as in Fig. 1). Removal of O₃ by deposition and chemical destruction has a smaller effect on O₃ concentration since, to a degree, the two terms cancel each other out; in high-NO_x areas, chemical loss increases by up to 0.1 Tg year⁻¹ and dry deposition decreases
420 by up to 0.05 Tg year⁻¹. In remote regions, there is considerable variation between models as indicated by the higher standard deviation in these areas (Fig. 5, column 2). GISS predicts decreases in O₃ production over remote regions of up to 0.1 Tg year⁻¹ and increases of up to 0.25 Tg year⁻¹ over high-NO_x regions, whereas MRI and UKESM1 predict increases in O₃ production across all regions except Saharan Africa. MRI predicts the largest increases in O₃ chemical production in remote areas of 0.25 Tg year⁻¹ (not shown).

425




Figure 5. The multi-model mean change in (a) surface O_3 concentration, (c) chemical production of O_3 , (e) chemical destruction of O_3 and (g) dry deposition of O_3 due to climate change. Panels (c), (e) and (g) show the change in O_3 in $Tg\ year^{-1}$ and chemical terms have been summed over a 1 km height. The inter-model standard deviations are shown in panels (b), (d), (f), (h).


Chemical loss and deposition changes become important in remote areas because these areas have the smallest increases in chemical production, but can have the largest changes in loss rate (Fig. S8) and deposition rate (Fig. S10). The rate of O₃ loss is correlated with the change in isoprene concentration, which is typical of a low-NO_x region due to reactions between isoprene and O₃ directly (Fig. S8). This leads to increases in the loss rate in most vegetated areas (Fig. S8). Conversely, the deposition rate decreases, presumably because the increased temperatures and lower relative humidity cause stomatal closure. Dry deposition varies between each model depending on stomatal response to temperature changes and boundary layer resistance changes (Fig. S10). In UKESM1, the increase in CO₂ also reduces stomatal conductance. UKESM1 shows a large decrease in deposition rate over the central Amazon, whereas MRI shows very little change regionally.

In high-NO_x areas, the increase in O₃ production is greater than the increase in loss leading to net chemical production of O₃. Evaluation of the sensitivity of O₃ chemical production rate to changes in isoprene emissions and NO_x concentration due to climate change for each model is shown in Fig. 6 and described below.

Isoprene emission rate increases in both GISS and UKESM1 on average, as expected in a warmer climate (MRI does not have interactive isoprene emission) (Fig. 6, row 3). UKESM1 predicts a substantial decrease in isoprene emission rate in the Northern Amazon and fractionally in West Africa (Fig. 6a). Decreases in isoprene emission and stomatal conductance have previously been simulated in the same area due to CO₂ inhibition (Pacifico et al., 2012; Chadwick et al., 2017; Turnock et al., 2020).

NO_x decreases in most areas in UKESM1 including high-NO_x areas, whereas GISS predicts increases of 2×10^{-11} to 6×10^{-11} mol mol⁻¹ in high-NO_x areas only and MRI predicts more uniform increases of 4×10^{-11} mol mol⁻¹ in all areas (Fig. 6, row 2). The magnitude of the background NO_x concentration in UKESM1 and the change due to climate change is also much larger than the other models. The negative NO_x concentration changes in UKESM1 and in some areas in GISS compared to MRI may be due to increased sequestration of NO_x into isoprene nitrates. This possibility is supported by evidence of anticorrelation between NO_x and isoprene in GISS and UKESM1 (Fig. 6). In GISS, the remote Amazon shows the largest isoprene increase and a decrease in NO_x concentration. UKESM1 also shows an increase in NO_x concentration downwind of the isoprene decrease in the Northern Amazon.

Despite differences in the magnitude and direction of the NO_x and isoprene changes, the change in O₃ chemical production rate has similar spatial patterns in all models (Fig. 6, row 4). Exceptions occur in central Africa where GISS predicts a decrease in production rate, and in Nigeria where UKESM1 is the only model that does not predict a large increase in O₃ production. These areas of Africa also exhibit differences in surface O₃ concentration between models, discussed in Sect. 3.3.



465 **Figure 6: (a), (b), (c) Surface NO_x concentrations in the absence of climate change and the average change due to climate change in (d), (e), (f) NO_x concentration, (g), (h), (i) isoprene emission rate and (j), (k), (l) O₃ production rate for the period 2090–2100 for (column 1) UKESM1, (column 2) GISS and (column 3) MRI.**

The O₃ production rate for GISS appears highly correlated with the change in NO_x concentration in Fig. 6e, whereas NO_x concentration decreases in many areas where O₃ production increases for UKESM1. Instead, isoprene may influence O₃ production in UKESM1. Areas with a decrease in isoprene emissions in UKESM1 also show a decrease or a smaller increase in O₃ production compared to other areas, suggesting isoprene is important for O₃ production in UKESM1 even in remote regions such as the Northern Amazon.

475 To determine the strength of the relationship between O₃ production rate and changes in precursors NO_x and isoprene, coefficients from a multiple linear regression are presented in Fig. 7. The monthly mean change in isoprene emission rate, NO_x concentration and O₃ production rate for each grid cell are shown graphically with locations and months of high-NO_x (above the 95th percentile) marked with stars. All three climate models produce coefficients between 0.33 and 0.41 for the relationship between changes in NO_x concentration and O₃ production rate (Fig. 7). However, the change in isoprene emissions using GISS and UKESM1 is a weaker predictor of O₃ production, even though increases in isoprene emission of over 100 % are predicted. All predictors are considered significant due to the large sample size, with r^2 values of 0.384, 0.732 0.590 for

UKESM1, GISS and MRI respectively (Table S3). The lower r^2 value for UKESM1 indicates that the changes in NOx concentration and isoprene emissions explain less than half of the change in O₃ production rate. Additional analysis shows that the O₃ production rate in UKESM1 is also related to the background NOx concentration (see S4: The relationship between NOx and ozone production).




Figure 7: Scatter plots of the monthly mean percentage change in surface NOx concentration, isoprene emission rate and O₃ production rate for each grid cell and each month for (a) UKESM1, (b) GISS and (c) MRI for the region 30° S–30° N, excluding Saharan Africa. Data for MRI has been randomly normally distributed along the y-axis. Grid cells and months where the background NOx concentration is greater than the 95th percentile for the region shown in Fig. 3.1 are marked with stars. The labelled intercept and coefficients refer to the results of a multiple linear regression $\Delta O_3 \text{ prod (\%)} \sim \Delta NOx (\%) + \Delta Isoprene (\%)$ using the plotted data. The second row contains the number of data points in each NOx concentration change range. The data are divided into 50 bins.

GISS simulates increases and decreases in NOx concentration of 50 %, compared to the smaller changes predicted by MRI, which fall mostly in the range 0–20 % (Fig. 7c). GISS therefore predicts decreases in O₃ production over remote regions (Fig. 6b) and seasons, whereas MRI predicts consistent increases (Fig. 6c). Additionally, high-NOx areas simulated by GISS experience an increase in O₃ production regardless of the NOx concentration change (Fig. 7b, stars). In high-NOx areas simulated by UKESM1, the percentage change in NOx concentration is small so there is not enough information to identify individual isoprene and NOx sensitivities, although areas with increased isoprene emission also show increases in O₃ production rate (Fig. 7a, stars).

The apparent relationship between O₃ and isoprene in UKESM1 (Fig. 6) does not show up using a linear model (Fig. 7). A relationship may be hidden by variation in the isoprene–O₃ production sensitivity in different grid cells, or by correlations

between isoprene and NO_x (discussed further in S4: The relationship between NO_x and ozone production). However, as isoprene also contributes to O₃ loss, the effect of isoprene on net chemical production (production – loss) is reduced by the two terms cancelling each other out, so the change in net O₃ chemical production is more clearly related to the change in percentage NO_x concentration (Fig. S7). In particular, the decrease in net O₃ production in the Northern Amazon (Fig. S7d) resembles the percentage change in NO_x concentration (Fig. S7a) more than the percentage change in isoprene emissions (Fig. S7b).

The change in O₃ production rate will be further affected by meteorological changes, temperature in particular. This is the reason that O₃ production increases in UKESM1 and MRI even in the absence of changes in NO_x and isoprene (the intercepts of the linear model are 19 % and 5 % respectively) and O₃ production increases in areas showing decreasing NO_x concentrations in UKESM1. Since the temperature change varies seasonally and regionally, with dry seasons experiencing the largest increase in temperature, some of the changes in O₃ production in Fig. 7 may be driven by temperature rather than NO_x or isoprene changes. If isoprene/NO_x and O₃ production are both influenced by the underlying meteorology, the identified correlations may be due to meteorology rather than the chemical species changes. We verify that the monthly mean temperature change in each gridcell is not significantly correlated with percentage NO_x change in any model, nor percentage isoprene change in UKESM1 (not shown). Therefore, NO_x and isoprene changes are likely controlled by many processes in addition to temperature, including background chemistry and emissions for NO_x, and vegetation type and cover for isoprene, as well as other meteorological variables. This indicates that the identified correlations between NO_x and O₃ production are unlikely to be the result of a spurious relationship driven by temperature, although it is still possible that the strength of the correlations may be inflated by confounding meteorological variables.

4. Discussion

When compared to in situ observations, the three climate models used in this study overestimate present-day surface O₃ in tropical regions by 14 ppb on average, including 11 ppb over the oceans. This is close to the global bias of 16 ppb calculated by Turnock et al. (2020) which included data from six climate models, including the three in this study. Therefore, the sources of error may not be unique to the tropics and subtropics. The major sources of variation between model and observations are related to differences in the area sampled and the heights of the stations relative to the lowest model grid cell (Pacifico et al., 2015).

In the tropics and subtropics, we expect in-canopy deposition and chemical processes to be the most important contributor to the positive bias because these processes create a steep O₃ gradient at the surface, whereas models aim to predict O₃ concentrations at 20+ m from the canopy top where these deposition processes are not included (Stroud et al., 2005; Gordon et al., 2014). Additionally, the volume of the model grid box is many times larger than the area sampled by measurement sites,

and also larger than the area of precursor emission sources such as fires. Therefore, the model inputs and predictions represent the average over a region that is not directly comparable with in situ measurements (Sinha et al., 2004). As a further validation, we also provide data from the TES satellite at 825 hPa, which records higher O₃ concentrations than the in situ sites since it measures at an altitude away from the canopy sink. Although this is much higher in altitude than the lowest grid box of any of the models, it should capture the above-canopy seasonal cycle at a resolution closer to the model grid resolution.

We find that the modelled surface O₃ bias compared to in situ observations is largest in biomass burning areas, although in South America the models capture the seasonal cycle well (Fig. 1). In situ sites, especially in the DR Congo (Fig. 1e), do not detect the large increases in O₃ predicted by models during biomass burning months although observed O₃ concentrations are also highest during biomass burning season (Adon et al., 2013). A high positive bias during the dry season has been found in previous studies (e.g. Turnock et al., 2020) although our study has covered several regions that did not previously have available data. It is likely that effective removal within the canopy that is not included in models softens the observed seasonal cycle. In this region, the trends captured by satellites are closer to the model predictions, which increases confidence that models are correctly identifying O₃ enhancements above the canopy due to fires. However, future studies assessing the risks to human and ecosystem health should be aware of this limitation in current models.

The remainder of the study focuses on the change in surface O₃ due to climate change. Although model biases increase uncertainty in the change due to climate change, we quantify the difference between two simulations, which should remove systematic biases, and we note that the models capture seasonal and regional trends that are explained by either in situ measurements or satellite measurements (Fig. 1). This gives confidence in the trends in future O₃ change presented in this study, but we highlight O₃ from biomass burning as an area for further study. The models have different chemistry schemes, land processes and temperature sensitivities which contribute to model variation (Stevenson et al., 2006; Wu et al., 2007; Archibald et al., 2020b). For this reason, we do not attempt to completely diagnose reasons for inter-model variation and instead the aim of this study is to identify robust predictions and areas of uncertainty for the change in O₃ due to climate change.

We find that while overall O₃ concentrations over the tropical land areas are reduced under the climate scenario examined here, climate change could lead to an ozone–climate penalty in areas which have a high background NO_x concentration. These high-NO_x areas already tend to have high O₃ concentrations in the absence of climate change (above 40 ppb), with climate change causing a further deterioration in air quality. Models predict that climate change will lead to seasonal mean increases in surface O₃ concentration of up to 12 ppb in tropical and subtropical areas with high-NO_x emissions (Fig. 4). The increase in surface O₃ in high-NO_x areas is robust, with seasonal mean increases of up to 15 ppb for UKESM1, 18 ppb for GISS and 12 ppb for MRI. These areas are defined by NO_x emission magnitudes above the 95th percentile for the region 40° S–40° N, which is dominated by anthropogenic contributions such as biomass burning or urban emissions. O₃ pollution in forested areas

has the potential to reduce forest productivity, decreasing the amount of carbon removed from the atmosphere and impairing forest resilience (e.g. Sitch et al., 2007; Grulke et al., 2020).

575 The ozone–climate penalty in high-NO_x regions is primarily driven by an increase in O₃ chemical production, which is largest in areas of high-NO_x (Figs 5, 6). This is in agreement with results from Doherty et al. (2013) and Archibald et al. (2020b) who showed that the rate of change of O₃ with temperature increases with NO_x concentration. Firstly, the major O₃ forming reaction NO + HO₂/RO₂, happens faster at higher temperatures and scaling up O₃ production in an area where O₃ production is already high will often lead to greater O₃ increases than an area with low O₃ precursor concentrations (Coates et al. 2016). Secondly, 580 VOC and NO_x concentrations can increase at higher temperatures. NO_x concentration can increase due to increased PAN decomposition at higher temperatures (Doherty et al., 2013), changes in lightning frequency, or changes to atmospheric chemistry and VOCs increase largely as a result of increased isoprene emissions. All models likely exhibit an increase in reaction rates, however there were differences between UKESM1 and the other models in their sensitivity to changes in NO_x concentration and isoprene emissions (Fig. 7).

585 Using a multiple linear regression, we find that changes in NO_x concentration are strongly correlated with changes in O₃ production rate for GISS and MRI ($r^2 = 0.732$ and 0.590 respectively) (Fig. 7). For UKESM1, we find that linear regression using changes in NO_x concentration and isoprene emission explains less than 50 % of the change in O₃ production rate. Including background NO_x concentration in the linear regression improves the r^2 value and suggests that the rate of O₃ 590 production increases in proportion to the background NO_x concentration (Fig. S5). UKESM1 has previously been identified as being among the least responsive to changes in precursor concentrations out of the CMIP6 models (Turnock et al., 2020), and indeed chemical production increases in many areas despite decreases in NO_x, so the increase in chemical O₃ production is more likely to be dominated by an increase in rate of reaction not changes in precursor concentration. The linear regression uses monthly means, so modelled O₃ increases during burning seasons (dry seasons) are likely compounded by the fact that 595 these seasons often show the greatest temperature increase due to climate change (Fig. S3). Therefore, some of the identified correlations may be due to meteorology changes rather than chemical changes.

As changes in NO_x concentrations are shown to be important for changes in O₃ production in GISS and MRI, we now discuss the inter-model differences in NO_x concentration changes in further detail. GISS and MRI agree that NO_x concentrations will 600 increase in high-NO_x regions, but disagree on the direction of change in remote regions. UKESM1 predicts a decrease in NO_x concentrations in many areas. GISS predicts a decrease in NO_x of 2×10^{-11} mol mol⁻¹ in remote areas (up to 50 %) while MRI predicts an increase of 2×10^{-11} to 6×10^{-11} mol mol⁻¹ in remote regions. This is likely causing the difference in O₃ production between GISS and MRI in remote regions. To reduce uncertainty in predictions for O₃ concentration changes due to climate change, further work to constrain future NO_x concentration changes is needed.

605

The NO_x concentration change depends on the balance of NO_x production and loss terms. A large contributor to increases in NO_x concentrations in all models is an increased decomposition of PAN into NO_x, which will be largest in source regions. Lightning NO_x is also a NO_x source, but its influence on surface NO_x remains unclear. Although lightning NO_x increases in all models during the wet season, the largest surface NO_x and O₃ increases occur in the dry season, so the ozone–climate penalty is unlikely to be driven by lightning NO_x changes. Nevertheless, the large increase in lightning NO_x in MRI may have a role in the increase in surface NO_x concentration in MRI, which is larger than the other models, and lightning NO_x decreases in the Northern Amazon during the dry season (Fig. S4a) may contribute to the decrease in NO_x and O₃ production in this region in UKESM1 (Fig. S7). A large contributor to the loss term will be reaction with isoprene derivatives, thus increased formation of isoprene nitrates. In both UKESM1 and GISS, isoprene and NO_x are anti-correlated in some areas, suggesting isoprene emissions changes have a notable effect on NO_x concentrations. For example, GISS predicts large isoprene increases in the remote Amazon, where the major NO_x decreases occur, and UKESM1 shows a small area of increased NO_x concentrations downwind of the Northern Amazon, where isoprene decreases. As MRI prescribes isoprene as a climatology, there is will be no significant change to NO_x loss via organic nitrate formation and this is likely a reason why NO_x increases over most areas of land. The reasons why NO_x loss dominates in UKESM1 whereas GISS shows a net NO_x increase requires further understanding of individual model details such as isoprene nitrate yield and NO_x recycling frequency. The sensitivity of NO_x concentration to changes in lightning, PAN and isoprene in each model is beyond the scope of this study, but would be useful to explore in further work. Further studies could also explore some temperature-sensitive sources of NO_x that were not included in the simulations such as soil NO_x emissions and changes in wildfire frequency.

NO_x is not the only driver of changes in O₃ production, and changes in temperature-dependent emissions of isoprene also influence the rate. This may be especially important in UKESM1, which predicts the highest concentration of background NO_x, because high-NO_x areas may experience a different chemical regime in which VOCs also increase O₃ production (e.g. Liu et al., 2013). For example, UKESM1 shows increases in O₃ production rate as isoprene increases in high-NO_x areas (Fig. 7, stars). However, calculating the net effect of isoprene on surface O₃ concentrations is complex; increasing isoprene emissions can increase both the rate of O₃ production and the rate of O₃ loss and, as described above, can decrease concentrations of NO_x and OH. Examining the percentage change in net O₃ chemical production in UKESM1 (Fig. S7) suggests the net effect of changing isoprene emissions on O₃ chemistry cancel out, and that the percentage change in net O₃ production is more closely related to percentage changes in NO_x concentration.

The increase in chemical loss is strongly correlated with isoprene concentration change in UKESM1 and GISS (Fig. S8). Therefore, the different isoprene schemes used by each model contributes to uncertainty in the loss rate over the continents. In particular, MRI used climatological isoprene resulting in no significant change in the loss rate (Fig. S8c) and UKESM1 includes CO₂ inhibition, which decreases the isoprene emission rate and loss rate in in Northern Amazonia (Fig. S8a). Overall,

this means that GISS has a higher loss rate, especially in low-NO_x, high isoprene regions such as the remote Amazon, which
640 may partly account for the larger decreases in O₃ in remote regions using this model (Fig. 3).

The global study by Zanis et al. (2022) employs two additional climate models and also finds climate benefits and uncertainties
in surface O₃ concentrations in the same remote regions. We excluded these two models from our own study as data for the
sensitivity study (NO_x concentration and isoprene emission rate) were not available at the time of writing. Zanis et al. (2022)
645 highlight the different isoprene emission schemes as a reason for model variation however our analysis (Fig. 7) finds NO_x to
be the most important precursor. The fact that the models contain positive correlations between the change in NO_x and O₃
production, and between the change in isoprene and O₃ loss, indicates the tropics and subtropics exhibit NO_x-limited
behaviour, although isoprene may be important in high-NO_x areas. We agree that isoprene is highly relevant for the change in
loss rate and for indirect effects on O₃ through changes in related atmospheric chemistry such as OH and NO_x concentrations.

650

The decrease in deposition rate is controlled by UKESM1 and GISS (MRI showed very little change), but there was spatial
variation in the magnitude of the change. This could be due to changes in meteorology between models (such as temperature
and precipitation), as well as model differences (UKESM1 includes CO₂ inhibition). Feedbacks such as O₃ damage to
vegetation were not considered in any model (Pacifico et al., 2015) but may be a useful addition to future simulations.

655

Models tend to predict a decrease in surface O₃ over regions strongly affected by ocean air such as North Brazil. This is due
to robust decreases in O₃ over the oceans from increases in atmospheric water vapour. Over land, increases in water vapour
and OH influence the concentrations and lifetimes of many species.

660 We finally note that Nigeria experiences substantial increases in O₃ production according to GISS and MRI, whilst a slight
decrease is predicted using UKESM1. This is an important geographical area for future research since poor air quality could
affect large numbers of people living in West African cities. In this area, the choice of emissions scenario is also important for
determining the O₃ response to climate change because NO_x emissions in Nigeria increase rapidly in the SSP3-7.0 scenario
compared to the present-day due to predicted urbanisation. Therefore, future studies should explore alternative emissions
665 pathways to better inform policy.

5. Conclusion

Using a multimodel mean of data from three Earth system models, we identify that by 2100, there will be an ozone–climate
penalty in high-NO_x areas, such as major cities and biomass burning areas (Fig. 4). This is not due to increased fire emissions,
but due to the increasing temperature, which speeds up the recycling of NO into NO_x and increases decomposition of PAN
670 into NO_x in source regions. It shows that the ozone–climate penalty is greatest in areas already experiencing high O₃, putting

forests in these areas at greater risk of O₃ damage and urban populations at increasing threat of health problems. This study adds to findings from the World Health Organisation World Air Quality Report (2021) that air pollution is an increasing issue across the tropics, and that there is a need for greater monitoring of air pollution across Africa and South America.

675 The Earth system models display NO_x-limited behaviour, including that higher NO_x concentrations lead to increased O₃ chemical production and therefore increased surface O₃ concentration (Figs 5–7). As the background concentrations of NO_x are largely anthropogenic, this suggests that without reduction in emissions, forested areas in urban and fire-prone locations are more at risk from increases in surface O₃ due to climate change than remote forests. As O₃ damage can reduce plant productivity, this has implications for the success of secondary forests and other human-modified forests which are mostly
680 located in agricultural areas, deforestation frontiers and forest edges (Heinrich et al., 2021), and may reduce their carbon sequestration potential (Sitch et al., 2007).

In remote regions, differences in the direction of O₃ concentration change between models creates uncertainty as to whether remote locations are at greater risk of O₃ damage in a warmer climate, although ocean influenced areas display robust climate
685 benefits (Fig. 4). Further work is needed to constrain the climate response of isoprene emissions and the temperature sensitivity of NO_x and O₃ chemistry.

Data availability statement

All CMIP6 model data used in the present study can be obtained from <https://esgf-node.llnl.gov/search/cmip6/>.

All TOAR I data used in the study can be obtained from <https://join.fz-juelich.de/services/rest/surfacedata/>.

690 All INDAAF data used in this study can be obtained from <http://www.indAAF.obs-mip.fr>

Author contributions

FB wrote the paper and led the data analysis with contributions from all authors. SS and GAF contributed to the interpretation of the data. MB contributed to the statistical analysis. IDS, HV, PB, MB and CGL contributed in situ data for model evaluation. SEB and KT contributed in the GISS-E2-1-G simulations; MD and NO contributed in the MRI-ESM2-0 simulations; JK and
695 FMO contributed in the UKESM1-0-LL simulations.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

FB was funded by the NERC GW4+ DTP (award no. NE/S007504/1) and the Met Office on a CASE studentship. LMM
acknowledges funding from the UK Natural Environment Research Council funding (UK Earth System Modelling Project,
UKESM, Grant no. NE/N017951/1). SS, LMM and AWC were supported by NERC funding (Grant no. NE/R001812/1). MD
and NO were supported by the Japan Society for the Promotion of Science KAKENHI (Grant no. JP18H03363, JP18H05292,
JP19K12312, JP20K04070 and JP21H03582), the Environment Research and Technology Development Fund
(JPMEERF20202003 and JPMEERF20205001) of the Environmental Restoration and Conservation Agency of Japan, the
Arctic Challenge for Sustainability II (ArCS II), Program Grant Number JPMXD1420318865, and a grant for the Global
Environmental Research Coordination System from the Ministry of the Environment, Japan (MLIT1753). JK was financially
supported by NERC through NCAS (Grant no. R8/H12/83/003), and the Met Office CSSP-China programme funded
POZsUM project. FMO'C and GAF were supported by the Met Office Hadley Centre Climate Programme funded by BEIS
and the EU Horizon 2020 Research Programme CRESCENDO project (Grant no. 641816). Resources supporting
the GISS simulations were provided by the NASA High-End Computing (HEC) Program through the NASA Center for
Climate Simulation (NCCS) at the Goddard Space Flight Center. GISS authors, SEB and KT acknowledge funding from the
NASA Modeling and Analysis program. IV was funded by the Fonds Wetenschappelijk Onderzoek Flanders (FWO; grant no.
G018319N). The INDAAF project is funded by the INSU/CNRS/IRD and is part of the ACTRIS-FR Research Infrastructure.
The authors are grateful to the field and chemistry analytical technicians who operate the African sites and help to provide
reliable data.

References

- Adon, M., Galy-Lacaux, C., Yoboué, V., Delon, C., Lacaux, J. P., Castera, P., Gardrat, E., Pienaar, J., Al Ourabi, H., Laouali,
D., Diop, B., Sigha-Nkamdjou, L., Akpo, A., Tathy, J. P., Lavenu, F., and Mougin, E.: Long term measurements of
sulfur dioxide, nitrogen dioxide, ammonia, nitric acid and ozone in Africa using passive samplers, *Atmos. Chem. Phys.*,
10, 7467–7487, <https://doi.org/10.5194/acp-10-7467-2010>, 2010.
- Adon, M., Galy-Lacaux, C., Delon, C., Yoboue, V., Solmon, F., & Kaptue Tchuenta, A. T.: Dry deposition of nitrogen
compounds (NO₂, HNO₃, NH₃), sulfur dioxide and O₃ in west and central African ecosystems using the inferential
method. *Atmospheric Chemistry and Physics*, 13, 11351–11374. <https://doi.org/10.5194/acp-13-11351-2013>, 2013
- Agathokleous, E., Belz, R. G., Calatayud, V., de Marco, A., Hoshika, Y., Kitao, M., Saitanis, C. J., Sicard, P., Paoletti, E., &
Calabrese, E. J.: Predicting the effect of O₃ on vegetation via linear non-threshold (LNT), threshold and hormetic dose-
response models. *Sci. Total Environ.*, 649, 61–74. <https://doi.org/10.1016/j.scitotenv.2018.08.264>, 2019
- Archibald, A. T., Jenkin, M. E., & Shallcross, D. E.: An isoprene mechanism intercomparison. *Atmos. Environ.*, 44, 5356–
5364. <https://doi.org/10.1016/j.atmosenv.2009.09.016>, 2010

- Archibald, A. T., O'Connor, F. M., Abraham, N. L., Archer-Nicholls, S., Chipperfield, M. P., Dalvi, M., Folberth, G. A.,
 730 Dennison, F., Dhomse, S. S., Griffiths, P. T., Hardacre, C., Hewitt, A. J., Hill, R. S., Johnson, C. E., Keeble, J., Köhler,
 M., Morgenstern, O., Mulcahy, J. P., Ordóñez, C., ... Zeng, G.: Description and evaluation of the UKCA stratosphere-
 troposphere chemistry scheme (StratTrop vn 1.0) implemented in UKESM1. *Geosci. Model Dev.*, *13*, 1223–1266.
<https://doi.org/10.5194/gmd-13-1223-2020>, 2020a
- Archibald, A. T., Turnock, S. T., Griffiths, P. T., Cox, T., Derwent, R. G., Knote, C., & Shin, M.: On the changes in surface
 735 O₃ over the twenty-first century: sensitivity to changes in surface temperature and chemical mechanisms: 21st century
 changes in surface O₃. *Philos. Trans. Royal Soc. A*, *378*. <https://doi.org/10.1098/rsta.2019.0329>, 2020b
- Bates, K. H. and Jacob, D. J.: A new model mechanism for atmospheric oxidation of isoprene: global effects on oxidants,
 nitrogen oxides, organic products, and secondary organic aerosol, *Atmos. Chem. Phys.*, *19*, 9613–9640,
<https://doi.org/10.5194/acp-19-9613-2019>, 2019.
- 740 Bela, M. M., Longo, K. M., Freitas, S. R., Moreira, D. S., Beck, V., Wofsy, S. C., Gerbig, C., Wiedemann, K., Andreae, M.
 O., & Artaxo, P.: O₃ production and transport over the Amazon Basin during the dry-to-wet and wet-to-dry transition
 seasons. *Atmos. Chem. Phys.*, *15*, 757–782. <https://doi.org/10.5194/acp-15-757-2015>, 2015
- Bonan, G. B.: Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests. *Science*, *320*(5882),
 1444–1449. <https://doi.org/10.1126/science.1155121>, 2008
- 745 Bond, D. W., Steiger, S., Zhang, R., Tie, X., & Orville, R. E.: The importance of NO_x production by lightning in the tropics.
Atmos. Environ., *36*, 1509–1519. [https://doi.org/https://doi.org/10.1016/S1352-2310\(01\)00553-2](https://doi.org/10.1016/S1352-2310(01)00553-2), 2002
- Chadwick, R., Douville, H., & Skinner, C. B.: Timeslice experiments for understanding regional climate projections:
 applications to the tropical hydrological cycle and European winter circulation. *Clim. Dyn.*, *49*, 3011–3029.
<https://doi.org/10.1007/s00382-016-3488-6>, 2017
- 750 Clark, S. K., Ward, D. S., & Mahowald, N. M.: Parameterization-based uncertainty in future lightning flash density. *Geophys.*
Res. Lett., *44*, 2893–2901. <https://doi.org/10.1002/2017GL073017>, 2017
- Clifton, O. E., Fiore, A. M., Massman, W. J., Baublitz, C. B., Coyle, M., Emberson, L., Fares, S., Farmer, D. K., Gentine, P.,
 Gerosa, G., Guenther, A. B., Helmig, D., Lombardozzi, D. L., Munger, J. W., Patton, E. G., Pusede, S. E., Schwede, D.
 B., Silva, S. J., Sörgel, M., Steiner, A., & Tai, A. P. K.: Dry Deposition of O₃ Over Land: Processes, Measurement, and
 755 Modeling. *Rev. Geophys.*, *58*. <https://doi.org/10.1029/2019RG000670>, 2020
- Coates, J., Mar, K. A., Ojha, N., & Butler, T. M.: The influence of temperature on O₃ production under varying NO_x conditions
 - A modelling study. *Atmos. Chem. Phys.*, *16*, 11601–11615. <https://doi.org/10.5194/acp-16-11601-2016>, 2016
- Cox, P. M., Betts, R. A., Collins, M., Harris, P. P., Huntingford, C., & Jones, C. D.: Amazonian forest dieback under climate-
 carbon cycle projections for the 21st century. *Theor. Appl. Climatol.*, *78*, 137–156. [https://doi.org/10.1007/s00704-004-](https://doi.org/10.1007/s00704-004-0049-4)
 760 [0049-4](https://doi.org/10.1007/s00704-004-0049-4), 2004

- Deushi, M., & Shibata, K.: Impacts of increases in greenhouse gases and O₃ recovery on lower stratospheric circulation and the age of air: Chemistry-climate model simulations up to 2100. *J. Geophys. Res.*, *116*, 7107. <https://doi.org/10.1029/2010JD015024>, 2011
- 765 Doherty, R. M., Wild, O., Shindell, D. T., Zeng, G., MacKenzie, I. A., Collins, W. J., Fiore, A. M., Stevenson, D. S., Dentener, F. J., Schultz, M. G., Hess, P., Derwent, R. G., & Keating, T. J.: Impacts of climate change on surface O₃ and intercontinental O₃ pollution: A multi-model study. *J. Geophys. Res. Atmos.*, *118*, 3744–3763. <https://doi.org/10.1002/jgrd.50266>, 2013
- Emberson, L.: Effects of O₃ on agriculture, forests and grasslands. *Philos. Trans. Royal Soc.*, *378*. <https://doi.org/10.1098/rsta.2019.0327>, 2000
- 770 Emberson, L., Kitwiroon, N., Beevers, S., Büker, P., & Cinderby, S.: Scorched earth: How will changes in the strength of the vegetation sink to O₃ deposition affect human health and ecosystems? *Atmos. Chem. Phys.*, *13*, 6741–6755. <https://doi.org/10.5194/acp-13-6741-2013>, 2013
- Finney, D. L., Doherty, R. M., Wild, O., Stevenson, D. S., Mackenzie, I. A., & Blyth, A. M.: A projected decrease in lightning under climate change. *Nat. Clim. Change.*, *8*, 210–213. <https://doi.org/10.1038/s41558-018-0072-6>, 2018
- 775 Franz, M., & Zaehle, S.: Competing effects of nitrogen deposition and O₃ exposure on Northern hemispheric terrestrial carbon uptake and storage, 1850–2099. *Biogeosci. Discuss.*, 1–42. <https://doi.org/10.5194/bg-2020-443>, 2020
- Fu, T. M., & Tian, H.: Climate Change Penalty to O₃ Air Quality: Review of Current Understandings and Knowledge Gaps. *Curr. Poll. Rep.*, *5*, 159–171. <https://doi.org/10.1007/s40726-019-00115-6>, 2019
- Fu, Y., & Liao, H.: Biogenic isoprene emissions over China: sensitivity to the CO₂ inhibition effect. *Atmos. Ocean Sci. Lett.*, *9*, 277–284. <https://doi.org/10.1080/16742834.2016.1187555>, 2016
- 780 Gordon, M., Vlasenko, A., Staebler, R. M., Stroud, C., Makar, P. A., Liggio, J., Li, S. M., & Brown, S.: Uptake and emission of VOCs near ground level below a mixed forest at Borden, Ontario. *Atmos. Chem. Phys.*, *14*, 9087–9097. <https://doi.org/10.5194/acp-14-9087-2014>, 2014
- Griffiths, P. T., Murray, L. T., Zeng, G., Shin, Y. M., Abraham, N. L., Archibald, A. T., Deushi, M., Emmons, L. K., Galbally, I. E., Hassler, B., Horowitz, L. W., Keeble, J., Liu, J., Moeini, O., Naik, V., O'Connor, F. M., Oshima, N., Tarasick, D., Tilmes, S., Turnock, S. T., Wild, O., Young, P. J., and Zanis, P.: Tropospheric ozone in CMIP6 simulations, *Atmos. Chem. Phys.*, *21*, 4187–4218, <https://doi.org/10.5194/acp-21-4187-2021>, 2021.
- 785 Guenther, A. B., Zimmerman, P. R., Harley, P. C., & Monson, R. K.: Isoprene and Monoterpene Emission Rate Variability? Model Evaluations and Sensitivity Analyses. *J. Geophys. Res. Atmos.*, *98*, 12609–12621. <https://doi.org/https://doi.org/10.1029/93JD00527>, 1993
- 790 Guenther, A., Nicholas Hewitt, ! C, Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., Mckay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., & Zimmerman, P.: A global model of natural volatile organic compound emissions. *J. Geophys. Res. Atmos.*, *100*, 8873–8892. <https://doi.org/https://doi.org/10.1029/94JD02950>, 1995

- 795 Hantson, S., Knorr, W., Schurgers, G., Pugh, T. A. M., & Arneth, A.: Global isoprene and monoterpene emissions under
changing climate, vegetation, CO₂ and land use. *Atmos. Environ.*, *155*, 35–45.
<https://doi.org/10.1016/j.atmosenv.2017.02.010>, 2017
- Heinrich, V. H. A., Dalagnol, R., Cassol, H. L. G., Rosan, T. M., Torres De Almeida, C., Silva Junior, C. H. L., Campanharo,
W. A., House, J. I., Stith, S., Hales, T. C., Adami, M., Anderson, L. O., Luiz, &, & Aragão, E. O. C.: Large carbon sink
800 potential of secondary forests in the Brazilian Amazon to mitigate climate change. *Nat. Commun.*, *12*, 1–11.
<https://doi.org/10.1038/s41467-021-22050-1>, 2021
- Jacob, D. J., & Winner, D. A. (2009). Effect of climate change on air quality. *Atmos. Environ.*, *43*, 51–63.
<https://doi.org/10.1016/j.atmosenv.2008.09.051>
- Jacobs, C. M. J.: Direct impact of atmospheric CO₂ enrichment on regional transpiration. *Wageningen University and*
805 *Research*. <https://doi.org/10.5194/gmd-12-2875-2019>, 1994
- Kawai, H., Yukimoto, S., Koshiro, T., Oshima, N., Tanaka, T., Yoshimura, H., and Nagasawa, R.: Significant improvement
of cloud representation in the global climate model MRI-ESM2, *Geosci. Model Dev.*, *12*, 2875–2897,
<https://doi.org/10.5194/gmd-12-2875-2019>, 2019.
- Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M.,
810 Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S.,
Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews,
E. E., McDermid, S., Mezuman, K., Miller, R. L., Murray, L. T., Oinas, V., Orbe, C., Pérez García-Pando, C., Perlwitz,
J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng,
E., Wu, J., and Yao, M.-S.: GISS-E2.1: Configurations and Climatology, *J. Adv. Model. Earth Sy.*, *12*,
815 e2019MS002025, <https://doi.org/10.1029/2019MS002025>, 2020.
- Kuhn, U., Ganzeveld, L., Thielmann, A., Dindorf, T., Schebeske, G., Welling, M., Sciare, J., Roberts, G., Meixner, F. X.,
Kesselmeier, J., Lelieveld, J., Kolle, O., Ciccioli, P., Lloyd, J., Trentmann, J., Artaxo, P., & Andreae, M. O.: Impact of
Manaus City on the Amazon Green Ocean atmosphere: O₃ production, precursor sensitivity and aerosol load. *Atmos.*
Chem. Phys., *10*, 9251–9282. <https://doi.org/10.5194/acp-10-9251-2010>, 2010
- 820 Leuning, R.: A critical appraisal of a combined stomatal-photosynthesis model for C₃ plants. *Plant, Cell Environ.*, *18*, 339–
355. <https://doi.org/https://doi.org/10.1111/j.1365-3040.1995.tb00370.x>, 1995
- Lewis, S. L.: Tropical forests and the changing earth system. *Philos. Trans. Royal Soc. B*, *361*, 195–210.
<https://doi.org/10.1098/rstb.2005.1711>, 2006
- Lin, M., Horowitz, L. W., Xie, Y., Paulot, F., Malyshev, S., Shevliakova, E., Finco, A., Gerosa, G., Kubistin, D., & Pilegaard,
825 K.: Vegetation feedbacks during drought exacerbate O₃ air pollution extremes in Europe. *Nat. Clim. Change*, *10*, 444–
451. <https://doi.org/10.1038/s41558-020-0743-y>, 2020a

- Liu, W., Hegglin, M. I., Checa-Garcia, R., Li, S., Gillett, N. P., Lyu, K., Zhang, X., & Swart, N. C.: Stratospheric ozone depletion and tropospheric ozone increases drive Southern Ocean interior warming. *Nat. Clim. Change*, 12, 365–372. <https://doi.org/10.1038/s41558-022-01320-w>, 2022
- 830 Liu, Y., Brito, J., Dorris, M. R., Rivera-Rios, J. C., Seco, R., Bates, K. H., Artaxo, P., Duvoisin, S., Keutsch, F. N., Kim, S., Goldstein, A. H., Guenther, A. B., Manzi, A. O., Souza, R. A. F., Springston, S. R., Watson, T. B., McKinney, K. A., & Martin, S. T.: Isoprene photochemistry over the Amazon rainforest. *Proc. Natl. Acad. Sci U.S.A.*, 113, 6125–6130. <https://doi.org/10.1073/pnas.1524136113>, 2016
- 835 Liu, Z., Doherty, R. M., Wild, O., Hollaway, M., & O'Connor, F. M.: Contrasting chemical environments in summertime for atmospheric O₃ across major Chinese industrial regions: the effectiveness of emission control strategies. *Atmospheric Chemistry and Physics*, 21, 10689–10706. <https://doi.org/10.5194/acp-21-10689-2021>, 2021
- Malhi, Y., Aragão O A, L. E. O. C., Galbraith, D., Huntingford, C., Fisher, R., Zelazowski, P., Sitch, S., Mcsweeney, C., & Meir, P.: Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest. *Proc. Natl. Acad. Sci.*, 106, 20610–20615. <https://doi.org/10.1073/pnas.0804619106>, 2009
- 840 Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500, *Geosci. Model Dev.*, 13, 3571–3605, <https://doi.org/10.5194/gmd-13-3571-2020>, 2020.
- 845 Morfopoulos, C., Müller, J. F., Stavrakou, T., Bauwens, M., de Smedt, I., Friedlingstein, P., Prentice, I. C., & Regnier, P.: Vegetation responses to climate extremes recorded by remotely sensed atmospheric formaldehyde. *Glob. Change. Biol.*, 28, 1809–1822. <https://doi.org/10.1111/gcb.15880>, 2021
- Mulcahy, J. P., Jones, C., Sellar, A., Johnson, B., Boutle, I. A., Jones, A., Andrews, T., Rumbold, S. T., Mollard, J., Bellouin, N., Johnson, C. E., Williams, K. D., Grosvenor, D. P., & McCoy, D. T. Improved Aerosol Processes and Effective
850 Radiative Forcing in HadGEM3 and UKESM1. *J. Adv. Model. Earth Syst.*, 10, 2786–2805. <https://doi.org/10.1029/2018MS001464>, 2018
- Mulcahy, J. P., Johnson, C., Jones, C. G., Povey, A. C., Scott, C. E., Sellar, A., Turnock, S. T., Woodhouse, M. T., Abraham, N. L., Andrews, M. B., Bellouin, N., Browse, J., Carslaw, K. S., Dalvi, M., Folberth, G. A., Glover, M., Grosvenor, D. P., Hardacre, C., Hill, R., Johnson, B., Jones, A., Kipling, Z., Mann, G., Mollard, J., O'Connor, F. M., Palmiéri, J.,
855 Reddington, C., Rumbold, S. T., Richardson, M., Schutgens, N. A. J., Stier, P., Stringer, M., Tang, Y., Walton, J., Woodward, S., and Yool, A.: Description and evaluation of aerosol in UKESM1 and HadGEM3-GC3.1 CMIP6 historical simulations, *Geosci. Model Dev.*, 13, 6383–6423, <https://doi.org/10.5194/gmd-13-6383-2020>, 2020.
- Murray, L. T.: Lightning NO_x and Impacts on Air Quality. *Curr. Poll. Rep.*, 2, 115–133. <https://doi.org/10.1007/s40726-016-0031-7>, 2016

- 860 Myhre, G., Aas, W., Cherian, R., Collins, W., Faluvegi, G., Flanner, M., Forster, P., Hodnebrog, Ø., Klimont, Z., Lund, M. T.,
Mülmenstädt, J., Lund Myhre, C., Olivié, D., Prather, M., Quaas, J., Samset, B. H., Schnell, J. L., Schulz, M., Shindell,
D., Skeie, R. B., Takemura, T., and Tsyro, S.: Multi-model simulations of aerosol and ozone radiative forcing due to
anthropogenic emission changes during the period 1990–2015, *Atmos. Chem. Phys.*, *17*, 2709–2720,
<https://doi.org/10.5194/acp-17-2709-2017>, 2017.
- 865 Niinemets U., Tenhunen J.D., Harley P.C., & Steinbrecher R.: A model of isoprene emission based on energetic requirements
for isoprene synthesis and leaf photosynthetic properties for Liquidambar and Quercus. *Plant, Cell Environ.*, *22*, 1319–
1335. [https://doi.org/https://doi.org/10.1046/j.1365-3040.1999.00505.x](https://doi.org/10.1046/j.1365-3040.1999.00505.x), 1999
- Oshima, N., Yukimoto, S., Deushi, M., Koshiro, T., Kawai, H., Tanaka, T. Y., & Yoshida, K.: Global and Arctic effective
radiative forcing of anthropogenic gases and aerosols in MRI-ESM2. 0. *Progress in Earth and Planetary Science*, *7*, 1–
870 21. <https://doi.org/10.1186/s40645-020-00348-w>, 2020
- Ossohou, M., Galy-Lacaux, C., Yoboué, V., Hickman, J. E., Gardrat, E., Adon, M., Darras, S., Laouali, D., Akpo, A., Ouafu,
M., Diop, B., & Opepa, C.: Trends and seasonal variability of atmospheric NO₂ and HNO₃ concentrations across three
major African biomes inferred from long-term series of ground-based and satellite measurements. *Atmos. Environ.*, *207*,
148–166. <https://doi.org/10.1016/j.atmosenv.2019.03.027>, 2019
- 875 Oswald, E. M., Dupigny-Giroux, L. A., Leibensperger, E. M., Poirot, R., & Merrell, J.: Climate controls on air quality in the
Northeastern U.S.: An examination of summertime O₃ statistics during 1993–2012. *Atmos. Environ.*, *112*, 278–288.
<https://doi.org/10.1016/j.atmosenv.2015.04.019>, 2015
- Pacifico, F., Harrison, S. P., Jones, C. D., Arneth, A., Sitch, S., Weedon, G. P., Barkley, M. P., Palmer, P. I., Serça, D.,
Potosnak, M., Fu, T. M., Goldstein, A., Bai, J., & Schurgers, G.: Evaluation of a photosynthesis-based biogenic isoprene
880 emission scheme in JULES and simulation of isoprene emissions under present-day climate conditions. *Atmos. Chem.*
Phys., *11*, 4371–4389. <https://doi.org/10.5194/acp-11-4371-2011>, 2011
- Pacifico, F., Folberth, G. A., Jones, C. D., Harrison, S. P., & Collins, W. J.: Sensitivity of biogenic isoprene emissions to past,
present, and future environmental conditions and implications for atmospheric chemistry. *J. Geophys. Res. Atmos.*, *117*.
<https://doi.org/10.1029/2012JD018276>, 2012
- 885 Pacifico, F., Folberth, G. A., Sitch, S., Haywood, J. M., Rizzo, L. v., Malavelle, F. F., & Artaxo, P.: Biomass burning related
O₃ damage on vegetation over the Amazon forest: A model sensitivity study. *Atmos. Chem. Phys.*, *15*, 2791–2804.
<https://doi.org/10.5194/acp-15-2791-2015>, 2015
- Pascoe, C., Lawrence, B. N., Guilyardi, E., Jukes, M., & Taylor, K. E.: Designing and Documenting Experiments in CMIP6.
Geosci. Model Dev. Discuss., *10*, 1–27. <https://doi.org/10.5194/gmd-2019-98>, 2019
- 890 Paulot, F., Henze, D. K., & Wennberg, P. O.: Impact of the isoprene photochemical cascade on tropical O₃. *Atmos. Chem.*
Phys., *12*, 1307–1325. <https://doi.org/10.5194/acp-12-1307-2012>, 2012
- Price, C., & Rind, D.: A Simple Lightning Parameterization for Calculating Global Lightning Distributions. *J. Geophys. Res.*,
97, 9919–9933. [https://doi.org/https://doi.org/10.1029/92JD00719](https://doi.org/10.1029/92JD00719), 2018

- Prosperi, P., Bloise, M., Tubiello, F. N., Conchedda, G., Rossi, S., Boschetti, L., Salvatore, M., & Bernoux, M.: New estimates
895 of greenhouse gas emissions from biomass burning and peat fires using MODIS Collection 6 burned areas. *Clim. Change*,
161, 415–432. <https://doi.org/10.1007/s10584-020-02654-0>, 2020
- Pugh, T. A. M., Mackenzie, A. R., Hewitt, C. N., Langford, B., Edwards, P. M., Furneaux, K. L., Heard, D. E., Hopkins, J. R.,
Jones, C. E., Karunaharan, A., Lee, J., Mills, G., Misztal, P., Moller, S., Monks, P. S., & Whalley, L. K.: Simulating
atmospheric composition over a South-East Asian tropical rainforest: performance of a chemistry box model. *Atmos.*
900 *Chem. Phys.*, *10*, 279–298. <https://doi.org/https://doi.org/10.5194/acp-10-279-2010>, 2010
- Rao, S., Klimont, Z., Smith, S. J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K., Amann, M., Bodirsky, B. L., van
Vuuren, D. P., Reis, L. A., Calvin, K., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M.,
Hasegawa, T., Heyes, C., Hilaire, J., Luderer, G., Masui, T., Stehfest, E., Strefler, J., van der Sluis, S., and Tavoni, M.:
Future air pollution in the Shared Socio-economic Pathways, *Global Environ. Chang.*, *42*, 346–
905 358, <https://doi.org/10.1016/j.gloenvcha.2016.05.012>, 2017.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko,
O., Lutz, W., Popp, A., Cuaresma, J. C., K.-C. S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K.,
Hasegawa, T., Havlik, P., Humpenöder, F., Silva, L. A. D., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D.,
Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L.,
910 Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni,
M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An
overview, *Global Environ. Change*, *42*, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- Romer, P. S., Duffey, K. C., Wooldridge, P. J., Edgerton, E., Baumann, K., Feiner, P. A., Miller, D. O., Brune, W. H., Koss,
A. R., de Gouw, J. A., Misztal, P. K., Goldstein, A. H., & Cohen, R. C.: Effects of temperature-dependent NO_x emissions
915 on continental O₃ production. *Atmos. Chem. Phys.*, *18*, 2601–2614. <https://doi.org/10.5194/acp-18-2601-2018>, 2018
- Sadiq, M., Tai, A. P. K., Lombardozzi, D., & Martin, M. V.: Effects of O₃-vegetation coupling on surface O₃ air quality via
biogeochemical and meteorological feedbacks. *Atmos. Chem. Phys.*, *17*, 3055–3066. <https://doi.org/10.5194/acp-17-3055-2017>, 2017
- Sanderson, M. G., Jones, C. D., Collins, W. J., Johnson, C. E., & Derwent, R. G.: Effect of climate change on isoprene
920 emissions and surface O₃ levels. *Geophys. Res. Lett.*, *30*. <https://doi.org/10.1029/2003GL017642>, 2003
- Schröder. TOAR Data Infrastructure. <https://doi.org/10.34730/4d9a287dec0b42f1aa6d244de8f19eb3>, 2021
- Schultz, M. G., Jacob, D. J., Wang, Y., Logan, J. A., Atlas, E. L., Blake, D. R., Blake, N. J., Bradshaw, J. D., Browell, E. v.,
Fenn, M. A., Flocke, F., Gregory, G. L., Heikes, B. G., Sachse, G. W., Sandholm, S. T., Shetter, R. E., Singh, H. B., &
Talbot, R. W.: On the origin of tropospheric O₃ and NO_x over the tropical South Pacific. *J. Geophys. Res.*, *104*, 5829–
925 5843. <https://doi.org/10.1029/98JD02309>, 1999
- Schultz, MG, Schröder, S, Lyapina, O, Cooper, OR, Galbally, I, Petropavlovskikh, I, von Schneidmesser, E, Tanimoto, H,
Elshorbany, Y, Naja, M, Seguel, RJ, Dauert, U, Eckhardt, P, Feigenspan, S, Fiebig, M, Hjellbrekke, A-G, Hong, Y-D,

- Kjeld, PC, Koide, H, Lear, G, Tarasick, D, Ueno, M, Wallasch, M, Baumgardner, D, Chuang, M-T, Gillett, R, Lee, M, Molloy, S, Moolla, R, Wang, T, Sharps, K, Adame, JA, Ancellet, G, Apadula, F, Artaxo, P, Barlasina, ME, Bogucka, M, Bonasoni, P, Chang, L, Colomb, A, Cuevas-Agulló, E, Cupeiro, M, Degorska, A, Ding, A, Fröhlich, M, Frolova, M, Gadhavi, H, Gheusi, F, Gilge, S, Gonzalez, MY, Gros, V, Hamad, SH, Helmig, D, Henriques, D, Hermansen, O, Holla, R, Hueber, J, Im, U, Jaffe, DA, Komala, N, Kubistin, D, Lam, K-S, Laurila, T, Lee, H, Levy, I, Mazzoleni, C, Mazzoleni, LR, McClure-Begley, A, Mohamad, M, Murovec, M, Navarro-Comas, M, Nicodim, F, Parrish, D, Read, KA, Reid, N, Ries, L, Saxena, P, Schwab, JJ, Scorgie, Y, Senik, I, Simmonds, P, Sinha, V, Skorokhod, AI, Spain, G, Spangl, W, Spoor, R, Springston, SR, Steer, K, Steinbacher, M, Suharguniyawan, E, Torre, P, Trickl, T, Weili, L, Weller, R, Xiaobin, X, Xue, L and Zhiqiang, M: Tropospheric O₃ Assessment Report: Database and metrics data of global surface O₃ observations. *Elm Sci Anth*, 58, <https://doi.org/10.1525/elementa.244>, 2017
- Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., Mora, L. d., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahahan, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *J. Adv. Modeling Earth Sy.*, 11, 4513–4558, <https://doi.org/10.1029/2019MS001739>, 2019.
- Shindell, D. T., Faluvegi, G., Unger, N., Aguilar, E., Schmidt, G. A., Koch, D. M., Bauer, S. E., & Miller, R. L.: Simulations of preindustrial, present-day, and 2100 conditions in the NASA GISS composition and climate model G-PUCCINI. *Atmos. Chem. Phys*, 6, 4427–4459. <https://doi.org/10.5194/acp-6-4427-2006>, 2006
- Shindell, D. T., Pechony, O., Voulgarakis, A., Faluvegi, G., Nazarenko, L., Lamarque, J. F., Bowman, K., Milly, G., Kovari, B., Ruedy, R., & Schmidt, G. A.: Interactive O₃ and methane chemistry in GISS-E2 historical and future climate simulations. *Atmos. Chem. Phys*, 13, 2653–2689. <https://doi.org/10.5194/acp-13-2653-2013>, 2013
- Shi, Y., Zang, S., Matsunaga, T., & Yamaguchi, Y.: A multi-year and high-resolution inventory of biomass burning emissions in tropical continents from 2001–2017 based on satellite observations. *Journal of Cleaner Production*, 270, 122511. <https://doi.org/10.1016/j.jclepro.2020.122511>, 2020
- Silva, S. J., & Heald, C. L.: Investigating Dry Deposition of O₃ to Vegetation. *J. Geophys. Res. Atmos.*, 123, 559–573. <https://doi.org/10.1002/2017JD027278>, 2018
- Sinha, P., Jaeglé, L., Hobbs, P. v, & Liang, Q.: Transport of biomass burning emissions from southern Africa. *J. Geophys. Res*, 109, 20204. <https://doi.org/10.1029/2004JD005044>, 2004
- Sitch, S., Cox, P. M., Collins, W. J., & Huntingford, C.: Indirect radiative forcing of climate change through O₃ effects on the land-carbon sink. *Nature*, 448, 791–794. <https://doi.org/10.1038/nature06059>, 2007

- Squire, O. J., Archibald, A. T., Abraham, N. L., Beerling, D. J., Hewitt, C. N., Lathière, J., Pike, R. C., Telford, P. J., & Pyle, J. A.: Influence of future climate and cropland expansion on isoprene emissions and tropospheric O₃. *Atmos. Chem. Phys.*, *14*, 1011–1024. <https://doi.org/10.5194/acp-14-1011-2014>, 2014
- 965 Squire, O. J., Archibald, A. T., Griffiths, P. T., Jenkin, M. E., Smith, D., & Pyle, J. A.: Influence of isoprene chemical mechanism on modelled changes in tropospheric O₃ due to climate and land use over the 21st century. *Atmos. Chem. Phys.*, *15*, 5123–5143. <https://doi.org/10.5194/acp-15-5123-2015>, 2015
- Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., Wild, O., Zeng, G., Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J.-F., Lawrence, M. G., Montanaro, V., Muller, J. F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Shindell, D. T., Strahan, S. E., Sudo, K., and Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.*, *111*, D08301, <https://doi.org/10.1029/2005JD006338>, 2006.
- 970 Stroud, C., Makar, P., Karl, T., Guenther, A., Geron, C., Turnipseed, A., Nemitz, E., Baker, B., Potosnak, M., Fuentes, J. D., Stroud, C. :, Makar, P., Karl, T., Guenther, A., Geron, C., Turnipseed, A., Nemitz, E., Baker, B., Potosnak, M., & Fuentes, J. D.: Role of canopy-scale photochemistry in modifying biogenic-atmosphere exchange of reactive terpene species: Results from the CELTIC field study. *J. Geophys. Res.*, *110*, 17303. <https://doi.org/10.1029/2005JD005775>, 2005
- 975 Szopa, S., V. Naik, B. Adhikary, P. Artaxo, T. Berntsen, W.D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler-Scharr, Z. Klimont, H. Liao, N. Unger, and P. Zanis, 2021: Short-Lived Climate Forcers. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. *Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA*, pp. 817–922, doi:10.1017/9781009157896.008.
- 980 Tsigaridis, K., & Kanakidou, M.: The present and future of secondary organic aerosol direct forcing on climate. *Current Climate Change Reports*, *4*, 84–98. <https://doi.org/10.1007/s40641-018-0092-3>, 2018
- Turnock, S. T., Allen, R. J., Andrews, M., Bauer, S. E., Deushi, M., Emmons, L., Good, P., Horowitz, L., John, J. G., Michou, M., Nabat, P., Naik, V., Neubauer, D., O'Connor, F. M., Olivie, D., Oshima, N., Schulz, M., Sellar, A., Shim, S., Takemura, T., Tilmes, S., Tsigaridis, K., Wu, T., and Zhang, J.: Historical and future changes in air pollutants from CMIP6 models, *Atmos. Chem. Phys.*, *20*, 14547–14579, <https://doi.org/10.5194/acp-20-14547-2020>, 2020.
- 990 Turnock, S. T., Wild, O., Sellar, A., & O'Connor, F. M: 300 years of tropospheric O₃ changes using CMIP6 scenarios with a parameterised approach. *Atmos. Environ.*, *213*, 686–698. <https://doi.org/10.1016/j.atmosenv.2019.07.001>, 2019

- Unger, N., Harper, K., Zheng, Y., Kiang, N. Y., Aleinov, I., Arneth, A., Schurgers, G., Amelynck, C., Goldstein, A., Guenther, A., Heinesch, B., Hewitt, C. N., Karl, T., Laffineur, Q., Langford, B., A. McKinney, K., Misztal, P., Potosnak, M., Rinne, J., Pressley, S., Schoon, N., and Serça, D.: Photosynthesis-dependent isoprene emission from leaf to planet in a global carbon-chemistry-climate model, *Atmos. Chem. Phys.*, 13, 10243–10269, <https://doi.org/10.5194/acp-13-10243-2013>, 2013.
- Weber, J., Archer-Nicholls, S., Abraham, N. L., Shin, Y. M., Bannan, T. J., Percival, C. J., Bacak, A., Artaxo, P., Jenkin, M., Khan, M. A. H., Shallcross, D. E., Schwantes, R. H., Williams, J., & Archibald, A. T.: Improvements to the representation of BVOC chemistry-climate interactions in UKCA (v11.5) with the CRI-Strat 2 mechanism: Incorporation and evaluation. *Geosci. Model Dev.*, 14, 5239–5268. <https://doi.org/10.5194/gmd-14-5239-2021>, 2021
- Wedow, J. M., Ainsworth, E. A., & Li, S.: Plant biochemistry influences tropospheric O₃ formation, destruction, deposition, and response. *Trends Biochem. Sci.*, 46, 992–1002. <https://doi.org/10.1016/j.tibs.2021.06.007>, 2021
- Wild, O., Voulgarakis, A., O'Connor, F., Lamarque, J. F., Ryan, E. M., & Lee, L.: Global sensitivity analysis of chemistry-climate model budgets of tropospheric O₃ and OH: Exploring model diversity. *Atmos. Chem. Phys.*, 20, 4047–4058. <https://doi.org/10.5194/acp-20-4047-2020>, 2020
- Williams, K. D., Copsey, D., Blockley, E. W., Bodas-Salcedo, A., Calvert, D., Comer, R., Davis, P., Graham, T., Hewitt, H. T., Hill, R., Hyder, P., Ineson, S., Johns, T. C., Keen, A. B., Lee, R. W., Megann, A., Milton, S. F., Rae, J. G. L., Roberts, M. J., Scaife, A. A., Schiemann, R., Storkey, D., Thorpe, L., Watterson, I. G., Walters, D. N., West, A., Wood, R. A., Woollings, T., and Xavier, P. K.: The Met Office Global Coupled Model 3.0 and 3.1 (GC3.0 and GC3.1) Configurations, *J. Adv. Model. Earth Sy.*, 10, 357–380, <https://doi.org/10.1002/2017MS001115>, 2018.
- Wiltshire, A. J., Duran Rojas, M. C., Edwards, J. M., Gedney, N., Harper, A. B., Hartley, A. J., Hendry, M. A., Robertson, E., and Smout-Day, K.: JULES-GL7: the Global Land configuration of the Joint UK Land Environment Simulator version 7.0 and 7.2, *Geosci. Model Dev.*, 13, 483–505, <https://doi.org/10.5194/gmd-13-483-2020>, 2020.
- Wiltshire, A. J., Burke, E. J., Chadburn, S. E., Jones, C. D., Cox, P. M., Davies-Barnard, T., Friedlingstein, P., Harper, A. B., Liddicoat, S., Sitch, S., and Zaehle, S.: JULES-CN: a coupled terrestrial carbon–nitrogen scheme (JULES vn5.1), *Geosci. Model Dev.*, 14, 2161–2186, <https://doi.org/10.5194/gmd-14-2161-2021>, 2021.
- World Health Organisation: World Air Quality Report, available at: <https://www.iqair.com/world-air-quality-report>, 2021
- Wu, S., Mickley, L. J., Jacob, D. J., Logan, J. A., Yantosca, R. M., & Rind, D.: Why are there large differences between models in global budgets of tropospheric O₃? *J. Geophys. Res. Atmos.*, 112. <https://doi.org/10.1029/2006JD007801>, 2007
- Yue, X., & Unger, N.: Fire air pollution reduces global terrestrial productivity. *Nat. Commun.*, 9, 1–9. <https://doi.org/10.1038/s41467-018-07921-4>, 2018
- Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yabu, S., Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y., & Ishii, M.: The meteorological research institute Earth system model version 2.0, MRI-ESM2.0: Description and basic evaluation of the physical component. *J. Meteorol. Soc. Japan*, 97, 931–965. <https://doi.org/10.2151/jmsj.2019-051>, 2019

- 1030 Zanis, P., Turnock, S., Naik, V., Szopa, S., K Georgoulas, A., Bauer, S. E., Deushi, M., Horowitz, L. W., Keeble, J., le Sager, P., O, F. M., Oshima, N., Tsigaridis, K., & van Noije, T.: *Climate change penalty and benefit on surface O3: A global perspective based on CMIP6 earth system models*, [preprint] <https://doi.org/10.1088/1748-9326/ac4a34> , 2022
- Zeng, G., Pyle, J. A., & Young, P. J.: Impact of climate change on tropospheric O3 and its global budgets. *Atmos. Chem. Phys.*, 8, 369–387. <https://doi.org/10.5194/acp-8-369-2008>, 2008
- 1035 Ziemke, J. R., Chandra, S., Duncan, B. N., Schoeberl, M. R., Torres, O., Damon, M. R., & Bhartia, P. K.: Recent biomass burning in the tropics and related changes in tropospheric O3. *Geophys. Res. Lett.*, 36. <https://doi.org/10.1029/2009GL039303>, 2009