



Technical note: The CAMS greenhouse gas reanalysis from 2003 to 2020

Anna Agustí-Panareda¹, Jérôme Barré¹, Sébastien Massart¹, Antje Inness¹, Ilse Aben², Melanie Ades¹, Bianca C. Baier^{3,4}, Gianpaolo Balsamo¹, Tobias Borsdorff², Nicolas Boussez¹, Souhail Boussetta¹,
5 Michael Buchwitz⁵, Luca Cantarello¹, Cyril Crevoisier⁶, Richard Engelen¹, Henk Eskes⁷, Johannes Flemming¹, Sébastien Garrigues¹, Otto Hasekamp², Vincent Huijnen⁷, Luke Jones¹, Zak Kipling¹, Bavo Langerock⁸, Joe McNorton¹, Nicolas Meilhac⁶, Stefan Noel⁵, Mark Parrington¹, Vincent-Henri Peuch¹, Michel Ramonet⁹, Miha Ratzinger¹, Maximilian Reuter⁵, Roberto Ribas¹, Martin Suttie¹, Colm Sweeney⁴, Jérôme Tarniewicz⁹, Lianghai Wu¹⁰

10

¹ European Centre for Medium Range Weather Forecasts, Shenfield Park, Reading RG2 9AX, United Kingdom

² SRON Netherlands Institute for Space Research, Utrecht, the Netherlands

³ Cooperative Institute for Research in Environmental Sciences, University of Colorado-Boulder, Boulder, CO, USA

⁴ NOAA, Global Monitoring Laboratory, Boulder, CO, USA

15 ⁵ Institute of Environmental Physics (IUP), University of Bremen, 28334 Bremen, Germany

⁶ Laboratoire de Météorologie Dynamique (LMD/IPSL), CNRS, Ecole polytechnique, 91128 Palaiseau Cedex, France

⁷ Royal Netherlands Meteorological Institute, Utrechtseweg 297, NL-3731 GA De Bilt, Netherlands

⁸ Royal Belgian Institute for Space Aeronomy, Avenue Circulaire 3, 1180 Uccle, Belgium

20 ⁹ Laboratoire des Sciences du Climat et de l'Environnement (LSCE-IPSL), CEA-CNRS-UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

¹⁰ Flemish Institute for Technological Research (VITO), Remote Sensing Unit, Boeretang 200, B-2400 Mol, Belgium

Correspondence to: Anna Agusti-Panareda (A.Agusti-Panareda@ecmwf.int)

25 **Abstract.** The Copernicus Atmosphere Monitoring Service has recently produced a greenhouse gases reanalysis (version egg4) that covers almost two decades from 2003 to 2020 and will be extended in the future. This reanalysis dataset includes carbon dioxide (CO₂) and methane (CH₄). The reanalysis procedure combines model data with satellite data into a globally complete and consistent dataset using the European Centre for Medium-range Weather Forecasts' Integrated Forecasting System (IFS). This dataset has been carefully evaluated against independent observations to ensure validity and point out deficiencies to the
30 user. The greenhouse gas reanalysis can be used to examine the impact of atmospheric greenhouse gases concentrations on climate change, such as global and regional climate radiative forcing, assess intercontinental transport, and also serve as boundary conditions for regional simulations, among other applications and scientific studies. The caveats associated with changes in assimilated observations and fixed underlying emissions are highlighted, as well as their impact on the estimation of trends and annual growth rates of these long-lived greenhouse gases.



35 1 Introduction

Atmospheric carbon dioxide (CO₂) and methane (CH₄) are the most abundant man-made greenhouse gases directly responsible for climate change (IPCC, 2021). Their long lifetime and increasing anthropogenic emissions near the surface account for their long-term trends (Friedlingstein et al., 2021). A lot of effort has been devoted to measuring the atmospheric concentrations from ground-based observatories (e.g. National Oceanic and Atmospheric Administration (NOAA), gml.noaa.gov; Integrated Carbon Observation System (ICOS), www.icos-cp.eu), which provide the gold standard for the estimation of trends, and more recently satellite data (Committee on Earth Observation Satellites (CEOS), Crisp et al., 2018) enhancing the spatial coverage of greenhouse gas observations at global scale. Atmospheric measurements also sample the variability of CO₂ and CH₄ coming from the weather and its associated atmospheric transport (e.g. Patra et al., 2008, 2011). For this reason, Numerical Weather Prediction (NWP) models have been extensively used to represent and reconstruct the variability of atmospheric concentrations of various tracers (e.g. Inness et al., 2019). Here we use the Integrated Forecasting System (IFS) of the European Centre for Medium-range Weather Forecasts (ECMWF) which has been adapted to include CO₂ and CH₄ in the weather forecast (Agustí-Panareda et al., 2017, 2019) to create a greenhouse gases (GHG) reanalysis. The reanalysis uses the data assimilation technique to combine CO₂ and CH₄ satellite data from the SCanning Imaging Absorption spectroMeter for Atmospheric CHartography (SCIAMACHY, www.sciamachy.org), the Infrared Atmospheric Sounding Interferometer (IASI, www.eumetsat.int/iasi) and The Thermal and Near Infrared Sensor for Carbon Observation (TANSO, www.eorc.jaxa.jp/GOSAT/instrument_1.html) instruments with IFS model simulations of CO₂ and CH₄ (Agustí-Panareda et al., 2022). The dataset is based on a consistent and stable model version to provide a homogenous, continuous and gapless record of the CO₂ and CH₄ in the entire atmosphere since 2003.

The IFS includes a forecasting model and a data assimilation system combined. The data assimilation system also integrates meteorological observations as in the fifth generation of ECMWF meteorological reanalyses, ERA5 (Hersbach et al., 2020), to best constrain the atmospheric variability of greenhouse gases (Massart et al., 2014, 2016). The forecasting model provides a 3-dimensional representation and evolution of the atmospheric CO₂ and CH₄ and meteorological variables (Agustí-Panareda et al., 2019). At the model surface the greenhouse gases are forced by a set of surface fluxes and emissions. Such modelling configuration allows to produce a realistic representation of the spatio-temporal variability of greenhouse gases in the atmosphere over a wide range of scales from hours to seasons and from local to global (Agustí-Panareda et al., 2022).

Figure 1 showcases the global evolution of CO₂ and CH₄ represented by the CAMS GHG reanalysis data set over the period 2003-2020 and the span of the used satellite data. The seasonal averages illustrate the spatial and temporal variability information contained in the reanalysis dataset which can be exploited for a range of applications in atmospheric sciences. A key potential use of the CAMS GHG reanalysis is to assess the impact of greenhouse gases on climate change. The reanalysis 3-dimensional fields could be used to investigate global and regional climate radiative forcing (e.g. atmosphere.copernicus.eu/climate-forcing), serve as boundary conditions for regional simulations, assess intercontinental



transport, and generally provide a reference for any other study focusing on atmospheric variability of CO₂ and CH₄. However, care should be taken when using the CAMS GHG reanalysis to estimate trends and annual growth rates of these long-lived
70 greenhouse gases by considering the caveats associated with the changes in the satellite retrievals of CO₂ and CH₄ and the fact that neither anthropogenic emissions nor natural fluxes are adjusted by the data assimilation system, unlike atmospheric inversions (e.g. Chevallier et al., 2019).

The objective of this technical report is to document the technical aspects of the method and input data used to produce the CAMS GHG reanalysis, and to provide guidance to potential users on the strengths and limitations of the dataset. Section 2
75 describes the processing chain to produce the reanalysis and its components. Section 3 focuses on the evaluation of the CAMS GHG reanalysis using independent observations from the TCCON and NDACC networks as well as surface in situ networks and AirCore profiles. A list of limitations and caveats of the CAMS GHG reanalysis associated with the changes in the assimilated data and the underlying model errors are compiled in Sect. 4. Finally, Sect. 5 provides a summary and outlook for future CAMS GHG reanalyses.




Figure 1. (a) Reanalysis timeseries of global column-averaged CO₂ (red) and CH₄ (purple) atmospheric mole fractions; (b) the span of the satellite data records for the corresponding species; (c) CO₂ and (d) CH₄ seasonal total column averages (DJF: December-January-February, MAM: March-April-May, JJA: June-July-August, SON: September, October, November) for the 2003-2020 period.



85




Figure 2. Schematic of the reanalysis cycling procedure. The flow diagram shows the steps and elements combined in the reanalysis. Surface fluxes are used as boundary condition for the atmospheric forecasts. Satellite data are combined with the forecast using data assimilation to produce an analysis (corrected 4D fields) to initialize the next forecast.

90



2 Methods

This section gives an overview of the different building blocks of the CAMS GHG reanalysis and the processing chain that integrates the different components to produce the reanalysis dataset.

2.1 The reanalysis cycling chain

95 The reanalysis production chain is illustrated in Fig. 2. It is a cycling procedure based on a 12-hour data assimilation window that involves four main parts:

- Satellite retrievals of CO₂ and CH₄ (see section 2.2) as well as NWP observations (Hersbach et al., 2020).
- Surface fluxes (see section 2.3) that constitute the sources and sinks of CO₂ and CH₄ in the atmosphere are compiled from various sources. They provide the surface boundary condition for the tracer transport model.
- 100 - A model forecast (see section 2.4) that provides a 4-dimensional representation of the state of the greenhouse gases over space and time, along with other meteorological variables, during the 12-hour analysis window (from 09:00 to 21:00 and 21:00 to 09:00 UTC). The forecasts are initialised with the previous analysis, except for the first forecast for the initial date, which is initialised with atmospheric molar fractions from the CAMS inversion dataset (Chevallier et al., 2020; Segers et al., 2020a).
- 105 - The above elements are combined using a data assimilation system (see section 2.5) to produce an analysis (Massart et al., 2014, 2016). The analysis will serve to initialise the following forecast over the subsequent 12 hourly cycle.

Details of these four different components of the reanalysis processing chain are provided in the subsections below, as well as the approach followed to monitor the assimilation of CO₂ and CH₄ satellite data.

110 2.2 Satellite GHG observations

The satellite measurements of radiances (L1 data) are processed by satellite retrievals developed by various data providers to derive information on the total and partial atmospheric column of CO₂ and CH₄ dry mole fraction (L2 data). In the CAMS GHG reanalysis only L2 products were used for CO₂ and CH₄. With nadir looking satellite instrument geometries the L2 data provide vertically integrated content with vertical sensitivity functions called either averaging kernel when an optimal estimation approach (Rogers, 2000) is used or weighting functions, that provide information on where the retrieval sensitivity is located along the vertical. The satellite products assimilated in this reanalysis are all provided with averaging kernel and prior information or weighting functions (Massart et al, 2014, 2016). Table 1 provides the specification for each of the assimilated satellite CO₂ and CH₄ products, selected as the state-of-the art retrievals at the beginning of 2017, when the CAMS GHG reanalysis production started. All of the L2 satellite products are freely available from the Copernicus Climate Change Service (C3S) Copernicus Climate Data Store (Alos et al., 2019) at [115](https://cds.climate.copernicus.eu/cdsapp#!/dataset/satellite-) [120](https://cds.climate.copernicus.eu/cdsapp#!/dataset/satellite-)



[carbon-dioxide](#) for CO₂ and <https://cds.climate.copernicus.eu/cdsapp#!/dataset/satellite-methane> for CH₄. The GHG reanalysis integrate the L2 GHG data from the following satellite instruments:

- 125 • **SCIAMACHY – Envisat:** The The SCanning Imaging Absorption spectroMeter for Atmospheric Cartography (SCIAMACHY) instrument onboard the Envisat satellite was launched by the European Space Agency (ESA) in March 2002 and it was developed by a consortium involving the Netherland Space Office, the German Aerospace Centre and the Belgian Federal Science Policy Office. It measures radiances variations from the ultraviolet to the near visible infrared. The GHG L2 products use the nadir spectra of reflected and scattered solar radiation in the near-infrared region. Satellite radiance observations in the near infrared spectral region with the nadir looking geometry are sensitive to changes in CO₂ and CH₄ down to the Earth's surface. The measurements provide total column information with sensitivity peaking near the surface. The ground pixel size is typically between 30 km and 60 km and the swath width is about 960 km. There are no across-track gaps between the ground pixels but there are gaps along-track as SCIAMACHY operates only part of the time (approx. 50%) in nadir observation mode. The CO₂ and CH₄ column products are retrieved by the University of Bremen (Reuter et al., 2011) and the Netherland Institute for Space Research (SRON) (Frankenberg et al., 2011), respectively. Both of L2 products are delivered by the ESA GHG-Climate Change Initiative (Buchwitz et al, 2015) and the C3S Climate Data Store (<https://cds.climate.copernicus.eu>).
130
- 135 • **TANSO-FTS – GOSAT:** The Thermal And Near infrared Sensor for carbon Observations - Fourier Transform Spectrometer (TANSO-FTS) instrument onboard the Greenhouse Gases Observing Satellite (GOSAT) satellite has been developed by the Japan Aerospace Exploration Agency (JAXA) and it was launched in January 2009. TANSO-FTS measures radiances in the short-wave infrared band that provide information of total-column CO₂ and CH₄ mole fractions. Similar to SCIAMACHY, the sensitivity of the total column information provided by L2 data is peaking near the surface due to the spectral band used. The ground pixel size is about 10 km, the swath is 750 km and it has a revisit time of 3 days. In contrast to SCIAMACHY, the GOSAT scan pattern consists of non-consecutive individual ground pixels, i.e., the scan pattern is not gap-free. For a general overview about GOSAT see also <http://www.gosat.nies.go.jp/en/>. The L2 retrieval product is engineered by the SRON (Schepers et al., 2012, 2016) and delivered by the ESA GHG-CCI and the C3S Climate Data Store (<https://cds.climate.copernicus.eu>).
140
145
- 150 • **IASI – Metop A and B:** The Infrared Atmospheric Sounding Interferometer (IASI) instruments are onboard the Meteorological Operational satellites (Metop-A and Metop-B) launched in October 2006 and September 2012 respectively. The French National Centre for Space Studies (CNES) lead the design and developments of the instruments in collaboration with the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT). The IASI instruments measure the thermal infrared band with high spectral resolution enabling it to detect a wide range of trace gas variations in the atmosphere, including CO₂ and CH₄ sensitive in the mid and upper tropospheric regions between 5 and 12 km of altitude. IASI is an across track scanning system with a swath width of 2200 km, providing global coverage twice a day. The field of view is sampled by 2×2 pixels whose ground resolution is 12 km at nadir. Both CO₂ and CH₄ are engineered and delivered by the Centre National de Recherche Scientifique



155 (CNRS)-Laboratoire de Météorologie Dynamique (LMD) (Crevoisier et al., 2009a, 2009b, 2014). The two L2 products are delivered by the ESA GHG-Climate Change Initiative (Buchwitz et al, 2015) and the C3S Climate Data Store (<https://cds.climate.copernicus.eu>).

160 **Table 1. Specifications of the satellite data used in the CAMS GHG reanalysis**

Gas	Instrument - Satellite	Period assimilated	Version (data provider)	Reference	Peaking sensitivity
CO ₂	SCIAMACHY – Envisat	20030101 – 20120324	CO2_SCI_BESD (v02.01.02, IUP-UB)	Reuter et al., (2011)	Near Surface
	IASI – Metop-A	20070701 - 20150531	CO2_IAS_NLIS (v8.0, CNRS-LMD)	Crevoisier et al. (2009a)	Middle and Upper troposphere
	IASI – Metop-B	20130201 – 20181130	CO2_IAS_NLIS (v4.2_nrt, CNRS-LMD)		Middle and upper troposphere
		20181201-20201231	CO2_IAS_NLIS (v4.0_nrt, CNRS-LMD)		
	TANSO-FTS - GOSAT	20090601-20131231	CO2_GOS_SRFP (V2.3.6, SRON)	Butz et al., (2011); Guerlet et al. (2013) ; Heymann et al. (2015)	Near Surface
		20140101-20181231	CO2_GOS_SRFP(V2.3.8, SRON)		
20190101-20201231		CO2_GOS_BESD (CAMS_NRT, IUP-UB)			
CH ₄	SCIAMACHY – Envisat	20030108-20100601	CH4_SCI_IMAP (v7.2, SRON)	Frankenberg et al., (2011)	Near Surface
	IASI – Metop-A	20070701-20150630	CH4_IAS_NLIS (V8.3, CNRS-LMD)	Crevoisier et al., (2009b, 2014)	Middle and Upper troposphere
	IASI – Metop-B	20130201- 20181130	CH4_IAS_NLIS (V8.1_nrt, CNRS-LDM)		Middle and upper troposphere
		20181201-20201231	CH4_IAS_NLIS (v4.0_nrt, CNRS-LDM)		
	TANSO-FTS - GOSAT	20090601-20131231	CH4_GOS_SRFP (V2.3.6, SRON)	Butz et al., (2010); Schepers et al., 2012	Near Surface
		20140101-20181231	CH4_GOS_SRFP (V2.3.8, SRON)		
20190101-20201231		CH4_GOS_SRPR (CAMS_NRT, SRON)			

2.3 Surface fluxes and prescribed sources/sinks

The emissions and surface fluxes provide the surface boundary conditions for the atmospheric concentrations of CO₂ and CH₄. They play a crucial role in determining the variability and growth rate of both greenhouse gases in the atmosphere. Errors in the budget of the total flux will result into systematic errors or biases in forecast of atmospheric CO₂ and CH₄. In the CAMS reanalysis the surface fluxes (including sources and sinks) are not optimized by the assimilation system. This lack of surface



flux optimization can lead to biases in the analysis when the observing system coverage is sparse in space and time or when the observation error is large, and the analysis is strongly influenced by the model forecast.

170 Table 2 lists the datasets used to produce the CAMS reanalysis. They include:


- Fire emissions derived using the CAMS Global Fire Assimilation System (GFAS) version 1.2 that assimilate fire radiative power observations from satellite-based sensors (Kaiser et al., 2012). GFAS produces daily estimates of wildfire and biomass burning emissions. The emissions are injected at the surface and distributed over the boundary layer by the model's convection and vertical diffusion scheme.
- 175 • Anthropogenic emissions from the Emission Database for Global Atmospheric Research (EDGAR) version 4.2FT2010 inventory (Janssens-Maenhout et al., 2011; Olivier and Janssens-Maenhout, 2012) excluding the short carbon cycle. The anthropogenic emissions are based on annual average values and include fossil fuel combustion and leakage, agricultural, landfill/waste emissions and aviation (based on the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP, Lamarque et al., 2013) nitric oxide (NO) emissions from aviation scaled to the annual CO₂ total emission from aviation from EDGAR). Anthropogenic emissions of CO₂ are extrapolated from 2010 to 2014 with the time series of country totals from EDGARv4.3 (Janssens-Maenhout et al., 2016) and from 2015 to 2020, a persistent growth based on the last available year (2014) is applied. CH₄ anthropogenic emissions are fixed with the last year of available gridded data (2010) from 2011 to 2020. Note that CO₂ and CH₄ emissions are not adjusted for the COVID emission reduction in 2020 (Le Quéré et al., 2020).
- 180 • Biogenic CO₂ fluxes are based on the online CHTESSEL module (Boussetta et al., 2013) that relates CO₂ biogenic fluxes with radiation, precipitation, temperature, humidity, and soil moisture. CHTESSEL is used in conjunction with the biogenic flux adjustment system (BFAS) that improves the continental budget of CO₂ fluxes by combining information from fluxes estimates by a global flux inversion system (Chevallier et al., 2010), land-use information and the CHTESSEL online fluxes (Agustí-Panareda et al., 2016).
- 185 • Wetland CH₄ monthly mean emissions come from a climatology (1990–2008) based on the LPJ-WHyMe model that is constrained by SCIAMACHY observations (Spanhi et al., 2011).
- A monthly modulation for CH₄ rice emissions is implemented based on the seasonal cycle of Matthews et al. (1991).
- The CH₄ chemical sink is represented by a monthly mean climatological loss rate from Bergamaschi et al. (2009) based on OH fields optimized with methyl chloroform (Bergamaschi et al., 2005; Houweling et al., 1998) and stratospheric radicals from the 2D photochemical Max-Planck-Institute (MPI) model (Brühl and Crutzen, 1993).
- 195 • Other sources and sinks include a CH₄ monthly soil sink (Ridgwell et al., 1991), CO₂ and CH₄ annual mean oceanic fluxes (Houweling et al., 1999; Lambert and Schmidt, 1993; Takahashi et al., 2009) and CH₄ monthly mean fluxes from termites (Sanderson, 1996) and wild animals (Houweling et al., 1999).



200

Table 2. Specifications of the emission and surface fluxes used in the CAMS GHG reanalysis

Gas	Emission/Flux type	Data provider - Version
CO ₂	CO ₂ and CH ₄ fire emissions	GFAS Version 1.2 (Kaiser et al., 2012)
	CO ₂ ocean fluxes	<u>Takahashi Climatology</u> (Takahashi et al., 2009)
	CO ₂ emissions from aviation	Based on ACCMIP NO emissions from aviation scaled to annual total CO ₂ from EDGAR aviation emissions (Olivier and Janssens-Maenhout, 2012)
	CO ₂ ecosystem fluxes bias corrected with BFAS	Based on CHTESSEL (modelled online in IFS) (Boussetta et al., 2013; Agustí-Panareda et al., 2016)
	CO ₂ anthropogenic emissions	EDGARv4.2FT2010 (2003-2010) (Olivier and Janssens-Maenhout, 2012)
CH ₄	CH ₄ total natural emissions	based on EDGARv4.2FT2010 (2003-2010) (Olivier and Janssens-Maenhout, 2012); LPJ-HYMN wetland climatology (Spahni et al., 2011); and other natural sources/sinks (Matthews et al., 1991; Ridgwell et al., 1999; Houweling et al., 1999; Lambert and Schmidt, 1993; Sanderson, 1996).
	CH ₄ chemical sink	Monthly mean climatology of CH ₄ loss rate from Bergamaschi et al. (2009)
	CH ₄ anthropogenic emissions	EDGARv4.2FT2010 (2003-2010) (Olivier and Janssens-Maenhout, 2012)



205

Figure 3. Monthly CO₂ and CH₄ surface fluxes. The CO₂ fluxes [PgCO₂/month] include modelled Net Ecosystem Exchange (NEE) fluxes (in green), anthropogenic emissions (in purple), ocean fluxes (in blue) and biomass burning emissions (in red). The total CH₄ fluxes [TgCH₄/month] excluding biomass burning emissions are shown by black line and CH₄ biomass burning emissions [TgCH₄/month] are depicted in red. The dash lines show the 1-year running mean for each of the fluxes.

210



2.4 Forecast model

The CAMS GHG reanalysis has been produced using the IFS model. The same model is used to produce operational numerical weather predictions (NWP) at ECMWF and the CAMS global forecast and analyses for reactive gas, aerosols and greenhouse gases at ECMWF (Fleming et al. 2015, Agustí-Panareda et al., 2017, Agustí-Panareda et al., 2022). The IFS model version used is IFS CY42R1, the same as in the CAMS reanalysis for reactive gases and aerosols (Inness et al., 2019). The forecasting model uses a reduced Gaussian grid with a resolution of T1255 corresponding to a horizontal resolution of approximately 80 km and 60 hybrid-sigma pressure vertical levels from the surface to 0.1hPa. The tracer advection is computed using a semi-implicit semi-Lagrangian scheme (Temperton et al., 2001; Diamantakis and Magnusson, 2016) that is not mass-conserving. This scheme leads to an error growth that can dominate the signal in the model simulations if it is not corrected. Thus, a mass fixer is required to ensure mass conservation at every time step (Diamantakis and Agustí-Panareda, 2017). The mass fixer is particularly important for long-lived greenhouse gases for which the interesting signals to monitor, e.g., trends or annual growth rates and large-scale spatial gradients, are weak compared to the large background values. The transport model also includes a turbulent mixing scheme (Sandu et al., 2013) and a convection scheme (Bechtold et al., 2014). For the CH₄ chemical sink in the troposphere and the stratosphere, climatological loss rates derived from the Max Planck Institute photochemical model are used (Bergamaschi et al., 2009). Full documentation of the IFS can be found at <https://www.ecmwf.int/en/forecasts/documentation-and-support/changes-ecmwf-model/ifs-documentation>.

2.5 Analysis procedure (data assimilation)

The IFS system is using an incremental formulation of the 4-dimensional variational technique (4D-Var). The 4D-Var technique consists of minimizing a cost function that combines the model information and the observation information in order to obtain the best possible state of the atmosphere (analysis) accounting for the model and observation errors. The incremental 4D-Var cost function is quadratic and is formulated as follows:

$$J(\delta\mathbf{x}) = \frac{1}{2}(\delta\mathbf{x} - \delta\mathbf{x}_b)^T \mathbf{B}^{-1}(\delta\mathbf{x} - \delta\mathbf{x}_b) + \frac{1}{2}(\mathbf{G}\delta\mathbf{x} - \mathbf{d})\mathbf{R}^{-1}(\mathbf{G}\delta\mathbf{x} - \mathbf{d}) \quad (1)$$

where $\delta\mathbf{x}$ is the increment i.e., the difference between the model state \mathbf{x} and the first guess \mathbf{x}_g , $\delta\mathbf{x}_b$ is the difference between the background (the forecast started from the previous analysis) and the first guess, \mathbf{B} the background error covariance matrix, \mathbf{R} the observation error covariance matrix, \mathbf{G} the observation operator or forward operator that translate the information from model space to observation space. The innovation vector is $\mathbf{d} = \mathbf{y} - \mathbf{G}\mathbf{x}_g$ with \mathbf{y} the observation vector and \mathbf{x}_g the first guess. When the minimization of the cost function is complete, $\delta\mathbf{x}$ is added to \mathbf{x}_g to provide the analysis.



$$x_a = x_g + \delta x \quad (2)$$

245 The analysis is performed over 12-hour assimilation windows from 9:00 to 21:00 and from 21:00 to 9:00 UTC. The incremental 4D-Var assimilation involves the stepwise minimization of the linearized cost function (equation 1) by updating the first guess x_g and increasing the resolution. In the CAMS reanalysis setup, two minimizations are completed successively at TL95 (approximately 210 km) and TL159 (approximately 110 km) spectral truncations. Once the assimilation procedure is completed an analysis is generated that will serve to initialize the next forecast at the full TL255 resolution.

250




Figure 4. Model background error for CO₂ and CH₄ used in the CAMS GHG reanalysis: (a,b) global mean standard deviation and (c,d) global mean error correlation length scale across the vertical levels.

255

The background errors for CO₂ and CH₄ were produced from an ensemble of data assimilations (Massart et al., 2016), which allows the calculation of differences between pairs of background fields which have the characteristics of the background errors. The background errors for the greenhouse gas species are univariate, which means that there is no correlation between the greenhouse gas species and the dynamical fields. Hence each species is assimilated independently from the others. The



260 background errors used for both the greenhouse gas species and the dynamical fields are also constant in time. In the ECMWF
data assimilation system, the background error covariance matrix is given in a wavelet formulation (Fisher, 2004, 2006). This
allows both spatial and spectral variations of the horizontal and vertical background error covariances globally. Figure 4 shows
the global mean of the standard deviation and average horizontal correlation length scales for both CH₄ and CO₂. Following
experimentation, the correlation length scales between the background errors were manually reduced in the atmospheric
265 boundary layer (1km from the surface).

2.6 Monitoring the data assimilation system

The time series of the departures (or differences) between the analysis (AN) and the assimilated satellite data (hereafter referred
270 to as observations, OBS), as well as those between the underlying model simulation (or background, BG) and the observations,
are used to monitor the performance of the analysis system and are shown in Figures 5 (for CO₂) and 6 (for CH₄). For each
satellite retrieval product, both the BG departures (OBS-BG, green lines) and the AN departures (OBS-AN, red lines) are
plotted (panel a: SCIAMACHY, panel b: IASI-A; panel c: IASI-B; panel d: GOSAT), together with the number of observations
assimilated monthly (blue lines). Overall, both the random (i.e., standard deviation, dashed lines) and the systematic
275 components of the departures (i.e., average values, solid lines) are shown to be reduced by the assimilation process, as
highlighted by the AN departures (red lines) being closer to zero than the BG departures (green lines). Note that the difference
between the BG and the AN departure is equal to the analysis increments associated with the related observations (i.e. AN-
BG).

280 The number of observations assimilated is different for each satellite instrument and varies with time: IASI generates the
largest number of data, with both instruments (IASI-A and IASI-B) providing between 150 000 and 200 000 XCO₂ or XCH₄
data per month; the observations taken by SCIAMACHY oscillate between 25 000 and 50 000 for CH₄ and between 5 000 and
10 000 for CO₂; the number of GOSAT XCO₂ data fluctuate around 2 500, whereas those from GOSAT XCH₄ are comprised
between 5 000 and 10 000 per month. It is also clear from Figs 5(a,d) and 6(a,d) that fewer XCO₂ and XCH₄ data from
285 SCIAMACHY, IASI and GOSAT-TANSO are assimilated during the winter months. A magenta vertical dashed line in Figs
5(c,d) and 6(c,d) indicates when the near-real time satellite products started to be assimilated in early 2019. This transition
produced an abrupt change in the quality and availability of both IASI and GOSAT retrievals.

The modelled XCO₂ is systematically larger than the observations (leading to overall negative BG departures) because of the
290 biases in the total fluxes (see section 2.3). Therefore, all instruments produced negative departures until 2013 (Fig. 5). From
2013 to 2019, the modelled values of XCO₂ became smaller than those measured by GOSAT (Fig. 5(d)), while the model
continued to (slightly) overestimate the IASI XCO₂ observations in the mid to upper troposphere. This overestimation is



consistent with a drift in the IASI CO₂ data towards a growing negative bias. After 2018, part of the drift is due to the fact that IASI (version v4.0) is saturating with increasing atmospheric CO₂. Note that this has been corrected with v9.1 (available on
295 the C3S datastore). A sudden change in the IASI-B XCO₂ departures is visible in Fig. 5(c) around December 2018, in
correspondence of the switch from the ESA-CCI reprocessed dataset to a near-real time LMD dataset used operationally in the
CAM5 GHG analysis. The transition to a new dataset was made necessary as the reanalysis production was running close to
real-time and reprocessed observations were not available. After the transition to near-real time observations, the IASI XCO₂
increments are reduced to almost zero, as hinted by the overlap between the red (AN departure) and green line (BG departure)
300 in Fig. 5(c). At the same time, a drop in the number of assimilated IASI XCO₂ observations is observed (blue line, same panel
and figure). Together with a drastic reduction in the magnitude of the increments, a large negative bias of approximately 5ppm
in both the AN and BG departures emerges, accompanied by a large standard deviation error (~20 ppm, cf. dashed lines in
same panel and figure). This degradation in the quality of the IASI-B XCO₂ observations in the near-real time dataset is due
to the change of the correction of the non-linearity of the detector of IASI-B that was made by CNES and EUMETSAT on
305 August 17th 2018 and that introduced a bias of ~0.2 K on the channels used to perform the CO₂ retrieval. This change has been
corrected in the versions of IASI-B MT-CO₂ that are available on the C3S datastore but were not used for this reanalysis. In
January 2019, there was also a transition from the ESA-CCI GOSAT XCO₂ retrievals to the near-real time IUP-UB retrieval
product (Heymann et al., 2015; Massart et al. 2016) as shown in Fig. 5(d). Consequently, the standard deviation of both the
AN and the BG departures increases (cf. dashed lines, same panel and figure), suggesting that the near-real time data is noisier
310 than the reprocessed dataset from ESA-CCI.

The mean XCH₄ departures (both AN and BG) of SCIAMACHY and IASI are relatively small (a few ppb) compared to
GOSAT (up to 10 ppb), throughout the entire time period (see solid red and green lines in Fig. 6). The XCH₄ SCIAMACHY
data was not used from 9 April 2012 onwards (Fig. 6(a)). The standard deviation of both the AN and BG departures are smaller
315 for GOSAT (around 10 ppb, dashed lines in Fig. 6(d)) than for SCIAMACHY (around 20 ppb, dashed lines in Fig.6(a)),
indicating that GOSAT provides less noisy observations. Similar to what was observed for CO₂, a discontinuity in the mean
AN and BG departures of GOSAT XCH₄ emerges in January 2019, in correspondence of the transition from the ESA-CCI
dataset and the NRT SRON retrievals (see dashed pink line in Fig. 6(d)). Both the AN and the BG departures change sign,
indicating that while up to 2019 both the analysis and model were underestimating the GOSAT observations, they start to
320 overestimate them since 2019. Since there was no modification to the model used for the reanalysis over this period, the cause
of this negative bias emerging in both the AN and the BG departures since 2019 can only be attributed to the NRT GOSAT
XCH₄ observations, and in particular to the fact that they are generated by using an extrapolated XCO₂ value in the proxy
retrieval. In addition, the number of assimilated NRT GOSAT XCH₄ observations approximately doubles (blue line in Fig.
6(d)). Note that the switch to the near-real time retrievals for IASI-B XCH₄ has a much more marginal impact on the system
325 (Fig. 6(c)).




Figure 5. Time series of global monthly number of XCO₂ satellite data (blue) and monthly mean CO₂ analysis (AN) and model background (BG) departures of the various observations (OBS) assimilated in the reanalysis (AN-OBS and BG-OBS in red and green respectively, see legend). The solid lines show the monthly average of the departures, and the dash lines the monthly standard deviations. The magenta dash line indicates the switch to the near-real time satellite products. Note that the range of values in y-axis varies depending on the satellite product.

330




Figure 6. Time series of global monthly number of XCH₄ satellite data (blue) and monthly mean CH₄ analysis (AN) and model background (BG) departures of the various observations (OBS) assimilated in the reanalysis (AN-OBS and BG-OBS in red and green respectively, see legend) for different satellite products. The solid lines show the monthly average of the departures, and the dash lines the monthly standard deviations. The magenta dash line indicates the switch to the near-real time satellite products. Note that the range of values in y-axis varies depending on the satellite product.

335



3 Evaluation with independent observations

340 Validation against a set of independent observations has been performed on the 18 years of the CAMS GHG reanalysis span. The independent data includes different types of observations: in situ near-surface continuous observations of CO₂ and CH₄ mole fractions from the collaborative ObsPack datasets (Schuldt et al., 2020; Sarra et al., 2021; NOAA Carbon Cycle Group ObsPack Team, 2019; see Table A1); dry-air column-averaged mole fractions from the Total Carbon Observing Network (TCCON, Wunch et al. 2011, 2015); tropospheric and stratospheric partial columns for CH₄ from the Network for the Detection
345 of Atmospheric Composition Change (NDACC, De Mazière et al., 2018) (see Table A2); AirCore vertical profiles of CO₂ and CH₄ mole fractions (Karion et al., 2010; Baier et al., 2021); and the NOAA global mean CO₂ and CH₄ mole fraction product based on the Greenhouse Gas Marine Boundary layer Reference (Conway et al., 1994, Dlugokencky et al., 1994, Massarie et al., 1995).

350 3.1 Surface and column data

3.1.1 Carbon dioxide


Overall, the error is within ± 10 ppm and ± 4 ppm for most of the near-surface and total column stations respectively for the whole 18-year period (Figs 7 and 8). Near the surface (Fig 7), there is a large variability in the CO₂ error between continental stations influenced by local fluxes (e.g., CDL, FSD, AMT, HUN, see Table A.1) and oceanic stations sampling well-mixed air
355 (ALT, BRW, MHD). Continental stations show large error variations with season (e.g., CDL, HUN), with an underestimation of CO₂ in the summer and an overestimation in the winter, indicating an underestimation of the amplitude of the CO₂ seasonal cycle largely driven by vegetation growth. Differences between stations will be determined by the footprint of observations having different contributions of fluxes from different biomes and from anthropogenic emissions. Accuracy of such fluxes can vary geographically.

360

Overall, there is positive bias of a few ppm between 2003 and 2015 in the baseline surface stations (e.g. BRW, SMO, SPO) which is consistent with the XCO₂ error at the TCCON sites (Fig. 8). This positive bias decreases from 2007 to 2015 when IASI-A CO₂ data are assimilated, with values lower than 2 ppm, and becomes negative from 2015 to 2019 (from 0 to -2 ppm). From 2019 onwards, there is a positive trend in the bias, and it becomes positive ($> +2$ ppm) in 2020. There is consistency
365 between the column and surface biases with a general positive bias at background stations before 2015 and a negative bias after 2015 (up to 2019) at the surface stations, although there is no data in 2020 from the surface stations.



370 The synoptic and large-scale variability of CO₂ is well represented by the reanalysis (lower panel in Fig 8). The root mean square error at TCCON stations is below 0.8 ppm for XCO₂. The normalised standard deviation is around 1.0 (+/- 0.3) and the Pearson correlation coefficient is larger than 0.8.



375 **Figure 7.** Mosaic plot of CO₂ weekly biases (in ppm) of the CAMS GHG reanalysis compared to surface continuous observations of CO₂ mole fraction obtained from GLOBALVIEWplus CO₂ ObsPack v6.0 (Schuldt et al., 2020) and listed in Table A1. Each coloured vertical line represents a weekly mean. Vertical yellow lines depict the changes in the assimilated data documented in Figs 1, 5 and 6. Grey shading indicates no observations are available.




385

3.1.2 Methane

The CH₄ reanalysis fields are generally in good agreement with surface and tropospheric column observations with typical weekly and monthly errors within ± 40 ppb and ± 25 ppb respectively (Figs 9, 10 and 11). Stratospheric partial columns compared to NDACC data reveal a positive bias that is of the same order as the reported measurement uncertainty of 7% (Fig. 10, upper panel). The averaged relative differences in the troposphere across all NDACC sites are -0.4% for the reanalysis (Fig. 10, lower panel), which is well within the measurement's uncertainty. The reanalysis overestimates the column-averaged CH₄ compared to TCCON observations (Fig. 11), for most mid- and high-latitude sites, with a relative difference of up to 2.5%, but shows a good agreement for the low latitude sites at Izaña, Darwin and Wollongong. At the surface the bias is overall positive up to 2007 (Fig. 9). With the introduction of IASI, the biases are reduced. However, with the switch to near-real time satellite data, the bias become negative at all sites reaching values lower than -20 ppb.

Differences between the surface and total column biases stem from the fact that the model suffers from large positive biases above the tropopause (between 100hPa and 10 hPa) of about 80-100 ppb during the months between September and November (Figs 5d and 6d of Verma et al., 2017) which affect the total column average. This stratospheric bias cannot be corrected systematically by CH₄ satellite data from SCIAMACHY, GOSAT and IASI.


For all observations (surface, partial and total columns) CH₄ shows a seasonality in the relative difference between observations and the reanalysis. The magnitude of the difference is site dependent. During local autumn/winter months the relative bias is increased (underestimation) at most surface sites and in the tropospheric columns. This underestimation is also seen in the TCCON time series. In the spring and summer there is an overestimation of CH₄ near the surface and in the total column. These biases are related to errors in the seasonal cycle of surface emissions, most likely from agriculture and wetlands, and the accuracy of the representation of the hydroxyl radical (OH) sink which overall have larger values in the climatology compared to CAMS IFS(CB05BASCOE) atmospheric chemistry model OH (Segers et al., 2020b, Williams et al., 2021). The XCH₄ root mean square error is around 1.4 ppb and the Pearson correlation coefficient is larger than 0.7 for XCH₄ except for some outliers (Fig. 11, lower panel), indicating a good representation of the synoptic variability, as for XCO₂.




410

Figure 9. Mosaic plot of CH₄ biases (in ppb) compared to surface continuous observations from GLOBALVIEWplus CH₄ ObsPack v1.0 data product (Cooperative Global Atmospheric Data Integration Project, 2019) listed in Table A1. Each coloured vertical line represents a weekly mean. Vertical yellow lines depict the changes in the assimilated data documented in Figs 1, 5 and 6. Grey shading indicates no observations are available.

415



420 **Figure 10.** Mosaic plot of seasonal relative CH₄ biases at all FTIR sites (see Table A2) for the stratospheric columns (top) and tropospheric columns (bottom) NDACC. Vertical yellow lines depict the changes in the assimilated data documented in Figs 1, 5 and 6. Grey shading indicates no observations are available.



425 **Figure 11 Top: Mosaic plot of monthly biases at all TCCON sites for the column-averaged mole fractions XCH₄ [ppb] averaged daily around local noon (+/- 2.5 hours). Vertical yellow lines depict the changes in the assimilated data documented in Figs 1, 5 and 6. Grey shading indicates no observations are available. Bottom: Taylor diagrams for the station dependent XCH₄ comparison of the CAMS GHG reanalysis against TCCON FTIR data.**

430



3.2 Vertical profiles


435

The uncertainty of CAMS GHG reanalysis varies with height and the accuracy of the analysis vertical profiles depends mostly on the underlying model uncertainty, as the satellite data assimilated in reanalysis only provide integrated total or partial atmospheric column. The reanalysis has been evaluated using observations of CO₂ and CH₄ vertical profiles (Karion et al., 2010; Baier et al., 2021) from the NOAA AirCore dataset v20210813. It includes 133 vertical profiles from the surface to the
440 lower stratosphere (up to around 40 hPa) from 2012 to 2020 at 7 sites listed in Table A.3.

445


Figure 12 shows that the largest mean error occurs (i) near the surface with a strong influence from surface fluxes; (ii) in Upper Troposphere/Lower Stratosphere (UTLS) region (between 500 hPa and 100 hPa) with a strong influence from long-range transport; and (iii) in the stratosphere (above 100 hPa) where uncertainties associated with chemical loss of CH₄ and the meteorology driving the tracer transport are largest, and the fact that satellite data used here are not able to constrain the stratospheric CO₂ and CH₄ in the reanalysis. Near the surface, there is a positive CO₂ bias associated with an overestimation of the total flux in the model and a negative CH₄ bias which stems from both errors in the emissions and the chemical loss rate in the troposphere. The negative CO₂ bias in the UTLS agree with the tendency of the model to underestimate fine-scale higher-valued CO₂ streamers associated with long-range transport. The large positive CH₄ bias in the stratosphere of around 200 ppb
450 is consistent with the positive biases with respect to NDACC stratospheric column (Fig. 10, upper panel) and the documented model biases with respect to MIPAS and ACE-FTS by Verma et al. (2017). The errors associated with the stratospheric chemical sink are thought to be the largest contributor to the stratospheric CH₄ bias as shown by tests using the IFS BASCOE-CB05 chemical loss rate (not shown here). In general, the reanalysis underestimates the CO₂ vertical gradient across the tropopause. This underestimation leads to a positive bias for CO₂ in the lower stratosphere of around 2 ppm. The analysis is
455 not able to remove the large errors near the surface by only adjusting the atmospheric mole fractions, i.e., without adjusting the emissions in the data assimilation process, nor it is able to reduce the stratospheric errors in the model (Massart et al. 2017, Verma et al. 2017). The vertical profiles have a large variability from day to day as shown in Figure 13 with a sequence of profiles at Traînou (France). The CAMS GHG reanalysis is able to capture these synoptic variations in the vertical profile, consistent with its skill to represent XCO₂ and XCH₄ synoptic variability (lower panels of Figs 8 and 11).

460



465

Figure 12: Vertical profiles of mean error (Model M - Observation O) of CAMS CO₂ (left) and CH₄ (right) reanalysis with respect to AirCore observations comprising 133 profiles at 7 sites (listed in Table A3) over the period from 2012 and 2020. The blue shading shows the +/- standard deviation of M-O with respect to the mean error. The vertical dash black line depicts the zero mean error.



470


Figure 13: Vertical mole fraction profiles of CO₂ and CH₄ from the CAMS GHG reanalysis (dash line) and AirCore observations (solid line) at Trainou (France, see Table A3) over the period in June 2019.


475



3.3 Trends

Although this reanalysis is using a consistent underlying model and re-processed observations of CO₂ and CH₄, the current system is not able to provide accurate enough atmospheric mole fraction that can be used to estimate trends and the atmospheric growth rate as computed by the changes in global mean CO₂ and CH₄ from one year to the next. The CO₂ and CH₄ global annual means based on Marine Boundary Layer (MBL) reference sites are compared to the NOAA Global Greenhouse Gas Reference Network (GGGRN) observations (<https://gml.noaa.gov/ccgg/about.html>, Andrews et al., 2014; Conway et al., 1994; Dlugokencky et al., 1994) in Fig. 14. Changes in the assimilated satellite data have a clear impact on the evolution of the global annual mean values of CO₂ and CH₄ in the CAMS GHG reanalysis. The reanalysis has a positive global bias in near-surface CO₂ and CH₄ of a few ppm and around 20 ppb respectively from 2003 to 2007. Note that this positive bias in the annual global mean does not imply that the bias will be positive everywhere, as shown by the negative surface CH₄ biases at the AirCore sites (Fig. 12) and the large temporal and geographical variability of the weekly bias illustrated in Figs 7 and 9. After the introduction of IASI in 2007 the global bias decreases and it is lowest during the period when the number of observations is largest in 2013 and 2014 (Figs 5 and 6). Finally, the change to the near-real time satellite retrievals in 2019 together with the incorrect trend in the emissions during the COVID slowdown period in 2020 (Le Quéré et al., 2020) lead to changes in the global bias from negative to positive for CO₂ and from positive to negative for CH₄. These changes in the global bias are consistent with the changes in the errors with respect to total-column and near-surface observations in Figs. 7 to 11. It is important to note that the changes in global bias associated with changes in the assimilated data are of the same order of magnitude as the observed atmospheric growth rate of CO₂ (gml.noaa.gov/ccgg/trends) and CH₄ (gml.noaa.gov/ccgg/trends_ch4). For this reason, this reanalysis product is not suitable for trend analysis.





500 **Figure 14: Global mean CO₂ [ppm] and CH₄ [ppb] from the CAMS GHG reanalysis (in red) and the NOAA global mean CO₂ and CH₄ (in black, <https://gml.noaa.gov/ccgg/trends/global.html>) based on the Greenhouse Gas Marine Boundary layer Reference (Conway et al., 1994, Dlugokencky et al., 1994, Massarie et al., 1995, Dlugokencky et al., 2021). The global mean of the CAMS GHG reanalysis has been computed based on the same NOAA Marine Boundary Layer (MBL) reference sites shown in the map (bottom panel, see <https://gml.noaa.gov/ccgg/mb/mb.html> for further details). The dash blue lines mark the years when there was a change in the observing system. The uncertainty associated with the computation of global mean using the MBL sites is estimated to be 0.1 ppm for CO₂ (Ed Dlugokencky and Pieter Tans, NOAA/GML, gml.noaa.gov/ccgg/trends/) and below 2 ppb for CH₄ (Ed Dlugokencky, NOAA/GML (gml.noaa.gov/ccgg/trends_ch4/)).**

505

510 4. Limitations and caveats

This section provides an overview of the shortcomings of the CAMS GHG reanalysis which users should consider when interpreting the data. The main issues documented in the previous sections are summarised below:

1. Emissions are prescribed and not adjusted by the data assimilation system in the CAMS reanalysis (Sect. 2.3). This leads to a growing model error for CO₂ and CH₄ that can be difficult to correct with a sparse observing system and 12-hour assimilation window. In addition, the prescribed emissions are not available in near-real time, which means they are either kept fixed since the last year available (e.g. 2010 for CH₄) or they are extrapolated with a climatological trend as done for CO₂ (see details in Sect. 2.3). Because of this, the CAMS GHG reanalysis is not suitable to investigate the impact of local emission changes, such as COVID impact studies, which require a large local emission adjustments to the prescribed inventories (e.g. Doumia et al., 2021) and atmospheric inversion systems to estimate the changes (e.g. McNorton et al, 2022).
 2. Changes of satellite data used with different temporal, horizontal and vertical coverage cause changes in the quality of the reanalysis. For example, winter seasons have a lower number of observations because of light conditions and the higher frequency of cloud presence. This affects the quality of the seasonal cycle and the inter-hemispheric gradient. Similarly, in regions where there is no observation coverage, such as the stratosphere, the reanalysis is based on the underlying model including its systematic errors (see discussion on stratospheric biases in Sect 3.2).
- 515
- 520
- 525



3. Changes in satellite retrievals affect the quality of the observations used in the CAMS GHG reanalysis. For example, the switch from the CCI re-processed satellite products to the near-real time products is associated with a marked change in the bias and random error (i.e. standard deviation) of the departures from XCO₂ and XCH₄ GOSAT observations, as well as in the bias of the departures from the XCO₂ IASI-B observations. This large increase in the bias of the assimilated CO₂ and CH₄ observations from 2019 onwards results into a large increase in the bias of the CAMS GHG reanalysis in 2019 and 2020 which has implications for the trend analysis (Sect. 3.3).
4. The fixed climatological chemical loss rate of CH₄ (Sect 2.3) has been shown to overestimate the atmospheric CH₄ chemical sink by Segers et al. (2020b). Preliminary tests coupling the IFS to the atmospheric loss rate derived from BASCOE-CB05 chemistry have indeed shown a large reduction in the CH₄ negative bias in mid-latitudes. Systematic errors in the CH₄ chemical sink used in this reanalysis may have contributed further to enhance the large negative CH₄ bias in the CAMS GHG reanalysis over the last period in 2020, when the increase in the CH₄ growth rate has been linked to a decrease in chemical loss rate (Stevenson et al., 2021).
5. The large CH₄ and CO₂ biases in the stratosphere are currently under investigation. The CH₄ stratospheric bias is mainly associated with the use of a climatological loss rate (Sect 2.3), as preliminary tests using a different chemical loss rate based on IFS CB05BASCOE simulations show that the bias in CH₄ is greatly reduced.
6. Changes in systematic errors with time due to model error and changes in observation coverage and quality will affect trend analysis (see Sect. 3.3).

545 An up-to-date list of the known issues of the CAMS reanalysis can be found in the online CAMS documentation website (<https://confluence.ecmwf.int/display/CKB/CAMS%3A+Reanalysis+data+documentation>). Some of these issues will also be addressed in the future CAMS reanalysis (planned to start production in 2024), including the improvement of the prescribed emission trends, the consistent use of satellite retrieval products and the use of variable CH₄ chemical loss rate.

550 **5. Summary and conclusions**

This technical report documents the first CAMS IFS reanalysis of CO₂ and CH₄ produced by ECMWF which complements the CAMS reanalysis of reactive gases and aerosols (Inness et al., 2019). The processing chain, assimilated satellite data and underlying model used are described and the resulting reanalysis is evaluated using independent in situ near-surface observations, total column retrievals and in situ atmospheric profile observations. The monthly systematic and random errors of CO₂ and CH₄ typically range within 1% from 2003 to 2020 with an overall good skill in the representation of synoptic spatial variability and seasonal cycle. The lowest systematic errors occur in the period with maximum number of observations in 2013 and 2014. In 2019 there was a switch from C3S pre-processed satellite products to the near-real time CAMS satellite products because at the time of production the C3S products had not reached 2019. This caused a jump in the quality of the



560 satellite data and the resulting CAMS GHG reanalysis. For this reason, a new re-run of the CAMS GHG reanalysis from 2019 onwards will be performed with consistent C3S satellite products in the near future.

565 The comparison of global mean values with observations shows variations in the bias that depend on changes in the assimilated satellite data of around 2 ppm and 10 ppb for CO₂ and CH₄ respectively, which have the same magnitude as the observed variations in their growth rate. For this reason, we do not recommend the use of this dataset to study changes in the atmospheric growth rate of CO₂ and CH₄. Similarly, large biases in stratospheric CO₂ and CH₄ should also be considered when analysing stratospheric signals and trends in the CAMS GHG reanalyses. A list of caveats and limitations that users need to be aware of is provided in Sect. 4.

570 The slow reduction of the lingering bias in the model background is associated with competing factors at play: (i) the error growth in the model background associated with the accumulation of systematic errors in emission and natural fluxes; (ii) the limited coverage of observations in time and space (both horizontal and vertical); (iii) the localised impact of observations associated with a short data assimilation window spanning 12 hours.

575 In order to improve the CAMS reanalysis in future releases we recommend the following actions: (i) increase the number and coverage of satellite data assimilated from additional satellite missions such as the Copernicus Sentinel-5 Precursor (S5P), Orbiting Carbon Observatory 2 and 3 (OCO-2, https://www.nasa.gov/mission_pages/oco2/; OCO-3, <https://www.jpl.nasa.gov/missions/orbiting-carbon-observatory-3-oco-3>) and Greenhouse gases Observing SATellite-2 (GOSAT-2, <https://global.jaxa.jp/projects/sat/gosat2>) as well as the latest re-processed satellite products from C3S; (ii) improve the underlying emissions, particularly the extrapolation in near-real time; (iii) couple the chemical loss rate with the CAMS reanalysis of chemical species (Inness et al. 2019); and (iv) use the IFS inversion capability (McNorton et al., 2022) being developed within the CoCO2 project (coco2-project.eu) for future re-analyses.

585



590 **Appendix A**

Table A.1 List of stations with in situ continuous observations of CO₂ and CH₄ from GLOBALVIEWplus CO₂ ObsPack v6.0 and CH₄ ObsPack v1.0 respectively used for the evaluation in Sect. 3.1.

Station, Country (site name)	Latitude/Longitude [degrees]	Elevation [m asl]	Data Reference
Alert, Canada (ALT)	82.45 62.51W	185	Worthy et al. (2003)
Barrow, AK, USA (BRW)	71.32N 156.61W	11	Peterson et al. (1986)
Candle Lake, Canada (CDL)	53.99N 105.12W	600	Worthy et al. (2003)
Mace Head, Ireland (MHD)	53.33N 9.90W	5	Ramonet et al. (2010)
Fraserdale, Canada (FSD)	49.88N 81.57W	210	Worthy et al. (2003)
Kasprowy Wierch, Poland (KAS)	49.23N 19.98E	1989	Rozanski et al. (2003)
Schauinsland, Germany (SSL)	47.92N 7.92E	1205	Schmidt et al. (2003)
Hegyhatsal, Hungary (HUN)	46.95N 16.65	248	Haszpra et al (2001)
Park Falls, WI, USA (LEF)	45.95N 90.27W	472	Andrews et al. (2014)
Puy de Dôme, France (PUY)	45.77N 2.97E	1465	Lopez et al. (2015); Colomb et al. (2020)
Argyle, ME, USA (AMT)	45.03N 68.68W	53	Andrews et al. (2014)
Sable Island, Canada (WSA)	43.93N 60.00W	5	Worthy et al. (2003)
Ryori, Japan (RYO)	39.03N 141.82E	260	Tsutsumi et al. (2005)
Moody, TX, USA (WKT)	31.31N 97.33W	251	Andrews et al. (2014)
Minamitorishima, Japan (MNM)	24.28N 153.98E	8	Tsutsumi et al. (2005)
Yonagunijima, Japan (YON)	24.47N 123.02E	30	Tsutsumi et al. (2005)
Tutuila, American Samoa (SMO)	14.25S 170.56W	42	Waterman et al. (1989)
Cape Point, South Africa (CPT)	34.35S 18.49E	230	Brunke et al. (2004)
Amsterdam Island, France (AMS)	37.80S 77.54E	55	Ramonet et al. (1996)
Baring Head Station, New Zealand (BHD)	41.41S 174.87E	85	Stephens et al. (2013)
Syowa Station, Antarctica, Japan (SYO)	69.01S 39.59E	14	Schuldt et al. (2020)
South Pole, Antarctica, USA (SPO)	89.98S 24.8W	2810	Conway et al. (1990)



Table A.2 List of total column stations used for the evaluation in Sect 3.1.

Station, country	Latitude/ Longitude [degrees]	Network	Data references
Eureka, Canada	80.05N 86.42W	TCCON+NDACC	Strong et al. (2019); Batchelor et al. (2009)
Ny Ålesund, Norway	78.9N 11.9E	TCCON+NDACC	Notholt et al., (2019)
Thule, Greenland	76.53N 68.74W	NDACC	Hannigan et al. (2009)
Kiruna, Sweden	67.84N 20.41E	NDACC	Bader et al. (2017)
Sodankylä, Finland	67.37N 26.63E	TCCON+NDACC	Kivi et al. (2014); Sha et al. (2021)
Harestua, Norway	60.2N 10.8E	NDACC	De Mazière et al. (2018)
St Petersburg, Russia	59.90N 29.80E	NDACC	Makarova et al. (2015)
East Trout Lake, Canada	54.35N 104.99W	TCCON	Wunch et al. (2018)
Bialystok, Poland	53.23N 23.02E	TCCON	Deutscher et al. (2015)
Bremen, Germany	53.1N 8.85E	TCCON	Notholt et al. (2014)
Karlsruhe, Germany	49.1N 8.44E	TCCON	Hase et al. (2015)
Paris, France	48.85N 2.36E	TCCON	Te et al. (2014)
Orléans, France	47.97N 2.11E	TCCON	Warneke et al. (2014)
Garmisch, Germany	47.48N 11.06E	TCCON+NDACC	Sussmann and Rettinger (2018a); Sussmann et al. (2012); Hausmann al. (2016)



Zugspitze, Germany	47.42N 10.98E	TCCON+NDACC	Sussmann and Rettinger (2018b)
Jungfrauoch, Switzerland	46.55N 7.98E	NDACC	Zander et al. (2008)
Park Falls, WI, USA	45.94N 90.27W	TCCON	Wennberg et al. (2017)
Rikubetsu, Japan	43.46N 143.77E	TCCON+NDACC	Morino et al. (2016); De Mazière et al. (2018)
Boulder, CO, USA	39.99N 105.26W	NDACC	Ortega et al. (2021)
Indianapolis, IN, USA	39.86N 86W	TCCON	Iraci et al. (2016)
Four Corners, USA	36.8N 108.48W	TCCON	Dubey et al. (2014)
Lamont, OK, USA	36.5N 97.49W	TCCON	Wennberg et al. (2016)
Anmeyondo, South Korea	36.54N 126.33E	TCCON	Goo et al. (2014)
Tsukuba, Japan	36.05N 140.12E	TCCON	Morino et al (2016)
Nicosia, Cyprus	35.14N 33.38E	TCCON	Petri et al. (2020)
Edwards, CA, USA	34.96N 117.88W	TCCON	Iraci et al. (2016)
JPL, CA, USA	34.2N 118.18W	TCCON	Wennberg et al. (2016)
Pasadena Caltech, CA, USA	34.14N 118.13W	TCCON	Wennberg et al. (2015)
Saga, Japan	33.24N 130.29E	TCCON	Kawakami et al. (2014)
Heifei, China	31.9 N 117.17E	TCCON	Liu et al. (2018)



Izaña, Spain	28.3N 16.48W	TCCON+NDACC	Blumenstock et al. (2014); García et al. (2021)
Mauna Loa, HI, United States	19.54N 155.58W	NDACC	Hannigan et al. (2009)
Altzomoni, Mexico	19.12N 98.66W	NDACC	De Mazière et al. (2018);
Burgos, Philippines	18.53N 120.65E	TCCON	Morino et al. (2018)
Manaus, Brazil	3.21S 60.6W	TCCON	Dubey et al. (2014)
Ascension Island, UK	7.92S 14.33W	TCCON	Feist et al. (2014)
Darwin, Australia	12.43S 130.89E	TCCON	Griffith et al. (2014)
Reunion St Denis, France	20.9S 55.49E	TCCON+NDACC	De Mazière et al. (2014)
Reunion Island, Maido, France	21.1S 55.4E	NDACC	Zhou et al. (2018)
Wollongong, Australia	34.41S 150.88E	TCCON+NDACC	Griffith et al. (2014); De Mazière et al. (2018);
Lauder, New Zealand	45.05S 168.68E	TCCON+NDACC	Sherlock et al. (2014a, 2014b); Pollard et al. (2019) Bader et al. (2017) ; Pollard et al. (2017)
Arrival Heights, Antarctica	77.83S 166.67E	NDACC	Bader et al. (2017)



605 **Table A.3 List of AirCore sites (from NOAA_AirCore_data_v20210813, Baier et al., 2021) used for the evaluation in Sect. 3.2.**

Site, country	Latitude /Longitude [degrees]
Boulder, CO, USA	40.03N 103.74W
Lamont, OK, USA	36.85N 98.21W
Lauder, New Zealand	45.50S 169.47E
Sodankylä, Finland	67.41N 26.31E
Park Falls, WI, USA	45.97N 90.32W
Edwards, CA, USA	34.65N 117.29W
Traînou, France	48.48N 1.16E

Code and data availability

The IFS forecasting and reanalysis system is not for public use as the ECMWF Member States are the proprietary owners. The resulting dataset is however freely available on the Copernicus Atmosphere Data Store. The CAMS GHG reanalysis can be accessed through the CAMS Atmosphere Data Store (ADS) at <https://doi.org/10.24380/8fck-9w87>. The format is available in both GRIB and NetCDF. The data record starts on 1 January 2003 00UTC and currently stops on 31 December 2020. Recent months will be added over time as soon as the reanalysis procedure and its validation are completed. The original data was available either as spectral coefficients with a triangular truncation of T255 or on a reduced Gaussian grid with a resolution of N128. But for the ease of the user, fields were interpolated from their native representation to a regular 0.75°x0.75° latitude longitude grid. For sub-daily data for the CAMS reanalysis is archived with a 3-hourly time step (0, 3, 6, 9, 12, 15, 18, 21 UTC). Pre-computed monthly averages are also directly available for all fields. The 3D fields are available on two different vertical resolution: 25 pressure levels (1000, 950, 925, 900, 850, 800, 700, 600, 500, 400, 300, 250, 200, 150, 100, 70, 50, 30, 20, 10, 7.5, 3, 2, 1 hPa) and 60 σ -hybrid model levels which are described at <https://www.ecmwf.int/en/forecasts/documentation-and-support/60-model-levels>. The data records have 18 2D radiation fields, 2 vertically integrated atmospheric content of CO₂ and CH₄ (column-mean mole fractions, 14 2D surface fluxes variables, 32 2D meteorological fields and 16 3D fields including meteorological and greenhouse gases fields. A complete listing of the variables included in the CAMS GHG reanalysis is provided in the ADS (<https://ads.atmosphere.copernicus.eu/cdsapp#!/home>).

625



Acknowledgements

The Copernicus Atmosphere Monitoring Service is operated by the European Centre for Medium-Range Weather Forecasts on behalf of the European Commission as part of the Copernicus Programme (<http://copernicus.eu>). The satellite data assimilated in the CAMS GHG reanalysis were obtained from the ESA GHG-CCI project
630 (<https://climate.esa.int/en/projects/ghgs/Data>) and the Copernicus Climate Change Service (C3S) Climate Data Store (<https://cds.climate.copernicus.eu>). The data used to evaluate the CAMS GHG reanalysis were obtained from: the Total Carbon Column Observing Network (TCCON) Data Archive hosted by CaltechDATA at <https://tccodata.org>; CH₄ from the Network for the Detection of Atmospheric Composition Change (NDACC, www.ndacc.org; Sussmann et al., 2011; 2013; De Mazière et al., 2018) (see Table A2); the Observation Package (ObsPack) GLOBALVIEWplus Data Products
635 <https://gml.noaa.gov/ccgg/obspack>; the NOAA AirCore dataset <https://gml.noaa.gov/ccgg/arc/?id=144>; and the NOAA Greenhouse Gas Marine Boundary Layer Reference <https://gml.noaa.gov/ccgg/mbl/mbl.html>.

For the NDACC data, the National Center for Atmospheric Research is sponsored by the National Science Foundation. The NCAR FTS observation programs at Thule, GR, Boulder, CO and Mauna Loa, HI are supported under contract by the
640 National Aeronautics and Space Administration (NASA). The Thule work is also supported by the NSF Office of Polar Programs (OPP). We wish to thank the Danish Meteorological Institute for support at the Thule site and NOAA for support of the MLO site. The multi-decadal monitoring program of University of Liège at the Jungfraujoch station has been primarily supported by the F.R.S.-FNRS and BELSPO (both in Brussels, Belgium) and by the GAW-CH programme of MeteoSwiss. The International Foundation High Altitude Research Stations Jungfraujoch and Gornergrat (HFSJG, Bern)
645 supported the facilities needed to perform the FTIR observations at Jungfraujoch.

The authors are also grateful to László Haszpra for his comments on the Hegyhátsál tall-tower station, Xin Lan and Ed Duglokencky for the feedback on the use of NOAA/GML MBL data; Debra Wunch for providing insight into the potential impact of polar vortex on the XCH₄ data at East Trout Lake TCCON site.

650 Author contributions

Anna Agustí-Panareda, Jérôme Barré and Sébastien Massart prepared the manuscript with the rest of the co-authors, monitored the simulation, testing the impact of the surface fluxes and assimilated observations, as well as fixing bugs. Anna Agustí-Panareda was responsible of the forecasting model implementation and performed the evaluation of the vertical profiles and trends. Antje Inness led the CAMS reanalysis work and was responsible of running and monitoring the reanalysis simulations.
655 Sébastien Massart developed the data assimilation of satellite products. Mark Parrington provided support on the biomass burning emissions and postprocessed the fire emission data in Fig. 3. Mel Ades worked on documenting the data assimilation aspects of the GHG reanalysis. Johannes Flemming, Zak Kipling and Luca Cantarello provided technical support for the diagnostics, bug fixes implemented during the testing stage and testing of the CAMS NRT satellite observations. Luke Jones provided technical support with the mass interpolation scheme and diagnostic tools. Miha Ratzinger was responsible for post-
660 processing the dataset into the Copernicus Atmosphere Data Store. Roberto Ribas and Martin Suttie performed the satellite data acquisition and pre-processing. Richard Engelen and Vincent-Henri Peuch coordinated the efforts on the CAMS reanalyses. Bavo Langerock, Henk Eskes, Michel Ramonet and Jérôme Tarniewicz were responsible of validating the reanalysis dataset with independent observations including the production of surface and column CO₂ and CH₄ evaluation plots. Ilse Aben, Tobias Borsdorff, Michael Buchwitz, Cyril Crevoisier, Otto Hasekamp, Nicolas Meilhac, Stefan Noel,



665 Maximilian Reuter and Lianghai Wu provided support for the use of the satellite retrieval products. All authors reviewed and edited the manuscript.

670 References

Agustí-Panareda, A., Diamantakis, M., Bayona, V., Klappenbach, F. & Butz, A.: Improving the inter-hemispheric gradient of total column atmospheric CO₂ and CH₄ in simulations with the ECMWF semi-Lagrangian atmospheric global model. *Geoscientific Model Development* **10**, <https://doi.org/10.5194/gmd-10-1-2017>, 1–18, 2017.

675

Agustí-Panareda, A., Diamantakis, M., Massart, S., Chevallier, F., Muñoz-Sabater, J., Barré, J., Curcoll, R., Engelen, R., Langerock, B., Law, R. M., Loh, Z., Morguí, J. A., Parrington, M., Peuch, V.-H., Ramonet, M., Roehl, C., Vermeulen, A. T., Warneke, T., and Wunch, D.: Modelling CO₂ weather – why horizontal resolution matters, *Atmos. Chem. Phys.*, **19**, 7347–7376, <https://doi.org/10.5194/acp-19-7347-2019>, 2019.

680

Agustí-Panareda, A., McNorton, J., Balsamo, G., Baier, B.C., Bousserez, N., Boussetta, S., Brunner, D., Chevallier, F., Choulga, M., Diamantakis, M., Engelen, R., Flemming, J., Granier, C. Guevara, M., Denier van der Gon, H., Elguindi, N., Haussaire, J.-M., Jung, M., Janssens-Maenhout, G., Kivi, R., Massart, S., Papale, D., Parrington, M., Razinger, M., Sweeney, C., Vermeulen, A., Walther, S. Global nature run data with realistic high-resolution carbon weather for the year of the Paris Agreement. *Sci Data* **9**, 160 (2022). <https://doi.org/10.1038/s41597-022-01228-2>, 2022.

685

Alos, A. L. *et al.* The Copernicus Climate Data Store: ECMWF's approach to providing online access to climate data and tools. in *AGU Fall Meeting Abstracts* vol. 2019 PA31A-05, 2019.

690 Andrews, A. E., Kofler, J. D., Trudeau, M. E., Williams, J. C., Neff, D. H., Masarie, K. A., Chao, D. Y., Kitzis, D. R., Novelli, P. C., Zhao, C. L., Dlugokencky, E. J., Lang, P. M., Crotwell, M. J., Fischer, M. L., Parker, M. J., Lee, J. T., Baumann, D. D., Desai, A. R., Stanier, C. O., De Wekker, S. F. J., Wolfe, D. E., Munger, J. W., and Tans, P. P.: CO₂, CO, and CH₄ measurements from tall towers in the NOAA Earth System Research Laboratory's Global Greenhouse Gas Reference Network: instrumentation, uncertainty analysis, and recommendations for future high-accuracy greenhouse gas monitoring efforts, *Atmos. Meas. Tech.*, **7**, 647–687, <https://doi.org/10.5194/amt-7-647-2014>, 2014.

695

Bader, W., Bovy, B., Conway, S., Strong, K., Smale, D., Turner, A. J., Blumenstock, T., Boone, C., Collaud Coen, M., Coulon, A., Garcia, O., Griffith, D. W. T., Hase, F., Hausmann, P., Jones, N., Krummel, P., Murata, I., Morino, I., Nakajima, H.,



O'Doherty, S., Paton-Walsh, C., Robinson, J., Sandrin, R., Schneider, M., Servais, C., Sussmann, R. and Mahieu, E.: The
700 recent increase of atmospheric methane from 10 years of ground-based NDACC FTIR observations since 2005, *Atmos. Chem.
Phys.*, 17(3), 2255–2277, doi:10.5194/acp-17-2255-2017, 2017.

Baier, B., Sweeney, C., Newberger, T., Higgs, J., Wolter, S., & NOAA Global Monitoring Laboratory: NOAA AirCore
705 atmospheric sampling system profiles (Version 20210813) [Data set]. NOAA GML. <https://doi.org/10.15138/6AV0-MY81>,
2021.

Batchelor, R. L., Strong, K., Lindenmaier, R., Mittermeier, R. L., Fast, H., Drummond, J. R. and Fogal, P. F.: A New Bruker
IFS 125HR FTIR Spectrometer for the Polar Environment Atmospheric Research Laboratory at Eureka, Nunavut, Canada:
Measurements and Comparison with the Existing Bomem DA8 Spectrometer, *J. Atmos. Ocean. Technol.*, 26(7), 1328–1340,
710 doi:10.1175/2009JTECHA1215.1, 2009.

Bechtold, P. *et al.* Representing Equilibrium and Nonequilibrium Convection in Large-Scale Models. *Journal of the
Atmospheric Sciences* **71**, 734–753, 2014.

715 Bergamaschi, P., M. Krol, F. Dentener, A. Vermeulen, F. Meinhardt, R. Graul, M. Ramonet, W. Peters, and E. J. Dlugokencky,
Inverse modelling of national and European CH₄ emissions using the atmospheric zoom model TM5, *Atmos. Chem. Phys.*, 5,
2431–2460, 2005.

Bergamaschi, P. *et al.* : Inverse modeling of global and regional CH₄ emissions using SCIAMACHY satellite retrievals.
720 *Journal of Geophysical Research: Atmospheres* **114**, 2009.

Blumenstock, T., Hase, F., Schneider, M., Garcia, O. & Sepulveda, E. TCCON data from Izana (ES), Release GGG2014.R0
(Version GGG2014.R0) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.IZANA01.R0/1149295>, 2014.

725 Boussetta, S. *et al.*: Natural land carbon dioxide exchanges in the ECMWF integrated forecasting system: Implementation and
offline validation. *Journal of Geophysical Research: Atmospheres* **118**, 5923–5946, 2013.

Brühl, C., and P. J. Crutzen, The MPIC 2D model, *NASA Ref. Publ.*, **1292**, 103–104, 1993.

730 Brunke, E-G., Labuschagne, C., Parker, B., Scheel, H.E. and Whittlestone, S. Baseline air mass selection at
Cape Point, South Africa: Application of 222Rn and other filter criteria to CO₂. *Atmospheric Environment* 38, 33, 5693 –
5702, 2004.



735 Buchwitz M, Reuter M, Schneising O, Boesch H, Guerlet S, Dils B, Aben I, Armante R, Bergamaschi P, Blumenstock T,
Bovensmann H, Brunner D, Buchmann B, Burrows J, Butz A, Chédin A, Chevallier F, Crevoisier C, Deutscher N, Frankenberg
C, Hase F, Hasekamp O, Heymann J, Kaminski T, Laeng A, Lichtenberg G, Demazière M, Noël S, Notholt J, Orphal J, Popp
C, Parker R, Scholze M, Sussmann R, Stiller G, Warneke T, Zehner C, Bril A, Crisp D, Griffith D, Kuze A, O'dell C,
Oshchepkov S, Sherlock V, Suto H, Wennberg P, Wunch D, Yokota T, Yoshida Y. The Greenhouse Gas Climate Change
740 data sets. REMOTE SENSING OF ENVIRONMENT 162; 2015. p. 344-362. JRC91437, 2015.

Butz, A., Hasekamp, O. P., Frankenberg, C., Vidot, J., and Aben, I., CH₄ retrievals from space-based solar backscatter
measurements: Performance evaluation against simulated aerosol and cirrus loaded scenes, J. Geophys. Res., Volume 115,
Issue D24, 27, DOI: 10.1029/2010JD014514, 2010.

745

Butz, A., Guerlet, S., Hasekamp, O., et al., Toward accurate CO₂ and CH₄ observations from GOSAT, Geophys. Res. Lett.,
doi:10.1029/2011GL047888, 2011.

Chevallier, F., Remaud, M., O'Dell, C. W., Baker, D., Peylin, P., and Cozic, A.: Objective evaluation of surface- and satellite-
750 driven carbon dioxide atmospheric inversions, Atmos. Chem. Phys., 19, 14233–14251, [https://doi.org/10.5194/acp-19-14233-](https://doi.org/10.5194/acp-19-14233-2019)
2019.

Chevallier, F., et al.: CO₂ surface fluxes at grid point scale estimated from a global 21-year reanalysis of atmospheric
measurements, J. Geo-phys. Res., doi:10.1029/2010JD013887, 2010.

755

Chevallier, F. *Description of the CO₂ inversion production chain 2020*, Copernicus Atmosphere Monitoring
Service. [https://atmosphere.copernicus.eu/sites/default/files/2020-](https://atmosphere.copernicus.eu/sites/default/files/2020-06/CAMS73_2018SC2_%20D5.2.12020_202004_%20CO2%20inversion%20production%20chain_v1.pdf)
[06/CAMS73_2018SC2_%20D5.2.12020_202004_%20CO2%20inversion%20production%20chain_v1.pdf](https://atmosphere.copernicus.eu/sites/default/files/2020-06/CAMS73_2018SC2_%20D5.2.12020_202004_%20CO2%20inversion%20production%20chain_v1.pdf), 2000.

760 Colomb, A., Delmotte, M., Pichon, J., Ramonet, M., Yver-Kwok, C., ICOS RI, 2020. ICOS ATC CO₂ Release, Puy de Dôm
e (10.0 m), 2016-08-25–2020-05-31, <https://hdl.handle.net/11676/xMV5-Hxu7bA9q872kedNJlAF>, 2020.

Conway, T. J. and K. W. Thoning. Short-term variations of atmospheric carbon dioxide at the South Pole, Antar
ctic Journal of the United States, 236-238, 1990.

765



- Conway, T.J., P.P. Tans, L.S. Waterman, K.W. Thoning, D.R. Kitzis, K.A. Masarie, and N. Zhang: Evidence for interannual variability of the carbon cycle from the NOAA/CMDL global air sampling network, *J. Geophys. Res.*, 99, 22831-22855, 1994.
- Crevoisier, C. *et al.*: First year of upper tropospheric integrated content of CO₂ from IASI hyperspectral infrared observations. 770 *Atmospheric Chemistry and Physics* **9**, 4797–4810, 2009a.
- Crevoisier, C. *et al.*: Tropospheric methane in the tropics – first year from IASI hyperspectral infrared observations. *Atmospheric Chemistry and Physics* **9**, 6337–6350, 2009b.
- 775 Crevoisier, C., Clerbaux, C., Guidard, V., Phulpin, T., Armante, R., Barret, B., Camy-Peyret, C., Chaboureau, J.-P., Coheur, P.-F., Crépeau, L., Dufour, G., Labonnote, L., Lavanant, L., Hadji-Lazaro, J., Herbin, H., Jacquinet-Husson, N., Payan, S., Péquignot, E., Pierangelo, C., Sellitto, P., and Stubenrauch, C.: Towards IASI-New Generation (IASI-NG): impact of improved spectral resolution and radiometric noise on the retrieval of thermodynamic, chemistry and climate variables, *Atmos. Meas. Tech.*, 7, 4367–4385, <https://doi.org/10.5194/amt-7-4367-2014>, 2014.
- 780 Crisp and Co-authors : A constellation architecture for monitoring carbon dioxide and methane from space. CEOS Atmospheric Composition Virtual Constellation Greenhouse Gas Team Rep., 173 pp., http://ceos.org/document_management/Virtual_Constellations/ACC/Documents/CEOS_AC-VC_GHG_White_Paper_Version_1_20181009.pdf, 2018.
- 785 De Mazière, M. *et al.* TCCON data from Réunion Island (RE), Release GGG2014.R0 (Version GGG2014.R0) [Data set]. <http://dx.doi.org/10.14291/tcon.ggg2014.reunion01.R0/1149288>, 2014.
- De Mazière, M. *et al.*: The Network for the Detection of Atmospheric Composition Change (NDACC): history, status and perspectives. *Atmospheric Chemistry and Physics* **18**, 4935–4964, 2018.
- 790 Deutscher, N. M. *et al.* TCCON data from Bialystok (PL), Release GGG2014.R1 (Version GGG2014.R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.BIALYSTOK01.R1/1183984>, 2015.
- Diamantakis, M. & Magnusson, L. : Sensitivity of the ECMWF Model to Semi-Lagrangian Departure Point Iterations. *Monthly* 795 *Weather Review* **144**, 3233–3250, 2016.
- Doumbia, T., Granier, C., Elguindi, N., Bouarar, I., Darras, S., Brasseur, G. *et al.* Changes in global air pollutant emissions during the COVID-19 pandemic: a dataset for atmospheric modeling. *Earth System Science Data*, 13, 4191-4206. doi:10.5194/essd-13-4191-2021, 2021.



800

Dlugokencky, E.J., L.P. Steele, P.M. Lang, and K.A. Masarie, The growth rate and distribution of atmospheric methane, *J. Geophys. Res.*, 99, 17,021-17,043, 1994.

Dlugokencky, E.J., J.W. Mund, A.M. Crotwell, M.J. Crotwell, and K.W. Thoning. Atmospheric Carbon Dioxide Dry Air
805 Mole Fractions from the NOAA GML Carbon Cycle Cooperative Global Air Sampling Network, 1968-2020, Version: 2021-07-30, <https://doi.org/10.15138/wkgj-f215>, 2021.

Dubey, M. K. et al. TCCON data from Manaus (BR), Release GGG2014.R0 (Version GGG2014.R0) [Data set].
<https://doi.org/10.14291/TCCON.GGG2014.MANAUS01.R0/1149274.1> 2014.

810

Dubey, M., R. Lindenmaier, B. Henderson, D. Green, N. Allen, C. Roehl, J.-F. Blavier, Z. Butterfield, S. Love, J. Hamelmann, D. Wunch. TCCON data from Four Corners, NM, USA, Release GGG2014R0. TCCON data archive, hosted by CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A. <https://doi.org/10.14291/tcon.ggg2014.fourcorners01.R0/1149272>, 2014.

815

Feist, D. G., Arnold, John, & Geibel. TCCON data from Ascension Island (SH), Release GGG2014.R0 (Version GGG2014.R0) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.ASCENSION01.R0/1149285>, 2014.

Fisher, M.: Generalized frames on the sphere with application to background error covariance modelling, Seminar on recent
820 developments in numerical methods for atmospheric and ocean modelling, 6–10 September 2004, Proceedings, ECMWF, pp. 87–101, available from ECMWF, Shinfield Park, Reading, Berkshire, RG2 9AX, UK, 2004.

Fisher, M.: Wavelet Jb – A new way to model the statistics of background errors, ECMWF Newsletter, 106, 23–28, available from ECMWF, Shinfield Park, Reading, Berkshire, RG2 9AX, UK, 2006.

825

Flemming, J., Huijnen, V., Arteta, J., Bechtold, P., Beljaars, A., Blechschmidt, A.-M., Diamantakis, M., Engelen, R. J., Gaudel, A., Inness, A., Jones, L., Josse, B., Katragkou, E., Marecal, V., Peuch, V.-H., Richter, A., Schultz, M. G., Stein, O., and Tsikerdekis, A.: Tropospheric chemistry in the Integrated Forecasting System of ECMWF, *Geosci. Model Dev.*, 8, 975–1003, <https://doi.org/10.5194/gmd-8-975-2015>, 2015.

830

Frankenberg, C., Aben, I., Bergamaschi, P., Dlugokencky, E. J., van Hees, R., Houweling, S., van der Meer, P., Snel, R., and Tol, P.: Global column-averaged methane mixing ratios from 2003 to 2009 as derived from SCIAMACHY: trends and variability, *J. Geophys. Res.-Atmos.*, 116, D04302, doi:10.1029/2010JD014849, 2011.



- 835 Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A. K., Jones, S. D., Kato, E., Kennedy, D.,
840 Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.:
845 Global Carbon Budget 2021, *Earth Syst. Sci. Data Discuss.* [preprint], <https://doi.org/10.5194/essd-2021-386>, in review, 2021.

- García, O. E., Schneider, M., Sepúlveda, E., Hase, F., Blumenstock, T., Cuevas, E., Ramos, R., Gross, J., Barthlott, S., Röhling, A. N., Sanromá, E., González, Y., Gómez-Peláez, Á. J., Navarro-Comas, M., Puentedura, O., Yela, M., Redondas, A., Carreño, V., León-Luis, S. F., Reyes, E., García, R. D., Rivas, P. P., Romero-Campos, P. M., Torres, C., Prats, N., Hernández, M. and
850 López, C.: Twenty years of ground-based NDACC FTIR spectrometry at Izaña Observatory-overview and long-term comparison to other techniques, *Atmos. Chem. Phys.*, 21(20), 15519–15554, doi:10.5194/acp-21-15519-2021, 2021.

- Goo, T.-Y., Oh, Y.-S. & Velazco, V. A. TCCON data from Anmeyondo (KR), Release GGG2014.R0 (Version GGG2014.R0) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.ANMEYONDO01.R0/1149284>, 2014.

855

- Griffith, D. W. T. et al. TCCON data from Darwin (AU), Release GGG2014.R0 (Version GGG2014.R0) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.DARWIN01.R0/1149290>, 2014.

- Griffith, D. W. T. et al. TCCON data from Wollongong (AU), Release GGG2014.R0 (Version GGG2014.R0) [Data set].
860 <http://dx.doi.org/10.14291/tcon.ggg2014.wollongong01.R0/1149291>, 2014.

- Guerlet, S., Butz, A., Schepers, D., Basu, S., Hasekamp, O. P., Kuze, A., Yokota, T., Blavier, J.-F., Deutscher, N. M., Griffith, D. W., Hase, F., Kyro, E., Morino, I., Sherlock, V., Sussmann, R., Galli, A., and Aben, I.: Impact of aerosol and thin cirrus on retrieving and validating XCO₂ from GOSAT shortwave infrared measurements, *Journal of Geophysical Research: Atmospheres*, 118, 4887–4905, <https://doi.org/10.1002/jgrd.50332>, 2013.
865



Hannigan, J. W., M. T. Coffey, and A. Goldman. Semiautonomous FTS Observation System for Remote Sensing of Stratospheric and Tropospheric Gases. *Journal of Atmospheric and Oceanic Technology* 26, pp. 1814–1828. doi: 10.1175/2009JTECHA1230.1, 2009.

870

Hase, F., Blumenstock, T., Dohe, S., Gros, J. & Kiel, M. TCCON data from Karlsruhe (DE), Release GGG2014.R1 (Version GGG2014.R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.KARLSRUHE01.R1/1182416>, 2015.

875 Haszpra, L., Barcza, Z., Bakwin, P. S., Berger, B. W., Davis, K. J., Weidinger, T. Measuring system for the long-term monitoring of biosphere/atmosphere exchange of carbon dioxide. *J. of Geophysical Research* 106D, 3057–3070, 2001.

880 Hausmann, P., Sussmann, R., and Smale, D.: Contribution of oil and natural gas production to renewed increase in atmospheric methane (2007–2014): top–down estimate from ethane and methane column observations, *Atmos. Chem. Phys.*, 16, 3227–3244, doi:10.5194/acp-16-3227-2016, 2016.

Hersbach, H. *et al.*: The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society* **146**, 1999–2049, 2020.

885 Heymann, J., M. Reuter, M. Hilker, M. Buchwitz, O. Schneising, H. Bovensmann, J. P. Burrows, A. Kuze, H. Suto, N. M. Deutscher, M. K. Dubey, D. W. T. Griffith, F. Hase, S. Kawakami, R. Kivi, I. Morino, C. Petri, C. Roehl, M. Schneider, V. Sherlock, R. Sussmann, V. A. Velazco, T. Warneke, and D. Wunch, Consistent satellite XCO₂ retrievals from SCIAMACHY and GOSAT using the BESD algorithm, *Atmos. Meas. Tech.*, 8, 2961–2980, 2015.

890 Houweling, S., F. Dentener, and J. Lelieveld, The impact of nonmethane hydrocarbon compounds on tropospheric photochemistry, *J. Geophys. Res.*, 103(D9), 10,673–10,696, doi:10.1029/97JD03582, 1998.

Houweling, S., Kaminski, T., Dentener, F., Lelieveld, J. & Heimann, M. Inverse modeling of methane sources and sinks using the adjoint of a global transport model. *Journal of Geophysical Research: Atmospheres* **104**, 26137–26160, 1999.

895

ICOS RI: ICOS Atmospheric Greenhouse Gas Mole Fractions of CO₂ for the period Sep 2015- May 2020, final quality controlled Level 2 data for 23 stations (62 vertical levels), pre-release 2020-1, , doi:10.18160/GZ5S-4GPR, 2020.

900 Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy,



S., Schulz, M., and Suttie, M.: The CAMS reanalysis of atmospheric composition, *Atmos. Chem. Phys.*, 19, 3515–3556, <https://doi.org/10.5194/acp-19-3515-2019>, 2019.

905 IPCC, *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp., 2014.

IPCC, Summary for Policymakers. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press., 2021.
910

Iraci, L., J. Podolske, P. Hillyard, C. Roehl, P. O. Wennberg, J.-F. Blavier, J. Landeros, N. Allen, D. Wunch, J. Zavaleta, E. Quigley, G. Osterman, E. Barrow, J. Barney. TCCON data from Indianapolis, Indiana, USA, Release GGG2014R1. TCCON data archive, hosted by CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A. <https://doi.org/10.14291/tcon.ggg2014.indianapolis01.R1/1330094>, 2016.
915

Iraci, L. T. et al. TCCON data from Edwards (US), Release GGG2014.R1 (Version GGG2014.R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.EDWARDS01.R1/1255068>, 2016.
920

Janssens-Maenhout, Greet; Crippa, Monica; Guizzardi, Diego; Muntean, Marilena; Schaaf, Edwin. Emissions Database for Global Atmospheric Research, version v4.2 (time-series). European Commission, Joint Research Centre (JRC) [Dataset] PID: <http://data.europa.eu/89h/jrc-edgar-emissiontimeseriesv42>, 2011.
925

Janssens-Maenhout, G., Crippa, M.; Guizzardi, D.; Muntean, M.; Schaaf, E. JRC-EDGARv431_AP_timeseries. European Commission, Joint Research Centre (JRC) [Dataset] PID: http://data.europa.eu/89h/jrc-edgar-edgar_v431_timeseries, 2016.

Kaiser, J. W. *et al.*: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power. *Biogeosciences* 9, 527–554, 2012.
930

Karion, A., Sweeney, C., Tans, P., Newberger, T. AirCore: An Innovative Atmospheric Sampling System, *J. Atmos. Ocean. Technol.*, 27, 1839-1843, <https://doi.org/10.1175/2010JTECHA1448.1>, 2010.



935 Kawakami, S. et al. TCCON data from Saga (JP), Release GGG2014.R0 (Version GGG2014.R0) [Data set].
<https://doi.org/10.14291/TCCON.GGG2014.SAGA01.R0/1149283>, 2014.

Kivi, R., Heikkinen, P. & Kyro, E. TCCON data from Sodankylä (FI), Release GGG2014.R0 (Version GGG2014.R0) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.SODANKYLA01.R0/1149280>, 2014.

940

Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, A., and Zeng, G.: The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP):
945 overview and description of models, simulations and climate diagnostics, 6, 179–206, <https://doi.org/10.5194/gmd-6-179-2013>, 2013.

Lambert, G. & Schmidt, S.: Re-evaluation of the oceanic flux of methane: Uncertainties and long-term variations. *Chemosphere* **26**, 579–589, 1993.

950

Le Quéré, C., Jackson, R.B., Jones, M.W. et al.: Temporary reduction in daily global CO₂ emissions during the COVID-19 forced confinement. *Nat. Clim. Chang.* 10, 647–653, <https://doi.org/10.1038/s41558-020-0797-x>, 2020.

Liu C., Wang, W., Sun, Y. TCCON data from Hefei, China, Release GGG2014R0. TCCON data archive, hosted by
955 CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A.
<http://dx.doi.org/10.14291/tcon.ggg2014.hefei01.R0>, 2018.

Lopez, M., M. Schmidt, M. Ramonet, J. L. Bonne, A. Colomb, V. Kazan, P. Laj and J. M. Pichon. Three years of semicontinuous greenhouse gas measurements at the Puy de Dôme station (central France). *Atmospheric Measurement*
960 *Techniques*(8): 3941-3958, 2015.

Makarova, M. V., Kirner, O., Timofeev, Y. M., Poberovskii, A. V., Imkhasin, K. K., Osipov, S. I. and Makarov, B. K.: Analysis of methane total column variations in the atmosphere near St. Petersburg using ground-based measurements and simulations, *Izv. Atmos. Ocean. Phys.*, 51(2), 177–185, doi:10.1134/S0001433815010089, 2015.

965

Masarie, K. A., Peters, W., Jacobson, A. R. & Tans, P. P.: ObsPack: a framework for the preparation, delivery, and attribution of atmospheric greenhouse gas measurements. *Earth System Science Data* **6**, 375–384, 2014.



- 970 Masarie, K.A. and Tans, P.P.: Extension and Integration of Atmospheric Carbon Dioxide Data into a Globally Consistent Measurement Record, *J. Geophys. Res.*, .100, 11593-11610, 1995.
- Massart, S., and Bonavita, M.: Ensemble of Data Assimilations applied to the CAMS' greenhouse gases analysis. ECMWF Technical Memorandum 780, available at: <https://www.ecmwf.int/en/elibrary/16438-ensemble-data-assimilations-applied-cams-greenhouse-gases-analysis> (last access: xxx), 2016.
- 975 Massart, S. *et al.*: Assimilation of atmospheric methane products into the MACC-II system: from SCIAMACHY to TANSO and IASI. *Atmospheric Chemistry and Physics* **14**, 6139–6158, 2014.
- Massart, S. *et al.*: Ability of the 4-D-Var analysis of the GOSAT BESD XCO₂ retrievals to characterize atmospheric CO₂ at large and synoptic scales. *Atmospheric Chemistry and Physics* **16**, 1653–167, 2016.
- 980 Massart, S., Agusti-Panareda, A. and Flemming, J.: Evidence of a stratospheric methane bias in the IFS against MIPAS data, ECMWF Technical Memorandum, 814, 21 pages, <https://www.ecmwf.int/sites/default/files/elibrary/2017/17698-evidence-stratospheric-methane-bias-ifs-against-mipas-data.pdf>, 2017.
- 985 Matthews, E., Fung, I. & Lerner, J.: Methane emission from rice cultivation: Geographic and seasonal distribution of cultivated areas and emissions. *Global Biogeochemical Cycles* **5**, 3–24, 1991.
- McNorton, J., Bousserez, N., Agustí-Panareda, A., Balsamo, G., Engelen, R., Huijnen, V., Inness, A., Kipling, Z., Parrington, M., and Ribas, R.: Quantification of methane emissions from hotspots and during COVID-19 using a global atmospheric inversion, *Atmos. Chem. Phys. Discuss.* [preprint], <https://doi.org/10.5194/acp-2021-1056>, in review, 2022.
- Morino, I., Yokozeki, N., Matsuzaki, T., & Horikawa. TCCON data from Rikubetsu (JP), Release GGG2014.R1 (Version GGG2014.R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.RIKUBETSU01.R1/1242265>, 2016.
- 995 Morino, I., Matsuzaki, T. & Horikawa, M. TCCON data from Tsukuba (JP), 125HR, Release GGG2014.R1 (Version GGG2014.R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.TSUKUBA02.R1/1241486>, 2016.
- Morino, I., V. A. Velasco, A. Hori, O. Uchino, D. W. T. Griffith. TCCON data from Burgos, Philippines, Release GGG2014R0. TCCON data archive, hosted by CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.burgos01.R0/1368175>, 2018.
- 1000



- 1005 NOAA Carbon Cycle Group ObsPack Team. (2019). Multi-laboratory compilation of atmospheric methane data for the period 1957-2017; obspack_ch4_1_GLOBALVIEWplus_v1.0_2019_01_08; [Data set]. NOAA Earth System Research Laboratory, Global Monitoring Division. <https://doi.org/10.25925/20190108>
- Notholt, J. et al. TCCON data from Ny Ålesund, Spitsbergen (NO), Release GGG2014.R1(Version R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.NYALESUND01.R1>, 2019.
- 1010 Notholt, J. et al. TCCON data from Bremen (DE), Release GGG2014.R0 (Version GGG2014.R0) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.BREMEN01.R0/1149275>, 2014.
- Olivier, J. and Janssens-Maenhout G.: CO₂ Emissions from Fuel Combustion – 2012 Edition, IEA CO₂ report 2012, Part III, Greenhouse-Gas Emissions, OECD Publishing, Paris, ISBN 978-92-64-17475-7, 2012.
- 1015 Ortega, I., J.W. Hannigan, R.R. Buchholz, G. Pfister. Long-term variability and source signature of gases emitted from oil & natural gas and cattle feedlot operations in the Colorado front range. *Atmospheric Environment*, Volume 263, 118663, ISSN 1352-2310, <https://doi.org/10.1016/j.atmosenv.2021.118663>, 2021.
- 1020 Peterson, J.T., W.D. Komhyr, L.S. Waterman, R.H. Gammon, K.W. Thoning, and T.J. Conway, Atmospheric CO₂ variations at Barrow, Alaska, 1973–1982, *J. Atmos. Chem.*, 4, 491–510, 1986.
- Patra, P. K., et al.: TransCom model simulations of hourly atmospheric CO₂: Analysis of synoptic-scale variations for the period 2002–2003, *Global Biogeochem. Cycles*, 22, GB4013, doi:10.1029/2007GB003081, 2008
- 1025 Patra, P. K., Houweling, S., Krol, M., Bousquet, P., Belikov, D., Bergmann, D., Bian, H., Cameron-Smith, P., Chipperfield, M. P., Corbin, K., Fortems-Cheiney, A., Fraser, A., Gloor, E., Hess, P., Ito, A., Kawa, S. R., Law, R. M., Loh, Z., Maksyutov, S., Meng, L., Palmer, P. I., Prinn, R. G., Rigby, M., Saito, R., and Wilson, C.: TransCom model simulations of CH₄ and related species: linking transport, surface flux and chemical loss with CH₄ variability in the troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 11, 12813–12837, <https://doi.org/10.5194/acp-11-12813-2011>, 2011.
- 1030 Petri, C., C. Rousogonous, T. Warneke, M. Vrekoussis, S. Sciare, J. Notholt. TCCON data from Nicosia, Cyprus, Release GGG2014R0. TCCON data archive, hosted by CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A. <http://doi.org/10.14291/tcon.ggg2014.nicosia01.R0>, 2020.
- 1035 Pollard, D. F., Sherlock, V., Robinson, J., Deutscher, N. M., Connor, B. and Shiona, H.: The Total Carbon Column Observing Network site description for Lauder, New Zealand, *Earth Syst. Sci. Data*, 9(2), 977–992, doi:10.5194/essd-9-977-2017, 2017.



- 1040 Ramonet, M. and P. Monfray. CO₂ baseline concept in 3-D atmospheric transport models. *Tellus B* 48(4): 502-520, 1996.
- Ramonet, M., P. Ciais, T. Aalto, C. Aulagnier, F. Chevallier, D. Cipriano, T.J. Conway, L. Haszpra, V. Kazan, F. Meinhardt, J.-D. Paris, M. Schmidt, P. Simmonds, I. Xueref-Remy, and J. Necki, A recent build-up of atmospheric CO₂ over Europe. Part 1: observed signals and possible explanations, *Tellus B*, DOI: 10.1111/j.1600-0889.2009.00442, 2010.
- 1045 Ramonet, M., Langerock, B., Warneke, T., Eskes, H.J.: Validation report of the CAMS greenhouse gas global re-analysis, years 2003-2020, Copernicus Atmosphere Monitoring Service (CAMS) report CAMS84_2018SC3_D5.1.2-2020_v0.1.pdf, April 2021, doi: 10.24380/438c-4597, 2021.
- Reuter, M., Bovensmann, H., Buchwitz, M., Burrows, J. P., Connor, B. J., Deutscher, N. M., Griffith, D. W. T., Heymann, J., 1050 Keppel-Aleks, G., Messerschmidt, J., Notholt, J., Petri, C., Robinson, J., Schneising, O., Sherlock, V., Velasco, V., Warneke, W., Wennberg, P. O., and Wunch, D.: Retrieval of atmospheric CO₂ with enhanced accuracy and precision from SCIAMACHY: Validation with FTS measurements and comparison with model results, *J. Geophys. Res.*, 116, D04301, <https://doi.org/10.1029/2010JD015047>, 2011.
- 1055 Ridgwell, A. J., Marshall, S. J. & Gregson, K.: Consumption of atmospheric methane by soils: A process-based model. *Global Biogeochemical Cycles* **13**, 59–70, 1999.
- Rodgers, C. D.: *Inverse Methods for Atmospheric Sounding: Theory and Practice*. World Scientific, 2000.
- 1060 Sanderson, M. G.: Biomass of termites and their emissions of methane and carbon dioxide: A global database. *Global Biogeochemical Cycles* **10**, 543–557, 1996.
- Sarra, A. G. D., Karion, A., Arlyn Andrews, Colomb, A., Scheeren, B., Viner, B., Myhre, C. L., Miller, C. E., Plass-Duelmer, C., Plass-Duelmer, C., Sloop, C. D., Sweeney, C., Kubistin, D., Jaffe, D., Dlugokencky, E., Vitkova, G., Manca, G., Huilin 1065 Chen, Lehner, I., ... Kazan, V. (2021). Multi-laboratory compilation of atmospheric carbon dioxide data for the years 2020-2021; obspack_co2_1_NRT_v6.1_2021-02-02 [Data set]. NOAA Earth System Research Laboratory, Global Monitoring Laboratory. <https://doi.org/10.25925/20210108>, 2021.
- Sandu, I., Beljaars, A., Bechtold, P., Mauritsen, T. & Balsamo, G.: Why is it so difficult to represent stably stratified conditions 1070 in numerical weather prediction (NWP) models? *Journal of Advances in Modeling Earth Systems* **5**, 117–133, 2013.



- Schepers, D. *et al.*: Methane retrievals from Greenhouse Gases Observing Satellite (GOSAT) shortwave infrared measurements: Performance comparison of proxy and physics retrieval algorithms. *Journal of Geophysical Research: Atmospheres* **117**, 2012.
- 1075 Schmidt, M., Graul, R., Sartorius, H., and Levin, L.: The Schauinsland CO₂ record: 30 years of continental observations and their implications for the variability of the European CO₂ budget, *J. Geophys. Res.-Atmos.*, 108, 4619-4626, 2003.
- Schuldt, K. N., Mund, J., Lujikx, I. T., Jacobson, A. R., Cox, A., Vermeulen, A., Manning, A., Beyersdorf, A., Manning, A., 1080 Karion, A., Hensen, A., Arlyn Andrews, Frumau, A., Colomb, A., Scheeren, B., Law, B., Baier, B., Munger, B., Paplawsky, B., ... Loh, Z. Multi-laboratory compilation of atmospheric carbon dioxide data for the period 1957-2019; obspack_co2_1_GLOBALVIEWplus_v6.0_2020-09-11 [Data set]. NOAA Earth System Research Laboratory, Global Monitoring Division. <https://doi.org/10.25925/20200903>, 2020.
- 1085 Segers, A., Tokaya, J. & Houweling, S. Description of the CH₄ Inversion Production Chain. https://atmosphere.copernicus.eu/sites/default/files/2021-01/CAMS73_2018SC3_D73.5.2.2-2020_202012_production_chain_Ver1.pdf, 2020a.
- Segers, A., Tokaya, J., Houweling, S.: Validation of the CH₄ surface flux inversion - reanalysis 1990-2019, Copernicus 1090 Atmosphere Monitoring Service EQC report, Dec 2020, https://atmosphere.copernicus.eu/sites/default/files/2021-02/CAMS73_2018SC2_D73.2.4.1-2020_202012_validation_CH4_1990-2019_v2.pdf, 2020b.
- Stevenson, D., Derwent, R., Wild, O., and Collins, W.: COVID-19 lockdown NO_x emission reductions can explain most of the coincident increase in global atmospheric methane, *Atmos. Chem. Phys. Discuss.* [preprint], [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-2021-604) 1095 2021-604, in review, 2021.
- Sha, M. K., Langerock, B., Blavier, J.-F. L., Blumenstock, T., Borsdorff, T., Buschmann, M., Dehn, A., De Mazière, M., Deutscher, N. M., Feist, D. G., García, O. E., Griffith, D. W. T., Grutter, M., Hannigan, J. W., Hase, F., Heikkinen, P., Hermans, C., Iraci, L. T., Jeseck, P., Jones, N., Kivi, R., Kumps, N., Landgraf, J., Lorente, A., Mahieu, E., Makarova, M. V., Mellqvist, 1100 J., Metzger, J.-M., Morino, I., Nagahama, T., Notholt, J., Ohyama, H., Ortega, I., Palm, M., Petri, C., Pollard, D. F., Rettinger, M., Robinson, J., Roche, S., Roehl, C. M., Röhling, A. N., Rousogonous, C., Schneider, M., Shiomi, K., Smale, D., Stremme, W., Strong, K., Sussmann, R., Té, Y., Uchino, O., Velasco, V. A., Vigouroux, C., Vrekoussis, M., Wang, P., Warneke, T., Wizenberg, T., Wunch, D., Yamanouchi, S., Yang, Y. and Zhou, M.: Validation of methane and carbon monoxide from Sentinel-5 Precursor using TCCON and NDACC-IRWG stations, *Atmos. Meas. Tech.*, 14(9), 6249–6304, doi:10.5194/amt-1105 14-6249-2021, 2021.



- Sherlock, V. et al. TCCON data from Lauder (NZ), 125HR, Release GGG2014.R0 (Version GGG2014.R0) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.LAUDER02.R0/11492981>, 2014a.
- 1110 Sherlock, V., Connor, B., Robinson, J., Shiona, H., Smale, D., and Pollard, D.: TCCON data from Lauder, New Zealand, 120HR, Release GGG2014R0. CaltechDATA, 2014b.
- Pollard, D., Robinson, J., and Shiona, H.: TCCON data from Lauder, New Zealand, 125HR, Release GGG2014R0. CaltechDATA, 2019.
- 1115 Spahni, R. *et al.* : Constraining global methane emissions and uptake by ecosystems. *Biogeosciences* **8**, 1643–1665, 2011.
- Stephens, B. B., G. W. Brailsford, A. J. Gomez, K. Riedel, S. E. Mikaloff Fletcher, S. Nichol, and M. Manning. Analysis of a 39-year continuous atmospheric CO₂ record from Baring Head, New Zealand. *Biogeosciences* 10, 4: 2683–2697, doi:10.5194/bg-10-2683-2013, 2013.
- 1120 Strong, K., Roche, S., Franklin, J. E., Mendonca, J., Lutsch, E., Weaver, D., Fogal, P. F., Drummond, J. R., Batchelor, R., & Lindenmaier, R. (2019). TCCON data from Eureka (CA), Release GGG2014.R3 (Version R3) [Data set]. CaltechDATA. <https://doi.org/10.14291/TCCON.GGG2014.EUREKA01.R3>, 2019.
- 1125 Sussmann, R., Forster, F., Rettinger, M., and Bousquet, P.: Renewed methane increase for five years (2007–2011) observed by solar FTIR spectrometry, *Atmos. Chem. Phys.*, 12, 4885–4891, doi:10.5194/acp-12-4885-2012, 2012.
- Sussmann, R., Ostler, A., Forster, F., Rettinger, M., Deutscher, N. M., Griffith, D. W. T., Hannigan, J. W., Jones, N., and Patra, P. K.: First intercalibration of column-averaged methane from the Total Carbon Column Observing Network and the Network for the Detection of Atmospheric Composition Change, *Atmos. Meas. Tech.*, 6, 397–418, doi:10.5194/amt-6-397-2013, 2013.
- 1130 Sussmann, R. & Rettinger, M. TCCON data from Garmisch (DE), Release GGG2014.R2 (Version R2) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.GARMISCH01.R2>, 2018a.
- 1135 Sussmann, R., M. Rettinger. TCCON data from Zugspitze, Germany, Release GGG2014R1. TCCON data archive, hosted by CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A. <http://dx.doi.org/10.14291/tcon.ggg2014.zugspitze01.R1>, 2018b.



- 1140 Takahashi, T. *et al.* : Climatological mean and decadal change in surface ocean pCO₂, and net sea–air CO₂ flux over the global oceans. *Deep Sea Research Part II: Topical Studies in Oceanography* **56**, 554–577, 2009.
- Te, Y., Jeseck, P. & Janssen, C. TCCON data from Paris (FR), Release GGG2014.R0 (Version GGG2014.R0) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.PARIS01.R0/1149279>, 2014.
- 1145 Temperton, C., Hortal, M. & Simmons, A. A.: two-time-level semi-Lagrangian global spectral model. *Quarterly Journal of the Royal Meteorological Society* **127**, 111–127, 2001.
- Tsutsumi, Y., H. Matsueda, S. Nishioka, Consistency of the CO₂ primary standards in JMA, 12th WMO/IAEA meeting of experts on carbon dioxide concentration and related tracers measurement techniques (Toronto, Canada, 15-18 September 2003), 1150 Global Atmosphere Watch Report No. 161 (WMO/TD-No.1275), 23-31, 2005.
- The Global Observing System For Climate: Implementation Needs, report number 200, World Meteorological Organization, 2016.
- 1155 Verma,S., Marshall,J., Parrington,M., Agustí-Panareda,A., Massart, S., Chipperfield, M. P. , Wilson, C., Gerbig, C.: Extending methane profiles from aircraft into the stratosphere for satellite total column validation using the ECMWF C-IFS and TOMCAT/SLIMCAT 3-D model, *Atmos. Chem. Phys.*, 17, 6663—6678, 2017.
- Warneke, T. et al. TCCON data from Orléans (FR), Release GGG2014.R0 (Version GGG2014.R0) [Data set]. 1160 <https://doi.org/10.14291/TCCON.GGG2014.ORLEANS01.R0/1149276>, 2014.
- Waterman, L.S., D. W. Nelson, W.D. Komhyr, T.B. Harris, and K.W. Thoning, Atmospheric carbon dioxide measurements at Cape Matatula, American Samoa, 1976 1984., *J. Geophys. Res.*, 94, 14817 14829, 1989.
- 1165 Wennberg, P. O. et al. TCCON data from Park Falls (US), Release GGG2014.R1 (Version GGG2014.R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.PARKFALLS01.R1>, 2017.
- Wennberg, P. O. et al. TCCON data from Lamont (US), Release GGG2014.R1 (Version GGG2014.R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.LAMONT01.R1/1255070>, 2016. 1170
- Wennberg, P. O. et al. TCCON data from Caltech (US), Release GGG2014.R1 (Version GGG2014.R1) [Data set]. <https://doi.org/10.14291/TCCON.GGG2014.PASADENA01.R1/1182415>, 2015.



Wennberg, P. O., D. Wunch, Y. Yavin, G. C. Toon, J.-F. Blavier, N. Allen, G. Keppel-Aleks. TCCON data from Jet Propulsion
1175 Laboratory, Pasadena, California, USA, Release GGG2014R0. TCCON data archive, hosted by CaltechDATA, California
Institute of Technology, Pasadena, CA, U.S.A. <https://doi.org/10.14291/tcon.ggg2014.jpl01.R0/1149163>, 2016.

Williams, J. E., Huijnen, V., Bouarar, I., Meziane, M., Schreurs, T., Pelletier, S., Marécal, V., Josse, B., and Flemming, J.:
Regional evaluation of the performance of the global CAMS chemical modeling system over the United States (IFS cycle
1180 47r1), *Geosci. Model Dev. Discuss.* [preprint], <https://doi.org/10.5194/gmd-2021-318>, in review, 2021.

Worthy, D.E., K.Higuchi, and D. Chan. North American influence on atmospheric carbon dioxide data collected at
Sable Island, Canada. *Tellus* 55B, 105-114, 2003.

1185 Wunch, D., G. C. Toon, J.-F. L. Blavier, R. A. Washenfelder, J. Notholt, B. J. Connor, D. W. T. Griffith, V. Sherlock, and P.
O. Wennberg. The Total Carbon Column Observing Network, *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 369(1943),
2087–2112, doi:10.1098/rsta.2010.0240, 2011.

Wunch, D., G. C. Toon, V. Sherlock, N. M. Deutscher, C. Liu, D. G. Feist, and P. O. Wennberg. The Total Carbon Column
1190 Observing Network's GGG2014 Data Version, Pasadena, California, 2015.

Wunch, D., J. Mendonca, O. Colebatch, N. Allen, J.-F. L. Blavier, S. Roche, J. Hedelius, G. Neufeld, S. Springett, D. Worthy,
R. Kessler, K. Strong. TCCON data from East Trout Lake, Canada, Release GGG2014R1. TCCON data archive, hosted by
CaltechDATA, California Institute of Technology, Pasadena, CA, U.S.A.
1195 <https://doi.org/10.14291/tcon.ggg2014.easttroutlake01.R1>, 2018.

Zander, R., Mahieu, E., Demoulin, P., Duchatelet, P., Roland, G., Servais, C., Mazière, M. De, Reimann, S. and Rinsland, C.
P.: Our changing atmosphere: Evidence based on long-term infrared solar observations at the Jungfraujoch since 1950, *Sci.*
Total Environ., 391(2–3), 184–195, doi:10.1016/j.scitotenv.2007.10.018, 2008.

1200

Zhou, M., Langerock, B., Vigouroux, C., Sha, M. K., Ramonet, M., Delmotte, M., Mahieu, E., Bader, W., Hermans, C., Kumps,
N., Metzger, J.-M., Dufлот, V., Wang, Z., Palm, M. and De Mazière, M.: Atmospheric CO and CH₄ time series and seasonal
variations on Reunion Island from ground-based in situ and FTIR (NDACC and TCCON) measurements, *Atmos. Chem. Phys.*,
18(19), 13881–13901, doi:10.5194/acp-18-13881-2018, 2018.