

Assessing stakeholder climate data needs for farm-level decision-making in the U.S. Corn Belt

Suzanna Clark¹, J. Felix Wolfinger², Melissa A. Kenney², Michael D. Gerst³, Heidi A. Roop¹

¹Department of Soil, Water, & Climate, University of Minnesota, St. Paul, MN, 55108, U.S.A.

²Institute on the Environment, University of Minnesota, St. Paul, MN, 55108, U.S.A.

³Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD, 20742, U.S.A.

Correspondence to: Suzanna Clark (suzclark@umn.edu)

Abstract. Across the Midwest region of the United States, agriculturalists make decisions on a variety of time scales, ranging from daily to weekly, monthly, and seasonally. Ever-improving forecasts and decision support tools could assist the decision-making process, particularly in the context of a changing and increasingly variable climate. To be usable, however, the information produced by these forecasts and tools should be salient, credible, legitimate, and iterative, qualities which are achieved through deliberate co-production with stakeholders. This study uses a document analysis approach to explore the climate information needs and priorities of stakeholders in the U.S. Corn Belt. Through the analysis of 50 documents, we find that stakeholders are primarily concerned with practical and tactical decision making, including from whom they get their information, the application of information to agricultural, water, and risk management, and desired economic outcomes. The information that stakeholders desire is less focused on social issues, environmental issues, or long-term climate resilience. These results can inform the development of future decision support tools, identify known gaps in climate information services to reduce stakeholder fatigue, and serve as an example to scientists trying to understand stakeholder needs in other regions and specialties.

Deleted:

Deleted: ¶

Deleted: ¶

Deleted: ¶

³Earth

Deleted: 5

Deleted: (suzclark@umn.edu)

Deleted:

Deleted: ¶
10

Deleted: stakeholder

Deleted: ¶

Deleted: ¶
15

Deleted: This study

1 Introduction

Across the Midwest region of the United States, agriculturalists make decisions on a variety of time scales, from weekly to seasonal, interannual, and decadal (Haigh et al., 2015b). These decisions can be classified as either operational or strategic (Haigh et al., 2015b; Prokopy et al., 2013), and farmers have been found to rely on either proprietary information obtained through a subscription service or free information from a company or university to inform their management decisions (Haigh et al., 2018). However, climate change across the Midwest is projected to lead to higher temperatures, a longer frost-free season, increased springtime rainfall, higher humidity, and an increased risk of flooding, and to alter the timing and variability of seasons (Angel et al., 2018, and references therein). Combined, these changes may make the information upon which agriculturalists rely obsolete, necessitating a new process for decision making.

In the context of a changing climate, it has been proposed that farmers could benefit from continuously improving climate models and decision support tools that incorporate environmental and climate information and forecasts (Klemm and McPherson, 2017). Forecasts can indicate the likelihood of an El Niño year (Ghil and Jiang, 1998; Jones et al., 2006), project temperature and precipitation extremes for the season (Andrys et al., 2015) or forecast impending events such as extreme storms (Chawla et al., 2018; Moya-álvarez et al., 2018). Many decision support tools developed by public and private for-profit entities already exist that assist agriculturalists in deciding, for example, whether or not to use cover crops or till the fields, when, and how much nutrient to apply, and whether to purchase crop insurance (Palutikof et al., 2019 and references therein; Haigh et al., 2018). The structure of these tools varies, with some guiding users step-by-step through necessary decision processes and trade-off choices, and others providing information or indicators that are relevant to a range of decisions but not customized to a single decision context (Kenney et al., 2016; Rose, 2015; Wiggins et al., 2018). To ensure that decision support tools and products are usable, their information needs to be shared at a time that is relevant to farmers' decision-making processes, and it should be informed by existing stakeholder needs and engagement with agriculturalists and agricultural advisors (Haigh et al., 2015b).

The gap between the information that scientists produce and the information that end users find usable is well documented (Dewulf et al., 2020; Kirchhoff et al., 2013; Lemos et al., 2012). To be useful and usable, science should be salient, credible, legitimate (Cash et al., 2003), and iterative (Dilling and Lemos, 2011; Sarkki et al., 2015), and scientists should consider both the information's potential use and the process by which it was created (Dilling and Lemos, 2011). Many researchers increasingly turn to stakeholder engagement and knowledge co-production (Stumpf et al., 2016) to achieve these goals. One example of this effort is the Useful 2 Usable project, a multi-institutional effort to transform existing climate data into

Deleted: 20 stakeholder needs in other regions and specialties.¶
1→

Deleted: ¶

Deleted: to

Deleted: ¶
25

Deleted: both

Deleted: ¶
30

Deleted: ¶
35

Deleted: for agriculture

Deleted: ¶
40

Deleted: nutrients

Deleted: ¶
45

Deleted: ¶
50→

Deleted: ¶

... 11

Deleted: ¶
55

55 usable agricultural products that incorporated stakeholder feedback through user surveys and data use statistics (Angel et al., 2017). Other stakeholder-led projects have led to usable science in regions as far reaching as California (Baker et al., 2020), Argentina (Podestá et al., 2013), Zambia (Arslan et al., 2015), the UK (Rose, 2015), and Australia (Hochman and Carberry, 2011).

Deleted: ¶
60

60 To explore stakeholder climate and environmental information needs and priorities in the U.S. Corn Belt, we used a document analysis approach (Bowen, 2009) modeled after the methods in Dilling and Berggren (2015) and Molino et al. (2020). We categorized and coded existing documents from a predetermined coding schema to allow for an easy inter-documental comparison of stakeholder needs. Through this approach we can recognize both data needs that are commonly expressed and expected data needs that have not been prioritized. Because much has already been published on the information needs of agriculturalists in the region, we use this document analysis approach to understand the existing stakeholder needs landscape. This reduces stakeholder fatigue and focuses future engagements on advancing the understanding of information translation.

Deleted: this study uses

Deleted: Existing documents are

Deleted: ¶
65

Deleted: need

70 Information collected from this study will be used to develop the Dashboard for Agricultural Water use and Nutrient management (DAWN). The DAWN project is co-creating sub-seasonal to seasonal forecasts that will be organized as decision-task-focused indicators and a decision support tool dashboard to support water and nutrient management decisions for food and energy crop production in the U.S. Midwest Corn Belt region. In addition, operationalized, predictive and downscaled seasonal climate outlooks present an opportunity to build open-access decision support systems that allow for more equitable access to relevant information.

Deleted: 70

Deleted: ¶
75

75 The methods section of this document outlines the study's design, including the methods and criteria for document retrieval and selection, the creation of a coding schema for the U.S. Corn Belt, and subsequent analysis of coded documents. The results section focuses on several main themes that were identified throughout the document coding, which relate to the climate and environmental information practitioners need, where they get their information, the decisions they focus on, and their desired outcomes. The discussion section interprets the results in the context of risk management, hypothesizes explanations for why some codes appeared more than others, and outlines the implications of this work for the DAWN project and scientists in other regions or sectors who are planning to conduct similar user-driven research and decision-support tool development.

Deleted: paper

Deleted: ¶
80

Deleted: ¶

... [2]

Deleted: focuses on

Deleted:

85 Deleted: 85 → 2 →

2 Methods

We used document analysis to assess stakeholders' perspectives without directly engaging with them (Saldana, 2013; Bowen, 2009). While this method is limited in its scope and relies on information and ideas that have already been shared, it benefits from being "stable, unobtrusive, exact, and available over a long span of time" (Yin, 2009; Dilling and Berggren, 2015). This method has also been used to identify stakeholder needs in other regions and with a variety of foci, such as to explore stakeholder needs with respect to climate change in the Mountain West (Dilling and Berggren, 2015) and the Northeastern United States (Molino et al., 2020). The following section outlines how documents were chosen, coded, and analyzed. The purpose of this study was to apply an existing schema to a new study period of interest, rather than to develop new methods.

2.1 Document selection

We found documents via a web search and "snowball sampling" (Goodman, 1961), beginning with recent research on Decision Calendars in the Midwest (e.g., Haigh et al., 2015). We defined a "document" as original, peer-reviewed, published research. Document retrieval focused on peer-reviewed literature, despite an abundant body of Extension literature, because Extension documents primarily focused on lending advice to agricultural practitioners, rather than surveying their needs. In addition, identified state and federal reports primarily summarized research that had been conducted elsewhere, so these documents were eliminated to reduce redundancy. The search was considered complete when no new documents were found. We did not include review documents in the coding, because they do not include original information, but we used them to identify other studies.

Documents were included if they met four criteria: (1) geographic scope, (2) date of publication, (3) input from stakeholders, and (4) focus on agricultural and natural resource management.

The first criterion for inclusion was geographic scope, which was motivated by the scope of the DAWN project (Figure 1). To be included, documents needed to focus on part or all of the following "Corn Belt" states: Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin (Hunt et al., 2020). Documents that stated a "Corn Belt" focus without specifying the state were also included in the initial identification and coded for specific states later (see Sect. 2.3). We did not include documents for which approximately 10% of the study area or less was made up by a Corn Belt state because their primary focus was outside the region of interest. Although agriculturalists in other regions beyond the Corn Belt also face challenges and have data needs in the face

Deleted: Document

Deleted: was used

Deleted: 90

Deleted: 95 → 2.1 →

Deleted: Selection

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Column Break

... [3]

Deleted: The search was considered complete when no new documents were found. Papers

Deleted: Review papers were not included in document coding, because they do not include original information, but they were used to identify other studies.

Deleted: Papers for which

of a changing climate, we excluded these documents both because of the scope of the DAWN project and because climatic processes and changes are region-specific.




Figure 1: The geographic scope of the DAWN project, showing the 11 states chosen for the study focus (dawn.umd.edu). CSU = Colorado State University; UNL = University of Nebraska, Lincoln; UMN = University of Minnesota; FFG = Family Farm Group; UIUC = University of Illinois - Urbana Champaign; UMD = University of Maryland; USDA ARS = U.S. Department of Agriculture - Agricultural Research Service; NCEI = National Centers for Environmental Information; HU = Howard University; LTAR = Long-Term Agroecosystem Research; UVB = Ultraviolet-B; PAR = Photosynthetically active radiation.

We included only documents that have been published since 2010, to focus the analysis on the most recent data possible. A ten-year time span ensured that enough documents were available to choose from and analyze, but also that the stated stakeholder perspectives were recent and reflective of the current social, political, and meteorological contexts in which decisions are being made. Note that, while the documents were constrained with respect to geography and publication year, documents were not constrained with regards to the kind of agriculture or the information time scale that they discussed.

The third and fourth criteria for inclusion were input from stakeholders and a focus on agricultural and natural resource management. "Input" could refer to quotes, survey results, product feedback, or the authors' interpretation of stakeholder needs. We chose the topics of interest inductively after a precursory literature search, resulting in crops and livestock as the two primary topical foci.

Deleted: Section Break (Next Page)

110

115

120

125

Column Break

... [4]

Deleted:

Deleted: Only papers

Deleted: were included

Deleted: papers

Deleted: The

Deleted: were chosen

135 We also identified some documents that discussed water resource management, which we added to the document set to increase the scope of potential applications of this study. They comprised about 10% of the final document set.

2.2 Coding

140 We conducted document analysis (Saldana, 2013) in MAXQDA, a software for qualitative data analysis. We analyzed and coded documents deductively to the extent possible, meaning that the coding schema was pre-determined based on a prior understanding of stakeholder needs, rather than inductively, in which codes are created in response to the specific document set. We chose the deductive approach both to create a coding schema that could be applied to other studies and to allow for analysis of expected needs that are not stated in addition to those that are. We added some codes inductively during the coding process if they showed up repeatedly and had not been included in the original coding schema. This is a common approach in qualitative analysis, whereby study design is modified inductively until most of the research is complete (Bickman and Rog, 2009).

150 In accordance with the study's goal of applying an existing coding schema to a new region, we adapted the coding schema from Molino et al. (2020), Dilling and Berggren (2015) and Dilling and Lemos (2011), with changes made to account for different geographic regions and stakeholders and to make the schema more intuitive. In the coding schema, a "code" was the most specific descriptor possible, and similar codes were grouped under "nodes," according to the method outlined in Molino et al. (2020). The final schema consisted of 6 nodes, each with 4-6 sub-nodes: (1) Coordination, Collaboration, and Communication; (2) Monitoring and Data Collection; (3) Policies, Programs, and Law; (4) Research Topics; (5) Social Issues; and (6) Case attributes (e.g., year published, author affiliation, funding source, etc.). Most sub-nodes included their own sub-sub-nodes, with an additional fourth layer where specificity was necessary. An overview of the final coding schema is illustrated in Figure 2, and the reader is referred to the Supplementary Material for the complete coding schema.

Deleted: Some papers were

Deleted: found

Deleted: were

Deleted: ¶

130 ¶

¶

¶

¶

¶

135 ¶

¶

¶

¶

¶

140 ¶

¶

¶

¶

145 ¶

¶

... [5]

Deleted: ¶

... [6]

Deleted: ¶

... [7]

Deleted: was conducted

Deleted: Documents were

Deleted: The

Deleted: was chosen

Deleted: Some

Deleted: were added

Deleted: was adapted

Deleted: (

Deleted: ,

Deleted:) and (

Deleted: ,

Deleted: ;

Deleted: ,

Deleted: -

Deleted:

Deleted: ¶

Deleted: a

Deleted: of codes including in instances

Deleted: code





Figure 2: The main nodes and associated sub-nodes of the coding schema. The order of nodes/sub-nodes does not suggest ranking.

We followed several principles during coding to ensure consistency across all coders and documents. First, we applied a code wherever it appeared, regardless of how frequently (or infrequently). Second, we only coded stakeholder input, whether it appeared as (1) a direct quote, (2) in the analysis from the document authors, or (3) as a quotation from another document when used as context for the study's analysis. Therefore, wherever "stakeholder" or "practitioner" input is mentioned throughout this document, it is taken from one of these three contexts. "Stakeholder" and "practitioner" can refer to agriculturalists, water managers, and rangeland managers. We did not code information from other documents if the geographic scope of the cited study did not include any locations within the Corn Belt. Finally, we weighted all codes evenly, with no added judgment about the code's perceived importance by the stakeholder.

Two team members initially coded several documents simultaneously and discussed their results until a consensus about coding was reached. Thereafter, for efficiency, documents were split between the two coders without double-coding. Document codes were inductively adjusted as necessary to ensure intercoder consistency. In most instances, codes appeared explicitly, such as with "precipitation", "forecast", and "data source." In some instances, however, such as with communication channels, coders applied codes based on interpretation.



- Deleted: Coding was guided by
- Deleted: a code was
- Deleted: was coded
- Deleted: paper
- Deleted: paper
- Deleted: Information
- Deleted: papers was not coded
- Deleted: was beyond
- Deleted: (Sect. 2.2).
- Deleted: were weighted

2.3 Analysis

We analyzed codes for code occurrence and code co-occurrence. Code occurrence was counted by the number of documents in which a code appeared, rather than the number of times a code itself was used, to avoid biasing toward longer documents. To establish themes across the document set, we analyzed the frequency of sub-nodes first, followed by sub-sub-nodes and specific codes, as needed, to analyze details. We analyzed the document set for basic statistics, including the number of documents published in each year, the number of documents published in each state, and the distribution of funding from grants, universities, private donors, or government budget.

To analyze code occurrence in such a large coding schema, we analyzed the document set node by node, rather than for all codes simultaneously. This method was appropriate because three nodes - Monitoring and Data Collection; Coordination, Collaboration, and Communication; and Policies, Programs, & Law - were used across the same total number of documents (45), while the node Research appeared in two thirds of the total documents (34 documents), so sub-nodes within these four nodes were analyzed with equal weight. We analyzed codes under the same node for co-occurrence, defined as occurring together within the same document. The node Social Issues was not used in any documents and was therefore not included in this analysis.

3 Results

3.1 Summary statistics

We identified 50 documents in total for analysis that met the criteria defined in Sect. 2.2. All but two of the documents had two or more authors, and most documents had at least one author from an academic institution (Figure 3). Among those academic institutions, the most frequently coded were the University of Nebraska - Lincoln (18 documents), Purdue University (17 documents), University of Wisconsin - Madison (12 documents), and Iowa State University (12 documents). Forty-five (45) out of the 50 documents were co-authored by an author from an institution (academic or otherwise) within the U.S. Corn Belt, and 25 documents were co-authored by an author from an institution outside of the geographic scope. Grants were the most dominant funding source (30 documents), followed by university programs (13 documents), government budgets (5 documents), and unlisted sources (2 documents). Eight documents described work that was funded as part of the Useful to Usable project. Some of the papers included survey results and others were the result of stakeholder engagements.

Deleted: 2.3

Deleted: Codes were

Deleted:

Deleted:

175

180

185

Column Break

Deleted: was analyzed

Deleted: The

Deleted: was analyzed

Deleted: was analyzed

Deleted: Codes

Deleted: were analyzed

Deleted:

Deleted: Statistics

Deleted: Section Break (Continuous)

190

195

Column Break

Deleted:)

Deleted: were found

Deleted: (8)

Deleted: were




Figure 3: Number of documents with at least one author affiliated with an organization or institution from an academic institution, the government, an NGO (non-governmental organization), or the private sector. Note that the sum of the documents is greater than 50, because most documents had multiple authors, who may be affiliated with different institutions.

All of the states in the Corn Belt were mentioned in at least one document, but their frequency varied widely. Kentucky was mentioned the least, in only 2 documents, while Iowa was mentioned in 28 documents (Figure 4). Most studies spanned state lines, with their geographic scope determined by natural features such as watersheds or anthropogenic boundaries such as individual farms and government jurisdictions. Some documents also included regions beyond the study's geographic scope, in other states and provinces, but those states and provinces are not discussed here.

More documents were published in the second half of the time period, from 2016 to 2021, than in the first half, from 2010 to 2016 (Figure 4). There was a peak in publications in 2017, but no apparent long-term trend.

Deleted:Section Break (Next Page).....

Author Affiliation	Number of Papers
Academic Institutions	49
Government	22
NGO	6
Private Sector	2

Deleted: ,

Deleted: , the government, or an academic institution

Deleted:

Deleted: paper

Deleted: -

Deleted:Section Break (Next Page)..... [10]





Figure 4: (left) States in the Corn Belt color-coded by the number of documents that included that state in their study area. (right) Number of publications per year.

3.2 Common themes

The most common codes fell under five overarching themes: where practitioners get their climate and environmental information; what information practitioners need; capacity and barriers that affect decision-making; which decisions practitioners can control; and the desired outcomes of information gathering and decision-making (Figure 5). Themes such as social issues, research standards, and collaboration standards were mentioned less frequently, or not at all.

Deleted: ¶
215 → 3.2 →

Deleted: Themes

Deleted: ¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

Column Break

Deleted: Together, these themes indicate that the most frequently mentioned concerns were those related to practical decision-making.




Figure 5.- The relative frequency (percent of documents) with which each major theme occurs throughout the document set.

3.2.1 Sources of information

The most frequently coded sub-nodes under "Coordination, Collaboration, and Communication" (Figure 2) were "Communication Channels" and "Training and Data Use". Codes related to communication/collaboration goals, proposed collaborations, and communication standards were used in fewer than five documents, if at all. Rather, when communication was mentioned, most documents discussed where practitioners get their climate and environmental information: from research agencies, other communities, private companies (e.g., seed and fertilizer suppliers), consultants, mass media, or Extension and boundary organizations. Most of the information sources mentioned were existing contacts. Communication channels between the government and private industry were mentioned about half as frequently as those between private companies and practitioners.

Deleted: <object>

Deleted: :

Deleted: papers

Deleted:Section Break (Next Page).....
Where practitioners get their

Deleted:

230

235

240

245

250

Column Break

Deleted: papers

Deleted: papers

Deleted: and

Deleted: /

There was little discussion of lines of communication between different organizations, companies, or agencies. In addition to a focus on who shares data with practitioners, there was frequent discussion of how the data are shared, i.e., whether they are usable. There was overall agreement that climate and environmental information should be readily available and easy to download (accessible), and that stakeholders will not use information that they are not aware of (available). There was little discussion, however, of whether information should be proprietary/paid or open source/free.

Deleted: , because the most common focus was the relationship between those who supply the information and those who use it.

Few documents mentioned developing an online "clearinghouse" of information. There were, however, several documents that mentioned evaluating current decision aids. This topic was often mentioned in the context of drought monitors such as the U.S. Drought Monitor (Dermer and Augustine, 2016; Haigh et al., 2021, 2019), the Natural Resources Conservation Service drought tool (Beeton and McNeeley, 2020), the South Dakota Drought Tool (Knutson and Haigh, 2013), and the National Integrated Drought Information System (Otkin et al., 2015), which were particularly prevalent in documents set in the western half of the Corn Belt, where drought is more common. There were also several agricultural tools that were mentioned, such as the Useful 2 Usable Growing Degree Day tool (Haigh et al., 2015b, a) and the USDA GrassCast Tool (Haigh et al., 2018). However, some documents mentioned that these tools go underutilized, and that their utility cannot replace the experience of a consultant (Ranjan et al., 2019).

Deleted: As most of the focus was on data delivery through personal contacts, there was little discussion of

3.2.2 Data and information types

Documents primarily discussed two data sources: forecast and observations. There was less mention of climatological data, historical data, or remote sensing (i.e., satellite) data. Forecast data were often mentioned on the timescale of days to months, with little mention of longer timescales. Observational data overwhelmingly mentioned collecting and using new field data, rather than gathering or analyzing existing historical data.

The most frequently mentioned data types were all related to water management and resources: precipitation, drought, soil moisture, and soil erosion. "Precipitation" was the most frequently coded data type, and it often co-occurred with "Forecast." Precipitation was often discussed in the context of extreme events, particularly drought and extreme rain events, while "drought" was most often coded in the context of learning about a certain event, having a drought response plan, or in the context of monitoring the event's progression.

Note that, in this coding schema, water resource codes refer to upstream resources, on how practitioners use water. We identified several documents related to water quality, but they were beyond the scope of the project because they focused on downstream effects rather than information use. Precipitation and drought

Deleted: papers

Deleted: papers

Deleted:Section Break (Next Page).....

255¶

255¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

.....Column Break.....

Information practitioners need¶

Deleted: - even if peripherally -

Moved (insertion) [1]

Deleted: were focused

Deleted: Several papers were found

275 data were mentioned most frequently, followed by soil moisture and erosion data. Soil health existed in the coding schema in a variety of forms, including soil temperature and compaction, but only moisture and erosion were coded in more than 10 documents, often in the context of rain-induced changes. In addition, although the coding schema included a variety of codes related to weather and climate, such as air temperature, air humidity, extreme heat, snow cover, hail, and wind, none of these codes were mentioned in more than 10% of documents. Similarly, codes related to pests and diseases were mentioned in fewer than 10% of documents, and codes related to air quality or land use were not used at all.

280 3.2.3 **Decision making**

To keep their businesses running, practitioners - agriculturalists, water managers, and rangeland managers - make countless decisions on a variety of time scales, from daily to weekly, seasonally, and annually (Haigh et al., 2015). Not all factors to decision-making are within practitioners' control (see Sect. 3.2.4), but among those that are, the most frequently mentioned management decisions were in one of three categories: agricultural management, water management, or risk management.

285 Certain agricultural management decisions were mentioned more than others, particularly whether to use cover crops, and whether or not to till the fields. No-till agriculture and cover crops both reduce soil erosion, increase soil biological activity, reduce nutrient leaching, and improve overall soil health (Agriculture, 2022; Creech, 2021), and several studies directly inquired whether stakeholders planned to adopt them. The other commonly mentioned agricultural management decisions were related to nutrient application and efficiency in the timing of nutrient application. The primary focus was on Nitrogen, with little discussion of Phosphorus, Potassium, Sulfur, or other nutrients.

Water management decisions most often related to irrigation, with some additional consideration for runoff and drainage, storage, and drought response timing. Irrigation was most often mentioned as a risk reduction or impact mitigation measure. Many of the agriculture and water management decisions mentioned in the documents overlapped with risk management, which appeared frequently. Practitioners mentioned both reducing the risk of natural events and making management decisions that are the least risky. Church et al. (2018) documented a subtle shift towards greater risk management over time.

300 Stakeholders mentioned multiple factors that affect their decision-making, whether by encouraging or discouraging a particular course of action, such as crop insurance and capacity, i.e., funding and infrastructure. Crop insurance was mentioned frequently as a factor that could discourage adopting new crops or farming strategies. Funding referred both to a practitioner's liquid funds and to the availability of financial assistance from the government and private entities. "Infrastructure" in the document set most

Deleted:Section Break (Continuous).....

Deleted: Making

Deleted: ¶

Deleted: ¶
280

Deleted:Section Break (Next Page).....

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

... [11]

Deleted: Nutrient

Deleted: were also often discussed

Deleted: the context of *when* nutrients should be applied, i.e.,

Deleted: /

Moved down [2]: Thus, only practitioners who thought water supply could become unstable were likely to utilize

Deleted: Nationwide, less than 20% of croplands are irrigated, and in most Corn Belt states, less than 10% of corn is irrigated (Hrozencik and Aillery, 2021). Installing ir... [12]

Deleted:

Deleted: Just because the decisions discussed above are within a practitioner's control does not mean that the decisions are without influence.

Deleted: .

Deleted: ¶

The other factors to decision-making that were mentioned most often were factors related to capacity, i.e., fund... [13]

typically referred to physical infrastructure, such as irrigation or machinery. Several other "Service and Capacity" codes were mentioned, but less often than funding and infrastructure, e.g., the ability to make decisions autonomously and flexibly, and structural barriers such as cover crop seed availability and limited market access (Roesch-Mcnally et al., 2018), which prevent the adoption of sustainable practices. Human capacity, such as staff time, training opportunities, leadership, and familiarity with decision support tools, was mentioned rarely.

3.2.4 Desired outcomes

The primary desired outcomes that were discussed were economic: increased crop yields and available markets to sell products. There was no mention of desired social outcomes, and environmental motivations were typically mentioned as either contrary to economic outcomes or as less important. In some instances, economic factors motivated decisions that could lead to a synergistic environmental benefit, such as with cover crops or contour farming. Several documents emphasized that, to promote particular environmental outcomes, governing bodies would have to provide incentives to offset potential economic losses.

4 Discussion

4.1 The decision-making process

The themes mentioned in Sect. 3.2 guide the decision-making process and provide important context for when practitioners need climate and environmental information, what information they need, why they need it, how they prefer to receive it, and from whom they prefer to receive it. Kuehne et al. (2017) created a model to predict the adoption and diffusion of new agricultural practices and identified a variety of factors that can affect the decision-making process. The main themes of the document set (see Sect. 3.2) are discussed in the context of Kuehne et al.'s model below.

In the context of information sources, the most frequently mentioned codes in the node "Coordination, Collaboration, and Communication" were related to communication channels, indicating that most practitioners are concerned with who delivers their information and through what means. In addition, most practitioners indicated that they get their information from a human source such as a trusted advisor, Extension agent, private company, or consultant, although this is highly variable by farm scale and type, because very large farms might have data scientists on staff (Lotton 2022, personal communication, April 2022). This suggests that the relationship between those who supply the information and those who use it is vital to information adoption. The lack of discussion of communication and collaboration standards or of creating new information sources suggests that practitioners place their trust in their information provider, not the information creator. The lack of desire for a "clearinghouse" of information echoes the emphasis on

Deleted: ¶

315 ¶

320 ¶

325 ¶

330 ¶

335 ¶

... [14]

Deleted: ¶

Deleted: ¶

Deleted: significant incentive

Deleted: ¶

Deleted: Decision-Making Process

Deleted: ¶

Deleted: and

Deleted: information

Deleted: grouped the different factors to adoption into those that affect the time to peak adoption and those that affect the time to initial adoption ... [15]

Deleted: The

Deleted: about

Deleted: . There was little

Deleted: , ¶

... [16]

Deleted: source

Deleted: original

Deleted: of the information

335 communication channels and personal communication. It also emphasizes the need for translation of data and information to specific on-farm decisions, such that information is ready to use once it is passed from its creator to the practitioner's trusted source. Practitioners need to receive translated and contextualized information from people who can help describe why something matters and what information is especially relevant to their particular farm.

340 Practitioners focused on information that is usable, available, and accessible. Data are deemed usable when they are salient, credible, and legitimate, but the value placed on each of these characteristics varies (Haigh et al., 2018). In some contexts, "usable" environmental and climate information referred to information that is "updated on a regular basis and [is] available on a grid that provides continuous coverage over large geographic domains with horizontal resolutions sufficient to capture local and regional differences in drought severity" (Otkin et al., 2018). In other contexts, stakeholders deemed information to be usable when it was trustworthy or reliable (Church et al., 2018; Lemos et al., 2014), familiar (Easton et al., 2017), transparent (Easton et al., 2017), and timely (Stuart et al., 2018).

350 Practitioners were focused not only on usable information, but on applied climate and environmental information that can be used to aid their decision making. For example, "soil moisture" and "soil erosion" are both practical applications of information about precipitation. "Air temperature" was coded less often than expected, which could indicate that practitioners care about derived temperature products more directly applicable to decision-making, such as first and last frost, extreme heat days, or temperature variability in the spring. Although the documents coded in this study rarely mentioned a forecast time scale, Haigh et al. (2015) found that management decisions are often made on seasonal time scales, in the fall and winter preceding planting season. Weekly and monthly forecasts may also be relevant for decisions related to the timing of fertilizer application (see below) (Easton et al., 2017; Haigh et al., 2015b; Kusunose et al., 2019; Mehta et al., 2010).

360 The external factors to whether or not practitioners incorporate their desired climate and environmental information include upfront costs such as advisory support, group involvement, and relevant existing skills and knowledge (Kuehne et al., 2017). In our coding schema, "upfront costs" is most closely analogous to financial, human, and physical capacity. These different forms of capacity affect the ability of a practitioner to implement a decision once it has been made, and their existence (or lack thereof) could persuade or dissuade a practitioner from using the requested information in the first place. Infrastructure and funding support were mentioned more frequently in the document set than human resources, but this does not necessarily suggest that such resources are not a priority; instead, these human resources, which are more

Deleted: Data usability, availability, and accessibility were frequently, in the context of how practitioners will find and use existing data.

Deleted: varied

Deleted: "[

Deleted:]

Deleted: /

Deleted: In addition to a focus on who delivers their

Deleted: practitioners were focused on several kinds of

Moved up [1]: "Precipitation" was the most frequently coded data type, and it often co-occurred with "Forecast."

Deleted: in particular.

Deleted: Precipitation was often discussed in the context of extreme events, particularly drought and extreme rain events, suggesting

Deleted: practitioners have either experienced these events in the recent past, or are wary of their increased frequency in the future. The other two frequently mentioned data types - soil moisture and soil erosion - are both closely related to precipitation, illustrating one application of precipitation data to practitioners'

Deleted: -

Deleted: The

Deleted: but

Deleted: Interestingly, "Air Temperature" was coded more than 50% less frequently than "Precipitation", despite the typical association of these two data types with weather. This does not necessarily indicate that air temperature is unimportant, however. Rather, practitioners might care about

Deleted:Section Break (Next Page)..... derived temperature products such as first and last frost, extreme heat days,

Deleted: temperature variability in the spring, that are more directly applicable to decision-making.

375 [17]

Deleted: is influenced by

Deleted: Upfront

Deleted: can also refer

Deleted: , as defined in our coding schema

Deleted: The most

Deleted: forms of capacity

Deleted: this

Deleted: were infrastructure and funding support. Th [18]

Deleted: such as staff time, staff members' existing [19]

focused on a project's continuation, could be secondary to the financial and infrastructural resources that enable a project's implementation.

In contrast, while capacity might enable a practitioner to make a decision, structural barriers prevent it. Challenges such as the structure of seed markets, laws governing water management, and uncooperative landlords were all given as reasons for why practitioners either could not or would not change their practices, even considering improved information or improved management strategies. Insurance can insulate against risk, allowing farmers to continue with "business as usual" and resist the adoption of conservation measures such as cover crops (Upadhaya and Arbuckle, 2021) or efficient nitrogen application (Stuart et al., 2014). Insurance regulations can also discourage trying new methods if the regulations are not flexible (Roesch-McNally et al., 2018).

Regardless of capacity or structural barriers, risk could ultimately affect how practitioners use information and make decisions. The relative advantage of using information or adopting a practice depends on both the practitioner's tolerance of risk and the risk of implementing the practice itself (Kuehne et al., 2017). Because this study was focused on information use but not the implementation of new conservation practices, it is difficult to assess the risk of the conservation practices themselves, and most of this discussion is centered on practitioners' perception of risk. Risk management took several forms in the document set: risk versus benefits of adopting new sustainable practices; reducing the risk associated with extreme events such as drought; perception of climate change risk; risk tolerance; using climate and environmental information in risk management; and financial risk. In many cases, a practitioner's perception of risk and existing risk reduction strategies affected their willingness to incorporate new information into decision-making. For example, installing irrigation requires up-front costs that may only be recouped during dry years, or if the climate becomes increasingly dry with time (Van Dop 2015). Thus, only practitioners who thought water supply could become unstable were likely to utilize irrigation (Church et al., 2018), while others filed it away as a practice they would never adopt (Bitterman et al., 2019).

Agricultural management and water management were both mentioned frequently in the context of risk management. Water management decisions, for example, might make a field or rangeland more resilient to drought risk, and the decision to implement irrigation was often mentioned simultaneously with practitioners' perceived risk of water shortages. Deciding when to apply nutrients is influenced by the balance between weather-induced and economic-induced risk, such that practitioners can maximize their yield. Stakeholders' willingness to adopt existing management strategies (irrigation, cover crops, etc.) and their interest in innovative strategies were affected by their perception of risk, but the reverse was also true - their perception of climate and weather risk was reduced if they already utilized risk management

Deleted:

Deleted: ¶

¶

¶

405 ¶

¶

¶

410 ¶

¶

¶

415 ¶

¶

¶

420 ¶

¶

¶

425 ¶

¶

¶

¶

¶

430 ¶

Column Break

Deleted: and understanding

Moved (insertion) [2]

400

strategies. Most of the stakeholders interviewed were concerned with near-term agricultural and water management decisions, such as cover crops, when ~~and~~ how to till, ~~and~~ nitrogen application. Their concerns were less focused on long-term trends, and historical ~~or~~ climatological data ~~were~~ only relevant in the context that ~~they~~ informed current decision-making.

405

As mentioned in Sect. 3.2.4, the information that practitioners need, who they get it from, and the decisions that they make were overwhelmingly motivated by desired economic outcomes. In the context of Kuehne et al.'s framework, the code "economic" could refer to profit orientation, profit benefit in the future, profit benefit in the years a practice is used, the time for profit benefits to be realized, or upfront costs (Kuehne et al., 2017). ~~Practitioners were primarily concerned with maximizing their yield and utilizing available~~ markets, buyers, and contracts to profit off their crop. Economic opportunities ~~and~~ markets were a concern for farmers considering adopting new crops or participating in government sustainable management programs.

410

4.2 Themes that were not discussed

415

Several topics were defined in the coding schema ~~but~~ not discussed in the document set. This does not mean that the topic is unimportant, but rather that it did not arise given how the code was defined; stakeholders could define it differently or assign it a different indicator than what was named in the schema. In some instances, stakeholders approach management decisions qualitatively and experientially rather than quantitatively or with data. For example, interviews with Extension agents on the DAWN project have revealed that farmers might determine soil moisture by kicking it, not by instrument-based measurements. The missing codes discussed below should be interpreted in this context, with the understanding that all of this study's available information is dependent on studies that have already been conducted.

420

First, collaboration goals were only mentioned in 6 documents, which could suggest that practitioners are more focused on data delivery than data creation. Given the prevalence of the code "Communication Channels", however, collaborations might occur in ways that are not explicitly mentioned by practitioners. For example, several documents discussed proposed collaborations, such as between government agencies and between research agencies, which could suggest a desire for improved efficiency in collaboration and data delivery.

425

Another topic that was expected but mentioned infrequently was the geospatial scale of information (6 documents), which contrasts with information from Extension specialists on the DAWN team that says that people often request field-scale information. This could be because the types of data most often discussed are already available at the scale that practitioners need. It could also be because information sources are

Deleted: /

Deleted: , etc

Deleted: /

Deleted: was

Deleted: it

Deleted: Overwhelmingly, when the code "economic" was used in our coding schema, the document discussed profit orientation and profit benefit in the year a practice was used.

Deleted: ¶

Deleted: ¶

¶

¶

435 ¶

¶

¶

¶

¶

440 ¶

Column Break

Deleted: Desired outcomes were rarely environmental or societal in this document set, except for when the researchers were investigating environmental issues directly, and even in those scenarios the practitioners indicated that it was not their primary concern. In some instances, economic factors motivated decisions that could lead to a synergistic environmental benefit, such as cover crops or contour farming. ¶

4.2 →

Deleted: ~~~~~Section Break (Continuous)~~~~~

¶

¶

¶

¶

445 ¶

¶

¶

¶

450 ¶

¶

¶

¶

¶

455 ¶

¶

... [20]

Deleted: that were

430 already localized: as exemplified under the sub-node, "Communication Channels", most practitioners get their information from trusted (local) sources such as consultants or crop advisors.

Finally, the node "Social Issues" was not used at all in the document set. This is not uncommon: social science and social issues are not often mentioned in stakeholder needs analysis documents (e.g. Dilling et al. 2015; Molino et al. 2020). This could be because most people do not link social and environmental issues when asked about climate information. In general, the themes that were not mentioned in the document set emphasize our conclusion that practitioners mentioned standards of practice far less often than they mentioned usable data, management challenges, and desired outcomes.

4.3 Research gaps, implications, and future work

440 Because only peer-reviewed academic literature was publicly available for the study, the focus of the document set was on what researchers found important to ask; as a result, some topics might have been missed that are important to stakeholders but are not frequently discussed in research. In addition, authors' affiliations were primarily academic institutions, and research funding was primarily grant-based. As a result, the documents were skewed towards those states and institutions which readily fund stakeholder and agriculture research.

445 The results outlined here have implications for both the DAWN project and further research. First, the fact that practitioners get their information from personal sources highlights the need to promote and explain the DAWN dashboard through existing channels, because potential users are not likely to find a new online dashboard otherwise. Second, for forecasts to be relevant, the DAWN dashboard and other decision support tools should seek to provide information on weekly and seasonal time scales that can directly inform management decisions. Third, the focus on risk management but not long-term forecasts or climatological data highlights an opportunity for education: some risk management decisions, such as infrastructure investments, must be made on time scales longer than annual. It is therefore important to communicate which factors contribute to potential risk and on which time scales.

455 Future research should include public fora with stakeholders where questions are more open-ended and less guided by existing research interests. It might also prove useful to conduct follow-up surveys with stakeholders who have already provided input to learn about changes in priority over time, in the context of new weather events and updated environmental information. If stakeholders are unavailable for follow-up research, similar goals can be achieved by interviewing organizations or Extension staff that work with stakeholders; valuable information can be gleaned this way without the need to interview practitioners directly.

Deleted: ¶ ... [21]

Deleted:Section Break (Continuous).....

¶

¶

465 ¶

¶

¶

¶

¶

470 ¶

.....Column Break.....

Deleted: papers

Deleted: ¶

4.3 →

Deleted: Gaps, Implications

Deleted: Future Work

Deleted:Section Break (Continuous).....

¶

¶

¶

475 ¶

¶

¶

480 ¶

¶

¶

485 ¶

.....Column Break.....

Deleted: ¶

490 ¶

¶

¶

¶

¶

495 ¶

¶

¶

... [22]

460 5 **Conclusion**

465 We analyzed 50 documents about stakeholder climate data needs in the U.S. Corn Belt. The most common themes considered practitioners' decision-making process: from whom they get their information, what information they need, the decisions they can control, what affects their decision-making, and what their desired outcomes are. Collaboration goals, social issues, and data geospatial scale were mentioned less often, indicating a lower priority, a knowledge gap, insufficient research methods, or some combination of these three. The conclusions presented here can inform the future development of decision support tools both within and beyond the DAWN project. Future research should seek to collect information that is motivated as much as possible by stakeholders' needs, rather than by scientists' research priorities. This study identifies the starting point for future studies, such that they are efficient and reduce stakeholder fatigue. It also serves as an example for the background research that scientists can and should do when initiating a project that requires stakeholder engagement: the method presented here can easily be applied to other geographies and sectors.

Deleted: ¶

Deleted: ¶
This paper

Deleted: ~~~~~Section Break (Next Page)~~~~~
¶
¶
510 ¶
¶
¶
¶
¶
515 ¶
¶
¶
¶
¶
520 ¶
¶
¶
¶
¶
525 ¶
~~~~~Column Break~~~~~

### Code/Data Availability

The full coding schema is available in the article's Supplementary Material.

### **Author Contributions**

Suzanna Clark – conceptualization, data curation, formal analysis, methodology, project administration, investigation, supervision, visualization, writing – original draft & preparation, writing – review and editing

Felix Wolfinger – data curation, formal analysis, investigation, writing – review & editing

Melissa Kenney – conceptualization, funding acquisition, methodology, supervision, writing – review & editing

Michael Gerst – conceptualization, writing – review & editing

Heidi Roop – conceptualization, writing – review & editing

### **Competing Interests**

Some authors are members of the editorial board of Geoscience Communication. The peer-review process was guided by an independent editor, and the authors have no other competing interests to declare.

### **Acknowledgements**

This work was supported by the United States Department of Agriculture-National Institute of Food and Agriculture grant number 20206801231674, awarded to the University of Maryland for the creation of a predictive decision support Dashboard for Agricultural Water use and Nutrient management (DAWN).

Deleted: ¶

Deleted:

Moved down [3]: Heidi Roop – conceptualization, writing – review & editing

Deleted:

Moved (insertion) [3]

Deleted: ¶

Deleted: ¶

Deleted: also

Deleted: ¶

Deleted: ¶

Deleted: ~~~~~Section Break (Next Page)~~~~~

¶

¶

¶

530 ¶

¶

¶

¶

535 ¶

¶

¶

¶

540 ¶

¶

¶

545 ¶

¶

¶

¶

550 ¶

¶

¶

¶

¶

555 ¶

¶

¶

¶

560 ¶

~~~~~Column Break~~~~~

References

Agriculture, U.S. Department of. 2022. Cover crops - keeping soil in place while providing other benefits

Deleted: ¶

Andrys, J., Lyons, T. J., and Kala, J.: Multidecadal evaluation of WRF downscaling capabilities over Western Australia in simulating rainfall and temperature extremes, *J. Appl. Meteorol. Climatol.*, 54, 370–394, <https://doi.org/10.1175/JAMC-D-14-0212.1>, 2015.

Deleted: ¶

Angel, J. R., Widhalm, M., Todey, D., Massey, R., and Biehl, L.: The U2U Corn Growing Degree Day tool: tracking corn growth across the U.S. Corn Belt, *Clim. Risk Manag.*, 15, 73–81, <https://doi.org/10.1016/j.crm.2016.10.002>, 2017.

Angel, J. R., Swanson, C., Boustead, B. M., Conlon, K., Hall, K. R., Jorns, J. L., Kunkel, K. E., Lemos, M. C., Lofgren, B. M., Ontl, T., Posey, J., Stone, K., Takle, E., and Todey, D.: Chapter 21: Midwest. Impacts, risks, and adaptation in the United States: the fourth national climate assessment, volume II, 872–940 pp., <https://doi.org/10.7930/NCA4.2018.CH21>, 2018.

Deleted: ¶

Arslan, A., McCarthy, N., Lipper, L., Asfaw, S., Cattaneo, A., and Kokwe, M.: Climate smart agriculture? Assessing the adaptation implications in Zambia, *J. Agric. Econ.*, 66, 753–780, <https://doi.org/10.1111/1477-9552.12107>, 2015.

Deleted: ¶

Baker, Z., Ekstrom, J. A., Meagher, K. D., Preston, B. L., and Bedsworth, L.: The social structure of climate change research and practitioner engagement: evidence from California, *Glob. Environ. Chang.*, 63, 102074, <https://doi.org/10.1016/j.gloenvcha.2020.102074>, 2020.

Beeton, T. A. and McNeeley, S. M.: Who, what, where, when, and how? A typology of drought decision-making on public and tribal lands in the north-central united states, *Weather. Clim. Soc.*, 12, 611–627, <https://doi.org/10.1175/WCAS-D-19-0137.1>, 2020.

Deleted: ¶

Bickman, L. and Rog, D. J.: *Handbook of Applied Social Research Methods*, 2nd ed., SAGE, 2009.

Bitterman, P., Bennett, D. A., and Secchi, S.: Constraints on farmer adaptability in the Iowa-Cedar River Basin, *Environ. Sci. Policy*, 92, 9–16, <https://doi.org/10.1016/j.envsci.2018.11.004>, 2019.

Bowen, G. A.: Document analysis as a qualitative research method, *Qual. Res. J.*, 9, <https://doi.org/https://doi.org/10.3316/QRJ0902027>, 2009.

Cash, D. W., Clark, W. C., Alcock, F., Dickson, N. M., Eckley, N., Guston, D. H., Jäger, J., and Mitchell, R. B.: Knowledge systems for sustainable development, *Proc. Natl. Acad. Sci. U. S. A.*, 100, 8086–8091, <https://doi.org/10.1073/pnas.1231332100>, 2003.

Chawla, I., Osuri, K. K., Mujumdar, P. P., and Niyogi, D.: Assessment of the weather research and forecasting (WRF) model for simulation of extreme rainfall events in the upper Ganga Basin, *Hydrol. Earth Syst. Sci.*, 22, 1095–1117, <https://doi.org/10.5194/hess-22-1095-2018>, 2018.

Church, S. P., Dunn, M., Babin, N., Mase, A. S., Haigh, T., and Prokopy, L. S.: Do advisors perceive climate change as an agricultural risk? An in-depth examination of Midwestern U.S. Ag advisors' views on drought, climate change, and risk management, *Agric. Human Values*, 35, 349–365, <https://doi.org/10.1007/s10460-017-9827-3>, 2018.

Derner, J. D. and Augustine, D. J.: Adaptive management for drought on rangelands, 38, 211–215, <https://doi.org/10.1016/j.rala.2016.05.002>, 2016.

Dewulf, A., Klenk, N., Wyborn, C., and Lemos, M. C.: Usable environmental knowledge from the perspective of decision-making: the logics of consequentiality, appropriateness, and meaningfulness, *Curr. Opin. Environ. Sustain.*, 42, 1–6, <https://doi.org/10.1016/j.cosust.2019.10.003>, 2020.

Dilling, L. and Berggren, J.: What do stakeholders need to manage for climate change and variability? A document-based analysis from three mountain states in the Western USA, *Reg. Environ. Chang.*, 15, 657–667, <https://doi.org/10.1007/s10113-014-0668-y>, 2015.

Dilling, L. and Lemos, M. C.: Creating usable science: opportunities and constraints for climate knowledge use and their implications for science policy, *Glob. Environ. Chang.*, 21, 680–689, <https://doi.org/10.1016/j.gloenvcha.2010.11.006>, 2011.

Easton, Z. M., Kleinman, P. J. A., Buda, A. R., Goering, D., Emberston, N., Reed, S., Drohan, P. J., Walter, M. T., Guinan, P., Lory, J. A., Sommerlot, A. R., and Sharpley, A.: Short-term forecasting tools for agricultural nutrient management, *J. Environ. Qual.*, 46, 1257–1269, <https://doi.org/10.2134/jeq2016.09.0377>, 2017.

Ghil, M. and Jiang, N.: Recent forecast skill for the El Nino/Southern Oscillation, *Geophys. Res. Lett.*, 25, 171–174, 1998.

Goodman, L. A.: Snowball sampling, *Ann. Math. Stat.*, 32, 148–170, 1961.

Deleted: ¶

Deleted: ¶

565 ¶

¶

¶

¶

¶

570 ¶

¶

¶

575 ¶

¶

¶

580 ¶

¶

¶

¶

585 ¶

¶

¶

590 ¶

¶

595 ¶

Column Break

Deleted: ¶
Saving money, time and soil: the economics of no-till farming: ¶

Deleted: ¶

Deleted: →

Deleted: →

Deleted: →

Deleted: →

Deleted: ¶

Deleted: ¶

Deleted: ¶

Deleted: ¶

Haigh, T., Morton, L. W., Lemos, M. C., Knutson, C., Prokopy, L. S., Lo, Y. J., and Angel, J.: Agricultural advisors as climate information intermediaries: exploring differences in capacity to communicate climate, *Weather. Clim. Soc.*, 7, 83–93, <https://doi.org/10.1175/WCAS-D-14-00015.1>, 2015a.

550 Haigh, T., Takle, E., Andresen, J., Widhalm, M., Carlton, J. S., and Angel, J.: Mapping the decision points and climate information use of agricultural producers across the U.S. Corn Belt, *Clim. Risk Manag.*, 7, 20–30, <https://doi.org/10.1016/j.crm.2015.01.004>, 2015b.

| [Haigh, T., Koundinya, V., Hart, C., Klink, J., Lemos, M., Mase, A. S., Prokopy, L., Singh, A., Todey, D., and Widhalm, M.: Provision of climate services for agriculture: public and private pathways to farm](#)
555 [decision-making, *Bull. Am. Meteorol. Soc.*, 99, 1781–1789, <https://doi.org/10.1175/BAMS-D-17-0253.1>, 2018.](#)

| [Haigh, T. R., Otkin, J. A., Mucia, A., Hayes, M., and Burbach, M. E.: Drought early warning and the timing of range managers' drought response, *Adv. Meteorol.*, 2019, <https://doi.org/10.1155/2019/9461513>, 2019.](#)

560 [Haigh, T., Hayes, M., Smyth, J., Prokopy, L., Francis, C., and Burbach, M.: Ranchers' use of drought contingency plans in protective action decision making, *Rangel. Ecol. Manag.*, 74, 50–62, <https://doi.org/10.1016/j.rama.2020.09.007>, 2021.](#)

Hochman, Z. and Carberry, P. S.: Emerging consensus on desirable characteristics of tools to support farmers' management of climate risk in Australia, *Agric. Syst.*, 104, 441–450,
565 <https://doi.org/10.1016/j.agry.2011.03.001>, 2011.

| [Hunt, E. D., Birge, H. E., Laingen, C., Licht, M. A., McMechan, J., Baule, W., and Connor, T.: A perspective on changes across the U.S. Corn Belt, *Environ. Res. Lett.*, 15, <https://doi.org/10.1088/1748-9326/ab9333>, 2020.](#)

Jones, R. N., Chiew, F. H. S., Boughton, W. C., and Zhang, L.: Estimating the sensitivity of mean annual runoff to climate change using selected hydrological models, *Adv. Water Resour.*, 29, 1419–1429,
570 <https://doi.org/10.1016/j.advwatres.2005.11.001>, 2006.

| [Kenney, M. A., Janetos, A. C., and Lough, G. C.: Building an integrated U.S. National Climate Indicators System, *Clim. Change*, 135, 85–96, <https://doi.org/10.1007/s10584-016-1609-1>, 2016.](#)

Deleted: ¶

Moved down [4]: Haigh, T., Hayes, M., Smyth, J., Prokopy, L., Francis, C., and Burbach, M.: Ranchers' use of drought contingency plans in protective action decision making, *Rangel. Ecol.*

Moved down [5]: *Manag.*, 74, 50–62,

Deleted: ¶

Deleted: ¶

600 ¶

¶

¶

¶

605 ¶

¶

¶

¶

610 ¶

¶

¶

¶

615 ¶

¶

¶

¶

620 ¶

¶

¶

¶

625 ¶

¶

¶

630 ¶

Column Break ... [23]

Moved (insertion) [4]

Deleted: ¶

Moved (insertion) [5]

Deleted: Hrozencik, R. A. and Aillery, M.: Trends in U.S. irrigated agriculture: increasing resilience under water supply scarcity, U.S. Department of Agriculture, Economic Research Service, 1–55 pp., <https://doi.org/10.2139/ssrn.3996325>, 2021. ¶

Deleted: ¶

575 Kirchhoff, C. J., Lemos, M. C., and Dessai, S.: Actionable knowledge for environmental decision making: broadening the usability of climate science, *Annu. Rev. Environ. Resour.*, 38, 393–414, <https://doi.org/10.1146/annurev-environ-022112-112828>, 2013.

Klemm, T. and McPherson, R. A.: The development of seasonal climate forecasting for agricultural producers, *Agric. For. Meteorol.*, 232, 384–399, <https://doi.org/10.1016/j.agrformet.2016.09.005>, 2017.

580 Knutson, C. and Haigh, T.: A drought-planning methodology for ranchers in the great plains, 35, 27–33, <https://doi.org/10.2111/RANGELANDS-D-12-00075.1>, 2013.

Kuehne, G., Llewellyn, R., Pannell, D. J., Wilkinson, R., Dolling, P., Ouzman, J., and Ewing, M.: Predicting farmer uptake of new agricultural practices: a tool for research, extension and policy, *Agric. Syst.*, 156, 115–125, <https://doi.org/10.1016/j.agsy.2017.06.007>, 2017.

585 Kusunose, Y., Ma, L., and VAN SANFORD, D.: User responses to imperfect forecasts: findings from an experiment with Kentucky wheat farmers, *Weather. Clim. Soc.*, 11, 791–808, <https://doi.org/10.1175/WCAS-D-18-0135.1>, 2019.

Lemos, M. C., Kirchhoff, C. J., and Ramprasad, V.: Narrowing the climate information usability gap, *Nat. Clim. Chang.*, 2, 789–794, <https://doi.org/10.1038/nclimate1614>, 2012.

590 Lemos, M. C., Kirchhoff, C. J., Kalafatis, S. E., Scavia, D., and Rood, R. B.: Moving climate information off the shelf: Boundary chains and the role of risas as adaptive organizations, *Weather. Clim. Soc.*, 6, 273–285, <https://doi.org/10.1175/WCAS-D-13-00044.1>, 2014.

Mehta, V. K., Knutson, C. L., Rosenberg, N. J., Olsen, J. R., Wall, N. A., Bernadt, T. K., and Hayes, M. J.: An assessment of decadal drought information needs of stakeholders and policymakers in the Missouri River Basin for decision support, 1–15 pp., 2010.

595 Molino, G. D., Kenney, M. A., and Sutton-Grier, A. E.: Stakeholder-defined scientific needs for coastal resilience decisions in the northeast U.S., *Mar. Policy*, 118, 103987, <https://doi.org/10.1016/j.marpol.2020.103987>, 2020.

Moya-álvarez, A. S., Gálvez, J., Holguín, A., Estevan, R., Kumar, S., Villalobos, E., Martínez- Castro, D., and Silva, Y.: Extreme rainfall forecast with the WRF-ARW model in the Central Andes of Peru, *Atmosphere (Basel)*, 9, <https://doi.org/10.3390/atmos9090362>, 2018.

600

Deleted: ¶

Deleted: →

Deleted: →

Deleted: →

Deleted: →

Deleted: →

Deleted: ¶

Deleted: ¶

Deleted: ¶

Deleted: ¶

Deleted: ¶

635 ¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

Column Break

Deleted: ¶

Deleted: ¶

Otkin, J. A., Shafer, M., Svoboda, M., Wardlow, B., Anderson, M. C., Hain, C., and Basara, J.: Facilitating the use of drought early warning information through interactions with agricultural stakeholders, *Bull. Am. Meteorol. Soc.*, 96, 1073–1078, <https://doi.org/10.1175/BAMS-D-14-00219.1>, 2015.

605 Otkin, J. A., Haigh, T., Mucia, A., Anderson, M. C., and Hain, C.: Comparison of agricultural stakeholder survey results and drought monitoring datasets during the 2016 U.S. Northern Plains flash drought, *Weather. Clim. Soc.*, 10, 867–883, <https://doi.org/10.1175/WCAS-D-18-0051.1>, 2018.

Palutikof, J. P., Street, R. B., and Gardiner, E. P.: Decision support platforms for climate change adaptation: an overview and introduction, *Clim. Change*, 153, 459–476, [https://doi.org/10.1007/s10584-](https://doi.org/10.1007/s10584-019-02445-2)
610 019-02445-2, 2019.

Podestá, G. P., Natenzon, C. E., Hidalgo, C., and Ruiz Toranzo, F.: Interdisciplinary production of knowledge with participation of stakeholders: a case study of a collaborative project on climate variability, human decisions and agricultural ecosystems in the Argentine Pampas, *Environ. Sci. Policy*, 26, 40–48, <https://doi.org/10.1016/j.envsci.2012.07.008>, 2013.

615 Prokopy, L. S., Haigh, T., Mase, A. S., Angel, J., Hart, C., Knutson, C., Lemos, M. C., Lo, Y. J., McGuire, J., Morton, L. W., Perron, J., Todey, D., and Widhalm, M.: Agricultural advisors: a receptive audience for weather and climate information?, *Weather. Clim. Soc.*, 5, 162–167, <https://doi.org/10.1175/WCAS-D-12-00036.1>, 2013.

620 Ranjan, P., Singh, A. S., Tomer, M. D., Lewandowski, A. M., and Prokopy, L. S.: Lessons learned from using a decision-support tool for precision placement of conservation practices in six agricultural watersheds in the US Midwest, *J. Environ. Manage.*, 239, 57–65, <https://doi.org/10.1016/j.jenvman.2019.03.031>, 2019.

Roesch-Mcnally, G. E., Basche, A. D., Arbuckle, J. G., Tyndall, J. C., Miguez, F. E., Bowman, T., and Clay, R.: The trouble with cover crops: farmers’ experiences with overcoming barriers to adoption, *Renew. Agric. Food Syst.*, 33, 322–333, <https://doi.org/10.1017/S1742170517000096>, 2018.

625 Rose, S.: The Inevitability of Climate Adaptation in U.S. Agriculture, *Choices Mag. Food, Farm, Resour. Issues*, 30, 1–5, <https://doi.org/10.22004/ag.econ.206238>, 2015.

Saldana, J.: The coding manual for qualitative researchers, <https://doi.org/10.1108/qrom-08-2016-1408>, 2013.

Deleted: ¶

Deleted: ¶

Deleted: ¶

Deleted: ¶

Deleted: ~~~~~Section Break (Next Page)~~~~~

¶

¶

675 ¶

¶

¶

¶

680 ¶

¶

¶

¶

685 ¶

¶

¶

¶

690 ¶

¶

¶

¶

695 ¶

~~~~~Column Break~~~~~

Deleted: ¶

655 Sarkki, S., Tinch, R., Niemelä, J., Heink, U., Waylen, K., Timaeus, J., Young, J., Watt, A., Neßhöver, C.,  
and van den Hove, S.: Adding “iterativity” to the credibility, relevance, legitimacy: a novel scheme to  
highlight dynamic aspects of science-policy interfaces, *Environ. Sci. Policy*, 54, 505–512,  
https://doi.org/10.1016/j.envsci.2015.02.016, 2015.

Deleted: ¶

660 Stuart, D., Schewe, R. L., and McDermott, M.: Reducing nitrogen fertilizer application as a climate  
change mitigation strategy: understanding farmer decision-making and potential barriers to change in the  
US, *Land use policy*, 36, 210–218, https://doi.org/10.1016/j.landusepol.2013.08.011, 2014.

Deleted: ¶

665 Stuart, D., Denny, R. C. H., Houser, M., Reimer, A. P., and Marquart-Pyatt, S.: Farmer selection of  
sources of information for nitrogen management in the US Midwest: Implications for environmental  
programs, *Land use policy*, 70, 289–297,  
https://doi.org/10.1016/j.landusepol.2017.10.047, 2018.

Deleted: →

Deleted:

Stumpf, R. P., Johnson, L. T., Wynne, T. T., and Baker, D. B.: Forecasting annual cyanobacterial bloom  
biomass to inform management decisions in Lake Erie, *J. Great Lakes Res.*, 42, 1174–1183,  
https://doi.org/10.1016/j.jglr.2016.08.006, 2016.

670 Upadhaya, S. and Arbuckle, J. G.: Examining factors associated with farmers’ climate-adaptive and  
maladaptive actions in the U.S. Midwest, *Front. Clim.*, 3, https://doi.org/10.3389/fclim.2021.677548,  
2021.

Deleted: ¶

Van Dop, M.A.: Irrigation adoption, groundwater demand and policy in the U.S. Corn Belt, 2040-2070,  
M.S. Thesis. Purdue University, 2016.

675 Wiggins, A., Young, A., and Kenney, M. A.: Exploring visual representations to support data re- use for  
interdisciplinary science, *Proc. Assoc. Inf. Sci. Technol.*, 55, 554–563,  
https://doi.org/10.1002/prat.2018.14505501060, 2018.

Yin, R. K.: Case study research: design and methods, 5th ed., Thousand Oaks, CA, 2009.

Deleted: ¶

Page 2: [1] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
Page 3: [2] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
Page 4: [3] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

Page 5: [4] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
Page 6: [5] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

Page 6: [6] Deleted      Suzi Clark      2/6/23 1:53:00 PM

.....

▼  
.....

1 . 1

Page 6: [7] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

Page 7: [8] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

Page 26: [9] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

Page 9: [10] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
Page 13: [11] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

Page 13: [12] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

Page 13: [13] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

Page 26: [14] Deleted      Suzi Clark      2/6/23 1:53:00 PM

▼  
.....

▼  
.....

**Page 26: [15] Deleted      Suzi Clark      2/6/23 1:53:00 PM**



**Page 26: [16] Deleted      Suzi Clark      2/6/23 1:53:00 PM**



**Page 15: [17] Deleted      Suzi Clark      2/6/23 1:53:00 PM**



**Page 15: [18] Deleted      Suzi Clark      2/6/23 1:53:00 PM**



**Page 15: [19] Deleted      Suzi Clark      2/6/23 1:53:00 PM**



**Page 17: [20] Deleted      Suzi Clark      2/6/23 1:53:00 PM**



**Page 18: [21] Deleted      Suzi Clark      2/6/23 1:53:00 PM**



**Page 18: [22] Deleted      Suzi Clark      2/6/23 1:53:00 PM**



**Page 23: [23] Deleted      Suzi Clark      2/6/23 1:53:00 PM**

