Authors' response to referee comments by Referee #1

Dear Editor,

Please find below the response on behalf of all authors to the comments posted by Referee #1 on our manuscript "Epidote dissolution–precipitation during viscous granular flow: a micro-chemical and isotope study" (MS id: egusphere-2022-311).

To make the response easier to read and to assess, we have posted our own comments next to/below each comment in the PDF review files posted by the Referee.

We thank Referee #1 for their constructive review, and hope that you will find this response and the revised manuscript satisfactory.

Yours sincerely, Veronica Peverelli, on behalf of all authors



EGUsphere, referee comment RC1 https://doi.org/10.5194/egusphere-2022-311-RC1, 2022 © Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Comment on egusphere-2022-311

Anonymous Referee #1

Referee comment on "Epidote dissolution-precipitation during viscous granular flow: a micro-chemical and isotope study" by Veronica Peverelli et al., EGUsphere, https://doi.org/10.5194/egusphere-2022-311-RC1, 2022

The study of Peverelli and co-authors entitled "Epidote dissolution-precipitation during viscous granular flow: a micro-chemical and isotope study" investigates an epidote and quartz vein from the Aar Massif with several analytical techniques. The authors conclude that during deformation quartz experienced dynamic recrystallization by subgrain rotation that decreased its grain size. Consequently, grain boundary sliding, creep cavitation, and viscous granular flow produced the investigated microstructure. During these processes, epidote experienced dissolution and (re)precipitation responsible for pinning quartz grain size at grain boundaries.

Although these sequence and association of deformation mechanisms is coherent with what has been popsed in literature in several rock types and geological settings, at the present the petrographic and microstructural description and data are not sufficient to support these interpretations. In my opinion, this can be fixed with major revisions, aimed at expanding this section of the manuscript. Furthermore, discussion should be expanded, adding more references of relevant2 tricles from other research groups and discussing novel data in the light of those works.

Below a few major comments, several comments are present in the attached pdf.

- Streamline of the text: interpretations of deformation mechanisms appear too early in the text and are not upported by documentation. I suggest moving those in the discussion section expanding this session in the light of relevant literature (at the moment largely missing)
- Section 3 need to be largely implemented, both for text and microphotos. At the moment the reader conflat assess what it is written and cannot place the different epidote vein genera s, shear zone and host rock in context

Summary of Comments on egusphere-2022-311-RC1_VP.pdf

Page: 1

	Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:20:42 AM
1	Petrographic and	microstructural data have be	en complemented with addi	tional figures, data and text.
	Number: 2	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:21:10 AM
1	More relevant refe	erences have been added		
	Number: 3	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/17/2022 8:00:47 AM
÷.	Moved all major in	nterpretations to the discuss	ion. However, a few interpret	ations that are meant to answer minor aspects of the
	manuscript have b	been left before the discussion	on session because they are r	not elaborated further and they are not crucial to the main
	message of the m	anuscript.		
Ţ	Number: 4	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:22:47 AM
	Done			

- EBSD documentation need to be expanded both in data and text, following comments in the attached PDF (e.g. complete EBSD map is present in appendix C, without low and high angle boundaries displayed in the map and text explaining it).
- The authors conclude that subgrain rotation recrystallization was the main deformation mechanism of quartz in results section. They state that CPO formed due to this mechanism. What we happened to this CPO when grain boundary sliding, creep cavitation, and viscous granular flow occurred? Would you expect an attenuation of this CPO, as several authors suggested? This discussion is completely absent in the manuscript. Please, add EBSD documentation, such as GROD or GOS or KAM grain size maps along your map in appendix C, pole figures also for <u>, misorientation angle distribution</u>. Afterward, please discuss more thoroughly all those deformation mechanisms in the light of relevant literature.
- Dissolution precipitation in epidote: in the conclusions the authors state: "we have demonstrated the occurrence of epidote dissolution-precipitation processes". However, I fear that diss-prec is just a proposition and that there could be other explanations. This is because the authors did not pretext any microstructural characterization of Epidote A grains pointing to diss-prec processes acting, such as lobate edges, microporosity,.... Please, refer to *Putnis 2009 Reviews in Mineralogy & Geochemistry* for microstructures indicative of diss-prec processes. Regardling another possible interpretation, as you imply the presence of an external fluid why you cannot just have new nucleation of epidote in creep cavities (thus syn-deformation) related to the percolating fluid, without dissolving previous Epidote A?

Concluding, I think that with major revisions aimed at including additional data, expanding results, expanding diagraphic section in the light of relevant literature, the authors could substantiate their integrations and conclusions.

Best wishes

Please also note the supplement to this comment: <u>https://egusphere.copernicus.org/preprints/egusphere-2022-311/egusphere-2022-311-RC</u> <u>1-supplement.pdf</u>

	Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:23:01 AM			
	Done, and moved	EBSD data to the main text					
P	Number: 2	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:24:39 AM			
	This is not someth	ing that we can comment or	h based on the present data.	To the authors' knowledge, this is really difficult to solve by			
	investigating natu	ral samples, and therefore it	is beyond the scope of the r	nanuscript.			
P	Number: 3	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:25:11 AM			
	Done						
	Number: 4	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:26:59 AM			
	We appreciate the	e suggestion and clarify the t	ext to exclude misunderstan	dings. The spatial distribution of Epidote-B in layers 2-3 is not			
	consistent with the	e microstructure being entire	ely acquired during crystallization	ation from a fluid. If a secondary fluid produced a second			
	veining event by cracking a previously formed vein, then we would still have a two-stage story: (1) precipitation of epidote from the fluid						
	and (2) deformation. The relative distribution of epidote and quartz, and the Zener relation argue that an additional process has to						
	overprint epidote	crystallization during veining	. If it is true that a previously	formed vein was cracked open leading to the formation of an			
	epidote band in layers 2-3, then the "primary" epidote in these layers is still "Epidote-A" as defined in the text.						

Number: 5	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:27:14 AM	
Done				





Epidote dissolution-precipitation during viscous granular flow: a micro-chemical and isotope study

Veronica Peverelli¹, Alfons Berger¹, Martin Wille¹, Thomas Pettke¹, Pierre Lanari¹, Igor M. Villa^{1,2}, Marco Herwegh¹

⁵ ¹Department of Geological Sciences, University of Bern, Bern, 3012, Switzerland ²Dipartimento di Scienze dell'Ambiente e della Terra, University of Milano-Bicocca, Milan, 20126, Italy *Correspondence to*: Veronica Peverelli (veronica.peverelli@geo.unibe.ch)

Abstract. Deformation of polymineralic aggregates can be accommodated by viscous granular flow, a process mediated by the interplay among intracrystalline plasticity and dissolution–precipitation, each active in specific minerals at given P–T

- 10 conditions. Common rock-forming minerals like quartz, feldspars and sheet silicates have been intensively studied in terms of deformation processes. Instead, the deformation behavior of epidote and its role during viscous granular flow is not well investigated, although this mineral is ubiquitous in granitic rocks deforming at greenschist-facies conditions. In this contribution, we provide microstructural and geochemical evidence for the occurrence of dissolution–precipitation of epidote during deformation of an epidote-quartz vein. The main part of the vein is deformed producing a fold, which is visible due to
- 15 relicts of primary-growth layering inside the vein. The deformation mechanisms active during deformation include dynamic recrystallization of quartz by subgrain rotation recrystallization, producing grain-size reduction of the primary vein quartz. This occurs contemporaneously with dissolution and (re)precipitation of epidote, and grain-boundary sliding, leading to a combined process described as viscous granular flow. The combination of intracrystalline plasticity, grain boundary sliding and dissolution locally and repeatedly produce creep cavities. These represent not only loci for nucleation of new epidote
- 20 grains at the expenses of dissolved one, but they also allow fluid-mediated transport of elements. The same trace element patterns between old epidote relicts and newly formed grains, with much narrower variability, indicate a process of chemical homogenization. The nature of the fluid mediating deformation is investigated using Pb–Sr isotope data of epidote, which suggest that deformation is assisted by internally recycled fluids with the addition of a syn-kinematic external fluid component.

25 **1 Introduction**

30

Deformation microstructures preserved in rocks are the result of the interplay of multiple microscale deformational processes, which may be identified and appreciated through careful petrographic studies (e.g., Passchier, 2005). These processes – which may be coupled to metamorphic reactions and diffusion (e.g., Wintsch and Yeh, 2013; Lanari and Duesterhoeft, 200^{11} are the response of the grains to varying physico-chemical conditions in order to minimize the system's internal energy (e.g., Evans et al., 2001; Herwegh and Berger, 2004; Passchier, 2005; Karato, 2008; Herwegh et al.,

Summary of Comments on egusphere-2022-311_Commented.pdf

Page: 1

Number: 1 Author: a Subject: Sticky Note Date: 22/06/2022 09:17:40 I would add here also ther relevant article that try to do so, such as Pearce et al., 2013_CMP, Bukowskà et al., 2016, JGR, Giuntoli et al., 2018_JMG...

For example those are more relevant than the review article of Lanari and Duesterhoeft, 2019

Summary of Comments on A9RD0F4.tmp

Page: 2

Number: 1 Done

Subject: Sticky Note Author: Veronica Peverelli

Date: 7/21/2022 12:48:05 PM





2011; Hobbs et al., 2010). Microstructures of monomineralic aggregates are determined by the deformation behavior of the constituent mineral and by the conditions (e.g., temperature, stress, availability of fluids) existing during deformation. For example, Stipp et al. (2002) showed that the microstructures resulting from dynamic recrystallization of quartz can be linked to different recrystallization mechanisms active at increasing temperatures. However, this type of interpretations is more 35 complicated when dealing with polymineralic aggregates, where the relative abundances and specific deformation mechanisms of each mineral have an effect on the deformation behavior of the bulk rock (e.g., Handy, 1990; 1994; Olgaard, 1990; Stünitz and Fitz Gerald, 1993; Kruse and Stünitz, 1999; Tullis, 2002; Herwegh and Berger, 2004; Passchier, 2005; Herwegh et al., 2011; Wehrens et al., 2017). As an example, it has been shown that the grain size of calcite in mylonitic carbonates is determined by the grain size and volume abundance of second phases, such as sheet silicates (Olgaard, 1990; 40 Herwegh and Berger, 2004; Herwegh et al., 2011). In this context, mass transfer processes like dissolution-precipitation play a fundamental role in that they enable the redistribution of material within the deforming system (e.g., Herwegh and Jenni, 2001; Konrad-Schmolke et al., 2018 circulating fluid may precipitate dissolved material into intragranular voids of an initially monomineralic aggregate during deformation. In this scenario, the process of dissolution-precipitation in combination with the presence of a second phase keeps the grain sizes of the deforming aggregate small in a process called 45 viscous granular flow (Olgaard, 1990; Fitz Gerald and Stünitz, 1993; Stünitz and Fitz Gerald, 1993; Paterson, 1995; Herwegh and Berger, 2004; Herwegh et al., 2011). Viscous granular flow entails dissolution-precipitation that dissolves mineral asperities, thus facilitating grain boundary sliding in polymineralic aggregates (e.g., Fitz Gerald and Stünitz, 1993; Stünitz and Fitz Gerald, 1993; Paterson, 1995). Grain boundary sliding promotes the formation of creep cavities, hence allowing the nucleation of second-phase minerals (e.g., Herwegh and Jenni, 2001; Fusseis et al., 2009; Gilgannon et al., 2021). Despite the important role of fluids involved in deformation processes, their direct characterization is rare (e.g., 50 Marquer and Burkhard, 1992; Marquer and Peucat, 1994), and it is often unclear whether the fluids are newly added to the

- system during deformation (i.e., of external origin) or if they are recycled (e.g., via dissolution of hydrous minerals). Hence, the interplay of recycled and newly added fluids, mass-transfer processes and deformation mechanisms to produce complex microstructures remains to be fully appreciated.
- 55 In this contribution, via combined microstructural observations and geochemical and Pb–Sr isotope geochemistry, we investigate the deformational processes affecting epidote (i.e., Ca₂Al₂(Al,Fe³⁺)Si₃O₁₂(OH)) and quartz defining a microfold in a deformed epidote-quartz hydrothermal vein. The data demonstrate that the present-day microstructure is the result of the interplay between dissolution–precipitation of epidote and dynamic recrystallization of quartz, both of which ensure grain size reduction in viscous granular flow. The nature of the fluid mediating dissolution–precipitation of epidote and the
- 60 dynamic recrystallization of quartz is investigated using Pb–Sr isotope data. Because epidote is a widespread rock-forming, hydrothermal and alteration mineral in granitic rocks (e.g., Bird and Spieler, 2004; Enami et al., 2004; Franz and Liebscher, 2004; Grapes and Hoskin, 2004; Schmidt and Poli, 2004; Morad et al., 2010; Hentschel et al., 2020), the occurrence of epidote dissolution–precipitation and its control on the deformation mechanisms of other rock-forming minerals (e.g., dislocation creep in quartz) has important implications for the structural evolution of the granitic continental crust.

Number: 1 Author: a Subject: Stic Note Date: 22/06/2022 09:10:24

I would state here that diss prec request a fluid phase and cite also other relevant articles e.g. some of Andrew Putnis articles. I see that fluid comes at the end of the paragraph, our would place it at the beginning, as it is essential for diss prec to act

) Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/21/2022 12:48:12 PM	
Done				





65 2 Geological Setting

The Grimsel Pass area (Fig. 1; central Swiss Alps, Switzerland) is in the southern part of the Aar Massif, one of the External Crystalline Massifs of the Alps (e.g. Schneeberger et al., 2019; Berger et al., 2017). The Aar Massif consists of a

- 70 polycyclic basement and Permian intrusives, including the post-Variscan Central Aar Granite. This granitoid was emplaced 299 ± 2 Ma (Schaltegger & Corfu, 1992) and mainly records Alpine deformation (e.g., Choukroune and Gapais, 1983; Bambauer et al., 2009; and references therein). This
- 75 Alpine deformation is expressed by a large number of ductile shear zones, and in the southern Aar Massif it can be subdivided into two major phases: 1) a reverse faulting phase (ca. 22–17 Ma; Challandes et al., 2008; Rolland et al., 2009; Wehrens et al., 2017) and 2) a strike-slip phase (from ca. 14
- 80 Ma onwards; Rolland et al., 2009; Wehrens et al., 2017; Herwegh et al., 2020). In the shear zones, green biotite is stable in the first phase at ≥ 400 °C, and it is progressively replaced



Figure 1: (a) Geological map of the Grimsel Pass area (modified from Wehrens et al., 2016). (b) Geographic location of the Grimsel Pass area (red star) in Switzerland (modified from map.geo.admin.ch).

by chlorite in the second one at lower temperatures (Rolland et al., 2009; Wehrens et al., 2017). The Alpine metamorphic overprint reached greenschist facies conditions in the area, with maximum temperatures and pressures of 450 ± 30 °C and 6
± 1 kbar, respectively (Challandes et al., 2008; Goncalves et al., 2012; Villa and Hanchar, 2013). Hydrothermal activity upon exhumation in Alpine times is recorded in the area primarily by the Grimsel Breccia Fault hydrothermal system (e.g., Hofmann et al., 2004; Belgrano et al., 2016; Diamond et al., 2018; Egli et al., 2018), and by Alpine cleft mineralization (e.g., Mullis et al., 1994; Janots et al., 2012; Berger et al., 2013; 2022; Rossi and Rolland, 2014; Bergemann et al., 2017; Ricchi et al., 2019). U–Pb geochronology of hydrothermal epidote in veins returned Miocene and Permian ages (Peverelli et al., 2021; accepted 1

3 Field relations and sample description

The investigated sample Grimsel-1 is an epidote-quartz (±biotite) vein that was collected inside Nagra's Grimsel Test Site (GTS; Figs. 1–2a), which is an underground rock laboratory in the Grimsel Pass area (see Schneeberger et al., 2019). In the GTS, the characteristic pattern of shear zones (e.g., Goncalves et al., 2012) is well exposed on the tunnel walls. The studied epidote vein runs oblic vein runs oblic vein respect to a steeply dipping shear zone of Alpine age hosted by the Central Aar Granite (Fig. 2b). The shear zone is ca. 10 cm wide, whereas the epidote vein reaches up to a couple of centimeters in wig Three other

Number: 1 1 Author:	a Subject: Sticky Note	Date: 22/06/2022 09:52:40
Please, 🙀 and summa	rize these findings here, like th	is the reader is not forced to jump to your other article. What are the differences
between and two events? A	lso add the precise absolute ag	ge with related error
Number: 2 2 Author:	a Subject: Sticky Note	Date: 22/06/2022 09:53:11
Please, highting t this on the	e image	
Number: 3 3uthor:	a Subject: Sticky Note	Date: 22/06/2022 09:56:00
What is their ength?		

	Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 8:20:10 PM	
1	Done				
	Number: 2	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/21/2022 12:47:55 PM	
1	It should be better visible in the updated figure				
	Number: 3	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:28:10 AM	
1	Detail added				







Figure 2 (a) Geological map of the Grimsel Test Site (GTS) of Nagra with the location of (b) shown in the red rectangle (modified from Schneeberger et al., 2019). (b) Field photograph of the location of the studied epidote-quartz vein (Grimsel-1) and the Alpine shear zone with which it is associated; both are in the Central Aar Granite, which shows increasing foliation towards the shear zone (modified from Peverelli et al., Pacepted).

epidote veins are associated (1) This shear zone (Fig. A1; not visible in Fig. 2b): (1) sample P2 is a weakly deformed epidote-quartz vein in a slightly deformed portion of the host; (2) sample P3 consists of a number of sheared epidote-quartz veins in a mylonitic portion; and (3) sample Gr0-c is a highly sheared epidote-quartz vein which subdivides the host rock into a mylonitic side and a slightly deformed one. Epidote in veins P2–3 returned Permian U–Pb ages (Peverelli et al., accepted). The host Central Aar Granite displays a gradation from slightly to highly deformed: this is expressed by an increasingly intense foliation and by a decreasing grain size moving towards the shear zone (Fig. 2b). In thin section (Fig. 3), the host rock mostly preserves its magmatic texture and it is made of ca. 55 vol. % of altered feldspar and 35 vol. % quartz. Feldspar grains are sometimes fractured, and quartz is mostly dynamically recrystallized by subgrain rotation
recrystallization (SG) Relict plagioclase and K-feldspar grains can be recognized by the different type and extent of

- alteration. Plagioclase (ca. 20 vol. %) is highly altered into epidote and white mica, to a larger extent in the cores than in the rims of the grain
- 110 Exsolution lamellae are preserved in K-feldspar grains (ca. 35 vol. %), which are moderately altered into sericite.Green biotite, epidote and minor chlorite define a weak foliation and
- 115 make up ca. 10 vol. %, with a few accessory titanite grains completing the host rock's mineral assemblage. The sharp boundary between host and vein is marked by a change in modal



Figure 3: Transmitted-light microscope scan of the studied epidote-quartz (±biotite) vein and the host Central Aar Granite. The numbers 1–3 refer to the vein layers described in Sect. 5.1. Dashed rectangles A–D indicate the locations of the microstructural domains shown in Fig. 4a–d. The black rectangle indicates the location of Fig. 10 (rotated by 90° clockwise). Plane-polarized light.

画 Number: 1	Autho1a Subject: Sticky Note	Date: 22/06/2022 09:40:02
Please, increase fo	on size and improve graphics of this	s sketch, maybe adding colors? At the moment it is not easy to read
画 Number: 2	Author: a Subject: Sticky Note	Date: 22/06/2022 10:04:24
Ok, but at the mo this information This is the case	oment it is not clear where those are 2 File Figure 3: it is not clear where t	located with respect to the shear zone, what are the relation among them, Please, add his host-vein are located with respect to the shear zone of Fig. 2b
画 Number: 3	Author: a Subject: Sticky Note	Date: 23/06/2022 09:23:08
You cannot state mechanism. I see that one ma EBSD map	this without showing microstructure p is present at be end of the append	s of it. Please, add thin section images and ideally EBSD maps that suggest SGR as def dix. Therefore, it is better to avoid tis sentence here and say so later in the text citing the
i Number: 4	Auth <u>or;</u> a Subject: Sticky Note	Date: 22/06/2022 10:05:27
Again, also this sh	hould be thown with microphotos, e	ither in the main text or in the supplementary.

Same for all the mainer ineralogical phases of host rock

画 Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/21/2022 12:47:13 PM		
Figure improved					
Number: 2	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:29:02 AM		
Removed referen	ce to extra samples, since the	ey are not discussed further			
回 Number: 3	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:31:18 AM		
Ok, moved to res	ults after having shown the f	igure, and added reference t	o Stipp et al. (2002) to support subgrain rotation		
recrystallization. EBSD data have been improved and added to the main text.					
Number: 4	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 8:29:52 PM		
It does not seem	It does not seem relevant in the context of the manuscript and the host rock is thoroughly described in Schneeberger et al., 2019. Added				

It does not seem relevant in the context of the manuscript and the host rock is thoroughly described in Schneeberger et al., 2019. reference to this paper





abundances.

The object of the present investigation is Grimsel-1 epidote-quartz vein (Fig. 3). This vein is subdivided into three layers: (1) layer with coarse-grained epidote, quartz and minor green biotite, (2) heavily deformed and finer-grained epidote-quartz layer, and (3) nearly pure quartz la Grain sizes, mineral proportions and dominant deformation mechanisms in each layer are given in Table 1. Layer 1 is the least deformed on hillst deformation is most intense in layers 2–3, where a fold is defined by the spatial distribution of epidote and quartz (Fig. 3). The reasons why layer 1 is less deformed than layers 2–3 125 may be linked to the closer proximity of layer 1 to the host rock, to a role of mechanically strong epidote clusters forming a load-bearing network, or to a combination of bo Landy, 1990; 1994; Masuda, 1990; 1995; Tullis, 2002; Passchier, 2005). The modal abundance of green biotite varies greatly among the three layers, with a sharp decrease from layer 1 to layer 2, and layer 3 being devoid of biotite (Table 1). The transition from layer 2 to layer 3 is marked by a change in the 130 epidote/quartz ratio. This is up to ca. 40/60 in layer 2, whereas layer 3 is characterized by the near absence of epidote. Peverelli et al. (2021) performed U-Pb dating by LA-ICP-MS of epidote grains in layer 1, obtaining a Tera-Wasserburg age of 19.2 ± 4.3 Ma. The Tera–Wasserburg regression revealed a single epidote generation at the current analytical precision (i.e., MSWD of 0.79), and the age is interpreted as the timing of epidote crystallization upon vein opening (Peverelli et al., 2021). The focus of this manuscript are the deformation mechanisms in layers 2–3.

135

Table 1: Main characteristics of the vein layers. Ep = epidote; SGR = subgrain rotation recrystallization; Qz = quartz.

Layer	Minerals other than Ep + Qz	Ep [vol. %]	Ep grain size	Qz deformation 4	Qz grain size	Qz relicts
1	Green biotite (ca. 5 vol. %)	ca. 70	0.025–1.6 mm	Dislocation glide + SGR	80 400 μm	0.6–1.2 mm
2	Green biotite (ca. 1 vol. %)	ca. 5–40	5–90 µm	SGR + dislocation glide	20 – 170 μm	1.4–2.5 mm
3	-	< 1	5–200 µm	SGR + dislocation glide	30–400 µm	0.6–2 cm

4 Methods

Analyses were performed at the Institute of Geological Sciences of University of Bern (Switzerland) unless specified otherwise. The petrographic characterization of the studied sample was done in a ca. 60 µm thick section on a Zeiss
140 Axioplan petrographic microscope. For backscattered electron (BSE), forescattered electron (FSE) and cathodoluminescence (CL) images, and for electron backscatter diffraction (EBSD), a Zeiss EVO50 scanning electron microscope (SEM) was used with a beam current of ca. 1 nA and accelerating voltage of 20 kV.

4.1 Grain-size analysis

Microstructural analysis was performed combining transmission light and scanning electron microscopy to determine average grain sizes of epidote and quartz, as well as their volume fractions in selected microstructural domains. Grain size

) Number: 1	Author: a Subject: Sticky Note	Date: 22/06/2022 10:24:04			
You definite	ely have to show these with higher magnifiq	ion pictures as insets of Fig.3. At the moment it is impossible to see what you are			
describing.		J			
I see that so	ome figures are coming in Fig. 5 and (🖓 t I	would place them much earlier. It is no point in describing them here and not showing			
them					
Mumber: 2	A hor: a Subject: Sticky Note	Date: 22/06/2022 10:12:55			
Ok, but def	Ok, but deform 🔐 ue to folds, shear bands? Please specify.				
👝 Number: 3	Author: a Soject: Sticky Note	Date: 22/06/2022 10:11:51			
This sounds	s interpretatione				
i 👝 Number: 4	Author: a Sub <mark>4</mark> ct: Sticky Note	Date: 22/06/2022 10:13:38			
Again, you must show data 🙀 support this! Add microphotos and ideally EBSD maps					

	Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 8:35:32 PM
1	Moved Fig. 4 here			
P	Number: 2	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 8:40:52 PM
	Done			
¢	Number: 3	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 8:41:00 PM
1	Moved to discussi	on		

 Number: 4
 Author: Veronica Peverelli
 Subject: Sticky Note
 Date: 8/5/2022 7:32:37 AM

 Removed from Table and added figures. However, we disagree that EBSD maps are needed to claim that subgrain rotation is the recrystallization mechanism active here (see Stipp et al., 2002)
 Date: 8/5/2022 7:32:37 AM





analysis was carried out in ImageJ (IJ 1.46r; Ferreira and Rasband, 2012) using sketches of the microstructures drawn on transmitted light, BSE and FSE images. In domains where the microstructure is not suitable for an automated grain size analysis by ImageJ (Figs. 4b and 7c), epidote and quartz grain sizes were directly measured on a transmitted light microscope. $\sqrt{1}$

150 4.2 Chemical maps

155

X-ray compositional maps in wave-length dispersive mode of Si, Fe, Al, Ca, Mn and Sr were obtained by electron probe micro-analyzer (EPMA) on a JEOL-8200 microprobe. Accelerating voltage was 15 keV, specimen current 100 nA, the step size 4 μ m, and dwell time was 190 msec. For calibration of the X-ray maps, spot analyses were acquired with a specimen current of 10 nA, and calibrated using the following standards: wollastonite (SiO₂), olivine (MgO), anorthite (CaO, Al₂O₃), magnetite (FeO), pyrolusite (MnO), tugtupite (Cl), rutile (TiO₂), and celestite (SrO). The processing of the X-ray

compositional maps, including map calibration, was done by using XMapTools (Lanari et al., 2014; 2019). Minerals were identified based on the concentrations of specific elements (i.e. Ca, Fe and Al for epidote; Si for quartz; K for micas) and classified. Maps of the structural formula of epidote were calculated on 12.5 oxygen basis.

4.3 Trace elements

- 160 For trace element measurements, a RESOlutionSE 193 nm excimer laser system (Applied Spectra, USA) equipped with an S-155 large-volume constant-geometry chamber (Laurin Technic, Australia) coupled with an Agilent 7900 ICP-QMS was employed. During ablation, a He atmosphere was used, and Ar was admixed to the carrier gas before reaching the plasma of the ICP-MS. NIST SRM612 was used for optimization of the analytical conditions, ensuring that the ThO production rate was < 0.2 % and the Th/U sensitivity ratio > 97 %. On-sample fluence was 5 J cm⁻² and repetition rate 5 Hz. The size of the
- 165 analysis spots ranged between 20–30 µm, and BSE images were used to plan the analyses to avoid locating any spots across zones with heterogeneous composition (e.g., chemical zoning) or on inclusions. External standardization was done with USGS GSD-1G standard, and the SRM612 standard was measured as an unknown for quality control in absence of a well characterized epidote standard. Bracketing standardization enabled a true-time linear drift correction. Data reduction was carried out with the software SILLS (Guillong et al., 2008), using the sum of total oxides minus H₂O (98.3 % for epidote and
- 170 100 % for SRM612) as internal standard (see Halter et al., 2002). The formulation of Pettke et al. (2012) was employed to calculate limits of detection for each element in every analysis.

4.4 U-Pb isotope data

Measurements of ²³⁸U/²⁰⁶Pb, ²⁰⁷Pb/²⁰⁶Pb and ⁸⁷Sr/⁸⁶Sr were made in epidote micro-separates following the procedure for sample digestion in acids and for column chemistry detailed by Peverelli et al. (2021; modified from Nägler and Kamber,

175 1996). Two epidote micro-separates – each mixing Epidote-A and Epidote-B (see Sect. 5.1) in unknown and different proportions – were prepared (Ep_A+B_1 and Ep_A+B_2). After hand-picking, the material was finely ground and washed

Number: 1 Author: a Subject: Sticky Note Date: 23/06/2022 09:24:25
I would like to know how ou define grains. If you define those based on EBSD maps and high angle boundaries >10°, then how do you asses this? Did you do it application of the provided of the pr

Number: 1 Author: Veronica Peverelli Subject: Sticky Note Specified in the methods section

Date: 8/5/2022 7:33:01 AM





with MilliQTM water. Two aliquots of each (Ep A+B 1a and Ep A+B 1b, and Ep A+B 2a and Ep A+B 2b) were weighed in as replicates for each micro-separate ensuring ca. 300 ng of Pb in each aliquot. During column chemistry with a Sr-specTM resin (Horwitz et al., 1992), the Sr and Pb fractions were collected in sequence (Haeusler et al., 2016). One large Epidote-A grain was also handpicked and ground, and an amount of the powder corresponding to 250 ng of Sr was digested 180 in acids before the extraction of the Sr fraction by column chemistry. The Pb fraction of this Epidote-A grain was not collected since Pb isotopic data are available from in-situ U-Pb isotope measurements by LA-ICP-MS (Peverelli et al., 2021). Procedural blank samples were used to assess contamination during work in the laboratories. Measurements of ²³⁸U and ²⁰⁶Pb concentrations for calculation of ²³⁸U/²⁰⁶Pb ratios were run at the Department of Geography of University of Bern 185 on a 7700x Agilent quadrupole ICP-MS. For these analyses, two aliquots of each digested epidote microseparate were separated and diluted to different volumes. Three aliquots were separated from the digested AGV-2 standard and diluted in 0.05, 0.25 and 0.7 ml HNO₃. The intensities in counts per second (cps) of ²³⁸U and ²⁰⁶Pb measured in the three aliquots of the AGV-2 standard were plotted against the real concentrations of these isotopes in ppb calculated for each aliquot using the reference data of Weis et al. (2006). The calibration curves thus obtained were used to calculate the real concentrations of 238 U and 206 Pb in the digested epidote material. The calculation of 2 standard errors (2 S.E.) on 238 U/ 206 Pb ratios – which are 190 not returned automatically by the measurement software -is detailed in Peverelli et al. (2021). Pb isotope ratios were measured on a Thermo Fisher Neptune Plus MC-ICP-MS in desolvated plasma mode equipped with a CETAC Aridus 2 desolvating system. Instrumental mass fractionation was corrected within-run by means of a Tl spike. External reproducibility of the measurements was quantified by measuring the NIST NBS 981 standard. The measured Pb isotope

ratios were identical to those obtained by Rehkämper and Mezger (2000; their Table 4).

4.5 Strontium isotope data

Strontium isotope ratios were measured on a ThermoFisher TritonTM thermal ionization mass spectrometer (TIMS) after loading 250 ng Sr diluted in 6.4 M HCl on Re filaments using 1.5 μ l Ta-oxide activator. The SRM 987 standard (200 ppm; Weis et al., 2006) was measured for quality control. The detected masses were 84, 85, 86 (center cup), 87 and 88. The interference of ⁸⁷Rb and within-run mass fractionation were corrected for by using the IUPAC ⁸⁷Rb/⁸⁵Rb and ⁸⁸Sr/⁸⁶Sr values of, respectively, 0.385617 and 8.735209. The SRM 987 standard returned a weighted average ⁸⁷Sr/⁸⁶Sr ratio of 0.710279 ± 0.000020 (2 standard deviation, S.D.; number of replicates, n = 12), which is higher than the reference preferred value of 0.710248 (see Weis et al., 2006). Standards AGV-2 (n = 1) and GSP-2 (n = 2) of USGS were measured as unknowns for quality control and the returned, respectively, ⁸⁷Sr/⁸⁶Sr ratios of 0.704041 ± 0.000018 (2 standard error, 2 S.E.), and 0.765396 ± 0.000010 and 0.765202 ± 0.000008. These values are also higher than the reference preferred values of 0.703981 ± 0.00009 (2 S.D.) and 0.765144 ± 0.000075 (2 S.D.) for AGV-2 and GSP-2, respectively (see Weis et al., 2006). A correction based on the reference materials returning higher ⁸⁷Sr/⁸⁶Sr values than their reference values would produce the same shift in all measured ⁸⁷Sr/⁸⁶Sr ratios. Hence, we did not correct our data because only the variability among the samples is relevant in this study, while the interpretation of the absolute Sr isotope ratios is beyond the scope of this work. Rubidium



400 µm



- 210 concentrations were not measured, as a correction for ⁸⁷Rb-derived ⁸⁷Sr is not necessary. This is valid а approach because the incompatibility of Rb in the 215
 - epidote crystal structure results in negligible Rb concentrations (see Frei et al., 2004; Feineman et al., 2007).

220 5 Results

230

5.1 Microstructural analysis

Figure 4: Transmitted light scans of the microstructured domains described in Sect. 5.1

Figure 4: Transmitted-light scans of the microstructural domains described in Sect. 5.1. Microstructures (a-c) are used for microstructural analysis (Fig. 7; Sect. 5.1.2). The blue rectangle in (a) indicates the location of the cathodoluminescence image of Fig. 11. Bt = biotite; Ep = epidote; Fsp = feldspar; Qz = quartz. Plane-polarized light.

The different characteristics of epidote and its microstructures in layers 1–3 (Figs. 3–4) allow the distinction of epidote into Epidote-A and Epidote-B as illustrated below. All microstructural domains indicate that vein opening with crystallization of epidote, quartz and minor biotite was followed by vein deformatio 2^2

225 5.1.1 Layer 1: veining and Epidote-A

This epidote layer is characterized by coarse (ca. 0.2-1.6 mm) epidote grains associated with smaller angular ones (ca. $20-200 \ \mu$ m). The coarser epidote grains (Figs. 4d and 5) form clusters with random shape orientations or are found as isolated crystals. Larger epidote grains are often surrounded by the smaller angular epidote grains (Fig. 5; red arrow) as a result of brittle grain-size reduction upon deformation with brittle deformation behavior of epidote. Epidote is euhedral to anhedral, and smaller epidote crystals are found in the gaps among the larger ones as well (see Fig. 2b of Peverelli et

al., 2021). Large quartz relicts are present displaying undulose extinction.

235 More commonly, quartz is dynamically recrystallized, forming core-mantle structures with subgrain rotation around quartz relicts refer to epidote in layer 1 as Epidote-A.



Figure 5: Transmitted light microscope photograph of one Epidote-A grain in layer 1 surrounded by small angular grains (red arrow). Bt = biotite; Ep = epidote; Qz = quartz. Plane-polarized light. 2^{4}

Number:	1 Author: a Subject: Sticky Note	1 Date: 22/06/2022 10:20:49	
Those ar	e definitely not plane polarised light polos	s, as Qtz has a birefringence.	
Number:	2 Author: a Suzject: Sticky Note	Date: 22/06/2022 10:22:19	
Again, w	hat evidence sug	ssing description	
👝 Number:	3 Authar: a Subject: Sticky Note	Date: 22/06/2022 12:13:22	
Again, yo	ou should say withis with a figure		
Number:	4 Author: a Subject: Sticky 4pte	Date: 22/06/2022 12:17:51	
Again, al	so here quartz has a slight birg the gence		

Numbe	: 1 Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 8:49:20 PM	
The sec	tion is ca. 60 um thick. Specified in t	he text, sorry.		
Numbe	: 2 Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/24/2022 9:52:13 PM	
Remov	ed early intepretation and moved it t	o discussion		
Numbe	: 3 Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/25/2022 11:43:30 AM	
Added	panels to Fig. 5			
Numbe	- 4 Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:33:55 AM	

Number: 4 Author: Veronica Peverelli Subject: Sticky Note Date: 8/5/2022 7:33:55 AM Again, this is due to the thickness of the thin section. Added in all captions, sorry for not mentioning it in the first place





5.1.2 Layers 2–3: microfold and Epidote-B

- In layers 2–3, the spatial distribution and the variable modal abundance of epidote define a fold (Figs. 3, 4b, 6 and A1).
 Smaller epidote grains with a shape-preferred orientation and quartz define axial planes and limbs (rectangle a in Fig. 3; Figs. 4a and 6). Epidote grain boundaries are mostly
- 245 curved and irregular, but a few euhedral epidote grains are also observed (Fig. 6). Epidote grains vary between 5–90 μm in size, and the modal abundance of epidote varies between ca. 5 to ca. 40 vol. % within the layer (Table 1), defining a quartz-supported microstructure. Where epidote
- 250 is most abundant, quartz and epidote define a "spong microstructure (Figs. 4b and 6). Small (ca. 1–20 μ m in size) oval epidote grains and fluid inclusions are observed within



Figure 6: Transmitted light microphotograph of layer 2 where oval epidote grains are hosted by dynamically recrystallized quartz with their long axes oriented consistently with the microfold axial planes. Bt = biotite; Ep = epidote; Qz = quartz. Overlapped plane-polarized and cross-polarized light images.

quartz grains (Figs. 4 (Fig. 7; Zener relation; Herwegh et al., 2010). Dynamic recrystallization of quartz by subgrain rotation prevails creating a crystallographic preferred orientation (FO), and quartz relicts with undulose extinction are miles Fig. 4a-c). However, a ca. 2 cm long relict quartz grain, whose orientation acquired during crystallization may have favored dislocation glide, marks the transition from layer 2 to layer 3 (Fig. A1, black arrow). Where minor epidote is present, anhedral epidote



Figure 7: Correlation between quartz grain size (D) and Zener paramete $(D_{\text{P}})^{\text{B}}$ of epidote as the second phase determined in the FS and BSE images or the microstructures in the microfold of layers 2–3. The green lines in the central BSE image contour epidote grains. Ep = epidote; Qz = quartz. Panels a, b and c correspond, respectively, to panels b, a and c of Fig. 4.

grains of few to ca. 10 µm in size are mostly interstitial and found at triple junctions among quartz subgrains (Fig. 9). We refer to subhedral to anhedral epidote grains defining the microfold in layers 2–3 as Epidote-B. While U–Pt ages by LA-ICP-MS of Epidote-A (Sect. 3) are available, no U-Pb dating is possible in Epidote-B because the small grain size of epidote entails contamination from epidote-epidote or epidote-quartz grain boundaries upon measurements.

回 Number: 1	Author: a Sadaja cl. Sticky Note	Date: 22/06/2022 12:39:32
Mark Ep-A and B	directly on m 🚰 photos	
回 Number: 2	Author: a Sul2ect: Sticky Note	Date: 22/06/2022 12:21:59
Meaning what exa	actly? Plense, refer to term that are used	by the community giving references for it
<u> Number:</u> 3	Author: a 3 bject: Sticky Note	Date: 22/06/2022 12:19:27
What do you mea	n with everlapped? I see birefringence of	only in quartz, like if you have polarizer and analizer at low angle (10°?)
pNumber: 4	Author: a Su4ect: Sticky Note	Date: 22/06/2022 12:22:58
Impossible to see	at this n 🗃 ification	
<u>()</u> Number: 5	thor: a Subject: Sticky Note	Date: 22/06/2022 12:56:58
Before jumping te	the deformation mechanism you have	to describe it. Then you interpret it, much better in the discussion section
what shp systemp	ould create this CPO?	
-Number: 6	Cthor: a Subject: Sticky Note	Date: 22/06/2022 12:33:05
Impossible to	t. Please add XPL photos, maybe also v	vith lambda plate inserted
- Numbor: 7	Aut 7 r: a Subject: Sticky Note	Data: 22/06/2022 12:57:52
This sentence see	eally out of place here	Date. 22/00/2022 12.37.32
	Authory a Subjects Sticky Note	Data: 22/06/2022 12:25:16
You shaw explai	n what Dp and fp are	Date. 22/00/2022 12:25:10

	Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:07:09 PM
	Done		· ·	
P	Number: 2	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:14:20 PM
	Rephrased			
Ģ	Number: 3	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:16:05 PM
	Epidote birefringe the thin section (ca	nce is poorly visible party be a. 60 um)	cause of the overlap betwee	en the two photographs and partly because of the thickness of
	Number: 4	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:22:53 PM
	Added panel in Fig	g. 5.6		
P	Number: 5	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/25/2022 11:45:38 AM
1	The identification	of machanisms of dynamic r	ecrystallization are thorough	ly described in Stipp et al. (2002), which we follow. We do not
	believe that indica	ting the dynamic recrystalliz	ation here is too early interp	retation. The slip system is addressed against the updated EBSD
	images.			
P	Number: 6	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:35:26 AM
	Added arrow in Fig	g. 5.4 and lessened the focus	s on these grain since they a	re not that relevant.
P	Number: 7	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:32:34 PM
	Agreed. Moved ea	rlier in the text		
P	Number: 8	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:26:25 PM
	Done			







Figure 8: Orientation of quartz C axis in layer 3 relative to the long side of the thin section; measured by electron backscatter diffraction.



Figure 9: Backscattered electron images showing epidote (Ep) grains along quartz (Qz) grain boundaries and at triple junctions among quartz grains.

5.2 Major and minor elements

The compositional map in Fig. 10 (location shown in Fig. 3)
covers layers 1–3. The compositions of FeO and Mn range between ca. 12.5–14 wt. % and ca. 2500–5000 μg g⁻¹, respectively, across all analyzed epidote grains. Larger Epidote-A grains in layer 1 are zoned, with FeO and Mn concentrations increasing from core to rim. Epidote in layers 2–3, on the other end, is characterized by uniform concentrations of FeO and Mn. It should be noted that, because the step size of the compositional maps is 4 μm, the chemical variability of epidote in layers 2–3 is better assessed among different grains across the overall microstructure rather than within each crystal (i.e., the majority of

285 grains do not contain enough $4 \times 4 \mu m$ pixels).

5.2 Trace elements

290

The minimum spot size used for measurements by LA-ICP-MS is 20 μ m, which is ca. four times larger than the smallest epidote grains. Therefore, chemical variability is assessed throughout each microstructural layer by relying on measurements in large-enough epidote grains to avoid contamination from grain boundaries.



Figure 10: X-ray compositional maps across layers 1–3 (Fig. 3; Sect. 5.1). White circles indicate spots for U–Pb dating by LA-ICP-MS of Peverelli et al. (2021). The pink line separates layer 1 from layers 2–3.

Number: 1 Author: a Subject: Please, add where X and Z are lo zet, number of grains, contouring parametres and at least <a> pole figure

Page: 1	9
---------	---

回 Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/26/2022 7:49:30 AM	
Done				





295

Epidote-A is also addressed here with no specific reference to intra-grain zoning but only across the overall Epidote-A microstructure (i.e., layer 1). Data of Epidote-B are collected in anhedral epidote grains in layer 2. The different extents of chemical variability between Epidote-A and Epidote-B noted in the compositional maps is reflected by trace element data (Table 2).



Figure 11: Epidote LA-ICP-MS data of (a) Pb and Sr, (b) CI-chondrite normalized (McDonough and Sun, 1995) rare earth element patterns (REE), and (c) trace elements normalized to the Central Aar Granite (ZAR; Schaltegger & Krähenbühl, 1990).

Table 2: Trace element composition of epidote in µg g⁻¹measured by LA-ICP-MS.

	Epidote-A																				
Spot	Ba	Sr	Pb	U	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Y	Но	Er	Tm	Yb	Lu	Zr	v
1	0.99	1400	130	79	5.6	14	2.2	12	6.3	3.9	14	2.8	22	200	5.5	19	2.7	19	2.7	3.6	140
2	1.2	1990	130	130	7.7	18	2.7	15	8.5	5.6	18	3.6	27	230	6.4	22	3.1	21	3.4	5.1	140
3	22	3290	150	350	4.8	8.8	1.2	5.6	2.5	2.0	5.7	1.2	12	120	3.6	14	2.5	21	3.8	22	110
4	21	2500	170	270	0.99	1.9	0.24	1.1	0.66	1.1	2.8	0.86	9.8	110	3.1	14	2.7	23	4.4	24	110
5	1.9	970	97	60	5.2	9.3	1.1	5.0	1.8	1.3	2.9	0.62	5.1	32	1.0	3.5	0.53	3.7	0.57	4.2	160
6	23	2590	170	130	10	15	1.6	7.0	2.8	1.9	5.6	1.3	9.9	71	2.3	9.2	1.6	15	3.1	51	130
7	0.98	740	87	140	5.0	9.3	1.2	5.3	2.7	1.6	5.5	1.1	8.4	53	1.8	6.0	0.88	5.8	0.77	4.5	200
8	21	2870	170	240	12	17	1.8	8.3	4.6	3.7	11	2.5	19	130	4.4	17	2.8	25	4.4	17	150
9	9.7	2000	150	100	8.5	12	1.4	5.9	3.2	2.7	8.0	1.8	14	82	2.9	8.7	1.1	7.8	1.3	9.6	140
10	24	2890	170	220	6.6	9.7	1.1	4.6	2.0	2.2	5.9	1.4	12	90	2.9	11	2.1	18	3.2	31	140
11	0.92	1660	93	100	3.4	7.2	0.90	3.3	1.2	1.4	2.7	0.59	5.1	41	1.3	4.1	0.42	2.5	0.28	2.2	110
12	34	3750	190	140	5.3	8.5	1.0	4.6	1.9	2.3	4.2	1.0	8.8	67	2.2	6.7	0.80	4.8	0.64	25	120
13	0.59	770	78	160	1.6	3.0	0.39	1.6	1.0	1.5	3.2	1.0	9.1	65	2.2	7.9	0.97	5.9	0.72	3.2	180
14	27	2820	180	260	1.3	2.5	0.29	1.2	1.2	1.5	4.6	1.3	12	120	3.5	14	2.8	28	4.6	42	130
15	5.8	1530	93	54	0.62	0.97	0.12	0.62	0.45	0.94	1.4	0.40	4.1	42	1.2	4.7	0.77	6.2	1.0	5.0	140
16	17	2810	150	140	1.9	3.0	0.35	1.4	0.55	1.3	1.7	0.66	7.8	77	2.3	9.3	1.6	14	2.3	18	130
17	5.2	1310	93	120	3.5	6.3	0.78	3.7	1.3	1.4	2.0	0.57	4.9	47	1.3	6.1	0.88	7.1	0.93	5.2	120
18	3.8	2340	110	110	1.5	2.9	0.43	1.6	0.76	1.2	1.6	0.47	4.9	47	5.5	5.5	0.78	5.8	0.90	3.1	120
										Epi	dote-B										
Spot	Ba	Sr	Pb	U	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Y	Но	Er	Tm	Yb	Lu	Zr	v
1	2.1	1640	171	130	4.7	10	1.4	7.1	3.4	3.0	7.3	1.7	16	150	4.4	14	2.1	14	2.3	5.2	130
2	2.9	1860	150	140	3.7	8.1	1.1	5.7	3.1	3.1	7.6	2.1	17	170	4.6	16	2.4	19	3.0	5.9	140
3	6.6	1950	120	190	1.7	4.2	0.55	2.9	1.2	1.6	3.1	0.81	8.1	85	2.1	8.6	1.5	9.7	1.7	8.2	120
4	5.0	1720	140	190	3.3	7.6	1.1	5.2	2.4	2.0	4.8	1.1	10	110	3.1	11	1.8	13	2.1	7.0	120
5	4.2	1700	130	170	3.0	7.4	0.97	5.0	2.3	2.1	4.2	1.2	11	120	3.0	11	2.0	14	2.5	7.7	120
6	5.1	1930	120	160	1.9	4.4	0.60	2.9	1.4	1.5	3.3	0.80	7.4	85	2.3	8.8	1.3	10	1.6	7.2	130
7	3.7	1530	130	160	2.9	6.6	0.88	4.5	2.1	1.7	4.1	0.73	7.9	84	2.3	8.3	1.3	9.0	1.6	5.2	130
8	6.5	1960	140	190	3.3	7.6	0.99	4.5	2.3	1.9	3.9	0.99	9.1	100	2.6	10	1.5	13	2.1	9.3	130
9	5.3	1850	130	190	3.4	8.2	1.1	4.7	2.2	1.9	3.7	0.91	8.4	100	2.7	9.9	1.5	13	2.1	8.6	130
10	5.4	1800	130	190	2.3	5.2	0.71	3.5	1.7	1.7	3.6	0.85	8.5	90	2.4	10	1.7	12	2.2	8.1	130
11	5.5	1800	150	180	3.9	8.5	1.2	5.6	2.5	1.9	4.9	0.99	9.8	100	2.7	10	1.6	12	1.9	8.0	130
12	4.8	1710	151	230	9.4	20	2.6	12	5.3	3.5	9.6	2.3	19	160	4.8	16	2.4	16	2.6	7.6	130
								Central	Aar Gran	ite (Schal	tegger ar	d Krähen	bühl, 199	D)							
KAW	Ва	Sr	Pb	U	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Y	Но	Er	Tm	Yb	Lu	Zr	v
2219	430	75	24	8	26	61	-	-	5.1	0.51	-	2.4	-	56	-	-	-	8.6	12	192	18





300

The concentrations of Sr and Pb in Epidote-B overlap with the trend defined by the same elements measured in

- Epidote-A, but they cover a more limited range of
- 305 values (Fig. 11a). The CI chondrite-normalized rare earth elements (REE)



Figure 12: Overlapped cathodoluminescence CL) and backscattered electron images. The only minerals in the image are epidote (Ep) and quartz (unlabeled grains). The different CL contrasts in quartz are due to variable trace element contents.

patterns (Fig. 11b) of Epidote-A and Epidote-B have similar trends, characterized by positive slopes (La_N/Yb_N of 0.1–0.4 in Epidote-A and 0.03–0.5 in Epidote-B) and variably positive Eu anomalies (1.5–2.4 in Epidote-A and 1.2–2.3 in Epidote-B).

310 The REE trends of Epidote-B fall within the range of Epidote-A and confirm the lesser extent of chemical variability of Epidote-B relative to Epidote-A. Selected elements are plotted as values normalized to the concentrations in the Central Aar Granite in Fig. 11d reinforce chemical affinity between Epidote-A and Epidote-B, as well as the lesser chemical variability of the latter relative to the former. Cathodoluminescence images of the recrystallized quartz grains (Fig. 12) qualitatively indicate trace element variability also in quartz grains.

315 5.3 Isotope data

320

Table 3: U-Pb and Sr isotopic data by solution ICP-MS and TIMS. Uncertainties are 2 standard errors (2 S.E.).

	²³⁸ U/ ²⁰⁶ Pb	2 S.E.	²⁰⁷ Pb/ ²⁰⁶ Pb	2 S.E.	⁸⁷ Sr/ ⁸⁶ Sr	2 S.E.
Ep_A+B_1a	6.49	0.11	0.79159	0.00001	0.727803	0.000011
Ep_A+B_1b	6.54	0.07	0.79108	0.00001	0.727807	0.000007
Ep_A+B_2a	6.44	0.16	0.79428	0.00001	0.726952	0.000010
Ep_A+B_2b	6.44	0.10	0.79391	0.00001	0.726830	0.000015
Epidote-A	4.421	0.091	0.78671	0.00581	0.726552	0.000007

¹ Datum-point #10 of Peverelli et al. (2021) by LA-ICP-MS.

The total ²³⁸U/²⁰⁶Pb and ²⁰⁷Pb/²⁰⁶Pb ratios measured by solution ICP-MS in the epidote microseparates (Table 3) are plotted in a Tera–Wasserburg diagram (Fig. 13a) together with the LA-ICP-MS data measured in Epidote-A. The ²³⁸U/²⁰⁶Pb ratios of the dry aliquots of each microseparate are within uncertainty of each other, whereas their ²⁰⁷Pb/²⁰⁶Pb ratios are slightly different. This is attributed to geological heterogeneity between the single aliquots of each microseparate: perfect



homogenization of the microseparates is not possible because total 325 ²⁰⁷Pb/²⁰⁶Pb ratios include uranogenic lead in addition initial Pb, and are to therefore influenced by the variability in U

concentrations.

ICP-MS

solution

330



^{yy the} Figure 13: U-Pb and Sr isotope data. (a) Tera-Wasserburg diagram with the LA-ICP-MS data measured in Epidote-A (Ep-A) by Peverelli et al. (2021) and those measured in epidote U microseparates (Ep-A+B) by solution ICP-MS; LA-ICP-MS error ellipses are 2σ. (b) Total ²⁰⁷Pb/²⁰⁶Pb and initial ⁸⁷Sr/⁸⁶Sr data of Epidote-A and the epidote microseparates; the ²⁰⁷Pb/²⁰⁶Pb ratio of Epidote-A is by LA-ICP-MS (data-point #10 of Peverelli et al., 2021); all other ²⁰⁷Pb/²⁰⁶Pb ratios are by solution ICP-MS; ⁸⁷Sr/⁸⁶Sr ratios are by TIMS; error bars are smaller than the symbols where not shown.

points plot above the regression through the LA-ICP-MS data in the diagram, and the isotopic ratios measured by the different techniques cannot be combined in one single regression.

Strontium (Table 3) and Pb isotopic data of the epidote microseparates and of pure epidote (Epidote-A) are plotted in Fig.

335 13b. To compare the Pb and Sr isotopic characteristics of Epidote-A with those of the microseparates, data obtained from the different techniques have to be combined in a ²⁰⁷Pb/²⁰⁶Pb versus ⁸⁷Sr/⁸⁶Sr plot. While Sr in epidote is entirely non-radiogenic, the ²⁰⁷Pb/²⁰⁶Pb ratios measured by solution ICP-MS contain both initial and radiogenic (i.e., U-derived) Pb. Therefore, to discuss Pb isotopic characteristics of Epidote-A (i.e., analyzed by LA-ICP-MS) against those of the epidote microseparates mixing Epidote-A and Epidote-B (i.e., analyzed by solution ICP-MS), we cannot simply use the initial ²⁰⁷Pb/²⁰⁶Pb ratio obtained from a Tera–Wasserburg plot because this value excludes the ingrown radiogenic Pb component of Epidote-A. Therefore, in a ²⁰⁷Pb/²⁰⁶Pb versus ⁸⁷Sr/⁸⁶Sr plot (Fig. 13b), the total ²⁰⁷Pb/²⁰⁶Pb ratio of Epidote-A is represented by analysis #10 of Peverelli et al. (2021; see their Table 5). This is the datum-point closest to the average (²³⁸U/²⁰⁶Pb = 4.62

and ${}^{207}Pb/{}^{206}Pb = 0.786$) and to the median (${}^{238}U/{}^{206}Pb = 4.35$ and ${}^{207}Pb/{}^{206}Pb = 0.787$) values of the U–Pb isotopic ratios measured in Epidote-A by LA-ICP-MS.







345 6 Discussion of the formation mechanisms for the epidote-quartz microfold

6.1 Interplay of epidote dissolution-precipitation and quartz dynamic recrystallization

The formation of a hydrothermal vein entails the crystallization of a mineral assemblage from a mineralizing fluid

- that fills a fracture (Bons et al., 2012).Although the original morphology of the studied epidote-quartz vein is obliterated by deformation, the euhedral shapes of Epidote-A grains
- 355 (Fig. 4d) suggest that the formation of Epidote-A in layer 1 is related to vein-filling mineralization, with crystallization occurring in equilibrium with a fluid. While the
- formation of Epidote-A is ascribed to veining, in the following we address the mechanisms affecting Epidote-B.
 The microstructures in layers 2–3 differing from those in layer 1 (Sect.
- 5.1), along with the lesser extent of chemical variability across epidote grains in layers 2–3 compared to Epidote-A in layer 1 (Figs. 10–11), suggest that the mechanism of



Figure 14: Sketch of the sequence of events affecting the studied epidote vein. (a) Original microstructure formed upon veining. (b) Detail of the fracturing occurring in Epidote-A in layer 1. (c) Dynamic recrystallization of quartz by subgrain rotation. (d) Folding of the epidote band in layer 2 and dissolution-precipitation of Epidote-B. (e) Detail of the viscous granular flow process and dynamic granular fluid pump allowing Epidote-B to precipitate in creep cavities along quartz grain boundaries. (f) Detail of quartz grain boundary sliding allowing dynamic granular fluid pump (modified from Fusseis et al., 2009). (g) Present-day microstructure. Not to scale.





- 370 formation of Epidote-B is different than that of Epidote-A. The alignment of Epidote-B grains along the microfold axial planes (Figs. 4a and 6), and the presence of minute Epidote-B along dynamically recrystallized quartz grains (Fig. 9) indicate that Epidote-B formed during deformation. The overlap in major and trace element compositions of Epidote-B and Epidote-A (Figs. 10–11), though, demonstrates that these epidote generations are chemically related. One way to reconcile this geochemical affinity with different formation mechanisms is a scenario in which Epidote-B formed via deformation-induced 375 dissolution of Epidote-A grains and (re)precipitation. In fact, a few euhedral epidote grains can still be recognized in layer 2, which are interpreted as Epidote-A relicts inherited from the original vein morphology. The gradation from epidote-
- rich/quartz-poor domains to virtually epidote-free/quartz dominated ones in layers 2–3 (Fig. 14g) suggests that the microstructure that formed upon veining was similar to that shown in Fig. 14a. In fact, shifts between epidote-rich and quartz-rich domains are frequent in epidote-quartz veins (e.g., Peverelli et al., 2021; their Figs. 1a and 1c). Further evidence
- for fluid-mediated mass transfer is also given by CL images of quartz (Fig. 12), which indicate that the grain boundaries of quartz interacted with an aqueous flut the presence of a fluid during deformation is also supported by fluid inclusions in dynamically recrystallized quartz.
- The transport of dissolved epidote-forming material towards the microfold axial planes (Fig. 14d) calls for a mechanism allowing the fluid to move throughout the deforming microstructural domain. The presence of Epidote-B grains along quartz grain boundaries and at triple junctions (Fig. 9) implies that dissolved epidote-forming material is transported and precipitated into nucleation loci. One mechanism enabling this mass transfer process is dynamic granular fluid pump (Fig. 14e-f; see Fusseis et al., 201 mong dynamically recrystallized quartz. A first deformation step in which quartz grain size is reduced by dynamic recrystallization via subgrain rotation (Fig. 14c) is suggested by a strong crystallographic preferred orientation in quartz-dominated domains (Fig. fig. fig. for the new quartz grains are formed, grain boundary sliding produces creep cavities (Fig. 14f; see Herwegh and Jenni, 2001; Fusseis et al., 2009; Gilgannon et al., 2017; 2021), thus creating nucleation loci for Epidote-B. A similar process is described by Gottardi and Hughes (2022) in quartzites deformed at the brittle-ductile transition in the crust, in which fluid inclusions are redistributed in the quartz matrix by dynamic recrystallization of quartz by subgrain rotation and grain-boundary migration. A similar microstructure and a similar interplay of processes are also discussed by Kruse and Stünitz (1999) for in mafic high-temperature mylonites. This first step

Number: 1	Authon 1 Subject: Sticky Note	Date: 22/06/2022 14:34:27
Based on what evi	de <mark>gree</mark> ' Please, state it here	
Number: 2	2µthor: a Subject: Sticky Note	Date: 22/06/2022 14:34:49
better remem <mark>per</mark>	he reader the size	
Number: 3	Author: a Subject: Sticky Note	Date: 22/06/2022 14:36:16
Another relevant r	eference is Menegon et al., 2015 Geolog	ЭУ
Number: 4 4	Author: a Subject: Sticky Note	Date: 22/06/2022 14:37:12
Again, ple <mark>use</mark> expa	nd this based on possible slip system a	cting and relevant references

Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:37:51 PM	
Rephrased and a	dded reference			
pNumber: 2	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:39:07 PM	
Done				
😑 Number: 3	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/21/2022 12:48:55 PM	
Added reference				
画 Number: 4	Author: Veronica Peverelli	Subject: Sticky Note	Date: 8/5/2022 7:36:00 AM	
Done				





is followed by one in which the grain size and volume abundance of the newly precipitated Epidote-B grains control the grain size of quartz (Fig. 7), hence exerting control on quartz deformation mechanisms by pinning the migrating quartz boundaries (see Olgaard, 1990; Herwegh and Berger, 2004; Herwegh et al., 20, 7, This allows to suggest a feedback process: (1) quartz grain boundary sliding creates creep cavities in which Epidote-B crystallizes, (2) Epidote-B keeps quartz grain size small, (3) more creep cavities are formed, and (4) more Epidote-B grains are formed. This process is referred to as viscous granular flow (e.g., Fitz Gerald and Stünitz, 1993;Stünitz and Fitz Gerald, 1993; Paterson, 1995; see also Kruse and Stünitz, 1999).

6.2 Open-system conditions and external fluids

Epidote dissolution-precipitation and the transport of dissolved epidote material to the loci of Epidote-B crystallization implies the presence of a fluid, whose nature can be assessed by Pb-Sr isotope data. We have mentioned that the epidote microseparates used for solution ICP-MS measurements mix Epidote-A and Epidote-B to unknown proportions, since the 405 mechanical separation of pure Epidote-B is not feasible due to its small grain size. The observation that ²³⁸U/²⁰⁶Pb and ²⁰⁷Pb/²⁰⁶Pb ratios of the microseparates do not lie on the same regression as those measured by LA-ICP-MS in Epidote-A (Fig. 13a) suggests that the epidote microseparates include different epidote generation, thus that Epidote-A and Epidote-B crystallized from fluids with different Pb isotope compositions. This is supported by the ⁸⁷Sr/⁸⁶Sr ratios of the microseparates being different than that measured in an Epidote-A grain (Fig. 13b; Table 2). There are two principal processes that can 410 induce changes in the Pb and Sr isotope compositions: (1) radioactive decay of ^{235,238}U into ^{207,206}Pb and ⁸⁷Rb into ⁸⁷Sr, or (2) open-system conditions allowing for advection of extraneous Pb and Sr with different isotopic compositions. Process (1) is unlikely in this case because ingrown radiogenic Pb and Sr of such young epidote are negligible. Consequently, the U-Pb-Sr isotope heterogeneity of the two epidote generations infers fluid-mediated addition of extraneous Sr and Pb during crystallization of Epidote-B. Because the vein microstructure carries evidence of a single deformation event (Fig. 14g), it is 415 likely that a second fluid type entered the system during vein deformation and formation of Epidote-B (Fig. 14e-f). Thus, the differences in U-Pb and Sr isotopic data between LA-ICP-MS and solution ICP-MS/TIMS data are interpreted to document mixing between different contributions: (1) one reflecting the isotopic composition of Epidote-A incorporated upon vein formation, and (2) one reflecting the mixing of Epidote-A material with extraneous Pb and Sr brought to the site of Epidote-

Number: 1 Authon Liet: Sticky Note Date: 22/06/2022 14:52:05 Please, add also other re references from other research groups

Page: 27

画 Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:42:23 PM	
Done				





B crystallization. The existence of open-system conditions bears the potential for fluid-mediated transfer of chemical 420 constituents (Figs. 14e-f). The fact that the chemical composition of Epidote-B is identical to, and less variable than, that of Epidote-A (Figs. 10–11) suggests rock-buffered conditions during fluid mediated crystallization of Epidote-A and Epidote-B at closely comparable physico-chemical (e.g., P, T, fO₂) conditions and thus fluid composition. The initial compositional variability of Epidote-A got thereby homogenized upon dissolution and crystallization into Epidote-B. Considering that the 425 299 ± 2 Ma old host granitoid (Schaltegger and Corfu, 1992) contains minerals with high concentrations of U, Th and Rb (e.g., allanite, biotite), even slightly variable contributions of radiogenic Pb and Sr leached from these minerals may produce a measurable shift in Pb-Sr isotope ratios without a resolvable effect on Sr and Pb fluid concentrations. Also, crystallization of Epidote-A and Epidote-B most likely occurred at comparable T conditions (ca. 400-450 °C as inferred from the presence of biotite in the vein; see Goncalves et al., 2012). Hence, fluid/epidote partition coefficients of all elements remained similar 430 between the two epidote crystallization events. Such a scenario can well account for the compositional uniformity but Pb and Sr isotopic heterogeneity between Epidote-A and Epidote-B, hence allowing that at least a fraction of the fluid involved in the deformation of layers 2-3 is of external origin. Peverelli et al. (accepted) measured the H isotope composition of epidote in this sample, obtaining values that can only be explained as a mixture of end-member waters (e.g., meteoric, seawater, etc.) as source for the epidote-forming fluids. In light of the present data, it is possible that their measured δD value reflects the mixing of internal and external fluids upon deformation of the epidote-quartz vein as described above.

435

6.3 Epidote geochemistry as a result of veining vs. granular fluid pump

The present major and trace element data show that Epidote-B is geochemically similar to, and less variable than, Epidote-A (Figs. 10–11), and this result bears important implications on epidote crystallization mechanisms. In a rock-dominated system, the larger geochemical variability of Epidote-A may be due to fluid distillation during epidote crystallization, slight

440

variations in physico-chemical conditions (e.g., P, T) throughout the veining processes, or pulsating fluid fluxes (i.e., multiple Epidote-A generations). The latter is unlikely because all laser spots for U-Pb dating define one statistically robust regression with MSWD of 0.79, providing no evidence for multiple epidote generations at the current analytical precision. Such a hypothesis is also inconsistent with trace element data (Fig. 11a-b) defining trends and not distinct epidote populations. These observations suggest that the geochemical variability of Epidote-A is due to an evolving epidote-forming





fluid chemistry along with more and more epidote precipitating, thus readily removing compatible elements from the fluid (see also Anenburg et al., 2015). The lesser extent of chemical variation in Epidote-B indicates homogenization of the whole trace element budget dissolved into the Epidote-B-forming fluid. Viscous granular flow is a dynamic process in which continuous feedback exists among dissolution of minerals, grain boundary sliding, creep cavitation, mass transfer and mineral precipitation (e.g., Fitz Gerald and Stünitz, 1993; Stünitz and Fitz Gerald, 1993; Paterson, 1995). This implies that repeated dissolution and reprecipitation of the same material is likely to occur, and the chemical budget gets compositionally homogenized with ongoing deformation. Consequently, in the case of the epidote-quartz microfold in layers 2–3, such a mechanism may account for the chemical homogeneity of Epidote-B. If this respect, granular fluid pump promotes recycling and homogenization of fluids in deforming polymineralic aggregates.

7 Consequences for epidote U-Pb ages

455 The inevitable question arising from the scenario developed above is what is the significance of the U–Pb age measured in Epidote-A. Temperature-driven resetting of the U-Pb system is excluded based on the peak temperature reached in the area (i.e. 450 ± 30 °C; Challandes et al., 2008; Goncalves et al., 2012) never exceeding the closure temperature for Pb diffusion in epidote (i.e. > 685 °C; Dahl, 1997). As a consequence, diffusional processes do not have the potential to affect Epidote-A U-Pb ages. However, the disturbance of the U-Pb isotope system may be affected by dissolution-precipitation mechanisms, as 460 has been reported for many minerals (e.g., monazite; Tartèse et al., 2011; Williams et al., 2011; Seydoux-Guillaume et al., 2012; Grand'Homme et al., 2018). Fluid-rock interaction has been shown to affect other isotopic systems as well (e.g., K-Ar, B; Halama et al., 2014). The microstructural relationships between Epidote-A and biotite in layer 1 are consistent with both minerals forming together in Alpine times and before vein deformation. Also, the U-Pb isotopic data presented in Peverelli et al. (2021) resolve a single generation of Epidote-A when plotted in a Tera–Wasserburg diagram. Moreover, the time-resolved ²⁰⁶Pb/²³⁸U ratios corrected for downhole fractionation display flat trends when corrected for zoning in initial 465 lead by applying a ²⁰⁸Pb correction (see Fig. 5 of Peverelli et al., 2021). This means that there is no resolvable isotopic zoning across the ca. 10–12 μm crater depth of the LA-ICP-MS measurements. The trace element data (Figs. 10c and 11; Table 2) reveal prominent chemical zoning. If any isotopic heterogeneity had been caused by interaction with the





deformation-related fluid, the crystal rims would be affected to a greater extent than the cores. Consequently, if resolvable isotopic zoning existed in the dated Epidote-A grains, plotting data-points from different domains of isotopic zoning would create scatter of the data-points in a Tera–Wasserburg diagram. However, U–Pb isotope measurements cover all geochemically variable zones (Fig. 10; white circles) and they define a single Tera–Wasserburg regression regardless of their proximity to cores/rims of the analyzed epidote grains. This supports that the dissolution–precipitation processes during Epidote-B formation did not appreciably disturb the U–Pb isotope system in Epidote-A relicts.

475 8 Conclusions and outlook

This study combines microstructural and geochemical methods to investigate the formation mechanisms of an epidote-quartz microfold within an epidote-quartz (\pm biotite) vein. For the first time, we have demony bed the occurrence of epidote dissolution-precipitation processes, and that this mechanism coexists with quartz dynamic recrystallization, both contributing to grain-size reduction of the system. The genetic link between epidote grains being dissolved and those being reprecipitated is revealed by the overlap of the major and trace element compositions of both generations. We have shown 480 that fluid circulation and the formation of epidote nucleation loci in the deforming system are enabled by dynamic granular fluid pump. Repeated dissolution and (re)precipitation processes of epidote in creep cavities formed among sliding quartz [2] Iso cause chemical homogenization of the epidote-dissolving/forming fluid, and hence of the new epidote generation gra forming during deformation. This demonstrates the importance of fluid recycling in deformational processes, although the application of Pb-Sr isotope geochemistry recognizes a role for the addition of externally derived fluids mediating mass 485 transfer processes in the studied epidote-quartz vein. Epidote plays an active role in determining the deformation behavior of other minerals, hence in the deformation style of polymineralic aggregates and in producing the resulting microstructures. To our knowledge, before this study, only brittle deformation behavior of epidote had been proven (Masuda et al., 1990; 1995), but no other detailed studies had addressed epidote deformation mechanisms. In this respect, epidote dynamic 490 recrystallization has never been either ruled out or demonstrated. However, since this mineral is widespread in crustal rocks (e.g., Bird and Spieler, 2004; Enami et al., 2004; Franz and Liebscher, 2004; Grapes and Hoskin, 2004; Schmidt and Poli,

Number: 1 Juthor: a Subject: Sticky Note Date: 23/06/2022 09:15:56
 demonstrate Strong word. You did not show mictrotextures typical of dissolution and precipitation processes (see for example Putnis 2009 Reviews in M logy & Geochemistry). I would rather use "suggest"
 Number 2 Author: a Subject: Sticky Note Date: 22/06/2022 15:11:24

<mark>⊜</mark> Number: 1	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/20/2022 9:44:12 PM	
Done				
_{) Number: 2}	Author: Veronica Peverelli	Subject: Sticky Note	Date: 7/21/2022 12:49:16 PM	
Corrected				





2004; Morad et al., 2010), the occurrence of epidote ductile deformation may have effects on the deformation of the continental crust and it calls for a better understanding of this mineral.

Appendix A

495 Transmitted light microscope images of the other epidote-quartz veins (samples P2, P3 and Gr0-c) associated with the studied epidote-quartz vein.



Figure A1: Transmitted light scans of (a) epidote vein P2, (b) epidote vein P3 and (c) epidote vein Gr0-c. Plane-polarized light.





Appendix B

515 Transmitted light photograph of the microfold in layers 2–3.



Figure B1: Transmitted light scan of the microfold in layers 2–3. Plane-polarized light.

525 Appendix C

Electron backscatter diffraction (EBSD) maps of dynamically recrystallized quartz by subgrain rotation in layer 3.



Figure C1: Electron backscatter diffraction (EBSD) map showing the orientation of quartz C axis in layer

Number: 1 Author: a Subject: Sticky Note Date: 22/06/2022 12:07:00
Please, add low and high and boundaries with a color. This is helpful to highlight subgrains, otherwise it is hard to substantiate your statement "dynamically recystallized quartz by subgrain rotation"

 Number: 1
 Author: Veronica Peverelli
 Subject: Sticky Note
 Date: 7/26/2022 7:50:26 AM

 Done. Subgrain rotation is confirmed by comparison with the literature (Stipp et al., 2002)





Author contribution

VP prepared the samples, carried out grain-size analysis and petrographic description, planned electron microprobe work, ran trace element measurements by LA-ICP-MS, measured Sr isotopes by TIMS, and prepared the manuscript. AB and MH supervised the work and greatly contributed to structuring the manuscript. MW supervised clean lab work and TIMS analyses, and measured Pb isotope data with IMV. TP granted access to the LA-ICP-MS laboratory and was closely involved in structuring the manuscript and in data evaluation. PL performed work at the electron microprobe and processed the X-ray maps. All authors read the manuscript and contributed to its improvement.

Acknowledgements

We thank Francesca Piccoli for technical assistance upon LA-ICP-MS measurements and feedback on the study, and Patrick
Neuhaus for carrying out the measurements of ²³⁸U/²⁰⁶Pb ratios at the Geography Department of University of Bern. We acknowledge funding of our new LA–ICP–MS facility through Swiss National Science Foundation, project 206021_170722, to Daniela Rubatto and Thomas Pettke. The solution ICP–MS isotope data were obtained on a Neptune MC–ICP mass spectrometer acquired with funds from the NCCR PlanetS supported by the Swiss National Science Foundation grant nr. 51NF40-141881. This work is part of the PhD thesis of Veronica Peverelli, who acknowledges SNF funding (project nr. 178785) granted to Alfons Berger.

Competing interests

The authors declare that they have no conflict of interest.

Code/data availability

All data are included in the manuscript (see tables).

550 References

Anenburg, M., Katzir, Y., Rhede, D., Jöns, N., and Bach, W.: Rare earth element evolution and migration in plagiogranites: a record preserved in epidote and allanite of the Troodos ophiolite, Contrib. to Mineral. Petrol., 169, https://doi.org/10.1007/s00410-015-1114-y, 2015.

Bambauer, H. U., Herwegh, M., and Kroll, H.: Quartz as indicator mineral in the Central Swiss Alps: The quartz
recrystallization isograd in the rock series of the northern Aar massif, Swiss J. Geosci., 102, 345–351, https://doi.org/10.1007/s00015-009-1319-z, 2009.



580



Barth, S., Oberli, F., Meier, M.: Th–Pb versus U–Pb isotope systematics in allanite from co-genetic rhyolite and granodiorite: implications for geochronology. Earth Planet. Sci. Lett. 124, 149–159, <u>https://doi.org/10.1016/0012-821X(94)00073-5</u>, 1994.

560 Belgrano, T. M., Herwegh, M., and Berger, A.: Inherited structural controls on fault geometry, architecture and hydrothermal activity: an example from Grimsel Pass, Switzerland, Swiss J. Geosci., 109, 345–364, https://doi.org/10.1007/s00015-016-0212-9, 2016.

Bergemann, C., Gnos, E., Berger, A., Whitehouse, M., Mullis, J., Wehrens, P., Pettke, T., Janots, E.: Th-Pb ion probe dating of zoned hydrothermal monazite and its implications for repeated shear zone activity: An example from the central alps, Switzerland. Tectonics 36, 671–689, <u>https://doi.org/10.1002/2016TC004407</u>, 2017.

 Switzerland. Tectonics 36, 671–689, <u>https://doi.org/10.1002/2016TC004407</u>, 2017.
 Berger, A., Gnos, E., Janots, E., Whitehouse, M., Soom, M., Frei, R., Waight, T.E.: Dating brittle tectonic movements with cleft monazite: Fluid-rock interaction and formation of REE minerals. Tectonics 32, 1176–1189, https://doi.org/10.1002/tect.20071, 2013.

Berger, A., Mercolli, I., Herwegh, M., Gnos, E.: Geological Map of the Aar Massif, Tavetsch and Gotthard Nappes. Geol.

570 spec. Map 1:100 000, explanatory notes 129. Federal Office of Topogaphy swisstopo, Bern, Switzerland, 2017 Berger, A., Wehrens, P., Lanari, P., Zwingmann, H., Herwegh, M.: Microstructures, mineral chemistry and geochronology of white micas along a retrograde evolution: An example from the Aar massif (Central Alps, Switzerland). Tectonophysics 721, 179–195, https://doi.org/10.1016/j.tecto.2017.09.019, 2017.

Berger, A., Egli, D., Glotzbach, C., Valla, P.G., Pettke, T., Herwegh, M.: Apatite low-temperature chronometry and
575 microstructures across a hydrothermally active fault zone. Chem. Geol. 588, 120633,
<u>https://doi.org/10.1016/j.chemgeo.2021.120633</u>, 2022.

Bird, D. K. and Spieler, A. R.: Epidote in Geothermal Systems, Rev. Mineral. Geochemistry 56, 235–300, https://doi.org/10.2138/gsrmg.56.1.235, 2004.

Bons, P.D., Elburg, M.A., Gomez-Rivas, E.: A review of the formation of tectonic veins and their microstructures. J. Struct. Geol. 43, 33–62, https://doi.org/10.1016/j.jsg.2012.07.005, 2012.

- Challandes, N., Marquer, D., Villa, I.M.: P-T-t modelling, fluid circulation, and 39Ar-40Ar and Rb-Sr mica ages in the Aar Massif shear zones (Swiss Alps). Swiss J. Geosci. 101, 269–288, <u>https://doi.org/10.1007/s00015-008-1260-6</u>, 2008. Choukroune, P. and Gapais, D.: Strain pattern in the Aar Granite (Central Alps): orthogneiss developed by bulk inhomogeneous flattening, 5, 411–418, https://doi.org/10.1016/b978-0-08-030273-7.50019-7, 1983.
- 585 Dahl, P. S.: A crystal-chemical basis for Pb retention and fission-track annealing systematics in U-bearing mineral, with implications for geochronology, Earth Planet. Sci. Lett., 150, 277–290, <u>https://doi.org/10.1016/S0012-821X(97)00108-8</u>, 1997.

Diamond, L. W., Wanner, C., and Waber, H. N.: Penetration depth of meteoric water in orogenic geothermal systems, Geology, 46, 1083–1066, https://doi.org/10.1130/G45394.1, 2018.



610



590 Egli, D., Baumann, R., Küng, S., Berger, A., Baron, L., and Herwegh, M.: Structural characteristics, bulk porosity and evolution of an exhumed long-lived hydrothermal system, 747–748, 239–258, https://doi.org/10.1016/j.tecto.2018.10.008, 2018.

Enami, M., Liou, J.G., Mattinson, C.G.: Epidote minerals in high P/T metamorphic terranes: Subduction zone and high- to ultrahigh-pressure metamorphism. Rev. Mineral. Geochemistry 56, 347–398, <u>https://doi.org/10.2138/gsrmg.56.1.347</u>, 2004.

595 Evans, B., Renner, J., and Hirth, G.: A few remarks on the kinetics of static grain growth in rocks, 90, 88–103, https://doi.org/10.1007/s005310000150, 2001.

Feineman, M. D., Ryerson, F. J., DePaolo, D. J., and Plank, T.: Zoisite-aqueous fluid trace element partitioning with implications for subduction zone fluid composition, 239, 250–265, https://doi.org/10.1016/j.chemgeo.2007.01.008, 2007.

- Ferreira, T., Rasband, W.: ImageJ User Guide User Guide ImageJ. Image J user Guid. 1.46r. 600 <u>https://doi.org/10.1038/nmeth.2019</u>, 2012.
 - Fitz Gerald, J.D., Stünitz, H.: Deformation of granitoids at low metamorphic grade. I: Reactions and grain size reduction. Tectonophysics 221, 269–297, <u>https://doi.org/10.1016/0040-1951(93)90163-E</u>, 1993.

Franz, G., Liebscher, A.: Physical and Chemical Properties of the Epidote Minerals-An Introduction-. Rev. Mineral. Geochemistry 56, 1–81, <u>https://doi.org/10.2138/gsrmg.56.1.1</u>, 2004.

605 Frei, D., Liebscher, A., Franz, G., Dulski, P.: Trace element geochemistry of epidote minerals. Rev. Mineral. Geochemistry 56, 553–605, <u>https://doi.org/10.2138/gsrmg.56.1.553</u>, 2004.

Gieré, R., Sorensen, S.S.: Allanite and other REE-rich epidote-group minerals. Rev. Mineral. Geochemistry 56, 431–493, https://doi.org/10.2138/gsrmg.56.1.431, 2004.

Gilgannon, J., Fusseis, F., Menegon, L., Regenauer-Lieb, K., and Buckman, J.: Hierarchical creep cavity formation in an ultramylonite and implications for phase mixing, 8, 1193–1209, https://doi.org/10.5194/se-8-1193-2017, 2017.

- Gilgannon, J., Waldvogel, M., Poulet, T., Fusseis, F., Berger, A., Barnhoorn, A., and Herwegh, M.: Experimental evidence that viscous shear zones generate periodic pore sheets, 12, 405–420, https://doi.org/10.5194/se-12-405-2021, 2021. Goncalves, P., Oliot, E., Marquer, D., Connolly, J.A.D.: Role of chemical processes on shear zone formation: An example
- from the Grimsel metagranodiorite (Aar massif, Central Alps). J. Metamorph. Geol. 30, 703–722, https://doi.org/10.1111/j.1525-1314.2012.00991.x, 2012.

Gottardi, R., Hughes, B.: Role of fluids on deformation in mid-crustal shear zones, Raft River Mountains, Utah. Geol. Magazine, 1 – 13, doi:10.1017/S0016756822000231, 2022.

Grand'Homme, A., Janots, E., Seydoux-Guillaume, A.M., Guillaume, D., Magnin, V., Hövelmann, J., Höschen, C., Boiron, M.C.: Mass transport and fractionation during monazite alteration by anisotropic replacement. Chem. Geol. 484, 51–68, https://doi.org/10.1016/j.chemgeo.2017.10.008.2018

620 <u>https://doi.org/10.1016/j.chemgeo.2017.10.008</u>, 2018.

Grapes, R.H., Hoskin, P.W.O.: Epidote group minerals in low-medium pressure metamorphic terranes. Rev. Mineral. Geochemistry 56, 301–345, <u>https://doi.org/10.2138/gsrmg.56.1.301</u>, 2004.



625



Guillong, M., Meier, D.L., Allan, M.M., Heinrich, C.A., Yardley, B.W.D., 2008. SILLS: A Matlab-Based Program for the Reduction of Laser Ablation ICP–MS Data of Homogeneous Materials and Inclusions. Mineral. Assoc. Canada Short Course 40, 328–333.

Haeusler M., Haas C., Lösch S., Moghaddam N., Villa I.M., Walsh S., Kayser M., Seiler R., Ruehli F., Janosa M.,
Papageorgopoulou C.: Multidisciplinary identification of the controversial freedom fighter Jörg Jenatsch, assassinated 1639
in Chur, Switzerland. PLOS ONE, 11(12): e0168014, 1-22, doi: 10.1371/journal.pone.0168014, 2016.

Halama, R., Konrad-Schmolke, M., Sudo, M., Marschall, H.R., Wiedenbeck, M.: Effects of fluid-rock interaction on
40Ar/39Ar geochronology in high-pressure rocks (Sesia-Lanzo Zone, Western Alps). Geochim. Cosmochim. Acta 126, 475–494, https://doi.org/10.1016/j.gca.2013.10.023, 2014.

Halter, W.E., Pettke, T., Heinrich, C.A., Rothen-Rutishauser, B.: Major to trace element analysis of melt inclusions by laserablation ICP-MS: Methods of quantification. Chem. Geol. 183, 63–86, <u>https://doi.org/10.1016/S0009-2541(01)00372-2</u>, 2002.

635 Handy, M. R.: The solid-state flow of polymineralic rocks, J. Geophys. Res.-Earth, 95, 8647–8661, https://doi.org/10.1029/JB095iB06p08647, 1990.

Handy, M. R.: Flow laws for rocks containing two non-linear viscous phases: A phenomenological approach, J. Struct. Geol., 16, 287–301, https://doi.org/10.1016/0191-8141(94)90035-3, 1994.

Hentschel, F., Janots, E., Trepmann, C.A., Magnin, V., Lanari, P., 2020. Corona formation around monazite and xenotime
during greenschist-facies metamorphism and deformation. Eur. J. Mineral. 32, 521–544. https://doi.org/10.5194/ejm-32-521-2020

Herwegh, M., & Berger, A.: Deformation mechanisms in second-phase affected microstructures and their energy balance. Journal of Structural Geology, 26(8), 1483–1498, <u>https://doi.org/10.1016/j.jsg.2003.10.006</u>, 2004.

Herwegh, M. and Jenni, A.: Granular flow in polymineralic rocks bearing sheet silicates: New evidence from natural examples, 332, 309–320, https://doi.org/10.1016/S0040-1951(00)00288-2, 2001.

- Herwegh, M., Linckens, J., Ebert, A., Berger, A., Brodhag, S.H.: The role of second phases for controlling microstructural evolution in polymineralic rocks: A review. J. Struct. Geol. 33, 1728–1750, <u>https://doi.org/10.1016/j.jsg.2011.08.011</u>, 2011.
 Herwegh, M., Berger, A., Baumberger, R., Wehrens, P., Kissling, E.: Large-Scale Crustal-Block-Extrusion During Late Alpine Collision. Sci. Rep. 7, 1–10, <u>https://doi.org/10.1038/s41598-017-00440-0</u>, 2017.
- Herwegh, M., Berger, A., Glotzbach, C., Wangenheim, C., Mock, S., Wehrens, P., Baumberger, R., Egli, D., and Kissling,
 E.: Late stages of continent-continent collision: Timing, kinematic evolution, and exhumation of the Northern rim (Aar Massif) of the Alps, Earth-Science Rev., 200, 102959, https://doi.org/10.1016/j.earscirev.2019.102959, 2020.
 Hobbs, B.E., Ord, A., Spalla, M.I., Gosso, G., Zucali, M.: The interaction of deformation and metamorphic reactions. Geol. Soc. Spec. Publ. 332, 189–223, https://doi.org/10.1144/SP332.12, 2010.



660

670



655 Hofmann, B. A., Helfer, M., Diamond, L. W., Villa, I. M., Frei, R., and Eikenberg, J.: Topography-driven hydrothermal breccia mineralization of Pliocene age at Grimsel Pass, Aar massif, Central Swiss Alps, Schweizerische Mineral. und Petrogr. Mitteilungen, 84, 271–302, 2004.

Horwitz, E. P., Dietz, M. L., and Chiarizia, R.: The application of novel extraction chromatographic materials to the characterization of radioactive waste solutions, J. Radioanal. Nucl. Ch., 161, 575–583, <u>https://doi.org/10.1007/bf02040504</u>, 1992.

Janots, E., Berger, A., Gnos, E., Whitehouse, M., Lewin, E., Pettke, T.: Constraints on fluid evolution during metamorphism from U-Th-Pb systematics in Alpine hydrothermal monazite. Chem. Geol. 326–327, 61–71, https://doi.org/10.1016/j.chemgeo.2012.07.014, 2012.

Karato, S.-I.: Deformation of Earth Materials: An Introduction to the Rheology of Solid Earth, Cambridge University Press, https://doi.org/10.1007/s00024-009-0536-8, 2009.

Kruse, R., Stünitz, H.: Deformation mechanisms and phase distribution in mafic high-temperature mylonites from the Jotun Nappe, southern Norway. Tectonophysics 303, 223–249, <u>https://doi.org/10.1016/S0040-1951(98)00255-8</u>, 1999.

Konrad-Schmolke, M., Halama, R., Wirth, R., Thomen, A., Klitscher, N., Morales, L., Schreiber, A., Wilke, F.D.H.: Mineral dissolution and reprecipitation mediated by an amorphous phase. Nat. Commun. 9, <u>https://doi.org/10.1038/s41467-018-03944-z</u>, 2018.

Lanari, P., Duesterhoeft, E.: Modeling Metamorphic Rocks Using Equilibrium Thermodynamics and Internally Consistent Databases: Past Achievements, Problems and Perspectives. J. Petrol. 60, 19–56, <u>https://doi.org/10.1093/petrology/egy105</u>, 2019.

Lanari, P., Vidal, O., De Andrade, V., Dubacq, B., Lewin, E., Grosch, E.G., Schwartz, S.: XMapTools: A MATLAB©-based

program for electron microprobe X-ray image processing and geothermobarometry. Comput. Geosci. 62, 227–240, https://doi.org/10.1016/j.cageo.2013.08.010, 2014.
 Lanari, P., Vho, A., Bovay, T., Airaghi, L., Centrella, S.: Quantitative compositional mapping of mineral phases by electron

probe micro-analyser. Geol. Soc. Spec. Publ. 478, 39–63, <u>https://doi.org/10.1144/SP478.4</u>, 2019.

Ludwig, K. R., 2012: On the treatment of concordant uranium-lead ages, Geochim. Cosmochim. Ac., 62, 665–676, https://doi.org/10.1016/S0016-7037(98)00059-3, 1998.

Marquer, D. and Burkhard, M.: Fluid circulation, progressive deformation and mass-transfer processes in the upper crust: the example of basement – cover relationships in the External Crystalline Massifs, Switzerland, J. Struct. Geol., 8–9, 1047–1057, 1992.

Marquer, D. and Peucat, J. J.: Rb-Sr systematics of recrystallized shear zones at the greenschist- amphibolite transition: examples from granites in the Swiss central Alps, 74, 343–358, 1994.

Masuda, T., Shibutani, T., Kuriyama, M., Igarashi, T.: Development of microboudinage: an estimate of changing differential stress with increasing strain. Tectonophysics 178, 379–387, <u>https://doi.org/10.1016/0040-1951(90)90160-A</u>, 1990.



690



Masuda, T., Shibutani, T., Yamaguchi, H.: Comparative rheological behaviour of albite and quartz in siliceous schists revealed by the microboudinage of piedmontite. J. Struct. Geol. 17, 1523–1533, <u>https://doi.org/10.1016/0191-8141(95)00060-Q</u>, 1995.

McClay, K. R.: Pressure solution and Coble creep in rocks and minerals: A review, J. Geol. Soc. London., 134, 57–70, https://doi.org/10.1144/gsjgs.134.1.0057, 1977.

McDonough, W.F., Sun, S.S.: The composition of the Earth. Chem. Geol. 120, 223–253, <u>https://doi.org/10.1016/0009-2541(94)00140-4</u>, 1995.

- Morad, S., El-Ghali, M.A.K., Caja, M.A., Sirat, M., Al-Ramadan, K., Manurberg, H.: Hydrothermal alteration of plagioclase in granitic rocks from Proterozoic basement of SE Sweden. Geol. J. 45, 105–116, <u>https://doi.org/10.1002/gj.1178</u>, 2010.
 Mullis, J., Dubessy, J., Poty, B., O'Neil, J.: Fluid regimes during late stages of a continental collision: Physical, chemical, and stable isotope measurements of fluid inclusions in fissure quartz from a geotraverse through the Central Alps, Switzerland. Geochim. Cosmochim. Acta 58, 2239–2267, <u>https://doi.org/10.1016/0016-7037(94)90008-6</u>, 1994.
- Passchier, C.W. and Trouw, R.A.: Microtectonics. Springer Science & Business Media, 2005.
 Paterson, M. S.: A theory for granular flow accommodated by material transfer via an intergranular fluid, 245, 135–151, https://doi.org/10.1016/0040-1951(94)00231-W, 1995.

Pettke, T., Oberli, F., Audétat, A., Guillong, M., Simon, A.C., Hanley, J.J., Klemm, L.M.: Recent developments in element concentration and isotope ratio analysis of individual fluid inclusions by laser ablation single and multiple collector ICP-MS.
705 Ore Geol. Rev. 44, 10–38, https://doi.org/10.1016/j.oregeorev.2011.11.001, 2012.

- Peverelli, V., Ewing, T., Rubatto, D., Wille, M., Berger, A., Villa, I.M., Lanari, P., Pettke, T., Herwegh, M.: U–Pb geochronology of epidote by laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) as a tool for dating hydrothermal-vein formation. Geochronology 3, 123–147, <u>https://doi.org/10.5194/gchron-3-123-2021</u>, 2021. Peverelli, V., Berger, A., Mulch, A., Pettke, T., Piccoli, F., and Herwegh, M.: Epidote U–Pb geochronology and H isotope
- geochemistry trace pre-orogenic hydration of mid-crustal granitoids, GEOLOGY, accepted.
 Putnis, A.: Mineral replacement reactions: from macroscopic observations to microscopic mechanisms. Mineral. Mag. 66, 689–708, <u>https://doi.org/10.1180/0026461026650056</u>, 2002.
 Putnis, A.: Mineral replacement reactions. Rev. Mineral. Geochemistry 70, 87–124, <u>https://doi.org/10.2138/rmg.2009.70.3</u>, 2009.
- Putnis, A., Putnis, C. V.: The mechanism of reequilibration of solids in the presence of a fluid phase. J. Solid State Chem. 180, 1783–1786, <u>https://doi.org/10.1016/j.jssc.2007.03.023</u>, 2007.
 Rehkämper, M., Mezger, K.: Investigation of matrix effects for Pb isotope ratio measurements by multiple collector ICP-MS: Verification and application of optimized analytical protocols. J. Anal. At. Spectrom. 15, 1451–1460, https://doi.org/10.1039/b005262k, 2000.





720 Ricchi, E., Bergemann, C.A., Gnos, E., Berger, A., Rubatto, D., Whitehouse, M.J.: Constraining deformation phases in the Aar Massif and the Gotthard Nappe (Switzerland) using Th-Pb crystallization ages of fissure monazite-(Ce). Lithos 342– 343, 223–238, <u>https://doi.org/10.1016/j.lithos.2019.04.014</u>, 2019.

Rolland, Y., Cox, S.F., Corsini, M.: Constraining deformation stages in brittle-ductile shear zones from combined field mapping and 40Ar/39Ar dating: The structural evolution of the Grimsel Pass area (Aar Massif, Swiss Alps). J. Struct. Geol. 31, 1377–1394, https://doi.org/10.1016/j.jsg.2009.08.003, 2009.

Rossi, M. and Rolland, Y.: Stable isotope and Ar/Ar evidence of prolonged multiscale fluid flow during exhumation of orogenic crust: Example from the mont blanc and Aar Massifs (NW Alps), 33, 1681–1709, https://doi.org/10.1002/2013TC003438, 2014.

Schaltegger, U., Krähenbühl, U.: Heavy rare-earth element enrichment in granites of the Aar Massif (Central Alps, Switzerland). Chem. Geol. 89, 49–63, https://doi.org/10.1016/0009-2541(90)90059-G, 1990.

Schaltegger, U., Corfu, F.: The age and source of late Hercynian magmatism in the central Alps: evidence from precise U-Pb ages and initial Hf isotopes. Contrib. to Mineral. Petrol. 111, 329–344, <u>https://doi.org/10.1007/BF00311195</u>, 1992.

Schmidt, M.W., Poli, S.: Magmatic epidote. Rev. Mineral. Geochemistry 56, 399–430, https://doi.org/10.2138/gsrmg.56.1.399, 2004.

- Schneeberger, R., Kober, F., Lanyon, G.W., M\u00e4der, U.K., Spillmann, T., Blechschmidt, I.: Grimsel Test Site: Revisiting the site-specific geoscientific knowledge, 2019.
 Seydoux-guillaume, A., Montel, J., Bingen, B., Bosse, V., Parseval, P. De, Paquette, J., Janots, E., and Wirth, R.: Low-temperature alteration of monazite : Fluid mediated coupled dissolution precipitation , irradiation damage , and disturbance of the U Pb and Th Pb chronometers, Chem. Geol., 330–331, 140–158, https://doi.org/10.1016/j.chemgeo.2012.07.031,
- 740 2012.

725

Stipp, M., Stünitz, H., Heilbronner, R., & Schmid, S. M.: Dynamic recrystallization of quartz: correlation between natural and experimental conditions. Geological Society, London, Special Publications, 200(1), 171-190. https://doi.org/10.1144/GSL.SP.2001.200.01.11, 2002.

Stünitz, H., Fitzgerald, J.D.F.: Deformation of granitoids at low metamorphic grade. II: Granular flow in albite-rich mylonites. Tectonophysics 221, 299–324, <u>https://doi.org/10.1016/0040-1951(93)90164-F</u>, 1993.

- Tartèse, R., Ruffet, G., Poujol, M., Boulvais, P., Ireland, T.R.: Simultaneous resetting of the muscovite K-Ar and monazite U-Pb geochronometers: A story of fluids. Terra Nov. 23, 390–398, <u>https://doi.org/10.1111/j.1365-3121.2011.01024.x</u>, 2011. Tullis, J.: Deformation of granitic rocks: Experimental studies and natural examples, Rev. Mineral. Geochemistry, 51, 51–95, https://doi.org/10.2138/gsrmg.51.1.51, 2002.
- 750 Villa, I.M., Hanchar, J.M.: K-feldspar hygrochronology. Geochim. Cosmochim. Acta 101, 24–33, https://doi.org/10.1016/j.gca.2012.09.047, 2013.





Wehrens, P., Berger, A., Peters, M., Spillmann, T., and Herwegh, M.: Deformation at the frictional-viscous transition: Evidence for cycles of fluid-assisted embrittlement and ductile deformation in the granitoid crust, Tectonophysics 693, 66–84, https://doi.org/10.1016/j.tecto.2016.10.022, 2016.

755 Wehrens, P., Baumberger, R., Berger, A., Herwegh, M.: How is strain localized in a meta-granitoid, mid-crustal basement section? Spatial distribution of deformation in the central Aar massif (Switzerland). J. Struct. Geol. 94, 47–67, <u>https://doi.org/10.1016/j.jsg.2016.11.004</u>, 2017.

Weis, D., Kieffer, B., Maerschalk, C., Barling, J., De Jong, J., Williams, G.A., Hanano, D., Pretorius, W., Mattielli, N., Scoates, J.S., Goolaerts, A., Friedman, R.M., Mahoney, J.B.: High-precision isotopic characterization of USGS reference

materials by TIMS and MC-ICP-MS. Geochemistry, Geophys. Geosystems 7, <u>https://doi.org/10.1029/2006GC001283</u>, 2006.
Williams, M.L., Jercinovic, M.J., Harlov, D.E., Budzyń, B., Hetherington, C.J.: Resetting monazite ages during fluid-related alteration. Chem. Geol. 283, 218–225, https://doi.org/10.1016/j.chemgeo.2011.01.019, 2011.
Wintsch, R.P., Yeh, M.W.: Oscillating brittle and viscous behavior through the earthquake cycle in the Red River Shear Zone: Monitoring flips between reaction and textural softening and hardening. Tectonophysics 587, 46–62,

765 <u>https://doi.org/10.1016/j.tecto.2012.09.019</u>, 2013.