Modelling floating riverine litter in the south-eastern Bay of Biscay: a regional distribution from a seasonal perspective

Irene Ruiz¹, Anna Rubio¹, Ana J. Abascal², Oihane C. Basurko¹

¹AZTI, Marine Research, Basque Research and Technology Alliance (BRTA), Pasaia, 20110, Spain

² IHCantabria - Instituto de Hidráulica Ambiental de la Universidad de Cantabria, Santander, 39011, Spain

Correspondence to: Irene Ruiz (iruiz@azti.es)

Abstract

5

Although rivers contribute to the flux of litter to the coastal and marine environment, estimates of riverine litter amounts are scarce and the behaviour of detailed studies on floating riverine litter behaviour once it has reached the sea are still scarce.at 10 river mouths and coastal waters is highly uncertain. This paper provides an-comprehensive overview analysis of the seasonal trends behaviour of floating riverine-marine released by riverslitter transport and fate within the south-eastern Bay of Biscay based on riverine litter characterization, drifters and high-frequency radars observations and Lagrangian simulations. Virtual particles were released close to the river mouths in the coastal area as a proxy of the floating litterfraction of riverine litter entering from rivers and reaching the open waters, ocean from rivers and Particles were parameterized with a wind drag coefficient (Cd) to represent their trajectories and fate according to the buoyancy of the litter items. They were forced with 15 numerical winds and measured currents provided by high-frequency radars covering selected seasonal week-long periods between 2009 and 2021. To gain a better insight on the type and buoyancy of the items, samples collected from a barrier placed at Deba river (Spain) were characterized at the laboratory. Items were grouped into two categories: low buoyant items (objects not exposed to wind forcing e.g., plastic bags) and highly buoyant items (objects highly exposed to wind forcing, 20 e.g., bottles). Overall, low buoyant items encompassed almost 90% by number and 68% by weight. Low buoyant items were parametrized with Cd=0%, and highly buoyant items with Cd=4%, this later one as a result of the joint analysis of modelled and observed trajectories of four satellite drifting buoys released at Adour (France), Deba (Spain) and Oria (Spain) river mouths. Results show that all regions in the study area are highly affected by rivers within or nearby the region itself. Simulations of riverine litterParticles parametrized with Cd=4% showed that particles drifted faster towards the coast by the 25 wind, notably during the first 24 hours. In summer, over the 97% of particles beached after one week of simulation. In autumn this value fell to 54%. In contrast, the low buoyant litter items take took longer to arrive to the coastlineshoreline, particularly during Spring with fewer than 25% of particles beached by the end of the simulations. When comparing coastline concentrations, Tthe highest concentrations of particles (>200 particles/km) were recorded during summer for Cd=4% in the French region of Pyrénées-Atlantiques for Cd=4%. Results showed that the regions in the study area were 30 highly affected by rivers within or nearby the region itself. These results coupled observations and a river-by-river

modelling approach and can assist policy and decision makers on setting emergency responses to high fluxes of floating

riverine litter-arrivals and on defining future monitoring strategies for heavy polluted regions within the south-eastern Bay of Biscaystudy area.

1 Introduction

- 35 Rivers act as key vectors bringing improperly disposed and mismanaged litter from land into coastal and the marine⁴ environments, especially in densely populated or highly industrialized river basins. Riverine litter poses a large threat not only to coastal and marine environments but also to freshwater systems by degrading aquatic life, impacting freshwater quality and increasing economic losses associated with human activities (van Emmerik and Schwarz, 2020; Al-Zawaidah et al., 2021). However, most of the litter research conducted to date has focused on marine environments (87%) when
- 40 compared to freshwaters systems (13%) (Blettler et al., 2018). Indeed, riverine litter contributions to oceans are still uncertain, and results vary depending on the approach applied such as the dataset or the model used (Lebreton et al., 2017; Schmidt et al., 2017; Meijer et al., 2021). Recent findings derived from extensive modelling efforts suggest that about 1,600 rivers worldwide account for 80% of plastic inputs to the ocean with small urban rivers among the most polluting (Meijer et al., 2021). However, most of the litter research conducted to date has focused on marine environments (87%) when
- 45 compared to freshwaters systems (13%), and only 7% of all scientific publications can be attributed to macroplastics (size > 2.5 cm) (Blettler et al., 2018). Riverine litter contributions to oceans are still uncertain, and results vary depending on the approach applied such as the dataset or the model used. Global estimates based on modelled amounts of mismanaged plastic waste (MPW) range between 0.5 to 2.7 million metric tonnes per year (Lebreton et al., 2017; Schmidt et al., 2017; Meijer et al., 2021); however, they can represent less than a tenth when methodology followed differ from MPW-based models (Mai
- 50 et al., 2020). Models require comprehensive field data and consistent and harmonized protocols to validate the amounts, type and size of riverine inputs, information that can then be used to implement tailor-made and effective measures at regional and local scale (González-Fernández and Hanke, 2017; Wendt-Potthoff et al., 2020; Margenat et al., 2021). Such comprehensive data was obtained in Europe thanks to the RIMMEL project (González-Fernández and Hanke, 2017). -and a network of visual observers of riverine macrolitter, which This research concluded that between 307 and 925 million of floating riverine litter items are annually transferred into the ocean, mainly through small rivers, streams and coastal run-off (González-Fernández et al., 2021).

60

Once at the river mouthit has reached the sea, floating riverine litter can accumulate close to the shorelinenearby or it can can be transported to open watersmove long distances, reaching remote areas far from the coastfrom river waters. Indeed, the distribution and fate of riverine-floating litter in the coastal and marine environment is conditioned affected by the metocean conditions (currents, turbulence, wind) but also by the buoyancy of the objects, defined by their composition, size and shape (Ryan, 2015; Lebreton et al., 2019; Maclean et al., 2021). Objects with low buoyancy are mainly driven by currents contrary to highly buoyant items which are pushed-driven along the water surface partially by winds. The wind effect ("windage") on floating marine litter behaviour is an important factor for pushing litter to shore and induce beaching, mainly for offshore-source litter, which is highly affected by winds, compared to coastal source macrolitter (Ko et al., 2020). Riverine litter

Con formato: Espacio Después: 12 pto

- 65 trapped in near shore areas is susceptible to beaching, settling and resurfacing episodes and reach open ocean mostly as small fragments (Morales Caselles et al., 2021), hampering cleanup efforts and contributing to the prevalence of litter in the marine environment. Adjustment for windage has been consequently investigated has been further investigated by in Lagrangian modelling studies in open ocean (Allshouse et al., 2017; Maximenko et al., 2018; Lebreton et al., 2019; Abascal et al., 2009) but also, although less mature, when compared to the in-coastal areas (Critchell and Lambrechts, 2016;
- 70 Utenhove, 2019; Tong et al., 2021). The lack of field-observational data to accurately parametrize the effect of wind and validate simulation results is one of the key limitations for parametrizing the windage effect and accurately predict floating marine litter behaviour-both in riverine and marine transport modelling. However, observations derived from drifting buoys, From decades, researchers have used real observations derived from drifting buoys, such as those provided for decades byin the Global Drifter program, which observations contribute to have been used to fill this gap.
- 75 Buoy data are used to fine tuning prediction models and provide a better description of the near surface circulation and its⁴ Lagrangian behaviour (Charria et al., 2013; Dagestad and Röhrs, 2019). They have also-allowed simulating more realistic floating marine_litter pathways from origin to fate by integrating experimental windage parametrizations and the corresponding comparison between observed and modeled trajectories (Duhec et al., 2015; Pereiro et al., 2018; Rizal et al., 2021). Nowadays, more affordable and environmentally friendly solutionsSatellite-tracked_drifting_buoys_and
- communication systems are costly, despite more economical and environmentally friendly solutions are gaining force among researchers <u>as</u>. Examples include-drifters built using biopolymers (Novelli et al., 2017; D'Asaro et al., 2020) <u>or and compact</u> and lightweight designs with a GPS-tracking component for an easy deployment (Meyerjürgens et al., 2019b; van Sebille et al., 2021), <u>Others have evolved to develop drifters shaped as real litter items (e.g., plastic bottles), which allow a more accurate tracking position of standard objects, accounting for wind effect at sea and on inland waterways (Duncan et al., 2020).
 </u>

At coastal scale, windage parametrization combined with realistic knowledge on coastal circulation become crucial to reduce the uncertainties of modelled trajectories (Van Sebille et al., 2020), Nowadays, coastal transport can be also characterized at high temporal and spatial resolution thanks to the use of ILand-based high frequency radar systems for the remote measurement of surface currents (hereafter HF radars (Rubio et al., 2017)). HF radars offer the opportunity to monitor

- 90 surface currents in coastal areas, where the transport processes are significantly more complex than open ocean waters due to the effect of coasts, bathymetry and other local forcings, like (e.g., river discharges or coastal upwellings). Given the highly dynamic and complexity nature of coastal waters, this realistic and useful knowledge on coastal circulation combined with the parametrization of key physical processes affecting litter transport (e.g., windage) become crucial to reduce the uncertainties of modelled trajectories of riverine and marine litter (Van Sebille et al., 2020).
- 95 -In the the south-eastern Bay of Biscay (hereafter SE Bay of Biscay), a HF radar provides, as part of the operational oceanography system EuskOOS (<u>https://www.euskoos.eus/</u>), near-real-time surface current fields at 5 km spatial and 1 hour temporal resolution, covering since 2009 a range up to 150 km from the coast. Theis system has already been already used in

Con formato: Espacio Después: 12 pto

1	Con formato: Español (España)
(Código de campo cambiado
(Con formato: Español (España)
(Código de campo cambiado

Con formato: Español (España)
Con formato: Español (España)
Con formato: Español (España)
Con formato: Español (España)
Código de campo cambiado

Con formato: Espacio Después: 12 pto

previously to study surface coastal transport processes in combination with multisource data (Manso-Narvarte et al., 2018, 2021; Rubio et al., 2011, 2013, 2018, 2020; Solabarrieta et al., 2014, 2015, 2016). The HF radar is also a good example of
 effective monitoring of surface currents with strong potential for floating marine litter management. The EuskOOS HF radar is part of JERICO RI (https://www.jerico-ri.eu/) and it is operated following JERICO-S3-project best practices, standards, and recommendations. Research conducted by Declerck et al., (2019) in the SE Bay of Biscay provided the first assessment of floating litter transport and distribution in the region, coupling surface currents observations from EuskOOS system,

Lagrangian modelling and riverine inputs. Nowadays, these observations are used by local authorities both in real time and

in hindcast in the framework of the operational service FML-TRACK (https://fmltrack.rivagesprotech.fr/)-to collect floating

- 105
 - marine litter in the area. However, the accurate modelling of transport and fate of both-floating marine and riverine litter need to consider the variety of floating objects and sources and additional physical processes as windage.

This paper aims at estimating the seasonal trends-behaviour of the floating fraction of marine litter released by rivers within the SE Bay of Biscay reaching open waters. floating riverine litter of transport and fate in the SE Bay of Biscay by modelling the Lagrangian behaviour of numerical particles released in the main rivers within the area. To do so, a

110 modelling the Lagrangian behaviour of numerical particles released in the main rivers within the area. To do so, a Lagrangian model was forced by real observations from the EuskOOS HF radar and particles were parameterized to represent the floating marineriverine litter trajectories of two groups of items according to their observed buoyancy. Riverine litter collected from a local barrier was characterized at the laboratory to explore the fraction of highly and low buoyant items. Since most of the items were low buoyant, simulations of particles considering only surface currents were performed as the reference. Complementary Lagrangian simulations for highly buoyant items (and less abundant in the area) were also performed. In this case, 4 low-cost buoys with similar buoyancy of certain highly buoyant objects were built and released at 3 different rivers. Drifter data were used to parameterize the wind effect on this type of items and consequently achieve more accurate results.

2 Study area

- 120 The study was conducted in the SE Bay of Biscay, between north-eastern NE-Spain (Basque Country) and south-western SW France (Landes). The study area extends from 43.27°N to 44.58°N and from 3.18°W to 1.27°W, falling within the coverage area of the HF radar station of the operational oceanography system EuskOOS (Fig 1). The study area comprises two Basque regions Bizkaia (Spain) and Gipuzkoa (Spain) -, two French departments Pyrénées-Atlantiques (France) and Landes (France) -, and eight rivers Deba (Spain), Urola (Spain), Oria (Spain), Urumea (Spain), Oiartzun (Spain), Bidasoa (Spain), Nivelle (France) and Adour (France) -. The mean annual river discharge varies widely between rivers 3.71 m³/s (Oiartzun)
- to 350 m³/s (Adour) (Sheppard, 2018) and the population density differs between the Spanish and French border 44.8 inhabitants/km² (Landes) to 303.7 inhabitants/km² (Basque Country) -(Eurostat, 2019). The bathymetry in the SE Bay of Biscay is characterized by the presence of a narrow continental shelf ranging between 7 and 24 km wide in the Basque area, gradually increasing along the French coast up to about 70 km (Bourillet et al., 2006; Rodríguez et al., 2021). The

Con formato: Sin Resaltar

130 continental shelf in the SE Bay of Biscay comprises two mainly areas, the Aquitaine shelf with a N-S orientation and Cantabrian shelf with an E-W orientation. The continental slope is very pronounced, with a slope of the order up to 10%-12% (Sheppard, 2018).

Over the continental shelf, the ocean circulation The circulation of the self-water masses is marked by a seasonal variability. At shorter temporal scales, circulation is mostly modulated by the bathymetry and the coastal orientation, the density-driven

- 135 currents, and winds (Le Boyer et al., 2013; Solabarrieta et al., 2014). Tidal currents in the area are quite week constrained by topography and <u>the</u> width o<u>f</u>n the continental shelf (Lavin et al., 2006; González et al., 2007; Karagiorgos et al., 2020). Along-shelf currents are more intense and persistent during winter and autumn (about 10–15 cm s⁻¹), contrary to the other seasons, especially in summer (about 2.5 cm s⁻¹)(Charria et al., 2013). In winter, the prevailing SW winds causes an E to N flow and the moderate to strong NW winds occurring in spring and summer induce S and SW surface currents circulation
- 140 over the French and Spanish coasts-accompanied by a greater variability (Solabarrieta et al., 2015). In winter, westerly winds in the Basque coast reinforce the slope current (named "*Iberian Poleward Current*" (IPC)), a warm and saline intrusion trapped within the 50 km of the shelf edge, achieving reaching its greatest velocities (up to 70 cm s-1) during this season. The IPC favours the along slope transport of water masses (Solabarrieta et al., 2014; Porter et al., 2016). The exchange between shelf and deep sea waters in winter is associated to the generation of eddies, from the interaction of currents with
- 145 the topography (Lavin et al., 2006; Rubio et al., 2018; Teles-Machado et al., 2016). Maximum run-offs combined with SW winds also allow river plumes spread northwards and along the French shore during winter. However, this path changes in spring, when river discharges are reduced and winds blow from NW (Lavin et al., 2006; Puillat et al., 2006), The main circulation features in the study area are summarized in the figure created by (Declerck et al., 2019).

First global modelling studies coupling ocean circulation and Lagrangian particle tracking models reported that the SE Bay

- 150 of Biscay is a hotspot for floating marine litter (Lebreton et al., 2012; van Sebille et al., 2012). Recent Lagrangian modelling studies combining measured and predicted surface currents by the HF radar and the IBI Copernicus model revealed that floating marine litter circulation in the SE Bay of Biscay is marked by a high seasonal variability. Floating marine litter distribution in the SE Bay of Biscay follows the general circulation in the area, Results showeding a higher retention during spring and summer and a northward dispersion along the French coast during autumn and winter (Declerck et al., 2019;
- 155 Rubio et al., 2020). Surface currents derived from Regional Ocean Modelling System (ROMS) and a particle-tracking model were combined by Pereiro et al., 2019 to track the numerical drifters representing floating marine litter in the Bay of Biscay. In this study, longer residence times and higher concentrations were observed in the SE Bay of Biscay when compared north-western Iberian coastal waters, particularly in winter. Rodríguez-Díaz et al., 2020 showed from numerical simulations run using HYCOM model that floating marine litter items with high windage (Cd=3%-5%) tend to accumulate in nearshore
- 160 areas of the Bay of Biscay or end up beached. These trend is consistent with recent numerical simulations combining surface currents from the operational Iberian Biscay Irish System (IBI) and the numerical model TESEO that also revealed the highly buoyant items (Cd=4%) rapidly beach in the SE Bay of Biscay, mainly in spring and summer (Ruiz et al., 2022a).Longer residence times and higher concentrations are observed in winter influenced by the run offs and the influx of

Con formato: Español (España) Con formato: Español (España) Código de campo cambiado Código de campo cambiado 165

170

floating litter from local but also from distant sources (Pereiro et al., 2019). In the Bay of Biscay, macrolitter with high windage tends to accumulate in nearshore areas (with a probability of around 90%) or are beached (with a probability higher than 60%) (Rodríguez Díaz et al., 2020). A similar trend was observed in the study area by (Ruiz et al., 2022) who concluded that macrolitter items with a high windage rapidly beach during spring and summer underlining the importance of windage effect on the coastal accumulation. Since June 2020, innovative detection and tracking solutions combining ocean modelling and remote observation systems are operating in the SE Bay of Biscay for supporting floating marine litter reduction strategies both downstream (interception at sea with collect vessels and on beaches with cleaning facilities) and upstream (source identification and reduction) (Delpey et al., 2021).



Fig 1. Study area with the release locations of the Satellite drifting buoys and the riverine barrier. Dots in light yellow represent the nodes of the HF Radar grid. Dots in orange represent the trajectories of the buoys. <u>Dots in light yellow represent the nodes of the HF Radar grid</u>. Numbers with stars in pink correspond to the particle releasing location for riverine-floating marine litter simulations: (1) Deba; (2) Urola; (3) Oria; (4) Urumea; (5) Oiartzun; (6) Bidasoa; (7) Nivelle; and (8) Adour River. Dots in light yellow represent the nodes of the HF Radar grid.

3 Methods and Data

175

3.1 Riverine Litter Sampling

In Spring 2018, a riverine barrier was placed in Deba river (Gipuzkoa) to retain and collect floating macro-riverine litter during low to moderate flows. <u>This barrier enabled a passive sampling for characterize litter items at lab.</u> The barrier, which consisted of a nylon artisanal net supported by hard floats (buoys), was 40 m long and 0.6 m high with a 60 mm mesh size (see photos in Appendix A). <u>The</u>A sampling was conducted weekly from April 2018 to June 2018. <u>I; in</u> total eight riverine litter samples were collected. Litter items were quantified, weighted, and categorized at lab according to the Master list included in the "Guidance on Monitoring of Marine Litter in European Seas" (Galgani et al., 2013) Items were grouped into

185 7 types of material (artificial polymer materials, rubber, cloth/textile, processed/worked wood, paper/cardboard, metal, and glass/ceramics) and further classified into 44 categories (see the classification in Appendix B). Riverine litter items were also categorized into two groups (low and highly buoyant items) considering their exposure to wind based on (Ruiz et al., 2022a).

3.2 Drifters Observations

Four satellite drifting buoys (herein after 'low-cost buoys') were built by the authors and deployed one-by-one in the river 190 mouths of Oria (1 buoy), Deba (1 buoy), and Adour (2 buoys) between April 2018 and November 2018 (Fig 1, Table 1). The 'low-cost buoys' provided positioning every 5 minutes using satellite technology. 'Low-cost buoys' were 9 cm in height, 9.5 cm in float diameter and weighed approximately 200 g (Fig 2). A GPS (SPOT Trace device) powered by 4 AAA cells was placed in the bottom of a high-density polyethylene (HDPE)-plastic container sealed to guarantee water tightness. They were chosen because of their capability to ensure a reasonable balance between an accurate signal emission and their 195 purchase and communication fees. SPOT Trace devices have been used over the past few years in coastal and open ocean applications in a wide range of studies. Studies range for calibrating HF Radars (Martínez Fernández et al., 2021), tracking drifting objects as icebergs (Carlson et al., 2020), pelagic Sargassum (Putman et al., 2020; van Sebille et al., 2021) or fishing vessels (Widyatmoko et al., 2021; Hoenner et al., 2022) to search and rescue training (Russell, 2017) and oil spill and litter monitoring (Novelli et al., 2018; Meyerjürgens et al., 2019b). Almost 2/3 of the buoy floated above the water surface thus 200 preventing any satellite signal losses. Buoys A and D and transmitted their positions on an ongoing basis until their landing. Buoys B and C stopped emitting while they were drifting. In all cases, battery lifetime was enough for an adequate performance of the buoys. Once on land, citizens collected the buoys and reported their corresponding position.

Transmission periods relied upon battery lifetime and buoys landing.



Fig 2. Main components of the "Low-cost buoy". The structure: (a) HDPE container and SPOT Trace device powered by 4 AAA cells. Assembly process: (b) final appearance once the buoy is sealed. The buoy is labelled with contact information both within and outside; (c) the SPOT Trace was fixed at the base of the container with adhesive tape to avoid twists and turns of the buoy.

Table 1.	Locations,	periods, and	1 distances	covered l	by the	drifting	buoys
----------	------------	--------------	-------------	-----------	--------	----------	-------

Buoy ID	River	Initial date	Final date	Distance covered (km)
А	Deba	16-Sept-2018 8:00	4-Oct-018 7:00	116.1
В	Oria	12- Apr-2018 16:00	18-Apr-2018 12:00	118.72
С	Adour	29-Jul-2018 20:00	2-Aug-2018 20:00	71.21
D	Adour	28-Nov-2018 9:00	30-Nov-2018 11:00	64.41

215 3.3 HF radar <u>c</u>urrent <u>o</u> bservations and wind data

Surface velocity current fields were obtained from the EuskOOS HF radar station composed by two antennas located at Matxitxako and Higer Capes and covering the SE Bay of Biscay since 2009 a range up to 150 km from the coast. The EuskOOS HF radar is part of JERICO-RI and it is operated following JERICO-S3 project best practices, standards, and recommendations (see (Solabarrieta et al., 2016; Rubio et al., 2018) for details). Data consist of hourly current fields with a 5 220 km spatial resolution obtained from using the gap-filling OMA methodology (Kaplan and Lekien, 2007; Solabarrieta et al., 2021). 85 OMA modes, built setting a minimum spatial scale of 20 km and applied to periods with data from the two antennas, were used to provide the maximum spatiotemporal continuity in the HFR current fields, which is a prerequisite to performing accurate Lagrangian simulations. The application of OMA methodology has been validated for the Lagrangian assessment of coastal ocean dynamics in the study area by Hernández-Carrasco et al., 2018. HF radar velocities were quality 225 controlled using procedures based on velocity and variance thresholds, signal-to-noise ratios, and radial and total coverage, following standard recommendations (Mantovani et al., 2020). Data subsets were built for the Lagrangian simulations avoiding periods with temporal gaps (still present in case of failure of one or the two antennas) of more than a few hours. Data used for the Lagrangian simulations were extracted considering the outputs from the standard QC (quality control) procedures for real-time HF radar data (Rubio et al., 2021). Once extracted, data were visually inspected to ensure a 230 complete radial coverage (i.e., ensuring optimal OMA reconstructed fields) and build data subsets for the Lagrangian simulations avoiding periods with temporal gaps of more than a few hours.

Hourly ERA5-U10-wind fields were obtained from the atmospheric reanalysis computed using the IFS model of the European Center for Medium-Range Weather Forecast (ECMWF) (see (C3S, 2019) for details). ERA5 atmospheric database covers the Earth on a 30 km horizontal grid using 137 vertical levels from the surface up to a height of 80 km and provides estimates of a large number of atmospheric, land and oceanic climate variables-on a 0.3° × 0.3° grid, currently from 1979 to within 3 months of real time. Both HF radar current observations and wind data cover the drifter's emission periods and the selected week-long periods between 2009 and 2021 for riverine litter simulations (see Appendix C for the selected periods).

3.5 Particle <u>t</u>Transport <u>m</u>Model

The application of the transport module of the TESEO particle-tracking model (Abascal et al., 2007, 2017a, b; Chiri et al., 2020) was twofold: (1)applied to-simulate the transport and fate of floating riverine-marine litter entering from rivers and reaching the open waters of the SE Bay of Biscay and (2) estimate a windage coefficient by calibrating the model according to the 'low-cost buoys' trajectories. This module allows for simulating passive particles driven by surface currents, wind and turbulent diffusion. Particle trajectories were calculated using the following equation:

$$\frac{d\vec{x}i}{dt} = \vec{u}_{a}(\vec{x}_{i,\underline{t}}) + \vec{u}_{d}(\vec{x}_{i,\underline{t}})$$
(1)

where $\overline{u_a}$ and $\overline{u_d}$ are the advective velocity and diffusive velocity, respectively, for the $\vec{x_i}$ point and t time. The advective velocity is calculated as the lineal combination of the wind and currents according to:

250

255

$$\overrightarrow{u_a} = \overrightarrow{u_c} + C_d \overrightarrow{u_w}$$
(2)

where $\vec{u_c}$ is the surface current velocity, $\vec{u_w}$ is the wind velocity at 10m over the sea surface and Cd is the wind drag coefficient. The turbulent diffusive velocity is obtained using Monte Carlo sampling in the range of velocities $[-\vec{u_d}, \vec{u_d}]$ which are assumed to be proportional to the diffusion coefficients (Hunter et al., 1993; Maier-Reimer and Sündermann, 1982). For each timestep Δt , the velocity fluctuation is defined as:

$$|\overrightarrow{u_d}| = \sqrt{\frac{6D}{\Delta t}}$$
(3)

where D is the diffusion coefficient, whose value is 1 m²/s in accordance to previously modelling work for floating marine litter (Pereiro et al., 2019; Ruiz et al., 2022). Simulations were forced by HF radar surface current velocity and wind data and interpolated at the particle's position for integrating the trajectories. Beaching along the coast was implemented by a simple approach: if the particle reaches the shoreline, it is identified as beached, and it is removed from the computational process.items from selected rivers once they arrive to the coastal area. Simulations were forced by HF radar surface current velocity and wind data. The transport module was also used to accurately estimate the windage coefficient by calibrating the model according to the 'low cost buoys' trajectories. TESEO has been calibrated and validated by comparing virtual particle trajectories to observed surface drifter trajectories at regional and local scale (Abascal et al., 2009, 2017a, b; Chiri et al., 2019). Although the TESEO is a 3D numerical model conceived to simulate the transport and degradation of hydrocarbons but₅ it has also been successfully applied to other applications such as the study of the transport and accumulation of marine litter in estuaries (Mazarrasa et al., 2019; Núñez et al., 2019) and in open waters (Ruiz et al., 2022a).

3.5.1 Wind drag estimation

Two simulation strategies were combined for (1) estimating the wind drag coefficient and (2) study the seasonal behaviour of <u>floatingriverine</u> litter items in the area (section 3.5.2) The wind drag coefficient (Cd) was determined by comparing the observed trajectories provided by the 'low-cost buoys' and the modelled trajectories performed with TESEO. The test was done through different parametrizations of the wind drag coefficient ranging from 0% to 7% (Table 2). This range was chosen based on previously -floating marine litter studies coupling Lagrangian modelling and observations from satellite drifting buoys (Carson et al., 2013; Stanev et al., 2019; Van Der Mheen et al., 2019). The coefficient providing the lowest error was considered the best coefficient to simulate highly buoyant litter. Due to the grid limitations of the surface currents and wind data in the coastal area, the comparison was not initialised at the launching position of the 'low-cost buoys' (river mouths) but instead it was initialised at the closest grid element that contained valid currents and wind data (Table 1). Observed positions were interpolated jento a uniform one-hour time, fitting the met-ocean temporal resolution. A release of 1,000 virtual particles was performed every 4 hours at the corresponding observed position (Table 2). Particles were tracked

280 over a 24-hour period and the trajectory of the center of mass of all the particles was computed at every time step to represent the track of the particle cloud. Observations were compared to modeled trajectories using the simple separation distance, which is the difference between the observed and the computed position of the center of mass at a time step t. Mean separation distance $\overline{D(t^{mod})}$ was calculated for every modelled position based on the simple separation distance following Eq. (41):

290

$$\overline{\mathcal{D}(t^{mod})} = \frac{1}{N} \sum_{i=1}^{N} \left| \vec{X}^{mod} \left(t^{mod} \right) - \vec{X}^{obs}(t^{obs}) \right| \quad (\underline{41})$$

where \vec{X}^{mod} (t^{mod})e and \vec{X}^{obs} (t^{obs}) are the modeled and observed trajectories for the simulation period i of a total of N periods. A mean separation distance curve was computed for every wind drag coefficient derived from the mean separation distance curves of the four buoys. The area beneath the mean separation distance curve was calculated to select the more suitable wind drag coefficient.—The area \tilde{D} was calculated as a numerical integration over the forecast period via the trapezoidal method following Eq. (52). This method approximates the integration over an interval by breaking the area down into trapezoids with more easily computable areas:

$$\widetilde{D} \approx \int_{t \mod = 1}^{t \mod = 24} \overline{\mathrm{D}(t^{\mod})} dt$$
 (52)

3.5.2 Lagrangian seasonal simulation of riverine litter items

Seasonal simulations were run for low and highly buoyant items to assess the seasonal differences on the transport and fate of floating riverine litter once it has reached the open waters of the SE Bay of Biscay. transport and fate. Particles were released around 2.5 nautical miles off the coastline_shoreline due to the complexity in resolving small-scale processes of both floating marine and riverine litter behaviour in and near_close to the river mouths. As parametrizations concerning wind effect linked to the object characteristics are scarce, the optimal wind drag coefficient estimated for the buoys (see section 3.5.1) was accounted for simulated the behaviour of the objects highly exposed to wind. No wind drag parametrization

300 (Cd=0%) was applied for low buoyant objects not subjected to wind effect. A total of ten periods per season uniformly distributed within the study period (2009-2021) were considered for the simulations based on the availability of HF radar Con formato: Inglés (Reino Unido)

surface current datasets (see Appendix C for the selected periods). In total, <u>80 simulations (40 for Cd=0% and 40 for Cd=4%) were run for 7 days. For each simulation, 4,000 particles were released in 8 rivers for each selected period (500 per river) assuming that river discharges are equal despite the seasonal variations and the morphological differences between 100 particles. Simulations were run for 7 days. The total number of particles modeled for Cd=0% was the same as Cd=4%. Particles were released around 2.5 nautical miles off the coastline due to the complexity in resolving small scale processes in and near the river mouths. A post-processing was carried out to compute by river: (1) the particles' evolution over the time from their release until their arrival to the coastlineshoreline; and (2) the particles' distribution on the coastlineshoreline, counting the number of beached particles per km of coastline and indicating the spatial concentration per the sease in and indicating the spatial concentration per the sease the number of particles per km of coastline and indicating the spatial concentration per the sease the sease of the sease the sease and indicating the spatial concentration per the sease the number of beached particles per km of coastline and indicating the spatial concentration per the sease th</u>

310 region.

	Simu	lation paramete	rs	Release p	arameters	Physical p	arameters
	Number of particles	Integration time	Time step	Release locations	Release time	Turbulent diffusion coefficient	Wind drag coefficient (Cd)
Simulations for wind drag estimation	1,000 per location	24 h	60 s	At the observed locations of the buoy	Over the emitting period of the buoy at spaced intervals of 4 hours	1 m²/s	0 %, 2%, 3%, 4%, 5%, 6%, 7%
Seasonal riverine litter simulations	500 per river	l week	60 s	At a distance of 2.5 nautical miles from the river mouth	At the beginning of the selected time period (10 periods per season)	1 m²/s	0 %, 4%

Table 2. Simulation, release, and physical parameter values for wind drag estimation and floating riverine litter simulations.

315 4 Results

4.1 Riverine litter characterization

In total 1,576 items and 11.597 kg of <u>floating</u> riverine litter were sampled and characterised (Fig 3). *Plastic* was the most common type of riverine litter in terms of number of items (95.1%) and in weight (67.9%); they were also frequent *Glass/ceramics* (16.1%) and *Cloth/textile* items (6.9%) when counted by weight. The top ten litter items accounted for

320 93.3% by number and 72.6% by weight of the total riverine litter (Table 3). *Plastic/polystyrene pieces between 2.5 cm and 50 cm* and *Other Plastic/polystyrene identifiable items* (e.g., food labelling) were the most abundant in terms of number (71.2%) and weight (16.9%). *Low buoyant items* encompassed almost 91% by number and 68% by weight of litter items (Fig 4).



325

Fig 3. Composition of riverine litter by type of material based on the in terms of number and weight of riverine litter. Litems were collected fromin the riverine barrier placed located in Deba river (Gipuzkoa) between April and June 2018.



330 Table 3. Top ten (X) riverine litter items collected in the riverine from the barrier located in Deba river (Gipuzkoa) between April and June 2018. Items have been ranked by abundance (left) and weight (right) according to the MSFD Master List Categories of Beach Litter Item and classified based on their exposure to wind effect.

Fig 4. Riverine litter items classification based on the exposure to wind effect. <u>from riverine litter litter litters</u> collected from in the riverine barrier located placed in Deba river (Gipuzkoa) between April and June 2018.

4.2 Wind drag coefficient for drifting buoys

Total distances covered by drifting buoys ranged from 62 km to 118 km (Table 1) and they all spread outscattered over the
rivers inside the HF radar coverage area. Buoys, spanning approximately 44°N and 2° 22'W. They provided their position data over 385 h before beached on Landes and Gipuzkoa coastlines. When compared with numerical trajectories obtained using different Cd parameterizations, the mean separation distance (D(t^{mod}), increased nearly linearly with time for all the parametrizations, achieving a maximum separation of almost 14 km at 24 hours for Cd=0% (Fig 5). Overall, using no windage parametrization gave provided the largest D. Simulations parametrized with Cd=4% gave-provided the best results
with an average ± standard deviation (SD) of 3.2 ± 1.25 km and a maximum value of 4.85 km at 24 h. When assessing the mean separation distance for Cd=4% was 2- 4 km (Fig 6). Hence, a wind drag coefficient of 4% was applied in the remaining analysis to estimate riverine litterthe behaviour of highly buoyant items.

Con formato: Inglés (Estados Unidos)

4.3 Seasonal trends on floating riverine litter transport and fate

- 350 Particle concentrations in the coastline varied between 0 and 258.46 particles/km (Fig 7). Particles parametrized with Cd=4% drifted faster towards the coast by the wind, notably during the first 24 hours. The highest concentrations (>200 particles/km) were recorded during summer in Pyrénées-Atlantiques for Cd=4%, probably due to the seasonal retention patterns within the study area (Appendix D). Although less intensely, Cd=4% also lead to a high particle concentration in Pyrénées-Atlantiques (106.86 particles/km) and Gipuzkoa (166.1 particles/km) during winter. The IL-owest concentrations
- 355 (0-20 particles/km) were recorded for Cd=0% at all seasons during the after the first 24 hours of simulation, and particularly during autumn. Overall, Bizkaia was the less affected impacted region by litter for both windage coefficients (<40 particles/km). During summer, over the 97% of particles parametrized with Cd=4% beached after one week of simulation (Fig 8). In autumn this value fell to 54%. In contrast, beached particles parametrized with Cd=0% were less abundant by the end of the simulations, particularly during spring with less than 25% of particles trapped in the shoreline.
- 360 When looking at the total amounts of beached particles per season, in summer over the 97% of particles parametrized with Cd=4% beached after one week of simulation (Fig 8). In autumn this value fell to 54%. In contrast, particles parametrized with Cd=0% take longer to arrive to the coastline, particularly during Spring with fewer than 25% of particles beached by the end of the simulations. Overall, the average of particles parametrized with Cd=0% was higher when comparing to Cd=4% (Fig 9). Particles released in French rivers and parametrized with Cd=0% were less abundant during summer, though
- 365 this trend was reversed in autumn. For Cd=0%, the number of particles released in Bidasoa river during summer were the least abundant after one week of simulation (<200 particles on average). The vast majority of particles released in Urumea river during winter were floating in the study area by the end of the simulations (479 particles on average). Particles parametrized with Cd=4% beached faster during the first 48 hours, mainly in summer and for those particles released in the French rivers. During this season, the average number of particles floating in the study area by the end of the simulation
- 370 ranged between 0 and 250 According to the temporal evolution of floating particles released per river, particles beached remarkably fast within the first 24 48 hours for Cd=4%, particularly those released during summer in French rivers. Similar behaviour pattern was observed within the same season between rivers, probably influenced by the vicinity of rivers and the spatiotemporal resolution of forcings (Fig 9).

Over 40% of the total particles parametrized with Cd=4% and almost 12% of parametrized with Cd=0% beached in 375 Gipuzkoa (Fig 10). During spring, almost 60% of beached particles parametrized with Cd=0% were located Bizkaia. For Cd=0%, particles released during summer in the rivers located in the western area of Gipuzkoa drifted longer distances and reached Landes shoreline. This trend changed during winter, when the vast majority of particles released in Gipuzkoa rivers beached mainly in Gipuzkoa and Bizkaia. Beached particles parametrized with Cd=0% experienced more seasonal variations derived from the surface current circulation patterns within the SE Bay of Biscay. For Cd=4%, particles beached in Gipuzkoa ranged between 51% in spring and 38% in winter and Bizkaia was the less affected region despite the season.

When looking the **seasonal trends** by river and region, beached particles were mainly found in Gipuzkoa for both Cd=4% and Cd=0% – 40.1% and 11.54% of the total particles released respectively –, particularly in winter after one week of simulations. For Cd=0%, beaching from particles released in Bidasoa, Nivelle and Adour River was higher in summer (9.01% particles released during summer) though this trend was reversed in autumn, when particles released in Basque rivers resulted in higher beaching. Overall, all regions were highly affected by rivers within or nearby the region itself (Fig 10).



Figure 5. Mean separation distance between modelled and observed trajectories for each wind drag coefficient. The dark line is the mean curve employedused for the trapezoidal integration.



Figure 6. Spatial mean distance between modeled and observed trajectories of buoy A, B, C and D with a drag coefficient Cd=4%. Particle trajectories were simulated during 24 h, with a re-initialization period every 4 hours. The modeled trajectories are shown in solid lines. Circles represents at the observed position the mean separation distance for all the modeled position





Figure 7. Particle concentration in Bizkaia, Gipuzkoa, Pyrénées Atlantiques and Landes coastlines. The seasonal distribution is shown for wind drag coefficient Cd=0% and Cd=4% after 24 hours and 168 hours of simulation

Figure 8. Total <u>Seasonal</u> amounts of beached particles <u>parametrized with Cd=0% and Cd=4% per season</u> after 168 hours of simulation. For wind drag coefficient Cd=0% and Cd=4%.





Figure 9. Temporal evolution of the particles <u>parametrized for Cd=0% and Cd=4% throughout the different seasons</u>. released by river during the simulation period for a wind drag coefficient Cd=0% and Cd=4%. The curves represent the <u>average</u> number of particles <u>floating</u> in the water surface by river and for every time step.





I

Figure 10. Seasonal analysis of the beached particles per region and river for wind drag coefficient Cd=0% and Cd=4% by the end of the simulation period. The nodes of the region correspond to the number of beached particles. The width of the node depicts the sum of the beached particles, and the links represent the number of particles beached per river

415 5 Discussion

5.1 Riverine litter composition

In this study, anAn artisanal net placed at the mouth of Deba river enabled sampling riverine litter provided a practical and tailored application for aggregating riverine-in the study area during Spring 2018. Short and narrow rivers prevail in the SE Bay of Biscay, particularly affected by a strong tidal regime, and very intense, stationary and persistent storms caused by a 420 combination of a warm sea, an unstable surface atmosphere and cold air at higher altitudes (Ocio et al., 2015). First field Setudies aiming at reporting the abundance and composition of floating riverine litter in European rivers date back less than 10 years and they were performed mainly in larger and more abundant rivers than Deba river. Despite the morphology and hydrological differences-between rivers, plastic was the predominat the distribution of items by type of material in Deba , river showed a clear predominance of plastic as observed in Siene (Gasperi et al., 2014), Danube (Lechner et al., 2014) or 425 Rhine River (van der Wal et al., 2015). Similarities were also found when comparing the Top ten list of riverine litter items to rivers located in the North-East Atlantic region. Plastic/polystyrene pieces between 2.5 cm and 50 cm (71.2%) top the list in terms of number of items, and their abundance was slightly higher when compared to North-East Atlantic rivers accounting for a greater proportion in Deba river (71.2%) than in North East Atlantic rivers (54.53%) (Bruge et al., 2018; Gonzalez-Fernandez et al., 2018). Lower abundances were observed in the Mediterranean (25.01%) and the Black Sea (13.74%). Riverine litter items trapped on vegetation or deposited on the riverbank can be degraded by weather conditions 430

- (15.1470) Average number of the intervention of the posted of the intervention of t
- than those of the Black Sea (13.74%) or the Mediterranean Sea (25.01%) can be attributed to a higher and faster fragmentation of riverine items along Deba river and the North East Atlantic basins. However, more detailed analyses on the physical characteristics of litter items (i.e., polymer type) are necessary to fully assess their impact on the occurrence of fragmented plastic pieces. Results are also in line with the ranking list of the Top ten beach litter items across the North-East Atlantic region revealing that Single Use Plastics (i.e. food containers, bottles and other packaging) are among the most
- 440 abundant riverine litter items together with plastic fragments (Addamo et al., 2017). These results differed from the analysis performed in sea small-scale convergence areas of floating marine litter ("*litter windrows*") on the coastal waters of the SE Bay of Biscay, where fishing-related items were the second most abundant sub-category in terms of number after *Plastic/polystyrene pieces between 2.5 cm and 50 cm* (Ruiz et al., 2020). Substantial differences also exist between riverine litter sampled in Deba river and floating marine litter assessed by visual observation from research vessels in open waters of
- the Bay of Biscay (Ruiz et al., 2022a). Differences might be related to the monitoring method and, also, to the size of the items, since small items, as plastic pieces, can be overlooked by the observer when visual counting method is applied, contrary to riverine litter samplings for later analysis at lab. Overall, riverine litter data acquisition is mainly focused on the floating fraction and the litter loads under the surface water are often ignore. Increasing the quantity of rivers sampled, the frequency and the riverine water compartments is necessary to establish the composition and trends of riverine litter in the SE Bay of Biscay.

5.2 Wind drag estimation

455

One of the largest uncertainties for simulating floating <u>riverine and marine</u> litter behaviour is the proper quantification of a wind drag coefficient. <u>Wind drag estimations conducted so far for floating marine litter items range between 0% and 6% (Ko et al., 2020; Critchell and Lambrechts, 2016; Neumann et al., 2014) with an upper limit of 10% (Yoon et al., 2010). <u>However, only a few of them have been validated using observational data (Maximenko et al., 2018; Callies et al., 2017).</u> <u>Empirical In this study</u>, data provided by "Low-cost buoys" combined with surface current measurements by HF radar were</u>

Con formato: Fuente: Sin Cursiva

can be used as a proxy for modelling predict the drift of floating litter objects with similar buoy characteristics (density, size and shape) in the study area. Results demonstrated that Cd=4% was the optimal wind drag coefficient for accurately represent the pathways of the "Low-cost buoys" in the study area. This value can be consistent with wind drag estimations 460 for the Bay of Biscay of the partially emerged Physalia physalis (Ferrer and Pastor, 2017) but it is almost three times higher than the maximum wind drag coefficient reported in the area by Pereiro et al., 2018 . This can be explained by the fact that buoys used in the experiment remained submerged beneath the sea surface and were less exposed to wind effect. The estimated wind drag coefficient was also greater than the Cd=3% observed for the Prestige oil spill accident (Abascal et al., 2009; Marta-Almeida et al., 2013). Indeed, oil spill studies refer to a range of wind drag coefficient between 2.5 to 4.4% of 165 the wind speed, with a mean value of 3 - 3.5% (e.g., ASCE, 1996; Reed et al., 1994). Oebject characteristics may change over the time due to the exposure to wind, waves, UV radiation, seawater and the attachment of organic material (Kooi et al., 2017; Min et al., 2020). Objects become breakable, and biofouling increases their density, overcoming the positive buoyancy and impacting on their trajectory. Investigations so far pinpointed longer time scales (weeks to months, and lager) than considered in this study (days) for a significant change on the behaviour of floating objects (Ryan, 2015; Fazey and Ryan, 470 2016). Consequently, physical variations on the buoy properties were not accounted for wind drag estimation. The separation distance between observed and modeled trajectories has been commonly used to evaluate the skill of particle-tracking models (Callies et al., 2017; Haza et al., 2019; Aksamit et al., 2020; Abascal et al., 2012), Commercial SPOT Trace devices have been used over the past few years in coastal and open ocean applications in a wide range of studies. Studies range from calibrating HF radars (Martínez Fernández et al., 2021), tracking drifting objects as icebergs (Carlson et al., 2020), pelagic 475 Sargassum (Putman et al., 2020; van Sebille et al., 2021) or fishing vessels (Widyatmoko et al., 2021; Hoenner et al., 2022), to search and rescue training (Russell, 2017) and oil spill and litter monitoring (Novelli et al., 2018; Meyerjürgens et al., 2019a; Mínguez et al., 2012; Abascal et al., 2015). In this study, the purpose was no to evaluate the model accuracy but estimated the wind drag coefficient for the "Low-cost buoys". However, the novel approach proposed by (Révelard et al., 2021) may be of particular interest for future experiments oriented to assess the wind drag coefficient of highly buoyant 180 items drifting during short time periods in the coastal area.-Nevertheless, object characteristics may change over the time due to the exposure to wind, waves, UV radiation, seawater and the attachment of organic material (Kooi et al., 2017; Min et al., 2020). Objects become breakable, and biofouling increases their density, overcoming the positive buoyancy and impacting on their trajectory. Investigations so far pinpointed longer time scales (weeks to months, and lager) than considered in this study (days) for a significant change on the behaviour of floating objects (Ryan, 2015; Fazey and Ryan, 2016). 185 Consequently, physical variations on the buoy properties were not accounted for wind drag estimation. The separation distance between observed and modeled trajectories has been commonly used to evaluate the skill of particle-tracking models (Callies et al., 2017; Haza et al., 2019; Aksamit et al., 2020; Abascal et al., 2012). In this study, the purpose was no to evaluate the model accuracy but estimated the wind drag coefficient for the "Low-cost buoys". However, the novel approach proposed by (Révelard et al., 2021) may be of particular interest for future experiments oriented to assess the wind 490 drag coefficient of highly buoyant items drifting during short time periods in the coastal area. The results obtained for Cd=4% can be consistent with wind drag estimations for the Bay of Biscay of the partially emerged Physalia physalis (Ferrer and Pastor, 2017) but greater than the Cd=3% observed for the Prestige oil spill-accident (Abascal et al., 2009; Marta-Almeida et al., 2013). Indeed, oil spill studies refer to a range of wind drag coefficient between 2.5 to 4.4% of the wind speed, with a mean value of 3 - 3.5% (e.g., ASCE, 1996; Reed et al., 1994). In this study, a wind drag value of 4% can 195 be expected due to the strong buoyancy of the "low-cost buoys" and can be applied for simulating the transport and fate of a specific group of litter items that share similar characteristics. However, due to the large heterogeneity of highly buoyant items, further experiments are needed to better parametrize the wind drag coefficient of different objects and consequently reduce the uncertainties on their behaviour.

Con formato: Inglés (Estados Unidos)
Con formato: Inglés (Estados Unidos)
Código de campo cambiado

Con formato: Inglés (Estados Unidos) Con formato: Inglés (Estados Unidos) Código de campo cambiado

5.3 Seasonal riverine litter distribution by region

500 It is broadly accepted that the SE Bay of Biscay is polluted with floating litter discarded or lost at the marine and coastal area but also with litter originated inland and transported via rivers and runoff. However, detailed studies on riverine litter contribution are still scarce and modelling efforts combining observations and physical parametrizations of floating riverine litter properties are non-existent. This study shows that the exposure to wind effect of riverine objects-largely control their transport and coastal accumulation of floating marine litter in the SE Bay of Biscay, with concentrations varying between 505 regions and over the time. Concentrations in Pyrénées-Atlantiques and Gipuzkoa regions diverged-differed widely from the other studied regions. Indeed, the highest concentrations occurred in both regions during summer for low buoyant (100-120 particle/km) and but also for highly buoyant items (>200 particles/km). Although largerA higher amounts of particles beached in Gipuzkoa during summer when compared to Pyrénées-Atlantiques but, concentrations wereare lower than Pyrénées Atlantiques since the coastline in the Basque shorelineregion is longer. Low buoyant pathways and fate reflect the 510 well-known surface water circulation patterns in the SE Bay of Biscay. The pathways and fate of low buoyant items reflect the seasonal surface water circulation patterns in the SE Bay of Biscay, Concentrations of floating riverine litter are therefore a direct consequence of the seasonal variability of floating drift and Results are in line with findings provided by (Declerck et al., 2019) who pinpointed a higher coastal retention in the area during spring and summer. Low buoyant objects not subjected to windage effects remain floating at the coastal waters and highly buoyant objects tended to beach remarkably 515 faster as reported in literature by (Rodríguez-Díaz et al., 2020). However, long-term data collected by in-situ observations of beached litter across the different regions are necessary to validate the large seasonal variations and to assess the reliability

of concentration levels for addressing riverine litter issue in priority regions with heavily polluted coastlines.

5.4 Rivers as key vectors of riverine litter

The interpretation of the spatial and temporal riverine litter distribution by river can be challenging since riverine litter fluxes in the study area are highly uncertain. In the study area, two major assumptions were made regarding the river systems: (1) same river discharge for all rivers and (2) same river discharge for all seasons. This means that same amounts of riverine litter were allocated for every river regardless the differences on the width and depth and the seasonal flow variations. Since each river basin has its own particularities, future modelling approaches should be adapted to the the morphology and hydrological conditions of the catchment area. Other drivers as the land use or socio-economic factors such economic status or population density can be a determining factor on the amount of mismanaged litter that could contribute to riverine litter fluxes (Schmidt et al., 2017; Schuyler et al., 2021). It is also necessary to further investigate if higher river flows in the area are directly related to an increased discharge of riverine litter since analysis already performed in different river basins show contradicting relations between the occurrence of riverine litter and river fluxes (van Emmerik and Schwarz, 2020). Along with the complex nature of qualifying riverine litter fluxes, litter behaviour in the coastal area of the SE Bay of Biscay is still in its early stage, and much has yet to be revealed. Particular attention should be paid to Pyrénées-Atlantiques and Gipuzkoa,

as main impacted regions in the studied area. The dominant number of rivers in this region can favour accumulation trends regardless the season. Regional coordination should be reinforced due to the transboundary movement of <u>floating</u> riverine litter in the study area and reasonable efforts oriented to retain or remove riverine litter as clean-up measures in the riverbanks should be investigated to avoid litter being transported to the coastal and marine environment.

535 5.5 Model limitations

The coastline of the SE Bay of Biscay is mainly covered by sand and muddy sand and characterized by the presence of moderate to high sea rocky cliffs, especially in the Basque region (ICES, 2019; Bilbao Lasa et al., 2020). The geomorphology can affect the retention of litter washing ashore. Sandy beaches tend to be more efficient at trapping and thus accumulating litter than rocky areas which favor litter fragmentation (Robbe et al., 2021; Weideman et al., 2020). Waves and

540	tides can also constrain coastal accumulation since they can resuspend litter and transport it back into the ocean (Brennan et
	al., 2018; Compa et al., 2022). Nevertheless, research on these processes is scarce and they cannot be resolved yet at a
	suitable resolution (Melvin et al., 2021). Consequently, in this study once particles beached, they were classified as it arrived
	to their final destination. It is, however, important to consider for future research in the study area the link between coastal
	accumulation, and the type of shoreline and resuspension, even though the model cannot yet simulate these processes. The
545	release location strongly influences where litter accumulates on the coastline. Litter items can beach rapidly when release
	locations are located near the coastline (Critchell et al., 2015). However, there is a big gap between the spatial resolution of
	ocean circulation models (up to 10 km spatial resolution) and the complex coastal accumulation processes. In this study, the
	release locations were located distant for the sources to avoid uncertainties on model performance at smaller scales.
	However, a greater model resolution with a finer grid can reinforce simulation results (NOAA, 2016). Nested models,
550	flowing from fine resolution near critical locations as the river mouths to open ocean resolution is a worthy issue for future
	consideration.

The interaction between floating litter and the shoreline is highly complex and relies in many processes including waves and tides. Indeed, waves and tides can constrain coastal accumulation since they can resuspend and transport litter back into the ocean (Brennan et al., 2018; Compa et al., 2022). The geomorphology can also affect the retention of litter washing ashore.

- Sandy beaches tend to be more efficient at trapping and accumulating litter than rocky areas, which favor litter fragmentation (Robbe et al., 2021; Weideman et al., 2020). How these processes contribute to the actual beaching is unknown and they cannot be resolved yet at a suitable resolution (Melvin et al., 2021). In this study, particles were released in open waters and once they reached the shoreline, they were classified as beached. The tidal effect and the wave-induced Stokes drift were not accounted to avoid introducing more uncertainties. However, further field and laboratory experiments to better understand on how these processes influence floating litter behaviour in the coastline is recommend. It is also important to consider for
- future research exploring the effect of the type of shoreline on coastal accumulation. In this study, a constant diffusion coefficient of 1 m²/s was considered as a pragmatic choice based on previously modelling work for floating marine litter. However, more field measurements are necessary to accurately assess the influence of the diffusion process on the transport of floating marine litter.

565 **5.6 Riverine litter collection and monitoring by a floating barrier**

Riverine litter quantities on a global scale urge countries to keep rivers pollution free, intercepting riverine litter before it reaches the ocean and minimizing the impact of marine pollution from land based sources. Research to date suggest that a significant reduction of marine litter in the ocean can be achieved with collection at rivers or with a combination of river barriers and clean up ocean devices (Hohn et al., 2020). Large scale and innovative removal initiatives (e.g., deployment of 570 interceptors at river mouths) are currently supporting cleanup actions worldwide on an experimental basis (Lindquist, 2016; Zhongming et al., 2019). At a smaller scale, oil spill booms or barriers have also been adapted to aggregate riverine litter in European river basins heavily exposed to the impacts of intense human activity, facilitating the collection and the analysis of litter composition (Gasperi et al., 2014). However, the efficiency of this type of devices is still not properly understood and can be conditioned by the wind, hydrology and morphological conditions of rivers (van Emmerik and Schwarz, 2020; 575 Andrés et al., 2021). Storms result in large flows of water and thus riverine litter fluxes to the coastal and marine environment. A well adapted device to storm specific events must be considered when deciding which tools implement for a cost effective plastic intervention strategy in the area. Further monitoring efforts are also required to account for seasonal variability on abundance and riverine litter typology. Within the LIFE LEMA project, two videometry systems were installed at the Oria and Adour river mouths and a detection algorithm was developed to monitor litter inputs in near real time 580 (Delpey et al., 2021; Ruiz et al., 2020b). Besides monitoring, information collected by the videometry systems can complement floating barriers collection and sampling and advise local authorities for a quick response on riverine litter

Con formato: Espacio Después: 12 pto

Con formato: Inglés (Estados Unidos)

contribution to coastal area during storm events. Monitoring tools based on visual observations as RIMMEL or CrowdWater apps (González Fernández, 2017; van Emmerik, 2020) can be also particularly helpful to build a database of riverine litter input to the SE Bay of Biscay so far remained limited or even non-existent, following a harmonized approach. Both data provided by cameras and visual observations can be crucial to evaluate the efficiency of mitigation measures as the installation of floating barriers as well as prevention measures applied inland the river basins for a successful reduction of litter inputs into the SE Bay of Biscay.

6 Conclusions

585

- The SE Bay of Biscay has been regarded described by global and regional models as an accumulation zone for floating 590 marine litter. However, detailed studies on floating riverine litter behaviour once items arrive to open waters are still scarce. but further improve understanding of floating macrolitter behaviour originated inland is required. Research on floating marine litter and pathways at sea are increasing but the understanding of the fate of floating macrolitter originated inland and transported through river systems is scarce and needs to be further studied. Based on HF radar current observations and wind 595 dataset for the period 2009 2021, this contribution tries to fill this gap by providing insights on how low and highly buoyant riverine litter released by several rivers of the SE Bay of Biscay may affect the nearby regions seasonally in terms of concentration and beaching. Analysis of riverine riverine litter_samples collected by a floating_barrier placed in the study area showed that low buoyant objects were predominant as riverine litter although highly buoyant objects were also relevant in terms of weight. Simulations for assessing the seasonal trends of floating riverine litter transport and fate were performed 600 with the Lagrangian model TESEO. To properly integrate the differences in litter buoyancy, simulations were parametrized with a wind drag coefficient for low and highly buoyant items. The wind drag for highly buoyant item was estimated by comparing the observed and the modelled positions of four drifters and turned out to be greater than the commonly assumed value for oil spill studies. The developed "Low-cost buoys" proved to be suitable to provide real time trajectories of highly buoyant objects exposed to wind. However,-but-drifters with different characteristics should be used in future studies for 605 accounting the windage effect on different type of items. The transport and fate of both highly and low buoyant items
- released by rivers was calculated by season. Highly buoyant items rapidly beached (in less than 48 hours), particularly in summer and winter; in contrast, despite the season over two thirds of low buoyant items remained floating after one week of being released. This highlights the discrepancy between behaviour for low and highly buoyant objects and the importance of parametrizing the windage effect in order to accurately predict riverine litter accumulation in the coastal area of the SE Bay
 of Biscay. Beached particles were mainly found in Gipuzkoa regardless the season and the wind drag coefficient. Overall, the less affected region was Bizkaia with the exception of Spring period for low buoyant items. Despite of the season, most

of the riverine litter remained in the study area and rivers polluted the regions within the river basin or surrounding. Investigating what beaches are most likely to accumulate large quantities and the contribution per river can provide relevant input to response operations after storm events in the short to medium term and can also support the identification of priority rivers for monitoring program, assisting in the future for an adapted intervention of riverine pollution regionally.

7 Appendices

Appendix A. Floating barrier for riverine litter collection



Appendix AFigure A1. Floating barrier (a) and installation in Deba river (Gipuzkoa) (b)

Appendix B. Riverine litter classification based on the exposure to wind effect

Appendix-Table B1. Data were gathered from surveys carried out during Spring 2018 in Deba river (Gipuzkoa)

TSG_ML General code	General name	Number of items	Weight (kg)
	Low buoyant items transported b	by currents	
G1	4/6-pack yokes, six-pack rings	1	3.3
G2	Bags	7	170.7
G3	Shopping bags incl. pieces	8	292.44
G4	Small plastic bags, e.g freezer bags	4	50.9
G5	What remains form rip-off plastic bags	21	186.31
G20-G24	Plastic caps and lids/Plastic rings	38	216.39
G26	Cigarrette lighters	1	9.7
G27	Cigarrette butts and filters	1	0.1
G30	Crisps packets/sweet wrappers	56	250.2
G31	Lolly sticks	1	2.4
G32	Toys and party poppers	2	97.5
G36	Fertilisers/animal feed bags	1	11.5
G48	Synthetic rope	2	6.7
G76	Plastic/polystyrene pieces 2.5 cm> < 50 cm	1122	1788.32
G77	Plastic/polystyrene > 50 cm	13	337.34
G96	Sanitary towels/panty liners/backing strips	35	1099.8
G100	Medical/Pharmaceutical containers/tubes	7	69.4

Con formato: Normal Con formato: Inglés (Estados Unidos)

Con formato: Centrado

Con formato: Inglés (Estados Unidos)

Con formato: Título 1

Tabla con formato

	Highly buoyant items transporte	ed by wind and o	currents	
		Total	91.12%	67.95%
G179	Bottle caps, lids & pull tabs		1	0
G177	Foil wrappers, aluminium foil		1	7
G173	Other (specify)		21	99.3
G159	Corks		4	21.2
G158	Other paper items		4	69.1
G156-157	Paper & Paper fragments		2	121.2
G148	Carboard (boxes & fragments)		3	85.7
G145	Other textiles (incl. rags)		7	320.5
G135	Clothing (clothes, shoes)		3	481.7
G134	Other rubber pieces		1	1.6
G125	Ballons and ballon sticks		5	1.1
G124	Other plastic/polystyrene items (identifiable)		58	1958.5
G101	Dog faeces bag		2	106

	inging buoyant items transported by wind an	ia currents	
G7	Drink bottles <= 0.5 l	5	142.6
G8	Drink bottles > 0.5 l	3	91.1
G9	Cleaner bottles & containers	2	105.7
G10	Food containers incl. Fast food containers	98	723.9
G11-12	Cosmetics bottles & other containers (shampoo, shower gel, deodorant)	4	100.3
G17	Injection gun containers	1	18.3
G33	Cups and cup lids	6	32.6
G150-151	Cartons/Tetrapack	2	121.2
G153	Cups, food trays, food wrappers, drink containers	4	69.1
G174	Aerosol/Spray cans industry	2	143.2
G175-176	Bottle caps, lids & pull tabs	2	5
G177	Bottles incl.Pieces	5	1832.3
G178	Light bulbs	1	31.7
	Total	8.88%	32.05 %

I

Tabla con formato

When		Periods selected	1 Detween 2002	9 and 2021 bas	ed on the avait	ability surface	current dataset	s provided by	une HF radar	
	Period 1	Period 2	Period 3	Period 4	Period 5	Period 6	Period 7	Period 8	Period 9	Period 10
Initial	07-Feb-2013	09-Mar-2021	23-Jan-2009	02-Jan-2013	18-Jan-2016	02-Jan-2014	17-Feb-2017	17-Jan-2012	22-Jan-2017	12-Jan-2021
date	08:00:00	22:00:00	01:00:00	11:00:00	17:00:00	15:00:00	06:00:00	00:00:60	17:00:00	23:00:00
Final	14-Feb-2013	16-Mar-2021	30-Jan-2009	09-Jan-2013	25-Jan-2016	09-Jan-2014	24-Feb-2017	24-Jan-2012	29-Jan-2017	19-Jan-2021
date	07:00:00	21:00:00	00:00:00	10:00:00	16:00:00	14:00:00	05:00:00	08:00:00	16:00:00	22:00:00
					Spring					
	Period 1	Period 2	Period 3	Period 4	Period 5	Period 6	Period 7	Period 8	Period 9	Period 10
Initial	14-Apr-2015	16-May-2012	16-Apr-2017	21-Apr-2012	05-Jun-2014	11-Apr-2021	06-May-2012	10-Apr-2015	08-May-2018	22-Apr-2016
date	23:00:00	00:00:00	14:00:00	08:00:00	06:00:00	20:00:00	00:00:90	08:00:00	22:00:00	11:00:00
Final	21-Apr-2015	22-May-2012	23-Apr-2017	28-Apr-2012	12-Jun-2014	18-Apr-2021	13-May-2012	17-Apr-2015	15-May-2018	29-Apr-2016
date	22:00:00	23:00:00	13:00:00	07:00:00	05:00:00	19:00:00	05:00:00	07:00:00	21:00:00	10:00:00
					Summer					
	Period 1	Period 2	Period 3	Period 4	Period 5	Period 6	Period 7	Period 8	Period 9	Period 10
Initial	19-Aug-2017	04-Jul-2015	15-Aug-2016	08-Aug-2012	14-Aug-2015	08-Sep-2013	11-Sep-2017	13-Sep-2015	08-Jul-2019	05-Aug-2014
date	01:00:00	16:00	18:00:00	11:00:00	00:00:00	23:00:00	11:00:00	02:00:00	4:00	20:00:00
Final	26-Aug-2017	11-Jul-2015	22-Aug-2016	15-Aug-2012	20-Aug-2015	15-Sep-2013	18-Sep-2017	20-Sep-2015	15/-Jul-2019	12-Aug-2014
date	00:00:00	15:00	17:00:00	10:00:00	23:00:00	22:00:00	10:00:00	01:00:00	3:00	19:00:00
					Autumn					
	Period 1	Period 2	Period 3	Period 4	Period 5	Period 6	Period 7	Period 8	Period 9	Period 10
Initial	16-Oct-2014	17-Oct-2011	24-Oct-2015	08-Nov-2011	10-Dec-2020	06/11/2015	23-Nov-2015	04-Oct-2017	04-Oct-2015	23-Nov-2020
date	22:00	8:00	11:00	17:00:00	10:00:00	1:00	21:00:00	23:00:00	20:00:00	04:00:00
Final	23-Oct-2014	24-Oct-2011	31-Oct-2015	15-Nov-2011	17-Dec-2020	13/11/2015	30-Nov-2015	11-Oct-2017	11-Oct-2015	30-Nov-2020
date	21:00	7:00	10:00	16:00:00	00:00:60	00:0	20:00:00	22:00:00	19:00:00	03:00:00

625 Appendix C. Selected seasonal week-long periods from the HF radar (2009-2021)

31

Con formato: Sangría: Izquierda: 0 cm

Con formato: Título 1



Appendix D. Seasonal mean current and wind fields (2009-2021)

1

L

I

AppendixFigure DI. Mean current (A) and wind fields (B) in the study area during each season for the selected periods between 2009 and 2021. The colour-bars represent the magnitude of current and wind speed. The arrows indicate the current and wind mean direction and are scaled with currents and wind speed (Data source: HFR – EuskOOS https://www.euskoos.eus/en/data/basque-ocean-meteorological-network/high-frequency-coastal-radars/ and ERA5 https://www.ecmwf.in/en/forecasts/faatasets/freand/size/seasts/faatasets/freand/seasts/faatasets/freand/seasts/faataset



635

8 Video supplement

Animations of the surface currents, winds and Lagrangian simulations area available for the study period 2009-2021. 9 Author contributions

Author contributions

IR: Investigation, formal analysis, visualization and writing – original draft preparation. AJ: Conceptualization, methodology, software, writing – review & editing. OCB: Conceptualization, supervision, resources, review and editing.
 AR: Conceptualization, methodology, supervision, resources, review and editing. All authors contributed to refining the manuscript for submission. This paper is part of the PhD research of IR supervised by OCB and AR.

10 Competing interests

The authors declare that they have no conflict of interest.

645 11 Financial support

This research has been partially funded through the EU's LIFE Program (LIFE LEMA project, grant agreement no. LIFE15 ENV/ES/000252) and by EU's H2020 Program (JERICO-S3 project, grant agreement No. 871153).

12 Acknowledgements

- We are grateful to the Emergencies and Meteorology Directorate Security department Basque Government for public data provision from the Basque Operational Oceanography System EuskOOS. Authors also acknowledge support from the JERICO-S3 project, funded by the European Union's Horizon 2020 Research and Innovation Program under grant agreement no. 871153. This study has been conducted using EU Copernicus Marine Service information. We thank Luis Ferrer for sharing his valuable knowledge on custom-built the drifters. We thank Cristina Barreau, Antoine Bruge, Igor Granado, Théo Destang, Alix McDaid and Jon Andonegi for their support with drifters' releasing. We thank the citizens who collected and reported drifters' arrival to Basque and French coasts. This is contribution number XXX of AZTI - Marine
- Research (BRTA).

13 Review statement

This paper was edited by XXX and reviewed by XXX anonymous referees.

14 References

Abascal, A., Castanedo, S., Gutierrez, A. D., Comerma, E., Medina, R., and Losada, I. J.: Teseo, an operational system for simulating oil spills trajectories and fate processes, Proc. Int. Offshore Polar Eng. Conf., 1751–1758, 2007.

Abascal, A. J., Castanedo, S., Mendez, F. J., Medina, R., and Losada, I. J.: Calibration of a Lagrangian Transport Model Using Drifting Buoys Deployed during the *Prestige* Oil Spill, J. Coast. Res., 2009, 11,80-90, 2009.

Abascal, A. J., Castanedo, S., Fernández, V., and Medina, R.: Backtracking drifting objects using surface currents from highfrequency (HF) radar technology, Ocean Dyn., 62, 1073–1089, https://doi.org/10.1007/s10236-012-0546-4, 2012.

Abascal, A. J., Castanedo, S., Minguez, R., Medina, R., Liu, Y., and Weisberg, R.: Stochastic Lagrangian Trajectory Modeling of Surface Drifters deployed during the Deepwater Horizon Oil Spill, in: Proceedings of the 38th AMOP Technical Seminar on Environmental Contamination and Response, 77–91, 2015.

 Abascal, A. J., Castanedo, S., Núñez, P., Mellor, A., Clements, A., Pérez, B., Cárdenas, M., Chiri, H., and Medina, R.: A
 high-resolution operational forecast system for oil spill response in Belfast Lough, Mar. Pollut. Bull., 114, 302–314, https://doi.org/https://doi.org/10.1016/j.marpolbul.2016.09.042, 2017a.

Abascal, A. J., Sanchez, J., Chiri, H., Ferrer, M. I., Cárdenas, M., Gallego, A., Castanedo, S., Medina, R., Alonso-Martirena, A., Berx, B., Turrell, W. R., and Hughes, S. L.: Operational oil spill trajectory modelling using HF radar currents: A northwest European continental shelf case study, Mar. Pollut. Bull., 119, 336–350, https://doi.org/https://doi.org/10.1016/j.marpolbul.2017.04.010, 2017b.

Addamo, A. M., Laroche, P., and Hanke, G.: Top marine beach litter items in Europe, Publ. Off. Eur. Union Luxembg., 2017.

Aksamit, N. O., Sapsis, T., and Haller, G.: Machine-Learning Mesoscale and Submesoscale Surface Dynamics from Lagrangian Ocean Drifter Trajectories, J. Phys. Oceanogr., 50, 1179–1196, https://doi.org/10.1175/jpo-d-19-0238.1, 2020.

680 Al-Zawaidah, H., Ravazzolo, D., and Friedrich, H.: Macroplastics in rivers: present knowledge, issues and challenges, Environ. Sci. Process. Impacts, 23, 535–552, https://doi.org/10.1039/d0em00517g, 2021.

Allshouse, M. R., Ivey, G. N., Lowe, R. J., Jones, N. L., Beegle-Krause, C. J., Xu, J., and Peacock, T.: Impact of windage on ocean surface Lagrangian coherent structures, Environ. Fluid Mech., 17, 473–483, https://doi.org/10.1007/s10652-016-9499-3, 2017.

685 ASCE: State-of-the-Art Review of Modeling Transport and Fate of Oil Spills, J. Hydraul. Eng., 122, 594–609, https://doi.org/10.1061/(ASCE)0733-9429(1996)122:11(594), 1996.

Blettler, M. C. M., Abrial, E., Khan, F. R., Sivri, N., and Espinola, L. A.: Freshwater plastic pollution: Recognizing research biases and identifying knowledge gaps, Water Res., 143, 416–424, https://doi.org/10.1016/j.watres.2018.06.015, 2018.

690 Bourillet, J. F., Zaragosi, S., and Mulder, T.: The French Atlantic margin and deep-sea submarine systems, Geo-Marine Lett., 26, 311–315, https://doi.org/10.1007/s00367-006-0042-2, 2006.

Le Boyer, A., Charria, G., Le Cann, B., Lazure, P., and Marié, L.: Circulation on the shelf and the upper slope of the Bay of Biscay, Cont. Shelf Res., 55, 97–107, 2013.

Brennan, E., Wilcox, C., and Hardesty, B. D.: Connecting flux, deposition and resuspension in coastal debris surveys, Sci.
 Total Environ., 644, 1019–1026, https://doi.org/10.1016/J.SCITOTENV.2018.06.352, 2018.

Bruge, A., Barreau, C., Carlot, J., Collin, H., Moreno, C., and Maison, P.: Monitoring Litter Inputs from the Adour River (Southwest France) to the Marine Environment, Mar. Sci. Eng., 6(1), 24, https://doi.org/10.3390/jmse6010024, 2018.

C3S: ERA5: Fifth Generation of ECMWF Atmospheric Reanalyses of the Global Climate., https://cds.climate.copernicus.eu/cdsapp#1/home, 2019.

700 Callies, U., Groll, N., Horstmann, J., Kapitza, H., Klein, H., Maßmann, S., and Schwichtenberg, F.: Surface drifters in the German Bight: model validation considering windage and Stokes drift, Ocean Sci., 13, 799–827, https://doi.org/10.5194/os-13-799-2017, 2017.

Carlson, D. F., Pavalko, W. J., Petersen, D., Olsen, M., and Hass, A. E.: Maker Buoy Variants for Water Level Monitoring and Tracking Drifting Objects in Remote Areas of Greenland, 20, 1254, 2020.

705 Carson, H. S., Lamson, M. R., Nakashima, D., Toloumu, D., Hafner, J., Maximenko, N., and McDermid, K. J.: Tracking the sources and sinks of local marine debris in Hawai 'i, Mar. Environ. Res., 84, 76–83, 2013.

Chamas, A., Moon, H., Zheng, J., Qiu, Y., Tabassum, T., Hee Jang, J., Abu-Omar, M., Scott, S. L., and Suh, S.: Degradation Rates of Plastics in the Environment, Cite This ACS Sustain. Chem. Eng, 8, 3511, https://doi.org/10.1021/acssuschemeng.9b06635, 2020.

710 Charria, G., Lazure, P., Le Cann, B., Serpette, A., Reverdin, G., Louazel, S., Batifoulier, F., Dumas, F., Pichon, A., and Morel, Y.: Surface layer circulation derived from Lagrangian drifters in the Bay of Biscay, J. Mar. Syst., 109, 60, https://doi.org/10.1016/j.jmarsys.2011.09.015, 2013.

Chiri, H., Abascal, A. J., Castanedo, S., and Medina, R.: Mid-long term oil spill forecast based on logistic regression modelling of met-ocean forcings, Mar. Pollut. Bull., 146, 962–976,
https://doi.org/https://doi.org/10.1016/j.marpolbul.2019.07.053, 2019.

Chiri, H., Abascal, A. J., and Castanedo, S.: Deep oil spill hazard assessment based on spatio-temporal met-ocean patterns, Mar. Pollut. Bull., 154, 111123, https://doi.org/10.1016/J.MARPOLBUL.2020.111123, 2020.

Compa, M., Alomar, C., Morató, M., Álvarez, E., and Deudero, S.: Spatial distribution of macro- and micro-litter items along rocky and sandy beaches of a Marine Protected Area in the western Mediterranean Sea, Mar. Pollut. Bull., 178, 113520, https://doi.org/10.1016/J.MARPOLBUL.2022.113520, 2022.

Critchell, K. and Lambrechts, J.: Modelling accumulation of marine plastics in the coastal zone; what are the dominant physical processes? , Estuar. Coast. Shelf Sci., 171, 111, 2016.

D'Asaro, E. A., Carlson, D. F., Chamecki, M., Harcourt, R. R., Haus, B. K., Fox-Kemper, B., Molemaker, M. J., Poje, A. C., and Yang, D.: Advances in Observing and Understanding Small-Scale Open Ocean Circulation During the Gulf of Mexico Research Initiative Era, Front. Mar. Sci., 7, https://doi.org/10.3389/fmars.2020.00349, 2020.

Davila, X., Rubio, A., Artigas, L. F., Puillat, I., Manso-Narvarte, I., Lazure, P., and Caballero, A.: Coastal submesoscale processes and their effect on phytoplankton distribution in the southeastern Bay of Biscay, Ocean Sci., 17, 849–870, https://doi.org/10.5194/os-17-849-2021, 2021.

Declerck, A., Delpey, M., Rubio, A., Ferrer, L., Basurko, O. C., Mader, J., and Louzao, M.: Transport of floating marine
 litter in the coastal area of the south-eastern Bay of Biscay: A Lagrangian approach using modelling and observations, J.
 Oper. Oceanogr., 1–15, https://doi.org/10.1080/1755876x.2019.1611708, 2019.

Delpey, M., Declerck, A., Epelde, I., Voirand, T., Manso-Navarte, I., Mader, J., Rubio, A., and Caballero, A.: Tracking floating marine litter in the coastal area by combining operational ocean modelling and remote observation systems., EGU Gen. Assem. 2021, online, 19–30 Apr 2021, EGU21-11465, 2021.

735 Duhec, A. V, Jeanne, R. F., Maximenko, N., and Hafner, J.: Composition and potential origin of marine debris stranded in the Western Indian Ocean on remote Alphonse Island, Seychelles, Mar. Pollut. Bull., 96, 76–86, https://doi.org/https://doi.org/10.1016/j.marpolbul.2015.05.042, 2015.

van Emmerik, T. and Schwarz, A.: Plastic debris in rivers, 7, e1398, https://doi.org/https://doi.org/10.1002/wat2.1398, 2020.

Eurostat: Population density by NUTS 3 region, https://appsso.eurostat.ec.europa.eu/nui/submitViewTableAction.do, 2019.

740 Fazey, F. M. C. and Ryan, P. G.: Biofouling on buoyant marine plastics: An experimental study into the effect of size on surface longevity, Environ. Pollut., 210, 354–360, https://doi.org/10.1016/J.ENVPOL.2016.01.026, 2016.

Ferrer, L. and Pastor, A.: The Portuguese man-of-war: Gone with the wind, Reg. Stud. Mar. Sci., 14, 53-62, https://doi.org/https://doi.org/10.1016/j.rsma.2017.05.004, 2017.

Galgani, F., Hanke, G., Werner, S., Oosterbaan, L., Nilsson, P., Fleet, D., Kinsey, S., Thompson, R. C., Van Franeker, J., and
 Vlachogianni, T.: Guidance on monitoring of marine litter in European Seas, Publications Office of the European Union, 2013.

Gasperi, J., Dris, R., Bonin, T., Rocher, V., and Tassin, B.: Assessment of floating plastic debris in surface water along the Seine River, Environ. Pollut., 195, 163–166, https://doi.org/https://doi.org/10.1016/j.envpol.2014.09.001, 2014.

González-Fernández, D. and Hanke, G.: Toward a Harmonized Approach for Monitoring of Riverine Floating Macro Litter Inputs to the Marine Environment, Front. Mar. Sci., 4, https://doi.org/10.3389/fmars.2017.00086, 2017.

González-Fernández, D., Cózar, A., Hanke, G., Viejo, J., Morales-Caselles, C., Bakiu, R., Barceló, D., Bessa, F., Bruge, A., Cabrera, M., Castro-Jiménez, J., Constant, M., Crosti, R., Galletti, Y., Kideys, A. E., Machitadze, N., Pereira de Brito, J., Pogojeva, M., Ratola, N., Rigueira, J., Rojo-Nieto, E., Savenko, O., Schöneich-Argent, R. I., Siedlewicz, G., Suaria, G., and Tourgeli, M.: Floating macrolitter leaked from Europe into the ocean, Nat. Sustain., 4, 474–483, https://doi.org/10.1038/s41893-021-00722-6, 2021.

- Gonzalez Fernandez, D., Hanke, G., Kideys, A., Navarrao Ortega, A., Sanchez Vidal, A., Bruge, A., Öztürk, B., Palma, C., Santelli, C., Duijsings, D., Barcelo, D., Dimitiriu, E., Rojo-Nieto, E., Ferreira, F., Bessa, F., Suaria, G., Siedlewicz, G., Castro Jimenez, J., Germano, J., Pereira De Brito, J., Rigueira, J., Pazdro, K., Cabrera, M., Pogojeva, M., Köck Schulmeyer, M., Constant, M., Canals Artigas, M., Paraboschi, M., Tourgeli, M., Machitadze, N., Ratola, N., Savenko, O., Kerherve, P.,
- 760 Sempere, R., Bakiu, R., Crosti, R., Schoeneich-Argent, R., Landry Levesque, S., Agostinho, T., Segal, Y., and Galletti, Y.: Floating Macro Litter in European Rivers - Top Items, EUR 29383 EN, Publications Office of the European Union, Luxembourg, 2018, ISBN 978-92-79-96373-5, doi:10.2760/316058, Luxembourg, https://doi.org/10.2760/316058, 2018.

González, M., Valencia, V., Mader, J., Fontán, A., Uriarte, A., and Caballero, A.: Operational Coastal Systems in the Basque Country Region: Modelling and Observations, Proc. Int. Offshore Polar Eng. Conf., 2007.

765 Haza, A. C., Paldor, N., Özgökmen, T. M., Curcic, M., Chen, S. S., and Jacobs, G.: Wind-Based Estimations of Ocean Surface Currents From Massive Clusters of Drifters in the Gulf of Mexico, J. Geophys. Res. Ocean., 124, 5844–5869, https://doi.org/https://doi.org/10.1029/2018JC014813, 2019.

770

Hernández-Carrasco, I., Solabarrieta, L., Rubio, A., Esnaola, G., Reyes, E., and Orfila, A.: Impact of HF radar current gapfilling methodologies on the Lagrangian assessment of coastal dynamics, Ocean Sci., 14, 827–847, https://doi.org/10.5194/os-14-827-2018, 2018.

Hoenner, X., Barlian, E., Ernawati, T., Hardesty, B. D., Kembaren, D. D., Mous, P. J., Sadiyah, L., Satria, F., and Wilcox, C.: Using anti-theft tracking devices to infer fishing vessel activity at sea, Fish. Res., 249, 106230, https://doi.org/https://doi.org/10.1016/j.fishres.2022.106230, 2022.

Hunter, J. R., Craig, P. D., and Phillips, H. E.: On the use of random walk models with spatially variable diffusivity, J.
 Comput. Phys., 106, 366–376, https://doi.org/10.1016/S0021-9991(83)71114-9, 1993.

Con formato: Español (España)

Kaplan, D. M. and Lekien, F.: Spatial interpolation and filtering of surface current data based on open-boundary modal analysis, J. Geophys. Res. Ocean., 112, https://doi.org/https://doi.org/10.1029/2006JC003984, 2007.

Karagiorgos, J., Vervatis, V., and Sofianos, S.: The Impact of Tides on the Bay of Biscay Dynamics, J. Mar. Sci. Eng., 8, 617, 2020.

780 Ko, C.-Y., Hsin, Y.-C., and Jeng, M.-S.: Global distribution and cleanup opportunities for macro ocean litter: a quarter century of accumulation dynamics under windage effects, Environ. Res. Lett., 15, 104063, https://doi.org/10.1088/1748-9326/abae29, 2020.

Kooi, M., Nes, E. H. van, Scheffer, M., and Koelmans, A. A.: Ups and Downs in the Ocean: Effects of Biofouling on Vertical Transport of Microplastics, Environ. Sci. Technol., 51, 7963–7971, https://doi.org/10.1021/acs.est.6b04702, 2017.

785 Lavin, A., Valdés, L., Sanchez, F., Abaunza, P., Forest, A., Boucher, J., Lazure, P., and Jegou, A.-M.: The Bay of Biscay: the encountering of the Ocean and the Shelf (18b, E), Combridge, MA, Harvard University Press, 2006.

Lebreton, L., Egger, M., and Slat, B.: A global mass budget for positively buoyant macroplastic debris in the ocean, Sci. Rep., 9, 1, 2019.

Lebreton, L. C. M., van der Zwet, J., Damsteeg, J.-W., Slat, B., Andrady, A., and Reisser, J.: River plastic emissions to the world's oceans, Nat. Commun., 8, 15611, https://doi.org/10.1038/ncomms15611, 2017.

Lechner, A., Keckeis, H., Lumesberger-Loisl, F., Zens, B., Krusch, R., Tritthart, M., Glas, M., and Schludermann, E.: The Danube so colourful: A potpourri of plastic litter outnumbers fish larvae in Europe's second largest river, Environ. Pollut., 188, 177–181, https://doi.org/10.1016/J.ENVPOL.2014.02.006, 2014.

Maclean, K., Weideman, E. A., Perold, V., and Ryan, P. G.: Buoyancy affects stranding rate and dispersal distance of floating litter entering the sea from river mouths, Mar. Pollut. Bull., 173, 113028, https://doi.org/https://doi.org/10.1016/j.marpolbul.2021.113028, 2021.

Maier-Reimer, E. and Sündermann, J.: On tracer methods in computational hydrodynamics, in: Engineering applications of computational hydraulics, 198–216, 1982.

 Manso-Narvarte, I., Caballero, A., Rubio, A., Dufau, C., and Birol, F.: Joint analysis of coastal altimetry and high-frequency
 (HF) radar data: Observability of seasonal and mesoscale ocean dynamics in the Bay of Biscay, Ocean Sci., 14, 1265–1281, https://doi.org/10.5194/os-14-1265-2018, 2018.

Manso-Narvarte, I., Rubio, A., Jordà, G., Carpenter, J., Merckelbach, L., and Caballero, A.: Three-Dimensional Characterization of a Coastal Mode-Water Eddy from Multiplatform Observations and a Data Reconstruction Method, Remote Sens., 13, 674, https://doi.org/10.3390/rs13040674, 2021.

805 Mantovani, C., Corgnati, L., Horstmann, J., Rubio, A., Reyes, E., Quentin, C., Cosoli, S., Asensio, J. L., Hermes, J., Mader, J., and Griffa, A.: Best Practices on High Frequency Radar Deployment and Operation for Ocean Current Measurement, Front. Mar. Sci., https://doi.org/10.3389/fmars.2020.00210, 2020.

Margenat, H., Ruiz-Orejón, L. F., Cornejo, D., Martí, E., Vila, A., Le Roux, G., Hansson, S., and Guasch, H.: Guidelines of field-tested procedures and methods for monitoring plastic litter in Mountain riverine systems, 2021.

810 Marta-Almeida, M., Ruiz-Villarreal, M., Pereira, J., Otero, P., Cirano, M., Zhang, X., and Hetland, R. D.: Efficient tools for marine operational forecast and oil spill tracking, Mar. Pollut. Bull., 71, 139–151, https://doi.org/10.1016/J.MARPOLBUL.2013.03.022, 2013.

Martínez Fernández, A., Redondo Caride, W., Alonso Pérez, F., Piedracoba Varela, S., Lorente Jiménez, P., Montero Vilar, P., Torres López, S., Fernández Baladrón, A., Varela Benvenuto, R. A., and Velo Lanchas, A.: SPOT and GPRS drifting buoys for HF Radar calibration, Instrum. Viewp., 48–49, 2021.

Maximenko, N., Hafner, J., Kamachi, M., and MacFadyen, A.: Numerical simulations of debris drift from the Great Japan Tsunami of 2011 and their verification with observational reports, Mar. Pollut. Bull., 132, 5, 2018.

Mazarrasa, I., Puente, A., Núñez, P., García, A., Abascal, A. J., and Juanes, J. A.: Assessing the risk of marine litter accumulation in estuarine habitats, Mar. Pollut. Bull., 144, 117–128, https://doi.org/https://doi.org/10.1016/j.marpolbul.2019.04.060, 2019.

Meijer, L. J. J., Emmerik, T. van, Ent, R. van der, Schmidt, C., and Lebreton, L.: More than 1000 rivers account for 80% of global riverine plastic emissions into the ocean, Sci. Adv., 7, eaaz5803, https://doi.org/doi:10.1126/sciadv.aaz5803, 2021.

Melvin, J., Bury, M., Ammendolia, J., Charles, M., and Liboiron, M.: Critical Gaps in Shoreline Plastics Pollution Research,

Front. Mar. Sci., 8, https://doi.org/10.3389/fmars.2021.689108, 2021.

825 Meyerjürgens, J., Badewien, T. H., Garaba, S. P., Wolff, J.-O., and Zielinski, O.: A State-of-the-Art Compact Surface Drifter Reveals Pathways of Floating Marine Litter in the German Bight, Front. Mar. Sci., 6, https://doi.org/10.3389/fmars.2019.00058, 2019a.

Meyerjürgens, J., Badewien, T. H., Zielinski, O., Braun, A., and Butter, M.: Track of GPS-Drifter North_Sea_Drifter7 in the German Bight, Southern North Sea, https://doi.pangaea.de/10.1594/PANGAEA.897995, 2019b.

830 Van Der Mheen, M., Pattiaratchi, C., and Van Sebille, E.: Role of Indian Ocean Dynamics on Accumulation of Buoyant Debris, J. Geophys. Res. Ocean., 124, 2571–2590, https://doi.org/10.1029/2018JC014806, 2019.

Min, K., Cuiffi, J. D., and Mathers, R. T.: Ranking environmental degradation trends of plastic marine debris based on physical properties and molecular structure, Nat. Commun., 11, 727, https://doi.org/10.1038/s41467-020-14538-z, 2020.

Mínguez, R., Abascal, • A J, Castanedo, • S, and Medina, • R: Stochastic Lagrangian trajectory model for drifting objects in the ocean, Stoch. Environ. Res. Risk Assess., 26, 1081–1093, https://doi.org/10.1007/s00477-011-0548-7, 2012.

Neumann, D., Callies, U., and Matthies, M.: Marine litter ensemble transport simulations in the southern North Sea, Mar. Pollut. Bull., 86, 219–228, https://doi.org/https://doi.org/10.1016/j.marpolbul.2014.07.016, 2014.

Novelli, G., Guigand, C. M., Cousin, C., Ryan, E. H., Laxague, N. J. M., Dai, H., Haus, B. K., and Özgökmen, T. M.: A Biodegradable Surface Drifter for Ocean Sampling on a Massive Scale, J. Atmos. Ocean. Technol., 34, 2509–2532, https://doi.org/10.1175/jtech-d-17-0055.1, 2017.

Novelli, G., Guigand, C. M., and Özgökmen, T. M.: Technological Advances in Drifters for Oil Transport Studies, Mar. Technol. Soc. J., 52, 53–61, https://doi.org/10.4031/mtsj.52.6.9, 2018.

Núñez, P., García, A., Mazarrasa, I., Juanes, J. A., Abascal, A. J., Méndez, F., Castanedo, S., and Medina, R.: A methodology to assess the probability of marine litter accumulation in estuaries, Mar. Pollut. Bull., 144, 309–324, https://doi.org/https://doi.org/10.1016/j.marpolbul.2019.04.077, 2019.

Ocio, D., Stocker, C., Eraso, A., and Cowpertwait, P.: Regionalized extreme flows by means of stochastic storm generation coupled with a distributed hydrological model. The case of the Basque Country., Proc. 36th IAHR World Congr. 28 June – 3 July, 2015., 1–13, https://doi.org/10.13140/RG.2.1.3924.4889, 2015.

Pereiro, D., Souto, C., and Gago, J.: Calibration of a marine floating litter transport model, J. Oper. Oceanogr., 11, 125–133, https://doi.org/10.1080/1755876x.2018.1470892, 2018.

Pereiro, D., Souto, C., and Gago, J.: Dynamics of floating marine debris in the northern Iberian waters: A model approach, J. Sea Res., 144, 57–66, https://doi.org/https://doi.org/10.1016/j.seares.2018.11.007, 2019.

Porter, M., Inall, M. E., Green, J. A. M., Simpson, J. H., Dale, A. C., and Miller, P. I.: Drifter observations in the summer time Bay of Biscay slope current, J. Mar. Syst., 157, 65–74, 2016.

Puillat, I., Lazure, P., Anne-marie, J., Lampert, L., and Miller, P.: Mesoscale hydrological variability induced by northwesterly wind on the French continental shelf of the Bay of Biscay, Sci. Mar. (Institut Ciències del Mar Barcelona, CSIC), 2006-06, Vol. 70, N. 1, P. 15-26, 70, 2006.

Putman, N. F., Lumpkin, R., Olascoaga, M. J., Trinanes, J., and Goni, G. J.: Improving transport predictions of pelagic Sargassum, J. Exp. Mar. Bio. Ecol., 529, 151398, https://doi.org/https://doi.org/10.1016/j.jembe.2020.151398, 2020.

860 Reed, M., Turner, C., and Odulo, A.: The role of wind and emulsification in modelling oil spill and surface drifter trajectories, Spill Sci. Technol. Bull., 1, 143–157, https://doi.org/10.1016/1353-2561(94)90022-1, 1994.

865

Révelard, A., Reyes, E., Mourre, B., Hernández-Carrasco, I., Rubio, A., Lorente, P., Fernández, C. D. L., Mader, J., Álvarez-Fanjul, E., and Tintoré, J.: Sensitivity of Skill Score Metric to Validate Lagrangian Simulations in Coastal Areas: Recommendations for Search and Rescue Applications, Front. Mar. Sci., 8, https://doi.org/10.3389/fmars.2021.630388, 2021.

Rizal, A., Gautama, B. G., Pranowo, W. S., Farhan, A. R., Siong, K., Harjono, M. R., Voisin, J. B., Maes, C., Dobler, D., Berlianty, D., Priyono, B., Dufau, C., Lucas, M., Fauny, O., and Rahmania, R.: Tracking the Stranded Area of Marine Debris in Indonesian coasts by using Floating Drifter, IOP Conf. Ser. Earth Environ. Sci., 925, 12034, https://doi.org/10.1088/1755-1315/925/1/012034, 2021.

870 Robbe, E., Woelfel, J., Arūnas Balčiūnas, •, and Schernewski, • Gerald: An Impact Assessment of Beach Wrack and Litter on Beach Ecosystem Services to Support Coastal Management at the Baltic Sea, Environ. Manage., 68, 835–859, https://doi.org/10.1007/s00267-021-01533-3, 2021.

Rodríguez-Díaz, L., Gómez-Gesteira, J. L., Costoya, X., Gómez-Gesteira, M., and Gago, J.: The Bay of Biscay as a trapping zone for exogenous plastics of different sizes, J. Sea Res., 163, 101929, https://doi.org/https://doi.org/10.1016/j.seares.2020.101929, 2020.

Rodríguez, J. G., Garmendia, J. M., Muxika, I., Gómez-Ballesteros, M., Quincoces, I., Díez, I., Arrese, B., Sánchez, F., and Galparsoro, I.: Macrofaunal variability in the continental shelf and canyons in the southeastern Bay of Biscay, Reg. <u>Stud.</u> Mar. Sci., 48, 102012, https://doi.org/https://doi.org/10.1016/j.rsma.2021.102012, 2021.

Rubio, A., Reverdin, G., Fontán, A., González, M., and Mader, J.: Mapping near-inertial variability in the SE Bay of Biscay
 from HF radar data and two offshore moored buoys, Geophys. Res. Lett., 38, 1–6, https://doi.org/10.1029/2011GL048783, 2011.

Rubio, A., Solabarrieta, L., Gonzalez, M., Mader, J., Castanedo, S., Medina, R., Charria, G., and Aranda, J. A.: Surface circulation and Lagrangian transport in the SE Bay of Biscay from HF radar data, Ocean. 2013 MTS/IEEE Bergen Challenges North. Dimens., https://doi.org/10.1109/OCEANS-Bergen.2013.6608039, 2013.

- 885 Rubio, A., Mader, J., Corgnati, L., Mantovani, C., Griffa, A., Novellino, A., Quentin, C., Wyatt, L., Schulz-Stellenfleth, J., Horstmann, J., Lorente, P., Zambianchi, E., Hartnett, M., Fernandes, C., Zervakis, V., Gorringe, P., Melet, A., and Puillat, I.: HF Radar Activity in European Coastal Seas: Next Steps toward a Pan-European HF Radar Network, Front. Mar. Sci., 4, https://doi.org/10.3389/fmars.2017.00008, 2017.
- Rubio, A., Caballero, A., Orfila, A., Hernández-Carrasco, I., Ferrer, L., González, M., Solabarrieta, L., and Mader, J.: Eddyinduced cross-shelf export of high Chl-a coastal waters in the SE Bay of Biscay, Remote Sens. Environ., 205, 290–304, https://doi.org/https://doi.org/10.1016/j.rse.2017.10.037, 2018.

Rubio, A., Hernández-Carrasco, I., Orfila, A., González, M., Reyes, E., Corgnati, L., Berta, M., Griffa, A., and Mader, J.: A Lagrangian approach to monitor local particle retention conditions in coastal areas, J. Oper. Oceanogr., 13:sup1, https://doi.org/10.1080/1755876X.2020.1785097, 2020.

895 Ruiz, I., Basurko, O. C., Rubio, A., Delpey, M., Granado, I., Declerck, A., Mader, J., and Cózar, A.: Litter Windrows in the South-East Coast of the Bay of Biscay: An Ocean Process Enabling Effective Active Fishing for Litter, Front. Mar. Sci., 7, https://doi.org/10.3389/fmars.2020.00308, 2020.

Ruiz, I., Abascal, A. J., Basurko, O. C., and Rubio, A.: Modelling the distribution of fishing-related floating marine litter within the Bay of Biscay and its marine protected areas, Environ. Pollut., 292, 118216, https://doi.org/https://doi.org/10.1016/j.envpol.2021.118216, 2022a.

Ruiz, I., Ana J, A., Basurko, O. C., and Rubio, A.: Modelling the distribution of fishing-related floating marine litter within the Bay of Biscay and its marine protected areas, Environ. Pollut., 292, 118216, https://doi.org/https://doi.org/10.1016/j.envpol.2021.118216, 2022b.

Russell, K.: Spain's Coastal Authority Uses Spot Trace for Search and Rescue Training, 905 https://www.satellitetoday.com/telecom/2017/06/20/spains-coastal-authority-uses-spot-trace-search-rescue-training/, 2017.

Ryan, P. G.: Does size and buoyancy affect the long-distance transport of floatingdebris?Â, Environ. Res. Lett., 10, 1, 2015.

Schmidt, C., Krauth, T., and Wagner, S.: Export of Plastic Debris by Rivers into the Sea, Environ. Sci. Technol., 51, 12246–12253, https://doi.org/10.1021/acs.est.7b02368, 2017.

 Schuyler, Q., Wilcox, C., Lawson, T. J., M K P Ranatunga, R. R., Hu, C.-S., Plastics Project Partners, G., and Denise
 Hardesty, B.: Human Population Density is a Poor Predictor of Debris in the Environment, Front. Environ. Sci., 9, https://doi.org/10.3389/fenvs.2021.583454, 2021.

van Sebille, E., Zettler, E., Wienders, N., Amaral-Zettler, L., Elipot, S., and Lumpkin, R.: Dispersion of Surface Drifters in the Tropical Atlantic, Front. Mar. Sci., 7, https://doi.org/10.3389/fmars.2020.607426, 2021.

Van Sebille, E., Aliani, S., Law, K. L., Maximenko, N., Alsina, J. M., Bagaev, A., Bergmann, M., Chapron, B., Chubarenko,
 I., and Cózar, A.: The physical oceanography of the transport of floating marine debris, Environ. Res. Lett., 15, 23003, 2020.

Sheppard, C.: World Seas: An Environmental Evaluation: Volume I: Europe, The Americas and West Africa, 2018.

Solabarrieta, L., Rubio, A., Castanedo, S., Medina, R., Charria, G., and Hernández, C.: Surface water circulation patterns in the southeastern Bay of Biscay: New evidences from HF radar data, Cont. Shelf Res., 74, 60–76, https://doi.org/https://doi.org/10.1016/j.csr.2013.11.022, 2014.

Con formato: Español (España)

- 920 Solabarrieta, L., Rubio, A., Cárdenas, M., Castanedo, S., Esnaola, G., Méndez, F. J., Medina, R., and Ferrer, L.: Probabilistic relationships between wind and surface water circulation patterns in the SE Bay of Biscay, Ocean Dyn., 65, 1289–1303, https://doi.org/10.1007/s10236-015-0871-5, 2015.
- Solabarrieta, L., Frolov, S., Cook, M., Paduan, J., Rubio, A., González, M., Mader, J., and Charria, G.: Skill Assessment of HF Radar–Derived Products for Lagrangian Simulations in the Bay of Biscay, J. Atmos. Ocean. Technol., 33, 2585–2597, https://doi.org/10.1175/jtech-d-16-0045.1, 2016.
 - Solabarrieta, L., Hernández-Carrasco, I., Rubio, A., Campbell, M., Esnaola, G., Mader, J., Jones, B. H., and Orfila, A.: A new Lagrangian-based short-term prediction methodology for high-frequency (HF) radar currents, Ocean Sci., 17, 755–768, https://doi.org/10.5194/os-17-755-2021, 2021.
- Stanev, E. V, Badewien, T. H., Freund, H., Grayek, S., Hahner, F., Meyerjürgens, J., Ricker, M., Schöneich-Argent, R. I.,
 Wolff, J. O., and Zielinski, O.: Extreme westward surface drift in the North Sea: Public reports of stranded drifters and Lagrangian tracking, Cont. Shelf Res., 177, 24–32, https://doi.org/https://doi.org/10.1016/j.csr.2019.03.003, 2019.

Teles-Machado, A., Peliz, Á., McWilliams, J. C., Dubert, J., and Cann, B. Le: Circulation on the Northwestern Iberian Margin: Swoddies, Prog. Oceanogr., 140, 116–133, https://doi.org/10.1016/J.POCEAN.2015.09.011, 2016.

Tong, X., Jong, M.-C., Zhang, J., You, L., and Gin, K. Y.-H.: Modelling the spatial and seasonal distribution, fate and transport of floating plastics in tropical coastal waters, J. Hazard. Mater., 414, 125502, https://doi.org/https://doi.org/10.1016/j.jhazmat.2021.125502, 2021.

Utenhove, E. van: Modelling the transport and fate of buoyant macroplastics in coastal waters, 2019.

van der Wal, M., van der Meulen, M., Tweehuijsen, G., Peterlin, M., Palatinus, A., Kovač Viršek, M., Coscia, L., and Kržan,
 A.: Identification and Assessment of Riverine Input of (Marine) Litter, Final Rep. Eur. Comm. DG Environ. under Framew.
 Contract No ENV.D.2/FRA/2012/0025, 1–208, 2015.

Weideman, E. A., Perold, V., Omardien, A., Smyth, L. K., and Ryan, P. G.: Quantifying temporal trends in anthropogenic litter in a rocky intertidal habitat, Mar. Pollut. Bull., 160, 111543, https://doi.org/10.1016/J.MARPOLBUL.2020.111543, 2020.

Wendt-Potthoff, K., Avellán, T., van Emmerik, T., Hamester, M., Kirschke, S., Kitover, D., and Schmidt, C.: Monitoring
 Plastics in Rivers and Lakes: Guidelines for the Harmonization of Methodologies, 2020.

Widyatmoko, A. C., Hardesty, B. D., and Wilcox, C.: Detecting anchored fish aggregating devices (AFADs) and estimating use patterns from vessel tracking data in small-scale fisheries, Sci. Rep., 11, 17909, https://doi.org/10.1038/s41598-021-97227-1, 2021.

Woods, J. S., Verones, F., Jolliet, O., Vázquez-Rowe, I., and Boulay, A.-M.: A framework for the assessment of marine litter950impactsinlifecycleimpactassessment,Ecol.Indic.,129,107918,https://doi.org/https://doi.org/10.1016/j.ecolind.2021.107918, 2021.

Yoon, J.-H., Kawano, S., and Igawa, S.: Modeling of marine litter drift and beaching in the Japan Sea, Mar. Pollut. Bull., 60, 448, 2010.