

1 Title page

2 **Back to the future? Conservative grassland management can**  
3 **preserve soil health in the changing landscapes of Uruguay**

4 Ina Säumel<sup>1</sup>, Leonardo R. Ramírez<sup>1</sup>, Sarah Tietjen<sup>2</sup>, Marcos Barra<sup>3</sup> and Erick Zagal<sup>4</sup>

5 <sup>1</sup>Integrative Research Institute THESys Transformation of Human-Environment-Systems  
6 Humboldt-Universität zu Berlin, Unter den Linden 6, Berlin, 10099, Germany.

7 <sup>2</sup>Leibniz Institute of Vegetable and Ornamental Crops (IGZ) e.V., Theodor-Echtermeyer-Weg  
8 1, 14979 Großbeeren, Germany

9 <sup>3</sup>Städtisches Klinikum Dessau, Auenweg 38, 06847 Dessau-Roßlau, Germany

10 <sup>4</sup>Departamento de Suelos y Recursos Naturales, Universidad de Concepción, Campus Chillán,  
11 Vicente Méndez 595, Chillán, Chile

12 *Correspondence to:* Ina Säumel (ina.saeumel@hu-berlin.de)

13 **Abstract.** The ‘soils of the anthropocene’ are predominately agricultural. To understand them,  
14 we analysed agri- and silvicultural intensification of Uruguayan grasslands in a country wide  
15 survey on fertility proxies, pH and trace metals in topsoils originating from different land uses  
16 across the whole country. Thus, our results reflect interactions of both the natural diversity of  
17 the Uruguayan soil formation and impacts of land use change. We observed a loss of nutrients,  
18 trace metals and organic matter from grassland, crops and timber plantations. As an example,  
19 the cation exchange capacity was 160 percent higher in native forests compared to grasslands  
20 and lowest in timber plantations, reaching only half of the CEC in grasslands. Acidification of  
21 topsoils continues as three fourth of all samples are 'extremely acidic' and 'very strongly acidic'.  
22 Topsoils of riverine forests accumulate more trace metals compared to the other uses. We

hat gelöscht: a

hat gelöscht: and lowest in timber plantations

25 assume an accumulation in the topsoils of riverine forests, where high levels of nutrients, trace  
26 metals and organic carbon are found. The translocation of nutrients and organic matter across  
27 the landscape to the erosion base depends on local land use trajectories. Increasing soil  
28 acidification is driving a positive feedback loop, and land use intensification is leading to  
29 degradation of local black soils within a few decades. Our data raises questions about the  
30 resilience and carrying capacity of Uruguayan soils with regard to currently implemented  
31 highly productive management forms, including the use of timber plantation for carbon  
32 sequestration, and supports more conservative forms of extensive management on the grassland  
33 biome.

## 34 1. Introduction

35 Human activities alter the bio- and pedosphere, leaving a footprint of such a magnitude that it  
36 can be verified stratigraphically (Waters et al., 2016). This unprecedented transformational  
37 force is intimately related to the expansion of societies and its productive frontiers, causing a  
38 loss of biodiversity, habitat and soil degradation and, consequently, to ecosystem modification  
39 (Foley et al., 2005, Borrelli et al., 2017). In this context, soil sciences have transitioned from  
40 studies on natural soil formation to the science of ‘anthropedogenesis’ (Richter, 2020),  
41 focussing on the ‘soils of the anthropocene’ that are predominately agricultural (51 Million  
42 km<sup>2</sup>) or urban (1.5 Million km<sup>2</sup>; FAO, 2019).

43 The temperate grasslands of South America have historically been characterised by rolling  
44 plains and low hills that have been extensively exploited for cattle production and its  
45 derivatives since the arrival of European colonization. The Río de la Plata grasslands are one  
46 of our planet’s four major black soil regions (Durán et al., 2011; Liu et al., 2012), and some of  
47 the most fertile soils in the world. Playing an important role in the global food production, these  
48 are characterized by a thick, humus and base cation rich and a high cation exchange capacity

hat gelöscht: Mio.

hat gelöscht: Mio

51 throughout their profile. Maintaining their properties are therefore crucial to developing  
52 sustainable and productive agriculture (Durán et al., 2011; Liu et al., 2012).

53 Today, the 'Uruguayan savanna' is part of the top-three most critically endangered biomes  
54 (Veldman et al., 2015). In recent decades, however, the grassland has decreased due to the  
55 expansion of cash crops and *Eucalyptus* plantations, both of which are promoted by national  
56 legislation including land grabbing and trans-nationalization (Piñeiro, 2012). This land use  
57 intensification, with its increased input of energy, nutrients and pesticides, leads to an overall  
58 loss of soil fertility and increasing toxicity related to acidification, salinization and  
59 contaminants (Liu et al., 2012; Borrelli et al., 2017). Ecological, economic and cultural  
60 functions of soils are severely degraded, and the degradation of black soils in South America  
61 is of particular concern because they have only been heavily exploited for a comparatively  
62 short period of time (Durán et al., 2011).

63 Since the first decades of the twentieth-century, compared to other disciplines, soil sciences  
64 have received an extraordinary amount of attention in Uruguayan academia, governance, the  
65 productive sector, and also in the general public, resulting in a national soil inventory program  
66 in 1965. The subsequent classification of Uruguayan soils and their productivity (CONEAT  
67 Index) remains an important source for today's land taxation and for management plans by the  
68 legal conservation regulations, and provides a detailed classification that takes into account soil  
69 type, texture, natural vegetation, altitude and geology (Lanfranco and Sapriza, 2011).

70 As soil degradation is extremely relevant for countries like Uruguay, which are  
71 socioeconomically dependent on their soils (Zubriggen et al. 2020), it is a topic of discussion  
72 for local farmers, academia, and the public. An actualization of the state of the art of soils and  
73 related processes is needed (García-Préchac et al. 2004; De Faccio et al. 2021), particularly as  
74 there has been little study of the impacts of the Uruguayan grassland intensification on soils  
75 properties (Beretta-Blanco et al., 2019). At the same time, while a new paradigm for grassland  
76 intensification with a wide set of means including fertilization has been proposed to increase

77 economic and environmental sustainability (Jaurena et al., 2021), it is urgent to get more  
78 insights into the dynamics of nutrients in soils of Uruguay and their availability for crops  
79 (Beretta-Blanco et al., 2019).

80 Soil classifications are mainly based on subsoils. However, we focus on topsoil as the most  
81 relevant and very responsive interface for ecological processes and farmer's management.

82 Understanding the state of the art of topsoils and its processes is crucial for developing  
83 recommendations for sustainable land management practices. Due to the diversity of  
84 perspectives on soil quality and health and related ecosystem services, operational procedures  
85 for evaluation of soil functioning are still lacking (Bonfante et al. 2021). We contribute to a  
86 better understanding of globally occurring degradation processes among often conflicting goals  
87 such as desired soil productivity, yield limits, especially in erosion sensitive soils, and  
88 necessary soil conservation.

89 We therefore explored soil parameters describing current chemical conditions of topsoils that  
90 are parts of different soil groups and orders, and Uruguayan soil categories. Specifically, in  
91 order to explore the gains and losses of macro and micro-nutrients and soil organic carbon  
92 across landscapes, and to determine the impact of land use change on acidification and trace  
93 metal presence and related trade-offs with soil degradation and conservation. In detail we  
94 address the following question: how do fertility proxies such as soil organic carbon and content  
95 of nutrients, acidification (pH) and trace metals accumulation in topsoils vary across different  
96 land uses (i.e. comparing grassland, timber plantation, native forest, and agricultural land)?

97 Thus, we expand the knowledge across land uses from more natural to strongly modified uses  
98 and discuss the results in light of different degradation processes such as erosion, depletion of  
99 nutrients or carbon, acidification and accumulation of pollutants and in the light on current  
100 debates on intensification.

hat gelöscht: .

hat gelöscht: since u

hat gelöscht: in

hat gelöscht: the field of tension

hat gelöscht: between

hat gelöscht: mobility

hat gelöscht: i)

## 108 2. Material and Methods

### 109 2.1 Study area and design

110 Uruguay covers about 176,000 km<sup>2</sup>, and has a population of 3.5 Million, mainly in urban areas.  
111 The Western half of Uruguay is dominated by Mollisols, developed on a wide range of  
112 sediments from the Devonian to the Cenozoic era (partially associated with Vertisols), while  
113 the Eastern plains and wetlands have, in addition to Mollisols, other soil types on the slopes,  
114 rocks and on flood plans (i.e. Alfisols, Ultisols, Entisols and Inceptisols; Durán et al. 2011).  
115 During a country-wide survey from December 2015 to March 2016, we collected 280 topsoil  
116 samples of 0-10 cm depth from 101 plots (50x50m in grassland and crops; 100x100m in native  
117 forests and timber plantations) distributed at 28 monitoring sites throughout Uruguay, South  
118 America (Fig. 1a-b; Table 1), using a stratified random design. In the first step, we randomly  
119 selected monitoring sites across the country. In the second step, we contacted landowners to  
120 explore their willingness to establish a long-term monitoring site. If the owner agreed, plot  
121 selection was stratified by different rural land use types: grassland, timber plantations of *Pinus*  
122 and *Eucalyptus* species, native forest, and crops. Native forests cover mainly riverine and park  
123 forests. The later are a savanna like transition zones between riverine forests and the open  
124 grasslands. We subdivided grassland plots according to the intensity of use: (i) undisturbed  
125 grassland (without grazing), (ii) partially grazed grasslands (with sporadic grazing and low  
126 animal charge), and (iii) highly grazed grassland (with high animal charge). Land use change  
127 from 1986 to 2017 follows basically three different trajectories: i) the expansion of timber  
128 plantations over grassland leading to a disaggregation of grassland by timber plantations; ii)  
129 cropland expansion where crop cover maintains the open landscape character of former  
130 grasslands, grassland conservation where large and regularly interconnected riverine forests in  
131 a landscape dominated by grasslands (Ramírez and Säumel 2021) and grassland intensification

hat gelöscht: m

133 changing from natural grassland to so called ‘improved’ or artificial grasslands (Modernel et  
134 al. 2016; Jaurena et al. 2021). Fertilization and application of other agrochemicals is standard  
135 procedure in timber plantations, artificial grasslands and industrial crops. We sampled top soil  
136 three times at each land use at the edges of the plot, and stored samples below 7°C until  
137 laboratory processing. The plots are placed in homogenous areas to avoid edge effects.

## 138 2.2 Analysis of Soil Samples

139 For gravimetric determination of soil moisture, the topsoils samples were dried at 105 °C until  
140 constant weight. Next, lumps in the samples were broken down and the remaining plant  
141 material was removed before sieving (2 mm) and ground.

142 We analysed 280 samples regarding macronutrients, pH and trace metals and 80 samples for  
143 soluble cations and micronutrients (Table 1; Sadzawka et al. 2006; Zagal & Sadzawka 2007).  
144 Among fertility-related variables, we measured the total amount of the macronutrients  
145 phosphorus (P), organic carbon (SOC) and nitrogen (N), so obtaining the C/N ratio. To  
146 determine total carbon and nitrogen, the samples were sieved again (0.5 mm) and analyzed  
147 using dry combustion with a LECO TruSpec CN (USA) at a combustion temperature of 950  
148 °C with ultra-pure oxygen. In addition, the presence of carbonates was tested for by adding  
149 concentrated hydrochloric acid, which was negative for all soil samples analysed.

150 We determined concentrations of the soluble cations calcium ( $\text{Ca}^{2+}$ ), magnesium ( $\text{Mg}^{2+}$ ),  
151 potassium ( $\text{K}^+$ ), sodium ( $\text{Na}^+$ ), and the micronutrients copper ( $\text{Cu}^{2+}$ ), zinc ( $\text{Zn}^{2+}$ ), manganese  
152 ( $\text{Mn}^{2+}$ ) and iron ( $\text{Fe}^{2+/3+}$ ) by atomic spectroscopy (Unicam AAS Solaar 969, Thermo Electron  
153 Corporation, USA), extracting with ammonium acetate (1 mol l<sup>-1</sup>, pH 7), and with DTPA-  
154  $\text{CaCl}_2$ -TEA at (pH 7.3). We calculated the cation-exchange capacity (CEC). Acidity was  
155 measured by adding calcium chloride (0.01M) to the samples in a 2.5:1 proportion, and after  
156 shaking and two hours rest, read with a pH meter (HI2550 meter, Hanna Instruments, USA)).

hat gelöscht: SOC

hat gelöscht: was

hat gelöscht: the presence of carbonates was

hat gelöscht: assessed to obtain the amount of SOC.

hat formatiert: Hochgestellt

hat formatiert: Hochgestellt

hat formatiert: Hochgestellt

hat formatiert: Hochgestellt

hat formatiert: Hochgestellt

hat formatiert: Hochgestellt

hat formatiert: Hochgestellt

hat formatiert: Hochgestellt

hat formatiert: Hochgestellt

161 For these variables, each tenth sample was duplicated. We categorized acidity using the USDA  
162 Natural Resources Conservation Service classification (Kellogg, 1993).

163 To determine the concentrations of arsenic (As), cadmium (Cd), chromium (Cr) and lead (Pb),  
164 samples were further sieved as before (0.5 mm), weighed out into a digesting container, and  
165 extracted with a mixture of nitric acid (70%) and hydrochloric acid (30%) in a 1:3 proportion.  
166 The digestion took place in a Titan MPS (Perkin Elmer, USA) microwave at a programme  
167 suitable for this digestion (Method 3051A). During the proceedings, only deionized water Type  
168 I ASTM1193 (EC max 0.06 to 0.1  $\mu\text{S}/\text{cm}$ ) was used. Reagents were used to eliminate traces of  
169 other materials and to avoid contamination of the samples. The trace metals were determined  
170 by inductively coupled plasma optical emission spectroscopy using the Optima 8000 (Perkin  
171 Elmer) with metal-associated wavelengths of 193.696 nm for As, 228.802 nm for Cd, 267.716  
172 nm for Cr and 220.353 nm for Pb.

173 Total P concentration of phosphorus was determined calorimetrically after microwave-assisted  
174 digestion with Unicam spectrometer at a wavelength of 660 nm. For all these variables, a  
175 repetition for each round of the microwave digestion was made.

### 176 **2.3 Soil classification**

177 We intersected the coordinates of the centre of the plots with maps containing geospatial  
178 information on the classification of the Uruguayan soils using ArcGIS 10.3 (ESRI, 2018). For  
179 Soil Groups classification, we used the of the World Reference Base for Soil Resources (WRB;  
180 IUSS Working Group, 2015); for Soil Orders, the USDA soil taxonomy (Soil Survey Staff,  
181 1999); and for the local Uruguayan classification, Soil CONEAT (Comisión Nacional de  
182 Estudio Agronómico de la Tierra) Groups categories, which include productive capacity of  
183 cattle and sheep (MGAP, 2020). The CONEAT groups are defined by their productive capacity  
184 in terms of beef, sheep and wool expressed by an index relative to the average productive

185 capacity of the country, to which the index 100 corresponds. The classification is based on  
186 photo-interpretation at a scale of 1:40,000, field verifications and physico-chemical analysis of  
187 the soils. The productivity indices correspond to soil groups. The CONEAT groups have been  
188 defined by the dominant and associated soils according to the Soil Classification of Uruguay.  
189 The groups are related to the units of the Soil Reconnaissance Chart of Uruguay at a scale of  
190 1:1,000,000. For each group, some important soil properties and associated landscape  
191 characteristics are indicated. The nomenclature of the CONEAT groups correlates with the Soil  
192 Use and Management Zones of Uruguay. The Soil Groups are superimposed on the rural parcel  
193 and are represented in the CONEAT cartography at a scale of 1:20,000 (for more details see  
194 MGAP, 2020).

## 195 **2.4 Data Analysis**

196 In a first step, we explored and prepared our database for further analysis. Exploring the  
197 distribution of the soil parameters in R (R Core Team 2021), ruled out the normality of the data  
198 using the Shapiro-Wilk Test and the homogeneity of variances with the Fligner-Killeen Test.  
199 We tested for outliers using the 1.5-3 IQR threshold and the function *outlierTest* from the R  
200 package *car* (Fox and Weisberg 2019), reviewing the flagged observations case by case in the  
201 experimental context. The outliers were removed (Supplementary Material: Fig.S2). The  
202 variables on soils characteristics showed generally positive skewed distributions, in some cases  
203 multimodal, and tests showed evidence contrary to assumptions of spatial autocorrelation,  
204 homoscedasticity and normality in most cases (Supplementary Material: Table S1-S2; Fig. S1-  
205 S2).  
206 Spearman's rank correlations ( $\rho$ ) were calculated to explore linear associations between soil  
207 parameters across all single samples and within different land uses. We used the *adonis*  
208 function of the R package *vegan* v2.5-7 (Oksanen et al., 2020) with a Euclidean dissimilarity



209 matrix from our normalized soil data to perform PERMANOVA tests with 9999 permutations  
210 to analyse the multivariate homogeneity of group dispersions based on differences on soil  
211 parameters between land uses.

212 To compare the general effects of land use type on topsoil parameters, non-parametric Kruskal-  
213 Wallis tests were carried out in R. When significant ( $p \leq 0.05$ ), we used Pairwise Wilcoxon  
214 Rank Sum tests with Benjamini & Hochberg correction (1995) to evaluate pairwise differences  
215 among land uses.

216 We used non-metric multidimensional scaling (NMDS) as a robust unconstrained ordination  
217 method to visualize patterns of top soil characteristics among all samples and within  
218 subsamples (intersected with different soil Orders) across different land uses using the Bray-  
219 Curtis dissimilarity matrix. The matrix was constructed with the *vegan* package to depict  
220 patterns of all soil parameters in two dimensions (Fig.3a, c-f) and for the dataset without soluble  
221 cations and micronutrients variables comparing subcategories within single land use types (i.e.  
222 within grasslands in 'undisturbed', 'partially grazed' and 'highly grazed' plots; timber  
223 plantations in '*Eucalyptus*' or '*Pinus*' plots; Fig. 3b).

### 224 3. Results

225 Nearly 75% of the sampled 101 plots intersected with the soil Group category 'Luvic Phaeozem'  
226 (WRB; IUSS Working Group, 2015). A quarter of our plots were distributed among four other  
227 Groups (i.e. Haplic Luvisols, Eutric Vertisols, Planosols, Lithic Leptisols; Fig. 1c). Half of our  
228 plots intersected with the Orders 'Argiudolls' or 'Argiudolls & Hapluderts' (Soil Survey Staff,  
229 1999), and about a quarter with 'Hapludalfs & Hapludults' or 'Argiudolls, Hapludolls &  
230 Hapludalfs' (Fig. 1d). Our plots intersected with 32 different CONEAT categories (MGAP,  
231 2021). The most frequent categories were '10.2', '9.3' and '2.12' (25% of our plots) and another  
232 quarter with '9.1', '8.8', '9.6' and 'G03.11' (Fig. 1e).

hat formatiert: Nicht Hervorheben

### 233 **3.1 General characteristics of Uruguayan topsoils**

234 The measured topsoil parameters vary widely across Uruguay, between the different land uses  
235 and classification to different soil orders (Table 1, Table A1-A4). The soil organic carbon in  
236 topsoils ranged between 0.8 to 16 percent, and was highest in native forests and lowest in timber  
237 plantations. The mean of the C/N ratio was about 13, and lowest in crop soils. Phosphorous  
238 ranged between 43 to 1009 mg kg<sup>-1</sup>. We also observed a high variability for the micro- and  
239 macronutrients and trace metals. The average cation-exchange capacity (CEC) was 12.41 cmol  
240 kg<sup>-1</sup>, highest in topsoils of native forest, followed by crops and grasslands and timber  
241 plantations (Table 1, Table A1-A2).

242 For the whole data set, high correlation was found between P with SOC and Zn ( $\rho=0.82$  and  
243  $0.76$ , respectively), and between Mg with Ca and Na ( $\rho=0.82$  and  $0.76$ , respectively; Fig. 2).  
244 Similar results were observed within particular land uses, although in native forests, a negative  
245 moderate correlation between Ca and Fe was observed (Fig. S3). There was high correlation  
246 between pH and Ca ( $\rho=0.89$ ; Fig. 2). In native forests, we also found similar correlation with  
247 other soluble cations like Mg and K (Fig. S3). In park forests topsoils, there was a negative  
248 correlation between pH and As and Pb ( $\rho=-0.81$  and  $\rho=-0.84$ , respectively). In highly grazed  
249 pastures and crops pH was highly correlated with Cr, and in crops also with As ( $\rho=0.93$ ; Fig.  
250 S3). Among trace metals, Spearman correlation was moderate between As with Cr and Pb and  
251 high between Cr and Pb (Fig. 2). These correlation trends increased in grasslands, timber  
252 plantations and crops. We also found a high correlation between cadmium and Cr in crops  
253 ( $\rho=0.81$ ). Phosphorus was highly correlated with Cr and As in pine plantations, while in  
254 soybean crops the correlation was between phosphorus with Cr and Pb (Fig. S3).

### 255 **3.2 Topsoil characteristics clustered by land use**

256 We found differences in multivariate distribution of samples according to the different land  
257 uses (i.e., grassland, timber plantations or native forests; Fig. 3a-f; Table S3-S8). All pairwise

258 comparison of land uses showed significant differences with all variables ( $p=0.0001$ ; Fig. 3a;  
259 Table S3). We analysed subcategories within a land use type using the dataset without soluble  
260 cations and micronutrients variables, only finding significant differences between *Eucalyptus*  
261 and *Pinus* stands ( $p=0.0001$ ; Fig. 3b; Table S4) but not among different grassland subtypes.  
262 We also found differences analysing subsamples of different soil order classification to the  
263 different land uses (Fig. 3c-f; Table S5-S8). The samples intersected with 'Argiudolls' included  
264 all plots on crops, and we found significant differences in all pairwise comparisons of land uses  
265 except between grasslands and timber plantation ( $p=0.0004$ ; Fig. 3c; Table S5). We further  
266 found differences between timber plantations and native forests at soils of the 'Argiudolls &  
267 Hapluderts' Orders ( $p=0.0009$ ; Fig. 3d; Table S6) and between timber plantations and  
268 grasslands or native forests in soils of the 'Argiudolls, Hapudolls & Hapludalf' Orders  
269 ( $p=0.0284$ ; Fig. 3e; Table S7). Results for samples in 'Hapludalfs & Hapludults' soils were  
270 similar to those obtained at country scale ( $p=0.0001$ ; Fig. 3f; Table S8).

### 271 **3.3 Differences in fertility proxies**

272 We found significantly higher fertility proxies for SOC, P, Ca, Mg and Zn in topsoils from  
273 native forests compared to grasslands and timber plantations (Fig. 4b-e, h). Phosphorous was  
274 significantly higher in topsoils of grassland compared to those of timber plantations (Fig. 4c).  
275 Potassium was significantly higher in topsoils of native forests compared to timber plantations  
276 (Fig. 4f). At subcategories level, we found significantly higher amounts of K in partially used  
277 grasslands in comparison to samples from highly grazed pastures, and higher SOC ( $p=0.002$ ),  
278 P ( $p=0.059$ ), Na ( $p=0.043$ ), K ( $p=0.012$ ) and Zn ( $p=0.048$ ) in *Eucalyptus* compared to *Pinus*  
279 plantations (Fig. S4). Among native forests, samples from riverine forests contain more Mg  
280 ( $p=0.023$ ) and Na ( $p=0.023$ ) in comparison to park forests.  
281 Considering all samples within the Order 'Argiudolls', C/N ratio in agricultural topsoils are  
282 lower compared to all other land uses. Phosphorus was higher in topsoils of native forests and

283 crops compared to grasslands and timber plantations. Soil organic carbon was highest in native  
284 forests (Fig. 5a-o; Table A3-A4).

### 285 **3.4 Soil Acidification**

286 Topsoil samples showed a markedly acid profile (median pH=4.66), with nearly 75% classified  
287 as 'extremely acidic' and 'very strongly acidic' (Fig. 6a). We found significant differences across  
288 land uses, with less acidic native forests, and lower pH in timber plantations (Fig. 6b).  
289 Comparing between land uses, we found more samples with neutral acidity in grasslands and  
290 more with higher acidity in timber plantations (Fig. 6b-c). In addition, we found lower pH in  
291 samples from *Pinus* compared to *Eucalyptus* stands ( $p=0.018$ ). Results of analysis inside soil  
292 Orders showed similar variations observed at country scale, with timber plantations being more  
293 acid and native forest closer to neutral pH (Fig. 6d-f).

### 294 **3.5 Trace metal accumulation across land uses**

295 For As, Cd, Cr and Pb, we found significantly higher concentrations in topsoils originating  
296 from native forests compared to the grassland and timber plantation samples (Fig. 4l-o and 5l-  
297 o; Table A1-A4). At the same time, samples from *Eucalyptus* plantations had higher  
298 concentrations of both Cr ( $p<0.005$ ) and Pb ( $p<0.005$ ) than *Pinus* topsoils, while the same was  
299 observed for Cr ( $p<0.001$ ) and Pb ( $p<0.05$ ) in riverine forests compared to park forests (Fig.  
300 S5).

## 301 **4. Discussion**

302 The vicious circle between the wish to stop soil degradation and concurrent increases in land  
303 productivity to satisfy the increasing demand for food, fibres and energy has not been broken  
304 since green revolution. Socio-economic and conventional management practices that drive soil  
305 degradation have generated several traps, such as the 'inputs trap' where a reduced yield per

306 area is followed by higher fertilizer application or the 'credit or poverty trap' where economic  
307 pressure forces the farmer to intensification (Gomiero, 2016). Caught in this loop, the soils of  
308 the temperate grasslands of Uruguay have suffered strong degradation from erosion,  
309 acidification, contamination, salinification and compaction (Zurbriggen et al., 2020). This is  
310 clearly reflected in the results of our topsoil survey, which also adds interesting insights from  
311 timber plantations, grasslands and native forests to an existing database consisting mainly of  
312 crops and pastures samples from 2002-2014, which demonstrated the loss of organic matter by  
313 25% and an increasing loss of nutrients (Beretta-Blanco et al. 2019). We contribute deeper  
314 insights on fertility, acidification and trace metals accumulation in topsoils from a wide range  
315 of different land uses, which is, to our knowledge, unique for the region since the CONEAT  
316 classification (CONEAT Index, 1976).

317

#### 318 **4.1 Translocation of elements in topsoils within landscape**

319 Our data demonstrate the high amounts of organic carbon, nutrients and trace metals in topsoil  
320 samples from riverine forests, suggesting transport of soil particles from the surrounding land  
321 uses (e.g., grasslands, crop or timber plantations) to the borders of rivers, streams and creeks.  
322 Therefore, we assume that organic carbon, nutrients and trace metals are displaced within the  
323 landscape and accumulate in the floodplains. Regional soil erosion models estimate the loss of  
324 2-5 tons ha<sup>-1</sup> year<sup>-1</sup> for a third of the country depending on precipitation, topography, soil  
325 erodibility and land management (Carrasco-Letelier and Beretta-Blanco 2017). One possible  
326 direct impact is the increasing eutrophication reported for larger local rivers, although the  
327 models used by these authors did not link Chlorophyll-a concentrations with agricultural land  
328 use (Beretta-Blanco and Carrasco-Letelier 2021 and replies). However, other vertical processes  
329 and differences across land uses can be hidden deeper in the soil profile, and have not been  
330 analysed in this study.

331 Organic carbon content and the exchangeable cations are strongly reduced in topsoils of  
332 grasslands, timber plantations and crops compared to native forests (Fig. 4b, d-h and 5b, d-h).  
333 Of all the fertility proxies assessed, phosphorus in topsoil was most significantly affected by  
334 different land uses, being highest in native forests (Fig. 4c and 5c). The cation exchange  
335 capacity (CEC) was highest in native forests (more than 160 percent of the CEC in grasslands)  
336 and lowest in timber plantations, reaching only half of the CEC in grasslands (Table A1-A2).  
337 Lower average nutrient concentrations and corresponding CEC reported for two timber  
338 plantations in the East and one in the Norwest of Uruguay (Hernández et al., 2009 and 2016;  
339 Céspedes-Payret et al., 2012) may result from a combined effect of topsoil degradation in  
340 timber plantations due to management practices and to soil texture (Sandoval-Lopez et al.,  
341 2020). The trees' uptake and the general export of nutrients from fast growing timber  
342 plantations through harvesting is higher than the natural input into those systems (Merino et  
343 al., 2005). This effect is particularly relevant because timber plantations in Uruguay are on  
344 'forestry priority soils', which are generally soils with low fertility, superficial to moderate  
345 depth and good drainage (OAS, 1994), so afforestation might reduce even more their soil  
346 fertility.

347 Comparing neighbouring sites of grassland and afforested grasslands, our data set shows no  
348 significant depletion of nutrients by timber plantations ( $p=0.208$ ) but a slightly higher average  
349 of CEC at grasslands (Table A1). In general, plants are expected to compensate the extraction  
350 of cations from upper soil by the 'uplift of nutrient' from deeper horizons (Jobbagy and Jackson,  
351 2004). This has been questioned for *Eucalyptus* plantations on sandy soils, where, for example,  
352 an increase of potassium in the topsoil was not observed compared to the neighbouring  
353 grassland (Céspedes-Payret et al., 2012). Phosphorous and cations decline with sand content,  
354 so physicochemical factors such as the percentage of sand and organic matter, influence soil  
355 fertility (Sandoval-Lopez et al., 2020). A study of *Eucalyptus* plantations in South-East Brazil  
356 did not show a significant depletion of nutrients and carbon; in contrast, carbon, potassium and

357 calcium increased in the topsoil after twelve years over one or two harvest and fertilization  
358 cycles (McMahon et al., 2019). To sum up, the patterns observed in the various sites in our  
359 survey and the other studies indicate very complex interactions of numerous factors. Removal  
360 of nutrients by high-yield timber plantations may exceed the capacity of nutrient exchange or  
361 turnover of forest litter and wastes. Both stand management and environmental conditions (e.g.,  
362 precipitation) influence nutrient and carbon stocks (McMahon et al., 2019; Sandoval-Lopez et  
363 al., 2020).

364 The C/N ratio in topsoils ranged within values reported for grassland and timber plantations of  
365 the region (see Berthrong et al., 2012). In contrast to that study, however, we observed no  
366 differences between grassland and timber plantations (Fig. 4a), but between topsoils of timber  
367 plantations, native forests and crops (Fig. 5a). Organic matter and nitrogen decrease in topsoils  
368 after grassland afforestation (Berthrong et al., 2009) and C/N ratio increases with plantation  
369 age and decreases with precipitation (Berthrong et al., 2012). The topsoils of timber plantations  
370 have, on average, lower N concentrations compared to other land uses (Table A2) and data in  
371 literature (Jobbagy and Jackson 2003).

372 Although in cropland, nutrients are regularly compensated for by increased application of  
373 fertilizers (Beretta-Blanco et al., 2019), timber plantations are usually fertilized only in the first  
374 year of planting (e.g., Binkley et al., 2017; Sandoval-Lopez et al., 2020). The CEC and the  
375 average ratio between  $K^+(Ca^{2+} + Mg^{2+})^{-1/2}$  in crops and grasslands are in the ranges reported by  
376 Beretta-Blanco et al. (2019) with lower availability of potassium for crops. A shortage of  
377 potassium and calcium is also most likely for future timber plantations after harvest, especially  
378 if the logging process do not include bark stripping on site (Hernández et al., 2009) and due to  
379 short time spans between harvest and new planting in the same area (Sandoval-Lopez et al.,  
380 2020), but see contrasting results of McMahon et al. (2019). The extraction of nutrients and  
381 biomass due to grazing (Fernandez et al., 2017), timber (Hernández et al., 2009 and 2016) and

382 crop production (Beretta-Blanco et al., 2019) lowers organic matter content and exchangeable  
383 cations, affects the physical, chemical and biological soil properties, and drives degradation.

#### 384 **4.2 Acidification in Uruguayan topsoils across land uses**

385 A further dimension of the soil degradation directly linked to the cation extraction is the  
386 acidification of soils. This has been demonstrated for crops and pastures (Beretta-Blanco et al.,  
387 2019), and a study site with *Eucalyptus* (Céspedes-Payret et al., 2012) and is now broadly  
388 supported by our topsoil samples originating from a wide range of different land uses across  
389 Uruguay. The pH of our topsoil samples are mainly in the category of very strongly to  
390 extremely acidic and lowest in timber plantations (Fig. 6), which are below the means reported  
391 so far (Jobbagy and Jackson, 2003; Céspedes-Payret et al., 2012). Moreover, as our data on  
392 topsoil pH fall short compared to the values estimated by the Food and Agriculture  
393 Organization of the United Nations and the Intergovernmental Technical Panel on Soils (Caon  
394 and Vargas, 2017), acidification and the deterioration of topsoil quality continues.  
395 Acidification results from intensified land uses with nitrogen fertilization, with biological N  
396 fixation by the legumes both used in the improved pastures (Modernel et al., 2016), or with  
397 cation extraction by crop or timber harvest (Jobbagy and Jackson, 2003). So far, although forest  
398 soils tend to be more acidic than agricultural soils due to acid-neutralizing treatments in the  
399 latter (Baize and van Oort, 2014), we found no differences between topsoils of crops and native  
400 forests (Fig. 6d) as the high organic carbon content in native forests buffers the process to a  
401 certain extent.

#### 402 **4.3 Riverine Forest soils as sink for trace metals**

403 In general, the average concentration of Zn, Cu, Cd, Cr, As and Pb in our topsoils were within  
404 the expected ranges of samples from crops, pastures and grassland in the region (Lavado et al.,  
405 1998, 2004; Roca, 2015). Some samples, especially from orchards and crops, exceeded the  
406 background values of copper, cadmium and arsenic (Table A2; Roca, 2015). Little is known



407 about the pedo-geochemical background of the Pampean soils (Roca, 2015): for example, high  
408 arsenic concentration in Uruguayan ground waters has been hypothesized to be due to  
409 quaternary ash deposits (Wu et al., 2021). In addition, the lateral heterogeneity of Pampean  
410 soils over short distances makes separating geochemical and anthropic signatures difficult  
411 (Roca, 2015). However, that the main risk of soil contamination in the region is from the  
412 application of fertilizers and agrochemicals is uncontested (Kaushal et al., 2018).

413 To our knowledge, there has been no regional study of trace metals in the native riverine forests  
414 or timber plantations. Our work thus expands the evidence base for these land uses. The topsoils  
415 of riverine forests accumulate more trace metals compared to those of timber plantations and  
416 crops (Fig. 4l-o). The higher amount of soil organic matter in riverine forests favours the  
417 retention of cations, including trace metals. Although the origin of potentially harmful elements  
418 in forest soils have been primarily attributed to atmospheric deposition (Baize and van Oort,  
419 2014), atmospheric deposition only plays a major role in vicinity of urban or industrial  
420 development, and our data from rural sites suggests a different entry path from the surrounding  
421 land uses to the riverine forests. High acidification and low amounts of organic matter reduce  
422 the retention of trace metals in the soil of timber plantations, and elements leach out of the soil  
423 towards the water table (Baize and van Oort, 2014). The acidification strongly contributes to  
424 the overall mobility of base cations into the ‘chemical cocktail of the Anthropocene’ (Kaushal  
425 et al., 2018), including trace metals. We thus observe positive feedback in already  
426 impoverished soils with high acidity favouring cations solubility, in addition to the uptake by  
427 trees intensifies this effect. Timber plantations extract trace metals from soil and also  
428 accumulate it in the bark or leaves, so they have been used for phytoremediation (Li et al.,  
429 2020). This may explain the higher concentration of cadmium in grassland compared to timber  
430 plantations (Fig. 4m). Differences between *Eucalyptus* and *Pinus* stands may be related to  
431 different age classes, as the later may have extracted more lead and chromium from the soil  
432 due to their older stand age with rotation periods of about 20 years (Li et al., 2020).

hat gelöscht:

hat gelöscht:

435 **4.4 Carbon storage in topsoils of *Eucalyptus* plantations?**

436 Our study provides evidence that the loss of soil organic carbon limits not only the productivity  
437 of the crops, but also potential carbon sequestration in the region (Beretta-Blanco et al., 2019).  
438 Grassland conversion to cropland decreases soil organic carbon storage compared to adjacent  
439 native grasslands (Hernandez-Ramirez et al., 2021).  
440 Afforestation of croplands has been also discussed as a carbon sequestration measure to  
441 proactively address and effectively mitigate ongoing climate change within a person's lifetime  
442 (Hernandez-Ramirez et al., 2021; Mayer et al. 2020). Although carbon stocks of four  
443 *Eucalyptus* stands in the Brazilian Cerrados increased but did not change in four *Eucalyptus*  
444 stands within the Atlantic Forest (McMahon et al., 2019), this may not be the case for short  
445 rotation of *Eucalyptus* plantation on Campos grasslands. In our topsoil samples originating  
446 from 28 different stands across Uruguay, organic carbon is lowest in topsoils of timber  
447 plantations (Fig. 4b). Similar amounts have been reported for a timber plantation in the East of  
448 Uruguay (Céspedes-Payret et al., 2012), and in North-Eastern Argentine (Sandoval-Lopez et  
449 al., 2020). Our data therefore provide clear evidence that rather than contributing to carbon  
450 sequestration in the topsoil, the carbon release from the transformation of native grasslands to  
451 plantation with these fast-growing species has several adverse effects depending on  
452 precipitation and soil type (reviewed by Mayer et al., 2020).  
453 Several trade-offs between carbon sequestration through afforestation and local water yield and  
454 soil fertility have been demonstrated, including nutrient and soil organic matter depletion,  
455 acidification, and biodiversity loss and corresponding challenges for landscape conservation  
456 (e.g., Jackson et al., 2005; Veldman et al., 2015; Friggens et al., 2020). Soil carbon changes  
457 dynamically during the first decade of afforestation. Remaining grassland carbon declines,  
458 while tree carbon gain starts (e.g. Paul et al., 2002). Although a net gain of carbon is expected  
459 when the new forests approached equilibrium after decades (Hernández-Ramirez et al., 2021),  
460 in contrast to long-lasting forests, *Eucalyptus* harvest in Uruguay takes place after less than a

461 decade (appr. 7-10 years). Soil organic carbon does not differ between *Eucalyptus* plantations  
462 and neighbouring grasslands (Appendix A; Table A1 and A2). Another study near our sites 12  
463 and 13 also did not find a significant difference between the soil organic carbon of the upper  
464 soil (0-30cm) compared to afforested versus native grasslands (Hernández et al., 2016). A study  
465 near our sites 7, 8, 10 reported higher top soil carbon of grassland compared with afforested  
466 sites in their vicinity only if sand content is lower than sixty percent (Sandoval-Lopez et al.,  
467 2020). McMahon et al. (2019) identified a greater carbon gain under *Eucalyptus* stands  
468 compared to (potential) carbon losses in neighbouring degraded Cerrado grasslands. There is  
469 evidence that the retention of residues after harvest increases the carbon stock in soil (Mayer  
470 et al., 2020). Long term monitoring of carbon stocks is needed to verify the influences of  
471 management and environmental changes. A regional study on *Eucalyptus* plantations across  
472 different biomes in Brazil shows both decreases and no changes depending on precipitation in  
473 the dry season, clay content and on the initial stock of carbon in the soil (Cook et al., 2016)  
474 The simplistic solution of huge tree plantations to compensate anthropogenic CO<sub>2</sub> emissions  
475 has been challenged in the last decade, and some crucial lessons learnt have been summarized  
476 (Di Sacco et al., 2020). Since grasslands are more sustainable carbon sinks compared to  
477 climate-vulnerable monocultures such as timber plantations (Dass et al. 2018), avoiding  
478 afforestation of previously non-forested lands is important. This is the case for *Eucalyptus*  
479 afforestation in the originally forestless grasslands of Uruguay: models on carbon sequestration  
480 and dynamics in Mollisols and Oxisols under South American grasslands estimated a higher  
481 carbon retention efficiency under grasslands compared to afforested sites, suggesting that  
482 silvopastoral systems are a potential solution for soil carbon sequestration in tropical soils  
483 (Berhongaray and Alvarez, 2019).  
484 Connecting or expanding existing forest and using native species for plantings (Di Sacco et al.,  
485 2020) is also recommended (see also Alvaro et al. 2020, Ortiz et al. 2020). Our topsoil data  
486 indicates that carbon sequestration occurs mainly in the topsoils of native riverine forests that

487 cover less than five percent of the Uruguayan territory. Consequently, the expansion of native  
488 forests and the use of native species in forestry project for long term establishment can reduce  
489 adverse effects of timber plantations. While there is preliminary evidence that N-fixing tree  
490 species can help increase local C stocks of afforestation, because of their potential for invasion,  
491 exotic N-fixers should be avoided (Mayer et al. 2020).

492 Recent studies indicated that novel techniques such as perennial grain cropping can help to turn  
493 around cropland degradation into beneficial cropland aggradation by using the advantage of  
494 perennial vegetation of conserving and even enhancing short term and long-term soil carbon  
495 storage and other ecosystems services (Kim et al. 2022).

#### 496 **4.5. Back to more conservative grassland management?**

497 Our soil survey data shows strong soil degradation of Uruguayan black soils from erosion,  
498 acidification and contamination, and suggests a translocation of nutrients and organic carbon  
499 across the landscape from grassland, timber and crop plantations to the riverine forests. The  
500 potential of grasslands as cropland reserve have been largely overestimated (Lambin et al.,  
501 2013), and they have already degraded during the last decades by inappropriate land  
502 management techniques (Jaurena et al., 2021, De Faccio et al., 2021), lack of mainstreaming  
503 soil conservative techniques (García-Préchac, 2004; Fernandez et al., 2017), decoupling of crop  
504 and livestock (De Faccio et al, 2021) plus climate change impacts with storm water events and  
505 drought, all of which trigger soil erosion (Wingeyer et al. 2015). From the although very limited  
506 point of view on topsoils, the concept of conserving ‘old growth grasslands’ with extensive use  
507 (Veldman et al. 2015) appears a more promising strategy to put the ‘grasslands at the core’ in  
508 the Campos region than the use intensification strategies envisioned by Jaurena et al. (2021).

## 509 **5. Conclusions**

510 The land use intensification in Uruguay associated with increasing inputs of energy, nutrients  
511 and pesticides leads to an overall loss of soil fertility and increasing toxicity related to  
512 acidification, salinization and trace metal contaminants. Our data demonstrate the high amounts  
513 of organic carbon, nutrients and trace metals in topsoil samples from riverine forests,  
514 suggesting transport of soil particles from the surrounding grasslands, crop or timber  
515 plantations to the borders of rivers, streams and creeks. Of all the fertility proxies assessed,  
516 phosphorus in topsoil was most significantly affected by different land uses, being highest in  
517 native forests. Cation exchange capacity was also highest in native forests and lowest in timber  
518 plantations, where only half that of grasslands was measured. Our study highlights that soil  
519 acidification is ongoing and probably also mobilizing trace metals and their accumulation in  
520 riverine forest topsoils.

## 521 **Data availability**

522 All data generated or analysed in this study are included in this article (and its supplementary  
523 materials). Further data are available from the corresponding authors upon reasonable request.

## 524 **Author contributions**

525 IS, LRR, ST and MB conceptualized the study, performed the soil sampling and data analysis.  
526 EZ advised the laboratory analyses. IS wrote the initial draft. LRR, ST, MB and EZ contributed  
527 to generating and reviewing the subsequent versions of the manuscript. IS received the funding  
528 for the study.

529 **Competing interests**

530 The authors declare that they have no conflict of interest

531 **Acknowledgement**

532 In alphabetical order, we thank Juan Barreneche, Lucia Gaucher, Sören Miehe, Nicolas Silvera  
533 and Matías Zarucki for their assistance in field work. We thank Manuel Garcia and Meica  
534 Valdivia for the pre-processing of samples. We thank the staff of the soil laboratory from the  
535 Department of Soil and Natural Resources, University of Concepción, Chile. We thank Diego  
536 de Panis for help with statistical analysis and visualization and valuable comments on a  
537 previous version of this manuscript. Thanks to Vera Krause, Serafina Bischoff, Sophia Reitzug,  
538 Rhea Rennert and Diego Nicolas Rojas for support with data analysis and plots and maps  
539 visualization. We also thank all landowners for access permission to establish our monitoring  
540 sites on their land, their hospitality and willingness to discuss land use goals concerning all  
541 dimension of sustainability. The study was funded by the German Federal Ministry of  
542 Education and Research (BMBF; 01LN1305A). Special thanks go to Amal Chatterjee for  
543 improving our English.

544 **References**

- 545 Alfaro, M., Dube, F. and Zagal, E.: The Influence of Overmature, Degraded Nothofagus  
546 Forests with Strong Anthropogenic Disturbance on the Quality of an Andisol and its Gradual  
547 Recovery with Silvopasture in Southwestern South America, in: *Agroforestry for  
548 Degraded Landscapes*, edited by: Dagar, J.C., Gupta, S.R., Teketay, D. Springer,  
549 Singapore. [https://doi.org/10.1007/978-981-15-6807-7\\_3](https://doi.org/10.1007/978-981-15-6807-7_3), 2020.
- 550 Baize, D. and van Oort, F.: Potentially Harmful Elements in Forest Soils, in: PHEs,

- 551 Environment and Human Health, edited by: Bini, C. and Bech, J., Springer Netherlands,  
552 Dordrecht, 151–198, [https://doi.org/10.1007/978-94-017-8965-3\\_4](https://doi.org/10.1007/978-94-017-8965-3_4), 2014.
- 553 [Benjamini Y. Hochberg. Y. Controlling the False Discovery Rate: A Practical and Powerful](#)  
554 [Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B](#)  
555 [\(Methodological\), 1995; 57\(1\), 289-300. https://www.jstor.org/stable/2346101.](#)
- 556 Beretta-Blanco A, Pérez O, Carrasco-Letelier L. Soil quality decrease over 13 years of  
557 agricultural production. *Nutr Cycling Agroecosyst.* 2019; 114: 45-55.
- 558 Beretta-Blanco, A.; Carrasco-Letelier, L. Relevant factors in the eutrophication of the Uruguay  
559 River and the Río Negro. *Sci. Total Environ.* 2021, 761,143299; DOI:  
560 10.1016/j.scitotenv.2020.143299
- 561 Berhongaray, G; Alvarez R. Soil carbon sequestration of Mollisols and Oxisols under grassland  
562 and tree plantations in South America - A review. *Geoderma Regional* 18 (2019)  
563 e00226.
- 564 Berthrong, S. T., Jobbágy, E. G., and Jackson, R. B.: A global meta-analysis of soil  
565 exchangeable cations, pH, carbon, and nitrogen with afforestation, *Ecological*  
566 *Applications*, 19, 2228–2241, <https://doi.org/10.1890/08-1730.1>, 2009.
- 567 Berthrong, S. T., Piñeiro, G., Jobbágy, E. G., and Jackson, R. B.: Soil C and N changes with  
568 afforestation of grasslands across gradients of precipitation and plantation age, *22, 76–*  
569 *86*, <https://doi.org/10.1890/10-2210.1>, 2012.
- 570 Binkley, Dan et al. The interactions of climate, spacing and genetics on clonal *Eucalyptus*  
571 plantations across Brazil and Uruguay. *Forest Ecology And Management*. Amsterdam:  
572 Elsevier Science Bv, v. 405, p. 271-283, [doi.org/10.1016/j.foreco.2017.09.050](https://doi.org/10.1016/j.foreco.2017.09.050)
- 573 Bonfante, A., Basile, A., and Bouma, J.: Targeting the soil quality and soil health concepts  
574 when aiming for the United Nations Sustainable Development Goals and the EU Green  
575 Deal, *SOIL*, 6, 453–466, <https://doi.org/10.5194/soil-6-453-2020>, 2020.
- 576 Borrelli, P., Robinson, D. A., Fleischer, L. R., Lugato, E., Ballabio, C., Alewell, C.,  
577 Meusburger, K., Modugno, S., Schütt, B., Ferro, V., Bagarello, V., Oost, K. V.,  
578 Montanarella, L., and Panagos, P.: An assessment of the global impact of 21st century  
579 land use change on soil erosion, *Nat Commun*, 8, 2013, [https://doi.org/10.1038/s41467-](https://doi.org/10.1038/s41467-017-02142-7)  
580 [017-02142-7](https://doi.org/10.1038/s41467-017-02142-7), 2017.
- 581 Caon, L., & Vargas, R. (2017). Threats to soils: Global trends and perspectives. A Contribution  
582 from the Intergovernmental Technical Panel on Soils, Global Soil Partnership Food and

hat gelöscht: :

584 Agriculture Organization of the United Nations. In G. Pierzynski & Brajendra (Eds.),  
585 Global land outlook working paper, 28 pp.

586 Carrasco-Letelier, L. and A. Beretta-Blanco. 2017. Soil erosion by water estimated for 99  
587 Uruguayan basins. *Cien. Inv. Agr.* 44(2): 184-194.

588 Céspedes-Payret et al. 2012 *Science of the Total Environment* 438 (2012) 549–557.  
589 <http://dx.doi.org/10.1016/j.scitotenv.2012.08.075>

590 Cook, R. L., Binkley, D., and Stape, J. L.: Eucalyptus plantation effects on soil carbon after  
591 20years and three rotations in Brazil, *Forest Ecology and Management*, 359, 92–98,  
592 <https://doi.org/10.1016/j.foreco.2015.09.035>, 2016.

593 Dass, P. et al 2018 Grasslands may be more reliable carbon sinks than forests in California.  
594 *Environ. Res. Lett.* 13 074027. <https://doi.org/10.1088/1748-9326/aacb39>

595 De Faccio Carvalho, P. C., Savian, J. V., Chiesa, T. D., Souza Filho, W. D., Terra, J. A., Pinto,  
596 P., Martins, A. P., Villarino, S., Da Trindade, J. K., De Albuquerque Nunes, P. A., and  
597 Piñeiro, G.: Land-use intensification trends in the Rio de la Plata region of South  
598 America: toward specialization or recoupling crop and livestock production, *Front.*  
599 *Agr. Sci. Eng.*, 8, 97, <https://doi.org/10.15302/J-FASE-2020380>, 2021.

600 Di Sacco, A.; Hardwick, K.; Blakesley, D.; Brancalion, P.H.S.; Breman, E.; Cecilio Rebola,  
601 L.; Chomba, S.; Dixon, K.; Elliott, S.; Ruyonga, G.; Shaw, K.; Smith, P.; Smith, R.J.;  
602 Antonelli, A. Ten Golden Rules for Reforestation to Optimise Carbon Sequestration,  
603 Biodiversity Recovery and Livelihood Benefits. <https://doi.org/10.1111/gcb.15498>.  
604 2020

605 Durán, A., Morrás, H., Studdert, G., and Liu, X.: Distribution, properties, land use and  
606 management of Mollisols in South America, *Chin. Geogr. Sci.*, 21, 511,  
607 <https://doi.org/10.1007/s11769-011-0491-z>, 2011.

608 ESRI 2018. ArcGIS Desktop: Release 10.3. Redlands, CA: Environmental Systems Research  
609 Institute.  
610 [https://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcs142p2\\_051232.pdf](https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051232.pdf)

611 FAO 2019. Land Use. Published online at OurWorldInData.org. Retrieved from:  
612 '<https://ourworldindata.org/land-use>' [Online Resource]

613 Fernández, R., Frasier, I., Noellemeyer, E., and Quiroga, A.: Soil quality and productivity  
614 under zero tillage and grazing on Mollisols in Argentina – A long-term study,  
615 *Geoderma Regional*, 11, 44–52, <https://doi.org/10.1016/j.geodrs.2017.09.002>, 2017.



- 616 Fox, J., Weisberg, S. (2019) An R Companion to Applied Regression, Third Edition, Sage.
- 617 Friggens, N. L., Hester, A. J., Mitchell, R. J., Parker, T. C., Subke, J., and Wookey, P. A.: Tree  
618 planting in organic soils does not result in net carbon sequestration on decadal  
619 timescales, *Glob Change Biol*, 26, 5178–5188, <https://doi.org/10.1111/gcb.15229>,  
620 2020.
- 621 García-Préchac, F., Ernst, O., Siri-Prieto, G., and Terra, J. A.: Integrating no-till into crop-  
622 pasture rotations in Uruguay, *Soil and Tillage Research*, 77, 1–13,  
623 <https://doi.org/10.1016/j.still.2003.12.002>, 2004.
- 624 Gomiero, Tiziano. 2016. "Soil Degradation, Land Scarcity and Food Security: Reviewing a  
625 Complex Challenge" *Sustainability* 8, no. 3: 281. <https://doi.org/10.3390/su8030281>
- 626 Hernández J, del Pino A, Salvo L, Arrarte G. Nutrient export and harvest residue  
627 decomposition patterns of a *Eucalyptus dunnii* Maiden plantation in temperate climate  
628 of Uruguay. *Forest Ecology Management* 2009; 258(2): 92-99.  
629 [10.1016/j.foreco.2009.03.050](https://doi.org/10.1016/j.foreco.2009.03.050)
- 630 Hernández, J., del Pino, A., Vance, E. D., Califra, Á., Del Giorgio, F., Martínez, L., and  
631 González-Barrios, P.: *Eucalyptus* and *Pinus* stand density effects on soil carbon  
632 sequestration, *Forest Ecology and Management*, 368, 28–38,  
633 <https://doi.org/10.1016/j.foreco.2016.03.007>, 2016.
- 634 Hernandez-Ramirez, G., Sauer, T. J., Chendev, Y. G., and Gennadiev, A. N.: Nonlinear  
635 turnover rates of soil carbon following cultivation of native grasslands and subsequent  
636 afforestation of croplands, *SOIL*, 7, 415–431, <https://doi.org/10.5194/soil-7-415-2021>,  
637 2021.
- 638 IUSS Working Group WRB. 2015. World Reference Base for Soil Resources 2014, update  
639 2015 International soil classification system for naming soils and creating legends for  
640 soil maps. World Soil Resources Reports No. 106. FAO, Rome.  
641 <https://www.fao.org/3/i3794en/i3794en.pdf>
- 642 Jackson et al 2005 Trading Water for Carbon with Biological Carbon Sequestration. *Science*  
643 310, 1944 (2005); DOI: 10.1126/science.1119282
- 644 Jackson, R.: Soil nutrient stocks are maintained over multiple rotations in Brazilian *Eucalyptus*  
645 plantations | Elsevier Enhanced Reader, 13, n.d.
- 646 Jaurena M, Durante M, Devincenzi T, Savian JV, Bendersky D, Moojen FG, Pereira M, Soca  
647 P, Quadros FLF, Pizzio R, Nabinger C, Carvalho PCF and Lattanzi FA (2021) Native  
648 Grasslands at the Core: A New Paradigm of Intensification for the Campos of Southern

- 649 South America to Increase Economic and Environmental Sustainability. *Front. Sustain.*  
650 *Food Syst.* 5:547834. doi: 10.3389/fsufs.2021.547834
- 651 Jobbagy, E. G., and R. B. Jackson. 2003. Patterns and mechanisms of soil acidification in the  
652 conversion of grasslands to forests. *Biogeochemistry* 64:205–229.
- 653 Jobbagy, E. G., and R. B. Jackson. 2004. The uplift of soil nutrients by plants: biogeochemical  
654 consequences across scales. *Ecology* 85:2380–2389.
- 655 Kaushal, S. S., Gold, A. J., Bernal, S., Johnson, T. A. N., Addy, K., Burgin, A., Burns, D. A.,  
656 Coble, A. A., Hood, E., Lu, Y., Mayer, P., Minor, E. C., Schroth, A. W., Vidon, P.,  
657 Wilson, H., Xenopoulos, M. A., Doody, T., Galella, J. G., Goodling, P., Haviland, K.,  
658 Haq, S., Wessel, B., Wood, K. L., Jaworski, N., and Belt, K. T.: Watershed ‘chemical  
659 cocktails’: forming novel elemental combinations in Anthropocene fresh waters,  
660 *Biogeochemistry*, 141, 281–305, <https://doi.org/10.1007/s10533-018-0502-6>, 2018.
- 661 Kellogg, C. E. "Soil survey division staff: soil survey manual. Chapter 3" United States  
662 Department of Agriculture, Washington (1993).
- 663 Kim, K., Daly, E.J., M. Gorzelak, Hernandez-Ramirez, G. 2022. Soil organic matter pools  
664 response to perennial grain cropping and nitrogen fertilizer. *Soil and Tillage Research.*  
665 220, 105376 <https://doi.org/10.1016/j.still.2022.105376>
- 666 Lambin, E.F.; Gibbs, H.K.; Ferreira, L.; Grau, R.; Mayaux, P.; Meyfroidt, P.; Morton, D.C.;  
667 Rudel, T.K.; Gasparri, I.; Munger, J. Estimating the world’s potentially available  
668 cropland using a bottom-up approach. *Glob. Environ. Chang.* 2013, 23, 892–901.
- 669 Lanfranco, B. and Sapriza, G.: El índice CONEAT como medida de productividad y valor de  
670 la tierra., 57, n.d.
- 671 Lavado, R.S., Rodríguez, M.B., Scheiner, S.D., Taboada, M.A., Rubio, G., Alvarez, R.,  
672 Alconada, M., Zubillaga, M.S., 1998. Heavy metals in soils of Argentina: comparison  
673 between urban and agricultural soils. *Communications in Soil Science and Plant*  
674 *Analysis* 29, 1913–1917.
- 675 Lavado, R.S., Zubillaga, M.S., Alvarez, R., Taboada, M.A., 2004. Baseline levels of potentially  
676 toxic elements in pampas soils. *Soil and Sediment Contamination: An International*  
677 *Journal* 13, 329–339.
- 678 Li, H., Jiang, L., You, C., Tan, B., and Yang, W.: Dynamics of heavy metal uptake and soil  
679 heavy metal stocks across a series of Masson pine plantations, *Journal of Cleaner*  
680 *Production*, 269, 122395, <https://doi.org/10.1016/j.jclepro.2020.122395>, 2020.

- 681 Liu, X., Lee Burras, C., Kravchenko, Y. S., Durán, A., Huffman, T., Morras, H., Studdert, G.,  
682 Zhang, X., Cruse, R. M., and Yuan, X.: Overview of Mollisols in the world:  
683 Distribution, land use and management, *Can. J. Soil. Sci.*, 92, 383–402,  
684 <https://doi.org/10.4141/cjss2010-058>, 2012.
- 685 Mayer M., Prescott C.E., Abaker W.E.A., Augusto L., Cécillon L., Ferreira G.W.D., James J.,  
686 Jandl R., Katzensteiner K., Laclau J.-P., Laganière J., Nouvellon Y., Paré D., Stanturf  
687 J.A., Vanguelova E.I., Vesterdal L. Influence of forest management activities on soil  
688 SOC stocks: A knowledge synthesis. *Forest Ecology and Management*, Volume 466,  
689 2020
- 690 McMahon, D. E., Vergütz, L., Valadares, S. V., Silva, I. R. da, and Jackson, R. B.: Soil nutrient  
691 stocks are maintained over multiple rotations in Brazilian Eucalyptus plantations,  
692 *Forest Ecology and Management*, 448, 364–375,  
693 <https://doi.org/10.1016/j.foreco.2019.06.027>, 2019.
- 694 Merino, A., Balboa, M. A., Rodríguez Soalleiro, R., and González, J. G. Á.: Nutrient exports  
695 under different harvesting regimes in fast-growing forest plantations in southern  
696 Europe, *Forest Ecology and Management*, 207, 325–339,  
697 <https://doi.org/10.1016/j.foreco.2004.10.074>, 2005.
- 698 MGAP (Ministerio de Ganadería, Agricultura y Pesca) 2020. CONEAT. Online:  
699 <https://www.gub.uy/ministerio-ganaderia-agricultura-pesca/politicas-y-gestion/coneat>.
- 700 Modernel, P., Rossing, W. A. H., Corbeels, M., Dogliotti, S., Picasso, V., and Tiftonell, P.:  
701 Land use change and ecosystem service provision in Pampas and Campos grasslands  
702 of southern South America, *Environ. Res. Lett.*, 11, 113002,  
703 <https://doi.org/10.1088/1748-9326/11/11/113002>, 2016.
- 704 OAS (Organization of American States, Department of Sustainable Development) 1994.  
705 Uruguay - Proyecto Regional de Alternativas para la Inversión Forestal. Available  
706 online: <http://www.oas.org/dsd/publications/unit/oea20s/oea20s.pdf>.
- 707 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.  
708 R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and  
709 Wagner, H.: *vegan: Community Ecology Package*, 2020.
- 710 Ortiz, J., Dube, F., Neira, P., Panichini, M., Stolpe, N.B., Zagal, E. and Martínez-Hernández,  
711 P.A. Soil Quality Changes within a (*Nothofagus obliqua*) Forest Under Silvopastoral  
712 Management in the Andes Mountain Range, South Central Chile. *Sustainability*;  
713 12(17):6815. <https://doi.org/10.3390/su12176815>, 2020.

hat formatiert: Spanisch

hat formatiert: Deutsch

- 714 Paul, K. I., Polglase, P. J., Nyakuengama, J. G., and Khanna, P. K.: Change in soil carbon  
715 following afforestation, *Forest Ecology and Management*, 168, 241–257,  
716 [https://doi.org/10.1016/S0378-1127\(01\)00740-X](https://doi.org/10.1016/S0378-1127(01)00740-X), 2002.
- 717 Piñeiro, D. E.: Land grabbing: concentration and “foreignisation” of land in Uruguay, *Canadian*  
718 *Journal of Development Studies / Revue canadienne d'études du développement*, 33,  
719 471–489, <https://doi.org/10.1080/02255189.2012.746216>, 2012.
- 720 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for  
721 Statistical Computing, Vienna, Austria, available at: <https://www.R-project.org/>, last  
722 access: 2 March 2021.
- 723 Richter, D. D.: Game Changer in Soil Science. The Anthropocene in soil science and  
724 pedology., *J. Plant Nutr. Soil Sci.*, 183, 5–11, <https://doi.org/10.1002/jpln.201900320>,  
725 2020.
- 726 Roca, N.: Heavy metal background levels in rural soils: a case study in Pampean soils  
727 (Argentina), 10, 2015.
- 728 [Sadzawka, A., Carrasco, R. M. A., Grez, Z. R., Mora, G. M. Flores, P. H. Neaman, A. 2006.](https://hdl.handle.net/20.500.14001/8541)  
729 [Métodos de análisis recomendados para los suelos de Chile. Revisión 2006. Santiago,](https://hdl.handle.net/20.500.14001/8541)  
730 [Chile: Serie Actas - Instituto de Investigaciones Agropecuarias. Available online:](https://hdl.handle.net/20.500.14001/8541)  
731 [https://hdl.handle.net/20.500.14001/8541.](https://hdl.handle.net/20.500.14001/8541)
- 732 Sandoval López, D.M., Arturi, M.F., Goya, J.F. et al. *Eucalyptus grandis* plantations: effects  
733 of management on soil carbon, nutrient contents and yields. *J. For. Res.* 31, 601–611  
734 (2020). <https://doi.org/10.1007/s11676-018-0850-z>.
- 735 Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and  
736 interpreting soil surveys. 2nd edition.  
737 [https://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcs142p2\\_051232.pdf](https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051232.pdf).
- 738 Veldman, J. W., Overbeck, G. E., Negreiros, D., Mahy, G., Le Stradic, S., Fernandes, G. W.,  
739 Durigan, G., Buisson, E., Putz, F. E., and Bond, W. J.: Where Tree Planting and Forest  
740 Expansion are Bad for Biodiversity and Ecosystem Services, *BioScience*, 65, 1011–  
741 1018, <https://doi.org/10.1093/biosci/biv118>, 2015.
- 742 Waters, C. N., Zalasiewicz, J., Summerhayes, C., Barnosky, A. D., Poirier, C., Ga uszka, A.,  
743 Cearreta, A., Edgeworth, M., Ellis, E. C., Ellis, M., Jeandel, C., Leinfelder, R., McNeill,  
744 J. R., Richter, D. d., Steffen, W., Syvitski, J., Vidas, D., Wagemich, M., Williams, M.,  
745 Zhisheng, A., Grinevald, J., Odada, E., Oreskes, N., and Wolfe, A. P.: The

Feldfunktion geändert

746 Anthropocene is functionally and stratigraphically distinct from the Holocene, *Science*,  
747 351, aad2622–aad2622, <https://doi.org/10.1126/science.aad2622>, 2016.

748 Wingeyer, A.B.; Amado, T.J.C.; Pérez-Bidegain, M.; Studdert, G.A.; Perdomo Varela, C.H.;  
749 Garcia, F.O.; Karlen, D.L. Soil Quality Impacts of Current South American  
750 Agricultural Practices. *Sustainability* 2015, 7, 2213–2242.


751 Wu, R., Alvareda, E., Polya, D., Blanco, G., and Gamazo, P.: Distribution of Groundwater  
752 Arsenic in Uruguay Using Hybrid Machine Learning and Expert System Approaches,  
753 *Water*, 13, 527, <https://doi.org/10.3390/w13040527>, 2021.





754 Zagal, E., Sadzawka, A. 2007. Protocolo de métodos de análisis para suelos y lodos. Available  
755 online: [https://www.sag.cl/sites/default/files/METODOS\\_LODOS\\_SUELOS.pdf](https://www.sag.cl/sites/default/files/METODOS_LODOS_SUELOS.pdf).

756 Zurbriggen, C., González-Lago, M., Baraibar, M., Baethgen, W., Mazzeo, N. and Sierra, M.,  
757 2020. Experimentation in the Design of Public Policies: The Uruguayan Soils  
758 Conservation Plans. *Iberoamericana – Nordic Journal of Latin American and*  
759 *Caribbean Studies*, 49(1), pp.52–62. DOI: <http://doi.org/10.16993/iberoamericana.459>.

hat formatiert: Deutsch



hat formatiert: Deutsch





763 **Figure 1:** Study design and sampling. (a) Location of the 28 monitoring sites across Uruguay  
764 including land use types sampled (grassland (b); timber plantations, native forest and  
765 agricultural land. Proportion of plots with particular category of soil classification according to  
766 the World Reference Base (WRB, c), Soil Taxonomy (d) and CONEAT (Comisión Nacional  
767 de Estudio Agronómico de la Tierra, e). [The Uruguayan CONEAT index provides a detailed](#)  
768 [classification that takes into account soil type, texture, natural vegetation, altitude and geology](#)  
769 [\(see details in chapter 2.3\)](#). Photos: RuralFutures.

770



hat gelöscht:


hat gelöscht: Colour intensity and the size of the circle are proportional to the correlation coefficients ( $\rho$ )

771

772 **Figure 2.** Spearman's rank correlations for parameters of topsoils (n=80) regarding (a) fertility  
 773 proxies, (b) acidity and (c) trace metals. The size of the circle refers to the absolute value of  
 774 the correlation coefficient ( $\rho$ ) and the color refers to the direction (positive or negative) of the  
 775 correlation. Empty slots show correlations with  $p > 0.05$ .


776






780


781 **Figure 3.** Non-metric multidimensional scaling showing significant clustering differences  
782 among samples from (a) grassland (GL), timber plantations (TP) and native forests (NF); (b)  
783 among samples from *Pinus* and *Eucalyptus* plantations; (c) among land uses (including  
784 agriculture AC) in Argiudolls; (d) among land uses in Argiudolls & Hapluderts; (e) among land  
785 uses in Argiudolls, Hapudolls & Hapludalf and (f) among land uses in Hapludalfs &  
786 Hapludults.  
787



789 **Figure 4.** Violin box plots for significant Kruskal-Wallis Tests across evaluated land uses (i.e.  
790 grassland (GL), timber plantation (TP) and native forest (NF)) for each variable. Significance  
791 in posterior Wilcoxon pairwise comparisons is depicted as  $p < 0.05$  (\*),  $p < 0.01$  (\*\*),  $p < 0.001$   
792 (\*\*\*),  $p < 0.0001$  (\*\*\*\*).  
793



795 **Figure 5.** Violin box plots for significant Kruskal-Wallis Tests in ‘Argiudolls’ Soil Taxonomy  
 796 category for fertility variables across available land uses (GL: Grassland, TP: Timber  
 797 plantation, NF: Native forest, AC: Agriculture). Significance in posterior Wilcoxon pairwise  
 798 comparisons is depicted as  $p < 0.05$  (\*),  $p < 0.01$  (\*\*),  $p < 0.001$  (\*\*\*),  $p < 0.0001$  (\*\*\*\*).



799 **Figure 6.** Acidity of all topsoils samples (a) and (b) according to different land use (GL:  
 800 Grassland, TP: Timber plantation, NF: Native forest, AC: Agriculture). Proportion of samples  
 801 with a given acidic category (Kellogg, 1993). pH of topsoil given as violin box plots for  
 802

803 significant Kruskal-Wallis Tests across different land uses independent from soil Order (c) and  
804 separated per soil Orders (d-f). Sample number (n) is given. Significance in posterior Wilcoxon  
805 pairwise comparisons is depicted as  $p < 0.05$  (\*),  $p < 0.01$  (\*\*),  $p < 0.001$  (\*\*\*),  $p < 0.0001$  (\*\*\*\*).  
806

807 **Table 1.** General characteristics of Uruguayan topsoils: descriptive statistics for the parameters  
 808 of all single samples (n) across different land uses and classification to different soil types (for  
 809 details on different land uses i.e. grassland (GL), timber plantations (TP), native forests (NF)  
 810 and agricultural land (AC) see Appendix A1-A2 and on different soil types see Appendix A3-  
 811 4). SD = standard derivation.

hat gelöscht: crops

| Variable   | n   | Mean±SD   | Min-Max   | Samples by land use |    |    |    |
|--|-----|-----------|-----------|---------------------|----|----|----|
|  |     |           |           | GL                  | TP | NF | AC |
| Soil moisture (%)  | 280 | 18.9±10.8 | 1.34-51.5 | 115                 | 93 | 57 | 15 |
| N <sub>total</sub> (%)   | 280 | 0.2±0.14  | 0.04-1.2  | 115                 | 93 | 57 | 15 |
| C <sub>total</sub> (%)   | 280 | 3±2.3     | 0.4-25.7  | 115                 | 93 | 57 | 15 |
| C/N ratio  | 279 | 13.3±1.8  | 9.1-20.6  | 115                 | 93 | 56 | 15 |
| SOC (%)  | 267 | 5.6±3     | 0.8-16.4  | 110                 | 90 | 52 | 15 |
| P (mg kg <sup>-1</sup> )   | 278 | 295±166   | 43-1008   | 114                 | 93 | 57 | 14 |
| Ca (cmol kg <sup>-1</sup> )  | 82  | 9.4±8.9   | 0.3-42.7  | 32                  | 27 | 18 | 5  |
| Mg (cmol kg <sup>-1</sup> )  | 82  | 2.5±2.2   | 0.1-9.7   | 32                  | 27 | 18 | 5  |
| K (cmol kg <sup>-1</sup> )   | 82  | 0.4±0.25  | 0.03-1.44 | 32                  | 27 | 18 | 5  |
| Na (cmol kg <sup>-1</sup> )  | 82  | 0.1±0.14  | 0-0.69    | 32                  | 27 | 18 | 5  |
| Zn (mg kg <sup>-1</sup> )  | 82  | 1.9±2.1   | 0.1-9.7   | 32                  | 27 | 18 | 5  |
| Cu (mg kg <sup>-1</sup> )  | 82  | 5.4±22.1  | 0.3-161.2 | 32                  | 27 | 18 | 5  |
| Fe (mg kg <sup>-1</sup> )  | 82  | 134±75    | 5-309     | 32                  | 27 | 18 | 5  |
| Mn (mg kg <sup>-1</sup> )  | 81  | 25±17     | 0.7-93    | 32                  | 27 | 17 | 5  |
| pH   | 279 | 4.8±0.8   | 3.6-7.34  | 114                 | 93 | 57 | 15 |
| As (mg kg <sup>-1</sup> )  | 279 | 3.9±3.6   | 0.6-30.7  | 115                 | 93 | 56 | 15 |
| Cd (mg kg <sup>-1</sup> )  | 274 | 0.4±0.2   | 0.2-1     | 111                 | 91 | 57 | 15 |
| Cr (mg kg <sup>-1</sup> )  | 280 | 13.6±12.6 | 2-78.9    | 115                 | 93 | 57 | 15 |
| Pb (mg kg <sup>-1</sup> )  | 280 | 7±3.4     | 2-27.6    | 115                 | 93 | 57 | 15 |
| CEC (cmol kg <sup>-1</sup> )   | 82  | 12.4±10.7 | 0.5-50.1  | 32                  | 27 | 18 | 5  |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> )                  | 82  | 0.07±0.08 | 0.01-0.43 | 32                  | 27 | 18 | 5  |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> +Na <sup>+</sup> ) | 82  | 0.07±0.07 | 0.01-0.42 | 32                  | 27 | 18 | 5  |

hat gelöscht: 5.6

hat gelöscht: 295

hat gelöscht: 9.4

hat gelöscht: 2.5

hat gelöscht: 0.4

hat gelöscht: 0.1

hat gelöscht: 1.9

hat gelöscht: 5.4

hat gelöscht: 134

hat gelöscht: 25

hat gelöscht: 4.8

hat gelöscht: 3.9

hat gelöscht: 0.4

hat gelöscht: 13.6

hat gelöscht: 7

hat gelöscht: 12.4

hat gelöscht: 0.07

hat gelöscht: 0.07

812

813



833

## Appendices

### 834 Appendix A

835 **Table A1.** Descriptive statistics of topsoil variables for all single soil types at grassland and  
 836 native forests. Number of samples (n), mean, standard deviation (SD), minimum (Min) and  
 837 maximum (Max) of each variable are given.

838

| Variable (unit)  | Grassland |      |      |      |      | Native forests |      |      |      |      |
|--|-----------|------|------|------|------|----------------|------|------|------|------|
|  | n         | Mean | SD   | Min  | Max  | n              | Mean | SD   | Min  | Max  |
| Soil moisture (%)  | 115       | 20.4 | 11.5 | 1.5  | 51.5 | 57             | 25.8 | 9.3  | 6.3  | 50.3 |
| N <sub>total</sub> (%)   | 115       | 0.22 | 0.13 | 0.06 | 0.80 | 57             | 0.30 | 0.19 | 0.04 | 1.20 |
| C <sub>total</sub> (%)   | 115       | 2.9  | 1.9  | 0.7  | 10.8 | 57             | 4.2  | 3.6  | 0.5  | 25.7 |
| C/N ratio  | 115       | 13.2 | 1.7  | 9.1  | 19.4 | 56             | 13.5 | 1.4  | 9.8  | 18.5 |
| SOC (%)  | 109       | 5.3  | 2.6  | 1.4  | 15.2 | 53             | 7.5  | 3.8  | 1.1  | 16.4 |
| P (mg/kg)  | 114       | 272  | 124  | 43   | 749  | 57             | 484  | 194  | 56   | 1009 |
| Ca (cmol kg <sup>-1</sup> )  | 32        | 9.5  | 10.0 | 0.3  | 42.7 | 18             | 15.1 | 10.0 | 0.3  | 37.3 |
| Mg (cmol kg <sup>-1</sup> )  | 32        | 2.2  | 2.1  | 0.18 | 9.7  | 18             | 4.8  | 2.6  | 0.12 | 8.9  |
| K (cmol kg <sup>-1</sup> )   | 32        | 0.40 | 0.26 | 0.08 | 1.44 | 18             | 0.50 | 0.28 | 0.03 | 1.29 |
| Na (cmol kg <sup>-1</sup> )  | 32        | 0.12 | 0.15 | 0.00 | 0.65 | 18             | 0.14 | 0.12 | 0.00 | 0.50 |
| Zn (mg kg <sup>-1</sup> )  | 32        | 1.4  | 1.2  | 0.10 | 4.7  | 18             | 4.0  | 2.7  | 0.80 | 9.60 |
| Cu (mg kg <sup>-1</sup> )  | 32        | 2.2  | 1.8  | 0.30 | 7.7  | 18             | 2.4  | 1.5  | 0.30 | 5.50 |
| Fe (mg kg <sup>-1</sup> )  | 32        | 121  | 81   | 5    | 279  | 18             | 149  | 76   | 10   | 309  |
| Mn (mg kg <sup>-1</sup> )  | 32        | 22   | 18   | 2    | 93   | 17             | 33   | 18   | 1    | 72   |
| pH   | 114       | 4.9  | 0.9  | 3.6  | 7.3  | 57             | 5.1  | 0.7  | 3.6  | 7.2  |
| As (mg kg <sup>-1</sup> )  | 115       | 4.3  | 4.5  | 0.6  | 30.7 | 56             | 4.1  | 2.5  | 1.2  | 13.6 |
| Cd (mg kg <sup>-1</sup> )  | 111       | 0.41 | 0.17 | 0.17 | 1.00 | 57             | 0.45 | 0.15 | 0.17 | 0.81 |
| Cr (mg kg <sup>-1</sup> )  | 115       | 12.3 | 9.2  | 2.6  | 49.5 | 57             | 22.8 | 17.3 | 2.0  | 70.7 |
| Pb (mg kg <sup>-1</sup> )  | 115       | 6.8  | 3.0  | 2.3  | 15.9 | 57             | 9.5  | 4.8  | 2.9  | 27.7 |
| CEC (cmol/ kg <sup>-1</sup> )  | 32        | 12.2 | 11.5 | 0.8  | 50.1 | 18             | 20.4 | 11.4 | 0.5  | 44.6 |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> )                  | 32        | 0.08 | 0.10 | 0.01 | 0.43 | 18             | 0.03 | 0.02 | 0.01 | 0.07 |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> +Na <sup>+</sup> ) | 32        | 0.08 | 0.10 | 0.01 | 0.42 | 18             | 0.03 | 0.02 | 0.01 | 0.07 |

839

840 **Table A2.** Descriptive statistics for of topsoil variables for all single soil types at timber  
 841 plantations and agricultural land. Number of samples (n), mean, standard deviation (SD),  
 842 minimum (Min) and maximum (Max) of each variable are given.

hat gelöscht: crops

843

| Variable (unit)  | Timber plantations |      |      |      |      | <u>Agricultural land</u> |      |      |      |       |
|--|--------------------|------|------|------|------|--------------------------|------|------|------|-------|
|  | n                  | Mean | SD   | Min  | Max  | n                        | Mean | SD   | Min  | Max   |
| Soil moisture (%)  | 93                 | 13.2 | 7.0  | 1.7  | 38.0 | 15                       | 18.1 | 12.0 | 1.3  | 50.5  |
| N <sub>total</sub> (%)   | 93                 | 0.18 | 0.10 | 0.04 | 0.61 | 15                       | 0.22 | 0.06 | 0.13 | 0.29  |
| C <sub>total</sub> (%)   | 93                 | 2.5  | 1.5  | 0.4  | 8.4  | 15                       | 2.6  | 0.7  | 1.4  | 3.3   |
| C/N ratio  | 93                 | 13.6 | 2.1  | 10.0 | 20.6 | 15                       | 11.7 | 0.7  | 10.5 | 12.8  |
| SOC (%)  | 90                 | 4.9  | 2.8  | 0.8  | 16.1 | 15                       | 5.2  | 1.3  | 2.8  | 6.7   |
| P (mg kg <sup>-1</sup> )   | 93                 | 206  | 91   | 56   | 551  | 14                       | 310  | 120  | 142  | 613   |
| Ca (cmol kg <sup>-1</sup> )  | 27                 | 5.2  | 4.8  | 0.6  | 18   | 5                        | 10.8 | 5.5  | 7.2  | 20.4  |
| Mg (cmol kg <sup>-1</sup> )  | 27                 | 1.4  | 1.0  | 0.16 | 4.1  | 5                        | 2.5  | 1.0  | 1.76 | 4.1   |
| K (cmol kg <sup>-1</sup> )   | 27                 | 0.32 | 0.21 | 0.09 | 0.93 | 5                        | 0.46 | 0.23 | 0.29 | 0.85  |
| Na (cmol kg <sup>-1</sup> )  | 27                 | 0.11 | 0.16 | 0.00 | 0.69 | 5                        | 0.06 | 0.02 | 0.03 | 0.08  |
| Zn (mg kg <sup>-1</sup> )  | 27                 | 0.94 | 0.74 | 0.10 | 2.8  | 5                        | 3.7  | 3.7  | 0.60 | 9.7   |
| Cu (mg kg <sup>-1</sup> )  | 27                 | 1.5  | 0.9  | 0.5  | 4.5  | 5                        | 57.7 | 78.8 | 0.8  | 161.0 |
| Fe (mg kg <sup>-1</sup> )  | 27                 | 143  | 69   | 16   | 290  | 5                        | 115  | 62   | 63   | 210   |
| Mn (mg kg <sup>-1</sup> )  | 27                 | 25   | 16   | 6    | 72   | 5                        | 17   | 5    | 10   | 23    |
| pH   | 93                 | 4.5  | 0.7  | 3.6  | 6.8  | 15                       | 5.1  | 0.4  | 4.2  | 5.7   |
| As (mg kg <sup>-1</sup> )  | 93                 | 3.4  | 3.0  | 0.7  | 25.5 | 15                       | 3.1  | 1.2  | 1.1  | 5.5   |
| Cd (mg kg <sup>-1</sup> )  | 91                 | 0.33 | 0.11 | 0.16 | 0.65 | 15                       | 0.41 | 0.14 | 0.21 | 0.66  |
| Cr (mg kg <sup>-1</sup> )  | 93                 | 10.6 | 11.3 | 2.0  | 78.9 | 15                       | 7.8  | 1.8  | 5.0  | 11.3  |
| Pb (mg kg <sup>-1</sup> )  | 93                 | 6.1  | 2.3  | 2.0  | 11.7 | 15                       | 5.1  | 1.2  | 3.4  | 7.3   |
| CEC (cmol kg <sup>-1</sup> )   | 27                 | 7.0  | 5.6  | 1.0  | 20.6 | 5                        | 13.8 | 6.5  | 9.5  | 25.0  |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> )                  | 27                 | 0.08 | 0.07 | 0.02 | 0.25 | 5                        | 0.04 | 0.03 | 0.02 | 0.09  |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> +Na <sup>+</sup> ) | 27                 | 0.08 | 0.07 | 0.02 | 0.25 | 5                        | 0.04 | 0.03 | 0.02 | 0.08  |

hat gelöscht: Crops

844

845

846

849 **Table A3.** Descriptive statistics of topsoil variables for Argiudolls and for grassland and native  
 850 forests within of Argiudolls. Number of single samples (n), mean, standard deviation (SD),  
 851 minimum (Min) and maximum (Max) of each variable are given.

852

| Variable (Unit)  | Total Argiudolls |      |      |      |      | Grassland |      |      |      |      | Native forests |      |      |      |      |
|--|------------------|------|------|------|------|-----------|------|------|------|------|----------------|------|------|------|------|
|  | n                | Mean | SD   | Min  | Max  | n         | Mean | SD   | Min  | Max  | n              | Mean | SD   | Min  | Max  |
| Soil moisture (%)  | 77               | 18.3 | 10.9 | 1.3  | 50.5 | 31        | 19.0 | 9.2  | 5.0  | 49.0 | 12             | 26.4 | 13.4 | 9.0  | 48.2 |
| N <sub>total</sub> (%)   | 77               | 0.24 | 0.15 | 0.07 | 1.20 | 31        | 0.21 | 0.11 | 0.07 | 0.66 | 12             | 0.34 | 0.28 | 0.13 | 1.20 |
| C <sub>total</sub> (%)   | 77               | 3.3  | 3.0  | 1.1  | 25.7 | 31        | 2.9  | 1.8  | 1.1  | 10.8 | 12             | 5.3  | 6.5  | 1.7  | 25.7 |
| C/N ratio  | 76               | 13.3 | 2.2  | 9.8  | 19.9 | 31        | 13.3 | 2.1  | 10.5 | 19.4 | 11             | 13.6 | 2.1  | 9.8  | 18.5 |
| SOC (%)  | 73               | 5.8  | 2.5  | 2.1  | 16.1 | 30        | 5.2  | 1.9  | 2.1  | 10.0 | 9              | 7.4  | 2.1  | 4.8  | 11.0 |
| P (mg kg <sup>-1</sup> )   | 76               | 280  | 122  | 93   | 693  | 30        | 234  | 78   | 93   | 353  | 12             | 438  | 155  | 155  | 693  |
| Ca (cmol kg <sup>-1</sup> )  | 21               | 11.4 | 8.0  | 2.6  | 37.1 | 8         | 11.5 | 6.7  | 3.2  | 20.5 | 4              | 16.2 | 14.3 | 5.1  | 37.1 |
| Mg (cmol kg <sup>-1</sup> )  | 21               | 2.3  | 1.4  | 0.5  | 6.1  | 8         | 2.0  | 0.8  | 0.9  | 3.2  | 4              | 3.5  | 2.5  | 1.0  | 6.1  |
| K (cmol kg <sup>-1</sup> )   | 21               | 0.45 | 0.26 | 0.1  | 1.3  | 8         | 0.4  | 0.14 | 0.16 | 0.61 | 4              | 0.69 | 0.41 | 0.34 | 1.29 |
| Na (cmol kg <sup>-1</sup> )  | 21               | 0.08 | 0.05 | 0.03 | 0.26 | 8         | 0.07 | 0.02 | 0.04 | 0.10 | 4              | 0.10 | 0.11 | 0.03 | 0.26 |
| Zn (mg kg <sup>-1</sup> )  | 21               | 2.4  | 3.0  | 0.1  | 10   | 8         | 1.3  | 0.8  | 0.1  | 2.2  | 4              | 5.1  | 4.3  | 1.3  | 10   |
| Cu (mg kg <sup>-1</sup> )  | 21               | 15.5 | 42.7 | 0.8  | 161  | 8         | 2.6  | 1.6  | 1.0  | 6.0  | 4              | 1.9  | 1.0  | 0.8  | 3.0  |
| Fe (mg kg <sup>-1</sup> )  | 21               | 117  | 62   | 10   | 232  | 8         | 104  | 50   | 28   | 171  | 4              | 117  | 84   | 10   | 216  |
| Mn (mg kg <sup>-1</sup> )  | 21               | 23.3 | 10.4 | 7.1  | 43.6 | 8         | 20.5 | 10.4 | 7.1  | 33.8 | 4              | 28.8 | 10.4 | 15.4 | 39.4 |
| pH   | 77               | 5.0  | 0.7  | 3.8  | 6.6  | 31        | 4.9  | 0.7  | 3.9  | 6.6  | 12             | 5.2  | 0.5  | 4.3  | 6.1  |
| As (mg kg <sup>-1</sup> )  | 76               | 3.2  | 1.4  | 1.0  | 12.2 | 31        | 2.9  | 0.9  | 1.5  | 4.9  | 11             | 4.0  | 2.8  | 1.5  | 12.2 |
| Cd (mg kg <sup>-1</sup> )  | 76               | 0.4  | 0.1  | 0.2  | 0.7  | 30        | 0.4  | 0.1  | 0.2  | 0.7  | 12             | 0.4  | 0.1  | 0.2  | 0.7  |
| Cr (mg kg <sup>-1</sup> )  | 77               | 9.3  | 6.2  | 2.2  | 28.3 | 31        | 8.1  | 4.1  | 3.6  | 18.5 | 12             | 13.2 | 9.5  | 2.2  | 28.3 |
| Pb (mg kg <sup>-1</sup> )  | 77               | 5.6  | 2.1  | 2.8  | 10.7 | 31        | 5.2  | 1.8  | 2.8  | 9.6  | 12             | 6.8  | 2.9  | 2.9  | 10.7 |
| CEC (cmol kg <sup>-1</sup> )   | 21               | 14.2 | 9.3  | 3.4  | 44.6 | 8         | 14.0 | 7.3  | 4.5  | 23.1 | 4              | 20.4 | 16.7 | 6.4  | 44.6 |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> )                  | 21               | 0.04 | 0.02 | 0.02 | 0.09 | 8         | 0.03 | 0.02 | 0.02 | 0.07 | 4              | 0.04 | 0.01 | 0.03 | 0.06 |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> +Na <sup>+</sup> ) | 21               | 0.04 | 0.02 | 0.02 | 0.08 | 8         | 0.03 | 0.02 | 0.02 | 0.07 | 4              | 0.04 | 0.01 | 0.03 | 0.06 |

853  
854

855 **Table A4.** Descriptive statistics of topsoil variables for Argiudolls and timber plantations and  
 856 Agricultural land within of Argiudolls. Number of single samples (n), mean, standard deviation  
 857 (SD), minimum (Min) and maximum (Max) of each variable are given.

hat gelöscht: crops

858

| Variable (unit)  | Total Argiudolls |      |      |      |      | Timber plantations |      |      |      |      | <u>Agricultural land</u> |      |      |      |      |
|--|------------------|------|------|------|------|--------------------|------|------|------|------|--------------------------|------|------|------|------|
|  | n                | Mean | SD   | Min  | Max  | n                  | Mean | SD   | Min  | Max  | n                        | Mean | SD   | Min  | Max  |
| Soil moisture (%)  | 77               | 18.3 | 10.9 | 1.3  | 50.5 | 22                 | 14.0 | 7.8  | 5.5  | 38.0 | 12                       | 16.5 | 13.0 | 1.3  | 50.5 |
| N <sub>total</sub> (%)   | 77               | 0.24 | 0.15 | 0.07 | 1.20 | 22                 | 0.22 | 0.10 | 0.10 | 0.45 | 12                       | 0.22 | 0.06 | 0.13 | 0.29 |
| C <sub>total</sub> (%)   | 77               | 3.3  | 3.0  | 1.1  | 25.7 | 22                 | 3.2  | 1.6  | 1.2  | 8.0  | 12                       | 2.6  | 0.7  | 1.4  | 3.3  |
| C/N ratio  | 76               | 13.3 | 2.2  | 9.8  | 19.9 | 22                 | 14.0 | 2.6  | 11.3 | 19.9 | 12                       | 11.8 | 0.6  | 10.7 | 12.7 |
| SOC (%)  | 73               | 5.8  | 2.5  | 2.1  | 16.1 | 22                 | 6.3  | 3.3  | 2.3  | 16.1 | 12                       | 5.2  | 1.4  | 2.8  | 6.7  |
| P (mg kg <sup>-1</sup> )   | 76               | 280  | 122  | 93   | 693  | 22                 | 234  | 70   | 111  | 353  | 12                       | 321  | 119  | 204  | 613  |
| Ca (cmol kg <sup>-1</sup> )  | 21               | 11.4 | 8.0  | 2.6  | 37.1 | 5                  | 6.9  | 4.1  | 2.6  | 11.4 | 4                        | 11.7 | 5.9  | 7.7  | 20.4 |
| Mg (cmol kg <sup>-1</sup> )  | 21               | 2.3  | 1.4  | 0.5  | 6.1  | 5                  | 1.5  | 0.8  | 0.5  | 2.7  | 4                        | 2.6  | 1.1  | 1.8  | 4.1  |
| K (cmol kg <sup>-1</sup> )   | 21               | 0.45 | 0.26 | 0.09 | 1.29 | 5                  | 0.34 | 0.21 | 0.09 | 0.60 | 4                        | 0.46 | 0.27 | 0.29 | 0.85 |
| Na (cmol kg <sup>-1</sup> )  | 21               | 0.08 | 0.05 | 0.0  | 0.3  | 5                  | 0.1  | 0.06 | 0.05 | 0.20 | 4                        | 0.06 | 0.02 | 0.04 | 0.08 |
| Zn (mg kg <sup>-1</sup> )  | 21               | 2.4  | 3.0  | 0.1  | 9.7  | 5                  | 0.5  | 0.4  | 0.2  | 1.2  | 4                        | 4.1  | 4.2  | 0.6  | 9.7  |
| Cu (mg kg <sup>-1</sup> )  | 21               | 15.5 | 42.7 | 0.8  | 161  | 5                  | 2.0  | 1.5  | 1.0  | 4.5  | 4                        | 71.8 | 83.4 | 0.8  | 161  |
| Fe (mg kg <sup>-1</sup> )  | 21               | 117  | 62   | 10   | 232  | 5                  | 158  | 73   | 69   | 232  | 4                        | 91   | 38   | 63   | 143  |
| Mn (mg kg <sup>-1</sup> )  | 21               | 23.3 | 10.4 | 7.1  | 43.6 | 5                  | 29.8 | 9.8  | 17.5 | 43.6 | 4                        | 15.3 | 3.7  | 9.9  | 18.1 |
| pH   | 77               | 5.0  | 0.7  | 3.8  | 6.6  | 22                 | 4.9  | 0.8  | 3.8  | 6.4  | 12                       | 5.2  | 0.3  | 4.9  | 5.7  |
| As (mg kg <sup>-1</sup> )  | 76               | 3.2  | 1.4  | 1.0  | 12.2 | 22                 | 3.0  | 1.0  | 1.0  | 4.4  | 12                       | 3.4  | 1.2  | 2.1  | 5.5  |
| Cd (mg kg <sup>-1</sup> )  | 76               | 0.39 | 0.13 | 0.18 | 0.74 | 22                 | 0.37 | 0.12 | 0.18 | 0.60 | 12                       | 0.43 | 0.13 | 0.27 | 0.66 |
| Cr (mg kg <sup>-1</sup> )  | 77               | 9.3  | 6.2  | 2.2  | 28.3 | 22                 | 9.5  | 7.3  | 2.7  | 26.4 | 12                       | 8.2  | 1.7  | 6.0  | 11.3 |
| Pb (mg kg <sup>-1</sup> )  | 77               | 5.6  | 2.1  | 2.8  | 10.7 | 22                 | 5.6  | 2.1  | 2.8  | 8.5  | 12                       | 5.3  | 1.2  | 3.9  | 7.3  |
| CEC (cmol kg <sup>-1</sup> )   | 21               | 14.2 | 9.3  | 3.4  | 44.6 | 5                  | 8.8  | 4.6  | 3.4  | 13.6 | 4                        | 14.8 | 7.0  | 9.9  | 25.0 |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> )                  | 21               | 0.04 | 0.02 | 0.02 | 0.1  | 5                  | 0.04 | 0.03 | 0.02 | 0.07 | 4                        | 0.04 | 0.03 | 0.02 | 0.09 |
| K <sup>+</sup> /(Ca <sup>2+</sup> +Mg <sup>2+</sup> +Na <sup>+</sup> ) | 21               | 0.04 | 0.02 | 0.02 | 0.08 | 5                  | 0.04 | 0.02 | 0.02 | 0.07 | 4                        | 0.04 | 0.03 | 0.02 | 0.08 |

hat gelöscht: Crops

859

hat gelöscht: .....Seitenbruch.....