

1 Title page

2 **Back to the future? Conservative grassland management can**
3 **preserve soil health in the changing landscapes of Uruguay**

4 Ina Säumel¹, Leonardo R. Ramírez¹, Sarah Tietjen², Marcos Barra³ and Erick Zagal⁴

5 ¹Integrative Research Institute THESys Transformation of Human-Environment-Systems
6 Humboldt-Universität zu Berlin, Unter den Linden 6, 10099 Berlin, Germany.

7 ²Leibniz Institute of Vegetable and Ornamental Crops (IGZ) e.V., Theodor-Echtermeyer-Weg
8 1, 14979 Großbeeren, Germany

9 ³Städtisches Klinikum Dessau, Auenweg 38, 06847 Dessau-Roßlau, Germany

10 ⁴Departamento de Suelos y Recursos Naturales, Universidad de Concepción, Campus Chillán,
11 Vicente Méndez 595, Chillán, Chile

12 *Correspondence to:* Ina Säumel (ina.saeumel@hu-berlin.de)

13 **Abstract.** The ‘soils of the anthropocene’ are predominately agricultural. To understand them,
14 we analysed agri- and silvicultural intensification of Uruguayan grasslands in a country wide
15 survey on fertility proxies, pH and trace metals in topsoils originating from different land uses
16 across the whole country. Thus, our results reflect interactions of both the natural diversity of
17 the Uruguayan soil formation and impacts of land use change. We observed a loss of nutrients,
18 trace metals and organic matter from grassland, crops and timber plantations. As an example,
19 the cation exchange capacity was 160 percent higher in native forests compared to grasslands
20 and lowest in timber plantations, reaching only half of the CEC in grasslands. Acidification of
21 topsoils continues as three fourth of all samples are 'extremely acidic' and 'very strongly acidic'.
22 Topsoils of riverine forests accumulate more trace metals compared to the other uses. We

23 assume an accumulation in the topsoils of riverine forests, where high levels of nutrients, trace
24 metals and organic carbon are found. The translocation of nutrients and organic matter across
25 the landscape to the erosion base depends on local land use trajectories. Increasing soil
26 acidification is driving a positive feedback loop, and land use intensification is leading to
27 degradation of local black soils within a few decades. Our data raises questions about the
28 resilience and carrying capacity of Uruguayan soils with regard to currently implemented
29 highly productive management forms, including the use of timber plantation for carbon
30 sequestration, and supports more conservative forms of extensive management on the grassland
31 biome.

32 **1. Introduction**

33 Human activities alter the bio- and pedosphere, leaving a footprint of such a magnitude that it
34 can be verified stratigraphically (Waters et al., 2016). This unprecedented transformational
35 force is intimately related to the expansion of societies and its productive frontiers, causing a
36 loss of biodiversity, habitat and soil degradation and, consequently, to ecosystem modification
37 (Foley et al., 2005, Borrelli et al., 2017). In this context, soil sciences have transitioned from
38 studies on natural soil formation to the science of ‘anthropedogenesis’ (Richter, 2020),
39 focussing on the ‘soils of the anthropocene’ that are predominately agricultural (51 Million
40 km²) or urban (1.5 Million km²; FAO, 2019).

41 The temperate grasslands of South America have historically been characterised by rolling
42 plains and low hills that have been extensively exploited for cattle production and its
43 derivatives since the arrival of European colonization. The Río de la Plata grasslands are one
44 of our planet’s four major black soil regions (Durán et al., 2011; Liu et al., 2012), and some of
45 the most fertile soils in the world. Playing an important role in the global food production, these
46 are characterized by a thick, humus and base cation rich and a high cation exchange capacity

47 throughout their profile. Maintaining their properties are therefore crucial to developing
48 sustainable and productive agriculture (Durán et al., 2011; Liu et al., 2012).

49 Today, the 'Uruguayan savanna' is part of the top-three most critically endangered biomes
50 (Veldman et al., 2015). In recent decades, however, the grassland has decreased due to the
51 expansion of cash crops and *Eucalyptus* plantations, both of which are promoted by national
52 legislation including land grabbing and trans-nationalization (Piñeiro, 2012). This land use
53 intensification, with its increased input of energy, nutrients and pesticides, leads to an overall
54 loss of soil fertility and increasing toxicity related to acidification, salinization and
55 contaminants (Liu et al., 2012; Borrelli et al., 2017). Ecological, economic and cultural
56 functions of soils are severely degraded, and the degradation of black soils in South America
57 is of particular concern because they have only been heavily exploited for a comparatively
58 short period of time (Durán et al., 2011).

59 Since the first decades of the twentieth-century, compared to other disciplines, soil sciences
60 have received an extraordinary amount of attention in Uruguayan academia, governance, the
61 productive sector, and also in the general public, resulting in a national soil inventory program
62 in 1965. The subsequent classification of Uruguayan soils and their productivity (CONEAT
63 Index) remains an important source for today's land taxation and for management plans by the
64 legal conservation regulations, and provides a detailed classification that takes into account soil
65 type, texture, natural vegetation, altitude and geology (Lanfranco and Sapriza, 2011).

66 As soil degradation is extremely relevant for countries like Uruguay, which are
67 socioeconomically dependent on their soils (Zubriggen et al. 2020), it is a topic of discussion
68 for local farmers, academia, and the public. An actualization of the state of the art of soils and
69 related processes is needed (García-Préchac et al. 2004; De Faccio et al. 2021), particularly as
70 there has been little study of the impacts of the Uruguayan grassland intensification on soils
71 properties (Beretta-Blanco et al., 2019). At the same time, while a new paradigm for grassland
72 intensification with a wide set of means including fertilization has been proposed to increase

73 economic and environmental sustainability (Jaurena et al., 2021), it is urgent to get more
74 insights into the dynamics of nutrients in soils of Uruguay and their availability for crops
75 (Beretta-Blanco et al., 2019).

76 Soil classifications are mainly based on subsoils. However, we focus on topsoil as the most
77 relevant and very responsive interface for ecological processes and farmer's management.
78 Understanding the state of the art of topsoils and its processes is crucial for developing
79 recommendations for sustainable land management practices. Due to the diversity of
80 perspectives on soil quality and health and related ecosystem services, operational procedures
81 for evaluation of soil functioning are still lacking (Bonfante et al. 2021). We contribute to a
82 better understanding of globally occurring degradation processes among often conflicting goals
83 such as desired soil productivity, yield limits, especially in erosion sensitive soils, and
84 necessary soil conservation.

85 We therefore explored soil parameters describing current chemical conditions of topsoils that
86 are parts of different soil groups and orders, and Uruguayan soil categories. Specifically, in
87 order to explore the gains and losses of macro and micro-nutrients and soil organic carbon
88 across landscapes, and to determine the impact of land use change on acidification and trace
89 metal presence and related trade-offs with soil degradation and conservation. In detail we
90 address the following question: how do fertility proxies such as soil organic carbon and content
91 of nutrients, acidification (pH) and trace metals accumulation in topsoils vary across different
92 land uses (i.e. comparing grassland, timber plantation, native forest, and agricultural land)?
93 Thus, we expand the knowledge across land uses from more natural to strongly modified uses
94 and discuss the results in light of different degradation processes such as erosion, depletion of
95 nutrients or carbon, acidification and accumulation of pollutants and in the light on current
96 debates on intensification.

97 **2. Material and Methods**

98 **2.1 Study area and design**

99 Uruguay covers about 176,000 km², and has a population of 3.5 Million, mainly in urban areas.
100 The Western half of Uruguay is dominated by Mollisols, developed on a wide range of
101 sediments from the Devonian to the Cenozoic era (partially associated with Vertisols), while
102 the Eastern plains and wetlands have, in addition to Mollisols, other soil types on the slopes,
103 rocks and on flood plans (i.e. Alfisols, Ultisols, Entisols and Inceptisols; Durán et al. 2011).
104 During a country-wide survey from December 2015 to March 2016, we collected 280 topsoil
105 samples of 0-10 cm depth from 101 plots (50x50m in grassland and crops; 100x100m in native
106 forests and timber plantations) distributed at 28 monitoring sites throughout Uruguay, South
107 America (Fig. 1a-b; Table 1), using a stratified random design. In the first step, we randomly
108 selected monitoring sites across the country. In the second step, we contacted landowners to
109 explore their willingness to establish a long-term monitoring site. If the owner agreed, plot
110 selection was stratified by different rural land use types: grassland, timber plantations of *Pinus*
111 and *Eucalyptus* species, native forest, and crops. Native forests cover mainly riverine and park
112 forests. The later are a savanna like transition zones between riverine forests and the open
113 grasslands. We subdivided grassland plots according to the intensity of use: (i) undisturbed
114 grassland (without grazing), (ii) partially grazed grasslands (with sporadic grazing and low
115 animal charge), and (iii) highly grazed grassland (with high animal charge). Land use change
116 from 1986 to 2017 follows basically three different trajectories: i) the expansion of timber
117 plantations over grassland leading to a disaggregation of grassland by timber plantations; ii)
118 cropland expansion where crop cover maintains the open landscape character of former
119 grasslands, grassland conservation where large and regularly interconnected riverine forests in
120 a landscape dominated by grasslands (Ramírez and Säumel 2021) and grassland intensification

121 changing from natural grassland to so called ‘improved’ or artificial grasslands (Modernel et
122 al. 2016; Jaurena et al. 2021). Fertilization and application of other agrochemicals is standard
123 procedure in timber plantations, artificial grasslands and industrial crops. We sampled top soil
124 three times at each land use at the edges of the plot, and stored samples below 7°C until
125 laboratory processing. The plots are placed in homogenous areas to avoid edge effects.

126 **2.2 Analysis of Soil Samples**

127 For gravimetric determination of soil moisture, the topsoils samples were dried at 105 °C until
128 constant weight. Next, lumps in the samples were broken down and the remaining plant
129 material was removed before sieving (2 mm) and ground.

130 We analysed 280 samples regarding macronutrients, pH and trace metals and 80 samples for
131 soluble cations and micronutrients (Table 1; Sadzawka et al. 2006; Zagal & Sadzawka 2007).
132 Among fertility-related variables, we measured the total amount of the macronutrients
133 phosphorus (P), organic carbon (SOC) and nitrogen (N), so obtaining the C/N ratio. To
134 determine total carbon and nitrogen, the samples were sieved again (0.5 mm) and analyzed
135 using dry combustion with a LECO TruSpec CN (USA) at a combustion temperature of 950
136 °C with ultra-pure oxygen. In addition, the presence of carbonates was tested for by adding
137 concentrated hydrochloric acid, which was negative for all soil samples analysed.

138 We determined concentrations of the soluble cations calcium (Ca^{2+}), magnesium (Mg^{2+}),
139 potassium (K^+), sodium (Na^+), and the micronutrients copper (Cu^{2+}), zinc (Zn^{2+}), manganese
140 (Mn^{2+}) and iron ($\text{Fe}^{2+/3+}$) by atomic spectroscopy (Unicam AAS Solaar 969, Thermo Electron
141 Corporation, USA), extracting with ammonium acetate (1 mol l⁻¹, pH 7), and with DTPA-
142 CaCl_2 -TEA at (pH 7.3). We calculated the cation-exchange capacity (CEC). Acidity was
143 measured by adding calcium chloride (0.01M) to the samples in a 2.5:1 proportion, and after
144 shaking and two hours rest, read with a pH meter (HI2550 meter, Hanna Instruments, USA)).

145 For these variables, each tenth sample was duplicated. We categorized acidity using the USDA
146 Natural Resources Conservation Service classification (Kellogg, 1993).
147 To determine the concentrations of arsenic (As), cadmium (Cd), chromium (Cr) and lead (Pb),
148 samples were further sieved as before (0.5 mm), weighed out into a digesting container, and
149 extracted with a mixture of nitric acid (70%) and hydrochloric acid (30%) in a 1:3 proportion.
150 The digestion took place in a Titan MPS (Perkin Elmer, USA) microwave at a programme
151 suitable for this digestion (Method 3051A). During the proceedings, only deionized water Type
152 I ASTM1193 (EC max 0.06 to 0.1 $\mu\text{S}/\text{cm}$) was used. Reagents were used to eliminate traces of
153 other materials and to avoid contamination of the samples. The trace metals were determined
154 by inductively coupled plasma optical emission spectroscopy using the Optima 8000 (Perkin
155 Elmer) with metal-associated wavelengths of 193.696 nm for As, 228.802 nm for Cd, 267.716
156 nm for Cr and 220.353 nm for Pb.
157 Total P concentration of phosphorus was determined calorimetrically after microwave-assisted
158 digestion with Unicam spectrometer at a wavelength of 660 nm. For all these variables, a
159 repetition for each round of the microwave digestion was made.

160 **2.3 Soil classification**

161 We intersected the coordinates of the centre of the plots with maps containing geospatial
162 information on the classification of the Uruguayan soils using ArcGIS 10.3 (ESRI, 2018). For
163 Soil Groups classification, we used the of the World Reference Base for Soil Resources (WRB;
164 IUSS Working Group, 2015); for Soil Orders, the USDA soil taxonomy (Soil Survey Staff,
165 1999); and for the local Uruguayan classification, Soil CONEAT (Comisión Nacional de
166 Estudio Agronómico de la Tierra) Groups categories, which include productive capacity of
167 cattle and sheep (MGAP, 2020). The CONEAT groups are defined by their productive capacity
168 in terms of beef, sheep and wool expressed by an index relative to the average productive

169 capacity of the country, to which the index 100 corresponds. The classification is based on
170 photo-interpretation at a scale of 1:40,000, field verifications and physico-chemical analysis of
171 the soils. The productivity indices correspond to soil groups. The CONEAT groups have been
172 defined by the dominant and associated soils according to the Soil Classification of Uruguay.
173 The groups are related to the units of the Soil Reconnaissance Chart of Uruguay at a scale of
174 1:1,000,000. For each group, some important soil properties and associated landscape
175 characteristics are indicated. The nomenclature of the CONEAT groups correlates with the Soil
176 Use and Management Zones of Uruguay. The Soil Groups are superimposed on the rural parcel
177 and are represented in the CONEAT cartography at a scale of 1:20,000 (for more details see
178 MGAP, 2020).

179 **2.4 Data Analysis**

180 In a first step, we explored and prepared our database for further analysis. Exploring the
181 distribution of the soil parameters in R (R Core Team 2021), ruled out the normality of the data
182 using the Shapiro-Wilk Test and the homogeneity of variances with the Fligner-Killeen Test.
183 We tested for outliers using the 1.5-3 IQR threshold and the function *outlierTest* from the R
184 package *car* (Fox and Weisberg 2019), reviewing the flagged observations case by case in the
185 experimental context. The outliers were removed (Supplementary Material: Fig.S2). The
186 variables on soils characteristics showed generally positive skewed distributions, in some cases
187 multimodal, and tests showed evidence contrary to assumptions of spatial autocorrelation,
188 homoscedasticity and normality in most cases (Supplementary Material: Table S1-S2; Fig. S1-
189 S2).

190 Spearman's rank correlations (ρ) were calculated to explore linear associations between soil
191 parameters across all single samples and within different land uses. We used the *adonis*
192 function of the R package *vegan v2.5-7* (Oksanen et al., 2020) with a Euclidean dissimilarity

193 matrix from our normalized soil data to perform PERMANOVA tests with 9999 permutations
194 to analyse the multivariate homogeneity of group dispersions based on differences on soil
195 parameters between land uses.

196 To compare the general effects of land use type on topsoil parameters, non-parametric Kruskal-
197 Wallis tests were carried out in R. When significant ($p \leq 0.05$), we used Pairwise Wilcoxon
198 Rank Sum tests with Benjamini & Hochberg correction (1995) to evaluate pairwise differences
199 among land uses.

200 We used non-metric multidimensional scaling (NMDS) as a robust unconstrained ordination
201 method to visualize patterns of top soil characteristics among all samples and within
202 subsamples (intersected with different soil Orders) across different land uses using the Bray-
203 Curtis dissimilarity matrix. The matrix was constructed with the *vegan* package to depict
204 patterns of all soil parameters in two dimensions (Fig.3a, c-f) and for the dataset without soluble
205 cations and micronutrients variables comparing subcategories within single land use types (i.e.
206 within grasslands in ‘undisturbed’, ‘partially grazed’ and ‘highly grazed’ plots; timber
207 plantations in ‘*Eucalyptus*’ or ‘*Pinus*’ plots; Fig. 3b).

208 **3. Results**

209 Nearly 75% of the sampled 101 plots intersected with the soil Group category 'Luvic Phaeozem'
210 (WRB; IUSS Working Group, 2015). A quarter of our plots were distributed among four other
211 Groups (i.e. Haplic Luvisols, Eutric Vertisols, Planosols, Lithic Leptisols; Fig. 1c). Half of our
212 plots intersected with the Orders 'Argiudolls' or 'Argiudolls & Hapluderts' (Soil Survey Staff,
213 1999), and about a quarter with 'Hapludalfs & Hapludults' or 'Argiudolls, Hapludolls &
214 Hapludalfs' (Fig. 1d). Our plots intersected with 32 different CONEAT categories (MGAP,
215 2021). The most frequent categories were '10.2', '9.3' and '2.12' (25% of our plots) and another
216 quarter with '9.1', '8.8', '9.6' and 'G03.11' (Fig. 1e).

217 **3.1 General characteristics of Uruguayan topsoils**

218 The measured topsoil parameters vary widely across Uruguay, between the different land uses
219 and classification to different soil orders (Table 1, Table A1-A4). The soil organic carbon in
220 topsoils ranged between 0.8 to 16 percent, and was highest in native forests and lowest in timber
221 plantations. The mean of the C/N ratio was about 13, and lowest in crop soils. Phosphorous
222 ranged between 43 to 1009 mg kg⁻¹. We also observed a high variability for the micro- and
223 macronutrients and trace metals. The average cation-exchange capacity (CEC) was 12.41 cmol
224 kg⁻¹, highest in topsoils of native forest, followed by crops and grasslands and timber
225 plantations (Table 1, Table A1-A2).

226 For the whole data set, high correlation was found between P with SOC and Zn ($\rho=0.82$ and
227 0.76 , respectively), and between Mg with Ca and Na ($\rho=0.82$ and 0.76 , respectively; Fig. 2).
228 Similar results were observed within particular land uses, although in native forests, a negative
229 moderate correlation between Ca and Fe was observed (Fig. S3). There was high correlation
230 between pH and Ca ($\rho=0.89$; Fig. 2). In native forests, we also found similar correlation with
231 other soluble cations like Mg and K (Fig. S3). In park forests topsoils, there was a negative
232 correlation between pH and As and Pb ($\rho=-0.81$ and $\rho=-0.84$, respectively). In highly grazed
233 pastures and crops pH was highly correlated with Cr, and in crops also with As ($\rho=0.93$; Fig.
234 S3). Among trace metals, Spearman correlation was moderate between As with Cr and Pb and
235 high between Cr and Pb (Fig. 2). These correlation trends increased in grasslands, timber
236 plantations and crops. We also found a high correlation between cadmium and Cr in crops
237 ($\rho=0.81$). Phosphorus was highly correlated with Cr and As in pine plantations, while in
238 soybean crops the correlation was between phosphorus with Cr and Pb (Fig. S3).

239 **3.2 Topsoil characteristics clustered by land use**

240 We found differences in multivariate distribution of samples according to the different land
241 uses (i.e., grassland, timber plantations or native forests; Fig. 3a-f; Table S3-S8). All pairwise

242 comparison of land uses showed significant differences with all variables ($p=0.0001$; Fig. 3a;
243 Table S3). We analysed subcategories within a land use type using the dataset without soluble
244 cations and micronutrients variables, only finding significant differences between *Eucalyptus*
245 and *Pinus* stands ($p=0.0001$; Fig. 3b; Table S4) but not among different grassland subtypes.
246 We also found differences analysing subsamples of different soil order classification to the
247 different land uses (Fig. 3c-f; Table S5-S8). The samples intersected with 'Argiudolls' included
248 all plots on crops, and we found significant differences in all pairwise comparisons of land uses
249 except between grasslands and timber plantation ($p=0.0004$; Fig. 3c; Table S5). We further
250 found differences between timber plantations and native forests at soils of the 'Argiudolls &
251 Hapluderts' Orders ($p=0.0009$; Fig. 3d; Table S6) and between timber plantations and
252 grasslands or native forests in soils of the 'Argiudolls, Hapudolls & Hapludalf' Orders
253 ($p=0.0284$; Fig. 3e; Table S7). Results for samples in 'Hapludalfs & Hapludults' soils were
254 similar to those obtained at country scale ($p=0.0001$; Fig. 3f; Table S8).

255 **3.3 Differences in fertility proxies**

256 We found significantly higher fertility proxies for SOC, P, Ca, Mg and Zn in topsoils from
257 native forests compared to grasslands and timber plantations (Fig. 4b-e, h). Phosphorous was
258 significantly higher in topsoils of grassland compared to those of timber plantations (Fig. 4c).
259 Potassium was significantly higher in topsoils of native forests compared to timber plantations
260 (Fig. 4f). At subcategories level, we found significantly higher amounts of K in partially used
261 grasslands in comparison to samples from highly grazed pastures, and higher SOC ($p=0.002$),
262 P ($p=0.059$), Na ($p=0.043$), K ($p=0.012$) and Zn ($p=0.048$) in *Eucalyptus* compared to *Pinus*
263 plantations (Fig. S4). Among native forests, samples from riverine forests contain more Mg
264 ($p=0.023$) and Na ($p=0.023$) in comparison to park forests.
265 Considering all samples within the Order 'Argiudolls', C/N ratio in agricultural topsoils are
266 lower compared to all other land uses. Phosphorus was higher in topsoils of native forests and

267 crops compared to grasslands and timber plantations. Soil organic carbon was highest in native
268 forests (Fig. 5a-o; Table A3-A4).

269 **3.4 Soil Acidification**

270 Topsoil samples showed a markedly acid profile (median pH=4.66), with nearly 75% classified
271 as 'extremely acidic' and 'very strongly acidic' (Fig. 6a). We found significant differences across
272 land uses, with less acidic native forests, and lower pH in timber plantations (Fig. 6b).
273 Comparing between land uses, we found more samples with neutral acidity in grasslands and
274 more with higher acidity in timber plantations (Fig. 6b-c). In addition, we found lower pH in
275 samples from *Pinus* compared to *Eucalyptus* stands ($p=0.018$). Results of analysis inside soil
276 Orders showed similar variations observed at country scale, with timber plantations being more
277 acid and native forest closer to neutral pH (Fig. 6d-f).

278 **3.5 Trace metal accumulation across land uses**

279 For As, Cd, Cr and Pb, we found significantly higher concentrations in topsoils originating
280 from native forests compared to the grassland and timber plantation samples (Fig. 4l-o and 5l-
281 o; Table A1-A4). At the same time, samples from *Eucalyptus* plantations had higher
282 concentrations of both Cr ($p<0.005$) and Pb ($p<0.005$) than *Pinus* topsoils, while the same was
283 observed for Cr ($p<0.001$) and Pb ($p<0.05$) in riverine forests compared to park forests (Fig.
284 S5).

285 **4. Discussion**

286 The vicious circle between the wish to stop soil degradation and concurrent increases in land
287 productivity to satisfy the increasing demand for food, fibres and energy has not been broken
288 since green revolution. Socio-economic and conventional management practices that drive soil
289 degradation have generated several traps, such as the 'inputs trap' where a reduced yield per

290 area is followed by higher fertilizer application or the 'credit or poverty trap' where economic
291 pressure forces the farmer to intensification (Gomiero, 2016). Caught in this loop, the soils of
292 the temperate grasslands of Uruguay have suffered strong degradation from erosion,
293 acidification, contamination, salinification and compaction (Zurbriggen et al., 2020). This is
294 clearly reflected in the results of our topsoil survey, which also adds interesting insights from
295 timber plantations, grasslands and native forests to an existing database consisting mainly of
296 crops and pastures samples from 2002-2014, which demonstrated the loss of organic matter by
297 25% and an increasing loss of nutrients (Beretta-Blanco et al. 2019). We contribute deeper
298 insights on fertility, acidification and trace metals accumulation in topsoils from a wide range
299 of different land uses, which is, to our knowledge, unique for the region since the CONEAT
300 classification (CONEAT Index, 1976).

301

302 **4.1 Translocation of elements in topsoils within landscape**

303 Our data demonstrate the high amounts of organic carbon, nutrients and trace metals in topsoil
304 samples from riverine forests, suggesting transport of soil particles from the surrounding land
305 uses (e.g., grasslands, crop or timber plantations) to the borders of rivers, streams and creeks.
306 Therefore, we assume that organic carbon, nutrients and trace metals are displaced within the
307 landscape and accumulate in the floodplains. Regional soil erosion models estimate the loss of
308 2-5 tons ha⁻¹ year⁻¹ for a third of the country depending on precipitation, topography, soil
309 erodibility and land management (Carrasco-Letelier and Beretta-Blanco 2017). One possible
310 direct impact is the increasing eutrophication reported for larger local rivers, although the
311 models used by these authors did not link Chlorophyll-a concentrations with agricultural land
312 use (Beretta-Blanco and Carrasco-Letelier 2021 and replies). However, other vertical processes
313 and differences across land uses can be hidden deeper in the soil profile, and have not been
314 analysed in this study.

315 Organic carbon content and the exchangeable cations are strongly reduced in topsoils of
316 grasslands, timber plantations and crops compared to native forests (Fig. 4b, d-h and 5b, d-h).
317 Of all the fertility proxies assessed, phosphorus in topsoil was most significantly affected by
318 different land uses, being highest in native forests (Fig. 4c and 5c). The cation exchange
319 capacity (CEC) was highest in native forests (more than 160 percent of the CEC in grasslands)
320 and lowest in timber plantations, reaching only half of the CEC in grasslands (Table A1-A2).
321 Lower average nutrient concentrations and corresponding CEC reported for two timber
322 plantations in the East and one in the Norwest of Uruguay (Hernández et al., 2009 and 2016;
323 Céspedes-Payret et al., 2012) may result from a combined effect of topsoil degradation in
324 timber plantations due to management practices and to soil texture (Sandoval-Lopez et al.,
325 2020). The trees' uptake and the general export of nutrients from fast growing timber
326 plantations through harvesting is higher than the natural input into those systems (Merino et
327 al., 2005). This effect is particularly relevant because timber plantations in Uruguay are on
328 'forestry priority soils', which are generally soils with low fertility, superficial to moderate
329 depth and good drainage (OAS, 1994), so afforestation might reduce even more their soil
330 fertility.

331 Comparing neighbouring sites of grassland and afforested grasslands, our data set shows no
332 significant depletion of nutrients by timber plantations ($p=0.208$) but a slightly higher average
333 of CEC at grasslands (Table A1). In general, plants are expected to compensate the extraction
334 of cations from upper soil by the 'uplift of nutrient' from deeper horizons (Jobbagy and Jackson,
335 2004). This has been questioned for *Eucalyptus* plantations on sandy soils, where, for example,
336 an increase of potassium in the topsoil was not observed compared to the neighbouring
337 grassland (Céspedes-Payret et al., 2012). Phosphorous and cations decline with sand content,
338 so physicochemical factors such as the percentage of sand and organic matter, influence soil
339 fertility (Sandoval-Lopez et al., 2020). A study of *Eucalyptus* plantations in South-East Brazil
340 did not show a significant depletion of nutrients and carbon; in contrast, carbon, potassium and

341 calcium increased in the topsoil after twelve years over one or two harvest and fertilization
342 cycles (McMahon et al., 2019). To sum up, the patterns observed in the various sites in our
343 survey and the other studies indicate very complex interactions of numerous factors. Removal
344 of nutrients by high-yield timber plantations may exceed the capacity of nutrient exchange or
345 turnover of forest litter and wastes. Both stand management and environmental conditions (e.g.,
346 precipitation) influence nutrient and carbon stocks (McMahon et al., 2019; Sandoval-Lopez et
347 al., 2020).

348 The C/N ratio in topsoils ranged within values reported for grassland and timber plantations of
349 the region (see Berthrong et al., 2012). In contrast to that study, however, we observed no
350 differences between grassland and timber plantations (Fig. 4a), but between topsoils of timber
351 plantations, native forests and crops (Fig. 5a). Organic matter and nitrogen decrease in topsoils
352 after grassland afforestation (Berthrong et al., 2009) and C/N ratio increases with plantation
353 age and decreases with precipitation (Berthrong et al., 2012). The topsoils of timber plantations
354 have, on average, lower N concentrations compared to other land uses (Table A2) and data in
355 literature (Jobbagy and Jackson 2003).

356 Although in cropland, nutrients are regularly compensated for by increased application of
357 fertilizers (Beretta-Blanco et al., 2019), timber plantations are usually fertilized only in the first
358 year of planting (e.g., Binkley et al., 2017; Sandoval-Lopez et al., 2020). The CEC and the
359 average ratio between $K^+(Ca^{2+} + Mg^{2+})^{-1/2}$ in crops and grasslands are in the ranges reported by
360 Beretta-Blanco et al. (2019) with lower availability of potassium for crops. A shortage of
361 potassium and calcium is also most likely for future timber plantations after harvest, especially
362 if the logging process do not include bark stripping on site (Hernández et al., 2009) and due to
363 short time spans between harvest and new planting in the same area (Sandoval-Lopez et al.,
364 2020), but see contrasting results of McMahon et al. (2019). The extraction of nutrients and
365 biomass due to grazing (Fernandez et al., 2017), timber (Hernández et al., 2009 and 2016) and

366 crop production (Beretta-Blanco et al., 2019) lowers organic matter content and exchangeable
367 cations, affects the physical, chemical and biological soil properties, and drives degradation.

368 **4.2 Acidification in Uruguayan topsoils across land uses**

369 A further dimension of the soil degradation directly linked to the cation extraction is the
370 acidification of soils. This has been demonstrated for crops and pastures (Beretta-Blanco et al.,
371 2019), and a study site with *Eucalyptus* (Céspedes-Payret et al., 2012) and is now broadly
372 supported by our topsoil samples originating from a wide range of different land uses across
373 Uruguay. The pH of our topsoil samples are mainly in the category of very strongly to
374 extremely acidic and lowest in timber plantations (Fig. 6), which are below the means reported
375 so far (Jobbagy and Jackson, 2003; Céspedes-Payret et al., 2012). Moreover, as our data on
376 topsoil pH fall short compared to the values estimated by the Food and Agriculture
377 Organization of the United Nations and the Intergovernmental Technical Panel on Soils (Caon
378 and Vargas, 2017), acidification and the deterioration of topsoil quality continues.
379 Acidification results from intensified land uses with nitrogen fertilization, with biological N
380 fixation by the legumes both used in the improved pastures (Modernel et al., 2016), or with
381 cation extraction by crop or timber harvest (Jobbagy and Jackson, 2003). So far, although forest
382 soils tend to be more acidic than agricultural soils due to acid-neutralizing treatments in the
383 latter (Baize and van Oort, 2014), we found no differences between topsoils of crops and native
384 forests (Fig. 6d) as the high organic carbon content in native forests buffers the process to a
385 certain extent.

386 **4.3 Riverine Forest soils as sink for trace metals**

387 In general, the average concentration of Zn, Cu, Cd, Cr, As and Pb in our topsoils were within
388 the expected ranges of samples from crops, pastures and grassland in the region (Lavado et al.,
389 1998, 2004; Roca, 2015). Some samples, especially from orchards and crops, exceeded the
390 background values of copper, cadmium and arsenic (Table A2; Roca, 2015). Little is known

391 about the pedo-geochemical background of the Pampean soils (Roca, 2015): for example, high
392 arsenic concentration in Uruguayan ground waters has been hypothesized to be due to
393 quaternary ash deposits (Wu et al., 2021). In addition, the lateral heterogeneity of Pampean
394 soils over short distances makes separating geochemical and anthropic signatures difficult
395 (Roca, 2015). However, that the main risk of soil contamination in the region is from the
396 application of fertilizers and agrochemicals is uncontested (Kaushal et al., 2018).

397 To our knowledge, there has been no regional study of trace metals in the native riverine forests
398 or timber plantations. Our work thus expands the evidence base for these land uses. The topsoils
399 of riverine forests accumulate more trace metals compared to those of timber plantations and
400 crops (Fig. 4l-o). The higher amount of soil organic matter in riverine forests favours the
401 retention of cations, including trace metals. Although the origin of potentially harmful elements
402 in forest soils have been primarily attributed to atmospheric deposition (Baize and van Oort,
403 2014), atmospheric deposition only plays a major role in vicinity of urban or industrial
404 development, and our data from rural sites suggests a different entry path from the surrounding
405 land uses to the riverine forests. High acidification and low amounts of organic matter reduce
406 the retention of trace metals in the soil of timber plantations, and elements leach out of the soil
407 towards the water table (Baize and van Oort, 2014). The acidification strongly contributes to
408 the overall mobility of base cations into the ‘chemical cocktail of the Anthropocene’ (Kaushal
409 et al., 2018), including trace metals. We thus observe positive feedback in already
410 impoverished soils with high acidity favouring cations solubility, in addition to the uptake by
411 trees intensifies this effect. Timber plantations extract trace metals from soil and also
412 accumulate it in the bark or leaves, so they have been used for phytoremediation (Li et al.,
413 2020). This may explain the higher concentration of cadmium in grassland compared to timber
414 plantations (Fig. 4m). Differences between *Eucalyptus* and *Pinus* stands may be related to
415 different age classes, as the later may have extracted more lead and chromium from the soil
416 due to their older stand age with rotation periods of about 20 years (Li et al., 2020).

417 **4.4 Carbon storage in topsoils of *Eucalyptus* plantations?**

418 Our study provides evidence that the loss of soil organic carbon limits not only the productivity
419 of the crops, but also potential carbon sequestration in the region (Beretta-Blanco et al., 2019).
420 Grassland conversion to cropland decreases soil organic carbon storage compared to adjacent
421 native grasslands (Hernandez-Ramirez et al., 2021).

422 Afforestation of croplands has been also discussed as a carbon sequestration measure to
423 proactively address and effectively mitigate ongoing climate change within a person's lifetime
424 (Hernandez-Ramirez et al., 2021; Mayer et al. 2020). Although carbon stocks of four
425 *Eucalyptus* stands in the Brazilian Cerrados increased but did not change in four *Eucalyptus*
426 stands within the Atlantic Forest (McMahon et al., 2019), this may not be the case for short
427 rotation of *Eucalyptus* plantation on Campos grasslands. In our topsoil samples originating
428 from 28 different stands across Uruguay, organic carbon is lowest in topsoils of timber
429 plantations (Fig. 4b). Similar amounts have been reported for a timber plantation in the East of
430 Uruguay (Céspedes-Payret et al., 2012), and in North-Eastern Argentine (Sandoval-Lopez et
431 al., 2020). Our data therefore provide clear evidence that rather than contributing to carbon
432 sequestration in the topsoil, the carbon release from the transformation of native grasslands to
433 plantation with these fast-growing species has several adverse effects depending on
434 precipitation and soil type (reviewed by Mayer et al., 2020).

435 Several trade-offs between carbon sequestration through afforestation and local water yield and
436 soil fertility have been demonstrated, including nutrient and soil organic matter depletion,
437 acidification, and biodiversity loss and corresponding challenges for landscape conservation
438 (e.g., Jackson et al., 2005; Veldman et al., 2015; Friggens et al., 2020). Soil carbon changes
439 dynamically during the first decade of afforestation. Remaining grassland carbon declines,
440 while tree carbon gain starts (e.g. Paul et al., 2002). Although a net gain of carbon is expected
441 when the new forests approached equilibrium after decades (Hernández-Ramirez et al., 2021),
442 in contrast to long-lasting forests, *Eucalyptus* harvest in Uruguay takes place after less than a

443 decade (appr. 7-10 years). Soil organic carbon does not differ between *Eucalyptus* plantations
444 and neighbouring grasslands (Appendix A; Table A1 and A2). Another study near our sites 12
445 and 13 also did not find a significant difference between the soil organic carbon of the upper
446 soil (0-30cm) compared to afforested versus native grasslands (Hernández et al., 2016). A study
447 near our sites 7, 8, 10 reported higher top soil carbon of grassland compared with afforested
448 sites in their vicinity only if sand content is lower than sixty percent (Sandoval-Lopez et al.,
449 2020). McMahon et al. (2019) identified a greater carbon gain under *Eucalyptus* stands
450 compared to (potential) carbon losses in neighbouring degraded Cerrado grasslands. There is
451 evidence that the retention of residues after harvest increases the carbon stock in soil (Mayer
452 et al., 2020). Long term monitoring of carbon stocks is needed to verify the influences of
453 management and environmental changes. A regional study on *Eucalyptus* plantations across
454 different biomes in Brazil shows both decreases and no changes depending on precipitation in
455 the dry season, clay content and on the initial stock of carbon in the soil (Cook et al., 2016)
456 The simplistic solution of huge tree plantations to compensate anthropogenic CO₂ emissions
457 has been challenged in the last decade, and some crucial lessons learnt have been summarized
458 (Di Sacco et al., 2020). Since grasslands are more sustainable carbon sinks compared to
459 climate-vulnerable monocultures such as timber plantations (Dass et al. 2018), avoiding
460 afforestation of previously non-forested lands is important. This is the case for *Eucalyptus*
461 afforestation in the originally forestless grasslands of Uruguay: models on carbon sequestration
462 and dynamics in Mollisols and Oxisols under South American grasslands estimated a higher
463 carbon retention efficiency under grasslands compared to afforested sites, suggesting that
464 silvopastoral systems are a potential solution for soil carbon sequestration in tropical soils
465 (Berhongaray and Alvarez, 2019).
466 Connecting or expanding existing forest and using native species for plantings (Di Sacco et al.,
467 2020) is also recommended (see also Alvaro et al. 2020, Ortiz et al. 2020). Our topsoil data
468 indicates that carbon sequestration occurs mainly in the topsoils of native riverine forests that

469 cover less than five percent of the Uruguayan territory. Consequently, the expansion of native
470 forests and the use of native species in forestry project for long term establishment can reduce
471 adverse effects of timber plantations. While there is preliminary evidence that N-fixing tree
472 species can help increase local C stocks of afforestation, because of their potential for invasion,
473 exotic N-fixers should be avoided (Mayer et al. 2020).

474 Recent studies indicated that novel techniques such as perennial grain cropping can help to turn
475 around cropland degradation into beneficial cropland aggradation by using the advantage of
476 perennial vegetation of conserving and even enhancing short term and long-term soil carbon
477 storage and other ecosystems services (Kim et al. 2022).

478 **4.5. Back to more conservative grassland management?**

479 Our soil survey data shows strong soil degradation of Uruguayan black soils from erosion,
480 acidification and contamination, and suggests a translocation of nutrients and organic carbon
481 across the landscape from grassland, timber and crop plantations to the riverine forests. The
482 potential of grasslands as cropland reserve have been largely overestimated (Lambin et al.,
483 2013), and they have already degraded during the last decades by inappropriate land
484 management techniques (Jaurena et al., 2021, De Faccio et al., 2021), lack of mainstreaming
485 soil conservative techniques (García-Préchac, 2004; Fernandez et al., 2017), decoupling of crop
486 and livestock (De Faccio et al, 2021) plus climate change impacts with storm water events and
487 drought, all of which trigger soil erosion (Wingeyer et al. 2015). From the although very limited
488 point of view on topsoils, the concept of conserving ‘old growth grasslands’ with extensive use
489 (Veldman et al. 2015) appears a more promising strategy to put the ‘grasslands at the core’ in
490 the Campos region than the use intensification strategies envisioned by Jaurena et al. (2021).

491 **5. Conclusions**

492 The land use intensification in Uruguay associated with increasing inputs of energy, nutrients
493 and pesticides leads to an overall loss of soil fertility and increasing toxicity related to
494 acidification, salinization and trace metal contaminants. Our data demonstrate the high amounts
495 of organic carbon, nutrients and trace metals in topsoil samples from riverine forests,
496 suggesting transport of soil particles from the surrounding grasslands, crop or timber
497 plantations to the borders of rivers, streams and creeks. Of all the fertility proxies assessed,
498 phosphorus in topsoil was most significantly affected by different land uses, being highest in
499 native forests. Cation exchange capacity was also highest in native forests and lowest in timber
500 plantations, where only half that of grasslands was measured. Our study highlights that soil
501 acidification is ongoing and probably also mobilizing trace metals and their accumulation in
502 riverine forest topsoils.

503 **Data availability**

504 All data generated or analysed in this study are included in this article (and its supplementary
505 materials). Further data are available from the corresponding authors upon reasonable request.

506 **Author contributions**

507 IS, LRR, ST and MB conceptualized the study, performed the soil sampling and data analysis.
508 EZ advised the laboratory analyses. IS wrote the initial draft. LRR, ST, MB and EZ contributed
509 to generating and reviewing the subsequent versions of the manuscript. IS received the funding
510 for the study.

511 **Competing interests**

512 The authors declare that they have no conflict of interest

513 **Acknowledgement**

514 In alphabetical order, we thank Juan Barreneche, Lucia Gaucher, Sören Miehe, Nicolas Silvera
515 and Matias Zarucki for their assistance in field work. We thank Manuel Garcia and Meica
516 Valdivia for the pre-processing of samples. We thank the staff of the soil laboratory from the
517 Department of Soil and Natural Resources, University of Concepción, Chile. We thank Diego
518 de Panis for help with statistical analysis and visualization and valuable comments on a
519 previous version of this manuscript. Thanks to Vera Krause, Serafina Bischoff, Sophia Reitzug,
520 Rhea Rennert and Diego Nicolas Rojas for support with data analysis and plots and maps
521 visualization. We also thank all landowners for access permission to establish our monitoring
522 sites on their land, their hospitality and willingness to discuss land use goals concerning all
523 dimension of sustainability. The study was funded by the German Federal Ministry of
524 Education and Research (BMBF; 01LN1305A). Special thanks go to Amal Chatterjee for
525 improving our English.

526 **References**

- 527 Alfaro, M., Dube, F. and Zagal, E.: The Influence of Overmature, Degraded Nothofagus
528 Forests with Strong Anthropic Disturbance on the Quality of an Andisol and its Gradual
529 Recovery with Silvopasture in Southwestern South America, in: *Agroforestry for
530 Degraded Landscapes*, edited by: Dagar, J.C., Gupta, S.R., Teketay, D. Springer,
531 Singapore. https://doi.org/10.1007/978-981-15-6807-7_3, 2020.
- 532 Baize, D. and van Oort, F.: Potentially Harmful Elements in Forest Soils, in: *PHEs*,

- 533 Environment and Human Health, edited by: Bini, C. and Bech, J., Springer Netherlands,
534 Dordrecht, 151–198, https://doi.org/10.1007/978-94-017-8965-3_4, 2014.
- 535 Benjamini Y, Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
536 Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*
537 (Methodological), 1995; 57(1), 289-300. <https://www.jstor.org/stable/2346101>.
- 538 Beretta-Blanco A, Pérez O, Carrasco-Letelier L. Soil quality decrease over 13 years of
539 agricultural production. *Nutr Cycling Agroecosyst.* 2019; 114, 45-55.
- 540 Beretta-Blanco, A.; Carrasco-Letelier, L. Relevant factors in the eutrophication of the Uruguay
541 River and the Río Negro. *Sci. Total Environ.* 2021, 761,143299; DOI:
542 10.1016/j.scitotenv.2020.143299
- 543 Berhongaray, G; Alvarez R. Soil carbon sequestration of Mollisols and Oxisols under grassland
544 and tree plantations in South America - A review. *Geoderma Regional* 18 (2019)
545 e00226.
- 546 Berthrong, S. T., Jobbágy, E. G., and Jackson, R. B.: A global meta-analysis of soil
547 exchangeable cations, pH, carbon, and nitrogen with afforestation, *Ecological*
548 *Applications*, 19, 2228–2241, <https://doi.org/10.1890/08-1730.1>, 2009.
- 549 Berthrong, S. T., Piñeiro, G., Jobbágy, E. G., and Jackson, R. B.: Soil C and N changes with
550 afforestation of grasslands across gradients of precipitation and plantation age, 22, 76–
551 86, <https://doi.org/10.1890/10-2210.1>, 2012.
- 552 Binkley, Dan et al. The interactions of climate, spacing and genetics on clonal *Eucalyptus*
553 plantations across Brazil and Uruguay. *Forest Ecology And Management.* Amsterdam:
554 Elsevier Science Bv, v. 405, p. 271-283, doi.org/10.1016/j.foreco.2017.09.050
- 555 Bonfante, A., Basile, A., and Bouma, J.: Targeting the soil quality and soil health concepts
556 when aiming for the United Nations Sustainable Development Goals and the EU Green
557 Deal, *SOIL*, 6, 453–466, <https://doi.org/10.5194/soil-6-453-2020>, 2020.
- 558 Borrelli, P., Robinson, D. A., Fleischer, L. R., Lugato, E., Ballabio, C., Alewell, C.,
559 Meusburger, K., Modugno, S., Schütt, B., Ferro, V., Bagarello, V., Oost, K. V.,
560 Montanarella, L., and Panagos, P.: An assessment of the global impact of 21st century
561 land use change on soil erosion, *Nat Commun*, 8, 2013, [https://doi.org/10.1038/s41467-](https://doi.org/10.1038/s41467-017-02142-7)
562 [017-02142-7](https://doi.org/10.1038/s41467-017-02142-7), 2017.
- 563 Caon, L., & Vargas, R. (2017). Threats to soils: Global trends and perspectives. A Contribution
564 from the Intergovernmental Technical Panel on Soils, Global Soil Partnership Food and

- 565 Agriculture Organization of the United Nations. In G. Pierzynski & Brajendra (Eds.),
566 Global land outlook working paper, 28 pp.
- 567 Carrasco-Letelier, L. and A. Beretta-Blanco. 2017. Soil erosion by water estimated for 99
568 Uruguayan basins. *Cien. Inv. Agr.* 44(2): 184-194.
- 569 Céspedes-Payret et al. 2012 *Science of the Total Environment* 438 (2012) 549–557.
570 <http://dx.doi.org/10.1016/j.scitotenv.2012.08.075>
- 571 Cook, R. L., Binkley, D., and Stape, J. L.: Eucalyptus plantation effects on soil carbon after
572 20years and three rotations in Brazil, *Forest Ecology and Management*, 359, 92–98,
573 <https://doi.org/10.1016/j.foreco.2015.09.035>, 2016.
- 574 Dass, P. et al 2018 Grasslands may be more reliable carbon sinks than forests in California.
575 *Environ. Res. Lett.* 13 074027. <https://doi.org/10.1088/1748-9326/aacb39>
- 576 De Faccio Carvalho, P. C., Savian, J. V., Chiesa, T. D., Souza Filho, W. D., Terra, J. A., Pinto,
577 P., Martins, A. P., Villarino, S., Da Trindade, J. K., De Albuquerque Nunes, P. A., and
578 Piñeiro, G.: Land-use intensification trends in the Rio de la Plata region of South
579 America: toward specialization or recoupling crop and livestock production, *Front.*
580 *Agr. Sci. Eng.*, 8, 97, <https://doi.org/10.15302/J-FASE-2020380>, 2021.
- 581 Di Sacco, A.; Hardwick, K.; Blakesley, D.; Brancalion, P.H.S.; Breman, E.; Cecilio Rebola,
582 L.; Chomba, S.; Dixon, K.; Elliott, S.; Ruyonga, G.; Shaw, K.; Smith, P.; Smith, R.J.;
583 Antonelli, A. Ten Golden Rules for Reforestation to Optimise Carbon Sequestration,
584 Biodiversity Recovery and Livelihood Benefits. <https://doi.org/10.1111/gcb.15498>.
585 2020
- 586 Durán, A., Morrás, H., Studdert, G., and Liu, X.: Distribution, properties, land use and
587 management of Mollisols in South America, *Chin. Geogr. Sci.*, 21, 511,
588 <https://doi.org/10.1007/s11769-011-0491-z>, 2011.
- 589 ESRI 2018. ArcGIS Desktop: Release 10.3. Redlands, CA: Environmental Systems Research
590 Institute.
591 https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051232.pdf
- 592 FAO 2019. Land Use. Published online at OurWorldInData.org. Retrieved from:
593 '<https://ourworldindata.org/land-use>' [Online Resource]
- 594 Fernández, R., Frasier, I., Noellemeyer, E., and Quiroga, A.: Soil quality and productivity
595 under zero tillage and grazing on Mollisols in Argentina – A long-term study,
596 *Geoderma Regional*, 11, 44–52, <https://doi.org/10.1016/j.geodrs.2017.09.002>, 2017.

- 597 Fox, J., Weisberg, S. (2019) An R Companion to Applied Regression, Third Edition, Sage.
- 598 Friggens, N. L., Hester, A. J., Mitchell, R. J., Parker, T. C., Subke, J., and Wookey, P. A.: Tree
599 planting in organic soils does not result in net carbon sequestration on decadal
600 timescales, *Glob Change Biol*, 26, 5178–5188, <https://doi.org/10.1111/gcb.15229>,
601 2020.
- 602 García-Préchac, F., Ernst, O., Siri-Prieto, G., and Terra, J. A.: Integrating no-till into crop–
603 pasture rotations in Uruguay, *Soil and Tillage Research*, 77, 1–13,
604 <https://doi.org/10.1016/j.still.2003.12.002>, 2004.
- 605 Gomiero, Tiziano. 2016. "Soil Degradation, Land Scarcity and Food Security: Reviewing a
606 Complex Challenge" *Sustainability* 8, no. 3: 281. <https://doi.org/10.3390/su8030281>
- 607 Hernández J, del Pino A, Salvo L, Arrarte G. Nutrient export and harvest residue
608 decomposition patterns of a *Eucalyptus dunnii* Maiden plantation in temperate climate
609 of Uruguay. *Forest Ecology Management* 2009; 258(2): 92-99.
610 [10.1016/j.foreco.2009.03.050](https://doi.org/10.1016/j.foreco.2009.03.050)
- 611 Hernández, J., del Pino, A., Vance, E. D., Califra, Á., Del Giorgio, F., Martínez, L., and
612 González-Barrios, P.: *Eucalyptus* and *Pinus* stand density effects on soil carbon
613 sequestration, *Forest Ecology and Management*, 368, 28–38,
614 <https://doi.org/10.1016/j.foreco.2016.03.007>, 2016.
- 615 Hernandez-Ramirez, G., Sauer, T. J., Chendev, Y. G., and Gennadiev, A. N.: Nonlinear
616 turnover rates of soil carbon following cultivation of native grasslands and subsequent
617 afforestation of croplands, *SOIL*, 7, 415–431, <https://doi.org/10.5194/soil-7-415-2021>,
618 2021.
- 619 IUSS Working Group WRB. 2015. World Reference Base for Soil Resources 2014, update
620 2015 International soil classification system for naming soils and creating legends for
621 soil maps. World Soil Resources Reports No. 106. FAO, Rome.
622 <https://www.fao.org/3/i3794en/i3794en.pdf>
- 623 Jackson et al 2005 Trading Water for Carbon with Biological Carbon Sequestration. *Science*
624 310, 1944 (2005); DOI: 10.1126/science.1119282
- 625 Jackson, R.: Soil nutrient stocks are maintained over multiple rotations in Brazilian Eucalyptus
626 plantations | Elsevier Enhanced Reader, 13, n.d.
- 627 Jaurena M, Durante M, Devincenzi T, Savian JV, Bendersky D, Moojen FG, Pereira M, Soca
628 P, Quadros FLF, Pizzio R, Nabinger C, Carvalho PCF and Lattanzi FA (2021) Native
629 Grasslands at the Core: A New Paradigm of Intensification for the Campos of Southern

- 630 South America to Increase Economic and Environmental Sustainability. *Front. Sustain.*
631 *Food Syst.* 5:547834. doi: 10.3389/fsufs.2021.547834
- 632 Jobbagy, E. G., and R. B. Jackson. 2003. Patterns and mechanisms of soil acidification in the
633 conversion of grasslands to forests. *Biogeochemistry* 64:205–229.
- 634 Jobbagy, E. G., and R. B. Jackson. 2004. The uplift of soil nutrients by plants: biogeochemical
635 consequences across scales. *Ecology* 85:2380–2389.
- 636 Kaushal, S. S., Gold, A. J., Bernal, S., Johnson, T. A. N., Addy, K., Burgin, A., Burns, D. A.,
637 Coble, A. A., Hood, E., Lu, Y., Mayer, P., Minor, E. C., Schroth, A. W., Vidon, P.,
638 Wilson, H., Xenopoulos, M. A., Doody, T., Galella, J. G., Goodling, P., Haviland, K.,
639 Haq, S., Wessel, B., Wood, K. L., Jaworski, N., and Belt, K. T.: Watershed ‘chemical
640 cocktails’: forming novel elemental combinations in Anthropocene fresh waters,
641 *Biogeochemistry*, 141, 281–305, <https://doi.org/10.1007/s10533-018-0502-6>, 2018.
- 642 Kellogg, C. E. "Soil survey division staff: soil survey manual. Chapter 3" United States
643 Department of Agriculture, Washington (1993).
- 644 Kim, K., Daly, E.J., M. Gorzelak, Hernandez-Ramirez, G. 2022. Soil organic matter pools
645 response to perennial grain cropping and nitrogen fertilizer. *Soil and Tillage Research.*
646 220, 105376 <https://doi.org/10.1016/j.still.2022.105376>
- 647 Lambin, E.F.; Gibbs, H.K.; Ferreira, L.; Grau, R.; Mayaux, P.; Meyfroidt, P.; Morton, D.C.;
648 Rudel, T.K.; Gasparri, I.; Munger, J. Estimating the world’s potentially available
649 cropland using a bottom-up approach. *Glob. Environ. Chang.* 2013, 23, 892–901.
- 650 Lanfranco, B. and Sapriza, G.: El índice CONEAT como medida de productividad y valor de
651 la tierra., 57, n.d.
- 652 Lavado, R.S., Rodríguez, M.B., Scheiner, S.D., Taboada, M.A., Rubio, G., Alvarez, R.,
653 Alconada, M., Zubillaga, M.S., 1998. Heavy metals in soils of Argentina: comparison
654 between urban and agricultural soils. *Communications in Soil Science and Plant*
655 *Analysis* 29, 1913–1917.
- 656 Lavado, R.S., Zubillaga, M.S., Alvarez, R., Taboada, M.A., 2004. Baseline levels of potentially
657 toxic elements in pampas soils. *Soil and Sediment Contamination: An International*
658 *Journal* 13, 329–339.
- 659 Li, H., Jiang, L., You, C., Tan, B., and Yang, W.: Dynamics of heavy metal uptake and soil
660 heavy metal stocks across a series of Masson pine plantations, *Journal of Cleaner*
661 *Production*, 269, 122395, <https://doi.org/10.1016/j.jclepro.2020.122395>, 2020.

- 662 Liu, X., Lee Burras, C., Kravchenko, Y. S., Durán, A., Huffman, T., Morras, H., Studdert, G.,
663 Zhang, X., Cruse, R. M., and Yuan, X.: Overview of Mollisols in the world:
664 Distribution, land use and management, *Can. J. Soil. Sci.*, 92, 383–402,
665 <https://doi.org/10.4141/cjss2010-058>, 2012.
- 666 Mayer M., Prescott C.E., Abaker W.E.A., Augusto L., Cécillon L., Ferreira G.W.D., James J.,
667 Jandl R., Katzensteiner K., Laclau J.-P., Laganière J., Nouvellon Y., Paré D., Stanturf
668 J.A., Vanguelova E.I., Vesterdal L. Influence of forest management activities on soil
669 SOC stocks: A knowledge synthesis. *Forest Ecology and Management*, Volume 466,
670 2020
- 671 McMahon, D. E., Vergütz, L., Valadares, S. V., Silva, I. R. da, and Jackson, R. B.: Soil nutrient
672 stocks are maintained over multiple rotations in Brazilian Eucalyptus plantations,
673 *Forest Ecology and Management*, 448, 364–375,
674 <https://doi.org/10.1016/j.foreco.2019.06.027>, 2019.
- 675 Merino, A., Balboa, M. A., Rodríguez Soalleiro, R., and González, J. G. Á.: Nutrient exports
676 under different harvesting regimes in fast-growing forest plantations in southern
677 Europe, *Forest Ecology and Management*, 207, 325–339,
678 <https://doi.org/10.1016/j.foreco.2004.10.074>, 2005.
- 679 MGAP (Ministerio de Ganadería, Agricultura y Pesca) 2020. CONEAT. Online:
680 <https://www.gub.uy/ministerio-ganaderia-agricultura-pesca/politicas-y-gestion/coneat>.
- 681 Modernel, P., Rossing, W. A. H., Corbeels, M., Dogliotti, S., Picasso, V., and Tittonell, P.:
682 Land use change and ecosystem service provision in Pampas and Campos grasslands
683 of southern South America, *Environ. Res. Lett.*, 11, 113002,
684 <https://doi.org/10.1088/1748-9326/11/11/113002>, 2016.
- 685 OAS (Organization of American States, Department of Sustainable Development) 1994.
686 Uruguay - Proyecto Regional de Alternativas para la Inversión Forestal. Available
687 online: <http://www.oas.org/dsd/publications/unit/oea20s/oea20s.pdf>.
- 688 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.
689 R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and
690 Wagner, H.: *vegan: Community Ecology Package*, 2020.
- 691 Ortiz, J., Dube, F., Neira, P., Panichini, M., Stolpe, N.B., Zagal, E. and Martínez-Hernández,
692 P.A. Soil Quality Changes within a (*Nothofagus obliqua*) Forest Under Silvopastoral
693 Management in the Andes Mountain Range, South Central Chile. *Sustainability*;
694 12(17):6815. <https://doi.org/10.3390/su12176815>, 2020.

- 695 Paul, K. I., Polglase, P. J., Nyakuengama, J. G., and Khanna, P. K.: Change in soil carbon
696 following afforestation, *Forest Ecology and Management*, 168, 241–257,
697 [https://doi.org/10.1016/S0378-1127\(01\)00740-X](https://doi.org/10.1016/S0378-1127(01)00740-X), 2002.
- 698 Piñeiro, D. E.: Land grabbing: concentration and “foreignisation” of land in Uruguay, *Canadian*
699 *Journal of Development Studies / Revue canadienne d'études du développement*, 33,
700 471–489, <https://doi.org/10.1080/02255189.2012.746216>, 2012.
- 701 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for
702 Statistical Computing, Vienna, Austria, available at: <https://www.R-project.org/>, last
703 access: 2 March 2021.
- 704 Richter, D. D.: Game Changer in Soil Science. The Anthropocene in soil science and
705 pedology., *J. Plant Nutr. Soil Sci.*, 183, 5–11, <https://doi.org/10.1002/jpln.201900320>,
706 2020.
- 707 Roca, N.: Heavy metal background levels in rural soils: a case study in Pampean soils
708 (Argentina), 10, 2015.
- 709 Sadzawka, A., Carrasco, R. M. A., Grez, Z. R., Mora, G. M, Flores, P. H. Neaman, A. 2006.
710 Métodos de análisis recomendados para los suelos de Chile. Revisión 2006. Santiago,
711 Chile: Serie Actas - Instituto de Investigaciones Agropecuarias. Available online:
712 <https://hdl.handle.net/20.500.14001/8541>.
- 713 Sandoval López, D.M., Arturi, M.F., Goya, J.F. et al. *Eucalyptus grandis* plantations: effects
714 of management on soil carbon, nutrient contents and yields. *J. For. Res.* 31, 601–611
715 (2020). <https://doi.org/10.1007/s11676-018-0850-z>.
- 716 Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and
717 interpreting soil surveys. 2nd edition.
718 https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051232.pdf.
- 719 Veldman, J. W., Overbeck, G. E., Negreiros, D., Mahy, G., Le Stradic, S., Fernandes, G. W.,
720 Durigan, G., Buisson, E., Putz, F. E., and Bond, W. J.: Where Tree Planting and Forest
721 Expansion are Bad for Biodiversity and Ecosystem Services, *BioScience*, 65, 1011–
722 1018, <https://doi.org/10.1093/biosci/biv118>, 2015.
- 723 Waters, C. N., Zalasiewicz, J., Summerhayes, C., Barnosky, A. D., Poirier, C., Ga uszka, A.,
724 Cearreta, A., Edgeworth, M., Ellis, E. C., Ellis, M., Jeandel, C., Leinfelder, R., McNeill,
725 J. R., Richter, D. d., Steffen, W., Syvitski, J., Vidas, D., Wagreich, M., Williams, M.,
726 Zhisheng, A., Grinevald, J., Odada, E., Oreskes, N., and Wolfe, A. P.: The

727 Anthropocene is functionally and stratigraphically distinct from the Holocene, *Science*,
728 351, aad2622–aad2622, <https://doi.org/10.1126/science.aad2622>, 2016.





729 Wingeyer, A.B.; Amado, T.J.C.; Pérez-Bidegain, M.; Studdert, G.A.; Perdomo Varela, C.H.;
730 Garcia, F.O.; Karlen, D.L. Soil Quality Impacts of Current South American
731 Agricultural Practices. *Sustainability* 2015, 7, 2213–2242.

732 Wu, R., Alvareda, E., Polya, D., Blanco, G., and Gamazo, P.: Distribution of Groundwater
733 Arsenic in Uruguay Using Hybrid Machine Learning and Expert System Approaches,
734 *Water*, 13, 527, <https://doi.org/10.3390/w13040527>, 2021.

735 Zagal, E, Sadzawka, A. 2007. Protocolo de métodos de análisis para suelos y lodos. Available
736 online: https://www.sag.cl/sites/default/files/METODOS_LODOS_SUELOS.pdf.

737 Zurbriggen, C., González-Lago, M., Baraibar, M., Baethgen, W., Mazzeo, N. and Sierra, M.,
738 2020. Experimentation in the Design of Public Policies: The Uruguayan Soils
739 Conservation Plans. *Iberoamericana – Nordic Journal of Latin American and*
740 *Caribbean Studies*, 49(1), pp.52–62. DOI: <http://doi.org/10.16993/iberoamericana.459>.


741



742

743 **Figure 1:** Study design and sampling. (a) Location of the 28 monitoring sites across Uruguay
744 including land use types sampled (grassland (b); timber plantations, native forest and
745 agricultural land. Proportion of plots with particular category of soil classification according to
746 the World Reference Base (WRB, c), Soil Taxonomy (d) and CONEAT (Comisión Nacional
747 de Estudio Agronómico de la Tierra, e). The Uruguayan CONEAT index provides a detailed
748 classification that takes into account soil type, texture, natural vegetation, altitude and geology
749 (see details in chapter 2.3). Photos: RuralFutures.







750




751

752 **Figure 2.** Spearman's rank correlations for parameters of topsoils (n=80) regarding (a) fertility
 753 proxies, (b) acidity and (c) trace metals. The size of the circle refers to the absolute value of
 754 the correlation coefficient (ρ) and the color refers to the direction (positive or negative) of the
 755 correlation. Empty slots show correlations with $p > 0.05$.


756




758 **Figure 3.** Non-metric multidimensional scaling showing significant clustering differences
759 among samples from (a) grassland (GL), timber plantations (TP) and native forests (NF); (b)
760 among samples from *Pinus* and *Eucalyptus* plantations; (c) among land uses (including
761 agriculture AC) in Argiudolls; (d) among land uses in Argiudolls & Hapluderts; (e) among land
762 uses in Argiudolls, Hapudolls & Hapludalf and (f) among land uses in HapludalFs &
763 Hapludults.
764



766 **Figure 4.** Violin box plots for significant Kruskal-Wallis Tests across evaluated land uses (i.e.
767 grassland (GL), timber plantation (TP) and native forest (NF)) for each variable. Significance
768 in posterior Wilcoxon pairwise comparisons is depicted as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$
769 (***), $p < 0.0001$ (****).
770



772 **Figure 5.** Violin box plots for significant Kruskal-Wallis Tests in ‘Argiudolls’ Soil Taxonomy
773 category for fertility variables across available land uses (GL: Grassland, TP: Timber
774 plantation, NF: Native forest, AC: Agriculture). Significance in posterior Wilcoxon pairwise
775 comparisons is depicted as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), $p < 0.0001$ (****).
776



778

779 **Figure 6.** Acidity of all topsoils samples (a) and (b) according to different land use (GL:

780 Grassland, TP: Timber plantation, NF: Native forest, AC: Agriculture). Proportion of samples

781 with a given acidic category (Kellogg, 1993). pH of topsoil given as violin box plots for

782 significant Kruskal-Wallis Tests across different land uses independent from soil Order (c) and

783 separated per soil Orders (d-f). Sample number (n) is given. Significance in posterior Wilcoxon

784 pairwise comparisons is depicted as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), $p < 0.0001$ (****).

785

786 **Table 1.** General characteristics of Uruguayan topsoils: descriptive statistics for the parameters
787 of all single samples (n) across different land uses and classification to different soil types (for
788 details on different land uses i.e. grassland (GL), timber plantations (TP), native forests (NF)
789 and agricultural land (AC) see Appendix A1-A2 and on different soil types see Appendix A3-
790 4). SD = standard derivation.

Variable	n	Mean±SD	Min-Max	Samples by land use			
				GL	TP	NF	AC
Soil moisture (%)	280	18.9±10.8	1.34-51.5	115	93	57	15
N _{total} (%)	280	0.2±0.14	0.04-1.2	115	93	57	15
C _{total} (%)	280	3±2.3	0.4-25.7	115	93	57	15
C/N ratio	279	13.3±1.8	9.1-20.6	115	93	56	15
SOC (%)	267	5.6±3	0.8-16.4	110	90	52	15
P (mg kg ⁻¹)	278	295±166	43-1008	114	93	57	14
Ca (cmol kg ⁻¹)	82	9.4±8.9	0.3-42.7	32	27	18	5
Mg (cmol kg ⁻¹)	82	2.5±2.2	0.1-9.7	32	27	18	5
K (cmol kg ⁻¹)	82	0.4±0.25	0.03-1.44	32	27	18	5
Na (cmol kg ⁻¹)	82	0.1±0.14	0-0.69	32	27	18	5
Zn (mg kg ⁻¹)	82	1.9±2.1	0.1-9.7	32	27	18	5
Cu (mg kg ⁻¹)	82	5.4±22.1	0.3-161.2	32	27	18	5
Fe (mg kg ⁻¹)	82	134±75	5-309	32	27	18	5
Mn (mg kg ⁻¹)	81	25±17	0.7-93	32	27	17	5
pH	279	4.8±0.8	3.6-7.34	114	93	57	15
As (mg kg ⁻¹)	279	3.9±3.6	0.6-30.7	115	93	56	15
Cd (mg kg ⁻¹)	274	0.4±0.2	0.2-1	111	91	57	15
Cr (mg kg ⁻¹)	280	13.6±12.6	2-78.9	115	93	57	15
Pb (mg kg ⁻¹)	280	7±3.4	2-27.6	115	93	57	15
CEC (cmol kg ⁻¹)	82	12.4±10.7	0.5-50.1	32	27	18	5
K ⁺ /(Ca ²⁺ +Mg ²⁺)	82	0.07±0.08	0.01-0.43	32	27	18	5
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	82	0.07±0.07	0.01-0.42	32	27	18	5

791

792

793

Appendices

794 Appendix A

795 **Table A1.** Descriptive statistics of topsoil variables for all single soil types at grassland and
 796 native forests. Number of samples (n), mean, standard deviation (SD), minimum (Min) and
 797 maximum (Max) of each variable are given.

798

Variable (unit)	Grassland					Native forests				
	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max
Soil moisture (%)	115	20.4	11.5	1.5	51.5	57	25.8	9.3	6.3	50.3
N _{total} (%)	115	0.22	0.13	0.06	0.80	57	0.30	0.19	0.04	1.20
C _{total} (%)	115	2.9	1.9	0.7	10.8	57	4.2	3.6	0.5	25.7
C/N ratio	115	13.2	1.7	9.1	19.4	56	13.5	1.4	9.8	18.5
SOC (%)	109	5.3	2.6	1.4	15.2	53	7.5	3.8	1.1	16.4
P (mg/kg)	114	272	124	43	749	57	484	194	56	1009
Ca (cmol kg ⁻¹)	32	9.5	10.0	0.3	42.7	18	15.1	10.0	0.3	37.3
Mg (cmol kg ⁻¹)	32	2.2	2.1	0.18	9.7	18	4.8	2.6	0.12	8.9
K (cmol kg ⁻¹)	32	0.40	0.26	0.08	1.44	18	0.50	0.28	0.03	1.29
Na (cmol kg ⁻¹)	32	0.12	0.15	0.00	0.65	18	0.14	0.12	0.00	0.50
Zn (mg kg ⁻¹)	32	1.4	1.2	0.10	4.7	18	4.0	2.7	0.80	9.60
Cu (mg kg ⁻¹)	32	2.2	1.8	0.30	7.7	18	2.4	1.5	0.30	5.50
Fe (mg kg ⁻¹)	32	121	81	5	279	18	149	76	10	309
Mn (mg kg ⁻¹)	32	22	18	2	93	17	33	18	1	72
pH	114	4.9	0.9	3.6	7.3	57	5.1	0.7	3.6	7.2
As (mg kg ⁻¹)	115	4.3	4.5	0.6	30.7	56	4.1	2.5	1.2	13.6
Cd (mg kg ⁻¹)	111	0.41	0.17	0.17	1.00	57	0.45	0.15	0.17	0.81
Cr (mg kg ⁻¹)	115	12.3	9.2	2.6	49.5	57	22.8	17.3	2.0	70.7
Pb (mg kg ⁻¹)	115	6.8	3.0	2.3	15.9	57	9.5	4.8	2.9	27.7
CEC (cmol/ kg ⁻¹)	32	12.2	11.5	0.8	50.1	18	20.4	11.4	0.5	44.6
K ⁺ /(Ca ²⁺ +Mg ²⁺)	32	0.08	0.10	0.01	0.43	18	0.03	0.02	0.01	0.07
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	32	0.08	0.10	0.01	0.42	18	0.03	0.02	0.01	0.07

799

800 **Table A2.** Descriptive statistics for of topsoil variables for all single soil types at timber
 801 plantations and agricultural land. Number of samples (n), mean, standard deviation (SD),
 802 minimum (Min) and maximum (Max) of each variable are given.

803

Variable (unit)	Timber plantations					Agricultural land				
	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max
Soil moisture (%)	93	13.2	7.0	1.7	38.0	15	18.1	12.0	1.3	50.5
N _{total} (%)	93	0.18	0.10	0.04	0.61	15	0.22	0.06	0.13	0.29
C _{total} (%)	93	2.5	1.5	0.4	8.4	15	2.6	0.7	1.4	3.3
C/N ratio	93	13.6	2.1	10.0	20.6	15	11.7	0.7	10.5	12.8
SOC (%)	90	4.9	2.8	0.8	16.1	15	5.2	1.3	2.8	6.7
P (mg kg ⁻¹)	93	206	91	56	551	14	310	120	142	613
Ca (cmol kg ⁻¹)	27	5.2	4.8	0.6	18	5	10.8	5.5	7.2	20.4
Mg (cmol kg ⁻¹)	27	1.4	1.0	0.16	4.1	5	2.5	1.0	1.76	4.1
K (cmol kg ⁻¹)	27	0.32	0.21	0.09	0.93	5	0.46	0.23	0.29	0.85
Na (cmol kg ⁻¹)	27	0.11	0.16	0.00	0.69	5	0.06	0.02	0.03	0.08
Zn (mg kg ⁻¹)	27	0.94	0.74	0.10	2.8	5	3.7	3.7	0.60	9.7
Cu (mg kg ⁻¹)	27	1.5	0.9	0.5	4.5	5	57.7	78.8	0.8	161.0
Fe (mg kg ⁻¹)	27	143	69	16	290	5	115	62	63	210
Mn (mg kg ⁻¹)	27	25	16	6	72	5	17	5	10	23
pH	93	4.5	0.7	3.6	6.8	15	5.1	0.4	4.2	5.7
As (mg kg ⁻¹)	93	3.4	3.0	0.7	25.5	15	3.1	1.2	1.1	5.5
Cd (mg kg ⁻¹)	91	0.33	0.11	0.16	0.65	15	0.41	0.14	0.21	0.66
Cr (mg kg ⁻¹)	93	10.6	11.3	2.0	78.9	15	7.8	1.8	5.0	11.3
Pb (mg kg ⁻¹)	93	6.1	2.3	2.0	11.7	15	5.1	1.2	3.4	7.3
CEC (cmol kg ⁻¹)	27	7.0	5.6	1.0	20.6	5	13.8	6.5	9.5	25.0
K ⁺ /(Ca ²⁺ +Mg ²⁺)	27	0.08	0.07	0.02	0.25	5	0.04	0.03	0.02	0.09
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	27	0.08	0.07	0.02	0.25	5	0.04	0.03	0.02	0.08

804

805

806

807 **Table A3.** Descriptive statistics of topsoil variables for Argiudolls and for grassland and native
808 forests within of Argiudolls. Number of single samples (n), mean, standard deviation (SD),
809 minimum (Min) and maximum (Max) of each variable are given.

810

Variable (Unit)	Total Argiudolls					Grassland					Native forests				
	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max
Soil moisture (%)	77	18.3	10.9	1.3	50.5	31	19.0	9.2	5.0	49.0	12	26.4	13.4	9.0	48.2
N _{total} (%)	77	0.24	0.15	0.07	1.20	31	0.21	0.11	0.07	0.66	12	0.34	0.28	0.13	1.20
C _{total} (%)	77	3.3	3.0	1.1	25.7	31	2.9	1.8	1.1	10.8	12	5.3	6.5	1.7	25.7
C/N ratio	76	13.3	2.2	9.8	19.9	31	13.3	2.1	10.5	19.4	11	13.6	2.1	9.8	18.5
SOC (%)	73	5.8	2.5	2.1	16.1	30	5.2	1.9	2.1	10.0	9	7.4	2.1	4.8	11.0
P (mg kg ⁻¹)	76	280	122	93	693	30	234	78	93	353	12	438	155	155	693
Ca (cmol kg ⁻¹)	21	11.4	8.0	2.6	37.1	8	11.5	6.7	3.2	20.5	4	16.2	14.3	5.1	37.1
Mg (cmol kg ⁻¹)	21	2.3	1.4	0.5	6.1	8	2.0	0.8	0.9	3.2	4	3.5	2.5	1.0	6.1
K (cmol kg ⁻¹)	21	0.45	0.26	0.1	1.3	8	0.4	0.14	0.16	0.61	4	0.69	0.41	0.34	1.29
Na (cmol kg ⁻¹)	21	0.08	0.05	0.03	0.26	8	0.07	0.02	0.04	0.10	4	0.10	0.11	0.03	0.26
Zn (mg kg ⁻¹)	21	2.4	3.0	0.1	10	8	1.3	0.8	0.1	2.2	4	5.1	4.3	1.3	10
Cu (mg kg ⁻¹)	21	15.5	42.7	0.8	161	8	2.6	1.6	1.0	6.0	4	1.9	1.0	0.8	3.0
Fe (mg kg ⁻¹)	21	117	62	10	232	8	104	50	28	171	4	117	84	10	216
Mn (mg kg ⁻¹)	21	23.3	10.4	7.1	43.6	8	20.5	10.4	7.1	33.8	4	28.8	10.4	15.4	39.4
pH	77	5.0	0.7	3.8	6.6	31	4.9	0.7	3.9	6.6	12	5.2	0.5	4.3	6.1
As (mg kg ⁻¹)	76	3.2	1.4	1.0	12.2	31	2.9	0.9	1.5	4.9	11	4.0	2.8	1.5	12.2
Cd (mg kg ⁻¹)	76	0.4	0.1	0.2	0.7	30	0.4	0.1	0.2	0.7	12	0.4	0.1	0.2	0.7
Cr (mg kg ⁻¹)	77	9.3	6.2	2.2	28.3	31	8.1	4.1	3.6	18.5	12	13.2	9.5	2.2	28.3
Pb (mg kg ⁻¹)	77	5.6	2.1	2.8	10.7	31	5.2	1.8	2.8	9.6	12	6.8	2.9	2.9	10.7
CEC (cmol kg ⁻¹)	21	14.2	9.3	3.4	44.6	8	14.0	7.3	4.5	23.1	4	20.4	16.7	6.4	44.6
K ⁺ /(Ca ²⁺ +Mg ²⁺)	21	0.04	0.02	0.02	0.09	8	0.03	0.02	0.02	0.07	4	0.04	0.01	0.03	0.06
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	21	0.04	0.02	0.02	0.08	8	0.03	0.02	0.02	0.07	4	0.04	0.01	0.03	0.06

811

812

813 **Table A4.** Descriptive statistics of topsoil variables for Argiudolls and timber plantations and
814 Agricultural land within of Argiudolls. Number of single samples (n), mean, standard deviation
815 (SD), minimum (Min) and maximum (Max) of each variable are given.

816

Variable (unit)	n	Total Argiudolls				n	Timber plantations				n	Agricultural land			
		Mean	SD	Min	Max		Mean	SD	Min	Max		Mean	SD	Min	Max
Soil moisture (%)	77	18.3	10.9	1.3	50.5	22	14.0	7.8	5.5	38.0	12	16.5	13.0	1.3	50.5
N_total (%)	77	0.24	0.15	0.07	1.20	22	0.22	0.10	0.10	0.45	12	0.22	0.06	0.13	0.29
C_total (%)	77	3.3	3.0	1.1	25.7	22	3.2	1.6	1.2	8.0	12	2.6	0.7	1.4	3.3
C/N ratio	76	13.3	2.2	9.8	19.9	22	14.0	2.6	11.3	19.9	12	11.8	0.6	10.7	12.7
SOC (%)	73	5.8	2.5	2.1	16.1	22	6.3	3.3	2.3	16.1	12	5.2	1.4	2.8	6.7
P (mg kg ⁻¹)	76	280	122	93	693	22	234	70	111	353	12	321	119	204	613
Ca (cmol kg ⁻¹)	21	11.4	8.0	2.6	37.1	5	6.9	4.1	2.6	11.4	4	11.7	5.9	7.7	20.4
Mg (cmol kg ⁻¹)	21	2.3	1.4	0.5	6.1	5	1.5	0.8	0.5	2.7	4	2.6	1.1	1.8	4.1
K (cmol kg ⁻¹)	21	0.45	0.26	0.09	1.29	5	0.34	0.21	0.09	0.60	4	0.46	0.27	0.29	0.85
Na (cmol kg ⁻¹)	21	0.08	0.05	0.0	0.3	5	0.1	0.06	0.05	0.20	4	0.06	0.02	0.04	0.08
Zn (mg kg ⁻¹)	21	2.4	3.0	0.1	9.7	5	0.5	0.4	0.2	1.2	4	4.1	4.2	0.6	9.7
Cu (mg kg ⁻¹)	21	15.5	42.7	0.8	161	5	2.0	1.5	1.0	4.5	4	71.8	83.4	0.8	161
Fe (mg kg ⁻¹)	21	117	62	10	232	5	158	73	69	232	4	91	38	63	143
Mn (mg kg ⁻¹)	21	23.3	10.4	7.1	43.6	5	29.8	9.8	17.5	43.6	4	15.3	3.7	9.9	18.1
pH	77	5.0	0.7	3.8	6.6	22	4.9	0.8	3.8	6.4	12	5.2	0.3	4.9	5.7
As (mg kg ⁻¹)	76	3.2	1.4	1.0	12.2	22	3.0	1.0	1.0	4.4	12	3.4	1.2	2.1	5.5
Cd (mg kg ⁻¹)	76	0.39	0.13	0.18	0.74	22	0.37	0.12	0.18	0.60	12	0.43	0.13	0.27	0.66
Cr (mg kg ⁻¹)	77	9.3	6.2	2.2	28.3	22	9.5	7.3	2.7	26.4	12	8.2	1.7	6.0	11.3
Pb (mg kg ⁻¹)	77	5.6	2.1	2.8	10.7	22	5.6	2.1	2.8	8.5	12	5.3	1.2	3.9	7.3
CEC (cmol kg ⁻¹)	21	14.2	9.3	3.4	44.6	5	8.8	4.6	3.4	13.6	4	14.8	7.0	9.9	25.0
K ⁺ /(Ca ²⁺ +Mg ²⁺)	21	0.04	0.02	0.02	0.1	5	0.04	0.03	0.02	0.07	4	0.04	0.03	0.02	0.09
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	21	0.04	0.02	0.02	0.08	5	0.04	0.02	0.02	0.07	4	0.04	0.03	0.02	0.08

817