



1 Title page

2 **Back to the future- Conservative grassland management for**
3 **Anthropocene soils in the changed landscapes of Uruguay?**

4 Ina Säumel¹, Leonardo R. Ramírez¹, Sarah Tietjen², Marcos Barra³ and Erick Zagal⁴

5 ¹Integrative Research Institute THESys Transformation of Human-Environment-Systems
6 Humboldt-Universität zu Berlin, Unter den Linden 6, Berlin, 10099, Germany.

7 ²Leibniz Institute of Vegetable and Ornamental Crops (IGZ) e.V., Theodor-Echtermeyer-Weg
8 1, 14979 Großbeeren, Germany

9 ³Städtisches Klinikum Dessau, Auenweg 38, 06847 Dessau-Roßlau, Germany

10 ⁴Departamento de Suelos y Recursos Naturales, Universidad de Concepción, Campus Chillán,
11 Vicente Méndez 595, Chillán, Chile

12 *Correspondence to:* Ina Säumel (ina.saeumel@hu-berlin.de)

13 **Abstract.** The ‘soils of the anthropocene’ are predominately agricultural. To understand them,
14 we analysed agri- and silvicultural intensification of Uruguayan grasslands in a country wide
15 survey on fertility proxies, pH and trace metals in topsoils originating from different land uses.
16 We observed a loss of nutrients, trace metals and organic matter from grassland, crops and
17 timber plantations, and its accumulation in the topsoils of riverine forests. The translocation of
18 nutrients and organic matter across the landscape to the erosion base depends on local land use
19 trajectories. Increasing soil acidification is driving a positive feedback loop, and land use
20 intensification is leading to degradation of local black soils within a few decades. Our data
21 raises questions about the resilience and carrying capacity of Uruguayan soils with regard to
22 currently implemented highly productive management forms, including the use of timber



23 plantation for carbon sequestration, and supports more conservative forms of extensive
24 management on the grassland biome.

25

26 **1. Introduction**

27 Human activities alter the bio- and pedosphere, leaving a footprint of such a magnitude that it
28 can be verified stratigraphically (Waters et al., 2016). This unprecedented transformational
29 force is intimately related to the expansion of societies and its productive frontiers, causing a
30 loss of biodiversity, habitat and soil degradation and, consequently, to ecosystem modification
31 (Foley et al., 2005, Borrelli et al., 2017). In this context, soil sciences have transitioned from
32 studies on natural soil formation to the science of ‘anthropogenesis’ (Richter, 2020),
33 focussing on the ‘soils of the anthropocene’ that are predominately agricultural (51 Mio. km²)
34 or urban (1.5 Mio km²; FAO, 2019).

35 The temperate grasslands of South America have historically been characterised by rolling
36 plains and low hills that have been extensively exploited for cattle production and its
37 derivatives since the arrival of European colonization. The Río de la Plata grasslands are one
38 of our planet’s four major black soil regions (Durán et al., 2011; Liu et al., 2012), and some of
39 the most fertile soils in the world. Playing an important role in the global food production, these
40 are characterized by a thick, humus and base cation rich and a high cation exchange capacity
41 throughout their profile. Maintaining their properties are therefore crucial to developing
42 sustainable and productive agriculture (Durán et al., 2011; Liu et al., 2012).

43 Today, the 'Uruguayan savanna' is part of the top-three most critically endangered biomes
44 (Veldman et al., 2015). In recent decades, however, the grassland has decreased due to the
45 expansion of cash crops and *Eucalyptus* plantations, both of which are promoted by national
46 legislation including land grabbing and trans-nationalization (Piñeiro, 2012). This land use



47 intensification, with its increased input of energy, nutrients and pesticides, leads to an overall
48 loss of soil fertility and increasing toxicity related to acidification, salinization and
49 contaminants (Liu et al., 2012; Borrelli et al., 2017). Ecological, economic and cultural
50 functions of soils are severely degraded, and the degradation of black soils in South America
51 is of particular concern because they have only been heavily exploited for a comparatively
52 short period of time (Durán et al., 2011).

53 Since the first decades of the twentieth-century, compared to other disciplines, soil sciences
54 have received an extraordinary amount of attention in Uruguayan academia, governance, the
55 productive sector, and also in the general public, resulting in a national soil inventory program
56 in 1965. The subsequent classification of Uruguayan soils and their productivity (CONEAT
57 Index) remains an important source for today's land taxation and for management plans by the
58 legal conservation regulations, and provides a detailed classification that takes into account soil
59 type, texture, natural vegetation, altitude and geology (Lanfranco and Sapriza, 2011).

60 As soil degradation is extremely relevant for countries like Uruguay, which are
61 socioeconomically dependent on their soils (Zubriggen et al. 2020), it is a topic of discussion
62 for local farmers, academia, and the public. An actualization of the state of the art of soils and
63 related processes is needed (García-Préchac et al. 2004; De Faccio et al. 2021), particularly as
64 there has been little study of the impacts of the Uruguayan grassland intensification on soils
65 properties (Beretta-Blanco et al., 2019). At the same time, while a new paradigm for grassland
66 intensification with a wide set of means including fertilization has been proposed to increase
67 economic and environmental sustainability (Jaurena et al., 2021), it is urgent to get more
68 insights into the dynamics of nutrient in soils of Uruguay and their availability for crops
69 (Beretta-Blanco et al., 2019).

70 Soil classifications are mainly based on subsoils. However, we focus on topsoil as the most
71 relevant and very responsive interface for ecological processes and farmers management, since
72 understanding the state of the art of topsoils and its processes is crucial for developing



73 recommendations for sustainable land management practices. Due to the diversity of
74 perspectives on soil quality and health and related ecosystem services, operational procedures
75 for evaluation of soil functioning are still lacking (Bonfante et al. 2021). We contribute to a
76 better understanding of globally occurring degradation processes in the field of tension between
77 desired soil productivity, yield limits, especially in erosion sensitive soils, and necessary soil
78 conservation.

79 We therefore explored soil parameters describing actual chemical conditions of topsoils that
80 are parts of different soil groups and orders, and Uruguayan soil categories. Specifically, in
81 order to explore the gains and losses of macro and micro-nutrients and soil organic matter
82 across landscapes and to determine the impact of land use change on acidification and trace
83 metal mobility and related trade-offs with soil degradation and conservation, we analysed i)
84 the variation of fertility proxies, ii) levels of acidification (pH), and iii) trace metals
85 accumulation.

86 **2. Material and Methods**

87 **2.1 Study area and design**

88 Uruguay covers about 176,000 km², and has a population of 3.5 million, mainly in urban areas.
89 The Western half of Uruguay is dominated by Mollisols, developed on a wide range of
90 sediments from the Devonian to the Cenozoic era (partially associated with Vertisols), while
91 the Eastern plains and wetlands have, in addition to Mollisols, other soil types on the slopes,
92 rocks and on flood plans (i.e. Alfisols, Ultisols, Entisols and Inceptisols; Durán et al. 2011).
93 During a country-wide survey from December 2015 to March 2016, we collected 280 topsoil
94 samples of 0-10 cm depth from 101 plots (50x50m in grassland and crops; 100x100m in native
95 forests and timber plantations) distributed at 28 monitoring sites throughout Uruguay, South



96 America (Fig. 1a-b), using a stratified random design. In the first step, we randomly selected
97 monitoring sites across the country. In the second step, we contacted landowners to explore
98 their willingness to establish a long-term monitoring site. If the owner agreed, plot selection
99 was stratified by different rural land use types: grassland, timber plantation, native forest, and
100 crops. We sampled top soil three times at each land use at the edges of the plot, and stored
101 samples below 7°C until lab processing.

102 **2.2 Analysis of Soil Samples**

103 For gravimetric determination of humidity, the topsoils samples were dried at 105 °C until
104 constant weight. Next, lumps in the samples were broken down and the remaining plant
105 material was removed before sieving (2 mm) and ground.

106 We analysed 280 samples regarding macronutrients, pH and trace metals and 80 samples for
107 soluble cations and micronutrients. Among fertility-related variables, we measured the total
108 amount of the macronutrients phosphorus (P), carbon (C) and nitrogen (N), so obtaining the
109 C/N ratio. To determine total carbon and nitrogen, the samples were sieved again (0.5 mm) and
110 analyzed using a LECO TruSpec CN (USA) at a combustion temperature of 950 °C with ultra-
111 pure oxygen. In addition, the presence of inorganic carbon was tested for by adding
112 concentrated hydrochloric acid, the presence of carbonates was assessed to obtain the amount
113 of organic carbon (C_{org}) and soil organic matter (SOM) as $C_{org} \times 2$ (Chenu et al. 2015).

114 We determined concentrations of the soluble cations calcium (Ca), magnesium (Mg),
115 potassium (K), sodium (Na), and the micronutrients copper (Cu), zinc (Zn), manganese (Mn)
116 and iron (Fe) by atomic spectroscopy (Unicam AAS Solaar 969, Thermo Electron Corporation,
117 USA), extracting with ammonium acetate (1 mol l⁻¹, pH 7), and with DTPA-CaCl₂-TEA at (pH
118 7.3). We calculated the cation-exchange capacity (CEC). Acidity was measured by adding
119 calcium chloride (0.01 mol) to the samples in a 2.5:1 proportion, and after shaking and two



120 hours rest, read with a pH meter (HI2550 meter, Hanna Instruments, USA)). For these
121 variables, each tenth sample was duplicated. We categorized the values using the USDA
122 Natural Resources Conservation Service classification (Kellogg, 1993).

123 To determine the concentrations of arsenic (As), cadmium (Cd), chromium (Cr) and lead (Pb),
124 samples were further sieved as before (0.5 mm), weighed out into a digesting container, and
125 extracted with a mixture of nitric acid (70%) and hydrochloric acid (30%) in a 1:3 proportion.
126 The digestion took place in a Titan MPS (Perkin Elmer, USA) microwave at a programme
127 suitable for this digestion (Method 3051A). During the proceedings, only deionized water Type
128 I ASTM1193 (EC max 0.06 to 0.1 $\mu\text{S}/\text{cm}$) was used. Reagents were used to eliminate traces of
129 other materials and to avoid contamination of the samples. The trace metals were determined
130 by inductively coupled plasma optical emission spectroscopy using the Optima 8000 (Perkin
131 Elmer) with metal-associated wavelengths of 193.696 nm for As, 228.802 nm for Cd, 267.716
132 nm for Cr and 220.353 nm for Pb.

133 Total P concentration of phosphorus was determined calorimetrically after microwave-assisted
134 digestion with Unicam spectrometer at a wavelength of 660 nm. For all these variables, a
135 repetition for each round of the microwave digestion was made.

136 **2.3 Soil classification**

137 We intersected the coordinates of the centre of the plots with maps containing geospatial
138 information on the classification of the Uruguayan soils using ArcGIS 10.3 (ESRI, 2018). For
139 Soil Groups classification, we used the of the World Reference Base for Soil Resources (WRB;
140 IUSS Working Group, 2015); for Soil Orders, the USDA soil taxonomy (Soil Survey Staff,
141 1999); and for the local Uruguayan classification, Soil CONEAT (Comisión Nacional de
142 Estudio Agronómico de la Tierra) Groups categories, which include productive capacity of
143 cattle and sheep (MGAP, 2021).



144 **2.4 Data Analysis**

145 In a first step, we explored and prepared our database for further analysis. Exploring the
146 distribution of the soil parameters in R (R Core Team 2021), ruled out the normality of the data
147 using the Shapiro-Wilk Test and the homogeneity of variances with the Fligner-Killeen Test.
148 We tested for outliers using the 1.5-3 IQR threshold and the function *outlierTest* from the R
149 package *car* (Fox and Weisberg 2019), reviewing the flagged observations case by case in the
150 experimental context. The variables on soils characteristics showed generally positive skewed
151 distributions, in some cases multimodal, and tests showed evidence contrary to assumptions of
152 spatial autocorrelation, homoscedasticity and normality in most cases (Supplementary
153 Material: Table S1-S2; Fig. S1-S2).

154 Spearman's rank correlations (ρ) were calculated to explore linear associations between soil
155 parameters across all single samples and within different land uses. We used the *adonis*
156 function of the R package *vegan* v2.5-7 (Oksanen et al., 2020) with a Euclidean dissimilarity
157 matrix from our normalized soil data to perform PERMANOVA tests with 9999 permutations
158 to analyse the multivariate homogeneity of group dispersions based on differences on soil
159 parameters between land uses.

160 To compare the general effects of land use type on topsoil parameters, non-parametric Kruskal-
161 Wallis tests were carried out in R. When significant ($p \leq 0.05$), we used Pairwise Wilcoxon
162 Rank Sum tests with Benjamini & Hochberg correction to evaluate pairwise differences among
163 land uses.

164 We used non-metric multidimensional scaling (NMDS) as a robust unconstrained ordination
165 method to visualize patterns of top soil characteristics among all samples and within
166 subsamples (intersected with different soil Orders) across different land uses using the Bray-
167 Curtis dissimilarity matrix. The matrix was constructed with the *vegan* package to depict
168 patterns of all soil parameters in two dimensions (Fig.3a, c-f) and for the dataset without soluble



169 cations and micronutrients variables comparing subcategories within single land use types (i.e.
170 within grasslands in ‘undisturbed’, ‘partially grazed’ and ‘highly grazed’ plots; timber
171 plantations in ‘*Eucalyptus*’ or ‘*Pinus*’ plots; Fig. 3b).

172 **3. Results**

173 Nearly 75% of the sampled 101 plots intersected with the soil Group category 'Luvic Phaeozem'
174 (WRB; IUSS Working Group, 2015). A quarter of our plots were distributed among four other
175 Groups (i.e. Haplic Luvisols, Eutric Vertisols, Planosols, Lithic Leptisols; Fig. 1c). Half of our
176 plots intersected with the Orders 'Argiudolls' or 'Argiudolls & Hapluderts' (Soil Survey Staff,
177 1999), and about a quarter with 'Hapludalfs & Hapludults' or 'Argiudolls, Hapludolls &
178 Hapludalfs' (Fig. 1d). Our plots intersected with 32 different CONEAT categories (MGAP,
179 2021). The most frequent categories were '10.2', '9.3' and '2.12' (25% of our plots) and another
180 quarter with '9.1', '8.8', '9.6' and 'G03.11' (Fig. 1e).

181 **3.1 General characteristics of Uruguayan topsoils**

182 The measured topsoil parameters vary widely across Uruguay, between the different land uses
183 and classification to different soil orders (Table 1, Table A1-A4). The soil organic matter in
184 topsoils ranged between 0.8 to 16 percent, and was highest in native forests and lowest in timber
185 plantations. The mean of the C/N ratio was about 13, and lowest in crop soils. Phosphorous
186 ranged between 43 to 1009 mg kg⁻¹. We also observed a high variability for the micro- and
187 macronutrients and trace metals. The average cation-exchange capacity (CEC) was 12.41 cmol
188 kg⁻¹, highest in topsoils of native forest, followed by crops and grasslands and timber
189 plantations (Table 1, Table A1-A2).

190 For the whole data set, high correlation was found between P with SOM and Zn ($\rho=0.82$ and
191 0.76, respectively), and between Mg with Ca and Na ($\rho=0.82$ and 0.76, respectively; Fig. 2).



192 Similar results were observed within particular land uses, although in native forests, a negative
193 moderate correlation between Ca and Fe was observed (Fig. S3). There was high correlation
194 between pH and Ca ($\rho=0.89$; Fig. 2). In native forests, we also found similar correlation with
195 other soluble cations like Mg and K (Fig. S3). In park forests topsoils, there was a negative
196 correlation between pH and As and Pb ($\rho=-0.81$ and $\rho=-0.84$, respectively). In highly grazed
197 pastures and crops pH was highly correlated with Cr, and in crops also with As ($\rho=0.93$; Fig.
198 S3). Among trace metals, Spearman correlation was moderate between As with Cr and Pb and
199 high between Cr and Pb (Fig. 2). These correlation trends increased in grasslands, timber
200 plantations and crops. We also found a high correlation between cadmium and Cr in crops
201 ($\rho=0.81$). Phosphorus was highly correlated with Cr and As in pine plantations, while in
202 soybean crops the correlation was between phosphorus with Cr and Pb (Fig. S3).

203 **3.2 Topsoil characteristics clustered by land use**

204 We found differences in multivariate distribution of samples according to the different land
205 uses (i.e., grassland, timber plantations or native forests; Fig. 3a-f; Table S3-S8). All pairwise
206 comparison of land uses showed significant differences with all variables ($p=0.0001$; Fig. 3a;
207 Table S3). We analysed subcategories within a land use type using the dataset without soluble
208 cations and micronutrients variables, only finding significant differences between *Eucalyptus*
209 and *Pinus* stands ($p=0.0001$; Fig. 3b; Table S4) but not among different grassland subtypes.
210 We also found differences analysing subsamples of different soil order classification to the
211 different land uses (Fig. 3c-f; Table S5-S8). The samples intersected with 'Argiudolls' included
212 all plots on crops, and we found significant differences in all pairwise comparisons of land uses
213 except between grasslands and timber plantation ($p=0.0004$; Fig. 3c; Table S5). We further
214 found differences between timber plantations and native forests at soils of the 'Argiudolls &
215 Hapluderts' Orders ($p=0.0009$; Fig. 3d; Table S6) and between timber plantations and
216 grasslands or native forests in soils of the 'Argiudolls, Hapudolls & Hapludalf' Orders



217 ($p=0.0284$; Fig. 3e; Table S7). Results for samples in 'Hapludalfs & Hapludults' soils were
218 similar to those obtained at country scale ($p=0.0001$; Fig. 3f; Table S8).

219 **3.3 Differences in fertility proxies**

220 We found significantly higher values for the fertility proxies for SOM, P, Ca, Mg and Zn in
221 topsoils from native forests compared to grasslands and timber plantations (Fig. 4b-e, h).
222 Phosphorous was significantly higher in topsoils of grassland compared to those of timber
223 plantations (Fig. 4c). Potassium was significantly higher in topsoils of native forests compared
224 to timber plantations (Fig. 4f). At subcategories level, we found significantly higher amounts
225 of K in partially used grasslands in comparison to samples from highly grazed pastures, and
226 higher values of SOM ($p=0.002$), P ($p=0.059$), Na ($p=0.043$), K ($p=0.012$) and Zn ($p=0.048$)
227 in *Eucalyptus* compared to *Pinus* plantations (Fig. S4). Among native forests, samples from
228 riverine forests contain more Mg ($p=0.023$) and Na ($p=0.023$) in comparison to park forests.
229 Considering all samples within the Order 'Argiudolls', C/N ratio in agricultural topsoils are
230 lower compared to all other land uses. Phosphorus was higher in topsoils of native forests and
231 crops compared to grasslands and timber plantations. Soil organic matter was highest in native
232 forests (Fig. 5a-o; Table A3-A4).

233 **3.4 Soil Acidification**

234 Topsoil samples showed a markedly acid profile (median pH=4.66), with nearly 75% classified
235 as 'extremely acidic' and 'very strongly acidic' (Fig. 6a). We found significant differences across
236 land uses, with less acidic values in native forests, and lower pH in timber plantations (Fig.
237 6b). Comparing between land uses, we found more samples with neutral acidity in grasslands
238 and more with higher acidity in timber plantations (Fig. 6b-c). In addition, we found lower pH
239 in samples from *Pinus* compared to *Eucalyptus* stands ($p=0.018$). Results of analysis inside soil
240 Orders showed similar variations observed at country scale, with timber plantations being more
241 acid and native forest closer to neutral pH (Fig. 6d-f).



242 **3.5 Trace metal accumulation across land uses**

243 For As, Cd, Cr and Pb, we found significantly higher values in topsoils originating from native
244 forests compared to the grassland and timber plantation samples (Fig. 4l-o and 5l-o; Table A1-
245 A4). At the same time, samples from *Eucalyptus* plantations had higher levels of both Cr
246 ($p < 0.005$) and Pb ($p < 0.005$) than *Pinus* topsoils, while the same was observed for Cr ($p < 0.001$)
247 and Pb ($p < 0.05$) in riverine forests compared to park forests (Fig. S5).

248 **4. Discussion**

249 The vicious circle between the wish to stop soil degradation and concurrent increases in land
250 productivity to satisfy the increasing demand for food, fibres and energy has not been broken
251 since green revolution. Socio-economic and conventional management practices that drive soil
252 degradation have generated several traps, such as the 'inputs trap' where a reduced yield per
253 area is followed by higher fertilizer application or the 'credit or poverty trap' where economic
254 pressure forces the farmer to intensification (Gomiero, 2016). Caught in this loop, the soils of
255 the temperate grasslands of Uruguay have suffered strong degradation from erosion,
256 acidification, contamination, salinification and compaction (Zurbriggen et al., 2020). This is
257 clearly reflected in the results of our topsoil survey, which also adds interesting insights from
258 timber plantations, grasslands and native forests to an existing database consisting mainly of
259 crops and pastures samples from 2002-2014, which demonstrated the loss of organic matter by
260 25% and an increasing loss of nutrients (Beretta-Blanco et al. 2019). We contribute deeper
261 insights on fertility, acidification and trace metals accumulation in topsoils from a wide range
262 of different land uses, which is, to our knowledge, unique for the region since the CONEAT
263 classification (CONEAT Index, 1976).

264



265 **4.1 Translocation of elements in topsoils within landscape**

266 Our data demonstrate the accumulation of SOM, nutrients and trace metals in topsoil samples
267 from riverine forests, suggesting transport of soil particles from the surrounding land uses (e.g.,
268 grasslands, crop or timber plantations) to the borders of rivers, streams and creeks. Regional
269 soil erosion models estimate the loss of 2-5 tons ha⁻¹ year⁻¹ for a third of the country depending
270 on precipitation, topography, soil erodibility and land management (Carrasco-Letelier and
271 Beretta-Blanco 2017). One possible direct impact is the increasing eutrophication reported for
272 larger local rivers, although the models used by these authors did not link Chlorophyll-a
273 concentrations with agricultural land use (Beretta-Blanco and Carrasco-Letelier 2021 and
274 replies).

275 Organic matter content and the exchangeable cations are strongly reduced in topsoils of
276 grasslands, timber plantations and crops compared to native forests (Fig. 4b, d-h and 5b, d-h).
277 Of all the fertility proxies assessed, phosphorus in topsoil was most significantly affected by
278 different land uses, being highest in native forests (Fig. 4c and 5c). The cation exchange
279 capacity (CEC) was highest in native forests (more than 160 percent of the CEC in grasslands)
280 and lowest in timber plantations, reaching only the half of the CEC in grasslands (Table A1-
281 A2). Lower average nutrient contents and corresponding CEC reported for two timber
282 plantations in the East and one in the Norwest of Uruguay (Hernández et al., 2009 and 2016;
283 Céspedes-Payret et al., 2012) may result from a combined effect of topsoil degradation in
284 timber plantations due to management practices and to soil texture (Sandoval-Lopez et al.,
285 2020). The trees' uptake and the general export of nutrients from fast growing timber
286 plantations through harvesting is higher than the natural input into those systems (Merino et
287 al., 2005). This effect is particularly relevant because timber plantations in Uruguay are on
288 'forestry priority soils', which are generally soils with low fertility, superficial to moderate
289 depth and good drainage (OAS, 1994), so afforestation might reduce even more their soil
290 fertility.



291 Comparing neighbouring sites of grassland and afforested grasslands, our data set shows no
292 significant depletion of nutrients by timber plantations ($p=0.208$) but a slightly higher average
293 of CEC at grasslands (Table A1). In general, plants are expected to compensate the extraction
294 of cations from upper soil by the 'uplift of nutrient' from deeper horizons (Jobbagy and Jackson,
295 2004). This has been questioned for *Eucalyptus* plantations on sandy soils, where, for example,
296 an increase of potassium in the topsoil was not observed compared to the neighbouring
297 grassland (Céspedes-Payret et al., 2012). Phosphorous and cations decline with sand content,
298 so physicochemical factors such as the percentage of sand and organic matter, influence soil
299 fertility (Sandoval-Lopez et al., 2020). A study of *Eucalyptus* plantations in South-East Brazil
300 did not show a significant depletion of nutrients and carbon; in contrast, carbon, potassium and
301 calcium increased in the topsoil after twelve years over one or two harvest and fertilization
302 cycles (McMahon et al., 2019). To sum up, the patterns observed in the various sites in our
303 survey and the other studies indicate very complex interactions of numerous factors. Removal
304 of nutrients by high-yield timber plantations may exceed the capacity of nutrient exchange or
305 turnover of forest litter and wastes. Both stand management and environmental conditions (e.g.,
306 precipitation) influence nutrient and carbon stocks (McMahon et al., 2019; Sandoval-Lopez et
307 al., 2020).

308 The C/N ratio in topsoils ranged within values reported for grassland and timber plantations of
309 the region (see Berthrong et al., 2012). In contrast to that study, however, we observed no
310 differences between grassland and timber plantations (Fig. 4a), but between topsoils of timber
311 plantations, native forests and crops (Fig. 5a). Organic matter and nitrogen decrease in topsoils
312 after grassland afforestation (Berthrong et al., 2009) and C/N ratio increases with plantation
313 age and decreases with precipitation (Berthrong et al., 2012). The topsoils of timber plantations
314 have, on average, lower N contents compared to other land uses (Table A2) and data in
315 literature (Jobbagy and Jackson 2003).



316 Although in cropland, nutrients are regularly compensated for by increased application of
317 fertilizers (Beretta-Blanco et al., 2019), timber plantations are usually fertilized only in the first
318 year of planting (e.g., Binkley et al., 2017; Sandoval-Lopez et al., 2020). The CEC and the
319 average ratio between $K^+(Ca^{2+} + Mg^{2+})^{-1/2}$ in crops and grasslands are in the ranges reported by
320 Beretta-Blanco et al. (2019) with lower availability of potassium for crops. A shortage of
321 potassium and calcium is also most likely for future timber plantations after harvest, especially
322 if the logging process do not include bark stripping on site (Hernández et al., 2009) and due to
323 short time spans between harvest and new planting in the same area (Sandoval-Lopez et al.,
324 2020), but see contrasting results of McMahan et al. (2019). The extraction of nutrients and
325 biomass due to grazing (Fernandez et al., 2017), timber (Hernández et al., 2009 and 2016) and
326 crop production (Beretta-Blanco et al., 2019) lowers organic matter content and exchangeable
327 cations, affects the physical, chemical and biological soil properties, and drives degradation.

328 **4.2 Acidification in Uruguayan topsoils across land uses**

329 A further dimension of the soil degradation directly linked to the cation extraction is the
330 acidification of soils. This has been demonstrated for crops and pastures (Beretta-Blanco et al.,
331 2019), and a study site with *Eucalyptus* (Céspedes-Payret et al., 2012) and is now broadly
332 supported by our topsoil samples originating from a wide range of different land uses across
333 Uruguay. The pH values of our topsoil samples are mainly in the category of very strongly to
334 extremely acidic with lowest values for timber plantations (Fig. 6), which are below the means
335 reported so far (Jobbagy and Jackson, 2003; Céspedes-Payret et al., 2012). Moreover, as our
336 data on topsoil pH fall short compared to the values estimated by the Food and Agriculture
337 Organization of the United Nations and the Intergovernmental Technical Panel on Soils (Caon
338 and Vargas, 2017), acidification and the deterioration of topsoil quality continues.
339 Acidification results from intensified land uses with nitrogen fertilization, with biological N
340 fixation by the legumes both used in the so-called “improved pastures” (Modernel et al., 2016),
341 or with cation extraction by crop or timber harvest (Jobbagy and Jackson, 2003). So far,



342 although forest soils tend to be more acidic than agricultural soils due to acid-neutralizing
343 treatments in the latter (Baize and van Oort, 2014), we found no differences between topsoils
344 of crops and native forests (Fig. 6d) as the high organic matter content in native forests buffers
345 the process to a certain extent.

346 **4.3 Riverine Forest soils as sink for trace metals**

347 In general, the average concentration of Zn, Cu, Cd, Cr, As and Pb in our topsoils were within
348 the expected ranges of samples from crops, pastures and grassland in the region (Lavado et al.,
349 1998, 2004; Roca, 2015). Some samples, especially from orchards and crops, exceeded the
350 background values of copper, cadmium and arsenic (Table A2; Roca, 2015). Little is known
351 about the pedo-geochemical background of the Pampean soils (Roca, 2015): for example, high
352 arsenic content in Uruguayan ground waters has been hypothesized to be due to quaternary ash
353 deposits (Wu et al., 2021). In addition, the lateral heterogeneity of Pampean soils over short
354 distances makes separating geochemical and anthropic signatures difficult (Roca, 2015).
355 However, that the main risk of soil contamination in the region is from the application of
356 fertilizers and agrochemicals is uncontested (Kaushal et al., 2018).

357 To our knowledge, there has been no regional study of trace metals in the native riverine forests
358 or timber plantations. Our work thus expands the evidence base for these land uses. The topsoils
359 of riverine forests accumulate trace metals compared to those of timber plantations and crops
360 (Fig. 4l-o). The higher amount of soil organic matter in riverine forests favours the retention of
361 cations, including trace metals. Although the origin of potentially harmful elements in forest
362 soils have been primarily attributed to atmospheric deposition (Baize and van Oort, 2014),
363 atmospheric deposition only plays a major role in vicinity of urban or industrial development,
364 and our data from rural sites suggests a different entry path from the surrounding land uses to
365 the riverine forests. High levels of acidification and low amounts of organic matter reduce the
366 retention of trace metals in the soil of timber plantations, and elements leach out of the soil
367 towards the water table (Baize and van Oort, 2014). The acidification strongly contributes to



368 the overall mobility of base cations into the ‘chemical cocktail of the Anthropocene’ (Kaushal
369 et al., 2018), including trace metals. We thus observe positive feedback in already
370 impoverished soils with high acidity favouring cations solubility, in addition to the uptake by
371 trees intensifies this effect. Timber plantations extract trace metals from soil and also
372 accumulate it in the bark or leaves, so they have been used for phytoremediation (Li et al.,
373 2020). This may explain the higher values of cadmium in grassland compared to timber
374 plantations (Fig. 4m). Differences between *Eucalyptus* and *Pinus* stands may be related to
375 different age classes, as the later may have extracted more lead and chromium from the soil
376 due to their older stand age with rotation periods of about 20 years (Li et al., 2020).

377 **4.4 Carbon storage in topsoils of *Eucalyptus* plantations?**

378 Our study provides evidence that the loss of soil organic matter limits not only the productivity
379 of the crops, but also potential carbon sequestration in the region (Beretta-Blanco et al., 2019).
380 Grassland conversion to cropland decreases soil organic carbon storage compared to adjacent
381 native grasslands (Hernandez-Ramirez et al., 2021).
382 Afforestation of croplands has been also discussed as a carbon sequestration measure to
383 proactively address and effectively mitigate ongoing climate change within a person's lifetime
384 (Hernandez-Ramirez et al., 2021; Mayer et al. 2020). Although carbon stocks of four
385 *Eucalyptus* stands in the Brazilian Cerrados increased but did not change in four *Eucalyptus*
386 stands within the Atlantic Forest (McMahon et al., 2019), this may not be the case for short
387 rotation of *Eucalyptus* plantation on Campos grasslands. In our topsoil samples originating
388 from 28 different stands across Uruguay, organic matter is lowest in topsoils of timber
389 plantations (Fig. 4b). Similar amounts have been reported for a timber plantation in the East of
390 Uruguay (Céspedes-Payret et al., 2012), and in North-Eastern Argentine (Sandoval-Lopez et
391 al., 2020). Our data therefore provide clear evidence that rather than contributing to carbon
392 sequestration in the topsoil, the carbon release from the transformation of native grasslands to



393 plantation with these fast-growing species has several adverse effects depending on
394 precipitation and soil type (reviewed by Mayer et al., 2020).

395 Several trade-offs between carbon sequestration through afforestation and local water yield and
396 soil fertility have been demonstrated, including nutrient and soil organic matter depletion,
397 acidification, and biodiversity loss and corresponding challenges for landscape conservation
398 (e.g., Jackson et al., 2005; Veldman et al., 2015; Friggens et al., 2020). Soil carbon changes
399 dynamically during the first decade of afforestation. Remaining grassland carbon declines,
400 while tree carbon gain starts (e.g. Paul et al., 2002). Although a net gain of carbon is expected
401 when the new forests approached equilibrium after decades (Hernández-Ramirez et al., 2021),
402 in contrast to long-lasting forests, *Eucalyptus* harvest in Uruguay takes place after less than a
403 decade (appr. 7-10 years). Soil organic matter does not differ between *Eucalyptus* plantations
404 and neighbouring grasslands (Appendix A; Table A1 and A2). Another study near our sites 12
405 and 13 also did not find a significant difference between the soil organic carbon of the upper
406 soil (0-30cm) compared to afforested versus native grasslands (Hernández et al., 2016). A study
407 near our sites 7, 8, 10 reported higher top soil carbon of grassland compared with afforested
408 sites in their vicinity only if sand content is lower than sixty percent (Sandoval-Lopez et al.,
409 2020). McMahon et al. (2019) identified a greater carbon gain under *Eucalyptus* stands
410 compared to (potential) carbon losses in neighbouring degraded Cerrado grasslands. There is
411 evidence that the retention of residues after harvest increases the carbon stock in soil (Mayer
412 et al., 2020). Long term monitoring of carbon stocks is needed to verify the influences of
413 management and environmental changes. A regional study on *Eucalyptus* plantations across
414 different biomes in Brazil shows both decreases and no changes depending on precipitation in
415 the dry season, clay content and on the initial stock of carbon in the soil (Cook et al., 2016)
416 The simplistic solution of huge tree plantations to compensate anthropogenic CO₂ emissions
417 has been challenged in the last decade, and some crucial lessons learnt have been summarized
418 (Di Sacco et al., 2020). Since grasslands are more sustainable carbon sinks compared to



419 climate-vulnerable monocultures such as timber plantations (Dass et al. 2018), avoiding
420 afforestation of previously non-forested lands is important. This is the case for *Eucalyptus*
421 afforestation in the originally forestless grasslands of Uruguay: models on carbon sequestration
422 and dynamics in Mollisols and Oxisols under South American grasslands estimated a higher
423 carbon retention efficiency under grasslands compared to afforested sites, suggesting that
424 silvopastoral systems are a potential solution for soil carbon sequestration in tropical soils
425 (Berhongaray and Alvarez, 2019).

426 Connecting or expanding existing forest and using native species for plantings (Di Sacco et al.,
427 2020) is also recommended (see also Alvaro et al. 2020, Ortiz et al. 2020). Our topsoil data
428 indicates that carbon sequestration occurs mainly in the topsoils of native riverine forests that
429 cover less than five percent of the Uruguayan territory. Consequently, the expansion of native
430 forests and the use of native species in forestry project for long term establishment can reduce
431 adverse effects of timber plantations. While there is preliminary evidence that N-fixing tree
432 species can help increase local C stocks of afforestation, because of their potential for invasion,
433 exotic N-fixers should be avoided (Mayer et al. 2020).

434 **4.5. Back to more conservative grassland management?**

435 Our soil survey data shows strong soil degradation of Uruguayan black soils from erosion,
436 acidification and contamination, and indicates a translocation of nutrients and organic matter
437 across the landscape from grassland, timber and crop plantations to the riverine forests. The
438 potential of grasslands as cropland reserve have been largely overestimated (Lambin et al.,
439 2013), and they have already degraded during the last decades by inappropriate land
440 management techniques (Jaurena et al., 2021, De Faccio et al., 2021), lack of mainstreaming
441 soil conservative techniques (García-Préchac, 2004; Fernandez et al., 2017), decoupling of crop
442 and livestock (De Faccio et al, 2021) plus climate change impacts with storm water events and
443 drought, all of which trigger soil erosion (Wingeyer et al. 2015). From the although very limited
444 point of view on topsoils, the concept of conserving ‘old growth grasslands’ with extensive use



445 (Veldman et al. 2015) appears a more promising strategy to put the ‘grasslands at the core’ in
446 the Campos region than the use intensification strategies envisioned by Jaurena et al. (2021).

447 **Data availability**

448 All data generated or analysed in this study are included in this article (and its supplementary
449 materials). Further data are available from the corresponding authors upon reasonable request.

450 **Author contributions**

451 IS, LRR, ST and MB conceptualized the study, performed the soil sampling and data analysis.
452 EZ advised the laboratory analyses. IS wrote the initial draft. LRR, ST, MB and EZ contributed
453 to generating and reviewing the subsequent versions of the manuscript. IS received the funding
454 for the study.

455 **Competing interests**

456 The authors declare that they have no conflict of interest

457 **Acknowledgement**

458 In alphabetical order, we thank Juan Barreneche, Lucia Gaucher, Sören Miehe, Nicolas Silvera
459 and Matias Zarucki for their assistance in field work. We thank Manuel Garcia and Meica
460 Valdivia for the pre-processing of samples. We thank the staff of the soil lab from the
461 Department of Soil and Natural Resources, University of Concepción, Chile. We thank Diego



462 de Panis for help with statistical analysis and visualization and valuable comments on a
463 previous version of this manuscript. Thanks to Vera Krause, Serafina Bischoff, Sophia Reitzug,
464 Rhea Rennert and Diego Nicolas Rojas for support with data analysis and plots and maps
465 visualization. We also thank all landowners for access permission to establish our monitoring
466 sites on their land, their hospitality and willingness to discuss land use goals concerning all
467 dimension of sustainability. The study was funded by the German Federal Ministry of
468 Education and Research (BMBF; 01LN1305A). Special thanks go to Amal Chatterjee for
469 improving our English.

470 **References**

- 471 Alfaro, M., Dube, F. and Zagal, E.: The Influence of Overmature, Degraded Nothofagus
472 Forests with Strong Anthropogenic Disturbance on the Quality of an Andisol and its Gradual
473 Recovery with Silvopasture in Southwestern South America, in: *Agroforestry for
474 Degraded Landscapes*, edited by: Dagar, J.C., Gupta, S.R., Teketay, D. Springer,
475 Singapore. https://doi.org/10.1007/978-981-15-6807-7_3, 2020.
- 476 Baize, D. and van Oort, F.: Potentially Harmful Elements in Forest Soils, in: *PHEs,
477 Environment and Human Health*, edited by: Bini, C. and Bech, J., Springer Netherlands,
478 Dordrecht, 151–198, https://doi.org/10.1007/978-94-017-8965-3_4, 2014.
- 479 Beretta-Blanco A, Pérez O, Carrasco-Letelier L. Soil quality decrease over 13 years of
480 agricultural production. *Nutr Cycling Agroecosyst.* 2019; 114:45-55.
- 481 Beretta-Blanco, A.; Carrasco-Letelier, L. Relevant factors in the eutrophication of the Uruguay
482 River and the Río Negro. *Sci. Total Environ.* 2021, 761,143299; DOI:
483 10.1016/j.scitotenv.2020.143299
- 484 Berhongaray, G; Alvarez R. Soil carbon sequestration of Mollisols and Oxisols under grassland
485 and tree plantations in South America - A review. *Geoderma Regional* 18 (2019)
486 e00226.
- 487 Berthrong, S. T., Jobbágy, E. G., and Jackson, R. B.: A global meta-analysis of soil
488 exchangeable cations, pH, carbon, and nitrogen with afforestation, *Ecological
489 Applications*, 19, 2228–2241, <https://doi.org/10.1890/08-1730.1>, 2009.



- 490 Berthrong, S. T., Piñeiro, G., Jobbágy, E. G., and Jackson, R. B.: Soil C and N changes with
491 afforestation of grasslands across gradients of precipitation and plantation age, 22, 76–
492 86, <https://doi.org/10.1890/10-2210.1>, 2012.
- 493 Binkley, Dan et al. The interactions of climate, spacing and genetics on clonal *Eucalyptus*
494 plantations across Brazil and Uruguay. *Forest Ecology And Management*. Amsterdam:
495 Elsevier Science Bv, v. 405, p. 271-283, doi.org/10.1016/j.foreco.2017.09.050
- 496 Bonfante, A., Basile, A., and Bouma, J.: Targeting the soil quality and soil health concepts
497 when aiming for the United Nations Sustainable Development Goals and the EU Green
498 Deal, *SOIL*, 6, 453–466, <https://doi.org/10.5194/soil-6-453-2020>, 2020.
- 499 Borrelli, P., Robinson, D. A., Fleischer, L. R., Lugato, E., Ballabio, C., Alewell, C.,
500 Meusburger, K., Modugno, S., Schütt, B., Ferro, V., Bagarello, V., Oost, K. V.,
501 Montanarella, L., and Panagos, P.: An assessment of the global impact of 21st century
502 land use change on soil erosion, *Nat Commun*, 8, 2013, <https://doi.org/10.1038/s41467-017-02142-7>, 2017.
- 504 Caon, L., & Vargas, R. (2017). Threats to soils: Global trends and perspectives. A Contribution
505 from the Intergovernmental Technical Panel on Soils, Global Soil Partnership Food and
506 Agriculture Organization of the United Nations. In G. Pierzynski & Brajendra (Eds.),
507 *Global land outlook working paper*, 28 pp.
- 508 Carrasco-Letelier, L. and A. Beretta-Blanco. 2017. Soil erosion by water estimated for 99
509 Uruguayan basins. *Cien. Inv. Agr.* 44(2): 184-194.
- 510 Céspedes-Payret et al. 2012 *Science of the Total Environment* 438 (2012) 549–557.
511 <http://dx.doi.org/10.1016/j.scitotenv.2012.08.075>
- 512 Chenu, C., Rumpel, C., and Lehmann, J.: Methods for studying soil organic matter: Nature,
513 dynamics, spatial accessibility, and Interactions with minerals, in: *Soil Microbiology,*
514 *Ecology and Biochemistry*, Academic Press, np, <https://doi.org/10.1016/B978-0-12-415955-6.00013-X>, 2015.
- 516 Cook, R. L., Binkley, D., and Stape, J. L.: Eucalyptus plantation effects on soil carbon after
517 20years and three rotations in Brazil, *Forest Ecology and Management*, 359, 92–98,
518 <https://doi.org/10.1016/j.foreco.2015.09.035>, 2016.
- 519 Dass, P. et al 2018 Grasslands may be more reliable carbon sinks than forests in California.
520 *Environ. Res. Lett.* 13 074027. <https://doi.org/10.1088/1748-9326/aac339>
- 521 De Faccio Carvalho, P. C., Savian, J. V., Chiesa, T. D., Souza Filho, W. D., Terra, J. A., Pinto,
522 P., Martins, A. P., Villarino, S., Da Trindade, J. K., De Albuquerque Nunes, P. A., and



- 523 Piñeiro, G.: Land-use intensification trends in the Rio de la Plata region of South
524 America: toward specialization or recoupling crop and livestock production, *Front.*
525 *Agr. Sci. Eng.*, 8, 97, <https://doi.org/10.15302/J-FASE-2020380>, 2021.
- 526 Di Sacco, A.; Hardwick, K.; Blakesley, D.; Brancalion, P.H.S.; Breman, E.; Cecilio Rebola,
527 L.; Chomba, S.; Dixon, K.; Elliott, S.; Ruyonga, G.; Shaw, K.; Smith, P.; Smith, R.J.;
528 Antonelli, A. Ten Golden Rules for Reforestation to Optimise Carbon Sequestration,
529 Biodiversity Recovery and Livelihood Benefits. <https://doi.org/10.1111/gcb.15498>.
530 2020
- 531 Durán, A., Morrás, H., Studdert, G., and Liu, X.: Distribution, properties, land use and
532 management of Mollisols in South America, *Chin. Geogr. Sci.*, 21, 511,
533 <https://doi.org/10.1007/s11769-011-0491-z>, 2011.
- 534 ESRI 2018. ArcGIS Desktop: Release 10.3. Redlands, CA: Environmental Systems Research
535 Institute.
536 https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051232.pdf
- 537 FAO 2019. Land Use. Published online at OurWorldInData.org. Retrieved from:
538 '<https://ourworldindata.org/land-use>' [Online Resource]
- 539 Fernández, R., Frasier, I., Noellemeyer, E., and Quiroga, A.: Soil quality and productivity
540 under zero tillage and grazing on Mollisols in Argentina – A long-term study,
541 *Geoderma Regional*, 11, 44–52, <https://doi.org/10.1016/j.geodrs.2017.09.002>, 2017.
- 542 Fox, J., Weisberg, S. (2019) *An R Companion to Applied Regression*, Third Edition, Sage.
- 543 Friggens, N. L., Hester, A. J., Mitchell, R. J., Parker, T. C., Subke, J., and Wokey, P. A.: Tree
544 planting in organic soils does not result in net carbon sequestration on decadal
545 timescales, *Glob Change Biol*, 26, 5178–5188, <https://doi.org/10.1111/gcb.15229>,
546 2020.
- 547 García-Préchac, F., Ernst, O., Siri-Prieto, G., and Terra, J. A.: Integrating no-till into crop–
548 pasture rotations in Uruguay, *Soil and Tillage Research*, 77, 1–13,
549 <https://doi.org/10.1016/j.still.2003.12.002>, 2004.
- 550 Gomiero, Tiziano. 2016. "Soil Degradation, Land Scarcity and Food Security: Reviewing a
551 Complex Challenge" *Sustainability* 8, no. 3: 281. <https://doi.org/10.3390/su8030281>
- 552 Hernández J, del Pino A, Salvo L, Arrarte G. Nutrient export and harvest residue
553 decomposition patterns of a *Eucalyptus dunnii* Maiden plantation in temperate climate



- 554 of Uruguay. *Forest Ecology Management* 2009; 258(2): 92-99.
555 10.1016/j.foreco.2009.03.050
- 556 Hernández, J., del Pino, A., Vance, E. D., Califra, Á., Del Giorgio, F., Martínez, L., and
557 González-Barrios, P.: *Eucalyptus* and *Pinus* stand density effects on soil carbon
558 sequestration, *Forest Ecology and Management*, 368, 28–38,
559 <https://doi.org/10.1016/j.foreco.2016.03.007>, 2016.
- 560 Hernandez-Ramirez, G., Sauer, T. J., Chendev, Y. G., and Gennadiev, A. N.: Nonlinear
561 turnover rates of soil carbon following cultivation of native grasslands and subsequent
562 afforestation of croplands, *SOIL*, 7, 415–431, <https://doi.org/10.5194/soil-7-415-2021>,
563 2021.
- 564 IUSS Working Group WRB. 2015. World Reference Base for Soil Resources 2014, update
565 2015 International soil classification system for naming soils and creating legends for
566 soil maps. World Soil Resources Reports No. 106. FAO, Rome.
567 <https://www.fao.org/3/i3794en/i3794en.pdf>
- 568 Jackson et al 2005 Trading Water for Carbon with Biological Carbon Sequestration. *Science*
569 310, 1944 (2005); DOI: 10.1126/science.1119282
- 570 Jackson, R.: Soil nutrient stocks are maintained over multiple rotations in Brazilian Eucalyptus
571 plantations | Elsevier Enhanced Reader, 13, n.d.
- 572 Jaurena M, Durante M, Devincenzi T, Savian JV, Bendersky D, Moojen FG, Pereira M, Soca
573 P, Quadros FLF, Pizzio R, Nabinger C, Carvalho PCF and Lattanzi FA (2021) Native
574 Grasslands at the Core: A New Paradigm of Intensification for the Campos of Southern
575 South America to Increase Economic and Environmental Sustainability. *Front. Sustain.*
576 *Food Syst.* 5:547834. doi: 10.3389/fsufs.2021.547834
- 577 Jobbagy, E. G., and R. B. Jackson. 2003. Patterns and mechanisms of soil acidification in the
578 conversion of grasslands to forests. *Biogeochemistry* 64:205–229.
- 579 Jobbagy, E. G., and R. B. Jackson. 2004. The uplift of soil nutrients by plants: biogeochemical
580 consequences across scales. *Ecology* 85:2380–2389.
- 581 Kaushal, S. S., Gold, A. J., Bernal, S., Johnson, T. A. N., Addy, K., Burgin, A., Burns, D. A.,
582 Coble, A. A., Hood, E., Lu, Y., Mayer, P., Minor, E. C., Schroth, A. W., Vidon, P.,
583 Wilson, H., Xenopoulos, M. A., Doody, T., Galella, J. G., Goodling, P., Haviland, K.,
584 Haq, S., Wessel, B., Wood, K. L., Jaworski, N., and Belt, K. T.: Watershed ‘chemical
585 cocktails’: forming novel elemental combinations in Anthropocene fresh waters,
586 *Biogeochemistry*, 141, 281–305, <https://doi.org/10.1007/s10533-018-0502-6>, 2018.




- 587 Kellogg, C. E. "Soil survey division staff: soil survey manual. Chapter 3" United States
588 Department of Agriculture, Washington (1993).
- 589 Lambin, E.F.; Gibbs, H.K.; Ferreira, L.; Grau, R.; Mayaux, P.; Meyfroidt, P.; Morton, D.C.;
590 Rudel, T.K.; Gasparri, I.; Munger, J. Estimating the world's potentially available
591 cropland using a bottom-up approach. *Glob. Environ. Chang.* 2013, 23, 892–901.
- 592 Lanfranco, B. and Sapriza, G.: El índice CONEAT como medida de productividad y valor de
593 la tierra., 57, n.d.
- 594 Lavado, R.S., Rodríguez, M.B., Scheiner, S.D., Taboada, M.A., Rubio, G., Alvarez, R.,
595 Alconada, M., Zubillaga, M.S., 1998. Heavy metals in soils of Argentina: comparison
596 between urban and agricultural soils. *Communications in Soil Science and Plant
597 Analysis* 29, 1913–1917.
- 598 Lavado, R.S., Zubillaga, M.S., Alvarez, R., Taboada, M.A., 2004. Baseline levels of potentially
599 toxic elements in pampas soils. *Soil and Sediment Contamination: An International
600 Journal* 13, 329–339.
- 601 Li, H., Jiang, L., You, C., Tan, B., and Yang, W.: Dynamics of heavy metal uptake and soil
602 heavy metal stocks across a series of Masson pine plantations, *Journal of Cleaner
603 Production*, 269, 122395, <https://doi.org/10.1016/j.jclepro.2020.122395>, 2020.
- 604 Liu, X., Lee Burras, C., Kravchenko, Y. S., Durán, A., Huffman, T., Morras, H., Studdert, G.,
605 Zhang, X., Cruse, R. M., and Yuan, X.: Overview of Mollisols in the world:
606 Distribution, land use and management, *Can. J. Soil. Sci.*, 92, 383–402,
607 <https://doi.org/10.4141/cjss2010-058>, 2012.
- 608 Mayer M., Prescott C.E., Abaker W.E.A., Augusto L., Cécillon L., Ferreira G.W.D., James J.,
609 Jandl R., Katzensteiner K., Laclau J.-P., Laganière J., Nouvellon Y., Paré D., Stanturf
610 J.A., Vanguelova E.I., Vesterdal L. Influence of forest management activities on soil
611 organic carbon stocks: A knowledge synthesis. *Forest Ecology and Management*,
612 Volume 466, 2020
- 613 McMahon, D. E., Vergütz, L., Valadares, S. V., Silva, I. R. da, and Jackson, R. B.: Soil nutrient
614 stocks are maintained over multiple rotations in Brazilian Eucalyptus plantations,
615 *Forest Ecology and Management*, 448, 364–375,
616 <https://doi.org/10.1016/j.foreco.2019.06.027>, 2019.
- 617 Merino, A., Balboa, M. A., Rodríguez Soalleiro, R., and González, J. G. Á.: Nutrient exports
618 under different harvesting regimes in fast-growing forest plantations in southern
619 Europe, *Forest Ecology and Management*, 207, 325–339,
620 <https://doi.org/10.1016/j.foreco.2004.10.074>, 2005.



- 621 Modernel, P., Rossing, W. A. H., Corbeels, M., Dogliotti, S., Picasso, V., and Tittonell, P.:
622 Land use change and ecosystem service provision in Pampas and Campos grasslands
623 of southern South America, *Environ. Res. Lett.*, 11, 113002,
624 <https://doi.org/10.1088/1748-9326/11/11/113002>, 2016.
- 625 OAS (Organization of American States, Department of Sustainable Development) 1994.
626 Uruguay - Proyecto Regional de Alternativas para la Inversión Forestal. Available
627 online: <http://www.oas.org/dsd/publications/unit/oea20s/oea20s.pdf>.
- 628 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.
629 R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and
630 Wagner, H.: *vegan: Community Ecology Package*, 2020.
- 631 Ortiz, J., Dube, F., Neira, P., Panichini, M., Stolpe, N.B., Zagal, E. and Martínez-Hernández,
632 P.A. Soil Quality Changes within a (*Nothofagus obliqua*) Forest Under Silvopastoral
633 Management in the Andes Mountain Range, South Central Chile. *Sustainability*;
634 12(17):6815. <https://doi.org/10.3390/su12176815>, 2020.
- 635 Paul, K. I., Polglase, P. J., Nyakuengama, J. G., and Khanna, P. K.: Change in soil carbon
636 following afforestation, *Forest Ecology and Management*, 168, 241–257,
637 [https://doi.org/10.1016/S0378-1127\(01\)00740-X](https://doi.org/10.1016/S0378-1127(01)00740-X), 2002.
- 638 Piñeiro, D. E.: Land grabbing: concentration and “foreignisation” of land in Uruguay, *Canadian*
639 *Journal of Development Studies / Revue canadienne d'études du développement*, 33,
640 471–489, <https://doi.org/10.1080/02255189.2012.746216>, 2012.
- 641 R Core Team: *R: A Language and Environment for Statistical Computing*, R Foundation for
642 Statistical Computing, Vienna, Austria, available at: <https://www.R-project.org/>, last
643 access: 2 March 2021.
- 644 Richter, D. D.: Game Changer in Soil Science. The Anthropocene in soil science and
645 pedology., *J. Plant Nutr. Soil Sci.*, 183, 5–11, <https://doi.org/10.1002/jpln.201900320>,
646 2020.
- 647 Roca, N.: Heavy metal background levels in rural soils: a case study in Pampean soils
648 (Argentina), 10, 2015.
- 649 Sandoval López, D.M., Arturi, M.F., Goya, J.F. et al. *Eucalyptus grandis* plantations: effects
650 of management on soil carbon, nutrient contents and yields. *J. For. Res.* 31, 601–611
651 (2020). <https://doi.org/10.1007/s11676-018-0850-z>




- 652 Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and
653 interpreting soil surveys. 2nd edition.
654 https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051232.pdf
- 655 Veldman, J. W., Overbeck, G. E., Negreiros, D., Mahy, G., Le Stradic, S., Fernandes, G. W.,
656 Durigan, G., Buisson, E., Putz, F. E., and Bond, W. J.: Where Tree Planting and Forest
657 Expansion are Bad for Biodiversity and Ecosystem Services, *BioScience*, 65, 1011–
658 1018, <https://doi.org/10.1093/biosci/biv118>, 2015.
- 659 Waters, C. N., Zalasiewicz, J., Summerhayes, C., Barnosky, A. D., Poirier, C., Ga uszka, A.,
660 Cearreta, A., Edgeworth, M., Ellis, E. C., Ellis, M., Jeandel, C., Leinfelder, R., McNeill,
661 J. R., Richter, D. d., Steffen, W., Syvitski, J., Vidas, D., Wagreich, M., Williams, M.,
662 Zhisheng, A., Grinevald, J., Odada, E., Oreskes, N., and Wolfe, A. P.: The
663 Anthropocene is functionally and stratigraphically distinct from the Holocene, *Science*,
664 351, aad2622–aad2622, <https://doi.org/10.1126/science.aad2622>, 2016.
- 665 Wingeyer, A.B.; Amado, T.J.C.; Pérez-Bidegain, M.; Studdert, G.A.; Perdomo Varela, C.H.;
666 Garcia, F.O.; Karlen, D.L. Soil Quality Impacts of Current South American
667 Agricultural Practices. *Sustainability* 2015, 7, 2213–2242.
- 668 Wu, R., Alvareda, E., Polya, D., Blanco, G., and Gamazo, P.: Distribution of Groundwater
669 Arsenic in Uruguay Using Hybrid Machine Learning and Expert System Approaches,
670 *Water*, 13, 527, <https://doi.org/10.3390/w13040527>, 2021.
- 671 Zurbriggen, C., González-Lago, M., Baraibar, M., Baethgen, W., Mazzeo, N. and Sierra, M.,
672 2020. Experimentation in the Design of Public Policies: The Uruguayan Soils
673 Conservation Plans. *Iberoamericana – Nordic Journal of Latin American and*
674 *Caribbean Studies*, 49(1), pp.52–62. DOI: <http://doi.org/10.16993/iberoamericana.459>
675



676
 677
 678
 679
 680
 681
 682


Figure 1: Study design and sampling. (a) Location of the 28 monitoring sites across Uruguay including land use types sampled (grassland (b); timber plantations, native forest and agricultural land. Proportion of plots with particular category of soil classification according to the World Reference Base (WRB, c), Soil Taxonomy (d) and CONEAT (Comisión Nacional de Estudio Agronómico de la Tierra, e).



683

684 **Figure 2.** Spearman's rank correlations for parameters of topsoils (n=80) regarding (a) fertility
 685 proxies, (b) acidity and (c) trace metals. Colour intensity and the size of the circle are
 686 proportional to the correlation coefficients (ρ). Empty slots show correlations with $p > 0.05$.

687






689 **Figure 3.** Non-metric multidimensional scaling showing significant clustering differences
690 among samples from (a) grassland (GL), timber plantations (TP) and native forests (NF); (b)
691 among samples from *Pinus* and *Eucalyptus* plantations; (c) among land uses (including
692 agriculture AC) in Argiudolls; (d) among land uses in Argiudolls & Hapluderts; (e) among land
693 uses in Argiudolls, Hapudolls & Hapludalf and (f) among land uses in Hapludalfs &
694 Hapludults.
695




696



697




698 **Figure 4.** Violin box plots for significant Kruskal-Wallis Tests across evaluated land uses (i.e.
699 grassland (GL), timber plantation (TP) and native forest (NF)) for each variable. Significance
700 in posterior Wilcoxon pairwise comparisons is depicted as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$
701 (***), $p < 0.0001$ (****).
702





704 **Figure 5.** Violin box plots for significant Kruskal-Wallis Tests in ‘Argiudolls’ Soil Taxonomy
 705 category for fertility variables across available land uses (GL: Grassland, TP: Timber
 706 plantation, NF: Native forest, AC: Agriculture). Significance in posterior Wilcoxon pairwise
 707 comparisons is depicted as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), $p < 0.0001$ (****).



708 **Figure 6.** Acidity of all topsoils samples (a) and (b) according to different land use (GL:
 709 Grassland, TP: Timber plantation, NF: Native forest, AC: Agriculture). Proportion of samples
 710 with a given acidic category (Kellogg, 1993). pH of topsoil given as violin box plots for
 711



712 significant Kruskal-Wallis Tests across different land uses independent from soil Order (c) and
713 separated per soil Orders (d-f). Sample number (n) is given. Significance in posterior Wilcoxon
714 pairwise comparisons is depicted as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), $p < 0.0001$ (****).
715



716 **Table 1.** General characteristics of Uruguayan topsoils: descriptive statistics for the parameters
 717 of all single samples (n) across different land uses and classification to different soil types (for
 718 details on different land uses i.e. grassland, timber plantations, native forests and crops see
 719 Appendix A1-A2 and on different soil types see Appendix A3-4).

Variable	n	Mean	S.D.	Minimum	Maximum
Humidity (%)	280	18.9	10.8	1.34	51.5
N _{total} (%)	280	0.2	0.14	0.04	1.2
C _{total} (%)	280	3.0	2.3	0.4	25.7
C/N ratio	279	13.3	1.8	9.1	20.6
SOM (%)	267	5.6	3.0	0.8	16.4
P (mg kg ⁻¹)	278	295	166	43	1008
Ca (cmol kg ⁻¹)	82	9.4	8.9	0.3	42.7
Mg (cmol kg ⁻¹)	82	2.5	2.2	0.1	9.7
K (cmol kg ⁻¹)	82	0.4	0.25	0.03	1.44
Na (cmol kg ⁻¹)	82	0.1	0.14	0	0.69
Zn (mg kg ⁻¹)	82	1.9	2.1	0.1	9.7
Cu (mg kg ⁻¹)	82	5.4	22.1	0.3	161.2
Fe (mg kg ⁻¹)	82	134	75	5	309
Mn (mg kg ⁻¹)	81	25	17	0.7	93
pH	279	4.8	0.8	3.6	7.34
As (mg kg ⁻¹)	279	3.9	3.6	0.6	30.7
Cd (mg kg ⁻¹)	274	0.4	0.2	0.2	1
Cr (mg kg ⁻¹)	280	13.6	12.6	2	78.9
Pb (mg kg ⁻¹)	280	7	3.4	2	27.6
CEC (cmol kg ⁻¹)	82	12.4	10.7	0.5	50.1
K ⁺ /(Ca ²⁺ +Mg ²⁺)	82	0.07	0.08	0.01	0.43
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	82	0.07	0.07	0.01	0.42

720

721



722

Appendices

723 Appendix A

724 **Table A1.** Descriptive statistics of topsoil variables for all single soil types at grassland and
 725 native forests. Number of samples (n), mean, standard deviation (SD), minimum (Min) and
 726 maximum (Max) of each variable are given.

Variable (unit)	n	Grassland				Native forests				
		Mean	SD	Min	Max	n	Mean	SD	Min	Max
Humidity (%)	115	20.4	11.5	1.5	51.5	57	25.8	9.3	6.3	50.3
N _{total} (%)	115	0.22	0.13	0.06	0.80	57	0.30	0.19	0.04	1.20
C _{total} (%)	115	2.9	1.9	0.7	10.8	57	4.2	3.6	0.5	25.7
C/N ratio	115	13.2	1.7	9.1	19.4	56	13.5	1.4	9.8	18.5
SOM (%)	109	5.3	2.6	1.4	15.2	53	7.5	3.8	1.1	16.4
P (mg/kg)	114	272	124	43	749	57	484	194	56	1009
Ca (cmol kg ⁻¹)	32	9.5	10.0	0.3	42.7	18	15.1	10.0	0.3	37.3
Mg (cmol kg ⁻¹)	32	2.2	2.1	0.18	9.7	18	4.8	2.6	0.12	8.9
K (cmol kg ⁻¹)	32	0.40	0.26	0.08	1.44	18	0.50	0.28	0.03	1.29
Na (cmol kg ⁻¹)	32	0.12	0.15	0.00	0.65	18	0.14	0.12	0.00	0.50
Zn (mg kg ⁻¹)	32	1.4	1.2	0.10	4.7	18	4.0	2.7	0.80	9.60
Cu (mg kg ⁻¹)	32	2.2	1.8	0.30	7.7	18	2.4	1.5	0.30	5.50
Fe (mg kg ⁻¹)	32	121	81	5	279	18	149	76	10	309
Mn (mg kg ⁻¹)	32	22	18	2	93	17	33	18	1	72
pH	114	4.9	0.9	3.6	7.3	57	5.1	0.7	3.6	7.2
As (mg kg ⁻¹)	115	4.3	4.5	0.6	30.7	56	4.1	2.5	1.2	13.6
Cd (mg kg ⁻¹)	111	0.41	0.17	0.17	1.00	57	0.45	0.15	0.17	0.81
Cr (mg kg ⁻¹)	115	12.3	9.2	2.6	49.5	57	22.8	17.3	2.0	70.7
Pb (mg kg ⁻¹)	115	6.8	3.0	2.3	15.9	57	9.5	4.8	2.9	27.7
CEC (cmol/ kg ⁻¹)	32	12.2	11.5	0.8	50.1	18	20.4	11.4	0.5	44.6
K ⁺ /(Ca ²⁺ +Mg ²⁺)	32	0.08	0.10	0.01	0.43	18	0.03	0.02	0.01	0.07
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	32	0.08	0.10	0.01	0.42	18	0.03	0.02	0.01	0.07

727



728 **Table A2.** Descriptive statistics for of topsoil variables for all single soil types at timber
 729 plantations and crops. Number of samples (n), mean, standard deviation (SD), minimum (Min)
 730 and maximum (Max) of each variable are given.

Variable (unit)	Timber plantations					Crops				
	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max
Humidity (%)	93	13.2	7.0	1.7	38.0	15	18.1	12.0	1.3	50.5
N _{total} (%)	93	0.18	0.10	0.04	0.61	15	0.22	0.06	0.13	0.29
C _{total} (%)	93	2.5	1.5	0.4	8.4	15	2.6	0.7	1.4	3.3
C/N ratio	93	13.6	2.1	10.0	20.6	15	11.7	0.7	10.5	12.8
SOM (%)	90	4.9	2.8	0.8	16.1	15	5.2	1.3	2.8	6.7
P (mg kg ⁻¹)	93	206	91	56	551	14	310	120	142	613
Ca (cmol kg ⁻¹)	27	5.2	4.8	0.6	18	5	10.8	5.5	7.2	20.4
Mg (cmol kg ⁻¹)	27	1.4	1.0	0.16	4.1	5	2.5	1.0	1.76	4.1
K (cmol kg ⁻¹)	27	0.32	0.21	0.09	0.93	5	0.46	0.23	0.29	0.85
Na (cmol kg ⁻¹)	27	0.11	0.16	0.00	0.69	5	0.06	0.02	0.03	0.08
Zn (mg kg ⁻¹)	27	0.94	0.74	0.10	2.8	5	3.7	3.7	0.60	9.7
Cu (mg kg ⁻¹)	27	1.5	0.9	0.5	4.5	5	57.7	78.8	0.8	161.0
Fe (mg kg ⁻¹)	27	143	69	16	290	5	115	62	63	210
Mn (mg kg ⁻¹)	27	25	16	6	72	5	17	5	10	23
pH	93	4.5	0.7	3.6	6.8	15	5.1	0.4	4.2	5.7
As (mg kg ⁻¹)	93	3.4	3.0	0.7	25.5	15	3.1	1.2	1.1	5.5
Cd (mg kg ⁻¹)	91	0.33	0.11	0.16	0.65	15	0.41	0.14	0.21	0.66
Cr (mg kg ⁻¹)	93	10.6	11.3	2.0	78.9	15	7.8	1.8	5.0	11.3
Pb (mg kg ⁻¹)	93	6.1	2.3	2.0	11.7	15	5.1	1.2	3.4	7.3
CEC (cmol kg ⁻¹)	27	7.0	5.6	1.0	20.6	5	13.8	6.5	9.5	25.0
K ⁺ /(Ca ²⁺ +Mg ²⁺)	27	0.08	0.07	0.02	0.25	5	0.04	0.03	0.02	0.09
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	27	0.08	0.07	0.02	0.25	5	0.04	0.03	0.02	0.08

731

732

733



734 **Table A3.** Descriptive statistics of topsoil variables for Argiudolls and for grassland and native
 735 forests within of Argiudolls. Number of single samples (n), mean, standard deviation (SD),
 736 minimum (Min) and maximum (Max) of each variable are given.

737

Variable (Unit)	Total Argiudolls					Grassland					Native forests				
	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max
Humidity (%)	77	18.3	10.9	1.3	50.5	31	19.0	9.2	5.0	49.0	12	26.4	13.4	9.0	48.2
N _{total} (%)	77	0.24	0.15	0.07	1.20	31	0.21	0.11	0.07	0.66	12	0.34	0.28	0.13	1.20
C _{total} (%)	77	3.3	3.0	1.1	25.7	31	2.9	1.8	1.1	10.8	12	5.3	6.5	1.7	25.7
C/N ratio	76	13.3	2.2	9.8	19.9	31	13.3	2.1	10.5	19.4	11	13.6	2.1	9.8	18.5
SOM (%)	73	5.8	2.5	2.1	16.1	30	5.2	1.9	2.1	10.0	9	7.4	2.1	4.8	11.0
P (mg kg ⁻¹)	76	280	122	93	693	30	234	78	93	353	12	438	155	155	693
Ca (cmol kg ⁻¹)	21	11.4	8.0	2.6	37.1	8	11.5	6.7	3.2	20.5	4	16.2	14.3	5.1	37.1
Mg (cmol kg ⁻¹)	21	2.3	1.4	0.5	6.1	8	2.0	0.8	0.9	3.2	4	3.5	2.5	1.0	6.1
K (cmol kg ⁻¹)	21	0.45	0.26	0.1	1.3	8	0.4	0.14	0.16	0.61	4	0.69	0.41	0.34	1.29
Na (cmol kg ⁻¹)	21	0.08	0.05	0.03	0.26	8	0.07	0.02	0.04	0.10	4	0.10	0.11	0.03	0.26
Zn (mg kg ⁻¹)	21	2.4	3.0	0.1	10	8	1.3	0.8	0.1	2.2	4	5.1	4.3	1.3	10
Cu (mg kg ⁻¹)	21	15.5	42.7	0.8	161	8	2.6	1.6	1.0	6.0	4	1.9	1.0	0.8	3.0
Fe (mg kg ⁻¹)	21	117	62	10	232	8	104	50	28	171	4	117	84	10	216
Mn (mg kg ⁻¹)	21	23.3	10.4	7.1	43.6	8	20.5	10.4	7.1	33.8	4	28.8	10.4	15.4	39.4
pH	77	5.0	0.7	3.8	6.6	31	4.9	0.7	3.9	6.6	12	5.2	0.5	4.3	6.1
As (mg kg ⁻¹)	76	3.2	1.4	1.0	12.2	31	2.9	0.9	1.5	4.9	11	4.0	2.8	1.5	12.2
Cd (mg kg ⁻¹)	76	0.4	0.1	0.2	0.7	30	0.4	0.1	0.2	0.7	12	0.4	0.1	0.2	0.7
Cr (mg kg ⁻¹)	77	9.3	6.2	2.2	28.3	31	8.1	4.1	3.6	18.5	12	13.2	9.5	2.2	28.3
Pb (mg kg ⁻¹)	77	5.6	2.1	2.8	10.7	31	5.2	1.8	2.8	9.6	12	6.8	2.9	2.9	10.7
CEC (cmol kg ⁻¹)	21	14.2	9.3	3.4	44.6	8	14.0	7.3	4.5	23.1	4	20.4	16.7	6.4	44.6
K ⁺ /(Ca ²⁺ +Mg ²⁺)	21	0.04	0.02	0.02	0.09	8	0.03	0.02	0.02	0.07	4	0.04	0.01	0.03	0.06
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	21	0.04	0.02	0.02	0.08	8	0.03	0.02	0.02	0.07	4	0.04	0.01	0.03	0.06

738

739



740 **Table A4.** Descriptive statistics of topsoil variables for Argiudolls and timber plantations and
 741 crops within of Argiudolls. Number of single samples (n), mean, standard deviation (SD),
 742 minimum (Min) and maximum (Max) of each variable are given.

Variable (unit)	Total Argiudolls					Timber plantations					Crops				
	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max	n	Mean	SD	Min	Max
Humidity (%)	77	18.3	10.9	1.3	50.5	22	14.0	7.8	5.5	38.0	12	16.5	13.0	1.3	50.5
N _{total} (%)	77	0.24	0.15	0.07	1.20	22	0.22	0.10	0.10	0.45	12	0.22	0.06	0.13	0.29
C _{total} (%)	77	3.3	3.0	1.1	25.7	22	3.2	1.6	1.2	8.0	12	2.6	0.7	1.4	3.3
C/N ratio	76	13.3	2.2	9.8	19.9	22	14.0	2.6	11.3	19.9	12	11.8	0.6	10.7	12.7
SOM (%)	73	5.8	2.5	2.1	16.1	22	6.3	3.3	2.3	16.1	12	5.2	1.4	2.8	6.7
P (mg kg ⁻¹)	76	280	122	93	693	22	234	70	111	353	12	321	119	204	613
Ca (cmol kg ⁻¹)	21	11.4	8.0	2.6	37.1	5	6.9	4.1	2.6	11.4	4	11.7	5.9	7.7	20.4
Mg (cmol kg ⁻¹)	21	2.3	1.4	0.5	6.1	5	1.5	0.8	0.5	2.7	4	2.6	1.1	1.8	4.1
K (cmol kg ⁻¹)	21	0.45	0.26	0.09	1.29	5	0.34	0.21	0.09	0.60	4	0.46	0.27	0.29	0.85
Na (cmol kg ⁻¹)	21	0.08	0.05	0.0	0.3	5	0.1	0.06	0.05	0.20	4	0.06	0.02	0.04	0.08
Zn (mg kg ⁻¹)	21	2.4	3.0	0.1	9.7	5	0.5	0.4	0.2	1.2	4	4.1	4.2	0.6	9.7
Cu (mg kg ⁻¹)	21	15.5	42.7	0.8	161	5	2.0	1.5	1.0	4.5	4	71.8	83.4	0.8	161
Fe (mg kg ⁻¹)	21	117	62	10	232	5	158	73	69	232	4	91	38	63	143
Mn (mg kg ⁻¹)	21	23.3	10.4	7.1	43.6	5	29.8	9.8	17.5	43.6	4	15.3	3.7	9.9	18.1
pH	77	5.0	0.7	3.8	6.6	22	4.9	0.8	3.8	6.4	12	5.2	0.3	4.9	5.7
As (mg kg ⁻¹)	76	3.2	1.4	1.0	12.2	22	3.0	1.0	1.0	4.4	12	3.4	1.2	2.1	5.5
Cd (mg kg ⁻¹)	76	0.39	0.13	0.18	0.74	22	0.37	0.12	0.18	0.60	12	0.43	0.13	0.27	0.66
Cr (mg kg ⁻¹)	77	9.3	6.2	2.2	28.3	22	9.5	7.3	2.7	26.4	12	8.2	1.7	6.0	11.3
Pb (mg kg ⁻¹)	77	5.6	2.1	2.8	10.7	22	5.6	2.1	2.8	8.5	12	5.3	1.2	3.9	7.3
CEC (cmol kg ⁻¹)	21	14.2	9.3	3.4	44.6	5	8.8	4.6	3.4	13.6	4	14.8	7.0	9.9	25.0
K ⁺ /(Ca ²⁺ +Mg ²⁺)	21	0.04	0.02	0.02	0.1	5	0.04	0.03	0.02	0.07	4	0.04	0.03	0.02	0.09
K ⁺ /(Ca ²⁺ +Mg ²⁺ +Na ⁺)	21	0.04	0.02	0.02	0.08	5	0.04	0.02	0.02	0.07	4	0.04	0.03	0.02	0.08

743