

Evaluation of the NAQFC Driven by the NOAA Global Forecast System Version 16: Comparison with the WRF-CMAQ During the Summer 2019 FIREX-AQ Campaign

Youhua Tang^{1,2}, Patrick C. Campbell^{1,2}, Pius Lee¹, Rick Saylor¹, Fanglin Yang³, Barry Baker¹, Daniel Tong^{1,2}, Ariel Stein¹, Jianping Huang^{3,4}, Ho-Chun Huang^{3,4}, Li Pan^{3,4}, Jeff McQueen³, Ivanka Stajner³, Jose Tirado-Delgado^{5,6}, Youngsun Jung⁵, Melissa Yang⁷, Ilann Bourgeois^{8,9}, Jeff Peischl^{8,9}, Tom Ryerson⁹, Donald Blake¹⁰, Joshua Schwarz⁹, Jose-Luis Jimenez⁸, James Crawford¹¹, Glenn Diskin⁷, Richard Moore⁷, Johnathan Hair⁷, Greg Huey¹¹, Andrew Rollins⁹, Jack Dibb¹², Xiaoyang Zhang¹³

1. NOAA Air Resources Laboratory, College Park, MD, USA.

2. Center for Spatial Information Science and Systems, George Mason University, Fairfax, VA, USA.

3. NOAA National Centers for Environmental Prediction, College Park, MD, USA

4. I.M. Systems Group Inc., Rockville, MD, USA

5. Office of Science and Technology Integration, NOAA National Weather Service, Silver Spring, MD, USA

6. Eastern Research Group Inc, USA

7. NASA Langley Research Center, Hampton, VA, USA

8. Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA

9. NOAA Chemical Sciences Laboratory, Boulder, CO, USA

10. Department of Chemistry, University of California at Irvine, Irvine, CA, USA

11. School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

12. Earth Systems Research Center, University of New Hampshire, Durham, NH, USA

13. Department of Geography & Geospatial Sciences, South Dakota State University, Brookings, SD, USA

Correspondence: Youhua Tang (youhua.tang@noaa.gov)

Abstract

The latest operational National Air Quality Forecasting Capability (NAQFC) has been advanced to use the Community Multi-scale Air Quality (CMAQ) model version 5.3.1 with CB6r3 (carbon bond 6 revision 3)-Aero7 (version 7 of the aerosol module) chemical mechanism and is driven by the Finite Volume Cubed-Sphere (FV3)-Global Forecast System, version 16 (GFSv16). This update has been accomplished by development of the meteorological preprocessor, NOAA-EPA Atmosphere-Chemistry Coupler (NACC), adapted from the existing Meteorology-Chemistry Interface Processor (MCIP). Differing from the typically used Weather Research and Forecasting (WRF)-CMAQ system in the air quality research community, the interpolation-based NACC can use various meteorological outputs to drive CMAQ (e.g., FV3-GFSv16) even though they are in

different grids. In this study, we compare and evaluate GFSv16-CMAQ vs. WRFv4.0.3-CMAQ using observations over the contiguous United States (CONUS) in summer 2019 verified with surface meteorological and AIRNow observations. During this period, the Fire Influence on Regional to Global Environments and Air Quality (FIREX-AQ) field campaign was performed and we compare the two models with airborne measurements from the NASA DC-8 aircraft. The GFS-CMAQ and WRF-CMAQ systems have overall similar performance with some differences for certain events, species and regions. The GFSv16 meteorology tends to have a stronger diurnal variability of planetary boundary layer height (higher during daytime, and lower at night) than WRF over the U.S. Pacific coast, and it also predicted lower nighttime 10-m winds. In summer 2019, GFS-CMAQ system showed better surface O₃ than WRF-CMAQ at night over the CONUS domain; however, their PM_{2.5} predictions showed mixed verification results: GFS-CMAQ yielded better mean biases but poorer correlations over the Pacific coast. These results indicate that using global GFSv16 meteorology with NACC to directly drive CMAQ via the interpolation is feasible and yields reasonable results compared to the commonly-used WRF driving approach.

1. Introduction

Traditionally, mesoscale meteorological models such as the Weather Research and Forecasting Model (WRF) (Powers et al., 2017) are used as the meteorological drivers for air quality models (AQMs) on the same (“native”) model grid, such as Community Multiscale Air Quality Model (CMAQ) (Byun & Schere, 2006). The NOAA National Weather Service’s (NWS) National Air Quality Forecasting Capability (NAQFC) has historically used a different approach, in which the hourly meteorological outputs from prior operational models, such as North American Mesoscale Model (NAM), are interpolated to the CMAQ grid to drive its air quality prediction. Prior to this work, a “PREMAQ” coupler (Otte et al, 2004) combined both meteorological processing and Sparse Matrix Operator Kernel Emissions (SMOKE) (Houyoux et al., 2000) processes, such as point source plume rise effects. However, since the release of CMAQ version 5, the meteorology-dependent plume rise, sea salt, and dust emission processes are included as inline modules in CMAQ, and thus the corresponding emission processes are no longer needed in PREMAQ. Furthermore, PREMAQ has no built-in interpolator, and thus relied on external interpolators to remap the non-native-grid meteorological inputs, such as NAM, to the targeted CMAQ domain, though it did perform vertical layer collapsing/interpolation to reduce vertical layers for CMAQ. The interpolation approach allows more flexibility for using different meteorological data (i.e., besides just WRF) to drive CMAQ; however, there is potential to raise mass-consistency issues between models. It should be noted that the mass-consistency issues may also exist using native-grid couplers (Byun, 1999a, 1999b), which can stem from the original mass-inconsistent meteorological model outputs or due to the temporal interpolation of the meteorological data. The well-developed offline AQMs, such as CMAQ, have already considered such mass-consistency treatments using different meteorological inputs (Byun et al., 1999c).

To upgrade the NAQFC system with the latest CMAQ model and NOAA operational meteorology, we developed an updated interpolation-based meteorological coupler, the NOAA-EPA Atmosphere-Chemistry Coupler (NACC) (Campbell et al., 2022) adapted from the U.S. EPA’s Meteorology-Chemistry Interface Processor (MCIP) version 5 (Otte and Pleim, 2010; <https://github.com/USEPA/CMAQ>). The NACC system replaced PREMAQ, and effectively couples the Finite-Volume Cubed-Sphere (FV3) Dynamical Core - Global Forecast System version 16 (GFSv16) (Yang et al., 2020; Harris et al, 2021) to CMAQ v5.3.1 (hereafter referred to as GFS-CMAQ). Campbell et al. (2022) described the development and application of the GFS-CMAQ system using NACC (in their work referred to as “NACC-CMAQ”) and a comprehensive comparison between the current (GFS-CMAQ since July 20, 2021) and previous (NAM-CMAQv5.0.2) operational NAQFC model performances.

In this study, we analyze the impacts of the meteorological model drivers, and compare GFS-CMAQ using NACC interpolation to the commonly-used native-grid WRF-CMAQ application and its impact on air quality predictions in summer 2019. Yu et al. (2012a, 2012b) had previously compared the CMAQ performance driven by WRF-NMM and WRF-ARW during the 2006 TexAQS/GoMACCS field campaign, and found that the NMM-CMAQ and ARW-CMAQ showed overall similar performance with some differences for certain events, chemical species, and regions. Similarly, this study focuses on the comparison of GFS-CMAQ versus WRF-CMAQ (see Section 2, Methodology), and verifies the model performance against the aircraft observations from the Fire Influence on Regional to Global Environments and Air Quality (FIREX-AQ) field experiment during summer 2019 (Section 4). Surface verification is also performed using AIRNow data for August 2019 (Section 3), serving as a benchmark for the new NAQFC versus the traditional WRF-CMAQ used in the air quality modeling community. This study focuses on the period of summer 2019, and Campbell et al. (2022) evaluated the GFS-CMAQ for longer periods.

2. Methodology

Here we compare the two CMAQ (version 5.3.1) runs driven by the interpolated GFSv16 meteorology (GFS-CMAQ) and WRF meteorology (WRF-CMAQ). All other settings, such as emission and lateral boundary conditions are the same. The meteorology-related physics is discussed in the following sections to address the models’ performance discrepancies. Both the GFS-CMAQ and WRF-CMAQ simulations are run from a period covering 12 July – 31 August, 2019, each using the last 10 days in July as the model spin-ups that are not included in the analyses.

2.1 GFS Meteorological Inputs

The GFSv16 is the current operational global forecast system in NOAA/NCEP using FV3 dynamical core. Its detailed configuration can be found in Campbell et al. (2022) and Yang et al.

(2020). Compared to the previous version (v15), GFSv16 updated many physical schemes (Table 1) and added the parameterization for sub-grid scale nonstationary gravity-wave drag. To use the GFS's meteorology to drive CMAQ, a meteorological coupler, NACC, is developed (Campbell et al., 2022). Differing from the original MCIP which was developed to process WRF/ARW meteorology for CMAQ, the NACC coupler interpolates non-native-grid meteorology to a user-defined grid and has parallel processing capability, which drastically reduces its run time for operational forecasts (Campbell et al., 2022). Currently, NACC employs two horizontal interpolation methods: bilinear and nearest-neighbor. In this study, we use the nearest-neighbor method to categorical (discontinuous) variables that include land use types, vegetation fraction, terrain elevation, Monin–Obukhov length, friction velocity, and soil temperatures, while the bilinear interpolation is used for mainly smoothly varying (continuous) meteorological variables that include wind fields, temperature, pressure, and specific humidity. The CMAQ model is defined in the Arakawa C-grid (Arakawa and Lamb, 1977), and thus the GFSv16 horizontal wind components (U, V) need to be interpolated to the perpendicular cell faces instead of the cell center (Otte and Pleim, 2010) after rotation to the defined map projection. The scalar variables are defined in the cell centers of the targeted grid, and thus their interpolations are more straightforward: from GFSv16 A-grid to CMAQ A-grid. The NACC coupler can either use the native layers or interpolate inputs to a set number of user-defined CMAQ vertical layers. The GFSv16 has 127 vertical layers with global coverage in roughly 13 km horizontal resolution, where the targeted CMAQ domain is in 12×12 km grids over the Contiguous United States (CONUS) with 35 vertical layers (Campbell et al, 2022). Here we use 24-hour GFSv16 forecasts starting at 12 UTC each day.

Most variables needed by CMAQ are directly interpolated from the GFSv16 hourly outputs. The NACC processor has options to calculate diagnostic variables, such as planetary boundary layer (PBL) height, if they are needed. In this study, we use the interpolated GFSv16's PBL height instead of the diagnostic one. It also has an option to import the externally provided land-surface variables. Here we import updated 2018–2020 climatological averaged leaf area index (LAI) and NOAA near-real-time (NRT) greenness vegetation fraction (GVF) from satellite-based Visible Infrared Imaging Radiometer Suite (VIIRS) retrievals (Campbell et al., 2022). The updated satellite-based LAI and GVF impact CMAQ's biogenic emissions and dry deposition processes, which were described in detail in Campbell et al. (2022).

2.2 WRF Meteorology

To compare with GFSv16 meteorology processed by NACC, a corresponding WRF version 4.0.3 (Skamarock et al, 2021) simulation is run covering the NAQFC's native grid, which is a 12 km horizontal resolution with Lambert conformal map projection over CONUS. Table 1 shows the WRF configuration, which is commonly employed in CONUS meteorological and air quality studies in the community, versus the current NOAA/NWS operational global model GFSv16. GFSv16 uses the NOAA/NCEP's Global Data Assimilation System (GDAS)

(https://www.emc.ncep.noaa.gov/data_assimilation/data.html) for its initial conditions and runs on its own global dynamics and physics without any other constraints. The regional WRF simulation uses GFSv16 for its initial conditions. In this study, GFSv16 was re-initialized with GDAS every 24 hours, and WRF conducted the continuous run after its spin-up period. Furthermore, WRF also takes its lateral boundary conditions from GFSv16 every 6 hours. For the WRF run, we have enabled the four-dimensional data assimilation (FDDA) for the horizontal winds, temperature, and humidity (Table 1) every 6 hours, thus nudging towards GFSv16. This nudging method used in WRF runs can help reduce the difference of two meteorological models, though its effect may vary depending on events as WRF and GFS use different physics.

WRF and GFSv16 have similar settings for the land surface model, surface layer and radiation schemes; however, their microphysics and PBL schemes are different (Table 1). Compared to the 35-layer WRF with a 100 hPa domain top, GFSv16 has a much higher domain top (0.2 hPa) and 127 vertical layers, which are interpolated by NACC to 35 sigma layers up to 14 km for CMAQ. We use NACC (inherited from MCIP version 5.0) to process WRF hourly meteorology, while maintaining the 35-layer vertical layer structure. Thus, in contrast to GFS-CMAQ, the WRF-CMAQ system uses the native grid without interpolation.

2.3 CMAQ Configuration

Here CMAQ version 5.3.1 (Appel et al., 2021) is used with the Carbon Bond 6 revision 3 (CB6r3; Yarwood et al., 2010, 2014; Luecken, et al., 2019) chemical mechanism and Aero7 treatment of secondary organic aerosols (CB6r3_AE7_AQ). CMAQ 5.3.1 includes a series of scientific updates from the previous version (Appel et al., 2021), including the updated air-surface exchange and deposition modules, which showed significant impact on ozone prediction compared to the previous NAQFC (Campbell et al., 2022). We also include the bi-directional NH₃ (BIDI-NH₃) exchange model for NH₃ surface fluxes. An updated Biogenic Emissions Landuse Dataset v5 (BELD5) is used in this study to drive the inline Biogenic Emissions Inventory System (BEIS) version 3.61. The anthropogenic emissions are provided by the National Emissions Inventory Collaborative (NEIC) with base year 2016 version 1 (NEIC 2016). We replace the U.S. EPA default CMAQ dust emissions model with an inline windblown dust model known as “FENGSHA” (Fu et al., 2014; Huang et al., 2015; Dong et al., 2016). The FENGSHA dust scheme uses the sediment supply map and magnitude of the friction velocity (USTAR) compared to a threshold friction velocity (UTHR) to calculate the potential of dust emission flux. The UTHR depends on the land cover, soil type (clay fraction), and soil moisture Campbell et al. (2022).

We have updated the wildfire emissions system in CMAQv5.3.1 based on the Blended Global Biomass Burning Emissions Product (GBBEPx) (Zhang and Kondragunta, 2006; Zhang et al., 2011). The GBBEPx uses satellite-detected fire radiative power (FRP) to estimate wildfire smoke emissions for a number of species: CO (carbon monoxide), NO_x (nitrogen oxides), SO₂

(sulfur dioxide), elemental carbon, primarily emitted organic aerosols, and PM_{2.5}. We derive the wildfire volatile organic compounds (VOC) emission from GBBEPx CO emission based on the emission ratios of FINN data (Wiedinmyer et al., 2011), and the splitting factors from the SMOKE model (Baker et al, 2016) are used to further split the total fire VOC emission to speciated hydrocarbon emissions. The satellite FRP is estimated from satellite brightness temperature anomaly, and the GBBEPx processor assumes that the wildfire emissions are proportional to the FRP over certain landuse types in certain regions. The GBBEPx emissions are based on polar orbiting satellites: MODIS (Aqua and Terra satellites) and VIIRS (Suomi-NPP and NOAA-20 satellites) instruments, which are updated once per day. A wildfire emission preprocessor converts the GBBEPx emissions to CMAQ-ready input files using emission speciation and diurnal profiles (high during daytime and low at night) (adopted from U.S. EPA-based profiles) (Baker et al., 2016), and a daily scaling factor. Here we classify the wildfire into either a long-lasting fire (longer than 24 hours) or short-term fire (shorter than 24 hours) based on land use types and regions. As historic statistics shows that most fires (>95%) in east of 110°W last less than 24 hours, only the fires west of 110°W that have a model grid cell total forest fraction > 0.4 are assumed to be long-lasting fires. All other short-term GBBEPx fires are assumed to have smoke emissions for 24 hours (i.e., day 1 only). Burning area could be highly uncertain as GBBEPx data do not include this information. One grid cell could have multiple fires and some big fire could appear in several grids. Here we carry the previous NAQFC's method, and apply a constant ratio: 10% of the grid cell, as the burning area (Pan et al., 2020) according to Rolph et al (2009). CMAQ treats wildfire emissions as point sources that undergo inline plume rise to distribute the smoke vertically. The default CMAQ plume rise used here is based on Briggs (1965), which is driven by fire heat flux (converted from FRP with a ratio of 1) and fixed burning area (assumed to be 10% of the 0.1°×0.1° grid cell).

3. Model Evaluations over the U.S. for August 2019

To first gain a general picture and compare the overall GFS-CMAQ and WRF-CMAQ model performances, in this section we evaluate the near-surface meteorological and air quality predictions during the FIREX-AQ August 2019 period against NOAA's METeorological Aerodrome Report (METAR; https://madis.ncep.noaa.gov/madis_metar.shtml) and the U.S. EPA's AirNow (<https://www.airnow.gov/>) observation networks. All the comparisons of meteorological variables are for those actually used in CMAQ. For GFS-CMAQ, it refers to the interpolated GFS data. Campbell et al (2022) included the detailed comparisons before and after interpolation, showing that the interpolated meteorology was very consistent with the original one. In this study, the model results are spatiotemporally interpolated to the corresponding observation locations for comparison.

3.1 Domain-Wide Meteorology against the METAR Network

Figure 1 shows the mean bias (MB) of interpolated GFS and WRF predicted surface meteorological variables compared to METAR data during August, 2019. Both meteorological models have a cool bias over the Western and Northeastern United States, and a warm bias over the western Rocky Mountain region and Southeastern United States (Figure 1a, 1b). Similar temperature predictions are expected since WRF uses the FDDA method nudging toward GFS data. However, GFS tends to be cooler than WRF over the Rocky Mountains and in the central and northeastern USA due to their different dynamics and physics. The GFSv16 cold bias in the lower troposphere is impacted by excessive evaporative cooling from rainfall (personal communication with NOAA/NCEP). Campbell et al. (2022) had detailed discussions about GFSv16 biases.

Both GFSv16 and WRF models have similar and rather significant dry biases for specific humidity (SH) predictions across CONUS (Figure 1c, 1d). Qian et al (2020) investigated this common dry bias in many models, and found that neglecting an irrigation contribution could cause this dry bias. Besides this issue, WRF's dry bias could also be affected by its nudging toward GFS, as GFS has widespread dry biases (Campbell et al. 2022). Their biases also have some noticeable differences over certain regions. For instance, WRF has less dry bias over Southern Texas than GFS.

Both models underestimate the mean 10-m wind speeds compared to METAR stations over the western U.S. WRF has stronger underpredictions over the Rocky Mountains and overpredictions over northeastern U.S., while GFS has stronger underpredictions over the Appalachian Mountains and overpredictions over Texas and Oklahoma. GFSv16's operational verification (https://www.emc.ncep.noaa.gov/gmb/emc.glopara/vsdb/v16rt2/g2o/g2o_00Z/index.html) also shows that it tends to underpredict the 10-m wind over the western U.S. during both daytime and nighttime, but shows overpredictions over the eastern U.S. Besides the difference of physical schemes etc. (Table 1), other possible reasons causing this surface wind difference could be effect of the gravity-wave drag (GFSv16 includes it, but the WRF run here does not), and vertical resolutions: GFS's 127 layers versus WRF's 35 layers, though they have similar vertical layers below 1km (Campbell et al, 2022). Some studies (Skamarock et al, 2019) revealed the necessity of fine vertical resolution for atmospheric simulations, especially within the PBL, near tropospheric top, and during convective events. Insufficient vertical resolution could also cause plume dilution on chemical transport modeling (Zhuang et al., 2018). The gravity-wave drag is also known to influence the synoptic-scale dynamics on the atmospheric flow over irregularities at the earth's surface such as mountains and valleys, and uneven distribution of diabatic heat sources associated with convective systems (Kim et al., 2003). Its parameterization is needed for large-scale models.

There are strong regional variabilities in the monthly mean PBL height differences between GFS and WRF during normal daytime (represented by 18 UTC) and nighttime (represented by 06 UTC) (Figure 2). During daytime, GFS has a higher PBL height compared to WRF over the U.S. Pacific coast, northern Rocky Mountains, northeastern and southeastern U.S., but it becomes lower over the central U.S. (e.g., Texas, Oklahoma, and Kansas). At night, however, most of these regional differences between GFS and WRF are reversed. This diurnal difference is mainly driven by the different PBL schemes employed in GFS (Han and Bretherton, 2019) and WRF (i.e., YSU) and the associated other physical suites, including the land surface data. So this PBL difference has strong regional variations depending on geographic differences. The GFS's PBL height has a strong diurnal variation over these regions, including the western and northeastern U.S., and it shows a sharp rise and collapse after sunrise and sunset, respectively (Campbell et al., 2022). These two selected times (18UTC and 06UTC) are not in the transition periods for PBL's fast diurnal changes.

3.2 Evaluation of Regional Meteorology and Air Quality against the AirNow Network

The U.S EPA AirNow network provides hourly observations of near-surface ozone, fine particulate matter (PM_{2.5}), and meteorology. Campbell et al. (2022) showed detailed verification of GFS-CMAQ with the surface AIRNow data. Here we focus on the difference between the interpolation-based GFSv16 versus WRF downscaling and the impacts on meteorological and chemical model performances. Figure 3 shows a comparison of these two models over two specific regions, the U.S. West (CA, OR and WA) and Northeast states (CT, DE, MA, MD, ME, NH, NJ, NY, PA, RI, VT and District of Columbia) (Figure S1), where the two models have relatively large differences for some meteorological variables. GFS and WRF predict very similar 2-m temperatures over the Pacific coast states: Washington, Oregon and California, and both of them had similar cool bias (around 1K), R and RMSE (Figure 3a). However, these two models show significant differences for the 10-m wind speed prediction over the Pacific coast (Figure 3c), where WRF overpredicts the wind speed, especially at night and in later August. Most AIRNow stations are located near urban or suburban areas, which generally have weaker 10-m wind speed than those at the METAR aviation weather stations near airports. For this reason, although Figure 1e and 1f shows that GFS and WRF underpredict monthly-mean wind speed over the METAR stations in the West, they still tend to overpredict wind (Figure 3c) over AIRNow stations, especially for the WRF 10-m wind speed at night. Considering that the model grid cells represent 12×12 km² averages, the true model-observation comparisons likely fall somewhere between the urban/suburban AIRNow stations and METAR stations, depending on the landuse fractions of each grid. Obviously the representation characteristics of observations could affect the verification results. Compared to AIRNow observations, GFSv16 has overall better scores for surface wind speed predictions over the U.S. West, where the WRF's larger surface wind speed overprediction is associated with its PBL height predictions (Figure 3e, 3f). During the nighttime, GFS has a lower PBL height (10–50% lower than WRF) and weaker vertical mixing, which tends to bring less momentum flux from the upper layers to the surface,

1 and lead to lower nighttime wind and better agreements with the AIRNow wind-speed
2 observation.

3
4 Over the northeast, the mean bias (MB) of GFS temperature is about -1K, while the WRF has a
5 smaller, slightly positive MB of about 0.22K (Figure 3b). However, the GFS's temperature
6 prediction has a better correlation coefficient, R, and RMSE, implying that it better captures
7 some events, such as the 28–29 of August. Both models overpredict 10-m wind speeds in the
8 northeast, but the GFS model yields better results due to a slightly lower PBLH at night (Figure
9 3f) than WRF that had significant overpredictions, especially during 25–29, August (Figure 3d)
10 when the tropical storm Erin approached this region. Especially on 28 of August, when the storm
11 was centered near the east coast of North Carolina, WRF significantly underpredicted 2-m
12 temperature (Figure 3b) and overpredicted 10-m wind speed (Figure 3d). In the west around the
13 same period, the tropical storm Ivo appeared in the southwest of the Baja California peninsula,
14 and brought heavy rainfall in Mexico. Associated with this storm, a low pressure system
15 expanded over most of the U.S. West. Differing from GFSv16, which is designed for the
16 operational meteorological forecast, the WRF configuration used in this study is normal for
17 driving CMAQ, but not tuned for storm weather prediction.

18
19 Figures 4a and 4b show the ozone predictions of the two models over these two regions, and
20 GFS-CMAQ yields predominantly lower O₃ than WRF-CMAQ, especially at night. Over the
21 west, the lower ozone in GFS-CMAQ is associated with their PBL height difference. First, with a
22 certain dry deposition velocity between the models, it is easier to deplete ozone given the smaller
23 volume of a shallower PBL. Second, the shallower PBL results in higher surface NO_x
24 concentrations (not shown) and ozone titration rates near NO_x source regions, and consequently
25 lower ozone there at night. Last, the lower PBL could decouple from the residual layer, and
26 result in weaker or no vertical ozone exchange with the residual layer at night (Caputi et al,
27 2019). All these factors contribute to the lower nighttime ozone of GFS-CMAQ compared to
28 WRF-CMAQ. Since GFS-CMAQ already underpredicts ozone due to combined meteorological
29 factors, such as the temperature underprediction (Figure 4a), the GFS-CMAQ's further ozone
30 reduction (possibly due to its lower PBLH at night) exacerbates its low bias. However, over the
31 Northeast, the similar impacts help the GFS-CMAQ yield much better MB due to its better
32 agreement with the observed nighttime low ozone over the Northeast. Over the entire CONUS
33 domain, the situation is similar: for an average August 2019, the GFS-CMAQ has a lower ozone
34 MB (1.1 ppb) compared to WRF-CMAQ (4.7 ppb). Figure 5 shows that both models have
35 similar daytime ozone prediction over CONUS. However, GFS-CMAQ better captures low
36 nighttime ozone over the U.S. East than WRF-CMAQ (Figure 5c, 5d).

37
38 GFS-CMAQ has substantially higher PM_{2.5} mean concentrations over the U.S. West, but lower
39 over the U.S. Northeast compared to WRF-CMAQ (Figures 4c, 4d). These model differences are
40 also related to their interpolated GFSv16 versus downscaled WRF meteorological drivers.

Because both models use the same emissions under relatively clean background conditions in the west (i.e., prevailing westerly flow from the Pacific Ocean), the PBL and wind speed differences have significant impacts on their near-surface pollutant concentrations, especially at night. Both models show strong PM_{2.5} diurnal variations (high at night and low during daytime), driven by the meteorological diurnal variation (e.g. PBL), which overcomes the emission diurnal variation (usually high during daytime and low at night). Compared to WRF-CMAQ, GFS-CMAQ has lower nighttime PBL height and weaker wind speed at night, which leads to weaker vertical mixing and venting, and increases the pollutant concentrations near the surface and yields higher surface PM_{2.5} over the U.S. West (Figure 4c). Its higher surface PM_{2.5} could also result in stronger local dry deposition. In contrast to local vertical mixing and venting effects on PM_{2.5} discussed above, there are strong (and potentially counterbalancing) impacts of model PBL and horizontal wind speed differences on downstream PM_{2.5} concentrations at night. WRF-CMAQ's deeper PBL and stronger wind speeds at night (Figures 3c–3f) tends to transport aerosols and their precursors more efficiently downstream via the dominant advection pathway. Figure 6 shows that these monthly mean background PM_{2.5} differences appear in East of Rocky Mountain (WRF-CMAQ is about 2 $\mu\text{g}/\text{m}^3$ higher) during both daytime and nighttime. This effect is very prominent in the Northeast region. Although both models predicted similar PM_{2.5} magnitude over the U.S. Northeast, GFS-CMAQ yields the overall PM_{2.5} underprediction, and its monthly-mean PM_{2.5} is 2.6 $\mu\text{g}/\text{m}^3$ lower than the WRF-CMAQ prediction (Figure 4d). Especially during 01–09 August, WRF-CMAQ had about 4 $\mu\text{g}/\text{m}^3$ higher surface PM_{2.5} background than that of GFS-CMAQ. In this case, the WRF-CMAQ model has a better agreement with observations (Figure 4d). It is possible that the GFS-CMAQ's nighttime PBL heights (wind speeds) are too shallow (weak) in this case, which does not allow enough transport of pollutants to the downstream (Eastern USA). Overall, GFS-CMAQ and WRF-CMAQ have mixed performances for PM_{2.5} predictions during the August 2019 period: GFS-CMAQ has better PM_{2.5} prediction over the U.S West, and WRF-CMAQ yields better results over east of Rocky mountain (Figure 6).

4 Model Comparisons against the FIREX-AQ Aircraft Data

From late July to early September, 2019, the joint NOAA-NASA FIREX-AQ field campaign (<https://csl.noaa.gov/projects/firex-aq/>) employed a suite of satellites, aircraft, vehicles and ground site platforms aimed to observe, analyze, and characterize air pollutants emitted from wildfire sources over the CONUS (Ye et al., 2021). The FIREX-AQ airborne measurements provide a three-dimensional dataset from various meteorological, gas, and aerosol instruments that can be used to verify the GFS-CMAQ and WRF-CMAQ model performance, while elucidating reasons for any model differences. Here the focus of the FIREX-AQ model comparison and verification is against observations taken primarily from the NASA DC-8 aircraft, which include meteorological variables, gaseous and aerosol concentrations, and aerosol optical properties merged in one-minute temporal resolution. The model data are spatiotemporally interpolated to the flight paths for comparison. The majority of the FIREX-AQ

flights were over the western United States, and sampled within environments that both were *and* were not (see section 4.1) influenced by wildfire emissions (https://daac.ornl.gov/MASTER/guides/MASTER_FIREX_AQ_JulySept_2019.html). During a cluster of major wildfire events (see Section 4.2), the DC-8 sampled both near-source and aged smoke plumes between 02–08 August, 2019 (i.e., the Williams Flats, Snow Creek, and Horsefly Fires) across the states of Idaho, Washington, and Montana.

4.1 Comparison of the July 22 non-wildfire event over the central California Valley

On 22 July, the DC-8 aircraft flew from California to Boise, Idaho, while maintaining a relatively low-altitude (<1 km) above sea level (ASL) over the California Central Valley (Figure 7). This flight was not impacted by any major wildfire event, and was mainly controlled by anthropogenic emissions and local meteorological conditions. Figure 7 shows that the GFSv16 and WRF models had similar meteorological temperature and humidity predictions, and that both models have dry and warm biases over the Central Valley at lower altitudes (Figures 7d–7e) (Yun et al., 2020). GFS’s horizontal wind speeds tend to have a stronger variability than WRF (Figure 7b), especially in high altitudes. For wind direction, WRF shows a better prediction than GFS around 20 and 24 UTC (Figure 7c).

Both GFS-CMAQ and WRF-CMAQ underestimate the vertical wind (*W*) variability by at least one order of magnitude, and WRF-CMAQ has weaker *W* variability than that of GFS-CMAQ, especially in high altitudes (Figure 7f). The model vertical velocities are not directly from the GFS or WRF model, but rather they are re-diagnosed in CMAQ to conserve mass (Otte and Pleim, 2010), and thus represent the whole layer’s vertical movement across the 12 km by 12 km grid cell. With its flight speed around 80 to 240 m/s, the DC-8 aircraft’s one-minute average sampling frequency results in an approximate 4.8 to 14 km horizontal scale, respectively, which is comparable to the 12 km CMAQ model resolution. The aircraft observations, however, include turbulence effects during its one-minute averages, which may not be temporally resolved by the models at this resolution. Thus, both the GFS-CMAQ and WRF-CMAQ vertical velocities are much lower and have almost no correlation with the aircraft observations.

Although both GFS-CMAQ and WRF-CMAQ have reasonable comparisons for most meteorological variables, including the horizontal winds, it continues to be a challenge to compare them with the observed vertical velocities. Thus to further elucidate the model vs. observation differences in vertical motions, Figure 8 shows a curtain plot of vertical velocities along the flight path from the two models. Since WRF-CMAQ remains in a native grid, its wind fields tend to be more balanced and have lower variability compared to the GFS-CMAQ wind fields. The stronger variability in *W* for GFS-CMAQ could be caused by GFS’s non-hydrostatic dynamics or CMAQ’s effort to counteract mass inconsistency effects from the interpolated horizontal wind fields (Byun, 1999b). Our comparison shows that the first factor should be the major one (Figure S2) for this event, as the GFS-CMAQ diagnosed *W* is very similar to that

1 from the original GFSv16 around 1km above the ground. As the original GFSv16 has similarly
2 stronger vertical velocities than the original WRF, the W difference between GFS-CMAQ and
3 WRF-CMAQ is unlikely due to interpolation error of horizontal winds.

4
5 GFS-CMAQ and WRF-CMAQ overall yield similar results for specific chemical species during
6 this DC-8 flight (Figure 9). Both models underestimate CO, O₃ and ethane (C₂H₆) concentrations
7 over the lower altitudes in the California Central Valley. Over the same flight segment, they have
8 better NO_x (NO + NO₂) and ethene (C₂H₄) predictions, implying that the emissions of these two
9 species have better accuracy than those of CO and ethane. Figure 9f shows that the two models
10 also underestimate NO_z (NO_y–NO_x), or the oxidized nitrogen species besides NO_x, indicating
11 that the photochemical ozone production may also be underestimated. NO_z is a good indicator of
12 the ozone photochemical formation (Sillman et al., 1997), where the O₃/NO_z ratio represents the
13 ozone photochemical efficiency per NO_x oxidation products. Thus, NO_z and O₃ are typically
14 highly correlated over regions with active photochemical production. Our later analysis shows
15 that the models tend to underestimate certain hydrocarbons, such as ethane, which is likely
16 linked to O₃ and NO_z underestimations, since the hydrocarbons are photochemical precursors of
17 O₃ and NO_z.

18
19 The two models show slight differences in peak values of CO, ethene, and NO_x around 23:30
20 UTC, where the GFS-CMAQ predicted concentrations are slightly higher and closer to
21 observations (Figure 9). These differences are due to their PBL predictions (both from the
22 corresponding meteorological model outputs), where GFS-CMAQ has a lower PBL height and
23 weaker emission vertical dilution compared to WRF-CMAQ (Figure 8). GFS-CMAQ tends to
24 underpredict O₃ more (Figure 9b), however, due to its higher NO_x titration. This implies that the
25 effects of the transport and non-local transformation of O₃ could be stronger than that of local
26 precursor emissions. WRF-CMAQ has higher NO_z (Figure 9f), but lower NO_x compared to
27 GFS-CMAQ due to the time lag of O₃ and NO_z photochemical formation. Consequently, the
28 peak O₃ values may not be well correlated with the emitted precursors, such as NO_x and VOC.
29 Furthermore, the modeled peak C₂H₆ and C₂H₄ concentrations do not occur at the same time
30 around 23:30 UTC, while observations indicate that these two species should be highly
31 correlated in this region. This model mismatch implies that the VOC speciation factors for a
32 certain area or emission sector need to be improved over Southern California.

33 34 **4.2 Comparison of the 6 August wildfire events over the U.S. Northwest**

35 On 06 August, the DC-8 observed a cluster of three wildfires: the Williams Flats Fire (47.98 °N,
36 118.624 °W, 80 km to the northwest of Spokane, Washington), Snow Creek Fire (47.703°N,
37 113.4°W, 32 km northeast of Condon, Montana), and Horsefly Fire (46.963 °N, 112.441°W, 24
38 km east of Lincoln, Montana). Figure 10a shows the flight path on that date, where the DC-8
39 aircraft departed from Boise, ID, flew over the Williams Flats Fire region, then flew to Montana
40 to sample the Snow Creek and Horsefly Fires (i.e., Montana Fires), and finally returned to the

Boise base. The aircraft flew below 8 km for most flight segments near the fire plumes. Figure S3 shows the corresponding GOES-16 satellite true color image, where these 06 August fires and associated smoke plumes are visible and can be distinguished from the cloud bands to the south that move northward later that day (Figure S3). The Williams Flats Fire was ignited by lightning, and was the largest fire event sampled during the FIREX-AQ campaign burning from about 02–08, August, 2019.

Both models significantly underpredicted CO (Figure 10c), submicron (aerosol diameter < 1 μ m) organic aerosol (Figure 10e) and aerosol optical extinction coefficient (AOE) (Figure 10f), which suggests an issue with the GBBEPx gas and aerosol emissions. The models performed well for NO₂ during the Williams Flats and Montana Fires Fire below 6 km ASL, but there were prominent underestimations for the high-altitude flight segments (Figure 10d). However, since the NO₂ instrument (The NOAA NOyO3 4-channel chemiluminescence) had its NO₂ detection limit around 0.01 ppb

(https://airbornescience.nasa.gov/sites/default/files/documents/NOAA%20NOyO3_SEAC4RS.pdf), the models might not truly underestimate NO₂ for these flight segments with extremely low NO₂. WRF-CMAQ predicted higher O₃ values than the GFS-CMAQ, which overall agreed better with observations for the Williams Flats Fire (Figure 10b). However, for the Montana Fires (~23–24 UTC), WRF-CMAQ has higher O₃ biases and GFS-CMAQ yields better results. The difference in O₃ is largely driven by the regional background concentration difference between the two models, where WRF-CMAQ tends to have higher domain-wide O₃ concentrations than GFS-CMAQ due to the meteorological effects discussed in section 3 even though they used the same lateral boundary conditions.

Figure S4 shows the spatial overlay comparison of vertically averaged GFS-CMAQ predictions at 21 UTC and the DC-8 flight observations for the altitude 1–3 km above ground level (AGL), on 6 August, 2019. The peak NO₂ observation around 118.5°W, 48°N indicates the general location of the Williams Flats fire. The GBBEPx emission and GFS-CMAQ prediction showed shifted peak-value locations driven by the westerly modeled winds. For this flight, the GBBEPx had stronger NO_x fire emissions over two Montana locations than that over Williams Flats. The model overpredicts the column averaged NO₂ concentrations, especially over the Montana fires, which can not be reflected by the point-by-point NO₂ comparison result in Figure 10d. For this flight, the mean GFS-CMAQ NO₂ along the flight path for 1–3km AGL is about 0.125 ppbv compared to the observed mean NO₂ of 0.169 ppbv, and the model indeed showed a NO₂ underprediction along the flight path. However, in this case, the flight path did not encounter the locations with modeled peak NO₂ values as the model misplaced the plumes, especially over the Montana fires leading to this inconsistency. For ozone comparison (Figure S4b), this inconsistency could also exist, though not as significant as for the high-gradient NO₂ concentrations. In the GFS-CMAQ prediction, the high ozone concentrations are almost co-located with high NO₂ concentration (Figure S4b), but the observation did not show this feature.

1 Instead, some high-O₃ flight segments had relatively low NO₂ concentrations, such as those
2 circled in the black rectangle box of Figure S4b. The observed NO_x titration was not able to be
3 produced by the 12 km models. Wang et al (2021) used a 100m-horizontal-resolution large eddy
4 simulation and demonstrated the capability of using such techniques to capture some high-
5 resolution features of fire plumes and associated chemical behavior. While such high resolution
6 techniques are not currently feasible for the operational NAQFC, they demonstrate the limitation
7 of using regional scale (12×12 km) models to capture such fine-scale features of the fire plume.
8

9 GFS-CMAQ has higher wildfire-related CO, NO₂, OA and AOE values that are closer to
10 observations than WRF-CMAQ for the Montana Fires between 23–24 UTC at flight altitudes of
11 ~ 4–5 km (Figure 10c–10f). Since these two models use the same GBBEPx emissions and
12 wildfire plume rise algorithm (Briggs, 1965), the differences should be due to other reasons. To
13 help explain these model differences, Figure 11a and 11b show the Differential Absorption High
14 Spectral Resolution Lidar (DIAL-HSRL) retrieved aerosol backscatter coefficients (ABC)
15 aboard the DC-8 aircraft without and with cloud screen, respectively. It shows that the major fire
16 plumes of the Williams Flats Fire were below 4 km (~ 19–22 UTC), but the Montana Fires (~23–
17 24 UTC) extended from the surface up to 6 km, with some detached plumes reaching 10 km. The
18 model predicted AOE have an overall similar pattern, with major plumes below 4 km for the
19 Williams Flats Fire (Figures 11c and 11d). Over the Montana Fires, the GFS-CMAQ predicts a
20 slightly higher PBL height, thus allowing for the fire plume to reach a higher height near the DC-
21 8 cruising altitude. In contrast, the WRF-CMAQ wildfire plumes are slightly lower than the
22 aircraft flight path around 23–24 UTC, which leads to underpredictions of the fire emitted
23 species (Figure 11d).
24

25 An interesting feature in the DIAL observations is the detached plume from 8 km to 10 km
26 altitude (Figure 11a), where some cirrus clouds existed, and the DIAL retrieval could not
27 distinguish whether they are pure clouds or clouds mixed with elevated aerosols above 8km. The
28 cloud-screening image (Figure 11b) mainly showed the enhanced aerosols below 7km and some
29 scattered signals near the high cloud edges. Cloud mixing with aerosols was usual for fire
30 induced clouds, or pyrocumulonimbus (Peterson et al., 2021). Although in this event, the middle-
31 size fires did not show evident of inducing the high-altitude clouds, the indicator of mixed clouds
32 and aerosols in high altitudes still existed: both in-situ measured OA (Figures 10e) and AOE
33 (Figure 10f, 11c, 11d) showed the enhanced aerosols around 25 UTC (or 1 UTC of the next day)
34 above 8km. This elevated plume was generally captured by the GFS-CMAQ simulation, while
35 underestimating its strength (Figure 11c); however, this feature was completely missed in WRF-
36 CMAQ (Figure 11d). Considering the altitude range of the detached plume, the major model
37 disparities are likely due to model convection differences in the free troposphere. To further
38 investigate this impact, Figures 11e and 11f show curtain plots of RH predicted by the two
39 models. GFS-CMAQ yields higher RH at such altitudes (10 km) compared to WRF-CMAQ
40 around 23–24 UTC, indicating that the GFS-CMAQ has a stronger convection. The CMAQ

1 model uses inputted meteorology to diagnose convection activity and drive its ACM2 convection
2 scheme. This convective activity is apparent in GOES-16 satellite images (Figure S3), as more
3 fractional clouds appeared ahead of the northward moving frontal band. Both the GFSv16 and
4 WRF models used here *do not* consider the fire heat feedback effect, and thus their predicted
5 convection and clouds are only driven by the synoptic weather conditions. If such synoptic-to-
6 mesoscale weather models consider wildfire heat feedback effects, their predictions may result in
7 stronger convection and help correct their underpredictions in PBL heights.
8

9 **4.3 Statistical Results of Model Performances for FIREX-AQ**

10 *4.3.1 Meteorological Statistics*

11

12 During the FIREX-AQ field campaign, the DC-8 aircraft performed more than 20 flights over
13 CONUS with detailed observations of various chemical compounds. Tables 2 and 3 show the
14 statistical results of mean bias (MB), normalized mean bias (NMB), root mean square error
15 (RMSE), correlation coefficient (R), and linear regression/slopes for the two models’
16 performance over the western U.S. (west of 110°W) only at low altitudes (<3km ASL) for both
17 non-fire and fire flight segments. The FIREX-AQ aircraft data included the smoke flag to mark
18 the sampling times associated with fire plumes, identified by carbon monoxide and aerosol
19 enhanced over background levels in downwind areas of specific fires. This smoke flag is used to
20 distinguish the flight segments with and without fire influences. Most of these flights departed
21 from Boise, ID, except the 22 July flight that flew from California to Idaho. As a result, they
22 mainly flew over Idaho and its surrounding regions. The GFS tends to have a slightly higher
23 wind speed with positive MB, while WRF has a small negative wind speed bias. Most of the DC-
24 8 flights are during the daytime, and the GFS has a higher daytime wind speed than WRF at low
25 altitudes. The GFS and WRF have very similar temperature predictions. For the RH, the GFS
26 predictions are slightly dryer than those of WRF, especially for non-fire events. The
27 meteorological models do not consider wildfire heat effects, and thus may have (in part) led to
28 slightly warm MB for the non-fire events (Table 2) and slightly cool MB for the fire events
29 (Table 3). Because both the GFSv16 and WRF models have similar MB shifts from an average
30 temperature overprediction (Table 2; non-fire events) to an underprediction (Table 3; wildfire
31 events), we can estimate that the fire effects make roughly a 1–2 Kelvin temperature
32 enhancement to the background along the DC-8 flight paths below 3 km. This estimate assumes
33 that the model temperature biases are generally representative of the western U.S. (west of
34 110°W), and are independent of the averaged flight segments that have different locations and
35 periods in Table 2 and Table 3. Correspondingly, the air masses are dryer in the sampled
36 wildfire plumes, as shown by the large reduction in the RH underpredictions (i.e., negative MBs)
37 from Table 2 to Table 3.
38

39 *4.3.2 Chemical Statistics for Flight Segments without Fire Influences*

40

1 For most chemical species, the two models also have similar performance, indicating that the
2 emissions and chemistry are major driving forces. For flight segments not encountering fire
3 plumes, both models overpredict NO_x, HNO₃, toluene, EC, and ammonium (NH₄⁺), but
4 underestimate PAN, benzene, C₂H₂, SO₂, and submicron sulfate and organic aerosols (OA)
5 (Table 2). The SO₂ and submicron sulfate underprediction may be impacted by underestimated
6 NEIC2016v1 SO₂ emissions over the western U.S. Since point sources, including power plant
7 emissions, are the SO₂ sources, this comparison implies that the point sources for 2019 events
8 have large uncertainties.

9
10 Although the models agree well with NO_y observations, they disproportionately underestimate
11 NO_z (non-NO_x reactive nitrogen species, or NO_y-NO_x) as shown by the regression slopes and
12 MBs. The NO_x and NO_y observations have different missing data, and NO_z is calculated when
13 both NO_x and NO_y observations are available at certain sampling times. Due to the different
14 sample number issues, their observed averages may not exactly match well (averaged NO_x+NO_z
15 ≠ NO_y in observations) though their corresponding modeled relationship are well balanced.
16 Gaseous NO_z species can be split into inorganic NO_z (HNO₃, HONO, HNO₄, NO₃, ClNO₃,
17 N₂O₅ et al.) and organic NO_z, such as PAN, methyl peroxy acetyl nitrate (MPAN) and RNO₃ (or
18 NTR, the other organic nitrate) et al. The precursors of organic NO_z include hydrocarbons. One
19 of the important organic NO_z species is PAN, and both models underestimate PAN for the flight
20 segments without fire influences (Table 2). PAN's carbonyl precursors include acetaldehyde
21 (CH₃CHO) (44% of the global source), methylglyoxal (30%), acetone (7%), and a suite of other
22 isoprene and terpene oxidation products (19%) (Fischer et al., 2014). CH₃CHO and acetone are
23 also underestimated (Table 2), which help explain PAN's underestimation. For the oxidized
24 hydrocarbons, like aldehydes (HCHO, CH₃CHO), their main atmospheric sources come from the
25 oxidation of highly reactive VOCs, including alkanes, alkenes, and aromatics, instead of direct
26 emissions (Parrish et al., 2012). So, the underestimations of HCHO and CH₃CHO are associated
27 with the underestimations of their precursor hydrocarbons, including anthropogenic and biogenic
28 VOCs. Our internal comparison with some limited surface VOC observations indicated that
29 BEIS tends to underpredict biogenic emissions over the U.S. West, e.g. isoprene in Table 2. In
30 this comparison, most anthropogenic hydrocarbons are disproportionately underestimated, except
31 toluene, implying the VOC speciation issue in the NEIC2016v1 anthropogenic emissions (Table
32 2). A previous work discovered that a model overprediction in toluene was also related to the
33 toluene speciation in the NEI emission inventory (Lu et al., 2020). In this comparison, both
34 models tend to underpredict organic NO_z, which is likely caused by the underestimation of
35 certain VOCs.

36
37 Submicron ammonium (NH₄⁺) and nitrate ion (NO₃⁻) are also underestimated by both models
38 during non-fire events (Table 2), suggesting there are NH₃ underestimates due to either
39 insufficient NH₃ emissions or exaggerated NH₃ removal processes. There are, however,
40 overpredictions in the intermediate species nitric acid (HNO₃), indicating the shift of gas-aerosol

equilibrium partition of the nitrate ion. It implies that the HNO_3 accumulates in the atmosphere because the modeled NO_x and inorganic NO_z (such as NO_3) pathways toward the nitrate ion and organic NO_z are reduced due to underestimations of their other precursors (NH_3 and VOCs).

There are underestimations in the VOC and CO concentrations, which contributes to the ozone underestimation during non-fire flight segments (Table 2). These non-fire comparisons also highlight that both models have similar biases due to similar meteorology (Section 4.3.1), and the use of the same anthropogenic emissions (NEIC2016v1), BEIS biogenic emission and chemical models/mechanisms (i.e CMAQv5.3.1). The differences in the two models' bias, error, and correlation/slope are much smaller than their individual magnitudes. As discussed above, VOC speciation in the emission inventory could be one issue, as the model tends to overpredict C_2H_4 , but underestimate C_2H_2 and C_2H_6 etc. Some common biases over certain regions could be related to certain common issues. For instance, some power plants supposed to shut down in the original NEIC 2016 inventory might still emit pollutants during the flight observations, leading to the disagreement of SO_2 .

4.3.3 Chemical Statistics for Flight Segments with Fire Influences

The WRF-CMAQ and GFS-CMAQ models significantly underestimate CO, VOC, HONO, and OA for fire-influenced flight segments at low altitudes (< 3 km) over the western U.S. (Table 3). In conjunction with underestimated GBBEPx emissions during these wildfire events, other possible causes for the average statistical underprediction are the CMAQ model's 12 km horizontal resolution and the flight sampling coverage. Most of the fires that are averaged in the statistics, such as the Horsefly (5.5 km^2 burning area) and Snow Creek Fires (7.3 km^2 burning area), are at a much finer scale than the model grid. Only the largest Williams Flats Fire, with a total burning area of 180 km^2 (Ye et al., 2021), had a comparable horizontal scale to the model resolution.

The DC-8 aircraft had many flight segments near wildfire sources during the fire events in Table 3, and thus dilution of the emissions due to the relatively coarse model resolution may lead to underestimations in the predicted slope for most wildfire emitted pollutants, such as CO and OA (Table 3). The O_3 concentrations are also underestimated; however, the O_3 underpredictions are reduced from the non-fire (Table 2) to fire events (Table 3). Abundant amounts of wildfire emitted NO_x can titrate ozone near the fire source region, and the models likely underestimate these titration effects due to the 12 km model resolution (Figure S4). Thus, the models cannot capture the strong spatial O_3 variability that is observed due to both reduction near source regions and enhancement in downstream areas. Again, for this fire event comparison, both models showed similar behavior and their differences were relatively smaller compared to the overall model biases.

5. Summary and Discussion

The operational NOAA/NWS National Air Quality Forecasting Capability (NAQFC) recently underwent a major upgrade on July 20, 2021. The advanced NAQFC includes the recent Community Multi-scale Air Quality (CMAQ) model version 5.3.1 with CB6 (carbon bond version 6)-Aero7 (version 7 of the aerosol module) chemical mechanism, and is driven by the latest operational Finite Volume Cubed-Sphere (FV3)-Global Forecast System, version 16 (GFSv16) (Campbell et al., 2022). Here we analyze the impacts of the driving meteorological models on CMAQ model performance with the new GFSv16 interpolation-based meteorology versus the commonly-used native-grid Weather Research and Forecasting (WRF) model version 4.0.3 meteorology. The meteorological and chemical analysis includes both 2D ground-based and 3D aircraft measurements during the summer 2019, which encompasses the joint NOAA-NASA Fire Influence on Regional to Global Environments and Air Quality (FIREX-AQ) campaign. As CMAQ has existing mass conservation via adjustments of the contravariant vertical velocity (Otte and Pleim, 2010), the NACC interpolated GFSv16 wind field can be well handled in CMAQ (i.e., GFS-CMAQ).

The different NWS/NOAA operational GFS and commonly chosen WRF physics schemes employed in this study (Table 1) clearly have impacts on temperature, horizontal/vertical wind fields, PBL heights, and the corresponding CMAQ model predictions. During this study period over the U.S. West, both models showed moisture dry bias and temperature warm bias in low altitudes, which could be due to the issue mentioned by Qian et al (2020): the irrigation contribution being neglected (section 3.1), and impacts from soil moisture deficits on surface fluxes in both models. Due to their different physics, GFS has a stronger diurnal variation of PBL height, lower at night and higher during daytime over the U.S. West and Northeast. The differences in the GFS and WRF physics result in a larger impact than the difference between interpolated and native grids on the models' meteorological and air quality predictions, even despite using FDDA to nudge WRF simulation toward the GFSv16 data. Nudging toward observations or including data assimilation may yield different results for the WRF run, though they are not used here. In this study, FDDA nudging used in WRF to avoid growing errors across a continuous 1-month simulation. We note that if this method is turned off, the differences between GFSv16 and WRF predictions would have been even greater. This would further substantiate the dominance of using different model physics and their impacts on CMAQ model predictions. Campbell et al (2022) have the detailed comparisons for interpolated and original fields, and they are very consistent. In this study, we further compare the CMAQ vertical velocity diagnosed from the interpolated GFS horizontal wind, which is very consistent with the original GFS vertical velocity. Overall, the results of this study further corroborate the use of the GFSv16 data and NACC interpolation-based methods (Campbell et al., 2022) for regional CMAQ model applications in the scientific community.

1 Over CONUS, GFS-CMAQ demonstrated lower mean surface ozone (by about 3 ppb) and
2 PM_{2.5} (by about 1 $\mu\text{g}/\text{m}^3$) than WRF-CMAQ in August 2019 (section 3). In the western U.S.,
3 the GFS has a stronger diurnal variability in PBL height and a better performance in nighttime
4 10-m wind speeds compared to WRF. The nighttime difference between these two models tends
5 to be more significant than the corresponding daytime difference. Their difference is also
6 impacted by both vertical/convective (mainly daytime) and upstream advective transport
7 differences in GFS-CMAQ and WRF-CMAQ, which somewhat confounds the impact of
8 different meteorological physics on chemical predictions from region to region. This transport
9 effect is more significant on PM_{2.5} than that on O₃, as O₃ has a shorter lifetime and is more
10 sensitive to local emissions in summer. In this study, neither GFS-CMAQ nor WRF-CMAQ
11 show overwhelming performance advantage over the other, similar to the NMM-CMAQ and
12 ARW-CMAQ comparison in Yu et al. (2012a, 2012b).

13
14 GFS-CMAQ and WRF-CMAQ demonstrated rather similar performance for major chemical
15 variables during both FIREX-AQ non-fire (Table 2) and fire events (Table 3). Both models
16 showed similar biases, indicating that other factors, including emissions, model resolution and
17 chemistry etc. could be more important for the model predictions compared to the meteorological
18 differences. The aircraft data comparison reveals many common issues in both model systems.
19 One critical issue is whether the flight sampling coverage is comparable to the 12 km model
20 resolution, especially for high-gradient fire emission, e.g. the case of 06 August flight (Figure
21 S4). The observation representation issue also exists in other places, such as the near-surface
22 meteorological comparison between AIRNow stations and METAR stations. Emission is the
23 driving force for atmospheric composition concentrations. The comprehensive aircraft
24 measurements help verify that the anthropogenic NEIC2016v1 inventory is overall reasonable,
25 except for SO₂, NH₃ and certain hydrocarbons. The wildfire emission has bigger uncertainties,
26 including the emission intensities, pollutant specification and plume rise, shown by the both
27 models' results.

28
29 The NACC interpolation method is advantageous as it enables using the original meteorological
30 driver directly via interpolation, and avoids running another model such as WRF to drive
31 regional CMAQ applications. It is also faster, and more consistent with the original
32 meteorological model (GFS) than using WRF even with nudging, since WRF's own physics
33 could have stronger impact. In the current NOAA/NCEP operational GFS-CMAQ system,
34 NACC only takes less than 5 minutes to process 72-hour data, which saves enough time for
35 CMAQ to forecast extra 24 hours. These aspects can simultaneously benefit real-time forecasting
36 and retrospective air quality applications in the scientific community. NACC can also adapt to
37 quickly use any regional domain globally, and may also use other global meteorological data
38 including reanalysis products. This helps mitigate the confounding factors of using different
39 model configurations across the myriad of WRF physics options, while alleviating the difficulty
40 in understanding their impacts on air quality predictions. The operational GFSv16 and associated

reanalysis products are well vetted and evaluated across different global agencies and laboratories, and thus are well suited for regional CMAQ applications using NACC. In fact, there is an ongoing project at NOAA to migrate both the GFSv16 data and NACC software to the Amazon Web Services (AWS) Cloud platform to provide a streamlined product for the user to generate the model-ready meteorological data for any regional CMAQ application globally.

Finally, we note that the current operational GFSv16 has all needed meteorological variables to drive CMAQ, and users have option to supply other data (fractional landuse, LAI etc). GFSv16's C768 grid has horizontal resolution from 10.21 km to 14.44 km, which is close to the NAQFC's 12 km horizontal resolution. One barrier of using this NACC approach is that the original-resolution GFS data files with all needed variables are very big even with compression (about 8GB per time step), and may not be accessible for community users. There is an ongoing effort of using cloud storages to solve this issue, and making this method available to the community. Traditional WRF-CMAQ usually starts from commonly available global meteorological data, such as NCEP or ECMWF reanalysis data, which are in relatively coarse resolution, and use WRF to generate all meteorological variables needed by CMAQ in native grid. In some cases, the WRF may become the only available method to drive a finer scale CMAQ model application. WRF's various physics can also be customized for CMAQ simulation over certain regions or under certain meteorological conditions. Both methods have its pros and cons. As shown in this study, GFS and WRF had mixed performance for driving CMAQ, though they are overall similar.

Code and Data Availability.

The FIREX-AQ field campaign data used in this study is in <https://www-air.larc.nasa.gov/cgi-bin/ArcView/firexaq> (last access, 16 May 2022). The NACC code used in this study is publicly available at <https://doi.org/10.5281/zenodo.5507489> and via GitHub at <https://github.com/noaa-oar-arl/NACC.git> (last access: 5 April 2022). The modified CMAQv5.3.1 for GFS-CMAQ is available at <https://doi.org/10.5281/zenodo.5507511> and via GitHub at <https://github.com/noaa-oar-arl/NAQFC> (last access: 5 April 2022).

Author contributions.

YT contributed to the project conceptualization, model run, software, data analysis, visualization, investigation, and writing of the original draft. PCC contributed to software, the model run, data analysis, investigation and draft revision. DT and XZ contributed to wildfire emissions data. BB contributed to software and funding acquisition. FY, JH and HH provided the GFS model data. LP provided the global aerosol model for the lateral boundary condition. PL, RS, AS, JF, IS, JT-D, YJ contributed to project supervision, project administration, and funding acquisition. MY, IB, JF, TR, DB, JS, J-LJ, JC, GD, RM, JH, GH, AR and JD contributed to the FIREX-AQ aircraft data.

Competing interests.

The contact author has declared that neither they nor their co-authors have any competing interests.

Acknowledgements

This research was funded by NOAA's National Air Quality Forecasting Capability (NAQFC) in the National Weather Service Office of Science and Technology Integration (NWS/OSTI).

References:

- Appel, K.W., Bash, J.O., Fahey, K.M., Foley, K.M., Gilliam, R.C., Hogrefe, C., Hutzell, W.T., Kang, D., Mathur, R., Murphy, B.N. and Napelenok, S.L.,. The Community Multiscale Air Quality (CMAQ) model versions 5.3 and 5.3. 1: system updates and evaluation. *Geoscientific Model Development*, 14(5), pp.2867-2897. 2021. <https://doi.org/10.5194/gmd-14-2867-2021>
- Arakawa, A. and Lamb, V.: Computational design of the basic dynamical processes of the UCLA general circulation model, *Methods in Computational Physics*, 17, 173–265, 1977.
- Baker, K.R., Woody, M.C., Tonnesen, G.S., Hutzell, W., Pye, H.O.T., Beaver, M.R., Pouliot, G. and Pierce, T., Contribution of regional-scale fire events to ozone and PM_{2.5} air quality estimated by photochemical modeling approaches. *Atmospheric Environment*, 140, 539-554. <https://doi.org/10.1016/j.atmosenv.2016.06.032>, 2016.
- Briggs, G. A., A Plume Rise Model Compared with Observations, *Journal of the Air Pollution Control Association*, 15:9, 433-438, doi:10.1080/00022470.1965.10468404, 1965.
- Byun, D. W., Dynamically Consistent Formulations in Meteorological and Air Quality Models for Multiscale Atmospheric Studies. Part I: Governing Equations in a Generalized Coordinate System, *Journal of the Atmospheric Science*, 56, 3789–3807. 1999a. [https://doi.org/10.1175/1520-0469\(1999\)056<3789:DCFIMA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<3789:DCFIMA>2.0.CO;2)
- Byun, D.W., Dynamically consistent formulations in meteorological and air quality models for multi-scale atmospheric applications: Part II. Mass conservation issues. *Journal of the Atmospheric Science* 56, 3808–3820, 1999b. [https://doi.org/10.1175/1520-0469\(1999\)056<3808:DCFIMA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<3808:DCFIMA>2.0.CO;2).
- Byun, D.W., Ching, J.K.S.. Science algorithms of the EPA models-3 Community Multiscale Air Quality (CMAQ) modeling system, EPA/600/R-99/030, U.S. EPA. 1999c
- Byun, D. and Schere, K.L.,. Review of the governing equations, computational algorithms, and other components of the Models-3 Community Multiscale Air Quality (CMAQ) modeling system. *Appl. Mech. Rev.* 59: 51–77. <https://doi.org/10.1115/1.2128636>. 2006.
- Campbell, P., Tang, Y., Lee, P., Baker, B., Tong, D., Saylor, R., Stein, A., Huang, J., Huang, H.-C., Strobach, E., McQueen, J., Pan, L., Stajner, I., Sims, J., Tirado-Delgado, J., Jung, Y., Yang, F., Spero, T., and Gilliam, R.: Development and evaluation of an advanced National

1 Air Quality Forecast Capability using the NOAA Global Forecast System version 16, *Geosci.*
2 *Model Dev.* 15, 3281–3313, <https://doi.org/10.5194/gmd-15-3281-2022>, 2022.

3 Chen, F., and Dudhia, J. Coupling an advanced land surface-hydrology model with the Penn
4 State-NCAR MM5 modeling system. Part I: Model implementation and sensitivity. *Monthly*
5 *Weather Review*, 129(4), 569–585. 2001. [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0493(2001)129<0569:CAALSH>2.0.CO;2)
6 [0493\(2001\)129<0569:CAALSH>2.0.CO;2](https://doi.org/10.1175/1520-0493(2001)129<0569:CAALSH>2.0.CO;2)

7 Chen J.-H and S.-J Lin. The remarkable predictability of inter-annual variability of atlantic
8 hurricanes during the past decade. *Geophysical Research Letters*, 38(L11804):6, 2011.
9 <https://doi.org/10.1029/2011GL047629>

10 Chen J-H.and S-J. Lin. Seasonal predictions of tropical cyclones using a 25-km-resolution
11 general circulation model. *J. Climate*, 26(2):380–398, 2013. [https://doi.org/10.1175/JCLI-D-](https://doi.org/10.1175/JCLI-D-12-00061.1)
12 [12-00061.1](https://doi.org/10.1175/JCLI-D-12-00061.1)

13 Clough, S. A., Shephard, M. W., Mlawer, J. E., Delamere, J. S., Iacono, M. J., Cady-Pereira, K.,
14 et al. (2005). Atmospheric radiative transfer modeling: A summary of the AER codes. *Journal*
15 *of Quantitative Spectroscopy & Radiative Transfer*, 91(2), 233–244.
16 <https://doi.org/10.1016/j.jqsrt.2004.05.058>

17 Caputi, D. J., Faloona, I., Trousdell, J., Smoot, J., Falk, N., and Conley, S.: Residual layer ozone,
18 mixing, and the nocturnal jet in California's San Joaquin Valley, *Atmos. Chem. Phys.*, 19,
19 4721–4740, <https://doi.org/10.5194/acp-19-4721-2019>, 2019

20 Dong, X., J. S. Fu, K. Huang, D. Tong, and G. Zhuang. Model development of dust emission
21 and heterogeneous chemistry within the Community Multiscale Air Quality modeling system
22 and its application over East Asia. *Atmos. Chem. Phys.*, 16, 8157–8180. 2016.
23 <https://doi.org/10.5194/acp-16-8157-2016>.

24 Ek, M. B., Mitchell, K. E., Lin, Y., Rogers, E., Grunmann, P., Koren, V., et al. (2003).
25 Implementation of Noah land surface model advances in the National Centers for
26 Environmental Prediction operational mesoscale Eta model. *Journal of Geophysical Research*,
27 108(D22), 8851. <https://doi.org/10.1029/2002JD003296>

28 Fischer, E. V., Jacob, D. J., Yantosca, R. M., Sulprizio, M. P., Millet, D. B., Mao, J., Paulot, F.,
29 Singh, H. B., Roiger, A., Ries, L., Talbot, R. W., Dzepina, K., and Pandey Deolal, S.:
30 Atmospheric peroxyacetyl nitrate (PAN): a global budget and source attribution, *Atmos.*
31 *Chem. Phys.*, 14, 2679–2698, <https://doi.org/10.5194/acp-14-2679-2014>, 2014.

32 Fu, X., S. X. Wang, Z. Cheng, J. Xing, B. Zhao, J. D. Wang, and J. M. Hao, 2014: Source,
33 transport and impacts of a heavy dust event in the Yangtze River Delta, China, in 2011.
34 *Atmos. Chem. Phys.*, 14, 1239–1254, doi:10.5194/acp-14-1239-2014.

35 Grell, G. A., Dudhia, J., and Stauffer, D. R. A description of the fifth-generation Penn
36 State/NCAR Mesoscale Model (MM5). NCAR tech. Note NCAR TN-398-1-STR, 117 pp.
37 1994.

38 Han, J., & Bretherton, C. S., TKE-Based Moist Eddy-Diffusivity Mass-Flux (EDMF)
39 Parameterization for Vertical Turbulent Mixing, *Weather and Forecasting*, 34(4), 869–886.
40 2019. <https://doi.org/10.1175/WAF-D-18-0146.1>

- 1 Harris, L., Chen, X., Putman, W., Zhou, L. and Chen, J.H., A Scientific Description of the GFDL
- 2 Finite-Volume Cubed-Sphere Dynamical Core. <https://doi.org/10.25923/6nhs-5897>. 2021
- 3 Hodyss, D. and N. Nichols, The error of representation: basic understanding, *Tellus A: Dynamic*
- 4 *Meteorology and Oceanography*, 67:1, <https://doi.org/10.3402/tellusa.v67.24822>, 2015.
- 5 Hong, Song-You, Yign Noh, Jimmy Dudhia, A new vertical diffusion package with an explicit
- 6 treatment of entrainment processes. *Mon. Wea. Rev.*, 134, 2318–2341. 2006.
- 7 doi:10.1175/MWR3199.1.
- 8 Houyoux, M.R., Vukovich, J.M., Coats, C.J., Wheeler, N.J.M., Kasibhatla, P.S.,. Emission
- 9 inventory development and processing for the seasonal model for regional air quality
- 10 (SMRAQ) project. *J. Geophys. Res.*, 105(D7), 9079-9090,
- 11 <https://doi.org/10.1029/1999JD900975>. 2000.
- 12 Huang, M., D. Tong, P. Lee, L. Pan, Y. Tang, I. Stajner, R. B. Pierce, J. McQueen, and J. Wang,
- 13 Toward enhanced capability for detecting and predicting dust events in the western United
- 14 States: The Arizona case study. *Atmos. Chem. Phys.*, 15, 12 595–12 610, doi:10.5194/acp-15-
- 15 12595-2015. 2015.
- 16 Kain, J. S., The Kain–Fritsch convective parameterization: An update. *J. Appl. Meteor.*, 43, 170–
- 17 181. doi:10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2, 2004.
- 18 Kim, Y.J., Eckermann, S.D. and Chun, H.Y., An overview of the past, present and future of
- 19 gravity-wave drag parameterization for numerical climate and weather prediction models.
- 20 *Atmosphere-Ocean*, 41(1), 65-98., doi: 10.3137/ao.410105, 2003.
- 21 Krueger, S. K., Q. Fu, K. N. Liou, and H-N. S. Chin. Improvement of an ice-phase microphysics
- 22 parameterization for use in numerical simulations of tropical convection. *Journal of Applied*
- 23 *Meteorology*, 34:281–287, January 1995. <https://doi.org/10.1175/1520-0450-34.1.281>
- 24 Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W.
- 25 D.. Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative
- 26 transfer models. *Journal of Geophysical Research*, 113, D13103. 2008.
- 27 <https://doi.org/10.1029/2008JD009944>
- 28 Jimenez, P. A., Dudhia, J., Gonzalez-Rouco, J. F., Navarro, J., Montavez, J. P., & Garcia-
- 29 Bustamante, E. (2012). A revised scheme for the WRF surface layer formulation. *Monthly*
- 30 *Weather Review*, 140(3), 898–918. <https://doi.org/10.1175/MWR-D-11-00056.1>
- 31 Lin, Y.-L., R. D. Farley, and H. D. Orville. Bulk parameterization of the snow field in a cloud
- 32 model. *J. Climate Appl. Meteor.*, 22:1065–1092, 1983. [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0450(1983)022<1065:BPOTSF>2.0.CO;2)
- 33 [0450\(1983\)022<1065:BPOTSF>2.0.CO;2](https://doi.org/10.1175/1520-0450(1983)022<1065:BPOTSF>2.0.CO;2)
- 34 Lord, S. J., H.E. Willoughby, and J.M. Piotrowicz. Role of a parameterized ice-phase
- 35 microphysics in an axisymmetric, nonhydrostatic tropical cyclone model. *J. Atmos. Sci.*,
- 36 41(19):2836–2848, October 1984. [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0469(1984)041<2836:ROAPIP>2.0.CO;2)
- 37 [0469\(1984\)041<2836:ROAPIP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1984)041<2836:ROAPIP>2.0.CO;2)
- 38 Lu, Q., Murphy, B. N., Qin, M., Adams, P. J., Zhao, Y., Pye, H. O. T., Efsthathiou, C., Allen, C.,
- 39 and Robinson, A. L.: Simulation of organic aerosol formation during the CalNex study:
- 40 updated mobile emissions and secondary organic aerosol parameterization for intermediate-

volatility organic compounds, *Atmos. Chem. Phys.*, 20, 4313–4332,
<https://doi.org/10.5194/acp-20-4313-2020>, 2020

Luecken, D. J., Yarwood, G., and Hutzell, W. T.: Multipollutant modeling of ozone, reactive nitrogen and HAPs across the continental US with CMAQ-CB6, *Atmos. Environ.*, 201, 62–72, <https://doi.org/10.1016/j.atmosenv.2018.11.060> , 2019.

Mlawer, E. J., Taubman S. J., Brown P. D., Iacono M. J., and Clough S. A., Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave. *J. Geophys. Res.*, 102, 16663–16682. 1997. <https://doi.org/10.1029/97JD00237>.

Monin, A. S., & Obukhov, A. M. Basic laws of turbulent mixing in the surface layer of the atmosphere (in Russian). Contribution Geophysics Institute, Academy of Sciences USSR, 151, 163–187. 1954.

Morrison, H., G. Thompson, V. Tatarskii, 2009: Impact of Cloud Microphysics on the Development of Trailing Stratiform Precipitation in a Simulated Squall Line: Comparison of One- and Two-Moment Schemes. *Mon. Wea. Rev.*, 137, 991–1007.
<https://doi.org/10.1175/2008MWR2556.1>

National Emissions Inventory Collaborative (2019). 2016v1 Emissions Modeling Platform. Retrieved from <http://views.cira.colostate.edu/wiki/wiki/10202>.

Ott, E., Hunt, B.R., Szunyogh, I., Zimin, A.V., Kostelich, E.J., Corazza, M., Kalnay, E., Patil, D.J. and Yorke, J.A., 2004. A local ensemble Kalman filter for atmospheric data assimilation. *Tellus A: Dynamic Meteorology and Oceanography*, 56(5), pp.415-428.
<https://doi.org/10.3402/tellusa.v56i5.14462>

Otte, T. L., J E. Pleim, and G Pouliot. PREMAQ: A new pre-processor to cmaq for air-quality forecasting. Presented at 2004 Models-3 Conference, Chapel Hill, NC, October 18-20, 2004.

Otte, T. L. and Pleim, J. E.: The Meteorology-Chemistry Interface Processor (MCIP) for the CMAQ modeling system: updates through MCIPv3.4.1, *Geosci. Model Dev.*, 3, 243–256, <https://doi.org/10.5194/gmd-3-243-2010>, 2010.

Pan, L., Kim, H., Lee, P., Saylor, R., Tang, Y., Tong, D., Baker, B., Kondragunta, S., Xu, C., Ruminski, M. G., Chen, W., McQueen, J., and Stajner, I.: Evaluating a fire smoke simulation algorithm in the National Air Quality Forecast Capability (NAQFC) by using multiple observation data sets during the Southeast Nexus (SENEX) field campaign, *Geosci. Model Dev.*, 13, 2169–2184, <https://doi.org/10.5194/gmd-13-2169-2020> , 2020.

Parrish, D. D., Ryerson, T. B., Mellqvist, J., Johansson, J., Fried, A., Richter, D., Walega, J. G., Washenfelter, R. A., de Gouw, J. A., Peischl, J., Aikin, K. C., McKeen, S. A., Frost, G. J., Fehsenfeld, F. C., and Herndon, S. C.: Primary and secondary sources of formaldehyde in urban atmospheres: Houston Texas region, *Atmos. Chem. Phys.*, 12, 3273–3288, <https://doi.org/10.5194/acp-12-3273-2012>, 2012.

Peterson, D.A., Fromm, M.D., McRae, R.H., Campbell, J.R., Hyer, E.J., Taha, G., Camacho, C.P., Kablick, G.P., Schmidt, C.C. and DeLand, M.T., Australia’s Black Summer pyrocumulonimbus super outbreak reveals potential for increasingly extreme stratospheric

smoke events. *npj Climate and Atmospheric Science*, 4(1), 2021.
<https://doi.org/10.1038/s41612-021-00192-9>

Poggio, L., de Sousa, L. M., Batjes, N. H., Heuvelink, G. B. M., Kempen, B., Ribeiro, E., and Rossiter, D.: SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty, *SOIL*, 7, 217–240, 2021. <https://doi.org/10.5194/soil-7-217-2021>

Powers, J.G., Klemp, J.B., Skamarock, W.C., Davis, C.A., Dudhia, J., Gill, D.O., Coen, J.L., Gochis, D.J., Ahmadov, R., Peckham, S.E. and Grell, G.A.,. The weather research and forecasting model: Overview, system efforts, and future directions. *Bulletin of the American Meteorological Society*, 98(8), pp.1717-1737. 2017. <https://doi.org/10.1175/BAMS-D-15-00308.1>

Putman, W. M. and S.-J. Lin, Finite-volume transport on various cubed-sphere grids, *Journal of Computational Physics*, 227(1), 55-78, <https://doi.org/10.1016/j.jcp.2007.07.022>, 2007.

Qian, Y., Yang, Z., Feng, Z., Liu, Y., Gustafson, W.I., Berg, L.K., Huang, M., Yang, B. and Ma, H.Y., Neglecting irrigation contributes to the simulated summertime warm-and-dry bias in the central United States. *npj Clim Atmos Sci* 3, 31. <https://doi.org/10.1038/s41612-020-00135-w>, 2020

Rolph, G. D., Draxler, R. R., Stein, A. F., Taylor, A., Ruminski, M. G., Kondragunta, S., Zeng, J., Huang, H. C., Manikin, G., McQueen, J. T., and Davidson, P. M.: Description and Verification of the NOAA Smoke Forecasting System: The 2007 Fire Season, *Weather Forecast.*, 24, 361–378. <https://doi.org/10.1175/2008waf2222165.1>, 2009.

Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Huang, X.-Yu.. A Description of the Advanced Research WRF Model Version 4. NCAR Tech Note, NCAR/TN–556+STR. doi:10.5065/1dfh-6p97, 2021.

Skamarock, W.C., Snyder, C., Klemp, J.B. and Park, S.H., vertical resolution requirements in atmospheric simulation. *Monthly Weather Review*, 147(7), 2641-2656. <https://doi.org/10.1175/MWR-D-19-0043.1>, 2019.

Shi, Y., Hu, F., Xiao, Z., Fan, G. and Zhang, Z., Comparison of four different types of planetary boundary layer heights during a haze episode in Beijing. *Science of the total environment*, 711, <https://doi.org/10.1016/j.scitotenv.2019.134928>, 2020.

Sillman, S., He, D., Cardelino, C., and Imhoff, R. E.: The use of photochemical indicators to evaluate ozone-NO_x –hydrocarbon sensitivity: Case studies from Atlanta, New York, and Los Angeles, *J. Air Waste Manage. Assoc.*, 47, 1030–1040, doi:10.1080/10962247.1997.11877500, 1997.

Sofiev, M., Ermakova, T., and Vankevich, R.: Evaluation of the smoke-injection height from wild-land fires using remote-sensing data, *Atmos. Chem. Phys.*, 12, 1995–2006, <https://doi.org/10.5194/acp-12-1995-2012>, 2012.

Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M. A., Mitchell, K., et al. Implementation and verification of the unified NOAA land surface model in the WRF model. Paper Presented at the 20th Conference on Weather Analysis and Forecasting/16th Conference on Numerical Weather Prediction, Seattle, WA, 2004.

- Wang, S., Coggon, M.M., Gkatzelis, G.I., Warneke, C., Bourgeois, I., Ryerson, T., Peischl, J., Veres, P.R., Neuman, J.A., Hair, J. and Shingler, T., Chemical Tomography in a Fresh Wildland Fire Plume: a Large Eddy Simulation (LES) Study. *Journal of Geophysical Research: Atmospheres*, 126(18), <https://doi.org/10.1029/2021JD035203>. 2021
- Wiedinmyer, C., Akagi, S.K., Yokelson, R.J., Emmons, L.K., Al-Saadi, J.A., Orlando, J.J. and Soja, A.J., 2011. The Fire INventory from NCAR (FINN): A high resolution global model to estimate the emissions from open burning. *Geoscientific Model Development*, 4(3), pp.625-641. doi:10.5194/gmd-4-625-2011.
- Yamartino, R. J., Nonnegative, conserved scalar transport using grid-cell-centered, spectrally constrained Blackman cubics for applications on a variable-thickness mesh. *Mon. Wea. Rev.* 121, 753-763, 1999. [https://doi.org/10.1175/1520-493\(1993\)121<0753:NCSTUG>2.0.CO;2](https://doi.org/10.1175/1520-493(1993)121<0753:NCSTUG>2.0.CO;2)
- Yang, F., Tallapragada, V., Kain, J. S., Wei, H., Yang, R., Yudin, V. A., Moorthi, S., Han, J., Hou, Y. T., Wang, J., Treadon, R., and Kleist, D. T. (2020). Model Upgrade Plan and Initial Results from a Prototype NCEP Global Forecast System Version 16. 2020 AMS Conference, Boston, MA. <https://ams.confex.com/ams/2020Annual/webprogram/Paper362797.html>
- Yarwood, G., Jung, J., Whitten, G.Z., Heo, G., Mellberg, J. and Estes, M., Updates to the Carbon Bond mechanism for version 6 (CB6). In 9th Annual CMAS Conference, Chapel Hill, NC (pp. 11-13). 2010.
- Yarwood, Y., Sakulyanontvittaya, T., Nopmongkol, O., and Koo, K.: Ozone depletion by bromine and iodine over the Gulf of Mexico, final report for the Texas Commission on Environmental Quality, available at: <https://www.tceq.texas.gov/assets/public/implementation/air/am/contracts/reports/pm/5821110365FY1412-20141109-enviro-bromine.pdf> (last access: 3 May 2021), November 2014.
- Ye, X., Arab, P., Ahmadov, R., James, E., Grell, G. A., Pierce, B., Kumar, A., Makar, P., Chen, J., Davignon, D., Carmichael, G. R., Ferrada, G., McQueen, J., Huang, J., Kumar, R., Emmons, L., Herron-Thorpe, F. L., Parrington, M., Engelen, R., Peuch, V.-H., da Silva, A., Soja, A., Gargulinski, E., Wiggins, E., Hair, J. W., Fenn, M., Shingler, T., Kondragunta, S., Lyapustin, A., Wang, Y., Holben, B., Giles, D. M., and Saide, P. E.: Evaluation and intercomparison of wildfire smoke forecasts from multiple modeling systems for the 2019 Williams Flats fire, *Atmos. Chem. Phys.*, 21, 14427–14469, <https://doi.org/10.5194/acp-21-14427-2021>, 2021.
- Yu, S., Mathur, R., Pleim, J., Pouliot, G., Wong, D., Eder, B., Schere, K., Gilliam, R. and Rao, S.T. Comparative evaluation of the impact of WRF/NMM and WRF/ARW meteorology on CMAQ simulations for PM_{2.5} and its related precursors during the 2006 TexAQs/GoMACCS study. *Atmospheric Chemistry and Physics*, 12(9), doi:10.5194/acp-12-4091-2012, 2012a.
- Yu, S., Mathur, R., Pleim, J., Pouliot, G., Wong, D., Eder, B., Schere, K., Gilliam, R. and Rao, S.T., 2012. Comparative evaluation of the impact of WRF–NMM and WRF–ARW meteorology on CMAQ simulations for O₃ and related species during the 2006

- 1 TexAQS/GoMACCS campaign. *Atmospheric Pollution Research*, 3(2), pp.149-162, doi:
2 10.5094/APR.2012.015, 2012b.
- 3 Zhang, X., and Kondragunta, S., 2006, Estimating forest biomass in the USA using generalized
4 allometric model and MODIS land data, *Geographical Research Letter*, 33, L09402,
5 doi:10.1029/2006GL025879.
- 6 Zhang, X., Kondragunta, S., and Quayle, B. Estimation of biomass burned areas using multiple-
7 satellite-observed active fires. *IEEE Transactions on Geosciences and Remote Sensing*, 49:
8 4469-4482, 10.1109/TGRS.2011.2149535, 2011.
- 9 Zhuang, J., Jacob, D. J., and Eastham, S. D.: The importance of vertical resolution in the free
10 troposphere for modeling intercontinental plumes, *Atmos. Chem. Phys.*, 18, 6039–6055,
11 <https://doi.org/10.5194/acp-18-6039-2018>, 2018.
- 12

1 Table 1. The two meteorological datasets used in this study

Model Settings	FV3-GFSv16/NACC	WRF-ARW/MCIP
Domain	Global C768L127 (~ 13 km horizontal resolution in 6 cubic spherical tiles , 127 vertical layers up to 80km), interpolated to the 12km CONUS domain with 35-layers up to about 14km (60hPa)	12km CONUS 35 vertical layers up to 100hPa
Dynamic core	Finite Volume 3, non-hydrostatic (Putman and Lin, 2007)	WRF-ARW dynamic in hybrid vertical coordinate (Skamarock et al., 2021)
Initial condition	FV3-GFSv16 analysis (GDAS) using the local ensemble Kalman filter (LETKF) (Ott et al., 2004) with 4-dimensional incremental analysis update (4D-IAU)	FV3-GFSv16 analysis (GDAS)
Lateral Boundary Condition	N/A	FV3-GFSv16 analysis (GDAS)
Cloud Microphysics	GFDL six-category cloud microphysics scheme (Lin et al., 1983; Lord et al., 1984; Krueger et al., 1995; Chen and Lin, 2011; Chen and Lin, 2013)	Morrison 2-moment scheme (Morrison et al., 2009)
PBL Physics Scheme	Scale-aware (sa) turbulent kinetic energy (TKE) -based moist eddy-diffusivity mass-flux (EDMF) (sa-TKE-EDMF) (Han and Bretherton, 2019)	Yonsei University Scheme (Hong et al., 2006)
Shallow/Deep Cumulus Parameterization	SAS Scheme (Han et al. 2011; 2017)	Kain Fritsch multiscale (Kain, 2004)
Shortwave and Longwave Radiation	RRTMG (Mlawer et al. 1997; Clough et al. 2005; Iacono et al. 2008)	RRTMG (Iacono et al. 2008).
Land Surface Model	Noah Land Surface Model (Chen and Dudhia 2001; Ek et al. 2003; Tewari et al. 2004)	Noah (Tewari et al., 2004)
Surface Layer	Monin-Obukhov (Monin-Obukhov 1954; Grell et al. 1994; Jimenez et al. 2012)	Revised MM5 Scheme (Jimenez et al., 2012)
Other treatment		FDDA nudging is enabled for temperature and specific humidity whole domain, and for wind components (U, V) outside the PBL.

2


Variables	Obs Mean	GFS-CMAQ					WRF-CMAQ				
		MB	NMB	RMSE	R	Slope	MB	NMB	RMSE	R	Slope
Temperature (K)	295	0.979	0.332	2.04	0.988	1.13	1.16	0.393	2.28	0.989	1.17
RH (%)	35.6	-7.3	-20.5	11.8	0.781	0.717	-6.05	-17	12.6	0.677	0.598
Wind Speed (m/s)	4.81	0.758	15.8	3.25	0.432	0.473	-1.11	-23.1	2.4	0.666	0.524
O ₃ (ppbv)	57.9	-10.7	-18.5	15	0.651	0.34	-10.4	-17.9	14.1	0.717	0.413
CO (ppbv)	134	-37.6	-28	53.2	0.654	0.573	-37.1	-27.7	52.9	0.652	0.572
NO _x (ppbv)	1.11	0.507	45.6	2.9	0.704	1.15	0.345	31.1	2.86	0.695	1.12
NO _y (ppbv)	2.56	-0.0418	-1.63	3.07	0.743	0.892	0.055	2.15	3.14	0.724	0.86
NO _z (ppbv)	1.63	-0.465	-28.6	1.17	0.782	0.553	-0.125	-7.66	1.08	0.788	0.721
HONO (ppbv)	0.00432	0.012	279	0.0438	0.379	0.444	0.0134	311	0.0487	0.358	0.48
HNO ₃ (ppbv)	0.291	0.154	53.1	0.421	0.683	1.34	0.337	116	0.65	0.708	1.89
PAN (ppbv)	0.399	-0.251	-63	0.416	0.675	0.221	-0.222	-55.6	0.386	0.681	0.284
NH ₃ (ppbv)	3.55	-0.801	-22.6	5.26	0.0481	0.038	-1.58	-44.5	4.37	0.304	0.155
C ₂ H ₄ (ppbv)	0.121	0.0582	48.1	0.189	0.702	0.869	0.0385	31.9	0.187	0.682	0.836
C ₂ H ₂ (ppbv)	0.146	-0.0734	-50.3	0.137	0.784	0.496	-0.0696	-47.7	0.137	0.771	0.494
SO ₂ (ppbv)	0.342	-0.235	-68.8	0.567	0.0238	0.00835	-0.221	-64.5	0.568	-1.26×10 ⁻³	-0.00047
Acetone (ppbv)	2.74	-2.28	-83.1	2.45	0.686	0.192	-2.2	-80.4	2.38	0.668	0.199
HCHO (ppbv)	2.1	-0.972	-46.4	1.26	0.559	0.447	-0.909	-43.4	1.25	0.513	0.442
CH ₃ CHO (ppbv)	0.736	-0.326	-44.2	0.538	0.647	0.386	-0.349	-47.4	0.554	0.643	0.38
Benzene (ppbv)	0.0449	-0.0193	-43	0.057	0.398	0.385	-0.0191	-42.6	0.0564	0.397	0.375
Toluene (ppbv)	0.039	0.0409	105	0.153	0.759	1.74	0.0352	90.1	0.14	0.762	1.63
Isoprene (ppbv)	0.073	0.0361	49.4	0.174	0.6	0.838	0.00661	9.06	0.145	0.648	0.797
EC (μg/std m ³)	0.108	0.191	177	0.572	0.518	2.09	0.228	211	0.609	0.455	1.88
OA (μg/std m ³)	10.9	-7.15	-65.7	9.72	0.565	0.263	-6.48	-59.5	9.45	0.495	0.243
Sulfate (μg/std m ³)	1.31	-0.781	-59.7	1.11	0.0856	0.0188	-0.773	-59	1.11	0.0322	0.00677
NH ₄ ⁺ (μg/std m ³)	0.745	-0.615	-82.5	0.805	0.416	0.103	-0.596	-79.9	0.778	0.509	0.145
Nitrate (μg/std m ³)	1.22	-1.08	-88.1	1.49	0.562	0.229	-1.04	-85.3	1.45	0.57	0.279
AOE (/Mm)	54.5	-29.3	-53.8	47	0.593	0.227	-27.4	-50.2	45.9	0.588	0.227

Table 2. Statistics of the two models compared to the observation for DC-8 flight segments without fire influences below 3km (ASL) over west of -100°W. All aerosols are in submicron. The normalized mean bias (NMB) is in unit %.


Variables	Obs Mean	GFS-CMAQ					WRF-CMAQ				
		MB	NMB	RMSE	R	Slope	MB	NMB	RMSE	R	Slope
Temperature (K)	287	-0.389	-0.135	0.702	0.995	1.01	-0.688	-0.24	0.863	0.997	1.04
RH (%)	27.8	-0.761	-2.74	7.84	0.712	0.553	4.3	15.5	11.1	0.556	0.534
Wind Speed (m/s)	5.42	0.766	14.1	2.16	0.612	0.616	-0.811	-15	2.12	0.604	0.556
O ₃ (ppbv)	55.7	-6.61	-11.9	11.8	0.587	0.262	-7.01	-12.6	11.5	0.653	0.346
CO (ppbv)	486	-377	-77.6	873	0.596	0.0347	-383	-78.8	883	0.442	0.0242
NO _x (ppbv)	2.63	0.06	2.28	6.41	0.465	0.231	-0.619	-23.5	7.02	0.31	0.153
NO _y (ppbv)	7.32	-4.19	-57.3	13.3	0.507	0.123	-4.66	-63.7	14.2	0.31	0.073
NO _z (ppbv)	5.7	-4.8	-84.3	10.2	-0.189	-0.0106	-4.68	-82	10.2	-0.204	-0.0121
HONO (ppbv)	0.283	-0.274	-96.8	1.18	0.355	0.0043	-0.274	-96.8	1.18	0.291	0.00457
HNO ₃ (ppbv)	0.148	0.148	99.7	0.256	0.532	1.07	0.179	121	0.28	0.402	0.768
PAN (ppbv)	0.971	-0.793	-81.7	1.63	0.27	0.0195	-0.765	-78.8	1.61	0.279	0.026
NH ₃ (ppbv)	17.7	-12.3	-69.3	28.3	0.379	0.0654	-13.7	-77.4	29.6	0.232	0.0386
C ₂ H ₄ (ppbv)	4.5	-4.34	-96.3	10.2	0.421	0.00498	-4.36	-96.8	10.2	0.14	0.0018
C ₂ H ₂ (ppbv)	1.04	-1.01	-96.9	2.08	0.534	0.00866	-1.01	-97	2.09	0.363	0.00623
SO ₂ (ppbv)	0.699	-0.322	-46.1	1.38	0.589	0.198	-0.392	-56.1	1.5	0.429	0.132
Acetone (ppbv)	3.54	-3.2	-90.3	4.56	0.13	0.00862	-3.18	-89.7	4.55	0.135	0.0112
HCHO (ppbv)	8.17	-7.13	-87.3	17.8	0.232	0.0062	-7.19	-88	17.8	0.119	0.00303
CH ₃ CHO (ppbv)	3.65	-3.18	-87.4	9.13	0.186	0.00547	-3.21	-88	9.2	-0.027	-0.00097
Benzene (ppbv)	0.683	-0.67	-98.1	1.84	0.54	0.00432	-0.672	-98.3	1.84	0.367	0.00275
Toluene (ppbv)	0.451	-0.436	-96.6	1.36	0.402	0.00491	-0.438	-97	1.36	0.195	0.00245
Isoprene (ppbv)	0.095	-7.9×10 ⁻³	-8.29	0.234	0.123	0.0579	-0.033	-34.7	0.242	-0.014	-0.00541
EC (μg/std m ³)	1.89	-0.53	-28	3.28	0.612	0.295	-0.787	-41.6	3.7	0.448	0.195
OA (μg/std m ³)	156	-146	-93.4	420	0.612	0.0174	-147	-94.2	423	0.472	0.0122
Sulfate (μg/std m ³)	0.791	-0.116	-14.7	0.676	0.415	0.184	-0.214	-27.1	0.728	0.322	0.13
NH ₄ ⁺ (μg/std m ³)	1	-0.591	-59.1	0.931	0.767	0.351	-0.615	-61.5	0.956	0.729	0.359
Nitrate (μg/std m ³)	1.7	-0.56	-32.9	1.47	0.805	0.613	-0.634	-37.2	1.59	0.774	0.599
AOE (/Mm)	391	-350	-89.3	994	0.688	0.027	-357	-91.1	1010.	0.532	0.0152

Table 3, same as Table 2 except for the wildfire affected flight segments.


a) GFS mean 2m temperature bias (K), 08/2019




b) WRF mean 2m temperature bias (K), 08/2019




c) GFS mean 2m specific humidity bias (g/kg), 08/2019



d) WRF mean 2m specific humidity bias (g/kg), 08/2019



e) GFS mean 10m wind speed bias (m/s), 08/2019



f) WRF mean 10m wind speed bias (m/s), 08/2019





Figure 1. GFS and WRF surface meteorological biases for METAR (METeorological Aerodrome Report) stations averaged over August, 2019

a) Monthly Mean PBL Difference (GFS-WRF) at 18UTC



b) Monthly Mean PBL Difference (GFS-WRF) at 06UTC





Figure 2. Monthly mean PBL height difference (GFS-WRF) for daytime (a) and nighttime (b), August, 2019.




Figure 3. The WRF and GFS time-series comparison over AIRNow stations over the U.S. West and Northeast for 2m temperature (a, b), 10m wind speed (c, d), and PBL height (e,f).




Figure 4. Same as Figure 3 but for ozone (a, b) and PM2.5 (c, d).




Figure 5. Monthly mean surface ozone predictions by GFS-CMAQ (left plots) and WRF-CMAQ (right plots) for daytime (top plots) and nighttime (bottom plots) compared to the corresponding AIRNow observations, August, 2019










Figure 6, same as figure 5 but for surface PM2.5


a) FIREX-AQ DC-8 Flight Path on 07/22




b)




c)



d)



e)



f)





Figure 7. Modeled meteorological variables compared with observations for the DC-8 flight on 22 July, 2019 (b to f). The plot a shows the flight path colored in altitudes above sea level with UTC time in red text. Base map credits: © OpenStreetMap contributors 2022. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.

a) GFS-CMAQ predicted vertical velocity along the DC-8 flight path, 22, July, 2019



b) WRF-CMAQ predicted vertical velocity along the DC-8 flight path, 22, July, 2019




Figure 8, Curtain plots of the vertical velocity (W) predicted by GFS-CMAQ (a) and WRF-CMAQ (b) along the DC-8 flight on 22 July, 2019. The colored dots showed the DC-8 measured vertical velocities. The solid lines showed the predicted PBL heights of these two models.




Figure 9. Model predicted chemical concentrations compared with observations along with the DC-8 flight on 22 July, 2019




Figure 10. The DC-8 flight path (a), and model-observation comparisons for O_3 (b), CO (c), NO_2 (d), submicron organic aerosol (OA) (e) and aerosol optical extinction coefficient (AOE) at wavelength of 550nm (f) on 06 August, 2019. Base map credits: © OpenStreetMap contributors 2022. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.




Figure 11, The Differential Absorption High Spectral Resolution Lidar (DIAL-HSRL) retrieved aerosol backscatter coefficient (ABC) at 532nm wavelength in unit /km/steradian (a) and cloud screened one (b); curtain plots of the AOE (b, c) and relative humidity (RH) (d, e) predicted by GFS-CMAQ (left) and WRF-CMAQ (right) along the DC-8 flight on 06 August, 2019. The colored dots showed the corresponding measured values. The solid lines showed the predicted PBL heights of these two models.