

# An update on dissolved methane distribution in the North subtropical Atlantic Ocean

Anna Kolomijeca<sup>1,2</sup>, Lukas Marx<sup>3</sup>, Sarah Reynolds<sup>3</sup>, Thierry Cariou<sup>1,4</sup>, Edward Mawji<sup>5</sup>, Cedric Boulart<sup>1</sup>

<sup>1</sup>UMR 7144 CNRS Sorbonne Université, Station Biologique de Roscoff, 29680 Roscoff, France

5 <sup>2</sup>MARUM, Center for Marine Environmental Sciences, D-28359 Bremen, Germany

<sup>3</sup>School of the Environment, Geography and Geosciences, University of Portsmouth, PO1 2UP Portsmouth, UK

<sup>4</sup>IRD, UAR191, Instrumentation, Moyens Analytiques, Observatoires en Géophysique et Océanographie (IMAGO), Technopôle de Brest-Iroise, 29280 Plouzané, France

<sup>5</sup>National Oceanography Centre, European Way, Southampton, SO14 3ZH, UK

10

*Correspondence to:* Anna Kolomijeca (akolomijeca@marum.de)

**Abstract.** Methane (CH<sub>4</sub>) is a potent greenhouse gas and plays a significant role in recent increasing global temperatures. The oceans are a natural source of methane contributing to atmospheric methane concentrations, yet our understanding of the oceanic methane cycle is poorly constrained. Accumulating evidence indicates that a significant part of oceanic CH<sub>4</sub> is produced in oxygenated surface waters as a by-product of phytoplanktonic activity. This study focussed on the subtropical North Atlantic Ocean (26°N 80°W and 26°N 18°W) where the distribution of dissolved CH<sub>4</sub> concentrations and associated air-sea fluxes during winter 2020 were investigated. Water samples from 64 stations were collected from the upper water column up to depths of 400 m. The upper oxic mixed-layer was oversaturated in dissolved CH<sub>4</sub> with concentrations ranging between 3-7 nmol/l with the highest concentrations of 7-10 nmol/l found to the east of the transect, consistent with other subtropical regions of the world's oceans. The high anomalies of dissolved CH<sub>4</sub> were found to be associated with phosphate depleted waters and regions where the abundance of the ubiquitous pico-cyanobacteria, *Synechococcus* and *Prochlorococcus* were elevated. Although other phytoplanktonic phyla cannot be excluded, this suggest that cyanobacteria contribute to the release of CH<sub>4</sub> in this region. The calculation of air-sea fluxes further confirmed that the subtropical North Atlantic Ocean as a source of CH<sub>4</sub>. This study provides evidence to corroborate the key role that pico-cyanobacteria play in helping to explain the oversaturation of CH<sub>4</sub> found in surface mixed layer of the open ocean, otherwise known as the 'Ocean Methane Paradox'.

15  
20  
25

## 1. Introduction

Since the industrial revolution, the average global temperature has increased at the fastest rate in recorded history, primarily driven by growing emissions of greenhouse gases (GHGs). Among them, methane (CH<sub>4</sub>) is considered the second largest contributor to Earth warming, after carbon dioxide (CO<sub>2</sub>), with an atmospheric concentration of 1,866 ppb (IPCC, 2021). Over the last 50 years, CH<sub>4</sub> concentrations have increased by 20 % (Karl et al., 2008); (Rhee et al., 2009) – and are expected to rise further, by approximately 2 % per year (Dang and Li, 2018).

30

Oceans were generally considered to be a minor contributor to the total global CH<sub>4</sub> budget, yet recent calculations indicate that oceans could emit 6 to 17 Tg CH<sub>4</sub>/yr, i.e. 1 to 10 % of the total natural emissions (Weber et al., 2019). This large variability reflects the great uncertainty on the contribution of natural sources due to a lack of data and understanding of the sources and controls on oceanic CH<sub>4</sub> emissions.

The marine flux of CH<sub>4</sub> results from the balance between production and oxidation processes, as for instance, the microbial anaerobic oxidation of CH<sub>4</sub> (AOM) in sediments significantly decreases CH<sub>4</sub> fluxes to the atmosphere, thus representing an important carbon sink in the ocean (Oppo et al., 2020). In fact, the marine methane release is dominated by shallow coastal environments including estuaries (up to 75 % (Weber et al., 2019)). In marine sediments CH<sub>4</sub> can be released via microbial anaerobic methanogenesis as a consequence of the degradation of organic matter; it accumulates in the sediment, eventually forming gas hydrates, which may then be released into the overlying water column. Under the influence of pressure and temperature CH<sub>4</sub> diffuses out of the sediment and ebullition carries CH<sub>4</sub> to the atmosphere (Weber et al., 2019).

In the open oxygenated waters, the primary mechanism controlling the CH<sub>4</sub> emissions is aerobic methanotrophy that converts CH<sub>4</sub> into CO<sub>2</sub> (Weber et al., 2019). However, this process could be overcome by *in-situ* production of CH<sub>4</sub> in upper oxic waters that can significantly contribute to marine CH<sub>4</sub> fluxes to the atmosphere. Typical CH<sub>4</sub> depth distribution in the open ocean indicates a general oversaturation in the mixed layer (Reeburgh, 2007a). In the surface waters of the Pacific Ocean (Weller et al., 2013), the Indian Ocean (Bui et al., 2018) and the Atlantic Ocean, values of 2-5 nmol/l and a maximum of 10 nmol/l were measured near the surface (Scranton and Brewer, 1977). These observations make the global ocean a net source of CH<sub>4</sub> to the atmosphere with a weighted supersaturation of 120 % (Kock and Bange, 2007). Exemptions are the Southern Ocean and the central Arctic Ocean, where surface waters are undersaturated in CH<sub>4</sub>, either due to extensive upwelling supplying CH<sub>4</sub>-depleted water to the surface (Bui et al., 2018), or the limitation of air-sea exchanges by ice cover (Weber et al., 2019).

The oversaturation of the surface mixed layer, commonly known as the “ocean methane paradox” (OMP), was explained by the result of methanogenic activity from *Archaea* living within anaerobic cavities of zooplankton guts and anaerobic micro-environments inside sinking particles of organic matter (Reeburgh, 2007a). Initially, only microbes from the *Archaea* domain were thought to have the capability of producing CH<sub>4</sub> under strict anaerobic conditions. Although one cannot exclude this process to explain the methane paradox (Schmale et al., 2018) (Stawiarski et al., 2019), an increasing number of studies have revealed relationships between CH<sub>4</sub> anomalies in surface waters and the presence of certain phytoplanktonic groups such as coccolithophores (Lenhart et al., 2016a) and cyanobacteria (Bižić et al., 2020)

Cyanobacteria are ubiquitous to every aqueous environment on Earth, both in illuminated and dark water bodies (Percival and Williams, 2013). In the open ocean, small-sized picophytoplankton of the genera *Prochlorococcus* spp. and *Synechococcus* spp. account for ~80 % of the total phytoplanktonic chlorophyll *a* (Hickman et al., 2010) and could represent up to 8.5 and 16.7 % of the ocean net primary production (ONPP), respectively (Flombaum et al., 2013). Generally, nutrient limitation sets the upper limit for primary production and the distribution of *Prochlorococcus* and *Synechococcus*; The oligotrophic subtropical North Atlantic in fact is nitrogen (N) - phosphorus (P) co-limited (Harvey et al., 2013), hence cyanobacteria need to acquire these nutrients from alternative sources. New nitrogen to the surface ocean is introduced via biological nitrogen

fixation by diazotrophs which can supply upto 163.2 Tg new nitrogen per year globally (Wang et al., 2019). With regard to P, *Prochlorococcus* and *Synechococcus* mostly depend on the remineralisation of dissolved organic phosphorus (DOP) via hydrolytic enzymes (e.g. alkaline phosphatase (Muñoz-Marín et al., 2020)). Additionally, evidence is mounting that cyanobacteria are major sources of semi-labile dissolved organic matter (DOM) phosphonates (Repeta et al., 2016). The bacterial degradation of methylphosphonates (MPn) releases CH<sub>4</sub> and therefore cyanobacteria are thought to play a key role in the global marine CH<sub>4</sub> flux. *In-situ* production of CH<sub>4</sub> is a repercussion from different metabolic pathways including the conversion of methylated substrates, and are induced by environmental stress (e.g. nutrient supply and variations in temperature, salinity and light attenuation). As such, CH<sub>4</sub> may be the by-product of the methylphosphonate remineralisation in phosphate-stressed surface waters, i.e. the MPn way (Karl et al., 2008, Bizic et al 2020). Due to the strong depletion of inorganic phosphorus in oligotrophic areas in the Atlantic or the Pacific Oceans, cyanobacteria use the organic phosphonates as a P source (Feingersch et al., 2012), leading to the release of methyl groups in the water that are rapidly converted into CH<sub>4</sub> (Beverdors et al., 2010). In contrast, nitrate availability might control CH<sub>4</sub> production in phosphate replete surface waters. While in P-limited waters cyanobacteria use methylphosphonate as a nutrient source and hence release CH<sub>4</sub>, in N-limited waters, CH<sub>4</sub> may result from the breakdown of DMSP (Dimethylsulfoniopropionate), a subsequent demethylation of DMS (Dimethyl Sulfide) and inherently the oxidation of dissolved methyl groups (Florez-Leiva et al., 2013). Other environmental parameters, such as variations in temperature or light attenuation may also influence CH<sub>4</sub> formation, although data is lacking to fully understand the metabolic pathways leading to CH<sub>4</sub> production.


In the current context of climate change and warming ocean waters, we are likely to expect a shift in the community of primary producers towards smaller sized cyanobacteria such as *Prochlorococcus* and *Synechococcus* (van de Waal and Litchman, 2020), with a concurrent decrease in total biomass in the open ocean (Marinov et al., 2010). Yet, with accumulating evidence on the importance of the cyanobacterial contribution towards CH<sub>4</sub> production and given the important role of CH<sub>4</sub> as a potent GHG, it is crucial to intensify the monitoring and investigating of CH<sub>4</sub> production and fluxes in surface oceanic waters in order to feed a global database (Bange et al., 2009). There is a lack of data to comprehend the current and future role of the Earth's oceans and primary production in relation to the global CH<sub>4</sub> budget.

In the present paper, we provide an update on the dissolved CH<sub>4</sub> concentrations and air-sea fluxes of CH<sub>4</sub> in surface waters of the subtropical North Atlantic Ocean as part of the JC191 hydrographical cruise (RRS James Cook, Jan-Mar 2020). Furthermore, we present data of the distribution of the two dominant cyanobacteria, *Prochlorococcus* and *Synechococcus* to highlight the contribution of the planktonic community to the CH<sub>4</sub> flux. More specifically, our objectives were to confirm the presence of CH<sub>4</sub> anomalies in surface waters and the associated air-sea fluxes, and to propose possible mechanisms and sources controlling the CH<sub>4</sub> distribution by examining the relationships between physico-chemical and biological parameters.

## 2. Methods

### 2.1 Water sampling and analysis for dissolved gas concentrations

Seawater samples were collected during the JC191 hydrographical cruise (as part of the GO-SHIP program, PI A. Sanchez-Franks (Sanchez-franks, 2020)) on board the RRS James Cook between January-March 2020 along a west-east transect in the subtropical North Atlantic from Fort Lauderdale, USA to Santa Cruz de Tenerife, Spain on the nominal 24°N parallel. 64 profiles (out of 135 stations occupied by CTD casts in total) from surface to 400 m depth (or full-depth for the shallower continental margin stations) were sampled (Figure 1).



105 **Figure 1. Cruise track of JC191 24N expedition from Fort Lauderdale, USA to Santa Cruz de Tenerife, Spain (January – March 2020). White numbers indicate some of the total 135 CTD stations.**

Water samples for dissolved CH<sub>4</sub> measurements were collected into 20-ml headspace vials using a 24-Niskin Bottle rosette equipped with a SBE911+ CTD (Conductivity, Temperature, Depth), a AquaTracka III fluorometer (Chelsea Technologies) and a dissolved oxygen Sensor (Seabird SBE 43). Dissolved gas samples were poisoned with Hg(Cl)<sub>2</sub>, then fitted with Teflon stopcocks and crimp-sealed under exclusion of any air bubbles. Samples were immediately stored at 4 °C after sampling until analysis on shore at the Station Biologique de Roscoff.

Gas extraction and analyses were performed using a Shimadzu Headspace Sampler (HS-20) connected to a gas chromatograph (Shimadzu GC-2030), fitted with a barrier discharge ionization detector (BID) and a 30-m SH-Rt-MSieve 5A column. With this set-up, headspace extraction is entirely automated with pressurization of the sample up to 2 bars, heating at 90 °C and equilibration for 7 minutes. An aliquot of the gas sample was transferred to a 1-ml injection loop, maintained at 150 °C and injected into a 50 °C heated column. Calibrations were made by injecting known volume of standard gas (Messer®, 1000 ppm, 500 ppm and 100 ppm (+/- 1ppm), CH<sub>4</sub> in helium and 500 ppb H<sub>2</sub> in helium). All analyses were made in duplicate and results are given as averaged values. The detection threshold of this method is 0.2 nmol for dissolved CH<sub>4</sub> and variation between duplicates was 5%.

## 2.2 Inorganic Nutrient Analysis

Samples for inorganic nutrient analysis ( $\text{NO}_2^-$ ,  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ ,  $\text{PO}_4^{3-}$ ) were collected unfiltered into sterile 15 ml centrifuge tubes (rinsed three times with water from the same Niskin). Samples were analysed directly from the collection tubes within 1-8  
125 hours and measured from the lowest to the highest concentration (surface to deep) to reduce any carry over effects. Nutrients were analysed on board using a 4-channel Seal Analytical AA3 segmented flow autoanalyzer following GO-SHIP protocols (Becker et al., 2020). In order to test the accuracy and precision of the analyses, certified reference materials (CRMs) from KANSO Technos Co. (lots lot CD, CJ, CI and lot BW) were measured in triplicate in every run.

## 2.3 Cyanobacteria sampling and analysis

38 of the stations occupied during JC191 were sampled at six depths in the upper water column (max. sampling depth 375 m) following the live fluorescence profiles, to determine the prevailing community of primary producers (unpublished data, Marx 2020). Bulk water samples (5 L) were collected from Niskin bottles, from which then subsamples for the flow cytometric determination of *Prochlorococcus* and *Synechococcus* abundances were collected. 4 ml of sample water were immediately fixed with 40  $\mu\text{l}$  Glutaraldehyde solution (50%) and stored at 4°C until transferred to a low temperature freezer (-80 °C) after  
135 12 hours.

Samples were analysed at the University of Portsmouth on a CyFlow Cube 8 (Sysmex) flow cytometer immediately after defrosting and at a flowrate of 1  $\mu\text{l/s}$ . The distinction between *Prochlorococcus* and *Synechococcus* was achieved by gating each group according to its fluorescence signals (red and orange fluorescence) against the size fractionation (forward and side scatter). For each of the stations sampled, the mixed layer depth (MLD, defined as the depth at which temperature decreased  
140 by 1 °C from the surface) was determined and the integrated average in abundance above said MLD was calculated.

## 2.4 Statistical analyses

In order to determine the biological and physico-chemical parameters that influence the distribution of dissolved methane in surface waters, a principal component analysis (PCA) was applied. This statistical tool simplifies the underlying structure of  
145 the multivariate dataset, converting a large number of variables into a smaller number of variables, i.e. components (PCs) with a minimum loss of information. Each PC is associated to an eigenvalue that indicates the variation in the data. Here, the PCA computes a singular value decomposition of the data matrix and does not incorporate eigenvalue of the covariance matrix to maintain numerical accuracy. Then, factor loadings were calculated, where a high factor loading indicates a significant correlation between variables. Additionally, Kendall rank correlations were computed to evaluate associations between two  
150 variables (e.g. methane concentrations and fluorescence or phytoplankton abundances). All statistics were performed with RStudio and the R 'stats' package' (Foundation for Statistical Computing, Vienna, 2013).

## 2.5 Flux calculation

The flux of air-sea CH<sub>4</sub> was calculated following established methods based on (Kelley and Jeffrey, 2002a; Wanninkhof, 2014)

155

$$F = k (C_w - C_a)$$

Where F is the CH<sub>4</sub> flux (mol m<sup>-2</sup> d<sup>-1</sup>) from seawater to air, with k as gas transfer coefficient under consideration of wind speed, C<sub>w</sub> as average CH<sub>4</sub> concentration measured in surface water and C<sub>a</sub> as equilibrated seawater-air CH<sub>4</sub> concentration. Wind speed  
160 data was obtained from on-ship measurements throughout the cruise transect and concentration of CH<sub>4</sub> in the air was assumed to be 1.9 ppm (based on NOAA, Global Monitoring Laboratory <https://www.esrl.noaa.gov> report). The gas transfer coefficient k was calculated following:

$$k = 0.251 \langle U^2 \rangle (Sc/660)^{-0.5}$$

165

where  $\langle U^2 \rangle$  represents the average of neutral stability of winds at 10 m height squared and Sc indicated the Schmidt number, taking into account the kinematic viscosity of water and the molecular diffusion coefficient of the gas. From inverse modelling using CCMP winds and the Modular Ocean Model – General Circulation model (MOM3 GCM), a value of 0.251 was obtained. Sc was calculated as

170

$$Sc = A + Bt + Ct^2 + dt^3 + Et^4$$

with coefficients A, B, C, D and E obtained from a least square fourth-order polynomial fit (Wanninkhof, 2014).

## 175 3. Results

### 3.1 Spatial distribution of physico-chemical parameters

CH<sub>4</sub> concentrations in surface waters of the subtropical North Atlantic along the 24°N were distributed non-uniformly between 3-10 nmol/l, i.e. systematically above saturation of ~2.7 nmol/l. Lowest concentrations of CH<sub>4</sub> of 3-4 nmol/l were found in the central gyre system above the mid-Atlantic ridge (~45°W), with increasing values of 8-10 nmol/l in both western  
180 and eastern boundaries closer to the continental shelf (Figure 2a).

Chlorophyll *a* (Chl *a*, Figure 2b) from real-time fluorescence profiles exhibit highest concentrations near both western and eastern shore systems and lowest concentrations in surface waters throughout the central gyre system (< 0.1 µg/l, Figure 2). A deep chlorophyll maxima (DCM) was observed between 100-130 m water depth and was consistently above the mixed layer

depth (MLD). Concentrations of Chl *a* increase towards the Mauritanian upwelling off the North African coast to above 0.4  $\mu\text{g/l}$ , indicating higher primary production due to enhanced nutrient supply. Accordingly, light transmission is decreased due to higher content of suspended particles in the water column (Figure 2h). Furthermore, fluorescence does align with patterns of dissolved oxygen in the water column of this transect. The surface waters in the subtropical North Atlantic are well oxygenated with concentrations above 200  $\mu\text{mol/kg}$  in the top 100 metres of the water column, subsequently decreasing with depth. Increased concentrations of dissolved oxygen are observed towards the eastern boundary following enhanced lateral transport from the coastal upwelling (Figure 2e).

Nitrate and phosphate concentrations were low ( $< 0.1 \mu\text{mol/kg}$ ,  $< 0.01 \mu\text{mol/kg}$ , respectively) in surface waters throughout the transect with nutriclines  $> 200 \text{ m}$  depth, shallowing ( $< 150 \text{ m}$  in eastern basin and  $< 100 \text{ m}$  in eastern boundary) towards the eastern boundary due to coastal upwelling off the eastern North African coast and associated enhanced mixing of deep, nutrient enriched waters into surface waters (Figure 2c and d).





Figure 2. Distribution of a) CH<sub>4</sub>, b) Chl *a*, c) NO<sub>3</sub>, d) PO<sub>4</sub>, e) dissolved oxygen, f) salinity, g) temperature and h) transmission along the cruise track. Chl *a*, dissolved oxygen, salinity, temperature and transmission data originate from sensors of and attached to the CTD. CH<sub>4</sub>, NO<sub>3</sub> and PO<sub>4</sub> were measured analytically. Part of the data were taken from CCHDO (Sanchez-franks, 2020).

### 3.2 Sea-to-air methane flux

200 At most stations, air-sea CH<sub>4</sub> flux was 1-2  $\mu\text{mol m}^{-2} \text{d}^{-1}$ , with maximum values of 5  $\mu\text{mol m}^{-2} \text{d}^{-1}$  in the area of 24°N 25°W (Figure 3). Only one station (~40°W) had a negative CH<sub>4</sub> flux, which indicates a sink of CH<sub>4</sub>. The overall average flux across the subtropical North Atlantic was 1.9  $\mu\text{mol m}^{-2} \text{d}^{-1}$ . However, it should be noted that the air-sea flux of CH<sub>4</sub> is dependant on the wind speed and can be locally different, varying depending on specific weather conditions.



205 **Figure 3. Sea-to-air CH<sub>4</sub> flux in  $\mu\text{mol m}^{-2} \text{d}^{-1}$  along the cruise track of JC191. Dots indicate stations occupied by CTD casts.**

Although sea-to-air flux of CH<sub>4</sub> presented here broadly agree with similar values of 1.6  $\mu\text{mol m}^{-2} \text{d}^{-1}$  measured in the oligotrophic North Pacific, it can differ substantially both temporarily and spatially. For instance, sea-to-air flux of CH<sub>4</sub> in the Sargasso Sea, also partly encompassed in the transect of JC191 is fluctuating between 1.6-4.4  $\mu\text{mol m}^{-2} \text{d}^{-1}$ , depending on the season and weather conditions (Holmes et al., 2014). Furthermore, the base ecosystem can impact the sea-to-air flux of CH<sub>4</sub> if compared to other oceanic regions such as Belgic coastal zones with 1-160  $\mu\text{mol CH}_4 \text{ m}^{-2} \text{d}^{-1}$  (Borges et al., 2016) or Red Sea mangroves with 13.3  $\mu\text{mol CH}_4 \text{ m}^{-2} \text{d}^{-1}$  (Sea et al., 2018), or surveys in the east China Sea (6.5-7.4  $\mu\text{mol m}^{-2} \text{d}^{-1}$  (Ye et al., 2015)) or the Gulf of Mexico (0.38  $\mu\text{mol m}^{-2} \text{d}^{-1}$  (Kelley and Jeffrey, 2002b)).

## 4. Discussion

### 4.1 Methane distribution in surface waters of the subtropical North Atlantic Ocean


215 Our data across the subtropical North Atlantic Ocean unambiguously indicates an oversaturation in CH<sub>4</sub> of the surface layer (400 m) of 110 to 370%, which is in agreement with previous observations describing concentrations varying between 2-5 nmol/l and maximum values of 10 nmol/l (Scranton and Brewer, 1977; Conrad and Seiler, 1988; Forster et al., 2009; de la Paz et al., 2015; Leonte et al., 2020). This is also in line with previous observations describing the upper layer in subtropical areas



of the global ocean as a source of CH<sub>4</sub> to the atmosphere (Reeburgh, 2007b; Dang and Li, 2018) and also in some regions of  
220 the Arctic Ocean (Kitidis et al., 2010; Kudo et al., 2018). To date, only the Southern Ocean is undersaturated in CH<sub>4</sub>, although  
limited by the scarcity of data collected, highlighting the need of increased surveying to be complemented (Bui et al., 2018).  
The distribution of dissolved CH<sub>4</sub> was variable across the subtropical North Atlantic with higher concentrations in the Eastern  
Basin (65-80 °W) and the Western Basin (15-30 °W) and lowest concentrations measured in the central gyre system of the  
transect (30-65 °W) (Figure 2). The vertical distribution of CH<sub>4</sub> appears to be associated with fluorescence; highest  
225 concentrations of CH<sub>4</sub> were found at ~100 m depth, where fluorescence and dissolved oxygen are highest and nutrients levels  
lowest, which is in agreement with previous findings (Kudo et al., 2018).


PCA was applied to the datasets collected to identify which environmental parameters (nutrients, fluorescence, dissolved  
oxygen, temperature, salinity, depth, transmission) are related to or influence the distribution of CH<sub>4</sub> along the transect. 54%  
of the variability could be explained by the first component which is primarily controlled by depth (Figure 4). CH<sub>4</sub> is mainly  
230 associated with oxygenated surface waters, characterised by low concentrations of nutrients and a higher fluorescence.  
However, the weak contribution of CH<sub>4</sub> to the first two components may be due to the heterogeneity of its distribution. To  
fully understand the relationship between the different parameters controlling the distribution of CH<sub>4</sub>, we therefore separated  
the transect into three main regions, i.e. i) the Western Basin, ii) the Central Gyre System and iii) the Eastern Basin. The PCA  
applied to the regionalized dataset (Figure 5) revealed, that CH<sub>4</sub> is clearly associated to the abundance of primary producers in  
235 surface waters (<100m) in the western basin, while in the central gyre and eastern basin CH<sub>4</sub> concentrations were also  
influenced either by *in-situ* physical processes such as mixing or mesoscale eddies (Kudo et al., 2018) or by external inputs  
such as the Mauritanian upwelling that supplies CH<sub>4</sub>-enriched waters to the upper layer (Conrad and Seiler, 1988).





245 **Figure 5: Principal Component Analysis on the regionalized dataset: west (80-65°W), central (65-30°W) and east (30-20°W). Numbers on the x and y axes indicate the factor loadings of each variable of each principal component (PC). The percentages show the explained variability in the dataset by each PC. CH<sub>4</sub> is highlighted in red.**

250 Although CH<sub>4</sub> appeared to have a uniform vertical distribution (Figure 2), selected profiles from two areas of interest, CTD 50 (69,5 °W, 24.9 °N) and CTD 122 (20,8 °W, 25,45 °N) of CH<sub>4</sub>, Chl *a*, phosphate and dissolved oxygen showed that the highest concentration of CH<sub>4</sub> in fact correspond to the maxima of Chl *a* and dissolved oxygen and the lowest concentrations of phosphate (Figure 6). The correlation between CH<sub>4</sub> and Chl *a* (Kendall rank correlation test,  $r^2=$ ,  $p<0.05$ ) further suggests that the primary producers play a role in the production of CH<sub>4</sub>.




255 Figure 6. Selected data from CTD 50 (69.5 °W, 24.9 °N) and CTD 122 (20.8 °W, 25.45 °N) of CH<sub>4</sub> (red circles), Chl *a* (green triangles), phosphate (purple squares) and dissolved oxygen (orange).

At station 50, the relationship between CH<sub>4</sub>, Chl *a* and phosphate appears to be linear, e.g. CH<sub>4</sub> concentration decreasing as Chl *a* decreases and phosphate increases; while at station 122 CH<sub>4</sub> concentration showed a nonlinear pattern. It is not clear why CH<sub>4</sub> concentrations are variable but zooplankton grazing could potentially have a substantial impact (Simon et al., 2012). A possible influence of gas seeps on the CH<sub>4</sub> concentration is negligible as gas seeps only influence CH<sub>4</sub> concentrations in the immediate water column 100 to 150 m above the seeps (Leonte et al. 2020). Below 250 m water depth, CH<sub>4</sub> concentrations are decreasing, corresponding to an increase in phosphate and minimal Chl *a* concentrations, again suggesting the influence of primary producers (Brown et al., 2014).


#### 265 4.2 Methane production linked to primary production

Autotrophic cyanobacteria *Prochlorococcus* and *Synechococcus* represent a major constituent of primary production in the subtropical North Atlantic, their distribution however differs greatly (Flombaum et al., 2013). Whereas *Synechococcus* solely occupies surface waters up to depths of ~ 100 m, *Prochlorococcus* occupies the whole water column, with deep-water maxima just above MLD, therefore responsible for fluorescence maxima at 100 to 130 m. The longitudinal distribution also differs between both taxa: The distribution of *Synechococcus* is limited to coastal, nutrient richer waters (Figure 7), whereas *Prochlorococcus* dominates the community throughout the transect (Figure 8).




275

Figure 7. Depth-integrated abundance of *Synechococcus* (in  $10^3$  cells/ml) above MLD. Black dots represent the 38 out of the total 135 stations sampled for cyanobacterial abundance.



280 Figure 8. Depth-integrated abundance of *Prochlorococcus* (in  $10^3$  cells/ml) above MLD. Black dots represent the 38 out of the total  
285 135 stations sampled for cyanobacterial abundance.

CH<sub>4</sub> distribution across the subtropical north Atlantic suggests the influence of the cyanobacterial community present; highest  
CH<sub>4</sub> concentrations were found 1) at DCM in the central gyre system, where *Prochlorococcus* is the predominant genus and  
2) at the gyre boundaries, where higher abundances of both *Prochlorococcus* and *Synechococcus* were found. A PCA applied  
on the dataset including abundances of both cyanobacteria strengthens the conclusion that CH<sub>4</sub> production appears to be  
associated to primary production (Figure 9).



290 **Figure 9. Principal Component Analysis (PCA) between dissolved CH<sub>4</sub> concentrations (red), *Prochlorococcus* and *Synechococcus* abundances (light blue) and other physico-chemical parameters (nutrients, depth, fluorescence, temperature, salinity and turbidity). Numbers on the x and y axes indicate the factor loadings of each variable of each principal component (PC). The percentages show the explained variability in the dataset by each PC.**

Not only do the bordering Gulf Stream to the West and the Canary Stream to the East provide a resupply of nutrients (Williams et al., 2011), and therefore support a greater abundance of cyanobacteria; the Canary Stream presents a pathway for the horizontal transfer of organic matter from the North African coast to the open ocean. Previous studies confirm high OM production in the eastern Subtropical North Atlantic as a result of the high rates of primary production fuelled by the Mauritanian upwelling (Reynolds et al., 2014). The transmission data (Figure 2h) suggests the export of organic matter throughout the eastern basin, which could contain MPn and thus a potential source of CH<sub>4</sub>. Elevated CH<sub>4</sub> concentrations are mainly limited to the highly productive boundary systems, whilst the central gyre which lacks a sufficient resupply of bioavailable nutrients due to its downwelling nature, results in a decreased abundance of both *Prochlorococcus* and *Synechococcus* and henceforth decreased CH<sub>4</sub> concentrations (Figure 2a). In this especially P limited region, alternative nutrient sources such as the degradation of DOM to access organic phosphorus compounds become increasingly important in order for the cyanobacteria to meet their nutrient needs. The degradation of dissolved organic phosphorus via alkaline phosphatases (AP) such as *phoA* or *phoX* and the overexpression of the phosphate binding protein (pstS) have been believed to be the main adaptations to P-stress (Luo et al., 2009; Cox and Saito, 2013; Sebastian and Ammerman, 2009). Recently however, evidence was brought forth that some strains of *Prochlorococcus* can also oxidise MPn and other higher phosphonate compounds while releasing formate and potentially CH<sub>4</sub> as a by-product (Sosa et al., 2019a). Phosphonates are notably abundant and enriched in DOM (Sosa et al., 2019b) and their degradation releases considerable amounts of CH<sub>4</sub>. *Prochlorococcus* and *Synechococcus* are the most abundant primary producers in the oligotrophic ocean, and as such produce considerable amounts of semi-labile DOM; both can synthesize phosphoenolpyruvate mutases (*pepM*) and therefore the DOM produced carries an enriched pool of MPn (Repeta et al., 2016; Sosa et al., 2019a). However, the metabolism of MPn is heavily regulated by bioavailable phosphate, thus the metabolic pathway of *pepM* might be heavily down regulated in the subtropical North Atlantic, whereas under replete conditions i.e. in the North Pacific *Prochlorococcus* can allocate up to 40 % of its internal P-quota towards phosphonate synthesis (Acker et al., 2022). Yet, the trait to produce phosphonates is located on genomic islands and is subject to horizontal gene transfer and can be frequently exchanged among marine microbial communities, hence also proteobacteria such *Pelagibacter spp.* in the SAR11 clade obtain *pepM* and are able to produce phosphonates (Acker et al., 2022). Similarly is the trait of phosphonate consumption subject to horizontal gene transfer and high-light strains of *Prochlorococcus* carry both production and consumption traits, therefore can also utilise MPn as alternative P source (Acker et al., 2022). The demethylation of MPn is mostly attributed to occur under P limiting conditions, such as in oligotrophic oceanic regions and further, most MPn oxidation and subsequent release of CH<sub>4</sub> is due to bacterial degradation of DOM and the breakdown of high energetic carbon-phosphorus bonds via C-P lyases, which are encoded by the *phn* operon, with transport systems including *phnC*, *phnD* and *phnE* and *phnJ* responsible for the cleavage of the C-P bond (Sosa et al., 2020). C-P lyases are abundant among *Pelagibacter spp.* and other alpha and gamma proteobacteria and can be found in ~ 50 % of organisms in the North Atlantic, where DOP concentrations are 4 fold lower in respect to the North Pacific (Sosa et al., 2020). Carini et al. (2014) estimated a potential production of 0.01-0.05 nM CH<sub>4</sub> per day by *Pelagibacterales* via



C-P cleavage within the MPn pathway (Carini et al. 2014). However, due to complex biogeochemical and physical processes involved, *in-situ* production of CH<sub>4</sub> is difficult to assess and quantifying the individual contribution of primary producers towards CH<sub>4</sub> production was not the primary scope of the work presented. Here, the focus was to link the distribution of CH<sub>4</sub> in the subtropical North Atlantic to two vastly abundant cyanobacteria as exemplary primary producers, *Prochlorococcus* and *Synechococcus*, in order to draw conclusions which might explain the ocean methane paradox. Many mechanisms and precursors of oceanic *in-situ* production of CH<sub>4</sub> remain unknown and it is unclear, whether cyanobacteria such as *Prochlorococcus* or *Synechococcus* are responsible for the release of CH<sub>4</sub> mainly via the degradation of MPn or indirectly by producing semi-labile DOM containing MPn cleaved by the bacterial community. Furthermore, recent laboratory-based studies also confirmed, that planktonic organisms produce CH<sub>4</sub> per se under high light intensities; *Prochlorococcus* showed a higher potential in CH<sub>4</sub> production than *Synechococcus*, with 0.8-110 pmol and 0.01-0.6 pmol CH<sub>4</sub> per 10<sup>6</sup> cells per hour respectively, larger cyanobacteria such as *Microcystis spp.* exceeded this by up to  $0.51 \pm 0.26 \mu\text{mol/g}$  per hour (Bižić et al., 2020). Nonetheless, *Prochlorococcus* alone produces a gross ~22 nM CH<sub>4</sub> per day and therefore contributes substantially to the supersaturation of oligotrophic oceanic regions (typically between 7-25 nM) and to the over saturated state measured here (up to 10 nM) in the subtropical North Atlantic (Bižić et al., 2020). Cyanobacteria can greatly influence and potentially control CH<sub>4</sub> production in the marine environment, although other marine organisms such as different bacterial clades or coccolithophores are believed to also contribute to overall CH<sub>4</sub> production. Nitrogen fixing diazotrophs such as *Trichodesmium spp.* or *Rhodopseudomonas palustris* may contribute even to a higher degree (Bižić-Ionescu et al., 2018), (Zheng et al., 2018) and although spatially limited with *Trichodesmium spp.* dominating the western basin and *Hemilaulus* associated *Richelia* symbionts more so in the eastern basin (Luo et al., 2009), they are also abundant throughout the subtropical North Atlantic. *Trichodesmium* has higher nutrient requirements and can therefore outcompete cyanobacteria in uptake of inorganic nutrients and degradation of alternative sources; yet, energy intensive diazotrophic nitrogen fixation is controlled by micronutrients such as iron (Macovei et al., 2019) and is primarily occurring in the western basin, an area of high iron input by aeolian plumes originating from the Saharan desert (Ratten et al., 2015; Reynolds et al., 2014). Lastly, coccolithophores such as the ubiquitous abundant *Emiliania huxleyi* can also produce CH<sub>4</sub> (between  $0.7\text{-}3.1 \pm 0.4$  CH<sub>4</sub> per gram POC per day (Lenhart et al., 2016b) (Klitzsch et al. 2019)) from OM degradation, and are believed to increase in abundance in the subtropical North Atlantic with increasing CO<sub>2</sub> concentrations at the air-sea interface due to further anthropogenic perturbation of the atmospheric CO<sub>2</sub> budget (Krumhardt et al., 2015).

Nonetheless, the data presented here suggest that the cyanobacterial community most likely plays a significant key role in the CH<sub>4</sub> flux in surface waters and the degradation of MPn from semi-labile DOM, helping to explain the methane paradox and henceforth the sea-air flux of CH<sub>4</sub> (Sosa et al., 2019a). Further investigation needs to focus on gathering *in-situ* data and should also include future scenarios, considering future climate and whole ecosystem community responses to consequences of altered climate conditions. The base ecosystem, specifically in P-limited regions is ever changing and highly adapted and horizontal transfer of genomic traits of P acquisition via various genomic pathways might become inherently important. Further work should also be considered in regards to mesocosm incubation studies with *in-situ* communities of natural composition to

improve our understanding of the various processes and interactions involved in oceanic CH<sub>4</sub> production linked to primary producers to explain the ocean methane paradox.

## 5 Conclusions

Our study demonstrates that the subtropical North Atlantic Ocean does indeed act as a source of CH<sub>4</sub> to the atmosphere, most likely controlled by cyanobacteria which are the dominant primary producers in the surface waters. Yet, anomalies found at depths below 200 m could also be attributed to the degradation of sinking organic material. The concentrations of dissolved CH<sub>4</sub> in this study were considerably higher near shelf regions and in the eastern boundary under the influence of the Mauritanian upwelling. The accumulation of organic matter and nutrients in these areas provide favourable conditions for both aerobic and anaerobic CH<sub>4</sub> production. It is expected that with increasing stratification and subsequent reduction in nutrient supply to the surface oligotrophic North Atlantic Ocean, the prevailing P-limitation will be further exacerbated, whereas coastal and shelf regions with increasing anthropogenic inputs of nutrients, could experience more frequent cyanobacterial blooms which, will in turn enhance CH<sub>4</sub> production (Dang and Li, 2018).

## 6. Author Contributions

Anna Kolomijeca – carried out dissolved gas sample collection, measurements and data analysis and drafted the final manuscript

Lukas Marx – carried out phytoplanktonic sample collection, measurements and data analysis and drafted the final manuscript

Sarah Reynolds – helped in preparation of the cruise, measurements and data analysis, supervised biological part of the survey

Thierry Cariou - carried out nutrient sample collection, measurements and data analysis

Edward Mawji - carried out nutrient sample collection, measurements and data analysis

Cedric Boulart – helped to plan and prepare for the cruise, result evaluation and provided overall project supervision

All authors contributed equally to the manuscript writing and had substantial input to the final version.

## 7. Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## 385 8. Funding

Ship time was funded by NERC as part of the Climate Linked Atlantic Sector Science (CLASS) research programme and as a UK contribution to the Global Ocean Ship-based Hydrographic Investigations Program (GO-SHIP).

A Kolomijeca's participation to the cruise and dissolved gas analyses were funded by the Region Bretagne SAD 'FULMAR' project, the Foundation Air Liquide SMIS-4M project and the CNRS (LEFE Memestra).

- 390 L. Marx's and Sarah Reynolds' participation was funded by the University of Portsmouth PhD bursary scheme and a UK NERC National Capability programme CLASS (Climate Linked Atlantic Sector Science) ECR fellowship.

## 9. Acknowledgments

- The Authors gratefully acknowledge the PI of the JC191 cruise Alejandra Sanchez-Franks (National Oceanography Centre, UK) for the great scientific support and providing cruise data. This work would not have been possible without the support and help of the Captain and crew of RRS James Cook. The authors would also like to acknowledge Eric Mace (Station Biologique de Roscoff) – the CNRS technician for his technical support in the preparation of the cruise.

## References

- 400 Acker, M., Hogle, S. L., Berube, P. M., Hackl, T., Coe, A., Stepanauskas, R., Chisholm, S. W., and Repeta, D. J.: Phosphonate production by marine microbes: Exploring new sources and potential function, *Proc. Natl. Acad. Sci. U. S. A.*, 119, [https://doi.org/10.1073/PNAS.2113386119/SUPPL\\_FILE/PNAS.2113386119.SD03.CSV](https://doi.org/10.1073/PNAS.2113386119/SUPPL_FILE/PNAS.2113386119.SD03.CSV), 2022.
- Bange, H. W., Bell, F. T. G., Cornejo, B. M., and Freing, C. A.: MEMENTO : A proposal to develop a database of marine nitrous oxide and methane measurements MEMENTO : a proposal to develop a database of marine nitrous oxide and methane measurements, 10–13, <https://doi.org/10.1071/EN09033>, 2009.
- 405 Beversdorf, L. J., White, A. E., Björkman, K. M., Letelier, R. M., and Karl, D. M.: Phosphonate metabolism by *Trichodesmium* IMS101 and the production of greenhouse gases, *Limnol. Oceanogr.*, 55, 1768–1778, <https://doi.org/10.4319/lo.2010.55.4.1768>, 2010.
- Bižić-Ionescu, M., Klintzsch, T., Ionescu, D., Hindiyeh, M. Y., Günthel, M., Muro-Pastor, A. M., Eckert, W., Keppler, F., and Grossart, H. P.: Widespread formation of methane by Cyanobacteria in aquatic and terrestrial environments, 398958, <https://doi.org/10.1101/398958>, 2018.
- 410 Bižić, M., Klintzsch, T., Ionescu, D., Hindiyeh, M. Y., Günthel, M., Muro-Pastor, A. M., Eckert, W., Urich, T., Keppler, F., and Grossart, H. P.: Aquatic and terrestrial cyanobacteria produce methane, *Sci. Adv.*, 6, 1–10, <https://doi.org/10.1126/sciadv.aax5343>, 2020.
- Borges, A. V., Champenois, W., Gypens, N., Delille, B., and Harlay, J.: Massive marine methane emissions from near-shore shallow coastal areas, *Sci. Rep.*, 6, 2–9, <https://doi.org/10.1038/srep27908>, 2016.
- 415 Brown, I. J., Torres, R., and Rees, A. P.: Dynamics of Atmospheres and Oceans The origin of sub-surface source waters define the sea – air flux of methane in the Mauritanian Upwelling , NW Africa, *Dyn. Atmos. Ocean.*, 67, 39–46, <https://doi.org/10.1016/j.dynatmoce.2014.06.001>, 2014.
- Bui, O. T. N., Kameyama, S., Yoshikawa-Inoue, H., Ishii, M., Sasano, D., Uchida, H., and Tsunogai, U.: Estimates of methane emissions from the Southern Ocean from quasi-continuous underway measurements of the partial pressure of methane in surface seawater during the 2012/13 austral summer, *Tellus, Ser. B Chem. Phys. Meteorol.*, 70, 1–15, <https://doi.org/10.1080/16000889.2018.1478594>, 2018.
- 420 Conrad, R. and Seiler, W.: Methane and hydrogen in seawater (Atlantic Ocean), *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 35, 1903–1917, [https://doi.org/10.1016/0198-0149\(88\)90116-1](https://doi.org/10.1016/0198-0149(88)90116-1), 1988.

- Cox, A. D. and Saito, M. A.: Proteomic responses of oceanic *Synechococcus* WH8102 to phosphate and zinc scarcity and cadmium additions, *Front. Microbiol.*, 4, <https://doi.org/10.3389/FMICB.2013.00387>, 2013.
- 425 Dang, H. and Li, J.: Climate tipping-point potential and paradoxical production of methane in a changing ocean, *Sci. China Earth Sci.*, 61, 1714–1727, <https://doi.org/10.1007/s11430-017-9265-y>, 2018.
- Feingersch, R., Philosof, A., Mejuch, T., Glaser, F., Alalouf, O., Shoham, Y., and Bèjja, O.: Potential for phosphite and phosphonate utilization by *Prochlorococcus*, *ISME J.*, 6, 827–834, <https://doi.org/10.1038/ismej.2011.149>, 2012.
- 430 Flombaum, P., Gallegos, J. L., Gordillo, R. A., Rincón, J., Zabala, L. L., Jiao, N., Karl, D. M., Li, W. K. W., Lomas, M. W., Veneziano, D., Vera, C. S., Vrugt, J. A., and Martiny, A. C.: Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*, *Proc. Natl. Acad. Sci. U. S. A.*, 110, 9824–9829, <https://doi.org/10.1073/pnas.1307701110>, 2013.
- Florez-Leiva, L., Damm, E., and Farías, L.: Methane production induced by dimethylsulfide in surface water of an upwelling ecosystem, *Prog. Oceanogr.*, 112–113, 38–48, <https://doi.org/10.1016/j.pocean.2013.03.005>, 2013.
- 435 Forster, G., Upstill-Goddard, R. C., Gist, N., Robinson, C., Uher, G., and Woodward, E. M. S.: Nitrous oxide and methane in the Atlantic Ocean between 50°N and 52°S: Latitudinal distribution and sea-to-air flux, *Deep. Res. Part II Top. Stud. Oceanogr.*, 56, 964–976, <https://doi.org/10.1016/j.dsr2.2008.12.002>, 2009.
- Foundation for Statistical Computing, Vienna, A. I. 3-900051-07-0: R Core Team, 2013.
- Harvey, B. P., Gwynn-Jones, D., and Moore, P. J.: Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming, *Ecol. Evol.*, 3, 1016–30, <https://doi.org/10.1002/ece3.516>, 2013.
- 440 Hickman, A. E., Dutkiewicz, S., Williams, R. G., and Follows, M. J.: Modelling the effects of chromatic adaptation on phytoplankton community structure in the oligotrophic ocean, *Mar. Ecol. Prog. Ser.*, 406, 1–17, <https://doi.org/10.3354/MEPS08588>, 2010.
- Holmes, M. E., Elizabeth, M., Sansone, J., Rust, M., and Popp, N.: Methane production , consumption , and air-sea exchange in the open ocean : An Evaluation based on carbon isotopic ratios Methane production , consumption , and air-sea exchange in the open ocean : An evaluation based on carbon isotopic ratios and via the, <https://doi.org/10.1029/1999GB001209>, 2014.
- 445 Karl, D. M., Beversdorf, L., Björkman, K. M., Church, M. J., Martinez, A., and Delong, E. F.: Aerobic production of methane in the sea, *Nat. Geosci.*, 1, 473–478, <https://doi.org/10.1038/ngeo234>, 2008.
- Kelley, C. A. and Jeffrey, W. H.: Dissolved methane concentration profiles and air-sea fluxes from 41°S to 27°N, *Global Biogeochem. Cycles*, 16, 13-1-13–6, <https://doi.org/10.1029/2001gb001809>, 2002a.
- Kelley, C. A. and Jeffrey, W. H.: Dissolved methane concentration profiles and air-sea fluxes from 41 ° S to 27 ° N Dissolved methane concentration profiles and air-sea fluxes from 41 ° S to 27 ° N, <https://doi.org/10.1029/2001GB001809>, 2002b.
- 450 Kitidis, V., Upstill-Goddard, R. C., and Anderson, L. G.: Methane and nitrous oxide in surface water along the North-West Passage, Arctic Ocean, *Mar. Chem.*, 121, 80–86, <https://doi.org/10.1016/j.marchem.2010.03.006>, 2010.
- Kock, A. and Bange, H. W.: Methane Measurements in Selected Ocean Areas: Eastern Tropical North Atlantic Ocean and Southwestern Labrador Sea, IFM-GEOMAR, Leibniz-Institut für Meereswissenschaften, Diploma, 107, 2007.
- 455 Krumhardt, K. M., Lovenduski, N. S., Freeman, N. M., Bates, N. R., Krumhardt, K. M., Lovenduski, N. S., Freeman, N. M., and Bates, N. R.: Increasing coccolithophore abundance in the subtropical North Atlantic from 1990 to 2014, 12, 18625–18660, <https://doi.org/10.5194/BGD-12-18625-2015>, 2015.
- Kudo, K., Yamada, K., Toyoda, S., Yoshida, N., Sasano, D., Kosugi, N., Ishii, M., Yoshikawa, H., Murata, A., Uchida, H., and Nishino, S.: Spatial distribution of dissolved methane and its source in the western Arctic Ocean, *J. Oceanogr.*, 74, 305–317, <https://doi.org/10.1007/S10872-017-0460-Y/TABLES/1>, 2018.
- 460 de la Paz, M., Huertas, I. E., Flecha, S., Ríos, A. F., and Pérez, F. F.: Nitrous oxide and methane in Atlantic and Mediterranean waters in the Strait of Gibraltar: Air-sea fluxes and inter-basin exchange, *Prog. Oceanogr.*, 138, 18–31, <https://doi.org/10.1016/j.pocean.2015.09.009>,

2015.

Lenhart, K., Klintzsch, T., Langer, G., Nehrke, G., Bunge, M., Schnell, S., and Keppler, F.: Evidence for methane production by the marine algae *Emiliania huxleyi*, 13, 3163–3174, <https://doi.org/10.5194/BG-13-3163-2016>, 2016a.

465 Lenhart, K., Klintzsch, T., Langer, G., Nehrke, G., Bunge, M., Schnell, S., and Keppler, F.: Evidence for methane production by the marine algae *Emiliania huxleyi*, 13, 3163–3174, <https://doi.org/10.5194/bg-13-3163-2016>, 2016b.

Luo, H., Benner, R., Long, R. A., and Hu, J.: Subcellular localization of marine bacterial alkaline phosphatases, *Proc. Natl. Acad. Sci. U. S. A.*, 106, 21219–21223, [https://doi.org/10.1073/PNAS.0907586106/SUPPL\\_FILE/SD3.XLS](https://doi.org/10.1073/PNAS.0907586106/SUPPL_FILE/SD3.XLS), 2009.

470 Macovei, V. A., Torres-Valdés, S., Hartman, S. E., Schuster, U., Moore, C. M., Brown, P. J., Hydes, D. J., and Sanders, R. J.: Temporal Variability in the Nutrient Biogeochemistry of the Surface North Atlantic: 15 Years of Ship of Opportunity Data, *Global Biogeochem. Cycles*, 33, 1674–1692, <https://doi.org/10.1029/2018GB006132>, 2019.

Marinov, I., Doney, S. C., and Lima, I. D.: Response of ocean phytoplankton community structure to climate change over the 21st century: Partitioning the effects of nutrients, temperature and light, 7, 3941–3959, <https://doi.org/10.5194/BG-7-3941-2010>, 2010.

475 Mihai Leonte, Carolyn D. Ruppel, Angel Ruiz-Angulo, John D. Kessler: Surface Methane Concentrations Along the Mid-Atlantic Bight Driven by Aerobic Subsurface Production Rather Than Seafloor Gas Seeps, *JGR Ocean.*, 125, 2020.

Muñoz-Marín, M. C., Gómez-Baena, G., López-Lozano, A., Moreno-Cabezuelo, J. A., Díez, J., and García-Fernández, J. M.: Mixotrophy in marine picocyanobacteria: use of organic compounds by *Prochlorococcus* and *Synechococcus*, *ISME J.* 2020 145, 14, 1065–1073, <https://doi.org/10.1038/s41396-020-0603-9>, 2020.

480 Oppo, D., De Siena, L., and Kemp, D. B.: A record of seafloor methane seepage across the last 150 million years, *Sci. Rep.*, 10, 1–12, <https://doi.org/10.1038/s41598-020-59431-3>, 2020.

Percival, S. L. and Williams, D. W.: Cyanobacteria, *Microbiol. Waterborne Dis. Microbiol. Asp. Risks Second Ed.*, 79–88, <https://doi.org/10.1016/B978-0-12-415846-7.00005-6>, 2013.

485 Ratten, J. M., LaRoche, J., Desai, D. K., Shelley, R. U., Landing, W. M., Boyle, E., Cutter, G. A., and Langlois, R. J.: Sources of iron and phosphate affect the distribution of diazotrophs in the North Atlantic, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 116, 332–341, <https://doi.org/10.1016/J.DSR2.2014.11.012>, 2015.

Reeburgh, W. S.: Oceanic methane biogeochemistry, <https://doi.org/10.1021/cr050362v>, February 2007a.

Reeburgh, W. S.: Oceanic methane biogeochemistry, *Chem. Rev.*, 107, 486–513, <https://doi.org/10.1021/cr050362v>, 2007b.

Repeta, D. J., Ferrón, S., Sosa, O. A., Johnson, C. G., Repeta, L. D., Acker, M., Delong, E. F., and Karl, D. M.: Marine methane paradox explained by bacterial degradation of dissolved organic matter, *Nat. Geosci.*, 9, 884–887, <https://doi.org/10.1038/ngeo2837>, 2016.

490 Reynolds, S., Mahaffey, C., Roussenov, V., and Williams, R. G.: Evidence for production and lateral transport of dissolved organic phosphorus in the eastern subtropical North Atlantic, *Global Biogeochem. Cycles*, 28, 805–824, <https://doi.org/10.1002/2013GB004801>, 2014.

Rhee, T. S., Kettle, A. J., and Andreae, M. O.: Methane and nitrous oxide emissions from the ocean: A reassessment using basin-wide observations in the Atlantic, *J. Geophys. Res. Atmos.*, 114, <https://doi.org/10.1029/2008JD011662>, 2009.

495 Sanchez-franks, A.: CRUISE REPORT : A05 Highlights National Oceanography Centre Cruise Report No . 70 RRS James Cook Cruise JC191 19 JANUARY - 1 MARCH 2020 Hydrographic sections from the Florida Straits to the Canaries Current across 24°N in the Atlantic Ocean, 44, 2020.

500 Schmale, O., Wäge, J., Mohrholz, V., Wasmund, N., Gräwe, U., Rehder, G., Labrenz, M., and Loick-Wilde, N.: The contribution of zooplankton to methane supersaturation in the oxygenated upper waters of the central Baltic Sea, *Limnol. Oceanogr.*, 63, 412–430, <https://doi.org/10.1002/LNO.10640>, 2018.

- Scranton, M. I. and Brewer, P. G.: Occurrence of methane in the near-surface waters of the western subtropical North-Atlantic, *Deep. Res.*, 24, 127–138, [https://doi.org/10.1016/0146-6291\(77\)90548-3](https://doi.org/10.1016/0146-6291(77)90548-3), 1977.
- Sea, M. A., Garcias-bonet, N., Saderne, V., and Duarte, C. M.: Carbon dioxide and methane fluxes at the air – sea interface of Red Sea mangroves, 5365–5375, 2018.
- 505 Sebastian, M. and Ammerman, J. W.: The alkaline phosphatase PhoX is more widely distributed in marine bacteria than the classical PhoA, *ISME J.* 2009 35, 3, 563–572, <https://doi.org/10.1038/ismej.2009.10>, 2009.
- Simon, E., Samuelsen, A., Bertino, L., and Dumont, D.: Estimation of positive sum-to-one constrained zooplankton grazing preferences with the DEnKF: A twin experiment, *Ocean Sci.*, 8, 587–602, <https://doi.org/10.5194/os-8-587-2012>, 2012.
- 510 Sosa, O. A., Casey, J. R., and Karl, D. M.: Methylphosphonate Oxidation in *Prochlorococcus* Strain MIT9301 Supports Phosphate Acquisition, Formate Excretion, and Carbon Assimilation into Purines, *Appl. Environ. Microbiol.*, 85, [https://doi.org/10.1128/AEM.00289-19/SUPPL\\_FILE/AEM.00289-19-S0001.PDF](https://doi.org/10.1128/AEM.00289-19/SUPPL_FILE/AEM.00289-19-S0001.PDF), 2019a.
- Sosa, O. A., Repeta, D. J., DeLong, E. F., Ashkezari, M. D., and Karl, D. M.: Phosphate-limited ocean regions select for bacterial populations enriched in the carbon–phosphorus lyase pathway for phosphonate degradation, *Environ. Microbiol.*, 21, 2402–2414, <https://doi.org/10.1111/1462-2920.14628>, 2019b.
- 515 Sosa, O. A., Burrell, T. J., Wilson, S. T., Foreman, R. K., Karl, D. M., and Repeta, D. J.: Phosphonate cycling supports methane and ethylene supersaturation in the phosphate-depleted western North Atlantic Ocean, *Limnol. Oceanogr.*, 65, 2443–2459, <https://doi.org/10.1002/LNO.11463>, 2020.
- 520 Stawiarski, B., Otto, S., Thiel, V., Gräwe, U., Loick-Wilde, N., Wittenborn, A. K., Schloemer, S., Wäge, J., Rehder, G., Labrenz, M., Wasmund, N., and Schmale, O.: Controls on zooplankton methane production in the central Baltic Sea, 16, 1–16, <https://doi.org/10.5194/BG-16-1-2019>, 2019.
- van de Waal, D. B. and Litchman, E.: Multiple global change stressor effects on phytoplankton nutrient acquisition in a future ocean, *Philos. Trans. R. Soc. B*, 375, <https://doi.org/10.1098/RSTB.2019.0706>, 2020.
- Wang, W. L., Moore, J. K., Martiny, A. C., and Primeau, F. W.: Convergent estimates of marine nitrogen fixation, *Nat.* 2019 5667743, 566, 205–211, <https://doi.org/10.1038/s41586-019-0911-2>, 2019.
- 525 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol. Oceanogr. Methods*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.
- Weber, T., Wiseman, N. A., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nat. Commun.*, 10, 1–10, <https://doi.org/10.1038/s41467-019-12541-7>, 2019.
- 530 Weller, D. I., Law, C. S., Marriner, A., Nodder, S. D., Chang, F. H., Stephens, J. A., Wilhelm, S. W., Boyd, P. W., and Sutton, P. J. H.: Progress in Oceanography Temporal variation of dissolved methane in a subtropical mesoscale eddy during a phytoplankton bloom in the southwest Pacific Ocean, *Prog. Oceanogr.*, 116, 193–206, <https://doi.org/10.1016/j.pocean.2013.07.008>, 2013.
- Williams, R. G., McDonagh, E., Roussenov, V. M., Torres-Valdes, S., King, B., Sanders, R., and Hansell, D. A.: Nutrient streams in the North Atlantic: Advective pathways of inorganic and dissolved organic nutrients, *Global Biogeochem. Cycles*, 25, n/a-n/a, <https://doi.org/10.1029/2010GB003853>, 2011.
- 535 Ye, W., Zhang, G., Zhu, Z., Huang, D., Han, Y., and Wang, L.: Methane Distribution and Sea-to-Air Flux in the East China Sea During the Summer of 2013 : Impact of Hypoxia Deep-Sea Research II Methane distribution and sea-to-air flux in the East China Sea during the summer of 2013 : Impact of hypoxia, *Deep. Res. Part II*, 124, 74–83, <https://doi.org/10.1016/j.dsr2.2015.01.008>, 2015.
- 540 Zheng, Y., Harris, D. F., Yu, Z., Fu, Y., Poudel, S., Ledbetter, R. N., Fixen, K. R., Yang, Z. Y., Boyd, E. S., Lidstrom, M. E., Seefeldt, L. C., and Harwood, C. S.: A pathway for biological methane production using bacterial iron-only nitrogenase, *Nat. Microbiol.*, 3, 281–286, <https://doi.org/10.1038/s41564-017-0091-5>, 2018.

