

Burned Area and Carbon Emissions Across Northwestern Boreal North America from 2001-2019

Stefano Potter¹, Sol Cooperdock^{1,2}, Sander Veraverbeke³, Xanthe Walker⁴, Michelle C. Mack⁴,
Scott J. Goetz⁵, Jennifer Baltzer⁶, Laura Bourgeau-Chavez⁷, Arden Burrell¹, Catherine Dieleman⁸
5 Nancy French⁷, Stijn Hantson⁹, Elizabeth E. Hoy^{10,17}, Liza Jenkins⁷, Jill F. Johnstone¹¹, Evan S.
Kane¹², Susan M. Natali¹, James T. Randerson¹³, Merritt R. Turetsky¹⁴, Ellen Whitman¹⁵,
Elizabeth Wiggins¹⁶, and Brendan M. Rogers¹

¹Woodwell Climate Research Center, Falmouth, MA, 02540 USA

²University of California, Los Angeles, CA, 90095 USA

10 ³Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

⁴Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ, 86011
USA

⁵School of Informatics, Computing, and Cyber Systems, Northern Arizona University, Flagstaff,
AZ, 86011 USA

15 ⁶Wilfrid Laurier University, Waterloo, ON, Canada

⁷Michigan Tech Research Institute, Ann Arbor, MI, 48105 USA

⁸University of Guelph, Guelph, ON, Canada

⁹Universidad del Rosario, Bogotá, Cundinamarca, Colombia

¹⁰NASA Goddard Space Flight Center, Greenbelt, MD, 20771 USA

20 ¹¹UAF: Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK, 99775 USA

¹²Michigan Tech University, Houghton, MI, 49931 USA

¹³Department of Earth System Science, University of California, Irvine, CA, 92697 USA

¹⁴Institute of Arctic and Alpine Research, University of Colorado, Boulder CO, 80309 USA

25 ¹⁵Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, AB,
Canada

¹⁶NASA Langley Research Center, Hampton, VA, 23666 USA.

¹⁷GST: NASA Goddard Space Flight Center/Global Science & Technology, Inc

Correspondence to: Stefano Potter (spotter@woodwellclimate.org)

30

35

40 **Abstract.** Fire is the dominant disturbance agent in Alaskan and Canadian boreal ecosystems and releases
large amounts of carbon into the atmosphere. Burned area and carbon emissions have been increasing with
climate change, which have the potential to alter the carbon balance and shift the region from a historic sink
to a source. It is therefore critically important to track the spatiotemporal changes in burned area and fire
carbon emissions over time. Here we developed a new burned area detection algorithm between 2001 -
45 2019 across Alaska and Canada at 500 meters (m) resolution that utilizes finer-scale 30 m Landsat imagery
to account for land cover unsuitable for burning. This method strictly balances omission and commission
errors at 500 m to derive accurate landscape- and regional-scale burned area estimates. Using this new
burned area product, we developed statistical models to predict burn depth and carbon combustion for the
same period within the NASA Arctic-Boreal Vulnerability Experiment (ABoVE) core and extended
50 domain. Statistical models were constrained using a database of field observations across the domain and
were related to a variety of response variables including remotely-sensed indicators of fire severity, fire
weather indices, local climate, soils, and topographic indicators. The burn depth and aboveground
combustion models performed best, with poorer performance for belowground combustion. We estimate
2.37 million hectares (Mha) burned annually between 2001-2019 over the ABoVE domain (2.87 Mha
55 across all of Alaska and Canada), emitting 79.3 +/- 27.96 (+/- 1 standard deviation) teragrams of carbon
(C) per year, with a mean combustion rate of 3.13 +/- 1.17 kilograms C m⁻². Mean combustion and burn
depth displayed a general gradient of higher severity in the northwestern portion of the domain to lower
severity in the south and east. We also found larger fire years and later season burning were generally
associated with greater mean combustion. Our estimates are generally consistent with previous efforts to
60 quantify burned area, fire carbon emissions, and their drivers in regions within boreal North America;
however, we generally estimate higher burned area and carbon emissions due to our use of Landsat imagery,
greater availability of field observations, and improvements in modeling. The burned area and combustion
data sets described here (the ABoVE Fire Emissions Database, or ABoVE-FED) can be used for local to
continental-scale applications of boreal fire science.

65

1 Introduction

Fire is the dominant disturbance agent in boreal forests (Stocks et al., 2003) and places large controls on ecosystem dynamics including vegetation composition and structure, nutrient cycling, permafrost, and carbon cycling (Bonan and Shugart, 1989; Bond-Lamberty et al., 2007; Walker et al., 2019). Fire frequency, intensity and burned area have been increasing in Alaskan and Canadian boreal forests over the last several decades (Hanes et al., 2018; Kasischke et al., 2010; Veraverbeke et al., 2017), and these trends are expected to continue throughout the 21st century due to a warmer and drier climate (Balshi et al., 2009; Boulanger et al., 2018; Young et al., 2017). Changes to the fire regime have been associated with more severe fires, which burn deeper into the organic soil profile and may be related to large fire years and seasonal timing of burn (Turetsky et al., 2011), although this has not been tested widely. Ultimately, changes in the fire regime have the potential to transition at least some North American boreal forests from a carbon sink to a source (Dieleman et al., 2020; Li et al., 2017; Walker et al., 2019; Wang et al., 2021). To better understand how changing boreal fire regimes influence carbon dynamics, it is critical to accurately map burned area and estimate resulting carbon emissions over time.

80

Burned area mapping in Alaska and Canada over long time frames (> 20 years) has primarily been based on digitized maps of fire observations (both by hand and in recent decades using GPS, aerial imagery, and satellite remote sensing) from the Alaska Large Fire Database (ALFD; Kasischke et al., 2002), the Canadian National Fire Database (CNFD; Amiro et al., 2001; Stocks et al., 2003), and more recently the Canadian National Burned Area Composite (NBAC; Hall et al., 2020). These databases are updated annually in Alaska and Canada, yet substantial uncertainty remains, particularly as the databases go further back in time and aerial and satellite imagery was less prevalent. Of particular importance is the possibility of commission errors because the databases do not typically account for unburned patches of vegetation and water bodies within the fire perimeters, leading to an overestimation of burned area (Skakun et al., 2021). At the same time, the databases are more likely to omit fires due to lost records or missed detections in

90

earlier decades (Kasischke et al., 2002; Stocks et al., 2003), leading to omissions. Mapping fire perimeters in recent decades has improved with the use of satellite remote sensing, particularly from 30 meter (m) Landsat (Epp and Lanoville, 1996) and 500 m Moderate Resolution Imaging Spectroradiometer (MODIS) imagery. While MODIS imagery is at coarser resolution than Landsat, its multiple acquisitions per day are highly amenable to burned area mapping, although there are known omission errors due to small (< 100 ha) burns as well as an overestimation of burned area at the pixel-level due to the relatively coarse 500 m resolution, which misses some unburned vegetation patches and water bodies (Giglio et al., 2018). Landsat imagery can largely bypass these issues of spatial resolution (Guindon et al., 2018; Walker et al., 2018), but the relatively infrequent overpass times and typical cloudy environments in the tundra and boreal biome result in data gaps, particularly prior to the launch of Landsat 7 (1999) due to data relay issues and limited tasking.

Traditionally, carbon emissions from wildfires have been calculated as a function of burned area, fuel consumption and emission factors (French et al., 2011; Seiler and Crutzen, 1980). Carbon emissions in these models are based on observed relationships between fuel consumption, fire weather, and fuel type. Current models that are built with this framework include the Wildland Fire Emissions Information System (WFEIS; French et al., 2011, 2014), the Fire Inventory from NCAR (FINN; Wiedinmyer et al., 2011) and the Global Fire Emission Database (GFED; van der Werf et al., 2017). In addition to these regional and global products, there are several model products that provide estimates in boreal ecosystems of Alaska (French et al., 2002; Kasischke & Hoy, 2012; Tan et al., 2007; Veraverbeke et al., 2015) and Canada (Amiro et al., 2001; de Groot et al., 2007). Researchers have also made improvements to process-based models' representation of fire occurrence and effects (Hantson et al., 2016; Rabin et al., 2017; Zhao et al., 2021). These models can be used to explore causal relationships and have the benefit of estimating how burn rates and carbon emissions may vary under differing future climate change scenarios.

In addition to simple empirical and process-based models of carbon combustion, several recent studies have implemented statistical techniques to model combustion based on field observations, satellite remote sensing imagery, and other geospatial data (Dieleman et al., 2020; Rogers et al., 2014; Veraverbeke et al., 2015, 2017; Walker et al., 2018). These advances are possible due to the increasing volume of field
120 observations of combustion, and have the advantages of unraveling complex relationships between combustion observations and geospatial information to extrapolate over space and time. Satellite imagery collected both pre-fire and post-fire has been particularly useful for these techniques, (Hudak et al., 2007; Key and Benson, 2006). Specifically, the differenced normalized burn ratio (dNBR) combines the near and short-wave infrared bands obtained before and after a fire, and the spectral information retained is sensitive
125 to reductions in vegetation and moisture content post-fire. Due to these qualities dNBR correlates relatively strongly with aboveground biomass loss, but there have been conflicting findings on the strength of the relationship with belowground fire severity, which are particularly important in boreal ecosystems (Kasischke and Hoy, 2012; McGuire et al., 2009). Additional environmental predictors have been combined with dNBR to statistically model above and belowground combustion across Alaska and Canada, including
130 quantified uncertainties (Dieleman et al., 2020; Rogers et al., 2014; Veraverbeke et al., 2015, 2017; Walker et al., 2018). Veraverbeke (2015) found topographic variables (elevation, slope, northness), pre-fire vegetation cover (% tree cover) and day of burning to be important predictors for both aboveground and belowground combustion, and more specifically the combination of dNBR, day of burning, elevation and tree cover to be the most informative in Alaska. Walker (2018) considered 71 variables associated with
135 topography, permafrost condition, fire severity, fire weather and soil properties, and found that dNBR, change in pre- and post-fire tree cover, terrain ruggedness, topographic wetness, percent black spruce and percent sand were the most informative for the 2014 Northwest Territories fires. Although these results have been encouraging, extrapolations have been limited to specific regions in Canada and Alaska, and often to specific fire years. It is likely that the inclusion of additional field data across a more representative
140 selection of field locations in Alaska and Canada would improve model fits and allow for extrapolation over a larger domain and longer time periods.

In this study we first derived a new 500 m burned area product for all of Alaska and Canada during 2001-
145 2019. Our approach builds on previous satellite-based burned area mapping efforts (Chen et al., 2020;
Dieleman et al., 2020; Loboda et al., 2018; van der Werf et al., 2017; Veraverbeke et al., 2015; Walker et
al., 2018) with 500 m MODIS data, but advances these by using 30 m Landsat imagery to both improve
accuracy and account for the presence of unburnable land cover. Using this burned area product, along with
a new comprehensive database of combustion observations in Alaska and central/western Canada (Walker
150 et al., 2020a), we used machine learning to estimate burn depth and fire carbon emissions across the ABoVE
domain. We compare our product to a suite of previous efforts, and use it to test previously hypothesized
relationships between fire severity, annual burned area, and seasonal timing of burning.

2 Methods

2.1 Study Area

The spatial domain of this study includes all of Alaska and Canada for our burned area product, and the
| Arctic Boreal Vulnerability Experiment (ABoVE) core and extended domain (hereafter the “ABoVE
domain”; Loboda et al., 2019) for our combustion and burn depth product (Figure 1). The combustion and
burn depth products were not derived beyond the ABoVE domain due to a lack of field observations in
160 eastern Canada. The temporal domain for all products is 2001-2019. Our study area includes all natural
boreal and arctic vegetation within the ABoVE domain, including boreal forests, boreal wetlands,
grasslands, tundra, and tundra wetlands. To determine these locations we derived a vegetation mask using
the 2005 Land Cover of North America product (250 m; CCRS, 2013; Pouliot and Latifovic, 2013; Pouliot
et al., 2014), MODIS land cover type with International Geosphere-Biosphere Programme (IGBP)
165 classification (Collection 6; year 2005; 500 m; Friedl and Sulla-Menashe, 2019), the Circumpolar Arctic
Vegetation Map (CAVM; Raynolds et al., 2019) and long-term climate (1970 – 2000; ~1 km; Fick and
Hijmans, 2017), all re-gridded to 500 m resolution on the MODIS sinusoidal projection (Figure S1). Boreal

vegetation was distinguished from temperate using a mean annual temperature threshold of 3°C, as recommended in Wolfe (1979) and implemented in Rogers et al. (2015). Pixels were designated as urban, crop, crop/natural vegetation mosaic, or water if they were represented as such in either the Land Cover of North America or MODIS land cover products. Pixels were designated as tundra if they were within the CAVM domain.




Figure 1. Study domain. Locations of combustion observations (red), the burned area product domain (light gray), and the combustion and burned depth product domain (dark gray).

2.2 Field Data

Field measurements of burn depth and combustion were derived from numerous data sources across different research groups that represent a major synthesis effort sponsored by the NASA ABoVE program (Boby et al., 2010; Dieleman et al., 2020; de Groot et al., 2009; Hoy et al., 2016; Rogers et al., 2014; Turetsky et al., 2011; Veraverbeke et al., 2015; Walker et al., 2018). Detailed descriptions of data collection methods can be found in the contributing publications. All field site information was standardized and

aggregated into a single publicly-available database (Walker et al., 2020a), which has been used to assess patterns and drivers of ecosystem structure and combustion across ecoregions (Walker et al., 2020b; Walker et al., 2020c). Although the field database only includes measurements from boreal ecosystems, our combustion and burn depth predictions include both boreal and tundra ecosystems. Of all the pixels for which we predicted combustion and burn depth, only 0.78% are in tundra landscapes.

2.3 Burned Area Mapping

The ABoVE Fire Emissions Database (ABoVE-FED) burned area product is derived from a dNBR thresholding approach, which has previously been successfully employed for burned area mapping in the region (Rogers et al., 2014; Veraverbeke et al., 2015; Walker et al., 2018). Our primary approach was to use Landsat imagery to separate burned from unburned pixels at 30 m. However, because Landsat imagery was not available for all regions and time periods, we used MODIS imagery to map burned pixels when necessary, and upscaled our Landsat-based product to 500 m MODIS resolution. More specifically we used pre- and post-fire near infrared (NIR) and shortwave infrared (SWIR) bands from Aqua (MYD09GA Collection 6; Vermote, 2015a) Terra (MOD09GA Collection 6; Vermote, 2015b) and Landsat 5-8, calculating dNBR as the difference in pre fire normalized burn ratio (NBR) and post fire NBR, where NBR is near infrared minus shortwave infrared divided by near infrared plus shortwave infrared.

This approach had the added advantage of accuracy; whereas a Landsat dNBR threshold tends to be surpassed at the site level in a diffuse manner across the landscape, due to stochastic site-level disturbances such as tree mortality, herbivory, flooding, or small-scale dieback, it is much less common for these small-scale disturbances to influence the majority of a 500 m pixel. We also minimized mapping non-fire disturbances by following the approach of Veraverbeke et al. (2015) and applying our dNBR approach to (1) mapped fire polygons from the ALFD and CNFD (93% of total burned pixels; hereafter collectively referred to as the National Large Fire Databases (NLFD)) and (2) MODIS active fire (MOD14A1 Collection 6 and MYD14A1 Collection 6; Giglio et al., 2018) acquisitions outside these polygons (7% of total burned

220 pixels). In each case we applied a 1 km buffer (Veraverbeke et al., 2015) to capture burned pixels immediately outside these areas. Finally, our approach is motivated by a desire to balance commission and omission errors at both the 30 m and 500 m scales, thereby providing an unbiased estimate of total burned area.

225 To map 30 m burned pixels, we first extracted dNBR at both burned and ~~unburned control sites in our~~ aggregated field database using available cloud-free Landsat 5, 7 and 8 Tier 1 surface reflectance images in Google Earth Engine (Gorelick et al., 2017). Landsat 5 and 7 were atmospherically corrected using the Landsat Ecosystem disturbance adaptive processing system (LEDPAS; Schmidt et al., 2013) while Landsat 8 was atmospherically corrected using Land Surface Reflectance Code (LaSRC; Vermote et al., 2016). Pre-
230 and post-fire normalized burn ratio (NBR) was calculated as the mean of all available Landsat observations between July and August. Pre-fire values were extracted one year before a given fire, and post-fire values were extracted one year after a fire. We then selected a 30 m Landsat dNBR threshold that most effectively separated burned and unburned control sites. Because there are many fewer unburned control sites in the Walker et al (2020a) combustion database, we derived additional control sites by extracting dNBR at burned
235 sites two years before a given fire, which had the advantage of controlling for any site-level spectral differences between burned and control sites represented in the database. This process generated a dNBR threshold of 0.084, which minimized 30 m site-level commission and omission errors to 6.6% (Figure S2).

We then created a mask at 30 m to account for unburnable land cover (i.e., non-vegetated pixels). This was
240 created using two sources: the Joint Research Center's yearly water history product (Pekel et al., 2016) and the North American Land Change Monitoring System's (NALCM) 2010 land cover product at 30 m resolution (Latifovic et al., 2012). The first product allowed us to capture transient water pixels in our time series while the NALCM land cover product classified each pixel into 19 different land cover classes, from which we masked out non-vegetated pixels including ice, water, barren and cropland. These two sources
245 were combined into separate masks for each year between 2001-2019. Because areas that burned in 2010

Deleted: paired

were often classified as barren lands in the 2010 NALCM product, we considered barren lands as vegetated in our mask for the year 2010.

250 Using the vegetation mask and the dNBR threshold, we created a binary burned/unburned 30 m Landsat product and upscaled this to the native MODIS 500 m resolution and projection. To determine whether or not a given 500 m pixel was classified as burned or unburned, we calculated the percentage of 30 m vegetated pixels that burned within its footprint. If more than 50% of the 30 m vegetated pixels within the larger 500 m pixel burned (i.e., were tripped by the dNBR threshold), the entire pixel was assigned as
255 burned, and the burned fraction was calculated as the percent of the burnable land cover (vegetation) in the 500 m pixel. Note we did not use the percent of burned 30 m pixels to determine burn fraction within a given 500 m pixel, primarily because of limitations imposed by frequently missing Landsat imagery (detailed below).

260 We used this approach whenever 500 m pixels contained 100% coverage by Landsat imagery at 30 m. When, however, there was less than 100% Landsat coverage, we needed to determine if it was more accurate to classify 500 m pixels using Landsat (with partial coverage) or MODIS Collection 6 imagery (Vermote, 2015a, Vermote, 2015b). To do so, we analyzed all MODIS pixels with complete Landsat coverage and masked out increasing numbers of Landsat pixel strips within the larger MODIS footprint (using increments
265 of 5%). After each removal of Landsat pixels, we compared the accuracy of the resulting burned/unburned classification using (i) Landsat imagery with partial coverage and (ii) MODIS imagery. This procedure suggested that using MODIS dNBR was more accurate than Landsat when less than 85% of a 500 m MODIS pixel was covered by Landsat imagery. We therefore used Landsat to classify burned pixels when at least 85% of a 500 m pixel was covered by Landsat imagery, and otherwise used MODIS. Burned pixels
270 were assigned a quality flag of zero when there was complete Landsat coverage, a quality flag of one when Landsat coverage was less than 100% but greater than 85%, and a quality flag of two when Landsat coverage was less than 85% and therefore MODIS imagery was used to classify burn status. Overall, 81%

of total burned pixels were derived using Landsat (66% from full coverage and 15% from partial coverage), although particular regions (notably Alaska and Newfoundland and Labrador) tend to rely more on MODIS
275 due to more limited availability of Landsat imagery (Figure S3).

We developed a correction factor for MODIS-based dNBR to account for differences between Landsat and MODIS NIR and SWIR spectra, as well as the influence of vegetation fraction on 500 m dNBR signals. To do so, we calculated pre- and post-fire NIR and SWIR bands from MODIS and Landsat (resampled to 500
280 m) for a 50% random sample of burned pixels. We then differenced the Landsat 500 m resampled bands from the 500 m MODIS bands and regressed them onto vegetation fraction to obtain a correction factor. The regression yielded an R^2 of 0.74 and an equation of $y = 0.94x + 0.01$, which was applied to all pixels where burn status was classified by MODIS. We then calculated a new dNBR threshold to classify pixels at 500 m in an unbiased manner. To do so, we determined the MODIS dNBR threshold that evenly split
285 omissions and commissions based on pixels mapped with complete Landsat coverage. This threshold was determined to be 0.0725, resulting in an omission/commission error of 14.2% at 500 m when using MODIS.

One issue with a burned area mapping approach such as ours that utilizes post-fire imagery one year after a fire is that it is difficult to determine the year(s) of burn where overlapping burns occurred in successive
290 years. To address these cases, we created a seasonal MODIS-based product following the methodology of Giglio et al. (2018). The dNBR for each day between January 15th and December 15th was calculated using the 30 preceding days as pre-fire NBR and the 30 days after as post-fire NBR. Any pixels with less than 10 valid observations in either window were masked out. We used a similar thresholding approach to that described above for mapping burned pixels with MODIS, resulting in a seasonal dNBR threshold of 0.23.
295 Any pixel mapped using the MODIS seasonal approach was assigned a quality flag of three.

In addition to determining fire locations, fire year, and the burned fraction, we also determined the day of burning for each pixel. When possible, day of burn was taken directly from the thermal anomaly active fire

300 detections from MOD14A1 Collection 6 and MYD14A1 Collection 6 (Giglio et al., 2018) active fire products. Where an active fire was registered, day of burn was assigned by taking the earliest active fire acquisition during the year. When an active fire was not registered for a given burned pixel, we utilized a multi-tiered approach to assign day of burn. When possible, we used a kriging technique to interpolate day of burn using the active fire detections within each fire polygon in the NLFD following Veraverbeke et al. (2015). To implement this, we required fire polygons to contain at least five active fire acquisitions within their boundaries and have some level of temporal variation (i.e., not all active fire acquisitions on the same day). When this was not the case, day of burn was assigned using the closest active fire pixel. Finally, when no active fire acquisitions were associated with a given fire polygon, we used our MODIS-based seasonal mapping approach to determine day of burn by locating the day of maximal dNBR within a given year. For fires that were detected by MODIS thermal anomalies but were not contained in the NLFD (7% of all burned area), we created our own polygons around the burned pixels (by converting pixels to vectors and buffering them) and used the same method to assign day of burn. Quality flags for our burn day product represent this tiered approach, with a flag of zero for pixels with direct active fire hits, a flag of one for pixels whose day of burn was determined by interpolation, and a flag of two for pixels whose day of burn was determined using the MODIS seasonal burned area product. A simplified flowchart of burned area processing methods is shown in Figure S4.

320 We compared ABoVE-FED burned area to several other products including the NLFD, NBAC, MCD64A1 Collection 5, MCD64A1 Collection 6, the Alaska Fire Emissions Database version 2 (AKFED; Veraverbeke et al., 2017), GFED4s (van der Werf et al., 2017), [a 500 m model by van Wees et al. 2022](#), and the Fire Model Intercomparison Project (FireMIP; Hantson et al., 2016; Rabin et al., 2017; Table S1). NBAC is a Canada-only product and is related to the CNFD, but improves upon it by incorporating multi-sensor remote sensing imagery (including Landsat) to account for water bodies and unburned vegetation patches. FireMIP includes simulations performed by coupled fire-vegetation models forced with a standardized set of input data. We also visually compared our product and others to high-resolution imagery

325 of fires from the Worldview 2 (1.84 m) satellite, available through DigitalGlobe, Inc., a Maxar Company
under the Nextview license agreement through the National Geospatial Intelligence Agency (Neigh et al.,
2013).

2.4 Combustion and Burn Depth Models

330 We built and applied statistical models of aboveground combustion, belowground combustion, and burn
depth to every mapped burned pixel in the ABoVE domain based on field observations across Alaska and
western Canada (Walker et al., 2020a). Because not all field sites included estimates of both above- and
belowground combustion, we created two separate combustion models, one utilizing all available
aboveground combustion measurements ($n = 515$) and one utilizing all available belowground combustion
335 measurements ($n = 769$). Our burn depth model utilized the same field sites as belowground combustion.
Further discussion of models implemented can be found in the Supplement.

2.4.1 Predictor variables

Combustion and burn depth measurements from Walker et al. (2020a) were related to a variety of spatial
340 predictors including remotely sensed indicators of fire severity, topography, soils, climate and fire weather.
We initially acquired 75 covariates associated with environmental conditions such as long-term climate,
fire weather, topography, vegetation type, soil type, remotely sensed vegetation indices (e.g. normalized
difference vegetation index; NDVI; Tucker, 1979) and permafrost condition (Table S2).

2.4.2 Climate Variables

Long-term climate was acquired from ClimateNA (CNA; Wang et al., 2016; Table S2), which provides
point estimates of mean climate from 1981 – 2010 based on the Climate Research Unit (CRU; Mitchell and
Jones, 2005). ClimateNA uses finer-resolution PRISM (Daly et al., 2002, 2008) and ANUSPLIN
(Hutchinson, 1989) climate normals to downscale coarse-resolution monthly climate data to a 4×4 km
350 grid, followed by bilinear interpolation and a locally-derived elevation adjustment to estimate point data.

CNA variables were represented as both annual and summer means (June – August), and were included to capture the influence of long-term climate on vegetation, fuel loads, and fuel moisture that drive combustion (Walker, et al., 2020b).

2.4.3 Fire Weather Indices

Fire weather indices (FWIs) represent the meteorology at the timing of fire occurrence and have been associated with fire behavior and carbon emissions due to their influence on fuel moisture and fire spread (e.g. Di Giuseppe et al., 2018; French et al., 2011; Ivanova et al., 2011; Veraverbeke et al., 2017). We acquired FWIs from the Global Fire Weather Emissions Database (GFWED v2.0; Field et al., 2015) at $0.5^\circ \times 0.66^\circ$ resolution. FWI information was extracted for the day of burn for all fires in the field database. Since FWI data were not available for all burned pixels in our fire product due to missing data in the shoulder seasons, we developed two versions of our aboveground combustion, belowground combustion, and burn depth models: a primary model that included FWIs in training and a secondary one that did not. Mapped pixels from the primary model were assigned a quality flag of zero, and pixels from the secondary model were assigned a flag of one. Of the 2,123,730 pixels that burned between 2001 – 2019, 4.4% did not have FWI data available and necessitated the use of these secondary models.

2.4.4 Environmental Variables

We acquired a variety of environmental covariates related to soils, topography, vegetation type, and permafrost occurrence (Table S2). Soil properties were taken from SoilGrids at 250 m resolution (Hengl et al., 2017), including percent clay (0 – 2 μm), silt (2 – 50 μm), sand (50 – 2000 μm), coarse material (> 2000 μm), bulk density (g cm^{-3}), soil organic carbon stock (tons ha^{-1}), and soil water pH. We integrated all variables across the top 30 cm of the soil profile.

Topographic variables, including elevation (m), aspect ($^\circ$), and slope ($^\circ$), were derived from a 10-meter digital elevation model (DEM) of the ABoVE domain which, in turn, was derived from a higher resolution

Arctic DEM (Porter et al. 2018) and gap-filled with additional DEM datasets (Burns et al., forthcoming). This 10 m DEM was resampled to 500 m and then aspect and slope were both calculated as the local gradient of the four connected neighbors of each pixel. After resampling to 500 m we also calculated a topographic wetness index (TWI) for each pixel that represents soil drainage patterns based on the slope and upslope area draining through a particular point (Beven & Kirkby, 1979).

Vegetation type was represented by the percent cover over seven broad classes including black spruce (*Picea mariana*), white spruce (*Picea glauca*), jack pine (*Pinus banksiana*), deciduous broadleaf species, other conifers, grasslands and non-vegetated (Beaudoin et al., 2014; Ottmar et al., 2007). We use pre-fire tree cover (Sexton et al., 2013) from either 2000, 2005, 2010, or 2015, depending on fire year.

Lastly, we acquired a permafrost zonation and a surface roughness index, which is a measure of terrain complexity (Gruber, 2012).

2.4.5 Remotely Sensed Variables

We derived numerous remotely-sensed vegetation indices from Landsat, including the NDVI, the normalized difference infrared index (NDII; Hardinsky & Smart, 1983), dNBR (Key and Benson, 2006), the relative difference normalized burn ratio (RdNBR; Miller and Thode, 2007), the relativized burn ratio (RBR; Parks et al., 2014), tasseled cap greenness, wetness and brightness (Kauth and Thomas, 1976), and pre-fire tree cover (Sexton et al., 2013). NDVI, NDII, and tasseled cap indices were acquired as a mean composite between May 15th and June 15th in the post-fire years, while dNBR, RdNBR and RBR were based on mean composites between June 1st and August 31st for both the pre- and post-fire years.

For model training all remotely sensed variables were extracted from Landsat 5-8 Tier 1 surface reflectance at 30 m with clouds, cloud shadows and snow masked out using the C Function of Mask (CFMask; Foga et al., 2017). We applied corrections due to spectral differences between Landsat 8 and 7 using a regression technique (Roy et al., 2016). Although our model was trained with Landsat imagery at 30 m, we predicted

combustion and burn depth at 500 m across the domain using MODIS imagery. All MODIS variables were extracted in Google Earth Engine at ideal MODIS quality flags (bit flag of 0). We then implemented a correction factor to account for sensor and spatial scaling issues in model predictions (section 2.4.7).

2.4.6 Feature Selection and Model Comparisons

We reduced our initial 75 covariates to an optimal number using recursive feature elimination (Guyon et al., 2002). Recursive feature elimination iteratively removes variables until a desired number remains, which in this case is defined by the number of covariates necessary to achieve the minimum root mean square error (RMSE). Recursive feature elimination achieves this by fitting a secondary machine learning model that can rank features by importance and discards the least important ones at each iteration. We used random forest (Breiman, 2001) as the measure of importance, and repeated our recursive feature elimination three times across a five-fold cross-validation to determine the optimal subset of covariates (Table S2). For the primary aboveground combustion, belowground combustion, and burn depth models, the optimal number of variables was 15, 45, and 40 (Figure S5), respectively, and for the secondary models the optimal number of variables was 15, 64, and 48. While it is possible a similar RMSE could have been achieved with reduced model complexity (reduced number of variables), we chose to directly use RMSE reduction as our threshold for feature selection.

We then tested a suite of statistical models across the selected feature space to compare predictive power. For each model, we searched for optimal model parameters using a 10-fold cross-validation repeated three times and a random search grid of length 10 (e.g. for any given model parameter, 10 random numbers were selected per parameter and tested for each parameter combination). After optimizing model parameters, we compared final model fits with a 10-fold cross-validation repeated 100 times. After comparing the median R^2 for each model across these 1,000 iterations, we selected the best performing model and chose it for the final model implementation. All model training took place in R (R Core Development Team, 2021). In all

cases the best performing model was a ranger random forest, although there were differences in the optimal parameters chosen (Table S3).

2.4.7 Spatial Scaling

Our combustion and burn depth models were developed using site-level data (most plots utilized a 30 × 30 m design) and geospatial predictors at their native resolution, including a variety of 30 m Landsat indices. However, our spatial model was applied at 500 m to match the resolution of our burned area product, ultimately because missing imagery prevented comprehensive burned area mapping at 30 m. To explore potential issues associated with implementing the model at these different spatial scales, we randomly sampled two hundred 500 m pixels from each year in 2004, 2006, 2012, 2014 and 2015, for a total of 1,000 pixels. We then implemented our combustion and burn depth models at both 30 m and 500 m to assess biases and errors introduced by both spatial and sensor differences. When models were assessed at 30 m, all predictor variables were acquired at their native resolutions (Table S2); when models were assessed at 500 m, all variables were resampled to 500 m. Any variables described in section 2.4.5 that were derived from Landsat were instead collected at 500 m from MODIS (using MOD09A1 Collection 6 and MYD09A1 Collection 6). We used MODIS provided quality flags to select for pixels that were corrected at ideal quality and masked out clouds and snow. All other variables were resampled to 500 m using bilinear interpolation if native resolution was >500 m, and using mean values within pixel boundaries if native resolution was <500 m. We then compared the predictions at 500 m resolution to the mean across all the 30 m sub pixels, and built type 2 linear regression models to correct for potential biases. The coefficients from these models were then used to adjust the final predictions for the combustion models across the full domain.

2.4.8 Combustion and Burned Depth Predictions and Quality Flags

Predictor variables for all burn pixels across the domain were collected in Google Earth Engine. Since the ideal MODIS quality flag criteria (section 2.4.5) left 0.31% of the total burned pixels missing, we collected predictors for these pixels with no MODIS quality flag applied, and assigned our own quality flag to

distinguish these samples. We provide four separate quality flags indicating whether our primary or secondary models (no FWIs) were implemented, and whether MODIS quality flags were applied. Our four flags have the following associations: Flag one – primary model with MODIS quality flag criteria (95.32 % of pixels); Flag two - primary model with no MODIS quality flag criteria (0.26 % of pixels); Flag three – secondary model with MODIS quality flag criteria (4.37 % of pixels); Flag four - secondary model with no MODIS quality flag criteria (0.05% of pixels).

2.4.9 Monte Carlo Analysis

To derive a measure of prediction uncertainty, we implemented a Monte Carlo analysis with 500 simulations that incorporated uncertainty from both the field-measured combustion and the random forest models. Our approach was based on techniques implemented in Rogers et al. (2014), Veraverbeke et al. (2015), Walker et al. (2018), and Dieleman et al. (2020). To account for uncertainty in field estimates of belowground combustion, we used the standard error of observed site-level combustion when it was available. In total, 271 field sites recorded standard error: 22 in Alaska, 47 in Saskatchewan, and 202 in the Northwest Territories. Standard error was estimated for both aboveground and belowground combustion in Alaska and Saskatchewan, and only for belowground in the Northwest Territories. For each Monte Carlo simulation, we derived an adjustment factor by multiplying a site's standard error by a random number from a normal distribution with a standard deviation of one and centered around zero. This resulting number was then added to the measured combustion.

Uncertainty in aboveground combustion in the Northwest Territories was calculated by first creating a random bias for the percent carbon content of trees (central estimate of 0.5), which varied randomly within a normal distribution with 3% standard deviation systematically across all trees measured for each Monte Carlo simulation (based on Rogers et al., 2014). We similarly included a 20% error in visual estimates of tree consumption (Dieleman et al., 2020; French, 2004; Walker et al., 2018), which also varied

480 systematically across all trees measured. Aboveground combustion in each simulation was then altered
using these adjustment terms (adding the carbon fraction adjuster and multiplying the tree consumption
adjuster).

Since these procedures only accounted for uncertainty of 271 of the possible samples, uncertainty for the
485 remaining 245 aboveground and 499 belowground samples was derived using an alternate approach. To do
so, we first linearly regressed the aboveground and belowground combustion standard error derived from
Monte Carlo simulations against measured aboveground and belowground combustion, respectively. The
coefficients from these two separate models were then used to predict the standard errors for all remaining
samples (Figure S6).

490 In addition to uncertainty in field measurements, there is also uncertainty in the random forest model used
to predict combustion across the ABoVE domain. To account for this, we leveraged the fact that model
residual errors tended to increase in proportion to combustion level, similar to Rogers et al. (2014) and
Dieleman et al. (2020). To estimate this relationship, we split the original model predictions (from the 10-
495 fold cross-validation repeated 100 times) into 15 bins based on quantiles of total combustion, and then
calculated the standard deviation of the residual error within each bin. We then used a general additive
model to smooth the standard deviation of the residuals across the bins (Figure S7). For each of the 500
Monte Carlo simulations using adjusted field estimates of combustion (derived from procedures described
above), new random forest model predictions were assigned a standard error based on total combustion
500 using the smoothed relationship. These standard errors were then multiplied by a random bias factor with
a standard deviation of one centered around zero, which was then added back onto the combustion
predictions to derive a final uncertainty estimate for each predicted combustion pixel across the ABoVE
domain.

505

We quantified uncertainty in our predictions in three ways: 1) pixel-level uncertainty, 2) uncertainty in mean combustion and 3) uncertainty in total emissions for a given region of interest. In each case, uncertainties derived from the Monte Carlo simulations were adjusted by the ratios of mean combustion from the primary model to that of the Monte Carlo simulations in order to account for different mean combustion levels, and hence emissions, between the models (which were minor). 1) Pixel-level uncertainty was calculated as the standard error of combustion for a given pixel across the Monte Carlo simulations. 2) Uncertainty in mean combustion for a given region was calculated as the standard error of mean combustion across the 500 Monte Carlo simulations for that region. In this case note that mean combustion was calculated by weighting pixels by their vegetated (burned) fractions. 3) Uncertainty in total emissions for a given a region of interest was calculated as the standard error of total emissions for that region across the 500 Monte Carlo simulations. A simplified flow chart of the combustion/burned depth modeling methodology is shown in Figure S8.

2.5 Relationships between belowground fire severity, annual burned area, and timing of burn

Turetsky et al. (2011) discovered a positive relationship between burn depth, annual burned area, and timing of burn (day of year) in black spruce forests and peatlands of interior Alaska, and also noted the influence of burn timing was more important in small fire years. To test if these relationships held true with a larger field database in Alaska ($n = 286$ for ABoVE-FED compared to $n = 178$ in Turetsky et al. (2011)), we performed a multiple regression of burn depth and belowground combustion using annual burned area and day of year as predictor variables. We also tested how burn depth and belowground combustion varied as a function of day of year within both small and large fire years. To do so, we split the field sites in Alaska into four quantiles based on annual burned area and then regressed burn depth and belowground combustion against day of year within each quantile. We also conducted this analysis using a sample of 500 ABoVE-FED pixels in Alaska instead of field observations, and then repeated both of these analyses using all available field observations and 500 random pixels within the broader ABoVE domain. In each case, we

sampled 500 pixels instead of using all available pixels to minimize the effect of large sample sizes on p-values.

3 Results

3.1 Burned Area

Temporally there was high variability in burned area year to year (Figure 2a). Across the domain, ABoVE-FED reported similar burned area totals compared to the NLFD (average of 2.87 Mha y^{-1} for ABoVE-FED compared to 2.90 Mha y^{-1} for NLFD; Figure 3), although there was variability in this relationship (NLFD estimated larger annual burned area in 11 years and smaller burned area in eight years between 2001-2019). This was the net result of two contrasting patterns: ABoVE-FED tended to report less burned area within mapped polygons, due to unmapped unburned patches and unburnable land cover (e.g., small water bodies) in the government fire databases, but detected additional burned areas associated with MODIS active fire acquisitions well outside mapped fire polygons (7% of total burned area in ABoVE-FED; 6% of total emissions; Figure S9). The state/territory with the most burned area detected outside the mapped polygons was British Columbia (31% of the 7% total burned area mapped outside NLFD polygons; Figure S9). Exploratory analysis revealed this was likely as a result of commission errors due to logging (i.e., logged areas tripping dNBR thresholds in conjunction with small fires registered by MODIS active fire hits). Across the domain, the mean fire size coincident with NLFD polygons was much larger (4,954 ha) than the mean fire size outside the polygons (166 ha). Because the NBAC product accounts for more of these unburned patches within polygons (Hall et al., 2020), it tended to report lower total burned area compared to ABoVE-FED (Figure S10). ABoVE-FED burned area was higher than MCD64A1 (Collection 5 and 6; Figure 3) in all years, which is consistent with known omissions in these global products for boreal North America (Giglio et al., 2018; Randerson et al., 2012; Figure S11). These large-scale patterns were corroborated by high-resolution imagery of particular fire events (Figure 4; Figure S12-S19). ABoVE-FED identified more burned pixels than MCD64A1 Collection 6 by being more sensitive to fire-induced spectral changes, but also accounted for unburnable portions of the landscape (Figure S20). GFED4s burned area

Deleted: substantially

was slightly higher (Figure S21; average of 2.38 Mha y⁻¹ during 2001-2016) than both MODIS products (Figure 3 ; average of 2.93 Mha y⁻¹ during 2001-2016), but lower than the NLFD and ABoVE-FED (Figure 3). The MCD64A1 Collection 5, Collection 6 and GFED4s databases underestimated burned area by 32, 23, and 18% compared to ABoVE-FED, respectively.

Deleted: , Figure S11; average of 2.93 Mha y⁻¹ for ABoVE-FED during 2001 – 2016 compared to 2.38 Mha y⁻¹ for GFED




Figure 2. Temporal variability in ABoVE-FED burned area (a), and emissions (b) from 2001-2019.




Figure 3. Comparison of ABoVE-FED burned area across Canada and Alaska to MODIS MCD64A1 Collection 5 (C5), MCD64A1 Collection 6 (C6), and the Alaskan and Canadian National Fire Databases (NFDB).

615

620

625

630

635




Figure 4. Comparison of high-resolution imagery and burned area products for a fire in Manitoba in 2014 (a). Panels show Worldview-2 imagery (b, fire shown in purple shades), ABoVE-FED (c), MODIS Collection 6 (d), MODIS Collection 5 (e), and the Canadian National Fire Database (f).

640

ABOVE-FED burned area was similar to AKFED where it is available (Alaska, the Northwest Territories and the Yukon Territory; Figure S22 ; average of 1.27 Mha y⁻¹ for ABOVE-FED during 2001 – 2015 compared to 1.22 Mha y⁻¹ for AKFED). All models participating in FireMIP simulated lower burned area than ABOVE-FED, and with a very high level of variability between models (1.34 ± 0.83 Mha y⁻¹ across Alaska and Canada during 2001 – 2012; Figure S23a).

645

Burned area was highly variable interannually, with the largest fire years occurring in 2004 in Alaska and the Yukon Territory; 2015 in Alaska, Saskatchewan, and Alberta; 2014 in the Northwest Territories; and 2013 in Manitoba and Quebec (Figure S21, S24). Across states, provinces, and territories, total burned area was highest in Alaska, the Northwest Territories and Saskatchewan. A total of 54 Mha burned across Alaska and Canada during all years, and 45 Mha in the ABOVE domain, with an annual mean of 2.87 Mha y⁻¹ across Alaska and Canada and 2.37 Mha y⁻¹ in the ABOVE domain.

650

Spatially ABOVE-FED estimated the most burned area in in Alaska, the Northwest Territories and Saskatchewan (Figure 5a; Figure S21, Figure S24)

655

660

665




Figure 5. Total burned area (a), total carbon emissions (b), mean combustion (c), and mean burn depth (d) between 2001-2019 aggregated to a 70 km grid. Note that burned area (a) covers all of Alaska and Canada, whereas all other metrics cover the ABoVE extended domain.

3.2 Combustion and Burn Depth Models

Our aboveground and belowground combustion models performed well, although the aboveground model performed significantly better across the suite of models examined (Figure 6a,b). A ranger random forest model (Wright and Ziegler, 2017) performed best for aboveground and belowground combustion, with median R^2 of 0.46 and 0.25, respectively, across the 10-fold cross validation repeated 100 times. Our secondary models that did not include information on FWIs (section 2.4.3) performed similarly to our primary models, with R^2 values for above and belowground combustion of 0.45 and 0.24, respectively. Although both the aboveground and belowground models performed reasonably well at predicting lower and moderate combustion values, which includes the majority of field observations, they both struggled to predict larger combustion values (Figure S25a,b). The burn depth model performed better than both combustion models, with a median R^2 of 0.53 using a ranger random forest model (Figure 6c).




Figure 6. Comparison of the spread and median R^2 values across a 10-fold cross validation repeated 100 times for our aboveground combustion (a), belowground combustion (b), and burn depth (c) models. Models compared include a ranger random forest (ranger), a quantile random forest (quantile), radial support vector machines (svmradial), polynomial support vector machines (svmpoly), linear support vector machines (svmlinear), ridge regression (ridge) and lasso regression (lasso).

745 There were notable differences in the feature importance of the aboveground and belowground models
(Figure S26a,b). The aboveground model was heavily influenced by its top predictor, pre-fire tree cover,
followed by metrics of relative humidity, with other variables including remotely-sensed fire severity and
vegetation moisture content having significant but relatively low importance. In contrast, the belowground
model was influenced strongly by a number of soil, terrain, climate, and tree cover variables. The most
750 important features for the burn depth model were similar to the belowground model, with soil properties,
tree cover and climate being the most influential (Figure S65c). Overall, the distribution of variables used
in the training dataset and predicting dataset were similar (Figure S27), with the exception of slope. Most
field sites were located in relatively flat terrain whereas the combustion predictions included locations with
steeper terrain.

755
Spatial patterns of mean burn depth and combustion tended to follow a gradient of higher burn depth and
mean combustion in the western part of the ABoVE domain (Alaska, Yukon Territory, and Alberta) to
lower mean combustion in central-western Canada (Saskatchewan, Northwest Territories, and Manitoba)
(Figure 5c,d, Figure S28). There was, however, considerable fine-scale variability at 500 m within these
760 regions (Figure 7), and spatial patterns were relatively consistent with previous combustion mapping
efforts.

765

770

775

780

785




Figure 7. Comparison of Alaskan fires in 2004 (a) for ABoVE-FED (b) and AKFED (c), the Northwest Territories fires in 2014 (d) for ABoVE-FED (e) and Walker et al., 2018 (f), and Saskatchewan fires in 2015 (g) for ABoVE-FED (h) and Dieleman et al., 2020 (i). Basemap Sources: Esri, © OpenStreetMap Contributors, HERE, Garmin, USGS, EPA, NPS, NRCran.

Across the ABoVE domain, 1.51 ± 0.53 Pg C was emitted over the 2001-2019 period, with a mean of 79.3 ± 27.96 Tg C per year. Mean combustion across all years and regions was 3.13 ± 1.17 kg C m⁻². Pixel-level uncertainty (Figure S29) tended to follow spatial patterns of mean combustion (Figure 5c) and was relatively consistent across years (Figure S30), with a mean value of 2.86 kg C m⁻². Seasonally, the majority of burned area occurred during June, July and August (Figure 8), although there were substantial regional differences, with some regions recording a large fraction of burned area outside this window (e.g., May fires in Alberta). In general, monthly patterns in emissions (Figure S31) followed patterns in burned

area. Overall, combustion tended to be highest in summer compared to spring and fall fires, although this pattern was most pronounced in the Yukon Territory, Northwest Territories, Saskatchewan and Alaska (Figure S32).




Figure 8. Monthly burned area across states, and Canadian provinces and territories between 2001-2019. January, February, November and December have been omitted due to low fire occurrence (less than 2% of total burned area between 2001-2019).

Estimates of total carbon emissions in ABoVE-FED were similar to AKFED (Figure S22; Table S4), with the notable exception of 2014 in the Northwest Territories: AKFED estimated 164 Tg C and ABoVE-FED estimated 89.7 Tg C. This was primarily a result of differences in mean modeled combustion in the Northwest Territories 2014 fires, with AKFED exhibiting its highest mean combustion in 2014 (Figure S21; 4.81 kg C m⁻² in AKFED compared to 2.89 kg C m⁻² in ABoVE-FED). In general, ABoVE-FED estimated slightly higher mean combustion levels than AKFED in Alaska (3.34 kg C m⁻² in ABoVE-FED and 3.03 kg C m⁻² in AKFED), lower combustion in the Northwest Territories (3.29 kg C m⁻² in ABoVE-FED and 3.44 kg C m⁻² in AKFED), and substantially higher combustion in the Yukon Territory (3.71 kg C m⁻² in ABoVE-FED and 2.26 kg C m⁻² in AKFED) (Figure S22; Table S4). ABoVE-FED carbon emissions were relatively similar to Walker et al. (2018) for the 2014 Northwest Territories fires, and to Dieleman et al. (2020) for the 2015 Saskatchewan fires (Figure 7, Table S4). Total carbon emissions from ABoVE-FED were substantially higher than GFED4s (Figure S33), with the largest differences occurring in Alaska. This was primarily a function of higher mean combustion values in ABoVE-FED compared to GFED4s (Figure S34). Between 2001-2016, ABoVE-FED estimated 80 Tg C y⁻¹ total emissions with a mean combustion value of 3.39 kg C m⁻², and GFED4s estimated 51 Tg C y⁻¹ total emissions with a mean combustion value of 2.30 kg C m⁻² (Table S4). However, more recently a 500 m model by van Wees et al. 2022 was completed and both emissions and combustion match more closely to ABoVE-FED. Between 2002-2019 this 500 m product estimates 73 Tg C y⁻¹ total emissions (Figure S33) with a mean combustion value of 3.38 kg C m⁻² (Figure S34). Meanwhile, between 2002-2019 ABoVE-FED estimates 83 Tg C y⁻¹ total emissions with a mean combustion value of 3.16 kg C m⁻². Compared to GFED4s these larger emission and combustion estimates in the 500 m product are largely due to increased estimates of belowground combustion, as the van Wees et al. 2022 model is informed by the same field measurements used in ABoVE-FED.

Differences in combustion and carbon emissions were very large between ABoVE-FED and fire-vegetation models participating in FireMIP (Figure S34b). ABoVE-FED estimated much higher emissions than FireMIP (70.1 Tg C y⁻¹ for ABoVE-FED during 2001-2012 compared to 4.0 Tg C y⁻¹ for FireMIP). This is likely because models in FireMIP mostly combust aboveground vegetation, whereas combustion from belowground sources (primarily soil organic matter) comprises 90% of total carbon emissions in ABoVE-FED (Figure S35) and 88% in the field plots from Walker et al. (2020a). ABoVE-FED mean aboveground combustion (7.84 Tg C y⁻¹ during 2001-2012) was much more similar to FireMIP's 4.0 Tg C y⁻¹.

We found multiple lines of evidence that belowground fire severity (burn depth and belowground combustion) is positively related to annual burned area and seasonal day of burn (Table S5; S6). In general, mean annual burned area had a stronger relationship with fire severity than did burn day of year using multiple linear regression. However, within quantiles of annual burned area (i.e., low vs. high fire years), day of year was strongly related to fire severity (particularly belowground combustion), and the slope of this relationship was generally larger in small fire years (Table S6). When assessed using domain-wide mean severity from mapped ABoVE-FED pixels, we found no significant relationship of burned depth with burned area or combustion (Figure S36).

There were also no significant (p-value ≤ 0.10) trends in burned area, combustion, or emissions across the 2001 - 2019 time series (Figure 2(a,b), Figure S37).

4 Discussion

4.1 Burned Area

Our approach to mapping burned area across boreal North America has several advantages compared to past approaches. Although our burned area product is at 500 m resolution, the majority of pixels (81%) were mapped using 30 m Landsat imagery. Using finer-scale 30 m imagery allowed us to directly calibrate dNBR thresholds to site-level information and account for unburnable fractions of 500 m pixels. We also

calibrated these dNBR thresholds for both 30 m Landsat and 500 m MODIS imagery to most effectively
875 balance omissions and commissions. This allowed us to provide an unbiased estimate of burned area, which
is a critical variable for understanding the impacts of fire on arctic-boreal ecosystems and climate.

In theory, ABoVE-FED burned area would be expected to be higher than other available products because
of its increased sensitivity to fire-induced spectral changes (compared to, for example, global MODIS
880 burned area products, via our focus on splitting omissions and commissions) and our accounting for active
fire acquisitions outside mapped fire polygons by the Alaskan and Canadian government agencies.
Alternatively, ABoVE-FED accounts for sub-pixel heterogeneity of burnable land surfaces, which would
otherwise result in lower burned area estimates compared to existing products. The net result is that
ABoVE-FED burned area tends to be higher than other products, but not exclusively.

885 We suggest future research efforts focused on burned area mapping in arctic-boreal environments could be
conducted at resolutions finer than 500 m. Doing so will allow for improved understanding of fire spread
and behavior patterns, and interactions between fire behavior and vegetation / land cover type. Finer-scale
mapping should also allow for more accurate assessments of burned area by accounting for the presence of
890 unburned patches of vegetation and water bodies, thereby facilitating increased understanding of the drivers
of fire spread and effects on ecosystem processes (Hall et al., 2020). Fires have typically been mapped at
landscape scales using 500 m MODIS imagery because of the frequent revisit times (multiple acquisitions
per day). With a resolution of 30 m, Landsat imagery has been less commonly used for mapping burned
area at landscape scales because the revisit time (16 days) is much longer, and because data coverage can
895 be highly variable regionally and spatially depending on available downlink stations and cloud cover
(Hilker et al., 2009; Ju and Masek, 2016; Figure S2), but this revisit frequency is improving with two
Landsat satellites (Landsat 8 and 9) and two Sentinel satellites (2a and 2b) in orbit, which provide much
more frequent overpasses (2-3 days when combined).

900 Similar to ABoVE-FED, approaches for mapping burned area using satellite imagery have typically relied
on image differencing of vegetation indices, particularly dNBR (French et al., 2015). This requires pre- and
post-fire image pairs, and thus compounds issues related to image availability at fine scales (30 m; (Chen
et al., 2021). Future burned area mapping at landscape scales could potentially be improved by using
machine learning. More specifically, deep learning approaches have been shown to be highly effective at
905 mapping wildfires across different landscapes and vegetation types (Jain et al., 2020, Knopp et al., 2020).
Convolutional Neural Networks, which use a spatial moving-window and therefore account for the spatial
characteristics of fire scars (Jain et al., 2020) are particularly promising. Finally, developing burned area
products in near real-time, as opposed to active fire-based assessments of hot pixel counts, would help
scientists, fire managers, and society contextualize and potentially mitigate rapidly progressing fire seasons
910 as they evolve.

4.2 Combustion and Burn Depth Models

Similar to previous studies (e.g., Veraverbeke et al., 2015), our aboveground combustion model performed
substantially better than our belowground model. This is due primarily to the challenge of estimating
915 belowground carbon consumption using remote sensing-based observations, which are more sensitive to
aboveground properties. For example, the ABoVE-FED aboveground combustion model was heavily
influenced by remotely-sensed properties such as pre-fire tree cover, fire severity (represented by dNBR),
and vegetation wetness (represented by NDII), whereas the belowground model was strongly influenced
by soil metrics, topography and solar radiation (Figure S16). This occurred despite our model utilizing
920 considerably more field observations ($n = 515$ for aboveground combustion and 769 for belowground) than
past efforts in boreal North America (e.g., Dieleman et al., 2020 ($n = 47$); Veraverbeke et al., 2015 ($n =$
 126); Walker et al., 2018 ($n = 211$)), suggesting an inherently limited capacity to model belowground
combustion using these techniques. Previous analysis of the field observations we used showed site-level
drainage is the dominant driver of combustion in the ABoVE domain, due in part to the large contribution
925 towards total combustion from belowground carbon stocks (Walker et al., 2018; 2020b). We therefore

suggest prioritizing the use of geospatial products that adequately capture drainage, and thereby its impact on belowground carbon stocks and vulnerability to combustion, for improving future estimates of carbon emissions from fire disturbance across boreal North America.

930 Despite these limitations, our model performance is similar to past efforts. For example, Veraverbeke et al., 2015 reported an aboveground combustion model fit of $R^2 = 0.53$ and a belowground fit of $R^2 = 0.29$ for Alaska. Walker et al. (2018) implemented a 10-fold cross validation approach and reported a model fit of $R^2 = 0.26$ for total (above and belowground) combustion in the Northwest Territories, Canada. Comparatively, we report a median R^2 of 0.46 and 0.25 for ABoVE-FED aboveground and belowground
935 combustion models, respectively. However, model performance was substantially higher in Dieleman et al. (2020), who reported a cross-validated R^2 of 0.73 for total combustion in Saskatchewan. This is likely due to the higher relative contribution from aboveground combustion in the younger and more productive boreal forests of southern Canada, combined with high-quality provincial spatial data sets such as logging history (Dieleman et al., 2020). In all these cases, spatial patterns from ABoVE-FED are generally consistent with
940 previous efforts (Figure 7), lending confidence to assessments of drivers and spatio-temporal patterns of combustion.

Somewhat surprisingly, our models of burn depth performed better than both aboveground and belowground combustion models (cross-validated $R^2 = 0.53$), which is considerably better than the R^2
945 model fit of 0.40 reported for the burn depth model in Veraverbeke (2015). This suggests substantial uncertainty in translating burn depth to carbon emissions in these boreal forests, which underscores the need for improved spatial layers of soil properties such as bulk density (Houle et al., 2017) and carbon fraction. The field and laboratory techniques used to calculate carbon emissions from burn depth also contain uncertainty, which is not always quantified. These errors are likely compounded when aggregating
950 data across field campaigns, ecozones, and research groups, such as we did here. Nevertheless, burn depth is a critical fire severity property in its own right, with applications ranging from understanding the

changing boreal carbon cycle (Walker et al., 2019) to post-fire succession and vegetation patterns (Baltzer et al., 2021; Johnstone et al., 2010). Our results suggest geospatial statistical modeling is well-suited for capturing and extrapolating depth of burn in organic soils, at least within the ABoVE domain.

955

Finally, we assessed the influence of spatial and sensor differences when building the combustion and burn depth models at 30 m but predicting them at 500 m. Overall, biases introduced by model nonlinearities, sub-grid heterogeneity, and vegetation fractions were found to be negligible (slope = 0.98 for aboveground and 0.97 for belowground combustion when regressing 500 m against aggregated 30 m predictions). This suggests that approaches to map fire carbon emissions at large scales using 500 m MODIS imagery are not fundamentally biased because of spatial scale.

960

The machine learning models we employed allow insights into the drivers of both aboveground and belowground combustion. Partial dependence plots indicated that aboveground combustion tended to increase when tree cover and dNBR increased, and when relative humidity and vegetation water content (NDII) decreased (Figure S38). These patterns are consistent with understanding of fire behavior and aboveground consumption dynamics, which are generally driven by aboveground fuels and climate conditions that facilitate fuel drying and fire spread (Beck et al., 2011; Rogers et al., 2014; Walker, 2020b). Alternatively, belowground combustion increased with higher silt (and lower sand) content, higher tree cover, and lower relative humidity (Figure S39). At moderate slopes (less than 20%), in which the majority of field observations were located, belowground combustion was higher in flatter landscapes. These relationships are consistent with current understanding about the drivers of soil organic matter accumulation and vulnerability to combustion (Walker et al., 2018, 2020b; Scholten et al., 2021). Drivers of burn depth were similar to those for belowground combustion, with the exception of higher burn depth occurring in areas with lower extreme maximum temperatures and Tasseled Cap Greenness (Figure S40). The former is likely related to deeper burn depths occurring in the northern portions of the ABoVE domain (Figure 5d),

965

970

975

where long-term maximum temperatures are generally lower. Tasseled Cap Greenness was assessed after a given fire, and can therefore be considered a metric of fire severity (low greenness = high severity).

980 Total emissions from ABoVE-FED are relatively consistent with past efforts, including AKFED and
GFED4s, but with some important differences. Total emissions and mean combustion (Figure S12) in
Alaska were similar between ABoVE-FED and AKFED, which is expected given the similar field
observations from Alaska used to develop these models. However, although AKFED was extended to the
Yukon and Northwest Territories (Veraverbeke et al., 2017), it did not incorporate field observations from
985 these regions. By utilizing 797 field plots across these provinces (albeit heavily dominated by the Northwest
Territories), our results suggest AKFED tended to underestimate combustion in the Yukon and overestimate
combustion in the Northwest Territories, especially during the large fire year of 2014. ABoVE-FED also
includes many more predictor variables than AKFED, and is based on a different statistical model. We did
not find large variations in mean combustion from year to year (Figure 2), which is likely related to both
990 the tendency of the random forest models to regress to the mean (Figure S15) as well as relatively consistent
observed mean combustion across large regions of the ABoVE domain (Walker et al., 2020a, c).

GFED4s is a widely-used data source for global and regional burned area and fire emissions. Our results
suggest GFED4s underestimates combustion across the ABoVE domain by roughly 1/3rd (32%; Figure
995 S24; mean of 3.39 kg C m⁻² in ABoVE-FED compared to 2.30 kg C m⁻² in GFED4s), leading to 36% lower
total emissions compared to ABoVE-FED (Figure S23). This is consistent with previous regional studies
noting a consistent underestimation for GFED4s emissions in Alaska (Veraverbeke et al., 2015) and the
Northwest Territories (Walker et al., 2018). This result has important implications for quantifying and
understanding the role of arctic-boreal fires on the global carbon cycle and climate. Regional- to
1000 continental-scale upscaling efforts such as ABoVE-FED, including the underlying field observation

database (Walker et al., 2020a), can help inform further versions of global fire models and thereby improve our quantification and understanding of the role of wildfire on the global carbon cycle.

1005 In contrast to AKFED and GFED4s, fire carbon emissions in FireMIP were an order of magnitude lower (94%) than ABoVE-FED (Figure S13b). This is likely due to the fact that most models in FireMIP only combust aboveground vegetation, whereas combustion of belowground soil organic matter constitutes the majority of emissions in boreal Alaska and Canada. This underscores the importance of developing algorithms that accumulate and burn soil organic matter within global fire models, which is important for both direct fire emissions as well as post-fire permafrost thaw and degradation (Genet et al., 2013; Jafarov et al., 2013; Natali et al., 2021; Treharne et al., 2022).

1015 ABoVE-FED confirms the high interannual variability of fire carbon emissions in the ABoVE domain, including the large fire years of 2004 in Alaska and the Yukon Territory, 2005 in Alaska, 2010 in Saskatchewan, 2014 in the Northwest Territories, and 2015 in Alaska and Saskatchewan. We also found general agreement with previous work (Turetsky et al., 2011) that large fire years and later seasonal fires facilitate deeper burning and higher belowground carbon emissions, including the phenomenon that burn timing has a stronger influence on severity in low fire years (i.e., extreme fire years result in high severity regardless of timing). However, these relationships varied depending on region and analysis technique, and were often confounded by site-level factors and fire weather at the time of burn. Overall, however, this 1020 underscores the influence that climate change (warming, drying, and longer fire seasons) has on boreal fire severity.

Consistent with previous studies (Rogers et al., 2014; Veraverbeke et al., 2015; Walker et al., 2018; Dieleman et al., 2020), ABoVE-FED includes high uncertainty in combustion at the pixel-level (2.86 kg C m⁻²). Much of this uncertainty likely arises from difficulty in predicting large combustion values, particularly from belowground sources (Figure S15b). This suggests ABoVE-FED is underpredicting 1025

emissions coming from the most severe fire events between 2001-2019. We attempted to correct for this bias in a number of ways, including testing a variety of models (Figure 6), tuning model parameters, assigning higher weights to the highest combustion values, and applying the Synthetic Minority
1030 Oversampling Technique (SMOTE; Chawla et al., 2002) to synthetically create more samples with higher combustion values. Ultimately, none of these approaches were able to correct for the low bias at high combustion levels without sacrificing performance for low combustion values. More field observations of high combustion combined with improved predictor variables (particularly drainage) may improve future model performance. Also consistent with previous studies, these pixel-level uncertainties were dampened
1035 through spatial averaging, such that domain-wide mean combustion had comparatively lower uncertainty ($3.13 \pm 1.17 \text{ kg C m}^{-2}$).

Conclusions

Here we used 30 m Landsat and 500 m MODIS imagery to map burned area across Alaska and Canada,
1040 and map fire carbon emissions across the ABoVE domain over a 19-year period between 2001-2019. We utilized a recent field database of combustion observations across the ABoVE domain (Walker et al., 2020a), which represents the largest of its kind for any biome on Earth. We found burned area and total emissions are highly variable by year, averaging 2.37 Mha of burned area and $79.26 \pm 28.65 \text{ Tg C}$ emitted per year across the ABoVE domain (2.87 Mha of burned area across all of Alaska and Canada), with a
1045 mean combustion level of $3.13 \pm 1.20 \text{ kg C m}^{-2}$. When compared to previous products we report more burned area than GFED4s and the MODIS MC64A1 Collection 5 and 6 products. We report similar carbon emissions to AKFED, but more emissions than both GFED4s and FireMIP. ABoVE-FED can be used to understand patterns of fire behavior and effects across central and western boreal North America, and to continue monitoring intensifying fire regimes in boreal forests.

1050

Code and Data Availability

1055 The burned area, combustion and burned depth databases associated with this publication can be
found at Potter et al. 2021 (<https://doi.org/10.3334/ORNLDAAAC/2063>
) . Code is available upon request from the corresponding author.

Competing Interests

All authors declare no competing interests.

1060

Author Contribution

S.P. and B.M.R. contributed to original draft writing. S.V., X.W., M.C.M., S.J.G., J.B., L.B.C., N.F.,
E.E.H., L.J., J.F.J., E.S.K., S.M.N., J.T.R., M.R.T., B.M.R. contributed to conceptualization. S.P., S.C.,
S.H., and B.M.R. contributed to formal analysis. S.V., X.W., M.C.M., S.J.G., L.B.G., N.F., S.M.N., J.T.R.,
1065 M.R.T., and B.M.R. contributed to funding acquisition. S.P., S.C., S.V., X.W., J.T.R. and B.M.R.
contributed to investigation. S.P., S.C., S.V., X.W., M.C.M., S.J.G., J.T.R. and B.M.R. contributed to
methodology. B.M.R. was project administrator. E.E.H. contributed project resources. S.V., X.W., M.C.M.,
S.J.G., S.M.N., J.T.R. and B.M.R. contributed to project supervision. S.P. and S.C. contributed to data
curation, validation and visualization and software development. All authors contributed to the writing-
1070 review process.

Acknowledgements

This work was funded by the National Aeronautics and Space Administration (NASA) Arctic-Boreal
Vulnerability Experiment (ABoVE grants: NNX15AU56A and NX15AT71A, B.M.R. and M.C.M.;
1075 NNX15AT83A and 80NSSC19M0107, L.L.B.C., N.H.F.F., L.K.J), the Gordon and Betty Moore
Foundation (grant #8414), the Woodwell Climate Research Center's Fund for Climate Solutions, and the
Department of Defense (DoD) Strategic Environmental Research and Development Program (SERDP
contract RC18-1183)

Computing resources for this work were provided by the NASA High-End Computing Program through
1080 the NASA Center for Climate Simulation at Goddard Space Flight Center. S.V. acknowledges support from
the Dutch Research Council through Vidi grant 016.Vidi.189.070 and from the European Research Council
under the European Union's Horizon 2020 research and innovation programme (grant agreement No
101000987). In kind support was provided through Bonanza Creek LTER with funding from the National
Science Foundation (DEB-1636476) and the USDA Forest Service, Pacific Northwest Research Station
1085 (RJVA-PNW-01-JV-11261952-231).

1090

1095

1100

1105 References

- Amiro, B. D., Todd, J. B., Wotton, B. M., Logan, K. A., Flannigan, M. D., Stocks, B. J., Mason, J. A., Martell, D. L., and Hirsch, K. G.: Direct carbon emissions from Canadian forest fires, 1959-1999. *Canadian Journal of Forest Research*, 31(3), 512–525. <https://doi.org/10.1139/cjfr-31-3-512>, 2001.
- Balshi, M. S., McGuire, A. D., Duffy, P., Flannigan, M., Kicklighter, D. W., and Melillo, J.: Vulnerability of carbon storage in North American boreal forests to wildfires during the 21st century. *Global Change Biology*, 15(6), 1491–1510. <https://doi.org/10.1111/j.1365-2486.2009.01877.x>, 2009.
- Baltzer, J. L., Day, N. J., Walker, X. J., Greene, D., Mack, M. C., Alexander, H. D., Arseneault, D., Barnes, J., Bergeron, Y., Boucher, Y., Bourgeau-Chavez, L., Brown, C. D., Carrière, S., Howard, B. K., Gauthier, S., Parisien, M.-A., Reid, K. A., Rogers, B. M., Roland, C., ... Johnstone, J. F.: Increasing fire and the decline of fire adapted black spruce in the boreal forest. *Proceedings of the National Academy of Sciences*, 118(45). <https://doi.org/10.1073/pnas.2024872118>, 2021.
- Beaudoin, A., Bernier, P., Guindon, L., Villemaire, P., Guo, X. J., Stinson, G., Magnussen, S., and Hall, R. J.: Mapping attributes of Canada's forests at moderate resolution through kNN and MODIS imagery. 44, 521–532. <https://doi.org/10.1139/cjfr-2013-0401>, 2014.
- Beck, P. S. A., Goetz, S. J., Mack, M. C., Alexander, H. D., Jin, Y., Randerson, J. T., and Loranty, M. M.: The impacts and implications of an intensifying fire regime on Alaskan boreal forest composition and albedo. *Global Change Biology*, 17(9), 2853–2866. <https://doi.org/10.1111/j.1365-2486.2011.02412.x>, 2011.
- Beaudoin, A., P. Y. Bernier, L. Guindon, P. Villemaire, X. J. Guo, G. Stinson, T. Bergeron, S. Magnussen, and R. J. Hall.: Mapping attributes of Canada's forests at moderate resolution through kNN and MODIS imagery, *Can. J. For. Res.*, 44(5), 521–532, doi:10.1139/cjfr-2013-0401, 2010.
- Beven, K. J., and Kirkby, M. J.: A physically based, variable contributing area model of basin hydrology. *Hydrological Sciences Bulletin*, 24(1), 43–69., 2014. <https://doi.org/10.1080/02626667909491834>, 1979.
- Boby, L. A., Schuur, E. A. G., Mack, M. C., Verbyla, D., and Johnstone, J. F.: Quantifying fire severity, carbon, and nitrogen emissions in Alaska's boreal forest. *Ecological Applications*, 20(6), 1633–1647. <https://doi.org/10.1890/08-2295.1>, 2010.
- Bonan, G. B., and Shugart, H. H.: Environmental Factors and Ecological Processes in Boreal Forests. *Annual Review of Ecology and Systematics*, 20, 1–28., 1989.
- Bond-Lamberty, B., Peckham, S. D., Ahl, D. E., and Gower, S. T.: Fire as the dominant driver of central Canadian boreal forest carbon balance. *Nature*, 450(7166), 89. <https://doi.org/10.1038/nature06272>, 2007.
- Boulanger, Y., Parisien, M.-A., and Wang, X.: Model-specification uncertainty in future area burned by wildfires in Canada. *International Journal of Wildland Fire*, 27(3), 164–175. <https://doi.org/10.1071/WF17123>, 2018.
- Breiman, L.: Random Forests. *Machine Learning*, 45(1), 5–32. <https://doi.org/10.1023/A:1010933404324>, 2001.
- Burns, PJ, Massey, RM, Shean, D, Husby, EB, Goetz, SG: A composite 10-m digital elevation model of Alaska and Canada for NASA ABoVE. In preparation for Environmental Research Letters, forthcoming.
- Chawla, N. V., Bowyer, K. W., Hall, L. O., and Kegelmeyer, W. P.: SMOTE: Synthetic Minority Over-sampling Technique. *Journal of Artificial Intelligence Research*, 16, 321–357. <https://doi.org/10.1613/jair.953>, 2002.
- Chen, D., Loboda, T. V., and Hall, J. V.: A systematic evaluation of influence of image selection process on remote sensing-based burn severity indices in North American boreal forest and tundra ecosystems. *ISPRS Journal of Photogrammetry and Remote Sensing*, 159, 63–77. <https://doi.org/10.1016/j.isprsjprs.2019.11.011>, 2020.

- 1155 Chen, D., Fu, C., Hall, J. V., Hoy, E. E., and Loboda, T. V.: Spatio-temporal patterns of optimal Landsat data for burn severity index calculations: Implications for high northern latitudes wildfire research. *Remote Sensing of Environment*, 258, 112393., 2021. <https://doi.org/10.1016/j.rse.2021.112393>
- 1160 Daly, C., Gibson, W., Taylor, G., Johnson, G., and Pasteris, P.: A knowledge-based approach to the statistical mapping of climate. *Climate Research*, 22, 99–113. <https://doi.org/10.3354/cr022099>, 2002.
- Daly, C., Halbleib, M., Smith, J. I., Gibson, W. P., Doggett, M. K., Taylor, G. H., Curtis, J., and Pasteris, P. P.: Physiographically sensitive mapping of climatological temperature and precipitation across the conterminous United States. *International Journal of Climatology*, 28(15), 2031–2064. <https://doi.org/10.1002/joc.1688>, 2008.
- 1165 de Groot, W. J., Pritchard, J. M. P. M., and Lynham, T. J. L. J.: Forest floor fuel consumption and carbon emissions in Canadian boreal forest fires. *Canadian Journal of Forest Research*. <https://doi.org/10.1139/X08-192>, 2009.
- 1170 de Groot, W. J., Landry, R., Kurz, W. A., Anderson, K. R., Englefield, P., Fraser, R. H., Hall, R. J., Banfield, E., Raymond, D. A., Decker, V., Lynham, T. J., Pritchard, J. M., Groot, W. J. de, Landry, R., Kurz, W. A., Anderson, K. R., Englefield, P., Fraser, R. H., Hall, R. J., ... Pritchard, J. M.: Estimating direct carbon emissions from Canadian wildland fires1. *International Journal of Wildland Fire*, 16(5), 593–606. <https://doi.org/10.1071/WF06150>, 2007.
- 1175 Didan, K. MOD13A3 MODIS/Terra vegetation Indices Monthly L3 Global 1km SIN Grid V006. NASA EOSDIS Land Processes DAAC. Accessed 2021-12-09 from <https://doi.org/10.5067/MODIS/MOD13A3.006>, 2015a.
- Didan, K. MOD13A1 MODIS/Terra vegetation Indices Monthly L3 Global 1km SIN Grid V006. NASA EOSDIS Land Processes DAAC. Accessed 2021-12-09 from <https://doi.org/10.5067/MODIS/MOD13A3.006>, 2015b.
- 1180 Di Giuseppe, F., Rémy, S., Pappenberger, F., and Wetterhall, F.: Using the Fire Weather Index (FWI) to improve the estimation of fire emissions from fire radiative power (FRP) observations. *Atmospheric Chemistry and Physics*, 18(8), 5359–5370. <https://doi.org/10.5194/acp-18-5359-2018>, 2018.
- 1185 Dieleman, C. M., Rogers, B. M., Potter, S., Veraverbeke, S., Johnstone, J. F., Laflamme, J., Solvik, K., Walker, X. J., Mack, M. C., and Turetsky, M. R.: Wildfire combustion and carbon stocks in the southern Canadian boreal forest: Implications for a warming world. *Global Change Biology*, 26(11), 6062–6079. <https://doi.org/10.1111/gcb.15158>, 2020.
- Epp, H., and Lanoville, R.: Satellite data and geographic information systems for fire and resource management in the Canadian arctic. *Geocarto International*, 11(2), 97–103. <https://doi.org/10.1080/10106049609354537>, 1996.
- 1190 Fick, S. E., and Hijmans, R. J.: WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, 37(12), 4302–4315. <https://doi.org/10.1002/joc.5086>, 2017.
- 1195 Field, R. D., Spessa, A. C., Aziz, N. A., Camia, A., Cantin, A., Carr, R., de Groot, W. J., Dowdy, A. J., Flannigan, M. D., Manomaiphiboon, K., Pappenberger, F., Tanpipat, V., and Wang, X.: Development of a Global Fire Weather Database. *Natural Hazards and Earth System Sciences*, 15(6), 1407–1423. <https://doi.org/10.5194/nhess-15-1407-2015>, 2015.
- Foga, S., Scaramuzza, P. L., Guo, S., Zhu, Z., Dilley, R. D., Beckmann, T., Schmidt, G. L., Dwyer, J. L., Hughes, M. J., Laue, B.: Cloud detection algorithm comparison and validation for operational Landsat data products. *Remote Sensing of Environment*, 194, 379–390. <http://doi.org/10.1016/j.rse.2017.03.026>, 2017.
- 1200 French, N. H. F., Groot, W. J. de, Jenkins, L. K., Rogers, B. M., Alvarado, E., Amiro, B., Jong, B. de, Goetz, S., Hoy, E., Hyer, E., Keane, R., Law, B. E., McKenzie, D., McNulty, S. G., Ottmar, R., Pérez-Salícru, D. R., Randerson, J., Robertson, K. M., and Turetsky, M.: Model comparisons for

- 1205 estimating carbon emissions from North American wildland fire. *Journal of Geophysical Research: Biogeosciences*, 116(G4). <https://doi.org/10.1029/2010JG001469>, 2011.
- French, N. H. F., Jenkins, L. K., Loboda, T. V., Flannigan, M., Jandt, R., Bourgeau-Chavez, L. L., and Whitley, M.: Fire in arctic tundra of Alaska: Past fire activity, future fire potential, and significance for land management and ecology. *International Journal of Wildland Fire*, 24(8), 1045. <https://doi.org/10.1071/WF14167>, 2015.
- 1210 French, N. H. F., Kasischke, E. S., and Williams, D. G.: Variability in the emission of carbon-based trace gases from wildfire in the Alaskan boreal forest. *Journal of Geophysical Research*, 108(D1), 8151. <https://doi.org/10.1029/2001JD000480>, 2002.
- French, N. H. F., McKenzie, D., Erickson, T., Koziol, B., Billmire, M., Endsley, K. A., Yager Scheinerman, N. K., Jenkins, L., Miller, M. E., Ottmar, R., and Prichard, S.: Modeling Regional-Scale Wildland Fire Emissions with the Wildland Fire Emissions Information System*. *Earth Interactions*, 18(16), 1–26. <https://doi.org/10.1175/EI-D-14-0002.1>, 2014.
- 1215 Friedl, M., Sulla-Menashe, D.: MCD12Q1 MODIS/Terra+Aqua Land Cover Type Yearly L3 Global 500m SIN Grid V006 [Data set]. NASA EOSDIS Land Processes DAAC. Accessed 2021-08-10 from <https://doi.org/10.5067/MODIS/MCD12Q1.006>, 2019.
- 1220 Genet, H., McGuire, A. D., Barrett, K., Breen, A., Euskirchen, E. S., Johnstone, J. F., Kasischke, E. S., Melvin, A. M., Bennett, A., Mack, M. C., Rupp, T. S., Schuur, A. E. G., Turetsky, M. R., and Yuan, F.: Modeling the effects of fire severity and climate warming on active layer thickness and soil carbon storage of black spruce forests across the landscape in interior Alaska. *Environmental Research Letters*, 8(4), 045016. <https://doi.org/10.1088/1748-9326/8/4/045016>, 2013.
- 1225 Giglio, L., Boschetti, L., Roy, D. P., Humber, M. L., and Justice, C. O.: The Collection 6 MODIS burned area mapping algorithm and product. *Remote Sensing of Environment*, 217, 72–85. <https://doi.org/10.1016/j.rse.2018.08.005>, 2018.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*, 202, 18–27. <https://doi.org/10.1016/j.rse.2017.06.031>, 2017.
- 1230 Gruber, S.: Derivation and analysis of a high-resolution estimate of global permafrost zonation. *The Cryosphere*, 6(1), 221–233. <https://doi.org/10.5194/tc-6-221-2012>, 2012.
- Guindon, L., Bernier, P., Gauthier, S., Stinson, G., Villemaire, P., and Beaudoin, A.: Missing forest cover gains in boreal forests explained. *Ecosphere*, 9(1), e02094. <https://doi.org/10.1002/ecs2.2094>, 2018.
- 1235 Guyon I, Weston J, Barnhill S, Vapnik V.: “Gene Selection for Cancer Classification Using Support Vector Machines.” *Machine Learning*, 46(1), 389–422., 2002.
- 1240 Hall, R. J., Skakun, R. S., Metsaranta, J. M., Landry, R., Fraser, R. H., Raymond, D., Gartrell, M., Decker, V., Little, J., Hall, R. J., Skakun, R. S., Metsaranta, J. M., Landry, R., Fraser, R. H., Raymond, D., Gartrell, M., Decker, V., and Little, J.: Generating annual estimates of forest fire disturbance in Canada: The National Burned Area Composite. *International Journal of Wildland Fire*, 29(10), 878–891. <https://doi.org/10.1071/WF19201>, 2020.
- 1245 Hanes, C. C., Wang, X., Jain, P., Parisien, M.-A., Little, J. M., and Flannigan, M. D.: Fire-regime changes in Canada over the last half century. *Canadian Journal of Forest Research*. <https://doi.org/10.1139/cjfr-2018-0293>, 2018.
- Hantson, S., Arneth, A., Harrison, S. P., Kelley, D. I., Prentice, I. C., Rabin, S. S., Archibald, S., Mouillot, F., Arnold, S. R., Artaxo, P., Bachelet, D., Ciais, P., Forrest, M., Friedlingstein, P., Hickler, T., Kaplan, J. O., Kloster, S., Knorr, W., Lasslop, G., ... Yue, C.: The status and challenge of global fire modelling. *Biogeosciences*, 13(11), 3359–3375. <https://doi.org/10.5194/bg-13-3359-2016>, 2016.
- 1250 Hardisky, M. A., Klemas, V., and Smart, R. M.: The influences of soil salinity, growth form, and leaf moisture on the spectral reflectance of *Spartina alterniflora* canopies. *Photogrammetric Engineering and Remote Sensing*, 49, 77–83., 1983.

- 1255 Hengl, T., Jesus, J. M. de, Heuvelink, G. B. M., Gonzalez, M. R., Kilibarda, M., Blagotić, A., Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: Global gridded soil information based on machine learning. *PLOS ONE*, 12(2), e0169748. <https://doi.org/10.1371/journal.pone.0169748>, 2017.
- 1260 Hilker, T., Wulder, M. A., Coops, N. C., Linke, J., McDermid, G., Masek, J. G., Gao, F., and White, J. C.: A new data fusion model for high spatial- and temporal-resolution mapping of forest disturbance based on Landsat and MODIS. *Remote Sensing of Environment*, 113(8), 1613–1627. <https://doi.org/10.1016/j.rse.2009.03.007>, 2009.
- Holden, Z. A., Morgan, P., and Evans, J. S.: A predictive model of burn severity based on 20-year satellite-inferred burn severity data in a large southwestern US wilderness area. *Forest Ecology and Management*, 258(11), 2399–2406. <https://doi.org/10.1016/j.foreco.2009.08.017>, 2009.
- 1265 Hoy, E. E., French, N. H. F., Turetsky, M. R., Trigg, S. N., and Kasischke, E. S.: Evaluating the potential of Landsat TM/ETM+ imagery for assessing fire severity in Alaskan black spruce forests. *International Journal of Wildland Fire*, 17(4), 500–514. <https://doi.org/10.1071/WF08107>, 2008.
- 1270 Hoy, E. E., Turetsky, M. R., and Kasischke, E.: More frequent burning increases vulnerability of Alaskan boreal black spruce forests. *Environmental Research Letters*, 11(9), 095001. <https://doi.org/10.1088/1748-9326/11/9/095001>, 2016.
- Hudak, A. T., Morgan, P., Bobbitt, M. J., Smith, A. M. S., Lewis, S. A., Lentile, L. B., Robichaud, P. R., Clark, J. T., and McKinley, R. A.: The Relationship of Multispectral Satellite Imagery to Immediate Fire Effects. *Fire Ecology*, 3(1), 64–90. <https://doi.org/10.4996/fireecology.0301064>, 2007.
- Houle, G. P., Kane, E. S., Kasischke, E. S., Gibson, C., and Turetsky, M. R.: Recovery of Carbon Pools a Decade Following Wildfire in Black Spruce Forests of Interior Alaska: Effects of Soil Texture and Landscape Position. *Canadian Journal of Forest Research*, 2017.
- 1280 Hutchinson, M. F.: A new objective method for spatial interpolation of meteorological variables from irregular networks applied to the estimation of monthly mean solar radiation, temperature, precipitation and windrun. *CSIRO Division of Water Resources Tech. Memo.*, 89(5), 95–104., 1989
- Ivanova, G. A., Conard, S. G., Kukavskaya, E. A., and McRae, D. J.: Fire impact on carbon storage in light conifer forests of the Lower Angara region, Siberia. *Environmental Research Letters*, 6(4), 045203. <https://doi.org/10.1088/1748-9326/6/4/045203>, 2011.
- 1285 Jafarov, E. E., Romanovsky, V. E., Genet, H., McGuire, A. D., and Marchenko, S. S.: The effects of fire on the thermal stability of permafrost in lowland and upland black spruce forests of interior Alaska in a changing climate. *Environmental Research Letters*, 8(3), 035030. <https://doi.org/10.1088/1748-9326/8/3/035030>, 2013.
- 1290 Jain, P., Coogan, S. C. P., Subramanian, S. G., Crowley, M., Taylor, S., and Flannigan, M. D.: A review of machine learning applications in wildfire science and management. *Environmental Reviews*, 28(4), 478–505. <https://doi.org/10.1139/er-2020-0019>, 2020.
- Johnstone, J. F., Chapin, F. S., Hollingsworth, T. N., Mack, M. C., Romanovsky, V., and Turetsky, M.: Fire, climate change, and forest resilience in interior Alaska This article is one of a selection of papers from The Dynamics of Change in Alaska's Boreal Forests: Resilience and Vulnerability in Response to Climate Warming. *Canadian Journal of Forest Research*, 40(7), 1302–1312. <https://doi.org/10.1139/X10-061>, 2010.
- 1295 Ju, J., and Masek, J. G.: The vegetation greenness trend in Canada and US Alaska from 1984–2012 Landsat data. *Remote Sensing of Environment*, 176, 1–16. <https://doi.org/10.1016/j.rse.2016.01.001>, 2016.
- 1300 Kantzas, E., Lomas, M., and Quegan, S.: Fire at high latitudes: Data-model comparisons and their consequences: BOREAL FIRES IN LAND MODELS. *Global Biogeochemical Cycles*, 27(3), 677–691. <https://doi.org/10.1002/gbc.20059>, 2013.
- Kasischke, E. S., and Hoy, E. E.: Controls on carbon consumption during Alaskan wildland fires. *Global Change Biology*, 18(2), 685–699. <https://doi.org/10.1111/j.1365-2486.2011.02573.x>, 2012.
- 1305

- Kasischke, E. S., Turetsky, M. R., Ottmar, R. D., French, N. H. F., Hoy, E. E., Kane, E. S., Kasischke, E. S., Turetsky, M. R., Ottmar, R. D., French, N. H. F., Hoy, E. E., and Kane, E. S.: Evaluation of the composite burn index for assessing fire severity in Alaskan black spruce forests. *International Journal of Wildland Fire*, 17(4), 515–526. <https://doi.org/10.1071/WF08002>, 2008.
- 1310 Kasischke, E. S., Verbyla, D. L., Rupp, T. S., McGuire, A. D., Murphy, K. A., Jandt, R., Barnes, J. L., Hoy, E. E., Duffy, P. A., Calef, M., and Turetsky, M. R.: Alaska's changing fire regime — implications for the vulnerability of its boreal forests This article is one of a selection of papers from *The Dynamics of Change in Alaska's Boreal Forests: Resilience and Vulnerability in Response to Climate Warming*. *Canadian Journal of Forest Research*, 40(7), 1313–1324. <https://doi.org/10.1139/X10-098>, 2010.
- 1315 Kasischke, E. S., Williams, D., and Barry, D.: Analysis of the patterns of large fires in the boreal forest region of Alaska. *International Journal of Wildland Fire*, 11(2), 131–144. <https://doi.org/10.1071/wf02023>, 2002.
- Kauth, R.J. and Thomas, G.S. 1976, The Tasseled Cap—A Graphic Description of the Spectral-Temporal Development of Agricultural Crops as Seen by LANDSAT. LARS Symposia, paper 159.
- 1320 Key, C. H., and Benson, N. C.: Landscape Assessment: Ground Measure of Severity, the Composite Burn Index; and Remote Sensing of Severity, the Normalized Burn Ratio. In D. C. Lutes, R. E. Keane, J. F. Caratti, C. H. Key, N. C. Benson, S. Sutherland, and L. J. Gangi (Eds.), *FIREMON: Fire Effects Monitoring and Inventory System*. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station, Gen. Tech. Rep., 2006.
- 1325 Knopp, L., Wieland, M., Rättich, M., and Martinis, S.: A Deep Learning Approach for Burned Area Segmentation with Sentinel-2 Data. *Remote Sensing*, 12(15), 2422. <https://doi.org/10.3390/rs12152422>, 2020.
- Latifovic, Rasim and Homer, Collin and Ressler, Rainer and Pouliot, D.A. and Hossian, S. and Colditz, Rene and Olthof, Ian and Chandra, Giri and Victoria, Arturo.: North American Land Change Monitoring System. *Remote Sensing of Land Use and Land Cover: Principles and Applications*. 303-324. 10.1201/b11964-24, 2012.
- 1330 Li, F., Lawrence, D. M., and Bond-Lamberty, B.: Impact of fire on global land surface air temperature and energy budget for the 20th century due to changes within ecosystems. *Environmental Research Letters*, 12(4), 044014. <https://doi.org/10.1088/1748-9326/aa6685>, 2017.
- 1335 Loboda, T.V., D. Chen, J.V. Hall, and J. He.: ABoVE: Landsat-derived Burn Scar dNBR across Alaska and Canada, 1985-2015. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAAC/1564>, 2018.
- Loboda, T.V., E.E. Hoy, and M.L. Carroll.: ABoVE: Study Domain and Standard Reference Grids, Version 2. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAAC/1527>, 2019.
- 1340 McGuire, A. D., Anderson, L. G., Christensen, T. R., Dallimore, S., Guo, L., Hayes, D. J., Heimann, M., Lorenson, T. D., Macdonald, R. W., and Roulet, N.: Sensitivity of the Carbon Cycle in the Arctic to Climate Change. *Ecological Monographs*, 79(4), 523–555, 2009.
- 1345 Miller, J. D., and Thode, A. E.: Quantifying burn severity in a heterogeneous landscape with a relative version of the delta Normalized Burn Ratio (dNBR). *Remote Sensing of Environment*, 109(1), 66–80. <https://doi.org/10.1016/j.rse.2006.12.006>, 2007.
- Mitchell, T. D., and Jones, P. D.: An improved method of constructing a database of monthly climate observations and associated high-resolution grids. *International Journal of Climatology*, 25(6), 693–712. <https://doi.org/10.1002/joc.1181>, 2005.
- 1350 Natali, S. M., Holdren, J. P., Rogers, B. M., Treharne, R., Duffy, P. B., Pomerance, R., and MacDonald, E.: Permafrost carbon feedbacks threaten global climate goals. *Proceedings of the National Academy of Sciences*, 118(21), e2100163118. <https://doi.org/10.1073/pnas.2100163118>
- 1355 Neigh, C. S. R., Masek, J. G., and Nickeson, J. E. (2013). High-Resolution Satellite Data Open for Government Research. *Eos, Transactions American Geophysical Union*, 94(13), 121–123. <https://doi.org/10.1002/2013EO130002>, 2021.

- Ottmar, R. D., Sandberg, D.V., Riccardi, C.L. and Prichard, S.J.: An overview of the Fuel Characteristic Classification System - Quantifying, classifying, and creating fuelbeds for resource planning. *Can. J. For. Res.*, 37(12), 2383–2393, doi:10.1139/X07-077, 2007.
- 1360 Parks, S., Dillon, G., and Miller, C.: A New Metric for Quantifying Burn Severity: The Relativized Burn Ratio. *Remote Sensing*, 6(3), 1827–1844. <https://doi.org/10.3390/rs6031827>, 2014.
- Pekel, J.-F., Cottam, A., Gorelick, N., and Belward, A. S.: High-resolution mapping of global surface water and its long-term changes. *Nature*, 540(7633), 418–422. <https://doi.org/10.1038/nature20584>, 2016.
- 1365 Porter, C., Morin, P., Howat, I., Noh, M., Bates, B., Peterman, K., Keesey, S., Schlenk, M., Gardiner, J., Tomko, K., Willis, M., Kelleher, C., Cloutier, M., Husby, E., Foga, S., Nakamura, H., Platson, M., Wethington, M., Williamson, C., Bauer, G., Enos, J., Arnold, G., Kramer, W., Becker, P., Doshi, Abhijit, D'Souza, C., Cummins, P., Laurier, F., Bojesen, M.: "ArcticDEM", <https://doi.org/10.7910/DVN/OHHUKH>, Harvard Dataverse, V1, 2018.
- 1370 Potter, S., S. Veraverbeke, X.J. Walker, M.C. Mack, S.J. Goetz, J.L. Baltzer, C. Dieleman, N.H.F. French, E.S. Kane, M.R. Turetsky, E.B. Wiggins, and B.M. Rogers.: ABoVE: Burned Area, Depth, and Combustion for Alaska and Canada, 2001–2019. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAAC/2063>, 2022
- 1375 Pouliot, D., and Latifovic, R.: Accuracy assessment of annual land cover time series derived from change-based updating. Banff, Canada: Institute of Electrical and Electronics Engineers. <https://doi.org/10.1109/Multi-Temp.2013.6866005>, 2013.
- Pouliot, D., Latifovic, R., Zabcic, N., Guindon, L., and Olthof, I.: Development and assessment of a 250m spatial resolution MODIS annual land cover time series (2000–2011) for the forest region of Canada derived from change-based updating. *Remote Sensing of Environment*, 140, 731–743. <https://doi.org/10.1016/j.rse.2013.10.004>, 2014.
- 1380 R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>, 2021.
- 1385 Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Kaplan, J. O., Li, F., Mangeon, S., Ward, D. S., Yue, C., Arora, V. K., Hickler, T., Kloster, S., Knorr, W., Nieradzick, L., Spessa, A., Folberth, G. A., Sheehan, T., ... Arneeth, A.: The Fire Modeling Intercomparison Project (FireMIP), phase 1: Experimental and analytical protocols with detailed model descriptions. *Geoscientific Model Development*, 10(3), 1175–1197. <https://doi.org/10.5194/gmd-10-1175-2017>, 2017.
- 1390 Randerson, J. T., Chen, Y., Werf, G. R. van der, Rogers, B. M., and Morton, D. C.: Global burned area and biomass burning emissions from small fires. *Journal of Geophysical Research: Biogeosciences*, 117(G4). <https://doi.org/10.1029/2012JG002128>, 2012.
- Raynolds, M. K., Walker, D. A., Balser, A., Bay, C., Campbell, M., Cherosov, M. M., Daniëls, F. J. A., Eidesen, P. B., Ermokhina, K. A., Frost, G. V., Jedrzejek, B., Jorgenson, M. T., Kennedy, B. E., Kholod, S. S., Lavrinenko, I. A., Lavrinenko, O. V., Magnússon, B., Matveyeva, N. V., Metúsalemsson, S., ... Troeva, E.: A raster version of the Circumpolar Arctic Vegetation Map (CAVM). *Remote Sensing of Environment*, 232, 111297. <https://doi.org/10.1016/j.rse.2019.111297>, 2019.
- 1395 Rogers, B. M., Soja, A. J., Goulden, M. L., and Randerson, J. T.: Influence of tree species on continental differences in boreal fires and climate feedbacks. *Nature Geoscience*, 8(3), 228–234. <https://doi.org/10.1038/ngeo2352>, 2015.
- 1400 Rogers, B. M., Veraverbeke, S., Azzari, G., Czimczik, C. I., Holden, S. R., Mouteva, G. O., Sedano, F., Treseder, K. K., and Randerson, J. T.: Quantifying fire-wide carbon emissions in interior Alaska using field measurements and Landsat imagery. *Journal of Geophysical Research: Biogeosciences*, 119(8), 1608–1629. <https://doi.org/10.1002/2014JG002657>, 2014.
- 1405 Roy, D. P., Kovalsky, V., Zhang, H. K., Vermote, E. F., Yan, L., Kumar, S. S., and Egorov, A.: Characterization of Landsat-7 to Landsat-8 reflective wavelength and normalized difference

- vegetation index continuity. *Remote Sensing of Environment*, 185, 57–70.
<https://doi.org/10.1016/j.rse.2015.12.024>, 2016.
- 1410 Scholten, R. C., Jandt, R., Miller, E. A., Rogers, B. M., & Veraverbeke, S.: Overwintering fires
 in boreal forests. *Nature*, 593(7859), 399–404. <https://doi.org/10.1038/s41586-021-03437-y>, 2021.
- Schmidt, G., Jenkerson, C. B., Masek, J., Vermote, E., and Gao, F.: Landsat ecosystem disturbance
 adaptive processing system (LEDAPS) algorithm description (Report No. 2013–1057; Open-File
 Report, p. 27). USGS Publications Warehouse. <https://doi.org/10.3133/ofr20131057>, 2013.
- 1415 Seiler, W., and Crutzen, P. J.: Estimates of gross and net fluxes of carbon between the biosphere and the
 atmosphere from biomass burning. *Climatic Change*, 2(3), 207–247.
<https://doi.org/10.1007/BF00137988>, 1980.
- Sexton, J. O., Song, X.-P., Feng, M., Noojipady, P., Anand, A., Huang, C., Kim, D.-H., Collins, K. M.,
 Channan, S., DiMiceli, C., and Townshend, J. R.: Global, 30-m resolution continuous fields of
 tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based
 estimates of error. *International Journal of Digital Earth*, 6(5), 427–448.
<https://doi.org/10.1080/17538947.2013.786146>, 2013.
- 1420 Skakun, R., Whitman, E., Little, J. M., and Parisien, M.-A.: Area burned adjustments to historical
 wildland fires in Canada. *Environmental Research Letters*, 16(6), 064014.
<https://doi.org/10.1088/1748-9326/abfb2c>, 2021.
- 1425 Stocks, B. J., Mason, J. A., Todd, J. B., Bosch, E. M., Wotton, B. M., Amiro, B. D., Flannigan, M. D.,
 Hirsch, K. G., Logan, K. A., Martell, D. L., and Skinner, W. R.: Large forest fires in Canada,
 1959–1997. *Journal of Geophysical Research: Atmospheres*, 107(D1), FFR 5-1-FFR 5-12.
<https://doi.org/10.1029/2001JD000484>, 2003.
- 1430 Tan, Z., Tieszen, L. L., Zhu, Z., Liu, S., and Howard, S. M.: An estimate of carbon emissions from 2004
 wildfires across Alaskan Yukon River Basin. *Carbon Balance and Management*, 2(1), 12.
<https://doi.org/10.1186/1750-0680-2-12>, 2007.
- Tucker, C. J.: Red and photographic infrared linear combinations for monitoring vegetation. *Remote
 Sensing of Environment*, 8(2), 127–150. [https://doi.org/10.1016/0034-4257\(79\)90013-0](https://doi.org/10.1016/0034-4257(79)90013-0), 1979.
- 1435 Turetsky, M. R., Kane, E. S., Harden, J. W., Ottmar, R. D., Manies, K. L., Hoy, E., and Kasischke, E. S.:
 Recent acceleration of biomass burning and carbon losses in Alaskan forests and peatlands.
Nature Geoscience, 4(1), 27–31. <https://doi.org/10.1038/ngeo1027>, 2011.
- Treharne, R., Rogers, B. M., Gasser, T., MacDonald, E., and Natali, S.: Identifying Barriers to Estimating
 Carbon Release From Interacting Feedbacks in a Warming Arctic. *Frontiers in Climate*, 3,
 716464. <https://doi.org/10.3389/fclim.2021.716464>, 2022.
- 1440 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C.,
 DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of
 deforestation, savanna, forest, agricultural, and peat fires (1997–2009). *Atmospheric Chemistry
 and Physics*, 10(23), 11707–11735. <https://doi.org/10.5194/acp-10-11707-2010>, 2010.
- 1445 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M.,
 van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global
 fire emissions estimates during 1997–2016. *Earth System Science Data*, 9(2), 697–720.
<https://doi.org/10.5194/essd-9-697-2017>, 2017.
- 1450 van Leeuwen, T. T., van der Werf, G. R., Hoffmann, A. A., Detmers, R. G., Rücker, G., French, N. H. F.,
 Archibald, S., Carvalho Jr., J. A., Cook, G. D., de Groot, W. J., Hély, C., Kasischke, E. S.,
 Kloster, S., McCarty, J. L., Pettinari, M. L., Savadogo, P., Alvarado, E. C., Boschetti, L., Manuri,
 S., ... Trollope, W. S. W.: Biomass burning fuel consumption rates: A field measurement
 database. *Biogeosciences*, 11(24), 7305–7329. <https://doi.org/10.5194/bg-11-7305-2014>, 2014.
- 1455 Veraverbeke, S., Rogers, B. M., Goulden, M. L., Jandt, R. R., Miller, C. E., Wiggins, E. B., and
 Randerson, J. T.: Lightning as a major driver of recent large fire years in North American boreal
 forests. *Nature Climate Change*, 7(7), 529–534. <https://doi.org/10.1038/nclimate3329>, 2017.

- Veraverbeke, S., Rogers, B. M., and Randerson, J. T.: Daily burned area and carbon emissions from boreal fires in Alaska. *Biogeosciences*, 12(11), 3579–3601. <https://doi.org/10.5194/bg-12-3579-2015>, 2015.
- 1460 Veraverbeke, S., Sedano, F., Hook, S. J., Randerson, J. T., Jin, Y., and Rogers, B. M.: Mapping the daily progression of large wildland fires using MODIS active fire data. *International Journal of Wildland Fire*, 23(5), 655–667. <https://doi.org/10.1071/WF13015>, 2014.
- Verbyla, D., and Lord, R.: Estimating post-fire organic soil depth in the Alaskan boreal forest using the Normalized Burn Ratio. *International Journal of Remote Sensing*. <https://doi.org/10.1080/01431160701802497>, 2008.
- 1465 Vermote, E., Wolfe, R.: MOD09GA MODIS/Terra Surface Reflectance Daily L2G Global 1km and 500m SIN Grid V006 [Data set]. NASA EOSDIS Land Processes DAAC. Accessed 2021-12-09 from <https://doi.org/10.5067/MODIS/MYD09GA.006>, 2015a.
- Vermote, E., Wolfe, R.: MOD09GA MODIS/Terra Surface Reflectance Daily L2G Global 1km and 500m SIN Grid V006 [Data set]. NASA EOSDIS Land Processes DAAC. Accessed 2021-12-09 from <https://doi.org/10.5067/MODIS/MOD09GA.006>, 2015b.
- 1470 Vermote, E., Justice, C., Claverie, M., and Franch, B.: Preliminary analysis of the performance of the Landsat 8/OLI land surface reflectance product. *Remote Sensing of Environment*. <http://dx.doi.org/10.1016/j.rse.2016.04.008>, 2016.
- 1475 Walker, X. J., Baltzer, J. L., Bourgeau-Chavez, L. L., Day, N. J., de Groot, W. J., Dieleman, C., Hoy, E. E., Johnstone, J. F., Kane, E. S., Parisien, M. A., Potter, S., Rogers, B. M., Turetsky, M. R., Veraverbeke, S. R., Whitman, E. and Mack, M. C.: ABoVE: Synthesis of Burned and Unburned Forest Site Data, AK and Canada, 1983–2016. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1744>, 2020a.
- 1480 Walker, X. J., Rogers, B. M., Veraverbeke, S., Johnstone, J. F., Baltzer, J. L., Barrett, K., Bourgeau-Chavez, L., Day, N. J., de Groot, W. J., Dieleman, C. M., Goetz, S., Hoy, E., Jenkins, L. K., Kane, E. S., Parisien, M. A., Potter, S., Schuur, E. A. G., Turetsky, M., Whitman, E., and Mack, M. C.: Fuel availability not fire weather controls boreal wildfire severity and carbon emissions. *Nature Climate Change*, 10(12), 1130–1136. <https://doi.org/10.1038/s41558-020-00920-8>, 2020b.
- 1485 Walker, X. J., Baltzer, J. L., Bourgeau-Chavez, L., Day, N. J., Dieleman, C. M., Johnstone, J. F., Kane, E. S., Rogers, B. M., Turetsky, M. R., Veraverbeke, S., and Mack, M. C.: Patterns of Ecosystem Structure and Wildfire Carbon Combustion Across Six Ecoregions of the North American Boreal Forest. *Frontiers in Forests and Global Change*, 3, 87. <https://doi.org/10.3389/ffgc.2020.00087>, 2020c.
- 1490 Walker, X. J., Baltzer, J. L., Cumming, S. G., Day, N. J., Ebert, C., Goetz, S., Johnstone, J. F., Potter, S., Rogers, B. M., Schuur, E. A. G., Turetsky, M. R., and Mack, M. C.: Increasing wildfires threaten historic carbon sink of boreal forest soils. *Nature*, 572(7770), 520–523. <https://doi.org/10.1038/s41586-019-1474-y>, 2019.
- 1495 Walker, X. J., Rogers, B. M., Baltzer, J. L., Cumming, S. G., Day, N. J., Goetz, S. J., Johnstone, J. F., Schuur, E. A. G., Turetsky, M. R., and Mack, M. C.: Cross-scale controls on carbon emissions from boreal forest megafires. *Global Change Biology*, 24(9), 4251–4265. <https://doi.org/10.1111/gcb.14287>, 2018.
- Wang, J. A., Baccini, A., Farina, M., Randerson, J. T. and Friedl, M. A.: Disturbance suppresses the aboveground carbon sink in North American boreal forests. *Nature Climate Change*, 11(5), 435–441, 2021
- 1500 Wang, T., Hamann, A., Spittlehouse, D., and Carroll, C.: Locally Downscaled and Spatially Customizable Climate Data for Historical and Future Periods for North America. *PLOS ONE*, 11(6), e0156720. <https://doi.org/10.1371/journal.pone.0156720>, 2016.
- 1505 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): A high resolution global model to estimate the emissions from open burning. *Geoscientific Model Development*, 4(3), 625–641. <https://doi.org/10.5194/gmd-4-625-2011>, 2011.

Wright, M., Ziegler, A.: ranger: A Fast Implementation of Random Forests for High Dimensional Data in C++ and R. *Journal of Statistical Software*, 77(1), 1-17. doi:10.18637/jss.v077.i01, 2017

1510 Wolfe, J. A.: Temperature parameters of humid to mesic forests of Eastern Asia and relation to forests of other regions of the Northern Hemisphere and Australasia: analysis of temperature data from more than 400 stations in Eastern Asia (U.S. Geological Survey Professional Paper No. 1106). Washington, DC: US Dep. Int., 1979.

1515 Young, A. M., Higuera, P. E., Duffy, P. A., and Hu, F. S. Climatic thresholds shape northern high-latitude fire regimes and imply vulnerability to future climate change. *Ecography*, 40(5), 606–617. <https://doi.org/10.1111/ecog.02205>, 12017.

Zhao, B., Zhuang, Q., Shurpali, N., Köster, K., Berninger, F., and Pumpanen, J.: North American boreal forests are a large carbon source due to wildfires from 1986 to 2016. *Scientific Reports*, 11(1), 7723. <https://doi.org/10.1038/s41598-021-87343-3>, 2021

1520

1525

1530

1535