



1    **Supplementary information**

2    Figure S1: Monthly means of  $\text{NH}_3$  total columns (molecules/cm<sup>2</sup>) derived from 10 years (2008-2017) of IASI  $\text{NH}_3$ -  
3    retrieved columns over the so-called Europe domain. The blue cross indicates Paris location.




4

5 Figure S2: Same than figure S1 but for the North America region.



6

7     Figure S3: Same than figure S1 but for the southern North America region.




8

9

10 Figure S4: Evolution of NH<sub>3</sub> with respect to land surface temperature (upper panels) and relative humidity (lower  
 11 panels) from ERA5 in the 3 study domains. The figure is done by averaging IASI NH<sub>3</sub> total columns per bins of  
 12 ERA5 skin temperatures, with an interval of 1°C. We do not consider bins that contain less than 5% of the  
 13 maximum number of measurements per bin; hence, the averages with not enough measurements per bins are  
 14 excluded. Blue dots: Yearly IASI NH<sub>3</sub> total columns (molecules/cm<sup>2</sup>) averaged per bins of ERA5 skin temperature  
 15 (°C, upper panels) and of relative humidity (%), lower panels), with an interval of 1°C between each consecutive  
 16 bin. The regions considered here are the regions presented in Table 1 above Europe (left panel), North America  
 17 (middle panel), and southern North America (right panel). The red line is a polynomial fit of second order, and  
 18 the relevant r<sup>2</sup> is shown on each panel.

19




20

21 The effect of relative humidity on NH<sub>3</sub> concentrations is different in each of the study domains, as we  
 22 can see in the lower panels of Figure 4. The highest correlation factor we observe is over Europe, and  
 23 it accounts to  $r^2 = 0.82$  (lower left panel). A study by Reynolds and Wolf (1987) concluded that the  
 24 relative humidity of the air does not play a major role in NH<sub>3</sub> volatilization unless the soil is dry. In fact,  
 25 soil is drier in Europe than both of the other regions, with North America being the most humid area.  
 26 This can explain why we see a good correlation in Europe and lower one in North America. In southern  
 27 North America, however, throughout the year we observe high temperatures and high humidity, which  
 28 can explain the low correlation factor  $r^2 = 0.30$  (lower right panel).

29 Despite the differences in the correlations, we can still see a decreasing trend of ammonia as the  
 30 relative humidity of the air increases. We looked at the times during which the NH<sub>3</sub> concentrations  
 31 where detected and we summarize them below:


- 32     • RH = 0 – 40%: Most of the NH<sub>3</sub> detected in all regions is during the spring season when the  
 33       concentrations are the highest. One note that when  $0 \leq RH \leq 25\%$ , April dominates in Europe  
 34       and May dominates in North America and southern North America.
- 35     • RH = 40 – 60%: The NH<sub>3</sub> detected is during summer and spring, hence the lower average as the  
 36       RH increases when the summer approaches.
- 37     • RH = 60 – 85%: The NH<sub>3</sub> is decreasing as RH increases and as the time approaches winter (when  
 38       RH is highest). In southern North America, however, these ammonia measurements  
 39       correspond to the spring season mostly.
- 40     • RH = 85 – 100%: Most of NH<sub>3</sub> detected are during winter in southern North America and  
 41       Europe, and evenly distributed throughout the year in North America.

42 Figure S5: Yearly IASI NH<sub>3</sub> total columns (molecules/cm<sup>2</sup>) averaged per bins of ERA5 skin temperatures (°C), with  
 43 an interval of 1°C between each consecutive bin. The red circles denote the growing seasons, at least 60% of the  
 44 NH<sub>3</sub> are detected during March-May and Sept-Nov periods. The regions considered here are the regions  
 45 presented in Table 1 in North America.




47 Figure S6: Yearly IASI NH<sub>3</sub> total columns (molecules/cm<sup>2</sup>) averaged per bins of ERA5 skin temperatures (°C), with  
 48 an interval of 1°C between each consecutive bin. The red circles denote the growing seasons, at least 60% of the  
 49 NH<sub>3</sub> are detected during March-May and Sept-Nov periods. The regions considered here are the regions  
 50 presented in Table 1 above southern North America: (a) sub-regions A to I, (b) sub-regions J to S.

51 (a)



52

53 (b)



54