Influences of manganese cycling on alkalinity in the redox stratified water column of Chesapeake Bay

Aubin Thibault de Chanvalon^{1,2}, George W. Luther², Emily R. Estes^{2,*}, Jennifer Necker², Bradley M. Tebo³, Jianzhong Su^{4,+}, Wei-Jun Cai⁴

 ¹Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM, Pau, France.
 ²School of Marine Science and Policy, University of Delaware, Lewes, Delaware, 19958 USA
 ³Division of Environmental and Biomolecular Systems, 3181 SW Sam Jackson Park Road, Portland, OR 97239, USA; Current address: Department of Chemistry, University of Washington, Seattle, WA 98195-1700
 ⁴School of Marine Science and Policy, University of Delaware, Newark, Delaware, 19716 USA

⁺State Key Laboratory of Marine Resources Utilization in South China Sea, Hainan University, Haikou, China *Texas A&M University, USA

Correspondence to: A. Thibault de Chanvalon (aubin.thibault-de-chanvalon@univ-pau.fr)

Abstract.

10

The alkalinity dynamics in coastal environments play a crucial role in controlling the global burial of carbonate minerals

- 15 and the ocean's capacity to sequester anthropogenic CO2. This study presents results from high vertical resolution profiles obtained during two summers in the temperate Chesapeake Bay estuary, enabling detailed investigation of carbonate dynamics over salinity and redox gradients, along with measurement of the speciation of most redox-sensitive elements. Under oxygenrich conditions, carbonate dissolution, primary production, and aerobic respiration explain the evolution of total alkalinity (TA) versus dissolved inorganic carbon (DIC), once adjusted for fresh and oceanic water mixing. A significant flooding event
- 20 in 2018 promoted carbonate dissolution. In oxygen-depleted waters, we observed a previously unreported 2.4-mole increase in DIC per 1 mole of TA production, which was consistent over the two years. Stoichiometry changes suggest that MnO2 reduction followed by Mn carbonate precipitation is responsible for this characteristic carbonate signature, likely produced in sediment pore water and then transferred to the water column along with other by-products of anoxic respiration as summer begins. Our findings highlight the critical role of Mn in alkalinity dynamics, especially in river-dominated environments where
- 25 it can limit H2S oxidation to SO42- and promote sulphur burial.

1 Introduction

About 30% of anthropogenic CO_2 emissions are rapidly trapped by dissolution in the ocean as dissolved inorganic carbon (DIC), which is dominated by bicarbonate ions (HCO_{3⁻}, Friedlingstein et al., 2019). At the century time scale, atmosphere-

- 30 ocean exchanges result in oceanic HCO_3^- enrichment not associated with a cationic enrichment, as measured by total alkalinity (TA). This is in contrast to silicate or carbonate weathering, which preponderates at timescales of thousands to million of years and increases TA simultaneously to DIC (Urey, 1952). This disequilibrium corresponds to an excess of proton release that can be tracked by an indicator such as TA minus DIC (Xue and Cai, 2020) and informs about the reduction of CO_2 buffering capacity of seawater (Gustafsson et al., 2019). The proton excess is only balanced in the deep ocean by the production of Ca^{2+}
- 35 from carbonate dissolution, a process named the chemical carbonate compensation (Boudreau et al., 2018). However, in most shallow waters, where carbonate precipitation largely predominates over dissolution and accounts for 2/3 of buried carbonate (Smith and Mackenzie, 2016), other localised processes may constrain carbonate dynamics (Borges et al., 2006; Lohrenz et al., 2010). For example, some calcifying species may slow down their carbonate precipitation in case of pH decrease (so called biological carbonate compensation; Boudreau et al., 2018). As another example, in estuaries, the seasonality of river flow,
- 40 temperature and continental erosion modulates CaCO₃ dissolution (e.g. Su et al., 2020b), atmospheric CO₂ exchanges (e.g. Borges et al., 2018), respiratory activity (e.g. Abril et al., 2003) and transfer of carbonate particles (Meybeck, 1987; Middelburg et al., 2020).

The global trend of human migration towards littoral areas associated with global warming favours eutrophication and a decrease in oxygen levels in coastal water (Breitburg et al., 2018; Rabalais et al., 2014). Sometimes driven by recurrent natural

- 45 process (e.g. Gupta et al., 2021), exceptional events (e.g. Hulot et al., 2023), stratification reinforced by global warming (e.g. Meire et al., 2013) or anthropogenic nutrient loading (e.g. Hagy et al., 2004); coastal deoxygenation multiplies the possibilities for anoxic conditions to build up as a permanent or seasonal feature. In anoxic environments, the drivers of the carbonate cycle change as the terminal electron acceptor for organic matter remineralisation shifts from O₂ to Mn oxides, Fe oxides then sulphate. Notably, the consumption of the negatively charged SO₄²⁻ by sulphate reducers, followed by sulphur burial, produces an significant alkalinity flux toward coastal water, accounting for approximately 10 % of the global alkalinity input to the
- ocean (Middelburg et al., 2020). Fe and Mn oxides play a critical role in sulphur burial as they prevent SO_4^{2-} regeneration and lead to the production of S⁰ or FeS instead (Findlay et al., 2014; Avetisyan et al., 2021), leaving a specific fingerprint in the TA and DIC concentrations.

To better constrain the carbonate cycle in temperate microtidal estuaries, we sampled a stratified water column in the 55 Chesapeake Bay eleven times over two campaigns with a high vertical resolution (down to 10 cm). This protocol allows a precise description of carbonate dynamics over a redox gradient along with the measurement of the speciation of most redox sensitive elements. Such sampling illustrates carbonate chemistry on transitioning from oxygenated waters to waters devoid of oxygen as usually only encountered in sediments or in anoxic lakes or seas (*e.g.* Black Sea). The original observed changes of alkalinity versus dissolved inorganic carbon changes are interpreted based on typical geochemical reactions occurring along

60 the redox gradient.

2 Materials and methods

2.1 Sampling

During two sampling campaigns from August 3rd to 9th, 2017 and July 28th to August 3rd 2018, eleven profile casts were conducted in a single station in the Chesapeake Bay with a water depth of 25 m (Station 858, 38°58.54'N; 076°22.22'W). The Susquehanna River is the main tributary of the bay representing on average 2/3 of the fresh water input (Zhang et al., 2015). Despite similar season, the two campaigns occurred at very different river flow with about 850 m³ s⁻¹ in 2017 versus 8500 m³ s⁻¹ in 2018 due to release of flood waters from the Conowingo Dam. The August 2018 condition corresponds to flooding which occurs on average every 3.5 years (return period of 3.5 years, USGS survey).

- Each CTD cast was performed during low or high tide slack periods. An oxygen sensor (Clark electrode, SBE Inc.; 70 detection limit of 1 μM) and fluorescence sensor (Eco-FL Fluorometer, WETLabs) were part of the CTD Rosette to take measurements during sampling. Also, a submersible all plastic pump profiler was attached with the pump near the sensor orifices allowing measurement and sampling at a resolution of a few centimetres over 25 m water depth. Water was pumped to the deck within 1 minute and water passed through a flow through voltammetry system measuring continuously O₂, Mn(II), Fe(II), organically complexed Fe(III), FeS clusters, H₂S and polysulfides (Hudson et al., 2019). When redox interfaces were
- 75 identified, samples were filtered through an acetate cartridge filter (pore size 0.45 µm) for pH and inorganic carbon parameters, which were processed onboard within a few hours after sampling in order to conserve chemical speciation. The pump profiler system was cleaned with deionised water (18 MΩ) onboard the deck of the ship after deployment. No coating effects were observed with the pump system.

2.2 Discrete Measurements

- For each sample, all redox species were determined in the through flow voltammetry system using cyclic voltammetry with a 100 μ m diameter Au/ Hg amalgam PEEK microelectrode prepared according to Luther et al. (2008) connected to a DLK-60 electrochemical analyser from Analytical Instrument Systems Inc. The detection limit of this method is 0.2 μ M for sulfide and polysulfides. Discrete samples for the determination of NO₂⁻, Fe and Mn species were filtered through nylon luer-lock syringe filters (Millipore, 0.20 μ m). Iron was measured on both bulk and filtered samples using the ferrozine method (Stookey, 1970):
- after HCl acidification (for Fe(II)) and an optional reduction step (for Fe(III)+Fe(II)) with hydroxylamine hydrochloric (final concentration 0.7 M for 1 hour), ammonium actetate (final concentration 0.5 M) and ferrozine (final concentration 1 mM). Absorbance at 562 nm was read with a diode array spectrophotometer (Hewlett Packard 8452B). Limit of detection is 100 nM for Fe(II) and Fe(III) with a 1-cm cell. Shipboard nitrite determination was performed using the method of Grasshoff

(1983). To 25 ml of sample, 0.5 ml of 58 mM sulfanilamide in 10% v/v HCl and 0.5 ml of a 4 mM N(1-naphthyl)ethylene

90 diamine hydrochloride solution were added. Samples with added reagents were shaken and left to sit for 15 min, followed by UV–Vis analysis at 540 nm using a 10-cm cell to increase detection limits. Calibration curves were constructed using sodium nitrite. Limit of detection is 10 nM for NO₂⁻.

Dissolved manganese was determined by displacement of a Cd(II)-porphyrin complex with Mn(II) to form the Mn(II)porphyrin complex (Ishii et al., 1982). Mn(III) species were identified based on slower reactivity with the Cd complex

- 95 (Madison et al., 2011) as modified in Thibault de Chanvalon and Luther (2019). Alternatively, Mn(III) species were identified after HCl treatment (down to pH=1.5) followed by filtration in order to flocculate and eliminate the dissolved manganese bound to humic material by filtration (Oldham et al., 2017b). Limit of detection is 50 nM for Mn(II) and Mn(III) in a 1-cm cell. MnOx was measured on 20 mL samples of suspended material retained on 0.2 µm filters by the Leucoberbelin blue (LBB) method (Jones et al., 2019). Four millilitres of a reagent solution ([LBB]= 78 µM, [acetic acid]=14mM) react with the
- 100 filter and the absorbance is read at 624 nm. KMnO₄ was used to calibrate the LBB method which allows the calculation of the electron equivalents obtained from particulate MnOx. Results are given in as MnO_2 equivalent with a limit of detection of 0.1 μ M and an uncertainty below 5%.

The DIC samples were preserved in 250-mL borosilicate glass bottle with 50 μ L saturated HgCl₂ solution. The total alkalinity (TA) samples were not poisoned to prevent HgS precipitation and H⁺ release in anoxic and low salinity waters (Cai

- 105 et al. 2017). Then TA was analysed by Gran titration in an open-cell setting (AS-ALK2, Apollo Scitech) within 24 h of collection (Cai et al. 2010a). The DIC samples were measured by a nondispersive infrared analyzer (AS-C3, Apollo Scitech) within a week (Huang et al., 2012). The precision for DIC and TA was about 0.1%. Both DIC and TA measurements were calibrated against certified reference materials (CRMs Batch 163 and Batch 173 provided by Andrew Dickson of the Scripps Institution of Oceanography). The pH samples were measured onboard at 25 °C within 1 h of collection using an Orion Ross
- 110 glass electrode, and calibrated with NIST standard buffers. The pCO_2 , calcite saturation and TA were calculated from measured DIC and pH via CO2sys program using Cai and Wang (1998) constants. The measured TA was found highly correlated to the calculated TA ($r^2 = 0.995$ and 0.998, slope = 0.995 and 1.017 for 2017 and 2018 campaign respectively), and their difference was always below 45 μ M with an average of 7.5 μ M for 2017 and of 22.2 μ M in 2018. These results suggest low contribution of non-carbonate species (e.g. nitrite, ammonium or organic matter (Cotovicz Jr. et al., 2016)) and measured TA was used for
- 115 the interpretation.

2.3 Models of biogeochemical process on TA and DIC

2.3.1 Identification of biogeochemical processes from scatter plots

This section identifies the required conditions to interpret a scatter plot of two species in term of chemical reaction stoichiometry: the "reaction driven" approximation. First, equation (1) describes the behaviour of a solute, in case of turbulent

120 diffusion mixing, with D_s the effective diffusion coefficient (sometimes called eddy diffusion or diffusive like mixing) superimposed on a chemical reaction as described in equation (2):

$$\frac{dC}{dt} = D_s \frac{d^2C}{dx^2} + \alpha_C v \tag{1}$$

$$\alpha_C C + \alpha_D D = \alpha_E E + \alpha_F F \tag{2}$$

Assuming steady state, *i.e.* dC/dt=0, and applying equation (1) to two different species sharing the same reactions at rate v on each point and $\alpha_D \neq 0$, we can express the rate of the reaction as:

$$-\frac{v\,\alpha_C}{D_s} = \frac{d^2C}{dx^2} = \frac{\alpha_C}{\alpha_D}\,\frac{d^2D}{dx^2} \tag{3}$$

Equation (3) is also true for any linear combination of a solute with a conservative element such as the salinity, S, because 125 $\alpha_S = 0$, hence:

$$\frac{d^2C + kS}{dx^2} = \frac{d^2C}{dx^2} + k \frac{d^2S}{dx^2} = \frac{d^2C}{dx^2}$$

By integration of equation (3) with respect to x on a portion of space where α_C/α_D is constant, it becomes equation 4:

$$\frac{dC}{dx} = \frac{\alpha_C}{\alpha_D} \frac{dD}{dx} + G \tag{4}$$

Here, C and D are concentrations of any elements produced or consumed during the reaction and can be TA, DIC, or a linear combination with salinity. G is the constant rising from the integration, and needs to be null for equation (5) to be valid.

$$\frac{dC}{dD} = \frac{\alpha_C}{\alpha_D} \tag{5}$$

130

Experimentally, in this case, the data points would be distributed along a straight line in a C versus D scatter plot. Theoretically, it can occur in three cases. First, a straight line occurs if the contribution from one endmember is negligible compared to the time scale of reactions, which corresponds also to a system with only one endmember. A second case where G=0 occurs when the slope of $\Delta C/\Delta D$ between the endmembers equal α_C/α_D which is achieved, in particular, if one endmember was previously generated from the other endmember by a chemical reaction with similar stoichiometry. Beside these two very particular cases, a general case can be depicted, in the case when the rate v can be described by a polynomial function of x:

$$\frac{d^2C}{dx^2} = -\frac{\alpha_C}{D_S}\sum_i a_i x^i$$

135 That can be solve analytically and gives the following equation:

$$\frac{dC}{dx} = \frac{\alpha_C}{D_S} \sum_i \left[\frac{a_i}{(i+1)(i+2)} \left(1 - (i+2) x^{i+1} \right) \right] + \Delta C \tag{6}$$

Here ΔC is the difference of concentration between the two endmembers. With similar reasoning on D and with $\Delta C = \Delta D$ =0, we find equation (4) with G=0. In this study, while $\Delta TA \neq \Delta DIC \neq 0$, the excess of TA (TAex) and the excess of DIC (DICex) are calculated by linear combination with salinity to be equal to zero for the upstream and downstream endmembers reaching the condition $\Delta TAex = \Delta DICex = 0$. Thus, if only one reaction occurs in-between, the slope $\Delta TAex/\Delta DICex$ would correspond to the stoichiometry of α_{TA}/α_{DIC} .

The oceanic endmember for both August campaigns was the one proposed by Su et al. (2020a). The oceanic endmember varies mainly with season (Cai et al., 2020) and a maximal change of 50 μM results in 5% uncertainty on the slope of the mixing line. For the upstream estuary, larger variations exist mainly due to changes of weathering intensity and river discharge (Meybeck, 2003; Joesoef et al., 2017), and a one-off endmember has to be determined by fit with the in situ measurements at the lowest measured salinity. Between the two campaigns, the upstream endmember changed by 77 μM generating 5% of change on the slope (see Fig. A1). The upstream endmember is not a river endmember (Su et al., 2020a) but corresponds to a salinity above 1.5 avoiding any interpretation for biological activity in the fresh water part of the estuary (Meybeck et al., 1988). However, it corresponds to a larger water mass pool, less sensitive to short term changes, with a residence time being higher than 240 days in the Chesapeake upstream part (Du and Shen, 2016), and thus is more likely to satisfy the condition of

150 stability of the endmember, which is a prerequisite of the steady state assumption.

The relative uncertainty on $\Delta TAex/\Delta DICex$ is equal to the sum of the relative uncertainty of $\Delta TAex$ and $\Delta DICex$. Posing ΔTA the change of TA measured, ΔS the change of salinity and sml_TA, the slope of the mixing line for TA, we have $\Delta TAex = \Delta TA - sml_TA \times \Delta S$. Uncertainty on ΔTA and ΔS are negligible to the relative uncertainty of the slope of the mixing line. Posing $\delta(x)$ as the uncertainty on x, we get equation (6) that describes the fact that the uncertainty is much lower on $\Delta DICex$ than on DICex because most the error associated with the calculation of the endmember is cancelled when calculating the

$$\frac{\delta \left(\Delta TAex / \Delta DICex\right)}{\Delta TAex / \Delta DICex} = \frac{\delta \left(\Delta TAex\right)}{\Delta TAex} + \frac{\delta \left(\Delta DICex\right)}{\Delta DICex} = \frac{\delta \left(sml_TA\right)}{sml_TA} + \frac{\delta \left(sml_DIC\right)}{sml_DIC} = 0.1$$
(6)

However, in a stratified water column, not only one but several successive reactions occur, requiring many integrations of equation (3). On the boundary of each space portion with constant α_C/α_D , specific local endmembers are defined with concentrations at steady state fixed due to the ongoing reactions and not due to the inertia of a large body of water. The general case is not straightforward to solve, but in the particular case where the C versus D plot represents a straight line in a portion of space, equation (4), still valid in each portion of space, indicates that G=0; thus, the local endmembers are maintained in steady state by chemical reactions with a similar apparent stoichiometry than the one that produced them, *i.e.* $\Delta C/\Delta D = \alpha_C/\alpha_D$.

Therefore, in a system defined between only two endmembers, away from atmospheric exchanges, in case of turbulent diffusion mixing, at steady-state and with negligible lateral mixing, the "reaction driven" approximation allows us to interpret

140

155

165 linear variations of TAex versus DICex as a sum of biogeochemical reactions spread all over the water column that can be broken into several discrete reaction zones. In each zone, if the local Δ TAex/ Δ DICex ratio is constant, it corresponds to the apparent stoichiometry of a combination of the biogeochemical reactions occurring in this zone. In the case of multiple simultaneous reactions in the same zone, by posing α_c^i the stoichiometry of the ith reaction concerning the reagent C and v^i the reaction rate of the ith reaction, we obtain:

$$\alpha_C v = \sum_i \alpha_C^i v^i$$

170 To maintain a global reaction rate independent to the species we have

$$v = \sum_{i} v^{i}$$

$$\alpha_{c} = \frac{1}{v} \sum_{i} \alpha_{c}^{i} v^{i}$$
(7)

Equation 7 indicates that the apparent stoichiometry in a given zone corresponds to the sum of the stoichiometric coefficients of each reaction weighted by the relative rate of each reaction. Therefore, to estimate the relative rate of each reaction to the observed local changes of TAex, DICex and AOU or H₂S, a linear combination of reactions is calculated. This combination has to fit 3 equations (one for each parameter), which allows a maximum of 3 reactions to be used to solve the system. A limited number of reactions is selected as candidates based on the discussion (see Table 1 and sections 4.2 and 4.3). Then, the system is solved with the minimum possible reactions, and the weighted coefficients, v^{i}/v , are calculated.

2.3.2 TA changes indicated by reaction stoichiometry

The simplest way to calculate the TA changes induced by an individual reaction is to do a direct look at reaction stoichiometry. Indeed, the total alkalinity (TA) corresponds to the quantity of acid added to titrate a solution down to pH 4.5 (Dickson, 1981). It can be described by equation (8), with the example of HCl as acid and B⁻ any titrated base.

$$HCl + B^{-} \to BH + Cl^{-} \tag{8}$$

Assuming a complete reaction, the quantity of acid added is equal to the negative charges initially present in the sample consumed plus the positive charges added to the species initially present in the sample. Thus, the total alkalinity corresponds to the loss of negative charges (or gain of positive charges) for species initially present in the sample produced by the pH change from the initial pH = pHini to pH = 4.5. Writing z_i^{pH} , the charges held by species *i*, initially present in the sample at a given pH, we get equation (9):

$$TA = \sum_{i} z_{i}^{pH=4.5} - \sum_{i} z_{i}^{pHini}$$
(9)

so

175

However, the electroneutrality of water induces:

$$\sum_{i} z_{i}^{pHini} = 0$$
$$TA = \sum_{i} z_{i}^{pH=4.5}$$
(10)

so

Equation (10) demonstrates than the total alkalinity is simply the sum of charges that each species present in the sample would have at pH 4.5. From Eq. (10), one can easily deduce the changes of alkalinity from any reaction stoichiometry as soon as the charges of all species at pH = 4.5 are known. For example, for a 0.0020 M NaOH solution, its TA is simply 0.0020 M as at pH=4.5, TA= ([Na⁺] + [H⁺] - [OH⁻]) = 0.002 + 10^{-4.5} - 10^{-14+4.5} = 0.0020 M. For natural waters, most of the time, the only acid/base species that are charged at pH = 4.5, are H₂PO₄⁻ and NH₄⁺. In that respect, whatever the initial pH and the acid-base equilibrium of species in the sample is, the sum of phosphate species will count negatively and the sum of ammonium species will count positively. Strictly speaking, at pH=4.5, acid species with pKa between 2.5 and 6.5, such as F⁻ and NO₂⁻, would be only partially titrated and the charge equals their concentration multiplied by a correction factor of (1+10^{pKa-4.5})⁻¹, but this correction can be neglected to a first approximation. Eq. (10) is equivalent to those published in Soetaert et al. (2007) or Wolf-Gladrow et al. (2007) whose equation 32 can be refined considering that :

$$\sum_{i} z_{i}^{pH=4.5} = [Na^{+}] + 2 [Mg^{2+}] + 2 [Ca^{2+}] + [K^{+}] + 2[Sr^{2+}] + \dots - [Cl^{-}] - [Br^{-}]$$
$$- [NO_{3}^{-}] - \dots TPO4 + TNH3 - 2TSO4 - THF - THNO2 - \dots$$

However, Eq. (10) is more general. For example, in suboxic water, specific species such as polysulfides (as HS_8^{2-} , Rickard and Luther, 2007) and in highly productive environments, carboxylic groups from DOC can be easily added as soon as the bearing charges at pH = 4.5 are known.

200 3 Results

3.1 Water column stratification

High-resolution profiles of salinity and temperature plotted in Fig. 1 show the stratified water column at station 858.
Carbonate and redox chemistry are plotted against salinity in Fig. 2. These data result from the 11 CTD casts performed over 1 week during each campaign and correspond to depths ranging from 0.7 to 25 meters depth. Plots against depth generate noisier profiles are shown in Fig. A2 while plots against salinity follow the water masses. Despite the overall lower salinity due to a near 7-fold river flow increase in 2018 than 2017 (5800 m s⁻¹ in 2018 versus 850 m³ s⁻¹ in 2017), similar zonation of the water column occurred.

A surface layer sampled only in 2018 is visible in the top 3 m depth (Fig. 1). It presents highly variable temperature and oxygen concentration and oversaturation of pCO2 (Fig. 2) indicating export to the atmosphere. Below, at 3 m depth, a

- subsurface layer (named primary production zone or PP in Fig. 2) is characterized by a high amount of O₂ (about or above 100% saturation), high pH (about 8; 8.11 ± 0.07 , n=13 in 2017 and 7.94 ± 0.08 , n=14 in 2018) and high day to day temperature variation (above 1 °C between different days). The layer presents relatively low pCO₂ (505 \pm 75 μ atm, n=13 in 2017 and 770 \pm 130 μ atm, n=14 in 2018) with minimal values at 110 μ atm in 2017 and 205 μ atm in 2018, which are below the atmospheric pCO₂ of 407 μ atm (Chen et al., 2020)). This signature corresponds to primary production (PP). Fluorescence (not shown)
 - 215 correlates with pH as expected for primary production (pH = Fluo (mV) x 13 +7.14, r^2 =0.8 in 2017 and r^2 =0.9 in 2018). DICex and TAex reach their minimal value in the surface layer. While DICex minimum is similar between the two campaigns (78 ± 17 µM, n= 13 in 2017 and 76 ± 10 µM, n=14 in 2018); TAex minimum is much lower in 2017 (7 ± 3 µM, n=12) than in 2018 (48 ± 7 µM, n=14).



Figure 1: Superimposed salinity (in white) and temperature (in black) profiles over the 11 casts for each campaign.

- Below, with increasing salinity, an important increase of pCO₂ accompanying the decrease of O₂, pH and temperature is
 visible. A relatively invariable low oxygen zone (called ILO in Fig. 2) is here defined by the salinity invariance of O₂ concentrations at about 24 ± 3 µM (n=15) in 2017 and 105 ± 5 µM (n =16) in 2018. Other species are also relatively stable for this depth such as pCO₂, at about 3250 ± 150 µatm (n=10) in 2017 and 2590 ± 100 µatm (n = 16) in 2018, and pH, about 7.28 ± 0.01 (n=13) in 2017 and 7.37 ± 0.02 (n=16) in 2018. Deeper, where the oxygen is not detectable (< ~ 1 µM) and in absence of free sulphide, the so-called anoxic zone corresponds to a pH minimum at 7.24 ± 0.01 (n=40), similar in 2017 and 2018 that
 generates a pCO₂ maximum. The deepest layer is a sulfidic layer, [H₂S] = 11.2 ± 2.8 µM (n=36), in which the pH seems quite
- 225 generates a pCO₂ maximum. The deepest layer is a sumfic layer, $[H_2S] = 11.2 \pm 2.8 \,\mu$ M (n=50), in which the pH seems quite stable at 7.33 ± 0.01 (n=29) and 7.32 ± 0.01 (n=9) in 2017 and 2018 respectively. The Ca²⁺ concentrations observed by Su et al. (2021) and during the 2018 cruise (data not shown) vary linearly with salinity (calcium excess stays below 200 μ M or 10% of total Ca). Assuming similar behaviour in 2017, calculations show that the whole water column (except 4 samples from the PP zone) is under saturated (0.36<Ωcal<1; mean=0.68) with respect to calcite in 2018, while undersaturation is only valid



Figure 2: Superimposed carbonate and redox chemistry profiles of 11 casts done in August 2017 and in August 2018 at the station 858 of the Chesapeake Bay.

230 below S=10 in 2017. The main changes between the two campaigns correspond to a greater oxygen penetration in 2018, especially visible in the ILO zone. Additionally, a surface layer (with salinity below 3) is visible above the primary production zone in 2018 but is related to the more superficial sampling in 2018.

3.2 Intermediate redox species

The development of the anoxic zone during summer (Su et al., 2021) and the regularity of this development over the years

- 235 (Sholkovitz et al., 1992; Trouwborst et al., 2006; Lewis et al., 2007; Cai et al., 2017; Oldham et al., 2017a), requires the presence of species able to rapidly oxidise the H₂S mixing upward and to reduce the O₂ mixing downward. The three main redox couples known to play this role, NO₃⁻/NO₂⁻, MnO_x/Mn²⁺ and Fe³⁺/Fe²⁺ are described in Fig. 3 by the superimposition of all cast results against salinity. Four representative casts are plotted in the Fig. A3. The primary production zone is depleted in dissolved Mn(II) and Mn(III), and shows an average value for solid MnOx (0.7 ± 0.1µM, n=4) and NO₂⁻ (2.1 ± 0.1 µM,
- n=18). In 2018, iron speciation was investigated in the primary production zone showing an important iron pool (1.6 ± 0.1 μM, n=9) dominated by solid Fe(II) (95 ± 2%). Below, in the ILO zone, NO₂⁻ reaches a maximum plateau at 5.3 ± 0.2 μM, n=11 in 2017 that is not visible in 2018, MnOx dominates the Mn pool in the ILO zone with concentration of 1.8 ± 0.1 μM, n=14 in 2017 and 1.3 ± 0.2 μM, n=3 in 2018). Fe(II) represents only 49% ± 13, n=4 of the iron pool (for a total iron concentration of 2.0 ± 0.4 μM, n=4) in 2017 while it represents 87 ± 5%, n=4 of the iron pool in 2018 (for a total concentration of 1.2 μM ± 0.2, n=4). Just below oxygen depletion, in the anoxic zone, MnOx reaches a maximum (2.0 ± 0.1 μM, n=25 in
- 2017 and 2.9 \pm 0.4 μ M, n=12 in 2018). During both campaigns, the sulfidic layer is characterised by the absence of NO₂⁻ and MnOx while dissolved manganese concentration increases up to 7.7 \pm 0.8 μ M, n=15 and the iron pool increases to 5.0 \pm 0.7



Figure 3: Mn, Fe and nitrite profiles build by superimposition of 11 casts for each campaign.

 μ M, n=13. No Mn(III) was detected with the porphyrin kinetics method (Thibault de Chanvalon and Luther, 2019) but about 30% of the total dissolved manganese flocculated after acidification down to pH 1.5 when analysed in 2017 indicating the

250 existence of Mn(III).

4. Discussion

In term of salinity, Station 858 of the Chesapeake Bay shows very similar water column zonation between summer 2017 and summer 2018 despite a 10-fold difference in freshwater discharge rates (Fig. 1 and Fig. 2). Major features are: first, a surface layer characterized by intense atmospheric exchange that was only sampled in 2018. Below, at about 3 meter depth, a 255 subsurface layer associated with high primary production (PP zone) with high pH about 8, oversaturation of dioxygen and low CO₂ partial pressure (down to 110 µatm). Below, a low oxygen layer with invariant concentration of most species surveyed (the invariant low oxygen (ILO) zone) is characterized by significant nitrite accumulation in 2017 (Fig. 2), probably due to oxidation of NH_4^+ diffusing upward and/or produced by *in situ* remineralisation. This feature is not visible in 2018 probably because the higher O₂ concentration in 2018 accelerates nitrite oxidation into nitrate and prevents any significant accumulation. 260 Below, the anoxic zone, with neither O₂ nor H₂S detectable, is characterized by an increase of MnOx concentration and a pH minimum. This MnOx maximum can be explained by the upward diffusing Mn²⁺ that is biologically oxidized by the downward diffusing O_2 , though at low, undetectable, concentration (Clement et al., 2009). Additionally, Mn^{2+} could be oxidized by the nitrite or the nitrate (not measured) diffusing downward (thermodynamically favourable (Luther, 2010)). Compared to the ILO zone, the anoxic MnOx maximum corresponds to an increase of 0.2 μ M in 2017, while it is much more marked in 2018 with an increase of 1.6 μ M (Fig. 3). This difference could come from a faster Mn²⁺ oxidation produced by the steeper oxygen 265 gradient above the anoxic zone and by the thinness of the anoxic layer in 2018. Finally, in the deeper sulfidic layer, the MnOx disappearance corresponds to the Mn^{2+} increases (Fig. 3) according to the reduction of settling MnOx by H₂S. The concentration increase of the manganese pool and of the iron pool with depth in the anoxic and sulfidic layers probably results from important sedimentary efflux.

270 **4.1 Validity of the "reaction driven" approximation**

The simultaneous and high-resolution sampling of multiple carbonate parameters and redox species gives us the rare opportunity to investigate in detail the interaction between carbonate species and redox sensitive elements. In the Chesapeake Bay, main changes of DIC and TA can be explained by mixing between upstream and oceanic endmembers (Appendix, Fig. A1). This "endmember driven" interpretation leads to the calculation of an excess of DIC and TA, DICex and TAex, relative to the mixing line as shown in Fig. 4a. At station 858, the steep gradient observed, for example the pH and pCO₂ gradients in the PP zone, the O₂ and NO₂⁻ gradients above the anoxic zone and the Mn, Fe and H₂S gradients at depth, suggest that ongoing in situ processes control the changes of concentrations and dominate the time-dependent endmember variability or the mixing with an unknown third endmember. Additionally, the TAex versus DICex plot (Fig. 4a) shows steep changes of direction, such

as in case of straight lines between local endmembers maintained by ongoing reactions while a preponderance of mixing would

280

produce more progressive changes. Finally, at depth, Fig. 4a shows a similar slope for both years studied ($\Delta TAex/\Delta DICex =$

2.4) rather than a similar TAex and DICex concentration, which reinforces the validity of the "reaction driven" approximation.



Figure 4: Description of the Taex/DICex/AOU system. (a) and (b) panels show samples measured along with interpretative slope discussed in the main text. Panel (c) shows the theoretical slope of TAex/DICex from combination of reactions presented in Table 1.

Assuming 1) that mixing is efficiently described by vertical turbulent diffusion mixing, 2) that the measured concentrations correspond to a steady state – no changes observed over the 1 week sampling, 3) that no additional endmember contributes
significantly to the excess calculation, in particular that the samples are isolated from atmospheric exchanges and 4) that lateral mixing is negligible, which is equivalent to the lateral invariance of the system – as in the stratified water column of station 858; the "reaction driven" approximation (section 2.3.1) permits interpretation of the concentration changes as a linear combination of the stoichiometry of several chemical reactions (equation 5). The weighted coefficients of each reaction are equal to the rate of each reaction relative to the sum of the rate of all occurring reactions, *vⁱ*/*v* (equation 7). This interpretative
framework describes the vertical stratification of the water column as the journey of a water mass slowly mixed deeper and deeper and whose DIC and TA are progressively enriched by all chemical reactions they undergo. Accordingly, this

interpretation does not identify reactions with minor impact on the carbonate cycle (equation 7) or reactions that cancelled each other later during the journey. For example, Δ TAex introduced by PP is frequently cancelled by a similar amount or excess of Δ TAex by AR. The starting point corresponds to the upstream (S=1.5) endmember at DICex=0; TAex=0. Figure 4a shows a slight DICex enrichment at TAex=0 which reflects a bias from the endmember calculation.

295

4.2 Identification of major reactions in oxygenated water column using $\Delta TAex/\Delta DICex/\Delta AOU$ signature.

Table 1: Summary of the main net reactions occurring among the different zones of a redox gradient with reactants starting in equilibrium with the atmosphere, adapted from Soetaert et al. (2007). The calculations assume Redfield ratio of the organic matter, i. e. $\gamma N = 0.156$ and $\gamma P = 0.0094$ (Soetaert et al., 2007). Redox zones describe the different conditions required for completing the reaction.

Name	Redox zones	Net Formula	ΔTA	ΔDIC
AR	Oxic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + O_2 \Rightarrow$ $\gamma^N NH_3 + \gamma^P H_3PO_4 + CO_2 + H_2O$		+1
CD	Any	$CaCO_3 + CO_2 \rightarrow Ca^{2+} + 2 HCO_3^{-}$	2	+1
Nit	Oxic	$NH_3 + 2 O_2 \rightarrow HNO_3 + H_2O$	-2	0
NR-N2	Anoxic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 0.8 \text{ HNO}_3 \Rightarrow \gamma^N NH_3 + \gamma^P H_3PO_4 + CO_2 + 0.4 N_2 + 1.4 H_2O$	0.95	+1
NR-NH3	Anoxic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 0.5 \text{ HNO}_3 \Rightarrow \gamma^N \text{ NH}_3 + \gamma^P H_3PO_4 + CO_2 + 0.5 \text{ NH}_3 + 0.5 \text{ H}_2O$	1.15	+1
MnR-MnC	Anoxic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 2 MnO_2 + CO_2 \Rightarrow \gamma^N NH_3 + \gamma^P H_3PO_4 + 2 MnCO_3 + H_2O$	0.15	-1
SR	Sulfidic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 0.5 H_2SO_4 \rightarrow \gamma^N NH_3 + \gamma^P H_3PO_4 + CO_2 + 0.5 H_2S + H_2O$	1.15	+1
SR-SMnC	Sulfidic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 1/2 H_2SO_4 + 0.5 MnO_2 \Rightarrow$ $\gamma^N NH_3 + \gamma^P H_3PO_4 + 1/2 CO_2 + 1/2 MnCO_3 + 0.5 S^0 + 1.5 H_2O$	1.15	+0.5
FeSR-FeS	Sulfidic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 0.44 \text{ FeOOH} + 0.44 H_2SO_4 \Rightarrow \gamma^N NH_3 + \gamma^P H_3PO_4 + CO_2 + 0.44 \text{ FeS} + 1.67 H_2O$	1.04	+1
SR-SFeS	Sulfidic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 0.33 \text{ FeOOH} + 0.5 H_2SO_4 \Rightarrow$ $\gamma^N NH_3 + \gamma^P H_3PO_4 + CO_2 + 0.33 \text{ FeS} + 0.17 \text{ S}^0 + 1.67 H_2O$	1.15	+1
SR-FeS ₂	Sulfidic	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 0.27 \text{ FeOOH} + 0.53 H_2SO_4 \Rightarrow \gamma^N NH_3 + \gamma^P H_3PO_4 + CO_2 + 0.27 \text{ FeS}_2 + 1.67 H_2O$	1.21	+1

In oxygenated water, the $\Delta TAex/\Delta DICex$ analysis can be supported and strengthened by looking also at the $\Delta AOU/\Delta DICex$ ratio (Fig. 4b) and establishing a $\Delta TAex/\Delta DICex/\Delta AOU$ signature for each water layer. Then, a linear combination of up to 3 reactions can be fit to the observed $\Delta TAex/\Delta DICex/\Delta AOU$ signature. The main candidate reactions to be combined are:

- aerobic respiration (AR), primary production (whose overall mass balance equation is here summarized as negative AR), carbonate dissolution (CD) and nitrification of ammonium (Nit) as presented in Table 1 and in Fig. 4c. In 2017, the whole oxygenated zone, including the PP zone and the ILO zone, is characterised by a $\Delta TAex/\Delta DICex/\Delta AOU = 0.1/1/1$ (Fig. 4a and 4b) which corresponds to the occurrence of only net aerobic respiration (AR) (theoretical values are 0.15/1/1 see Table 1 and Fig. 4c). Note a - that "net aerobic respiration" indicates that primary production is possible at a significant rate, but slower
- 305 than AR; and b that this $\Delta TAex/\Delta DICex/\Delta AOU$ signature indicates weak nitrification following respiration. In case of full nitrification, the theoretical slopes should be $\Delta TAex/\Delta DICex/\Delta AOU = -0.167/1/1.31$ as proposed by Zeebe and Wolf-Gladrow (2001). A combination of 1 AR and 0.025 Nit (nitrification of 16% of the produced NH₃) improves the modelled value to 0.1/1/1.05. The weakness of nitrification can be explained by slow kinetics of NH₄⁺ oxidation, with a half-life time estimated between a few days in estuaries (Horrigan et al., 1990) to multiple years in coastal environments (Heiss and Fulweiler, 2016).
- 310 As a comparison, other NH_4^+ fates, such as adsorption leads to an ammonium half life time of about a few minutes (Alshameri et al., 2018) to a few hours (Raaphorst and Malschaert, 1996) depending on the concentration of fine particles. Additionally, algae are known to use NH_4^+ as a N source (Raven et al., 1992) and NH_4^+ can be directly assimilated by heterotrophic organisms.
- In 2018, fresh water masses brought by the exceptional flood modified the carbonate system equilibrium. First, a low 315 salinity layer with pCO₂ at 1540 µatm overlays the primary production layer (Fig. 2), preventing the uptake of atmospheric CO₂ by primary production as was observed in 2016 (Chen et al., 2020). Just below the surface layer, in the PP zone, the lock down of atmospheric exchange by the low salinity layer produces supersaturation of trapped O₂ (Fig. 2). In Fig. 4a and 4b, this process translates into a vertical distribution at DICex = 40 µM with $\Delta TAex/\Delta DICex/\Delta AOU = 1.37/0/-1$. This original signature can be modelled by the combination of simultaneous carbonate dissolution (CD), the water column being
- 320 undersaturated, and PP, in equal proportion (2nd line in Table 2); the carbonate dissolution buffers the DIC consumption by the PP. Note that the ratio between $\Delta TAex/\Delta AOU$ implicates significant nitrification.

In 2018, below the PP zone down to DICex = 240μ M, the beginning of the ILO zone, the TAex increases significantly with a Δ TAex/ Δ DICex/ Δ AOU = 0.67/1/0.67 (Fig. 4a and 4b), incompatible with AR. To explain this signature a contribution of CD superimposed on AR seems most likely. A linear combination fitting leads 0.5 CD for 1 AR (3rd line in Table 2) and

- 325 results in $\Delta TAex/\Delta DICex/\Delta AOU = 0.77/1/0.67$. An explanation of its occurrence solely in 2018 could be the increase in carbonate rich suspended material at high flow conditions (Su et al., 2021). Excess of Ca²⁺ compared to the mixing line with oceanic end member (not shown) indicates that up to 200 µM of Ca²⁺ is produced in the oxygenated layer. Deeper, in the ILO zone, $\Delta TAex/\Delta DICex/\Delta AOU = 0.2/1/1.25$ (Fig. 4a and 4b), which results mainly from AR (0.15/1/1) with possible addition of CD and Nit, the exact signature being fitted for 0.54 CD and 0.46 Nit for 1 AR (4th line in Table 2), as a continuum of AR
- 330 and CD relative rates from the overlaying layer. This important nitrification is also in good agreement with the nitrification observed in the PP zone during this campaign, the lack of nitrite build up in the ILO zone and the relatively high oxygen concentration (at 105 μM) in the ILO zone in 2018 which is able to sustain nitrification. Overall, our results show that the higher river flow of 2018 increases carbonate dissolution for the top 5-10 m water depth that is superimposed on primary

production or aerobic respiration. Additionally, in 2017 the $\Delta TAex/\Delta DICex/\Delta AOU$ system indicates weak nitrification, while

335 in 2018 significant nitrification in the ILO and PP zones are suggested by the "reaction driven" approximation. The role of nitrification in explaining TAex depletion is only hypothetical since no direct measurement of NH_4^+ and NO_3^- were not performed. In particular, TAex depletion is particularly intense during high flow, high suspended particles season and could be produced by NH₄⁺ adsorption to the particles rather than by nitrification.

4.3 Identification of major reactions in the anoxic water column using $\Delta TAex/\Delta DICex/\Delta H_2S$ signature.

- 340 In the absence of oxygen, the $\Delta TAex/\Delta DICex/\Delta H_2S$ will be used for rates calculation. In the anoxic zone, the signature is similar in summer 2017 and 2018 at $\Delta TAex/\Delta DICex/\Delta H_2S = 2.4/1/0$. The "reaction driven" approximation of this signature is more difficult than in oxidized water because more reactions are known to occur simultaneously in absence of oxygen. However, to fit with the "reaction driven" approximation, it is not necessary to describe each reaction step but only the overall changes concerning the journey of a water mass over different redox conditions resulting in a combined result or reaction. This 345 approach has recently been proposed for FeS burial by Hiscock and Millero (2006), Rassmann et al. (2020) or Su et al. (2020b). A scenario combining sulfate reduction (SR) is particularly attractive since SR represents the main carbon remineralisation pathway in absence of oxygen. However, a combination of SR with CD would result in a $\Delta TAex/\Delta DICex$ between 1.15 and 2
- (see Table 1 or Fig. 4c) and fails to reach the $\Delta TAex/\Delta DICex$ of 2.4. Moreover, SR alone underestimates the importance of the H₂S oxidation pathway that can consume all the alkalinity produced during SR. For example, SR follow by oxygenated 350 oxidation results in $\Delta TAex/\Delta DICex/\Delta AOU$ signature equal to AR only. In the Chesapeake Bay, H₂S oxidation is critical since no H_2S is measurable in the anoxic zone while the gradient at the sediment/water interface indicates high H_2S sedimentary efflux (Fig. 2).

Generalizing these observations, recent efforts to build an alkalinity budget on the global scale (Hu and Cai, 2011; Middelburg et al., 2020) highlight that the alkalinity produced by anaerobic respiration corresponds to the uncharged species produced, mostly in solid or gaseous phases. Indeed, the alkalinity changes produced during a natural reaction equal the "charge transfer" from species having some charge at pH = 4.5, such as NO₃⁻ and SO₄²⁻, to species that would lose its charges at pH =4.5, mainly HCO₃, that is not counted in the alkalinity calculation (see Eq. (12)). Although correct, this approach tends to

neglect the roles of Fe and Mn oxides (Middelburg et al., 2020) since their transformation from (oxyhydr)oxides into sulphur or carbonate solid species does not involve any charge transfer. When looking in detail at these processes, the metal oxides are

360

355

critical since they are the main H₂S oxidation pathway that does not regenerate H₂SO₄ but rather produces S⁰, S_n²⁻ or FeS (Findlay et al., 2014; Avetisyan et al., 2021) which limits alkalinity consumption.

To build a pool of candidate reactions for the fitting, first, dissolved species at too low a concentration (e.g. Mn^{2+}_{aq} , Fe^{2+}_{aq}) to be a net reagent to affect the carbon cycle at steady state are not taken into account. These species are usually recycled rapidly and hold a role of catalyser or electron shuttle between other redox species and did not reach 10 µM during the 365 campaigns (Fig. 3). Second, many minerals are expected to be at low concentration or thermodynamically not favoured, and their associated reactions are neglected (e.g., iron phosphate, ferrous or manganous oxide, metal sulphur clusters, MnS, FeCO₃,

adsorption processes, reverse weathering). Therefore, only aqueous species with important stock concentrations (that can exceed 0.1 mM in anoxic water) are taken into account, *i. e.*, SO_4^{2-} , Ca^{2+} , H_2S , NH_4^+ together with gaseous (N₂, CO₂) and main solid phases (FeS₂, FeS, S⁰, MnCO₃, FeOOH, MnO₂). Third, the combination of many carbon remineralisation reactions with re-oxidation reactions or the net result produces a net chemical equation equal to another remineralisation reaction. As an example, the chemical equation of SR followed by H₂S oxidation with oxygen is equal to the equation of aerobic respiration: the proposed model confounds both pathways because the resulting chemical changes are similar.

Table 1 lists the resulting combined reactions, and the calculated $\Delta TAex/\Delta DICex$ slopes are represented in Fig. 4c. For the anoxic zone, the nitrate respiration can be associated with N₂ production (NR-N2) or with NH₃ production (NR-NH₃), and the **Table 2: Linear combination of reactions from Table 1 that fit the observations (see text for details, H₂O molecules are omitted).**

			Linear combi	mation CH2O	ABLCE ACX ADU	ADICe ⁵ Net Formula (for 1 CH ₂ O)
			AR+0.025Nit	0.1	1.05	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + O_2 \Rightarrow$ 0.16 γ^N HNO3+ 0.84 γ^N NH ₃ + γ^P H ₃ PO ₄ + CO ₂
Observed in 2017 Observed in 2018			CD-AR+0.65Nit	x	-∞	$(0.65+\gamma^{N}) \text{ NH}_{3} + \gamma^{P} \text{ H}_{3}\text{PO}_{4} + \text{CaCO}_{3} + 2 \text{ CO}_{2} \Rightarrow$ $(\text{CH}_{2}\text{O})(\text{NH}_{3})_{\gamma N}(\text{H}_{3}\text{PO}_{4})_{\gamma P} + \text{O}_{2} + \text{Ca}^{2+} + 2 \text{ HCO}_{3}^{-} + 0.65 \text{ HNO}_{3}$
	in 2018	Oxic	AR+0.5CD	0.77	0.67	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + O_2 + 0.5 CaCO_3 \rightarrow \gamma^N NH_3 + \gamma^P H_3PO_4 + 0.5 CO_2 + 0.5 Ca^{2+} + HCO_3^{-1}$
	Dbserved		AR + 0.54 CD + 0.46 Nit	0.2	1.25	$(CH_{2}O)(NH_{3})_{\gamma N}(H_{3}PO_{4})_{\gamma P} + 1.93 O_{2} + 0.54 CaCO_{3} + 2\gamma_{N} NH_{3} \rightarrow 3\gamma^{N} HNO_{3} + \gamma^{P} H_{3}PO_{4} + 0.46 CO_{2} + 0.54 Ca^{2+} + 1.1 HCO_{3}^{-}$
		Anoxic	0.98 SR-SMnC + 0.02 MnR-MnC	2.4	0	$(CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 0.49 H_2SO_4 + 0.53 MnO_2 \Rightarrow$ $\gamma^N NH_3 + \gamma^P H_3PO_4 + 0.47 CO_2 + 0.53 MnCO_3 + 0.49 S^0$
			0.65 SR-FeS + 0.35 MnR-MnC	2.4	0	$(CH_{2}O)(NH_{3})_{\gamma N}(H_{3}PO_{4})_{\gamma P} + 0.3 H_{2}SO_{4} + 0.7 MnO_{2} + 0.3 FeOOH \Rightarrow \gamma^{N} NH_{3} + \gamma^{P} H_{3}PO_{4} + 0.3 CO_{2} + 0.7 MnCO_{3} + 0.29 FeS$
			6.4 CD + MnR-MnC	2.4	0	$(CH_{2}O)(NH_{3})_{\gamma N}(H_{3}PO_{4})_{\gamma P} + 6.4 CaCO_{3} + 2 MnO_{2} + 7.4 CO_{2} \Rightarrow \gamma^{N} NH_{3} + \gamma^{P} H_{3}PO_{4} + 12.8 HCO_{3} + 2 MnCO_{3} + 6.4 Ca^{2+}$
		dic	0.38 MnR-MnC + 0.76 SR - 0.15 SR-SMnC	2.4	1.2	$\begin{array}{c} (CH_2O)(NH_3)_{\gamma N}(H_3PO_4)_{\gamma P} + 0.31 \ H_2SO_4 + 0.68 \ MnO_2 + 0.07 \ S^0 \clubsuit \\ \gamma^N \ NH_3 + \gamma^P \ H_3PO_4 + 0.32 \ CO_2 + 0.68 \ MnCO_3 + 0.38 \ H_2S \end{array}$
		Sulfi	0.64 MnR-MnC + 1.36 SR - SR-SMnC	2.4	3.2	$(CH_{2}O)(NH_{3})_{\gamma N}(H_{3}PO_{4})_{\gamma P} + 0.18 H_{2}SO_{4} + 0.79 MnO_{2} + 0.5 S^{0} \rightarrow \gamma^{N} NH_{3} + \gamma^{P} H_{3}PO_{4} + 0.21 CO_{2} + 0.79 MnCO_{3} + 0.68 H_{2}S$

- 375 manganese oxides respiration produces carbonate precipitation (MnR-MnC). For the sulfidic zone, SR can occur alone producing a build-up of H₂S. However, at a certain point, H₂S gets significantly oxidised either by MnO₂ which produces MnCO₃ and S₀ (SR-SMnC) or by FeOOH, producing FeS and S₀ (SR-SFeS) and ultimately FeS₂ (SR-FeS2). Direct respiration of FeOOH is also taken into account, but as the only final Fe product in the model is FeS or FeS₂, it has to be accompanied by some SR (FeSR-FeS).
- 380 Figure 4c demonstrates that the slope of $\Delta TAex/\Delta DICex = 2.4$ can be obtained for any reaction in combination with MnR-MnC. Combinations without MnR-MnC, however, lead to a negative SR whose overall equation could be interpreted as a possible small participation of anoxygenic phototrophic (purple) bacteria (Findlay et al., 2015, 2017) but are not considered further as the amount of $\Delta TAex$ involved would be tiny. Therefore, in the absence of nitrate, oxygen and H₂S, only a combination of MnR-MnC with SR-SMnC (producing S⁰, 5th line in Table 2), SR-FeS (producing FeS, 6th line in Table 2) or
- 385 CD (releasing Ca²⁺, 7th line in Table 2) gives the particularly high Δ TAex/ Δ DICex of 2.4. S₀ was not measured during our campaign, but it has been previously reported at this site (Findlay et al., 2014), and the S₀ produced by SR-SMnC can react with FeS to form FeS₂. All the three identified combinations require a critical role of MnO₂. Since Figure 3 does not indicate any clear reaction for iron while the steep gradient of MnOx in proximity to the H₂S rich layer suggests reaction between MnOx and H₂S; the combination with SR-SMnC is the most likely (5th line in Table 2). Deeper, the vertical gradient of sulphide
- 390 suggests that part of the H₂S came by diffusion from the sediment's porewater (Fig. 2). The assumptions required for the "reaction driven" approximation are still valid as soon as steady state is maintained by ongoing reactions, even if one of the local endmembers has not been sampled since it is probably located in the sediment. In the presence of sulphide, the $\Delta TAex/\Delta DICex/\Delta H_2S$ signature is 2.4/1/1.2 in 2017 and 2.4/1/3.2 in 2018 (Fig. 4a and 4d) and can be explained by the same combination of reactions without complete oxidation of H₂S from SR to take into account the build-up of H₂S (Table 2).

4.4 Comparison with other studies

- The high observed ratio of ΔTAex/ΔDICex = 2.4 seems very specific to the Chesapeake Bay. Moreover, the "reaction driven" interpretation can be applied to other published datasets for which the ΔTA/ΔDIC/ΔH₂S system can be calculated (Table 3). In the water column, most of the available datasets are not suitable for the "reaction driven" approximation since either they focus on surface water where DIC and TA are strongly impacted by atmospheric exchange or the water masses change too fast
 to consider that reactions dominate over water mixing. However, porewater measured in the Gulf of Mexico has ΔTAex/ΔDICex/ΔH₂S = 1.15/1/0.53 (Hu et al., 2010) as expected when sulfate reduction is associated with H₂S accumulation (SR reaction; ΔTAex/ΔDICex/ΔH₂S = 1.15/1/0.5; Table 1). Similarly, Hiscock and Millero (2006) report ΔTAex/ΔDICex/ΔH₂S = 1.3/1/0.5 in the Western Black Sea close to the SR signature. In the Baltic Sea sediment, ΔTAex/ΔDICex/ΔH₂S = 1.3/1/0.07 was reported (Lukawska-Matuszewska, 2016), which is close to the expected signature in the case of important H₂S consumption by Fe oxides and consequent precipitation as pyrite (SR-FeS2 reaction; ΔTAex/ΔDICex/ΔH₂S = 1.2/1/0). In Rhone river prodelta sediments, the reported ΔTAex/ΔDICex/ΔH₂S is 1/1/0 that can be
 - related to the 1/1/0 signature of FeSR-FeS (Table 1) that is expected in iron rich sediments with high sedimentation rates

preventing FeS₂ formation in the pore water (Rassmann et al., 2020). In permeable, carbonate rich sediments, the reported signature of $\Delta TAex/\Delta DICex/\Delta H_2S$ in Hawaii sands is 0.86/1/ND (Drupp et al., 2016). The lack of salinity and oxygen datasets

Publication	Sample type	a $\Delta TA/\Delta DIC/ \Delta H2S$ or b $\Delta TA/\Delta DIC/ \Delta AOU$	ΔΤΑ	Reaction driven interpretation
Hu et al., 2010	Gulf of Mexico sediment (slope)	1.15/1/0.53 a	+ 15 mM	SR
Lukawska-Matuszewska, 2016	Baltic Sea sediment	1.3/1/0.07 a	+ 13 mM	SR-FeS2
Rassmann et al., 2020	Rhone prodelta sediment	1/1/0 a	+ 5.6 mM	FeSR-FeS
Cai et al. 1998	Satilla estuary	2/1/ND	+ 0.2 mM	CD
Drupp et al. 2016	Oxygenated Hawai carbonate reef sands	0.86-0.91/1/ND	+ 1.5 mM	Not applicable*
Su et al. 2021	Chesapeake Bay water column	2/1/1.5 b	+0.4 mM	Not applicable**
		0.2/1/1 b	+0.05 mM	AR
		0.8/1/ND	+0.1 mM	NR-N2 or CD+Nit
Hiscock and Millero 2006	Western Black Sea water column	1.3/1/0.5 a	+1.2 mM	SR+?CD

Table 3: Overview of the $\Delta TA/\Delta DIC/\Delta H2S$ signature observed in different environments

*Lack of data on oxygen concentration prevent any interpretation. ** Important air-water exchange prevent any "reaction driven" approximation.

- 410 prevents further model fits, but the $\Delta TAex/\Delta DICex$ is below 2.4. For oxygen depleted data from the whole Chesapeake Bay described in Su et al. (2020), the signatures are $\Delta TAex/\Delta DICex/\Delta AOU = 0.2/1/1$ in presence of oxygen (typical signature of AR) and 0.8/1/ND in absence of oxygen that could correspond to NR-N2 or a combination of CD+Nit. Overall, this bibliographic survey highlights the effectiveness of the "reaction driven" approximation to identify preponderant reactions controlling the carbon cycle and puts in perspective the originality of the $\Delta TAex/\Delta DICex/\Delta AOU$ signature of 2.4/1/0 observed
- 415 in the Chesapeake Bay.

4.5 Local and global budget

While the "reaction driven" approximation indicates a dominant role of the SR-SMnC reaction; this possibility needs to be validated looking at the saturation state of rhodocrosite (the main MnCO₃ mineral) and looking at the mass budget between MnOx consumed and TAex produced. The rhodocrosite saturation (Luo and Millero, 2003) is always below 0.3 in our samples,
which stands against the occurrence of *in situ* SR-SMnC reaction. When inspecting the mass budget, the 88 µM to 155 µM of MnO₂ required to produce the 100 µM TAex increase observed (Fig. 4a) is one order of magnitude higher than the observed MnOx or Mn²⁺ concentration (Fig. 3). This mass budget discrepancy cannot be solved invoking suspended material since the 88 µM of MnO₂ would require a suspended material concentration of about 4.4 g L⁻¹ (assuming an average concentration of 20 µmol g⁻¹ of Mn), which is again one or two orders of magnitude higher than the discrepancy between water and solid concentration. But another process dephasing aqueous from solid reaction products is also possible at station 858.

since the dissolved phase could have moved up, rather than the particles settling down. In this case, the SR-SMnC reaction

was not happening only in the water column of the Chesapeake Bay and part of the TAex and DICex pool could have been produced in the sediment during the previous year, then diffused out of the sediment simultaneously with other reduced

- 430 elements as the summer begins. Indeed, many previous studies at station 858 (*e.g.*, Sholkovitz et al., 1992) explained the seasonality of anoxia with an upward move of the redox front from the sediments to bottom waters during the start of summer. Important sedimentary efflux of H₂S, Fe_{aq} and Mn_{aq} were still visible during both of our August campaigns. Therefore, the 100 μ M TA increase does not fit with the ambient Mn²⁺ or MnO₂ in the water column but rather with the MnCO₃ deposited in the sediment. The sedimentary solid Mn stock of the Chesapeake Bay is particularly important, up to 70 μ mol g⁻¹ at station 858
- 435 (Sinex and Helz, 1981) compared to an average value of 15 µmol g⁻¹ for the upper continental crust (Rudnick and Gao, 2003). Indeed semi-enclosed basins are known to concentrate manganese at the deeper sediment (Thamdrup and Dalsgaard, 2000; Lenstra et al., 2020). Recent investigations at a close station (ET 5.1; 38°48.36'N; 75°54.66'W) in the Chesapeake Bay (Lenstra et al., 2021) report that about 60% of the surface sedimentary Mn pool is MnOx (acid ascorbic extractable) and 25 % is Mn carbonate (1M HCl extraction). Assuming a porosity of 0.8, a bulk solid density of 2.6, the sedimentary pool corresponds to
- 440 35 mM of manganese which largely exceeds the 88 μ M required to produce the 100 μ M TAex increase. Therefore, the Chesapeake Bay sediment is particularly rich in manganese and could host important SR-SMnC reactions in the surface pore water whose soluble products diffuse up to the water column during summer and could bear with them the high Δ TAex/ Δ DICex signature observed.
- The assumptions detailed in section 4.1 permit the "reaction driven" approach to be reconsidered in regard to this sediment efflux scenario. Indeed, the sediment efflux does not need to be considered as an additional endmember, that would violate the third assumption, since its salinity and the pore water concentrations results from the upstream and oceanic endmembers superimposed to geochemical reactions. In section 2.3.1, we point out that equation (4) was valid in between each local endmembers and that the straight lines on the $\Delta TAex/\Delta DICex$ plot between them indicates that the local endmembers are maintained in steady state by a chemical reaction with a similar stoichiometry. These results indicate that even if most of the
- 450 MnCO₃ was produced when the local endmembers were localised in the sediment, a- their migration does not alter the TAex/DICex signature and b- the chemical reaction that produced them is still ongoing at sufficient rate to maintain a steady state characterised by the steep changes of direction observed in Fig. 4a.

Conclusion

The "reaction driven" approximation proposed in this study is a powerful interpretative framework that can identify the

- 455 major reactions controlling the carbonate cycle. In the Chesapeake Bay, similar redox stratification can support varying intensity of carbonate dissolution, absent in 2017 or important as in 2018. In 2018, the "reaction driven" approximation also suggests an important role of the N cycle to consume alkalinity while, in 2017, nitrification was limited to 16% of the nitrogen mineralized. The summer anoxia observed in the Chesapeake Bay is characterized by an exceptionally high ΔTAex/ΔDICex of 2.4 which has never been reported in anoxic water columns or sediment pore waters. The "reaction driven" approximation
- suggests it comes from sulphate reduction with almost complete hydrogen sulphide oxidation by MnOx followed by MnCO₃ precipitation. This interpretation is supported by the important manganese dynamics observed in this and previous papers (Oldham et al, 2017a; Sholkovitz et al 1992; Trouwborst et al, 2006). However, most of the reactions would have occurred in the sediment from which the components of the ΔTAex/ΔDICex of 2.4 diffused into the bottom water with the redox front during the set up of summer water column anoxia. Our study demonstrates that the Manganese cycle can have a strong impact on alkalinity, as it prevents H₂S oxidation to SO₄²⁻ and favour sulphur burial.

Appendix



Figure A1: Variation of Total Alkalinity (TA) during oceanic and river mixing. Dashed lines represent the theoretical DIC and TA if only mixing occurs.



Figure A2: Superimposed carbonate and redox chemistry profiles over 11 casts against depth.



Figure A3: Examples of redox and metal chemistry profiles. Note that $Fe(III)_s$ and $Fe(III)_{aq}$ scales are zoomed in by 3 and O₂ scale is zoomed out by 100.

Data availability

470 The data used in this paper is available on request to the corresponding author.

Author contribution

ATC, ERE, JN, BMT and SJ performed the data analysis. ATC and SJ process the data. ATC, GWL, SJ and WJC interpreted the results. GWL, BMT and WJC get the funding. ATC wrote the paper with contributions from all authors.

Competing interests

475 The authors declare that they have no conflict of interest.

Acknowledgements

We gratefully acknowledge the support of the captain and crew of *R/V Hugh R. Sharp*. This work was funded by grants from the Chemical Oceanography program of the National Science Foundation (OCE-1558738 to GWL; OCE-1558692 to BMT and OCE-1756815 to WJC)

480 **References**

Abril, G., Etcheber, H., Delille, B., Frankignoulle, M., and Borges, A. V.: Carbonate dissolution in the turbid and eutrophic Loire estuary, Mar. Ecol. Prog. Ser., 259, 129–138, 2003.

Alshameri, A., He, H., Zhu, J., Xi, Y., Zhu, R., Ma, L., and Tao, Q.: Adsorption of ammonium by different natural clay minerals: Characterization, kinetics and adsorption isotherms, Appl. Clay Sci., 159, 83–93, https://doi.org/10.1016/j.clay.2017.11.007, 2018.

Avetisyan, K., Zweig, I., Luther, G. W., and Kamyshny, A.: Kinetics and mechanism of polysulfides and elemental sulfur formation by a reaction between hydrogen sulfide and δ -MnO2, Geochim. Cosmochim. Acta, 313, 21–37, https://doi.org/10.1016/j.gca.2021.08.022, 2021.

Borges, A. V., Schiettecatte, L.-S., Abril, G., Delille, B., and Gazeau, F.: Carbon dioxide in European coastal waters, Estuar. Coast. Shelf Sci., 70, 375–387, https://doi.org/10.1016/j.ecss.2006.05.046, 2006.

Borges, A. V., Abril, G., and Bouillon, S.: Carbon dynamics and CO2 and CH4 outgassing in the Mekong delta, Biogeosciences, 15, 1093–1114, https://doi.org/10.5194/bg-15-1093-2018, 2018.

Boudreau, B. P., Middelburg, J. J., and Luo, Y.: The role of calcification in carbonate compensation, Nat. Geosci., 11, 894–900, https://doi.org/10.1038/s41561-018-0259-5, 2018.

- 495 Breitburg, D., Levin, L. A., Oschlies, A., Grégoire, M., Chavez, F. P., Conley, D. J., Garçon, V., Gilbert, D., Gutiérrez, D., Isensee, K., Jacinto, G. S., Limburg, K. E., Montes, I., Naqvi, S. W. A., Pitcher, G. C., Rabalais, N. N., Roman, M. R., Rose, K. A., Seibel, B. A., Telszewski, M., Yasuhara, M., and Zhang, J.: Declining oxygen in the global ocean and coastal waters, Science, 359, eaam7240, https://doi.org/10.1126/science.aam7240, 2018.
- 500 Cai, W.-J., Huang, W.-J., Luther, G. W., Pierrot, D., Li, M., Testa, J., Xue, M., Joesoef, A., Mann, R., Brodeur, J., Xu, Y.-Y., Chen, B., Hussain, N., Waldbusser, G. G., Cornwell, J., and Kemp, W. M.: Redox reactions and weak buffering capacity lead to acidification in the Chesapeake Bay, Nat. Commun., 8, 369, https://doi.org/10.1038/s41467-017-00417-7, 2017.

Cai, W.-J., Xu, Y.-Y., Feely, R. A., Wanninkhof, R., Jönsson, B., Alin, S. R., Barbero, L., Cross, J. N.,
Azetsu-Scott, K., Fassbender, A. J., Carter, B. R., Jiang, L.-Q., Pepin, P., Chen, B., Hussain, N., Reimer,
J. J., Xue, L., Salisbury, J. E., Hernández-Ayón, J. M., Langdon, C., Li, Q., Sutton, A. J., Chen, C.-T. A.,
and Gledhill, D. K.: Controls on surface water carbonate chemistry along North American ocean margins,
Nat. Commun., 11, https://doi.org/10.1038/s41467-020-16530-z, 2020.

Cerco, C. F., Kim, S.-C., and Noel, M. R.: Management modeling of suspended solids in the Chesapeake 510 Bay, USA, Estuar. Coast. Shelf Sci., 116, 87–98, https://doi.org/10.1016/j.ecss.2012.07.009, 2013.

Chen, B., Cai, W.-J., Brodeur, J. R., Hussain, N., Testa, J. M., Ni, W., and Li, Q.: Seasonal and spatial variability in surface pCO2 and air–water CO2 flux in the Chesapeake Bay, Limnol. Oceanogr., 65, 3046–3065, https://doi.org/10.1002/lno.11573, 2020.

Clement, B. G., Luther, G. W., and Tebo, B. M.: Rapid, oxygen-dependent microbial Mn(II) oxidation
 kinetics at sub-micromolar oxygen concentrations in the Black Sea suboxic zone, Geochim. Cosmochim.
 Acta, 73, 1878–1889, https://doi.org/10.1016/j.gca.2008.12.023, 2009.

Cotovicz Jr., L. C., Libardoni, B. G., Brandini, N., Knoppers, B. A., and Abril, G.: Comparisons between real-Time PCO2 measurements with indirect estimates in two contrasting Brazilian estuaries: The eutrophic guanabara bay (RJ) and the oligotrophic sao francisco River estuary (AL), Quím. Nova, 39, 1206–1214, https://doi.org/10.21577/0100-4042.20160145, 2016.

Dickson, A. G.: An exact definition of total alkalinity and a procedure for the estimation of alkalinity and total inorganic carbon from titration data, Deep Sea Res. Part Oceanogr. Res. Pap., 28, 609–623, https://doi.org/10.1016/0198-0149(81)90121-7, 1981.

Drupp, P. S., De Carlo, E. H., and Mackenzie, F. T.: Porewater CO2–carbonic acid system chemistry in permeable carbonate reef sands, Mar. Chem., 185, 48–64, https://doi.org/10.1016/j.marchem.2016.04.004, 2016.

Du, J. and Shen, J.: Water residence time in Chesapeake Bay for 1980–2012, J. Mar. Syst., 164, 101–111, https://doi.org/10.1016/j.jmarsys.2016.08.011, 2016.

Findlay, A. J., Gartman, A., MacDonald, D. J., Hanson, T. E., Shaw, T. J., and Luther, G. W.: Distribution
and size fractionation of elemental sulfur in aqueous environments: The Chesapeake Bay and Mid-Atlantic Ridge, Geochim. Cosmochim. Acta, 142, 334–348, 2014.

Findlay, A. J., Bennett, A. J., Hanson, T. E., and Luther, G. W.: Light-Dependent Sulfide Oxidation in the Anoxic Zone of the Chesapeake Bay Can Be Explained by Small Populations of Phototrophic Bacteria, Appl. Environ. Microbiol., 81, 7560–7569, https://doi.org/10.1128/AEM.02062-15, 2015.

535 Findlay, A. J., Di Toro, D. M., and Luther, G. W.: A model of phototrophic sulfide oxidation in a stratified estuary, Limnol. Oceanogr., 62, 1853–1867, https://doi.org/10.1002/lno.10539, 2017.

Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B., Anthoni, P., Barbero, L., Bastos, A., Bastrikov, V., Becker, M., Bopp, L., Buitenhuis, E., Chandra, N., Chevallier,

- 540 F., Chini, L. P., Currie, K. I., Feely, R. A., Gehlen, M., Gilfillan, D., Gkritzalis, T., Goll, D. S., Gruber, N., Gutekunst, S., Harris, I., Haverd, V., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Joetzjer, E., Kaplan, J. O., Kato, E., Klein Goldewijk, K., Korsbakken, J. I., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Marland, G., McGuire, P. C., Melton, J. R., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Neill, C., Omar, A. M., Ono, T., Peregon, A., Pierrot,
- D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Séférian, R., Schwinger, J., 545 Smith, N., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Werf, G. R., Wiltshire, A. J., and S.: Global Carbon Budget 2019. Earth Syst. Sci. Data, 11. 1783-1838. Zaehle, https://doi.org/10.5194/essd-11-1783-2019, 2019.

Grasshoff, K.: Determination of nitrite, nitrate, oxygen, thiosulphate, in: Methods of seawater analysis, New York, 61–72, 1983.

Gupta, G. V. M., Jyothibabu, R., Ramu, C. V., Reddy, A. Y., Balachandran, K. K., Sudheesh, V., Kumar, S., Chari, N. V. H. K., Bepari, K. F., Marathe, P. H., Reddy, B. B., and Vijayan, A. K.: The world's largest coastal deoxygenation zone is not anthropogenically driven, Environ. Res. Lett., 16, 054009, https://doi.org/10.1088/1748-9326/abe9eb, 2021.

555 Gustafsson, E., Hagens, M., Sun, X., Reed, D. C., Humborg, C., Slomp, C. P., and Gustafsson, B. G.: Sedimentary alkalinity generation and long-term alkalinity development in the Baltic Sea, Biogeosciences, 16, 437–456, https://doi.org/10.5194/bg-16-437-2019, 2019.

Hagy, J. D., Boynton, W. R., Keefe, C. W., and Wood, K. V.: Hypoxia in Chesapeake Bay, 1950-2001: Long-Term Change in Relation to Nutrient Loading and River Flow, Estuaries, 27, 634–658, 2004.

⁵⁶⁰ Heiss, E. M. and Fulweiler, R. W.: Coastal water column ammonium and nitrite oxidation are decoupled in summer, Estuar. Coast. Shelf Sci., 178, 110–119, https://doi.org/10.1016/j.ecss.2016.06.002, 2016.

Hiscock, W. T. and Millero, F. J.: Alkalinity of the anoxic waters in the Western Black Sea, Deep Sea Res. Part II Top. Stud. Oceanogr., 53, 1787–1801, https://doi.org/10.1016/j.dsr2.2006.05.020, 2006.

Horrigan, S. G., Montoya, J. P., Nevins, J. L., McCarthy, J. J., Ducklow, H., Goericke, R., and Malone,
T.: Nitrogenous nutrient transformations in the spring and fall in the Chesapeake Bay, Estuar. Coast. Shelf
Sci., 30, 369–391, https://doi.org/10.1016/0272-7714(90)90004-B, 1990.

Hu, X. and Cai, W.-J.: An assessment of ocean margin anaerobic processes on oceanic alkalinity budget, Glob. Biogeochem. Cycles, 25, https://doi.org/10.1029/2010GB003859, 2011.

Hu, X., Cai, W.-J., Wang, Y., Luo, S., and Guo, X.: Pore-water geochemistry of two contrasting brinecharged seep sites in the northern Gulf of Mexico continental slope, Mar. Chem., 118, 99–107, https://doi.org/10.1016/j.marchem.2009.11.006, 2010.

Huang, W.-J., Wang, Y., and Cai, W.-J.: Assessment of sample storage techniques for total alkalinity and dissolved inorganic carbon in seawater, Limnol. Oceanogr. Methods, 10, 711–717, https://doi.org/10.4319/lom.2012.10.711, 2012.

575 Hudson, J. M., MacDonald, D. J., Estes, E. R., and Luther, G. W.: A durable and inexpensive pump profiler to monitor stratified water columns with high vertical resolution, Talanta, 199, 415–424, https://doi.org/10.1016/j.talanta.2019.02.076, 2019.

Hulot, V., Metzger, E., Thibault De Chanvalon, A., Mouret, A., Schmidt, S., Deflandre, B., Rigaud, S., Beneteau, E., Savoye, N., Souchu, P., Le Merrer, Y., and Maillet, G. M.: Impact of an exceptional winter

⁵⁸⁰ flood on benthic oxygen and nutrient fluxes in a temperate macrotidal estuary: Potential consequences on summer deoxygenation, Front. Mar. Sci., 10, https://doi.org/10.3389/fmars.2023.1083377, 2023.

Ishii, H., Koh, H., and Satoh, K.: Spectrophotometric determination of manganese utilizing metal ion substitution in the cadmium- α , β ,- γ , δ -tetrakis (4-carboxyphenyl) porphine complex, Anal. Chim. Acta, 136, 347–352, 1982.

585 Joesoef, A., Kirchman, D. L., Sommerfield, C. K., and Cai, W.-J.: Seasonal variability of the inorganic carbon system in a large coastal plain estuary, Biogeosciences, 14, 4949–4963, https://doi.org/10.5194/bg-14-4949-2017, 2017.

Jones, M. R., Luther, G. W., Mucci, A., and Tebo, B. M.: Concentrations of reactive Mn(III)-L and MnO2 in estuarine and marine waters determined using spectrophotometry and the leuco base, leucoberbelin blue, Talanta, 200, 91–99, https://doi.org/10.1016/j.talanta.2019.03.026, 2019.

Lenstra, W. K., Séguret, M. J. M., Behrends, T., Groeneveld, R. K., Hermans, M., Witbaard, R., and Slomp, C. P.: Controls on the shuttling of manganese over the northwestern Black Sea shelf and its fate in the euxinic deep basin, Geochim. Cosmochim. Acta, 273, 177–204, https://doi.org/10.1016/j.gca.2020.01.031, 2020.

- 595 Lenstra, W. K., Klomp, R., Molema, F., Behrends, T., and Slomp, C. P.: A sequential extraction procedure for particulate manganese and its application to coastal marine sediments, Chem. Geol., 120538, https://doi.org/10.1016/j.chemgeo.2021.120538, 2021.
- Lewis, B. L., Glazer, B. T., Montbriand, P. J., Luther, G. W., Nuzzio, D. B., Deering, T., Ma, S., and Theberge, S.: Short-term and interannual variability of redox-sensitive chemical parameters in 600 hypoxic/anoxic bottom waters of the Chesapeake Bay, Mar. Chem., 105, 296–308, https://doi.org/10.1016/j.marchem.2007.03.001, 2007.

Lohrenz, S. E., Cai, W.-J., Chen, F., Chen, X., and Tuel, M.: Seasonal variability in air-sea fluxes of CO 2 in a river-influenced coastal margin, J. Geophys. Res., 115, 2010.

Lukawska-Matuszewska, K.: Contribution of non-carbonate inorganic and organic alkalinity to total measured alkalinity in pore waters in marine sediments (Gulf of Gdansk, S-E Baltic Sea), Mar. Chem., 186, 211–220, https://doi.org/10.1016/j.marchem.2016.10.002, 2016.

Luo, Y. and Millero, F. J.: Solubility of Rhodochrosite (MnCO3) in NaCl Solutions, J. Solut. Chem., 12, 2003.

Luther, G. W.: The role of one-and two-electron transfer reactions in forming thermodynamically unstable intermediates as barriers in multi-electron redox reactions, Aquat. Geochem., 16, 395–420, 2010.

Madison, A. S., Tebo, B. M., and Luther, G. W.: Simultaneous determination of soluble manganese(III), manganese(II) and total manganese in natural (pore)waters, Talanta, 84, 374–381, https://doi.org/10.1016/j.talanta.2011.01.025, 2011.

Meire, L., Soetaert, K. E. R., and Meysman, F. J. R.: Impact of global change on coastal oxygen dynamics and risk of hypoxia, Biogeosciences, 10, 2633–2653, https://doi.org/10.5194/bg-10-2633-2013, 2013.

Meybeck, M.: Global chemical weathering of surficial rocks estimated from river dissolved loads, Am. J. Sci., 287, 401–428, 1987.

Meybeck, M.: Global occurrence of major elements in rivers, Treatise Geochem., 5, 605, 2003.

Meybeck, M., Cauwet, G., Dessery, S., Somville, M., Gouleau, D., and Billen, G.: Nutrients (organic C,
P, N, Si) in the eutrophic River Loire (France) and its estuary, Estuar. Coast. Shelf Sci., 27, 595–624, https://doi.org/10.1016/0272-7714(88)90071-6, 1988.

Middelburg, J. J., Soetaert, K., and Hagens, M.: Ocean Alkalinity, Buffering and Biogeochemical Processes, Rev. Geophys., 58, e2019RG000681, https://doi.org/10.1029/2019RG000681, 2020.

Oldham, V. E., Jones, M. R., Tebo, B. M., and Luther, G. W.: Oxidative and reductive processes contributing to manganese cycling at oxic-anoxic interfaces, Mar. Chem., 195, 122–128, https://doi.org/10.1016/j.marchem.2017.06.002, 2017a.

Oldham, V. E., Miller, M. T., Jensen, L. T., and Luther, G. W.: Revisiting Mn and Fe removal in humic rich estuaries, Geochim. Cosmochim. Acta, 209, 267–283, https://doi.org/10.1016/j.gca.2017.04.001, 2017b.

630 Raaphorst, W. V. and Malschaert, J. F. P.: Ammonium adsorption in superficial North Sea sediments, Cont. Shelf Res., 16, 1415–1435, https://doi.org/10.1016/0278-4343(95)00081-X, 1996.

Rabalais, N. N., Cai, W.-J., Carstensen, J., Conley, D. J., Fry, B., Hu, X., Quiñones-Rivera, Z., Rosenberg, R., Slomp, C. P., Turner, R. E., Voss, M., Wissel, B., and Zhang, J.: Eutrophication-Driven Deoxygenation in the Coastal Ocean, Oceanography, 27, 172–183, 2014.

635 Rassmann, J., Eitel, E. M., Lansard, B., Cathalot, C., Brandily, C., Taillefert, M., and Rabouille, C.: Benthic alkalinity and dissolved inorganic carbon fluxes in the Rhône River prodelta generated by decoupled aerobic and anaerobic processes, Biogeosciences, 17, 13–33, https://doi.org/10.5194/bg-17-13-2020, 2020.

Raven, J. A., Wollenweber, B., and Handley, L. L.: A comparison of ammonium and nitrate as nitrogen sources for photolithotrophs, New Phytol., 121, 19–32, https://doi.org/10.1111/j.1469-8137.1992.tb01088.x, 1992. Rickard, D. and Luther, G. W.: Chemistry of Iron Sulfides, Chem. Rev., 107, 514–562, https://doi.org/10.1021/cr0503658, 2007.

Rudnick, R. L. and Gao, S.: Composition of the continental crust, Treatise Geochem., 3, 1–64, 2003.

645 Sholkovitz, E. R., Shaw, T. J., and Schneider, D. L.: The geochemistry of rare earth elements in the seasonally anoxic water column and porewaters of Chesapeake Bay, Geochim. Cosmochim. Acta, 56, 3389–3402, 1992.

Sinex, S. A. and Helz, G. R.: Regional geochemistry of trace elements in Chesapeake Bay sediments, Environ. Geol., 3, 315–323, https://doi.org/10.1007/BF02473521, 1981.

650 Smith, S. V. and Mackenzie, F. T.: The Role of CaCO3 Reactions in the Contemporary Oceanic CO2 Cycle, Aquat. Geochem., 22, 153–175, https://doi.org/10.1007/s10498-015-9282-y, 2016.

Soetaert, K., Hofmann, A. F., Middelburg, J. J., Meysman, F. J. R., and Greenwood, J.: The effect of biogeochemical processes on pH, Mar. Chem., 105, 30–51, https://doi.org/10.1016/j.marchem.2006.12.012, 2007.

655 Stookey, L. L.: Ferrozine—a new spectrophotometric reagent for iron, Anal. Chem., 42, 779–781, 1970.

Su, J., Cai, W.-J., Brodeur, J., Chen, B., Hussain, N., Yao, Y., Ni, C., Testa, J. M., Li, M., Xie, X., Ni, W., Scaboo, K. M., Xu, Y., Cornwell, J., Gurbisz, C., Owens, M. S., Waldbusser, G. G., Dai, M., and Kemp, W. M.: Chesapeake Bay acidification buffered by spatially decoupled carbonate mineral cycling, Nat. Geosci., 13, 441–447, https://doi.org/10.1038/s41561-020-0584-3, 2020a.

660 Su, J., Cai, W., Brodeur, J., Hussain, N., Chen, B., Testa, J. M., Scaboo, K. M., Jaisi, D. P., Li, Q., Dai, M., and Cornwell, J.: Source partitioning of oxygen-consuming organic matter in the hypoxic zone of the Chesapeake Bay, Limnol. Oceanogr., 65, 1801–1817, https://doi.org/10.1002/lno.11419, 2020b.

Su, J., Cai, W., Testa, J. M., Brodeur, J. R., Chen, B., Scaboo, K. M., Li, M., Shen, C., Dolan, M., Xu, Y., Zhang, Y., and Hussain, N.: Supply-controlled calcium carbonate dissolution decouples the seasonal
dissolved oxygen and PH minima in Chesapeake Bay, Limnol. Oceanogr., lno.11919, https://doi.org/10.1002/lno.11919, 2021.

Thamdrup, B. and Dalsgaard, T.: The fate of ammonium in anoxic manganese oxide-rich marine sediment, Geochim. Cosmochim. Acta, 64, 4157–4164, https://doi.org/10.1016/S0016-7037(00)00496-8, 2000.

670 Thibault de Chanvalon, A. and Luther, G. W.: Mn speciation at nanomolar concentrations with a porphyrin competitive ligand and UV-vis measurements, Talanta, 200, 15–21, https://doi.org/10.1016/j.talanta.2019.02.069, 2019.

Trouwborst, R. E., Clement, B. G., Tebo, B. M., Glazer, B. T., and Luther, G. W.: Soluble Mn(III) in Suboxic Zones, Science, 313, 1955–1957, https://doi.org/10.1126/science.1132876, 2006.

675 Urey, H. C.: On the Early Chemical History of the Earth and the Origin of Life, Proc. Natl. Acad. Sci., 38, 351–363, https://doi.org/10.1073/pnas.38.4.351, 1952.

Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Körtzinger, A., and Dickson, A. G.: Total alkalinity: The explicit conservative expression and its application to biogeochemical processes, Mar. Chem., 106, 287–300, https://doi.org/10.1016/j.marchem.2007.01.006, 2007.

Kue, L. and Cai, W.-J.: Total alkalinity minus dissolved inorganic carbon as a proxy for deciphering ocean acidification mechanisms, Mar. Chem., 222, 103791, https://doi.org/10.1016/j.marchem.2020.103791, 2020.

Zeebe, R. E. and Wolf-Gladrow, D.: CO2 in seawater: equilibrium, kinetics, isotopes, Gulf Professional Publishing, 2001.

685 Zhang, Q., Brady, D. C., Boynton, W. R., and Ball, W. P.: Long-Term Trends of Nutrients and Sediment from the Nontidal Chesapeake Watershed: An Assessment of Progress by River and Season, JAWRA J. Am. Water Resour. Assoc., 51, 1534–1555, https://doi.org/10.1111/1752-1688.12327, 2015.