

# Droughts can Reduce the Nitrogen Retention Capacity of Catchments

Carolin Winter<sup>1</sup>, Tam V. Nguyen<sup>1</sup>, Andreas Musolff<sup>1</sup>, Stefanie R. Lutz<sup>2</sup>, Michael Rode<sup>3</sup>, Rohini Kumar<sup>4</sup>, Jan H. Fleckenstein<sup>1,5</sup>

5 <sup>1</sup>Department for Hydrogeology, Helmholtz Centre for Environmental Research - UFZ, Germany

<sup>2</sup>Copernicus Institute of Sustainable Development, Utrecht University, Netherlands

<sup>3</sup>Department Aquatic Ecosystem Analysis and Management, Helmholtz Centre for Environmental Research - UFZ, Germany

<sup>4</sup>Department of Computational Hydrosystems, Helmholtz Centre for Environmental Research - UFZ, Germany

10 <sup>5</sup>Hydrologic Modelling Unit, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Bayreuth, Germany

*Correspondence to:* Carolin Winter (carolin.winter@ufz.de)

**Abstract.** In 2018–2019, Central Europe experienced an unprecedented two-year drought with severe impacts on society and ecosystems. In this study, we analyzed the impact of this drought on water quality by comparing long-term (1997–2017) nitrate export with 2018–2019 export in a heterogeneous mesoscale catchment. We combined data-driven analysis with process-based  
15 modeling to analyze nitrogen retention and the underlying mechanisms in the soils and during subsurface transport. We found a drought-induced shift in concentration-discharge relationships, reflecting exceptionally low riverine nitrate concentrations during dry periods and exceptionally high concentrations during subsequent wet periods. Nitrate loads were up to 73% higher compared to the long-term load-discharge relationship. Model simulations confirmed that this increase was driven by decreased denitrification and plant uptake and subsequent flushing of accumulated nitrogen during rewetting. Fast transit times (<2  
20 months) during wet periods in the upstream sub-catchments enabled a fast water quality response to drought. In contrast, longer transit times downstream (>20 years) inhibited a fast response but potentially contribute to a long-term drought legacy. Overall, our study reveals that severe droughts, which are predicted to become more frequent across Europe, can reduce the nitrogen retention capacity of catchments, thereby intensifying nitrate pollution and threatening water quality.

25 **Key words:** drought, water quality, nitrate, catchment, biogeochemical cycling, nitrogen retention, data-driven analysis, process-based modeling

## 1 Introduction

In 2018–2019, large parts of Europe experienced a severe drought that was unprecedented in the last 250 years (Hari et al., 2020; Rakovec et al., 2022). This drought, caused by exceptionally low precipitation concurring with high temperatures, had  
30 detrimental impacts on vegetation during the growing season and caused massive forest diebacks (Hari et al., 2020; Schuldt et

al., 2020). Besides the scarcity of water and its direct impact on ecosystems and society (Delpla et al., 2009; Fu et al., 2020; Stahl et al., 2010), there is first evidence that this drought could also have impacted freshwater quality in regard to nitrate concentrations. The Nitrate Report 2020 of the Netherlands (RIVM, 2021), for example, found an increase in nitrogen (N) surplus in agricultural areas across the country and, with it, an increase in leachate nitrate concentrations below the root zone.

35 This increase in N surplus and leachate nitrate concentrations in response to drought has been explained by the low water availability that might reduce crop growth and, thus, N plant uptake (Cramer et al., 2009; RIVM, 2021). However, high nitrate concentrations in the leachate do not necessarily reach the stream network, because catchments act as a filter and reactor that can delay the transit of N to the receiving stream or permanently remove it via denitrification (Van Meter & Basu, 2015). The extent of delays and removal strongly depends on the catchment characteristics and boundary conditions (e.g., Ehrhardt et al.,

40 2021; Jawitz et al., 2020; Winter et al., 2021). Moreover, different N sources and their spatial distribution within a catchment can impact nitrate export at the catchment outlet (Casquin et al., 2021; Dupas et al., 2019). Therefore, both catchment characteristics and their spatial configuration might shape the response of riverine nitrate export to drought.

Diverse responses of stream water nitrate concentrations have been reported in different catchments for previous droughts and subsequent post-drought periods (Morecroft et al., 2000; Mosley et al., 2015). Several studies have found decreasing nitrate

45 concentrations during droughts, which have been attributed either to disconnected shallow flow paths that normally allow for efficient transport of nitrate to the stream (J. Yang et al., 2018) or to increased in-stream retention efficiency and to increased uptake along with higher temperatures (Morecroft et al., 2000; Mosley, 2015; Oelsner et al., 2007). However, also increases or no changes in stream concentrations have been reported during droughts, mainly due to high nitrate concentrations in the baseflow or due to the presence of point sources, which increase in relative importance under low flow conditions (e.g.,

50 Andersen et al., 2004; Jarvie et al., 2003; Sprague, 2005; van Vliet & Zwolsman, 2008). With rewetting after the drought, many studies have reported high nitrate concentration peaks as a consequence of accumulated nitrate in the soil zone being flushed from the catchment via fast and shallow flow paths (Górski et al., 2019; Loecke et al., 2017; Morecroft et al., 2000; Mosley, 2015; Outram et al., 2014). This pattern can be explained by both remobilization of accumulated nitrate and stimulation of mineralization with the rewetting of dry soils (Campbell & Biederbeck, 1982; Haynes, 1986; Loecke et al.,

55 2017). Together, these findings imply that droughts can have profound impacts on nitrate availability and transport to the stream network and that the catchment response to droughts seems to vary between catchments and potentially also with drought magnitude. Furthermore, recent studies have highlighted the role of different sub-catchments with different response times to changes composing the integrated signal of nitrate export at the catchment outlet (e.g., Ehrhardt et al., 2019; Nguyen et al., 2022; Winter et al., 2021). It can therefore be important to account for the spatial heterogeneity of a catchment and to

60 look at sub-catchment-specific contributions to better understand the overall catchment response to drought in terms of nitrate export.

To identify drought impacts on nitrate export, data-driven approaches have the advantage of giving a direct reflection of real observations that are an integrated result of the complex biogeochemical and hydrological processes within the catchment. Data-driven approaches thus provide observation-based understanding without strong assumptions on the underlying processes

65 while allowing to build hypotheses on these processes. For example, the relationships of nitrate concentrations and discharge (C-Q) and of nitrate loads and discharge (L-Q) can serve as a robust characterization of catchment-specific nitrate export patterns under different flow conditions and allow for conclusions on N source availability and distribution and on catchment specific response times (e.g., Bieroza et al., 2018; Minaudo et al., 2019; Musolff et al., 2015). Moreover, a comparison of N input and output from a catchment allows for the quantification of catchment N retention resulting from hydrological and/or  
70 biogeochemical N legacies and denitrification (Ehrhardt et al., 2019; Van Meter et al., 2016; Van Meter & Basu, 2015; Winter et al., 2021). Tools complementary to data-driven analyses are mechanistic and process-based models, which explicitly aim at a physical description of the underlying processes causing the observed concentrations and loads and therefore provide detailed insights into catchment-internal N dynamics. For example, the mesoscale Hydrological Model with StorAge Selection functions (mHM-SAS; Nguyen et al., 2022) allows quantifying the rates of N uptake and removal in the catchment soils, lateral  
75 N transport at the sub-catchment scale and time-variant transit time distributions (TTDs). However, resulting simulations also rely on fixed assumptions and the distinct processes that these models entail. Therefore, combining data-driven analyses and mechanistic process-based modeling has several advantages: The data-driven analysis allows for robust identification of drought impacts on nitrate export and a discussion on the underlying processes, while the process-based modeling allows testing if these processes can actually explain the observed behavior.

80 In this study, we used data-driven analysis and process-based modeling to analyze the impact of the 2018-2019 drought on nitrate concentrations and loads compared to previous years (1997-2017) in a heterogeneous mesoscale catchment with three nested gauges located in Germany. This setup allows us to disentangle sub-catchment-specific drought responses while obtaining results at a larger and integral spatial scale relevant to water quality management. We hypothesize that droughts can cause a change in the nitrogen retention capacity of catchments, but with different impacts on riverine nitrate export at  
85 contrasting sub-catchments. Our specific objectives were to i) identify changes in riverine nitrate concentrations and loads at the sub-catchment scale via data-driven analyses (using C-Q and L-Q relationships) and ii) quantify changes in N cycling in the catchment soils and in the time scales of lateral N transport from the soil leachates to the stream network via process-based modeling (using mHM-SAS). This approach allows us to gain knowledge on drought impacts on catchment functioning in terms of retaining and releasing N, which is crucial for our ability to adapt to climate change and effectively mitigate nitrate  
90 pollution.

The 2018-2019 drought was an unprecedented event, but with accelerating climate change, such prolonged droughts are likely to become more frequent (Hari et al., 2020; IPCC, 2018; Rakovec et al., 2022; Samaniego et al., 2018). In this context, this study is one step towards a better understanding of the impacts of such droughts on nitrate export dynamics and the underlying mechanisms within a catchment.

2.1 Study Site

This study was conducted in the mesoscale Selke catchment, which is located in the Harz Mountains and Harz foreland in Saxony-Anhalt, Germany (Figure 1a). As a sub-catchment of the Bode basin, it is also part of the network of TERrestrial ENvironmental Observatories (TERENO; Wollschläger et al., 2017). The Selke catchment is gauged with three nested stations:  
100 Silberhütte (SH), Meisdorf (MD), and Hausneindorf (HD, Figure 1a). The two upstream sub-catchments form the upper Selke with similar characteristics, such as forest being the dominant land use and also in terms of relatively short TTs and comparable nitrate export dynamics (Nguyen et al., 2022; Winter et al., 2021). The downstream part forms the lowland area of the catchment, which is dominated by agricultural land use. Compared to the upper Selke, TTs are longer, and the variability of export dynamics is reduced (Nguyen et al., 2022; Winter et al., 2021).




Figure 1: Study site and long-term hydro-meteorological conditions. a) Land use map of the Selke catchment with its three gauging stations (SH, MD, and HD). b) Climatic anomalies in the Selke catchment in terms of precipitation, temperature, and discharge for the years (starting in May) from 1990 to 2019.

Table 1: Sub-catchment specific characteristics of the Selke catchment

Sub-catchment characteristic	Unit	Upper Selke	Upper Selke	Lower Selke
		Silberhütte (SH)	Meisdorf (MD)	Hausneindorf (HD)

<b>Area</b>	(km <sup>2</sup> )	100.9	78.9	277.6
<b>Mean Elevation</b>	(m.a.s.l.)	448.9	370	164.8
<b>Mean Slope</b>	(%)	6.8	11.5	2.6
<b>Mean annual precipitation</b>	(mm)	718.6	646.9	537
<b>Mean annual temperature</b>	(°C)	8	8.4	9.9
<b>Land use</b>	(%)			
Forest		61.3	87.7	12.1
Agriculture		36.1	10	76.2
Others		2.6	2.3	10.7
<b>Dominant soil type</b>		Dystic/spodic cambisols	Haplic chernozem	
<b>Dominant geology</b>		Paleozoic greywacke/ Denovian shale	Sedimentary	

Note. Catchment characteristics refer to spatially separated, not nested sub-catchments. Precipitation and temperature data were taken from the period 1997–2020.

## 2.2 Characterization of Different Hydro-Meteorological conditions and anomalies

- 115 We adopted the definition of annual wet, drying, dry, and wetting periods from J. Yang et al. (2018), based on the catchment subsurface storage condition in a headwater catchment of the Selke catchment. Accordingly, wet periods last from January to April, drying periods (i.e., the transition from wet to dry) last from May to June, dry periods last from July to October, and wetting periods (i.e., the transition from dry to wet) last from November to December. Instead of annual averages starting in January, we calculated annual averages of discharge and N fluxes over 12-month periods starting with the drying period in
- 120 May and ending with the wet period end of April. This was done under the rationale that nitrate starts accumulating in a catchment over the drying and dry period, when fast flow paths are deactivated (J. Yang et al., 2018), and that, subsequently, accumulated nitrate is more efficiently exported from the catchment during wetter conditions. Under this rationale, comparing annual statistics of nitrate export is more meaningful if comparing 12-month periods starting in May instead of January, considering the seasonality in climatic conditions in central Germany. Throughout the manuscript, we therefore defined years

125 by drying-wetting cycles starting in May, with the consecutive numbering being based on the starting date. For example, the year 2018 started on 1<sup>st</sup> May 2018 and ended on 30<sup>th</sup> April 2019. In the same manner, we refer to the two-year drought as 2018-2019, spanning a period from May 2018 until the end of April 2020.

To compare the hydro-meteorological conditions during the drought years (2018 and 2019) with the ones during previous years (1990–2017), we calculated their hydro-meteorological anomalies compared to the long-term average. To this end, we  
130 calculated the annual averages in observed temperature, precipitation, and discharge and modeled soil moisture (see section 2.5) for the study catchment. We then calculated the long-term average over all years and subtracted the single annual averages from those long-term averages. The divergence of annual hydro-meteorological conditions from the long-term average is considered the hydro-meteorological anomaly. In the Selke catchment for the 2018-2019 drought, the years starting in May 2018 and May 2019 are characterized by exceptionally low precipitation (anomaly of -205mm yr<sup>-1</sup> and -110 mm yr<sup>-1</sup> in 2018  
135 and 2019, respectively, compared to the long-term average of 602 mm yr<sup>-1</sup> over the period 1997-2020), high temperatures (+1.6°C and +1.4°C in 2018 and 2019, compared to the long-term average of 9.0°C) and low discharge (-37.0 mm yr<sup>-1</sup> and -39.2 mm yr<sup>-1</sup> compared to the long-term average of 98.5 mm yr<sup>-1</sup>) (Figure 1b). In terms of soil moisture, these two years were the driest in the Selke catchment since the start of our time series in May 1997, with 2018 being even drier than 2019 (Figure S1). Consequently, the two-year drought that affected large parts of Central Europe (Hari et al., 2020) can also be characterized  
140 as an exceptional drought in the Selke catchment.

### 2.3 Data

Daily long-term temperature and precipitation data (1997–2020) were provided by the German Meteorological Service (Deutscher Wetterdienst, DWD). N input data (i.e., fertilizer, manure, and plant residues) and land use management (i.e., crop rotation), and atmospheric deposition were based on agricultural authority data obtained from X. Yang et al. (2018) and Jomaa  
145 et al. (2018). Accordingly, N input to agricultural fields in mHM-SAS follows a two- or three-year crop rotation, as typical in this area and as implemented in X. Yang et al. (2018) and Nguyen et al. (2022). Daily discharge data and biweekly-monthly grab samples of nitrate concentrations were provided by the State Office of Flood Protection and Management of Saxony-Anhalt (LHW; Figure 2a-c). Sensor measurements (using TriOS Pro-UV sensors; Rode et al., 2016) of nitrate concentrations at a 15-minute resolution (2012–2020) were provided by the TERENO facilities of the Helmholtz Centre for Environmental  
150 Research (UFZ). To match the temporal resolution of long-term data, nitrate concentrations were aggregated to daily averages (Figure 2a-c). Part of this data was previously used by (Musolff et al., 2021; Rode et al., 2016; Winter et al., 2021, 2022; X. Yang et al., 2018). Therefore, for the detailed processing of nitrate concentration data, we refer to Rode et al. (2016) and the other references above. With a coefficient of determination ( $R^2$ ) between 0.8 for MD and HD and 0.9 for SH, processed high-frequency nitrate concentration data from sensor measurements showed a good agreement with concentrations from grab  
155 samples analyzed in the lab.

## 2.4 Data-Driven Analysis Using Concentration-Discharge Relationships

To characterize nitrate export in the Selke catchment, we performed a data-driven analysis using concentration-discharge (C-Q) relationships from daily averages and load-discharge (L-Q) relationships from annual averages (starting in May). A simple but efficient way to describe the C-Q relationship is a power-law relationship of the form

$$C(t) = \alpha Q(t)^\beta \quad (1)$$

with  $t$  standing for the respective time step. The parameters  $\alpha$  and  $\beta$  describe the intercept ( $\alpha$ ) and the slope of the relationship in the log-log space ( $\beta$ ), also termed C-Q slope (Musolff et al., 2015; Thompson et al., 2011). A positive C-Q slope indicates an increase of nitrate concentrations with discharge (enrichment pattern), whereas a negative C-Q slope indicates decreasing nitrate concentrations with increasing discharge (dilution pattern). Both patterns imply a directional relationship between concentrations and discharge, with nitrate concentrations either increasing or decreasing with increasing discharge. On the contrary, a C-Q slope around zero indicates that nitrate concentrations are not or are poorly correlated with the dynamics of discharge. Since nitrate loads (L) are the product of nitrate concentrations (C) and discharge (Q), the L-Q slope can be described using  $\beta+1$ , with the differentiation in this study that the C-Q slope and the L-Q slope were calculated with data of different temporal resolutions and are thus not directly comparable.

The C-Q relationship was calculated from daily averages of measured data only and is therefore restricted to the period 2012–2020. To test if the C-Q slope for the two-year drought was different than the long-term average, we compared both slopes and their standard errors. To account for the different sample sizes between the pre-drought and drought periods, we applied additional block sampling across all possible combinations of two consecutive years and compared the resulting pre-drought C-Q slopes with the one from the two-year drought.

Instead of restricting our analysis to observed daily data, as done for C-Q relationships, we calculated L-Q relationships with the annual sums of daily load and discharge data starting in May 1997. To this end, we used the continuous daily discharge measurements and filled the gap in daily nitrate concentration measurements by interpolating from biweekly–monthly grab samples via Weighted Regression on Time Discharge and Season (WRTDS; Hirsch et al., 2010). The fit between daily loads calculated from interpolated and measured nitrate concentration data (2012–2019) was high, with an  $R^2$  between 0.93 and 0.96 and a small percentage bias between -0.5% and -1.1% (Figure S2).

## 2.5 Process-Based Nitrogen Export Modeling with Storage Selection Functions

To get a deeper insight into catchment processes that cause the observed nitrate export patterns during and after the two-year drought, we simulated daily nitrate concentrations at the three gauging stations using the mesoscale Hydrological Model with StorAge Selection functions (mHM-SAS, Nguyen et al., 2021, 2022). The mHM-SAS model is a deterministic model with a strong physical basis, explained in detail in Nguyen et al. (2022) and in the supplements (Text S1). Briefly, mHM-SAS provides a spatially distributed (1x1 km<sup>2</sup>) representation of hydro-climatic inputs, N pools, and fluxes in the soil zone based on a combination of mHM and the soil nitrogen model (X. Yang et al., 2018). Nitrate pools and fluxes in the saturated and

unsaturated zone below the soil are represented for each sub-catchment using the nitrate transport model with StorAge Selection (SAS) functions (van der Velde et al., 2012, Nguyen et al., 2022). SAS functions describe the selective removal of water from a subsurface storage with different water ages and nitrate concentrations, which allows for a nitrate transport formulation based on time-variant TTDs. The SAS function in mHM-SAS is described using a beta function ( $\beta(a,b)$ ), with  $a$  and  $b$  being two fitted parameters that vary in time (see Supplement, Eq. (6)). The derived  $a/b$  ratio represents the selection schemes for discharge, e.g., preference for young water ( $a/b$  ratio  $< 1$ ) or old water ( $a/b$  ratio  $> 1$ ) (Nguyen et al., 2022). Nguyen et al. (2022) calibrated the model in the Selke sub-catchments for the years 2012–2015 with a Nash-Sutcliffe Efficiency (NSE) of 0.68, 0.66, and -0.13 over the calibration and 0.81, 0.81, and 0.57 for SH, MD, and HD over the validation period (2016–2019), which includes parts of the two-year drought. The lower NSE in HD can be explained by the lower seasonality in nitrate concentrations (Nguyen et al., 2022; Schaefli & Gupta, 2007). Using the same setup, here we extended the model simulations to the 1997–2020 time period (Figure 2a-c). To create an initial age distribution in the storage before 1997 and to minimize the effect of initial conditions, we replicated model input data between 1993 and 1996 ten times as a warm-up period to obtain initial conditions for our actual model runs (1997–2020). We used these extended simulations to contrast average conditions with the 2018–2019 drought-induced changes in the C-Q relationships in the different sub-catchments of the Selke catchment. Despite distinctly different climatic conditions during the 2018–2019 drought period, nitrate concentrations simulated with mHM-SAS showed an even better fit to the measured sensor data (NSE of 0.89, 0.88, and 0.79 in SH, MD, and HD respectively) than for the calibration period. (Figure 2a-c). This good fit over the drought years indicates that the model is also applicable under very dry hydro-meteorological conditions. With this setup, the mHM-SAS model allows for a separation of the contribution of each sub-catchment to the overall catchment responses to account for sub-catchment specific N cycling in the soil zone (denitrification, plant uptake, mineralization, and leaching), instream uptake, denitrification along the groundwater flow paths and for dynamic transit times (TTs).

## 2.6 Catchment Retention Capacity

We estimated the capacity of a catchment to retain N ( $N_{ret}$ ) during one year (starting in May) by the ratio of average nitrate-N load export ( $N_{OUT}$ ) against annual average atmospheric deposition and long-term average N inputs from fertilizer and manure ( $N_{IN}$ ):

$$N_{ret} = 1 - \frac{N_{OUT}}{N_{IN}} \quad (2)$$


We used the long-term average N input across crop rotations as precise information on which crop is applied to which field in which year is not available, and thus a long-term average is more robust. This approach is justified by the fact that N input data did not show any trends and no significant deviation during the two-year drought. Additionally, we assessed the sensitivity of our results to uncertainty in N input through crop rotation by varying N inputs by  $\pm 20\%$ , which confirmed the overall robustness of our results (Figure S3).

To estimate changes in  $N_{\text{ret}}$  relative to the hydrological conditions, we fitted a non-linear regression between  $N_{\text{ret}}$  and observed discharge ( $Q$ ) for the years previous to the two-year drought (1997-2017). To this end, we assume that  $Q$  is log-normally distributed and related to nitrate loads (i.e.,  $N_{\text{OUT}}$ ) in the form of a power law relationship ( $\alpha Q^{\beta+1}$ ), with  $\beta+1$  being the L-Q slope. Consequently,  $N_{\text{ret}}$  can be described as a function of  $Q^{\beta+1}$ , which would be linear if the L-Q slope equals 1. The result is an  $N_{\text{ret}}$ - $Q$  relationship that asymptotically approaches 1 (i.e., 100% retention) at zero discharge, and that is zero if  $N_{\text{OUT}}$  equals  $N_{\text{IN}}$ .  $N_{\text{ret}}$  is a combined measure of biogeochemical N retention in the catchment soils and its consecutive transport via hydrological flow paths to the stream network, which is affected by TTs and denitrification along the flow paths. To acquire a more direct estimate of soil N retention, we additionally fitted the  $N_{\text{ret}}$ - $Q$  relationship for simulated nitrate leachates (Figure S4).

### 3 Results

#### Nitrate Concentrations

Daily nitrate-N concentrations (i.e., sensor measurements) differed between the upper (SH and MD) and the lower Selke (HD) and between normal (i.e., 2012–2017) and drought years (2018–2019; Figure 2a-c). Median concentrations in the upper Selke measured before the two-year drought ranged from 0.6 mg L<sup>-1</sup> and 0.7 mg L<sup>-1</sup> during dry periods in SH and MD, respectively, and 2.6 mg L<sup>-1</sup> and 2.4 mg L<sup>-1</sup> during wet periods. Median nitrate-N concentrations measured at HD were higher and less variable, with a median of 2.2 mg L<sup>-1</sup> during dry periods and 3.1 mg L<sup>-1</sup> during wet periods. During the two-year drought, nitrate-N concentrations generally showed a higher seasonal variability (Figure 2a-c). During dry periods in 2018 and 2019, nitrate-N concentrations were lower than during previous dry periods, with a median of 0.2 mg L<sup>-1</sup> and, 0.4 mg L<sup>-1</sup> and 1.4 mg L<sup>-1</sup> in SH, MD, and HD, respectively. During the subsequent wet periods (January-April 2019 and 2020), nitrate-N concentrations were exceptionally high, with a median of 4.2 mg L<sup>-1</sup>, 3.8 mg L<sup>-1</sup>, and 3.7 mg L<sup>-1</sup>. In the upper Selke (SH and MD), nitrate-N concentrations reached the highest value observed since the start of measurements in 1983, with 6.4 mg L<sup>-1</sup> in January 2019. Peak concentration in the lower Selke (HD) during that time was 5.9 mg L<sup>-1</sup>, which was also among the highest values measured at this gauge (Fig. 2c).




**Figure 2: Nitrate as nitrogen concentrations and discharge at the gauging stations of the three Selke sub-catchments (1997–2020) with low-frequency grab samples (orange dots), simulated concentrations via mHM-SAS (grey line), daily averages of high-frequency sensor measurements from 2012 on (red line), and daily discharge (bluish lines). Dashed lines indicate the mHM-SAS calibration (2012–2015) and validation (2016–2020) periods, comparing simulated nitrate-N concentrations with daily averages of sensor measurements.**

### 3.1 Concentration-Discharge and Load-Discharge Relationship

Nitrate concentrations during the two-year drought show an accelerated seasonality (see section 2.3) that is reflected in a more chemodynamic C-Q relationship (Figure 3a-c). All three sub-catchments show a positive C-Q relationship for daily averages of pre-drought (January 2012 – April 2018) nitrate-N concentrations and discharge, with the highest slope in SH, followed by

MD, and the lowest slope in HD. During the two-year drought, the C-Q slope for all sub-catchments increased by values that are multiples of the standard error of the pre-drought regressions for the entire period (Figure 3a-c), but also for block sampled C-Q slopes that account for the smaller samples size of the drought period (Figure S5).



**Figure 3: Concentration-discharge (C-Q) and load-discharge (L-Q) relationships for the three sub-catchments of the Selke catchment (SH, MD, and HD). a-c) Daily averaged nitrate concentration and discharge data with log-transformed axes. The lines show the C-Q slope for daily averages before the two-year drought (grey) and since the start of the two-year drought (dark red). d-f) Annual averages for loads and discharge in the log-log space since 1997 (years starting in May). Black lines show the L-Q slope previous to the two-year drought. Boxplots at the right side of each panel (d-f) show the distribution of data points within the pre-drought load range and drought cycles indicated as colored dots.**


Median nitrate-N loads per area over the years 1997–2017 were 6.6 kg yr<sup>-1</sup> ha<sup>-1</sup>, 5.7 kg yr<sup>-1</sup> ha<sup>-1</sup>, and 3.2 kg yr<sup>-1</sup> ha<sup>-1</sup> in SH, MD, and HD, respectively. During the two-year drought, nitrate-N loads were in a similar range in SH (6.0 and 7.2 kg yr<sup>-1</sup> ha<sup>-1</sup>, in 2018 and 2019, respectively) and lower but still within the interquartile range in MD (4.5 and 4.9 kg yr<sup>-1</sup> ha<sup>-1</sup>) (Figure 3d-f). They were clearly lower in HD with a load of 2.1 kg yr<sup>-1</sup> ha<sup>-1</sup> in both years (Figure 3d-f).

L-Q relationships showed a good fit ( $R^2$  0.9 – 0.96) with an L-Q slope close to 1 in all sub-catchments, reflecting that nitrate-N loads increased with increasing discharge (Figure 3d-f). During the drought cycles 2018 and 2019, loads exported from the

upper Selke were clearly above the loads expected from the long-term L-Q relationship. So, relative to discharge that was naturally low during the drought, loads were unexpectedly high. More specifically, exported loads at SH were 2.2 and 2.9 kg ha<sup>-1</sup> yr<sup>-1</sup> higher in 2018 and 2019 than expected from the long-term L-Q relationship, and at MD, loads were 1.9 kg ha<sup>-1</sup> yr<sup>-1</sup> higher in both years. In relative numbers, this is an increase of 57%-73% compared to the predicted export from the L-Q relationship from previous years. In the lower Selke (HD), on the contrary, the difference between observed and expected nitrate-N export was marginal during the two-year drought (0.2 kg ha<sup>-1</sup> yr<sup>-1</sup> in both years, equivalent to an increase of 10%). However, exported nitrate loads in the lower Selke have generally decreased since 2011 (Winter et al., 2021). Therefore, 2013–2017 loads are the ones plotting clearly below the long-term L-Q slope (Figure 3f); as such, the L-Q relationships from the two-year drought can be seen as slightly increased if compared to the most recent years only. To illustrate, exported nitrate loads at HD during the two-year drought are around 0.5 kg ha<sup>-1</sup> yr<sup>-1</sup> higher than expected from the 2013–2017 L-Q relationship.

### 3.2 Simulated Internal Nitrogen Fluxes

The sub-catchment-specific N fluxes, simulated via mHM-SAS (with a good model fit for in-stream nitrate concentrations) and averaged over the years starting in May, are depicted in Figure 4. They show that particularly in 2018, which was the driest year of the two-year drought (Figure 1b), N fluxes clearly differed from the long-term average (1997–2017). Notably, mineralization rates in 2018 increased by 39 %, 36 %, and 66 % in SH, MD, and HD, respectively. In the same year, denitrification in the soils of the sub-catchments was 27 %–34 % lower than the long-term average, whereas plant uptake was reduced by around 10 % in the upper Selke (SH and MD) but not in the lower Selke (HD), likely due to differences in the soil type (Table 1). N in the leachates was relatively low in both cycles (2018 and 2019), especially in MD and HD, due to the dry soil moisture content (Figure S1).




**Figure 4: N input and N fluxes simulated via mHM-SAS, separately for the three Selke sub-catchments (SH, MD, and HD). N entering the catchment soils are shown as positive values, N fluxes leaving the catchment soils are shown as negative values. Error bars represent the standard deviation between annual averages from 1997 to 2017.**

### 3.3 Transit Times and Storage Selection Preference

Similar to nitrate concentrations and loads, the simulated a/b ratio for SAS functions (indicative of young versus old water preference) and median TTs showed a different behavior during the two-year drought compared to previous years, with clear contrasts between upper (SH and MD) and lower Selke (HD; Figure 5). Upper Selke sub-catchments showed a young water preference (a/b ratio < 1) with shorter median TTs during the wet periods (January–April; median of 42 and 56 days in SH and MD, previous to the drought and median of 23 and 39 days during the drought). During dry periods (July–October) previous to the drought, the median of median TTs in the upper Selke was 2.5 and 7.0 yrs in SH and MD, respectively (Figure 5d–e). Nevertheless, more than half of all years still showed a young water preference, even during dry periods (Figure 5a–b). However, during the dry periods of the two-year drought, median TTs were very long (median of median TTs was 66 yrs in both sub-catchments) with a pronounced old water preference, particularly in 2018. Note that the maximum TTs in the simulations are restricted to the number of years since the start of simulations plus the warm-up period, which explains the July–December plateau (Figure 5). Long median TTs during the dry season can therefore be interpreted as > 66 yrs. This cutoff likely causes an underestimation of the median of previous years as well and creates some additional uncertainty in the absolute numbers. However, this does not affect the general result of exceptionally high median TTs during the dry periods in 2018–2020 compared to previous years (1997–2017).

In the lower Selke sub-catchment (HD), there is a clear selection preference for old water throughout all years and periods (Figure 3c). Even during wet periods, median TTs were relatively long (median 20 yrs) compared to the upper Selke sub-catchments (SH and MD; Figure 5d–f). Similar to the upper Selke sub-catchments, median TTs during the dry periods in 2018 and 2019 were longer than normal, with a median of 30 yrs compared to 24 yrs in previous years.




**Figure 5: Simulated sub-catchment-specific water age selection preference (a-c) as the ratio of the fitted parameters  $a$  and  $b$ , and (d-f) median transit times are given in years. Grey areas indicate the range of all years previous to the drought (1997–2017, with a year starting in May), and their interquartile range and white lines are the median of all pre-drought years. Colored lines indicate the two-year drought, with the two years starting in May 2018 and 2019.**

### 3.4 Catchment Retention Capacity

In all cases, the sub-catchments retained the largest part of  $N$  input ( $\geq 0.79$ ). This was most pronounced in HD, where retention capacity was always  $\geq 0.89$  (note that these values account for the nested catchment). The fit of the  $N_{\text{ret}}-Q$  relationship (1997–2017) was high, with an  $R^2$  of 0.90, 0.94, and 0.91 in SH, MD, and HD, respectively. In all sub-catchments,  $N_{\text{ret}}$  shows a clear decrease with decreasing annual discharge in all sub-catchments, which is steepest and most sensitive at low  $N_{\text{ret}}$  and high discharge (Figure S3) and flattening towards low discharge conditions (Figure 6). Similar patterns could also be observed for the  $N_{\text{ret}}-Q$  relationship with simulated nitrate leachates from the catchment soils. However, uncertainty for the simulated soil  $N$  retention is larger, due to uncertainty in the model parametrization (Figure S4).

In the upper Selke (SH and MD), the  $N_{\text{ret}}-Q$  relationship during the two-year drought (2018–2019) was clearly lower than that of previous cycles (1997–2017).  $N_{\text{ret}}$  dropped by around 0.04–0.05 compared to the long-term regression line, which can be

translated into 1.8-2.8 kg N ha<sup>-1</sup> yr<sup>-1</sup> that are not retained but exported. In the lower part of the catchment (HD),  $N_{ret}$  was very close to the long-term regression line (difference of 0.002).



**Figure 6: Relationship between the N retention capacity of catchments ( $N_{ret}$ ) and log-scaled discharge ( $Q$ ) at the nested catchment scale, given as annual averages (12-month period starting in May). Black lines represent the non-linear relationship between  $N_{ret}$  and discharge ( $Q$ ) prior to the two-year drought (1997–2017), with log-transformed x-axes.**

## 4 Discussion

### 4.1 Intra-Annual Variability of Nitrate Concentrations

The presented results show that the two-year drought spanning the years 2018 and 2019 and across Central Europe had strong impacts on catchment water quality in terms of nitrate export. We found an increased intra-annual variability of nitrate export, with lower concentrations during the dry periods (July–October) and exceptionally high concentrations during the subsequent wet periods (January–April). This shows that the drought did not only affect nitrate fluxes in the soil leachates, as discussed in the 2020 Nitrate Report from the Netherlands (RIVM, 2021), but it can also propagate through catchments affecting in-stream nitrate concentrations at the catchment outlet within a relatively short time. These results are in close agreement with previous studies that similarly reported low nitrate concentrations during a drought and high concentrations during subsequent rewetting (Górski et al., 2019; Loecke et al., 2017; Morecroft et al., 2000; Mosley, 2015). For example, Loecke et al. (2017) reported that the shift from very dry to wet conditions resulted in an increase in flow-weighted nitrate export compared to previous years, and Davis et al. (2014) suggested that the antecedent soil moisture conditions play an important role in the response of nitrate export during runoff events.

The shift in the C-Q relationship towards a steeper C-Q slope (Figure 3a-c) reflects that the intensification in the seasonality of nitrate export was not solely driven by low discharge due to the drought. Instead, nitrate concentrations during dry periods were even lower than expected from the C-Q relationship and higher during wet periods, showing an increased intra-annual variability compared to that of discharge. This increased concentration variability indicates that biogeochemical and

hydrological processes (e.g., runoff generation processes; Lange & Haensler, 2012) within the catchment changed during the drought, affecting the availability and transport of nitrate. The exceptionally low nitrate concentrations during the dry periods of the two-year drought can be explained by the strong old water preference during these periods (Figure 5a-c). Nguyen et al. (2022) showed that old water in the upstream Selke catchment is considerably affected by denitrification (Damköhler number > 10), which can explain the relatively low nitrate concentrations. The pronounced old water preference in the upstream catchment during the two-year drought is in agreement with a study by X. Yang et al. (2021). Using stable water isotopes as age tracers in a small (1.44 km<sup>2</sup>) headwater catchment of the Selke catchment, they found a large increase in stream water ages, driven by a decrease in younger surface runoff and stream discharge. Differences in the median TTs between this and our study (8 yrs and 46 yrs) can be explained by the difference in the catchment area of around two orders of magnitude and by uncertainty in the estimation of longer TTs, especially as annually cycling isotope tracers show only limited applicability towards long TTs (Seeger & Weiler, 2014).

In the downstream sub-catchment, the potential for denitrification in groundwater is very low (Damköhler number < 1; Nguyen et al., 2022). Hannappel et al. (2018) looked at evidence for groundwater denitrification in the area of our study site and found such evidence only in the upstream but not in the downstream area, which can be explained by differences in the geology (Table 1) and a lack of electron donors in the aquifer which are needed for denitrification (Rivett et al., 2008). Therefore nitrate concentrations in the downstream sub-catchment do not significantly decrease with water ages and could, instead, still show signs of historically higher N inputs (Winter et al., 2021). Hence the low nitrate concentrations during the dry periods are likely driven by the upstream catchment area. However, also the efficiency of instream N uptake is enhanced with higher temperatures (Nguyen et al., 2022), which is an additional factor explaining part of the low nitrate concentrations during drought, even more so in the lowland part of the catchment where light availability is higher and flow velocity is reduced (Rode et al., 2016). Therefore, a combination of both dilution of old and largely denitrified water from upstream and increased in-stream uptake efficiency, mainly downstream, are responsible for the low nitrate concentrations during the dry periods of 2018 and 2019.

While predominately old (pre-drought) water was exported during dry periods, the rates of denitrification and plant N uptake from the soils during 2018 decreased across the catchment. This decrease can be attributed to the very low soil moisture during the drought that inhibits denitrification (as implemented in the soil routines in HYPE; Lindström et al., 2010) and plant growth. Together with the deactivation of shallow flow paths, the reduced N removal can lead to an accumulation of inorganic N in the catchment soils. Similarly, the higher accumulation of organic material during summer and the rewetting of dry soils in autumn can cause a peak in mineralization that transforms organic material into mobile inorganic N (Campbell & Biederbeck, 1982; Haynes, 1986), in agreement with the simulated high mineralization rates for 2018 (Table 2). With the shift towards younger water fractions and median TTs < 2 months during the wetting and wet periods in the upstream sub-catchments, the accumulated and mineralized N pool can be rapidly transported to the stream network, which can explain the high nitrate-N concentration peaks (Figure 2). In contrast, in the downstream sub-catchment, old water fractions dominate all year round

(Figure 5f), and therefore most of the accumulated N cannot reach the stream network within the subsequent wetting- and wet period.

Note that the ‘wetting-’ and ‘wet periods’ 2018 and 2019 are part of the two-year drought and, in relative terms, were also  
385 exceptionally dry compared to previous wetting and wet periods, but they typically show a higher catchment wetness compared to ‘dry periods’ due to pronounced hydro-meteorological seasonality over the study region (Sinha et al., 2016; J. Yang et al., 2018).

The changes in N cycling were only evident for the year starting in May 2018, not for the one starting in May 2019, which was dry but not as dry as in 2018 (Figure 1b, Table 2). This indicates that the described perturbation in N cycling does only  
390 occur under severe drought conditions. In a small (0.6 km<sup>2</sup>) catchment, Burt et al. (2015) found evidence that post-drought mineralization can supply sufficient N to sustain increased nitrate concentrations through the next high-flow season. Hence, such drought legacies might have also built up in the larger Selke catchment and impacted nitrate export in subsequent years. One indication for this is that although the year 2018 was drier and had a stronger impact on soil-N fluxes (Table 2), nitrate export dynamics for the year 2019 were comparable to the ones observed in 2018 (Figure 3).

395 In comparison to other regions of the world, irrigation is not a common practice in the study area (EEA, 2018), but this might change in the future (Riediger et al., 2014). Crop irrigation would increase the soil moisture content and might therefore buffer drought impacts, such as the decrease in plant uptake and denitrification, and might lead to mobilization of nitrate, which would otherwise be retained in the upper soil until rewetting in autumn. As such, crop irrigation might counterbalance the reduction of N retention in the catchment soils and the accumulation of N during summer. However, irrigation would also  
400 increase the pressure on the available water resources and might enhance greenhouse gas emissions from agricultural soils (Sapkota et al., 2020)

#### **4.2 Sub-Catchment-Specific Contributions to the Integral Nitrate Response to the Drought**

The spatial configuration of land use and other characteristics within a catchment can play an important role in nitrate export from the entire catchment and its temporal variability (Casquin et al., 2021; Dupas et al., 2019; Winter et al., 2021). In the  
405 Selke catchment, considering the pronounced differences between the upstream and the downstream area is crucial to understand the integrated signal of nitrate export at the catchment outlet (Winter et al., 2021). Previous studies showed that the forested upstream area contributed disproportionally to annual nitrate loads, despite having a lower N input, whereas the downstream part contributed most to nitrate export during low-flow periods (Nguyen et al., 2022; Winter et al., 2021). During the two-year drought, this difference in the sub-catchment-specific contributions became even more pronounced. Water from  
410 the upstream catchment area during the dry periods was comparably low in nitrate and therefore had the potential to dilute the higher concentrations from the downstream agricultural areas. Nonetheless, the contribution from the downstream area maintained nitrate-N concentrations at levels >1 mg L<sup>-1</sup> even under low flow conditions during summer, when aquatic ecosystems are most vulnerable to eutrophication (Jeppesen et al., 2010; Whitehead et al., 2009). With rewetting, the ability of the forested upstream area to dilute downstream nitrate concentrations has been almost entirely lost, as nitrate concentrations


415 reached similar or even higher levels than the downstream area (Figure 1c, 2a-c). Therefore, the observed changes in immediate response to the drought with regard to the seasonal dynamic of nitrate concentrations at the catchment outlet were almost entirely controlled by the upstream area due to its shorter TTs and young water preference. The sub-catchment differences in median TTs and storage selection preferences can be explained by differences in landscape characteristics within the catchment. The upstream area is located in the Harz Mountains with higher precipitation, steeper slopes, and shallower soils and, in turn, a faster transit of water and shorter flow paths to the streams, which typically results in faster TTs and is reflected in a selection preference for younger water (Jiang et al., 2014; Tetzlaff et al., 2009; Table 1). In contrast, thick sedimentary aquifers, flat topography, and very sparse tile drainage in the downstream area favor long TTs and a preference for old water (Winter et al., 2021).

Considering the important role of the upstream, largely forested (87.7 %; Table 1), part of the catchment for the overall nitrate export, one should also consider the potential long-term impacts of the two-year drought on forest ecosystems. Schuldt et al. (2020) showed that especially the dry summer in 2018 caused severe tree mortality in Central Europe, whereas Schnabel et al. (2021) could show in a German floodplain forest that the drought impact on trees was even stronger in 2019 due to an accumulated drought effect. Such forest dieback can cause increased nitrate concentrations (Kong et al., 2022; Mikkelsen et al., 2013), but its effect on water quality can again be delayed for several years (Huber, 2005). Therefore, forest dieback should be considered as an additional drought-induced threat to stream water quality that might impact nitrate concentrations in the future.

### 4.3 Exported Nitrate Loads and Catchment Retention Capacity

The overall discharge during the two-year drought was very low. Nitrate loads, however, were only low in the downstream part of the catchment (HD). In the upstream area (SH and MD), nitrate loads were up to 73 % higher than expected from the long-term L-Q relationship. This can be explained by the exceptionally high nitrate concentrations during the wetting- and wet periods of the two-year drought (November-April, Figure 2). As discussed above, these high nitrate concentrations are the result of reduced plant uptake and denitrification of N in the soils during the previous dry periods and short TTs during wet periods. Hence the increase of exported loads relative to discharge, but also relative to N input to the catchment, show that under severe drought, a catchment can lose a part of its functionality to retain N. The long-term L-Q relationship as well as the  $N_{\text{ret}}$ -Q relationship, showed good fits for the pre-drought year, indicating a strong discharge control. Scatter in these relationships might have been induced by other factors, such as temperature, controlling biogeochemical processes, and N availability (Nogueira et al., 2021) or specific runoff event characteristics (Winter et al., 2022). With climate change, temperature and other associated factors as well as runoff event characteristics were not stable between 1997 and 2017 (Figure 1b; IPCC, 2018; Winter et al., 2022). Nevertheless, none of these years has shown a deviation from the L-Q relationship comparable to that in 2018 and 2019. Indicating that the two-year drought considerably altered catchment functioning, whereas the overall L-Q relationship can be considered relatively stable over previous years.

We identified two drivers of a decrease in  $N_{\text{ret}}$ , i.e., an increase in discharge and the two-year drought (Figure 7). The discharge-driven decrease in the  $N_{\text{ret}}$  can be explained by hydrologic mobilization and transport dominating over biogeochemical processes such as N uptake and removal. In contrast, under dry (but not drought) conditions, the role of nitrate uptake and removal gains in relative importance, and with that, the retention capacity of the catchment increases. However, extreme hydro-meteorological events, such as the 2018–2019 drought, can cause a perturbation of those biogeochemical processes if soils are too dry to maintain functionality in terms of N cycling. The decrease of N uptake and removal can result in a divergence from the retention-discharge relationship, also for other years with very low discharge (Figure 6, 7). Until the end of the data record available for this study, no recovery from this loss in  $N_{\text{ret}}$  in the upstream catchment area could be observed, but there has not yet been a specifically wet year since the drought in 2018. Therefore, the resilience of the sub-catchments (Hashimoto et al., 1982), in terms of their ability to recover from a loss in catchment N retention capacity after the two-year drought, remains uncertain.



**Figure 7 Conceptual framework of the capacity of a catchment to retain N in relation to average discharge. The framework illustrates two potential drivers that cause a reduction in the catchment retention capacity. Those are either an increase in discharge (blue arrow) or a drought event (yellow arrow). Data was taken from the gauge at MD that gives an integrated signal of the upper Selke catchment, which is characterized by a relatively fast reaction in riverine nitrate export to drought.**

When looking at N export at the sub-catchment gauges, the decrease in the catchment retention capacity is only evident in the upstream sub-catchments. This difference in catchment retention capacity can be explained by different sub-catchment-specific TTs. Whereas predominantly short TTs in the upstream sub-catchments during the wet period allow for a rapid response of stream nitrate concentrations to drought, long median TTs in the downstream sub-catchment (even under wet conditions) dampened such a fast drought response. Instead, increased N in the soil leachates, together with long median TTs potentially generate a hydrological N legacy that might become visible at the catchment outlet years to decades later (Van Meter et al.,

2016; Van Meter & Basu, 2015), especially under the assumption of low denitrification potential in groundwater (Nguyen et al., 2022). One indication is high nitrate concentrations in 2019 and 2020 in a groundwater observation well in the lower Selke catchment (Figure S6). Moreover, Jutglar et al. (2021) reported immediate and delayed drought responses in the form of increasing nitrate concentrations in the groundwater in South-West Germany after the drought in 2003. Besides such first evidence, estimating the potential biogeochemical and hydrological nitrogen legacies induced by severe drought will be a task for future research.

## 5 Conclusion

The presented study is among the first to assess the impact of the 2018-2019 drought in Central Europe on nitrate concentrations in a heterogeneous mesoscale catchment. We found that such an exceptional drought can have significant impacts on water quality in terms of nitrate concentrations, load export, and catchment N retention capacity.

C-Q relationships revealed an increased intra-annual variability in nitrate concentrations, with low concentrations during dry periods and exceptionally high concentrations during wet periods, mainly driven by the more responsive upstream part of the catchment. Low concentrations could be explained by a selection preference for old and largely denitrified water, whereas high concentrations reflect a reactivation of shallow and young flow paths that transport accumulated N from the catchment soils. The increased intra-annual variability was not only driven by a stronger temporal shift in N transport but also by a decrease in N uptake by plants and removal via denitrification during dry periods. Thus also, the overall provision of exportable nitrate increased, which was reflected in a decrease in the capacity of the catchment soils to retain N and an increase in nitrate loads at the (sub-)catchment outlets relative to the loads expected from the long-term L-Q relationship. We identified two drivers for a decrease in the catchment N retention capacity i) a decrease with increasing discharge that reduces the relative importance of soil N cycling compared to hydrological transport, or ii) a severe drought that decreases N cycling by drying out the catchment soils. The subsequent transport of increased nitrate leaching from the soil zone to the stream network is dependent on the sub-catchment-specific TTs and the denitrification potential in the groundwater. In catchments with short median TTs, the catchment-retention capacity can decrease within the observation period. Instead, long TTs can dampen such a short-term response but potentially form a hydrological N legacy that might become visible at the catchment outlet years to decades later.

Hotter multi-year droughts are likely to become more frequent and more prolonged with climate change (Hari et al., 2020; IPCC, 2018). Our study shows that such climatic extremes are a threat not only to water quantity but also to water quality in terms of nitrate pollution, as they can reduce the capacity of the catchment soils and entire catchments to retain N. Moreover, such droughts have the potential to override positive effects of measures to improve water quality (e.g., two-stage ditches; Bierozza et al., 2019). Consequently, droughts need to be considered as an additional risk to water quality that can intensify the existing anthropogenic pressures. To counteract the additional risk, one should consider intensified restrictions on manure and fertilizer applications. Furthermore, our study emphasizes the role of catchment heterogeneity and TTs for a catchment's

vulnerability to drought impacts on nitrate export and the timing of such impacts. Whereas fast-reacting sub-catchments with short TTs can contribute to immediate drought responses, slowly-reacting sub-catchments (long TTs) might build up drought-induced N legacies that could impact future water quality on the long term, depending on the subsurface denitrification potential. We could show that a severe drought can potentially amplify such sub-catchment specific differences. The increased variability of nitrate export on both temporal and spatial scales calls for an increased spatiotemporal frequency of water quality monitoring and more site-specific management plans for site-specific problems. This also means that more studies on drought effects on water quality in different catchments and also for other pollutants are needed to assess the additional risk that is posed by longer and hotter droughts.

## 510 **Data availability statement**

The source code and input data for mHM-SAS are available at <https://doi.org/10.5281/zenodo.7228149>. The source code for mHM-nitrate (X. Yang et al., 2018), on which mHM-SAS is build up upon, is available at <https://git.ufz.de/yangx/mHM-Nitrate>. The raw discharge data can be downloaded from the State Office of Flood Protection and Water Quality of Saxony Anhalt (LH) at <https://gld.lhw-sachsen-anhalt.de/>. The raw meteorological data can be downloaded from Germanys National Meteorological Service (DWD) at [https://opendata.dwd.de/climate\\_environment/CDC/grids\\_germany/daily/regnie/](https://opendata.dwd.de/climate_environment/CDC/grids_germany/daily/regnie/), and gridded products based on Zink et al. (2017) are available under <https://www.ufz.de/index.php?en=41160>. Raw nitrate-N concentration data are archived in the TERENO database, available upon request through the TERENO-Portal ([www.tereno.net/ddp](http://www.tereno.net/ddp)).

## **Author contributions**

520 Conceptualization and main investigation, writing of the first draft and visualization of this study was conducted by CW under supervisions of AM, JF and SL. The analysis was carried out by CW and TN. Data was provided by MR and RK. Project administration and funding acquisition was done by JF. All authors reviewed and edited the manuscript.

## **Acknowledgments**

525 This study was funded by the Helmholtz Research Program, Topic 5 Landscapes of the Future, subtopic 5.2 Water resources and the environment, and by the Helmholtz International Research School TRACER (HIRS-0017). A cordial thank goes to the State Office of Flood Protection and Water Quality of Saxony-Anhalt (LHW) and Germany's National Meteorological Service (Deutscher Wetterdienst, DWD) for providing the discharge data and the meteorological data sets. We thank the two anonymous reviewers for their constructive feedback, which helped improve the original manuscript.

## References

- 530 Andersen, C. B., Lewis, G. P., & Sargent, K. A. (2004). Influence of wastewater-treatment effluent on concentrations and  
fluxes of solutes in the Bush River, South Carolina, during extreme drought conditions. *Environmental Geosciences*,  
11(1), 28–41. <https://doi.org/10.1306/eg.10200303017>
- Bieroza, M. Z., Bergström, L., Ulén, B., Djodjic, F., Tonderski, K., Heeb, A., et al. (2019). Hydrologic Extremes and Legacy  
Sources Can Override Efforts to Mitigate Nutrient and Sediment Losses at the Catchment Scale. *Journal of*  
535 *Environmental Quality*, 48(5), 1314–1324. <https://doi.org/10.2134/jeq2019.02.0063>
- Burt, T. P., Worrall, F., Howden, N. J. K., & Anderson, M. G. (2015). Shifts in discharge-concentration relationships as a  
small catchment recover from severe drought. *Hydrological Processes*, 29(4), 498–507.  
<https://doi.org/10.1002/hyp.10169>
- Campbell, C. A., & Biederbeck, V. O. (1982). Changes in mineral N and numbers of bacteria and actinomycetes during two  
540 years under wheat-fallow in Southwestern Saskatchewan. *Canadian Journal of Soil Science*, 62(1), 125–137.  
<https://doi.org/10.4141/cjss82-014>
- Casquin, A., Dupas, R., Gu, S., Couic, E., Gruau, G., & Durand, P. (2021). The influence of landscape spatial configuration  
on nitrogen and phosphorus exports in agricultural catchments. *Landscape Ecology*, 36(12), 3383–3399.  
<https://doi.org/10.1007/s10980-021-01308-5>
- 545 Cramer, M. D., Hawkins, H.-J., & Verboom, G. A. (2009). The importance of nutritional regulation of plant water flux.  
*Oecologia*, 161(1), 15–24. <https://doi.org/10.1007/s00442-009-1364-3>
- Davis, C. A., Ward, A. S., Burgin, A. J., Loecke, T. D., Riveros-Iregui, D. A., Schnoebelen, D. J., et al. (2014). Antecedent  
Moisture Controls on Stream Nitrate Flux in an Agricultural Watershed. *Journal of Environmental Quality*, 43(4),  
1494–1503. <https://doi.org/10.2134/jeq2013.11.0438>
- 550 Delpla, I., Jung, A.-V., Baures, E., Clement, M., & Thomas, O. (2009). Impacts of climate change on surface water quality in  
relation to drinking water production. *Environment International*, 35(8), 1225–1233.  
<https://doi.org/10.1016/j.envint.2009.07.001>
- Dupas, R., Abbott, B. W., Minaudo, C., & Fovet, O. (2019). Distribution of landscape units within catchments influences  
nutrient export dynamics. *Frontiers in Environmental Science*, 7, 43. <https://doi.org/doi:10.3389/fenvs.2019.00043>
- 555 EEA, (European Environment Agency). (2018). Corine Land COVer. European Environment Agency, Copenhagen. Retrieved  
from <https://land.copernicus.eu/pan-european/corine-land-cover>
- Ehrhardt, S., Kumar, R., Fleckenstein, J. H., Attinger, S., & Musolff, A. (2019). Trajectories of nitrate input and output in three  
nested catchments along a land use gradient. *Hydrology and Earth System Sciences*, 23(9), 3503–3524.  
<https://doi.org/doi:0.5194/hess-23-3503-2019>

- 560 Ehrhardt, S., Ebeling, P., Dupas, R., Kumar, R., Fleckenstein, J. H., & Musolff, A. (2021). Nitrate transport and retention in Western European catchments are shaped by hydroclimate and subsurface properties. *Water Resources Research*, 57(10), e2020WR029469. <https://doi.org/10.1029/2020WR029469>
- 565 Fu, Z., Ciais, P., Bastos, A., Stoy, P. C., Yang, H., Green, J. K., et al. (2020). Sensitivity of gross primary productivity to climatic drivers during the summer drought of 2018 in Europe. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375(1810), 20190747. <https://doi.org/10.1098/rstb.2019.0747>
- Górski, J., Dragon, K., & Kaczmarek, P. M. J. (2019). Nitrate pollution in the Warta River (Poland) between 1958 and 2016: trends and causes. *Environmental Science and Pollution Research*, 26(3), 2038–2046. <https://doi.org/10.1007/s11356-017-9798-3>
- 570 Hannappel, S., Köpp, C., & Bach, T. (2018). Charakterisierung des Nitratabbauvermögens der Grundwasserleiter in Sachsen-Anhalt. *Grundwasser*, 23(4), 311–321.
- Hari, V., Rakovec, O., Markonis, Y., Hanel, M., & Kumar, R. (2020). Increased future occurrences of the exceptional 2018–2019 Central European drought under global warming. *Scientific Reports*, 10(1), 1–10. <https://doi.org/10.1038/s41598-020-68872-9>
- 575 Hashimoto, T., Stedinger, J. R., & Loucks, D. P. (1982). Reliability, resiliency, and vulnerability criteria for water resource system performance evaluation. *Water Resources Research*, 18(1), 14–20. <https://doi.org/10.1029/WR018i001p00014>
- Haynes, R. J. (1986). The decomposition process: Mineralization, immobilization, humus formation. *Mineral Nitrogen in the Plant-Soil Systems*, 52–126.
- 580 Hirsch, R. M., Moyer, D. L., & Archfield, S. A. (2010). Weighted regressions on time, discharge, and season (WRTDS), with an application to Chesapeake Bay river inputs 1. *JAWRA Journal of the American Water Resources Association*, 46(5), 857–880. <https://doi.org/doi:10.1111/j.1752-1688.2010.00482.x>
- Huber, C. (2005). Long lasting nitrate leaching after bark beetle attack in the highlands of the Bavarian Forest National Park. *Journal of Environmental Quality*, 34(5), 1772–1779. <https://doi.org/10.2134/jeq2004.0210>
- 585 IPCC. (2018). *Global warming of 1.5° C: an IPCC special report on the impacts of global warming of 1.5° C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty* [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)]. Intergovernmental Panel on Climate Change.
- 590 Jarvie, H. P., Neal, C., Withers, P. J., Robinson, A., & Salter, N. (2003). Nutrient water quality of the Wye catchment, UK: exploring patterns and fluxes using the Environment Agency data archives. *Hydrology and Earth System Sciences*, 7(5), 722–743. <https://doi.org/10.5194/hess-7-722-2003>

- Jawitz, J. W., Desormeaux, A. M., Annable, M. D., Borchardt, D., & Dobberfuhl, D. (2020). Disaggregating landscape-scale nitrogen attenuation along hydrological flow paths. *Journal of Geophysical Research: Biogeosciences*, 125(2), e2019JG005229. <https://doi.org/10.1029/2019JG005229>
- Jeppesen, E., Moss, B., Bennion, H., Carvalho, L., DeMeester, L., Feuchtmayr, H., et al. (2010). Interaction of Climate Change and Eutrophication. In M. Kernan, R. W. Battarbee, & B. Moss (Eds.), *Climate Change Impacts on Freshwater Ecosystems* (pp. 119–151). Oxford, UK: Wiley-Blackwell. <https://doi.org/10.1002/9781444327397.ch6>
- Jiang, S., Jomaa, S., & Rode, M. (2014). Modelling inorganic nitrogen leaching in nested mesoscale catchments in central Germany. *Ecohydrology*, 7(5), 1345–1362. <https://doi.org/doi:10.1002/eco.1462>
- Jomaa, S., Aboud, I., Dupas, R., Yang, X., Rozemeijer, J., & Rode, M. (2018). Improving nitrate load estimates in an agricultural catchment using Event Response Reconstruction. *Environmental Monitoring and Assessment*, 190(6), 330.
- Jutglar, K., Hellwig, J., Stoelzle, M., & Lange, J. (2021). Post-drought increase in regional-scale groundwater nitrate in southwest Germany. *Hydrological Processes*, 35(8), e14307. <https://doi.org/10.1002/hyp.14307>
- Kong, X., Ghaffar, S., Determann, M., Friese, K., Jomaa, S., Mi, C., et al. (2022). Reservoir water quality deterioration due to deforestation emphasizes the indirect effects of global change. *Water Research*. 221, 118721, <https://doi.org/10.1016/j.watres.2022.118721>
- Lange, J., & Haensler, A. (2012). Runoff generation following a prolonged dry period. *Journal of Hydrology*, 464–465, 157–164. <https://doi.org/10.1016/j.jhydrol.2012.07.010>
- Loecke, T. D., Burgin, A. J., Riveros-Iregui, D. A., Ward, A. S., Thomas, S. A., Davis, C. A., & Clair, M. A. St. (2017). Weather whiplash in agricultural regions drives deterioration of water quality. *Biogeochemistry*, 133(1), 7–15. <https://doi.org/10.1007/s10533-017-0315-z>
- Mikkelsen, K. M., Bearup, L. A., Maxwell, R. M., Stednick, J. D., McCray, J. E., & Sharp, J. O. (2013). Bark beetle infestation impacts on nutrient cycling, water quality and interdependent hydrological effects. *Biogeochemistry*, 115(1–3), 1–21. <https://doi.org/10.1007/s10533-013-9875-8>
- Morecroft, M. D., Burt, T. P., Taylor, M. E., & Rowland, A. P. (2000). Effects of the 1995-1997 drought on nitrate leaching in lowland England. *Soil Use and Management*, 16(2), 117–123. <https://doi.org/10.1111/j.1475-2743.2000.tb00186.x>
- Mosley, L. M. (2015). Drought impacts on the water quality of freshwater systems; review and integration. *Earth-Science Reviews*, 140, 203–214. <https://doi.org/10.1016/j.earscirev.2014.11.010>
- Musolff, A., Zhan, Q., Dupas, R., Minaudo, C., Fleckenstein, J. H., Rode, M., et al. (2021). Spatial and Temporal Variability in Concentration-Discharge Relationships at the Event Scale. *Water Resources Research*, 57(10), e2020WR029442. <https://doi.org/10.1029/2020WR029442>
- Musolff, Andreas, Schmidt, C., Selle, B., & Fleckenstein, J. H. (2015). Catchment controls on solute export. *Advances in Water Resources*, 86, 133–146. <https://doi.org/10.1016/j.advwatres.2015.09.026>

- National Institute for Public Health and the Environment (RIVM). (2021). Agricultural practices and water quality in the Netherlands; status (2016-2019) and trend (1992-2019). <https://doi.org/10.21945/RIVM-2020-0184>
- Nguyen, T. V., Kumar, R., Lutz, S. R., Musolff, A., Yang, J., & Fleckenstein, J. H. (2021). Modeling Nitrate Export From a Mesoscale Catchment Using StorAge Selection Functions. *Water Resources Research*, 57(2), e2020WR028490. <https://doi.org/10.1029/2020WR028490>
- Nguyen, T. V., Kumar, R., Musolff, A., Lutz, S. R., Sarrazin, F., Attinger, S., & Fleckenstein, J. H. (2022). Disparate Seasonal Nitrate Export From Nested Heterogeneous Subcatchments Revealed With StorAge Selection Functions. *Water Resources Research*, 58(3), e2021WR030797. <https://doi.org/10.1029/2021WR030797>
- Nogueira, G. E. H., Schmidt, C., Brunner, P., Graeber, D., & Fleckenstein, J. H. (2021). Transit-Time and Temperature Control the Spatial Patterns of Aerobic Respiration and Denitrification in the Riparian Zone. *Water Resources Research*, 57(12), e2021WR030117. <https://doi.org/10.1029/2021WR030117>
- Oelsner, G. P., Brooks, P. D., & Hogan, J. F. (2007). Nitrogen Sources and Sinks Within the Middle Rio Grande, New Mexico. *JAWRA Journal of the American Water Resources Association*, 43(4), 850–863. <https://doi.org/10.1111/j.1752-1688.2007.00071.x>
- Outram, F. N., Lloyd, C. E. M., Jonczyk, J., Benskin, C. M. H., Grant, F., Perks, M. T., et al. (2014). High-frequency monitoring of nitrogen and phosphorus response in three rural catchments to the end of the 2011–2012 drought in England. *Hydrology and Earth System Sciences*, 18(9), 3429–3448. <https://doi.org/10.5194/hess-18-3429-2014>
- Rakovec, O., Samaniego, L., Hari, V., Markonis, Y., Moravec, V., Thober, S., et al. (2022). The 2018–2020 Multi-Year Drought Sets a New Benchmark in Europe. *Earth's Future*, 10(3), e2021EF002394. <https://doi.org/10.1029/2021EF002394>
- Riediger, J., Breckling, B., Nuske, R. S., & Schröder, W. (2014). Will climate change increase irrigation requirements in agriculture of Central Europe? A simulation study for Northern Germany. *Environmental Sciences Europe*, 26(1), 18. <https://doi.org/10.1186/s12302-014-0018-1>
- Rivett, M. O., Buss, S. R., Morgan, P., Smith, J. W., & Bemment, C. D. (2008). Nitrate attenuation in groundwater: a review of biogeochemical controlling processes. *Water Research*, 42(16), 4215–4232.
- Rode, M., Halbedel née Angelstein, S., Anis, M. R., Borchardt, D., & Weitere, M. (2016). Continuous in-stream assimilatory nitrate uptake from high-frequency sensor measurements. *Environmental Science & Technology*, 50(11), 5685–5694. <https://doi.org/10.1021/acs.est.6b00943>
- Samaniego, L., Thober, S., Kumar, R., Wanders, N., Rakovec, O., Pan, M., et al. (2018). Anthropogenic warming exacerbates European soil moisture droughts. *Nature Climate Change*, 8(5), 421–426. <https://doi.org/10.1038/s41558-018-0138-5>
- Sapkota, A., Haghverdi, A., Avila, C. C. E., & Ying, S. C. (2020). Irrigation and Greenhouse Gas Emissions: A Review of Field-Based Studies. *Soil Systems*, 4(2), 20. <https://doi.org/10.3390/soilsystems4020020>
- Schaeffli, B., & Gupta, H. V. (2007). Do Nash values have value? *Hydrological Processes*, 21(ARTICLE), 2075–2080.

- 660 Schnabel, F., Purruicker, S., Schmitt, L., Engelmann, R. A., Kahl, A., Richter, R., et al. (2021). *Cumulative growth and stress responses to the 2018–2019 drought in a European floodplain forest* (p. 2021.03.05.434090). <https://doi.org/10.1101/2021.03.05.434090>
- Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., et al. (2020). A first assessment of the impact of the extreme 2018 summer drought on Central European forests. *Basic and Applied Ecology*, 45, 86–103. <https://doi.org/10.1016/j.baae.2020.04.003>
- 665 Seeger, S., & Weiler, M. (2014). Reevaluation of transit time distributions, mean transit times and their relation to catchment topography. *Hydrology and Earth System Sciences*, 18(12), 4751–4771. <https://doi.org/10.5194/hess-18-4751-2014>
- Sinha, S., Rode, M., & Borchardt, D. (2016). Examining runoff generation processes in the Selke catchment in central Germany: Insights from data and semi-distributed numerical model. *Journal of Hydrology: Regional Studies*, 7, 38–
- 670 54. <https://doi.org/10.1016/j.ejrh.2016.06.002>
- Sprague, L. A. (2005). Drought Effects on Water Quality in the South Platte River Basin, Colorado1. *JAWRA Journal of the American Water Resources Association*, 41(1), 11–24. <https://doi.org/10.1111/j.1752-1688.2005.tb03713.x>
- Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L., Van Lanen, H., Sauquet, E., et al. (2010). Streamflow trends in Europe: evidence from a dataset of near-natural catchments. <https://doi.org/10.5194/hess-14-2367-2010>
- 675 Tetzlaff, D., Seibert, J., McGuire, K. J., Laudon, H., Burns, D. A., Dunn, S. M., & Soulsby, C. (2009). How does landscape structure influence catchment transit time across different geomorphic provinces? *Hydrological Processes*, 23(6), 945–953. <https://doi.org/10.1002/hyp.7240>
- Thompson, S. E., Basu, N. B., Lascurain, J., Aubeneau, A., & Rao, P. S. C. (2011). Relative dominance of hydrologic versus biogeochemical factors on solute export across impact gradients. *Water Resources Research*, 47(10). <https://doi.org/10.1029/2010WR009605>
- 680 Van Der Velde, Y., Torfs, P., Van Der Zee, S., & Uijlenhoet, R. (2012). Quantifying catchment-scale mixing and its effect on time-varying travel time distributions. *Water Resources Research*, 48(6). <https://doi.org/10.1029/2011WR011310>
- Van Meter, K. J., & Basu, N. B. (2015). Catchment legacies and time lags: A parsimonious watershed model to predict the effects of legacy storage on nitrogen export. *PLoS One*, 10(5), e0125971. <https://doi.org/10.1371/journal.pone.0125971>
- 685 Van Meter, K. J., Basu, N. B., Veenstra, J. J., & Burras, C. L. (2016). The nitrogen legacy: emerging evidence of nitrogen accumulation in anthropogenic landscapes. *Environmental Research Letters*, 11(3), 035014. <https://doi.org/10.1088/1748-9326/11/3/035014>
- Van Vliet, M. T. H., & Zwolsman, J. J. G. (2008). Impact of summer droughts on the water quality of the Meuse river. *Journal of Hydrology*, 353(1), 1–17. <https://doi.org/10.1016/j.jhydrol.2008.01.001>
- 690 Whitehead, P. G., Wilby, R. L., Battarbee, R. W., Kernan, M., & Wade, A. J. (2009). A review of the potential impacts of climate change on surface water quality. *Hydrological Sciences Journal*, 54(1), 101–123. <https://doi.org/10.1623/hysj.54.1.101>

- 695 Winter, C., Lutz, S. R., Musolff, A., Kumar, R., Weber, M., & Fleckenstein, J. H. (2021). Disentangling the impact of  
catchment heterogeneity on nitrate export dynamics from event to long-term time scales. *Water Resources Research*,  
57(1), e2020WR027992. <https://doi.org/10.1029/2020WR027992>
- Winter, C., Tarasova, L., Lutz, S. R., Musolff, A., Kumar, R., & Fleckenstein, J. H. (2022). Explaining the Variability in High-  
Frequency Nitrate Export Patterns Using Long-Term Hydrological Event Classification. *Water Resources Research*,  
58(1), e2021WR030938. <https://doi.org/10.1029/2021WR030938>
- 700 Wollschläger, U., Attinger, S., Borchardt, D., Brauns, M., Cuntz, M., Dietrich, P., et al. (2017). The Bode hydrological  
observatory: a platform for integrated, interdisciplinary hydro-ecological research within the TERENO Harz/Central  
German Lowland Observatory. *Environmental Earth Sciences*, 76(1), 29. <https://doi.org/10.1007/s12665-016-6327-5>
- Yang, J., Heidbüchel, I., Musolff, A., Reinstorf, F., & Fleckenstein, J. H. (2018). Exploring the dynamics of transit times and  
705 subsurface mixing in a small agricultural catchment. *Water Resources Research*, 54(3), 2317–2335.  
<https://doi.org/10.1002/2017WR021896>
- Yang, X., Jomaa, S., Zink, M., Fleckenstein, J. H., Borchardt, D., & Rode, M. (2018). A New Fully Distributed Model of  
Nitrate Transport and Removal at Catchment Scale. *Water Resources Research*, 54(8), 5856–5877.  
<https://doi.org/10.1029/2017WR022380>
- 710 Yang, X., Tetzlaff, D., Soulsby, C., Smith, A., & Borchardt, D. (2021). Catchment Functioning Under Prolonged Drought  
Stress: Tracer-Aided Ecohydrological Modeling in an Intensively Managed Agricultural Catchment. *Water Resources  
Research*, 57(3), e2020WR029094. <https://doi.org/10.1029/2020WR029094>