

Satellite-derived Constraints on the Effect of Drought Stress on Biogenic Isoprene Emissions in the Southeast US

Yuxuan Wang¹, Nan Lin¹, Wei Li¹, Alex Guenther², Joey C. Y. Lam³, and Amos P. K. Tai^{3,4}, Mark J. Potosnak⁵, Roger Seco⁶

¹ Department of Earth and Atmospheric Sciences, University of Houston, Houston, Texas, USA.

² Earth System Science, University of California, Irvine, Irvine, California, USA

³ Earth and Environmental Sciences Programme, Faculty of Science, The Chinese University of Hong Kong, Hong Kong SAR, China

⁴ State Key Laboratory of Agrobiotechnology and Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong, Hong Kong SAR, China

⁵ Environmental Science and Studies, DePaul University, Chicago, IL, USA

⁶ Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Carrer Jordi Girona 18-26, 08034 Barcelona, Spain

Corresponding author: Yuxuan Wang (ywang246@central.uh.edu)

Abstract. While substantial progress has been made to improve our understanding of biogenic isoprene emissions under unstressed conditions, there remain large uncertainties in isoprene emissions under stressed conditions. Here we use the US Drought Monitor (USDM) as a weekly drought severity index and tropospheric columns of formaldehyde (HCHO), the key product of isoprene oxidation, retrieved from the Ozone Monitoring Instrument (OMI) to derive top-down constraints on the response of summertime isoprene emissions to drought stress in the Southeast U.S. (SE US), a region of high isoprene emissions and prone to drought. OMI HCHO column density is found to be 6.7% (mild drought) - 23.3% (severe drought) higher than that in no-drought conditions. A global chemical transport model, GEOS-Chem, with the MEGAN2.1 emission algorithm can simulate this direction of change, but the simulated increases at the corresponding drought levels are 1.1-1.5 times of OMI HCHO, suggesting the need for a drought-stress algorithm in the model. By minimizing the model-to-OMI differences in HCHO to temperature sensitivity under different drought levels, we derived a top-down drought stress factor (γ_{d_OMI}) in GEOS-Chem that parameterizes using water stress and temperature. The algorithm led to an 8.6% (mild drought) - 20.7% (severe drought) reduction in isoprene emissions in the SE US relative to the simulation without it. With γ_{d_OMI} the model predicts a non-linear increasing trend in isoprene emissions with drought severity that is consistent with OMI HCHO and a single site's isoprene flux measurements. Compared with a previous drought stress algorithm derived from the latter, the satellite-based drought stress factor performs better in capturing the regional scale drought-isoprene responses as indicated by the close-to-zero mean bias between OMI and simulated HCHO columns under different drought conditions. The drought stress algorithm also reduces the model's high bias in organic aerosols (OA) simulations by 6.60% (mild drought) to 11.71% (severe drought) over the SE US compared to the no-stress simulation. The simulated ozone response to the drought stress factor displays a spatial disparity due to the isoprene suppressing effect on oxidants, with an <1 ppb increase in O₃ in high-isoprene regions and a 1-3 ppbv decrease in O₃ in low-isoprene regions. This

study demonstrates the unique value of exploiting long-term satellite observations to develop empirical stress algorithms on biogenic emissions where in situ flux measurements are limited.

1. Introduction

Biogenic nonmethane volatile organic compounds (BVOCs) emitted by terrestrial ecosystems are of great importance to air quality, tropospheric chemistry, and climate due to their effects on atmospheric oxidants and aerosols (Atkinson, 2000; Claeys et al., 2004; Pacifico et al., 2009). The dominant BVOC is isoprene ($\text{CH}_2=\text{C}(\text{CH}_3)\text{CH}=\text{CH}_2$), comprising 70% of the global total BVOC emitted from vegetation (Sindelarova et al., 2014). Isoprene emissions depend on vegetation/plant type, physiological status, leaf age, and meteorological conditions such as radiation, temperature, and soil moisture. These relationships provide the basic framework of isoprene emission models that are capable of coupling with meteorology and the land biosphere, the most widely used being the Model of Emissions of Gases and Aerosols from Nature (MEGAN) (Guenther et al., 1993, 2006, 2012, 2017). Recent work has shown stressed conditions - such as drought, heatwaves, and high winds - can induce large changes in isoprene emissions different from model predictions in the absence of those stress factors (Potosnak et al., 2014; Huang et al., 2015; Kravitz et al., 2016; Seco et al., 2015; Otu-Larbi et al., 2020; Seco et al., 2022). As stressed conditions are rarely sampled by field campaigns due to their infrequent and irregular nature and hence poorly constrained, stress impacts on isoprene emissions are among the least understood aspects in our predictivity of BVOC-chemistry-climate interactions.

A common stress for terrestrial vegetation worldwide is drought, characterized by low precipitation, high temperature, and low soil moisture (Trenberth et al., 2014). These conditions are primary abiotic stresses that will cause physiological impacts on plants affecting photosynthesis, stomatal conductance, transpiration, and leaf area. During short-term or mild droughts, the photosynthetic rate of plants quickly decreases due to limited stomatal conductance, while isoprene is not immediately impacted because of the availability of stored carbon and because the photosynthetic electron transport is not inhibited. Isoprene can even increase by several factors due to warm leaf temperatures which increases isoprene synthase activity (Potosnak et al., 2014; Ferracci et al., 2020). During prolonged or severe drought stress, after a lag related to photosynthesis reduction, isoprene emission eventually declines because of inadequate carbon availability. This conceptualized non-monotonic response of isoprene emission to drought has been demonstrated at the Missouri Ozarks AmeriFlux (MOFLUX) field site in Missouri (Potosnak et al., 2014; Seco et al., 2015), the only available drought-relevant whole canopy isoprene flux measurements to date, and qualitatively supported by ambient isoprene concentrations monitored by regional surface networks (Wang et al., 2017). It is noteworthy that the MOFLUX data covered only two drought events (summer 2011 and summer 2012), while the surface sites are sparsely distributed with an urban focus. More recently, the isoprene concentration measurements during the Wytham Isoprene iDirac Oak Tree Measurements (WiSDOM) campaign showed that isoprene was up to four times higher than normal in responses to a combined heatwave and drought episode (June-October 2018) over a mid-latitude temperate forest in the UK (Ferracci et al., 2020; Otu-Larbi et al., 2020), which supports the enhanced isoprene emissions at the MOFLUX site under mild droughts. However, these observations offer only limited constraints on drought stress impacts on isoprene emissions.

With wide spatiotemporal coverage, satellite provides arguably the best platform to capture drought development and impacts. Satellite observations of tropospheric formaldehyde (HCHO) columns have been used as a proxy of isoprene emissions for more than a decade (Abbot et al., 2003; Palmer et al., 2003), as HCHO is formed promptly and in high yield from isoprene oxidation (Sprengnether et al., 2002). Previous applications of satellite HCHO products provided “top-down” estimates on seasonality, magnitude, spatial distribution, and interannual variability of isoprene emissions globally and regionally (e.g., Marais et al., 2016; Kaiser et al., 2018; Stavrakou et al., 2018). While most of these studies focused on *unstressed* conditions, recent efforts have shown that satellite HCHO registered drought signals on a monthly scale (Zheng et al., 2017; Naimark et al., 2021; Li et al., 2022; Opacka et al., 2022). These signals are yet to be exploited to constrain isoprene response to drought.

The present study aims at improving the current quantification of satellite HCHO response to drought by accounting for sub-monthly variability of drought severity. We use a weekly time scale, the finest temporal scale of drought indices available, and separate five levels of drought severity defined by the US Drought Monitor. By comparison, previous investigations used binary classification (drought or not) on a monthly time scale. Our improvement in scale is expected to better capture the nonlinear response of isoprene emissions to drought severity as described above. The study region is the Southeast United States (SE US), which has large isoprene emissions due to substantial forest coverage and is also prone to drought due to large interannual variability in precipitation (Seager et al., 2009). In addition, the MOFLUX site is located in the SE US, which will allow us to evaluate if satellite-derived drought responses of HCHO are consistent with those from isoprene flux measurements at MOFLUX. Finally, we use these HCHO signals in conjunction with models to identify the model gaps in predicting isoprene responses to drought.

2. Data and Method

2.1 Drought index

There are many types of drought indices focusing on different factors, including precipitation, temperature, evaporation, runoff, and the impact of drought on ecosystems and vegetation (Palmer, 1965; McKee et al., 1993; Guttman, 1999; Vicente-Serrano et al., 2010; Chang et al., 2018). Drought indices also differ by time scale. As drought by definition is a prolonged period of water deficit, the shortest time scale of drought is weekly. Here we chose the United States Drought Monitor (USDM) drought index to identify drought periods. USDM’s weekly timescale and multiple drought severity levels (Svoboda et al., 2002) provide a better delineation of drought variability than the monthly or seasonal scale used in the previous analysis of drought signals in HCHO and isoprene (Wang et al., 2017; Naimark et al., 2021).

(a) USDM released on July 12,2012 (b) USDM time series for SE US




Figure 1. (a) Drought distribution for the second week of July 2012 based on USDM. The black star indicates the location of MOFLUX site. (b) Time series of drought frequency in the study area (black box in Figure 1a) for JJA from 2005 to 2017. N0 (white) for wet and normal, D0 (light yellow) for abnormal drought, D1 (yellow) for moderate drought, D2 (orange) for severe drought, D3 (red) for extreme drought, and D4 (brown) for exceptional drought.

The USDM is a composite drought index based on six key physical indicators including the Palmer Drought Severity Index (PDSI, Palmer, 1965), CPC Soil Moisture Model Percentiles (Huang et al., 1996), U.S. Geological Survey (USGS) Daily Streamflow Percentiles (<http://water.usgs.gov.waterwatch/>), Percent of Normal Precipitation (Willeke et al., 1994), Standardized Precipitation Index (SPI, McKee et al., 1993), and remotely sensed Satellite Vegetation Health Index (Kogan, 1995). Opinions of local experts are also considered (Svoboda et al., 2002). The USDM website (<https://droughtmonitor.unl.edu/>) provides weekly ArcGIS shapefiles of the polygons covering the whole US under five drought levels: D0 for abnormal drought, D1 for moderate drought, D2 for severe drought, D3 for extreme drought, and D4 for exceptional drought. We used the method of Chen et al. (2019) to rasterize and convert USDM shapefiles to $0.5^\circ \times 0.5^\circ$ gridded indices with -1 indicating non-drought (N0) and 0-4 for D0-D4 drought, respectively. **Figure 1a** displays the spatial distribution of gridded USDM indices for the second week of July 2012, which clearly depicts the extent and severity of the infamous 2012 Great Plains drought (Hoerling et al., 2014). **Figure 1b** shows the weekly time series of USDM indices averaged over SE US ($75\text{--}100^\circ\text{W}$, $25\text{--}40^\circ\text{N}$, black box in Figure 1a) for the summer months (June, July, August; JJA) of 2005 -2017, our study period. During this period, abnormal drought (D0) appeared every summer, while extreme and exceptional drought (D3-D4) were mainly concentrated in 2006-2008 and 2010-2012. This pattern is consistent with the long-term drought statistics from other drought indices such as SPEI and PDSI (Svoboda et al., 2015).

2.2 OMI HCHO and NO₂ product

We used the Ozone Monitoring Instrument (OMI) v003 level 3 tropospheric formaldehyde (HCHO) column density (OMHCHOd) as described by Chance (2019). OMI was launched on NASA's Aura satellite in 2004 and has since provided daily global measurements of ozone (O₃) and its precursors with a nadir spatial resolution of $24 \times 13 \text{ km}^2$. Since January 2009, OMI has been suffering from a major row anomaly. OMHCHOd data processing explored all level 2 OMHCHO observations to filter out pixels with bad formaldehyde retrievals, high cloud fractions ($>30\%$), high SZA ($>70^\circ$), and pixels affected by OMI's row anomaly (Chance, 2019). The spatial resolution is $0.1^\circ \times 0.1^\circ$. Zhu et al. (2016) verified the OMHCHOd data using high-precision HCHO aircraft observations obtained during

NASA SEAC4RS activities in SE US from August to September 2013. They showed that OMI retrievals have accurate spatial and temporal distribution but were biased low by 37% relative to the aircraft. We corrected this underestimation by applying a uniform and constant factor of 1.5 to the OMHCHOd data, as did by Shen et al. (2019) in their long-term analysis of OMI HCHO. **Figure 2a** presents the corrected OMHCHOd for the SE US averaged over JJA 2005–2017, where higher levels of HCHO are clearly seen over forested regions in Missouri, Georgia, Arkansas, and Texas. OMHCHOd values shown hereafter are those with the correction factor applied. Although it is not known if the correction factor has temporal spatial variations during our study period, its application produced a good match between OMI and simulated HCHO columns under non-drought (N0) conditions (Figure 2c). To examine the concurrent changes of nitrogen oxides ($\text{NO}_x = \text{NO}_2 + \text{NO}$) under droughts, we also used the level 3 tropospheric column of NO_2 from OMI during the same period (Nickolay et al., 2019).




Figure 2. Mean 2005–2017 HCHO columns for June – August over the SE US of (a) OMI observation (OMHCHOd) and (b) GEOS-Chem simulation (GCHCHO_NoStress). (c) Scatterplot of spatial correlation between the two. The dashed line indicates the 1:1 agreement.

2.3 GEOS-Chem chemical transport model

We used the long-term simulation of the nested-grid GEOS-Chem global chemical transport model (version 12-02, <http://www.geos-chem.org>) to obtain daily mean results of modeled formaldehyde columns and isoprene emissions for North America during JJA 2005 – 2017. The simulation was driven by the Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2) meteorological data from NASA’s Global Modeling and Assimilation Office (GMAO) with a horizontal resolution at $0.5^\circ \times 0.625^\circ$. Biogenic emissions were calculated using MEGAN2.1, which is the prevailing version of MEGAN implemented in most chemical and climate models. MEGAN2.1 has a soil dependence algorithm whose parameterization is based on plant wilting point threshold and soil moisture (Guenther et al., 2017). However, this factor is disabled in GEOS-Chem as in many other CTMs due to the unavailability of the required driving variables, such as wilting point and soil moisture, which cannot be simulated well in most models (Trugman et al., 2018). Thus, outputs from the standard GEOS-Chem simulations do not have

drought effects on isoprene emissions and these outputs are referred to as NoStress_GC. Anthropogenic emissions over North America were from the 2011 National Emissions Inventory (NEI2011, <http://www.epa.gov/air-emissions-inventories>) for the United States, with historical scale factors applied to each simulated year. Open fire emissions were from GFED4 (Giglio et al., 2013) for 2005–2017.

To better match with OMI overpassing time, model HCHO outputs at 13:30 local time were sampled (GCHCHO_NoStress). **Figure 2b** shows GCHCHO_NoStress averaged over the same domain and period as OMHCHOd in **Figure 2a**. The scatter plot (**Figure 2c**) shows a good spatial correlation between the two ($R^2 = 0.88$). This correlation is consistent with other studies comparing GEOS-Chem and OMI HCHO columns in SE US during non-drought periods (Kaiser et al., 2018).

2.4 Observations of ozone, organic aerosol, LAI, and isoprene flux

To evaluate how the drought stress factor changes the simulations of surface O_3 and organic aerosol (OA), we adopted the gridded ($1^\circ \times 1^\circ$) hourly O_3 observations created by Schnell et al. (2014) using the modified inverse distance weighting method. The dataset aggregates several networks of O_3 measurements including the US Environmental Protection Agency's (EPA) Air Quality System (AQS), Clean Air Status and Trends Network (CASTNET), and Environment Canada's National Air Pollution Surveillance Program (NAPS). Following the same method, we created a gridded organic aerosol (OA) dataset using the organic carbon (OC) observations from the Interagency Monitoring of Protected Visual Environments (IMPROVE) network. A factor of 2.1 was used to convert OC to OA as suggested by other studies (Pye et al., 2017; Schroder et al., 2018). To examine the changes of leaf area index (LAI) under droughts, the MODerate resolution Imaging Spectroradiometer (MODIS) Collection 5 LAI products reprocessed by Yuan et al. (2011) with a resolution of $0.25^\circ \times 0.25^\circ$ was used. These three datasets were further remapped through bilinear interpolation to match the spatial resolution of the USDM. The isoprene flux measurements at the MOFLUX site during 2012 May–September were used to derive a site-based drought stress algorithm. The site is located in the Ozarks region of central Missouri ($38.74^\circ N$, $92.20^\circ W$, black star in Figure 1a). It is surrounded by a deciduous forest dominated by isoprene-emitting white and red oak species. The dataset is widely used to investigate isoprene emissions response to droughts (Potosnak et al., 2014; Seco et al., 2015; Jiang et al., 2018; Opacka et al., 2022).

3. Observational Evidence of Drought Stress on Isoprene Emissions

3.1 Changes of HCHO column densities with drought

To reveal drought responses of HCHO, we sampled weekly-mean HCHO columns onto the gridded spatial and temporal locations of each USDM category and generated average HCHO distributions at each drought level over the SE US. The outputs are shown in **Figure 3a** for OMI and **3b** for NoStress_GC, respectively. The processing of weekly-mean HCHO data corresponds to the timing of USDM: a whole week includes Wednesday of the previous week to Tuesday of the present week. There are 12 consecutive weeks from June to August in each year of 2005–2017, giving a total of 156 weeks' gridded HCHO data to be assigned to individual USDM categories by week and location. **Figure 3d** shows the number of weeks underlying the gridded averages of HCHO for each USDM category. As severe

droughts are less frequent than mild droughts, some locations in SE US did not experience D2-D4 droughts during the study period and hence are shown as white in **Figure 3**.




Figure 3. The mean spatial distributions of (a) OMI HCHO column density; (b) NoStress_GC HCHO column density, (c) NoStress_GC isoprene emissions, and (d) the number of weeks during JJA 2005 to 2017 in the southeast US under different USDm drought levels (N0, D0-D4).

OMI HCHO column density increases with increasing drought severity in almost all locations in the SE US (**Figure 3a**). Relative to no-drought condition (N0), the mean HCHO column from OMI is 6.7%, 12.6%, 16.5%, 21.2%, and 23.2% higher under D0 - D4 drought in the entire SE US, respectively. These HCHO changes are statistically significant at a 95% confident interval, indicating that the OMI HCHO products contain significant drought signals. The increasing rate of OMI HCHO with USDM is not linear, faster under mild droughts (D0-D2) and flattening under more severe droughts (D2-D4). This is qualitatively consistent with the conceptualized model of the nonlinear response of isoprene emissions to drought described before (Potosnak et al., 2014).

Model HCHO column density also increases with increasing drought severity (**Figure 3b**). GCHCHO_NoStress is 9.90%, 15.1%, 19.5%, 21.8%, and 29.1% higher under D0-D4 drought than that of N0, respectively. These increases are 1.1-1.5 times those of OMI under all drought levels. The model comparison against OMI HCHO also changes with drought severity. GCHCHO_NoStress has a minimal bias (0.05×10^{16} molec cm^{-2}) under N0. As drought severity increases, the mean bias over the entire SE US increases to 0.10×10^{16} molec cm^{-2} , 0.09×10^{16} molec cm^{-2} , 0.11×10^{16} molec cm^{-2} , 0.08×10^{16} molec cm^{-2} , and 0.15×10^{16} molec cm^{-2} under D0 - D4 levels, respectively. The spatial correlation between OMI and NoStress_GC degrades with USDM, with R^2 being smaller than 0.65 under D0 - D4 levels compared to R^2 of 0.70 under N0. Worsening model performance with increasing drought severity suggests the model lacks a process that changes with drought. As isoprene accounts for more than 80% of the contribution of non-methane VOCs to the HCHO column in the southeast US (Palmer et al., 2003; Millet et al., 2006), the missing process is most likely drought-induced changes in isoprene emissions.




Figure 4. (a) Relative changes of regional-mean OMI HCHO column, NoStress_GC simulated HCHO column, isoprene emissions, anthropogenic benzene emission, anthropogenic toluene emission, and MODIS leaf area index (LAI) under different drought levels in the southeast US. All data are scaled to their respective values at N0. The dotted lines are the arithmetic mean of all grids, and the solid lines are the corrected mean excluding the missing area. (b) Regional-mean tropospheric NO₂ columns from OMI and NoStress_GC (solid lines), and their respective changes from non-drought (N0) conditions (dashed lines). Note the different scales between the solid and dashed lines. The calculation is based on the grids with the presence of all USDM levels.

Figure 4a displays the relative changes in the regional mean HCHO column from OMI and NoStress_GC, emissions of isoprene and select anthropogenic VOCs from NoStress_GC, and MODIS LAI as a function of USDM indices, each scaled by its respective value at N0. The dotted line is the arithmetic mean of all available grids under each dryness category, and the solid line is the mean for those grids with valid data in all dryness categories (i.e., removing

white areas shown in Figure 3). In either calculation, NoStress_GC overestimates the relative increase of HCHO under D0-D4 by 10-50% compared to OMI. After correcting for no data areas at D2-D4, isoprene emissions in NoStress_GC are 22.7%, 29.6%, 40.3%, 54.5%, and 56.0% higher in D0-D4 than N0. Note that LAI is observed to decrease by 5-10% per USDM level (**Figure 4a**), which makes the predicted increase of isoprene emissions with drought severity even more remarkable. This is likely caused by the increasingly higher temperature under droughts, given the exponential relationship of isoprene emissions with temperatures in MEGAN (Guenther et al., 2006).

By comparison, the modeled increase of the HCHO column with drought is 12-25%, more buffered than that of isoprene emissions. This is mainly caused by the loss of HCHO to photolysis, which is expected to increase under droughts with clearer skies (Wang et al., 2017; Naimark et al., 2021). In addition, HCHO formation also depends on the abundance of oxidants, such as hydroxyl radicals (OH) and NO_x, that oxidize isoprene. High isoprene emissions can suppress OH under the low-NO_x conditions that prevail in part of the SE US (Wells et al., 2020), leading to the buffered response in HCHO. Previous studies (Travis et al., 2016; Kaiser et al., 2018) showed that the NEI2011 anthropogenic inventory in the model were biased high in the SE US and a reduction of 60% of NO_x emission was suggested. By comparing to OMI NO₂ column, we found NoStress_GC indeed overestimates NO₂ columns by ~42% in the SE US (**Figure 4b**), but the absolute bias in NO₂ is nearly constant from N0 to D4 (solid lines in Figure 4b). NO₂ column also shows an increasing trend from N0 to D4, yet with a much smaller rate (less than 9%) than HCHO. The model captures the relative change in NO₂ column with USDM (dashed lines in Figure 4b), despite the high bias due to the NEI2011 inventory, which indicates that the changes in natural sources of NO_x (e.g., biomass burning and soil NO_x) with droughts are well represented by NoStress_GC. To further examine the effect of high bias of NO_x on simulated HCHO, we conducted a sensitivity simulation of reducing the NEI2011 NO_x emissions by 50% over the SE US during JJA 2011-2013. Most of the SE US was under droughts during the summertime of 2011-2012, while 2013 was a less drought-stricken year (**Figure 1**). The sensitivity simulation resulted in a small reduction of the simulated HCHO column and the change was nearly constant among the USDM levels (**Figure S1a-b**), ranging from -0.04×10^{16} molec cm⁻² (2.6%) to -0.05×10^{16} molec cm⁻² (3.5%). This rules out the possibility that the high NO_x bias in the model is the reason for the overestimation of HCHO under droughts. Given the suppression effect of isoprene on OH and the well-captured NO₂ relative changes under droughts, the overestimation of HCHO columns by the model is unlikely to be caused by model chemistry, and more likely by the overestimation of isoprene emissions under drought conditions.

While oxidation of anthropogenic VOCs also produces HCHO, using benzene and toluene as indicator species, we found no change in anthropogenic VOC emissions with drought in the model (**Figure 4a**). This insensitivity rules out anthropogenic VOCs as a key driver of model overestimation of HCHO under drought conditions. If anything, we expect anthropogenic VOC emissions to increase during drought due to higher evaporative emissions driven by higher temperature and more fossil fuel consumption driven by more demand for space cooling. Wildfires are another important source that can lead to high HCHO levels, but their contributions to HCHO are more likely to be underpredicted in GEOS-Chem partly due to insufficient hydrocarbon emissions and the underrepresented fire plume chemistry (Alvarado et al., 2020; Liao et al., 2021; Zhao et al., 2022). A deeper planetary boundary layer (PBL) is

expected under droughts primarily caused by a larger sensible height flux released from dry soil (Miralles et al., 2014). Indeed, the MERRA-2 PBL height used in our simulation increases by 12.42%, 17.79%, 20.99%, 26.21%, and 29.52% from D0 to D4 relative to the value of 1589 m at N0 in the SE US during the midday (13:30 LT). Considering the PBL heights in MERRA-2 agree well with observations with only an overall 200 m low bias (Guo et al., 2021), we do not expect mixing heights to be the main cause of the high bias of HCHO column under drought conditions. To further quantify the effects of wildfires and PBL on the changes of HCHO column with drought, we conducted two additional sensitivity tests: (1) turning off the GFED4 wildfire emission inventory during 2011-2013 JJA, and (2) keeping PBL constant as in 2013 (normal year) during 2011-2012 (drought years) JJA. The results in **Figure S1c-d** show overall negligible changes in HCHO column in the SE US, which verifies our assumptions above.

In summary, the model overestimates HCHO increases during drought as compared to OMI. This overestimation is attributed to the model overestimation of isoprene emissions during drought. Drought stress effect on isoprene emissions is thus required in GEOS-Chem to resolve the discrepancy in HCHO responses to drought between OMI and the model.

3.2 Isoprene flux measurement

To further evaluate isoprene emissions in NoStress_GC, we compared the isoprene flux measurements at the MOFLUX site (Potosnak et al., 2014; Seco et al., 2015) with predicted isoprene emissions at the model grid that contains the site. At the time of writing, the MOFLUX site is the only long-term, canopy-level, biogenic isoprene flux measurement site in the Northern midlatitude that sampled droughts. The site experienced multiple drought levels in the summer of 2012, which allows for the model-observation comparison across different drought severities as shown in **Figure 5**. The abnormal dry conditions (D0) started in early June, which developed to moderate drought (D1) in late June, worsened to severe drought (D2) and extreme drought (D3) in July-August, and bounced back to D2 in September (**Figure 5a**). The model generally captures the daily variability of isoprene emissions with a statistically significant correlation coefficient (R) of 0.67, but its biases differ by USDM levels. The model underestimates isoprene flux from N0 (bias of $-1.81 \text{ mg/m}^2/\text{hr}$) to D1 (bias of $-2.89 \text{ mg/m}^2/\text{hr}$), has a minimal bias ($-0.47 \text{ mg/m}^2/\text{hr}$) at D2, and changes to an overestimate at D3 (bias of $1.2 \text{ mg/m}^2/\text{hr}$) (**Figure 5b**). While differences are expected when comparing a single-point flux measurement with the grid-mean model prediction, such differences most likely result in a systematic bias that should not relate to the temporal variability of drought. The fact that the model bias changes from being underpredicting to overpredicting as drought severity increases further confirms the importance of the model lack of a drought suppression effect on isoprene emissions during severe to exceptional droughts (D3 and D4). This is qualitatively consistent with that of the HCHO biases described above.




Figure 5. (a) Comparison of daily time series of isoprene emissions observed at the MOFLUX site (OBS) and simulated by MEGAN2.1 in GEOS-Chem (GC). The background is color-coded according to the USDM drought severity. R and MB at the upright corner show the correlation coefficient and mean bias, respectively. (b) Boxplot of isoprene emissions separated by USDM drought levels. The upper and lower whiskers represent the 90% and 10% quantiles, respectively.

4. Drought Stress Algorithm

The MEGAN2.1 isoprene emission routines in GEOS-Chem use a simplified mechanistic representation of the major environmental factors controlling biogenic emissions (Guenther et al., 2012), in which the isoprene emission factor $\gamma_{2.1}$ is the product of a canopy-related normalization factor (C_{FAC}) multiplied by other factors representing light (γ_{PAR}), temperature (γ_T), leaf age (γ_{AGE}), LAI (γ_{LAI}), carbon dioxide (CO_2) inhibition (γ_{CO2}), and soil moisture (γ_{SM}):

$$\gamma_{2.1} = C_{FAC} \gamma_{PAR} \gamma_T \gamma_{AGE} \gamma_{LAI} \gamma_{CO2} \gamma_{SM} = \gamma_0 \gamma_{SM} \quad (1)$$

where γ_0 is the product of the non-drought factors. Because of the lack of reliable soil moisture databases, γ_{SM} is always set to be one in GEOS-Chem as in many other chemical transport models, which means no water stress term in the standard model configuration (i.e., NoStress_GC). We show above that NoStress_GC overestimates isoprene emissions and consequently HCHO column densities under drought conditions in the SE US. In this section, we describe the approach whereby observational constraints from the MOFLUX isoprene flux measurement and OMI HCHO were separately used to derive a drought stress factor γ_d which replaces γ_{SM} in Equation (1) to simulate the response of isoprene emissions to drought in MEGAN2.1 implementation of GEOS-Chem (hereafter referring to as GC/MEGAN2.1). The drought stress factor γ_d derived from the MOFLUX isoprene flux measurement is denoted as γ_{d_MOFLUX} and that from OMI HCHO as γ_{d_OMI} . Their corresponding simulations are referred to as MOFLUX_Stress_GC and OMI_Stress_GC, respectively. In either algorithm, the underlying assumption is that the GEOS-Chem model has no significant bias in predicting isoprene fluxes or HCHO columns due to factors other than isoprene emissions under drought conditions. The assumption is reasonable because the GEOS-Chem model uses

reanalysis meteorology, state-of-the-science isoprene oxidation schemes, time-specific anthropogenic emissions and fire emissions, and natural emissions calculated online using model meteorology as described in Section 2.3. The discussion in Section 3.1 validated some aspects of the assumption such as NO_x emissions, wildfire emissions, and PBL.

4.1 MOFLUX-based Drought Stress Algorithm

The γ_{d_MOFLUX} was derived following Jiang et al. (2018) by implementing photosynthesis and water stress parameters with a formula of:

$$\gamma_{d_MOFLUX} = \gamma_0 \gamma_{d_isoprene} \begin{cases} \gamma_{d_isoprene} = 1 & (\beta_t \geq 0.3) \\ \gamma_{d_isoprene} = V_{cmax}/\alpha & (\beta_t < 0.3, \alpha = 77) \end{cases} \quad (2)$$

where V_{cmax} is the maximum carboxylation rate by photosynthetic Rubisco enzyme and β_t represents the water stress ranging from zero (fully stressed) to one (no stress). This simplified method intends to use the decreased photosynthetic enzyme activity to physiologically represent the variation in isoprene emissions under drought stress via dividing V_{cmax} by an empirical parameter α when the water stress is below a threshold.

Since the default GEOS-Chem does not have these photosynthetic parameters, we adopted the ecophysiology module created by Lam et al. (2022) that is based on the photosynthesis calculation in the Joint UK Land Environmental Simulator (JULES; Best et al., 2011; Clark et al., 2011) as an online component in GEOS-Chem so that it simulates photosynthesis rate and bulk stomatal conductance dynamically and consistently with the underlying meteorology that drives GEOS-Chem. The outputs of V_{cmax} and β_t from the ecophysiology module were passed to MEGAN2.1 in GEOS-Chem to parameterize the drought stress according to Equation 2. In addition to GEOS-Chem meteorology, the ecophysiology module uses soil parameters from the Hadley Centre Global Environment Model version 2 – Earth System Model (HadGEM2-ES). In general, the implementation of the ecophysiology module much improved the simulated stomatal conductance and dry deposition velocity relative to site observations on average for seasonal timescales, but the β_t computed has not been calibrated to intermittent drought conditions. Instead of adopting the values of V_{cmax} and β_t from Jiang et al. (2018) which were based on the Community Land Model, we need to determine the β_t threshold and the α value specific to GEOS-Chem with the ecophysiology module of Lam et al. (2022). To calibrate β_t , we first examined the statistical distribution of β_t at the MOFLUX grid (**Figure S2**) during May-September 2011 and 2012 when multiple USDM drought categories occurred. Then we decided on a value of 0.3 as the threshold β_t below which the drought stress will be triggered in the model because this value is greater than 75% quantile of all the β_t values from D0 to D3, thus capturing most of the drought conditions.

We note the observed isoprene flux at MOFLUX is consistently higher than predicted values during the non-drought period (e.g., N0 in Figure 5a). This systematic bias is expected because we compare the single-point observations with grid-mean isoprene emission fluxes. To correct the systemic bias, we scaled down the model isoprene emissions at the MOFLUX grid by a factor of 1.42, which is the ratio of the average hourly isoprene fluxes between observations and simulations at the MOFLUX grid during non-drought conditions ($\beta_t > 0.3$). The factor of 1.42 was applied to

downscale modeled isoprene fluxes at the MOFLUX grid during the entire time series, including drought conditions. The resulted time series are shown in **Figure 6a**. Based on the downscaled model prediction, we derived that $\alpha=77$ under drought conditions ($\beta_t < 0.3$), which minimized the mean bias under drought conditions between the modeled and observed isoprene fluxes at the MOFLUX grid.

Figure 6b shows the comparison of the hourly NoStress_GC and MOFLUX_Stress_GC isoprene emissions with observations in May-September 2012. The overall mean bias is reduced from 2.05 mg/m²/hr to 0.01 mg/m²/hr despite the fact that the stress factor is only applied to drought conditions. The correlation coefficient (R) and index of agreement (IOA) also increase from 0.77 to 0.85 and from 0.80 to 0.93, respectively. All the changes in the comparison metrics indicate the model simulations are improved considerably based on the single-point measurement.




Figure 6. (a) Hourly time series of isoprene emissions at the MOFLUX site from observations (black line) and simulations with (MOFLUX_Stress_GC; blue line) and without drought stress (NoStress_GC; red line; after downscaling). The dots color-coded by USDM levels represent the daily values of β_t (right axis). The dashed line indicates the threshold of 0.3. **(b)** Comparison of isoprene emissions between observations (Obs) and simulations with (MOFLUX_Stress_GC; blue-bordered triangle) and without (NoStress_GC; black-bordered circle) drought stress. Data are color-coded by USDM levels.

4.2 OMI-based Drought Stress Algorithm




Figure 7. Response of HCHO/LAI ratio ($10^{16} \text{ molec cm}^{-2}$) to temperature (K) in different drought levels averaged over JJA 2005-2017. The colored solid line is the modelled NoStress_GC HCHO/LAI ratio, and the black dashed line is the observed HCHO/LAI ratio from OMI. The exponentially fitted formulas and the resulted coefficient of determination (R^2) are labelled in each subplot.

Isoprene emission increases exponentially with temperatures below $\sim 310 \text{ K}$ (Guenther et al., 2006) in the absence of other stress factors such as drought. Indeed, an exponential relationship between biogenic isoprene emission per unit LAI and temperature is predicted by MEGAN2.1 at all USDM levels (**Figure S3**). However, the predicted temperature sensitivity is found to increase substantially with drought severity with no sign of plateauing or slow-down even under the most severe drought conditions when MOFLUX measurements measured a decrease in isoprene emissions (c.f. Figure 5). Similarly, we found NoStress_GC overestimates HCHO sensitivities to high temperatures ($> 300 \text{ K}$) under drought conditions (D0-D4) (**Figure 7**), but no such overestimation is seen under non-drought (N0) or low temperature conditions during drought ($< 300 \text{ K}$). This indicates the role of drought stress on isoprene emissions is likely through suppressing the dependence of emissions on temperatures during drought. Leaf level measurements conducted during the 2012 drought at the MOFLUX site provide independent evidence of drought suppression of the isoprene response to increasing temperature for less drought-resilient tree species (Geron et al., 2016). Taking advantage of these empirical observations, we derived the OMI-based drought stress algorithm by minimizing the differences in HCHO column sensitivities to temperatures between OMI and GEOS-Chem under drought conditions as shown in **Figure 7**. When calculating the relationships between HCHO column densities and temperatures, we first scaled HCHO column by LAI on a grid-by-grid basis to account for the regional differences in isoprene emissions due to different vegetation coverage as well as the effect of LAI changes with drought (c.f. **Figure 4**). Each point in **Figure 7** represents the mean

HCHO/LAI ratio, denoted as Ω , within each 1K temperature interval. We used exponential functions ($\ln\Omega = kT + b$) to separately fit the temperature (T) dependence of HCHO/LAI ratio (Ω) under different drought levels (**Figure 7**) for both the model and OMI. The resulting formulas were listed in **Table 1** and the R^2 of most fitting lines is greater than 0.9.

Table 1. Fitted exponential formulas of NoStress_GC and OMI HCHO/LAI ratio (Ω , 10^{16} molec cm^{-2}) to surface air temperature (T, K), and fitted value of HCHO/LAI ratio at 290K, 300K, and 310K.

USDM	NoStress_GC HCHO/LAI (Ω , 10^{16} molec cm^{-2})				OMI HCHO/LAI (Ω , 10^{16} molec cm^{-2})			
	Fitting Formula	290K	300K	310K	Fitting Formula	290K	300K	310K
N0	$\ln\Omega = 0.104T - 31.42$	0.25	0.72	2.03*	$\ln\Omega = 0.101T - 30.78$	0.26	0.72	1.97*
D0	$\ln\Omega = 0.091T - 27.83$	0.27	0.67	1.66	$\ln\Omega = 0.085T - 25.92$	0.26	0.60	1.40
D1	$\ln\Omega = 0.108T - 32.83$	0.24	0.71	2.10	$\ln\Omega = 0.100T - 30.56$	0.23	0.64	1.74
D2	$\ln\Omega = 0.110T - 33.33$	0.24	0.71	2.14	$\ln\Omega = 0.098T - 29.97$	0.24	0.65	1.75
D3	$\ln\Omega = 0.118T - 35.72$	0.24	0.78	2.56	$\ln\Omega = 0.121T - 36.62$	0.20	0.67	2.23
D4	$\ln\Omega = 0.125T - 37.59$	0.26	0.90	3.13	$\ln\Omega = 0.115T - 34.62$	0.26	0.83	2.60

* Asterisk indicates that the temperature does not reach this value in actual data and is an extrapolated value.

As the fitting equations suggest, both NoStress_GC and OMI HCHO/LAI ratios increase with temperature under all conditions, but the former shows a higher sensitivity to temperature under drought conditions. This can be clearly seen from the higher HCHO/LAI ratios of NoStress_GC (Ω_{GC} ; solid lines) than those of OMI (Ω_{OMI} ; dashed lines) especially when the temperature is greater than 300 K under D0-D4. To better explain this, we also calculated the fitted value of HCHO/LAI at three temperatures of 290K, 300K, and 310K in **Table 1**. Since it is difficult for the N0 condition to reach a temperature of 310K, the values were extrapolated and marked with an asterisk in the table. The results show that the model overestimates the temperature dependence at all drought levels. At 290K, all biases between Ω_{OMI} and Ω_{GC} are less than 0.05×10^{16} molec cm^{-2} . At 310K, the bias between the two is 0.06×10^{16} molec cm^{-2} (3.0%) at N0 but increases by more than a factor of 4 to 0.26×10^{16} molec cm^{-2} (18.6%), 0.36×10^{16} molec cm^{-2} (20.7%), 0.39×10^{16} molec cm^{-2} (22.3%), 0.33×10^{16} molec cm^{-2} (14.8%), and 0.53×10^{16} molec cm^{-2} (20.4%) at D0-D4 drought, respectively. As isoprene emission is a fixed function of temperature in MEGAN2.1, the overdependence of HCHO column on temperature is caused by the previous two weeks' temperatures being higher under drought, which leads to a higher value of γ_T reflecting the temperature “memory” effects on isoprene emissions (**Figure S4**).

Based on the fitted formulas in **Table 1**, the ratio between $\frac{\Omega_{OMI}}{\Omega_{GC}}$ under each level from D0 to D4 can be derived by:

$$\frac{\Omega_{OMI}}{\Omega_{GC}} = \frac{e^{k_{OMI}T + b_{OMI}}}{e^{k_{GC}T + b_{GC}}} = e^{(k_{OMI} - k_{GC})T} e^{(b_{OMI} - b_{GC})} \quad (3)$$

where k_{OMI} (k_{GC}) and b_{OMI} (b_{GC}) represent the slopes and intercepts of the formulas in **Table 1** for OMI (GC) HCHO column; T is surface temperature, and e is the exponential constant. By averaging the values of $k_{OMI}-k_{GC}$ and $b_{OMI}-b_{GC}$ from D0 to D4, we can obtain:

$$\frac{\Omega_{OMI}}{\Omega_{GC}} = 380.10e^{-0.02T} (\beta_t < 0.6, T > 300 K) \quad (4)$$

where $\beta_t < 0.6$ represents the 75% quantile of the β_t values from D0 to D4 for the whole SE US study region in JJA 2005-2017 (**Figure S2**).

The formula of γ_{d_OMI} is thus:

$$\gamma_{d_OMI} = \gamma_0 \gamma_{d_isoprene} \begin{cases} \gamma_{d_isoprene} = 1 (\beta_t \geq 0.6 \text{ or } T \leq 300K) \\ \gamma_{d_isoprene} = \frac{\Omega_{OMI}}{\Omega_{GC}} = 380.10e^{-0.02T} (\beta_t < 0.6, T > 300K) \end{cases} \quad (5)$$

Note the threshold of β_t in equation 5 is different from the value used by γ_{d_MOFLUX} because all the SE US grids were considered in deriving β_t for γ_{d_OMI} . Another difference is that the factor is activated only if the temperature is higher than 300K when significant biases between Ω_{OMI} and Ω_{GC} are found (**Figure 7**).




Figure 8. Boxplot of HCHO column statistical distributions for OMI observations (black) and different GEOS-Chem simulations: without drought stress (NoStress_GC; red) and with drought stress factors derived from MOFLUX observations (MOFLUX_Stress_GC; blue) and from OMI HCHO constraints (OMI_Stress_GC; pink).

Figure 8 compares the statistical distributions of HCHO column densities from OMI, NoStress_GC, MOFLUX_Stress_GC, and OMI_Stress_GC during May-September 2012 over the SE US. Compared to OMI, NoStress_GC simulation has a mean high bias of 0.02×10^{16} molec cm^{-2} - 0.24×10^{16} molec cm^{-2} during D0-D4. The γ_{d_OMI} algorithm reduces the high bias to -0.05×10^{16} molec cm^{-2} - 0.11×10^{16} molec cm^{-2} . By contrast, the γ_{d_MOFLUX} algorithm reduces the HCHO simulations too much over the SE US and causes an overall underestimation of 0.02×10^{16}

molec cm⁻² - 0.25×10^{16} molec cm⁻². The γ_{d_MOFLUX} algorithm also narrows the statistical distribution of HCHO as indicated by the smaller boxes and shorter whiskers compared to OMI. This suggests that the γ_{d_MOFLUX} algorithm based on the single-site observations is incapable of representing the drought stress over the SE US, possibly because the MOFLUX site has thin soil layers and thus is vulnerable to water stress (Opacka et al., 2022). Isoprene emissions measured here are therefore more sensitive to droughts and the same extent of drought stress is likely too strong to be applied to other regions in the SE US. As a result, the γ_{d_OMI} algorithm is used in the next section to further evaluate how this algorithm would change the responses of atmospheric compositions to droughts.

5. Changes in Simulated Biogenic Isoprene Emissions, HCHO, O₃, and OA

In this section, we evaluated the changes in biogenic isoprene emissions and HCHO column densities by running a long-term (2005-2017, JJA) simulation, after adding the OMI-based drought stress factor for isoprene emissions γ_{d_OMI} in GEOS-Chem. Since isoprene is an important precursor for the formation of tropospheric O₃ and OA, maximum daily 8-hour average (MDA8) O₃, and OA changes were also examined. We used the ComplexSOA mechanism in GEOS-Chem (Pye et al., 2010; Marais et al., 2016) which includes more detailed pathways of isoprene to secondary organic aerosols such as aqueous-phase reactive uptake and the formation of organo-nitrates.

Figure 9 shows the changes in biogenic isoprene emissions resulting from adding γ_{d_OMI} drought stress in GEOS-Chem. Here we expanded the maps to the entire contiguous US to examine whether the drought stress algorithm can impose large changes on other US regions although such changes need to be interpreted with caution. The numbers at each panel indicate the means of isoprene emissions of NoStress_GC and the mean differences (MD) relative to the OMI_Stress_GC over the SE US. As expected, the biggest decrease in isoprene emissions is found in the SE US with the regional-mean emissions reduced by 0.17×10^{-10} kg m⁻² s⁻¹ (8.60%), 0.35×10^{-10} kg m⁻² s⁻¹ (14.24%), 0.43×10^{-10} kg m⁻² s⁻¹ (16.57%), 0.49×10^{-10} kg m⁻² s⁻¹ (17.49%), 0.58×10^{-10} kg m⁻² s⁻¹ (18.66%), and 0.65×10^{-10} kg m⁻² s⁻¹ (20.74%) from N0 to D4, respectively (**Figure 9c**). Despite lowering emissions relative to NoStress_GC, OMI_Stress_GC simulates an increase of isoprene emissions under drought conditions compared to non-drought in the SE US; the respective increases are 0.28×10^{-10} kg m⁻² s⁻¹ (15.20%), 0.34×10^{-10} kg m⁻² s⁻¹ (18.40%), 0.49×10^{-10} kg m⁻² s⁻¹ (26.47%), 0.69×10^{-10} kg m⁻² s⁻¹ (37.46%), and 0.65×10^{-10} kg m⁻² s⁻¹ (35.23%) from D0 to D4 relative to N0 (**Figure 9c**). This increase results from the top-down constraints by the corresponding changes in OMI HCHO column densities with USDM and consequently exhibits the behavior of non-uniform increases with drought severity (e.g., peak increase of 37.5% at D3, followed by a ~2% reduction at D4), which is consistent with the MOFLUX flux measurements.

For other regions, such as California and Minnesota, biogenic isoprene emissions decreased slightly by less than 0.5×10^{10} kg m⁻² s⁻¹. The smaller effect of the drought stress factor imposed on regions other than the SE US is understandable because of the lower isoprene emissions.




Figure 9. Simulated biogenic isoprene emissions during JJA 2005-2017 by USDM dryness category by NoStress_GC (a), OMI_Stress_GC minus NoStress_GC (b), and statistical distributions of SE US isoprene emissions between the two simulations (c). Numbers at the bottom-left corner of each panel indicate the SE US (black box) regional mean of biogenic isoprene emissions for NoStress_GC (left column), and mean differences (MD) between OMI_Stress_GC and NoStress_GC (middle column).




Figure 10. Mean HCHO column densities during JJA 2005-2017 by USDM dryness category for OMI (a), NoStress_GC minus OMI (b), and OMI_Stress_GC minus NoStress_GC (c). Numbers at the bottom-left corner of each panel indicate the SE US (black box) regional mean of OMI HCHO column (left column), mean bias (MB), and root mean square error (RMSE) in HCHO column densities between NoStress_GC and OMI (middle column), and mean differences (MD) and root mean square deviation (RMSD) between OMI_Stress_GC and NoStress_GC (right column). MD and RMSD are calculated in the same way as MB and RMSE; the different names are used to distinguish between model-to-model comparison and model-to-observation comparison, respectively.

The changes in the HCHO column are shown in **Figure 10**. Different from the overestimation in the SE US, NoStress_GC underestimates HCHO column densities in the western US compared to OMI (**Figure 10b**). This negative bias should be interpreted with care because the scaling factor of 1.5 (c.f. section 2.2) is derived over the SE US and may not hold in other regions. For the SE US overall, the drought stress factor reduces modeled HCHO columns by 0.08×10^{16} molec cm^{-2} (5.43%), 0.10×10^{16} molec cm^{-2} (6.46%), 0.12×10^{16} molec cm^{-2} (7.22%) and 0.13×10^{16} molec cm^{-2} (7.62%), 0.16×10^{16} molec cm^{-2} (8.91%) under D0-D4, respectively, relative to NoStress_GC (**Figure 10c**). This leads to a better agreement with OMI as OMI_Stress_GC has nearly zero MB under D0-D4 (**Figure S5**; MB = -0.05×10^{16} molec cm^{-2} $\sim$ -0.02×10^{16} molec cm^{-2}). The RMSE is also reduced by 3%-13% relative to the NoStress_GC simulation compared to observations. The changes in both metrics indicate that the drought algorithm

considerably improves the model performance in capturing the biogenic isoprene response to drought as evidenced by HCHO column. Similar to the changes in biogenic isoprene emissions, the OMI_Stress_GC only slightly decreases HCHO column densities (<5%) compared to the NoStress_GC simulation in other US regions.




Figure 11. Same as Figure 10 but for surface maximum daily 8-hour average (MDA8) O₃.

Figure 11a displays the observed MDA8 O₃ changes with USDM. Similar to the changes of the HCHO column with USDM levels, O₃ in the SE US exhibits a gradual increase, relative to the mean of 41.74 ppbv at N0, of 4.70 ppbv, 7.26 ppbv, 9.01 ppbv, 10.26 ppb, and 10.36 ppbv under D0-D4, respectively. This is consistent with our previous study (Li et al., 2022; Lei et al., 2022) which investigated O₃ changes with drought severity in more detail. The NoStress_GC simulation has a high bias in MDA8 O₃ across all USDM categories (**Figure 11b**). High positive bias is a common issue of surface O₃ simulations in chemical transport models, which is a research question and can be attributed to the uncertainties in various processes, such as NO_x emissions, isoprene oxidation pathways, O₃ dry deposition velocity, boundary layer dynamics (Fiore et al., 2005; Lin et al., 2008; Squire et al., 2015; Travis et al., 2016; Travis and Jacob, 2019). Despite the systematic high bias, NoStress_GC captures the increasing trend of MDA8 O₃ with increasing dryness but with a respectively smaller increment (relative to N0) of 3.62 ppbv, 5.67 ppbv, 7.01 ppbv, 7.41 ppbv, and 7.41 ppbv under D0 to D4. This discrepancy between NoStress_GC and observations can also

be inferred from the fact that the MB between model and observations decreases from 14.53 ppbv at N0 to 11.67 ppbv at D4 (**Figure 11b**). **Figure 11c** shows the difference in MDA8 O₃ between OMI_Stress_GC and NoStress_GC. In the SE US where isoprene emissions are the highest and reduced the most by the drought stress algorithm, OMI_Stress_GC shows a small increase in MDA8 O₃ of less than 1 ppbv. This increase in O₃ can be explained by an increase of OH resulting from reducing isoprene emissions under low-NO_x conditions in the SE US (Wells et al., 2020). For the SE US study domain as a whole, the change in MDA8 ozone was negligible but negative (regional mean of -0.5 ppbv). Although the drought factor does not reduce the overall high bias, it makes the model more consistent with the observed increment in MDA8 O₃ for the subregion with increased O₃ (e.g., 90–94°W, 32–35°N) as drought severity increases. Since NO_x has a high positive bias from the NEI2011 inventory (**Figure 4**), the improvement of MDA8 in these regions is likely to be underestimated. Over northeastern Texas, Oklahoma, and Kansas where isoprene emission is also reduced by the drought algorithm yet from a much lower emission base compared to other SE US areas, OMI_Stress_GC simulates 1-3 ppbv lower MDA8 O₃ under drought conditions (D0-D4), leading to a better agreement with observations. For regions with lower isoprene and higher NO_x concentrations, O₃ formation is more sensitive to the changes in isoprene, which explains the reduction in MDA8 O₃ caused by the drought stress factor.




Figure 12. Same as Figure 10 but for organic aerosol (OA).

The changes in OA with USDM are shown in **Figure 12**. Observed OA in the SE US shows an average increase (relative to N0) of 0.12 $\mu\text{g}/\text{m}^3$, 0.32 $\mu\text{g}/\text{m}^3$, 0.34 $\mu\text{g}/\text{m}^3$, 0.31 $\mu\text{g}/\text{m}^3$, and 0.45 $\mu\text{g}/\text{m}^3$ under D0 to D4, respectively. The extremely high values over the northwest states (e.g., Washington and Montana) are likely associated with higher wildfire emissions under droughts (Wang et al., 2017). The NoStress_GC simulation considerably overestimates OA in the SE US with an MB of 1.52 $\mu\text{g}/\text{m}^3$ (50.83%) at N0 and the overestimation becomes even higher to 2.02-2.90 $\mu\text{g}/\text{m}^3$ (64.95%-85.58%) at D0-D4 (**Figure 12b**), thus causing an overprediction of the drought-OA relationship. Zheng et al (2020) reported a similar level of overestimation and attributed this to the overdependence of isoprene-derived secondary organic aerosol (SOA) on sulfate. As isoprene is one of the dominant sources of OA in the SE US (Xu et al., 2015; Budisulistiorini et al., 2016), our analysis suggests that the model overestimation of isoprene emissions under drought conditions is another reason for this high OA bias in the SE US. Indeed, the drought stress factor greatly improves the OA simulation by reducing the MB by 0.30 $\mu\text{g}/\text{m}^3$ (6.60%), 0.46 $\mu\text{g}/\text{m}^3$ (8.98%) 0.60 $\mu\text{g}/\text{m}^3$ (10.07%), 0.67 $\mu\text{g}/\text{m}^3$ (10.85%), 0.62 $\mu\text{g}/\text{m}^3$ (10.88%), 0.74 $\mu\text{g}/\text{m}^3$ (11.71%) under N0 to D4 over the SE US relative to NoStress_GC, thus lowering the MB to be within 1.22-2.18 $\mu\text{g}/\text{m}^3$ (40.82% - 65.52%; **Figure S5**) compared with observations. We also examined the change of three major SOA components in **Figure S6**. Anthropogenic SOA (ASOA) barely changes; isoprene SOA (ISOA) decreases the most as expected since the drought stress factor is applied to isoprene emissions only. Interestingly, terpene SOA (TSOA) also shows a slight decrease, suggesting positive feedback between ISOA and TSOA.

In summary, the OMI-based drought stress factor shows good performance in correcting the overestimation of biogenic isoprene in default GEOS-Chem simulations under drought conditions. The drought stress factor was constrained by the observed exponential fitting between the HCHO to LAI ratio and temperature, not by observed HCHO columns directly. It nearly eliminates the high HCHO bias compared with OMI observations in the SE US under drought conditions, which consequently improves the simulation of OA. MDA8 O₃ slightly increases in the areas with high isoprene emissions, leading to no improvement in model bias but a better agreement with the observed O₃ increment with drought severity. Places with lower isoprene emissions show an MDA8 O₃ reduction of 1-3 ppbv, indicating the region-specific O₃ responses to the changes of isoprene due to the nonlinearity of O₃ chemistry.

6. Conclusions

Using long-term (JJA 2005-2017) weekly USDM drought index and OMI HCHO column data over the SE US, we revealed a step-increase pattern of HCHO by 6.7%, 12.6%, 16.5%, 21.2%, and 23.2% from D0 to D4 relative to non-drought conditions (N0), respectively, which indicates the increasingly higher isoprene emissions with drought on a regional scale although the rate of increase decreases under severe droughts. Compared with OMI observations, the GEOS-Chem simulated HCHO column density exhibits a similar pattern, but the changes are 1.1-1.5 times higher with a respective increase of 9.90%, 15.1%, 19.5%, 21.8%, and 29.1% from D0 to D4. Since there are no big changes in anthropogenic VOCs under droughts, biogenic isoprene emissions are the key drivers for the increase of HCHO, and a drought stress factor is missing in the MEGAN2.1 biogenic inventory in the default GEOS-Chem simulations causing the overestimation of the HCHO changes in response to droughts.

The MOFLUX site provides the only long-term ground-based isoprene flux observations covering multiple drought severities. We developed a drought stress algorithm based on the MOFLUX site following Jiang et al. (2018), and the algorithm improves the HCHO simulation at the MOFLUX grid while underestimating HCHO after all the SE US grids are included. By comparison, the OMI-based drought stress algorithm derived from the different HCHO-temperature sensitivities between OMI and GEOS-Chem can reflect better spatial coverage and nearly removes the positive bias between OMI and the default simulations seen from a test simulation in May-September 2012 over the SE US.

The long-term simulation with the OMI-based drought stress factor can significantly reduce the biogenic isoprene emissions by $0.35 \times 10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$ (14.24%), $0.43 \times 10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$ (16.57%), $0.49 \times 10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$ (17.49%), $0.58 \times 10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$ (18.66%) and $0.65 \times 10^{-10} \text{ kg m}^{-2} \text{ s}^{-1}$ (20.74%) from D0 to D4, respectively, which consequently leads to a better agreement between OMI and simulated HCHO column. Despite lowering emissions relative to the no-stress simulation, OMI_Stress_GC simulates a non-uniform trend of increasing isoprene emissions with drought severity that is consistent with OMI HCHO and MOFLUX. Relative to N0, the simulated increase in isoprene emissions is 15-18% under D0-D1, increasing to 26% at D2 and peaking at 37% at D3, followed by a slight decrease to 35% at D4.

The observed MDA8 O₃ and OA over the SE US show a similar increase pattern with HCHO. The OMI-based drought stress algorithm also helps reduce the mean bias of OA by 0.30 $\mu\text{g/m}^3$ (6.60%), 0.46 $\mu\text{g/m}^3$ (8.98%), 0.60 $\mu\text{g/m}^3$ (10.07%), 0.67 $\mu\text{g/m}^3$ (10.85%), 0.62 $\mu\text{g/m}^3$ (10.88%), 0.74 $\mu\text{g/m}^3$ (11.71%) from N0 to D4 over the SE US compared with the high positive bias of more than 2.02 $\mu\text{g/m}^3$ (50.83%) without the drought stress. By contrast, the MDA8 O₃ response to the reduced biogenic isoprene caused by the drought stress factor presents a spatial disparity due to the nonlinear O₃ chemistry. Places with high isoprene emissions show an increase of MDA8 O₃ by less than 1 ppbv, which slightly improves the simulated drought-O₃ relationship. For the regions with low isoprene emissions in the SE US, the drought stress factor reduces MDA8 O₃ by 1-3 ppbv.

This study reveals an increasingly higher level of biogenic isoprene under drought conditions over the regions with high vegetation coverage. As drought is predicted to become more frequent in a warming climate (Cook et al., 2018), it is essential to update current biogenic emission inventories by adding a drought stress factor and to improve the constraints of isoprene chemistry in the climate chemistry models in order to have a better projection of air quality in the future. We demonstrate the feasibility of applying satellite data to the development of drought stress algorithms when ground-based measurements are limited. Our attempt here is a top-down approach and used temperature as the only parameter to adjust isoprene emissions under drought conditions. The water stress threshold in our algorithm is used only as a triggering parameter; that is, it is used to determine whether a grid is in drought or not and thus can be replaced with other drought-identifying approaches. One issue with our approach is the type of temperature data to be used in the algorithm. Ideally, it should be leaf temperature because this is what regulates stomata at the process level. However, leaf temperature is not readily available from meteorological fields that drive CTMs. MEGAN uses 2 m air temperature to parameterize isoprene emissions, and thus our algorithm uses the same temperature. More biogenic

emission flux observations covering different vegetation types and drought severities will be helpful to better depict the relationships between biogenic VOCs and drought stress.

Acknowledgment

This work was supported by NASA Atmospheric Composition Modeling and Analysis Program (80NSSC19K0986). The development of the ecophysiology module in GEOS-Chem has also been supported by the General Research Fund (14306220) granted by the Hong Kong Research Grants Council. The authors thank NASA Langley Research Center for the OMI HCHO column data and the National Drought Mitigation Center for making and providing the USDM maps. Roger Seco was supported by grants RYC2020-029216-I and CEX2018-000794-S funded by MCIN/AEI/10.13039/501100011033 and by the European Social Fund “ESF Investing in your future”.

Data Availability

GEOS-Chem model is publicly available at <http://www.geos-chem.org>. LAI is obtained from http://geoschemdata.wustl.edu/ExtData/HEMCO/Yuan_XLAI/v2021-06/. O₃ and organic carbon observational data can be downloaded via https://aqs.epa.gov/aqsweb/documents/data_mart_welcome.html. Observational isoprene measurements at MOFLUX are from Potosnak et al. 2014 and Seco et al. 2015 and are available upon request from co-author Alex Guenther. Satellite Ω HCHO is available publicly at https://cmr.earthdata.nasa.gov/search/concepts/C1626121562-GES_DISC.html.

Competing interests

The authors declare that they have no conflict of interest.

Author contributions

YW conceived the research idea. NL and WL conducted the model simulation and data analysis. JCYL and APKT created the ecophysiology module. AG, MJP and RS provided the field observations. All authors contributed to the interpretation of the results and the preparation of the manuscript

References

- Abbot, D. S., Palmer, P. I., Martin, R. V., Chance, K. V., Jacob, D. J., and Guenther, A.: Seasonal and interannual variability of North American isoprene emissions as determined by formaldehyde column measurements from space, *Geophys. Res. Lett.*, 30, <https://doi.org/10.1029/2003GL017336>, 2003.
- Alvarado, L. M. A., Richter, A., Vrekoussis, M., Hilboll, A., Kalisz Hedegaard, A. B., Schneising, O., and Burrows, J. P.: Unexpected long-range transport of glyoxal and formaldehyde observed from the Copernicus Sentinel-5 Precursor satellite during the 2018 Canadian wildfires, *Atmospheric Chem. Phys.*, 20, 2057–2072, <https://doi.org/10.5194/acp-20-2057-2020>, 2020.

Atkinson, R.: Atmospheric chemistry of VOCs and NO_x, *Atmos. Environ.*, 34, 2063–2101, [https://doi.org/10.1016/S1352-2310\(99\)00460-4](https://doi.org/10.1016/S1352-2310(99)00460-4), 2000.

Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes, *Geosci. Model Dev.*, 4, 677–699, <https://doi.org/10.5194/gmd-4-677-2011>, 2011.

Budisulistiorini, S. H., Baumann, K., Edgerton, E. S., Bairai, S. T., Mueller, S., Shaw, S. L., Knipping, E. M., Gold, A., and Surratt, J. D.: Seasonal characterization of submicron aerosol chemical composition and organic aerosol sources in the southeastern United States: Atlanta, Georgia, and Look Rock, Tennessee, *Atmospheric Chem. Phys.*, 16, 5171–5189, <https://doi.org/10.5194/acp-16-5171-2016>, 2016.

Chance, K.: OMI/Aura Formaldehyde (HCHO) Total Column Daily L3 Weighted Mean Global 0.1 deg Lat/Lon Grid V003, Goddard Earth Sci. Data Inf. Serv. Cent. GES DISC Greenbelt MD USA, 2019.

Chang, K.-Y., Xu, L., Starr, G., and Paw U, K. T.: A drought indicator reflecting ecosystem responses to water availability: The Normalized Ecosystem Drought Index, *Agric. For. Meteorol.*, 250–251, 102–117, <https://doi.org/10.1016/j.agrformet.2017.12.001>, 2018.

Chen, L. G., Gottschalck, J., Hartman, A., Miskus, D., Tinker, R., and Artusa, A.: Flash Drought Characteristics Based on U.S. Drought Monitor, *Atmosphere*, 10, 498, <https://doi.org/10.3390/atmos10090498>, 2019.

Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M. O., Artaxo, P., and Maenhaut, W.: Formation of Secondary Organic Aerosols Through Photooxidation of Isoprene, *Science*, 303, 1173–1176, <https://doi.org/10.1126/science.1092805>, 2004.

Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics, *Geosci. Model Dev.*, 4, 701–722, <https://doi.org/10.5194/gmd-4-701-2011>, 2011.

Cook, B. I., Mankin, J. S., and Anchukaitis, K. J.: Climate Change and Drought: From Past to Future, *Curr. Clim. Change Rep.*, 4, 164–179, <https://doi.org/10.1007/s40641-018-0093-2>, 2018.

Ferracci, V., Bolas, C. G., Freshwater, R. A., Staniaszek, Z., King, T., Jaars, K., Otu-Larbi, F., Beale, J., Malhi, Y., Waine, T. W., Jones, R. L., Ashworth, K., and Harris, N. R. P.: Continuous Isoprene Measurements in a UK Temperate Forest for a Whole Growing Season: Effects of Drought Stress During the 2018 Heatwave, *Geophys. Res. Lett.*, 47, e2020GL088885, <https://doi.org/10.1029/2020GL088885>, 2020.

Fiore, A. M., Horowitz, L. W., Purves, D. W., Levy II, H., Evans, M. J., Wang, Y., Li, Q., and Yantosca, R. M.: Evaluating the contribution of changes in isoprene emissions to surface ozone trends over the eastern United States, *J. Geophys. Res. Atmospheres*, 110, <https://doi.org/10.1029/2004JD005485>, 2005.

Geron, C., Daly, R., Harley, P., Rasmussen, R., Seco, R., Guenther, A., Karl, T., and Gu, L.: Large drought-induced variations in oak leaf volatile organic compound emissions during PINOT NOIR 2012, *Chemosphere*, 146, 8–21, <https://doi.org/10.1016/j.chemosphere.2015.11.086>, 2016.

Giglio, L., Randerson, J. T., and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *J. Geophys. Res. Biogeosciences*, 118, 317–328, <https://doi.org/10.1002/jgrg.20042>, 2013.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chem. Phys.*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.

653 Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and monoterpene emission
654 rate variability: Model evaluations and sensitivity analyses, *J. Geophys. Res. Atmospheres*, 98, 12609–12617,
655 <https://doi.org/10.1029/93JD00527>, 1993.

656 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The
657 Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated
658 framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, [https://doi.org/10.5194/gmd-5-](https://doi.org/10.5194/gmd-5-1471-2012)
659 1471-2012, 2012.

660 Guenther, A. B., Shah, T., and Huang, L.: A next generation modeling system for estimating Texas biogenic VOC
661 emissions [M], AQR Proj., 16–011, 2017.

662 Guo, J., Zhang, J., Yang, K., Liao, H., Zhang, S., Huang, K., Lv, Y., Shao, J., Yu, T., Tong, B., Li, J., Su, T., Yim,
663 S. H. L., Stoffelen, A., Zhai, P., and Xu, X.: Investigation of near-global daytime boundary layer height using high-
664 resolution radiosondes: first results and comparison with ERA5, MERRA-2, JRA-55, and NCEP-2 reanalyses,
665 *Atmospheric Chem. Phys.*, 21, 17079–17097, <https://doi.org/10.5194/acp-21-17079-2021>, 2021.

666 Guttman, N. B.: Accepting the Standardized Precipitation Index: A Calculation Algorithm1, *JAWRA J. Am. Water*
667 *Resour. Assoc.*, 35, 311–322, <https://doi.org/10.1111/j.1752-1688.1999.tb03592.x>, 1999.

668 Hoerling, M., Eischeid, J., Kumar, A., Leung, R., Mariotti, A., Mo, K., Schubert, S., and Seager, R.: Causes and
669 Predictability of the 2012 Great Plains Drought, *Bull. Am. Meteorol. Soc.*, 95, 269–282,
670 <https://doi.org/10.1175/BAMS-D-13-00055.1>, 2014.

671 Huang, J., Dool, H. M. van den, and Georgarakos, K. P.: Analysis of Model-Calculated Soil Moisture over the
672 United States (1931–1993) and Applications to Long-Range Temperature Forecasts, *J. Clim.*, 9, 1350–1362,
673 [https://doi.org/10.1175/1520-0442\(1996\)009<1350:AOMCSM>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<1350:AOMCSM>2.0.CO;2), 1996.

674 Huang, L., McGaughey, G., McDonald-Buller, E., Kimura, Y., and Allen, D. T.: Quantifying regional, seasonal and
675 interannual contributions of environmental factors on isoprene and monoterpene emissions estimates over eastern
676 Texas, *Atmos. Environ.*, 106, 120–128, <https://doi.org/10.1016/j.atmosenv.2015.01.072>, 2015.

677 Jiang, X., Guenther, A., Potosnak, M., Geron, C., Seco, R., Karl, T., Kim, S., Gu, L., and Pallardy, S.: Isoprene
678 emission response to drought and the impact on global atmospheric chemistry, *Atmos. Environ.*, 183, 69–83,
679 <https://doi.org/10.1016/j.atmosenv.2018.01.026>, 2018.

680 Kaiser, J., Jacob, D. J., Zhu, L., Travis, K. R., Fisher, J. A., González Abad, G., Zhang, L., Zhang, X., Fried, A.,
681 Crounse, J. D., St. Clair, J. M., and Wisthaler, A.: High-resolution inversion of OMI formaldehyde columns to
682 quantify isoprene emission on ecosystem-relevant scales: application to the southeast US, *Atmospheric Chem.*
683 *Phys.*, 18, 5483–5497, <https://doi.org/10.5194/acp-18-5483-2018>, 2018.

684 Kogan, F. N.: Droughts of the Late 1980s in the United States as Derived from NOAA Polar-Orbiting Satellite Data,
685 *Bull. Am. Meteorol. Soc.*, 76, 655–668, [https://doi.org/10.1175/1520-0477\(1995\)076<0655:DOTLIT>2.0.CO;2](https://doi.org/10.1175/1520-0477(1995)076<0655:DOTLIT>2.0.CO;2),
686 1995.

687 Kravitz, B., Guenther, A. B., Gu, L., Karl, T., Kaser, L., Pallardy, S. G., Peñuelas, J., Potosnak, M. J., and Seco, R.:
688 A new paradigm of quantifying ecosystem stress through chemical signatures, *Ecosphere*, 7, e01559,
689 <https://doi.org/10.1002/ecs2.1559>, 2016.

690 Lam, J. C. Y., Tai, A. P. K., Ducker, J. A., and Holmes, C. D.: Development of an ecophysiology module in the
691 GEOS-Chem chemical transport model version 12.2.0 to represent biosphere-atmosphere fluxes relevant for ozone
692 air quality, *EGUsphere*, 1–32, <https://doi.org/10.5194/egusphere-2022-786>, 2022.

693 Lei, Y., Yue, X., Liao, H., Zhang, L., Zhou, H., Tian, C., Gong, C., Ma, Y., Cao, Y., Seco, R., Karl, T., and
 694 Potosnak, M.: Global Perspective of Drought Impacts on Ozone Pollution Episodes, *Environ. Sci. Technol.*, 56,
 695 3932–3940, <https://doi.org/10.1021/acs.est.1c07260>, 2022.

696 Li, W., Wang, Y., Flynn, J., Griffin, R. J., Guo, F., and Schnell, J. L.: Spatial Variation of Surface O₃ Responses to
 697 Drought Over the Contiguous United States During Summertime: Role of Precursor Emissions and Ozone
 698 Chemistry, *J. Geophys. Res. Atmospheres*, 127, e2021JD035607, <https://doi.org/10.1029/2021JD035607>, 2022.

699 Liao, J., Wolfe, G. M., Hannun, R. A., St. Clair, J. M., Hanisco, T. F., Gilman, J. B., Lamplugh, A., Selimovic, V.,
 700 Diskin, G. S., Nowak, J. B., Halliday, H. S., DiGangi, J. P., Hall, S. R., Ullmann, K., Holmes, C. D., Fite, C. H.,
 701 Agastra, A., Ryerson, T. B., Peischl, J., Bourgeois, I., Warneke, C., Coggon, M. M., Gkatzelis, G. I., Sekimoto, K.,
 702 Fried, A., Richter, D., Weibring, P., Apel, E. C., Hornbrook, R. S., Brown, S. S., Womack, C. C., Robinson, M. A.,
 703 Washenfelder, R. A., Veres, P. R., and Neuman, J. A.: Formaldehyde evolution in US wildfire plumes during the
 704 Fire Influence on Regional to Global Environments and Air Quality experiment (FIREX-AQ), *Atmospheric Chem.*
 705 *Phys.*, 21, 18319–18331, <https://doi.org/10.5194/acp-21-18319-2021>, 2021.

706 Lin, J.-T., Youn, D., Liang, X.-Z., and Wuebbles, D. J.: Global model simulation of summertime U.S. ozone diurnal
 707 cycle and its sensitivity to PBL mixing, spatial resolution, and emissions, *Atmos. Environ.*, 42, 8470–8483,
 708 <https://doi.org/10.1016/j.atmosenv.2008.08.012>, 2008.

709 Marais, E. A., Jacob, D. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Krechmer, J., Zhu, L., Kim, P.
 710 S., Miller, C. C., Fisher, J. A., Travis, K., Yu, K., Hanisco, T. F., Wolfe, G. M., Arkinson, H. L., Pye, H. O. T.,
 711 Froyd, K. D., Liao, J., and McNeill, V. F.: Aqueous-phase mechanism for secondary organic aerosol formation from
 712 isoprene: application to the southeast United States and co-benefit of SO₂ emission controls, *Atmospheric Chem.*
 713 *Phys.*, 16, 1603–1618, <https://doi.org/10.5194/acp-16-1603-2016>, 2016.

714 McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and duration to time scales, in:
 715 *Proceedings of the 8th Conference on Applied Climatology*, 179–183, 1993.

716 Millet, D. B., Jacob, D. J., Turquety, S., Hudman, R. C., Wu, S., Fried, A., Walega, J., Heikes, B. G., Blake, D. R.,
 717 Singh, H. B., Anderson, B. E., and Clarke, A. D.: Formaldehyde distribution over North America: Implications for
 718 satellite retrievals of formaldehyde columns and isoprene emission, *J. Geophys. Res. Atmospheres*, 111,
 719 <https://doi.org/10.1029/2005JD006853>, 2006.

720 Miralles, D. G., Teuling, A. J., van Heerwaarden, C. C., and Vilà-Guerau de Arellano, J.: Mega-heatwave
 721 temperatures due to combined soil desiccation and atmospheric heat accumulation, *Nat. Geosci.*, 7, 345–349,
 722 <https://doi.org/10.1038/ngeo2141>, 2014.

723 Naimark, J. G., Fiore, A. M., Jin, X., Wang, Y., Klovnski, E., and Braneon, C.: Evaluating Drought Responses of
 724 Surface Ozone Precursor Proxies: Variations With Land Cover Type, Precipitation, and Temperature, *Geophys. Res.*
 725 *Lett.*, 48, e2020GL091520, <https://doi.org/10.1029/2020GL091520>, 2021.

726 Nickolay A. Krotkov, Lok N. Lamsal, Sergey V. Marchenko, Edward A. Celarier, Eric J. Bucsela, William H.
 727 Swartz, Joanna Joiner and the OMI core team (2019), OMI/Aura NO₂ Cloud-Screened Total and Tropospheric
 728 Column L3 Global Gridded 0.25 degree x 0.25 degree V3, NASA Goddard Space Flight Center, Goddard Earth
 729 Sciences Data and Information Services Center (GES DISC), Accessed: [last access: 4 October 2022],
 730 <https://doi.org/10.5067/Aura/OMI/DATA3007>.

731 Opacka, B., Müller, J.-F., Stavrakou, T., Miralles, D. G., Koppa, A., Pagán, B. R., Potosnak, M. J., Seco, R., De
 732 Smedt, I., and Guenther, A. B.: Impact of Drought on Isoprene Fluxes Assessed Using Field Data, Satellite-Based
 733 GLEAM Soil Moisture and HCHO Observations from OMI, *Remote Sens.*, 14, 2021,
 734 <https://doi.org/10.3390/rs14092021>, 2022.

735 Otu-Larbi, F., Bolas, C. G., Ferracci, V., Staniaszek, Z., Jones, R. L., Malhi, Y., Harris, N. R. P., Wild, O., and
 736 Ashworth, K.: Modelling the effect of the 2018 summer heatwave and drought on isoprene emissions in a UK
 737 woodland, *Glob. Change Biol.*, 26, 2320–2335, <https://doi.org/10.1111/gcb.14963>, 2020.

738 Pacifico, F., Harrison, S. P., Jones, C. D., and Sitch, S.: Isoprene emissions and climate, *Atmos. Environ.*, 43, 6121–
 739 6135, <https://doi.org/10.1016/j.atmosenv.2009.09.002>, 2009.

740 Palmer, P. I., Jacob, D. J., Fiore, A. M., Martin, R. V., Chance, K., and Kurosu, T. P.: Mapping isoprene emissions
 741 over North America using formaldehyde column observations from space, *J. Geophys. Res. Atmospheres*, 108,
 742 <https://doi.org/10.1029/2002JD002153>, 2003.

743 Palmer, W. C.: *Meteorological Drought*, U.S. Department of Commerce, Weather Bureau, 68 pp., 1965.

744 Potosnak, M. J., LeSturgeon, L., Pallardy, S. G., Hosman, K. P., Gu, L., Karl, T., Geron, C., and Guenther, A. B.:
 745 Observed and modeled ecosystem isoprene fluxes from an oak-dominated temperate forest and the influence of
 746 drought stress, *Atmos. Environ.*, 84, 314–322, <https://doi.org/10.1016/j.atmosenv.2013.11.055>, 2014.

747 Pye, H. O. T., Chan, A. W. H., Barkley, M. P., and Seinfeld, J. H.: Global modeling of organic aerosol: the
 748 importance of reactive nitrogen (NO_x and NO_3), *Atmospheric Chem. Phys.*, 10, 11261–11276,
 749 <https://doi.org/10.5194/acp-10-11261-2010>, 2010.

750 Pye, H. O. T., Murphy, B. N., Xu, L., Ng, N. L., Carlton, A. G., Guo, H., Weber, R., Vasilakos, P., Appel, K. W.,
 751 Budisulistiorini, S. H., Surratt, J. D., Nenes, A., Hu, W., Jimenez, J. L., Isaacman-VanWertz, G., Misztal, P. K., and
 752 Goldstein, A. H.: On the implications of aerosol liquid water and phase separation for organic aerosol mass,
 753 *Atmospheric Chem. Phys.*, 17, 343–369, <https://doi.org/10.5194/acp-17-343-2017>, 2017.

754 Schnell, J. L., Holmes, C. D., Jangam, A., and Prather, M. J.: Skill in forecasting extreme ozone pollution episodes
 755 with a global atmospheric chemistry model, *Atmospheric Chem. Phys.*, 14, 7721–7739, <https://doi.org/10.5194/acp-14-7721-2014>, 2014.

757 Schroder, J. C., Campuzano-Jost, P., Day, D. A., Shah, V., Larson, K., Sommers, J. M., Sullivan, A. P., Campos, T.,
 758 Reeves, J. M., Hills, A., Hornbrook, R. S., Blake, N. J., Scheuer, E., Guo, H., Fibiger, D. L., McDuffie, E. E., Hayes,
 759 P. L., Weber, R. J., Dibb, J. E., Apel, E. C., Jaeglé, L., Brown, S. S., Thornton, J. A., and Jimenez, J. L.: Sources and
 760 Secondary Production of Organic Aerosols in the Northeastern United States during WINTER, *J. Geophys. Res.*
 761 *Atmospheres*, 123, 7771–7796, <https://doi.org/10.1029/2018JD028475>, 2018.

762 Seager, R., Tzanova, A., and Nakamura, J.: Drought in the Southeastern United States: Causes, Variability over the
 763 Last Millennium, and the Potential for Future Hydroclimate Change, *J. Clim.*, 22, 5021–5045,
 764 <https://doi.org/10.1175/2009JCLI2683.1>, 2009.

765 Seco, R., Karl, T., Guenther, A., Hosman, K. P., Pallardy, S. G., Gu, L., Geron, C., Harley, P., and Kim, S.:
 766 Ecosystem-scale volatile organic compound fluxes during an extreme drought in a broadleaf temperate forest of the
 767 Missouri Ozarks (central USA), *Glob. Change Biol.*, 21, 3657–3674, <https://doi.org/10.1111/gcb.12980>, 2015.

768 Seco, R., Holst, T., Davie-Martin, C. L., Simin, T., Guenther, A., Pirk, N., Rinne, J., and Rinnan, R.: Strong
 769 isoprene emission response to temperature in tundra vegetation, *Proc. Natl. Acad. Sci.*, 119, e2118014119,
 770 <https://doi.org/10.1073/pnas.2118014119>, 2022.

771 Shen, L., Jacob, D. J., Zhu, L., Zhang, Q., Zheng, B., Sulprizio, M. P., Li, K., De Smedt, I., González Abad, G., Cao,
 772 H., Fu, T.-M., and Liao, H.: The 2005–2016 Trends of Formaldehyde Columns Over China Observed by Satellites:
 773 Increasing Anthropogenic Emissions of Volatile Organic Compounds and Decreasing Agricultural Fire Emissions,
 774 *Geophys. Res. Lett.*, 46, 4468–4475, <https://doi.org/10.1029/2019GL082172>, 2019.

775 Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrakou, T., Müller, J.-F., Kuhn, U., Stefani, P.,
776 and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years,
777 *Atmospheric Chem. Phys.*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014.

778 Sprengnether, M., Demerjian, K. L., Donahue, N. M., and Anderson, J. G.: Product analysis of the OH oxidation of
779 isoprene and 1,3-butadiene in the presence of NO, *J. Geophys. Res. Atmospheres*, 107, ACH 8-1-ACH 8-13,
780 <https://doi.org/10.1029/2001JD000716>, 2002.

781 Squire, O. J., Archibald, A. T., Griffiths, P. T., Jenkin, M. E., Smith, D., and Pyle, J. A.: Influence of isoprene
782 chemical mechanism on modelled changes in tropospheric ozone due to climate and land use over the 21st century,
783 *Atmospheric Chem. Phys.*, 15, 5123–5143, <https://doi.org/10.5194/acp-15-5123-2015>, 2015.

784 Stavrakou, T., Müller, J.-F., Bauwens, M., De Smedt, I., Van Roozendael, M., and Guenther, A.: Impact of Short-
785 Term Climate Variability on Volatile Organic Compounds Emissions Assessed Using OMI Satellite Formaldehyde
786 Observations, *Geophys. Res. Lett.*, 45, 8681–8689, <https://doi.org/10.1029/2018GL078676>, 2018.

787 Svoboda, M., LeComte, D., Hayes, M., Heim, R., Gleason, K., Angel, J., Rippey, B., Tinker, R., Palecki, M.,
788 Stooksbury, D., Miskus, D., and Stephens, S.: THE DROUGHT MONITOR, *Bull. Am. Meteorol. Soc.*, 83, 1181–
789 1190, <https://doi.org/10.1175/1520-0477-83.8.1181>, 2002.

790 Svoboda, M. D., Fuchs, B. A., Poulsen, C. C., and Nothwehr, J. R.: The drought risk atlas: Enhancing decision
791 support for drought risk management in the United States, *J. Hydrol.*, 526, 274–286,
792 <https://doi.org/10.1016/j.jhydrol.2015.01.006>, 2015.

793 Travis, K. R. and Jacob, D. J.: Systematic bias in evaluating chemical transport models with maximum daily
794 8 h average (MDA8) surface ozone for air quality applications: a case study with GEOS-Chem v9.02,
795 *Geosci. Model Dev.*, 12, 3641–3648, <https://doi.org/10.5194/gmd-12-3641-2019>, 2019.

796 Travis, K. R., Jacob, D. J., Fisher, J. A., Kim, P. S., Marais, E. A., Zhu, L., Yu, K., Miller, C. C., Yantosca, R. M.,
797 Sulprizio, M. P., Thompson, A. M., Wennberg, P. O., Crounse, J. D., St. Clair, J. M., Cohen, R. C., Laughner, J. L.,
798 Dibb, J. E., Hall, S. R., Ullmann, K., Wolfe, G. M., Pollack, I. B., Peischl, J., Neuman, J. A., and Zhou, X.: Why do
799 models overestimate surface ozone in the Southeast United States?, *Atmospheric Chem. Phys.*, 16, 13561–13577,
800 <https://doi.org/10.5194/acp-16-13561-2016>, 2016.

801 Trenberth, K. E., Dai, A., van der Schrier, G., Jones, P. D., Barichivich, J., Briffa, K. R., and Sheffield, J.: Global
802 warming and changes in drought, *Nat. Clim. Change*, 4, 17–22, <https://doi.org/10.1038/nclimate2067>, 2014.

803 Trugman, A. T., Medvigy, D., Mankin, J. S., and Anderegg, W. R. L.: Soil Moisture Stress as a Major Driver of
804 Carbon Cycle Uncertainty, *Geophys. Res. Lett.*, 45, 6495–6503, <https://doi.org/10.1029/2018GL078131>, 2018.

805 Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global
806 Warming: The Standardized Precipitation Evapotranspiration Index, *J. Clim.*, 23, 1696–1718,
807 <https://doi.org/10.1175/2009JCLI2909.1>, 2010.

808 Wang, Y., Xie, Y., Dong, W., Ming, Y., Wang, J., and Shen, L.: Adverse effects of increasing drought on air quality
809 via natural processes, *Atmospheric Chem. Phys.*, 17, 12827–12843, <https://doi.org/10.5194/acp-17-12827-2017>,
810 2017.

811 Wells, K. C., Millet, D. B., Payne, V. H., Deventer, M. J., Bates, K. H., de Gouw, J. A., Graus, M., Warneke, C.,
812 Wisthaler, A., and Fuentes, J. D.: Satellite isoprene retrievals constrain emissions and atmospheric oxidation,
813 *Nature*, 585, 225–233, <https://doi.org/10.1038/s41586-020-2664-3>, 2020.

814 Willeke, G., Hosking, J. R. M., Wallis, J. R., and Guttman, N. B.: The national drought atlas, *Inst. Water Resour.*
815 *Rep.*, 94, 1994.

816 Xu, L., Guo, H., Boyd, C. M., Klein, M., Bougiatioti, A., Cerully, K. M., Hite, J. R., Isaacman-VanWertz, G.,
817 Kreisberg, N. M., and Knote, C.: Effects of anthropogenic emissions on aerosol formation from isoprene and
818 monoterpenes in the southeastern United States, *Proc. Natl. Acad. Sci.*, 112, 37–42, 2015.

819 Yuan, H., Dai, Y., Xiao, Z., Ji, D., and Shangguan, W.: Reprocessing the MODIS Leaf Area Index products for land
820 surface and climate modelling, *Remote Sens. Environ.*, 115, 1171–1187, <https://doi.org/10.1016/j.rse.2011.01.001>,
821 2011.

822 Zhao, T., Mao, J., Simpson, W. R., De Smedt, I., Zhu, L., Hanisco, T. F., Wolfe, G. M., St. Clair, J. M., González
823 Abad, G., Nowlan, C. R., Barletta, B., Meinardi, S., Blake, D. R., Apel, E. C., and Hornbrook, R. S.: Source and
824 variability of formaldehyde (HCHO) at northern high latitudes: an integrated satellite, aircraft, and model study,
825 *Atmospheric Chem. Phys.*, 22, 7163–7178, <https://doi.org/10.5194/acp-22-7163-2022>, 2022.

826 Zheng, Y., Unger, N., Tadić, J. M., Seco, R., Guenther, A. B., Barkley, M. P., Potosnak, M. J., Murray, L. T.,
827 Michalak, A. M., Qiu, X., Kim, S., Karl, T., Gu, L., and Pallardy, S. G.: Drought impacts on photosynthesis,
828 isoprene emission and atmospheric formaldehyde in a mid-latitude forest, *Atmos. Environ.*, 167, 190–201,
829 <https://doi.org/10.1016/j.atmosenv.2017.08.017>, 2017.

830 Zheng, Y., Thornton, J. A., Ng, N. L., Cao, H., Henze, D. K., McDuffie, E. E., Hu, W., Jimenez, J. L., Marais, E. A.,
831 Edgerton, E., and Mao, J.: Long-term observational constraints of organic aerosol dependence on inorganic species
832 in the southeast US, *Atmospheric Chem. Phys.*, 20, 13091–13107, <https://doi.org/10.5194/acp-20-13091-2020>,
833 2020.

834 Zhu, L., Jacob, D. J., Kim, P. S., Fisher, J. A., Yu, K., Travis, K. R., Mickley, L. J., Yantosca, R. M., Sulprizio, M.
835 P., De Smedt, I., González Abad, G., Chance, K., Li, C., Ferrare, R., Fried, A., Hair, J. W., Hanisco, T. F., Richter,
836 D., Jo Scarino, A., Walega, J., Weibring, P., and Wolfe, G. M.: Observing atmospheric formaldehyde (HCHO) from
837 space: validation and intercomparison of six retrievals from four satellites (OMI, GOME2A, GOME2B, OMPS)
838 with SEAC⁴RS aircraft observations over the southeast US, *Atmospheric Chem. Phys.*, 16, 13477–13490,
839 <https://doi.org/10.5194/acp-16-13477-2016>, 2016.

840