

Simulating marine neodymium isotope distributions using Nd v1.0 coupled to the ocean component of the FAMOUS-MOSES1 climate model: sensitivities to reversible scavenging efficiency and benthic source distributions

Suzanne Robinson¹, Ruza F. Ivanovic¹, Lauren J. Gregoire¹, Julia Tindall¹, Tina van de Fliedert², Yves Plancherel², Frerk Pöppelmeier³, Kazuyo Tachikawa⁴, Paul J. Valdes⁵

¹School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK
²Department of Earth Science and Engineering, Imperial College London, London, SW7 2AZ, UK
³Climate and Environmental Physics, Physics Institute and Oeschger Center for Climate Change Research, University of Bern, 3012 Bern, Switzerland
⁴Aix Marseille Univ, CNRS, IRD, INRAE, Coll France, CEREGE, Aix-en-Provence, France
⁵School of Geographical Sciences, University of Bristol, Bristol, UK

Correspondence to: Suzanne Robinson (ee14s2r@leeds.ac.uk)

Abstract. The neodymium (Nd) isotopic composition of seawater is a widely used ocean circulation tracer. However, uncertainty in quantifying the global ocean Nd budget, particularly constraining elusive non-conservative processes, remains a major challenge. A substantial increase in modern seawater Nd measurements from the GEOTRACES programme coupled with recent hypotheses that a seafloor-wide benthic Nd flux to the ocean may govern global Nd isotope distributions (ϵ_{Nd}) presents an opportunity to develop a new scheme specifically designed to test these paradigms. Here, we present the implementation of Nd isotopes (^{143}Nd and ^{144}Nd) into the ocean component of the FAMOUS coupled atmosphere-ocean general circulation model (Nd v1.0), a tool which can be widely used for simulating complex feedbacks between different Earth system processes on decadal to multi-millennial timescales.

Using an equilibrium pre-industrial simulation tuned to represent the largescale Atlantic Ocean circulation, we perform a series of sensitivity tests evaluating the new Nd isotope scheme. We investigate how Nd source/sink and cycling parameters govern global marine ϵ_{Nd} distributions, and provide an updated compilation of 6,048 Nd concentrations and 3,278 ϵ_{Nd} measurements to assess model performance. Our findings support the notions that reversible scavenging is a key process for enhancing the Atlantic-Pacific basinal ϵ_{Nd} gradient, and is capable of driving the observed increase in Nd concentration along the global circulation pathway. A benthic flux represents a major source of Nd to the deep ocean. However, model-data disparities in the North Pacific highlight that under a uniform benthic flux, the source of ϵ_{Nd} from seafloor sediments is too unradiogenic in our model to be able to accurately represent seawater measurements. Additionally, model-data mismatch in the northern North Atlantic alludes to the possibility of preferential contributions from 'reactive' unradiogenic detrital sediments.

- Deleted: n
- Deleted: indiscriminate
- Deleted: with a constant benthic flux
- Deleted: suggests a missing source of Nd that is much more unradiogenic than the bulk sediment,
- Deleted: ing
- Deleted: under a benthic flux driven model of marine Nd cycling

40 The new Nd isotope scheme forms an excellent tool for exploring global marine Nd cycling and the interplay between climatic and oceanographic conditions under both modern and palaeoceanographic contexts.

1 Introduction

The Neodymium (Nd) isotope composition of seawater shows a clear provinciality between different ocean basins and is often used as a water mass provenance tracer (e.g., Frank, 2002; Goldstein and Hemming, 2003). The measured ¹⁴³Nd/¹⁴⁴Nd ratio is
45 denoted relative to the bulk earth standard:

$$\epsilon_{Nd} = \left(\frac{(^{143}Nd/^{144}Nd)_{sample}}{(^{143}Nd/^{144}Nd)_{CHUR}} - 1 \right) \times 10^4, \quad (1)$$

where (¹⁴³Nd/¹⁴⁴Nd)_{CHUR} relates to the Chondritic Uniform Reservoir (CHUR; 0.512638; Jacobsen and Wasserburg, 1980). Distinct variations in the Nd isotope signal of water masses originate from different continental regions and their isotopic fingerprints, and subsequent influence by ocean circulation, water mass mixing and particle cycling, as well as interaction
50 with sediments (e.g. Tachikawa et al., 2017; van de Flierdt et al., 2016 for recent reviews). Neodymium in the deep ocean has an estimated mean ocean residence time between 360-800 years (Arsouze et al., 2009; Rempfer et al., 2011; Gu et al., 2019; Pöppelmeier et al., 2020a, 2021b), which is shorter than the global overturning of the deep ocean. Unlike other tracers of ocean circulation (e.g. δ¹³C, Δ¹⁴C), Nd is not fractionated by marine biological cycling, giving rise to its promise as a carbon cycle independent ocean circulation tracer (Blaser et al., 2019a). Yet, a fundamental caveat in the application of ε_{Nd} as a
55 reliable oceanographic tracer is that there remains incomplete knowledge of the Nd fluxes in and out of the ocean, and that while internal cycling must occur, the efficiency of it with different particle types is poorly constrained (Abbott et al., 2015a; Haley et al., 2017; van de Flierdt et al., 2016).

Numerical models are useful tools for investigating Nd cycling since they can specify the processes that govern the
60 spatiotemporal variability in Nd isotope distributions in the ocean. Neodymium isotopes have been simulated in a range of different modelling studies testing specific hypotheses relating to Nd fluxes and thermohaline redistribution (Pöppelmeier et al., 2020a; Arsouze et al., 2009; Rempfer et al., 2011; Siddall et al., 2008; Jones et al., 2008; Roberts et al., 2010; Tachikawa et al., 2003; Ayache et al., 2016; Gu et al., 2019; Oka et al., 2021, 2009; Pöppelmeier et al., 2022; Ayache et al., 2022; Pasquier et al., 2022; Du et al., 2020). However, recent work suggests that a seafloor wide benthic flux, resulting from early diagenetic
65 reactions, may dominate the marine Nd cycle (Haley et al., 2017; Abbott, 2019; Abbott et al., 2015a, b, 2019; Du et al., 2016). These observations, alongside an ever-growing body of high-quality and highly-resolved measurements of dissolved seawater Nd concentrations ([Nd]) and ε_{Nd} from the GEOTRACES programme (GEOTRACES Intermediate Data Product Group, 2021), present an opportunity to re-evaluate, revise and explore constraints on the marine Nd cycle.

- Deleted: W
- Deleted: of
- Deleted: .
- Deleted: that
- Deleted: (Arsouze et al., 2009; Rempfer et al., 2011; Gu et al., 2019; Pöppelmeier et al., 2020a, 2021b).
- Deleted: the measured
- Deleted: isotope composition of seawater
- Deleted: actively
- Deleted: involved
- Deleted: of Nd
- Deleted: this
- Deleted: that a universal understanding of the exact mechanisms controlling marine geochemical Nd cycling remains incomplete

Initially, the predominant lithogenic fluxes of Nd to the ocean were believed to be mostly at the surface (aeolian dust and riverine fluxes; Goldstein et al., 1984; Goldstein and Jacobsen, 1987). Early modelling studies applying surface fluxes reproduced reasonable ϵ_{Nd} in the North Atlantic (Bertram and Elderfield, 1993; Tachikawa et al., 1999). However, considering only dust and river fluxes alone led to an unrealistic calculated residence time of Nd in seawater on the order of 5,000 years (Bertram and Elderfield, 1993; Jeandel et al., 1995). It was then found that considering dust and river surface inputs alone failed to balance both [Nd] and ϵ_{Nd} , thus indicating a ‘missing source’ of Nd to the ocean that accounted for $\approx 90\%$ of the Nd flux to the ocean (Tachikawa et al., 2003). This led to a new hypothesis relating to other Nd sources to seawater that could account for this ‘missing source’, including submarine groundwater discharge (SGD) (Johannesson and Burdige, 2007), and exchange of Nd between seawater and sediment deposited on the continental margins (Lacan and Jeandel, 2005). The term ‘boundary exchange’ was then coined to describe significant modification of Nd isotopic composition by the co-occurrence of Nd release from sediment and boundary scavenging, without substantially changing [Nd] (Lacan and Jeandel, 2005). Arsouze et al. (2007) simulated realistic global ϵ_{Nd} distributions using sedimentary fluxes constrained to the continental margins as the only source-sink term, and since then, particle-seawater exchange along the margins has represented the major flux of Nd to seawater in recent global Nd isotope models (Arsouze et al., 2009; Rempfer et al., 2011; Gu et al., 2019).

Nonetheless, seafloor-sediment interactions along the continental margins alone cannot fully reconcile the global marine Nd cycle. Specifically, it cannot explain the observed vertical profiles of [Nd], which are decoupled from ϵ_{Nd} (i.e., the ‘Nd paradox’: Goldstein and Hemming, 2003), with low concentrations near the surface increasing with depth. This is a common characteristic of isotopes/elements (e.g. thorium) that are reversibly scavenged (i.e., where the element is scavenged onto sinking particles at the surface and is subsequently remineralised in the deep ocean) (Bertram and Elderfield, 1993; Bacon and Anderson, 1982). Siddall et al. (2008) first addressed numerically a hypothesis that the ‘Nd paradox’ can be explained by a combination of lateral advection and reversible scavenging by applying the reversible scavenging model pioneered by Bacon and Anderson (1982) to Nd cycling. In their study, both [Nd] and ϵ_{Nd} were modelled simultaneously and explicitly to explore internal cycling of Nd in the deep ocean. Their findings demonstrated that scavenging and remineralisation processes are important active components in the marine cycling of Nd, driving the increase of [Nd] with depth, but still allowing ϵ_{Nd} to act as an effective water mass tracer.

Although inclusion of reversible scavenging can explain aspects of marine Nd cycling, the use of over-simplified fixed surface [Nd] and ϵ_{Nd} boundary conditions in the model by Siddall et al. (2008) limited what could be determined about the full marine cycling of Nd and hence the ‘Nd paradox’. The most comprehensive Nd isotope enabled ocean models to date now explicitly represent and quantify a wider range of distinct Nd fluxes that are both external and internal to the marine realm. For example, Arsouze et al. (2009) used a fully prognostic coupled dynamic and biogeochemical model to simulate [Nd] and ϵ_{Nd} , considering dust fluxes, dissolved riverine sources, sediment sources along the continental margins, and reversible scavenging. In their study, a sediment source from the continental margins represented the major source of Nd to seawater ($\approx 95\%$ of the total

Deleted: only

Deleted: Through applying a simple box model to calculate the oceanic Nd budget, i

Deleted: input from the dissolution of

Deleted: strong

Deleted: interactions

Deleted: though

Deleted: The term ‘boundary exchange’ was coined to describe strong Nd isotopic interactions between continental margins and water masses though the co-occurrence of sediment dissolution and boundary scavenging (Lacan and Jeandel, 2005).

Formatted: Not Strikethrough

Deleted: boundary

Deleted: exchange

Deleted:

Deleted: boundary

Deleted: along continental margins

Deleted: boundary exchange

Deleted: boundary exchange

Deleted: boundary

135 source). Rempfer et al. (2011) continued this work, undertaking a more detailed and comprehensive investigation of Nd sources and particle scavenging using a coarse resolution intermediate complexity model (Bern3D ocean model) and extensive sensitivity experiments. This later scheme was then closely followed by Gu et al. (2019) for the implementation of Nd isotopes in the ocean component of a more comprehensive Earth System Model (ESM, specifically the Community Earth System Model: CESM1.3) to explore in detail the changes to end-member ϵ_{Nd} signatures in response to ocean circulation and climate changes. Overall, these comprehensive models, capable of quantifying the major sources implicated in marine Nd cycling indicated that dust and river fluxes were important for representing [Nd] and ϵ_{Nd} distributions in the surface, but that the main flux of Nd to seawater is via a sediment source operating along the continental margins.

Recent pore fluid concentration profiles measured on the Oregon margin in the Pacific Ocean indicate that there may be a benthic flux of Nd from sedimentary pore fluids, presenting a potential major seafloor-wide (i.e., no longer limited to the continental margins) source of Nd to seawater (Abbott et al., 2015b, a); although there are observations of low [Nd] from southeastern equatorial Pacific pore waters, implying possibly heterogeneous benthic flux in relation to deep-sea sediment composition (Paul et al., 2019). Additional measurements from the Tasman Sea inferred the likely presence of a benthic source of similar magnitude to that estimated for the North Pacific, which may indicate that regions with dominantly calcareous sediment also contribute a significant benthic source of Nd (Abbott, 2019). Evidence of this previously overlooked benthic sedimentary source of Nd has led to a shifting paradigm which contends that the dominant addition of Nd to the ocean is from a diffuse sedimentary source at depth, rather than surface point sources from rivers and dust and the shallow-intermediate continental margins (Haley et al., 2017).

Simple box models have been employed to investigate, to a first order, the non-conservative effects from a benthic flux (Du et al., 2016, 2018, 2020; Haley et al., 2017; Pöppelmeier et al., 2020b), suggesting overprinting of deep water mass ϵ_{Nd} is linked to benthic flux exposure time and the difference between the Nd isotope composition of the benthic flux and the bottom water (Abbott et al., 2015a; Du et al., 2018). However, these models lack comprehensive descriptions of both the marine Nd cycle and of physical ocean circulation and climate interactions, limiting a clear interpretation of precisely how (and under what physical/environmental conditions) the benthic flux may determine global marine Nd distributions. Applying an intermediate complexity model, Pöppelmeier et al. (2021) investigated the benthic flux hypothesis in more detail by updating the Nd isotope enabled Bern3D model (Rempfer et al., 2011) to represent recent observations that indicate a Nd flux from bottom waters could occur across the entire seafloor. This was done by removing the depth limitation of the sediment source (previously 3 km) and invoking a constant benthic flux that escapes from all sediment-water interfaces. The scheme was further extended by revising key source-sink parameterisations, for subsequent investigation of non-conservative Nd isotope behaviour under different ocean circulation states (Pöppelmeier et al., 2022). The authors demonstrate substantial non-conservative effects occur even under strong circulation regimes with low benthic flux exposure times, and are not strictly limited to the deep ocean. This work highlights the importance of downward vertical fluxes via reversible scavenging alongside the benthic flux

Deleted: boundary

Deleted: ,

Deleted: a new,

Deleted: ly

Deleted:

Deleted: ,

Deleted:

Deleted: ity of

Deleted: suggest

Deleted: inferred

Deleted:

Deleted: (Haley et al., 2017). The bottom up model

Deleted: the shallow

Deleted: The benthic flux hypothesis provides a compelling yet unproved mechanism to explain deep water ϵ_{Nd} alteration alongside vertical [Nd] gradients in the North Pacific in the absence of modern deep-water formation, via exposure of old bottom water to a substantial benthic flux (Abbott et al., 2015a; Du et al., 2016).

Deleted: boundary exchange

185 to describe non-conservative marine Nd isotope behaviour. Nonetheless, the low horizontal resolution of the intermediate complexity model limits full resolution of key circulation patterns such as distinct deep-water formation in the Labrador and in the Nordic Seas, inhibiting the capabilities of the scheme to fully capture and explore water mass end member ϵ_{Nd} distributions.

190 Thus, despite substantial progress to explicitly describe seawater Nd budgets, it is apparent that outstanding questions remain, alongside divergent lines of argument amongst subject specialists, limiting a full, quantitative description of marine Nd cycling. The decoupled nature of marine [Nd] and ϵ_{Nd} , which is yet to be fully understood, and new emerging observations (e.g., Stichel et al., 2020; Lagarde et al., 2020a; Paffrath et al., 2021; Deng et al., 2022) emphasise the critical need to progress our understanding of modern marine Nd cycling, in the light of continued use of ϵ_{Nd} as a valuable tracer of ocean circulation. In this context, Nd isotope enabled ocean models remain an effective way to understand seawater measurements and to progress with testing and constraining processes governing marine Nd cycling, which can then feedback key information to the wider GEOTRACES and ocean-tracer modelling community.

200 With this in mind, there is a current gap in the toolkit for modelling marine Nd cycling between the class of high complexity, state-of-the-art ESMs (e.g., Gu et al., 2019) and the more efficient intermediate complexity models (e.g., Rempfer et al., 2011; Pöppelmeier et al., 2020a). To bridge this gap, there is a need for a model with the full complexity of an Atmosphere-Ocean General Circulation Model (AOGCM), allowing the exploration of how Nd isotopes vary under changing climate conditions (including extensive palaeoceanographic applications), that is also capable of running very quickly to facilitate efficient parameter space exploration, performance optimisation and long integrations. Owing to a winning combination of speed and complexity, the FAMOUS GCM fills this gap, and is well suited to exploring Earth system interactions (Smith et al., 2008; Jones et al., 2005; Smith, 2012; Jones et al., 2008).

205 Here we present the new global marine Nd isotope scheme (Nd_{v1.0}) implemented in FAMOUS. We utilise the new sediment ϵ_{Nd} maps from Robinson et al. (2021) as boundary conditions for a mobile global sediment Nd flux, with the end-goal of further constraining the major sources, sinks and cycling of Nd isotopes, and exploring instances of non-conservative behaviour related to changes in Nd source distributions and scavenging processes. We develop sensitivity experiments (Sect. 3) to isolate the physical effects of varying two key parameters that detail major Nd fluxes and cycling to the global ocean, verifying foremost that the new isotope scheme is responding to Nd source/sink and biogeochemical processes as expected, and contextualising how, and where, reversible scavenging processes and the benthic flux can govern marine [Nd] and ϵ_{Nd} distributions. Finally (Sect. 3 and 4), we evaluate overall model performance through comparison with modern measurements and assess the model's ability to simulate observed spatial and vertical gradients between ocean basins, encompassing underlying structural uncertainty/model bias.

Deleted: (e.g., Stichel et al., 2020)

Deleted: ,

Deleted: its

Deleted: T

Deleted:

Deleted: D

2 Methods

2.1 Model description

We use the FAMOUS GCM (Smith et al., 2008; Jones et al., 2005; Smith, 2012), a fast coupled AOGCM, derived from the Met Office's Hadley Centre Coupled Model V3 (HadCM3) AOGCM (Gordon et al., 2000). The atmospheric component of FAMOUS is based on quasi-hydrostatic primitive equations and has a horizontal resolution of 5° latitude by 7.5° longitude, 11 vertical levels on a hybrid sigma-pressure coordinate system and a 1-hour timestep. The ocean component is a rigid-lid model, with a horizontal resolution of 2.5° latitude by 3.75° longitude and 20 vertical levels, spaced unequally in thickness from 10 m at the near-surface ocean to over 600 m at depth, and a 12-hour timestep. The ocean and atmosphere are coupled once per day.

FAMOUS's parameterisations of physical and dynamical processes are almost identical to those of HadCM3, its parent model, but it has approximately half the spatial resolution and a longer timestep, allowing it to run ten times faster. Thus, FAMOUS achieves a current speed of up to 650 model years per wall clock day on 16 processors, making it ideal for running large ensembles (Gregoire et al., 2011, 2016), more bespoke sensitivity studies (Smith and Gregory, 2009; Gregoire et al., 2015), and multi-millennial simulations, e.g. to examine ice sheet behaviours (Gregoire et al., 2012, 2016, 2015), ocean drifts (Dentith et al., 2019) and biogeochemistry (Dentith et al., 2020). FAMOUS is calibrated to the performance of HadCM3, taking the philosophy that this is the most appropriate evaluation target and it is unrealistic to expect the lower resolution, lower complexity model to out-perform its parent model (Valdes et al., 2017; Smith and Gregory, 2009).

We added Nd isotopes (^{143}Nd and ^{144}Nd) as optional passive tracers into the ocean component of FAMOUS, using a version of the model with the Met Office Surface Exchange Scheme (MOSES) version 1 (Cox et al., 1999: FAMOUS-MOSES1). It was a pragmatic choice to avoid the more recent FAMOUS-MOSES2.2 configuration (Essery and Clark, 2003; Valdes et al., 2017; Williams et al., 2013; Essery et al., 2001), because evaluation of our new Nd scheme would be hindered by the collapsed Atlantic Ocean convection and strong deep Pacific MOC produced by FAMOUS-MOSES2.2 under pre-industrial boundary conditions (see Dentith et al., 2019). Nonetheless, the Nd isotope code implementation presented here is directly transferable to other versions of the UK Met Office Unified Model (UM) version 4.5, including HadCM3/L, or FAMOUS-MOSES2.2.

2.2 A new reference simulation for this study

The use of Nd isotopes as a water provenance tracer comes from measurements of distinct ϵ_{Nd} signatures in different water masses. This is perhaps best demonstrated in the south Atlantic 'zig-zag' depth profiles, where ϵ_{Nd} displays large heterogeneity and distinguishes southward flowing North Atlantic Deep Water (NADW) from northward flowing Antarctic Intermediate Water (AAIW) and Antarctic Bottom Water (AABW) (Goldstein and Hemming, 2003; Jeandel, 1993). As such, it is desirable

Deleted: its parent model.

Deleted:

to implement the Nd isotope scheme in a version of FAMOUS that best positions these basinal water masses in the correct locations.

255 The standard pre-industrial FAMOUS setup (XFHCC; Smith, 2012) has certain known limitations, including over-ventilated
abyssal Atlantic waters characterised by strong, over-deep NADW formation, with ‘North Atlantic’ convection occurring only
in the Norwegian-Greenland Sea (there is no deep water formation in the Labrador Sea), and insufficient Atlantic sector AABW
formation (Dentith et al., 2019; Smith, 2012). This known physical bias would dominate simulated Nd distributions, thus
hampering validation of the new Nd isotope scheme against modern measurements. To mitigate this, we chose to employ a
260 new reference simulation with improved basin-scale physical ocean circulation. This simulation was obtained from a perturbed
parameter ensemble varying 13 physical tuning parameters (see Supplementary Information: Text S1 and Table S1 for brief
description and for a list of perturbed parameters from this multi-sweep ensemble of FAMOUS MOSES1: Smith, 1990, 1993;
Crossley and Roberts, 1995). We screened the resulting 549 simulations based on a set of pre-defined targets, with a particular
focus on Atlantic Ocean structure and water mass composition. Specifically, we sought a simulation with appropriate modern
265 AMOC strength (14-19 Sverdrup, Sv; where $1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$) (Frajka-Williams et al., 2019), AMOC structure (Talley et al.,
2011), regions of AMOC convection as indicated by mixed layer depth (specifically in both the Labrador Sea and the Nordic
Seas as these represent key regions for NADW formation and the resultant end member Nd isotope compositions; Lambelet
et al., 2016), depth of maximum overturning ($\approx 1,000 \text{ m}$), and presence of AABW in the abyssal Atlantic (Frajka-Williams et
al., 2019; Talley et al., 2011; Kuhlbrodt et al., 2007; Ferreira et al., 2018). Furthermore, we assessed the capabilities of the
270 simulation to represent appropriate water mass structure in the Pacific (Talley et al., 2011). Through this approach, we
identified four possible candidate simulations from the large ensemble as the basis for the new Nd isotope scheme: XPDAA,
XPDAB, XPDAC and XPDEA, here denoted by their unique five-letter Met Office UM identifier. See Supplementary
Information: Table S2 for initial boundary conditions, and Figures S1-4 for Atlantic meridional stream function and mixed
layer depth for the four experiments.

275 These simulations were integrated for a further 5,000 years to ensure adequate spin-up of the physical circulation. They were
then assessed for their ability to simulate global modern observations of salinity and temperature (compared to the NOAA
World Ocean Atlas salinity and temperature database; Locarnini et al., 2019; Zweng et al., 2019) as an objective and
quantitative basis for selecting the best-performing parameter configuration to be our *control* (Fig. 1).




Figure 1: Taylor diagrams summarising the performance of the four control-candidate simulations (XPDA, XPDAB, XPDAC, XPDEA) in terms of their correlation, centred root mean square error, and ratio of their variances to the NOAA World Ocean Atlas (a) salinity and (b) temperature databases (Zweng et al., 2019). The simulations were selected from a large FAMOUS pre-industrial perturbed parameter ensemble following the circulation performance screening described in Sect. 2.2. XPCQX represents the initial experiment, upon which the four control-candidates aim to improve.

From this analysis, XPDA returned the lowest root mean square error (RSME) (and hence best performance using this metric;

Fig. 1) for simulating both salinity (RSME of 0.93 PSU) and temperature (RSME of 2.21 °C) and so was used as our control simulation (Fig. 2). For comparison, HadCM3, the parent model of FAMOUS (Gordon et al., 2000), and the Model for Interdisciplinary Research on Climate 4m (MIROC4m) AOGCM (Sherriff-Tadano et al., 2021) have a RMSE of 1.06 PSU and 1.47 °C and 0.59 PSU and 1.42 °C.

Formatted: Font: Times New Roman, Bold

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Field Code Changed

Formatted: Normal

Deleted: ; Pope et al., 2000

Field Code Changed

Deleted: that to similar/higher-order models, which is and givesto for understandings given our pragmatic choice to undertake the Nd-isotope scheme development with a fast GCM





Figure 2: (a) Salinity and (b) temperature profiles for the *control* simulation (centennial mean from final 100 years of a 5,000-year simulation) along a transect crossing the Pacific-Southern-Atlantic Ocean.

The simulated steady state AMOC strength in FAMOUS under fixed pre-industrial boundary conditions is approximately 17 Sv (Fig. 3a), which we consider to be in excellent agreement with direct modern AMOC observations from the RAPID AMOC array at 26.5° N of 17.2 Sv from April 2004-October 2012, and the depth of maximum meridional stream function at 26.5° N is around 800 m (Fig. 3b), slightly shallower than RAPID observations of 900-1100 m (McCarthy et al., 2015; Sinha et al., 2018). In terms of the Atlantic circulation structure, the overturning cell of NADW descends to depths of 3,000 m as it bridges into the South Atlantic, and AABW fills the bottom of the Atlantic basin with southern-sourced waters up to 20° N. For the Nd isotope implementation, all sensitivity studies and model tuning described subsequently are based upon this simulation, XPDAA.

Formatted: Normal

Deleted: ¶



305 **Figure 3:** Atlantic Meridional streamfunction for a 5,000-year simulation using *control* (XPDA), showing (a) the maximum
 310 streamfunction in time series (red dashed line indicates RAPID-AMOC 2004-2012 averaged AMOC strength at 26.5° N of 17.2 Sv;
 McCarthy et al., 2015), and (b) the zonal integration calculated from the last one hundred years.

2.3 Neodymium isotope implementation in FAMOUS

In our simulated Nd isotope scheme (Nd v1.0), we represent three different global sources of Nd into seawater: aeolian dust
 310 flux, dissolved riverine input and seafloor sediment. Neodymium (*Nd*) here refers to the sum of ^{143}Nd and ^{144}Nd , and the
 isotopic ratio (*IR*) relates to the ratio of ^{143}Nd to ^{144}Nd , as shown:

$$Nd = {}^{143}\text{Nd} + {}^{144}\text{Nd} \quad (2)$$

$$IR = \frac{{}^{143}\text{Nd}}{{}^{144}\text{Nd}} \quad (3)$$

By rearranging Eq. (2) and using the isotopic ratio (*IR*: Eq. (3)), individual fluxes of each isotope can be calculated (Eq. (4)
 315 and Eq. (5)) using information about Nd fluxes from each specific source and their associated *IR* distributions:

Deleted: D
 Deleted: dissolution of
 Deleted:

320

$$^{143}\text{Nd} = \frac{\text{Nd}}{(1+(1/IR))}$$

(4)

325

$$^{144}\text{Nd} = \frac{\text{Nd}}{(IR+1)}$$

(5)

330

Neodymium isotopes are thus simulated and transported individually and independently as two separate tracers in our scheme, explicitly resolving the concentration and distribution of each Nd isotope, allowing for [Nd] and ϵ_{Nd} to be calculated offline from the model output.

335

The implementation of each source/sink term is described in detail in the following sections. To summarise these different components, Nd sources from aeolian dust fluxes and dissolved riverine input enter the ocean only via the uppermost near-surface ocean layer of the model. Seafloor sedimentary fluxes, an umbrella term that refers to a multitude of processes encompassing input from sediment deposited on the continental margins (Lacan and Jeandel, 2005), submarine groundwater discharge (Johannesson and Burdige, 2007), and an abyssal (i.e., below 3 km) benthic flux released from pore waters (Abbott et al., 2015a), are simulated via a combination of a sedimentary source applied across sediment-water interfaces together with a separate sink occurring via particle scavenging. Removal of Nd from the ocean model occurs when Nd scavenged/adsorbed onto sinking biogenic and lithogenic (dust) particles reaches the seafloor via vertical fluxes and undergoes sedimentation, acting as an active sink by removing the particle-associated Nd from the ocean.

340

In the model numerics, fluxes of each Nd isotope into the ocean ($\text{kg m}^{-3} \text{s}^{-1}$) are multiplied by a factor of 10^{18} . This technique is a common heuristic to assist human comprehension. It minimises the mathematical error associated with carrying small numbers (e.g. when manually checking the code inputs and outputs). Accordingly, concentrations of each Nd isotope in model units are in ($10^{18} \text{ kg m}^{-3}$). A full list of all variables described in the text and their abbreviations are given in Table 1.

340

Table 1: Nd scheme model parameters, abbreviations, fixed model parameter values, and units.

Variable	Abbreviation	Fixed parameter value	Unit
Total Nd concentration	$[\text{Nd}]_t$		- pmol kg^{-1}
Dissolved Nd	$[\text{Nd}]_d$		- pmol kg^{-1}
Particle-associated Nd	$[\text{Nd}]_p$		- pmol kg^{-1}
Nd source, total	f_{total}		- g Nd yr^{-1}
Nd source, density	S_{total}		- g Nd $\text{m}^{-3} \text{yr}^{-1}$
Dust source, total	f_{dust}	0.33	Gg Nd yr^{-1}
Dust source, density	S_{dust}		- g Nd $\text{m}^{-3} \text{yr}^{-1}$

Deleted: It should be noted that of the Nd isotopes, ^{143}Nd and ^{144}Nd together only account for 36% of total Nd. As such, obtaining absolute fluxes of ^{143}Nd and ^{144}Nd from total Nd fluxes requires a scaling of 0.36, and correspondingly, to compare the total simulated Nd from Eq. (2) with observed total Nd requires scaling by 1/0.36. Due to this counteraction, unscaled fluxes can be used, i.e. total Nd fluxes are used in the model and as such, the sum of the simulated $^{143}\text{Nd} + ^{144}\text{Nd}$ (Eq. (2)) can be easily converted to observed [Nd] for direct comparison, as carried out previously by Gu et al. (2019); Pöppelmeier et al. (2020b); Rempfer et al. (2011).

Deleted: intense sediment-seawater interactions occurring along the boundary exchange the continental margins (Lacan and Jeandel, 2005),

Deleted: .

Deleted: s

Deleted: , and so

Deleted: (Nd)

Formatted Table

Deleted: .3

Deleted: $\times 10^8$

Flux of dust	F_{dust}	2D horizontal global field from Hopcroft <i>et al.</i> , 2015	$\text{g m}^{-2} \text{ yr}^{-1}$	
Nd concentration dust	\bar{C}_{dust}	20	$\mu\text{g g}^{-1}$	
Nd dust dissolution	β_{dust}	0.02		
Riverine source, total	f_{river}	0.44	Gg Nd yr^{-1}	Deleted: .
Riverine source, density	S_{river}	-	$\text{g Nd m}^{-3} \text{ yr}^{-1}$	Deleted: $\times 10^8$
River discharge	R_{RIVER}	-	$\text{g m}^{-2} \text{ yr}^{-1}$	Deleted: m^3
Riverine scaling factor	α_{river}	1		
Nd concentration river	C_{river}	-	$\mu\text{g g}^{-1}$	
Nd removal, estuaries	$\bar{\gamma}_{\text{river}}$	0.7		
Sediment source, total	f_{sed}	-	g Nd yr^{-1}	
Sediment source, density	S_{sed}	-	$\text{g Nd m}^{-3} \text{ yr}^{-1}$	
Total sediment surface	A_{total}	-	m^2	
Gridbox sediment surface	$A(i, k)$	-	m^2	
Gridbox volume	$V(i, k)$	-	m^3	
Thickness of euphotic layer	z_{eu}	81	m	
Penetration depth of opal	l_{opal}	10,000	m	
Penetration depth of CaCO3	l_{calcite}	3,500	m	
Particle settling velocity	ω	1,000	m yr^{-1}	
Ratio $[\text{Nd}]_{\text{p}}$ to $[\text{Nd}]_{\text{d}}$	$[\text{Nd}]_{\text{p}}/[\text{Nd}]_{\text{d}}$	-		
Global average density of seawater	ρ	1,024.5	kg m^{-3}	Deleted: p
Reversible scavenging, density	S_{rs}	-	$\text{g Nd m}^{-3} \text{ yr}^{-1}$	
Ratio between average POC concentration and density of seawater	R_{POC}	2.93×10^{-9}		
Ratio between average CaCO3 concentration and density of seawater	R_{CaCO3}	6.27×10^{-9}		
Ratio between average opal concentration and density of seawater	R_{opal}	5.21×10^{-9}		
Ratio between average dust concentration and density of seawater	R_{dust}	1.73×10^{-9}		
Total Nd inventory after equilibrium	$Nd(I)$	-	Tg	Deleted: 10^{12}

365 **2.3.1 Dust source**

Surface dust deposition (F_{dust}) is prescribed in the model from the annual mean dust deposition for the pre-industrial as simulated by the atmosphere component of the Hadley Centre Global Environmental Model version 2 (HadGEM2-A) GCM (Collins et al., 2011) (Fig. 4a). The dust deposition scheme (described by Woodward, 2011) has been shown to be in generally good agreement with observations, with concentrations in the Atlantic well simulated across the whole of the Saharan dust plume, although some discrepancies occur, including an overestimation at some Pacific sites during spring (Collins et al., 2011). The simulation of pre-industrial climate conditions within HadGEM2 are described by Hopercroft and Valdes (2015), and the dust results specifically are described in full by Hopercroft et al. (2015). Based on these simulated dust fluxes, we apply an Nd source per volume (S_{dust} : $\text{g m}^{-3} \text{yr}^{-1}$) in the uppermost layer of the ocean model, assuming a global mean concentration of Nd in dust \bar{C}_{dust} ($\bar{C}_{dust} = 20 \mu\text{g g}^{-1}$) (Goldstein et al., 1984; Grousset et al., 1988, 1998), from which only a certain fraction β_{dust} ($\beta_{dust} = 0.02$; Greaves et al., 1994) dissolves in seawater.


$$S_{dust}(i, k) = \frac{F_{dust}(i, 1) \times \bar{C}_{dust} \times \beta_{dust}}{dz(i, 1)} \quad (6)$$

Here i, k represents the horizontal and vertical indexing of model grid cell and dz is the grid cell's thickness (10 m in the uppermost surface layer where $k=1$). The total global flux of Nd from surface aeolian dust deposition to seawater (f_{dust}) is 0.33 Gg yr^{-1} . Although we use an updated dust deposition field compared to previous studies, prescribed \bar{C}_{dust} and β_{dust} are broadly consistent with earlier Nd isotope schemes, providing a comparable total Nd dust source ($0.1\text{--}0.5 \text{ Gg yr}^{-1}$; Arsouze et al., 2009; Gu et al., 2019; Pöppelmeier et al., 2020b; Rempfer et al., 2011; Tachikawa et al., 2003).

Deleted: k

- Deleted: .
- Deleted: $\times 10^8$
- Deleted: (Nd)
- Deleted: .0
- Deleted: $\times 10^8 \text{ g}$
- Deleted: .0
- Deleted: $\times 10^8$
- Deleted:
- Deleted: (Nd)

a) Annual dust flux



b) Dust ϵ_{Nd} regions




Figure 4: (a) Annual dust deposition taken from the pre-industrial annual mean dust deposition simulated by HadGEM2-A GCM (Hopcroft et al., 2015), and (b) ϵ_{Nd} signal from dust deposition following Tachikawa et al. (2003) and updated with information from Mahowald et al. (2006) and Blanchet (2019).

For the Nd isotope compositions of the dust flux, we started with the first-order estimate by Tachikawa et al. (2003), as follows: North Atlantic $> 50^\circ$ N: $\epsilon_{Nd} = -15$, Atlantic $< 50^\circ$ N: $\epsilon_{Nd} = -12$, North Pacific $> 44^\circ$ N: $\epsilon_{Nd} = -5$, Indopacific $< 44^\circ$ N: $\epsilon_{Nd} = -7$, and remainder: $\epsilon_{Nd} = -8$. This was revised with additional constraints, accounting for the dust plume expansions as reported by Mahowald et al. (2006), in combination with detrital Nd isotope data by Blanchet (2019) (Fig. 4b). For this, the mean detrital

Deleted: the mean Nd isotope signatures of the respective source regions as determined by the global compilation of

405 isotopic signature of a source region (e.g., the Sahara) was calculated and then applied to the region where this dust is deposited over the ocean based on the plume expansion by Mahowald et al. (2006) (for the example of the Sahara this is the North Atlantic). Regions in between major dust plumes were linearly interpolated. For the polar regions no constraints are available, and a default value has been assigned instead, but the lack of dust deposition in these regions ensures that this has no impact on the Nd cycle.

2.3.2 Dissolved riverine source


410 To represent the Nd source from dissolved river fluxes, we used the river outflow to the ocean simulated by FAMOUS (*RIVER*) as our river water discharge ($\text{g m}^{-2} \text{yr}^{-1}$) and combined this outflow with both the Nd concentration (C_{river} ; $\mu\text{g g}^{-1}$) and isotopic concentration (used to calculate the flux from each Nd isotope using Eq. (4) and Eq. (5)) of river water dissolved material, as estimated by Goldstein and Jacobsen (1987; see Table 3). All river source Nd fields are shown in Fig. 5.

Formatted: Normal

Deleted: k

Deleted: s

Deleted: ¶



415 Figure 5: (a) Simulated river outflow (RIVER) in FAMOUS, (b) major river ϵ_{Nd} , (c) major river [Nd], and (d) the resulting riverine Nd source. The coastal grids in (b) and (c) are prescribed following average [Nd] and ϵ_{Nd} estimates of dissolved river runoff to each of the oceans by Goldstein and Jacobsen (1987; see Table 3).

Deleted: and
 Deleted: ;
 Deleted: t
 Deleted:
 Deleted: ^{CO2} see
 Deleted: Table 3)
 Deleted: .
 Field Code Changed

425 River outflow in FAMOUS is based on a routing scheme that instantaneously delivers terrestrial runoff (precipitation - [evaporation + soil moisture]) from the location that precipitation falls to the designated coastal grid cell that the runoff would reach due to river routing (i.e., the river mouth). Relating the riverine Nd flux to the model's prognostic river discharge allows the Nd river source to respond to different climatic conditions, making it a more dynamic and predictive tool for examining the impact of changes in the global hydrological balance, such as wetting/drying events, or shifts in the monsoons. Essentially, river outflow plumes are 'tagged' with the estimated ε_{Nd} (which is provided as an input map along the coasts; Fig. 5b) according to the model's projected water discharge at that location and C_{river} (also provided as an input map along the coasts; Fig. 5c). For paleo or future climate applications where river routing is significantly different to today, the input maps controlling C_{river} tagging at the coast would need to be updated to reflect Nd dissolution from the modified fluvial pathway over land, the model should predict the rest.

435 Estuaries are important biogeochemical reactors of rare earth elements (REEs), with sea salt driving flocculation of river-dissolved organic matter, which results in estuarine REE removal (Elderfield et al., 1990). This removal is known to be important in balancing the marine REE budgets, however the complex processes involved are still not fully constrained (Elderfield et al., 1990). Rousseau et al. (2015) summarised published observations of Nd removal (%) in estuaries from observations of estuarine dissolved Nd dynamics (Rousseau et al., 2015; their Table 1). Published values (n = 17) range from 40% in Tamar Neaps (Elderfield et al., 1990) to 97% in the Amazon (Sholkovitz, 1993), with a mean Nd removal of $71 \pm 16\%$ (s.d.). Based upon this, and parallel to previous model schemes (e.g., Rempfer et al., 2011; Gu et al., 2019; Arsouze et al., 2009), we assume that 70% of dissolved Nd from river systems are removed in estuaries (i.e., $\bar{\gamma}_{river} = 0.7$).

The dissolved riverine source per unit volume of Nd (S_{river} ; $g\ m^{-3}\ yr^{-1}$) in the uppermost layer of the ocean is therefore calculated as:

445
$$S_{river}(i, k) = \frac{RIVER(i, 1) \times C_{river} \times (1.0 - \bar{\gamma}_{river})}{dz(i, 1)} \quad (7)$$

The total global flux of river sourced dissolved Nd to seawater in the model (f_{river}) is $0.44\ Tg\ yr^{-1}$. Previous Nd isotope schemes have applied either fixed annual mean continental river discharge estimates from Goldstein and Jacobsen (1987) and Dai and Trenberth (2002), as applied in the Bern3D Nd isotope scheme (Rempfer et al., 2011; Pöppelmeier et al., 2021a), or, similar to this study, using the model's own river routing and discharge schemes, as with NEMO-OPA (Arsouze et al., 2009) and CESM1 (Gu et al., 2019). Our estimated global total dissolved riverine Nd source to the oceans sits within previous model estimates ($0.26 - 1.7\ Tg\ yr^{-1}$; Arsouze et al., 2009; Gu et al., 2019; Pöppelmeier et al., 2020b; Rempfer et al., 2011; Tachikawa et al., 2003).

Deleted: a

Deleted: ε_{Nd} and

Deleted: k

Deleted: .

Deleted: $\times 10^8$

Deleted: (Nd)

Deleted: (

Deleted: .

Deleted: $\times 10^8$

Deleted: .

Deleted: \times

Deleted: 10^9

Deleted: (Nd)

Deleted: .

There is a larger range in estimated riverine Nd flux to the ocean relative to the estimated dust flux ranges. It should be noted that the largest simulated Nd river source amongst these studies (1.7 Gg yr^{-1} ; Pöppelmeier et al., 2020b) in the updated Bern3D Nd isotope scheme applied a river scaling factor, used as a tuning parameter and based on findings by Rousseau et al. (2015), who suggest a globally significant release of Nd to seawater by dissolution of river sourced lithogenic suspended sediments grounded upon observations in the Amazon estuary. Our model does not attempt to fully resolve all complex estuarine processes, and in this ~~study~~, we chose to represent the dissolved riverine flux as a single source to seawater. Furthermore, our sediment Nd source to the ocean (described Sect. 2.3.3), which occurs across sediment-water interfaces, utilises the continental margin and seafloor ϵ_{Nd} distribution maps by Robinson et al. (2021), thus using the most recent compilation of published global observations of Nd isotope compositions of river sediment samples deposited on the continental shelf and slopes (alongside geological outcrops and marine sediment samples). It is therefore likely that this margin source encompasses at least in part the Nd isotope fingerprint from a river particle flux.

2.3.3 Continental margin and seafloor sediment source

The sediment source describes the flux of Nd into seawater entering each model grid cell adjacent to sediment, this source is not restricted to the uppermost surface layers and can be implemented across all vertical and horizontal sediment-water interfaces, as is the case for all experiments described in this study. As such, the sediment source represents (1) ~~input from sediment deposited on the continental margins (Lacan and Jeandel, 2005)~~; (2) submarine groundwater discharge, as suggested by Johannesson and Burdige (2007), which releases Nd to seawater via discharge of fresh groundwater to coastal seas and is mainly limited to the upper 200 m; and (3) a benthic flux, which specifically refers to a transfer of Nd from sediment pore water to seawater resulting from early diagenetic reactions and is not depth limited (Abbott et al., 2015b, a; Haley et al., 2017).

The total sediment Nd source per unit volume (S_{sed} : $\text{g m}^{-3} \text{ yr}^{-1}$) into any given ocean grid cell is dependent on the fraction of the surface area of that cell that is in contact with sediment. Similar to previous schemes (e.g., Gu et al., 2019; Pöppelmeier et al., 2020b; Rempfer et al., 2011), the total globally integrated sediment-associated Nd source to the ocean (f_{sed} : g Nd yr^{-1}) is used as a tuning parameter.

$$S_{\text{sed}}(i, k) = f_{\text{sed}} \times \frac{A(i, k)}{A_{\text{total}}} \times \frac{1}{V(i, k)}, \quad (8)$$

where $A(i, k)$ is the total area of the sediment surface in contact with seawater per ocean grid cell (m^2), A_{total} is the total global area of the sediment surface where a sediment source occurs (m^2) and $V(i, k)$ is the volume of water per ocean grid cell (m^3).

There remains no true consensus on whether (and how) to apply spatial variability to sediment Nd sources, as reflected in the way previous schemes have adopted different approaches. Arsouze et al. (2009) limited the sediment flux to the upper 3,000 m of the ocean, imposing a depth scaling factor, and considering estimated [Nd] distributions across the continental margins.

Deleted: $\times 10^9$

Deleted:

Deleted: (Nd)

Deleted: study

Deleted: -Early findings by Goldstein and Jacobsen (1987) document that the Nd isotope composition of dissolved and suspended river loads can vary by up to four epsilon units. Observations presented by Rousseau et al. (2015) showed the measured Amazon dissolved river end-member value ($\epsilon_{\text{Nd}} = -8.9$) was more radiogenic than the typical suspended river material ($\epsilon_{\text{Nd}} = -10.6$) and as such, combining dissolved and particulate sources in river ϵ_{Nd} budgets is non-trivial.

Formatted: Not Strikethrough

Deleted: Notwithstanding, our model does permit a scaling of the Nd river flux (a_{river}), which, although out of the scope of this study, could be applied in future model development to explore in more detail a particulate river source as a major Nd flux to seawater.

Deleted: boundary exchange, as defined by Lacan and Jeandel (2005b), which describes the flux of Nd from particle-seawater exchange occurring predominantly across the continental margins

Formatted: Font: (Default) Times

Deleted: k

Deleted: (to represent 'boundary exchange' processes)

Deleted: using the earlier marine margin Nd compilation by Jeandel et al. (2007)

Both Rempfer et al. (2011) and Gu et al. (2019) simplified this method by applying spatially uniform sediment Nd fluxes, also limited to the upper 3,000 m. In more recent work, Pöppelmeier et al. (2020b) removed the depth limitation (as we have done), and incorporated a geographically-varying scaling factor to try to capture more localised features through increased benthic fluxes (Abbott et al., 2015b; Blaser et al., 2020; Grenier et al., 2013; Lacan and Jeandel, 2005; Rahlf et al., 2020). Pasquier et al. (2022) presented the first inverse model of global marine cycling of Nd, and the strength of the sediment Nd flux to the ocean was imposed with an exponential depth function, resembling eddy kinetic energy and particulate organic matter fluxes, which are characteristically larger near surface and coastal regions.

For our scheme, we do not assume spatial variations in the sediment source of Nd (f_{sed}), essentially avoiding making explicit inferences on the nature of the sedimentary Nd source. It has been proposed that preferential mobilisation of certain components of the sediment drive spatial variations in sediment fluxes (e.g., Abbott et al., 2015a; Wilson et al., 2013; Du et al., 2016; Abbott, 2019; Abbott et al., 2019), and that both detrital and authigenic phases likely exchange Nd within pore water during early diagenesis (Du et al., 2016; Blaser et al., 2019b). However, the elusiveness of marine Nd cycling alongside our limited knowledge of the specific mechanisms controlling sediment-water Nd interactions mean that determining generalisable rules for where and under what conditions (e.g., redox environments or fresh labile detrital material) preferential mobilisation may occur is unknown and challenging to resolve. Therefore, in accordance with Pöppelmeier et al. (2020a), Du et al. (2020), Gu et al. (2019) and Rempfer et al. (2011), we adopt a constant detrital sediment flux as a first-order approximation. In fact, we contend that applying this simpler method, as opposed to constructing a more complex source term that is arguably just as arbitrary (given the uncertainty in Nd cycling), allows for a more explicit quantification of differences between observed and simulated Nd distributions. As such, without overfitting our model, we allow for the clearest indication of those parts of the system that are well understood (and represented), and those which prove deficient. Under this framework, we may separate out and test the effect of many of the major Nd sources/sinks with dedicated sensitivity simulations, including the possibility, in future work, of incrementally modifying the sediment source distributions to increase the complexity of the scheme and assess the impact of our various assumptions.

The isotopic ratio of the sediment Nd flux to seawater is prescribed using the recent updated global gridded map of bulk detrital ϵ_{Nd} at the continental margins and seafloor (Robinson et al., 2021, Fig. 6a). Using Eq. (4) and Eq. (5), fluxes of each Nd isotope are calculated from this condition and bi-linearly regridded to the model's native resolution (Fig. 6b).

Deleted: that extrapolated modern Nd flux observations

Deleted: Pasquier et al. (2021)

Deleted: (

Deleted: ,

Deleted: in this latest scheme,

Deleted: . The flux per unit area is uniform with depth, latitude and longitude

Deleted: exchange

Deleted: means

Deleted: Previous studies used the predecessor continental ϵ_{Nd} map by Jeandel et al. (2007) (e.g. Arsouze et al., 2009, 2007; Gu et al., 2019; Rempfer et al., 2011); see Fig. 5 in Robinson et al. (2021) for a comparison of the differences). Marginal ϵ_{Nd} distributions in our improved map are broadly more radiogenic across the Arctic Shield, Northern Eurasia, South America, north-eastern Africa and Antarctica and more unradiogenic over southern Greenland, north-eastern Europe, western and eastern Africa and parts of the Americas. Crucially, the newer map of Robinson et al. (2021) provides the first estimate of global seafloor sediment Nd isotope composition, which is necessary for considering a global (e.g., rather than depth limited) benthic flux.




Figure 6: (a) Map of the global ϵ_{Nd} distributions at the sediment-ocean interface from Robinson et al. (2021), and (b) as used as a model input in this study (bi-linearly regridded from (a) onto the coarser FAMOUS ocean grid).

Pasquier et al. (2022) first applied the sedimentary ϵ_{Nd} map from Robinson et al. (2021) in their recent global marine Nd isotope scheme, imposing positive (i.e., radiogenic) modifications to the Pacific sedimentary ϵ_{Nd} and using a reactivity scaling factor (linked to sediment lability) that favours more extreme ϵ_{Nd} signals. Here, we again adopt a simpler approach, imposing the unmodified sediment ϵ_{Nd} distributions from Robinson et al. (2021), allowing for a presentation of our new scheme based on what we confidently know about Nd cycling without the complication of over-conditioning our model inputs. Future research may then use this work as a foundation to robustly explore different choices for model inputs (i.e., boundary conditions) to revisit these fundamental questions about Nd sediment source.

Deleted: (

Deleted: ,

Deleted: Pasquier et al. (2021)

2.3.4 Internal cycling via reversible scavenging

Vertical cycling and removal of Nd from the water column via sinking particles ('reversible scavenging') is parameterised using the same approach as previous Nd isotope implementations (Siddall et al., 2008; Arsouze et al., 2009; Rempfer et al., 2011; Gu et al., 2019; Pöppelmeier et al., 2020a; Oka et al., 2021). Based on the original scheme of Bacon and Anderson (1982), the method captures the physical process of absorption/incorporation and desorption/dissolution of Nd onto particle surfaces in seawater. The scheme assumes that particle associated Nd is in dynamic equilibrium with falling particles throughout the water column, with continuous exchange between the particle and dissolved pools. This process redistributes Nd within the water column, acting as a net sink of dissolved Nd at shallower depths as they adsorb onto particle surfaces, and a net source at greater depths where dissolution of particles releases dissolved Nd back to seawater. The scavenging is the only sink of Nd in our model; Nd associated to particles (particulate organic carbon, POC; calcium carbonate, CaCO_3 ; opal; dust) reaching the seafloor is removed from the system, operating under a steady state assumption that acts to balance the three external input sources (marine sediments, dust and rivers). Previous studies (e.g., Siddall et al., 2008; Arsouze et al., 2009; Rempfer et al., 2011; Gu et al., 2019; Wang et al., 2021) have demonstrated that reversible scavenging is an active and important component in global marine Nd cycling, and is necessary for successfully simulating both $[Nd]_d$ and ϵ_{Nd} distributions.

Updating the approach employed by Siddall et al. (2008), we prescribed individual biogenic particle export fields based on satellite-derived primary production. FAMOUS does contain an optional ocean biogeochemistry module (Hadley Centre Ocean Carbon Cycle; HadOCC), which includes simplified nutrient-phytoplankton-zooplankton-detritus (NPZD) classes (Palmer and Totterdell, 2001) and could instead be used as the basis for predicting vertical particle fluxes in the ocean (which was the approach adopted by Arsouze et al., 2009; Gu et al., 2019; and Rempfer et al., 2011). We favoured satellite-derived estimates in order to improve the accuracy of particle-associated cycling of Nd and reduce biases inherent to the intermediate complexity biogeochemistry model (Dentith et al., 2020; Palmer and Totterdell, 2001). This approach also optimises the computational efficiency of our scheme.

In our scheme, biogenic particle fields (POC, CaCO_3 , and opal) are prescribed using gridded, global satellite-derived particle export productivity from Dunne et al. (2012, 2007). The euphotic zone (z_{eu}), is set to a globally uniform depth in FAMOUS of 81 m, the closest bottom grid box depth to match that defined in the OCMIP II protocol (75 m) (Najjar and Orr, 1998). Below z_{eu} , appropriate depth-dependent dissolution profiles, derived from assumptions of particle degradability and sinking speed (Martin et al., 1987; Laws et al., 2000; Behrenfeld and Falkowski, 1997), and widely used to model flux attenuation in ocean models (e.g. all models used in the Ocean-Carbon Cycle Model Intercomparison Project; Doney et al., 2004; Sarmiento and LeQuéré, 1996), were applied to the biogenic export fluxes.

Downward fluxes of POC (F_{POC}) follow the power-law profile of Martin et al. (1987):

Deleted: , since the added ecosystem and geochemistry tracer fields in HadOCC slows the model down. However, future work could modify our implementation to instead associate Nd cycling with the prognostic particle fields in HadOCC.

$$F_{POC}(Z) = F_{POC}(Z_{eu}) \times \left(\frac{Z}{Z_{eu}}\right)^{-\alpha} \text{ (for } Z > Z_{eu} \text{),} \quad (9)$$

615 where z is depth (m), and α represents a dimensionless dissolution constant for POC set to 0.9 (Najjar and Orr, 1998). Although a widely used parameterisation of dissolution, it should be noted this so-called ‘Martin curve’ is known to underestimate the flux to the sediment in the off-equatorial tropics and subtropics and overestimates the flux in subpolar regions, indicating particles penetrate deeper than the Martin curve in the tropics and shallower in sub-polar regions (Dunne et al., 2007).

620 Downward fluxes of opal (F_{opal}) and CaCO_3 (F_{CaCO_3}) follow exponential dissolution profiles with particle-specific length scales l_{opal} (10,000 m) and l_{calcite} (3,500 m) (Maier-Reimer, 1993; Henderson et al., 1999; Najjar and Orr, 1998):

$$F_{opal}(Z) = F_{opal}(Z_{eu}) \times \exp\left(\frac{Z_{eu}-Z}{l_{opal}}\right), \quad (10)$$

$$F_{\text{CaCO}_3}(Z) = F_{\text{CaCO}_3}(Z_{eu}) \times \exp\left(\frac{Z_{eu}-Z}{l_{\text{CaCO}_3}}\right), \quad (11)$$

625 The sinking of dust is prescribed according to the pre-industrial annual mean dust deposition simulated by Hopcroft et al. (2015) (see Sect. 2.3.1). We assume that dust does not dissolve significantly with depth and so dust export fluxes are constant throughout the water column. In line with previous schemes (e.g. Arsouze et al., 2009; Pöppelmeier et al., 2020b; Rempfer et al., 2011; Siddall et al., 2008), a uniform settling velocity (ω) of 1,000 m yr⁻¹ is applied to all particle fields, although we acknowledge this is a simplification, the uniform settling rate applied captures the mean particle flux to the seafloor.

630 Annual averaged export of biogenic fields are shown in Fig. 7 and the total annual export production of POC (9.6 Gt(C) yr⁻¹), CaCO_3 (0.45 Gt(C) yr⁻¹), and opal (90 Tmol(Si) yr⁻¹) are comparable with previous estimates, although export of CaCO_3 and opal are at the lower end, as highlighted in Table 3 by Dunne et al. (2007). Note, the annual export of CaCO_3 reflects the new optimised surface calcite parameterisation as described in Dunne et al. (2012).

Deleted: ,

Deleted: ing

Deleted: .

Deleted:




Figure 7: Particle export fields from the ocean surface ($\text{g m}^{-2} \text{yr}^{-1}$). Biogenic particle fields are prescribed using satellite-derived export productivity fields for (a) POC, (b) CaCO_3 and (c) opal (Dunne et al., 2007, 2012). The dust input fields (d) are annual mean dust deposition simulated for the pre-industrial by the HadGEM2-A GCM (Hopcroft et al., 2015). Note the different scale used for panel (a).

Reversible scavenging (Nd_{rs}) considers total Nd for each isotope (j) as the dissolved Nd (Nd_d) and particulate Nd (Nd_p) associated with the different particle fields (χ ; where χ = POC, CaCO_3 , opal and dust).

$$[Nd]_c^j = [Nd]_d^j + [Nd]_p^j = [Nd]_d^j + \sum_{\chi} [Nd]_{p,\chi}^j \quad (12)$$

Nd_p sinks in the water-column with the particles due to gravitational force. Dissolution of biogenic particles with increasing depth below the euphotic zone releases Nd incorporated/adsorbed onto particles back to seawater (i.e., the Nd is dissolved/desorbed). Thus, particles act as internal sinks for marine Nd_d in shallower depths and as sources at greater depths. This combined process for reversible scavenging (S_{rs}) in the model can be described by:

$$S_{rs}(i, k) = -\frac{\partial(\omega[Nd]_p(i, k))}{\partial z(i, k)} \quad (13)$$

where $[Nd]_p$ can be determined within the model from total Nd for each isotope:

$$[Nd]_p^j(i, k) = [Nd]_c^j(i, k) \times \left(1 - \frac{1}{1 + \sum_{\chi} R_{\chi}(i, k) \times K_{\chi}^j}\right) \quad (14)$$

Deleted: w

655 $R_{\chi}(i,k)$ describes the dimensionless ratio between particle concentration for each particle type ($C_{\chi}(i,k)$) and the average density of seawater ($\rho = 1024.5 \text{ kg m}^{-3}$), input as fixed boundary conditions in our scheme. C_{χ} is calculated from the prescribed particle fluxes (F_{χ} ; Fig. 7) by assuming a globally uniform settling velocity ($\omega = 1,000 \text{ m yr}^{-1}$; Arsouze et al., 2009; Dutay et al., 2009; Gu et al., 2019; Kriest and Oeschies, 2008; Rempfer et al., 2011), i.e., $C_{\chi} = F_{\chi} / \omega$. The equilibrium between dissolved concentration ($[Nd]_d$) and concentration associated to particle-type ($[Nd]_p$) can be described by the equilibrium partition coefficient (K_{χ}^j):

$$K_{\chi}^j = \frac{[Nd]_{p,\chi}}{[Nd]_d} \times \frac{1}{R_{\chi}}, \tag{15}$$

660 here $[Nd]_p/[Nd]_d$ represents the scavenging efficiency in the model. It is independent of particle type and is used as a tuning parameter. R_{χ} , however, is dependent on particle type (where $R_{\chi} = \bar{C}_{\chi}/\rho$) and thus K_{χ} is different for different particles. Our global mean particle concentrations (\bar{C}_{χ} ; Table 2) are similar to those reported in previous schemes.

665 **Table 2: Global mean particle concentrations for each particle type used to calculate equilibrium scavenging coefficients following Eq. (15). In summary export fluxes of POC, CaCO_3 and opal in this study are from Dunne et al. (2007; 2012) and dust fluxes are from Hopcroft et al. (2015). Global mean particle concentrations reported in previous Nd isotope schemes are shown for comparison.**

Global mean particle concentration (kg m^{-3})				
Particle type	Acronym	this study	Rempfer et al. (2011)	Gu et al. (2019)
POC	\bar{C}_{POC}	3.00×10^{-6}	3.30×10^{-6}	2.66×10^{-6}
CaCO_3	\bar{C}_{CaCO_3}	6.43×10^{-6}	1.60×10^{-6}	9.53×10^{-6}
Opal	\bar{C}_{opal}	5.33×10^{-6}	5.90×10^{-6}	8.30×10^{-6}
Dust	\bar{C}_{dust}	1.78×10^{-6}	1.30×10^{-6}	1.23×10^{-6}

670 Isotopic fractionation during absorption/incorporation and desorption/dissolution is neglected due to similar masses of ^{143}Nd and ^{144}Nd , avoiding undue complexity arising from any assumption about preferential scavenging (Siddall et al., 2008). Adsorption occurs everywhere that particles are present and we do not allow for preferential scavenging onto different particle types, consistent with previous models (e.g. Rempfer et al., 2011).

Therefore, and by including advection and diffusion processes (*Transport*), it follows that the total conservation equation for each Nd isotope in the model scheme can be written as:

Deleted: p

Deleted: π

Deleted: ; see Supplementary Information: Table S3 for a comparison

Formatted: Subscript

Deleted: Average

Deleted: Particle type

... [1]

Deleted: average

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted Table

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Deleted: ¶

Deleted: In contrast, Siddall et al. (2008) optimised K_{χ} to fit observed $[Nd]_d$ for each particle type. Their optimised solution effectively implied no scavenging of Nd by POC, this result was considered tentative due to similar dissolution profiles of CaCO_3 and POC, which likely biased scavenging to CaCO_3 that may have been more correctly attributable to POC.

690

$$\frac{\delta [Nd]_d^j}{\delta t} = S_{dust}^j + S_{river}^j + S_{sed}^j + S_{rs}^j + Transport,$$

(16)

2.4 Evaluation methods & data sets

To validate the new Nd isotope scheme (Nd v1.0) and to assess the model’s performance, we compare the simulated $[Nd]_d$ and ϵ_{Nd} to modern seawater measurements, with a focus on describing the ability of the model to represent key spatial and vertical distributions across ocean basins.

695

As part of this assessment, a basic indication of model skill is returned by the mean absolute error (MAE):

$$MAE = \frac{1}{N} \sum_{k=1}^N |obs_k - sim_k|,$$

(17)

where obs_k and sim_k are measured and simulated $[Nd]_d$ or ϵ_{Nd} respectively. N is the number of observations, and k is an index over all observational data. For each measurement – based on its longitude, latitude, and depth – the value predicted by the model is extracted and the mean deviation of simulated and observed $[Nd]_d$ and ϵ_{Nd} is presented in pmol kg⁻¹ and epsilon units respectively. Here we chose specifically not to apply a grid box volume weighting to the MAE, which would act to emphasise abyssal Pacific results in our assessment of model skill, where there are few observations and relatively low variability in Nd distributions. The advantage of using an unweighted MAE is that the assessment metric better scrutinises regions with larger (spatial) gradients in both $[Nd]_d$ or ϵ_{Nd} ; i.e., at the surface and high latitudes. We also report root mean square error (RMSE) for each simulation as an additional indicator of model skill and to allow for additional comparability with other studies:

705

$$RMSE = \sqrt{\frac{1}{N} \sum_{k=1}^N \frac{(obs_k - sim_k)^2}{N}}$$

(18)

The observational data used in this assessment are from the seawater REE compilation used by Osborne et al. (2017, 2015), augmented with more recent measurements including data in the GEOTRACES Intermediate Data Product 2021 (GEOTRACES Intermediate Data Product Group, 2021) from GEOTRACES cruises (GA02, GA08, GP12, GN02, GN03 and GIPY05). Combined, our observational database represents a total of 6,048 $[Nd]_d$ and 3,278 ϵ_{Nd} measurements, making it the largest compilation of seawater Nd data used to date to validate the performance of an Nd isotope enabled model. Notably, we omit measurements of $[Nd]_d > 100$ pmol kg⁻¹ from the model data comparison because they represent very localised signals which we do not attempt to resolve. These include extreme surface concentrations present in restricted basins such as fjords, the Baltic Sea and the Gulf of Alaska (Chen et al., 2013; Haley et al., 2014), or input from hydrothermal activity (Chavagnac et al., 2018), which is not believed to govern global marine Nd distributions due to the immediate removal of hydrothermal Nd at the vent site (Goldstein and O’Nions, 1981). The location and spatial distribution of all observational records used in this study are shown in Fig. 8, and full details of the seawater compilation including a full list of all the references for the data sources are provided in Supplementary Information: Table S3.

Deleted: D

Deleted:

Formatted: Font: Italic

Deleted: , because/since...

Deleted: better

Deleted: studies.The


Formatted: Space Before: 6 pt, After: 6 pt

Formatted: Font: 10 pt


Formatted: Font: (Default) +Body (Times New Roman)

Deleted: 4

a) [Nd] observations



b) ϵ_{Nd} observations



725 **Figure 8:** Location of marine observational records used in this study, (a) filled orange triangles show the location of dissolved Nd concentration records, and (b) filled sky-blue circles show the location of dissolved ϵ_{Nd} records.

Neither the horizontal nor vertical distribution of global seawater $[Nd]_d$ and ϵ_{Nd} observational data are even. Most $[Nd]_d$ measurements are in the Pacific Ocean, in comparison to ϵ_{Nd} measurements where they are most frequent in the Atlantic Ocean.

Deleted: followed by the Atlantic Ocean (representing 37% and 35% of measurements respectively, noting that the Pacific has more than twice the volume of the Atlantic), with far fewer measurements in the Indian Ocean (12%) and the Southern Ocean (5%).

Deleted: I

Deleted: ,

735 Both $[Nd]_d$ and ϵ_{Nd} observational data are biased towards shallower depths. It is therefore important to highlight that, as with previous studies, skew in the distribution of seawater Nd measurements will act to bias our assessment of model performance.

In some instances, near land grid cells, the location of the compared measured and modelled Nd may not match due to the coarseness of the model grid. In such cases, we employ a nearest neighbour algorithm to extract the modelled value from the closest ocean model grid cell. Furthermore, if multiple measurements occur within one model grid cell, the arithmetic mean of the values is used for our comparison to model results, and as such, $n = 3,471$ and $2,136$ for the calculation of both $MAE_{[Nd]}$ and $MAE_{\epsilon_{Nd}}$ respectively.

To ensure that our evaluation is not overly reliant on the cost-function analysis alone, and hence evaluated the spatial biases in our assessment from the geographically uneven spread of measured $[Nd]_d$ and ϵ_{Nd} , we also assess the capability of the model to reproduce appropriate global Nd inventories and to simulate large-scale horizontal and vertical gradients. We compare our results briefly with findings from previous modelling studies, but we highlight that the purpose of this study is to understand the behaviour of our model, and not to undertake a comprehensive calibration of its performance. Optimisation (or ‘tuning’) of the Nd scheme will follow this work, and this needs to be considered when comparing the cost function performance to previous schemes.

2.5 Sensitivity experiment design

750 We designed a number of sensitivity experiments (Table 3) to systematically vary individual model parameters describing the reversible scavenging efficiency ($[Nd]_p/[Nd]_d$) and the main Nd source (f_{sed}). These two parameters ($[Nd]_p/[Nd]_d$ and f_{sed}) were chosen primarily as they represent important and largely unconstrained non-conservative processes that are understood to govern simulated global distributions of both seawater $[Nd]_d$ and ϵ_{Nd} (Rempfer et al., 2011; Pöppelmeier et al., 2020a; Gu et al., 2019; Arsouze et al., 2009; Siddall et al., 2008). By isolating individual effects, the primary aim was to understand in detail the model’s sensitivity to different forcings, identify which parameters are important for $[Nd]_d$ and/or ϵ_{Nd} patterns, and to identify assumptions within the explored parameters that require further constraining (through further field campaign, laboratory analysis or model experimentation).

Deleted: (45%), followed by Pacific Ocean (29%), and again there are fewest measurements from the Indian Ocean (8%) and Southern Ocean (7%).

Deleted: , with median depths of 496 m and 505 m, with only 10% and 11% of measurements at depths below 3,000 m respectively.

Deleted: ; $MAE_{[Nd]}$ towards the Pacific Ocean and Atlantic Ocean, $MAE_{\epsilon_{Nd}}$ towards the Atlantic Ocean, and both towards shallower depths

Deleted: reduce

Deleted: evaluate

Deleted: For example, we compare $[Nd]_d$ and ϵ_{Nd} patterns across the global thermohaline circulation, through inter basin gradients, depth profiles and between distinct water masses, critically assessing the contributions of distinct Nd sources and cycling processes

Deleted: in order to better understand our model’s behaviour and performance.

Deleted: across different ocean basins and ocean depths

Deleted: ¶

Deleted: Firstly, $[Nd]_p/[Nd]_d$ is systematically varied in six sensitivity simulations ($[Nd]_p/[Nd]_d$ ranging 0.001-0.006), these values are based upon results from similar modelling schemes (Rempfer et al., 2011; Arsouze et al., 2009; Gu et al., 2019) and considers the few direct observations of $[Nd]_p/[Nd]_d$ (Jeandel et al., 1995; Stichel et al., 2020; Zhang et al., 2008; Paffrath et al., 2021; Bertram and Elderfield, 1993; Tachikawa et al., 1999). Here, and based upon simulations undertaken when validating the scheme, alongside estimates from previous optimised Nd isotope schemes (Arsouze et al., 2009; Rempfer et al., 2011; Gu et al., 2019; Pöppelmeier et al., 2020a), f_{sed} is fixed at 4.5×10^9 g yr⁻¹ throughout. Secondly, f_{sed} is varied in four sensitivity simulations (f_{sed} ranging 1.5×10^9 - 6.0×10^9 g yr⁻¹), using values based upon previous recent estimates of a global sediment flux to seawater (3.3×10^9 - 5.5×10^9 g yr⁻¹; Gu et al., 2019; Pöppelmeier et al., 2020b; Rempfer et al., 2011), and encompassing a larger parameter space in order to explore the sensitivity of Nd distributions. Notably, our f_{sed} sensitivity studies alter the percentage contribution of the sediment Nd flux to the total Nd flux to seawater from 66% (where $f_{sed} = 1.5 \times 10^9$ g yr⁻¹) to 89 % (where $f_{sed} = 6.0 \times 10^9$ g yr⁻¹). In all simulations $[Nd]_p/[Nd]_d$ is fixed at 0.003, which is in the middle of the range (0.001-0.006) discussed above based upon reasons outlined for the $[Nd]_p/[Nd]_d$ sensitivity simulations. Following experiment it is apparent that, ¶ Table 3: Suite of FAMOUS simulations designed to assess the sensitivity of simulated $[Nd]_d$ and ϵ_{Nd} distributions to two systematically varied parameters: reversible scavenging efficiency ($[Nd]_p/[Nd]_d$) and the global rate of direct Nd transfer from sediment to ocean water (f_{sed}). Simulation name refers to the title given to each sensitivity simulation in this manuscript, and the simulation identifier refers to the unique five-letter Met Office identifier (which, for example, can be used to call down full experiment details from the NERC PUMA facility: puma.nerc.ac.uk).

Formatted: Font: Times

Deleted: Simulation name

[... [21]

815

Table 3: Suite of FAMOUS simulations designed to assess the sensitivity of simulated $[Nd]_d$ and ϵ_{Nd} distributions to two systematically varied parameters: reversible scavenging efficiency ($[Nd]_p/[Nd]_d$) and the global rate of direct Nd transfer from sediment to ocean water (f_{sed}). Simulation name refers to the title given to each sensitivity simulation in this manuscript, and the simulation identifier refers to the unique five-letter Met Office identifier (which, for example, can be used to call down full experiment details from the NERC PUMA facility: puma.nerc.ac.uk). Global mean absolute error (MAE) and root mean square error (RMSE) for $[Nd]_d$ and ϵ_{Nd} are shown as an indicator of model skill.

Simulation name	Simulation identifier	$f_{sed} (Gg\ yr^{-1})$	$[Nd]_p/[Nd]_d$	Total Nd flux (Gg yr ⁻¹)	Nd inventory (Tg)	Residence time (yrs)	MAE _{Nd}	RMSE _{Nd}	% within 10 pmol kg ⁻¹	MAE _{εNd}	RMSE _{εNd}	% within 3 ‰
Varying $[Nd]_p/[Nd]_d$												
EXPT_RS1	XPDAI	4.5	0.001	5.27	16.00	3036	44.46	48.7	6	3.34	4.12	48
EXPT_RS2	XPDAD	4.5	0.002	5.27	8.91	1691	15.96	19.78	41	3.11	3.89	53
EXPT_RS3	XPDAH	4.5	0.003	5.27	6.03	1145	10.13	13.42	58	2.88	3.67	57
EXPT_RS4	XPDAE	4.5	0.004	5.27	4.51	856	9.66	13.01	62	1.67	3.45	60
EXPT_RS5	XPDAF	4.5	0.005	5.27	3.58	679	10.52	13.77	56	2.46	3.19	64
EXPT_RS6	XPDAG	4.5	0.006	5.27	2.95	559	11.7	14.63	50	2.27	3.00	70
Varying f_{sed}												
EXPT_SED1	XPDAL	1.5	0.003	2.27	2.5	1110	10.71	13.82	53	2.71	3.55	60
EXPT_SED2	XPDAM	3.0	0.003	3.77	3.89	1032	7.96	11.40	73	2.82	3.68	62
EXPT_SED3	XPDAH	4.5	0.003	5.27	6.03	1145	10.13	13.42	58	2.88	3.67	57
EXPT_SED4	XPDAN	6.0	0.003	6.77	7.79	1150	13.23	17.19	47	2.93	3.71	55

820

Due to time constraints, all sensitivity experiments presented here were run in parallel. Firstly, $[Nd]_p/[Nd]_d$ is systematically varied in six sensitivity simulations ($[Nd]_p/[Nd]_d$ ranging 0.001-0.006), these values are based upon results from similar modelling schemes (Rempfer et al., 2011; Arsouze et al., 2009; Gu et al., 2019) and considers the few direct observations of $[Nd]_p/[Nd]_d$ (Jeandel et al., 1995; Stichel et al., 2020; Zhang et al., 2008; Paffrath et al., 2021; Bertram and Elderfield, 1993; Tachikawa et al., 1999). Here, and based upon simulations undertaken when validating the scheme, alongside estimates from previous optimised Nd isotope schemes (Arsouze et al., 2009; Rempfer et al., 2011; Gu et al., 2019; Pöppelmeier et al., 2020a), f_{sed} is fixed at 4.5 Gg yr⁻¹ throughout.

825

Secondly, f_{sed} is varied in four sensitivity simulations (f_{sed} ranging 1.5 -6.0 Gg yr⁻¹), using values based upon previous recent estimates of a global sediment flux to seawater (3.3–5.5 Gg yr⁻¹; Gu et al., 2019; Pöppelmeier et al., 2020b; Rempfer et al., 2011), and encompassing a larger parameter space in order to explore the sensitivity of Nd distributions. Notably, our f_{sed} sensitivity studies alter the percentage contribution of the sediment Nd flux to the total Nd flux to seawater from 66% (where f_{sed} = 1.5 Gg yr⁻¹) to 89 % (where f_{sed} = 6.0 Gg yr⁻¹). In all simulations $[Nd]_p/[Nd]_d$ is fixed at 0.003, which is in the middle of the range (0.001-0.006) discussed above. Following the $[Nd]_p/[Nd]_d$ sensitivity experiment (Sect. 3.1), it is apparent that 0.004

830

would have been a better choice for $[Nd]_p/[Nd]_d$, and propose this as a starting point for further performance refinement in subsequent work.

835 Dissolved seawater Nd in all simulations are initialised from zero and integrated for at least 9,000 years under constant pre-industrial boundary conditions to allow the deep ocean circulation and marine Nd cycle to reach steady state, which we define as being when the Nd inventory becomes [near] constant with time ($< 0.02\%$ change per 100 years). All the presented results refer to or show the centennial mean from the end of the 9,000-year simulations.

3 Results and discussion

840 3.1 Model sensitivity to reversible scavenging efficiency ($[Nd]_p/[Nd]_d$)

The first set of sensitivity experiments test the response of simulated $[Nd]_d$ and ϵ_{Nd} to a systematic variation of the reversible scavenging tuning parameter ($[Nd]_p/[Nd]_d$). Despite total Nd flux to seawater being fixed, varying the scavenging efficiency ($[Nd]_p/[Nd]_d$) leads to different Nd inventories (Fig. 9 and Supplementary Information: Fig. S7 for the percentage rate of change per 100 years) and residence times (Table 3), consistent with previous studies (Siddall et al., 2008; Arsouze et al., 2009; Rempfer et al., 2011; Gu et al., 2019). A higher $[Nd]_p/[Nd]_d$ increases both the Nd scavenging efficiency and removal via sedimentation by enabling a larger fraction of seawater Nd to adsorb onto particles, in turn leading to a lower Nd inventory and a lower residence time (where; residence time = Nd inventory/total Nd flux).

Deleted: 0
Deleted: 5

Formatted: Normal
Deleted: while all other parameters are kept constant

Deleted: In the first 2,000 years, the Nd inventory increases exponentially in all simulations (Fig. 9). *EXPT_RS1* continues to increase rapidly until the end of the experiment, but after 2,000 years for simulations *EXPT_RS2* to *EXPT_RS6*, the rate of global Nd accumulation begins to slow as the inventories tend towards equilibrium. By year 6,000, all experiments with a global Nd inventory $< 8.0 \times 10^{12}$ g reach steady state and remain so until the end of the experiment (9,000 years). We therefore deem these simulations to have reached an acceptable equilibrium state. We explain why *EXPT_RS1* and *EXPT_RS2* do not reach equilibrium below, but otherwise, because of their unrealistic condition after 9,000 years (i.e. they reach an Nd inventory far past the target inventory of 4.2×10^{12} g; Tachikawa et al. (2003)), these two simulations are largely omitted from further discussion and analysis.

Figure 9: Global Nd inventory (g) simulated with different values for the reversible scavenging tuning parameter, $[Nd]_p/[Nd]_d$, as indicated. Dashed line represents the estimated global marine Nd inventory of 4.2×10^{12} g from Tachikawa et al. (2003) used as an approximate target for our simulations.

Deleted: kept constant

Deleted: 4

Deleted: Table 4: Overview of simulations exploring model sensitivity to the reversible scavenging tuning parameter $[Nd]_p/[Nd]_d$. Displaying global mean absolute error (MAE) for $[Nd]_d$ and ϵ_{Nd} .

Deleted: Simulation




Figure 9: Global Nd inventory (g) simulated with different values for the reversible scavenging tuning parameter, $[Nd]_p/[Nd]_d$, as indicated. Dashed line represents the estimated global marine Nd inventory of 4.2×10^{12} g by Tachikawa et al. (2003) used as a first-order target for our simulations.

By year 6,000, experiments *EXPT_RS3-EXPT_RS5* reach steady state. We therefore deem these simulations to have reached an acceptable equilibrium state. We explain why *EXPT_RS1*, *EXPT_RS2* and *EXPT_RS6* do not reach equilibrium below, but otherwise, because of their unrealistic condition after 9,000 years (i.e. *EXPT_RS1* and *EXPT_RS2* reach an Nd inventory far past the target inventory of 4.2 Tg; Tachikawa et al. (2003) and the high sink in *EXPT_RS6* causes imbalances in surface $[Nd]_d$ which tends towards zero), these simulations are largely omitted from further discussion and analysis.

Neodymium in *EXPT_RS1*, which has the lowest $[Nd]_p/[Nd]_d$, has a residence time of 3,036 years. This is much larger than the global ocean overturning time of 1,500 years, resulting in an Nd inventory of 16 Tg after 9,000 years. The $[Nd]_p/[Nd]_d$ and consequently the sink in *EXPT_RS1*, is too small to balance the input of Nd from the sources, causing such high Nd to accumulate in the simulation, which, as a result, does not reach steady state (rate of change 0.16% per 100 years at the end of the simulation; Fig. 9). Thus, *EXPT_RS1* returns the largest $MAE_{[Nd]}$ and MAE_{eNd} from these simulations, indicating the worst model-data fit, especially for representing $[Nd]_d$ measurements. *EXPT_RS6*, the simulation with the largest $[Nd]_p/[Nd]_d$, returns the lowest MAE_{eNd} . However, the sink term is too strong, resulting in a Nd inventory of 2.95 Tg, and the simulation fails to reach steady state (rate of change 0.07% per 100 years by the end of the run). On balance, we surmise that *EXPT_RS4* has the most reasonable combination of prognostic skill in terms of simulated Nd inventory (4.51 Tg) and residence time (856 years).

Deleted: Figure

Deleted: from

Deleted: n

Deleted: approximate

Deleted: and remain so until the end of the experiment (9,000 years)...

Deleted: 037

Deleted: $.06 \times 10^{13}$

Deleted:

Deleted: ,almost a factor of four larger than that of the estimated Nd budget in the oceans of 4.2×10^{12} g from Tachikawa et al. (2003).

Deleted: 2.11

Deleted: *EXPT_RS2*, although tending towards equilibrium at the end of the 9,000-year simulation (Fig. 9), just fails to reach steady state as defined above, still experiencing a rate of change of 0.003% per 100 years, highlighting that the rate of Nd removal from the ocean in this model configuration is still too low.

Deleted: experiment

Deleted: reaching only 2.95×10^{12} g, the final Nd inventory is too low to be considered as capturing Nd cycling well.

Deleted: $\times 10^{12}$

Deleted:

920

This simulation, with $[Nd]_p/[Nd]_d$ of 0.004, shows the best balance between Nd accumulation and removal, and hence returns the lowest $MAE_{[Nd]}$ alongside a moderately well performing $MAE_{\epsilon Nd}$ that falls in the middle of the range of results.

925

In contrast to our model, the schemes by Rempfer et al. (2011) and Gu et al. (2019) required a lower $[Nd]_p/[Nd]_d$ of 0.001 and 0.0009 respectively for their optimised experiment with Nd inventories of ≈ 4.2 Tg. Despite having similar scavenging schemes, a direct comparison of the parameter values used in the different Nd isotope modelling studies is difficult to make. This is because the divergence in sensitivity to reversible scavenging efficiency can be attributed to a combination of the differing magnitude and spatial distributions of model biogeochemical particle fields and Nd inputs, which are also partly controlled by the different architecture and horizontal resolution of the physical models. We thus propose that a future modelling protocol for intercomparing different global Nd isotope schemes would be well suited to exploring these differing sensitivities comprehensively.

930

The different values of $MAE_{[Nd]}$ and $MAE_{\epsilon Nd}$ across our sensitivity experiments (Table 3) illustrates the distinctive and uncoupled behaviour of $[Nd]_d$ and ϵ_{Nd} within the ocean, as broadly described by the Nd paradox, indicating that different processes govern the global distributions of each.

935

Generally, simulated $[Nd]_d$ distributions in *EXPT_RS4* match observational data well (Fig. 10). The lowest concentrations occur in the surface layers, and deep water $[Nd]_d$ increases along the global circulation pathway and with increasing age of water masses (Bertram and Elderfield, 1993; van de Flierdt et al., 2016; Tachikawa et al., 2017). Thus, we may infer that the model scheme in FAMOUS does have the broad capability of representing the physical and biogeochemical processes governing global marine $[Nd]_d$ distributions. However, the scheme does tend to overestimate the global vertical $[Nd]_d$ gradient with depth (Fig. 10 and Supplementary Information: Fig. S9 for major ocean basin averaged depth profiles), a feature reported in previous similar model schemes (e.g. Arsouze et al., 2009; Gu et al., 2019), indicating that the representation of processes governing vertical $[Nd]_d$ does not yet fully capture all processes occurring in the ocean. This overestimated vertical $[Nd]_d$ gradient with depth could be a result of the simplified uniform particle settling rate applied, which may be too high in the upper layers (due to lower settling velocities in the surface mixed layer resulting from enhanced turbulence and complexation with organic ligands; Chamecki et al., 2019; Noh et al., 2006) and too low at depth (due to higher settling velocities from particle aggregation).

945

Deleted: $\times 10^{12}$

Deleted:

Deleted: In other words, every study has run a different experiment, and so the results are not directly comparable.

Deleted: 4

Deleted: demonstrates

Deleted: In our results, increasing the efficiency of vertical cycling improves model-data fit for ϵ_{Nd} in a global sense, reducing $MAE_{\epsilon Nd}$ to a minimum of 2.27 where $[Nd]_p/[Nd]_d$ is 0.006. However, the story is more complicated for $[Nd]_d$ performance, where increasing $[Nd]_p/[Nd]_d$ from 0.001 up to 0.004 also reduces $MAE_{[Nd]}$ (to a minimum of 9.66 pmol kg⁻³), but then subsequent increases in $[Nd]_p/[Nd]_d$ (0.005 and 0.006) worsen model-data fit for $[Nd]_d$ because the sink term becomes too strong, removing too much Nd from the ocean, as reflected in the low accumulated Nd inventories (3.58×10^{12} g and 2.95×10^{12} g respectively).

Deleted: ¶

Deleted: , with lowest $[Nd]_d$ in the North Atlantic Ocean and the highest concentrations in the deep North Pacific Ocean

Deleted: These $[Nd]_d$ distributions are consistent with previous schemes that also take into account reversible scavenging (Arsouze et al., 2009; Gu et al., 2019; Oka et al., 2021; Rempfer et al., 2011; Siddall et al., 2008).

Deleted: simulate a too pronounced

Deleted:

Deleted:

Deleted: 7

Deleted: , leading to an underestimation of $[Nd]_d$ in the surface ocean and overestimation at abyssal depths.

Deleted: may

Deleted: i.e., there may be

Deleted: (

Deleted: i.e.,

Deleted: due to

Deleted: at depth




Figure 10: Maps of $[Nd]_d$ (left) and ϵ_{Nd} (right) in simulation *EXPT_RS4* split into four different depth bins, (a-b) shallow (0-200 m), (c-d) intermediate (200-1,000 m), (e-f) deep (1,000-3,000 m), and (g-h) deep abyssal ocean (>3,000 m). Water column measurements from within each depth bin (Osborne et al., 2017, 2015; GEOTRACES Intermediate Data Product Group, 2021) are superimposed as filled circles using the same colour scale.

Too high simulated $[Nd]_d$ at depth may also be caused by the benthic sediment source being too strong, coupled with the simplified assumption of a globally uniform sediment flux acting across the entire seafloor. This suggests that perhaps too much emphasis has been put on a widespread abyssal benthic flux to resolve the Nd paradox, and it may be that shallower and marginal regions pose a larger in magnitude and more distinct Nd source to seawater as opposed to an open-ocean benthic source. Recent pore water measurements by Deng et al. (2022) on the East China continental shelf revealed larger benthic

Deleted: 1

Deleted: Global volume-weighted distributions

Deleted: Overestimation of the

Deleted: to seawater

Deleted: points to the suggestion

Deleted: P

Deleted: (

Deleted: ,

000 fluxes across the dynamic hydraulic environment of the continental shelf compared to deep pore water, which acted as a sink,
likely due to authigenic precipitation. As such, spatial variations of the benthic flux are likely affected by multiple complex
depth-related processes, although deep ocean regions may still dominate the total area-integrated benthic flux. A future
evolution of the scheme could explore depth dependent benthic flux rates.

Deleted: in

Deleted: like those in the recent scheme by

Moreover, the overestimation of $[Nd]_d$ at depth could be a result of biases within the simulated biogenic particle fields.
Comprehensive global observations of sinking particles fluxes – the central driver of ocean biogeochemical cycling – remain
fundamentally difficult to obtain (Dunne et al., 2007), and our particle fluxes may be inaccurate as a consequence (see Sect.
2.3.4 for assumptions and respective limitations). It also seems likely that the reversible scavenging parameterisations, which
are simple by design due to incomplete understanding (also Sect. 2.3.4), restrict the model's ability to precisely capture all
aspects of the measured Nd distributions. For example, we know that the assumption of a globally uniform $[Nd]_p/[Nd]_d$ is
oversimplified, as particle measurements suggest $[Nd]_p/[Nd]_d$ is not uniform in seawater (Stichel et al., 2020). High
remineralsation rates, inducing increased exchange, have been observed in the Labrador Sea (Lagarde et al., 2020), whilst
vast expanses of low biological productivity in the Pacific gyres mean $[Nd]_p$ are generally much lower here compared to the
North Atlantic. Thus, the global $[Nd]_p/[Nd]_d$ in our scheme may be skewed towards lower values. Nonetheless, it is necessary
to adopt a pragmatic approach for the model development, and hence, similar to other model schemes, the globally uniform
 $[Nd]_p/[Nd]_d$ remains in our implementation. Future work could explore regionally varied $[Nd]_p/[Nd]_d$, which would benefit
from additional measurement constraints to sufficiently justify a more complex distribution. Additionally other particles such
as Fe and Mn oxides and hydroxides, not considered here, may play an important role for scavenging of Nd (e.g., Bayon et al.,
2004; Lagarde et al., 2020; Du et al., 2022). Further observational evidence of the processes involved and their importance for
Nd cycling by combined particulate and dissolved measurements and laboratory experiments (e.g., Stichel et al., 2020; Pearce
et al., 2013; Rousseau et al., 2015; Wang et al., 2021; Paffrath et al., 2021; Lagarde et al., 2020), plus experimentation within
a modelling framework, may help to improve this limitation.

Deleted: Direct and

Deleted: c

Deleted: which is a pragmatic choice in the model development,

Deleted: As such

Deleted: also

Deleted: b

However, the largest model-data disparities for $[Nd]_d$ occur in the shallow ocean (above 200 m) at specific locations close to
continental margins where Nd is input to the ocean through major point sources that are not well resolved by the model. This
includes continental margins in the Labrador Sea (where simulated $[Nd]_d$ is 3 pmol kg⁻¹ compared to measured $[Nd]_d$ of
70 pmol kg⁻¹) and the Sea of Japan (where simulated $[Nd]_d$ is 2 pmol kg⁻¹ compared to measured $[Nd]_d$ of 50 pmol kg⁻¹), for
example. Such low $[Nd]_d$ in the surface layers may be exacerbated by operational constraints in the scheme, such as the
extensive and immediate dilution of point sourced Nd across the whole of its containing grid cell combined with the
instantaneous nature of simulated reversible scavenging, which may be much faster in the model than would occur normally.
Furthermore, the scheme may be underestimating the magnitude of surface sources of Nd, for example the Nd sourced from
dust may be too low in certain regions, possibly resulting from the globally fixed (2 %) solubility of Nd from dust sources
applied. In the preliminary optimisation of the Nd isotope scheme in the GNOM v1.0 model, Pasquier et al. (2022) found that

Deleted: which may be a

Deleted: of

Deleted: (

Deleted: ,

Deleted: Pas(2020)

045 dust solubility needed to be increased to unrealistically high levels to best fit measurements, for example optimised parameters
for North American dust solubility was 83%. A future exploration of the regional solubility and amount of Nd coming from
dust sources would be needed to explore this in detail. On the other hand, recent evidence suggests that there may be a major
particulate riverine flux of Nd to the ocean (Rousseau et al., 2015; Ralhf et al., 2021), which implies that the magnitude of the
sediment Nd sourced to seawater near river mouths could be too weak. Specific to our scheme, the vertical convection of water
from the physical ocean model in FAMOUS may also be too intense in the North Atlantic, leading to surface underestimations
050 of $[Nd]_d$ and overestimations at depth; a re-tuning of the physical model would help to ascertain the extent to which this
limitation contributes towards the simulated Nd bias.

Nonetheless, consistent with compilations of water column measurements (Tachikawa et al., 2017; van de Flierdt et al., 2016),
overall global distributions of simulated ϵ_{Nd} are broadly most unradiogenic in the North Atlantic and more radiogenic in the
North Pacific (Fig. 10), with intermediate values in the Southern and Indian Oceans. The most unradiogenic ϵ_{Nd} occurs in the
1055 surface layers of the Hudson Bay and Labrador Sea regions, and they closely match measured data ($\epsilon_{Nd} = -18$). However, the
most radiogenic ϵ_{Nd} , simulated in the surface layers of marginal regions in the North and equatorial western Pacific ($\epsilon_{Nd} = -3$),
is significantly lower than measured ($\epsilon_{Nd} = +3$), and in the central and North Pacific; particularly above 1,000 m, simulated ϵ_{Nd}
is -7, but measurements are closer to -1. At the basin scale, the magnitude of the ϵ_{Nd} gradient from Pacific to Atlantic is
underestimated by the model, and presents a familiar bias as seen in previous Nd isotope schemes (Arsouze et al., 2009;
1060 Rempfer et al., 2011; Pöppelmeier et al., 2020a; Jones et al., 2008; Gu et al., 2019). This is mainly due to the simulated Pacific
being too unradiogenic (basinal mean ϵ_{Nd} of -7.5) compared to measured water samples ($\epsilon_{Nd} = -4$), while simulated and
measured basinal mean Atlantic ϵ_{Nd} values are in much better agreement ($\epsilon_{Nd} = -11$ and -12.5 respectively; Supplementary
Information Fig. S9). Possible explanations for why our simulated Pacific ϵ_{Nd} is so much lower than what has been recorded
in modern seawater include that the model boundary conditions, specifically the marine sediment source of Nd (taken directly
1065 from Robinson et al., 2021), may not be sufficiently radiogenic, or the sediment flux from the seafloor is not uniform; points
we will return to later (Sect. 3.2).

Simulated $[Nd]_d$ with depth at distinct locations in all the reversible scavenging sensitivity experiments (Fig. 11 a-f and h-m)
generally (though not always) exhibit similar depth profiles to the observational data. The best model-data fit is seen within
depth profiles in the Pacific and Southern Ocean, and notably more so under higher $[Nd]_p/[Nd]_d$. The largest model-data offsets
1070 across all sensitivity experiments in terms of both the magnitude and the depth gradients in $[Nd]_d$ occur in the North Atlantic
Ocean. In particular, the large near-surface concentrations are not resolved, with the largest disparities occurring under higher
scavenging parameterisations.

Deleted: the
Deleted: the
Deleted:
Deleted: and the magnitude of the Nd sourced from dust
Deleted: would
Deleted: In addition
Deleted: as a major marine
Deleted: source
Deleted: (
Deleted: REF0 Rouss 2015 and Ralf 2021),
Deleted: as such
Deleted: may
Deleted: T
Deleted:

Deleted: ϵ_{Nd}
Deleted: In fact, these specific comparisons, which demonstrate a
good match in the North Atlantic and weaker performance in the
North Pacific, are congruent with a more general trend in the
simulations. That is,
Deleted: 7
Deleted: ϵ_{Nd}
The discrepancy in simulated and measured Pacific ϵ_{Nd} values may be
amplified by sparse sampling that is biased towards shallow
radiogenic continental regions and volcanic island areas, in which
case, model performance is better than we are able to assess. In
addition, and as highlighted in Sect. 2.4, $MAE_{\epsilon_{Nd}}$ is biased towards
the Atlantic (45% of all observational data), compared to the Pacific
(29%). Consequently, due to biases in measurement density, a
simulation better representing the Atlantic is more strongly favoured
in the adopted cost function than one better representing the Pacific at
the expense of the Atlantic. One further explanation for why
Deleted: of
Deleted: o
Deleted:
Deleted: measurements is
Deleted: a
Deleted: profiles
Deleted: s
Deleted: depth
Deleted: especially
Deleted: , where simulated $[Nd]_d < 10 \text{ pmol kg}^{-1}$ in the upper 1,000
m are too low compared to observed concentrations around 18-
21 pmol kg^{-1}




Figure 11: Central panel (g) displays $[Nd]_d$ at the seafloor (i.e., lowermost box of the ocean model), in simulation *EXPT_RS4* (100-year mean from the end of the run), with superimposed water column measurements (Osborne et al., 2017, 2015; GEOTRACES Intermediate Data Product Group, 2021) from $\geq 3,000$ m shown by filled coloured circles on the same colour scale. Surrounding panels (a-f) and (h-m) display depth profiles of simulated (coloured lines, one per sensitivity simulation with varied $[Nd]_p/[Nd]_d$) and measured (filled circles) $[Nd]_d$. Larger shifts in the $[Nd]_d$ between simulations highlight regions most sensitive to the efficiency of reversible scavenging.

For ϵ_{Nd} , the response to varying scavenging efficiency has varied effects across depths and between ocean regions, indicating a more complex relationship between how reversible scavenging can delineate global ϵ_{Nd} distributions in comparison to $[Nd]_d$. Increased scavenging efficiency traps regional ϵ_{Nd} provenance signals, resulting in larger inter-basin ϵ_{Nd} gradients, the strength of which are favoured by the combination of strong scavenging efficiency and a short seawater residence time. Although the simulated ϵ_{Nd} profiles with depth do not closely follow measurements, it is valuable to discuss the sensitivity of simulated ϵ_{Nd} to variations in scavenging efficiency. The North Atlantic is the most sensitive basin to changes in reversible scavenging (registered by the greater ϵ_{Nd} profile shifts between different experiments, Fig. 12a-d), particularly at depths below 2,000 m, indicating that reversible scavenging, alongside convection at sites of deep-water formation, are important processes for

Deleted:

Deleted: Interestingly, we see a different sensitivity to varying $[Nd]_p/[Nd]_d$ in $[Nd]_d$ (Fig. 11) compared with ϵ_{Nd} (Fig. 12) within different ocean basins and depths. For $[Nd]_d$, we find that all ocean basins are sensitive to the parameterisation of reversible scavenging efficiency (i.e., wider divergence between the sensitivity for each depth profile)

governing the simulated ϵ_{Nd} signal of deep-water masses. Thus, a stronger reversible scavenging efficiency is needed to set the unradiogenic deep ocean signature in the North Atlantic, and to trap these regional ϵ_{Nd} provenance signals locally.




Figure 12: Central panel (g) displays ϵ_{Nd} at the seafloor (i.e., lowermost box of the ocean model), in simulation EXPT_RS4 (100-year mean from the end of the run), with superimposed water column measurements (Osborne et al., 2017, 2015; GEOTRACES Intermediate Data Product Group, 2021) from $\geq 3,000$ m shown by filled coloured circles on the same colour scale. Surrounding panels (a-f) and (h-m) display depth profiles of simulated (coloured lines, one per sensitivity simulation with varied $[Nd]_p/[Nd]_d$) and measured (filled circles) ϵ_{Nd} . Larger shifts in the ϵ_{Nd} between simulations highlight regions most sensitive to the efficiency of reversible scavenging.

Typical depth profiles from measurements in the sub-tropical North Atlantic show contrasts in ϵ_{Nd} that co-vary with the presence of major water masses (coloured circles in Fig. 12a-b). Across all sensitivity experiments, there is relative consistency with depth for the profiles in the sub-tropical North Atlantic (e.g., Fig. 12b), ranging from -10 to -12 under increasing scavenging efficiency, which acts to drive more localised unradiogenic signals. This simulated uniformity may be due to the indiscriminate seafloor sedimentary flux here being too high, and so dominating the deep seawater ϵ_{Nd} signal. An additional

Deleted: , particularly at depths below 1,000 m, which is broadly consistent with the findings reported in previous work by Rempfer et al. (2011). However, Siddall et al. (2008) showed a strong sensitivity of Pacific $[Nd]_p$ alongside a weak sensitivity of Atlantic $[Nd]_p$ to the reversible scavenging efficiency, the differences were attributed by the authors to the dominance of vigorous Atlantic advective lateral transport, whereas we demonstrate similar sensitivities in the Pacific, Atlantic, Indian and Southern Oceans (Fig. 11). The more simplified fixed surface boundary conditions applied in Siddall et al. (2008) are not influenced by changing $[Nd]_p/[Nd]_d$, unlike the fluxes in the scheme presented here and by Rempfer et al. (2011), which may explain the contradictory response of the Atlantic basin across these studies.

Deleted:

Deleted: For ϵ_{Nd} , the response to varying scavenging efficiency has varied effects across depths and between ocean regions, indicating a more complex relationship between how reversible scavenging can delineate global ϵ_{Nd} distributions in comparison to $[Nd]_d$. Increased scavenging efficiency traps regional ϵ_{Nd} provenance signals, and this results in larger inter-basin ϵ_{Nd} gradients, the strength of which are favoured by the combination of strong scavenging efficiency and a short seawater residence time; as shown by the relationship between $MAE_{\epsilon_{Nd}}$ and $[Nd]_p/[Nd]_d$ (Table 4). The North Atlantic is the most sensitive basin to changes in reversible scavenging (registered by the greater ϵ_{Nd} profile shifts between different experiments, Fig. 12a-d), particularly at depths below 2,000 m, indicating that reversible scavenging is important, there, for governing the simulated ϵ_{Nd} signal of deep-water masses. We attribute this sensitivity to the fact that the Atlantic Ocean experiences strong convection and is surrounded by continental margins. As such, the basin is subject to substantial continental Nd inputs to the upper layers of the ocean, and the shallow ϵ_{Nd} signal is transferred to the ocean interior, setting the unradiogenic deep ocean signature. Thus, a stronger reversible scavenging efficiency is needed to trap regional ϵ_{Nd} provenance signals locally.

Deleted: , here

explanation may be due to the lack of abyssal (i.e., below 3 km) AABW, which does not extend past $\sim 20^\circ$ N (Fig. 3). This insufficient AABW production and penetration into the Atlantic are known limitations of FAMOUS (Dentith et al., 2019; Smith, 2012), although these biases are reduced in our *control* compared to the previous studies (Sect. 2.2). Thus, the seawater below the surface mixed layer comprises only North Atlantic water, and in so doing, the model will not resolve the AABW signal inferred from measurements at depth in higher latitudes.

Higher up the water column, the more unradiogenic NADW seen in the seawater measurements is conspicuously absent (note the ϵ_{Nd} minima in subtropical North Atlantic measurements $\sim 1,000$ - $2,000$ m deep, Fig. 12a-b), even though we know NADW does reside there in the model (see Fig. 3 for verification). We may relate this to the high latitude North Atlantic ϵ_{Nd} being too radiogenic to tag NADW with its characteristically unradiogenic ϵ_{Nd} signal, particularly around the mouth of the Labrador Sea (Fig. 10a-b). This could be exacerbated by a dampening of the unradiogenic NADW ϵ_{Nd} from a relatively radiogenic seafloor benthic flux along the water flow path (Fig. 6). Consequently, despite a close correlation to seawater ϵ_{Nd} in the subtropics and lower-NADW (where measurements of ϵ_{Nd} are -12.4 and simulated is -12), upper-NADW end member ϵ_{Nd} is not sufficiently unradiogenic, even under the highest reversible scavenging efficiency ($[Nd]_p/[Nd]_d = 0.006$) where simulated ϵ_{Nd} is -12 in comparison with seawater records -13.2 ϵ_{Nd} (Lambelet et al., 2016).

In contrast, ϵ_{Nd} in the Pacific Ocean is least sensitive to changes in $[Nd]_p/[Nd]_d$, particularly below 500 m (Fig. 12e-f, h-i). This is expected, due to the absence of major ocean convection and ventilation, meaning the Pacific contains an older, more homogenised pool of water in comparison to the Atlantic, and thus, water masses are far less distinct, and the localised ϵ_{Nd} signal does not get dispersed via convection as rapidly. The deep seafloor wide sediment source governs simulated ϵ_{Nd} distributions in the intermediate-deep Pacific, due to its larger flux relative to that transported vertically via particle scavenging and dissolution, or horizontally by the sluggish convection of the Pacific. One further aspect to note, is that under a high sink ($[Nd]_p/[Nd]_d = 0.006$), in some Pacific regions $[Nd]_d$ in the surface layers tends towards zero. This causes numerical instabilities in modelled Nd ratios and thus, where $[Nd]_d < 0.2$ pmol kg^{-1} , ϵ_{Nd} is meaningless and so is masked-out from the results shown by Fig. 12, to avoid misinterpretation.

The simulated depth profile in all experiments in the Indian Ocean matches the observed intermediate ϵ_{Nd} signal of -8 between 500 - $2,000$ m (Fig. 12m). However, the more radiogenic ϵ_{Nd} signal of -6 in the surface layers is not captured, nor the shift in ϵ_{Nd} below $2,000$ m to more unradiogenic values reaching a minimum of -10.5 at $3,000$ m, with all experiments simulating a relatively uniform ϵ_{Nd} with depth. Under higher reversible scavenging parameters for this depth profile (*EXPT_RS4-EXPT_RS6*), the surface concentrations are too low (simulated $[Nd]_d = 3$ pmol kg^{-1} and observed $[Nd]_d = 8$ pmol kg^{-1}), indicating that the model is not fully resolving either the concentration or the ϵ_{Nd} from a surface flux here, and scavenging may be too intense at the surface. In the deeper ocean ($\approx 3,000$ m), these simulations show the model represents the $[Nd]_d$ profiles better but is missing an unradiogenic exchange of ϵ_{Nd} at depth. This could point to an unradiogenic sediment source or exchange

Deleted: can be partly explained by

Deleted: basin

Deleted: Shifts in ϵ_{Nd} between the sensitivity studies in this region therefore relate to reversible scavenging efficiency changing the ϵ_{Nd} of NADW.

Deleted: Depth variability is greatest in the equatorial Pacific, where ϵ_{Nd} decreases from -2.5 at the surface to -5 at depth, but, overall, simulated vertical ϵ_{Nd} gradients are small in the Pacific, consistent with seawater measurements.

Deleted: This does make it possible for reversible scavenging to convey a more localised surface signal into the interior of the Pacific under a lower $[Nd]_p/[Nd]_d$ than can be achieved in the Atlantic, because

Deleted: Our less sensitive response of Pacific ϵ_{Nd} to reversible scavenging efficiency contrasts with results from Rempfer et al. (2011), who found a greater response of Pacific ϵ_{Nd} (compared to the Atlantic). We attribute this difference primarily to the spatial variation in the sediment Nd flux, which, in the study from 2011, is constrained to shallower radiogenic continental sources with no external marine Nd source below $3,000$ m. In comparison, our

Deleted: On the whole, the relatively greater sensitivity of ϵ_{Nd} in the surface Pacific, compared with the deep Pacific, to the reversible scavenging efficiency produces a more radiogenic signal closer to that of the radiogenic measurements. This is likely caused by the downward transportation of radiogenic surface inputs to subsurface layers is consistent with previous modelling (e.g. Gu et al., 2019), and occurs due to the inherent presence of larger particle fluxes and therefore greater influence of reversible scavenging here compared to the deeper ocean, where particle dissolution acts to reduce particle concentrations and their associated scavenging mechanisms.

Formatted: Strikethrough

Deleted: better, but

Deleted: , where there is an insensitivity to reversible scavenging

that is misrepresented by the bulk sediment boundary conditions and sediment source assumptions (e.g., our application of a global seafloor sediment source of Nd irrespective of sedimentary characteristics).

In the Southern Ocean, simulations with $[Nd]_s/[Nd]_d \geq 0.003$ broadly match the general measured ϵ_{Nd} at depths above 1,000 m (decreasing with depth from -7.8 at the surface to -8.2 at 1,000 m). Below this, and down to 3,000 m, observed diversions in ϵ_{Nd} are not captured in any of the sensitivity experiments. This discrepancy can be attributed to the simulation of quite homogenous AABW throughout the water column in the region, which represents a physical bias of FAMOUS (Dentith et al., 2019; Smith, 2012). Specifically, in the Pacific sector of the Southern Ocean (Fig. 12j), the model cannot resolve the measured unradiogenic spike at 1,500 m, which captures the distinct presence of lower Circumpolar Deep Water (CDW, $\epsilon_{Nd} = -8.4 \pm 1.6$: Lambelet et al., 2018) formed from mixing of Atlantic, Pacific and Indian sourced waters.

Thus, we conclude that scavenging and removal via sedimentation is necessary to balance the simulated input sources and enables the scheme to reach equilibrium around reasonable Nd inventories. We find that reversible scavenging is an important process that enhances the ϵ_{Nd} gradient between oceans by maintaining localised basinal ϵ_{Nd} signals throughout the water column. The strength of this process is particularly important for maintaining the simulated unradiogenic ϵ_{Nd} in the well ventilated North Atlantic Ocean (alongside deep-water formation), but less important for the more stagnant modern Pacific Ocean.

Nonetheless, parameterising reversible scavenging efficiency alone cannot account for the correct trends and magnitude within ϵ_{Nd} gradients observed between basins and in depth profiles. Currently, the scheme assumes that all particles reaching the seafloor via reversible scavenging are buried in the sediment, and as such are decoupled from the seafloor sediment source; there is no Nd release and the particles are removed from the model. A future evolution of the scheme could explore the dissolution of authigenic sedimentary phases on the seafloor during diagenesis, to investigate how this may influence the ϵ_{Nd} distributions of the benthic flux. Moreover, our results demonstrate the importance of further constraining other aspects of marine Nd cycling under a holistic framework. Furthermore, we note that the scheme described here may not be fully resolving the end member ϵ_{Nd} of different water masses due to an imperfect representation of the sources of Nd to seawater (i.e., the model boundary conditions and strength of the source fluxes). Thus, in some instances, the scheme carries inappropriate/dampened ϵ_{Nd} signals, coupled alongside particular structural model biases in the physical circulation (e.g., limited AABW intrusion in the North Atlantic), although in the case of the latter point, we note that this is not a limitation of the presented scheme, and that the implementation can be useful for identifying such physical biases.

3.2 Model sensitivity to Nd flux from the sediment (f_{sed})

The second set of sensitivity simulations tests the response of $[Nd]_d$ and ϵ_{Nd} to systematically varying the total Nd flux from the sediment (f_{sed}). In this experiment, Nd accumulates rapidly from the start of the simulations (Fig. 13 and Supplementary

Deleted: Interestingly, apart from the Indian Ocean, a model configuration with more efficient scavenging generally tends to produce results closest to the observational data (Table 4). In the Indian Ocean, seawater ϵ_{Nd} distributions are subject to monsoon systems, which facilitate the [seasonal] delivery of large riverine fluxes of Nd to seawater, for example from the Ganges and Brahmaputra Delta (Gupta and Naqvi, 1984). Large freshwater fluxes also deliver large amounts of freshly eroded and labile sediment to the continental margins, which likely contribute significantly to governing marine ϵ_{Nd} distributions through boundary particle-seawater exchange processes. ¶

Deleted: The main conclusion to be made regarding model sensitivity to varying the reversible scavenging tuning parameter $[Nd]_s/[Nd]_d$ is

Deleted: from dust, rivers and seafloor sediment, and

Deleted: physical

Deleted:

Deleted: -

Deleted: , including surface inputs from dust and river sources, sediment-seawater exchanges and the redistribution and mixing via physical ocean circulation which also governs global $[Nd]_d$ and ϵ_{Nd} distributions

Deleted:

Deleted: tranche

Deleted: , while all other parameters were kept constant

Information: Fig. S8 for the percentage rate of change per 100 years) and tapers off thereafter to varying degrees depending on the rate of accumulation. By year 6,000 all f_{sed} sensitivity experiments have reached steady state ($< 0.02\%$ change per 100 years).




Figure 13: Global Nd inventory (Tg) simulated with different values for the total sediment flux tuning parameter (f_{sed}) as indicated. Dashed line represents the estimated global marine Nd inventory of 4.2 Tg from Tachikawa et al. (2003) used as an approximate target for the simulations. Note the logarithmic y-axis scale.

Modifying the sediment flux while keeping the sink term $[Nd]_p/[Nd]_d$ constant returns a varied equilibrium Nd inventory across the suite of simulations (Fig. 13). However, without changing the scavenging efficiency (i.e. Nd sink), there is a more moderate range of 40 years difference in the f_{sed} simulations' Nd residence time (Table 3). In general, the relatively low scavenging efficiency of all of the f_{sed} sensitivity simulations ($[Nd]_p/[Nd]_d = 0.003$, see Sect. 3.1 for context) yields long residence times greater than 1,000 years.

EXPT_SED1, which has the lowest f_{sed} , corresponding to a sediment flux (per area) of $1.17 \text{ pmol cm}^{-2} \text{ yr}^{-1}$, and hence the lowest total Nd flux to the ocean, returns the smallest total Nd inventory of 2.5 Tg (Table 3). Conversely, *EXPT_SED4*, which has the largest f_{sed} , corresponding to a sediment flux of $4.69 \text{ pmol cm}^{-2} \text{ yr}^{-1}$, results in the greatest Nd inventory (7.8 Tg), producing both the worst $MAE_{[Nd]}$ and MAE_{cNd} .

We note that, compared to varying the reversible scavenging efficiency, varying f_{sed} drives relatively small changes in Nd distributions, as demonstrated by the minor differences in MAE_{cNd} between sensitivity experiments. This makes sense, since varying f_{sed} only acts to change the fractional contribution from each specific Nd source, e.g., an enhanced f_{sed} reduces the

Deleted: . For *EXPT_SED1* and *EXPT_SED2*, the rate of increase in the Nd inventory begins to reduce by 1,000 years, as these simulations, which have the smallest sediment source (f_{sed}), approach steady state. By year 2,000 the rate of Nd accumulation in *EXPT_SED3* and *EXPT_SED4* reduces, and these simulations also reach steady state

Deleted: , albeit above that of the target global inventory reference ($4.2 \times 10^{12} \text{ g}$).

Deleted: B

Deleted: 0

Deleted: 5

Deleted: $\times 10^{12}$

Deleted:

Deleted: Table 5 summarises the final global mean marine Nd inventory and residence time of the simulations, and reports their skill according to our global cost functions, $MAE_{[Nd]}$ and MAE_{cNd} .

Deleted: scavenging efficiency

Deleted: , consistent with the findings of Rempfer et al. (2011).

Deleted: Table 5: Overview of simulations exploring model sensitivity to the total sediment Nd flux tuning parameter (f_{sed}). Displaying global mean absolute error (MAE) for $[Nd]_d$ and c_{Nd} . Simulation ... [4]

Deleted:

Formatted: Superscript

Formatted: Superscript

Deleted: $\times 10^{12}$

Deleted:

Deleted: (only half the total estimated by Tachikawa et al., 2003), but did well with c_{Nd} distributions, returning the lowest global MAE_{cNd} .

Deleted: (per area)

Deleted: $\times 10^{12}$

Deleted:

Deleted: here

Deleted: C

Deleted: discrete

Deleted: has no direct impact on the relative distribution of ^{143}Nd and ^{144}Nd once it is in the water, it

1360 fraction of total Nd flux coming from dust and rivers (which are inputs constrained to the surface and point sources close to the continents), concentrating the flux across the global seafloor, and vice versa. However, in all simulations, and consistent with previous studies, the sediment Nd source to seawater remains the major source, and f_{sed} would need to be reduced much more to greatly influence $MAE_{\epsilon Nd}$.

365 Overall, from the simulations in this study, *EXPT_SED2* demonstrates the best skill for $[Nd]_d$ compared to measurements, returning the lowest $MAE_{[Nd]}$ of 7.96 pmol kg⁻¹ (Table 3) and achieving the most balanced Nd source and sink terms. Although the simulation does not represent the lowest $MAE_{\epsilon Nd}$, the range across the f_{sed} experiment is small (2.71 to 2.93; Table 3) and it does simulate the highest percentage of simulated ϵNd within 3 ϵNd units of measurements (62%). The sediment flux in *EXPT_SED2* of 2.35 pmol cm² yr⁻¹ is considered similar to the few existing measurements of a benthic flux across different sedimentary environments, which are mostly on the order of 10 pmol cm² yr⁻¹ (Abbott, 2019; Abbott et al., 2015a; Haley et al., 2004; Du et al., 2018). In comparison with model estimates of the benthic flux, the sediment flux in *EXPT_SED2* is most comparable to the global benthic flux estimated by Pöppelmeier et al. (2020a) of 5.4 pmol cm² yr⁻¹, but an order of magnitude lower than the value of 20-30 pmol cm² yr⁻¹ estimated by Du et al. (2020). Alternatively, the depth scaling of the benthic flux in the modelling study by Pasquier et al. (2022) resulted in surface fluxes of 83.7 pmol cm² yr⁻¹ and deep ocean fluxes of 1.11 pmol cm² yr⁻¹.

375 The target global Nd inventory of 4.2 Tg, derived from the canonical work by Tachikawa et al. (2003), is based on a sparser array of seawater measurements, particularly in the Southern Ocean, compared to what is available today. Using our newer compilation of $[Nd]_d$ to identify the best performing simulation for $[Nd]_d$ (*EXPT_SED2*), we can report an updated estimate of the global Nd inventory from the model of 3.89 Tg (Table 4), noting that this is likely an underestimate of the total, since the model does not represent marginal seas. Furthermore, we can examine the basinal distribution of $[Nd]_d$ (also Table 4), whereby 380 the largest amount is found in the stagnant North Pacific, where the greatest area integrated sediment flux occurs. Smaller Nd inventories are found in the more convective North Atlantic and Southern Ocean regions, which have less expansive seafloors and larger biogenic particle fluxes resulting in more Nd removal via reversible scavenging and sedimentation. We propose future work should take advantage of the greater abundance of seawater Nd measurements (e.g. as compiled in this study) to produce a detailed update to the estimate of oceanic inventories of Nd. These more accurate Nd budgets across diverse ocean 385 regions and depths would help in improving constraints on the marine Nd cycle and would provide excellent model evaluation tools.

Table 4: Nd inventory (Tg) for *EXPT_SED2* within different ocean regions

Ocean Region	Nd inventory (Tg)
Global	3.89

Deleted: at reaching the target Nd inventory (3.89×10^{12} g compared to the 4.2×10^{12} g target). It returns the lowest $MAE_{[Nd]}$ because it achiev

Deleted: es

Deleted:

Deleted: used as a target for analysis from

Deleted: (

Deleted: ,

Deleted: is now nearly 20 years old, and is limited by a lack of observations at the time

Deleted: notably

Deleted: Here we take advantage of using the most up-to-date

Deleted: used in this study to validate the scheme to propose a new global marine Nd inventory estimate of 3.89 Tg (*EXPT_SED2*: Table 4). Of the total Nd simulated in *EXPT_SED2*,

Deleted: largest

Deleted: Alternatively, r

Deleted: elatively

Deleted: s

Deleted: with

Deleted: smaller

Deleted: areas

Deleted: Although out of the scope of this study, w

Deleted: d

Deleted: inventories within the oceans

Deleted: ,

Deleted: t

Deleted: future

Deleted: , and

Deleted: Nd inventory

Arctic Ocean	0.05
North Atlantic	0.33
South Atlantic	0.45
North Pacific	0.96
South Pacific	0.76
Indian Ocean	0.63
Southern Ocean	0.28

1420

Altogether, simulated $[Nd]_d$ in *EXPT_SED2* matches the general observational data trend of Nd concentration. However, in the deep layers of the North Pacific (below 3,000 m) simulated $[Nd]_d$ is underestimated (36 pmol kg⁻¹ compared to 50 pmol kg⁻¹ from seawater measurements). This underestimation of $[Nd]_d$ at depth can be explained by a combination of having a seafloor sediment source at the lower end of our range (f_{sed} is 3.0 Tg yr⁻¹) and slow release from reversible scavenging ($[Nd]_d/[Nd]_d$ of 0.003; our experiments suggest 0.004 may be a more suitable efficiency to use, Sect 3.1), highlighting the importance of both non-conservative processes in governing deep $[Nd]_d$ distributions.

Deleted: ¶

Deleted: increasing with depth (Fig. 14), with good model-data fit especially in the upper 1,000 m across all ocean basins (except the North Atlantic, discussed below).

Deleted:

Deleted: × 10⁹

Deleted:

Deleted: , especially in the deep North Pacific




Figure 14: Maps of $[Nd]_d$ (left) and ϵ_{Nd} (right) in simulation *EXPT_SED2* split into four different depth bins, (a-b) shallow (0-200 m), (c-d) intermediate (200-1,000 m), (e-f) deep (1,000-3,000 m), and (g-h) deep abyssal ocean (>3,000 m). Water column measurements from within each depth bin (Osborne et al., 2017, 2015; GEOTRACES Intermediate Data Product Group, 2021) are superimposed as filled circles using the same colour scale.

Moreover, the longer lifetime of simulated Nd in the *EXPT_SED2* ocean (1,032 years) compared to that estimated by previous work (360-800 years; Gu et al., 2019; Rempfer et al., 2011; Siddall et al., 2008; Tachikawa et al., 2003) means that Nd becomes well mixed in the deep ocean (below 1,000 m), homogenising the ϵ_{Nd} signal. This causes the observed inter-basin gradients (a critical feature in the use of Nd as an ocean circulation tracer) to become severely damped, particularly away from direct input of fresh reactive phases with distinctive ϵ_{Nd} (Robinson et al., 2021; Abbott et al., 2019). Consistent with earlier studies (e.g.,

Deleted: Global volume-weighted distributions

Deleted: However, where including a reduced sediment source (e.g. compared to *EXPT_SED3* and *EXPT_SED4*) increases the relative importance of dust and river inputs, such as in the shallow Atlantic (Lambelet et al., 2016), simulated ϵ_{Nd} matches unradiogenic measurements with reasonable skill.

Arsouze et al., 2009; Jones et al., 2008), the largest model-data disparities occur in the North and equatorial Pacific, where simulated ϵ_{Nd} is far too unradiogenic compared to the observational data, pointing to a number of processes that may be better optimised in our Nd scheme. For example, a larger (and also more radiogenic) sediment source (i.e., greater f_{sed}) may be needed, particularly around shallow-intermediate marginal settings; a suggestion also supported by the too low $[Nd]_d$.

Total Nd concentrations and isotopic distributions show different responses to varying f_{sed} . We find that $[Nd]_d$ is sensitive to f_{sed} across all ocean basins (i.e., a wide divergence in the depth profiles show in Fig. 15), mostly at depths below 500 m where there is no direct influence from river and dust inputs and the relatively large area of the deep abyssal seafloor as an Nd interface becomes important (particularly with low reversible scavenging).

Deleted: Additionally, increased scavenging and a lower simulated residence time would improve the representation of localised Nd isotope signatures.




Figure 15: Central panel (g) displays $[Nd]_d$ at the seafloor (i.e., lowermost box of the ocean model) in simulation *EXPT_SED2* (100-year mean from the end of the run), with superimposed water column measurements (Osborne et al., 2017, 2015; GEOTRACES Intermediate Data Product Group, 2021) from $\geq 3,000$ m shown by filled coloured circles on the same colour scale. Surrounding panels (a-f) and (h-m) display depth profiles of simulated (coloured lines, one per sensitivity simulation with varied f_{sed}) and measured (filled circles) $[Nd]_d$. Larger shifts in the $[Nd]_d$ between simulations highlight regions most sensitive to the magnitude of the seafloor sediment source.

Deleted: -year

Typically, the best model-data fit for $[Nd]_d$ depth-profiles is achieved under lower f_{sed} (1.5 to 3.0 Gg yr^{-1}), and particularly in the Pacific, Indian and Southern Ocean (Fig. 15). Correspondingly, under high f_{sed} and particularly in the deeper interior of the ocean (below 3,000 m), simulated $[Nd]_d$ diverges to higher concentrations than measured due to too much Nd being sourced to the deep ocean from the sediment. Additionally, in these high f_{sed} scenarios, the strength of the deep sediment source obscures horizontal seafloor $[Nd]_d$ gradients across basins (see the near-uniform seafloor $[Nd]_d$ in Fig. 15g), masking the influence of reversible scavenging, which is controlled by the location and dissolution of particle fields and is important for governing $[Nd]_d$ patterns (Sect. 3.1).

Similar to the reversible scavenging sensitivity experiment (Sec. 3.1), the largest simulated offsets between simulated and measured $[Nd]_d$ under all f_{sed} experiments occur in the North Atlantic and sub-tropical Atlantic at depths above 1,000 m, where, even under the largest sediment fluxes, simulated $[Nd]_d$ is too low. The depth profile south of Greenland (Fig. 15c) shows the greatest sensitivity to varying f_{sed} in the upper 1,000 m, with *EXPT_SED4*'s simulated surface concentrations of 10 pmol kg^{-1} ($f_{sed} = 6.0 \text{ Gg yr}^{-1}$) being closest to the measured concentrations of 22 pmol kg^{-1} . By implication, the accurate representation of sediment fluxes is required to reproduce upper ocean $[Nd]_d$ in this region, and an enhanced sediment flux alone cannot account fully for the observed high surface concentrations. Either, a combination of the surface and near-surface fluxes resolved here are too diluted in the model (possibly due to grid box resolution), or the model is missing a significant surface/near-surface Nd source. Previous schemes have likewise simulated too low surface $[Nd]_d$ in the North Atlantic, likely also due to difficulties in representing highly localised and variable surface features in global models (Gu et al., 2019; Rempfer et al., 2011); more direct observations of surface fluxes would benefit future constraints in this region.

Whereas $[Nd]_d$ is sensitive to f_{sed} globally, the height of ϵ_{Nd} sensitivity is more regional (Fig. 16). The Atlantic, Indian and Southern Ocean ϵ_{Nd} are significantly more sensitive to changes in the bulk seafloor sediment Nd flux than the Pacific Ocean. Overall, differences in f_{sed} tend to drive whole depth profile shifts of low magnitude in ϵ_{Nd} , this contrasts the findings of Rempfer et al. (2011), who varied a margin constrained f_{sed} and reported deep water ϵ_{Nd} (below 1,000 m) were affected less than in our results. If this conflicting difference in deep ocean sensitivity to f_{sed} is because of the applied spatial distributions

Deleted: $\times 10^9$

Deleted: $\times 10^9$

Deleted: This indicates that under a relatively low reversible scavenging parameter, a corresponding smaller flux of Nd from the sediment is needed to balance the marine Nd budget, and by inference, a higher f_{sed} would be more appropriate with stronger reversible scavenging fluxes.

Deleted: Rempfer et al. (2011) found the opposite. In their study, doubling and halving f_{sed} both resulted in well pronounced nutrient-like profiles of $[Nd]_d$. This highlights the impact of having a global sediment source unrestricted in depth (our study) compared to limiting that flux to the upper 3,000 m (Rempfer et al., 2011) paired with a more efficient reversible scavenging (Rempfer et al., 2011) that allows the biogenic particle adsorption and desorption processes to dominate the deep-water distributions.

Deleted: $\times 10^9$

Deleted:

Deleted: .

Deleted:

of a sediment Nd source to seawater ($< 3,000$ m for Rempfer et al. (2011), all depths for us), then further constraints of sediment Nd fluxes across space and time are crucial when interpreting ocean circulation from ϵ_{Nd} .




Figure 16: Central panel (g) displays ϵ_{Nd} at the **seafloor (i.e., lowermost box of the ocean model)** in simulation *EXPT_SED2* (100-year mean from the end of the run), with superimposed water column measurements (Osborne et al., 2017, 2015; GEOTRACES Intermediate Data Product Group, 2021) from $\geq 3,000$ m shown by filled coloured circles on the same colour scale. Surrounding panels (a-f) and (h-m) display depth profiles of simulated (coloured lines, one per sensitivity simulation with varied f_{sed}) and measured (filled circles) ϵ_{Nd} . Larger shifts in the ϵ_{Nd} between simulations highlight regions most sensitive to the magnitude of the seafloor sediment source.

Once again, the most notable ϵ_{Nd} model-data mismatch occurs within the depth profiles of the North Atlantic. Due to major rivers delivering a high [Nd] load to the Atlantic (Fig. 5c), in their vicinity, simulations with low f_{sed} are conditioned towards riverine ϵ_{Nd} , providing the typical unradiogenic ϵ_{Nd} signature of NADW (Fig. 16a-d). Conversely, enhancing f_{sed} increases the fraction of Nd supplied to the ocean from the bulk seafloor sediment, which has more uniform, intermediate ϵ_{Nd} values in the central North Atlantic (-12.5) in contrast to the more unradiogenic ϵ_{Nd} signals of the continental margins and riverine source

Deleted: Interestingly, there is not a linear/direct response of ϵ_{Nd} to changing f_{sed} . Broadly and globally, the largest f_{sed} value of 6.0×10^9 g yr⁻¹ (*EXPT_SED4*) leads to more radiogenic ϵ_{Nd} and an intermediate f_{sed} value of 3.0×10^9 g yr⁻¹ (*EXPT_SED2*) leads to the most unradiogenic ϵ_{Nd} shift. The largest response in ϵ_{Nd} to changing f_{sed} occurs with a sediment flux between 3.0×10^9 and 4.5×10^9 g yr⁻¹. Increasing the flux beyond this produces a very weak response, indicating a threshold in ϵ_{Nd} sensitivity that results from an already dominant sediment source encompassing over 85% of the total Nd flux to seawater.

Deleted:

Deleted: sea

Deleted: and demonstrating the important balance between surface inputs and a sediment flux for simulating ϵ_{Nd} in the deep North Atlantic ...

in the Labrador (-28) and West Atlantic basins (-15.6), with localised near-surface sediment extreme minimums of (-34) in the northern Labrador Sea (Robinson et al., 2021). Greater f_{sed} thus acts to overprint and hence mix away the more distinct surface ϵ_{Nd} signal of NADW gained at their sites of deep-water formation in favour of a more general intermediate ϵ_{Nd} signal as it becomes exposed to Nd fluxes along its southward seafloor flow path. Model-measurement disparity at the mouth of the Labrador Sea and south of Greenland suggests that a specific fraction of the bulk sediment with more unradiogenic ϵ_{Nd} than is captured by Robinson et al. (2021) may be driving particle-seawater interactions (Fig. 17). Mechanical glacial erosion, which exposes large amounts of highly unradiogenic and labile fine grained crystalline detritus in the region surrounding the Labrador Sea likely drives enhanced localised extreme unradiogenic benthic fluxes (Von Blanckenburg and Nägler, 2001; Anderson, 2005). Moreover, marine particle ϵ_{Nd} measured in the North Atlantic has been shown to reflect the strong heterogeneity of the surrounding source rock, for example the provenance of extremely radiogenic particles (+6 to +9) at the Irminger Basin has been associated with mafic rocks found in East Greenland and Iceland (Stichel et al., 2020). It therefore follows that better spatial constraints on sediment fluxes are required for providing accurate ϵ_{Nd} tagging of Atlantic seawater.




Figure 17: Volume-weighted distributions of ϵ_{Nd} in simulation *EXPT_SED2* split into two different depth bins, (a) shallow (0-200 m), and (b) intermediate (200-1,000 m) within the North Atlantic and Labrador Sea basins. Water column measurements from within each depth bin (Osborne et al., 2017, 2015; GEOTRACES Intermediate Data Product Group, 2021) are superimposed as filled circles on the same colour scale. Note that Iceland is not resolved in FAMOUS (Jones, 2003).

In contrast to the Atlantic, the Pacific Ocean ϵ_{Nd} is the least sensitive to varying f_{sed} (Fig. 16e-f, h-i). Characteristically, the Pacific encompasses vast open ocean oligotrophic expanses with low biogenic particle export, in tandem with smaller relative dust and river sources, which means the seafloor sediment source already dominates the simulated Nd fluxes and distributions, even under lower sediment fluxes. In the North Pacific, the ϵ_{Nd} signal from the bulk seafloor sediment cannot explain the radiogenic ϵ_{Nd} observed here (Fig. 16), and so the model likely is not fully capturing the correct source (or the prescribed boundary condition is not correctly representing the distribution and fraction of the sediment phase contributing to the sediment flux) of Nd to seawater. Authigenic precipitation of clay minerals during reverse weathering is prevalent in the pelagic sediment throughout the North Pacific, these top centimetres of sediment are typically assumed to be unreactive, and act as an active

Deleted: marginal

Deleted: in all simulations at shallow and intermediate depths strongly

Deleted: is

Deleted: interacting with

Deleted:

Deleted: exchange

Deleted: in place of the bulk sediment ϵ_{Nd} signal

Deleted: This is further supported by recent core-top particle-seawater interaction investigations of the region (Blaser et al., 2016). It therefore follows that either existing seafloor sediment measurements do not characterise this region well, or a deep ocean benthic flux, or at least, an indiscriminate whole-ocean floor flux, is neither reasonable nor necessary (in fact, it is counterproductive) for providing

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Deleted: Furthermore, the bulk detrital ϵ_{Nd} (-8) released from the seafloor sediment in the Northwest Pacific (Fig. 6) is the same as the signal from the dust flux off the Asian continent, which contributes significant terrigenous material to the seafloor here (Nakai et al., 1993; Han et al., 2011), which makes it hard to distinguish between these two sources.

Deleted: the simulations with higher f_{sed} match the observed $[Nd]_d$ better than those with lower f_{sed} (Fig. 15), but

Deleted: Consistent with previous literature, the greatest imbalances in the Nd isotope budget occur in the reservoirs of the vast deep Pacific (Jones et al., 2008).

sink for REE (Kato et al., 2011). Moreover, precipitation at hydrothermal vents also acts as an effective localised REE sink (German et al., 1990). Therefore, the red clays of the deep Pacific may act as an effective Nd sink and not a source. Alternatively, a recent reactive transport model by Du et al. (2022) suggests reverse weathering maintains under-saturation of primary silicates in deep Pacific pore water, driving enhanced silicate weathering and favouring preferential dissolution of highly radiogenic volcanic sediments (e.g., basaltic glass and clinopyroxene). Here, we highlight the need for more precise constraints on complex benthic processes in the Pacific, including exploring the environmental conditions which would drive spatially elevated and 'reactivity weighted' sediment fluxes.

To conclude, we surmise that a sediment-seawater flux represents a key major source of $[Nd]_d$ that is particularly fundamental to the intermediate and deep ocean Nd budgets, and as such plays an important role in governing marine Nd cycling. Indeed, a strong seafloor-wide uniform sediment source pushes the ocean towards a more globally uniform and too high $[Nd]_d$ than measurements from the deep ocean suggest. This could be interpreted as evidence against a globally widespread benthic flux driven model of the marine Nd cycle with a spatially constant flux across diverse sedimentary environments, in favour of the more distinct $[Nd]_d$ distributions that may be achieved under a model of marine Nd cycling with larger and more heterogeneous surface and near surface Nd sources, and a greater dominance of reversible scavenging. Future work should explore employing a more horizontally nuanced benthic flux tied to local environmental and sedimentary conditions, in line with recent modelling studies (Pöppelmeier et al., 2020a; Pasquier et al., 2022), to introduce more spatial patterning in simulated $[Nd]_d$.

Certainly, these sensitivity simulations demonstrate that at least under a spatially uniform benthic flux, the bulk ϵ_{Nd} of seafloor detrital sediment (Robinson et al., 2021) cannot be considered fully representative of the ϵ_{Nd} composition of the sediment that is interacting with seawater in all instances. They highlight the need for observational and experimental quantification of the broad mobile Nd phases globally, and their ϵ_{Nd} signal, as well as constraints on the spatial distribution of such a benthic flux (e.g., identifying where and under what environmental conditions a benthic flux occurs, and at what strength). The model's response to f_{sed} sets the stage for further testing of global sedimentary ϵ_{Nd} on marine Nd cycling, providing the foundation for resolving the inherent complex multitude of processes.

4 Summary and conclusions

In this study, we describe the implementation of Nd isotopes (^{143}Nd and ^{144}Nd) into the ocean component of the FAMOUS GCM (Nd v1.0), providing a powerful tool designed for comprehensively exploring global marine Nd cycling, especially through representing explicit non-conservative processes to explore the extent of their influence on global seawater Nd distributions. Our Nd isotope scheme starts from previous Nd isotope implementations (Rempfer et al., 2011; Gu et al., 2019; Pöppelmeier et al., 2020a; Arsouze et al., 2009; Siddall et al., 2008), but revisits and updates Nd sources, sinks and tracer transformation in line with increased observations and recent findings relating to global marine Nd cycling.

Deleted: (

Deleted: ,

Deleted: This could imply that a 'reactivity weighted' sedimentary flux is most important for the deep North Pacific compared with other basins, providing evidence to support the preferential dissolution of reactive Nd sediment phases in the Pacific under a model of marine Nd cycling dominated by a seafloor sediment source.

Deleted: Notably, we find that ϵ_{Nd} is much less sensitive than $[Nd]_d$ to changing the rate of this flux (f_{sed}), and a high f_{sed} coupled with a weak reversible scavenging cannot account for $[Nd]_d$ gradients of increasing concentration along the thermohaline circulation in the deep ocean.

Deleted: indiscriminate

Deleted: 'bottom up' model

Deleted: (Haley et al., 2017)

Deleted: benthic

Deleted: the whole

Deleted: eafloor

Deleted: 'top-down'

Deleted: driven model with

Deleted: As such, although a benthic flux can represent the depth profiles of linearly increasing $[Nd]_d$ with depth, reversible scavenging is necessary and should not be considered a secondary process in controlling the global thermohaline variations in deep water $[Nd]_d$. Nonetheless, we acknowledge that

Deleted: may

Deleted: , reducing the importance of the reversible scavenging terms.

Deleted: openly

Deleted: D

Deleted: We present a new reference equilibrium pre-industrial simulation (XPDA) with improved basin scale physical ocean circulation performance relative to a standard version (XFHCC; Smith, 2012), though we note that for this study it was not important to evaluate the performance of the surface climate.

Deleted: Our model represents the main features of $[Nd]$ and ϵ_{Nd} well, although there is a tendency to simulate an over-pronounced vertical $[Nd]$ gradient, and to produce a too unradiogenic ϵ_{Nd} signal in the Pacific.

Model sensitivity to reversible scavenging efficiency demonstrates its importance for determining the increase in Nd concentration along the circulation pathway, and acts to enhance regional basinal gradients in simulated ϵ_{Nd} by maintaining the localised provenance signal. On the other hand, a widespread seafloor sedimentary flux presents a major deep ocean source of Nd, the magnitude of which governs horizontal seafloor Nd concentrations across ocean basins. In the deep North Pacific, simulated ϵ_{Nd} is too unradiogenic compared to measurements, highlighting that the ϵ_{Nd} of the sediment flux, as captured by the bulk ϵ_{Nd} , is not a true representation of the highly radiogenic labile sediment phase interacting with seawater. Moreover, authigenic precipitation may pose a missing effective Nd sink. Model-data mismatch at the mouth of the Labrador Sea suggests that detrital contributions from highly unradiogenic sediments disproportionately govern particle-seawater interactions. Overall, our results demonstrate that more precise constraints are needed to resolve the complex benthic processes which would drive spatially elevated and reactivity weighted sediment fluxes.

Exploring in detail the behaviour of simulated $[Nd]_d$ and ϵ_{Nd} distributions also highlighted some of the structural limitations of the model (e.g., difficulties representing highly localised and surface features) and influential biases in the physical ocean circulation (e.g., limited northward intrusion of AABW in the North Atlantic).

This study provides the groundwork for a future comprehensive optimisation of the marine Nd isotope scheme in FAMOUS. In the first instance, we suggest calibration of the key tuning parameters ($[Nd]_p/[Nd]_d$ and f_{sed}) to best represent modern seawater measurements. Additionally, it would be beneficial to obtain additional observational constraints on the broad labile sediment ϵ_{Nd} interacting with seawater across different seafloor regions, including constraining the exchange between authigenic and detrital sediment phases during early diagenesis. Future sensitivity studies could also focus on the influence of river particulate and continental marginal sources on marine Nd to provide further insight to (and possibly constrain) the relative importance of these inputs, as opposed to a predominantly benthic seafloor-wide source.

This new model scheme can aid in the delivery of more robust applications of ϵ_{Nd} as a modern and palaeo-tracer. It provides a platform for dynamic modelling under different modes of marine Nd cycling or varied climatic and oceanographic conditions, enabling current hypotheses to be tested rigorously with the aim of constraining Nd cycling under a complex, partially understood marine geochemical system.

Code Availability

The code detailing the advances for the marine Nd isotope scheme (Nd_v1.0) described in this manuscript is available via the Research Data Leeds Repository (Robinson 2022: <https://doi.org/10.5518/1136>) under a Creative Commons Attribution 4.0 International (CCBY 4.0) license. These files are known as code modification (i.e. 'mod') files and should be applied to the original FAMOUS model code, which is protected under UK Crown Copyright and can be obtained from the National Centre

Deleted: The presented sensitivity experiments demonstrate that the Nd isotope scheme in the FAMOUS GCM is sensitive to Nd source and biogeochemical processes.

Formatted: Strikethrough

Deleted: with depth and

Deleted: Moreover, reversible scavenging

Deleted: homogenise ϵ_{Nd} vertically in the water column,

Deleted: ing

Deleted: benthic flux

Deleted: , a term referring to a multitude of processes encompassing boundary particle-seawater exchange along the continental margins (Lacan and Jeandel, 2005), submarine groundwater discharge (Johannesson and Burdige, 2007), and a benthic flux released from pore waters (Abbott et al., 2015a),

Deleted: , the magnitude of which governs horizontal seafloor Nd concentrations across ocean basins. The weak sensitivity of simulated ϵ_{Nd} in the deep North Pacific implies that with a seafloor-wide benthic flux of marine Nd, the ϵ_{Nd} of the sediment flux, as captured by the bulk ϵ_{Nd} , is not a true representation in all instances of the labile sediment phase interacting with seawater in this basin. Alternatively, it may also indicate that there is a significant missing source of radiogenic Nd to seawater, likely of volcanic origin; or possibly a combination of both explanations. Furthermore, model-data mismatch at the mouth of the Labrador Sea suggests that the labile sediment Nd phases interacting with seawater in the northern North Atlantic are considerably more unradiogenic than captured by the bulk sediment.

Deleted: These

Deleted: results

Deleted: achieve target Nd inventories and residence times

Deleted: This would improve the boundary conditions we can feed into the model, giving it the best chance to simulate Nd distributions accurately and making the interpretation of model performance (and thus also of the relative importance of different sources and sinks) more straightforward.

Deleted: in order to

Formatted

Deleted: Implementing Nd isotopes in a fast GCM provides a useful tool for exploring model-scheme sensitivities and uncertain marine biogeochemical processes. This framework allows for performing the long integrations necessary to spin-up ocean physics and biogeochemistry, which can be simulated for multiple time periods (e.g. in palaeo studies) in equilibrium (i.e. with fixed forcing, Haywood et al., 2016; Lunt et al., 2017) and transient (i.e. with temporally evolving forcing, e.g. Ivanovic et al., 2016; Menviel et al., 2019) scenarios. Alternatively, or in addition, large ensembles of simulations can be run for [re]calibrating (i.e., 'tuning') the model or quantifying uncertainty in the inputs (boundary conditions, parameter values etc.). In this way, it becomes possible to build on new ... [5]

Deleted: (e.g., the balancing of 'top down' versus 'bottom up' fluxes) ...

Deleted: D

for Atmospheric Science (NCAS) Computational Modelling Services (CMS): <https://cms.ncas.ac.uk/>, with specific FAMOUS documentation available at: <https://cms.ncas.ac.uk/miscellaneous/um-famous/>. All files and corresponding information that needs to be applied to configure each individual simulation presented here are available from the same DOI (above); note that to complete the setup of these simulations, line 4909 in each simulation's tracer.f file needs updating with the corresponding RS_TUNE value as listed in Table 3. A complete version of the modified code for the EXPT_RS4 simulation using FAMOUS-MOSES1, including Nd isotope implementation, is archived at the Research Data Leeds Repository, and linked from the DOI above.

Control candidate simulations for new FAMOUS reference

- XPDAA control simulation (0-5,000 years)
- XPDAB control candidate simulation (0-5,000 years)
- XPDAC control candidate simulation (0-5,000 years)
- XPDEA control candidate simulation (0-5,000 years)

Reversible scavenging efficiency ($[Nd]_p/[Nd]_d$) sensitivity simulations

- XPDAl $[Nd]_p/[Nd]_d=0.001$ (0-9,000 years)
- XPDAD $[Nd]_p/[Nd]_d=0.002$ (0-9,000 years)
- XPDAl $[Nd]_p/[Nd]_d=0.003$ (0-9,000 years)
- XPDAl $[Nd]_p/[Nd]_d=0.004$ (0-9,000 years)
- XPDAl $[Nd]_p/[Nd]_d=0.005$ (0-9,000 years)
- XPDAl $[Nd]_p/[Nd]_d=0.006$ (0-9,000 years)

Total Nd source from sediment (f_{sed}) sensitivity simulations

- XPDAl $f_{sed}=1.5 \text{ Gg yr}^{-1}$ (0-9,000 years)
- XPDAl $f_{sed}=3.0 \text{ Gg yr}^{-1}$ (0-9,000 years)
- XPDAl $f_{sed}=4.5 \text{ Gg yr}^{-1}$ (0-9,000 years)
- XPDAl $f_{sed}=6.0 \text{ Gg yr}^{-1}$ (0-9,000 years)

Deleted: $\times 10^9$

Deleted:

Deleted: $\times 10^9$

Deleted: $\times 10^9$

Deleted: $\times 10^9$

Data Availability

The data are available via the Research Data Leeds Repository (Robinson 2022a: <https://doi.org/10.5518/1136>)

1795 **Supplementary Material**

The supplement related to this article is to be associated with a DOI, and currently attached as a .zip (Supplementary_Robinson.et.al_inPrep.zip), containing:

- Supplementary Information:** Documented supplementary text, figures, and tables (Text S1, Table S1-S2, Figures S1-S11)
Table S3: Spreadsheet of seawater Nd concentration and isotope measurements and references used to validate model scheme
simulation_files/: Folder with all the simulation files needed to run each simulation
standard_famous_mods/: Folder containing all standard FAMOUS GCM modification files

Author contributions

RFI designed the project with SMR, and supervised its completion. SMR wrote and implemented the code with technical input from JCT, RFI and LJG. SMR completed the experiment design, ran the simulations, analysed results, and prepared the manuscript with input from all co-authors. All co-authors contributed scientific expertise throughout. PIV designed and ran the pre-industrial FAMOUS GCM perturbed physics ensemble. FP updated the dust source ϵ_{Nd} boundary conditions. YP provided the seawater REE compilation, which was updated by SMR.

Competing interests

The authors declare that they have no conflict of interest.

1810 **Acknowledgements**

SMR was funded by the Natural Environment Research Council (NERC) SPHERES Doctoral Training Partnership (grant number: NE/ L002574/1). LJG was funded by UKRI Future Leaders Fellowship ‘SMBGen’ (grant number: MR/S016961/1). JCT was supported through the Centre for Environmental Modelling and Computation (CEMAC), University of Leeds. FP was supported by the European Union’s Horizon 2020 program (grant number: 101023443). TvdF acknowledges funding from the NERC grant NE/ P019080/1. Our heartfelt thanks go to Dr Jamie D Wilson for valuable discussions when implementing the reversible scavenging scheme, and to Professor Andy Ridgwell for helpful discussions during the design of code developments. We would like to thank all Reviewers for taking the time and effort necessary to review the manuscript. We sincerely appreciate all valuable comments and suggestions, which helped us to improve the quality of the manuscript.

References

1820 Abbott, A. N.: A benthic flux from calcareous sediments results in non-conservative neodymium behavior during lateral transport: A study from the Tasman Sea, Geology, 47, 363–366, <https://doi.org/10.1130/g45904.1>, 2019.

Deleted: 3

Deleted: 9

Deleted: 4

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) Times New Roman, Font colour: Auto, Pattern: Clear

Formatted: Font: (Default) Times New Roman, Not Bold, Font colour: Auto, Pattern: Clear

Formatted: Font: (Default) Times New Roman, Not Bold, Font colour: Auto, Pattern: Clear

- 1825 Abbott, A. N., Haley, B. A., and McManus, J.: Bottoms up: Sedimentary control of the deep North Pacific Ocean's ϵNd signature, *Geology*, 43, 1035–1038, <https://doi.org/10.1130/G37114.1>, 2015a.
- Abbott, A. N., Haley, B. A., McManus, J., and Reimers, C. E.: The sedimentary flux of dissolved rare earth elements to the ocean, *Geochim. Cosmochim. Acta*, 154, 186–200, <https://doi.org/10.1016/j.gca.2015.01.010>, 2015b.
- Abbott, A. N., Löhr, S., and Trethrew, M.: Are clay minerals the primary control on the oceanic rare earth element budget?, *Front. Mar. Sci.*, 6, 1–19, <https://doi.org/10.3389/fmars.2019.00504>, 2019.
- 1830 Anderson, S. P.: Glaciers show direct linkage between erosion rate and chemical weathering fluxes, 67, 147–157, <https://doi.org/10.1016/j.geomorph.2004.07.010>, 2005.
- Arsouze, T., Dutay, J.-C. C., Lacan, F., and Jeandel, C.: Modeling the neodymium isotopic composition with a global ocean circulation model, *Chem. Geol.*, 239, 165–177, <https://doi.org/10.1016/j.chemgeo.2006.12.006>, 2007.
- 1835 Arsouze, T., Dutay, J. C., Lacan, F., and Jeandel, C.: Reconstructing the Nd oceanic cycle using a coupled dynamical-Biogeochemical model, 6, 2829–2846, <https://doi.org/10.5194/bg-6-2829-2009>, 2009.
- Ayache, M., Dutay, J. C., Arsouze, T., Révillon, S., Beuvier, J., and Jeandel, C.: High-resolution neodymium characterization along the Mediterranean margins and modelling of Nd distribution in the Mediterranean basins, 13, 5259–5276, <https://doi.org/10.5194/bg-13-5259-2016>, 2016.
- 1840 Ayache, M., Dutay, J., Tachikawa, K., Arsouze, T., and Jeandel, C.: Neodymium budget in the Mediterranean Sea : Evaluating the role of atmospheric dusts using a high-resolution dynamical-biogeochemical model, *Biogeosciences Discuss.*, 1–35, 2022.
- Bacon, M. P. and Anderson, R. F.: Distribution of Thorium Isotopes Between Dissolved and Particulate Forms in the Deep Sea, *J. Geophys. Res.*, 87, 2045–2056, <https://doi.org/10.1029/JC087iC03p02045>, 1982.
- 1845 Bayon, G., German, C. R., Burton, K. W., Nesbitt, R. W., and Rogers, N.: Sedimentary Fe-Mn oxyhydroxides as paleoceanographic archives and the role of aeolian flux in regulating oceanic dissolved REE, *Earth Planet. Sci. Lett.*, 224, 477–492, <https://doi.org/10.1016/j.epsl.2004.05.033>, 2004.
- Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration, *Limnol. Oceanogr. Lmnnl. Oceanogr.*, 42, 1997.
- 1850 Bertram, C. J. and Elderfield, H.: The geochemical balance of the rare earth elements and neodymium isotopes in the oceans, *Geochim. Cosmochim. Acta*, 57, 1957–1986, [https://doi.org/10.1016/0016-7037\(93\)90087-D](https://doi.org/10.1016/0016-7037(93)90087-D), 1993.
- Blanchet, C. L.: A database of marine and terrestrial radiogenic Nd and Sr isotopes for tracing earth-surface processes, *Earth Syst. Sci. Data*, 11, 741–759, <https://doi.org/10.5194/essd-11-741-2019>, 2019.
- Von Blanckenburg, F. and Nägler, T. F.: Weathering versus circulation-controlled changes in radiogenic isotope tracer composition of the Labrador Sea and North Atlantic Deep Water, *Paleoceanography*, 16, 424–434, <https://doi.org/10.1029/2000PA000550>, 2001.
- 1855 Blaser, P., Frank, M., and van de Fliedert, T.: Revealing past ocean circulation with neodymium isotopes, *Past Glob. Chang. Mag.*, 27, 54–55, <https://doi.org/10.22498/pages.27.2.54>, 2019a.
- Blaser, P., Pöppelmeier, F., Schulz, H., Gutjahr, M., Frank, M., Lippold, J., Heinrich, H., Link, J. M., Hoffmann, J., Szidat, S., and Frank, N.: The resilience and sensitivity of Northeast Atlantic deep water ϵNd to overprinting by detrital fluxes over the past 30,000 years, *Geochim. Cosmochim. Acta*, 245, 79–97, <https://doi.org/10.1016/j.gca.2018.10.018>, 2019b.
- 1860 Blaser, P., Gutjahr, M., Pöppelmeier, F., Frank, M., Kaboth-Bahr, S., and Lippold, J.: Labrador Sea bottom water provenance and REE exchange during the past 35,000 years, *Earth Planet. Sci. Lett.*, 542, 116299, <https://doi.org/10.1016/j.epsl.2020.116299>, 2020.
- Chamecki, M., Chor, T., Yang, D., and Meneveau, C.: Material Transport in the Ocean Mixed Layer: Recent Developments Enabled by Large Eddy Simulations, <https://doi.org/10.1029/2019RG000655>, 2019.
- 1865

- Chavagnac, V., Saleban Ali, H., Jeandel, C., Leleu, T., Destrigneville, C., Castillo, A., Cotte, L., Waeles, M., Cathalot, C., Laes-Huon, A., Pelleter, E., Nonnotte, P., Sarradin, P. M., and Cannat, M.: Sulfate minerals control dissolved rare earth element flux and Nd isotope signature of buoyant hydrothermal plume (EMSO-Azores, 37°N Mid-Atlantic Ridge), *Chem. Geol.*, 499, 111–125, <https://doi.org/10.1016/j.chemgeo.2018.09.021>, 2018.
- 1870 Chen, T. Y., Stumpf, R., Frank, M., Beldowski, J., and Staubwasser, M.: Contrasting geochemical cycling of hafnium and neodymium in the central Baltic Sea, *Geochim. Cosmochim. Acta*, 123, 166–180, <https://doi.org/10.1016/j.gca.2013.09.011>, 2013.
- Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.:
1875 Development and evaluation of an Earth-System model - HadGEM2, *Geosci. Model Dev.*, 4, 1051–1075, <https://doi.org/10.5194/gmd-4-1051-2011>, 2011.
- Cox, P. M., Betts, R. A., Bunton, C. B., Essery, R. L. H., Rowntree, P. R., and Smith, J.: The impact of new land surface physics on the GCM simulation of climate and climate sensitivity, *Clim. Dyn.*, 15, 183–203, <https://doi.org/10.1007/s003820050276>, 1999.
- 1880 Crossley, J. F. and Roberts, D. L.: THE THERMODYNAMIC / DYNAMIC SEA ICE MODEL, Unified Model Doc. Pap. No 45, 1995.
- Dai, A. and Trenberth, K. E.: Estimates of Freshwater Discharge from Continents : Latitudinal and Seasonal Variations, *J. Hydrometeorol.*, 3, 660–687, 2002.
- Deng, K., Yang, S., Du, J., Lian, E., and Vance, D.: Dominance of benthic flux of REEs on continental shelves: implications
1885 for oceanic budgets, *Geochemical Perspect. Lett.*, 22, 26–30, <https://doi.org/10.7185/geochemlet.2223>, 2022.
- Dentith, J. E., Ivanovic, R. F., Gregoire, L. J., Tindall, J. C., and Smith, R. S.: Ocean circulation drifts in multi-millennial climate simulations: the role of salinity corrections and climate feedbacks, *Clim. Dyn.*, 52, 1761–1781, <https://doi.org/10.1007/s00382-018-4243-y>, 2019.
- Dentith, J. E., Ivanovic, R. F., Gregoire, L. J., Tindall, J. C., and Robinson, L. F.: Simulating stable carbon isotopes in the
1890 ocean component of the FAMOUS general circulation model with MOSES1 (XOAVI), *Geosci. Model Dev.*, 13, 3529–3552, <https://doi.org/10.5194/gmd-13-3529-2020>, 2020.
- Doney, S. C., Lindsay, K., Caldeira, K., Campin, J. M., Drange, H., Dutay, J. C., Follows, M., Gao, Y., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Madec, G., Maier-Reimer, E., Marshall, J. C., Matear, R. J., Monfray, P., Mouchet, A., Najjar, R., Orr, J. C., Plattner, G. K., Sarmiento, J., Schlitzer, R., Slater, R., Totterdell, I. J., Weirig, M. F., Yamanaka, Y., and Yool, A.: Evaluating global ocean carbon models: The importance of realistic physics, *Global Biogeochem. Cycles*, 18, 3017, <https://doi.org/10.1029/2003GB002150>, 2004.
- 1895 Du, J., Haley, B. A., and Mix, A. C.: Neodymium isotopes in authigenic phases, bottom waters and detrital sediments in the Gulf of Alaska and their implications for paleo-circulation reconstruction, *Geochim. Cosmochim. Acta*, 193, 14–35, <https://doi.org/10.1016/j.gca.2016.08.005>, 2016.
- 1900 Du, J., Haley, B. A., Mix, A. C., Walczak, M. H., and Praetorius, S. K.: Flushing of the deep Pacific Ocean and the deglacial rise of atmospheric CO₂ concentrations, *Nat. Geosci.*, 11, 749–755, <https://doi.org/10.1038/s41561-018-0205-6>, 2018.
- Du, J., Haley, B. A., and Mix, A. C.: Evolution of the Global Overturning Circulation since the Last Glacial Maximum based on marine authigenic neodymium isotopes, *Quat. Sci. Rev.*, 241, 106396, <https://doi.org/10.1016/j.quascirev.2020.106396>, 2020.
- 1905 Du, J., Haley, B. A., Mix, A. C., Abbott, A. N., McManus, J., and Vance, D.: Reactive-transport modeling of neodymium and its radiogenic isotope in deep-sea sediments: The roles of authigenesis, marine silicate weathering and reverse weathering, *Earth Planet. Sci. Lett.*, 596, 117792, <https://doi.org/10.1016/j.epsl.2022.117792>, 2022.
- Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling

- through the ocean interior and on the seafloor, *Global Biogeochem. Cycles*, 21, 1–16, <https://doi.org/10.1029/2006GB002907>, 2007.
- 1910 Dunne, J. P., Hales, B., and Toggweiler, J. R.: Global calcite cycling constrained by sediment preservation controls, *Global Biogeochem. Cycles*, 26, 1–14, <https://doi.org/10.1029/2010GB003935>, 2012.
- Dutay, J. C., Lacan, F., Roy-Barman, M., and Bopp, L.: Influence of particle size and type on ^{231}Pa and ^{230}Th simulation with a global coupled biogeochemical-ocean general circulation model: A first approach, *Geochemistry, Geophys. Geosystems*, 10, 1–26, <https://doi.org/10.1029/2008GC002291>, 2009.
- 1915 Elderfield, H., Upstill-Goddard, R., and Sholkovitz, E. R.: The rare earth elements in rivers, estuaries, and coastal seas and their significance to the composition of ocean waters, *Geochim. Cosmochim. Acta*, 54, 971–991, [https://doi.org/10.1016/0016-7037\(90\)90432-K](https://doi.org/10.1016/0016-7037(90)90432-K), 1990.
- Essery, R. and Clark, D. B.: Developments in the MOSES 2 land-surface model for PILPS 2e, *Glob. Planet. Change*, 38, 161–164, [https://doi.org/10.1016/S0921-8181\(03\)00026-2](https://doi.org/10.1016/S0921-8181(03)00026-2), 2003.
- 1920 Essery, R. L. H., Best, M. J., and Cox, P. M.: MOSES 2.2 technical documentation. Hadley Centre Technical Note 30, 2001.
- Ferreira, D., Cessi, P., Coxall, H. K., De Boer, A., Dijkstra, H. A., Drijfhout, S. S., Eldevik, T., Harnik, N., McManus, J. F., Marshall, D. P., Nilsson, J., Roquet, F., Schneider, T., and Wills, R. C.: Atlantic-Pacific Asymmetry in Deep Water Formation, <https://doi.org/10.1146/annurev-earth-082517-010045>, 2018.
- 1925 van de Flierdt, T., Griffiths, A. M., Lambelet, M., Little, S. H., Stichel, T., and Wilson, D. J.: Neodymium in the oceans: A global database, a regional comparison and implications for palaeoceanographic research, <https://doi.org/10.1098/rsta.2015.0293>, 2016.
- Frajka-Williams, E., Ansorge, I. J., Baehr, J., Bryden, H. L., Chidichimo, M. P., Cunningham, S. A., Danabasoglu, G., Dong, S., Donohue, K. A., Elipot, S., Heimbach, P., Holliday, N. P., Hummels, R., Jackson, L. C., Karstensen, J., Lankhorst, M., Le Bras, I. A., Susan Lozier, M., McDonagh, E. L., Meinen, C. S., Mercier, H., Moat, B. I., Perez, R. C., Piecuch, C. G., Rhein, M., Srokosz, M. A., Trenberth, K. E., Bacon, S., Forget, G., Goni, G., Kieke, D., Koelling, J., Lamont, T., McCarthy, G. D., Mertens, C., Send, U., Smeed, D. A., Speich, S., van den Berg, M., Volkov, D., and Wilson, C.: Atlantic meridional overturning circulation: Observed transport and variability, <https://doi.org/10.3389/fmars.2019.00260>, 2019.
- 1930 Frank, M.: Radiogenic isotopes: Tracers of past ocean circulation and erosional input, *Rev. Geophys.*, 40, 1001, <https://doi.org/10.1029/2000RG000094>, 2002.
- 1935 GEOTRACES Intermediate Data Product Group: The GEOTRACES Intermediate Data Product 2021 (IDP2021), <https://doi.org/5/cf2d9ba9-d51d-3b7c-e053-8486abc0f5fd>, 2021.
- German, C. R., Klinkhammer, G. P., Edmond, J. M., Mitrat, A., and H. Elderfield: Hydrothermal scavenging of rare-earth elements in the ocean, *Nature*, 345, 516–518, 1990.
- 1940 Goldstein, S. J. and Jacobsen, S. B.: The Nd and Sr isotopic systematics of river-water dissolved material: Implications for the sources of Nd and Sr in seawater, *Chemical Geology (Isotope Geoscience Section)*, 245–272 pp., [https://doi.org/10.1016/0168-9622\(87\)90045-5](https://doi.org/10.1016/0168-9622(87)90045-5), 1987.
- Goldstein, S. L. and Hemming, S. R.: Long-lived Isotopic Tracers in Oceanography, Paleoceanography, and Ice-Sheet Dynamics, 453–483, <https://doi.org/10.1016/B0-08-043751-6/06179-X>, 2003.
- 1945 Goldstein, S. L. and O’Nions, R. K.: Nd and Sr isotopic relationships in pelagic clays and gerromanganese deposits, *Nature*, 292, 324–327, 1981.
- Goldstein, S. L., O’Nions, R. K., and Hamilton, P. J.: A Sm-Nd isotopic study of atmospheric dusts and particulates from major river systems, *Earth Planet. Sci. Lett.*, 70, 221–236, [https://doi.org/10.1016/0012-821X\(84\)90007-4](https://doi.org/10.1016/0012-821X(84)90007-4), 1984.
- 1950 Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, T. C., Mitchell, J. F. B., and Wood, R. A.: The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux

- adjustments, *Clim. Dyn.*, 16, 147–168, <https://doi.org/10.1007/s003820050010>, 2000.
- Greaves, M. J., Statham, P. J., and Elderfield, H.: Rare earth element mobilization from marine atmospheric dust into seawater, *Mar. Chem.*, 46, 255–260, [https://doi.org/10.1016/0304-4203\(94\)90081-7](https://doi.org/10.1016/0304-4203(94)90081-7), 1994.
- 1955 Gregoire, L. J., Valdes, P. J., Payne, A. J., and Kahana, R.: Optimal tuning of a GCM using modern and glacial constraints, *Clim. Dyn.*, 37, 705–719, <https://doi.org/10.1007/s00382-010-0934-8>, 2011.
- Gregoire, L. J., Payne, A. J., and Valdes, P. J.: Deglacial rapid sea level rises caused by ice-sheet saddle collapses, *Nature*, 487, 219–222, <https://doi.org/10.1038/nature11257>, 2012.
- Gregoire, L. J., Valdes, P. J., and Payne, A. J.: The relative contribution of orbital forcing and greenhouse gases to the North American deglaciation, *Geophys. Res. Lett.*, 42, 9970–9979, <https://doi.org/10.1002/2015GL066005>, 2015.
- 1960 Gregoire, L. J., Otto-Bliesner, B., Valdes, P. J., and Ivanovic, R.: Abrupt Bolling warming and ice saddle collapse contributions to the Meltwater Pulse 1a rapid sea level rise, *Geophys. Res. Lett.*, 43, 9130–9137, <https://doi.org/10.1002/2016GL070356>, 2016.
- Grénier, M., Jeandel, C., Lacan, F., Vance, D., Venchiarutti, C., Cros, A., and Cravatte, S.: From the subtropics to the central equatorial Pacific Ocean: Neodymium isotopic composition and rare earth element concentration variations, *J. Geophys. Res. Ocean.*, 118, 592–618, <https://doi.org/10.1029/2012JC008239>, 2013.
- 1965 Grousset, F. E., Biscaye, P. E., Zindler, A., Prospero, J., and Chester, R.: Neodymium isotopes as tracers in marine sediments and aerosols: North Atlantic, *Earth Planet. Sci. Lett.*, 87, 367–378, [https://doi.org/10.1016/0012-821X\(88\)90001-5](https://doi.org/10.1016/0012-821X(88)90001-5), 1988.
- Grousset, F. E., Parra, M., Bory, A., Martinez, P., Bertrand, P., Shimmield-M, G., and Ellam, R. M.: Saharan Wind Regimes Traced by the Sr-Nd Isotopic Composition of Subtropical Atlantic Sediments: Last Glacial Maximum vs Today, *Quaternary Science Reviews*, 395–409 pp., [https://doi.org/https://doi.org/10.1016/S0277-3791\(97\)00048-6](https://doi.org/https://doi.org/10.1016/S0277-3791(97)00048-6), 1998.
- 1970 Gu, S., Liu, Z., Jahn, A., Rempfer, J., Zhang, J., and Joos, F.: Modeling Neodymium Isotopes in the Ocean Component of the Community Earth System Model (CESM1), *J. Adv. Model. Earth Syst.*, 11, 624–640, <https://doi.org/10.1029/2018MS001538>, 2019.
- Haley, B. A., Klinkhammer, G. P., and McManus, J.: Rare earth elements in pore waters of marine sediments, *Geochim. Cosmochim. Acta*, 68, 1265–1279, <https://doi.org/10.1016/j.gca.2003.09.012>, 2004.
- 1975 Haley, B. A., Frank, M., Hathorne, E., and Pias, N.: Biogeochemical implications from dissolved rare earth element and Nd isotope distributions in the Gulf of Alaska, *Geochim. Cosmochim. Acta*, 126, 455–474, <https://doi.org/10.1016/j.gca.2013.11.012>, 2014.
- Haley, B. A., Du, J., Abbott, A. N., and McManus, J.: The Impact of Benthic Processes on Rare Earth Element and Neodymium Isotope Distributions in the Oceans, *Front. Mar. Sci.*, 4, 426, <https://doi.org/10.3389/fmars.2017.00426>, 2017.
- 1980 Henderson, G. M., Heinze, C., Anderson, R. F., and Winguth, A. M. E.: Global distribution of the ^{230}Th flux to ocean sediments constrained by GCM modelling, *Deep. Res. Part I Oceanogr. Res. Pap.*, 46, 1861–1893, [https://doi.org/10.1016/S0967-0637\(99\)00030-8](https://doi.org/10.1016/S0967-0637(99)00030-8), 1999.
- Hopcroft, P. O. and Valdes, P. J.: Last glacial maximum constraints on the Earth System model HadGEM2-ES, *Clim. Dyn.*, 45, 1657–1672, <https://doi.org/10.1007/s00382-014-2421-0>, 2015.
- 1985 Hopcroft, P. O., Valdes, P. J., Woodward, S., and Joshi, M. M.: Last glacial maximum radiative forcing from mineral dust aerosols in an Earth system model, *J. Geophys. Res.*, 120, 8186–8205, <https://doi.org/10.1002/2015JD023742>, 2015.
- Jacobsen, S. B. and Wasserburg, G. J.: Sm-Nd evolution of chondrites, *Earth Planet. Sci. Lett.*, 50, 139–155, [https://doi.org/10.1016/0012-821X\(80\)90125-9](https://doi.org/10.1016/0012-821X(80)90125-9), 1980.
- 1990 Jeandel, C.: Concentration and isotopic composition of Nd in the South Atlantic Ocean, *Earth Planet. Sci. Lett.*, 117, 581–591, [https://doi.org/10.1016/0012-821X\(93\)90104-h](https://doi.org/10.1016/0012-821X(93)90104-h), 1993.

- Jeandel, C., Bishop, J. K., and Zindler, A.: Exchange of neodymium and its isotopes between seawater and small and large particles in the Sargasso Sea, *Geochim. Cosmochim. Acta*, 59, 535–547, [https://doi.org/10.1016/0016-7037\(94\)00367-U](https://doi.org/10.1016/0016-7037(94)00367-U), 1995.
- 1995 Jeandel, C., Arsouze, T., Lacan, F., Téchiné, P., and Dutay, J. C.: Isotopic Nd compositions and concentrations of the lithogenic inputs into the ocean: A compilation, with an emphasis on the margins, *Chem. Geol.*, 239, 156–164, <https://doi.org/10.1016/j.chemgeo.2006.11.013>, 2007.
- Johannesson, K. H. and Burdige, D. J.: Balancing the global oceanic neodymium budget: Evaluating the role of groundwater, *Earth Planet. Sci. Lett.*, 253, 129–142, <https://doi.org/10.1016/j.epsl.2006.10.021>, 2007.
- 2000 Jones, C.: A fast ocean GCM without flux adjustments, *J. Atmos. Ocean. Technol.*, 20, 1857–1868, [https://doi.org/10.1175/1520-0426\(2003\)020<1857:AFOGWF>2.0.CO;2](https://doi.org/10.1175/1520-0426(2003)020<1857:AFOGWF>2.0.CO;2), 2003.
- Jones, C., Gregory, J., Thorpe, R., Cox, P., Murphy, J., Sexton, D., and Valdes, P.: Systematic optimisation and climate simulation of FAMOUS, a fast version of HadCM3, *Clim. Dyn.*, 25, 189–204, <https://doi.org/10.1007/s00382-005-0027-2>, 2005.
- 2005 Jones, K. M., Khatiwala, S. P., Goldstein, S. L., Hemming, S. R., and van de Flierdt, T.: Modeling the distribution of Nd isotopes in the oceans using an ocean general circulation model, *Earth Planet. Sci. Lett.*, 272, 610–619, <https://doi.org/10.1016/j.epsl.2008.05.027>, 2008.
- Kato, Y., Fujinaga, K., Nakamura, K., Takaya, Y., Kitamura, K., Ohta, J., Toda, R., Nakashima, T., and Iwamori, H.: Deep-sea mud in the Pacific Ocean as a potential resource for rare-earth elements, *Nat. Geosci.*, 4, 535–539, <https://doi.org/10.1038/ngeo1185>, 2011.
- 2010 Kriest, I. and Oschlies, A.: On the treatment of particulate organic matter sinking in large-scale models of marine biogeochemical cycles, 5, 55–72, <https://doi.org/10.5194/bg-5-55-2008>, 2008.
- Kuhlbrodt, T., Griesel, A., Montoya, M., Levermann, A., Hofmann, M., and Rahmstorf, S.: On the driving processes of the Atlantic meridional overturning circulation, *Rev. Geophys.*, 45, 2001, <https://doi.org/10.1029/2004RG000166>, 2007.
- 2015 Lacan, F. and Jeandel, C.: Neodymium isotopes as a new tool for quantifying exchange fluxes at the continent-ocean interface, *Earth Planet. Sci. Lett.*, 232, 245–257, <https://doi.org/10.1016/j.epsl.2005.01.004>, 2005.
- Lagarde, M., Lemaitre, N., Planquette, H., Grenier, M., Belhadj, M., Lherminier, P., and Jeandel, C.: Particulate Rare Earth Element behavior in the North Atlantic (GEOVIDE cruise), 17, 5539–5561, <https://doi.org/10.5194/bg-17-5539-2020>, 2020.
- Lambelet, M., van de Flierdt, T., Crocket, K., Rehkämper, M., Kreissig, K., Coles, B., Rijkenberg, M. J. A., Gerringa, L. J. A., de Baar, H. J. W., and Steinfeldt, R.: Neodymium isotopic composition and concentration in the western North Atlantic Ocean: Results from the GEOTRACES GA02 section, *Geochim. Cosmochim. Acta*, 177, 1–29, <https://doi.org/10.1016/j.gca.2015.12.019>, 2016.
- 2020 Lambelet, M., van de Flierdt, T., Butler, E. C. V., Bowie, A. R., Rintoul, S. R., Watson, R. J., Remenyi, T., Lannuzel, D., Warner, M., Robinson, L. F., Bostock, H. C., and Bradtmiller, L. I.: The Neodymium Isotope Fingerprint of Adélie Coast Bottom Water, *Geophys. Res. Lett.*, 45, 11,247–11,256, <https://doi.org/10.1029/2018GL080074>, 2018.
- 2025 Laws, E. A., Falkowski, P. G., Smith, W. O., Ducklow, H., and McCarthy, J. J.: Temperature effects on export production in the open ocean, *Global Biogeochem. Cycles*, 14, 1231–1246, <https://doi.org/10.1029/1999GB001229>, 2000.
- Mahowald, N. M., Muhs, D. R., Levis, S., Rasch, P. J., Yoshioka, M., Zender, C. S., and Luo, C.: Change in atmospheric mineral aerosols in response to climate: Last glacial period, preindustrial, modern, and doubled carbon dioxide climates, *J. Geophys. Res. Atmos.*, 111, <https://doi.org/10.1029/2005JD006653>, 2006.
- 2030 Maier-Reimer, E.: Geochemical cycles in an ocean general circulation model. Preindustrial tracer distributions, *Global Biogeochem. Cycles*, 7, 645–677, <https://doi.org/10.1029/93GB01355>, 1993.
- Martin, J. H., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: VERTEX: carbon cycling in the northeast Pacific, *Deep Sea*

- Res. Part A, *Oceanogr. Res. Pap.*, 34, 267–285, [https://doi.org/10.1016/0198-0149\(87\)90086-0](https://doi.org/10.1016/0198-0149(87)90086-0), 1987.
- 2035 McCarthy, G. D., Smeed, D. A., Johns, W. E., Frajka-Williams, E., Moat, B. I., Rayner, D., Baringer, M. O., Meinen, C. S., Collins, J., and Bryden, H. L.: Measuring the Atlantic Meridional Overturning Circulation at 26°N, *Prog. Oceanogr.*, 130, 91–111, <https://doi.org/10.1016/j.pocean.2014.10.006>, 2015.
- Najjar, R. G. and Orr, J. C.: Design of OCMIP-2 simulations of chlorofluorocarbons, the solubility pump and common biogeochemistry, *Intern. OCMIP Rep.*, 25, 1998.
- 2040 Noh, Y., Kang, I. S., Herold, M., and Raasch, S.: Large eddy simulation of particle settling in the ocean mixed layer, *Phys. Fluids*, 18, 85109, <https://doi.org/10.1063/1.2337098>, 2006.
- Oka, A., Hasumi, H., Obata, H., Gamo, T., and Yamanaka, Y.: Study on vertical profiles of rare earth elements by using an ocean general circulation model, *Global Biogeochem. Cycles*, 23, <https://doi.org/10.1029/2008GB003353>, 2009.
- Oka, A., Tazoe, H., and Obata, H.: Simulation of global distribution of rare earth elements in the ocean using an ocean general circulation model, *J. Oceanogr.*, 1, 3, <https://doi.org/10.1007/s10872-021-00600-x>, 2021.
- 2045 Osborne, A. H., Haley, B. A., Hathorne, E. C., Plancherel, Y., and Frank, M.: Rare earth element distribution in Caribbean seawater: Continental inputs versus lateral transport of distinct REE compositions in subsurface water masses, *Mar. Chem.*, 177, 172–183, <https://doi.org/10.1016/j.marchem.2015.03.013>, 2015.
- Osborne, A. H., Hathorne, E. C., Schijf, J., Plancherel, Y., Boning, P., and Frank, M.: The potential of sedimentary foraminiferal rare earth element patterns to trace water masses in the past, *Geochemistry Geophys. Geosystems*, 18, 1–26, <https://doi.org/10.1002/2016GC006782>, 2017.
- 2050 Paffrath, R., Pahnke, K., Böning, P., Rutgers van der Loeff, M., Valk, O., Gdaniec, S., and Planquette, H.: Seawater-Particle Interactions of Rare Earth Elements and Neodymium Isotopes in the Deep Central Arctic Ocean, *J. Geophys. Res. Ocean.*, 126, e2021JC017423, <https://doi.org/10.1029/2021JC017423>, 2021.
- 2055 Palmer, J. R. and Totterdell, I. J.: Production and export in a global ocean ecosystem model, *Deep. Res. Part I Oceanogr. Res. Pap.*, 48, 1169–1198, [https://doi.org/10.1016/S0967-0637\(00\)00080-7](https://doi.org/10.1016/S0967-0637(00)00080-7), 2001.
- Pasquier, B., Hines, S. K. V. V., Liang, H., Wu, Y., John, S. G., Goldstein, S. L., and John, S. G.: GNOM v1.0 : An optimized steady-state model of the modern marine neodymium cycle, *Geosci. Model Dev.*, 147, 4625–4656, <https://doi.org/10.5194/gmd-15-4625-2022>, 2022.
- 2060 Paul, S. A. L., Haeckel, M., Bau, M., Bajracharya, R., and Koschinsky, A.: Small-scale heterogeneity of trace metals including rare earth elements and yttrium in deep-sea sediments and porewaters of the Peru Basin, southeastern equatorial Pacific, 16, 4829–4849, <https://doi.org/10.5194/bg-16-4829-2019>, 2019.
- Pearce, C. R., Jones, M. T., Oelkers, E. H., Pradoux, C., and Jeandel, C.: The effect of particulate dissolution on the neodymium (Nd) isotope and Rare Earth Element (REE) composition of seawater, *Earth Planet. Sci. Lett.*, 369–370, 138–147, <https://doi.org/10.1016/j.epsl.2013.03.023>, 2013.
- 2065 Pope, V. D., Gallani, M. L., Rowntree, P. R., and Stratton, R. A.: The impact of new physical parametrizations in the Hadley Centre climate model: HadAM3, *Clim. Dyn.*, 16, 123–146, <https://doi.org/10.1007/s003820050009>, 2000.
- Pöppelmeier, F., Scheen, J., Blaser, P., Lippold, J., Gutjahr, M., and Stocker, T. F.: Influence of Elevated Nd Fluxes on the Northern Nd Isotope End Member of the Atlantic During the Early Holocene, *Paleoceanogr. Paleoclimatology*, 35, e2020PA003973, <https://doi.org/10.1029/2020PA003973>, 2020a.
- Pöppelmeier, F., Blaser, P., Gutjahr, M., Jaccard, S. L., Frank, M., Max, L., and Lippold, J.: Northern Sourced Water dominated the Atlantic Ocean during the Last Glacial Maximum, *Geology*, 48, <https://doi.org/10.1130/G47628.1>, 2020b.
- Pöppelmeier, F., Janssen, D. J., Jaccard, S. L., and Stocker, T. F.: Modeling the marine chromium cycle: new constraints on global-scale processes, 18, 5447–5463, <https://doi.org/10.5194/bg-18-5447-2021>, 2021a.
- 2075 Pöppelmeier, F., Scheen, J., Jeltsch-Thömmes, A., and F. Stocker, T.: Simulated stability of the Atlantic Meridional

- Overturning Circulation during the Last Glacial Maximum, *Clim. Past*, 17, 615–632, <https://doi.org/10.5194/cp-17-615-2021>, 2021b.
- Pöppelmeier, F., Lippold, J., Blaser, P., Gutjahr, M., Frank, M., and Stocker, T. F.: Neodymium isotopes as a paleo-water mass tracer: A model-data reassessment, *Quat. Sci. Rev.*, 279, 107404, <https://doi.org/10.1016/j.quascirev.2022.107404>, 2022.
- 2080 Rahlf, P., Hathorne, E., Laukert, G., Gutjahr, M., Weldeab, S., and Frank, M.: Tracing water mass mixing and continental inputs in the southeastern Atlantic Ocean with dissolved neodymium isotopes, *Earth Planet. Sci. Lett.*, 530, 115944, <https://doi.org/10.1016/j.epsl.2019.115944>, 2020.
- Rahlf, P., Laukert, G., Hathorne, E. C., Vieira, L. H., and Frank, M.: Dissolved neodymium and hafnium isotopes and rare earth elements in the Congo River Plume: Tracing and quantifying continental inputs into the southeast Atlantic, *Geochim. Cosmochim. Acta*, 294, 192–214, <https://doi.org/10.1016/j.gca.2020.11.017>, 2021.
- 2085 Rempfer, J., Stocker, T. F., Joos, F., Dutay, J.-C., and Siddall, M.: Modelling Nd-isotopes with a coarse resolution ocean circulation model: Sensitivities to model parameters and source/sink distributions, *Geochim. Cosmochim. Acta*, 75, 5927–5950, <https://doi.org/10.1016/j.gca.2011.07.044>, 2011.
- Roberts, N. L., Piotrowski, A. M., McManus, J. F., and Keigwin, L. D.: Synchronous deglacial overturning and water mass source changes, *Science* (80-.), 327, 75–78, <https://doi.org/10.1126/science.1178068>, 2010.
- Robinson, S., Ivanovic, R., van de Flierdt, T., Blanchet, C. L., Tachikawa, K., Martin, E. E., Cook, C. P., Williams, T., Gregoire, L., Plancherel, Y., Jeandel, C., and Arsouze, T.: Global continental and marine detrital ϵNd : An updated compilation for use in understanding marine Nd cycling, *Chem. Geol.*, 567, 120119, <https://doi.org/10.1016/j.chemgeo.2021.120119>, 2021.
- 2095 Rousseau, T. C. C., Sonke, J. E., Chmieleff, J., Van Beek, P., Souhaut, M., Boaventura, G., Seyler, P., and Jeandel, C.: Rapid neodymium release to marine waters from lithogenic sediments in the Amazon estuary, *Nat. Commun.*, 6, <https://doi.org/10.1038/ncomms8592>, 2015.
- Sarmiento, J. L. and LeQuéré, C.: Oceanic carbon dioxide uptake in a model of century-scale global warming, *Science* (80-.), 274, 1346–1350, <https://doi.org/10.1126/science.274.5291.1346>, 1996.
- Sherriff-Tadano, S., Abe-Ouchi, A., and Oka, A.: Impact of mid-glacial ice sheets on deep ocean circulation and global climate, *Clim. Past*, 17, 95–110, <https://doi.org/10.5194/cp-17-95-2021>, 2021.
- 2100 Sholkovitz, E. R.: The geochemistry of rare earth elements in the Amazon River estuary, *Geochim. Cosmochim. Acta*, 57, 2181–2190, [https://doi.org/10.1016/0016-7037\(93\)90559-F](https://doi.org/10.1016/0016-7037(93)90559-F), 1993.
- Siddall, M., Khatriwala, S., van de Flierdt, T., Jones, K., Goldstein, S. L., Hemming, S., and Anderson, R. F.: Towards explaining the Nd paradox using reversible scavenging in an ocean general circulation model, *Earth Planet. Sci. Lett.*, 274, 448–461, <https://doi.org/10.1016/j.epsl.2008.07.044>, 2008.
- 2105 Sinha, B., Smeed, D. A., McCarthy, G., Moat, B. I., Josey, S. A., Hirschi, J. J. M., Frajka-Williams, E., Blaker, A. T., Rayner, D., and Madec, G.: The accuracy of estimates of the overturning circulation from basin-wide mooring arrays, *Prog. Oceanogr.*, 160, 101–123, <https://doi.org/10.1016/j.pcean.2017.12.001>, 2018.
- Smith, R. N. B.: A scheme for predicting layer clouds and their water content in a general circulation model, *Q. J. R. Meteorol. Soc.*, 116, 435–460, <https://doi.org/10.1002/qj.49711649210>, 1990.
- 2110 Smith, R. N. B.: Subsurface, surface and boundary layer processes, Unified Model Doc. Pap. No 24, 1993.
- Smith, R. S.: The FAMOUS climate model (versions XFXWB and XFHC): Description update to version XDBUA, *Geosci. Model Dev.*, 5, 269–276, <https://doi.org/10.5194/gmd-5-269-2012>, 2012.
- Smith, R. S. and Gregory, J. M.: A study of the sensitivity of ocean overturning circulation and climate to freshwater input in different regions of the North Atlantic, *Geophys. Res. Lett.*, 36, L15701, <https://doi.org/10.1029/2009GL038607>, 2009.
- 2115 Smith, R. S., Gregory, J. M., and Osprey, A.: A description of the FAMOUS (version XDBUA) climate model and control run, *Geosci. Model Dev.*, 1, 53–68, <https://doi.org/10.5194/gmd-1-53-2008>, 2008.

- Stichel, T., Kretschmer, S., Geibert, W., Lambelet, M., Plancherel, Y., Rutgers van der Loeff, M., and van de Flierdt, T.: Particle-Seawater Interaction of Neodymium in the North Atlantic, *ACS Earth Sp. Chem.*, <https://doi.org/10.1021/acsearthspacechem.0c00034>, 2020.
- 2120 Tachikawa, K., Jeandel, C., Vangriesheim, A., and Dupré, B.: Distribution of rare earth elements and neodymium isotopes in suspended particles of the tropical Atlantic Ocean (EUMELI site), *Deep. Res. Part 1*, 46, 733–755, [https://doi.org/10.1016/S0967-0637\(98\)00089-2](https://doi.org/10.1016/S0967-0637(98)00089-2), 1999.
- 2125 Tachikawa, K., Athias, V., and Jeandel, C.: Neodymium budget in the modern ocean and paleo-oceanographic implications, *J. Geophys. Res.*, 108, 3254, <https://doi.org/10.1029/1999JC000285>, 2003.
- Tachikawa, K., Arsouze, T., Bayon, G., Bory, A., Colin, C., Dutay, J. C., Frank, N., Giraud, X., Gourlan, A. T., Jeandel, C., Lacan, F., Meynadier, L., Montagna, P., Piotrowski, A. M., Plancherel, Y., Pucéat, E., Roy-Barman, M., and Waelbroeck, C.: The large-scale evolution of neodymium isotopic composition in the global modern and Holocene ocean revealed from seawater and archive data, *Chem. Geol.*, 457, 131–148, <https://doi.org/10.1016/j.chemgeo.2017.03.018>, 2017.
- 2130 Talley, L. D., Pickard, G. L., Emery, W. J., and Swift, J. H.: *Descriptive physical oceanography: An introduction: Sixth edition*, Academic Press, 1–555 pp., <https://doi.org/10.1016/C2009-0-24322-4>, 2011.
- Valdes, P. J., Armstrong, E., Badger, M. P. S., Bradshaw, C. D., Bragg, F., Crucifix, M., Davies-Barnard, T., Day, J., Farnsworth, A., Gordon, C., Hopcroft, P. O., Kennedy, A. T., Lord, N. S., Lunt, D. J., Marzocchi, A., Parry, L. M., Pope, V., Roberts, W. H. G., Stone, E. J., Tourte, G. J. L., and Williams, J. H. T.: The BRIDGE HadCM3 family of climate models: HadCM3@Bristol v1.0, *Geosci. Model Dev.*, 10, 3715–3743, <https://doi.org/10.5194/gmd-10-3715-2017>, 2017.
- 2135 Wang, R., Clegg, J. A., Scott, P. M., Larkin, C. S., Deng, F., Thomas, A. L., Zheng, X. Y., and Piotrowski, A. M.: Reversible scavenging and advection – Resolving the neodymium paradox in the South Atlantic, *Geochim. Cosmochim. Acta*, 314, 121–139, <https://doi.org/10.1016/j.gca.2021.09.015>, 2021.
- 2140 Williams, J. H. T., Smith, R. S., Valdes, P. J., Booth, B. B. B., and Osprey, A.: Optimising the FAMOUS climate model: inclusion of global carbon cycling, *Geosci. Model Dev.*, 6, 141–160, <https://doi.org/10.5194/gmd-6-141-2013>, 2013.
- Wilson, D. J., Piotrowski, A. M., Galy, A., and Clegg, J. A.: Reactivity of neodymium carriers in deep sea sediments: Implications for boundary exchange and paleoceanography, *Geochim. Cosmochim. Acta*, 109, 197–221, <https://doi.org/10.1016/j.gca.2013.01.042>, 2013.
- Woodward, S.: Mineral Dust in HadGEM2, *Hadley Cen*, 8, 2011.
- 2145 Zhang, Y., Lacan, F., and Jeandel, C.: Dissolved rare earth elements tracing lithogenic inputs over the Kerguelen Plateau (Southern Ocean), *Deep. Res. Part II Top. Stud. Oceanogr.*, 55, 638–652, <https://doi.org/10.1016/j.dsr2.2007.12.029>, 2008.
- Zweng, M. ., Reagan, J. R., Seidov, D., Boyer, T. P., Locarnini, R. A., Garcia, H. E., Mishonov, A. ., Baranova, . O.K., Weathers, K. W., Paver, C. R., and Smolyar, I. V.: *World Ocean Atlas 2018*, <http://www.node.noaa.gov/>, 2019.

2150

Page 24: [1] Deleted	Suzanne Robinson [ee14s2r]	11/11/2022 07:35:00
Page 27: [2] Deleted	Suzanne Robinson [ee14s2r]	19/11/2022 08:41:00
Page 29: [3] Deleted	Suzanne Robinson [ee14s2r]	19/11/2022 08:45:00
Page 39: [4] Deleted	Suzanne Robinson [ee14s2r]	19/11/2022 08:48:00
Page 48: [5] Deleted	Suzanne Robinson [ee14s2r]	23/11/2022 18:22:00