

Local moisture recycling across the globe

Jolanda J.E. Theeuwes^{1,2}, Arie Staal¹, Obbe A. Tuinenburg¹, Bert V.M. Hamelers^{2,3}, Stefan C. Dekker¹

¹Copernicus Institute of Sustainable Development, Utrecht University, Utrecht, 3584 CB, The Netherlands

²Wetsus, European Centre of Excellence for Sustainable Water Technology, Leeuwarden, 8911 MA, The Netherlands

³Department of Environmental Technology, Wageningen University and Research, Wageningen, 6708 PB, The Netherlands

Correspondence to: Jolanda J.E. Theeuwes (j.j.e.theeuwes@uu.nl)

Abstract. Changes in evaporation over land affect terrestrial precipitation via atmospheric moisture recycling and consequently freshwater availability. Although global moisture recycling at regional and continental scales are relatively well understood, the patterns of local moisture recycling and the main variables that impact it remain unknown. [For the first time,](#) we calculate the local moisture recycling ratio (LMR) as the fraction of evaporated moisture that precipitates within a distance of 0.5° (typically 50 km) from its source, identify variables that correlate with it over land globally and study its model dependency. We derive seasonal and annual LMR using a 10-year climatology (2008–2017) of monthly averaged atmospheric moisture connections at a scale of 0.5° obtained from a Lagrangian atmospheric moisture tracking model. We find that, annually, on average 1.7% (st.dev. = 1.1%) of evaporated moisture returns as precipitation locally, but with large temporal and spatial variability, where LMR peaks in summer and over wet and mountainous regions. Our results show that wetness, orography, latitude, convective available potential energy, wind speed, and total cloud cover correlate clearly with LMR, indicating that especially wet regions with little wind and strong ascending air are favourable for high LMR. Finally, we find that spatial patterns of local recycling are consistent between different models, yet the magnitude of recycling varies. Our results can be used to study impacts of evaporation changes on local precipitation, with implications for, for example, greening and water management.

1 Introduction

Atmospheric moisture connections redistribute water from evaporation sources to precipitation sinks, affecting climates globally, regionally, and locally. These connections are key in the global hydrological cycle and are used to understand the importance of terrestrial evaporation for water availability. As evaporated moisture can travel up to thousands of kilometres in the atmosphere, changes in evaporation can affect precipitation in a large area. An evaporation shed (Van der Ent and Savenije, 2013) describes where evaporated moisture from a specific source region precipitates and therefore, can be used to study (1) the changes in precipitation on a global scale following a change in evaporation in the source region and (2) atmospheric moisture recycling. Globally, more than half of terrestrial evaporated moisture precipitates over land (Van der Ent et al., 2010; Tuinenburg et al., 2020), which is a process called terrestrial moisture recycling. About half of terrestrial precipitation originates from land (Tuinenburg et al., 2020). Hence, terrestrial moisture recycling has an important contribution

31 to water availability. For example, 80% of China's water resources originates from evaporation over Eurasia (Van der Ent et
32 al., 2010). Furthermore, areas can also feed precipitation to themselves through regional moisture recycling. In the Amazon
33 basin, 63% of the evaporated moisture precipitates within the basin itself (Tuinenburg et al., 2020). Terrestrial moisture
34 recycling is considered an ecosystem service (Falkenmark et al., 2019; Keys et al., 2016) as globally, almost 20% of terrestrial
35 precipitation originates from vegetation-regulated moisture recycling (Keys et al., 2016). How this ecosystem service is
36 affected by, for instance, deforestation, can be studied using atmospheric moisture connections.

37

38 Moisture recycling has been used to study downwind impacts of land-use changes (e.g. Bagley et al., 2012; Keys et al., 2012;
39 Wang-Erlandsson et al., 2018), which can affect both the magnitude and pattern of moisture recycling (Van der Ent et al.,
40 2014), and the impact of ecosystems on other ecosystems (e.g. O'Connor et al., 2021). Hence, atmospheric moisture
41 connections can be used for freshwater governance to understand and manage the impacts of land-use changes downwind such
42 as changes in freshwater availability for irrigation and plants. (te Wierik et al, 2021; Te Wierik et al, 2020). For example,
43 previous research showed that for 45% of the land surface, an increase in vegetation is beneficial for downwind water
44 availability (Cui et al., 2022).

45

46 So far, analytical recycling models and moisture tracking models have been used to study terrestrial recycling and downwind
47 impacts of land cover change on global and regional levels (Burde & Zangvil, 2001; Van der Ent et al., 2010). Multiple studies
48 focus on the regional recycling for specific regions, with a spatial scale ranging from 500 km up to several thousands of
49 kilometres (e.g., Burde, 2006; Dominguez et al., 2006; Lettau et al, 1979; Staal et al., 2018; Trenberth, 1999). Furthermore,
50 regional recycling on a spatial scale of 1.5° has been studied globally using a Eulerian moisture tracking model, assuming a
51 well-mixed atmosphere (Van der Ent and Savenije, 2011). It was debated that regional recycling ratios are difficult to compare
52 due to differences in the shape and size of the studied regions (Van der Ent and Savenije, 2011). Therefore, Van der Ent &
53 Savenije (2011) defined the typical length scale of evaporation recycling, which can be used to compare between different
54 regions because it is independent of the size and shape of a regions. This length scale decreases with increasing regional
55 recycling and, therefore, is a proxy for an area's regional recycling. However, it does not allow for the quantification of the
56 amount of water that recycles within the defined region and therefore does not provide quantitative insight into the regional
57 impacts of evaporation changes induced by land-cover changes.

58

59 In regions with a high regional recycling, reforestation can enhance freshwater availability and for regions with a low recycling,
60 reforestation may cause local drying (Hoek van Dijke et al., 2022) due to reductions in streamflow as a result of enhanced
61 evaporation locally (Brown et al., 2005; Jackson et al., 2005). To physically understand, for instance, the role of local wetting
62 or drying due to reforestation, deforestation, or the use of groundwater or surface water for irrigation, local moisture recycling
63 is key. We argue that local impacts need to be studied explicitly as they may have a crucial role in future water governance,
64 e.g., to prevent tree restoration projects causing local drying.

65

66 The state-of-the-art high-resolution atmospheric moisture connections obtained with the Lagrangian atmospheric moisture
67 tracking model “UTrack” allows us to calculate the evaporation recycling ratio at higher spatial resolution (0.5°) (Tuinenburg
68 et al., 2020; Tuinenburg and Staal, 2020). We define this as the local moisture recycling ratio (LMR) as this high resolution
69 allows us to study local-scale land-atmosphere feedbacks, which will help us better understand hydrological impacts of land-
70 use change. LMR describes which fraction of evaporated moisture recycles within its source grid cell and its eight surrounding
71 grid cells. Moisture recycling has not been studied before on this high-resolution scale globally. To get a better physical
72 understanding of this metric we identify which factors correlate with it. We analyse this for different latitude classes to account
73 for different cell sizes across latitude. Factors included in this analysis are: orography, precipitation, precipitation type,
74 evaporation, shear, convective available potential energy, atmospheric moisture flux, wind speed, total cloud cover, boundary
75 layer height and surface net solar radiation. These variables relate to either convection, local wetness, or moisture transport
76 away from the source location, which we identified as important factors for local moisture recycling. Furthermore, we study
77 how LMR varies over the globe and throughout the year for a 10-year climatology (2008-2017), as well as its scaling and
78 model dependency.

79 **2 Methods**

80 We use global atmospheric moisture connections obtained from Tuinenburg et al., (2020) to calculate LMR worldwide. These
81 moisture connections are a 10-year climatology (2008–2017) of monthly averages and have a spatial resolution of 0.5° . These
82 UTrack-atmospheric-moisture data are derived using a Lagrangian atmospheric moisture tracking model by Tuinenburg &
83 Staal (2020) that tracks evaporated moisture at a spatial scale of 0.25° , [and stored at a spatial resolution of \$0.5^\circ\$](#) . In this model,
84 for each grid cell of 0.25° , each mm of evaporation is represented by one hundred released moisture parcels. The wind
85 transports these parcels horizontally and vertically through the atmosphere. Additionally, a probabilistic scheme describes the
86 vertical movement of the moisture parcels over 25 atmospheric layers. In this scheme, the parcels are randomly distributed
87 across the vertical moisture profile of each grid cell. At each time step (0.1 h), the moisture budget is made using evaporation,
88 precipitation and total precipitable water. Parcels are tracked for up to 30 days or up to the point at which only 1% of their
89 original moisture is still present. On average, the lifetime of atmospheric moisture is 8-10 days (Sodemann, 2020). However,
90 some moisture might still remain in the parcels after 10 days. After 30 days for most of the parcels all of the original moisture
91 has rained out (Tuinenburg and Staal, 2020). Input data for UTrack consist of evaporation, precipitation, precipitable water,
92 and wind speed obtained from the ERA5 dataset (Hersbach et al., 2020). We refer to Tuinenburg & Staal (2020) for a complete
93 description of the model settings and the tests and assumptions underlying them.

94

95 LMR is the fraction of evaporated moisture that precipitates locally. To study the scale dependency of local moisture recycling,
96 we examine three definitions of LMR (Fig. A1): the fraction of evaporated moisture that precipitates in $f(1)$ its source grid

cell, i.e., r_l , (2) its source grid cell and its eight neighbouring grid cells, i.e., r_9 , and (3) its source grid cell and its 24 neighbouring grid cells, i.e., r_{25} . Equations 1-3 describe the three definitions of LMR, in which $E_{i,j}$ is the amount of moisture evaporated from source grid cell i,j . The fraction of $E_{i,j}$ that precipitates within its source grid cell and its (8 or 24) neighbouring grid is indicated by $P_{E,i+l,j+k}$ ($i+l,j+k$, with $l = 0$ and $k = 0$ for r_l , $l = -1,0,1$ and $k = -1,0,1$ for r_9 and $l = -2,-1,0,1,2$ and $k = -2,-1,0,1,2$ for r_{25}).

$$r_1 = \frac{P_{E,i,j}}{E_{i,j}} \quad (1)$$

$$r_9 = \frac{\sum_{l=-1}^1 \sum_{k=-1}^1 P_{E,i+l,j+k}}{E_{i,j}} \quad (2)$$

$$r_{25} = \frac{\sum_{l=-2}^2 \sum_{k=-2}^2 P_{E,i+l,j+k}}{E_{i,j}} \quad (3)$$

r_l , r_9 , and r_{25} result in different local moisture recycling ratios across the globe (Fig. A2). r_l peaks over the ocean where precipitation is relatively low and evaporation is relatively large, which results in relatively large recycling ratios. In addition, we find exceptionally low values over mountain peaks, yet not over all elevated terrain. This result is inconsistent with the patterns found for r_9 and r_{25} , as these patterns include peaks over mountainous and low recycling over the oceans. These patterns can be explained by enhanced convection over mountains due to orographic lift and strong winds over the ocean that carry moisture away from its source. The patterns found for r_9 and r_{25} seem to capture multiple physical processes that are important for moisture transport and formation of precipitation better than the pattern of r_l . In our study we do not focus on r_l , as r_l does not include all small-scale flows of <50 km. This is because moisture can evaporate from cell i,j , and precipitate in the adjacent cell, while transport length is <50 km. Furthermore, as the patterns of r_9 and r_{25} are similar and agree with our understanding of relevant processes, we decided to define the local moisture recycling ratio (LMR) as r_9 to keep the spatial scale as small as possible. For r_9 , the distance ~~between from~~ the center of the source grid cell and its surrounding grid cells describes the typical length of the local moisture flow, which is approximately 0.5°. We calculated this typical length across the globe by calculating the average of the average zonal length, meridional length, and diagonal length of all terrestrial grid cells. The total average equals 50.1 km (st.dev. = 15.5 km), so, the average moisture flow length is approximately 50 km.

Furthermore, the LMR derived with the Lagrangian approach using output from UTrack is compared with the output from the Eulerian moisture tracking model WAM2-layers (Link et al., 2020), to study the model dependency of LMR. For this comparison, the resolution of the UTrack data is reduced to 1.5° to match the output of the WAM2-layers model. To do so, all evaporationsheds over land were multiplied with their source evaporation. Then, the recycling within cells of 1.5° was calculated for all terrestrial surfaces. A detailed description of the atmospheric moisture connections obtained with WAM2-layers and the model itself are provided by Link et al. (2020) and Van der Ent et al. (2013).

129 We study the relations between multiple variables and the 10-year climatology (2008-2017) of local moisture recycling to
 130 identify factors that [affect-relate to recycling to assess what factors might affect recycling](#). To calculate this 10-year climatology
 131 of LMR, for each month, we weighted the multi-year (2008-2017) monthly LMR by multi-year monthly evaporation in the
 132 same period:

$$133 \quad LMR_{annual\ average} = \sum_{i=jan}^{dec} LMR \frac{E_{month\ i}}{E_{year}}$$

134 (4)

135 in which E_{year} is the sum of the evaporation of the 12 months. To identify factors that [might](#) affect LMR, variables that relate
 136 to atmospheric moisture and vertical displacement of air, as both higher atmospheric moisture content and ascending air
 137 promote precipitation are selected. All these variables are obtained, either directly or indirectly from ERA5 reanalysis data
 138 (Hersbach et al., 2020). We downscaled the original resolution from 0.25° to 0.5° by centrally averaging the data. [For all](#)
 139 [variables we calculated the climatological mean for the years 2008-2017](#).

140

141 In total 13 variables are selected (Fig. A3): (1) elevation (z) which we expect to enhance LMR through orographic lift. (2)
 142 Precipitation, which we expect to correlate positively with LMR given that in Lagrangian moisture tracking models, the amount
 143 of moisture that leaves the parcel (i.e., precipitates) scales with precipitation. (3) Total evaporation as it enhances the
 144 atmospheric moisture content and we, therefore, expect it to promote precipitation locally. (4) Wetness (precipitation minus
 145 evaporation), as with increasing wetness the downward flux of moisture increases and evaporated water becomes more likely
 146 to precipitate, possibly promoting LMR. (4) Convective precipitation and (5) large-scale precipitation, as they scale with
 147 precipitation, by definition. Both are included to study whether the type of precipitation is an important factor explaining LMR.
 148 (6) Latitude, which is a proxy for processes related to the Hadley cell circulation, which is characterized by strong ascent and
 149 descent of air at specific latitudes, which we expect to have an important contribution to LMR, because they respectively
 150 enhance and reduce the formation of precipitation (Wang and Yang, 2022). (7) The vertical integral of the atmospheric
 151 moisture flux (in northward and eastward directions and the total flux) as it carries the moisture away from its source and could
 152 thus reduce LMR. (8) Convective available potential energy ($CAPE$), which feeds convection and therefore promotes
 153 precipitation locally, which could enhance LMR. (9) Vertical wind shear between 650 and 750 hPa of both meridional and
 154 zonal winds, as it affects moisture transport in multiple directions and, therefore, we expect it to impact LMR. (10) Total wind
 155 speed, as it carries the wind, and therefore, we expect it to correlate negatively with LMR. (11) Total cloud cover as a proxy
 156 for condensation processes which possibly enhance LMR (Richards and Arkin, 1998). (12) Boundary layer height, because
 157 thinner boundaries need less evaporation to reach saturation of air, and therefore, we expect it will promote precipitation
 158 locally. Finally, (13) net surface solar radiation [as a proxy for the energy source of convection, and other processes, which we](#)
 159 [expect to be important for LMR](#). We calculate shear (τ) using Equation (5).


$$160 \quad \tau = \frac{\ln \frac{v_2}{v_1}}{\ln \frac{z_2}{z_1}} \quad (5)$$

161 In this equation, v_1 and v_2 are the wind speed (in zonal and meridional directions) at two different heights (z_1 and z_2). We
162 identified significant correlations using Spearman rank correlations. It should be noted that a correlation does not imply
163 causality. We exclude oceans, seas and Antarctica from this analysis using the land-sea mask from ERA5. We classify the data
164 based on latitude to account for decreasing grid cell size with increasing latitude. Each class has a range of 15° and includes
165 the grid cells on both the Northern and Southern Hemispheres (see Table A1). Between 60° and 90° south, the grid cells do
166 not contain land besides Antarctica, and are therefore not included in the classes. Additionally, we used the Ecoregions 2017
167 data (<https://ecoregions.appspot.com/>) to study the spatially averaged local moisture recycling of 14 biomes across the globe
168 (Fig. A4). We study variation amongst biomes, as biomes include information on both biotic factors such as vegetation type,
169 and abiotic factors such as climate.

170 **3 Results**

171 **3.1 LMR obtained from output of UTrack**

172 Annually, on average about 1.7% (st. dev. = 1.1%) of terrestrial evaporated moisture recycles locally. LMR shows spatio-
173 temporal variation (Fig. 1) with peaks over elevated (e.g., the Atlas Mountains and Ethiopian Highlands) and wet areas (e.g.,
174 Congo Basin and Southeast Asia) and minima over arid regions (e.g., Australia and the Sahara Desert). Additionally, we find
175 peaks in LMR during summer (i.e., during DJF for the Southern Hemisphere and during JJA for the Northern Hemisphere).
176 This seasonality is especially strong over mountainous and wet areas. For the mid-latitudes, especially the Mediterranean Basin
177 shows seasonality with peaks in summer (JJA). However, seasonality is largest at low latitudes. Within the tropics we find
178 some spatial differences. First, LMR in the Congo Basin and Southeast Asia exceed LMR in the Amazon Basin. Second,
179 recycling in the Congo Basin and Southeast Asia peaks in JJA and recycling in the Amazon Basin peaks in DJF, which is the
180 wet season for a large part of the Amazon.



181
 182 **Figure 1. 10-year climatology (2008–2017) of the seasonal averages of local moisture recycling across the global land surface. Here,**
 183 **local moisture recycling is defined as the fraction of evaporated moisture that precipitates in its source grid cell and its eight**
 184 **neighbouring grid cells (r_9). Different seasons are DJF: December–February, MAM: March–May, JJA: June–August, and SON:**
 185 **September–November.**

186 We calculated recycling on a 1.5° grid using both the dataset by Link et al. (2020), which we refer to as $r_{WAM2-layers}$, and the
 187 dataset by Tuinenburg et al. (2020) (upscaled to 1.5°), which we refer to as r_{UTrack} , to study the model dependency of local
 188 recycling (Fig. 2). We find that the global spatial patterns of r_{UTrack} and $r_{WAM2-layers}$ agree (Fig. 2 & Fig. A5). However, the
 189 magnitude of $r_{WAM2-Layers}$ is larger than r_{UTrack} over mountains, the tropics, and the high latitudes. r_{UTrack} is larger than r_{WAM2-}
 190 $layers$ over drylands and deserts (e.g., the Sahel region and Western Asia) (Fig. 2). However, over drylands and deserts recycling
 191 ratios are relatively small and therefore, the relative difference as presented in Fig. 2 has less meaning here. Globally, the
 192 difference between r_{UTrack} and $r_{WAM2-layers}$ and its variation is largest around the equator (Fig. A6). On average, the relative
 193 difference between UTrack and WAM2-Layers ($(UTrack - WAM2-Layers) / UTrack$) equals -1.5 (st.dev. = 3.4).




Figure 2. r_{Utrack} , the fraction of evaporated moisture that recycles within its source grid cell of 1.5° (top). The relative deviation between r_{Utrack} and $r_{WAM2-layers}$ (bottom). This deviation is calculated using the recycling within one grid cell at a resolution of 1.5° obtained from the datasets of Tuinenburg et al. (2020) and Link et al. (2020).

3.2 Factors underlying LMR

For each latitude class we calculated the Spearman rank correlation coefficient (ρ) (Table 1). Below we discuss only statistically significant ($p < 0.05$) correlations with $\rho \geq 0.4$, that indicate a moderate correlation. These correlations are emboldened in Table 1. We find that LMR correlates positively with total precipitation (\mathbf{P}) and wetness ($\mathbf{P-E}$) for all classes between 15° and 75° . In addition, between 15° and 30° , LMR correlates strongly with $\mathbf{P_{total}}$ ($\rho = 0.80$). Furthermore, Besides total precipitation, large scale precipitation (between 15° and 45° and between 60° and 75°) and convective precipitation (between 15° and 45°) correlate positively with LMR. Overall, there is a moderate correlation between precipitation and LMR for the mid-latitudes. The highest correlation between LMR and convective precipitation is found between 15° and 30° latitude. For this latitude class, Here LMR also correlates positively with evaporation and CAPE, which enhances convective precipitation. Despite the low correlation between LMR and CAPE, for most of the latitude classes, high CAPE clearly relates to LMR, as the skewed profile in the scatter density plot indicates that only a small amount of the grid cells with a relatively high CAPE have a low LMR (Fig 3). Furthermore, the presence of clouds also correlates with LMR. Between LMR and total cloud cover, a positive correlation holds between 15° and 45° , and a negative correlation holds

211 between 60° and 75°. ~~Furthermore,~~ the vertical integrals of the eastward and northward moisture fluxes correlate less with
 212 LMR compared to ~~the~~ vertical fluxes (e.g., precipitation). ~~as for~~ For the higher latitudes, the northward moisture flux correlates
 213 positively with LMR (between 60° and 75°) and the eastward moisture flux correlates negatively with LMR (between 75° and
 214 90°). ~~The moisture flux depends on wind speed.~~ ~~However,~~ yet, wind speed correlates negatively with LMR for the lower
 215 latitudes (between 0° and 45°). Furthermore, LMR correlates positively with orography between 30° and 75°. We find that for
 216 high elevation, LMR is always relatively high (Fig A7). Additionally, LMR correlates negatively with boundary layer height
 217 between 45° and 60°. Finally, LMR correlates negatively with ~~wind~~ shear at 650 hpa in the meridional direction (between 75°
 218 and 90°) and ~~with~~ latitude (between 60° and 75°). However, we find an oscillating relation between LMR and latitude (Fig
 219 4), which is not captured by the Spearman rank correlation coefficients. This pattern indicates high ~~values of~~ LMR over the
 220 equator (0°) and 60° north, and low ~~values~~ LMR around 30° north and south. Orography seems to disrupt the relation between
 221 latitude and LMR causing peaks in LMR around 35° north and 20° south (Fig 4). LMR does not correlate to surface net solar
 222 radiation for any latitude. However, for low surface net solar radiation ($<0.75 \times 10^6 \text{ J/m}^2$) holds that LMR increases with
 223 increasing surface net solar radiation (Fig 3).





Figure 3: Scatter plots of the 10-year climatology (2008–2017) of the local moisture recycling ratio over land and precipitation (top left), evaporation (top right), convective available potential energy (CAPE) (bottom left), and solar net surface radiation (bottom right). Each dot represents a 0.5° resolution grid cell over land.



228

229 **Figure 4. Scatter plot of the 10-year climatology (2008-2017) of LMR and latitude. The colour scale indicates elevation, with blue**
 230 **being low elevation and yellow being high elevation. The black line represents the zonal average of LMR. Each dot represents a**
 231 **0.5° resolution grid cell over land.**

232

233 **Table 1. Spearman rank correlation coefficients between LMR and all tested variables. “*” indicates a significant correlation ($p < 0.05$)**
 234 **and moderate and strong relations ($\rho > 0.4$) are emboldened. The classes including latitudes between 0° and 60° include grid cells of**
 235 **the Northern Hemisphere and Southern Hemisphere. The classes including latitudes exceeding 60° include grid cells of the Northern**
 236 **Hemisphere only.**

Variable	Spearman rank correlation coefficient					
	$0^\circ\text{-}15^\circ$	$15^\circ\text{-}30^\circ$	$30^\circ\text{-}45^\circ$	$45^\circ\text{-}60^\circ$	$60^\circ\text{-}75^\circ$	$75^\circ\text{-}90^\circ$
Total precipitation (P)	0.15*	0.80*	0.47*	0.40*	0.45*	0.37*
Total evaporation (E)_	-0.05*	0.63*	0.19*	-0.12*	0.19*	0.20*
Wetness (P-E)	0.18*	0.59*	0.52*	0.48*	0.43*	0.27*
Convective precipitation	0.20*	0.79*	0.46*	0.29*	0.35*	0.33*
Large scale precipitation	-0.06*	0.75*	0.46*	0.38*	0.40*	0.36*
Fraction of cp	0.36*	-0.35*	-0.13*	-0.14*	0.19*	0.28*
Latitude	0.24*	-0.18*	0.22*	0.14*	-0.40*	-0.18*
Eastward moisture flux	0.15*	0.00	-0.30*	-0.38*	-0.20*	-0.49*
Northward moisture flux	-0.03*	0.22*	0.29*	-0.03*	0.48*	0.23*
Total moisture flux	-0.28*	0.30*	-0.29*	-0.33*	-0.03	-0.16*
CAPE	0.31*	0.58*	0.37*	0.06*	0.12*	-0.02
Zonal shear	0.15*	-0.12*	0.02	-0.31*	0.00	0.24*
Meridional shear	-0.22*	0.15*	-0.08*	-0.01	0.05*	-0.46*
Orography	0.31*	0.29*	0.49*	0.54*	0.68*	-0.13*
Total cloud cover	0.28*	0.78*	0.43*	0.09*	-0.56*	0.08*
Surface net solar radiation	-0.16*	0.10*	-0.30*	-0.08*	0.28*	0.21*
Boundary layer height	-0.31*	-0.32*	-0.39*	-0.53*	-0.18*	-0.06*
Total wind speed	-0.46*	-0.55*	-0.47*	-0.26*	-0.26*	-0.30*

237 **4 Discussion**

238 **4.1 Factors underlying LMR**

239 Moisture recycling affects humanity by influencing water security, agriculture, forestry, regional climate stability and Earth
 240 system resilience (Keys et al., 2019; Wang-Erlandsson et al., 2022). Different types of moisture recycling were subject to
 241 research used for different applications (e.g., Bagley et al., 2012; Pranindita et al., 2022; Van der Ent et al., 2010), ~~but for the~~
 242 ~~first time, we~~ We analysed the local moisture recycling ratio (LMR) (of evaporated moisture) across the globe at 0.5° resolution,
 243 and which factors ~~might~~ affect it. We find that LMR, defined as the fraction of evaporated moisture that precipitates within a

distance of 0.5° (typically 50 km) from its source, varies over time and space, peaking in summer and over elevated and wet regions. First, ~~we identified~~ latitude, elevation, and Convective Available Potential Energy (CAPE) ~~seem to be~~ important factors influencing LMR (Fig. 5). These variables all promote convection (Roe, 2005; Scheff and Frierson, 2012; Wallace and Hobbs, 2006), strongly suggesting a dependency of LMR on convection. Convective storms develop due to unstable conditions resulting in precipitation locally (Eltahir, 1998) and a higher CAPE results in more rainfall (Eltahir and Pal, 1996; Williams and Renno, 1993). The pattern of LMR across latitudes also coincides with updraft and downdraft of air caused by the Hadley cell circulation (Wallace and Hobbs, 2006). Around the equator and 60° north and south, air ascends, where we find a high LMR. Around 30° north and south, air descends, where we find a low LMR. Deviations from this pattern correspond to higher elevations which promote LMR through orographic lift. Overall, our results suggest a positive relation between convection and LMR.




Figure 5. Conceptual model of the most important factors influencing local moisture recycling around the globe. Rainy clouds indicate variables that increase LMR and clouds without raindrops indicate variables that decrease LMR. Blue indicates wet regions, yellow indicates arid regions.

Second, ~~we find that~~ wetness ~~is~~ ~~seems to be~~ an important factor underlying LMR as LMR significantly correlates with precipitation and $P-E$ (precipitation minus evaporation). Furthermore, both large-scale and convective precipitation significantly correlate with LMR. This is surprising, as convection promotes precipitation locally (Eltahir, 1998); therefore, we expected a stronger correlation between LMR and convective precipitation than between LMR and large-scale precipitation. As both correlations are similar, this suggests that the type of precipitation ~~does-might~~ not affect LMR. Although convection is a local-scale process (i.e., having a spatial scale below 100 km) (Miyamoto et al., 2013), remotely evaporated

264 moisture can be transported to a region with high convective activity and then precipitate as convective precipitation (Jana et
265 al., 2018; Liberato et al., 2012). In that way, the precipitation type is independent of the distance between moisture source and
266 target location and therefore does not relate to LMR. Total cloud cover correlates both positively (between 15° and 45°) and
267 negatively (between 60° and 75°) with LMR. Total cloud cover correlates with precipitation, convective precipitation, and
268 large-scale precipitation for all latitudes except between 60° and 75° (Tab. A2). Due to the positive correlation between LMR
269 and precipitation and the absence of a correlation between precipitation and total cloud cover at these latitudes we can
270 statistically explain the negative correlation between total cloud cover and LMR. Physically, this result is harder to explain.
271 Our results describe the importance of convection underlying LMR at lower latitudes, where total cloud cover correlates with
272 convective precipitation. For higher latitudes, the importance of convection underlying LMR decreases, and we therefore
273 expected also the correlation between total cloud cover and LMR to decrease but not to become negative. Likely, another
274 process that we cannot identify with our analysis causes the correlation between total cloud cover and LMR to be negative.
275 Overall, we find that wetness enhances LMR independent of the precipitation type.

276

277 Unexpectedly, we do not find a clear correlation between the vertical integral of the atmospheric moisture flux and LMR.
278 However, for the lower latitudes (between 0° and 45° latitude), LMR correlates to wind speed (at 10 and 100 m) which carries
279 evaporated moisture away from its source location, enhancing the moisture flux. Therefore, horizontal moisture fluxes at
280 specific altitudes are better for our analysis than the vertical integral of the moisture flux. However, since wind carries moisture
281 away from its source, we expected that wind speed and LMR would also correlate for the higher latitudes (latitude above 45°).
282 It could be that for the higher latitudes, a more significant amount of moisture is present at higher latitudes, explaining why
283 LMR and wind at 10m do not correlate. However, wind speeds at 650 hPa and 750 hPa also do not correlate to LMR for these
284 latitudes (Tab. A2).

285

286 Despite the importance of vertical shear in atmospheric moisture tracking models (Van der Ent et al., 2013), we do not find a
287 correlation between local moisture recycling and vertical shear between 650 and 750 hPa. Shear is the friction between air
288 layers that minimizes complete mixing, which for some regions around the world is strongest between 650 and 750 hPa
289 (Dominguez et al., 2016). A possible explanation is that due to its small spatial scale, the temporal scale of LMR is also small,
290 which may prevent the air reaching 700 hPa within the spatial scale of LMR. Furthermore, it is possible that our study design
291 is insufficient to capture the relation between LMR and shear throughout the year over the globe. We aimed for a general
292 analysis to identify the main factors that might influence LMR. A more detailed study that distinguishes between different
293 seasons and isolates different climate zones is necessary to identify more factors that influence LMR as some factors might be
294 more important during a specific season. For example, convection occurs more during summer than during winter, and
295 therefore, might have a stronger correlation with LMR during summer. Besides, some factors are shape and size dependent
296 similar to LMR, while other factors are not dependent on grid cell size and shape. This might cause bias in the results of the
297 Spearman analysis. Furthermore, due to the many interactions within the Earth system and, consequently, between the variables

298 included in our study, it is impossible to determine the true drivers of LMR. However, the correlations do indicate how changes
299 in the environment might affect LMR.

300 **4.2 regional patterns**

301 To zoom in on the importance of each of the different factors underlying LMR for various areas across the globe, we determined
302 LMR for the major global biomes (Fig. A8). LMR is highest for the wet tropics (between 0° and 15° north and south) and
303 montane grasslands and lowest for desert-like biomes in both the Northern and Southern Hemisphere (between 30° and 45°
304 north and south), confirming the importance of wetness, orography, and latitude. However, in the tropics (between 0° and 15°
305 latitude), we do not find any correlation between LMR and precipitation, evaporation, wetness, or orography. Possibly, due to
306 the abundance of water and energy to evaporate, there is LMR under all circumstances, except for when the wind speed is
307 high. Comparing LMR for each biome between both hemispheres indicates that some of the factors underlying LMR are more
308 robust than other ones for some biomes. In the Mediterranean biomes, located between 30–40° north and south, air generally
309 descends due to the Hadley cell circulation. As a result, these biomes are expected to have low LMR. Although we find a low
310 LMR for the Mediterranean biomes in the Southern Hemisphere, we find a relatively high LMR for the Mediterranean biomes
311 in the Northern Hemisphere. The Spearman rank analysis indicates that at these latitudes, wind speed correlates with LMR,
312 which may explain the difference between both hemispheres.

313
314 Although LMR is the highest in the wet tropics, we find different results among the various tropical regions (Amazon Basin,
315 Congo Basin & Southeast Asia). LMR in the Congo Basin exceeds LMR in the Amazon Basin (Fig. 1), despite larger amounts
316 of precipitation in the Amazon Basin (Hersbach et al., 2020). In the tropics, current deforestation results in drying (Bagley et
317 al., 2014; Staal et al., 2020), reducing evaporation. As LMR in the Congo Basin exceeds LMR in the Amazon Basin,
318 deforestation has a relatively large impact on local precipitation in the Congo Basin, suggesting a larger impact on droughts
319 locally. This is further exacerbated by the fact that the Congo Basin, in comparison with the Amazon Basin, has many small-
320 scale moisture feedback loops (Wunderling et al., 2022). Unlike LMR, basin recycling is similar for both basins (Tuinenburg
321 et al., 2020). Suggesting, the impact of deforestation on precipitation in the entire basin is similar for both basins, indicating
322 both basins would experience similar overall drying. However, drought conditions can also enhance recycling ratios (Bagley
323 et al., 2014), possibly promoting LMR. Further research is necessary to understand the impact of deforestation on LMR in the
324 tropics in more detail.

325 **4.3 The spatial scale of the local moisture recycling ratio**

326 We study local moisture recycling on a spatial scale of 0.5°, which is approximately 55 km around the equator and 50 km on
327 average globally for all land cells. Instead of recycling within one grid cell (r_1), we studied the recycling of evaporated moisture
328 within its source grid cell and its 8 surrounding grid cells. Compared to r_1 , this r_9 includes all moisture flows with a length

scale of typically 50 km (0.5°). For r_1 , moisture flows with a length smaller than 50 km can occur close to the border of grid cells and therefore, r_1 by definition underestimates the actual recycling. These moisture flows are accounted for in r_9 .

However, defining LMR on a grid scale gives complications. First, the longitudinal distance for a grid cell size decreases with latitude, resulting in different sizes and shapes, which makes it difficult to compare LMR among all grid cells. For the low- and mid-latitudes, the variation in grid cell size affects LMR only slightly, as confirmed when LMR for each grid cell was scaled to a single area (Fig. A9). Therefore, we believe that the variation in grid size causes only a small bias in the statistical analysis, as the largest fraction of the land surface is at the low- and mid-latitudes, and moisture recycling is less important for the higher latitudes. However, it should be noted that for similar wind speed, LMR will be lower in smaller grid cells than larger grid cells. Second, the spatial scale of recycling is strongly dependent on regional differences such as biome type, the dominating winds, and the proximity to mountains. For instance, with increasing distance to the Andes mountains the median travelling distance of transpired moisture from the Amazon forest increases (Staal et al., 2018) and for the Ganges basin, evaporated moisture is blocked by the Himalayas, limiting upward moisture flow and inducing precipitation (Tuinenburg et al., 2012). Further, precipitation can be triggered by micrometeorological processes (e.g. Knox et al., 2011; Taylor et al., 2012) making it unknown at what spatial scale moisture recycling is the dominant process for precipitation. Therefore, we believe that a grid-based approach to systematically study LMR globally is a solid approach to define and study the physical processes at a spatial scale >50 km through, for instance, the Spearman analysis to study the underlying processes. However, our definition of LMR is not sufficient to identify processes on a spatial scale smaller than 50 km that might be relevant.

4.4 Model and definition dependencies

It is important to note that the typical length scale of moisture recycling, as defined by Van der Ent & Savenije (2011), allows for a comparison of regional moisture recycling for different regions around the world due to its independence of the region's size and shape (Fig A10). The typical length scale of evaporated moisture recycling decreases with increasing recycling. It peaks over deserts and is small over the tropics and mountainous regions (Fig A9), overlapping with the spatial pattern of LMR. However, this typical length scale does not allow for the quantification of the amount of recycled moisture and therefore, it is difficult to apply this metric to study the impact of evaporation changes due to land-use change. Therefore, studies that aim to quantify moisture recycling locally may best use recycling ratios. However, studies that aim to compare recycling among different regions can best use the typical length scale of recycling.

In this article, we focus on model dependency as we calculated the differences in magnitude of recycling within one grid cell of 1.5° obtained from output of the UTrack and WAM2-layers models (Link et al., 2020; Tuinenburg et al., 2020). The spatial patterns are similar, yet the different magnitudes indicate a large model dependency, and, therefore, an uncertainty in moisture recycling. Furthermore, Van der Ent et al. (2010) calculated recycling within a grid cell of 1.5° for the years 1999–2008 using WAM2-layers and found a similar spatial pattern with high recycling over mountainous and tropical regions and low recycling

362 over desert-like regions. These recycling ratios also have a larger magnitude than LMR. However, it is not straightforward to
363 interpret the differences in recycling ratios as both models use different input data (i.e., ERA5 and ERA-Interim). To assess
364 the possible role of the models in causing the difference in moisture recycling, we describe the main differences between the
365 models. First, WAM2-layers calculates the atmospheric moisture recycling on a larger temporal and spatial scale than UTrack,
366 A larger grid cell size and time step increases the likelihood of evaporation and precipitation taking place within the same
367 small amount of time, which might result in an overestimation of recycling within one grid cell. Second, WAM2-layers
368 generates moisture flows using two vertical layers; therefore, strong winds at specific vertical levels will be described in less
369 detail, reducing estimated moisture transport and enhancing estimated moisture recycling within a single grid cell. Differences
370 between r_{UTrack} and $r_{WAM2-layers}$ are highly visible over mountainous regions where wind experiences relatively strong friction,
371 highly impacting the wind. Finally, different approaches are used to include vertical mixing in the two models. Vertical mixing
372 causes the greatest error in moisture tracking models, but it is unknown to what extent vertical mixing is underestimated (Stohl
373 et al., 2005; Tuinenburg & Staal, 2020).

374

375 Besides studies using atmospheric moisture tracking (e.g., Bagley et al., 2014; Keys et al., 2014; Van der Ent et al., 2010),
376 some previous studies used different methods to calculate regional moisture recycling for a specific area, such as isotope
377 measurements (e.g., An et al., 2017) and bulk recycling models (e.g., Burde & Zangvil, 2001). The most common recycling
378 models are modifications of Budyko's model (Budyko, 1974; Burde and Zangvil, 2001), which are 1D or 2D analytical models.
379 These models assume that the atmosphere is completely mixed, meaning that evaporated water directly mixes perfectly with
380 advected water throughout the entire water column. Because of this assumption, first, these models overlook fast recycling,
381 which describes local showers that yield precipitation before the evaporated water is fully mixed. Excluding fast recycling
382 causes models to underestimate terrestrial moisture recycling for some regions (e.g., Amazon Basin) (Burde et al., 2006b).
383 Second, these models ignore the influence of vertical shear, which causes a significant error (Dominguez et al., 2020). Our
384 method minimises the errors due to fast recycling and vertical shear because of two model aspects. First, at each time step,
385 each parcel has a small chance of getting mixed, causing each parcel to move approximately once in the vertical direction
386 every 24 hours, additional to the displacement based on reanalysis data of vertical winds. This process minimizes complete
387 mixing and reduces the error due to shear and fast recycling. Second, the error due to fast recycling also becomes smaller
388 because lower atmospheric levels contribute more to the total precipitation than higher levels due to the skewed vertical
389 moisture profile. WAM2-layers accounts for vertical shear as it models two vertical atmospheric layers of which the interface
390 is located at the height at which shear typically occurs. These two layers are both completely mixed and therefore, compared
391 to bulk models, WAM2-layers better represents the distribution of moisture throughout the atmospheric column. As an
392 alternative method, moisture flows can be calculated on a smaller time step to increase the interactions between different wind
393 components, resulting in a better representation of turbulence (Keune et al., 2022). Despite the error reduction, the
394 representation of fast recycling in UTrack should be studied in more detail, as fast recycling is expected to influence LMR
395 significantly.

LMR is calculated as a ten-year average. This period of ten years might miss multi-year climate variability such as the El Niño Southern Oscillation and the North Atlantic Oscillation. The time series of atmospheric moisture connections provided by Link et al. (2020) allowed to study inter-annual variation in relatively local recycling. This shows that recycling is dependent on multi-year atmospheric phenomena. During the major El Niño event of 2015-2016, the northeast of South Africa had a lower-than-average local recycling ratio (Fig. A11) for 2015. This pattern coincides with the impact of wetness during El Niño years, consistent with the hypothesis that wetness enhances LMR. Furthermore, strong events such as heat waves and droughts might affect the multi-year annual mean. For example, we clearly find lower recycling over Russia during 2010, which may relate to the 2010 heatwave in eastern Europe and Russia. Overall, for these multi-year and strong events we find that, for regions that face wetter-than-normal conditions, LMR is enhanced, and for regions that face drier-than-normal conditions, LMR is reduced. Hence, drought events might result in a decrease in LMR as seen for the 2010 heat wave event in Europe and Russia. However, not for all inter-annual climate variability modes we find a clear impact on moisture recycling. It may be that these phenomena do not affect wetness throughout the entire year, and therefore, annual means might not represent them well.

4.5 Implications/applications of LMR

LMR could be applied in the field of water management. The spatial pattern of LMR shows some overlap with global agricultural water management (Molden, 2007; Salmon et al., 2015). Generally, the tropics have a high LMR and agriculture is mainly rainfed (Salmon et al., 2015; Costa et al., 2019), indicating that these agricultural regions are self-dependent to some extent regarding precipitation. Also, agriculture in the Mediterranean Basin and South Australia is mainly rainfed. For semi-arid regions that dependent on rainfed agriculture, changes in precipitation may have a significant impact (Keys et al., 2016). LMR in the Mediterranean basin exceeds LMR in southern Australia, indicating that a larger fraction of evaporated moisture returns locally. Thus, when evaporation is maintained in the Mediterranean Basin, part of the precipitation will sustain here, which holds to a lesser extent for southern Australia. Besides LMR (i.e., local evaporation recycling), local precipitation recycling can help to understand the precipitation dependence on local evaporation for each region. Irrigated agriculture is important in India and China (Salmon et al., 2015; Döll and Siebert, 2002), which are regions with a relatively low LMR, indicating that only a small amount of the evaporated moisture returns as precipitation locally. For irrigated agriculture in regions that are characterized by a high LMR, a relatively large amount of the evaporated water returns to its source, which reduces the amount of water that is necessary for irrigation. Terrestrial evaporation is an important source for precipitation and freshwater availability (Keune and Miralles, 2019). Therefore, spatial planning using LMR might improve agricultural water management.


Global climate change likely affects atmospheric moisture connections due to changes in atmospheric dynamics. For example, due to global warming, tropical atmospheric circulation may weaken (Vecchi et al., 2006), and the Hadley cells may move poleward (Shaw, 2019), which will affect the updraft and downdraft of air around the globe, which we found to be important processes underlying LMR. Furthermore, climate change has different opposing impacts on storm tracks which have an

important role in moisture transport by transporting latent heat poleward (Shaw et al., 2016). Furthermore, in a warmer climate continental recycling is predicted to decrease and precipitation over land would be more dependent on evaporation over the ocean (Findell et al., 2019). However, our study does not account for any impacts of climate change. As our results indicate that wetness and convection enhance LMR, LMR may change due to, for example, drying and wetting of regions, changes in Hadley cell circulation, and circulation in the tropics. Furthermore, climate change enhances the risk of droughts (Rasmijn et al., 2018; Teuling, 2018) and LMR might be used to study drought resilience globally. As for a high LMR a local drought might drastically impact the local water cycle.


We expect that LMR can be helpful also in other ways. Specifically, we expect the concept of LMR can be used to study how changes in evaporation, due to for example afforestation, affect the local water cycle beyond merely a loss of moisture. However, besides evaporation, land-use changes also influence the energy balance and other factors that might alter the atmospheric moisture connections and thus, LMR. Using future land use scenarios as input for moisture tracking models, it will be possible to study the impact of land-use changes on atmospheric moisture connections. However, future scenarios often include other changes besides land use, which makes it possible to study the changes of land use specifically. However, LMR can help us better predict the impact of land cover changes on the local water cycle. It might help us identify regions where reforestation would not cause local drying due to enhanced evaporation (Hoek van Dijke et al., 2022; Tuinenburg et al., 2022). Overall, LMR gives us better insight into the atmospheric part of the local water cycle and terrestrial evaporation as a source for local freshwater availability.

5 Conclusions

We calculated the local moisture recycling ratio (LMR) from atmospheric moisture connections at a spatial scale of 0.5° . LMR is the fraction of evaporated moisture that precipitates within a distance of 0.5° (typically 50 km) from its source. On average, 1.7% (st.dev. = 1.1%) of global terrestrial evaporation returns as precipitation locally, with peaks of approximately 6%. LMR peaks in summer and in wet and elevated regions. We find [a correlation between LMR and topography, precipitation, wetness, convective available potential energy, and wind](#) ~~that~~ [suggesting these variables might affect LMR](#). In addition, latitude correlates with LMR, which likely indicates the importance of the ascending air and descending air related to the Hadley cell circulation. Furthermore, by comparing LMR calculated using different models we found that the spatial pattern of LMR is not model-dependent, yet, the magnitude of LMR is strongly dependent on the model. LMR defines the local impacts of enhanced evaporation on precipitation and thus its role as a source for local freshwater availability. Therefore, LMR can be used to evaluate which locations may be suitable for greening without largely disrupting the local water cycle.



459
460 **Figure A1.** Three definitions of the local moisture recycling ratio (LMR) from left to right: r_1 describes the fraction of evaporated
461 moisture that returns as precipitation in its source grid cell, r_9 describes the fraction of evaporated moisture that returns as
462 precipitation in its source grid cell and 8 neighbouring grid cells, and r_{25} describes the fraction of evaporated moisture that returns
463 as precipitation in its source grid cell and 24 neighbouring grid cells. LMR is calculated on a spatial scale of 0.5° and the first three
464 plots do not have a similar resolution. The plot on the right shows LMR on a spatial scale of 0.5° which is the resolution at which we
465 calculate all definitions (r_1 , r_9 and r_{25}).




466

467 **Figure A2.** 10-year climatology (2008–2017) of the three definitions of the local moisture recycling ratio (LMR). The top panel
 468 indicates the fraction of evaporated moisture that precipitates within its source grid cell (r_1), the middle panel shows the fraction of
 469 evaporated moisture that precipitates within its source grid cell and its 8 neighbouring grid cells (r_9), and the lower panel shows the
 470 fraction of evaporated moisture that precipitates within its source grid cell and its 24 neighbouring grid cells (r_{25}).


471 **Table A1:** Defined classes for spearman rank correlation analysis.

<i>Class</i>	Latitude ranges
1	-15°:15°
2	-30°:-15° and 15°:30°
3	-45°:-30° and 30°:45°
4	-60°:-45° and 45°:60°
5	60°:75°
6	75°:90°



473 **Figure A3.** Global 10-year climatology (2008–2017) of (from top to bottom and left to right) precipitation, evaporation, precipitation
 474 – evaporation, convective precipitation, large-scale precipitation, fraction of convective precipitation, vertical integral of moisture
 475 flux in eastward direction, vertical integral of moisture flux in northward direction, CAPE, orography, vertical shear (between 650
 476 and 750 hPa) of zonal wind, and vertical shear (between 650 and 750 hPa) of meridional wind.

477



478

479 **Figure A4.** Major global biomes Ecoregions 2017 (<https://ecoregions.appspot.com/>)_©NASA Terra Metrics, @Google INEGI
 480 [Imagery](#)

481




Figure A5. The 10-year climatology (2008-2017) of the recycling within one grid cell of 1.5° calculated with the dataset by Link et al. (2020), i.e., the output from the Eulerian moisture tracking model WAM2-layers (top) and the difference with the 10-year climatology (2008-2017) of the recycling within one grid cell of 1.5° calculated with the dataset by Tuinenburg et al. (2020).





Figure A6. The zonal mean of the absolute difference (left) and relative difference (right) between r_{UTrack} and $r_{WAM2-layers}$ (calculated as r_{UTrack} minus $r_{WAM2-layers}$, indicated by the blue line) and its standard deviation (blue area).




490

491 **Figure A7. Scatter plots of the 10-year climatology (2008–2017) of the local moisture recycling ratio and (from top to bottom and**
 492 **left to right) precipitation – evaporation, convective precipitation, large-scale precipitation, fraction of convective precipitation,**
 493 **vertical integral of moisture flux in eastward direction, vertical integral of moisture flux in northward direction, orography, vertical**
 494 **shear (between 650 and 750 hPa) of zonal wind, vertical shear (between 650 and 750 hPa) of meridional wind, boundary layer height,**
 495 **total cloud cover, and wind speed. Each scatter represents one grid cell.**

496 Table A2. Spearman rank correlation coefficients for additional variables at different latitude classes. “**” indicates a significant
 497 correlation ($p<0.05$) and moderate and strong relations ($\rho>0.4$) are emboldened. The classes including latitudes between 0° and 60°
 498 include grid cells of the Northern Hemisphere and Southern Hemisphere. The classes including latitudes exceeding 60° include grid
 499 cells of the Northern Hemisphere only.


Variables	Spearman rank correlation coefficient					
	$0^\circ\text{-}15^\circ$	$15^\circ\text{-}30^\circ$	$30^\circ\text{-}45^\circ$	$45^\circ\text{-}60^\circ$	$60^\circ\text{-}75^\circ$	$75^\circ\text{-}90^\circ$
Total cloud cover and wind speed	-0.58	-0.41	-0.23	0.08	0.16	-0.51
Large-scale precipitation and wind speed	-0.30	-0.46	-0.37	0.06	0.11	-0.28
Convective precipitation and wind speed	-0.63	-0.50	-0.33	-0.13	-0.41	-0.61
Total cloud cover and precipitation	0.85	0.92	0.76	0.58	-0.08	0.46
Total cloud cover and convective precipitation	0.85	0.90	0.63	0.23	-0.09	0.67
Total cloud cover and large-scale precipitation	0.71	0.90	0.81	0.70	-0.02	0.43
LMR and wind speed at 650 hpa	0.26	-0.18	-0.37	-0.16	-0.15	-0.27
LMR and wind speed at 750 hpa	-0.09	0.023	-0.39	-0.19	-0.09	-0.31

500



501


502 Figure A8. Time series of the local moisture recycling ratio for global biomes on the Northern (left) and Southern (right)
 503 Hemispheres. The plots show the 10-year climatology (2008–2017).



504

505 **Figure A9:** The local moisture recycling ratio scaled to a grid cell size of 50 km x 50 km. The plot shows the 10-year climatology


506 (2008-2017). We divided the original local moisture recycling ratio by the area of the grid cell and multiplied it with 2500 km²



507

508 **Figure A10:** Evaporation recycling length scale as defined by Van der Ent and Savenije (2011) for each grid cell of 0.5°x0.5°. The

509 plot shows the average of 2008-2017.



510
 511 **Figure A11. Inter-annual variation of recycling within a single grid cell of 1.5° between 2001-2018. Each plot shows the**
 512 **difference between annual averaged recycling and the climatological mean of recycling. Data obtained from Link et al.**
 513 **(2020).**

514 **Code availability**

515 ~~The code that was used to calculate the local moisture recycling ratio and for the analyses is available from the corresponding~~
516 ~~author upon reasonable request.~~The code that was used to calculate the local moisture recycling ratio and to plot the local
517 moisture recycling ratio is provided on GitHub (<https://github.com/jtheeu/LocalMoistureRecycling>, last access 28 February
518 [2023](#)).

519 **Data availability**

520 [The local moisture recycling ratios are available from the Zenodo archive at 0.5 and 1.5 degrees resolution](https://doi.org/10.5281/zenodo.7684640)
521 [\(https://doi.org/10.5281/zenodo.7684640\)](https://doi.org/10.5281/zenodo.7684640).
522 The atmospheric moisture connections from Tuinenburg et al., (2020) are available from the PANGAEA archive at 0.5 and
523 1.0 degrees resolution (<https://doi.pangaea.de/10.1594/PANGAEA.912710>).
524 The atmospheric moisture connections from Link et al., (2020) are available from the PANGAEA archive at 1.5 degrees
525 resolution (<https://doi.pangaea.de/10.1594/PANGAEA.908705>).

526 **Author contributions**

527 JT designed the study with contributions from all authors. JT carried out the research. JT wrote the first draft of the manuscript
528 in close collaboration with AS. All authors contributed to the discussion and the final version of the manuscript.

529 **Acknowledgements**

530 [We want to thank Patrick Keys, Ruud van der Ent and the anonymous reviewer for commenting on earlier versions of this](#)
531 [manuscript](#). This work was performed in the cooperation framework of Wetsus, European Centre of Excellence for Sustainable
532 Water Technology (www.wetsus.eu). Wetsus is co-funded by the Dutch Ministry of Economic Affairs and Climate Policy, the
533 Northern Netherlands Provinces the Province of Fryslân. The authors would like to thank the participants of the natural water
534 production theme for the fruitful discussions and financial support. AS acknowledges support from the Talent Programme
535 grant VI.Veni.202.170 by the Dutch Research Council (NWO). OT acknowledges support from the research programme
536 Innovational Research Incentives Scheme Veni (016.veni.171.019), funded by the Dutch Research Council.

537 **References**

538 An, W., Hou, S., Zhang, Q., Zhang, W., Wu, S., Xu, H., Pang, H., Wang, Y., and Liu, Y.: Enhanced recent local moisture
539 recycling on the northwestern Tibetan Plateau deduced from ice core deuterium excess records, J. Geophys. Res.-Atmos., 122,

12541–12556, <https://doi.org/10.1002/2017JD027235>, 2017.

Bagley, J. E., Desai, A. R., Dirmeyer, P. A., and Foley, J. A.: Effects of land cover change on moisture availability and potential crop yield in the worlds breadbaskets, *Environ. Res. Lett.*, 7, 014009, <https://doi.org/10.1088/1748-9326/7/1/014009>, 2012.

Bagley, J. E., Desai, A. R., Harding, K. J., Snyder, P. K., and Foley, J. A.: Drought and deforestation: Has land cover change influenced recent precipitation extremes in the Amazon?, *J. Clim.*, 27, 345–361, <https://doi.org/10.1175/JCLI-D-12-00369.1>, 2014.

Brown, A. E., Zhang, L., McMahon, T. A., Western, A. W., and Vertessy, R. A.: A review of paired catchment studies for determining changes in water yield resulting from alterations in vegetation, *J. Hydrol.*, 310, 28–61, <https://doi.org/10.1016/j.jhydrol.2004.12.010>, 2005.

Budyko, M. I.: *Climate and life*, Academic Press: New York, 1974.

Burde, G. I.: Bulk recycling models with incomplete vertical mixing. Part II: Precipitation recycling in the Amazon Basin, *J. Clim.*, 19, 1461–1472, <https://doi.org/10.1175/JCLI3687.1>, 2006.

Burde, G. I. and Zangvil, A.: The estimation of regional precipitation recycling. Part I: Review of recycling models, *J. Clim.*, 14, 2509–2527, <https://doi.org/10.1175/1520-0442>, 2001.

Costa, M. H., Fleck, L. C., Cohn, A. S., Abrahão, G. M., Brando, P. M., Coe, M. T., Fu, R., Lawrence, D., Pires, G. F., Pousa, R., and Soares-Filho, B. S.: Climate risks to Amazon agriculture suggest a rationale to conserve local ecosystems, *Front. Ecol. Environ.*, 17, 584–590, <https://doi.org/10.1002/fee.2124>, 2019.

Cui, J., Lian, X., Huntingford, C., Gimeno, L., Wang, T., Ding, J., He, M., Xu, H., Chen, A., Gentine, P., and Piao, S.: Global water availability boosted by vegetation-driven changes in atmospheric moisture transport, *Nat. Geosci.*, 15, 982–988, <https://doi.org/10.1038/s41561-022-01061-7>, 2022.

Döll, P. and Siebert, S.: Global modeling of irrigation water requirements, *Water Resour. Res.*, 38, 1037, <https://doi.org/10.1029/2001wr000355>, 2002.

Dominguez, F., Kumar, P., Liang, X. Z., and Ting, M.: Impact of atmospheric moisture storage on precipitation recycling, *J. Clim.*, 19, 1513–1530, <https://doi.org/10.1175/JCLI3691.1>, 2006.

Dominguez, F., Miguez-Macho, G., and Hu, H.: WRF with water vapor tracers: A study of moisture sources for the North American Monsoon, *J. Hydrometeorol.*, 17, 1915–1927, <https://doi.org/10.1175/JHM-D-15-0221.1>, 2016.

Dominguez, F., Hu, H., and Martinez, J. A.: Two-Layer Dynamic Recycling Model (2L-DRM): Learning from moisture tracking models of different complexity, *J. Hydrometeorol.*, 21, 3–16, <https://doi.org/10.1175/jhm-d-19-0101.1>, 2020.

Eltahir, E. A. B.: A soil moisture-rainfall feedback mechanism, *Water Resour. Res.*, 34, 765–776, 1998.

Eltahir, E. A. B. and Pal, J. S.: Relationship between surface conditions and subsequent rainfall in convective storms, *J. Geophys. Res.-Atmos.*, 101, 26237–26245, <https://doi.org/10.1029/96jd01380>, 1996.

Van der Ent, R. J. and Savenije, H. H. G.: Length and time scales of atmospheric moisture recycling, *Atmos. Chem. Phys.*, 11, 1853–1863, <https://doi.org/10.5194/acp-11-1853-2011>, 2011.

Van der Ent, R. J. and Savenije, H. H. G.: Oceanic sources of continental precipitation and the correlation with sea surface

574 temperature, *Water Resour. Res.*, 49, 3993–4004, <https://doi.org/10.1002/wrcr.20296>, 2013.

575 Van der Ent, R. J., Savenije, H. H. G., Schaefli, B., and Steele-Dunne, S. C.: Origin and fate of atmospheric moisture over
576 continents, *Water Resour. Res.*, 46, W09525, <https://doi.org/10.1029/2010WR009127>, 2010.

577 Van der Ent, R. J., Tuinenburg, O. A., Knoche, H. R., Kunstmann, H., and Savenije, H. H. G.: Should we use a simple or
578 complex model for moisture recycling and atmospheric moisture tracking?, *Hydrol. Earth Syst. Sc.*, 17, 4869–4884,
579 <https://doi.org/10.5194/hess-17-4869-2013>, 2013.

580 Van der Ent, R. J., Wang-Erlandsson, L., Keys, P. W., and Savenije, H. H. G.: Contrasting roles of interception and
581 transpiration in the hydrological cycle - Part 2: Moisture recycling, *Earth Syst. Dynam.*, 5, 471–489,
582 <https://doi.org/10.5194/esd-5-471-2014>, 2014.

583 Falkenmark, M., Wang-Erlandsson, L., and Rockström, J.: Understanding of water resilience in the Anthropocene, *J. Hydrol.*,
584 2, 100009, <https://doi.org/10.1016/j.hydroa.2018.100009>, 2019.

585 Findell, K. L., Keys, P. W., van der Ent, R. J., Lintner, B. R., Berg, A., and Krasting, J. P.: Rising temperatures increase
586 importance of oceanic evaporation as a source for continental precipitation, *J. Clim.*, 32, 7713–7726,
587 <https://doi.org/10.1175/jcli-d-19-0145.1>, 2019.

588 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers,
589 D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara,
590 G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L.,
591 Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P.,
592 Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–
593 2049, <https://doi.org/10.1002/qj.3803>, 2020.

594 Hoek van Dijke, A. J., Herold, M., Mallick, K., Benedict, I., Machwitz, M., Schlerf, M., Pranindita, A., Theeuwens, J. J. E.,
595 Bastin, J.-F., and Teuling, A. J.: Shifts in regional water availability due to global tree restoration, *Nat. Geosci.*, 15, 363–368,
596 <https://doi.org/10.1038/s41561-022-00935-0>, 2022.

597 Jackson, R. B., Jobbágy, E. G., Avissar, R., Roy, S. B., Barrett, D. J., Cook, C. W., Farley, K. A., Le Maitre, D. C., McCarl,
598 B. A., and Murray, B. C.: Atmospheric science: Trading water for carbon with biological carbon sequestration, *Science* (80-.
599), 310, 1944–1947, <https://doi.org/10.1126/science.1119282>, 2005.

600 Jana, S., Rajagopalan, B., Alexander, M. A., and Ray, A. J.: Understanding the dominant sources and tracks of moisture for
601 summer rainfall in the southwest united states, *J. Geophys. Res.-Atmos.*, 123, 4850–4870,
602 <https://doi.org/10.1029/2017JD027652>, 2018.

603 Keune, J. and Miralles, D. G.: A precipitation recycling network to assess freshwater vulnerability: Challenging the watershed
604 convention, *Water Resour. Res.*, 55, 9947–9961, <https://doi.org/10.1029/2019WR025310>, 2019.

605 Keune, J., Schumacher, D. L., and Miralles, D. G.: A unified framework to estimate the origins of atmospheric moisture and
606 heat using Lagrangian models, *Geosci. Model Dev.*, 15, 1875–1898, <https://doi.org/10.5194/gmd-15-1875-2022>, 2022.

607 Keys, P. W., Van der Ent, R. J., Gordon, L. J., Hoff, H., Nikoli, R., and Savenije, H. H. G.: Analyzing precipitationsheds to

understand the vulnerability of rainfall dependent regions, *Biogeosciences*, 9, 733–746, [https://doi.org/10.5194/bg-9-733-](https://doi.org/10.5194/bg-9-733-2012)
2012, 2012.

Keys, P. W., Barnes, E. A., van der Ent, R. J., and Gordon, L. J.: Variability of moisture recycling using a precipitationshed
framework, *Hydrol. Earth Syst. Sc.*, 18, 3937–3950, <https://doi.org/10.5194/hess-18-3937-2014>, 2014.

Keys, P. W., Wang-Erlandsson, L., and Gordon, L. J.: Revealing invisible Water: Moisture recycling as an ecosystem service,
PLoS One, 11, e0151993, <https://doi.org/10.1371/journal.pone.0151993>, 2016.

Keys, P. W., Porkka, M., Wang-Erlandsson, L., Fetzer, I., Gleeson, T., and Gordon, L. J.: Invisible water security: Moisture
recycling and water resilience, *Water Secur.*, 8, 57–89, <https://doi.org/10.1016/j.wasec.2019.100046>, 2019.

Knox, R., Bisht, G., Wang, J., and Bras, R.: Precipitation variability over the forest-to-nonforest transition in Southwestern
Amazonia, *J. Clim.*, 24, 2368–2377, <https://doi.org/10.1175/2010JCLI3815.1>, 2011.

Lettau, H., K., L., and Molion, L. C. B.: Amazonia’s hydrologic cycle and the role of atmospheric recycling in assessing
deforestation effects., *Mon. Weather Rev.*, 107, 227–238, 1979.

Liberato, M. L. R., Ramos, A. M., Trigo, R. M., Trigo, I. F., Durán-Quesada, A. M., Nieto, R., and Gimeno, L.: Moisture
sources and large-scale dynamics associated with a flash flood event, *Geoph. Monog. Ser.*, 200, 111–126,
<https://doi.org/10.1029/2012GM001244>, 2012.

Link, A., Van Der Ent, R., Berger, M., Eisner, S., and Finkbeiner, M.: The fate of land evaporation - A global dataset, *Earth
Syst. Sci. Data*, 12, 1897–1912, <https://doi.org/10.5194/essd-12-1897-2020>, 2020.

Molden, D.: *A Comprehensive Assessment of Water Management in Agriculture; Summary*, 40 pp., 2007.

O’Connor, J. C., Dekker, S. C., Staal, A., Tuinenburg, O. A., Rebel, K. T., and Santos, M. J.: Forests buffer against variations
in precipitation, *Glob. Chang. Biol.*, 27, 4686–4696, <https://doi.org/10.1111/gcb.15763>, 2021.

Pranindita, A., Wang-Erlandsson, L., Fetzer, I., and Teuling, A. J.: Moisture recycling and the potential role of forests as
moisture source during European heatwaves, *Clim. Dynam.*, 58, 609–624, <https://doi.org/10.1007/s00382-021-05921-7>, 2022.

Rasmijn, L. M., Van Der Schrier, G., Bintanja, R., Barkmeijer, J., Sterl, A., and Hazeleger, W.: Future equivalent of 2010
Russian heatwave intensified by weakening soil moisture constraints, *Nat. Clim. Chang.*, 8, 381–385,
<https://doi.org/10.1038/s41558-018-0114-0>, 2018.

Richards, F. and Arkin, P.: On the Relationship between Satellite-Observed Cloud Cover and Precipitation, *Mon. Weather
Rev.*, 109, 1081–1093, 1998.

Roe, G. H.: Orographic precipitation, *Annu. Rev. Earth. Pl. Sc.*, 33, 645–671,
<https://doi.org/10.1146/annurev.earth.33.092203.122541>, 2005.

Salmon, J. M., Friedl, M. A., Froking, S., Wissler, D., and Douglas, E. M.: Global rain-fed, irrigated, and paddy croplands: A
new high resolution map derived from remote sensing, crop inventories and climate data, *Int. J. Appl. Earth. Obs.*, 38, 321–
334, <https://doi.org/10.1016/j.jag.2015.01.014>, 2015.

Scheff, J. and Frierson, D.: Twenty-First-Century multimodel subtropical precipitation declines are mostly midlatitude shifts,
J. Clim., 25, 4330–4347, <https://doi.org/10.1175/JCLI-D-11-00393.1>, 2012.

642 Shaw, T. A.: Mechanisms of future predicted changes in the zonal mean mid-latitude circulation, *Curr. Clim. Chang. Reports*,
643 5, 345–357, <https://doi.org/10.1007/s40641-019-00145-8>, 2019.

644 Shaw, T. A., Baldwin, M., Barnes, E. A., Caballero, R., Garfinkel, C. I., Hwang, Y. T., Li, C., O’Gorman, P. A., Rivière, G.,
645 Simpson, I. R., and Voigt, A.: Storm track processes and the opposing influences of climate change, *Nat. Geosci.*, 9, 656–664,
646 <https://doi.org/10.1038/ngeo2783>, 2016.

647 Sodemann, H.: Beyond turnover time: Constraining the lifetime distribution of water vapor from simple and complex
648 approaches, *J. Atmos. Sci.*, 77, 413–433, <https://doi.org/10.1175/JAS-D-18-0336.1>, 2020.

649 Staal, A., Tuinenburg, O. A., Bosmans, J. H. C., Holmgren, M., Van Nes, E. H., Scheffer, M., Zemp, D. C., and Dekker, S. C.:
650 Forest-rainfall cascades buffer against drought across the Amazon, *Nat. Clim. Chang.*, 8, 539–543,
651 <https://doi.org/10.1038/s41558-018-0177-y>, 2018.

652 Staal, A., Flores, B. M., Aguiar, A. P. D., Bosmans, J. H. C., Fetzer, I., and Tuinenburg, O. A.: Feedback between drought and
653 deforestation in the Amazon, *Environ. Res. Lett.*, 15, 044024, <https://doi.org/10.1088/1748-9326/ab738e>, 2020.

654 Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model
655 FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, <https://doi.org/10.5194/acp-5-2461-2005>, 2005.

656 Taylor, C. M., de Jeu, R. A. M., Guichard, F., Harris, P. P., and Dorigo, W. A.: Afternoon rain more likely over drier soils,
657 *Nature*, 489, 423–426, <https://doi.org/10.1038/nature11377>, 2012.

658 Teuling, A. J.: A hot future for European droughts, *Nat. Clim. Chang.*, 8, 364–365, [https://doi.org/10.1038/s41558-018-0154-](https://doi.org/10.1038/s41558-018-0154-5)
659 5, 2018.

660 Trenberth, K. E.: Atmospheric moisture recycling: Role of advection and local evaporation, *J. Clim.*, 12, 1368–1381,
661 [https://doi.org/10.1175/1520-0442\(1999\)012<1368:amrroa>2.0.co;2](https://doi.org/10.1175/1520-0442(1999)012<1368:amrroa>2.0.co;2), 1999.

662 Tuinenburg, O. A. and Staal, A.: Tracking the global flows of atmospheric moisture and associated uncertainties, *Hydrol. Earth*
663 *Syst. Sc.*, 24, 2419–2435, <https://doi.org/10.5194/hess-24-2419-2020>, 2020.

664 Tuinenburg, O. A., Hutjes, R. W. A., and Kabat, P.: The fate of evaporated water from the Ganges basin, *J. Geophys. Res.-*
665 *Atmos.*, 117, 1–17, <https://doi.org/10.1029/2011JD016221>, 2012.

666 Tuinenburg, O. A., Theeuwes, J. J. E., and Staal, A.: High-resolution global atmospheric moisture connections from
667 evaporation to precipitation, *Earth Syst. Sci. Data*, 12, 3177–3188, <https://doi.org/10.5194/essd-12-3177-2020>, 2020.

668 Tuinenburg, O. A., Bosmans, J. H. C., and Staal, A.: The global potential of forest restoration for drought mitigation, *Environ.*
669 *Res. Lett.*, 17, 034045, <https://doi.org/10.1088/1748-9326/ac55b8>, 2022.

670 Vecchi, G. A., Soden, B. J., Wittenberg, A. T., Held, I. M., Leetmaa, A., and Harrison, M. J.: Weakening of tropical Pacific
671 atmospheric circulation due to anthropogenic forcing, *Nature*, 441, 73–76, <https://doi.org/10.1038/nature04744>, 2006.

672 Wallace, J. M. and Hobbs, P. V.: *Atmospheric science: an introductory survey*, 2nd ed., Elsevier, 2006.

673 Wang-Erlandsson, L., Fetzer, I., Keys, P. W., van der Ent, R. J., Savenije, H. H. G., and Gordon, L. J.: Remote land use impacts
674 on river flows through atmospheric teleconnections, *Hydrol. Earth Syst. Sc.*, 22, 4311–4328, [https://doi.org/10.5194/hess-22-](https://doi.org/10.5194/hess-22-4311-2018)
675 4311-2018, 2018.

676 Wang-Erlandsson, L., Tobian, A., van der Ent, R. J., Fetzer, I., te Wierik, S., Porkka, M., Staal, A., Jaramillo, F., Dahlmann,
 677 H., Singh, C., Greve, P., Gerten, D., Keys, P. W., Gleeson, T., Cornell, S. E., Steffen, W., Bai, X., and Rockström, J.: A
 678 planetary boundary for green water, *Nat. Rev. Earth Environ.*, 3, 380–392, <https://doi.org/10.1038/s43017-022-00287-8>, 2022.
 679 Wang, C. and Yang, K.: Changes in the moisture contribution over global arid regions, *Clim. Dynam.*, 1–15,
 680 <https://doi.org/10.1007/s00382-022-06600-x>, 2022.
 681 te Wierik, S. A., Cammeraat, E. L. H., Gupta, J., and Artzy-Randrup, Y. A.: Reviewing the impact of land use and land-use
 682 change on moisture recycling and precipitation patterns, *Water Resour. Res.*, 57, e2020WR029234,
 683 <https://doi.org/10.1029/2020WR029234>, 2021.
 684 Te Wierik, S. A., Gupta, J., Cammeraat, E. L. H., and Artzy-Randrup, Y. A.: The need for green and atmospheric water
 685 governance, *Wiley Interdiscip. Rev. Water*, 7, e1406, <https://doi.org/10.1002/wat2.1406>, 2020.
 686 Williams, E. and Renno, N.: An analysis of the conditional instability of the tropical atmosphere, *Mon. Weather Rev.*, 121,
 687 21–35, 1993.
 688 Wunderling, N., Wolf, F., Tuinenburg, O. A., and Staal, A.: Network motifs shape distinct functioning of Earth’s moisture
 689 recycling hubs, *Nat. Commun.*, 13, 6574, <https://doi.org/10.1038/s41467-022-34229-1>, 2022.
 690