

Towards an ensemble-based evaluation of land surface models in light of uncertain forcings and observations

Vivek. K. Arora¹, Christian Seiler², Libo Wang², and Sian Kou-Giesbrecht¹

¹Canadian Centre for Climate Modelling and Analysis, Climate Research Division, Environment Canada,
Victoria, BC, Canada

²Climate Processes Section, Climate Research Division, Environment and Climate Change Canada, Toronto, ON,
Canada

³
⁴ *Correspondence to:* Vivek K. Arora (vivek.arora@ec.gc.ca)

Abstract

Quantification of uncertainty in fluxes of energy, water, and CO₂ simulated by land surface models (LSMs) remains a challenge. LSMs are typically driven with, and tuned for, a specified meteorological forcing data set and a specified set of geophysical fields. Here, using two data sets each for meteorological forcing and land cover representation (in which the increase in crop area over the historical period is implemented in the same way), as well as two model structures (with and without coupling of carbon and nitrogen cycles), the uncertainty in simulated results over the historical period is quantified for the Canadian Land Surface Scheme Including Biogeochemical Cycles (CLASSIC) model. The resulting eight (2 x 2 x 2) model simulations are evaluated using an in-house model evaluation framework that uses multiple observations-based data sets for a range of quantities. The simulated area burned, fire CO₂ emissions, soil carbon mass, vegetation carbon mass, runoff, heterotrophic respiration, gross primary productivity, and sensible heat flux show the largest spread across the eight simulations relative to their mean. Simulated net atmosphere-land CO₂ flux, a critical determinant of the performance of LSMs, is found to be largely independent of the simulated pre-industrial vegetation and soil carbon mass although our framework represents the historical increase in crop area in the same way in both land cover representations. This indicates that models can provide reliable estimates of the strength of the land carbon sink despite some biases in carbon stocks. Results show that evaluating an ensemble of model results against multiple observations disentangles model deficiencies from uncertainties in model inputs, observation-based data, and model configuration.

1. Introduction

The current generation land surface models (LSMs) explicitly simulate the fluxes of energy, water, momentum, and trace gases (including CO₂, CH₄, and N₂O) between the atmosphere and the land surface. These models have become an essential tool in understanding what role the land surface plays in the global climate system under current and projected future changes in environmental conditions, including atmospheric CO₂ concentration (Bonan and Doney, 2018). LSMs are also an essential component of climate and Earth system models (ESMs), together with their ocean and atmosphere components. Within the framework of ESMs, LSMs are coupled interactively to their atmospheric components through the fluxes of energy, momentum, and matter.

The complexity of LSMs has increased over time as more physical and biogeochemical processes have been included in their framework (Fisher and Koven, 2020; Kyker-Snowman et al., 2022). This increased complexity combined with the uncertainty in our understanding of physical and biogeochemical processes implies that different models respond differently even when driven with the same external forcings. One estimate of the uncertainty in our understanding of land surface physical and biogeochemical processes is obtained by evaluating the inter-model spread in a given quantity when models are forced in the same manner. Other than the uncertainty among models due to differences in their model structures and parameterizations of various processes, uncertainty also exists due to at least three other

reasons. These include uncertainty 1) in parameter values¹ of represented processes, 2) in driving meteorological data, and 3) in the specification of the geophysical fields. LSMs are typically driven with meteorological data consisting of seven primary variables (incoming long and shortwave radiation, temperature, precipitation, specific humidity, wind speed, and pressure). In addition, the geophysical fields of land cover, soil texture, and soil permeable depth are also required. Driving data for LSMs also consist of atmospheric CO₂ concentration and other model-specific external forcings such as nitrogen deposition and fertilizer application rates for models that include a representation of the terrestrial nitrogen cycle, and lightning, population density, and gross domestic product (GDP) for models that simulate wildfires.

Every year more than 15 land surface modelling groups participate in the TRENDY (trends in net land-atmosphere carbon exchanges) project where they perform a set of simulations that are driven with specified external forcings. The simulations are performed from the year 1700 to the present day. These simulations contribute to the annual Global Carbon Project's (GCP) analysis of the land carbon sink together with its analysis of anthropogenic CO₂ emissions and the ocean carbon sink (Friedlingstein et al., 2019). The external forcings used to drive LSMs in the TRENDY intercomparison include, 1) six hourly meteorological data from 1901 to the present day (the most recent 2020 TRENDY intercomparison used the CRU-JRA forcing obtained by blending the climate research unit (CRU) monthly data and the Japanese reanalysis (JRA)); 2) atmospheric CO₂ concentration; and 3) information about changes in crop area and other land use changes (LUC) from the land use harmonization (LUH) product (Hurtt et al., 2020a). The information about

¹ Changes in parameter values do not constitute different parameterizations. For example, two models may use the same parameterization, say $y=mx+b$, but different values of its parameters m and b . However, $y=mx + b$ and $y = mx^2$ are considered to be two different parameterizations.

changes in crop area and other LUC is used by land surface modelling groups to reconstruct historical land cover from the year 1700 to the present day consistent with the number of the plant functional types (PFTs) a given model represents. The protocol also provides nitrogen deposition and fertilization application rates for models including nitrogen cycling.

Models participating in the TRENDY simulations are thus driven with common meteorological and LUC forcings as part of its protocol. The resulting spread across models participating in the TRENDY project thus provides a measure of inter-model uncertainty, as mentioned earlier. Traditionally the uncertainty associated with model structure has gained the most attention and the scientific community has responded to this by performing model intercomparison projects (MIPs) where models are driven according to a common protocol. The coupled model intercomparison project (CMIP) in the climate community together with its various sub-projects (Eyring et al., 2016) is another prominent example. MIPs now routinely form the basis of evaluating models against observations and multi-model means of various quantities. Multi-model means are also considered the best estimate for a given quantity (Tebaldi and Knutti, 2007).

The modelling community has been long aware of the uncertainty associated with parameter values, since a large fraction of physical and biogeochemical model processes are parameterized, and such uncertainty analysis dates back to the early hydrological models (e.g. Hornberger and Spear, 1981; Beven and Binley, 1992). More recent examples of parameter value uncertainty in the context of a given LSM include Poulter et al. (2010), Booth et al. (2012), and Li et al. (2018a). The land surface modelling community, however, has only recently begun to address and quantify uncertainty related with driving meteorological data. Wu et al. (2017), for

example, illustrate the uncertainty in gross primary productivity (GPP) simulated by the Lund-Potsdam-Jena General Ecosystem Simulator (LPJ-GUESS) model when driven by six different meteorological data sets. Bonan et al. (2019) analyze the uncertainty in simulated carbon cycle related variables using three versions of the Community Land Model (CLM) when driven with two meteorological data sets over the historical period. Slevin et al. (2017) assess the uncertainty in simulated GPP by the JULES land model when driven by three different meteorological data sets. Studies that evaluate the effect of different land cover representations on model performance are even fewer. Tian et al. (2004) and Lawrence and Chase (2007) study the effect of new land surface boundary conditions, including leaf area index and fractional vegetation cover, based on the MODIS satellite data as implemented in CLM2 in the Community Atmosphere Model (CAM2) and CLM3 in the Community Climate System Model (CCSM 3.0), respectively.

Here, we drive the Canadian Land Surface Scheme Including Biogeochemical Cycles (CLASSIC) with two sets of historical meteorological forcings and also two land cover representations to quantify the uncertainty associated with both these forcings. Other than these, we also use two versions of the CLASSIC model: one that represents the interactions between the carbon (C) and nitrogen (N) cycles and the other in which these interactions are turned off. CLASSIC has contributed to the simulations for the TRENDY intercomparison, and the GCP, since 2016 (formerly under the CLASS-CTEM name). Seiler et al. (2021a) have evaluated how well the CLASSIC model performs when forced with three different meteorological data sets using the model version without the N cycle. Using the two meteorological forcing data sets, two representations of land cover, and two versions of the model we perform eight simulations over the historical period since 1700. All of these simulations may be considered equally likely

representations of the modelled state of the land surface over the historical period. Yet, they all have their own distinct biases since simulated land surface states and fluxes are different. We use these simulations to illustrate the uncertainty associated with meteorological forcing and the two different representations of land cover that are used to drive the model. We also use an in-house open-source benchmarking system (see code/data availability section) to evaluate these different simulations against observations-based data sets: AMBER (Automated Benchmarking R Package) (Seiler et al., 2021b) uses gridded and in-situ observation-based estimates of 19 energy, water, and C cycle related variables to evaluate LSMs.

Section 2 of this paper describes the framework of the CLASSIC land model and the forcing data that are required to drive the model. Section 3 describes the two meteorological data sets, the two representations of land cover that are used to drive the model, and the simulations performed for this study. Section 4 analyses the results from the simulations to illustrate their different states and reports results from the AMBER benchmarking exercise. Finally, the discussion and conclusions are presented in Section 5. The use of more than one meteorological forcing data sets and land cover representation yields a conundrum since tuning model parameters for a given forcing data set is not a useful exercise anymore. We also report a new finding that despite different land C states (characterized in terms of vegetation and soil C mass) in the eight simulations considered here, the net atmosphere-land CO₂ flux over the historical period in these simulations is consistent with estimates from the GCP. This and the discussion about the broader question of model tuning are also presented in Section 5.

2. The CLASSIC land modelling framework

2.1 The physical and carbon biogeochemical processes

The CLASSIC land model is the successor to, and based on, the coupled Canadian Land Surface Scheme (CLASS; (Verseghy, 1991; Verseghy et al., 1993)) and the Canadian Terrestrial Ecosystem Model (CTEM; (Arora and Boer, 2005; Melton and Arora, 2016b)). CLASSIC also serves as the land component in the family of Canadian Earth System Models (Arora et al., 2009, 2011; Swart et al., 2019). Melton et al. (2019) provide an overview of the CLASSIC land model and launched it as a community model. The basis of the modelling of physical and biogeochemical processes in CLASSIC comes from CLASS and CTEM, respectively, both of which have a long history of development. CLASSIC simulates land-atmosphere fluxes of water, energy, and momentum based on its physics, and fluxes of CO₂, CH₄, N₂O, NO_x, and NH₃ based on its biogeochemical process. The representation of the terrestrial N cycle is a new addition to CLASSIC (Asaadi and Arora, 2021; Kou-Giesbrecht and Arora, 2022) and allows for the simulation of the interactions between the C and N cycles explicitly.

The CLASSIC model simulations can be performed over a spatial domain, which may be global or regional, using gridded data or at a point scale, e.g. using meteorological and geophysical data from a FluxNet site. The primary physical and biogeochemical processes of CLASSIC are briefly summarized in the next two sections.

2.1.1 Physical processes

The calculations for physical processes in CLASSIC are performed over vegetated, snow, and bare fractions at a time step of 30 minutes. In the version used here, the fractional coverage of the four plant functional types (PFTs) (needleleaf trees, broadleaf trees, crops, and grasses)

characterizes vegetation for each grid cell. The fractional coverage of these four PFTs is specified over the historical period in this study. The structure of vegetation is characterized by leaf area index (LAI), vegetation height, canopy mass, and rooting distribution through the soil layers all of which are dynamically simulated by the biogeochemical module of CLASSIC. Twenty ground layers represent the soil profile, starting with 10 layers of 0.1 m thickness. The thickness of layers gradually increases to 30 m for a total ground depth of over 61 m. The depth of permeable soil layers and thus the depth to bedrock varies geographically and is specified based on the SoilGrids250m data set (Hengl et al., 2017). Liquid and frozen soil moisture contents, and soil temperature, are determined prognostically for permeable soil layers. The temperature, albedo, mass, and density of a single-layer snow pack (when the climate permits snow to exist) are also prognostically modelled. The result of physics calculations yields fluxes of energy (primarily net radiation, ground heat flux, and latent and sensible heat fluxes) and water (primarily evapotranspiration and runoff) at the land-atmosphere boundary.

2.1.2 Biogeochemical processes

The biogeochemical processes in CLASSIC, based on CTEM, are described in detail in the appendix of Melton and Arora (2016). The biogeochemical processes simulate the land-atmosphere exchange of CO₂ and as a result simulate vegetation as a dynamic component depending on the environmental conditions.

The biogeochemical module of CLASSIC prognostically calculates the amount of C in the model's three live (leaves, stem, and root) and two dead (litter and soil) C pools for each PFT. The live vegetation pools are separated into their structural and non-structural components. The C

174 amount in these pools is represented per unit land area (kg C/m^2). The amount of C in the live
175 and dead C pools and all terrestrial ecosystem processes in the biogeochemical module in this
176 study are modelled for nine PFTs that map directly onto the four base PFTs used in the physics
177 module of CLASSIC. Needleleaf trees are divided into their deciduous and evergreen phenotypes,
178 broadleaf trees are divided into cold deciduous, drought deciduous, and evergreen phenotypes,
179 and crops and grasses are divided based on their photosynthetic pathways into C_3 and C_4
180 versions. The sub-division of PFTs is essential for modelling biogeochemical processes. For
181 instance, simulating the onset and offset of leaves is different between evergreen and deciduous
182 phenotypes of needleleaf and broadleaf trees. However, once the leaf area index (LAI) is known,
183 a physical process does not need information about the underlying deciduous or evergreen
184 nature of leaf phenology. For example, the interception of rain and snow by canopy leaves (that
185 is typically modelled as a function of LAI and a PFT-dependent parameter that accounts for leaf
186 orientation and shape) does not depend on the underlying evergreen or deciduous nature of the
187 leaf phenology. In general, biogeochemical processes benefit more in terms of realism than
188 physical processes when the number of PFTs is increased. For example, in CLASSIC, large changes
189 in leaf area index (LAI) do not change total latent heat flux considerably since the partitioning of
190 evapotranspiration into its sub-components (transpiration, soil evaporation, and
191 evaporation/sublimation of intercepted rain/snow) adjusts. A decrease in transpiration and
192 evaporation of intercepted precipitation, due to a decrease in LAI, is compensated by an increase
193 in soil evaporation. This is expected since water and energy fluxes are determined largely by
194 available energy and precipitation.

The litter and soil C pools are tracked for each soil layer but the movement of C between the soil layers is not yet modelled. Other than photosynthesis and leaf respiration which are modelled at a time step of 30 minutes all other biogeochemical processes are modelled at a daily time step. These include: 1) allocation of C from leaves to stem and roots, 2) autotrophic respiration from the live C pools and heterotrophic respirations from the dead C pools, 3) leaf phenology, 4) turnover of live vegetation components that generates litter, 5) mortality, 6) LUC, and 7) fire (Arora and Melton, 2018). Competition between PFTs for space is not modelled in this study and fractional coverage of the nine PFTs is specified based on the representation of the land cover as explained in the next section.

When the N cycle is turned on, land-atmosphere fluxes of N_2O , NO_x , and NH_3 , and N leaching are also modelled in response to biological N fixation, N fertilizer inputs, and N deposition from the atmosphere. In particular, when the N cycle interacts with the C cycle, the maximum photosynthetic capacities of model PFTs ($V_{c,\text{max}}$) are determined prognostically as a function of their leaf N content (Asaadi and Arora, 2021; Kou-Giesbrecht and Arora, 2022). When the N cycle is turned off, prescribed PFT-specific $V_{c,\text{max}}$ rates are used (Melton and Arora, 2016a) and an empirical downregulation parameterization is used to emulate the effect of nutrient constraints as atmospheric CO_2 increases (Arora et al., 2009). N in all model components (leaves, stem, roots, litter, and soil organic matter) is prognostically tracked, and therefore C:N ratio of all components is prognostically modelled except for soil organic matter for which a C:N ratio of 13 is specified. In addition, N in the soil mineral pools of nitrate (NO_3^-) and ammonium (NH_4^+) is also prognostically modelled.

3. Driving data for CLASSIC and model simulations

3.1 Land cover

Land cover is one of the most important geophysical fields that is required by LSMs and at its most basic level provides information about fractional vegetation cover in each grid cell for a given regional or global domain. Vegetation in LSMs is typically represented in terms of PFTs. Models may choose to represent a basic set of a few PFTs (trees, grasses, shrubs, and crops) or a more elaborate set that distinguishes PFTs based on their stature (trees, grasses, or shrubs), leaf form (needleleaf or broadleaf), leaf phenology (evergreen or deciduous), photosynthetic pathway (C_3 or C_4), and geographical location (tropical, temperate, or boreal). The version of CLASSIC in this study uses a somewhat smaller set of nine PFTs for biogeochemical processes as described in the previous section. The fractional coverage of PFTs in a model may be dynamically simulated based on competition between PFTs or prescribed based on observation-based land cover information. While CLASSIC does have a parameterization of competition between its PFTs (Arora and Boer, 2006; Melton and Arora, 2016b), for the historical simulations considered here and for the simulations that contribute to the TRENDY ensemble, prescribed fractional coverage of PFTs is used.

For the process of generating a historical reconstruction of land cover, consisting of time-varying fractional coverage of a model's PFTs, two types of observation-based data sets are used. The first is a remotely-sensed land cover product that represents the geographical distribution of land cover for the present day for a short period. Examples of this include the GLC 2000 land cover product which represents the November 1999 to December 2000 period (<https://forobs.jrc.ec.europa.eu/products/glc2000/glc2000.php>) and the more recent European Space Agency (ESA) Climate Change Initiative (CCI) land cover product for the period 1992-2018

(ESA, 2017). The second type of data set required to reconstruct historical land cover is that of a spatially and temporally varying cropland (and pasture) area for a much longer period, which in this case is based on the data set provided by the land use harmonization (LUH) product as part of the TRENDY protocol for the period 850-2018. The LUH product is comprehensive (Hurt et al., 2020b). For example, not all models use the pasture area and other information provided in the LUH product.

The process of generating land cover for a given model's PFTs is a three-step process. First, the fractional coverage of model PFTs is obtained from a remotely sensed land cover product that represents the snapshot of land cover for the present day. This requires typically mapping 20 – 40 land cover classes that exist in a remotely-sensed land cover product to a given model's PFTs. This step introduces the largest uncertainty in the entire process. The original land cover in the CLASSIC model is based on the GLC 2000 land cover product. Table 2 of Wang et al. (2006) summarizes the mapping/reclassification of the 22 GLC 2000 land cover categories to the nine PFTs used in CLASSIC. Each land cover class was split into one or more of the nine CLASSIC PFTs based on the class description and knowledge of global biomes. For example, the discrete "broadleaf deciduous open tree cover" category of the GLC 2000 product is assumed to consist of 60% broadleaf deciduous trees, 20% grasses, and 20% bare ground. This first step yields a snapshot of land cover expressed in terms of the fractional coverage of CLASSIC's nine PFTs. The second step of generating fractional coverage of PFTs for a given snapshot in time requires replacing the fractional area of crop categories with values from the LUH data set for the same year. For example, when using the GLC 2000 land cover product, the area of C₃ and C₄ crops from the LUH data set for the year 2000 are used, and the fractional coverage of the other seven non-

crop CLASSIC PFTs is adjusted such that the total vegetation fraction in each grid cell stays the same. Finally, in the last step, the temporally varying crop area from the LUH product is used to go backward in time to 1700 from the year 2000 with typically decreasing crop area while the area of other non-crop PFTs is adjusted in proportion to their existing fractional coverage such that the total vegetation fraction in each grid cell stays the same. Similarly, the area of C_3 and C_4 crops from the LUH product is used from the year 2000 onwards to the present day. All these steps yield a reconstruction of historical land cover, expressed in terms of fractional coverage of CLASSIC's nine PFTs (as interpreted from the GLC 2000 land cover product), from 1700 to 2018, in which crop area changes spatially and temporally according to the LUH product.

GLC 2000 is an older land cover product and more recent land cover products are now available. Here, in addition to the GLC 2000 based land cover, we also use the European Space Agency (ESA) Climate Change Initiative (CCI) land cover product. The ESA CCI land cover product is available at 300 m spatial resolution for the period 1992-2018 and contains 37 land cover categories (ESA, 2017). We use the land cover from the year 1992 to create a snapshot of CLASSIC PFTs for the present day. Although there is some interannual variability overall the total vegetated area doesn't change substantially from 1992-2018 in the ESA-CCI land cover. A default mapping/reclassification table for converting the ESA CCI classes into PFTs is provided in its user guide (ESA, 2017). However, it overestimates tree cover along the taiga-tundra transition zone and underestimates it elsewhere in Canada (Wang et al., 2018, 2019). Wang et al. (2022) have developed a new reclassification table for converting the 37 ESA CCI land cover categories to CLASSIC's nine PFTs which is used in this study. A high-resolution land cover map over Canada and a tree cover fraction data at 30 m resolution are used to compute the sub-pixel fractional

composition of each class in the ESA CCI dataset, which is then used to inform the cross-walking reclassification procedure (Wang et al., 2022).




Figure 1: Comparison of zonally summed areas of total vegetation (a), grass (b), and tree (c) cover used in the CLASSIC model based on GLC 2000 (blue line) and ESA CCI (dark red line) land cover products to each other, to selected other models that participated in the 2020 TRENDY intercomparison (grey lines) for which land cover information was available, and to Li et al. (2018) (dotted black line) who analyzed the ESA CCI data. All data correspond to the 1992-2018 period. CLASSIC does not yet explicitly represents shrub PFTs. Tall shrubs are merged into tree PFTs in CLASSIC. For the Li et al. (2018) data plotted here, the shrub PFTs are combined with the tree PFTs for a consistent comparison to CLASSIC.

The above process yields two representations of land cover in which the geographical distribution of CLASSIC PFTs is based on GLC 2000 and ESA CCI land cover products. Both these

representations include the same reconstruction of crop area over the historical period. Figure 1 illustrates the uncertainty in land cover by comparing zonally summed areas of total vegetation, tree, and grass cover in CLASSIC, averaged over the period 1992-2018, when model land cover is based on the GLC 2000 (blue line) and ESA CCI (dark red line) land cover products. These two estimates are also compared to selected other models that participated in the 2020 TRENDY intercomparison (grey lines), also for the period 1992-2018, for which land cover information was available, and to Li et al. (2018b) (dotted black line) who analyzed the ESA CCI data based on the default reclassification table from the ESA CCI user guide. Figure 1 shows while there is relatively good agreement across TRENDY models in terms of total vegetation cover there's a much larger uncertainty in its split between tree and grass PFTs. There are two reasons for the spread in total vegetated, treed, and grassed areas across TRENDY models. First, modelling groups use different remotely sensed land cover products for obtaining fractional cover of their model PFTs. Second, the current process of mapping/reclassifying 20-40 land cover classes of a land cover product to a model's PFTs is mainly based on the class description and expert judgment which introduces subjectiveness in the process. Compared to the GLC 2000 based land cover in the CLASSIC model, the newer ESA CCI based land cover yields a somewhat higher total vegetation cover, a higher grass cover, and a somewhat lower tree cover area. Unlike the older GLC 2000 based land cover used in CLASSIC, the newer ESA CCI based grass and tree cover area are within the range of the TRENDY models reported here. Finally, Figure 1 also allows us to compare the results from the analysis of Li et al. (2018b) for the ESA CCI land cover (dotted black line) to the ESA CCI reclassification for CLASSIC (dark red line) by (Wang et al., 2022). Li et al. (2018b) used the default mapping/reclassification table for converting the ESA CCI classes into PFTs. This comparison

illustrates that the remapping of the ESA CCI land cover classes to CLASSIC's PFTs yields total vegetation, tree, and grass coverage that is broadly comparable to Li et al. (2018b) although some differences remain for the grasses.

Our framework accounts for the uncertainty in land cover representation. However, since both land cover representations in our study account for the increase in crop area over the historical period in the same way by adjusting the area of non-crop PFTs in proportion to their existing coverage using the LUH product, our framework is unable to account for the uncertainty associated with the implementation of LUC. Di Vittorio et al. (2018) quantify this uncertainty by implementing several approaches to account for the increase in crop area over the historical period in the framework of an integrated assessment model: by preferentially converting grasses and shrubs, by preferentially converting forests, and by proportionally adjusting areas of non-crop PFTs in a way similar to ours. LUC emissions are higher if the increase in crop area is preferentially obtained by converting forests. A similar uncertainty analysis for LUC emissions is performed by Peng et al. (2017) using the ORCHIDEE land model who analyze the effect of using different rules to incorporate the changes in crop and pasture area over the historical period. The uncertainty related to incorporating LUC information to modify a model's land cover is further illustrated in Di Vittorio et al. (2014) and Meiyappan and Jain (2012).

3.2 Meteorological data

As a land surface component of an ESM, CLASSIC requires meteorological forcing at a sub-daily temporal resolution. In the offline simulations reported here, the model is run with half-

hourly values of meteorological data (incoming long and shortwave radiation, temperature, precipitation, specific humidity, wind speed, and pressure). The first meteorological data set used to drive CLASSIC is from the TRENDY protocol for the year 2020, CRU-JRA v2.1.5, which provides 6 hourly values of the seven variables from the Japanese reanalysis (JRA) with monthly values adjusted to the climate research unit's data (CRU, <https://crudata.uea.ac.uk/cru/data/hrg/>). This yields a blended product from year January 1901 to December 2019 with the 6-hourly temporal resolution of a reanalysis but without the biases that may be present in reanalysis data (Harris, 2020). The second meteorological data set used here to drive CLASSIC is from the Global Soil Wetness Project 3 (GSWP3). The GSWP3 forcing data are based on a dynamical downscaling of the 20th century reanalysis (Compo et al., 2011) using a Global Spectral Model (GSM) run at about 50 km resolution. GSM is nudged towards the vertical structures of 20th century (20CR) zonal and meridional air temperature and winds so that the synoptic features are retained at their higher spatial resolution. Additional bias corrections are also performed as explained in van den Hurk et al. (2016). The GSWP3 forcing is available for the 1901-2016 period. The 6-hourly values from both the CRU-JRA and GSWP3 forcings are further disaggregated to half-hourly values for use by CLASSIC.

Figure A1 compares the two meteorological forcings data sets, over the 1997-2016 period, to illustrate that although these two data sets are very similar there are differences between the two. Global precipitation over land (excluding Greenland and Antarctica) in the GSWP3 data set (857 mm/year) is somewhat higher than in the CRU-JRA data set (820 mm/year). The global near-surface air temperature over land (excluding Greenland and Antarctica) is also slightly higher in the GSWP3 data set (14.22 °C) compared to the CRU-JRA data set (14.08 °C). The

largest temperature difference occurs between the two data sets over the northern tropics (panel h) where the GSWP3 data set is about 0.93 °C warmer than the CRU-JRA data set. The geographical distribution of mean annual temperature is very similar between the two data sets but there are some differences in the geographical distribution of precipitation (not shown). Despite very similar total precipitation amounts and their seasonality over large global regions in the two data sets, differences exist in the frequency distribution of precipitation. Figure A2 illustrates this over three broad regions, the Amazon, the Sahel, and the Midwest United States, which shows the frequency distribution of daily precipitation amounts (mm/day) over the 1997-2016 period from the two data sets. Figure A2 shows that the frequency of precipitation events greater than about 5-10 mm/day is higher in the GSWP3 data set compared to the CRU-JRA data set for the Amazonian, the Sahel, and the Midwest United States regions.

3.3 Other forcings

Other than the land cover and meteorological forcings CLASSIC requires globally averaged atmospheric CO₂ concentration, geographically varying time-invariant soil texture and soil permeable depth, population density, time-invariant monthly lightning, and geographically and time-varying N fertilizer application rates and atmospheric N deposition rates. The atmospheric CO₂ concentration values are provided by the TRENDY protocol. The soil texture information consists of the percentage of sand, clay, and organic matter and is derived from Shangguan et al. (2014). N fertilizer is specified according to the TRENDY protocol and based on Lu and Tian (2017). N deposition is also specified according to the TRENDY protocol and based on model forcings provided for the sixth phase of CMIP (CMIP6) through input4MIPs (Hegglin et al., 2016). N deposition for the historical (1850-2014) period is used as is provided while that for

the period 2015-2018 is specified based on N deposition from the SSP5-85 scenario. For the period 1700-1849, N deposition values from the year 1850 are used.

Table 1: Summary of simulations performed with two representations of the historical land cover, two sets of meteorological data, and two versions of the CLASSIC land model.

Simulation	Land cover reconstruction	Meteorological forcing	N cycle interactions with the C cycle
A	based on GLC 2000	CRU-JRA v2.1.5	On
B	based on GLC 2000	GSWP3	On
C	based on GLC 2000	CRU-JRA v2.1.5	Off
D	based on GLC 2000	GSWP3	Off
E	based on ESA CCI	CRU-JRA v2.1.5	On
F	based on ESA CCI	GSWP3	On
G	based on ESA CCI	CRU-JRA v2.1.5	Off
H	based on ESA CCI	GSWP3	Off

3.4 Model simulations

Using the two representations of the historical land cover (based on the GLC 2000 and ESA CCI land cover products), the two sets of meteorological data (CRU-JRA and GSWP3), and the two versions of the CLASSIC model (with and without interactions between the C and N cycles) we perform eight sets of pre-industrial and historical simulations as summarized in Table 1. Pre-industrial simulations that correspond to the year 1700 are required before doing the historical simulations (from which we analyze the model results) so that model pools can be spun up to near equilibrium for each combination of land cover, meteorological forcing, and model version. The pre-industrial simulations use 1901-1925 meteorological data repeatedly since this period shows little trends in meteorological variables. Global thresholds of net atmosphere-land C flux of 0.05 Pg/yr and net atmosphere-land N flux of 0.5 Tg N/yr, in simulations with the

N cycle turned on, are used to ensure the model pools have reached equilibrium. Each historical simulation is then initialized from its corresponding pre-industrial simulation after it has reached equilibrium. Simulations driven with the CRU-JRA meteorological data are performed for the period 1701-2018, and the period 1701-2016 for simulations driven with the GSWP3 meteorological data. Similar to the pre-industrial simulations, meteorological data from 1901-1925 is used repeatedly for the period 1701-1900. The global model simulations are performed at a spatial resolution of about 2.81° (about 312 km at the equator) and the size of the spatial longitude-latitude grid is 128×64 . All model forcings are regridded to this common spatial resolution. The model is run over about 1900 land grid cells at this resolution excluding glacial cells in Greenland and Antarctica.

3.5 Automated benchmarking

The results from the eight CLASSIC simulations reported here are evaluated using an in-house model benchmarking system called the Automated Model Benchmarking R package (AMBER) (Seiler et al., 2021b). AMBER is based on a skill score system originally developed by (Collier et al., 2018) which is used to quantify model performance and explained in detail in the appendix. Five scores are used that assess a model's bias (S_{bias}), root-mean-square error (S_{rmse}), seasonality (S_{phase}), interannual variability (S_{iav}), and spatial distribution (S_{dist}) against globally gridded and in-situ data set(s) of observation-based estimates for a given quantity. A score is computed by first calculating a dimensionless statistical metric, which is then scaled onto a unit interval, and finally calculating its spatial mean. Scores range from 0 to 1 and are dimensionless.

428 Higher values indicate better performance. Finally, an overall score $S_{overall}$ is calculated as follows
429 by giving twice as much weight to S_{rmse}

$$430 \quad S_{overall} = \frac{S_{bias} + 2S_{rmse} + S_{phase} + S_{iav} + S_{dist}}{1 + 2 + 1 + 1 + 1}. \quad (1)$$

431 The decision to give extra weight to S_{rmse} is entirely subjective but follows Collier et al. (2018).

432 The scores are calculated by comparing gridded and in-situ observation-based estimates,
433 referred to as reference data sets in Seiler et al. (2021b), for 19 energy (surface albedo, net
434 shortwave and longwave radiation, total net radiation, latent heat flux, sensible heat flux, ground
435 heat flux), water (soil moisture, snow, and runoff), and C cycle (GPP, net ecosystem exchange,
436 net biome productivity, aboveground biomass, soil C, LAI, area burnt, and fire CO₂ emissions)
437 related variables to model simulated quantities. Table 2 summarizes the source of these
438 observation-based data sets. The resulting model scores express to what extent simulated and
439 observation-based data agree. A low score does not necessarily indicate poor model
440 performance. Uncertainties in the meteorological forcing data and geophysical fields used to
441 drive the model, and/or in the observation-based data itself are possible reasons for the lack of
442 agreement. One way to assess uncertainties in observation-based data sets is to quantify the skill
443 score by comparing two independently-derived observation-based data sets (Seiler et al., 2022).
444 The resulting scores are referred to as benchmark scores and quantify the level of agreement
445 among the observation-based data sets themselves provided, of course, there are at least two
446 sets of observation-based data for a given quantity. The comparison of model scores against
447 benchmark scores then shows how well a model-simulated quantity compares to the reference
448 data sets relative to the agreement between the observation-based data sets themselves.

449 Table 2: Observation-based data sets used for model evaluation in AMBER.

Globally gridded variable(s)	Source	Approach used	Reference
Leaf area index	AVHRR	Artificial neural network	Claverie et al. (2016)
Net biome productivity	CAMS	Atmospheric inversion	Agustí-Panareda et al. (2019)
Net biome productivity	Carboscope	Atmospheric inversion	Rödenbeck et al. (2018)
Surface albedo, net shortwave and longwave radiation, net radiation	CERES	Radiative transfer model	Kato et al. (2013)
Net radiation, latent and sensible heat flux, ground heat flux, runoff	CLASSr	Blended product	Hobeichi et al. (2019)
Leaf area index	Copernicus	Artificial neural network	Verger et al. (2014)
Net biome productivity	CT2019	Atmospheric inversion	Jacobson et al. (2020)
Snow amount	ECCC	Blended product	Mudryk (2020)
Liquid soil moisture	ESA	Land surface model	Liu et al. (2011)
Area burnt	ESA CCI	Burned area mapping	Chuvieco et al. (2018)
Latent and sensible heat flux, gross primary productivity	FLUXCOM	Machine learning	Jung et al. (2019, 2020)
Above ground biomass	GEOCARBON	Machine learning	Avitabile et al. (2016); Santoro et al. (2015)
Surface albedo, net shortwave and longwave radiation, net radiation	GEWEXSRB	Radiative transfer model	Stackhouse et al. (2011)
Area burnt	GFED 4s	Burned area mapping	Giglio et al. (2010)
Gross primary productivity	GOSIF	Statistical model	Li and Xiao (2019)
Soil carbon	HWSD	Soil inventory	Wieder (2014); Todd-Brown et al. (2013)
Surface albedo	MODIS	Bidirectional Reflectance Distribution Function	Strahler et al. (1999)
Gross primary productivity	MODIS	Light use efficiency model	Zhang et al. (2017)
Leaf area index	MODIS	Radiative transfer model	Myneni et al. (2002)
Soil carbon	SGS250m	Machine learning	Hengl et al. (2017)
Above ground biomass	Zhang	Data fusion	Zhang and Liang (2020)
In situ variable(s)	Source	Approach used (number of sites)	Reference
Leaf area index	CEOS	Transfer function (141)	Garrigues et al. (2008)
Latent, sensible, and ground heat flux, gross primary productivity, ecosystem respiration, net ecosystem exchange	FLUXNET 2015	Eddy covariance (204)	Pastorello et al. (2020)
Above ground biomass	FOS	Allometry (274)	Schepaschenko et al. (2019)
Runoff	GRDC	Gauge records (50)	Dai and Trenberth (2002)
Snow amount	Mortimer	Gravimetry (3271)	Mortimer et al. (2020)
Above ground biomass	Xue	Allometry (1974)	Xue et al. (2017)

450

451

452 4. Results

Figures 2 through 9 show the time series and/or zonally-averaged values of annual values of a variable of interest when averaged across four ensemble members each according to whether the N cycle is turned on or not, whether the GLC 2000 or ESA CCI based land cover is used, and whether model simulations are driven by the CRU-JRA or GSWP3 meteorological data. Figures A3, A4, A6, A7, A9, and A11 in the appendix, which are complementary to the above-mentioned figures, show the physical and biogeochemical states of the land surface and primary physical fluxes of water and energy, and primary biogeochemical fluxes of CO₂ simulated by CLASSIC at the land-atmosphere boundary for all the eight simulations considered here. While the figures in the appendix illustrate the range in simulated physical and biogeochemical states and fluxes across the eight simulations, Figures 2 through 9 evaluate the effect of model structure, meteorological forcing, and land cover on a given quantity. We also quantify the spread across the eight simulations using the coefficient of variation (cv= standard deviation/mean) calculated using annual global values for a given quantity averaged over the 1997-2016 20-year period of each simulation. This time period is also used for other reported results.

4.1 Physical land surface state and fluxes

Figure A3, panels a and b, shows the globally-averaged simulated soil moisture and temperature in the top 1 m soil layer. While simulated soil temperature in the top 1 m is fairly similar across the eight simulations, the simulated soil moisture is distinctly separated into two groups. The separation into these two groups is caused by the driving meteorological data as shown in Figure 2. The coefficient of variation for soil moisture and temperature values averaged over the 1997-2016 period of each simulation are 0.02 and 0.004, respectively, indicating that overall the variation in these quantities is relatively small compared to their means. The use of

the GSWP3 meteorological dataset yields slightly higher (~4%) globally-averaged soil moisture compared to the CRU-JRA meteorological data set (236.5 mm vs. 227.1 mm, Figure 2c).




Figure 2: Time series of annual globally-averaged soil moisture in the top 1m averaged over the four ensemble members that are driven with and without an interactive N cycle (panel a), driven with the GLC 2000 and ESA CCI based land cover representations (panel b), and driven with the GSWP3 and CRU-JRA meteorological data (panel c). The thin lines show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, for each ensemble averaged over its set of four simulations are also shown.




Figure 3: Time series of annual global evapotranspiration and runoff (over all land area excluding Greenland and Antarctica) averaged over the four ensemble members that are driven with and without an interactive N cycle (panels a, b), driven with the GLC 2000 and ESA CCI based land cover (panels c, d), and driven with the GSWP3 and CRU-JRA meteorological data (panels e, f). The thin lines show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, for each ensemble averaged over its set of four simulations are also shown.

Figure A3, panels c and d, shows the simulated fluxes of global evapotranspiration and runoff across the eight simulations. Similar to soil moisture, evapotranspiration and runoff also fall broadly into two groups and the reason for this again is the driving meteorological data. Figure 3 shows that while the biggest factor that affects evapotranspiration and runoff is the difference in driving meteorological data the interactive N cycle also affects these water fluxes. Neither evapotranspiration nor runoff is significantly affected by the choice of land cover. The reason an interactive N cycle affects evapotranspiration is that the N cycle in CLASSIC affects the rate of photosynthesis through the prognostic determination of leaf N content. Photosynthesis in turn affects canopy conductance, which affects transpiration through the canopy leaves. Average evapotranspiration over the 1997-2016 period of the simulations driven with GSWP3 meteorological data is about 9% lower than in simulations driven with CRU-JRA meteorological data (65.8 vs. 72.1 $\times 1000 \text{ km}^3/\text{year}$, Figure 3, panel e). An interactive N cycle reduces evapotranspiration by about 2% due to lower photosynthesis rates as shown later (Figure 3, panel a). Average runoff is about 27% higher in simulations driven with GSWP3 compared to simulations driven with CRU-JRA meteorological data (52.6 vs 41.3 $\times 1000 \text{ km}^3/\text{year}$, Figure 3, panel f). This is due to slightly high precipitation in the GSWP3 meteorological data set (Figure A1) but is more so due to the simulated lower evapotranspiration when using the GSWP3 data (Figure 3, panel e). The coefficient of variation for evapotranspiration and runoff values averaged over the last 20 years of each simulation are 0.05 and 0.13, respectively.




Figure 4: Time series of annual global latent and sensible heat fluxes (over all land area excluding Greenland and Antarctica) averaged over the four ensemble members that are driven with and without an interactive N cycle (panels a, b), driven with the GLC 2000 and ESA CCI based land cover (panels c, d), and driven with GSWP3 and CRUJRA meteorological data (panels e, f). The thin lines show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, for each ensemble averaged over its set of four simulations are also shown.

Figure A4 shows the primary energy fluxes from the eight simulations. These include net downward shortwave and longwave radiation, and latent and sensible heat fluxes. Incoming shortwave and longwave radiation are part of the driving meteorological data. Similar to water fluxes, the differences in energy fluxes in CLASSIC are also primarily driven by differences in meteorological data (Figure A4, A5, and Figure 4). Net shortwave radiation (Figure A4, panel a) is equal to incoming shortwave radiation minus the fraction that is reflected back. Net longwave radiation (Figure A4, panel b) is equal to incoming longwave radiation minus the longwave radiation emitted by the land based on its surface temperature following the Stefan-Boltzmann law. The difference in net shortwave radiation is also affected by simulated vegetation biomass and leaf area index. The latter affects surface albedo which determines what fraction of incoming shortwave radiation is reflected. This is the reason why an interactive N cycle affects net shortwave radiation since the N cycle affects photosynthesis, and in turn, simulated vegetation biomass and leaf area index (Figure A5, panel b). Latent heat flux is affected primarily by meteorological data (Figure 4) but also if the N cycle is interactive or not since it is essentially evapotranspiration but in energy units. Finally, differences in sensible heat fluxes are strongly affected by differences in driving meteorological data (Figure 4). Globally-averaged sensible heat flux in the simulations driven with GSWP3 data is ~14% higher compared to CRU-JRA driven simulations (40.2 vs. 35.0 W/m²). The coefficient of variation for latent and sensible heat flux values averaged over the last 20 years of each simulation are 0.05 and 0.07, respectively. Net shortwave (cv=0.006) and longwave (cv=0.03) radiative fluxes vary little across the eight simulations.




Figure 5: Time series of annual global vegetation C mass (over all land area excluding Greenland and Antarctica) (panels a, c, and e) and zonally-averaged values of vegetation C mass over land (panels b, d, and f) averaged over the four ensemble members that are driven with and without an interactive N cycle (panels a, b), driven with the GLC 2000 and ESA CCI based land cover (panels c, d), and driven with GSWP3 and CRU-JRA meteorological data (panels e, f). The thin lines for the time series show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown in panels a, c, and e.

4.2 Biogeochemical land surface state and fluxes

4.2.1 Primary CO₂ fluxes and C pools

Figure A6 shows the simulated C state of the land surface expressed in terms of vegetation and soil C pools. Panels a and b show the annual time series of global vegetation and soil C mass from the eight simulations, and panels c and d show their zonally-averaged distributions averaged over the last 20 years of each simulation. The biggest difference in the time series of global vegetation ($cv=0.16$) and soil ($cv=0.21$) C mass compared to soil moisture and temperature, which characterized the physical land surface state, is the large spread across the eight simulations as indicated by their high cv values. The zonally-averaged values further provide insight into the reasons for this spread and show that the largest differences between simulated vegetation and soil C occur at northern high latitudes (north of about 40°N). Panels c and d of Figure A6 also show observation-based zonally-averaged values of vegetation and soil C mass based on the Reusch and Gibbs (2008) and the Harmonized World Soils Database (v1.2) (Fischer et al., 2008), respectively, to provide a reference. A more thorough comparison with observations is provided in Section 4.3.

Differences in vegetation C mass are caused primarily when the N cycle is interactive or not (Figure 5). Both land cover and the driving meteorological data play a smaller role in the simulated spread of vegetation C mass (Figure 5). The ESA CCI based land cover has a larger vegetated area but most of this increase comes from an increase in the area of grasses that do not store a lot of C in their vegetation C mass. The spread in simulated soil C is caused due to the





Figure 6: Time series of annual global soil carbon mass (over all land area excluding Greenland and Antarctica) (panels a, c, and e) and zonally-averaged values of soil carbon mass over land (panels b, d, and f) averaged over the four ensemble members that are driven with and without an interactive N cycle (panels a, b), driven with the GLC 2000 and ESA CCI based land cover (panels c, d), and driven with GSWP3 and CRU-JRA meteorological data (panels e, f). The thin lines for the time series show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown in panels a, c, and e.



586

587 Figure 7: Time series of annual global gross primary productivity (over all land area excluding
 588 Greenland and Antarctica) (panels a, c, and e) and zonally-averaged values of gross primary
 589 productivity over land (panels b, d, and f) averaged over the four ensemble members that are
 590 driven with and without an interactive N cycle (panels a, b), driven with the GLC 2000 and ESA
 591 CCI based land cover (panels c, d), and driven with GSWP3 and CRU-JRA meteorological data
 592 (panels e, f). The thin lines for the time series show the individual years and the thick lines show
 593 their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and
 594 present-day (1997-2016) time periods, and their difference, are also shown in panels a, c, and e.

N cycle but also the choice of land cover (Figure 6). Since CLASSIC assumes that litter from grasses is more recalcitrant than that from trees, the choice of ESA CCI based land cover leads to a higher soil C mass because it has a higher grass area than the GLC 2000 based land cover (Figure 6, panels c and d). The choice of meteorological data does not affect the magnitude of simulated globally-summed soil C mass significantly but does affect its change over the historical period. In Figure 6 (panel c) the decrease in soil C mass from the 1851-1860 period to the 1997-2016 period is higher when using the GSWP3 (29.9 Pg C) compared to when using the CRU-JRA (14.8 Pg C) meteorological data.

The reason why an interactive N cycle in CLASSIC affects vegetation C and soil C mass, and why the ESA CCI based land cover yields high soil C, is seen in Figures A7 and 7. Figure A7 shows the spread of primary C fluxes including gross primary productivity (GPP) ($cv=0.07$), and autotrophic ($cv=0.04$) and heterotrophic ($cv=0.10$) respiratory fluxes, across the eight simulations. Since GPP is lower in the runs with the N cycle, both vegetation (Figure 5a) and soil C mass (Figure 6a) are also lower. The lower GPP in the runs with the N cycle is due primarily to lower GPP at high latitudes (Figure 7b) which yields low vegetation C mass at high latitudes (Figure 5b). Low GPP at high latitudes translates to even larger relative differences in soil C given the longer turnover time scales of soil C at high latitudes (Figure 6b). The use of the ESA CCI based land cover which has a higher grass area than the GLC 2000 based land cover leads to higher GPP (Figure 7d) and therefore higher soil C at all latitudes (Figure 6d). In Figure A8, global heterotrophic and autotrophic respiratory fluxes are most affected by land cover and the inclusion or absence of an interactive N cycle but not by the driving meteorological data.




Figure 8: Time series of annual area burned (over all land area excluding Greenland and Antarctica) (panels a, c, and e) and zonally-averaged values of area burned (panels b, d, and f) averaged over the four ensemble members that are driven with and without an interactive N cycle (panels a, b), driven with the GLC 2000 and ESA CCI based land cover (panels c, d), and driven with GSWP3 and CRU-JRA meteorological data (panels e, f). The thin lines for the time series show the individual years and the thick lines show their 11-year moving average in panels (a), (c), and (e). Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown for panels (a), (c), and (e).

Overall, while the primary biogeochemical fluxes (cv values vary from 0.04 to 0.10) vary as much as the water and energy fluxes, the resulting spread in vegetation C mass (cv=0.16) and soil C mass (cv=0.21) across the eight simulations is much larger and driven primarily by the inclusion or absence of an interactive N cycle and the difference in land cover.

4.2.2 Area burned and fire CO₂ emissions

Figure A9 shows the time series of global area burned and global fire CO₂ emissions, and their zonally-averaged values. We chose the area burned (cv=0.24) and fire CO₂ emissions (cv=0.21) in addition to the primary biogeochemical fluxes since fire shows large variability both in space and in time, and both these variables yield the largest spread across the eight simulations, among all the fluxes and simulated quantities considered here. Figures A9 (panels c and d) also show observation-based estimates for area burned and fire CO₂ emissions based on GFED 4s (Giglio et al., 2013) to provide an observation-based context. Figures 8 and A10 help us understand which factors contribute to this large variability. The variability in the area burned is caused primarily by the choice of land cover and meteorological data and the variability is higher in the southern hemisphere (Figure 8, panels d and f). An interactive N cycle does not affect the zonal distribution of area burned and fire CO₂ emissions (Figures 8 and A10) as much. The reason both area burned and fire CO₂ emissions are affected by the choice of land cover is because the ESA CCI land cover has higher grass area and, as a result, it yields higher area burned and fire CO₂ emissions since a larger area is burned for grasses than for trees in the model. The choice of driving meteorological data is a factor in the area burned and our simulations show that the use of GSWP3 meteorological forcing yields a higher area burned than the CRU-JRA data. In particular

wind speed, which determines the rate of spread of fire in CLASSIC, is much higher in the GWSP3 than in the CRU-JRA meteorological data. Globally-averaged land wind speed (excluding Greenland and Antarctica) in GSWP3 data is 6.1 m/s compared to 3.4 m/s in the CRU-JRA data for the period 2000-2016.

Table 3: Simulated energy, water, and carbon cycle quantities considered in this study sorted according to their coefficient of variation. The quantities are listed from the most variable at the top to the least variable at the bottom. The coefficient of variation is based on annual values averaged over the 1997-2016 period across the eight simulations. The last column shows the dominant source of variability for each model simulated quantity.

Energy, water, or carbon cycle quantities	Coefficient of variation	Dominant source of variability
Area burned (million km ²)	0.24	Land cover
Fire CO ₂ emissions (Pg C/year)	0.21	Land cover
Soil carbon mass (Pg C)	0.21	The inclusion or the absence of the N cycle
Vegetation carbon mass (Pg C)	0.16	The inclusion or the absence of the N cycle
Runoff (1000 km ³ /year)	0.13	Meteorological forcing
Leaf area index (m ² /m ²)	0.11	The inclusion or the absence of the N cycle
Heterotrophic respiration (Pg C/year)	0.10	Land cover
Gross primary productivity (Pg C/year)	0.07	Land cover
Sensible heat flux (W/m ²)	0.07	Meteorological forcing
Autotrophic respiration (Pg C/year)	0.04	Land cover
Latent heat flux (W/m ²) / Evapotranspiration (1000 km ³ /year)	0.05	Meteorological forcing
Net longwave radiation (W/m ²)	0.03	Meteorological forcing
Soil moisture in the top 1m soil layer (mm)	0.02	Meteorological forcing
Albedo for shortwave radiation (fraction)	0.008	The inclusion or the absence of the N cycle
Net shortwave radiation (W/m ²)	0.006	Meteorological forcing
Soil temperature in the top 1m soil layer (°C)	0.004	Meteorological forcing


4.2.3 Coefficient of variation summary

Table 3 shows the energy, water, and C-related quantities considered so far but also leaf area index and albedo and lists them from the most variable at the top to the least variable at the bottom according to their coefficient of variation. The area burned is found to be the most variable quantity and soil temperature is the least variable quantity. Table 3 also shows the most dominant source of variability for each simulated quantity: land cover, meteorological forcings, or the inclusion or absence of an interactive N cycle. Net atmosphere-land CO₂ flux (or net biome productivity) and ground heat flux are not included in Table 3 because these fluxes are calculated as the difference of larger fluxes and as a result, their values are closer to zero which yields a large value of the coefficient of variation.

Overall, the results presented so far illustrate that different model simulated quantities are sensitive to different forcings and model versions. As such it is not advisable to tune a model to match observations when driven with a specific forcing data set.

4.2.4 Net biome productivity

Figure A11 shows the spread in the time series of annual global net atmosphere-land CO₂ flux and their zonally-averaged values across the eight simulations averaged over the 1995-2016 period from each simulation. The global net atmosphere-land CO₂ flux or net biome productivity (NBP) is considered a critical determinant of the performance of LSMs, and is treated as such by TRENDY, because this flux ultimately affects the changes in the atmospheric CO₂ burden. TRENDY requires that LSMs simulate a terrestrial C sink for the decades of the 1990s to the present to be considered for inclusion in the TRENDY ensemble.



680 Figure 9: Time series of global net atmosphere-land CO₂ flux (over all land area excluding
 681 Greenland and Antarctica) (panels a, c, and e) and its zonally-averaged values (panels b, d, and f)
 682 averaged over the four ensemble members that are driven with and without an interactive N
 683 cycle (panels a, b), driven with the GLC 2000 and ESA CCI based land cover (panels c, d), and
 684 driven with GSWP3 and CRU-JRA meteorological data (panels e, f). The thin lines for the time
 685 series show the individual years and the thick lines show their 11-year moving average. Model
 686 values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods,
 687 and their difference, are also shown for panels (a), (c), and (e).

Figure A11 also shows the estimates of global net atmosphere-land CO₂ flux from the participating TRENDY models in grey boxes with mean and shaded ranges for the decades from the 1960s to 2010s from the Global Carbon Project (Friedlingstein et al., 2022). Positive values in Figure A11 indicate a C sink over land and negative values a C source to the atmosphere. In Figure A11a, all eight simulations reported here would qualify for inclusion in the TRENDY ensemble since they all simulate a terrestrial C sink from the 1990s to the present day. Before 1960, since the atmospheric CO₂ concentration is not high enough, the model yields both a land C sink and source in response to interannual variability in meteorological data. In addition, the time series of global NBP from all eight simulations lie within the uncertainty range of reported estimates from the Global Carbon Project. Figure A11a suggests that based on global NBP, at least, it is not possible to exclude any of the eight simulations. In Figure A11b, zonally-averaged NBP averaged over the 1997-2016 period from each of the eight simulations mostly lie within the range of NBP simulated by models that participated in TRENDY 2020. CLASSIC simulates a C sink at northern high latitudes consistent with TRENDY models but it simulates a C sink on the stronger side of TRENDY models in the southern tropics (0° - 20°S). This is likely because CLASSIC is known to simulate low C emissions associated with LUC most of which are generated in tropical regions (Asaadi and Arora, 2021).


Figure 9 provides additional insights into the effect of different forcings on the simulated NBP. In Figure 9, averaged over the 1997-2016 period, an interactive N cycle leads to a somewhat weaker C sink (panel a, 0.98 vs. 1.11 Pg C/yr), the choice of the ESA CCI based land cover leads to a somewhat stronger C sink (panel c, 1.14 vs 0.94 Pg C/yr), and the choice of the GSWP3 meteorological data leads to a much weaker C sink (panel e, 0.74 vs 1.33 Pg C/yr) than the CRU-

JRA meteorological data. In Figure 9, panels a and b, the largest difference between the model versions with and without the N cycle occurs in the tropics ($\sim 5^{\circ}\text{N} - 20^{\circ}\text{S}$) where an interactive N cycle leads to a weaker C sink. There are differences in zonally-averaged NBP with and without the N cycle south of 45°S but the land area below this latitude is small so the averages are calculated over only a few grid cells. The choice of the land cover (Figure 9, panels c and d) does not substantially change the distribution of the zonally-averaged values of NBP although, as noted above, the choice of ESA CCI based land cover leads to a somewhat stronger C sink. Finally, the choice of the GSWP3 meteorological forcing leads to a weaker C sink at most latitudes (Figure 9, panels e and f).

4.3 Automated benchmarking

Figure 10 plots the overall score, S_{overall} , against benchmark scores for several of the energy, water, and C cycle related variables. AMBER does not yet evaluate N cycle related variables for which observations are more scarce than for C cycle related variables. The whiskers show the range in the overall score both for the benchmark and model scores. The vertical whiskers show the range of eight model scores when a given variable from all eight model simulations is compared to an observation-based data set. The horizontal whiskers show the range when three or more observation-based datasets are compared to each other. When only two observation-based data sets are compared to each other there is only one benchmark score, and therefore there is no range. The range in model scores comes from the eight simulations, and the range in benchmark scores comes from the different observation-based data sets. Figure 10 shows that typically as the benchmark scores increase so do the overall model scores for a given quantity.

731 This indicates that uncertainty in observation-based estimates themselves leads to a poor
 732 agreement between observations and model-simulated quantities.




733
 734 Figure 10: Comparison of benchmark scores with model overall scores for a range of energy-,
 735 water-, and carbon-related quantities. The whiskers indicate the range for benchmark scores
 736 across different observation-based data sets and the range across the eight model simulations
 737 for the overall model scores. The quantities in panel (a) are ALBS (surface albedo), RSS (net
 738 shortwave radiation), RLS (net longwave radiation), and RNS (net radiation). Quantities in panel
 739 (b) are HFLS (latent heat flux), HFSS (sensible heat flux), HFG (ground heat flux), MRRO (runoff),
 740 and SNW (snow water equivalent). Quantities in panel (c) are GPP (gross primary productivity),
 741 NEE (net ecosystem exchange), NBP (net biome productivity), AGB (aboveground biomass), CSOIL
 742 (soil carbon mass), BURNT (area burned), and LAI (leaf area index).

743
 744 For energy and water fluxes scores (panels a and b) the model overall scores lie around the 1:1
 745 line indicating that model scores are generally as good as the benchmark scores, except for

surface albedo (ALBS), runoff (MRRO), ground heat flux (HFG), and comparison against one observation-based estimate of snow water equivalent which lie below the 1:1 line. For C cycle related variables most scores lie somewhat below the 1:1 line indicating that simulated quantities do not agree as well with observations as observations agree among themselves. The lower benchmark score for soil C (panel c) is because the SoilGrids250m (SG250m) data and the Harmonized World Soil Database (HWSD) do not agree well amongst themselves because the SG250m soil C data includes peatlands and permafrost C at high latitudes while the HWSD data does not (see Figure 11b). Since the version of CLASSIC used here does not represent peatlands and permafrost C it compares better with the HWSD data than with the SG250m data. In the case of soil C, the choice of HWSD data for comparison against model values is obvious. However, for other variables, it may not always be obvious which observation-based estimate is more appropriate or better for comparison against model results. The uncertainty in forcing data sets and in observation-based estimates, against which model results are evaluated, implies that even a perfect model cannot be evaluated to its fullest extent.

Figure 11 shows the zonal distribution of vegetation C mass, LAI, area burnt, GPP, and fire CO₂ emissions (which constitute standard output from AMBER) and illustrates how AMBER compares the spread across the simulations indicated by 50%, 80%, and 100% shading against observation-based estimates. The black and shades of grey indicate the model mean and the spread across the eight model simulations, respectively, and the thick lines in other colours show the mean values of observation-based estimates. The time period over which observations and model quantities are averaged is chosen to be the same. In Figure 11a, for aboveground biomass, the GEOCARBON data set uses one product for the extratropics and another for the tropics to

create a global aboveground biomass product. The Zhang product (Zhang and Liang, 2020) is based on the fusion of multiple gridded biomass datasets for generating a global product. Both products are described in detail in Seiler et al. (2022). The model results generally compare better with the Zhang product outside the 10°N to 10°S region but with the GEOCARBON product within this region. The values to the south of 40°S are generally less reliable because of the little vegetated land area below this latitude. In Figure 11b, the model simulated values for soil organic C compare better with the HWSD dataset compared to the SG250m data for reasons mentioned in the previous paragraph. Simulated leaf area index (Figure 11c) and gross primary productivity (Figure 11e) generally compare well their observation-based estimates. The simulated area burned (Figure 11d) and fire emissions (Figure 11f) also compare well with observation-based estimates except that the model is not able to capture the small area burned and emissions at northern high latitudes between around 50°N to 70°N. Figures A12 and A13 compare zonally averaged values of other simulated quantities with observation-based estimates used in the AMBER framework. Together Figures 11, A12, and A13 illustrate that the model is overall able to capture the latitudinal distribution of most land surface quantities.



785 Figure 11: Zonally-averaged values of aboveground biomass (a), soil carbon mass (b), leaf area
786 index (c), fractional area burnt (d), gross primary productivity (e), and fire CO₂ emissions (f) from
787 the eight simulations summarized in Table 1. The model results are shown as their mean (black)
788 and the spread across the eight simulations indicated by 50%, 80%, and 100% ranges in different
789 shades of grey. The observation-based estimates used in AMBER to calculate scores are shown
790 in coloured lines.




Figure 12: Summary of difference in overall scores for model simulated quantities and combinations for which the differences are statistically significant. The scores in parentheses for each quantity are the average scores across the eight simulations and provide context. The error bars denote the 95% confidence interval as explained in the text.

Since overall scores are available for all eight simulations for model quantities that are compared to observations it is possible to evaluate how an interactive N cycle, and the choice of meteorological data and land cover data affect model performance. Figure 12 summarizes the difference in overall scores for model quantities and combinations for which the differences are statistically significant at the 5% level based on Tukey's test (Tukey, 1977). The score indicated in parentheses for each quantity is the average score across the eight simulations and provides context. For example, when evaluating the effect of change in land cover for NEE the use of the GLC 2000 based land cover, compared to the use of the ESA CCI based land cover, degrades the average score for net ecosystem exchange by about 0.02 given that the average score for net

ecosystem exchange in 0.53. The error bars on the value 0.02 denote the 95% confidence interval and in this case are calculated by differencing four simulations that use the GLC 2000 based land cover versus four simulations that use the ESA CCI based land cover. The use of the GLC 2000 based land cover on the other hand slightly improves scores for ecosystem respiration and liquid soil moisture. The use of GSWP3 data improves model scores for net shortwave, longwave, and total radiation, for sensible and ground heat flux but degrades the overall score for area burned, soil moisture, and more so for snow water equivalent. Finally, an interactive N cycle slightly improves model performance for area burned and fire CO₂ emissions (due to improved aboveground biomass in the tropics) but degrades it for ecosystem respiration, GPP, and net ecosystem exchange. The inclusion of an interactive N cycle changes $V_{c,max}$ to a prognostic variable for each PFT as opposed to being specified based on observations. This is analogous to running an atmospheric model with a fully dynamic 3-dimensional ocean as opposed to using specified sea surface temperatures (SST) and sea ice concentrations (SIC). Using a dynamic ocean allows future projections (since future SSTs and SICs are not known) but invariably degrades a model's performance for the present day since simulated SSTs and SICs will have their biases. Similarly, using an interactive N cycle allows to project future changes in $V_{c,max}$ (based on changes in N availability) but also degrades CLASSIC's performance for the present day since simulated $V_{c,max}$ has its own biases. Overall, the model performance is most affected by the choice of the driving meteorological data for water and energy fluxes, and by the inclusion or absence of an N cycle and by the choice of land cover for carbon-cycle related state variables and fluxes.

5. Conclusions

The response of the terrestrial biosphere over the historical period has been driven primarily by four global change drivers – increasing atmospheric CO₂, changing climate, LUC, and N deposition and fertilizer application. Our framework allows us to evaluate how a land surface model responds to increasing atmospheric CO₂, changing climate, and anthropogenic N additions to the coupled soil-vegetation system is dependent on two driving meteorological data sets, two land cover representations, and the two model variations (with and without an interactive N cycle). However, the framework used here does not quantify the uncertainty associated with LUC over the historical period since we use only one reconstruction of increasing crop area over the historical period. These results help draw three primary conclusions. First, even if the observations and models were perfect (including their structure and their parameterizations) the uncertainty associated with driving meteorological data and geophysical fields makes it difficult to evaluate LSMs. The uncertainty in global scale driving data implies that a model can never be truly evaluated to its fullest extent. Model results can only be as good as the data that are used to force them and therefore even a perfect model cannot yield perfect results.

Second, model tuning when driving the model with a single set of forcings and evaluating it against a single set of observations is likely not a fruitful exercise. Models should not be tuned to a single set of driving data and rather their performance must be evaluated against a range of available observations in light of the uncertainty associated with driving data and the uncertainty associated with observations. A model's ability to reproduce a given single set of observations when driven with a single set of driving data is not a true measure of its success. Here again, a

perfect model driven by perfect forcing data cannot be truly evaluated to its fullest extent since observations themselves have uncertainties.

Third, with the caveat that our framework uses only one reconstruction of increase in crop area over the historical period, the response of a model expressed in terms of net atmosphere-land CO₂ flux to perturbation in meteorological, CO₂, and LUC forcing over the historical period appears to be largely independent of its pre-industrial state as simulated here. The pre-industrial soil and vegetation C mass for the eight simulations considered here vary between 1035 ± 195 Pg C and 405 ± 58 Pg C (mean \pm standard deviation), respectively. Both pre-industrial and present-day vegetation and soil C pools explain only about 2% to 7% of the variability in simulated net atmosphere-land CO₂ flux (Figure A11) over the 1997-2016 period of each of the eight simulations. The net atmosphere-CO₂ flux from all eight simulations for the period the 1960s to 2000s is found to lie within the uncertainty range provided by the GCP (Friedlingstein et al., 2022). Given the current uncertainty in net atmosphere-land CO₂ flux, it is therefore not possible to exclude any of the eight simulations at least on this basis. The finding that a transient response of a model is independent of its pre-industrial state is also consistent with land components of CMIP6 models. Arora et al. (2020) analyzed results from CMIP6 simulations in which atmospheric CO₂ increases at a rate of 1% per year from the year 1850 until CO₂ quadruples from ~285 to ~1140 ppm. They found that the C-concentration and C-climate feedback parameters for the land component of CMIP6 models do not depend on the absolute values of their vegetation and soil C pools but rather how a given model responds to changes in atmospheric CO₂ and the associated change in temperature. This conclusion is perhaps somewhat comforting in that while pre-industrial states of LSMs may be different from their true

871 observed states they still have the ability to reproduce net atmosphere-land CO₂ flux over the
872 historical period that is consistent with current observation-based estimates. Clearly, this
873 reasoning does not apply if pre-industrial vegetation or soil C mass are zero. One reason why
874 present day net atmosphere-land CO₂ flux is independent of a LSM's pre-industrial state is
875 because the model is first spun up to equilibrium conditions and then forced with time-variant
876 forcings. However, successful reproduction of atmosphere-land CO₂ fluxes over the historical
877 period is no guarantee that future projections from LSMs are reliable.

878 The ensemble-based approach used here also allows for the evaluation of the effect of a
879 given meteorological forcing and land cover, and the effect of an interactive N cycle on model
880 simulated quantities in a robust manner. Ensemble averages of simulations that use the CRU-JRA
881 and GSWP3 meteorological forcing show that the use of the GSWP3 meteorological forcing yields
882 lower evapotranspiration (latent heat flux), higher runoff, higher sensible heat flux, higher
883 burned area, and a weaker land C sink for the present day compared to when the CRU-JRA
884 meteorological forcing is used. The use of the ESA CCI land cover leads to higher soil C, higher
885 GPP, and higher area burned primarily because of the larger grass area when land cover is based
886 on the ESA CCI product compared to the GLC 2000 product. The use of the ESA CCI based land
887 cover also leads to a slightly weaker land C sink for the present day. Finally, the comparison of
888 simulations with and without the N cycle averaged over all meteorological data and land cover
889 combinations allows us to identify the effect of the N cycle. Simulated vegetation C mass and GPP
890 are lower in the model version with the interactive N cycle. In particular, we found that the
891 somewhat low productivity at high latitudes, when the N cycle is turned on, leads to relatively
892 large differences in soil C at high latitudes regardless of the meteorological data or land cover

being used to drive the model. Although, this is not the reason for differences in net atmosphere-land CO₂ flux between models with and without N cycling: as mentioned above present-day net atmosphere-land CO₂ flux is independent of both the pre-industrial and present-day vegetation and soil C pools. Given the knowledge about the effect of N cycling on model behaviour, the reasons can now be investigated to further improve the N cycle component of CLASSIC.

It is logical to assume that the results presented here are sensitive to the horizontal resolution of the model. Both forcing data that are used to drive the model, and observations against which model results are compared, are regridded to be consistent with the model's spatial resolution. For example, at the scale of a few meters, meteorological variables measured at a given site will indeed be less uncertain than their spatially-averaged values say for a 2.81° grid cell. Similarly, observations at a scale of a few meters for soil C and/or vegetation C mass will also likely be more certain than their values at large spatial scales. This is one reason why AMBER uses both gridded and in-situ observation-based estimates to calculate its scores. Fluxes of latent and sensible heat, on the other hand, may not be any more certain at a given site than over large spatial scales. This is because of the problems associated with energy budget closure (Mauder et al., 2020) which, at the point scale, prevent the sum of annual latent and sensible heat flux to be equal to net radiation (average of ground heat fluxes is close to zero at an annual time scale).

LSMs have become increasingly complex over the years and so has the requirement for forcing data to drive these models. The evaluation of LSMs has also become complex as the models now generate a multitude of variables that must be evaluated against their observation-based estimates. Estimates of observation-based data to evaluate models, and the availability of forcing data, have also increased. Given the uncertainties associated with model inputs, model

915 structure, and observation-based data, it is unrealistic to expect LSMs to perfectly reproduce
916 observations for large-scale global simulations. It is not known *a priori* which model structure,
917 forcing data sets, and observation data sets are better. Driving data including meteorological data
918 sets and land cover representations may be more realistic in some parts of world and less in
919 others. Observation-based data sets also have their limitations and attributes which may make
920 them better or ill-suited for comparison with a given model. A more robust model evaluation
921 must therefore take into account the uncertainties both in the forcing and observation-based
922 data. A comprehensive and robust model evaluation can be performed by comparing multiple
923 model realizations against multiple observation-based data sets.

924

925 **References**

- 926 Agustí-Panareda, A., Diamantakis, M., Massart, S., Chevallier, F., Muñoz-Sabater, J., Barré, J., Curcoll, R.,
 927 Engelen, R., Langerock, B., Law, R. M., Loh, Z., Morguí, J. A., Parrington, M., Peuch, V.-H., Ramonet, M.,
 928 Roehl, C., Vermeulen, A. T., Warneke, T., and Wunch, D.: Modelling CO₂ weather – why horizontal
 929 resolution matters, *Atmos Chem Phys*, 19, 7347–7376, 2019.
- 930 Arora, V. K. and Boer, G. J.: A parameterization of leaf phenology for the terrestrial ecosystem
 931 component of climate models, *Glob. Change Biol.*, 11, 39–59, [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2004.00890.x)
 932 2486.2004.00890.x, 2005.
- 933 Arora, V. K. and Boer, G. J.: Simulating Competition and Coexistence between Plant Functional Types in a
 934 Dynamic Vegetation Model, *Earth Interact.*, 10, 1–30, 2006.
- 935 Arora, V. K. and Melton, J. R.: Reduction in global area burned and wildfire emissions since 1930s
 936 enhances carbon uptake by land, *Nat. Commun.*, 9, 1326, <https://doi.org/10.1038/s41467-018-03838-0>,
 937 2018.
- 938 Arora, V. K., Boer, G. J., Christian, J. R., Curry, C. L., Denman, K. L., Zahariev, K., Flato, G. M., Scinocca, J.
 939 F., Merryfield, W. J., and Lee, W. G.: The Effect of Terrestrial Photosynthesis Down Regulation on the
 940 Twentieth-Century Carbon Budget Simulated with the CCCma Earth System Model, *J. Clim.*, 22, 6066–
 941 6088, <https://doi.org/10.1175/2009JCLI3037.1>, 2009.
- 942 Arora, V. K., Scinocca, J. F., Boer, G. J., Christian, J. R., Denman, K. L., Flato, G. M., Kharin, V. V., Lee, W.
 943 G., and Merryfield, W. J.: Carbon emission limits required to satisfy future representative concentration
 944 pathways of greenhouse gases, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2010GL046270>, 2011.
- 945 Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp,
 946 L., Boucher, O., Cadule, P., Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, R. A., Hajima, T., Ilyina,
 947 T., Joetzjer, E., Kawamiya, M., Koven, C. D., Krasting, J. P., Law, R. M., Lawrence, D. M., Lenton, A.,
 948 Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire, A., Wu, T., and
 949 Ziehn, T.: Carbon–concentration and carbon–climate feedbacks in CMIP6 models and their comparison
 950 to CMIP5 models, *Biogeosciences*, 17, 4173–4222, <https://doi.org/10.5194/bg-17-4173-2020>, 2020.
- 951 Asaadi, A. and Arora, V. K.: Implementation of nitrogen cycle in the CLASSIC land model, *Biogeosciences*,
 952 18, 669–706, <https://doi.org/10.5194/bg-18-669-2021>, 2021.
- 953 Avitabile, V., Herold, M., Heuvelink, G. B. M., and others: An integrated pan tropical biomass map using
 954 multiple reference datasets, *Glob Chang Biol*, 2016.
- 955 Beven, K. and Binley, A.: The future of distributed models: Model calibration and uncertainty prediction,
 956 *Hydrol. Process.*, 6, 279–298, <https://doi.org/10.1002/hyp.3360060305>, 1992.
- 957 Bonan, G. B. and Doney, S. C.: Climate, ecosystems, and planetary futures: The challenge to predict life
 958 in Earth system models, *Science*, 359, eaam8328, <https://doi.org/10.1126/science.aam8328>, 2018.
- 959 Bonan, G. B., Lombardozzi, D. L., Wieder, W. R., Oleson, K. W., Lawrence, D. M., Hoffman, F. M., and
 960 Collier, N.: Model Structure and Climate Data Uncertainty in Historical Simulations of the Terrestrial

961 Carbon Cycle (1850–2014), *Glob. Biogeochem. Cycles*, 33, 1310–1326,
 962 <https://doi.org/10.1029/2019GB006175>, 2019.

963 Booth, B. B. B., Jones, C. D., Collins, M., Totterdell, I. J., Cox, P. M., Sitch, S., Huntingford, C., Betts, R. A.,
 964 Harris, G. R., and Lloyd, J.: High sensitivity of future global warming to land carbon cycle processes,
 965 *Environ. Res. Lett.*, 7, 024002, <https://doi.org/10.1088/1748-9326/7/2/024002>, 2012.

966 Chuvieco, E., Lizundia-Loiola, J., Pettinari, M. L., Ramo, R., Padilla, M., Tansey, K., Mouillot, F., Laurent,
 967 P., Storm, T., Heil, A., and Others: Generation and analysis of a new global burned area product based on
 968 MODIS 250 m reflectance bands and thermal anomalies, *Earth Syst. Sci. Data*, 10, 2015–2031, 2018.

969 Claverie, M., Matthews, J. L., Vermote, E. F., and Justice, C. O.: A 30+ Year AVHRR LAI and FAPAR Climate
 970 Data Record: Algorithm Description and Validation, *Remote Sens.*, 8, 263, 2016.

971 Collier, N., Hoffman, F. M., Lawrence, D. M., Keppel-Aleks, G., Koven, C. D., Riley, W. J., Mu, M., and
 972 Randerson, J. T.: The International Land Model Benchmarking (ILAMB) System: Design, Theory, and
 973 Implementation, *J. Adv. Model. Earth Syst.*, 10, 2731–2754, <https://doi.org/10.1029/2018MS001354>,
 974 2018.

975 Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason, B. E., Vose, R.
 976 S., Rutledge, G., Bessemoulin, P., Brönnimann, S., Brunet, M., Crouthamel, R. I., Grant, A. N., Groisman,
 977 P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J., Maugeri, M., Mok, H. Y., Nordli, Ø., Ross, T.
 978 F., Trigo, R. M., Wang, X. L., Woodruff, S. D., and Worley, S. J.: The Twentieth Century Reanalysis Project,
 979 *Q. J. R. Meteorol. Soc.*, 137, 1–28, <https://doi.org/10.1002/qj.776>, 2011.

980 Dai, A. and Trenberth, K. E.: Estimates of Freshwater Discharge from Continents: Latitudinal and
 981 Seasonal Variations, *J. Hydrometeorol*, 3, 660–687, 2002.

982 Di Vittorio, A. V., Chini, L. P., Bond-Lamberty, B., Mao, J., Shi, X., Truesdale, J., Craig, A., Calvin, K., Jones,
 983 A., Collins, W. D., Edmonds, J., Hurtt, G. C., Thornton, P., and Thomson, A.: From land use to land cover:
 984 restoring the afforestation signal in a coupled integrated assessment–earth system model and the
 985 implications for CMIP5 RCP simulations, *Biogeosciences*, 11, 6435–6450, [https://doi.org/10.5194/bg-11-](https://doi.org/10.5194/bg-11-6435-2014)
 986 6435-2014, 2014.

987 Di Vittorio, A. V., Mao, J., Shi, X., Chini, L., Hurtt, G., and Collins, W. D.: Quantifying the Effects of
 988 Historical Land Cover Conversion Uncertainty on Global Carbon and Climate Estimates, *Geophys. Res.*
 989 *Lett.*, 45, 974–982, <https://doi.org/10.1002/2017GL075124>, 2018.

990 ESA: Land Cover CCI Product User Guide Version 2 Technical Report, European Space Agency. Available
 991 at http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf, 2017.

992 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of
 993 the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization,
 994 *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

995 Fischer, G., Nachtergaele, F., Prieler, S., van Velthuizen, H. T., Verelst, L., and Wiberg, D.: Global Agro-
 996 ecological Zones Assessment for Agriculture (GAEZ 2008), IIASA and FAO, Laxenburg, Austria and Rome,
 997 Italy, 2008.

998 Fisher, R. A. and Koven, C. D.: Perspectives on the Future of Land Surface Models and the Challenges of
999 Representing Complex Terrestrial Systems, *J Adv Model Earth Syst*, 12, 2020.

1000 Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W.,
1001 Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B., Anthoni, P.,
1002 Barbero, L., Bastos, A., Bastrikov, V., Becker, M., Bopp, L., Buitenhuis, E., Chandra, N., Chevallier, F.,
1003 Chini, L. P., Currie, K. I., Feely, R. A., Gehlen, M., Gilfillan, D., Gkritzalis, T., Goll, D. S., Gruber, N.,
1004 Gutekunst, S., Harris, I., Haverd, V., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Joetzjer, E., Kaplan, J.
1005 O., Kato, E., Klein Goldewijk, K., Korsbakken, J. I., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A.,
1006 Lienert, S., Lombardozzi, D., Marland, G., McGuire, P. C., Melton, J. R., Metzl, N., Munro, D. R., Nabel, J.
1007 E. M. S., Nakaoka, S.-I., Neill, C., Omar, A. M., Ono, T., Peregon, A., Pierrot, D., Poulter, B., Rehder, G.,
1008 Resplandy, L., Robertson, E., Rödenbeck, C., Séférian, R., Schwinger, J., Smith, N., Tans, P. P., Tian, H.,
1009 Tilbrook, B., Tubiello, F. N., van der Werf, G. R., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget
1010 2019, *Earth Syst. Sci. Data*, 11, 1783–1838, <https://doi.org/10.5194/essd-11-1783-2019>, 2019.

1011 Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C.,
1012 Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni,
1013 P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M.,
1014 Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T.,
1015 Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Houghton, R. A., Hurtt,
1016 G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer,
1017 J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J.,
1018 Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., Ono,
1019 T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M.,
1020 Schwinger, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H.,
1021 Tilbrook, B., Tubiello, F., van der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J.,
1022 Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global Carbon Budget 2021,
1023 *Earth Syst. Sci. Data*, 14, 1917–2005, <https://doi.org/10.5194/essd-14-1917-2022>, 2022.

1024 Garrigues, S., Lacaze, R., Baret, F., Morisette, J. T., Weiss, M., Nickeson, J. E., Fernandes, R., Plummer, S.,
1025 Shabanov, N. V., Myneni, R. B., and Others: Validation and intercomparison of global Leaf Area Index
1026 products derived from remote sensing data, *J. Geophys. Res. Biogeosciences*, 113, 2008.

1027 Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries,
1028 R. S.: Assessing variability and long-term trends in burned area by merging multiple satellite fire
1029 products, 7, 1171–1186, 2010.

1030 Giglio, L., Randerson, J. T., and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area
1031 using the fourth-generation global fire emissions database (GFED4), *J. Geophys. Res. Biogeosciences*,
1032 118, 317–328, <https://doi.org/10.1002/jgrg.20042>, 2013.

1033 Harris, I. C.: CRU JRA v2.1: A forcings dataset of gridded land surface blend of Climatic Research Unit
1034 (CRU) and Japanese reanalysis (JRA) data; Jan. 1901 - Dec. 2019, Centre for Environmental Data Analysis,
1035 University of East Anglia Climatic Research Unit,
1036 <https://catalogue.ceda.ac.uk/uuid/10d2c73e5a7d46f4ada08b0a26302ef7>, 2020.

1037 Hegglin, M., Kinnison, D., and Lamarque, J.-F.: Wet and dry NHx and NOy deposition data,
1038 input4MIPs.CMIP6.CMIP.NCAR. Version 2016-11-15, Earth System Grid Federation,
1039 <https://doi.org/10.22033/ESGF/input4MIPs.10448>, 2016.

1040 Hengl, T., Mendes de Jesus, J., Heuvelink, G. B. M., Ruiperez Gonzalez, M., Kilibarda, M., Blagotić, A.,
 1041 Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan,
 1042 R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m:
 1043 Global gridded soil information based on machine learning, *PLOS ONE*, 12, 1–40,
 1044 <https://doi.org/10.1371/journal.pone.0169748>, 2017.

1045 Hobeichi, S., Abramowitz, G., and Evans, J.: Conserving Land-Atmosphere Synthesis Suite (CLASS), *J Clim*,
 1046 2019.

1047 Hornberger, G. M. and Spear, R. C.: Approach to the preliminary analysis of environmental systems, *J*
 1048 *Env. Manage U. S.*, 12:1, 1981.

1049 van den Hurk, B., Kim, H., Krinner, G., Seneviratne, S. I., Derksen, C., Oki, T., Douville, H., Colin, J.,
 1050 Ducharne, A., Cheruy, F., Viovy, N., Puma, M. J., Wada, Y., Li, W., Jia, B., Alessandri, A., Lawrence, D. M.,
 1051 Weedon, G. P., Ellis, R., Hagemann, S., Mao, J., Flanner, M. G., Zampieri, M., Matera, S., Law, R. M., and
 1052 Sheffield, J.: LS3MIP (v1.0) contribution to CMIP6: the Land Surface, Snow and Soil moisture Model
 1053 Intercomparison Project – aims, setup and expected outcome, *Geosci. Model Dev.*, 9, 2809–2832,
 1054 <https://doi.org/10.5194/gmd-9-2809-2016>, 2016.

1055 Hurtt, G. C., Chini, L., Sahajpal, R., Froking, S., Boudirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori,
 1056 S., Klein Goldewijk, K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F., Jungclaus, J., Kaplan, J.
 1057 O., Kennedy, J., Krisztin, T., Lawrence, D., Lawrence, P., Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter,
 1058 B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello, F. N., van Vuuren, D. P., and Zhang, X.:
 1059 Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6,
 1060 *Geosci Model Dev*, 13, 5425–5464, 2020a.

1061 Hurtt, G. C., Chini, L., Sahajpal, R., Froking, S., Boudirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori,
 1062 S., Klein Goldewijk, K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F., Jungclaus, J., Kaplan, J.
 1063 O., Kennedy, J., Krisztin, T., Lawrence, D., Lawrence, P., Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter,
 1064 B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello, F. N., van Vuuren, D. P., and Zhang, X.:
 1065 Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6,
 1066 *Geosci. Model Dev.*, 13, 5425–5464, <https://doi.org/10.5194/gmd-13-5425-2020>, 2020b.

1067 Jacobson, A. R., Schuldt, K. N., Miller, J. B., and Oda, T.: CarbonTracker Documentation CT2019 release,
 1068 2020.

1069 Jung, M., Koirala, S., Weber, U., Ichii, K., Gans, F., Camps-Valls, G., Papale, D., Schwalm, C., Tramontana,
 1070 G., and Reichstein, M.: The FLUXCOM ensemble of global land-atmosphere energy fluxes, *Sci Data*, 6, 74,
 1071 2019.

1072 Jung, M., Schwalm, C., Migliavacca, M., Walther, S., Camps-Valls, G., Koirala, S., Anthoni, P., Besnard, S.,
 1073 Bodesheim, P., Carvalhais, N., and others: Scaling carbon fluxes from eddy covariance sites to globe:
 1074 synthesis and evaluation of the FLUXCOM approach, *Biogeosciences*, 17, 1343–1365, 2020.

1075 Kato, S., Loeb, N. G., Rose, F. G., Doelling, D. R., Rutan, D. A., Caldwell, T. E., Yu, L., and Weller, R. A.:
 1076 Surface Irradiances Consistent with CERES-Derived Top-of-Atmosphere Shortwave and Longwave
 1077 Irradiances, *J Clim*, 26, 2719–2740, 2013.

1078 Kou-Giesbrecht, S. and Arora, V. K.: Representing the Dynamic Response of Vegetation to Nitrogen
 1079 Limitation via Biological Nitrogen Fixation in the CLASSIC Land Model, *Glob. Biogeochem. Cycles*, 36,
 1080 e2022GB007341, <https://doi.org/10.1029/2022GB007341>, 2022.

1081 Kyker-Snowman, E., Lombardozzi, D. L., Bonan, G. B., Cheng, S. J., Dukes, J. S., Frey, S. D., Jacobs, E. M.,
 1082 McNellis, R., Rady, J. M., Smith, N. G., Thomas, R. Q., Wieder, W. R., and Grandy, A. S.: Increasing the
 1083 spatial and temporal impact of ecological research: A roadmap for integrating a novel terrestrial process
 1084 into an Earth system model, *Glob. Change Biol.*, 28, 665–684, <https://doi.org/10.1111/gcb.15894>, 2022.

1085 Lawrence, P. J. and Chase, T. N.: Representing a new MODIS consistent land surface in the Community
 1086 Land Model (CLM 3.0), *J. Geophys. Res. Biogeosciences*, 112, <https://doi.org/10.1029/2006JG000168>,
 1087 2007.

1088 Li, J., Duan, Q., Wang, Y.-P., Gong, W., Gan, Y., and Wang, C.: Parameter optimization for carbon and
 1089 water fluxes in two global land surface models based on surrogate modelling, *Int. J. Climatol.*, 38,
 1090 e1016–e1031, <https://doi.org/10.1002/joc.5428>, 2018a.

1091 Li, W., MacBean, N., Ciais, P., Defourny, P., Lamarche, C., Bontemps, S., Houghton, R. A., and Peng, S.:
 1092 Gross and net land cover changes in the main plant functional types derived from the annual ESA CCI
 1093 land cover maps (1992–2015), *Earth Syst. Sci. Data*, 10, 219–234, [https://doi.org/10.5194/essd-10-219-](https://doi.org/10.5194/essd-10-219-2018)
 1094 2018, 2018b.

1095 Li, X. and Xiao, J.: Mapping Photosynthesis Solely from Solar-Induced Chlorophyll Fluorescence: A Global,
 1096 Fine-Resolution Dataset of Gross Primary Production Derived from OCO-2, *Remote Sens.*, 11, 2563,
 1097 2019.

1098 Liu, Y. Y., Parinussa, R. M., Dorigo, W. A., De Jeu, R. A. M., Wagner, W., Van Dijk, A., McCabe, M. F.,
 1099 Evans, J., and Others: Developing an improved soil moisture dataset by blending passive and active
 1100 microwave satellite-based retrievals, 2011.

1101 Lu, C. and Tian, H.: Global nitrogen and phosphorus fertilizer use for agriculture production in the past
 1102 half century: shifted hot spots and nutrient imbalance, *Earth Syst. Sci. Data*, 9, 181–192,
 1103 <https://doi.org/10.5194/essd-9-181-2017>, 2017.

1104 Mauder, M., Foken, T., and Cuxart, J.: Surface-Energy-Balance Closure over Land: A Review, *Bound.-*
 1105 *Layer Meteorol.*, 177, 395–426, <https://doi.org/10.1007/s10546-020-00529-6>, 2020.

1106 Meiyappan, P. and Jain, A. K.: Three distinct global estimates of historical land-cover change and land-
 1107 use conversions for over 200 years, *Front. Earth Sci.*, 6, 122–139, [https://doi.org/10.1007/s11707-012-](https://doi.org/10.1007/s11707-012-0314-2)
 1108 0314-2, 2012.

1109 Melton, J. R. and Arora, V. K.: Competition between plant functional types in the Canadian Terrestrial
 1110 Ecosystem Model (CTEM) v. 2.0, *Geosci. Model Dev.*, 9, 323–361, 2016a.

1111 Melton, J. R. and Arora, V. K.: Competition between plant functional types in the Canadian Terrestrial
 1112 Ecosystem Model (CTEM) v. 2.0, *Geosci Model Dev*, 9, 323–361, [https://doi.org/10.5194/gmd-9-323-](https://doi.org/10.5194/gmd-9-323-2016)
 1113 2016, 2016b.

1114 Melton, J. R., Arora, V. K., Wisernig-Cojoc, E., Seiler, C., Fortier, M., Chan, E., and Teckentrup, L.: CLASSIC
 1115 v1.0: the open-source community successor to the Canadian Land Surface Scheme (CLASS) and the
 1116 Canadian Terrestrial Ecosystem Model (CTEM) – Part 1: Model framework and site-level performance,
 1117 *Geosci. Model Dev. Discuss.*, 2019, 1–40, <https://doi.org/10.5194/gmd-2019-329>, 2019.

1118 Mortimer, C., Mudryk, L., Derksen, C., Luoju, K., Brown, R., Kelly, R., and Tedesco, M.: Evaluation of
 1119 long-term Northern Hemisphere snow water equivalent products, *The Cryosphere*, 14, 1579–1594,
 1120 2020.

1121 Mudryk, L.: Historical gridded snow water equivalent and snow cover fraction over Canada from remote
 1122 sensing and land surface models, available at [http://climate-scenarios.canada.ca/?page=blended-snow-](http://climate-scenarios.canada.ca/?page=blended-snow-data)
 1123 [data](http://climate-scenarios.canada.ca/?page=blended-snow-data) (last accessed Oct 2022), 2020.

1124 Myneni, R. B., Hoffman, S., Knyazikhin, Y., Privette, J. L., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y.,
 1125 Smith, G. R., Lotsch, A., Friedl, M., Morisette, J. T., Votava, P., Nemani, R. R., and Running, S. W.: Global
 1126 products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, *Remote Sens*
 1127 *Env.*, 83, 214–231, 2002.

1128 Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J.,
 1129 Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Ribeca, A., van Ingen, C., Zhang, L., Amiro, B.,
 1130 Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M.,
 1131 Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C.,
 1132 Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D.,
 1133 Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns,
 1134 S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti,
 1135 A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S., D’Andrea, E., da Rocha,
 1136 H., Dai, X., Davis, K. J., De Cinti, B., de Grandcourt, A., De Ligne, A., De Oliveira, R. C., Delpierre, N., Desai,
 1137 A. R., Di Bella, C. M., di Tommasi, P., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E.,
 1138 Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B.,
 1139 Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno,
 1140 M., Gharun, M., Gianelle, D., et al.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for
 1141 eddy covariance data, *Sci Data*, 7, 225, 2020.

1142 Peng, S., Ciais, P., Maignan, F., Li, W., Chang, J., Wang, T., and Yue, C.: Sensitivity of land use change
 1143 emission estimates to historical land use and land cover mapping, *Glob. Biogeochem. Cycles*, 31, 626–
 1144 643, <https://doi.org/10.1002/2015GB005360>, 2017.

1145 Poulter, B., Hattermann, F., Hawkins, E., Zaehle, S., Sitch, S., Restrepo-Coupe, N., Heyder, U., and
 1146 Cramer, W.: Robust dynamics of Amazon dieback to climate change with perturbed ecosystem model
 1147 parameters, *Glob. Change Biol.*, 16, 2476–2495, <https://doi.org/10.1111/j.1365-2486.2009.02157.x>,
 1148 2010.

1149 Reusch, A. and Gibbs, H. K.: New IPCC Tier-1 Global Biomass Carbon Map For the Year 2000, Oak Ridge
 1150 National Laboratory, Oak Ridge, Tennessee, 2008.

1151 Rödenbeck, C., Zaehle, S., Keeling, R., and Heimann, M.: How does the terrestrial carbon exchange
 1152 respond to inter-annual climatic variations? A quantification based on atmospheric CO₂ data,
 1153 *Biogeosciences*, 15, 2481–2498, 2018.

1154 Santoro, M., Beaudoin, A., Beer, C., Cartus, O., Fransson, J. E. S., Hall, R. J., Pathe, C., Schmullius, C.,
1155 Schepaschenko, D., Shvidenko, A., Thurner, M., and Wegmüller, U.: Forest growing stock volume of the
1156 northern hemisphere: Spatially explicit estimates for 2010 derived from Envisat ASAR, *Remote Sens*
1157 *Env.*, 168, 316–334, 2015.

1158 Schepaschenko, D., Chave, J., Phillips, O. L., Lewis, S. L., Davies, S. J., Réjou-Méchain, M., Sist, P., Scipal,
1159 K., Perger, C., Herault, B., Labrière, N., Hofhansl, F., Affum-Baffoe, K., Aleinikov, A., Alonso, A., Amani, C.,
1160 Araujo-Murakami, A., Armston, J., Arroyo, L., Ascaranz, N., Azevedo, C., Baker, T., Ba\lazy, R., Bedeau,
1161 C., Berry, N., Bilous, A. M., Bilous, S. Y., Bissiengou, P., Blanc, L., Bobkova, K. S., Braslavskaya, T., Brienén,
1162 R., Burslem, D. F. R. P., Condit, R., Cuni-Sanchez, A., Danilina, D., Del Castillo Torres, D., Derroire, G.,
1163 Descroix, L., Sotta, E. D., d'Oliveira, M. V. N., Dresel, C., Erwin, T., Evdokimenko, M. D., Falck, J.,
1164 Feldpausch, T. R., Folli, E. G., Foster, R., Fritz, S., Garcia-Abril, A. D., Gornov, A., Gornova, M., Gothard-
1165 Bassébé, E., Gourlet-Fleury, S., Guedes, M., Hamer, K. C., Susanty, F. H., Higuchi, N., Coronado, E. N. H.,
1166 Hubau, W., Hubbell, S., Ilstedt, U., Ivanov, V. V., Kanashiro, M., Karlsson, A., Karminov, V. N., Killeen, T.,
1167 Koffi, J.-C. K., Konovalova, M., Kraxner, F., Krejza, J., Krisnawati, H., Krivobokov, L. V., Kuznetsov, M. A.,
1168 Lakyda, I., Lakyda, P. I., Licona, J. C., Lucas, R. M., Lukina, N., Lussetti, D., Malhi, Y., Manzanera, J. A.,
1169 Marimon, B., Junior, B. H. M., Martinez, R. V., Martynenko, O. V., Matsala, M., Matyashuk, R. K., Mazzei,
1170 L., Memiaghe, H., Mendoza, C., Mendoza, A. M., Moroziuk, O. V., Mukhortova, L., Musa, S., Nazimova, D.
1171 I., Okuda, T., Oliveira, L. C., Ontikov, P. V., et al.: The Forest Observation System, building a global
1172 reference dataset for remote sensing of forest biomass, *Sci Data*, 6, 198, 2019.

1173 Seiler, C., Melton, J. R., Arora, V. K., and Wang, L.: CLASSIC v1.0: the open-source community successor
1174 to the Canadian Land Surface Scheme (CLASS) and the Canadian Terrestrial Ecosystem Model (CTEM) –
1175 Part 2: Global benchmarking, *Geosci. Model Dev.*, 14, 2371–2417, 2021a.

1176 Seiler, C., Melton, J. R., Arora, V. K., and Wang, L.: CLASSIC v1.0: the open-source community successor
1177 to the Canadian Land Surface Scheme (CLASS) and the Canadian Terrestrial Ecosystem Model (CTEM) –
1178 Part 2: Global benchmarking, *Geosci. Model Dev.*, 14, 2371–2417, [https://doi.org/10.5194/gmd-14-](https://doi.org/10.5194/gmd-14-2371-2021)
1179 [2371-2021](https://doi.org/10.5194/gmd-14-2371-2021), 2021b.

1180 Seiler, C., Melton, J. R., Arora, V. K., Sitch, S., Friedlingstein, P., Anthoni, P., Goll, D., Jain, A. K., Joetzjer,
1181 E., Lienert, S., Lombardozzi, D., Luyssaert, S., Nabel, J. E. M. S., Tian, H., Vuichard, N., Walker, A. P., Yuan,
1182 W., and Zaehle, S.: Are Terrestrial Biosphere Models Fit for Simulating the Global Land Carbon Sink?, *J.*
1183 *Adv. Model. Earth Syst.*, 14, e2021MS002946, <https://doi.org/10.1029/2021MS002946>, 2022.

1184 Shangguan, W., Dai, Y., Duan, Q., Liu, B., and Yuan, H.: A global soil data set for earth system modeling, *J.*
1185 *Adv. Model. Earth Syst.*, 6, 249–263, <https://doi.org/10.1002/2013MS000293>, 2014.

1186 Slevin, D., Tett, S. F. B., Exbrayat, J.-F., Bloom, A. A., and Williams, M.: Global evaluation of gross primary
1187 productivity in the JULES land surface model v3.4.1, *Geosci. Model Dev.*, 10, 2651–2670,
1188 <https://doi.org/10.5194/gmd-10-2651-2017>, 2017.

1189 Stackhouse, P. W., Jr, Gupta, S. K., Cox, S. J., Zhang, T., Mikovitz, J. C., and Hinkelman, L. M.: The
1190 NASA/GEWEX surface radiation budget release 3.0: 24.5-year dataset, *Gewex News*, 21, 10–12, 2011.

1191 Strahler, A. H., Muller, J., Lucht, W., Schaaf, C., and others: MODIS BRDF/albedo product: algorithm
1192 theoretical basis document version 5.0, MODIS, 1999.

1193 Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V.,
 1194 Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A.,
 1195 Sigmond, M., Solheim, L., von Salzen, K., Yang, D., and Winter, B.: The Canadian Earth System Model
 1196 version 5 (CanESM5.0.3), *Geosci. Model Dev.*, 12, 4823–4873, [https://doi.org/10.5194/gmd-12-4823-](https://doi.org/10.5194/gmd-12-4823-2019)
 1197 2019, 2019.

1198 Tebaldi, C. and Knutti, R.: The use of the multi-model ensemble in probabilistic climate projections,
 1199 *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.*, 365, 2053–2075, <https://doi.org/10.1098/rsta.2007.2076>,
 1200 2007.

1201 Tian, Y., Dickinson, R. E., Zhou, L., and Shaikh, M.: Impact of new land boundary conditions from
 1202 Moderate Resolution Imaging Spectroradiometer (MODIS) data on the climatology of land surface
 1203 variables, *J. Geophys. Res. Atmospheres*, 109, <https://doi.org/10.1029/2003JD004499>, 2004.

1204 Todd-Brown, K. E. O., Randerson, J. T., Post, W. M., Hoffman, F. M., Tarnocai, C., Schuur, E. A. G., and
 1205 Allison, S. D.: Causes of variation in soil carbon simulations from CMIP5 Earth system models and
 1206 comparison with observations, 10, 1717–1736, 2013.

1207 Tukey, J. W.: *Exploratory Data Analysis*, Addison-Wesley, Reading, MA, 1977.

1208 Verger, A., Baret, F., and Weiss, M.: Near real-time vegetation monitoring at global scale, *IEEE J. Sel. Top.*
 1209 *In*, 2014.

1210 Versegny, D. L.: Class—A Canadian land surface scheme for GCMS. I. Soil model, *Int. J. Climatol.*, 11,
 1211 111–133, <https://doi.org/10.1002/joc.3370110202>, 1991.

1212 Versegny, D. L., McFarlane, N. A., and Lazare, M.: Class—A Canadian land surface scheme for GCMS, II.
 1213 Vegetation model and coupled runs, *Int. J. Climatol.*, 13, 347–370,
 1214 <https://doi.org/10.1002/joc.3370130402>, 1993.

1215 Wang, A., Price, D. T., and Arora, V.: Estimating changes in global vegetation cover (1850–2100) for use
 1216 in climate models, *Glob. Biogeochem. Cycles*, 20, <https://doi.org/10.1029/2005GB002514>, 2006.

1217 Wang, L., Bartlett, P., Chan, E., and Xiao, M.: Mapping of Plant Functional Type from Satellite-Derived
 1218 Land Cover Datasets for Climate Models, in: *IGARSS 2018 - 2018 IEEE International Geoscience and*
 1219 *Remote Sensing Symposium*, 3416–3419, <https://doi.org/10.1109/IGARSS.2018.8518046>, 2018.

1220 Wang, L., Bartlett, P., Pouliot, D., Chan, E., Lamarche, C., Wulder, M. A., Defourny, P., and Brady, M.:
 1221 Comparison and Assessment of Regional and Global Land Cover Datasets for Use in CLASS over Canada,
 1222 *Remote Sens.*, 11, 2286, <https://doi.org/10.3390/rs11192286>, 2019.

1223 Wang, L., Arora, V. K., Bartlett, P., Chan, E., and Curasi, S. R.: Mapping of ESA-CCI land cover data to plant
 1224 functional types for use in the CLASSIC land model, *EGUsphere*, 2022, 1–43,
 1225 <https://doi.org/10.5194/egusphere-2022-923>, 2022.

1226 Wieder, W.: *Regridded Harmonized World Soil Database v1.2*, ,
 1227 <https://doi.org/10.3334/ORNLDAAAC/1247>, 2014.

1228 Wu, Z., Ahlström, A., Smith, B., Ardö, J., Eklundh, L., Fensholt, R., and Lehsten, V.: Climate data induced
 1229 uncertainty in model-based estimations of terrestrial primary productivity, *Environ. Res. Lett.*, 12,
 1230 064013, <https://doi.org/10.1088/1748-9326/aa6fd8>, 2017.

1231 Xue, B.-L., Guo, Q., Hu, T., Wang, G., Wang, Y., Tao, S., Su, Y., Liu, J., and Zhao, X.: Evaluation of modeled
 1232 global vegetation carbon dynamics: Analysis based on global carbon flux and above-ground biomass
 1233 data, *Ecol Modell*, 355, 84–96, 2017.

1234 Zhang, Y. and Liang, S.: Fusion of Multiple Gridded Biomass Datasets for Generating a Global Forest
 1235 Aboveground Biomass Map, *Remote Sens.*, 12, 2559, 2020.

1236 Zhang, Y., Xiao, X., Wu, X., Zhou, S., Zhang, G., Qin, Y., and Dong, J.: A global moderate resolution dataset
 1237 of gross primary production of vegetation for 2000–2016, *Sci. Data*, 4, 170165, 2017.

1238

1239 **Code/data availability**

1240

1241 More information about the CLASSIC land surface model and its Fortran code are available at
 1242 https://cccma.gitlab.io/classic_pages/.

1243

1244 AMBER source code as well as the scripts required for reproducing the computational environment,
 1245 including all dependencies on other R-packages, can be found at
 1246 <https://doi.org/10.5281/zenodo.5670387>.

1247

1248 The full suite of results from AMBER for the eight simulations presented in this study can be
 1249 found at <https://cseiler.shinyapps.io/ShinyCLASSIC/>.

1250

1251 **Author contribution**

1252

1253 VA and SKG performed the simulations, and VA wrote the majority of the manuscript. CS performed the
 1254 AMBER related analysis. LW put together the ESA CCI land cover. CS, LW, and SKG provided comments
 1255 on the entire manuscript and also wrote their respective sections.

1256

1257 **Competing interests**

1258

1259 There are no competing interests.

1260

1261 **Acknowledgment**

1262

1263 We thank Joe Melton for providing comments on an earlier version of this paper. We also thank
 1264 Benjamin Bond-Lamberty for taking this paper on as an Associate Editor, and the two anonymous
 1265 reviewers for providing helpful comments which greatly improved this paper.

Appendix

A1: Automated Model Benchmarking R Package (AMBER)

The Automated Model Benchmarking R package quantifies model performance using five scores that assess a model's bias (S_{bias}), root-mean-square-error (S_{rmse}), seasonality (S_{phase}), inter-annual variability (S_{iav}), and spatial distribution (S_{dist}). All scores are dimensionless and range from zero to one, where increasing values imply better performance. The exact definition of each skill score is provided below.

A1.1 Bias Score (S_{bias})

The bias is defined as the difference between the time-mean values of model and reference data:

$$bias(\lambda, \phi) = \overline{v_{mod}}(\lambda, \phi) - \overline{v_{ref}}(\lambda, \phi), \quad (A1)$$

where $\overline{v_{mod}}(\lambda, \phi)$ and $\overline{v_{ref}}(\lambda, \phi)$ are the mean values in time (t) of a variable v as a function of longitude λ and latitude ϕ for model and reference data, respectively. Nondimensionalization is achieved by dividing the bias by the standard deviation of the reference data (σ_{ref}):

$$\varepsilon_{bias}(\lambda, \phi) = \frac{|bias(\lambda, \phi)|}{\sigma_{ref}(\lambda, \phi)} \quad (A2)$$

Note that ε_{bias} is always positive, as it uses the absolute value of the bias. For evaluations against stream flow measurements, the bias is divided by the annual mean rather than the standard deviation of the reference data. This is because we assess streamflow on an annual rather than monthly basis, implying that the corresponding standard deviation is small. The same approach is applied to soil C and vegetation C mass, whose reference data provide a static snapshot in time. For both of these cases, $\varepsilon_{bias}(\lambda, \phi)$ becomes:

$$\varepsilon_{bias}(\lambda, \phi) = \frac{|bias(\lambda, \phi)|}{\overline{v_{ref}}(\lambda, \phi)} \quad (A3)$$

1289

1290 A bias score that ranges from zero to one is calculated next:

$$s_{bias}(\lambda, \phi) = e^{-\varepsilon_{bias}(\lambda, \phi)} \quad (A4)$$

1292 While small relative errors yield score values close to one, large relative errors cause score values
1293 to approach zero. Taking the mean of s_{bias} across all latitudes and longitudes, denoted by a double
1294 bar over a variable, leads to the scalar score:

$$S_{bias} = \overline{\overline{s_{bias}(\lambda, \phi)}} \quad (A5)$$

1296

1297 **A1.2 Root-Mean-Square-Error Score (S_{rmse})**

1298 While the bias assesses the difference between time-mean values, the root-mean-square-error
1299 ($rmse$) is concerned with the residuals of the modeled and observed time series:

$$rmse(\lambda, \phi) = \sqrt{\frac{1}{t_f - t_0} \int_{t_0}^{t_f} (v_{mod}(t, \lambda, \phi) - v_{ref}(t, \lambda, \phi))^2 dt} \quad (A6)$$

1301

1302 where t_0 and t_f are the initial and final time steps, respectively. A similar metric is the centralized
1303 $rmse$ ($crmse$), which is based on the residuals of the anomalies:

1304

$$crmse(\lambda, \phi) = \sqrt{\frac{1}{t_f - t_0} \int_{t_0}^{t_f} [(v_{mod}(t, \lambda, \phi) - \overline{v_{mod}}(\lambda, \phi)) - (v_{ref}(t, \lambda, \phi) - \overline{v_{ref}}(\lambda, \phi))]^2 dt} \quad (A7)$$

1306

1307 The *crmse*, therefore, assesses residuals that have been bias-corrected. Since we already
 1308 assessed the model's bias through S_{bias} , it is convenient to assess the residuals using *crmse* rather
 1309 than *rmse*. In a similar fashion to the bias, we then compute a relative error:

$$1310 \quad \varepsilon_{rmse}(\lambda, \phi) = \frac{crmse(\lambda, \phi)}{\sigma_{ref}(\lambda, \phi)} \quad (A8)$$

1311 scale this error onto a unit interval:

$$1312 \quad s_{rmse}(\lambda, \phi) = e^{-\varepsilon_{rmse}(\lambda, \phi)} \quad (A9)$$

1313 and compute the spatial mean:

$$1314 \quad S_{rmse} = \overline{\overline{s_{rmse}(\lambda, \phi)}} \quad (A10)$$

1315 **A3 Phase Score (S_{phase})**

1316 The skill score S_{phase} assesses how well the model reproduces the seasonality of a variable by
 1317 computing the time difference $\theta(\lambda, \phi)$ between modeled and observed maxima of the
 1318 climatological mean cycle:

$$1319 \quad \theta(\lambda, \phi) = \max(c_{mod}(t, \lambda, \phi)) - \max(c_{ref}(t, \lambda, \phi)) \quad (A11)$$

1320 where c_{mod} and c_{ref} are the climatological mean cycle of the model and reference data,
 1321 respectively. This time difference is then scaled from zero to one based on the consideration that
 1322 the maximum possible time difference is 6 months:

$$1323 \quad s_{phase}(\lambda, \phi) = \frac{1}{2} \left[1 + \cos\left(\frac{2\pi \theta(\lambda, \phi)}{365}\right) \right] \quad (A12)$$

1324 The spatial mean of s_{phase} then leads to the scalar score:

$$1325 \quad S_{phase} = \overline{\overline{s_{phase}(\lambda, \phi)}} \quad (A13)$$

1326

1327 **A4 Inter-Annual Variability Score (S_{iav})**

1328 The skill score S_{iav} quantifies how well the model reproduces patterns of inter-annual variability.

1329 This score is based on data where the seasonal cycle (c_{mod} and c_{ref}) has been removed:

$$1330 \quad iav_{mod}(\lambda, \phi) = \sqrt{\frac{1}{t_f - t_0} \int_{t_0}^{t_f} (v_{mod}(t, \lambda, \phi) - c_{mod}(t, \lambda, \phi))^2 dt} \quad (A14)$$

$$1331 \quad iav_{ref}(\lambda, \phi) = \sqrt{\frac{1}{t_f - t_0} \int_{t_0}^{t_f} (v_{ref}(t, \lambda, \phi) - c_{ref}(t, \lambda, \phi))^2 dt} . \quad (A15)$$

1332

1333 The relative error, nondimensionalization, and spatial mean are computed next:

$$1334 \quad \varepsilon_{iav}(\lambda, \phi) = |iav_{mod}(\lambda, \phi) - iav_{ref}(\lambda, \phi)| / iav_{ref}(\lambda, \phi) \quad (A16)$$

$$1335 \quad s_{iav}(\lambda, \phi) = e^{-\varepsilon_{iav}(\lambda, \phi)} \quad (A17)$$

$$1336 \quad S_{iav} = \overline{\overline{s_{iav}(\lambda, \phi)}} \quad (A13)$$

1337 **A5 Spatial Distribution Score (S_{dist})**

1338 The spatial distribution score S_{dist} assesses how well the model reproduces the spatial pattern of

1339 a variable. The score considers the correlation coefficient R and the relative standard deviation σ

1340 between $\overline{v_{mod}}(\lambda, \phi)$ and $\overline{v_{ref}}(\lambda, \phi)$. The score S_{dist} increases from zero to one, the closer R and

1341 σ approach a value of one. No spatial integration is required as this calculation yields a single

1342 value:

$$1343 \quad S_{dist} = 2(1 + R) \left(\sigma + \frac{1}{\sigma} \right)^{-2} \quad (A19)$$

1344 where σ is the ratio between the standard deviation of the model and reference data:

1345
$$\sigma = \sigma_{\overline{v_{mod}}} / \sigma_{\overline{v_{ref}}} \quad (\text{A20})$$

1346 and $\sigma_{\overline{v_{mod}}}$ and $\sigma_{\overline{v_{ref}}}$ are the standard deviations of the annual mean values from the model and
1347 reference/observation-based data, respectively, and therefore are scalars.

1348 **A6 Overall Score ($S_{overall}$)**

1349 As a final step, scores are averaged to obtain an overall score:

1350
$$S_{overall} = \frac{S_{bias} + 2 S_{rmse} + S_{phase} + S_{iav} + S_{dist}}{1+2+1+1+1} \quad (\text{A21})$$

1351 Note that S_{rmse} is weighted by a factor of two and is an entirely subjective decision but follows
1352 Collier et al. (2018).

1353
1354




Figure A1: Comparison of monthly precipitation (upper panel) and temperature (lower panel) for five global regions (global, north of 25°N, northern and southern tropics, and south of 25°S) from the CRU-JRA and GSWP3 meteorological forcing data sets that are used to drive the CLASSIC model. The global and regional averages exclude Greenland and Antarctica. The legend entries show the annual mean values averaged over the 1997-2016 period. The thin lines show individual years and the thick line is their average.




Figure A2: Comparison of the frequency distribution of daily precipitation between the CRU-JRA and GSWP3 meteorological data sets for three broad regions and the period 1997-2016: a) the Amazonian region, b) the Sahel region, and c) the Midwest United States. The frequency is represented as a percentage of time daily precipitation is between x and $x+1$ mm/day, where x is the value on the x-axis. Panel (d) shows the location of these broad regions. The underlying map in panel (d) is from Google Maps.




Figure A3: Time series of simulated globally-averaged annual soil moisture (a) and soil temperature (b) in the top 1m, global annual evapotranspiration (c), and runoff (d) from the eight simulations summarized in Table 1. The thin lines show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown.




Figure A4: Time series of simulated globally-averaged annual energy fluxes from the eight simulations summarized in Table 1. Panel (a) shows net downward shortwave radiation, panel (b) shows net downward longwave radiation, panel (c) shows latent heat flux, and panel (d) shows sensible heat flux. The thin lines show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown for individual simulations.




Figure A5: Time series of globally-averaged annual net downward longwave and shortwave radiation (over all land area excluding Greenland and Antarctica) averaged over the four ensemble members each that are driven with and without N cycle (panels a, b), driven with GLC 2000 and ESA CCI based land cover (panels c, d), and driven with GSWP3 and CRU-JRA meteorological data (panels e, f). The thin lines show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown.




Figure A6: Time series of simulated global annual vegetation carbon mass (a) and soil carbon (b) from the eight simulations summarized in Table 1. The global totals exclude Greenland and Antarctica. Panels (c) and (d) show the zonally-averaged values of vegetation carbon mass and soil carbon mass over land from the eight simulations averaged over the 1997-2016 period. The thin lines show the individual years and the thick lines show their 11-year moving average in panels (a) and (b). Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown in panels (a) and (b).




Figure A7: Time series of simulated global annual gross primary productivity (GPP) (a), autotrophic respiration (b), and heterotrophic respiration (c) from the eight simulations summarized in Table 1. Panel (d) shows the zonally-averaged values of GPP from the eight simulations averaged over the 1997-2016 period for each simulation. The thin lines show the individual years and the thick lines show their 11-year moving average in panels (a) to (c). Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown in panels (a) to (c).




Figure A8: Time series of global heterotrophic and autotrophic respiration (over all land area excluding Greenland and Antarctica) averaged over the four ensemble members each that are driven with and without an interactive N cycle (panels a, b), driven with the GLC 2000 and ESA CCI based land cover (panels c, d), and driven the with GSWP3 and CRU-JRA meteorological data (panels e, f). The thin lines show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown.





Figure A9: Time series of simulated global annual area burned (a) and fire CO₂ emissions (b) from the eight simulations summarized in Table 1. Panels (c) and (d) show the zonally-averaged area burned and fire CO₂ emissions from the eight simulations averaged over the 1997-2016 period. The thin lines for the time series show the individual years and the thick lines show their 11-year moving average. Model values averaged over the pre-industrial (1851-1860) and present-day (1997-2016) time periods, and their difference, are also shown for panels (a) and (b).



1463

1464 Figure A10: Time series of global fire CO₂ emissions (over all land area excluding Greenland and
 1465 Antarctica) (panels a, c, and e) and their zonally-averaged values (panels b, d, and f) averaged
 1466 over the four ensemble members each that are driven with and without an interactive N cycle
 1467 (panels a, b), driven with the GLC 2000 and ESA CCI based land cover (panels c, d), and driven
 1468 with GSWP3 and CRU-JRA meteorological data (panels e, f). The thin lines for the time series
 1469 show the individual years and the thick lines show their 11-year moving average in panels (a),
 1470 (c), and (e). Model values averaged over the pre-industrial (1851-1860) and present-day (1997-
 1471 2016) time periods, and their difference, are also shown for panels (a), (c), and (e).




Figure A11: Time series of simulated global annual net atmosphere-land CO₂ flux (a) and its zonally-averaged values from the eight simulations summarized in Table 1 averaged over the 1997-2016 period. In panel (a) simulated annual net atmosphere-land CO₂ flux values are compared to the estimates from the Global Carbon Project (Friedlingstein et al., 2022). The thin lines for the time series in panel (a) show the individual years and the thick lines show their 11-year moving average. In panel (b) the simulated zonally-averaged values are compared to the range from 11 models that contributed to the TRENDY 2020 intercomparison.





Figure A12: Zonally-averaged values of soil moisture (a), runoff (b), latent heat flux (c), and sensible heat flux (d) from the eight simulations summarized in Table 1. The model results are shown as their mean (black) and the spread across the eight simulations indicated by 50%, 80%, and 100% ranges in different shades of grey. The observation-based estimates used in AMBER to calculate scores are shown in coloured lines.

1502

1503



1504

1505

1506

1507

1508

1509

1510

1511

1512

Figure A13: Zonally-averaged values of surface albedo (a), snow water equivalent (b), net surface radiation (c), net longwave radiation (d), and net shortwave radiation (e) from the eight simulations summarized in Table 1. The model results are shown as their mean (black) and the spread across the eight simulations indicated by 50%, 80%, and 100% ranges in different shades of grey. The observation-based estimates used in AMBER to calculate scores are shown in coloured lines.