

Hydrologic implications of projected changes in rain-on-snow melt for Great Lakes Basin watersheds

Daniel T. Myers¹, Darren L. Ficklin¹, and Scott M. Robeson¹

¹Department of Geography, Indiana University, Bloomington, Indiana 47405, USA

5 *Correspondence to:* Daniel T. Myers (danmyers901@gmail.com)

Abstract. Rain-on-snow (ROS) melt events reduce the amount of water stored in the snowpack while also exacerbating flooding. The hydrologic implications of changing ROS events in a warming climate, however, are still uncertain. This research used a calibrated and validated Soil and Water Assessment Tool (SWAT) hydrologic model, modified with energy budget equations to simulate ROS melt and forced with a climate model ensemble representing moderate greenhouse-gas concentrations, to simulate changes to ROS melt in the North American Great Lakes Basin from 1960-2069. The changes to ROS events between the historic period (1960-1999) and mid-century (2040-2069) represent an approximately 30% reduction in melt in warmer, southern subbasins, but less than 5% reduction in melt in colder, northern subbasins. Additionally, proportionally more rainfall reduces the formation of snowpacks, with area-weighted winter+spring rain-to-snow ratios rising from approximately 1.5 historically to 1.9 by the mid-21st century. Areas with historic mean winter+spring air temperatures lower than -2 °C have ROS regimes that are resilient to mid-21st century warming projections, but ROS occurrence in areas that have mean winter+spring temperatures near the freezing point are sensitive to changing air temperatures. Also, relationships between changes in the timing of ROS melt and water yield endure throughout the spring but become weak by summer. As the influence of ROS melt events on hydrological systems is being altered in a changing climate, these conclusions are important to inform adaptive management of freshwater ecosystems and human uses in regions of the globe that are sensitive to changes in ROS events.

10
15
20

1 Introduction

Rain-on-snow (ROS) melt events can have important implications for winter floods because of the combined impacts of rainwater and snowmelt runoff (Suriano and Leathers, 2018; Leathers et al., 1998). In places where ROS events are common, they have contributed to the majority of extreme floods, including locations in the United States Northwest, Upper Midwest, Northeast, and Appalachians (Li et al., 2019). ROS events can occur across a wide swath of North America and Eurasia in areas that have substantial snowpack (Pomeroy et al., 2016; Rennert et al., 2009; Rössler et al., 2014; Sui and Koehler, 2001; Ye et al., 2008; Musselman et al., 2018), but their impact on hydrology extends beyond the cold season because snowpack conditions throughout the winter and spring influence the availability of groundwater and stream water later in the year (Blahušíaková et al., 2020; Jenicek et al., 2016; Myers et al., 2021b). Compared to thermally driven snowmelt rates, rainfall-

25

30 based melt events are often more short-lived and intense. As a result, ROS melt produces proportionally more runoff compared to temperature-based snowmelt, with lower rates of infiltration and groundwater recharge (Wilson et al., 1980; Earman et al., 2006). Thus, ROS melt events can lead to snow droughts and reduced water availability after the snow season because of the lost water storage (Harpold et al., 2017; Hatchett and McEvoy, 2018; Blahušiaková et al., 2020; Myers et al., 2021b).

In the North American Great Lakes Basin, ROS melt is associated with over 25% of the most extreme snowmelt 35 events (Suriano, 2020) and has been shown to influence hydrological droughts later in the year (Myers et al., 2021b). ROS events melt an average of 4 cm of snow per event in the Great Lakes Basin, but decreased in frequency by 37% from 1960-2009 (Suriano and Leathers, 2018). ROS melt typically occurs when a mid-latitude cyclone takes a more northerly track, transporting warm, moist air into the basin (Suriano, 2018). At the same time, the snow-water equivalent available in the snowpack is critical and average snow depths in the Basin decreased by 25% from 1960-2009 (Suriano et al., 2019). However, 40 the hydrological impacts of changing ROS melt amounts and frequencies in a transient climate are uncertain, as a decrease in snowpack could limit the amount of ROS melt, but also increase surface runoff from rain on bare ground during cold seasons.

This research combines outputs from an ensemble of downscaled climate models with a version of the Soil and Water Assessment Tool (SWAT) hydrologic model (Arnold et al., 1998) that incorporates a ROS melt modification (Myers et al., 2021b) to simulate climate change impacts to watersheds in the Great Lakes Basin. Our research asks, “How does ongoing 45 climate change alter ROS melt and hydrology in the Great Lakes Basin by the mid-21st century?” This research contributes to scientific knowledge by advancing our understanding of climate change impacts to watersheds, particularly concerning the impacts of ROS melt, improving our ability to manage water quantity and quality into the future. It is important to understand these climate change impacts to sustainably manage rivers and prepare for risks, both within the Great Lakes Basin and in ROS prone regions around the world.

50 **2 Materials and methods**

2.1 Study area

The North American Great Lakes Basin is the Earth’s largest fresh surface water system (Environment Canada and USEPA, 1995), including portions of eight U.S. states and one Canadian province (Figure 1). To the north near Lake Superior, snow cover lasts an average of 180 days, but is as low as 107 days around Lake Erie (Suriano et al., 2019). The Great Lakes 55 Basin is experiencing a rapidly changing climate (Lehner et al., 2006; Environment Canada and USEPA, 1995), with average annual air temperatures having already risen nearly 1 °C since the early 20th century while annual total precipitation has risen approximately 10% (Wuebbles et al., 2019).

2.2 Hydrology simulation

We used the SWAT hydrological model (Arnold et al., 1998) with a modified snowmelt routine to simulate ROS 60 melt. SWAT simulates hydrology with a water balance of inputs (precipitation), exports (evapotranspiration, surface runoff,

groundwater flow, and lateral flow), and soil water storage. SWAT partitions precipitation into rainfall or snowfall based on whether air temperature is above or below a temperature threshold (Fontaine et al., 2002). Groundwater flow was simulated with a shallow aquifer water balance and evapotranspiration was simulated using the Penman-Monteith method (Monteith, 1965; Ritchie, 1972).

65 A full description of the ROS modification, hydrology simulation, calibration, and evaluation can be found in Myers et al. (2021b). In short, the SWAT source code was modified to include an energy budget equation for snowmelt from the SNOW-17 model (Anderson, 2006, 1973) that simulates ROS melt based on a function of air temperature, precipitation, wind, saturated vapor pressure, and atmospheric pressure. For the SWAT ROS model, air temperature and precipitation are based on existing SWAT model inputs, wind effects on ROS are simulated using an average function for ROS melt from turbulent
70 energy transfer, saturation vapor pressure is based on air temperature, and atmospheric pressure is based on elevation, so no additional data inputs are required. Previously, SWAT would simulate snowmelt using a snowpack temperature that was based on air temperature (Fontaine et al., 2002), which would not consider daily ROS melt events. Using a snowmelt module that can simulate ROS melt, such as our SWAT ROS model, reduces error and leads to more accurate hydrological simulations in the Great Lakes Basin due to the more accurate simulation of the timing of snowmelt (Myers et al., 2021b).

75 This study used a calibrated version of the SWAT hydrological model for the Great Lakes Basin previously developed in Myers et al. (2021b), based on historic climate inputs (Maurer et al., 2007). In Myers et al. (2021b), a sensitivity analysis was performed using the PAWN method (Pianosi and Wagener, 2015) that identified 24 sensitive parameters. The model was then calibrated at the daily time step with the A Multi-Algorithm Genetically Adaptive Multiobjective (AMALGAM) algorithm (Vrugt and Robinson, 2007) using 99 stations for streamflow and 50 stations for snowpack snow water equivalent
80 (SWE). SWE was estimated from the gridded North American snow depth dataset (Mote et al., 2018) using a function of snow depth, precipitation, temperature, and time of year (Hill et al., 2019). Nash Sutcliffe Efficiency (NSE; Nash and Sutcliffe, 1970) and the revised Index of Agreement (d_r ; Willmott et al., 2012) were the objective functions for streamflow, while mean absolute error (MAE; Willmott and Matsuura, 2005) was the objective function for snowpack SWE.

The SWAT ROS model for the Great Lakes Basin simulated historic streamflow at the daily time step with an average
85 NSE of 0.38 (with 29% of stations greater than 0.5, 48% greater than 0.4, and a maximum NSE of 0.71) and an average d_r of 0.62 (Myers et al., 2021b). Stations with streamflow NSE > 0.50 at the daily time step were spread throughout the Great Lakes Basin, suggesting that the model was representative of the spatial tendencies in climate forcings and hydrological responses (Figure 2a). Stations that performed well with $d_r > 0.60$ were also distributed across the Basin, which is important because d_r is not as influenced by extremes as NSE (Willmott et al., 2012) and can be a more interpretable indicator of overall model
90 performance (Willmott et al., 2015). Further, the presence of stations not performing as well by NSE could be at least partly explained by the diversity of spatially-distributed hydrological behaviors of the Basin having been simplified in the model's multi-site and multi-objective calibration (Zhang et al., 2008), as well as uncertainties in gridded climate forcings (Maurer et al., 2010; Muche et al., 2020; Stern et al., 2022) and snowpack calibration data (Mote et al., 2018; Hill et al., 2019). Thus, we chose to focus on average amounts of ROS melt over different time periods for this study, without focusing on specific events

95 such as extreme (e.g., 0.95 quantile) water yields. Those extremes may not be represented as reliably in future climate projections given that many of the stations had lower NSE values than we deemed adequate for that purpose, largely because NSE values are sensitive to extremes (Legates and McCabe, 1999; Willmott et al., 2009). Further, projections of climate change impacts on hydrologic extremes are best analyzed using models that focus on extreme flows specifically (Willems et al., 2014).

The model simulated historic snowpack SWE at the daily time step with an MAE of 26 mm (Figure 2a-c). Daily
100 snowpack SWE error across the Basin ranged from <20 mm MAE throughout the southwest subbasins to approximately 40-70 mm MAE in the northeast (Figure 2c). This spatial variation in MAE scaled with the average observed daily snowpack SWE during winter and spring across the Basin, which ranged from approximately 50-100 mm in the southwest subbasins to over 150 mm in the northeast, described in Section 3.4. Thus, subbasins with lower amounts of observed SWE would also have smaller errors than the average of 26 mm. We find this measure of absolute error to be acceptable, particularly considering
105 the errors inherent to the gridded snowpack data we compare against (e.g., spatial averaging and simplification of accumulation and ablation processes during conversion from snow depth; Myers et al., 2021b; Ensor and Robeson, 2008; Hill et al., 2019). Mean absolute errors in modeling the gridded snowpack SWE have been found to vary temporally as well, for instance being 12.7 mm MAE on January 1, 45.1 mm MAE on February 1, 26.8 mm MAE on March 1, and 9.6 mm MAE for April 1, 1978, across the Great Lakes Basin in comparison with station measurements, and proportional to the amount of snowpack on the
110 ground during those days (Myers et al., 2021b). Previous work by Kalin et al. (2010) has stated that arbitrary interpretations of performance metrics for models at small temporal scales should be relaxed compared to what would be expected for models at coarse (e.g., monthly) time steps. Calibrated parameters for this model can be found in Table S1 in Myers et al. (2021b). We also investigated seasonal model performance for simulating snowmelt (water equivalent) during only days when ROS melt was occurring, and found that the SWAT ROS model we use had an MAE of 8.6 mm, 9.4 mm, and 5.8 mm for simulating
115 snowmelt on those days in the winter, spring, and fall, respectively.

2.3 Climate projections

The calibrated hydrological model was forced with 1950-2099 climate projections from downscaled and bias-corrected outputs of the Coupled Model Intercomparison Project Phase 5 (CMIP5) multi-model ensemble (Taylor et al., 2012; US Bureau of Reclamation, 2013; Maurer et al., 2007). These models were downscaled to a 1° latitude/longitude grid (Figure
120 1) and bias-corrected using the Bias-Correction Constructed Analogues method (Maurer et al., 2010), which corrects bias by quantile mapping with historic data (US Bureau of Reclamation, 2013). One-degree grid resolution was chosen for these projections because it matched the resolution of our snowpack data for calibration (Mote et al., 2018). It was important to have our snow data and climate projections at the same resolution so that we could calibrate the model for snow processes at the same scale as the response to air temperature and precipitation. These processes could be sensitive to differences between grids
125 (Rajulapati et al., 2021; Winchell et al., 2013; Myers et al., 2021a).

Global climate models (GCMs) can be a major source of uncertainty when modeling the hydrological impacts of climate change (Wang et al., 2020; Chegwiddden et al., 2019). Thus, nineteen climate models for the Representative

Concentration Pathway (RCP) 4.5 were used to account for variation in climate projections (Table 1). We chose to include nineteen climate models because that was the total number of models using the RCP 4.5 scenario that had been downscaled and bias corrected in the multi-model ensemble (Maurer et al., 2007). RCP 4.5 is a moderate greenhouse-gas scenario that considers long-term changes in emissions, land cover change, the global economy, and climate change mitigation (Thomson et al., 2011). The mean of this multi-model ensemble was used to represent our projection (Christensen et al., 2010), with the standard deviation of the GCM ensemble shown in Figure 3.

2.4 Analyses

Hydrological outputs from the SWAT model were aggregated to the boundaries of regulatory river basins for the United States (Hydrologic Unit Code HUC 8; USGS, 2022) and Canada (Tertiary-Level watersheds; Government of Ontario, 2022) using spatial averaging. Aggregating our subbasins into the regulatory major river basins from the USGS HUC 8 and Ontario Tertiary Level watersheds data allowed us to compare SWAT outputs with the existing basin structure and facilitates comparisons and discussions of our results with other studies.

Results were analyzed by comparing averages and extreme high events among historic (1960-1999) and mid-21st century (2040-2069) time periods at the subbasin scale, based on water years (1 October to 30 September). The mid-21st century period was the focus for informing water resources management and because of better agreement among the models. Simulations of ROS changes generally agree across CMIP5 RCP's (4.5 and 8.5) until mid-century, but then diverge by late-century (Musselman et al., 2021a). Calculations of basin-wide averages were weighted by subbasin area. Seasons were defined as winter (December, January, February), spring (March, April, May), summer (June, July, August), and fall (September, October, November). ROS melt events were defined as days with >1 mm rainfall on >1 mm snowpack SWE and snowmelt occurring, which is a definition that has been previously used to model ROS events and project climate change impacts (Jeong and Sushama, 2018). The ROS center of volume statistic, defined as the day of the water year when half the total volume of ROS melt has passed, was used to examine changes to the timing of ROS melt events, adapted from Hodgkins et al. (2003).

Pearson's correlation was used to evaluate the strength of linear relationships, with "significant" relationships defined as $p < 0.05$. For comparisons between time periods, statistical significance was evaluated by comparing annual area-weighted ensemble-average values for the Great Lakes Basin between the historic (1960-1999, $n=40$ years) and mid-21st century (2040-2069, $n=30$ years) periods using two-tailed unpaired t-tests. Boxplots and percentiles were spatially weighted by major river basin area (Willmott et al., 2007). Finally, the SWAT model outputs for water yield represent the area-averaged water export through the outlet in mm.

3 Results

3.1 Precipitation and air temperature projections

Across the Great Lakes Basin, CMIP5 ensemble average annual precipitation and air temperatures are projected to increase between the historic 1960-1999 period and mid-21st century using RCP 4.5. Spatially averaged annual precipitation increases by 53 mm (6.3%) from 839±63 mm (mean and standard deviation of GCM ensemble) during 1960-1999 to 892±77 mm by the mid-21st century ($p<0.001$), while spatially averaged annual air temperatures increase 2.7 °C from 5.2±0.7 °C during the 1960-1999 period to 7.9±1.0 °C ($p<0.001$; Figure 3a and b). Changes in ensemble mean winter+spring air temperatures are most prominent in northern parts of the Basin, where winter+spring air temperatures are projected to rise approximately 3 °C between the historic 1960-1999 period and mid-21st century using the RCP 4.5 scenario (Figure 4a and b). Using the GCM ensemble mean, changes in mean winter+spring rainfall are strongest in the Lake Superior region, where the amount of rainfall is projected to increase over 40% (Figure 4c and d; Figure 3d). Northern areas experience the least change in ensemble mean winter+spring snowfall amount between the historic 1960-1999 period and mid-21st century, while southern parts of the Basin have a decrease in snowfall over 10% (Figure 4e and f; Figure 3e). Further, our model shows that winter+spring rain to snow ratios over the basin (calculated by dividing the total winter+spring rainfall by total winter+spring snowfall) increase from around 1.5 historically to 1.9 by mid-century ($p<0.001$), which means that proportionally more rainfall could contribute to the declines in snowmelt and snowpack SWE (Figure 3c).

3.2 Snowpack and snowmelt projections

Between the historic 1960-1999 period and mid-21st century, winter months generally see an increase in the amount of ROS melt for individual subbasins due to the increased amount of rainfall (not snowfall) under RCP 4.5 projections. For instance, the area-weighted median (value of a ranked set where half the total area is ranked lower; Willmott et al., 2007) amount of January ROS melt among subbasins increases 59% from 3.2 mm historically to 5.1 mm by mid-21st century (Figure 5a). Similarly, area-weighted median February ROS melt rises 50% from 7.0 mm historically to 10.5 mm in the mid-21st century. In the spring, the amount of ROS melt decreases due to the reduction in snowpack. For instance, the area-weighted median amount of April ROS melt among subbasins decreases 56% from 52.3 mm historically to 22.9 mm by the mid-21st century.

Mean monthly snowmelt (including temperature-based melt and ROS melt), among individual Great Lakes Basin subbasins, is projected to experience a decrease and shift to earlier timing in the spring by the mid-21st century (Figure 5b) using the RCP 4.5 pathway. Historically the maximum snowmelt overall has been in April with an area-weighted median of 85.3 mm, while the March median snowmelt has been less at 44.8 mm. By mid-21st century the median amount of monthly snowmelt among subbasins reaches a maximum at 44.8 mm in March, but drops to only 39.5 mm in April, which is a 54% decrease during April between the two periods.

Changes in the amount of monthly snowmelt among individual subbasins are affected by changes to ROS melt amounts because days that have ROS melt occurrences account for greater than 50% of total snowmelt for most subbasins from December through April (Figure 5c). Temperature-based snowmelt is usually a slower process, while ROS melt events combined with temperature-based melt on these days can rapidly melt snowpack. However, the proportion of melt occurring during December ROS days (compared with all December melt) decreases from an area-weighted median of 71% historically (1960-1999) to 59% by mid-21st century (a decrease of 12%). With warmer temperatures, temperature-based melt can have more of an influence on total snowmelt. The proportion of total annual snowmelt from ROS tends to increase in the northern and eastern parts of the Great Lakes Basin, but decrease in the south and west, between the historic 1960-1999 period and mid-21st century, by about 5% in each direction as temperatures warm (Figure 6a and b). Additionally, snowpack SWE decreases throughout the winter and spring. For instance, by March in the mid-21st century, only 61.6 mm of area-weighted median snowpack SWE is left in the Basin, compared with a median of 104.0 mm historically (a decrease of 41 %) (Figure 5d).

3.3 Rain-on-snow melt projections

ROS melt is affected by the changing climate and there will be different intensities and frequencies of ROS melt events in the future using the RCP 4.5 pathway. Northernmost subbasins near Lake Superior experience the least changes in the annual amount of ROS melt between the historic 1960-1999 period and mid-21st century, with less than a 5% change. However, the central and southern areas of the Basin experience large decreases in annual ROS melt, with the greatest reduction in southern subbasins in Michigan and southern Ontario with a >30% decrease in the amount of annual ROS melt, and a >20% decrease in the frequency of ROS events (Figure 6c-f). Overall, at the major river basin scale using the RCP 4.5 scenario, the ensemble average amount of annual snowmelt during ROS events changes by -42% to +1%, with a basinwide area-weighted average of -22% ($p < 0.001$). Meanwhile, the range among climate projections in the ensemble for this change in basinwide average annual snowmelt during ROS events is -50% to -3%, suggesting that the climate models agree that there will be a reduction in annual ROS melt for the basin overall.

Additionally, northern and central subbasins around eastern Lake Huron and the southern shore of Lake Superior tend to see a slight increase in the annual frequency of ROS events of +5%, while more southern subbasins experience the greatest decrease in frequency around -25% (Figure 6e and f). This is because the northern subbasins, which maintain substantial snowpack throughout the winter and spring and have temperatures well below freezing, have ROS frequencies that are resilient to increases in air temperature (through mid-century), while ROS frequencies in southern subbasins with winter+spring air temperatures around the freezing and melting points are sensitive to even small perturbations in air temperature, with threshold-like responses around these temperature points to the partitioning between rainfall and snowfall.

Following the earlier timing of ROS melt, the center of volume for ROS melt (the day of the water year when half of the total annual ROS melt is passed) decreases between the historic 1960-1999 period and mid-21st century. Historically, the ROS melt center of volume ranged from day 145 (23 February) in the southern part of the Great Lakes Basin to day 207 (26

April) in the northern part (Figure 7a). By the mid-21st century, the ROS melt center of volume becomes earlier and ranges
220 from day 134 (12 February) to day 198 (16 April), which is approximately two weeks earlier (Figure 7b).

3.4 Relationships with climate and snowpack

The cause of the reduction in annual ROS melt across the Basin is largely from a reduction in snowpack SWE due to
the rising air temperatures. Although an increase in ensemble average winter+spring precipitation in major river basins by
+7% to +15% contributes to ROS melt (Figure 3a, Figure 4c-f), its influence is negated by a decrease in the annual amount of
225 winter snowpack SWE in river basins of -10% to -52% (Figure 7c and d). This is due to proportionally more winter rainfall,
as the winter+spring rain to snow ratio increases from 1.55 ± 0.32 (mean and standard deviation of GCM ensemble) during the
historic 1960-1999 period to 1.91 ± 0.31 by mid-21st century (Figure 3c).

Changes in the annual amount of ROS melt are strongly correlated with historic winter+spring snowpack SWE
($r=0.87$, $p<0.001$) and also with the frequency of ROS events ($r=0.82$, $p<0.001$). These relationships are also related to location,
230 with higher latitudes experiencing less of a change in ROS melt and frequency (Figure 8a and b). A decrease in winter+spring
SWE is consistently larger in the southern subbasins, where warming mean temperatures above the freezing and melting points
reduce the ability for snow to accumulate (Figure 7d). The lack of snowpack means that ROS melt events may be unable to
occur as often or as intensely in these southern subbasins as they were able to historically. Changes to the amount of annual
ROS melt and frequency of ROS events are not correlated with historic winter and spring total precipitation amounts (Figure
235 8c and d), as the type of precipitation is more influential, and depends on air temperatures (and thus latitude).

Mean historic winter and spring air temperatures have a strong relationship with changes to ROS melt. Subbasins that
have colder winter+spring air temperatures during the historic 1960-1999 period have weaker changes to the amount of ROS
melt ($r=-0.95$, $p<0.001$; Figure 8e) and frequency of ROS events ($r=-0.72$, $p<0.001$; Figure 8f). For example, subbasins in the
Lake Superior watershed historically have mean winter+spring air temperatures of around -5 °C (Figure 4a). Even with an
240 increase in mean winter+spring air temperatures of $+3$ °C (Figure 4b), temperatures remain cold enough to have a reduced
influence on ROS occurrences for much of the winter and spring. Areas where ROS is most sensitive to the changing climate
are the central and southern river basins of the Great Lakes Basin, where historic mean winter+spring air temperatures were
around 0 °C. Here, perturbations to air temperatures due to climate change can have the greatest effects on ROS melt, leading
to decreases in the amount of annual ROS melt stronger than -30% , as the locations that experience these temperatures are
245 often near the freezing and melting points (Figure 6c and d).

3.5 Relationships with seasonal hydrology

The earlier center of volume for ROS melt has a lagged response on monthly water yields for the major river basins
that lasts throughout the spring but becomes obscured by summer. There is a positive correlation between changes in the ROS
melt center of volume and March ($r=0.42$ ($p<0.001$)) and April ($r=0.70$, $p<0.001$) water yields between the historic 1960-1999
250 period and mid-21st century, due to the influence of ROS melt on daily water yields when snowpacks are actively melting

(Figure 9a and b). In May, this relationship abruptly switches to negative ($r=-0.33$, $p<0.001$), as the reduced impact of ROS means that less water is rapidly exported from the watershed during large melt events, and there is a delayed contribution to water yield (Figure 9c). However, by the summer months of June, July, and August, correlations are weak if any (Figure 9d-f). Changes in summer water yields have much stronger relationships with summer precipitation than they do with ROS, for instance in July ($r=0.74$, $p<0.001$) and August ($r=0.84$, $p<0.001$), obscuring the influence of the timing of ROS melt on summer water yields over time. This suggests that the earlier timing of ROS melt events (by center of volume) is related to hydrology through the spring, although the relationship can be obscured by summer by other factors such as summer precipitation.

4 Discussion

4.1 Implications

It is important to understand the factors affecting spatial variability of ROS changes in the Great Lakes Basin to realize the impacts of this variability on aquatic resources. Spatial variability in ROS melt changes has previously been shown to occur because of differences in latitude, elevation, and atmospheric processes (Pan et al., 2018; Ye et al., 2008; Jeong and Sushama, 2018; Cohen et al., 2015), although there were not large elevation changes in the Great Lakes Basin to observe elevation-based variability at our scale. There are particularly large increases in ROS melt runoff predicted for northeastern North America, but decreases in more southern latitudes due to a decrease of snow cover (Jeong and Sushama, 2018). The frequency of ROS melt events can be affected by latitude because of its association with air temperature, precipitation type (rain or snow), and snow cover (Suriano, 2022). This aligns with our findings for the Great Lakes Basin that the change in ROS melt amounts decreases by mid-century and is strongly related to latitude, with the greatest decreases in southern subbasins where snowpack becomes exhausted, as latitude affects whether mean winter air temperatures will be near the threshold around the freezing and melting points where ROS is most sensitive to changes in climate.

Similarly, an understanding of the temporal factors affecting variability in ROS changes can provide insights to the timing of hydrological impacts. Temporally, the frequency of ROS events in a warming climate has the potential to increase as more rain falls on snowpack, but decline after a warming threshold is reached and the snowpack becomes scarce (Beniston and Stoffel, 2016). For instance, in eastern Russia, the frequency of ROS events has historically increased with warming air temperatures because of more winter rainfall, at a rate of 0.5 to 2.5 events per °C air temperature increase, but future increases could be limited by a lack of snow in warmer regions (Ye et al., 2008). Suriano (2022) has found snowfall amounts to be a dominant control on the frequency of North American ROS melt events.

The earlier timing of ROS melt (and earlier passage of its annual center of volume) has the potential to influence other parts of the hydrologic cycle. In the eastern United States, since 1940, the timing of the center of streamflow volume passing through gages in the winter and spring has become earlier at a rate of 1.6 days per decade, due to increasing air temperatures in snowmelt-driven regions and earlier snowmelt occurrences (Dudley et al., 2017). A similar trend of earlier snowmelt timing has been found for the western United States as well (Stewart et al., 2004; Musselman et al., 2021b). The

earlier timing of snowmelt aligns with what our projections show for the Basin overall, due to winter precipitation increases and ROS, supporting that the earlier center of volume for ROS melt could influence spring water yields. Also, following a 2
285 °C warming scenario, prior research has found that the frequency of ROS melt events are expected to become approximately one month earlier in the eastern United States as cold-season snowfall switches to rainfall (Li et al., 2019), which aligns with our findings for the earlier center of volume for ROS melt over the water year. As the contributions of snowmelt to peak spring water yields become weaker due to the earlier timing of ROS events, vegetation green-up can have a more dominant influence on spring hydrographs in a changing climate (Khodaei et al., 2022).

290 Changing rain to snow ratios can have meaningful impacts to hydrological systems. The rain to snow ratio is important because it was previously found to be the primary avenue for changing air temperatures to affect snowpack in the Sierra Nevada of California, USA, exacerbating runoff during early-season flooding (Huang et al., 2018). As more precipitation falls as rain rather than snow, the size of floods from rainfall and ROS events can far exceed the size of typical snowmelt-driven floods, due to the rapid contributions of rainfall and ROS runoff, with the largest increases being over 2.5 times in size for the western
295 United States (Davenport et al., 2020). The rain to snow ratio can also influence the size and timing of spring snowmelt and summer baseflow (Huntington et al., 2004). Thus, the rain to snow ratio could help explain the earlier COV of ROS melt for the Great Lakes Basin by mid-21st century, since we found that as the basin-average rain to snow ratio increases from approximately 1.5 historically to 1.9 by mid-21st century, the COV of ROS melt occurs two weeks earlier.

The proportion of total snowmelt from ROS or temperature-based melt also has important hydrological impacts.
300 Research in the western United States has found that climate change can decrease the speed at which snowpack melts, as warmer air temperatures mean that there will be bare ground later in the snowmelt season while radiative energy fluxes are high, and more snowmelt occurs during the colder part of winter when energy fluxes are low, causing snowmelt to be a slower process (Musselman et al., 2017). This increase in the proportion of temperature-based melt to total snowmelt reflects a transition to slower, earlier snowmelt, and helps explain the decrease in high spring streamflow that historically influenced
305 hydrological regimes in the Great Lakes Basin (Hodgkins et al., 2007). The decrease in the proportion of total snowmelt from ROS in the Great Lakes Basin could also contribute to groundwater recharge (Earman et al., 2006; Wilson et al., 1980) and the increases in May water yields in subbasins where the annual amount of ROS decreases (Figure 9c), as the water would not have been rapidly exported from the subbasins in earlier ROS events.

310 **4.2 Historic ROS melt**

Previous work by Jeong and Sushama (2018), whose definition of ROS we adopted, produced estimates of historic frequencies of ROS events comparable to ours, approximately 10-20 ROS days per year in the Great Lakes Basin. Also, Jeong and Sushama (2018) report a 1976-2005 average annual amount of ROS runoff of approximately 100 mm or greater throughout the Basin, which is similar to our historic (1960-1999) estimates that were approximately 75 mm annual ROS melt in the
315 southwest part of the Basin and 175 mm in the northeast. Jeong and Sushama (2018) evaluated their model results using

observations and found that spatial patterns in ROS were captured reasonably well, although some errors likely were due to data uncertainties rather than model errors. Using a different definition of an ROS event (air temperature >0 °C and precipitation >5 mm during 2-day extreme snowmelt events), Welty and Zeng (2021) produced far fewer ROS events than we did. Additionally, Suriano (2022) defined an ROS event as a snow depth decrease of at least 1 cm with average daily temperature >0 °C, at least 0.01 cm precipitation, and no more than 2.54 cm snowfall (by depth) the previous day, over a 1960-2009 historic period. With this definition, Suriano (2022) reported a historic frequency of approximately 5 to 15 ROS events per year in the Great Lakes Basin, which is slightly less than our estimate of approximately 10-20 annual ROS events during 1960-1999.

To verify our historic estimates, we identified ROS amounts and frequencies in observed data using the same approach and definition as our GCM-forced SWAT model. The historic climate observations were from Maurer et al. (2007), used in Myers et al. (2021b), and our historic SWE observations were from Myers et al. (2021b), which had been estimated from the daily gridded North American snow depth dataset (Mote et al., 2018), both using the same 1° latitude/longitude grid with 50 evaluation points over the Great Lakes Basin. We found that for historic annual estimates of ROS melt, the mean among the gridded evaluation points for our GCM ensemble was 120 mm, while the mean calculated from observations was 118 mm, which was not a significant difference (two-sample t-test, $p=0.90$). For individual evaluation points, the estimates of annual ROS melt were positively related and had an MAE of 33 mm (Fig. S1a). This suggests that our GCM ensemble produced reasonable estimates of historic ROS melt amounts in the Basin. We also found that the historic observations produced an average annual ROS frequency of 20 days across the evaluation points, which was greater than the mean of 12 days estimated by our GCM ensemble for the points over the historic 1960-1999 period ($p<0.001$). This was because our ROS definition included historic observed events that were the result of natural stochasticity in snowpack SWE amounts (i.e., sporadic daily increases or decreases in the SWE data due to factors such as the timing of measurements at different stations, rather than “clean” modeled melt; Suriano and Leathers, 2017; Suriano, 2022). Thus, our definition underestimated the frequency of ROS days when applied to model data, due to the lack of additional stochastic small melt events identified by the criteria, producing an MAE of 8 days between observations and modeled ROS frequency (Fig. S1b). However, when ROS amounts are accumulated over the season, this issue is remedied (Fig. S1a).

5 Conclusions

Climate change is disrupting rain-on-snow patterns globally, potentially impacting ecosystems, communities, and economies in regions where these events are prevalent. This study used the Soil and Water Assessment Tool (SWAT) rain-on-snow melt (ROS) model, which builds upon SWAT by incorporating energy budget equations to simulate ROS melt (Myers et al., 2021b), to study the impacts of climate change on ROS melt due to altered snowpack, air temperatures, and precipitation. An ensemble of RCP 4.5 climate projections (representing moderate greenhouse gas concentrations) was used to study relationships. Although winter+spring precipitation increases in the Great Lakes Basin by the mid-21st century, compared with

historic 1960-1999 amounts, its influence on ROS melt is limited by an exhausted snowpack with warmer air temperatures, particularly for southern subbasins. Winter+spring rain to snow ratios from the climate input data rise from around 1.5
350 historically to 1.9 by mid-21st century, so proportionally more rainfall decreases snowpack SWE. Changes in ROS melt are positively correlated with snowpack snow water equivalent and winter+spring precipitation.

We find that relationships with ROS patterns and latitude are strong in the Great Lakes Basin, with northern subbasins having air temperatures that remain well below freezing most of the winter+spring and are resilient to temperature increases, while southern subbasins that had mean winter+spring temperatures around freezing historically are more sensitive to changes
355 in air temperatures. The changing temperature directly affects whether snowpack would form or melt, or whether precipitation would be snow or rain. We expect this result of increased sensitivity for ROS changes to apply to cold regions around the globe with average winter+spring air temperatures around 0 °C. With increasing air temperatures, temperature-based snowmelt can have more of an influence on total monthly snowmelt, as the proportion of monthly melt from ROS decreases (e.g. -12% in December) between the historic 1960-1999 period and mid-21st century.

We also find there are temporal relationships with ROS melt timing in the Great Lakes Basin by mid-21st century, as the center of volume (the day of the water year when at least half the total ROS melt volume has passed) becomes earlier by approximately two weeks compared to the historic 1960-1999 period. The temporal scale of impacts from this earlier timing on monthly water yields lasts through the spring (positively correlated in March and April, but negative in May), although these relationships can be obscured by summer because of changing summer precipitation. Investigations projecting the
365 response of extreme water yields to changing ROS conditions in future climates are an additional avenue for future research with meaningful implications for water resources management.

Finally, it is important that future work involve collaborations outside the academic realm so that the findings of climate change impacts to ROS melt can inform management of aquatic resources (Meadow and Owen, 2021) and engage communities with the research (Serreze et al., 2021). Future work could also investigate how changing ROS conditions affect
370 other components of the water balance including groundwater and soil water storage in the Great Lakes Basin. The implications of this work, specifically involving the influence of changing ROS melt on extreme hydrological events and future water availability, as well as the climate-related sensitivities to changing ROS melt, could help prepare managers of ecosystems and human water uses for the climatic changes of the mid-21st century.

Supplementary material

375 Supplementary material for this article is available online for Table S1 and Figure S1.

Funding

This work was supported by the Indiana University Geography Department, William R. Black Fellowship, Indiana University Sustainability Research Development Grant, National Science Foundation [grant numbers DBI-1564806 and CNS-0521433], Indiana University Pervasive Technology Institute, Lilly Endowment, Inc., Indiana METACyt Initiative, and Shared

380 University Research Grants from IBM, Inc. to Indiana University. Any opinions, findings, and conclusions or
recommendations expressed are those of the authors and do not necessarily reflect the views of the National Science
Foundation.

Data availability

The data and SWAT ROS model used in this study are publicly available from Mendeley Data at
385 <http://dx.doi.org/10.17632/bfypd4wpcn.1>.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We thank Ram Neupane, Alejandra Botero-Acosta, and Dan Li for guidance with this study. We also thank Indiana
390 University's University Information Technology Services High Performance Computing team for technical support. We
acknowledge the World Climate Research Program's Working Group on Coupled Modelling, which is responsible for CMIP,
and we thank the climate modeling groups (listed in Table S1 of this paper) for producing and making available their model
output. For CMIP the U.S. Department of Energy's Program for Climate Model Diagnosis and Intercomparison provides
coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth
395 System Science Portals. We also thank Sandra Akkermans and the Integrated Topics in Earth & Environment course at
Wageningen University for valuable comments to improve our manuscript.

Author contribution

Daniel T. Myers: Conceptualization, Methodology, Software, Validation, Formal Analysis, investigation, Data Curation,
Writing – Original Draft, Writing – Review & Editing, Visualization. Darren L. Ficklin: Conceptualization, Methodology,
400 Software, Resources, Writing – Original Draft, Writing – Review & Editing, Supervision, Project Administration, Funding
Acquisition. Scott M. Robeson: Conceptualization, Methodology, Software, Resources, Writing – Original Draft, Writing –
Review & Editing, Visualization.

References

- Anderson, E. A.: National Weather Service river forecast system-snow accumulation and ablation model, in: National Oceanic
405 and Atmospheric Administration Technical Memorandum NWS-HYDR0-17, Washington, DC, USA, 229 pages,
1973.
- Anderson, E. A.: Snow Accumulation and Ablation Model–SNOW-17. NWSRFS User Documentation. 61 pages.

<https://www.wcc.nrcs.usda.gov/ftpref/wntsc/H&H/snow/AndersonSnow17.pdf> (accessed April 10, 2021), 61 pages. pp., 2006.

- 410 Arnold, J. G., Srinivasan, R., Mutiah, R. S., and Williams, J. R.: Large Area Hydrologic Modeling and Assessment Part I: Model Development, *J. Am. Water Resour. Assoc.*, 34, 73–89, <https://doi.org/10.1111/j.1752-1688.1998.tb05961.x>, 1998.
- Arora, V. K., Scinocca, J. F., Boer, G. J., Christian, J. R., Denman, K. L., Flato, G. M., Kharin, V. V, Lee, W. G., and Merryfield, W. J.: Carbon emission limits required to satisfy future representative concentration pathways of
415 greenhouse gases, *Geophys. Res. Lett.*, 38, 2011.
- Beniston, M. and Stoffel, M.: Rain-on-snow events, floods and climate change in the Alps: Events may increase with warming up to 4 °C and decrease thereafter, *Sci. Total Environ.*, 571, 228–236, <https://doi.org/10.1016/j.scitotenv.2016.07.146>, 2016.
- Bentsen, M., Bethke, I., Debernard, J. B., Iversen, T., Kirkevåg, A., Seland, Ø., Drange, H., Roelandt, C., Seierstad, I. A., and
420 Hoose, C.: The Norwegian Earth System Model, NorESM1-M–Part 1: description and basic evaluation of the physical climate, *Geosci. Model Dev.*, 6, 687–720, 2013.
- Bi, D., Dix, M., Marsland, S. J., O’Farrell, S., Rashid, H., Uotila, P., Hirst, A. C., Kowalczyk, E., Golebiewski, M., and Sullivan, A.: The ACCESS coupled model: description, control climate and evaluation, *Aust. Meteorol. Ocean. J.*, 63, 41–64, 2013.
- 425 Blahušáková, A., Matoušková, M., Jenicek, M., Ledvinka, O., Kliment, Z., Podolinská, J., and Snopková, Z.: Snow and climate trends and their impact on seasonal runoff and hydrological drought types in selected mountain catchments in Central Europe, *Hydrol. Sci. J.*, 65, 2083–2096, <https://doi.org/10.1080/02626667.2020.1784900>, 2020.
- Chegwidden, O. S., Nijssen, B., Rupp, D. E., Arnold, J. R., Clark, M. P., Hamman, J. J., Kao, S. C., Mao, Y., Mizukami, N., Mote, P. W., Pan, M., Pytlak, E., and Xiao, M.: How Do Modeling Decisions Affect the Spread Among Hydrologic
430 Climate Change Projections? Exploring a Large Ensemble of Simulations Across a Diversity of Hydroclimates, *Earth’s Futur.*, 7, 623–637, <https://doi.org/10.1029/2018EF001047>, 2019.
- Christensen, J. H., Kjellström, E., Giorgi, F., Lenderink, G., and Rummukainen, M.: Weight assignment in regional climate models, *Clim. Res.*, 44, 179–194, <https://doi.org/10.3354/cr00916>, 2010.
- Cohen, J., Ye, H., and Jones, J.: Trends and variability in rain-on-snow events, *Geophys. Res. Lett.*, 42, 7115–7122,
435 <https://doi.org/10.1002/2015GL065320>, 2015.
- Davenport, F. V., Herrera-Estrada, J. E., Burke, M., and Diffenbaugh, N. S.: Flood Size Increases Nonlinearly Across the Western United States in Response to Lower Snow-Precipitation Ratios, *Water Resour. Res.*, 56, 1–19, <https://doi.org/10.1029/2019WR025571>, 2020.
- Dudley, R. W., Hodgkins, G. A., McHale, M. R., Kolian, M. J., and Renard, B.: Trends in snowmelt-related streamflow timing
440 in the conterminous United States, *J. Hydrol.*, 547, 208–221, <https://doi.org/10.1016/j.jhydrol.2017.01.051>, 2017.
- Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., and

- Benshila, R.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, *Clim. Dyn.*, 40, 2123–2165, 2013.
- 445 Dunne, J. P., John, J. G., Adcroft, A. J., Griffies, S. M., Hallberg, R. W., Shevliakova, E., Stouffer, R. J., Cooke, W., Dunne, K. A., and Harrison, M. J.: GFDL’s ESM2 global coupled climate–carbon earth system models. Part I: Physical formulation and baseline simulation characteristics, *J. Clim.*, 25, 6646–6665, 2012.
- Earman, S., Campbell, A. R., Phillips, F. M., and Newman, B. D.: Isotopic exchange between snow and atmospheric water vapor: Estimation of the snowmelt component of groundwater recharge in the southwestern United States, *J. Geophys. Res. Atmos.*, 111, 1–18, <https://doi.org/10.1029/2005JD006470>, 2006.
- 450 Ensor, L. A. and Robeson, S. M.: Statistical characteristics of daily precipitation: Comparisons of gridded and point datasets, *J. Appl. Meteorol. Climatol.*, 47, 2468–2476, <https://doi.org/10.1175/2008JAMC1757.1>, 2008.
- Environment Canada and USEPA: The Great Lakes: An Environmental Atlas and Resource Book, <https://doi.org/10.1038/184665a0>, 1995.
- Fontaine, T. A., Cruickshank, T. S., Arnold, J. G., and Hotchkiss, R. H.: Development of a snowfall-snowmelt routine for mountainous terrain for the soil water assessment tool (SWAT), *J. Hydrol.*, 262, 209–223, [https://doi.org/10.1016/S0022-1694\(02\)00029-X](https://doi.org/10.1016/S0022-1694(02)00029-X), 2002.
- 455 Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., and Vertenstein, M.: The community climate system model version 4, *J. Clim.*, 24, 4973–4991, 2011.
- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., and 460 Fieg, K.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *J. Adv. Model. Earth Syst.*, 5, 572–597, 2013.
- Government of Ontario: Ontario Watershed Boundaries - Datasets - Ontario Data Catalogue, <https://data.ontario.ca/dataset/ontario-watershed-boundaries>, 2022.
- Harpold, A. A., Dettinger, M., and Rajagopal, S.: Defining snow drought and why it matters, *Eos (Washington. DC.)*, 98, 465 <https://doi.org/10.1029/2017eo068775>, 2017.
- Hatchett, B. J. and McEvoy, D. J.: Exploring the origins of snow drought in the northern Sierra Nevada, California, *Earth Interact.*, 22, 1–13, <https://doi.org/10.1175/EI-D-17-0027.1>, 2018.
- Hill, D. F., Burakowski, E. A., Crumley, R. L., Keon, J., Michelle Hu, J., Arendt, A. A., Wikstrom Jones, K., and Wolken, G. J.: Converting snow depth to snow water equivalent using climatological variables, 13, 1767–1784, 470 <https://doi.org/10.5194/tc-13-1767-2019>, 2019.
- Hodgkins, G. A., Dudley, R. W., and Huntington, T. G.: Changes in the timing of high river flows in New England over the 20th Century, *J. Hydrol.*, 278, 244–252, [https://doi.org/10.1016/S0022-1694\(03\)00155-0](https://doi.org/10.1016/S0022-1694(03)00155-0), 2003.
- Hodgkins, G. A., Dudley, R. W., and Aichele, S. S.: Historical Changes in Precipitation and Streamflow in the U.S. Great Lakes Basin, 1915–2004, United States Geol. Surv. Sci. Investig. Rep. 2007-5118, 37 pages, 2007.
- 475 Hourdin, F., Grandpeix, J.-Y., Rio, C., Bony, S., Jam, A., Cheruy, F., Rochetin, N., Fairhead, L., Idelkadi, A., and Musat, I.:

- LMDZ5B: the atmospheric component of the IPSL climate model with revisited parameterizations for clouds and convection, *Clim. Dyn.*, 40, 2193–2222, 2013.
- Huang, X., Hall, A. D., and Berg, N.: Anthropogenic Warming Impacts on Today’s Sierra Nevada Snowpack and Flood Risk, *Geophys. Res. Lett.*, 45, 6215–6222, <https://doi.org/10.1029/2018GL077432>, 2018.
- 480 Huntington, T. G., Hodgkins, G. A., Keim, B. D., and Dudley, R. W.: Changes in the proportion of precipitation occurring as snow in New England (1949–2000), *J. Clim.*, 17, 2626–2636, 2004.
- Jenicek, M., Seibert, J., Zappa, M., Staudinger, M., and Jonas, T.: Importance of maximum snow accumulation for summer low flows in humid catchments, *Hydrol. Earth Syst. Sci.*, 20, 859–874, <https://doi.org/10.5194/hess-20-859-2016>, 2016.
- 485 Jeong, D. Il and Sushama, L.: Rain-on-snow events over North America based on two Canadian regional climate models, *Clim. Dyn.*, 50, 303–316, <https://doi.org/10.1007/s00382-017-3609-x>, 2018.
- Kalin, L., Isik, S., Schoonover, J. E., and Lockaby, B. G.: Predicting Water Quality in Unmonitored Watersheds Using Artificial Neural Networks, *J. Environ. Qual.*, 39, <https://doi.org/10.2134/jeq2009.0441>, 2010.
- Khodaei, M., Hwang, T., Ficklin, D. L., and Duncan, J. M.: With warming, spring streamflow peaks are more coupled with vegetation green-up than snowmelt in the northeastern United States, *Hydrol. Process.*, 36, e14621, <https://doi.org/10.1002/HYP.14621>, 2022.
- 490 Leathers, D. J., Kluck, D. R., and Kroczyński, S.: The Severe Flooding Event of January 1996 across North-Central Pennsylvania, *Bull. Am. Meteorol. Soc.*, 75, 785–797, [https://doi.org/10.1175/1520-0477\(1998\)079<0785:TSFEOJ>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0785:TSFEOJ>2.0.CO;2), 1998.
- 495 Legates, D. R. and McCabe, G. J.: Evaluating the use of “goodness-of-fit” Measures in hydrologic and hydroclimatic model validation, *Water Resour. Res.*, 35, 233–241, <https://doi.org/10.1029/1998WR900018>, 1999.
- Lehner, B., Verdin, K., and Jarvis, A.: *HydroSHEDS Technical Documentation*, World Wildlife Fund, Washington, DC, 2006.
- Li, D., Lettenmaier, D. P., Margulis, S. A., and Andreadis, K.: The Role of Rain-on-Snow in Flooding Over the Conterminous United States, *Water Resour. Res.*, 55, 8492–8513, <https://doi.org/10.1029/2019WR024950>, 2019.
- 500 Long, M. C., Lindsay, K., Peacock, S., Moore, J. K., and Doney, S. C.: Twentieth-century oceanic carbon uptake and storage in CESM1 (BGC), *J. Clim.*, 26, 6775–6800, 2013.
- Maurer, E. P., Brekke, L., Pruitt, T., and Duffy, P. B.: Fine-resolution climate projections enhance regional climate change impact studies, *Eos, Trans. Am. Geophys. Union*, 88, 504, <https://doi.org/10.1029/2007eo470006>, 2007.
- Maurer, E. P., Hidalgo, H. G., Das, T., Dettinger, M. D., and Cayan, D. R.: The utility of daily large-scale climate data in the assessment of climate change impacts on daily streamflow in California, *Hydrol. Earth Syst. Sci.*, 14, 1125–1138, <https://doi.org/10.5194/hess-14-1125-2010>, 2010.
- 505 Meadow, A. M. and Owen, G.: *Planning and Evaluating the Societal Impacts of Climate Change Research Projects: A guidebook for natural and physical scientists looking to make a difference*, 2021.
- Monteith, J. L.: Evaporation and environment., *Symp. Soc. Exp. Biol.*, 19, 205–234, 1965.

- 510 Mote, T. L., Estilow, T. W., Henderson, G. R., Leathers, D. J., Robinson, D. A., and Suriano, Z. J.: Daily gridded north American snow, temperature, and precipitation, 1959-2009, version 1, Boulder, Color. USA NSIDC Natl. Snow Ice Data Center. N5028PQ3. <https://nsidc.org/data/G10021/versions/1> (accessed 30 June, 2021), 2018.
- Muche, M. E., Sinnathamby, S., Parmar, R., Knightes, C. D., Johnston, J. M., Wolfe, K., Purucker, S. T., Cyterski, M. J., and Smith, D.: Comparison and Evaluation of Gridded Precipitation Datasets in a Kansas Agricultural Watershed Using
515 SWAT, *JAWRA J. Am. Water Resour. Assoc.*, 56, 486–506, <https://doi.org/10.1111/1752-1688.12819>, 2020.
- Musselman, K. N., Clark, M. P., Liu, C., Ikeda, K., and Rasmussen, R.: Slower snowmelt in a warmer world, *Nat. Clim. Chang.*, 7, 214–219, <https://doi.org/10.1038/nclimate3225>, 2017.
- Musselman, K. N., Lehner, F., Ikeda, K., Clark, M. P., Prein, A. F., Liu, C., Barlage, M., and Rasmussen, R.: Projected increases and shifts in rain-on-snow flood risk over western North America, *Nat. Clim. Chang.*, 8, 808–812,
520 <https://doi.org/10.1038/s41558-018-0236-4>, 2018.
- Musselman, K. N., Lehner, F., Eidhammer, T., Pendergrass, A., and Gutmann, E. D.: Assessing the Predictability and Probability of 21 st Century Rain-on-Snow Flood Risk for the Conterminous US, in: AGU Fall Meeting 2021, 2021a.
- Musselman, K. N., Addor, N., Vano, J. A., and Molotch, N. P.: Winter melt trends portend widespread declines in snow water resources, *Nat. Clim. Chang.*, 2021, 418–424, <https://doi.org/10.1038/S41558-021-01014-9>, 2021b.
- 525 Myers, D. T., Ficklin, D. L., Robeson, S. M., Neupane, R. P., Botero-Acosta, A., and Avellaneda, P. M.: Choosing an arbitrary calibration period for hydrologic models: How much does it influence water balance simulations?, *Hydrol. Process.*, 35, e14045, <https://doi.org/10.1002/hyp.14045>, 2021a.
- Myers, D. T., Ficklin, D. L., and Robeson, S. M.: Incorporating rain-on-snow into the SWAT model results in more accurate simulations of hydrologic extremes, *J. Hydrol.*, 603, 126972, <https://doi.org/10.1016/J.JHYDROL.2021.126972>,
530 2021b.
- Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I - A discussion of principles, *J. Hydrol.*, 10, 282–290, [https://doi.org/10.1016/0022-1694\(70\)90255-6](https://doi.org/10.1016/0022-1694(70)90255-6), 1970.
- Pan, C. G., Kirchner, P. B., Kimball, J. S., Kim, Y., and Du, J.: Rain-on-snow events in Alaska, their frequency and distribution from satellite observations, *Environ. Res. Lett.*, 13, 1–15, <https://doi.org/10.1088/1748-9326/aac9d3>, 2018.
- 535 Pianosi, F. and Wagener, T.: A simple and efficient method for global sensitivity analysis based on cumulative distribution functions, *Environ. Model. Softw.*, 67, 1–11, <https://doi.org/10.1016/j.envsoft.2015.01.004>, 2015.
- Pomeroy, J. W., Fang, X., and Marks, D. G.: The cold rain-on-snow event of June 2013 in the Canadian Rockies — characteristics and diagnosis, *Hydrol. Process.*, 30, 2899–2914, <https://doi.org/10.1002/hyp.10905>, 2016.
- Rajulapati, C. R., Papalexiou, S. M., Clark, M. P., and Pomeroy, J. W.: The Perils of Regridding: Examples Using a Global
540 Precipitation Dataset, *J. Appl. Meteorol. Climatol.*, 60, 1561–1573, <https://doi.org/10.1175/JAMC-D-20-0259.1>, 2021.
- Rennert, K. J., Roe, G., Putkonen, J., and Bitz, C. M.: Soil thermal and ecological impacts of rain on snow events in the circumpolar arctic, *J. Clim.*, 22, 2302–2315, <https://doi.org/10.1175/2008JCLI2117.1>, 2009.

- Ritchie, J. T.: Model for predicting evaporation from a row crop with incomplete cover, *Water Resour. Res.*, 8, 1204–1213, <https://doi.org/10.1029/WR008i005p01204>, 1972.
- 545
- Rössler, O., Froidevaux, P., Börst, U., Rickli, R., Martius, O., and Weingartner, R.: Retrospective analysis of a nonforecasted rain-on-snow flood in the Alps-A matter of model limitations or unpredictable nature?, *Hydrol. Earth Syst. Sci.*, 18, 2265–2285, <https://doi.org/10.5194/hess-18-2265-2014>, 2014.
- Rotstayn, L. D., Jeffrey, S. J., Collier, M. A., Dravitzki, S. M., Hirst, A. C., Syktus, J. I., and Wong, K. K.: Aerosol-and greenhouse gas-induced changes in summer rainfall and circulation in the Australasian region: a study using single-forcing climate simulations, *Atmos. Chem. Phys.*, 12, 6377–6404, 2012.
- 550
- Serreze, M. C., Gustafson, J., Barrett, A. P., Druckenmiller, M. L., Fox, S., Voveris, J., Stroeve, J., Sheffield, B., Forbes, B. C., and Rasmus, S.: Arctic rain on snow events: bridging observations to understand environmental and livelihood impacts, *Environ. Res. Lett.*, 16, 105009, 2021.
- 555
- Stern, M. A., Flint, L. E., Flint, A. L., Boynton, R. M., Stewart, J. A. E., Wright, J. W., and Thorne, J. H.: Selecting the Optimal Fine-Scale Historical Climate Data for Assessing Current and Future Hydrological Conditions, *J. Hydrometeorol.*, 23, 293–308, <https://doi.org/10.1175/JHM-D-21-0045.1>, 2022.
- Stewart, I. T., Cayan, D. R., and Dettinger, M. D.: Changes in Snowmelt Runoff Timing in Western North America under a 'Business as Usual' Climate Change Scenario, *Clim. Chang.* 2004 621, 62, 217–232, <https://doi.org/10.1023/B:CLIM.0000013702.22656.E8>, 2004.
- 560
- Sui, J. and Koehler, G.: Rain-on-snow induced flood events in southern Germany, *J. Hydrol.*, 252, 205–220, [https://doi.org/10.1016/S0022-1694\(01\)00460-7](https://doi.org/10.1016/S0022-1694(01)00460-7), 2001.
- Suriano, Z. J.: On the role of snow cover ablation variability and synoptic-scale atmospheric forcings at the sub-basin scale within the Great Lakes watershed, *Theor. Appl. Climatol.*, 35, 607–621, <https://doi.org/10.1007/s00704-018-2414-8>, 2018.
- 565
- Suriano, Z. J.: Synoptic and meteorological conditions during extreme snow cover ablation events in the Great Lakes Basin, *Hydrol. Process.*, 34, 1949–1965, <https://doi.org/10.1002/hyp.13705>, 2020.
- Suriano, Z. J.: North American rain-on-snow ablation climatology, *Clim. Res.*, 87, 133–145, <https://doi.org/10.3354/CR01687>, 2022.
- 570
- Suriano, Z. J. and Leathers, D. J.: Spatio-temporal variability of Great Lakes basin snow cover ablation events, *Hydrol. Process.*, 31, 4229–4237, <https://doi.org/10.1002/hyp.11364>, 2017.
- Suriano, Z. J. and Leathers, D. J.: Great lakes basin snow-cover ablation and synoptic-scale circulation, *J. Appl. Meteorol. Climatol.*, 57, 1497–1510, <https://doi.org/10.1175/JAMC-D-17-0297.1>, 2018.
- Suriano, Z. J., Robinson, D. A., and Leathers, D. J.: Changing snow depth in the Great Lakes basin (USA): Implications and trends, 26, 1–11, <https://doi.org/10.1016/j.ancene.2019.100208>, 2019.
- 575
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bull. Am. Meteorol. Soc.*, 93, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.

- Thomson, A. M., Calvin, K. V., Smith, S. J., Kyle, G. P., Volke, A., Patel, P., Delgado-Arias, S., Bond-Lamberty, B., Wise, M. A., Clarke, L. E., and Edmonds, J. A.: RCP4.5: A pathway for stabilization of radiative forcing by 2100, *Clim. Change*, 109, 77–94, <https://doi.org/10.1007/S10584-011-0151-4/FIGURES/12>, 2011.
- 580 US Bureau of Reclamation: Downscaled CMIP3 and CMIP5 Hydrology Climate Projections: Release of Hydrology Projections, Comparison with Preceding Information, and Summary of User Needs, https://gdo-dcp.ucllnl.org/downscaled_cmip_projections/techmemo/downscaled_climate.pdf, 2013.
- USGS: Watershed Boundary Dataset, <https://www.usgs.gov/national-hydrography/watershed-boundary-dataset>, 2022.
- 585 Voltaire, A., Sanchez-Gomez, E., y Méliá, D. S., Decharme, B., Cassou, C., Sénési, S., Valcke, S., Beau, I., Alias, A., and Chevallier, M.: The CNRM-CM5. 1 global climate model: description and basic evaluation, *Clim. Dyn.*, 40, 2091–2121, 2013.
- Volodin, E. M., Dianskii, N. A., and Gusev, A. V.: Simulating present-day climate with the INMCM4. 0 coupled model of the atmospheric and oceanic general circulations, *Izv. Atmos. Ocean. Phys.*, 46, 414–431, 2010.
- 590 Vrugt, J. A. and Robinson, B. A.: Improved evolutionary optimization from genetically adaptive multimethod search, *Proc. Natl. Acad. Sci.*, <https://doi.org/10.1073/pnas.0610471104>, 2007.
- Wang, H. M., Chen, J., Xu, C. Y., Zhang, J., and Chen, H.: A Framework to Quantify the Uncertainty Contribution of GCMs Over Multiple Sources in Hydrological Impacts of Climate Change, *Earth's Futur.*, 8, 1–17, <https://doi.org/10.1029/2020EF001602>, 2020.
- 595 Watanabe, M., Suzuki, T., O'ishi, R., Komuro, Y., Watanabe, S., Emori, S., Takemura, T., Chikira, M., Ogura, T., and Sekiguchi, M.: Improved climate simulation by MIROC5: Mean states, variability, and climate sensitivity, *J. Clim.*, 23, 6312–6335, 2010.
- Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., Nozawa, T., Kawase, H., Abe, M., and Yokohata, T.: MIROC-ESM 2010: Model description and basic results of CMIP5-20c3m experiments, *Geosci. Model Dev.*, 4, 845–872, 2011.
- 600 Welty, J. and Zeng, X.: Characteristics and causes of extreme snowmelt over the conterminous United States, *Bull. Am. Meteorol. Soc.*, 102, E1526–E1542, 2021.
- Willems, P., Mora, D., Vansteenkiste, T., Taye, M. T., and Van Steenbergen, N.: Parsimonious rainfall-runoff model construction supported by time series processing and validation of hydrological extremes—Part 2: Intercomparison of models and calibration approaches, *J. Hydrol.*, 510, 591–609, 2014.
- 605 Willmott, C. J. and Matsuura, K.: Advantages of the mean absolute error (MAE) over the root mean square error (RMSE) in assessing average model performance, *Clim. Res.*, 30, 79–82, <https://doi.org/10.3354/cr030079>, 2005.
- Willmott, C. J., Robeson, S. M., and Matsuura, K.: Geographic Box Plots, 28, 331–344, <https://doi.org/10.2747/0272-3646.28.4.331>, 2007.
- 610 Willmott, C. J., Matsuura, K., and Robeson, S. M.: Ambiguities inherent in sums-of-squares-based error statistics, *Atmos. Environ.*, 43, 749–752, <https://doi.org/10.1016/J.ATMOENV.2008.10.005>, 2009.

- Willmott, C. J., Robeson, S. M., and Matsuura, K.: Short Communication A refined index of model performance, *Int. J. Climatol.*, 33, 1053–1056, <https://doi.org/10.1002/joc.2419>, 2012.
- Willmott, C. J., Robeson, S. M., Matsuura, K., and Ficklin, D. L.: Assessment of three dimensionless measures of model performance, *Environ. Model. Softw.*, 73, 167–174, <https://doi.org/10.1016/J.ENVSOFT.2015.08.012>, 2015.
- Wilson, L. G., DeCook, K. J., and Neuman, S. P.: Regional recharge research for Southwest alluvial basins, Water Resour. Res. Center, Univ. Arizona, Tuscon, 1980.
- Winchell, M., Srinivasan, R., Di Luzio, M., and Arnold, J.: ArcSWAT Interface For SWAT2012: User’s Guide, Texas Agricultural Experiment Station and United States Department of Agriculture, Temple, TX., 2013.
- 620 Wu, T., Song, L., Li, W., Wang, Z., Zhang, H., Xin, X., Zhang, Y., Zhang, L., Li, J., and Wu, F.: An overview of BCC climate system model development and application for climate change studies, *J. Meteorol. Res.*, 28, 34–56, 2014.
- Wuebbles, D., Cardinale, B., Cherkauer, K., Davidson-Arnott, R., Hellmann, J., Infante, D., Johnson, L., de Loe, R., Lofgren, B., and Packman, A.: An assessment of the impacts of climate change on the Great Lakes, *Environ. Law Policy Cent.*, 2019.
- 625 Ye, H., Yang, D., and Robinson, D.: Winter rain on snow and its association with air temperature in northern Eurasia, *Hydrol. Process.*, 22, 2728–2736, <https://doi.org/10.1002/hyp.7094>, 2008.
- Yukimoto, S., Adachi, Y., Hosaka, M., Sakami, T., Yoshimura, H., Hirabara, M., Tanaka, T. Y., Shindo, E., Tsujino, H., and Deushi, M.: A new global climate model of the Meteorological Research Institute: MRI-CGCM3—Model description and basic performance—, *J. Meteorol. Soc. Japan. Ser. II*, 90, 23–64, 2012.
- 630 Zhang, X., Srinivasan, R., and Van Liew, M.: Multi-site calibration of the SWAT model for hydrologic modeling, *Trans. ASABE*, 51, 2039–2049, 2008.

Table 1. Climate models for the Representative Concentration Pathway (RCP) 4.5 scenario used in the research. Full names for each modeling center can be found in Table S1.

Climate Model	Modeling Center	Country	Citation
ACCESS1-0	CSIRO-BOM	Australia	(Bi et al., 2013)
BCC-CSM1-1	BCC	China	(Wu et al., 2014)
CanESM2	CCCMA	Canada	(Arora et al., 2011)
CCSM4	NCAR	USA	(Gent et al., 2011)
CESM1-BGC	NSF-DOE-NCAR	USA	(Long et al., 2013)
CNRM-CM5	CNRM-CERFACS	France	(Voltaire et al., 2013)
CSIRO-Mk3-6-0	CSIRO-QCCCE	Australia	(Rotstayn et al., 2012)
GFDL-ESM2G	NOAA GFDL	USA	(Dunne et al., 2012)
GFDL-ESM2M	NOAA GFDL	USA	(Dunne et al., 2012)
INM-CM4	INM	Russia	(Volodin et al., 2010)
IPSL-CM5A-LR	IPSL	France	(Hourdin et al., 2013)
IPSL-CM5A-MR	IPSL	France	(Dufresne et al., 2013)
MIROC-ESM	MIROC	Japan	(Watanabe et al., 2011)
MIROC-ESM-CHEM	MIROC	Japan	(Watanabe et al., 2011)
MIROC5	MIROC	Japan	(Watanabe et al., 2010)
MPI-ESM-LR	MPI-M	Germany	(Giorgetta et al., 2013)
MPI-ESM-MR	MPI-M	Germany	(Giorgetta et al., 2013)
MRI-CGCM3	MRI	Japan	(Yukimoto et al., 2012)
NorESM1-M	NCC	Norway	(Bentsen et al., 2013)

Figure captions

- 640 **Figure 1.** Study area map of the Great Lakes Basin, showing historical and projected climate data grid points and study river systems.
- Figure 2.** Evaluation statistics for simulating historic, a) streamflow using Nash Sutcliffe Efficiency (NSE), b) streamflow using revised Index of Agreement (d_r), and c) snowpack using mean absolute error (MAE) at the daily time step. Adapted from Myers et al., (2021b).
- Figure 3.** Basinwide ensemble-average a) annual total precipitation, b) annual air temperature, c) winter+spring rain to snow ratio, d) winter+spring rainfall, and e) winter+spring snowfall from the climate input data, with 10-year averages (red lines), based on the RCP 4.5 pathway. Shading indicates historic 1960-1999 (red) and mid-21st century 2040-2069 (blue) periods, as well as ensemble standard deviations (grey).
- 645 **Figure 4.** Historic and projected climate changes for the Great Lakes Basin between the historic 1960-1999 period and mid-21st century (2040-2069) for RCP 4.5 ensemble mean winter+spring: (a and b) air temperatures, (c and d) rainfall, and (e and f) snowfall, based on water years. Left panels show historic amounts while right panels show absolute or percent changes.
- 650 **Figure 5.** Changes in RCP 4.5 ensemble mean monthly a) rain-on-snow (ROS) melt, b) total snowmelt (including temperature-based melt and rain-on-snow), c) the proportion of monthly snowmelt from rain-on-snow events, and d) snowpack snow water equivalent (SWE) for 158 individual major river basins of the Great Lakes Basin between the historic 1960-1999 and mid-21st century periods. Boxplots display the median and interquartile range of results for major river basins and are weighted by river basin area (Willmott et al., 2007).
- Figure 6.** Changes in (a and b) proportion of total snowmelt from rain-on-snow, (c and d) subbasin ROS melt, and (e and f) frequency of ROS events between the historic 1960-1999 period and mid-21st century. Left panels indicate historic conditions, while the right panels indicate absolute or percent changes. Projections represent the RCP 4.5 ensemble mean.
- 655 **Figure 7.** Changes in (a and b) the center of volume (COV) for ROS events (the day of the water year [DOY] when half the total annual rain-on-snow melt is passed), and (c and d) daily mean winter+spring snowpack SWE between the historic 1960-1999 period and mid-21st century. Left panels indicate historic conditions, while right panels indicate percent changes. Projections represent the RCP 4.5 ensemble mean. The water year lasts from 1 October to 30 September.
- 660 **Figure 8.** Relationships between the historic 1960-1999 conditions and the percent change from the historic period to mid-21st century for RCP 4.5 ensemble means. A) historic winter+spring SWE and ROS melt amount change, b) historic winter+spring SWE and ROS frequency change, c) historic winter+spring precipitation and ROS melt change, d) historic winter+spring precipitation and ROS frequency change, e) historic winter+spring air temperature and ROS melt change, and f) historic winter+spring air temperature and ROS frequency change. Colors are coded with the latitudinal gradient.
- 665 **Figure 9.** Correlations of the change in the COV for rain-on-snow melt events between the historic 1960-1999 period and mid-21st century, with the percent change in monthly water yield during that time, for spring and summer months. Markers represent projections of the RCP 4.5 ensemble mean for individual major river basins. Colors are coded with the latitudinal gradient. The water year lasts from 1 October to 30 September.

670