

Limits and CO₂ equilibration of near-coast alkalinity enhancement

Jing He^{1,*} and Michael D. Tyka^{2,*}

¹MIT-WHOI Joint Program in Oceanography and Applied Ocean Science and Engineering, Cambridge, MA, 02139, USA

²Google Inc., 601 N 34th St, Seattle, WA 98103, USA

*These authors contributed equally to this work.

Correspondence: Michael Tyka(mike.tyka@gmail.com)

Abstract. Ocean Alkalinity Enhancement (OAE) has recently gained attention as a potential method for carbon dioxide removal (CDR) at gigatonne scale, with near-coast OAE operations being economically favorable due to proximity to mineral and energy sources. In this paper we study critical questions which determine the scale and viability of OAE: Which coastal locations are able to sustain a large flux of alkalinity at minimal pH and Ω_{Arag} (aragonite saturation) changes ? What is the 5 interference distance between adjacent OAE projects ? How much CO₂ is absorbed per unit of alkalinity added ? How quickly does the induced CO₂ deficiency equilibrate with the atmosphere ? Choosing relatively conservative constraints on ΔpH or $\Delta\Omega_{\text{Omega}}$, we examine the limits of OAE using the LLC270 (0.3deg) ECCO global circulation model. We find that the sustainable OAE rate varies over 1-2 orders of magnitude between different coasts and exhibits complex patterns and non-local dependencies which vary from region to region. In general, OAE in areas of strong coastal currents allow for the largest fluxes 10 and depending on the direction of these currents, neighboring OAE sites can exhibit dependencies as far as 400 km or more. At these steady state fluxes most regional stretches of coastline are able to accommodate on the order of tens to hundreds of megatonnes of negative emissions within 300 km of the coast. We conclude that near-coastal OAE has the potential to scale globally to several GtCO₂/yr of drawdown with conservative pH constraints, if the effort is spread over the majority of available coastlines. Depending on the location, we find a diverse set of equilibration kinetics, determined by the interplay of gas 15 exchange and surface residence time. Most locations reach an uptake-efficiency plateau of 0.6-0.8mol CO₂ per mol of alkalinity after 3-4 years, after which there is only slow additonal CO₂ uptake. Regions of significant downwelling (e.g. around Iceland) should be avoided by OAE deployments, as in such locations up to half of the CDR potential of OAE can be lost to bottom waters. The most ideal locations, reaching an uptake of around 0.8, include north Madagascar, San Francisco, Brazil, Peru and locations close to the southern ocean such as Tasmania, Kerguelen and Patagonia, where the gas exchange appears to 20 occur faster than the surface residence time. However, some locations (e.g. Hawaii) take significantly longer to equilibrate (up to 8-10 years) but can still eventually achieve high uptake ratios.

1 Introduction

To mitigate the worst effects of climate change, the Paris Agreement aims to limit global temperature warming to below 2°C. This requires not only rapid decarbonization, but also negative CO₂ emissions technologies (NET) (Rogelj et al., 2018).

25 About 150-800 GtCO₂ of net negative emissions are needed in the IPCC SSP1-1.9 to SSP1-2.6 scenarios (in addition to

decarbonization) to limit global warming to 2°C by 2100, and this scenario further assumes net negative annual emissions towards the end of the century (Rogelj et al., 2018; Metz and Intergovernmental Panel on Climate Change, 2005; Masson-Delmotte et al., 2021).

On geological timescales, Earth regulates atmospheric CO₂ concentrations by the combined action of surface rock weathering and ocean CO₂ uptake (Penman et al., 2020). High-CO₂ conditions lead to elevated temperatures and an intensified hydrological cycle, which increases silicate rock weathering (Archer et al., 2009). The subsequently dissolved alkalinity increases the ocean's capacity for CO₂ and the excess atmospheric CO₂ dissolves into the ocean, largely reacting to form (bi)carbonate ions (Zeebe and Wolf-Gladrow, 2001). Indeed the ocean's total dissolved inorganic carbon (DIC) exceeds that of the current atmosphere by 50 fold (Sarmiento and Gruber, 2006).


This mechanism operates on a 10-100ka timescale (Archer et al., 2009), limited by the slow intrinsic kinetics of silicate rock dissolution and the slow introduction of unweathered rock. Exposure of fresh igneous rocks has been linked to rapid cooling of the Earth's past climate (Gernon et al., 2021). Unfortunately, this natural homeostat operates too slowly to mitigate anthropogenic climate change this century. Ocean alkalinity enhancement (OAE) (Renforth and Henderson, 2017) is a proposed approach to accelerate this process in order to increase the ocean's capacity for CO₂ and draw down some of the anthropogenic atmospheric CO₂.

The kinetics of rock dissolution can be accelerated in a number of ways. The simplest approach is to increase the rock's surface area through grinding. Powdered rocks such as olivine can then be added to the ocean and will dissolve over the course of years to decades, adding alkalinity (Hangx and Spiers, 2009; Schuiling and de Boer, 2011; Renforth, 2012; Montserrat et al., 2017; Rigopoulos et al., 2018; Meysman and Montserrat, 2017). Alternatively some rocks (e.g. CaCO₃) may be preprocessed by calcining, transforming them into more-rapidly dissolving substances such as CaO (Kheshgi, 1995). Major concerns with these approaches are the risk of CaCO₃ precipitation (Moras et al., 2022; Hartmann et al., 2022), which would remove alkalinity from the ocean, and the introduction of co-contaminants into the ocean. Iron, abundant in most olivine minerals, could inadvertently fertilize the ocean and cause significant ecological effects (Bach et al., 2019). Silicates would likely shift the phytoplankton species composition towards diatoms (Bach et al., 2019). The impact of heavy metals such as Nickel(Guo et al., 2022) is likely complex (Ferdeger et al., 2022) and species-specific. Finally, changes in turbidity and large energy costs of grinding (Li and Hitch, 2015) make deployment of particles <10μm impractical, while deployment of coarser particles is limited to shallow waters, as the dissolution is much slower (Montserrat et al., 2017). The long dissolution times also delay the beneficial effects on atmospheric CO₂ concentrations, likely by decades.

An alternative to direct addition of rock mass to the ocean are electrochemical methods which effectively remove acidity from seawater and neutralize it using rocks on land. Acid could be neutralized by using mine tailings and other industrial wastes or by pumping it into underground basalt formations(Matter et al., 2009; McGrail et al., 2006; Goldberg et al., 2008). Several variants has been proposed based on electrolysis (House et al., 2007; Rau, 2009; Davies et al., 2018) or bipolar electrodialysis (Eisaman et al., 2018; de Lannoy et al., 2018; Digdaya et al., 2020), all essentially producing either pure NaOH or a basified seawater stream, which would be returned to the ocean to increase the pH and elicit CO₂ drawdown. The disadvantage is the significant electrical energy requirement and the fact that the produced alkalinity is relatively dilute (1mol/kg (de Lannoy

et al., 2018)), exacerbating transport costs out to sea, compared to shipping powdered rock. Prior assessments of shipping costs (Renforth, 2012) when using dedicated fleets have focused on transport of rock-based, solid alkalinity (notably olivine), which have a high molality of alkalinity (25mol/kg).

Regardless of the alkalinity source, OAE methods leverage the marine carbonate system (Renforth and Henderson, 2017), a
65 multiple equilibrium state (Zeebe and Wolf-Gladrow, 2001) described by the equation



Dissolved inorganic carbon (DIC) is the combined concentration of all carbonate species. Addition of alkalinity (e.g. OH^-) shifts the above equilibrium to the right by consuming H^+ ions, thus lowering the partial pressure of CO_2 in the ocean and driving further ocean CO_2 uptake (Middelburg et al., 2020; Zeebe and Wolf-Gladrow, 2001). As the sea-surface CO_2 exchange
70 is rate limiting (surface water experiences an equilibration time scale on the order of weeks to years (Jones et al., 2014)), the addition of alkalinity causes a local increase in pH and aragonite saturation (Ω_{Arag}) and a decrease in pCO₂, all of which could potentially affect the local ecology (Subhas et al., 2022; Ferderer et al., 2022; Bach et al., 2019). Furthermore, increases in aragonite saturation could lead to precipitation of calcium carbonate, which removes alkalinity from the surface water and is counter productive with respect to CO_2 uptake (Moras et al., 2022; Hartmann et al., 2022). A number of previous studies
75 have used ocean circulation models combined with a carbon cycle model to estimate the carbon uptake potential of various hypothetical OAE scenarios (Köhler et al., 2013; González and Ilyina, 2016; Feng et al., 2017; Ilyina et al., 2013; Keller et al., 2014; Burt et al., 2021; Tyka et al., 2022). Some of these studies investigate very high rates of alkalinity injection to test the limits of OAE. Ilyina et al. (2013) simulated alkalinity addition on the order of 2.8Pmol/yr (for an approximate uptake of 50 GtCO₂/yr). González and Ilyina (2016) added enough alkalinity to remove around 44GtCO₂/yr. Both these studies found
80 drastic changes in pH and the carbonate saturation state.

Most of these simulations consider globally uniform alkalinity injection patterns, which is unrealistic for practical deployment and provides little insight into which geographical locations are ideal for conducting OAE. An ideal region (for purposes of negative emissions) minimizes the effect of added alkalinity on the local carbonate system and ecology, while maximizing the CO_2 uptake per unit alkalinity added.

85 Several authors have conducted scenario-driven and locally-resolved simulations. Köhler et al. (2013) investigated finely ground olivine addition from ship tracks for a total uptake of 3.2 GtCO₂/yr, simulation the distribution of alkalinity via ballast water of commercial ships. These ship tracks span the full ocean extent from 40°S to 60°N, although heavily weighted to the area between 20°N to 50°N.

Feng et al. (2017) simulated adding olivine along global coastlines where continental shelves are shallower than 200 m. They
90 found that to stay below aragonite saturation levels of $\Omega_{\text{Arag}}=3.4$ and $\Omega_{\text{Arag}}=9$, coastal olivine addition can remove around 12 GtCO₂/yr and 36 GtCO₂/yr, respectively. Some finer-grained studies have been undertaken: Burt et al. (2021) tested regional alkalinity addition based on eight hydrodynamic regimes in a 1.5° model, and Tyka et al. (2022) simulated alkalinity addition

at individual points in a 6° lat-lng grid. Both studies revealed that the pH sensitivity and the efficiency of CO₂ uptake vary geographically and temporally.

95 Here, we also study alkalinity addition through a practical and economic lens, focusing on electrochemical methods, which produce NaOH or other rapidly-dissolving forms of alkalinity. We begin with the assumption that the optimal places for electrochemical alkalinity production would be on the coast, with access both to seawater and low-cost renewable electricity. To minimize risks to coastal ecosystems and ensure adequate spreading and quick dilution, the alkalinity would be transported some distance off-shore. This is increasingly critical for larger scale deployments, to avoid high concentrations of alkalinity.
100 We wish to determine how far off coast and over what area alkalinity can be added to the surface ocean while staying within conservative biological and geochemical limits. While these issues are less relevant to initial small-scale OAE our goal is to examine the limits of the technology's potential scale. A judicious amount of OAE may also be beneficial, by stabilizing or reversing the anthropogenic acidification of the surface ocean (Albright et al., 2016; Feng et al., 2016). Specific implementations of OAE may also be subject to additional limitations such as trace metal contamination (Guo et al., 2022; Bach et al., 2019),
105 which we do not address here.

Increases in alkalinity change the activities of all forms of CO₂ in the carbonate system (Middelburg et al., 2020; Zeebe and Wolf-Gladrow, 2001), many of which are relevant to marine organisms (Riebesell and Tortell, 2011). Both the direct impact on marine species and the risk of triggering calcium carbonate precipitation must be considered (Bach et al., 2019; Hartmann et al., 2022). Given the complexity of the carbonate system and the variety of responses to each parameter there is no single
110 "correct" choice of proxy (Fassbender et al., 2021) by which to quantify the shift in carbonate state, although the parameters are strongly correlated with each other. Further, what constitutes a safe limit for any given ocean parameter is under debate and likely varies significantly between regions, thus a blanket hard limit is difficult to establish. Here we use two proxies to quantify changes in the carbonate system: ΔpH and ΔΩ_{Arag}.

Prior studies simulated the addition of uniform amounts of alkalinity over some defined area and measured the varying
115 response of ocean parameters. However, because the sensitivity of these parameters varies over more than an order of magnitude we designed our experiment in reverse, i.e we adjust the alkalinity addition rate in each grid cell to result in a uniform and relatively small change of a given parameter. We can then examine how the injection rate varies and construct maps that indicate regions of high suitability for OAE.

Finally, to assess the effectiveness and timescale of CO₂ uptake due to an OAE deployment in a given region of interest we
120 can define the uptake efficiency η_{CO2} as

$$\eta_{CO2}(t) = \frac{\Delta DIC(t)}{\Delta Alk}, \quad (1)$$

where t denotes the time since alkalinity was added. η_{CO2} is a unitless molar ratio. Following the addition of some quantity ΔAlk to seawater, the ocean will begin taking up CO₂, eventually reaching a maximum η_{CO2}($t = \infty$) ≈ 0.8 (Renforth and Henderson, 2017; Tyka et al., 2022). The exact value depends on the parameters of the carbonate system, i.e. Alk, DIC,
125 temperature etc., with a typical range of 0.75-0.85 (Tyka et al., 2022). However, the equilibration kinetics of this equilibration

are known to vary spatially due to differences in the gas exchange timescales and the surface residence time of CO₂ deficient water (Jones et al., 2014; Burt et al., 2021). We thus conducted simulated experiments with short, localized pulse injections, followed by tracking of the total excess alkalinity and DIC relative to a reference simulation as done previously with a much coarser model (Tyka et al., 2022). This gives an accurate picture of where alkalinity from a particular injection point is advected
130 to, how much alkalinity is lost to the deep ocean and how much and when CO₂ uptake can be expected.

2 Methods

2.1 The Model

We use the ECCO (Estimating the Circulation and Climate of the Ocean) LLC270 physical fields (Zhang et al., 2018), to simulate the transport of alkalinity by currents and model alkalinity addition in near-coast areas globally. We inject alkalinity
135 to the simulation in strips along all global coastlines, 37 km wide and larger. ECCO is an ocean state estimate based on the MIT general circulation model (MITgcm) (Marshall et al., 1997) that also integrates all available ocean data since the onset of satellite altimetry in 1992. ECCO uses the adjoint method to iteratively adjust the initial conditions, boundary conditions, forcing fields, and mixing parameters to minimize the model-data errors (Wunsch et al., 2009; Wunsch and Heimbach, 2013). This produces a three-dimensional continuous ocean state estimate that agrees well with observational data. We use the LLC270
140 configuration with a 1/3° degree horizontal resolution (Zhang et al., 2018). All input and forcing files needed to reproduce the ECCO state estimates and the source code are freely available online, and we use them to reproduce the LLC270 flow fields. The LLC270 configuration uses a lat-lng-cap (LLC) horizontal grid, which uses 5 faces to cover the globe. The horizontal resolution ranges from 7.3 km at high latitudes to 36.6 km at low latitudes, and has 50 vertical layers whose grid thickness goes from 10m near the ocean surface to 458m at the bottom (Zhang et al., 2018). We use the iteration-42 state estimate described
145 in Carroll et al. (2020), which spans the years 1992-2017.

To represent the ocean carbonate system, we used the gchem and dic packages within MITgcm. The ocean carbon model was based on Dutkiewicz et al. 2005 and uses 5 biogeochemical tracers (DIC, Alkalinity, phosphate, dissolved organic phosphorus, and oxygen) to simulate the carbonate system. In this model, DIC is advected and mixed by the physical flow fields from the MITgcm, and the sources and sinks of DIC are: CO₂ flux between the ocean and atmosphere, freshwater flux, biological
150 production, and the formation of calcium carbonate shells. The biogeochemical tracers were initialized with contemporary data from GLODAPv2 mapped climatologies (Lauvset et al., 2016; Olsen et al., 2017) where possible, or using data from Dutkiewicz et al. (2005) and were allowed to relax locally by running 100yrs of forward simulation (looping the ECCO forcing fields). Atmospheric CO₂ concentrations were held constant at 415μatm, rather than trying to anticipate future emission scenarios. The surface carbonate tracers were found to stabilize during this time.

155 Since we are not simulating a full Earth system, our model does not account for feedbacks of other carbon sinks which reduce the impact of moving CO₂ from the atmosphere to the ocean (Keller et al., 2018). Wind speeds, used to calculate the gas exchange, are imported from the LLC270 forcing data and the air-sea exchange of CO₂ is parameterized with a uniform gas transfer coefficient (Wanninkhof, 1992). To simulate ocean OAE, we forced the simulation by adding pure alkalinity to the

surface ocean in specified locations to the top grid cell (10m depth) and at a parameterized rate; this assumes that alkalinity is
160 of an effectively instantly dissolving nature, such as an NaOH solution. This avoids complicating factors arising from slower-dissolving materials such as fine olivine powder, for which dissolution rates vary with ocean conditions and may sink out of the surface layers before complete dissolution (Fakhraee et al., 2022). We focused on alkalinity addition in coastal bands following shorelines because that is economically most viable and accessible for shipping or pipelines. Feedbacks of elevated alkalinity on the rate of surface calcification are also not explicitly modeled.

165 Six coastal strips are examined with widths of approximately 37, 74, 111, 185, 296 and 592 km, although it varies slightly due to the varying grid cell sizes in the LLC270 grid. Feng et al. 2017 used a coarser 3.6° lon by 1.8° lat model, and their injection pattern roughly corresponds to our 296 km coastal strip. The much finer LLC model allows us to resolve coastal features in greater detail and to test thinner injection strips. We also examined injection in discrete locations spaced 200 km or 400 km apart along the coastline, in circular patches \approx 120km wide.

170 All runs presented in this paper use the same approach: First a reference simulation is run (spanning 20 ECCO years 1994-2019). Then a second run is conducted starting at the same starting conditions, with an alkalinity forcing added, which perturbs the system in some way. We then analyze the difference in the carbonate systems (ΔpH , $\Delta\Omega_{\text{Omega}}$, ΔDIC , etc.) between these two runs. Since the carbonate model does not influence the flow field, there is no divergence in the flowfields over the 175 25 simulation years and the two trajectories are directly comparable. These simulations are run on a small MPI cluster (13 machines, 59 processes each) on Google Cloud Engine, and take about 6 hours of wall time per simulation year.

2.2 pH and Omega limits

The carbonate chemistry in different regions varies in its sensitivity to alkalinity injection, owing to local differences in ocean circulation, gas exchange and carbonate chemistry. The goal of our experiments is to determine the maximal alkalinity addition rate which can be sustained at any given grid point which limits the change in one of two surface parameters, pH and the
180 aragonite saturation Ω_{Arag} to some chosen value.

Here we chose target constraints $\Delta\text{pH}_{tgt}=0.1$ or $\Delta\Omega_{tgt}=0.5$. These values are somewhat arbitrary and serve simply to calculate the relative sensitivity of different regions. However, as an intuitive point of reference, the already incurred anthropogenic surface acidification since preindustrial times (Doney et al., 2009) is $\Delta\text{pH} \approx -0.1$. Likewise a change of $\Delta\Omega_{\text{Arag}}=+0.5$ is unlikely to trigger carbonate precipitation according to Moras et al. (2022) who established an Ω_{Arag} threshold of 5.

185 An alternative approach would have been to set absolute thresholds for pH and/or Omega, however we did not pursue this for the following reasons. For the estimation of biological impact a relative change to current conditions seems most appropriate as the local ecosystem is adapted to the local conditions and many biologically relevant stressors change proportionally to the relative concentration change. For example the energy expenditure of an organism to maintain its intracellular pH is approximately proportional to the logarithm of the proton concentration difference. For purposes of estimating the precipitation limit
190 an absolute threshold would indeed be more appropriate. However, in polar latitudes where Ω_{Arag} is currently low, the change in alkalinity required to reach the limit (e.g. $\Omega_{\text{Arag}} > 5$, (Moras et al., 2022)) would be so large that it would no longer repre-

sent a realistic scenario, easily exceed the above relative pH limits and likely exceed the bounds of the simulation's predictive domain. We note that our choice of a relative Omega change we obtain a lower bound on the true limit.

In practice which limits are acceptable is subject to debate and likely different in different locations and we do not attempt to anticipate them here, focusing merely on the relative capacity of different ocean regions with respect to these ocean parameters. Our approach is as follows. Each surface gridpoint that is part of the coastal injection strip is given a particular baseline injection rate r (in mol/m²/s). At every timestep and for every gridpoint, the local (in time and space) ΔpH is calculated using the carbonate model and a reference value obtained from an unperturbed reference simulation ($\Delta\text{pH} = \text{pH} - \text{pH}_{ref}$). If this value is lower than ΔpH_{tgt} then alkalinity is added according to the baseline rate, if not then addition is skipped for this timestep.

This mechanism is insufficient to ensure the pH does not exceed the maximal value, as the change in pH is determined not only by the local alkalinity addition, but also by advection of alkalinity from neighboring cells and seasonally varying biological activity. We thus iteratively adjusted the baseline rate for each grid cell to empirically determine a rate which gives rise to approximately the desired ΔpH in the following way.

First a pilot simulation was run where the baseline rate was set uniformly to an extreme value of $r=400$ mol/m²/yr, (higher than any region can accommodate). We ran this simulation for 3 years and recorded the observed amount of addition at each grid cell (generally much lower than the baseline as the above algorithm prevents excessive addition). Second, we reran the simulation using a new, position-dependent baseline rate calculated from the amounts actually added from the pilot simulation using a linear extrapolation to our desired pH maximum. We ran this second simulation for 8 years. We found that the observed ΔpH is now generally very close to the desired ΔpH_{tgt} , however some regions still exceed the target value while others undershot. We thus performed a third simulation where we adjusted the addition rate at any grid point inversely proportional to the observed pH deviation yielding a final third simulation which was allowed to run for 20 years using the ECCO forcing fields from 1995 to 2014. We found that this procedure yielded a relatively narrow distribution of ΔpH or $\Delta\Omega_{Arag}$ for all grid points in the injection strip although some variability remains (Fig. S1). A separate iterative optimization was performed for every injection pattern. Grid points outside of the injection strip showed much smaller changes in pH and never exceeded the target ΔpH . The same procedure was used in a separate set of experiments for $\Delta\Omega_{Arag}$.

Once the rate of OAE is stable and acceptable, we can measure how much alkalinity is being added at each grid point. Note that because of the considerable interdependence between nearby grid points there is no one unique injection pattern that satisfies the ΔpH or $\Delta\Omega_{Arag}$ condition, however multiple independent optimization runs started at different ECCO years yield injection patterns that match very closely.

220 2.3 Pulse additions

When alkalinity is added to the surface ocean it lowers the partial pressure of CO₂ (pCO₂) and thus increases the rate at which CO₂ dissolves in the surface ocean. The effectiveness η_{CO_2} of this uptake is determined by a number of factors which vary significantly by location. The timescale of gas exchange τ_{CO_2} is approximately 3-9 months and varies by location (Jones et al., 2014) while the residence time τ_{res} of water parcels in the mixed layer varies over shorter time, between 2 and 20 weeks.

225 Thus the resultant equilibration efficiency ratio τ_{res}/τ_{CO_2} was found to be significantly below 1.0 in 95% of ocean locations
(Jones et al., 2014). However, CO₂-deficient water parcels initially lost from the mixed layer can re-mix into the mixed layer
at some later time and thus drive further equilibration elsewhere and over longer timescales. This longer term effect was not
explicitly modeled in previous work (Jones et al., 2014) and results in a complex equilibration curve which is not well captured
by a single exponential function. As the kinetics of this longer-term equilibration depend on the deep transport and mixing of
230 the lost alkalinity it has to be simulated explicitly.

We thus extend the work of Jones et al. (2014) by simulating pulse injections of alkalinity in a variety of locations using
the ECCO flow fields. These simulations thus explicitly include all the relevant aspects together (gas-exchange, Revelle factor,
surface transport, mixed layer-depth, residence time and remixing), by measuring the actual excess CO₂ uptake of the ocean
relative to the unperturbed reference simulation. However, because the alkalinity is also distributed horizontally over great
235 distances and mixes from different origins it is impossible to disambiguate the CO₂ uptake timescale of different injection
points from a single simulation. One solution to this problem is to use a Lagrangian approach (van Sebille et al., 2018) which
allows for the tracking of stochastic particles. Here we chose a simpler approach. For a select number of coastal locations we
run a separate simulation and inject a 1-month pulse of alkalinity. Following the pulse we monitor the total excess DIC in the
ocean relative to a reference simulation, the distribution of alkalinity across the depth layers and the pCO₂ deficit at the surface
240 over time. Ideally the length of the pulse would be a single timestep, however this would either necessitate an extreme addition
rate or a tiny total quantity of added alkalinity, which would lead to a poor signal to noise ratio during analysis. The choice of
pulse length thus represents a compromise, as this length is still much shorter than the overall relaxation time. Ideally such a
pulse injection experiment could be conducted for every grid point (as was done with a coarse model in Tyka et al. (2022) and
245 at different times of the year, however, because each pulse requires a whole separate simulation, this exceeded our computation
capacity with the high-resolution ECCO model. Thus we chose 17 individual locations of interest along most major coastlines
with pulses occurring in January.

2.4 Alkalinity injection from ships

In addition to the steady-state perturbation of ocean parameters over large areas of OAE deployment, it is critical to examine
the short term impacts that arise right at the injection site which will temporarily take the local carbonate system into an
250 extremely alkaline regime. This is unlikely to be a concern for gradually dissolving alkalinity such as ground olivine (Hangx
and Spiers, 2009; Schuiling and de Boer, 2011), but highly relevant for rapidly dissolving alkalinity such as NaOH solution or
other solubilized alkaline media.

Of interest is the dilution speed of the added alkalinity (we assume here a solution of 1M NaOH) against the timescale at
which homogeneous nucleation of aragonite is triggered and needs to be avoided. The most natural approach, used also in
255 waste disposal, would be to inject directly into the turbulent wake of the ship to mix the discharge with seawater as rapidly as
possible (Renforth and Henderson, 2017). The dilution kinetics have been studied and modeled in previous work (Chou, 1996;
IMCO, 1975) and large discrepancies exist between the published models. The IMCO model uses the following empirical form
for the unitless dilution factor as function of time t

$$D(t) = \frac{C}{Q} U^{1.4} L^{1.6} t^{0.4} \quad (2)$$

260 where t is time (seconds), Q is the release rate (m^3/s), U is the speed of the ship (m/s), L is the waterline length (m). C is an empirical constant, set to 0.003 for release from a single orifice and 0.0045 for release from multiple ones. Intuitively larger speeds and longer ships have more turbulent wakes producing faster dilution. Chou (1996) used the following, similar model which instead of waterline length uses the width of the ship, B , in units of meters, to account for the ship size.

$$D(t) = \frac{0.2108}{Q} U^{1.552} B^{1.448} t^{0.552} \quad (3)$$

265 Chou's formula gives much faster dilution rates and was verified against field testing data. Other work (Lewis, 1985; Byrne et al., 1988; Lewis and Riddle, 1989) used even higher exponents on the time t so we take the IMCO formula to be an upper limit though in general no universally applicable law can be expected as the dispersion timescale will inevitably depend on local conditions. For a given starting concentration C_0 of the alkaline effluent (e.g. 1mol/L NaOH) we can calculate the resultant alkalinity by considering the dilution with seawater with alkalinity Alk_0

$$270 \quad \text{Alk}(t) = \frac{1}{D(t)} C_0 + \left(1 - \frac{1}{D(t)}\right) \text{Alk}_0 \quad (4)$$

We can then determine the $\text{pH}(t)$ and the carbonate saturation state $\Omega(t)$ by solving the carbonate system at any given $\text{Alk}(t)$. We used PyCO2SYS (Humphreys et al., 2020) to solve the carbonate system numerically, assuming PyCO2SYS default values and $\text{Alk}_0=2300\text{mol/m}^3$ and $\text{DIC}=2050\text{mol/m}^3$. These analytical models are valid only on timescales smaller than one hour after which dilution kinetics are driven by the local background turbulence rather than the immediate influence of the wake 275 turbulence. Thus longer-scale dilution effects will vary substantially from location to location. As the timescale considered here is much shorter than the typical CO_2 gas-exchange timescale we do not explicitly model CO_2 uptake during this initial dilution.

2.5 Estimation of transport costs

Alkalinity prepared on land must be transported out to sea, which adds to the total cost of the achieved negative emissions 280 (in \$/ tCO_2). While at very small scale it could be released right at the coast, for larger overall OAE deployment the alkalinity will need to be spread over greater areas (to avoid excessive local pH or Omega changes) which increases the cost for every additional unit of alkalinity added. Having obtained maps of the allowable rate of OAE in any given area we can estimate the transport costs for each of our simulated scenarios. Large-scale maritime shipping costs are around $\$0.0016-\$0.004 \text{ t}^{-1}\text{km}^{-1}$ (Renforth, 2012). For each grid point where alkalinity is injected, we calculate the distance D to the nearest coast, and take 285 double that to be the minimum round trip distance for a ship to travel. Realistically, a ship will have to travel farther than D since it needs to go to the nearest port or NaOH factory, so this is the lower bound on the transport distance. This allows us




Figure 1. Shown is a coastal injection in a strip 296 km wide, subject to $\Delta p\text{H}_{tgt}=0.1$. All time-averages ran over years 5-20. (a) Averaged pH change compared to a reference simulation with no added alkalinity. (b) Average alkalinity flux at each grid point, from the coast. (c) and (d) Injection flux for two example regions. Annually averaged surface currents are overlaid as a vector field.

to calculate the lower bound of the shipping cost per tCO₂ for each grid point. The scenario model provides the alkalinity injection rate for each grid point, and assuming an eventual uptake efficiency $\Delta_{CO_2} \approx 0.8$ we can obtain the total shipping cost for every grid point. Summing the total cost over all grid points in which injection occurs and dividing by the total expected
290 global CO₂ uptake yields a lower bound of the average effective global transport cost per tonne CO₂.

3 Results and Discussion

3.1 Injection capacity

In all our simulations the alkalinity flux in the injection grid cells was iteratively adjusted to elicit a change in pH or the change
295 in Omega (though not simultaneously) to a value of either $\Delta p\text{H}=+0.1$ or $\Delta\Omega_{Arag}=+0.5$ respectively. Due to the correlation between neighboring grid cells, seasonal and year-to-year changes in currents and biotic activity these constraints are not perfectly satisfied, but we were able to confine them to a narrow range around the desired value (Fig S1). The injection rate, as well as $\Delta p\text{H}$, $\Delta\Omega_{Arag}$, stabilize within the first 5-6 years of the simulation and remain stable for the remainder of the simulation
300 (Fig. S1), indicating that a steady state is reached where the alkalinity addition rate is matched by outflowing alkalinity into open ocean areas and by neutralization by atmospheric CO₂. Note that while the time-averaged $\Delta p\text{H}$ is close to 0.1, there is significant temporal variability that leads to $\Delta p\text{H}$ slightly exceeding 0.1 at some parts of the year (Fig. S1).

As expected, $\Delta p\text{H}$ or $\Delta\Omega_{Arag}$ outside the injection grid cells is much lower and never exceeds the target value. However, the effect on adjacent areas outside of the injection grid points is variable and depends on the pattern of ocean currents that




Figure 2. (a) Total CO₂ flux (GtCO₂/yr), subject to $\Delta\text{pH}_{tgt}=0.1$, broken down into flux through the injection area and outside of the injection area. (b) Dependence of the averaged injection flux (mol/m²/yr) on the width of the coastal injection strip for different coastal regions and strip widths. The black dashed line averages all coastal regions.

sweep alkalinity away from injection areas. For instance, western boundary currents carry the coastal excess alkalinity far out into the open ocean, so we see elevated changes in the North Atlantic and Indian oceans, even outside the injection areas.

For each injection pattern and chosen limit we can now obtain a global map of steady-state alkalinity flux (mol/m²/yr), which shows the variability of capacity for OAE (Fig 1 b,c and d). We note substantial variability on multiple scales. Firstly on a large scale, some coastal areas have a fundamentally greater capacity for distributing and neutralizing alkalinity flux than others (see also Fig. 2b). Large capacities are found around islands which sit in or near ocean currents, as those rapidly sweep the alkalinity away from near-coast areas. Examples include Kerguelen, Easter Island and Hawaii. Continental coasts which exhibit large capacities are found around south and east Africa, off the coast of Peru and Brazil, south-east Australia and the west coast of Japan. Finally an area of very large tolerance for alkalinity addition is found in the northern Atlantic, however, as will be shown later, this is due to downwelling and deep water formation, which is highly undesirable for OAE as the alkalinity cannot efficiently equilibrate with the atmosphere before being lost. Conversely inland seas and partially enclosed seas exhibit the smallest capacity for OAE, notably the Red Sea, the Mediterranean and the Baltic Sea. An interesting counterexample is the Gulf of Mexico and the Caribbean Sea which, owing to the traversing Gulf Stream, have significant capacity for OAE. We found no correlation between background pH and OAE rate, as the influence of local ventilation due to currents dominates the capacity for alkalinity addition.

As expected, the large scale patterns obtained by limiting $\Delta\text{pH}=0.1$ or $\Delta\Omega_{Arag}=0.5$ are very similar (Fig. S2) up to a linear factor and highly correlated on a per grid-cell basis. This is consistent with the observation that OAE capacity is primarily influenced by local currents. However we note that the fundamental sensitivities of pH and Ω_{Arag} with respect to Alk change

quite differently from pole to equator. In particular $\partial pH/\partial Alk$ decreases from $2.6 \times 10^3 \text{ L mol}^{-1}$ at the poles to $1.0 \times 10^3 \text{ L mol}^{-1}$ at the equator, while $\partial \Omega/\partial Alk$ increases from $7.6 \times 10^3 \text{ L mol}^{-1}$ at the poles to $11 \times 10^3 \text{ L mol}^{-1}$ at the equator (computed from GLODAPv2(Lauvset et al., 2016; Olsen et al., 2017) data and pyCO2SYS(Humphreys et al., 2020; Lewis and Wallace, 1998)). Furthermore, from the perspective of preventing precipitation due to excessive Ω_{Arag} , the available
325 headroom is much smaller at tropical latitudes, where Ω_{Arag} is already close to 4, than near the poles where Ω_{Arag} is as low as 1-2(Lauvset et al., 2016; Olsen et al., 2017). Thus, as our relative $\Delta\Omega_{Arag}$ limit was set to 0.5, the OAE limits obtained here should be considered a lower bound with respect to calcite precipitation, i.e. polar regions could tolerate a much larger addition rate. For pH the actual ecologically tolerable limits will vary from coast to coast, and we do not attempt to anticipate them here. We note however that for our examined constraints (both of which are very conservative) a significant amount of
330 negative emissions can be obtained even in very narrow coastal strips as the transport out to open sea is very efficient. This observation is consistent with prior work by Feng et al. (2017).

On a fine scale we find that the sustainable injection flux varies over 2-3 orders of magnitude between nearby grid points (Fig. S1) with a distribution that is approximately log-normal. The variance is even larger for thin injection strips (Fig. S1) and very large fluxes can be sustained in some locations if the ventilation out of the strip is high enough. Depending on the
335 prevailing currents, the highest injection rate can be found both on the outside and inside of the injection strip.

For all regions we observe that while widening the injection strip increases the total allowable rate of alkalinity injection, the increase is sublinear (Fig. 2a). Every additional unit of alkalinity added needs to be transported further offshore and the average injection rate decreases per unit area. This is consistent with the view that the majority of the neutralization of the added alkalinity by invading CO₂ occurs outside the injection strip and the local pH is primarily determined by the rate of transport
340 of alkalinity into other areas (Burt et al., 2021). This is confirmed by integrating the total CO₂ flux over the injection strip and over the rest of the ocean surface, relative to the reference simulation. Figure 2a shows that especially for thin coastal injection strips, direct gas exchange through the strip surface accounts for only a minor component of the induced CO₂ uptake and the majority occurs outside of the injection areas. As the strip widens however, this fraction increases significantly. Especially in regions with weak transport we observe that often a larger quantity of alkalinity can be added right at the border of the strip
345 than in the middle, as alkalinity can dilute to bordering areas that are not directly receiving alkalinity (Fig. 3). Indeed, the largest alkalinity fluxes observed occur when the strip widths are very thin (Fig. S1). The consequence is that any particular coastal region can increase its capacity for injection by going further out to sea, but that there are diminishing returns of doing so, i.e. the increase in capacity is sublinear with width.

The influence of widening the injection area is also shown in detail for three regions in Figure 3 and a larger number of
350 regional details are included in the supplementary material in Figure S4. Widening strips allow more alkalinity to be added overall, however saturation of near-coast areas occurs in most regions (esp evident in Northwest Australia). The largest injection flux is often found directly at the strip boundary, owing to easy diffusion out to open sea. However, it is not always the case that the highest injection flux occurs at the strip edge, as seen in the Japan and East Africa examples. These findings illustrate the highly non-local nature of the injection capacity.




Figure 3. Injection flux (mol/m²/yr) shown for different strip widths in 3 different regions (from top to bottom: Japan, East Africa and Northwest Australia). The inset text indicates the total alkalinity added per year in the shown area.

355 In general we find that the sensitivity of pH and Ω_{Arag} at any given grid point is highly dependent on the surrounding pattern of injection. We conducted two additional experiments in which rather than a contiguous strip, injection occurred at discrete points placed either 200 km or 400 km apart. At 200 km apart we observe much higher injection fluxes in each injection patch, but the total global injection flux barely changes (Fig. 4). Placing injection patches 400 km apart instead of 200 km apart did not further increase the sustainable flux in each injection patch, and reduced the overall injection capacity by 25%. This suggests
 360 that at 200 km there is still significant cross-correlation between neighboring patches, which is apparent when looking at the pH changes observed: the pH impact bleeds into neighboring patches (Fig. 4b). At 400 km apart, however, there is much less interference between adjacent patches and the injection limits are simply dominated by local current patterns (Fig. 4c). These observations are consistent with prior work (Jones et al., 2012) which found a global median pCO₂ autocorrelation length is about 400 km. Thus in order to maximize any particular coastline's injection potential, injection areas should be placed at most




Figure 4. Alkalinity enhancement in three different patterns, exemplified at the west coast of South America: (a) Injection in a contiguous strip (b) Injection in 200 km separated patches (c) Injection in 400 km separated patches The total globally area-integrated OAE rate for the three injection patterns was (a) 336 Tmol/yr, (b) 312 Tmol/yr and (c) 233 Tmol/yr respectively. The sub panels show: (i) Time averaged pH change (ii) Alkalinity flux (iii) Distribution of alkalinity flux (globally).

365 200-400 km apart, however the exact optima will depend on the local current patterns. The location of ports, infrastructure, access to electricity and/or alkaline minerals will dictate the choice of locations. The correlation between neighboring injection locations has ramifications for the planning, monitoring and governance of different OAE projects, as they will affect each other in downstream coastal areas. For monitoring and verification purposes it will be impossible to disambiguate CO₂ drawdown caused by different OAE projects by measurement alone. Any plans to add alkalinity to the ocean will need to be simulated
 370 specifically, ideally with regionally optimized models, and take into account already occurring OAE projects nearby.

3.2 CO₂ uptake timescales

To measure localized CO₂ uptake timescales, we conducted a total of 17 pulse injections (as described in the methods section), placed near all major coastlines. We generally chose locations previously determined as areas of high alkalinity tolerance. Firstly we observed a very large variety of CO₂ uptake timescales (Fig. 5) both in the short term (after 1 year) and in the
 375 medium term (after 10 years). After 1 year the molar uptake fraction $\eta_{CO_2} = \Delta DIC / \Delta Alk$ varied between 0.2 and 0.85 and after 10 years most locations resulted in an uptake fraction of 0.65-0.80 consistent with previous work (Tyka et al., 2022; Burt et al., 2021).




Figure 5. CO₂ uptake relative to alkalinity addition (molar ratio $\eta_{CO_2} = \Delta DIC / \Delta Alk$) following pulse additions at 17 different locations. (a) Locations which equilibrate fast and reach close to the theoretical maximum of CO₂ uptake (0.8) (b) Locations which equilibrate fast but reach a lower plateau of relative CO₂ absorption (0.6-0.8) with slow further progression. (c) Locations with slow equilibration or significant loss of alkalinity to the deep. Note that in some cases despite the slower initial equilibration high eventual uptake ratios can be achieved eventually.

A typical behavior is observed for example when releasing alkalinity on the northern coast of Brazil (Fig. 6a). Here the alkalinity remains long enough at the surface to realize its full CO₂ uptake potential within 2-3 years. A number of other tested
380 locations follow this general pattern and are efficient for OAE deployment (Fig. 5a). Here the equilibration follows roughly a single exponential relaxation.




Figure 6. Pulse additions of alkalinity in 3 representative locations: (a) Brazil, (b) Japan, (c) Iceland. (i) Excess quantity of CO_2 absorbed (relative to the reference simulation) over time following a 1 month pulse injection at various near coast locations, expressed as a molar fraction of the amount of alkalinity added (η_{CO_2}). A significant variation of uptake timescales is observed, depending on speed at which excess alkalinity is removed from surface layers, both in the short term and in the long term. (ii)-(iv) Spatial detail of surface pCO_2 deficit and depth residence of excess alkalinity for the same 3 locations shown in (i). The initial injection location is indicated by a black area. The surface pCO_2 deficit is plotted over time (note the log scale of the colormap), indicating areas which are absorbing extra CO_2 (or emitting less CO_2) compared to the reference simulation. Below, the relative excess alkalinity is plotted against the depth of the water column (averaged over all lat/ln^g grid points for each depth). Further locations are shown in detail in Figure S5 in the supplementary material.

Another set of locations appear to lose a significant amount of alkalinity to deeper layers before atmospheric equilibration is achieved (Fig. 5b,c). For example injection off the coast of Japan resulted in slow initial uptake as a portion of the alkalinity is subducted quickly (Fig. 6b). However in the following decade re-mixing with surface waters gradually returns this alkalinity back towards the surface resulting in slow but steady CO₂ uptake (Fig. 6b). Other examples of this delayed CO₂ equilibration are shown in (Fig. 5b). These locations exhibit a short mixed layer residence time and have a poor equilibration efficiency (Jones et al., 2014) but equilibration is eventually achieved in the following decades. Changes in surface pCO₂ and alkalinity will also potentially affect biological CO₂ uptake and calcification rates, however these dependencies are not captured by the carbonate model used here (Dutkiewicz et al., 2005) and will need to be examined by future studies.

Finally some extreme examples of poor long-term equilibration efficiency are found in areas of deep water formation, such as the northern Atlantic. Here a CO₂ uptake ratio of just over 0.4 is achieved and even after 20 years very little further progress is made (Fig. 6c). Around half of the added alkalinity is subducted very deep and will likely remain out of contact with the atmosphere until the global overturning circulation returns these waters to the surface, on the timescale of many hundreds of years. We did not examine the dependence of the time of year (all of our pulses occurred in January) nor were we able to conduct an exhaustive set of locations as was done previously with a coarser global circulation model (Tyka et al., 2022). We note that for all cases the alkalinity-induced CO₂ deficit spreads over a very large area within one year and a significant fraction of the CO₂ uptake occurs after the deficits have diluted to the sub μatm range. This makes direct monitoring and verification of OAE extremely challenging and will likely need to rely on modelling and indirect experimental verification.

3.3 Alkalinity injection from ships

Other than potential ecological impacts on marine life intersecting the caustic release wake of an OAE ship, one concern is that a short Ω_{Arag} spike could induce precipitation of CaCO₃ (Renforth, 2012). Once nucleated, the CaCO₃ particles could continue to grow, even when the pH has returned to normal ocean levels (≈ 8.1) because the ocean is supersaturated with respect to calcite ($\Omega_{Calc} \approx 2.5 - 6$) and aragonite ($\Omega_{Arag} \approx 1.5 - 4$) (Lauvset et al., 2016; Olsen et al., 2017). While the nucleation of CaCO₃ is strongly inhibited by the presence of magnesium in seawater (Sun et al., 2015; Pan et al., 2021), the growth of existing crystals may not be (Moras et al., 2022; Hartmann et al., 2022). Only once the CaCO₃ particle has reached a size and density that causes it to sink would it stop removing alkalinity from the surface ocean. Thus, depending on the number of particles nucleated, the alkalinity removed from the surface ocean be larger than the alkalinity added (Moras et al., 2022; Hartmann et al., 2022; Fuhr et al., 2022).

Since the immediate dilution dynamics of alkalinity injected into the wake of a ship is far below the resolution of the ECCO LLC270 global circulation model we examine this process analytically, as described in the methods section. The relevant timescales are compared in Figure 7. In blue are shown the predicted pH and Ω_{Arag} as a function of time, based on the dilution formulas given by IMCO (1975) and Chou (1996).

Shown also are homogeneous nucleation times of CaCO₃ for comparison. Several studies (Pokrovsky, 1994, 1998) have measured the homogeneous nucleation of CaCO₃ in seawater at different saturation states down to $\Omega_{Arag}=9$. Three of these models are plotted in shades of orange and black in Figure 7. Theoretical studies (Sun et al., 2015) suggest that for Mg:Ca




Figure 7. The expected time evolution of pH (left scale) and Ω_{Arag} (right scale) due to dilution for alkalinity injection into a ship wake. Here we assume a large tanker (275m long, 50m wide, traveling at 6m/s) releasing 1.0M NaOH at a rate of 5m³/s. Ships of this size have a capacity of 100 to 200 kilotons of cargo which would take 6-12hrs to discharge. Two previously published dilution models are shown in blue with a large variance apparent. For comparison, the timescales of homogeneous nucleation are also shown (yellow, brown and black). The dashed and dotted lines indicate the estimated Ω limits for precipitation. Despite the substantial uncertainty in the existing models, dilution can proceed at least 1-2 orders of magnitude faster than precipitation is expected to occur.

ratios of 5.2, as found in seawater, no nucleation of Aragonite occurs at all below $\Omega_{Calc} = 18$ (equivalent to $\Omega_{Arag} \approx 12$), due to inhibition by Magnesium. This is consistent with Morse and He (1993), however timescales only up to a few hours were examined. Moras et al. (2022) suggested a safe limit of $\Omega_{Arag}=5$ based on alkalinity addition using CaO. Figure 7 shows that ship-wake dilution proceeds at least one order of magnitude faster than the homogeneous nucleation time, thus we can expect
420 that CaCO₃ particles will not be induced to nucleate.

At the immediate injection site where the pH exceeds 9.5-10.0 the temporary precipitation of Mg(OH)₂ is expected, which redissolves readily upon dilution (Pokrovsky and Savenko, 1995) and buffers the pH against further increase (not accounted for in Figure 7). The temporary reduction in the Mg:Ca ratio could make CaCO₃ nucleation more favorable, but the time spent in this state (<1minute) is likely still well below the required nucleation time.




Figure 8. Total carbon uptake potential for a variable alkalinity addition strategy that results in a maximal pH change of 0.1 (see Figure 1 for an example). Carbon uptake is estimated at $\eta_{CO_2} = 0.8$. The strip widths necessary are indicated for each point. Shading denotes the shipping cost uncertainty, which ranges from \$0.0016-\$0.004t⁻¹km⁻¹ (Renforth, 2012)

425 3.4 Transport costs

Alkalinity prepared on land must be transported out to sea, which incurs costs. Having obtained maps of the sustainable rate of OAE in any given area we can estimate the transport costs for each of our simulated scenarios as described in the methods section. Furthermore, the quantity of alkalinity per tonne depends on the molality of the material moved. For rock-based methods (Hangx and Spiers, 2009; Schuiling and de Boer, 2011; Renforth, 2012; Montserrat et al., 2017; Rigopoulos et al., 430 2018; Meysman and Montserrat, 2017), the molality of solid alkaline materials such as olivine is 25mol/kg. For electrochemical alkalinity methods (House et al., 2007; Rau, 2009; Davies et al., 2018; Eisaman et al., 2018; de Lannoy et al., 2018; Digdaya et al., 2020) that produce alkaline liquids it will be closer to 1mol/kg depending on the industrial processes used and effort spent 435 concentrating the alkaline solution. Figure 8 shows how the transportation costs are influenced by the total desired negative emissions (larger scale requires transport further offshore). For concentrated alkalinity (such as ground rocks) transport costs are not a major contributor to cost, consistent with prior work (Renforth, 2012). However for dilute alkalinity, such as obtained 440 from electrochemical processes, transportation could become a significant contributor to the overall cost. A potential low-cost solution could be, for example, the near-coast precipitation(Thorsen et al., 2000; Davies et al., 2018; Sano et al., 2018) of Mg(OH)₂ with subsequent redissolution of the solid Mg(OH)₂ after transport out to sea. However economic tradeoffs between 445 the cost of concentrating the alkalinity and the cost of transportation will need to be made.

In this paper we examined the suitability and effectiveness of near-coast regions of the ocean for alkalinity enhancement (OAE). We conducted a series of high resolution (0.3°) global circulation simulations in which alkalinity was added to coastal strips of varying width under the constraint of limited ΔpH or $\Delta\Omega_{Arag}$. We found that the resultant steady-state rate at which alkalinity can be added at any given location exhibits complex patterns and non-local dependencies which vary from region to region. The allowable injection rate is highly dependent on the surrounding injection pattern and varies over time, responding to external seasonal factors which are not always predictable. This makes it difficult to prevent occasional short spiking beyond the specified limit, thus potentially requiring that the limit is set conservatively in practice. These difficulties are also expected to arise in practice and have repercussions on how such OAE would be performed in reality and how it would be monitored, reported and verified (MRV). The non-local nature of the pH effect also likely requires different adjacent countries to coordinate their OAE efforts.

We found that even within the relatively conservative constraints set, most regional stretches of coastline are able to accommodate on the order of a tens to hundreds of megatonnes of negative emissions, with areas with access to fast currents being able to accommodate more, such as East Africa or the coast of Peru. Globally we conclude that near-coastal OAE has the potential to scale to a few gigatonnes of CO₂ drawdown, if the effort is spread over the majority of available coastlines. However, given that many other factors will determine suitable locations (such as availability of appropriate alkaline minerals, low-cost energy and geopolitical suitability) the global potential may be lower in practice. We also examine the cost of transport of alkalinity, which increases with global deployment size as the alkaline material needs to be spread over greater distances from the shore. For alkalinity schemes based on dry minerals the transport costs remain minor, but for electrochemical methods, which produce more dilute alkalinity, this may present limits to scaling.

We also examined the effectiveness and timescale of alkalinity enhancement on uptake of CO₂, through pulsed injections and subsequent tracking of surface water equilibration. Depending on the location, we find a complex set of equilibration kinetics. Most locations reach a plateau of 0.6-0.8mol CO₂ per mol of alkalinity after 3-4 years, after which there is little further CO₂ uptake. The plateau efficiency depends on the amount of alkalinity lost to the deep ocean which will not equilibrate with the atmosphere until it returns to the surface, on the timescale of 100-500 years or more. The most ideal locations, reaching close to the theoretical maximum of ≈ 0.8 , include north Madagascar, Brazil, Peru and locations close to the southern ocean such as Tasmania, Kerguelen and Patagonia, where the gas exchange appears to occur faster than the surface residence time. The variation of the achievable CO₂ drawdown per unit alkalinity on timescales relevant to the climate change crisis and the speed at which equilibration is reached poses further difficulty for verification of CDR credits.

Further study to determine these uptake efficiencies, at a finer location sample resolution and ideally with model ensembles, are needed for optimal placement of OAE deployments. While our results give an overall picture and are indicative of the complexity, more sophisticated biogeochemistry models (e.g., Carroll et al., 2022) and higher-resolution regional coupled biogeophysical models (e.g., Sein et al., 2015; Wang et al., 2022) will be essential for simulation and deployment of real-world OAE projects. It would also be interesting to examine how the CDR efficiency and OAE limits change in different future

emission scenarios. At higher pCO₂ the surface ocean would be more acidic and thus a larger OAE rate could be sustained
475 without exceeding pre-industrial surface pH. Increased stratification may increase surface residence times, thus decreasing the equilibration time. Furthermore, changes in biological activity, general circulation patterns and atmospheric dynamics further complicate the picture. Such effects and their interplay could potentially be studied in a full earth system model under different emissions scenarios.

Code and data availability. Code and data will be made available on a publicly available repository upon final publication

480 *Author contributions.* JH and MDT designed the experiments, wrote the code, ran the simulations and wrote the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We would like to thank Chris Van Arsdale, Lennart Bach, Brendan Carter, Matt Eisaman and Matthew Long for many helpful comments on the manuscript.

References

- 485 Albright, R., Caldeira, L., Hosfelt, J., Kwiatkowski, L., Maclare, J. K., Mason, B. M., Nebuchina, Y., Ninokawa, A., Pon-
gratz, J., Ricke, K. L., and et al.: Reversal of ocean acidification enhances net coral reef calcification, *Nature*, 531, 362–365,
<https://doi.org/10.1038/nature17155>, 2016.
- Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K., Matsumoto, K., Munhoven, G., Montenegro, A., and Tokos, K.: Atmospheric Lifetime of Fossil Fuel Carbon Dioxide, *Annu. Rev. Earth Planet. Sci.*, 37, 117–134,
490 <https://doi.org/10.1146/annurev.earth.031208.100206>, 2009.
- Bach, L. T., Gill, S. J., Rickaby, R. E. M., Gore, S., and Renforth, P.: CO₂ Removal With Enhanced Weathering and
Ocean Alkalinity Enhancement: Potential Risks and Co-benefits for Marine Pelagic Ecosystems, *Frontiers in Climate*, 1, 7,
<https://doi.org/10.3389/fclim.2019.00007>, 2019.
- Burt, D. J., Fröb, F., and Ilyina, T.: The Sensitivity of the Marine Carbonate System to Regional Ocean Alkalinity Enhancement, *Frontiers in
495 Climate*, 3, <https://doi.org/10.3389/fclim.2021.624075>, 2021.
- Byrne, C., Law, R., Hudson, P., Thain, J., and Fileman, T.: Measurements of the dispersion of liquid industrial waste discharged into the
wake of a dumping vessel, *Water Research*, 22, 1577–1584, [https://doi.org/10.1016/0043-1354\(88\)90171-6](https://doi.org/10.1016/0043-1354(88)90171-6), 1988.
- Carroll, D., Menemenlis, D., Adkins, J. F., Bowman, K. W., Brix, H., Dutkiewicz, S., Fenty, I., Gierach, M. M., Hill, C., Jahn,
500 O., Landschützer, P., Lauderdale, J. M., Liu, J., Manizza, M., Naviaux, J. D., Rödenbeck, C., Schimel, D. S., Van der Stocken,
T., and Zhang, H.: The ECCO-Darwin Data-Assimilative Global Ocean Biogeochemistry Model: Estimates of Seasonal to Mul-
tidecadal Surface Ocean pCO₂ and Air-Sea CO₂ Flux, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001888,
<https://doi.org/https://doi.org/10.1029/2019MS001888>, e2019MS001888 2019MS001888, 2020.
- Carroll, D., Menemenlis, D., Dutkiewicz, S., Lauderdale, J. M., Adkins, J. F., Bowman, K. W., Brix, H., Fenty, I., Gierach, M. M., Hill,
C., Jahn, O., Landschützer, P., Manizza, M., Mazloff, M. R., Miller, C. E., Schimel, D. S., Verdy, A., Whitt, D. B., and Zhang, H.:
505 Attribution of Space-Time Variability in Global-Ocean Dissolved Inorganic Carbon, *Global Biogeochemical Cycles*, 36, e2021GB007162,
<https://doi.org/https://doi.org/10.1029/2021GB007162>, e2021GB007162 2021GB007162, 2022.
- Chou, H.-T.: On the dilution of liquid waste in ships' wakes, *Journal of Marine Science and Technology*, 1, 149–154,
<https://doi.org/10.1007/BF02391175>, 1996.
- Davies, P. A., Yuan, Q., and de Richter, R.: Desalination as a negative emissions technology, *Environ. Sci.: Water Res. Technol.*, 4, 839–850,
510 <https://doi.org/10.1039/C7EW00502D>, 2018.
- de Lannoy, C.-F., Eisaman, M. D., Jose, A., Karnitz, S. D., DeVaul, R. W., Hannun, K., and Rivest, J. L.: Indirect ocean capture of at-
mospheric CO₂: Part I. Prototype of a negative emissions technology, *International Journal of Greenhouse Gas Control*, 70, 243–253,
<https://doi.org/10.1016/j.ijggc.2017.10.007>, 2018.
- Digdaya, I. A., Sullivan, I., Lin, M., Han, L., Cheng, W.-H., Atwater, H. A., and Xiang, C.: A direct coupled electrochemical system for
515 capture and conversion of CO₂ from oceanwater, *Nat Commun*, 11, 362–365, <https://doi.org/10.1038/s41467-020-18232-y>, 2020.
- Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean Acidification: The Other CO₂ Problem, *Annual Review of Marine Science*,
1, 169–192, <https://doi.org/10.1146/annurev.marine.010908.163834>, pMID: 21141034, 2009.
- Dutkiewicz, S., Sokolov, A., Scott, J., and Stone, P.: A three-dimensional ocean-seaice-carbon cycle model and its coupling to a two dimen-
520 sional atmospheric model: Uses in climate change studies. Report 122, MIT Joint Program on the Science and Policy of Global Change ,
http://mit.edu/globalchange/www/MITJPSPGC_Rpt122.pdf, [Online; accessed 20-April-2021], 2005.

- Eisaman, M. D., Rivest, J. L., Karnitz, S. D., de Lannoy, C.-F., Jose, A., DeVaul, R. W., and Hannun, K.: Indirect ocean capture of atmospheric CO₂: Part II. Understanding the cost of negative emissions, *International Journal of Greenhouse Gas Control*, 70, 254–261, <https://doi.org/10.1016/j.ijgac.2018.02.020>, 2018.
- Fakhraee, M., Li, Z., Planavsky, N., and Reinhard, C.: Environmental impacts and carbon capture potential of ocean alkalinity enhancement, 525 *Nature Portfolio Under Review*, <https://doi.org/10.21203/rs.3.rs-1475007/v1>, 2022.
- Fassbender, A. J., Orr, J. C., and Dickson, A. G.: Technical note: Interpreting pH changes, *Biogeosciences*, 18, 1407–1415, <https://doi.org/10.5194/bg-18-1407-2021>, 2021.
- Feng, E. Y., Keller, D. P., Koeve, W., and Oschlies, A.: Could artificial ocean alkalinization protect tropical coral ecosystems from ocean acidification?, *Environ. Res. Lett.*, 11, 074008, <https://doi.org/10.1088/1748-9326/11/7/074008>, 2016.
- Feng, E. Y., Koeve, W., Keller, D. P., and Oschlies, A.: Model-Based Assessment of the CO₂ Sequestration Potential of Coastal Ocean Alkalization, *Earth's Future*, 5, 1252–1266, [https://doi.org/https://doi.org/10.1002/2017EF000659](https://doi.org/10.1002/2017EF000659), 2017.
- Ferdeiner, A., Chase, Z., Kennedy, F., Schulz, K. G., and Bach, L. T.: Assessing the influence of ocean alkalinity enhancement on a coastal phytoplankton community, *Biogeosciences Discussions*, 2022, 1–36, <https://doi.org/10.5194/bg-2022-17>, 2022.
- Fuhr, M., Geilert, S., Schmidt, M., Liebetrau, V., Vogt, C., Ledwig, B., and Wallmann, K.: Kinetics of Olivine Weathering in Seawater: An 535 Experimental Study, *Frontiers in Climate*, 4, <https://doi.org/10.3389/fclim.2022.831587>, 2022.
- Gernon, T. M., Hincks, T. K., Merdith, A. S., Rohling, E. J., Palmer, M. R., Foster, G. L., Bataille, C. P., and Müller, R. D.: Global chemical weathering dominated by continental arcs since the mid-Palaeozoic, *Nature Geoscience*, 14, 690–696, <https://doi.org/10.1038/s41561-021-00806-0>, 2021.
- Goldberg, D. S., Takahashi, T., and Slagle, A. L.: Carbon dioxide sequestration in deep-sea basalt, *Proceedings of the National Academy of 540 Sciences*, 105, 9920–9925, <https://doi.org/10.1073/pnas.0804397105>, 2008.
- González, M. F. and Ilyina, T.: Impacts of artificial ocean alkalinization on the carbon cycle and climate in Earth system simulations, *Geophys. Res. Lett.*, 43, 6493–6502, <https://doi.org/10.1002/2016gl068576>, 2016.
- Guo, J. A., Strzepek, R., Willis, A., Ferderer, A., and Bach, L. T.: Investigating the effect of nickel concentration on phytoplankton growth to assess potential side-effects of ocean alkalinity enhancement, *Biogeosciences*, 19, 3683–3697, 2022.
- Hangx, S. J. and Spiers, C. J.: Coastal spreading of olivine to control atmospheric CO₂ concentrations: A critical analysis of viability, 545 *International Journal of Greenhouse Gas Control*, 3, 757–767, <https://doi.org/10.1016/j.ijgac.2009.07.001>, 2009.
- Hartmann, J., Sutner, N., Lim, C., Schneider, J., Marín-Samper, L., Arístegui, J., Renforth, P., Taucher, J., and Riebesell, U.: Stability of alkalinity in Ocean Alkalinity Enhancement (OAE) approaches – consequences for durability of CO₂ storage, *Biogeosciences Discussions*, 2022, 1–29, <https://doi.org/10.5194/bg-2022-126>, 2022.
- House, K. Z., House, C. H., Schrag, D. P., and Aziz, M. J.: Electrochemical Acceleration of Chemical Weathering as an Energetically Feasible Approach to Mitigating Anthropogenic Climate Change, *Environ. Sci. Technology*, 41, 8464–8470, <https://doi.org/10.1021/es0701816>, 2007.
- Humphreys, M. P., Gregor, L., Pierrot, D., van Heuven, S. M. A. C., Lewis, E. R., and Wallace, D. W. R.: PyCO2SYS: marine carbonate system calculations in Python, <https://doi.org/10.5281/ZENODO.3744275>, 2020.
- Ilyina, T., Wolf-Gladrow, D., Munhoven, G., and Heinze, C.: Assessing the potential of calcium-based artificial ocean alkalinization to mitigate rising atmospheric CO₂ and ocean acidification, *Geophysical Research Letters*, 40, 5909–5914, 2013.
- IMCO: Procedures and arrangements for the discharge of noxious liquid substances. Method for calculation of dilution capacity in the ship's wake. IMCO document MEPC III-7, May 27, 1975.

- Jones, D. C., Ito, T., Takano, Y., and Hsu, W.-C.: Spatial and seasonal variability of the air-sea equilibration timescale of carbon dioxide, 560 Global Biogeochemical Cycles, 28, 1163–1178, [https://doi.org/https://doi.org/10.1002/2014GB004813](https://doi.org/10.1002/2014GB004813), 2014.
- Jones, S. D., Le Quéré, C., and Rödenbeck, C.: Autocorrelation characteristics of surface ocean pCO₂and air-sea CO₂ fluxes, Global Biogeochemical Cycles, 26, [https://doi.org/https://doi.org/10.1029/2010GB004017](https://doi.org/10.1029/2010GB004017), 2012.
- Keller, D. P., Feng, E. Y., and Oschlies, A.: Potential climate engineering effectiveness and side effects during a high carbon dioxide-emission scenario, Nature Communications, 5, 3304, 2014.
- 565 Keller, D. P., Lenton, A., Littleton, E. W., Oschlies, A., Scott, V., and Vaughan, N. E.: The Effects of Carbon Dioxide Removal on the Carbon Cycle, Current Climate Change Reports, 4, 250–265, <https://doi.org/10.1007/s40641-018-0104-3>, 2018.
- Kheshgi, H. S.: Sequestering atmospheric carbon dioxide by increasing ocean alkalinity, Energy, 20, 915–922, 570 [https://doi.org/https://doi.org/10.1016/0360-5442\(95\)00035-F](https://doi.org/10.1016/0360-5442(95)00035-F), 1995.
- Köhler, P., Abrams, J. F., Völker, C., Hauck, J., and Wolf-Gladrow, D. A.: Geoengineering impact of open ocean dissolution of olivine on atmospheric CO₂, surface ocean pH and marine biology, Environ. Res. Lett., 8, 014009, <https://doi.org/10.1088/1748-9326/8/1/014009>, 2013.
- Lauvset, S. K., Key, R. M., Olsen, A., van Heuven, S., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F. F., Suzuki, T., and Watelet, S.: A new global interior ocean mapped climatology: the 1°×1° GLODAP version 2, Earth System Science Data, 8, 325–340, <https://doi.org/10.5194/essd-8-325-2016>, 2016.
- 575 Lewis, E. R. and Wallace, D. W. R.: Program Developed for CO₂ System Calculations, <https://doi.org/10.15485/1464255>, 1998.
- Lewis, R.: The dilution of waste in the wake of a ship, Water Research, 19, 941–945, [https://doi.org/10.1016/0043-1354\(85\)90360-4](https://doi.org/10.1016/0043-1354(85)90360-4), 1985.
- Lewis, R. E. and Riddle, A. M.: Sea disposal: Modelling studies of waste field dilution, Marine Pollution Bulletin, 20, 124–129, 580 [https://doi.org/10.1016/0025-326x\(88\)90817-x](https://doi.org/10.1016/0025-326x(88)90817-x), 1989.
- Li, J. and Hitch, M.: Ultra-fine grinding and mechanical activation of mine waste rock using a high-speed stirred mill for mineral carbonation, Int J Miner Metall Mater, 22, 1005–1016, <https://doi.org/10.1007/s12613-015-1162-3>, 2015.
- Marshall, J., Adcroft, A., Hill, C., Perelman, L., and Heisey, C.: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers, Journal of Geophysical Research: Oceans, 102, 5753–5766, <https://doi.org/https://doi.org/10.1029/96JC02775>, 1997.
- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, 585 K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., eds.: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, for the Intergovernmental Panel on Climate Change, Cambridge, 2021.
- Matter, J. M., Broecker, W., Stute, M., Gislason, S., Oelkers, E., Stefánsson, A., Wolff-Boenisch, D., Gunnlaugsson, E., Axelsson, G., and Björnsson, G.: Permanent Carbon Dioxide Storage into Basalt: The CarbFix Pilot Project, Iceland, Energy Procedia, 1, 3641–3646, 590 <https://doi.org/10.1016/j.egypro.2009.02.160>, 2009.
- McGrail, B. P., Schaef, H. T., Ho, A. M., Chien, Y.-J., Dooley, J. J., and Davidson, C. L.: Potential for carbon dioxide sequestration in flood basalts, J. Geophys. Res., 111, n/a–n/a, <https://doi.org/10.1029/2005jb004169>, 2006.
- Metz, B. and Intergovernmental Panel on Climate Change, eds.: IPCC special report on carbon dioxide capture and storage, Cambridge University Press, for the Intergovernmental Panel on Climate Change, Cambridge, oCLC: ocm64949778, 2005.
- 595 Meysman, F. J. R. and Montserrat, F.: Negative CO₂ emissions via enhanced silicate weathering in coastal environments, Biology Letters, 13, 20160 905, <https://doi.org/10.1098/rsbl.2016.0905>, 2017.

- Middelburg, J. J., Soetaert, K., and Hagens, M.: Ocean Alkalinity, Buffering and Biogeochemical Processes, *Rev. Geophys.*, 58, <https://doi.org/10.1029/2019rg000681>, 2020.
- Montserrat, F., Renforth, P., Hartmann, J., Leermakers, M., Knops, P., and Meysman, F. J. R.: Olivine Dissolution in Seawater: Implications for CO₂Sequestration through Enhanced Weathering in Coastal Environments, *Environ. Sci. Technology*, 51, 3960–3972, <https://doi.org/10.1021/acs.est.6b05942>, 2017.
- Moras, C. A., Bach, L. T., Cyronak, T., Joannes-Boyau, R., and Schulz, K. G.: Ocean alkalinity enhancement – avoiding runaway CaCO₃ precipitation during quick and hydrated lime dissolution, *Biogeosciences*, 19, 3537–3557, 2022.
- Morse, J. W. and He, S.: Influences of T, S and PCO₂ on the pseudo-homogeneous precipitation of CaCO₃ from seawater: implications for whiting formation, *Marine Chemistry*, 41, 291–297, [https://doi.org/https://doi.org/10.1016/0304-4203\(93\)90261-L](https://doi.org/10.1016/0304-4203(93)90261-L), 1993.
- Olsen, A., Key, R. M., Lauvset, S. K., Kozyr, A., Tanhua, T., Hoppema, M., Ishii, M., Jeansson, E., Van Heuven, S. M. A. C., Jutterström, S., Schirnick, C., Steinfeldt, R., Suzuki, T., Lin, X., Velo, A., and Pérez, F. F.: Global Ocean Data Analysis Project, Version 2 (GLODAPv2) (NCEI Accession 0162565), <https://doi.org/10.7289/V5KW5D97>, 2017.
- Pan, Y., Li, Y., Ma, Q., He, H., Wang, S., Sun, Z., Cai, W.-J., Dong, B., Di, Y., Fu, W., and Chen, C.-T. A.: The role of Mg²⁺ in inhibiting CaCO₃ precipitation from seawater, *Marine Chemistry*, 237, 104 036, 2021.
- Penman, D. E., Caves Rugenstein, J. K., Ibarra, D. E., and Winnick, M. J.: Silicate weathering as a feedback and forcing in Earth's climate and carbon cycle, *Earth-Science Reviews*, 209, 103 298, <https://doi.org/https://doi.org/10.1016/j.earscirev.2020.103298>, 2020.
- Pokrovsky, O.: Precipitation of calcium and magnesium carbonates from homogeneous supersaturated solutions, *Journal of Crystal Growth*, 186, 233–239, [https://doi.org/https://doi.org/10.1016/S0022-0248\(97\)00462-4](https://doi.org/https://doi.org/10.1016/S0022-0248(97)00462-4), 1998.
- 615 Pokrovsky, O. and Savenko, V.: The role of magnesium at homogeneous precipitation of calcium carbonate from seawater, *Oceanology*, 34, 493–497, 1995.
- Pokrovsky, O. S.: Kinetics of CaCO₃ Homogeneous Precipitation in Seawater, *Mineralogical Magazine*, 58A, 738–739, [https://doi.org/https://doi.org/10.1180/minmag.1994.58a.2.121](https://doi.org/10.1180/minmag.1994.58a.2.121), 1994.
- Rau, G. H.: Electrochemical CO₂ capture and storage with hydrogen generation, *Energy Procedia*, 1, 823–828, <https://doi.org/https://doi.org/10.1016/j.egypro.2009.01.109>, *greenhouse Gas Control Technologies* 9, 2009.
- 620 Renforth, P.: The potential of enhanced weathering in the UK, *International Journal of Greenhouse Gas Control*, 10, 229–243, <https://doi.org/10.1016/j.ijgge.2012.06.011>, 2012.
- Renforth, P. and Henderson, G.: Assessing ocean alkalinity for carbon sequestration, *Reviews of Geophysics*, 55, 636–674, <https://doi.org/10.1002/2016rg000533>, 2017.
- 625 Riebesell, U. and Tortell, P. D.: Effects of Ocean Acidification on Pelagic Organisms and Ecosystems, in: *Ocean Acidification*, Oxford University Press, <https://doi.org/10.1093/oso/9780199591091.003.0011>, 2011.
- Rigopoulos, I., Harrison, A. L., Delimitis, A., Ioannou, I., Efstathiou, A. M., Kyratsi, T., and Oelkers, E. H.: Carbon sequestration via enhanced weathering of peridotites and basalts in seawater, *Applied Geochemistry*, 91, 197–207, <https://doi.org/https://doi.org/10.1016/j.apgeochem.2017.11.001>, 2018.
- 630 Rogelj, J., Popp, A., Calvin, K. V., Luderer, G., Emmerling, J., Gernaat, D., Fujimori, S., Strefler, J., Hasegawa, T., Marangoni, G., Krey, V., Kriegler, E., Riahi, K., van Vuuren, D. P., Doelman, J., Drouet, L., Edmonds, J., Fricko, O., Harmsen, M., Havlík, P., Humpenöder, F., Stehfest, E., and Tavoni, M.: Scenarios towards limiting global mean temperature increase below 1.5 °C, *Nature Climate Change*, 8, 325–332, <https://doi.org/10.1038/s41558-018-0091-3>, 2018.

- Sano, Y., Hao, Y., and Kuwahara, F.: Development of an electrolysis based system to continuously recover magnesium from seawater.,
635 Heliyon, 4, e00923, 2018.
- Sarmiento, J. L. and Gruber, N.: Ocean biogeochemical dynamics, Princeton University Press, Princeton, oCLC: ocm60651167, 2006.
- Schulind, R. D. and de Boer, P. L.: Rolling stones; fast weathering of olivine in shallow seas for cost-effective CO₂ capture and mitigation
of global warming and ocean acidification, Earth System Dynamics Discussions, 2, 551–568, <https://doi.org/10.5194/esdd-2-551-2011>,
2011.
- 640 Sein, D. V., Mikolajewicz, U., Gröger, M., Fast, I., Cabos, W., Pinto, J. G., Hagemann, S., Semmler, T., Izquierdo, A., and Jacob, D.:
Regionally coupled atmosphere-ocean-sea ice-marine biogeochemistry model ROM: 1. Description and validation, Journal of Advances
in Modeling Earth Systems, 7, 268–304, [https://doi.org/https://doi.org/10.1002/2014MS000357](https://doi.org/10.1002/2014MS000357), 2015.
- Subhas, A. V., Marx, L., Reynolds, S., Flohr, A., Mawji, E. W., Brown, P. J., and Cael, B. B.: Microbial ecosystem responses to alkalinity
enhancement in the North Atlantic Subtropical Gyre, Frontiers in Climate, 4, 2022.
- 645 Sun, W., Jayaraman, S., Chen, W., Persson, K. A., and Ceder, G.: Nucleation of metastable aragonite CaCO₃ in seawater,
Proceedings of the National Academy of Sciences, 112, 3199–3204, <https://doi.org/10.1073/pnas.1423898112>, 2015.
- Thorsen, T. G., Hagen, R. I., Wærnes, O., and Langseth, B.: Method for production of magnesium hydroxide from sea water, <https://patentscope.wipo.int/search/en/detail.jsf?docId=WO2000029326>, 2000.
- Tyka, M. D., Van Arsdale, C., and Platt, J. C.: CO₂ capture by pumping surface acidity to the deep ocean, Energy Environ. Sci., 15, 786–798,
650 <https://doi.org/10.1039/D1EE01532J>, 2022.
- van Sebille, E., Griffies, S. M., Abernathey, R., Adams, T. P., Berloff, P., Biastoch, A., Blanke, B., Chassignet, E. P., Cheng, Y., Cotter, C. J.,
Deleersnijder, E., Döös, K., Drake, H. F., Drijfhout, S., Gary, S. F., Heemink, A. W., Kjellsson, J., Koszalka, I. M., Lange, M., Lique, C.,
MacGilchrist, G. A., Marsh, R., Mayorga Adame, C. G., McAdam, R., Nencioli, F., Paris, C. B., Piggott, M. D., Polton, J. A., Rühs, S.,
Shah, S. H., Thomas, M. D., Wang, J., Wolfram, P. J., Zanna, L., and Zika, J. D.: Lagrangian ocean analysis: Fundamentals and practices,
655 Ocean Modelling, 121, 49–75, <https://doi.org/https://doi.org/10.1016/j.ocemod.2017.11.008>, 2018.
- Wang, H., Pilcher, D. J., Eisaman, M. D., and Carter, B. R.: 2022.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, Journal of Geophysical Research: Oceans, 97, 7373–
7382, <https://doi.org/https://doi.org/10.1029/92JC00188>, 1992.
- Wunsch, C. and Heimbach, P.: Chapter 21 - Dynamically and Kinematically Consistent Global Ocean Circulation and Ice State Estimates, in:
660 Ocean Circulation and Climate, edited by Siedler, G., Griffies, S. M., Gould, J., and Church, J. A., vol. 103 of *International Geophysics*,
pp. 553–579, Academic Press, <https://doi.org/https://doi.org/10.1016/B978-0-12-391851-2.00021-0>, 2013.
- Wunsch, C., Heimbach, P., Ponte, R. M., Fukumori, I., and Members, T. E.-G. C.: The Global General Circulation of the Ocean Estimated
by the ECCO-Consortium, Oceanography, <https://doi.org/10.5670/oceanog.2009.41>, <p>Following on the heels of the World Ocean Cir-
665 culation Experiment, the Estimating the Circulation and Climate of the Ocean (ECCO) consortium has been directed at making the best
possible estimates of ocean circulation and its role in climate. ECCO is combining state-of-the-art ocean general circulation models with
the nearly complete global ocean data sets for 1992 to present. Solutions are now available that adequately fit almost all types of ocean
observations and that are, simultaneously, consistent with the model. These solutions are being applied to understanding ocean variability,
biological cycles, coastal physics, geodesy, and many other areas.</p>, 2009.
- Zeebe, R. E. and Wolf-Gladrow, D. A.: CO₂ in seawater: Equilibrium, kinetics, isotopes: Volume 65, Elsevier Oceanography Series, Elsevier
670 Science, London, England, 2001.
- Zhang, H., Menemenlis, D., and Fenty, I.: ECCO LLC270 Ocean-Ice State Estimate, <https://dspace.mit.edu/handle/1721.1/119821>, 2018.