

Dansgaard-Oeschger events in climate models: Review and baseline MIS3 protocol

Irene Malmierca-Vallet¹, Louise C. Sime¹, and the D-O community members⁺

¹British Antarctic Survey, Cambridge, U.K.

⁺A full list of authors appears at the end of the paper

Correspondence: Irene Malmierca-Vallet (irealm37@bas.ac.uk)

Abstract.

Dansgaard-Oeschger (D-O) events, millennial-scale climate oscillations between stadial and interstadial conditions (of up to 10-15°C in amplitude at high northern latitudes), occurred throughout the Marine Isotope Stage 3 (MIS3; 27.8 – 59.4 ka) period. The climate modelling community up to now has not been able to answer the question: Are our climate models too stable to simulate D-O events? To address this, this manuscript lays the ground-work for a MIS3 D-O protocol for general circulation models which are used in the International Panel for Climate Change (IPCC) assessments. We review: D-O terminology, community progress on simulating D-O events in these IPCC-class models (processes and published examples), and evidence about the boundary conditions under which D-O events occur. We find that no model exhibits D-O like behaviour under pre-industrial conditions. Some, but not all, models exhibit D-O like oscillations under MIS3 and/or full glacial conditions. Greenhouse gases and ice-sheet configurations are crucial. However most models have not run simulations of long enough duration to be sure which models show D-O like behaviour, under either MIS3 or full glacial states. We propose a MIS3 baseline protocol at 34 ka, which features low obliquity values, medium-to-low MIS3 greenhouse gas values and the intermediate ice-sheet configuration which our review suggests are most conducive to D-O like behaviour in models. We also provide a protocol for a second freshwater (Heinrich-Event preconditioned) experiment, since previous work suggests that this variant may be helpful in preconditioning a state in models which is conducive to D-O events. This review provides modelling groups investigating MIS3 D-O oscillations with a common framework, which is aimed at 1) maximising the chance of the occurrence of D-O like events in the simulations; 2) allowing more precise model-data evaluation and; 3) providing an adequate central point for modellers to explore model stability.

1 Introduction

During a Dansgaard-Oeschger (D-O) event, Greenland transitions between cold stadial (GS) and warmer Greenland Interstadial (GI) conditions. The warming can occur within a decade (Kindler et al., 2014; Huber et al., 2006), whilst cooling occurs over a much longer period that is typically several centuries in length. During a warming phase, surface air temperatures over Greenland increase by 10-15°C (Andersen et al., 2006; Kindler et al., 2014; Huber et al., 2006). D-O events are best documented during Marine Isotope Stage 3 (MIS3; between 27.8 – 59.4 thousand of years BP, hereafter ka Goni and Harrison,

25 2010), including being recorded in several ice cores from Greenland (Fig. 1 Johnsen et al., 2001). Whilst the D-O event
recorded in these cores are renowned, the events are global in nature (Voelker et al., 2002; Sanchez Goñi and Harrison, 2010;
Sánchez Goñi et al., 2017), with known climate signatures including imprints in surface temperature and the hydrological
cycle at high northern latitudes (Andersen et al., 2004; Thomas et al., 2009; Seierstad et al., 2014), in the tropics (Deplazes
30 2017; Rousseau et al., 2017), and in North and South America (Wang et al., 2004; Wagner et al., 2010; Asmerom et al., 2010;
Deplazes et al., 2013; Vanneste et al., 2015). While there are no Greenland ice core records of the previous glacial (MIS6
around 140-190 ka), speleothems and Antarctic ice cores indicate that it is extremely likely that D-O events also occurred
during MIS6 and earlier glacial periods (Lang et al., 1999; Uriarte, 2019; Landais et al., 2004; Turner and Marshall, 2011;
Barker et al., 2011; Lambert et al., 2012). This observational evidence shows that D-O events do not occur under interglacial
35 or full Last Glacial Maximum conditions (Galaasen et al., 2014; Tzedakis et al., 2018; Galaasen et al., 2020).

In 2011, Valdes (2011) argued that climate models used in the assessments of the Intergovernmental Panel on Climate
Change (IPCC) have not proved their ability to simulate D-O events. This has several implications for the delivery of accurate
projections of climate change, within the context of tipping points and abrupt climate change (Brovkin et al., 2021). Whilst
in the intervening years a number of models have captured key features of D-O events through AMOC hysteresis behaviour
40 and/or produced D-O type millennial-scale variability under a range of forcings (Brown and Galbraith, 2016; Galbraith and
de Lavergne, 2019; Klockmann et al., 2018; Peltier et al., 2020; Armstrong et al., 2021; Zhang et al., 2021; Vettoretti et al.,
2022), we still do not know if climate models are too stable because too few models have run and published an appropriate
simulation. This deficiency is related to both the computational expense which prevents models from being run for the longer
time periods needed for investigating D-O events and to the lack of an agreed appropriate experimental set-up. The limited
45 knowledge of pre-Last Glacial Maximum (LGM) boundary conditions, in particular in the case of the ice sheet height and
distribution, makes it challenging to generate an appropriate MIS3 experimental set-up.

An important question is if model stability is caused by the model parameters and MIS3 conditions are such that the models
are in a mono-stable state, in an oscillatory state or if the models exhibits bi-modality where noise-induced transitions are not
induced due to too low model variability (Ditlevsen and Johnsen, 2010). Previous studies have questioned the significance
50 of the periodic occurrence of DO events in MIS3 (~ 1470) (Ditlevsen et al., 2007). If the full glacial period is included, the
distribution of waiting time between DO-events is consistent with a random process (Ditlevsen et al., 2005). Durations of
stadials vs. interstadials indicate correlations with global ice volume and orbital parameters (Lohmann and Ditlevsen, 2018;
Mitsui and Crucifix, 2017), thus underpinning the decision to focus on MIS3 boundary configurations.

Whether models can simulate abrupt changes is a crucial research question: if the current IPCC-class models are too stable
55 to simulate D-O events, their ability to predict future abrupt transitions, and their use in identifying tipping points is doubtful.
For example, a tipping point may have been recently reached in the Arctic's Barents Sea (Barton et al., 2018; Tesi et al., 2021);
sea ice loss in the area is linked with enhanced heat transport via an intensified throughflow, or "Atlantification" (Årthun et al.,
2012; Polyakov et al., 2017). In addition, future enhanced precipitation, decline in Arctic sea ice and melting of glaciers and ice
sheets could intensify the supply of freshwater to the North Atlantic and Arctic which could lead to the reorganization of the

60 Atlantic circulation and tip the energy distribution between South and North in a similar way as occurred during D-O events (Lenton et al., 2008). If climate models do not reliably simulate past tipping events, it suggests that simulations of the coming century may be giving us a false sense of security.

Coupled Model Intercomparison Project (CMIP) coordinates and designs climate model protocols for the past, present and future climates, and has become an indispensable tool to facilitate our understanding of climate change (IPCC, 2013; Eyring et al., 2016). The Paleoclimate Model Intercomparison Project 4 (PMIP4) is one of the individual Model Intercomparison Projects which took part in CMIP6 (Kageyama et al., 2018). The design of a common MIS3 experimental protocol would allow the modelling community to address the questions posed above.

This manuscript compiles current information about unforced D-O like oscillations in CMIP5/CMIP6 models and discusses the boundary conditions and mechanisms responsible for these oscillations. Given the nomenclature on D-O events varies throughout the literature. Firstly, Table 1 and Figure 1 provide a framework for a more consistent terminology for use within this proposed MIS3 DO protocol. Secondly, we review the literature to ascertain whether models reproduce D-O like events under MIS3, or other, climate conditions. We then use this information to develop a protocol for the simulations of D-O events. This protocol focuses on Marine Isotope Stage 3 (MIS3) partly because of the excellent records of D-O events during this period (Schulz et al., 1999) but also because, as our synthesis shows, MIS3 conditions are also conducive to promoting D-O like events in some models. Given that D-O events did not occur under full glacial conditions in the last glacial period, the proposed modelling protocol is an important improvement on the use of an LGM PMIP protocol. It will undoubtedly help to shed light on the mechanism and processes involved in millennial-scale oscillations during MIS3. The common MIS3 climate modelling protocol is aimed at: 1) maximising the chance of the occurrence of D-O like events in the simulations; 2) improving model-data evaluation and; 3) providing an adequate central point for modellers to also explore model stability. In addition to the protocol for a baseline simulation, we also outline a protocol for a Heinrich-Event preconditioned (freshwater) experiment. These protocols provide a common framework for model experiments to explore cold-period instabilities using commonly specified greenhouse gas (GHG), ice sheet, insolation, and freshwater-related forcings.




Figure 1. MIS3 ice core records and nomenclature. Stable water isotope and CO₂ measurements from Antarctic and Greenland ice cores (Bauska et al., 2021; NGRIP Project Members, 2004; Kindler et al., 2014). See also Table 1 for D-O nomenclature. The "cnt" red box indicates the 38 to 32 ka period proposed for the MIS3 baseline experiment.

Table 1: Terminology

Term	Description
Abrupt change	We follow the IPCC Assessment Report 4 (IPCC AR4) definition of abrupt event/change (Meehl et al., 2007; Brovkin et al., 2021). This term refers to a large-scale change, which is much faster than the change in the pertinent forcing (e.g. rising atmospheric CO ₂ concentrations).
Tipping point	This term refers to a critical threshold at which a small perturbation can qualitatively modify the development or state of a system (Lenton et al., 2008).
Tipping element	This term describes large-scale components of the Earth system that could pass a tipping point (Lenton et al., 2008). Earth system components are the ocean, atmosphere, cryosphere, anthroposphere and biosphere, which have further important sub-components e.g. the meridional ocean circulation, the monsoon systems, sea ice, and various ecosystems (Brovkin et al., 2021).
D-O event	During the Last Glacial period, a series of dramatic climatic fluctuations occurred in the North Atlantic. These are known as D–O events, during which atmospheric and oceanic conditions alternated between relatively mild (interstadial) and full glacial (stadial) conditions (Dansgaard et al., 1982; Johnsen et al., 1992). Around 25 abrupt transitions (each completed within a decade) from stadial to interstadial conditions occurred during the Last Glacial period and their amplitude vary from 5 to 16°C (Landais et al., 2004; Huber et al., 2006; Kindler et al., 2014). The duration of interstadials varies from approximately a century to many millennia (Rasmussen et al., 2014).
D-O type oscillations	For the purpose of this MIS3 DO protocol, the term of D-O type oscillation refers to D-O scale climate variability reproduced by climate models, comparable to the D-O events observed in the Greenland ice core record.
Greenland Stadial and Interstadial	We follow the INTIMATE (Integration of Ice core, Marine and Terrestrial records of the North Atlantic) definition of stadial/interstadial terms (Rasmussen et al., 2014). The Greenland Interstadials (GI) and Greenland Stadials (GS) periods terms are the Greenland expressions of the D-O events and represent warm and cold phases of the NA area, respectively.
Heinrich events	These are defined by the presence of layers of ice-rafted debris (IRD) of primarily (not exclusively) Laurentide origin in North Atlantic sediment cores (e.g. Heinrich, 1988; Hemming, 2004). Heinrich events have been observed during some of the longer stadials, but likely do not cover the entire period of these longer stadials (Roche et al., 2004; Marcott et al., 2011).
Heinrich Stadial (HS)	This term refers to a stadial containing a specific Heinrich Event. Rasmussen et al. (2014) indicates that the term of HS can refer to the complete stadial period, or to part of a stadial only, characterized by changes shown in proxies of IRD, AMOC or SST (Barker et al., 2009).
Bond cycle	D-O events tend to follow a pattern of diminishing amplitude (or a general cooling trend of the GSs) following each HE (Bond et al., 1992; Alley, 1998; Alley et al., 1999; Clark et al., 2007; Rousseau et al., 2022). These cycles of HE grouped D-O events were named Bond cycles by Broecker (1994) and Alley (1998). The average gap between HEs is around 7 ka, so this is the average length of a Bond Cycle (Clark et al., 2007).

2 Review of spontaneous D-O type quasi-oscillations in coupled climate models

85 We compile published evidence of long unforced quasi-oscillations (in the Atlantic Meridional Circulation; AMOC) in IPCC-class models under all climate states in Table A1, alongside glacial boundary condition simulations which do not show D-O

type oscillations (Table A3). This permits us to explore the questions of: what proportion of models exhibit D-O like behaviour; which boundary conditions are most conducive to this; and what mechanisms are common to the modelled D-O behaviours. A number of PI/present-day model simulations exhibit spontaneous centennial-length cold events (Table A1), however, they do not appear to be D-O like events. We deal with these first.

Under pre-industrial greenhouse gas (GHG) forcing and present-day ice sheets, spontaneous centennial-length cold events that last around 100-200 years occur in four IPCC-class models (Table A1). EC-Earth and Community Climate System Model version 4 (CCSM4) show high atmospheric blocking over the eastern subpolar gyre that causes a cold event under pre-industrial boundary conditions (Drijfhout et al., 2013; Kleppin et al., 2015, Table A1). ECHAM6-FESOM also produces cooling events under pre-industrial conditions due to sudden reductions of deep water convection and increase of sea ice cover in the Labrador Sea (Sidorenko et al., 2015). Changes in convection also occur in the Kiel Climate Model (KCM; Martin et al., 2015), however here centennial-scale variability of the AMOC is linked to variability in Southern Ocean convection. Unlike the CCSM4 and the EC-Earth models, the KCM and ECHAM6-FESOM studies do not indicate an active role of the atmosphere. Although these four models all show abrupt spontaneous cooling events under pre-industrial boundary conditions, these events do not have the typical saw-tooth characteristics, or longer timescales, of D-O type events.

Regular cycles of D-O type quasi-oscillations are found in UofT CCSM4 under LGM boundary conditions (Peltier and Vettoretti, 2014). The initiation of the abrupt D-O type warming events is associated with the opening of a large polynya over the Irminger Sea (Vettoretti and Peltier, 2016) (Table A1). The AMOC spontaneously exhibits D-O like quasi-oscillations (Peltier et al., 2020). The Peltier et al. (2020) salt oscillator is maintained by the salinity gradient between the subtropical gyre and the Northern North Atlantic. Although UofT CCSM4 is the only model to show long unforced quasi-oscillations in the AMOC under full glacial conditions, most of the other PMIP4 LGM simulations (Kageyama et al., 2021a) have not been run long enough to be sure that such oscillations would not arise if they were run for longer (see Table B1). Having said that, ideally models should not show oscillatory D-O type behaviour when configured under a full glacial climate state, given that in reality D-O events do not occur under full glacial conditions (Huber et al., 2006; Galaasen et al., 2014; Kindler et al., 2014; Tzedakis et al., 2018).

D-O type quasi-oscillations are also found in MIROC4m under mid-glacial conditions (Kuniyoshi et al., 2022). Some aspects of the D-O warming mechanism observed in the UofT-CCSM4, in particular the spatial location of the opening of a big polynya in the Irminger Sea, determining the stadial-interstadial transition, is also identified in MIROC4m (Kuniyoshi et al., 2022) (Table A1).

Under late glacial conditions, at 30 ka, a quasi-oscillating AMOC is produced by the HadCM3 model (Armstrong et al., 2021) and results from a North Atlantic salt oscillator mechanism similar to that in UofT CCSM4 (Peltier and Vettoretti, 2014; Vettoretti and Peltier, 2016; Peltier et al., 2020). The HadCM3 model also shows millennial-scale climate oscillations triggered by deglacial meltwater discharge in LGM simulations (Romé et al., 2022). Under intermediate glacial conditions (MIS3: 40-32 ka), the COSMOS model shows spontaneous millennial-scale climate oscillations triggered solely by orbitally driven insolation changes (Zhang et al., 2021). Variations in either obliquity or eccentricity-modulated precession lead to climate variations over the tropical and subpolar North Atlantic which exert opposite effects on AMOC strength, and hence result in an oscillatory

climate regime (Zhang et al., 2021). The CM2Mc model also produces somewhat smoothed quasi-oscillating AMOC under intermediate MIS3-like boundary conditions, with a present-day ice sheet distribution in combination with a CO₂ concentration of 180 ppm and low obliquity (22°) (Brown and Galbraith, 2016; Galbraith and de Lavergne, 2019) (Table A1). The MPI-ESM model exhibits more abrupt D-O like quasi-oscillations with a present-day ice sheet distribution in combination with CO₂ concentrations ranging between 190-217 ppm (Table A1; Klockmann et al., 2018, 2020).

In contrast to the above, neither NorESM nor CCSM3 produce D-O type events or quasi-oscillations under MIS3 conditions (38 ka) (Table A3; Guo et al. (2019b); Zhang and Prange (2020)). The NorESM - with a reasonably simulated AMOC and Arctic sea ice distribution in the PI and historical simulations (as documented by Guo et al. (2019a)) - simulates a MIS3 climate that is in a stable regime with relatively strong convections in the Norwegian and Labrador seas. Indeed, NorESM sensitivity experiments including large reductions in atmospheric CO₂ levels and Laurentide Ice Sheet heights, aimed at perturbing the system into a cold stadial-like climate, indicate that the model state appears to be far from a possible threshold (Guo et al., 2019b). Zhang and Prange (2020) use the LGM ICE-5G ice sheet configuration (Peltier, 2004), with a high Laurentide Ice Sheet (at just over 4000 m) which may have contributed to a strong AMOC in the CCSM3 simulation, alongside its particular background climate.

In summary, IPCC-class models set up with pre-industrial or present-day conditions do not exhibit D-O type warming events, but can feature shorter centennial length cooling and warming events. This model behaviour is consistent with observations, since millennial timescale D-O events do not occur under interglacial conditions but periods of centennial-scale AMOC variability are present throughout several interglacials (Galaasen et al., 2014; Tzedakis et al., 2018; Galaasen et al., 2020). Some models which are set up with more MIS3 like conditions exhibit D-O type warming events, but some do not. Under full LGM conditions only one model (UoT-CCSM4) out of ten (PMIP4 LGM simulations: Kageyama et al. (2021a)) show spontaneous D-O type oscillations (Tables A1 and B1; Kageyama et al., 2021a; Peltier and Vettoretti, 2014).

Since it can take some time for D-O type oscillations to evolve, it is unclear if some models would develop such oscillations if they were run for longer (at least for 2000 model years). Of the forty LGM/MIS3-like simulations (Table B1; Kageyama et al., 2021a; Armstrong et al., 2021; Klockmann et al., 2018), sixteen simulations have been run for less than 2000 years (Table B1), which makes it difficult to tell whether any of these simulations are capable of, or likely to, exhibit D-O like behaviour under specific boundary conditions. In addition the duration of LGM/MIS3 simulations is currently inadequate, we note that the majority of CMIP6 models appear not to have performed any form of glacial period simulation (Table C1). Thus, it is difficult to ascertain what proportion, or indeed which, models are capable of capturing D-O like behaviour, under any form of glacial period state (Table C1).

2.1 The role of ocean and sea-ice feedbacks

Changes in the AMOC are crucial to the correct simulation of D-O events (Broecker and Peteet, 1985). The AMOC features stabilising positive feedbacks: a strong AMOC transports warm and salty water into the subpolar North Atlantic, thus weakening the stratification and also keeping the sea ice cover reduced (*e.g.* Rahmstorf, 2002; Clark et al., 2002). As a consequence, there is a large transport of heat northward across the hemispheres (*e.g.* Feulner et al., 2013; Buckley and Marshall, 2016),




Figure 2. Schematic depicting the transition from GS to GI conditions *i.e.* a D-O warming event.




Figure 3. Schematic depicting the transition from GI to GS conditions *i.e.* a D-O cooling event.

strong heat loss in the North Atlantic and Arctic, and active deep convection that sustains the strong AMOC. A weak AMOC, on the other hand, is associated with a weaker northward transport of salt and heat. This increases the stratification in the subpolar North Atlantic and thus favors the expansion of sea ice. The weak northward heat transport and the insulating effect of the sea ice keep the density gain due to heat loss small and the AMOC in a weak state (*e.g.* Klockmann et al., 2018). This weak AMOC state is stable when Antarctic Bottom Water becomes dense and salty enough to replace North Atlantic Deep Water (NADW) in the deep North Atlantic.

Sea ice can act as both a slow and fast positive feedback on AMOC-induced changes in climate. Extensive stadial sea ice cover during a weak AMOC state cools Greenland and suppresses atmosphere-ocean exchange of heat and oceanic convection in the North Atlantic (Li et al., 2005, 2010). This also leads to a slow build up of heat in the North Atlantic subsurface. Foraminifera from marine sediment cores offer evidence to back-up that this sub-surface warming occurred before the onset of fast D-O warming events (Rasmussen and Thomsen, 2004; Singh et al., 2014; Dokken et al., 2013). This heat build-up sets up the conditions for subsequent fast losses of GS sea ice.

Wind-driven, AMOC, and sea-ice linked salinity changes also play a crucial role in D-O positive and negative feedbacks. Indeed the net freshwater transport in the Atlantic basin by the AMOC can be used to assess the stability regime of the AMOC (Rahmstorf, 1995; Huisman et al., 2010). The interaction of subpolar and tropical salinity anomalies at the surface and in the subsurface (Jackson and Vellinga, 2013), and possible roles of the intertropical convergence zone and freshwater export through the Fram Strait, are also important in D-O related salinity feedbacks. Klockmann et al. (2018) note that if the subtropical gyre shifts northward and the sub polar gyre contracts, an inflow of salty subtropical water extends over the entire Atlantic basin east of the Mid-Atlantic Ridge. This inflow can supply salty water to the deep-convection sites in the Iceland Basin and Irminger Sea, and help maintain continuous deep convection and a strong AMOC even at low CO₂ concentrations (Brown and Galbraith, 2016; Klockmann et al., 2018; Guo et al., 2019b; Muglia and Schmittner, 2015; Sherriff-Tadano et al., 2018), thus preventing the initiation of GS-like conditions. Where the AMOC does enter a weak state for a prolonged period, and the climate enters a GS, a build-up of heat in subsurface waters and salt in the tropical Atlantic can enable the very rapid resumption of the AMOC (Lynch-Stieglitz, 2017), with the upward mixing of heat from the subsurface and importation of salt from the tropic Atlantic via gyre mechanisms (Peltier and Vettoretti, 2014).

The importance of vertical (diapycnal) mixing in the ocean for these long timescale, D-O type, instabilities has long been recognised (Welander, 1982). However, we note that the different ocean- and climate-models (Table A1) parameterise diapycnal mixing in very different ways (*e.g.* Nilsson et al., 2003; de Lavergne et al., 2019). The lack of a single consistent parameterisation and differences in the strength of diapycnal mixing across climate models means it is to be expected that some models will produce D-O like oscillations “out of the box” under MIS3 boundary conditions, but others may require changes, or tuning, to their diapycnal mixing parameters. Even within the same model, a large range of diapycnal diffusivities may yield steady states that satisfy common plausibility constraints such as AMOC transport and sea ice distribution (see *e.g.* Holden et al., 2010). This is partly because wind-driven Southern Ocean upwelling plays a complementary role to diapycnal mixing in setting the steady state overturning (Samelson, 2004), and partly because surface buoyancy forcing controls the relative strength of the upper (Atlantic) and lower (Antarctic) overturning cells (Oliver and Edwards, 2008). A realistic AMOC

transport may be obtained due to compensating biases in these processes, which has serious implications for whether AMOC feedbacks (necessary for capturing D-O behaviour) are represented in an adequate manner within these models.

Figure 2 and 3 show some of the key states, processes, and ocean sea-ice feedbacks that enable D-O events. Following Lohmann and Ditlevsen (2019), D-O events can be broken down into four periods: (1) cold stadial state (Fig. 2a), (2) rapid warming phase governed by very fast-time-scale mechanisms (Fig. 2b), (3) warm interstadial state (Fig. 2c and Fig. 3a) and, (4) gradual cooling phase (Fig. 3b) followed by a faster abrupt transition into a cold stadial phase (Fig. 3c). For some of the D-O events, the magnitude of the warming transitions are on the order of ten degrees in a decade, while the slow cooling in the interstadials is on the order of a few degrees in a millennium (the sawtooth shape) (Lohmann and Ditlevsen, 2019). This picture of rapid retreat of North Atlantic sea ice (Spolaor et al., 2016; Dokken et al., 2013) associated with the resumption of convection and the AMOC, alongside an upwards mixing of salt and heat, followed by a slower cooling phase back into stadial conditions matches accumulation, temperature, and water isotopes retrieved from Greenland ice core records of D-O warming events (Li et al., 2005, 2010; Sime et al., 2019).

2.2 The role of Northern Hemisphere Ice Sheets

Section 2 and Table A1 suggest that large Northern Hemisphere Ice Sheets and the wind regime associated with these can contribute to a strong AMOC which stabilises the North Atlantic and prevents D-O events. Thus ice sheets have a critical role to play in setting up the conditions for D-O events (Zhang et al., 2014; Klockmann et al., 2018; Brown and Galbraith, 2016; Muglia and Schmittner, 2015; Sherriff-Tadano et al., 2018). Figure 4 and 5 show some of the key mechanisms and feedbacks that are behind a state of reduced likelihood for D-O events and a potentially D-O type oscillating state, respectively.

The Northern Hemisphere Eurasian ice sheet was most probably limited to mountainous areas during mid-MIS3 (Helmens, 2014; Hughes et al., 2016), and its impact on D-O dynamics was probably relatively small. However, the size and presence (or absence) of the Laurentide ice sheet (LIS), which has elevations reaching a maximum of approximately 3000 m (Abe-Ouchi et al., 2015) at the LGM, does appear to cause important and robust (across multiple models) changes to Northern Hemisphere atmospheric circulation and resultant wind forcing of the ocean. LIS-dependent wind changes influence the subpolar gyre and the stability of the atmosphere-ice-ocean coupled system (Li and Born, 2019; Zhang et al., 2014).

A larger LIS (especially its height) causes stronger Northern Hemisphere winds (Li and Battisti, 2008; Pausata et al., 2011; Hofer et al., 2012; Ullman et al., 2014; Löfverström et al., 2014; Merz et al., 2015); an amplified stationary wave over North America (Manabe and Broccoli, 1985; Cook and Held, 1988); the North Atlantic glacial jet to be more stable due to differences in wave-mean flow feedbacks (Riviere et al., 2010); and alters variability of the large-scale atmospheric circulation, especially in the North Atlantic (Justino and Peltier, 2005; Pausata et al., 2009; Riviere et al., 2010). In addition, LIS height could control the sea-ice coverage and gyre circulation by shifting the westerlies over the North Atlantic region Zhang et al. (2014).

LIS altered winds that have wide implications for D-O relevant tipping elements (Seager and Battisti, 2007; Wunsch, 2006). Li and Born (2019) note that, first, the presence of a large LGM-type LIS is linked to a strong, more zonal and equatorward-shifted North Atlantic jet which weakens atmospheric heat transport into the North Atlantic (van der Schrier et al., 2010)




Figure 4. Schematic showing a state of reduced likelihood for D-O events.

225 and favours episodes of Greenland blocking (Madonna et al., 2017). Both could trigger the atmosphere-ice-ocean feedbacks that cause abrupt climate change in this area. Second, a steadier and stronger North Atlantic jet strengthens the wind-driven component of the subpolar gyre (Li and Born, 2019). Given that at latitudes north of about 45N, the subpolar gyre, which is essentially wind-driven, plays a crucial role in the northward transport of heat and salt, and is strongly linked to the AMOC (e.g. Jungclaus et al., 2013), wind-driven changes in this gyre have a strong impact on the density gain in the North Atlantic.

230 In many simulations with a large LIS (LGM-like ice sheets), the subtropical gyre can shift northward and cause an inflow of salty subtropical water over deep-convection sites, contributing to continuous deep convection and a strong AMOC even at low CO₂ concentrations (Brown and Galbraith, 2016; Klockmann et al., 2018; Guo et al., 2019b; Muglia and Schmittner, 2015; Sherriff-Tadano et al., 2018; Zhang et al., 2014). Similarly, Zhang et al. (2014) note that a higher LIS can promote less South Labrador Sea sea ice export to northeastern North Atlantic (which reduces sea ice concentration) to permit deep convection and shift the core of westerlies northwards, strengthening subtropical gyre for heat and salt transport (Zhang et al., 2014).

235 For these reasons, large LGM-type ice sheets, particular a large LIS, tend to lead to a density gain over the North Atlantic and the northward salt transport is enhanced with respect to the PI ice sheet case. For many, but not all models, this tends to lead to more active convection in the North Atlantic and a strong AMOC (across a wide range of CO₂ concentrations). That said, the AMOC in many LGM simulations is likely too strong (Klockmann et al., 2018; Kageyama et al., 2021b). Thus the




Figure 5. Schematic of a potentially D-O type oscillating state.

240 AMOC is far away from a tipping point with LGM-size ice sheets for many models (Zhang et al., 2014; Klockmann et al., 2018; Guo et al., 2019b).

In some simulations with reduced ice sheets, the jet stream shifts northwards, leading to regional cooling and a rise in seasonal sea ice concentration over the subpolar gyre region (Armstrong et al., 2021). This freshens the area and lowers deep-water formation, which weakens the subpolar gyre and as a result the simulations are more prone to enter a weak convection, 245 weak AMOC mode which is conducive to D-O type oscillations (Klockmann et al., 2018; Armstrong et al., 2021). Thus, with intermediate MIS3 LIS, *i.e.* reduced in its height compared to the LGM, multiple AMOC states are more likely (Zhang et al., 2014; Kawamura et al., 2017; Zhang and Prange, 2020; Armstrong et al., 2021; Klockmann et al., 2018).

3 Contours of a baseline MIS3 experiment protocol

250 Although the choice of a time within MIS3 for a D-O baseline experiment should be unimportant, given that in reality D-O events occurred during the whole of the MIS3, our analysis of existing simulations, boundary conditions and mechanisms above suggests that there are periods which may be particularly conducive to D-O events occurring in models. Oscillatory D-O type behaviour appears to be more likely, but not guaranteed (Guo et al., 2019b; Zhang and Prange, 2020), when models are

run with intermediate or low MIS3 CO₂ values and ice-sheets, *i.e.* reduced in size compared to the LGM (Brown and Galbraith, 2016; Kawamura et al., 2017; Klockmann et al., 2018; Zhang and Prange, 2020; Galbraith and de Lavergne, 2019; Zhang et al., 255 2014; Vettoretti et al., 2022), and particularly without a high LIS.

The impact of orbital parameters has been investigated in less detail than the role of GHGs and ice sheets. Using the model COSMOS, Zhang et al. (2021) demonstrated that under intermediate glacial conditions, obliquity appears to play a significant role in the occurrence of D-O type behaviour. In particular, the orbital parameters at 40 ka do not produce D-O type behaviour, whilst at 34 ka lower obliquity (22.6°) leads to D-O type behaviour (see Figure 2 from Zhang et al. (2021)). Additionally, 260 the MIROC4m model produces D-O-type oscillations (under mid-glacial conditions) and low obliquity (22.9°) (Kuniyoshi et al., 2022). From these COSMOS and MIROC4m results, we deduce that low obliquity seems conducive to D-O behaviour in models.

These considerations suggest that the interval starting at 38 ka to 32 ka is a good choice for the proposed baseline experiment: it is characterised by (1) a rather regular sequence of D-O events (Fig. 1), and (2) has the ideal intermediate MIS3 ice-sheet 265 configuration conducive to generating D-O-type quasi-oscillations (Section 2).

A baseline simulation needs to be run for a sufficient duration to allow the strong positive feedbacks, together with long time-scale negative feedbacks, that enable D-O type oscillations. The analysis of existing simulations (Section 2) suggests this should be a minimum of 5000 years (Peltier and Vettoretti, 2014; Kleppin et al., 2015; Sidorenko et al., 2015; Brown and Galbraith, 2016; Klockmann et al., 2018, 2020). However, given computational constraints, a minimum duration of around 270 2000 years, with a spin-up period of 1000 years, may be a more practical minimum requirement for most modeling groups. It would, however, be important to examine and document key metrics for model drift (such as top-of-atmosphere radiation imbalance, deep ocean or global mean ocean temperature) during the initial spin-up. The exact length of spin-up is thus subject to discretion of each modelling group based on these key metrics.

There are two obvious possibilities for spinning up the MIS3 control experiment (MIS3-cnt). The baseline experiment could 275 be initialised from either the end of a well spun-up LGM or PI experiment. Other possibilities could be to spin up from a linear combination of LGM and PI states (as done in Klockmann et al., 2016, 2018) or spinning up from present day's observations (as done in Guo et al., 2019b). Modelling groups are encouraged to choose whichever option is more feasible/convenient for them. In case that several spin-up options are available, short spin-ups with diagnosed top-of-atmosphere (TOA) imbalance or global mean ocean temperature could help distinguish the faster spin-up option. It is worth noting that initial ocean state (*i.e.* 280 Atlantic salinity stratification) does play a role in abrupt AMOC change and associated feedbacks (Zhang et al., 2013; Knorr et al., 2021), of which impacts shall be considered and evaluated in the future.

We suggest performing a MIS3-cnt experiment centered at 34 ka, using GHG and orbital conditions for 34 ka (Fig. 6); and ice sheet configuration as outlined below (sections 3.1 and 3.2).

We acknowledge that some models might not oscillate under the proposed 34 ka baseline scenario. Indeed, this is expected 285 for NorESM, which under 38 ka conditions, is in a stable regime and the model state seems to be far from a possible tipping point. In spite of that, standardised MIS3 simulations which do not show D-O like behaviour are still highly valuable, for exactly the same reasons that LGM simulations are relevant to the wider modelling community. These standardised MIS3




Figure 6. Monthly zonal-mean MIS3 (34ka) - PI anomalies of the top-of-atmosphere short-wave incoming radiation (W m^{-2}).

simulations could contribute to progress on the overarching CMIP6 questions 1 and 2 (Eyring et al., 2016): “How does the Earth System respond to forcing?”, and “What are the origins and consequences of systematic model biases?” With a larger
 290 number of standardised MIS3 simulations, we would be able to answer questions such as:

- Are state-of-the-art climate models capable of representing D-O events under more realistic MIS3 conditions? Benchmarking these simulations will deliver a measure of how well models simulate abrupt changes, and tipping events.
- Standardized MIS3 simulations can help explore the existence of a theoretical sweet spot for millennial activity in current climate models (Barker and Knorr, 2021). As close to or within the sweet spot, the AMOC is characterized
 295 by high sensitivity to transient and/or noisy climatic forcing (Zhang et al., 2014; Lohmann and Ditlevsen, 2018) or by self-oscillating behaviors (Zhang et al., 2021).
- If models are too stable to simulate abrupt transitions, what are the processes that contribute to relative levels of model stability?
- In addition, a larger number of standardised MIS3 simulations could encourage the creation of new data sets, improving
 300 model-data evaluation.

3.1 Atmospheric trace gases

MIS3 atmospheric CO₂ values varied between a maximum of ~ 233 ppm to a minimum of ~ 187.5 ppm (Table 2; Figure 1; Bauska et al., 2021). Interestingly, increases of around 5 ppm happened during the abrupt warming of most D-O events and increases of up to 10 ppm happened within some Heinrich stadials (Bauska et al., 2021). GHG forcing is critical to model stability. Low (LGM-like) to intermediate (MIS3) CO₂ concentrations tend to be associated with abrupt D-O type AMOC transitions in models (Section 2 and Klockmann et al., 2018; Zhang et al., 2017, 2014; Brown and Galbraith, 2016; Vettoretti et al., 2022). We thus suggest to perform the MIS3-cnt experiment using the GHGs values specified in Table 2 and keep these values fixed for the whole duration of the simulation including the spin-up.

3.2 Northern Hemisphere Ice Sheets

Constraining MIS3 ice-sheet boundary conditions is a challenge. Scarcity and fragmentation of evidence (Kleman et al., 2010; Batchelor et al., 2019) is an issue. In particular, it is difficult to determine the size and shape of the ice sheets during MIS3 because subsequent larger LGM configurations have overridden and destroyed evidence of the position of the margins of these smaller ice sheets.

Global sea level fluctuations during the mid-MIS3 were driven nearly exclusively by the LIS (Gowan et al., 2021). Global average sea level remained above -55 m for the period between 30-55 ka. From glacial isostatic modelling and geological constraints, a global mean sea level between -30 m and -50 m is inferred (Dalton et al., 2022). For much of MIS3, since the Eurasian ice sheets and the Cordilleran Ice Sheet were likely restricted to mountain-based caps (Helmens, 2014; Hughes et al., 2016; Clague and Ward, 2011), the primary control on ice volume is assumed to be from the LIS. Recent work in the area of the Hudson Bay (Dalton et al., 2016, 2019; McMartin et al., 2019; Dalton et al., 2022) suggests ice-free conditions may have occurred during mid-MIS3. This implies climatic conditions in this region similar to present (Dalton et al., 2017), and a LIS margin removed from the southern Hudson Bay. Similarly Tarasov et al. (2012) show a considerably lower and less extensive LIS compared to ICE-5G and ICE-6G (Peltier, 2004; Peltier et al., 2015) LGM ice sheet reconstructions. Pico et al. (2017) sea-level curves are consistent with the estimated MIS3 ice-sheet volumes from Batchelor et al. (2019). Using Glacio Isostatic Adjustment modeling, Pico et al. (2017) also show that a small LIS can explain high MIS3 sea-level estimates alongside the eastern coast of the United States. Batchelor et al. (2019)'s synthesis of numerical modelling results and empirical data provides additional support for a considerable reduction in the MIS3 LIS extent and very minimal European ice sheet.

The recent MIS3 ice sheet reconstruction, PaleoMIST 1.0 (Paleo Margins, Ice Sheets, and Topography), was developed independently of far-field sea-level records and indirect proxy records by Gowan et al. (2021). This reconstruction is based on trying to fit the evolution of ice flow indicators, as well as chronological constraints of ice-free conditions.

Gowan et al. (2021) provide a maximum and minimal MIS3 reconstruction, specifically for the Laurentide Ice Sheet. The maximum scenario is more consistent with recently discovered eastward oriented, pre-LGM ice flow direction indicators found in southeastern Manitoba (Gauthier and Hodder, 2020), so we currently consider it to be more likely. However, at 35 ka, the difference between the two scenarios is minor. The difference is primarily with the thickness (and therefore also topography)

of the ice sheet, rather than extent, but it amounts to less than 1 m of sea level equivalent. The 35 ka time slice represents conditions after Heinrich Event 4 (Andrews and Voelker, 2018), and the ice margin in Hudson Strait is retreated about 350 km from the edge of the continental shelf. The ice margin elsewhere for the Laurentide Ice Sheet is based on chronological constraints, most that are documented in the compilation by Dalton et al. (2019). The Cordillera Ice Sheet extent is based on evidence of relatively restricted ice cover during MIS 3 (Clague and Ward, 2011). The Greenland Ice Sheet margin is set to be intermediate of the LGM and present day extent. The Eurasian ice extent at 35 ka includes an advance of ice into the Baltic Sea, which happened after Heinrich Event 4 (Hughes et al., 2016). For East Antarctica, the margin is set to be the same as present. In West Antarctica, the margin at 35 ka is close to the shelf edge, as the maximum extent may have been achieved by 30 ka (Larter et al., 2014).

Given its strong evidence basis, we thus suggest the use of the maximum 35 ka Gowan et al. (2021) PaleoMIST ice sheet configuration. We note the LIS is considerably reduced in size, compared to the ICE-6G LGM reconstruction in the southeastern margin (Fig. 7a,d); the EIS is also significantly smaller (Fig. 7a,d).

Whilst the implementation of the ice sheet will differ between models, the steps of Kageyama et al. (2017) describe how to implement a glacial state ice sheet in the IPSL climate model. For consistency, we likewise recommend the same steps should be followed as far as possible. Since a reduced sea-level can modify river courses, Kageyama et al. (2017) recommend that as a minimum, rivers should reach the oceans. Also, the ocean should be initialized with a salinity 0.6 psu higher than the PI experiment, to account for the sea-level difference between MIS3 and PI experiment (freshwater stored as ice on land) (Guo et al., 2019b).

The single ice sheet reconstruction MIS3 set-up summarized above contrasts with the PMIP4 LGM protocol, which provides three different possible ice sheet configurations (PMIP3, ICE6G-C and GLAC-1D) for the tier 1 LGM experiment (Kageyama et al., 2017). This partially reflects the more limited knowledge of ice sheet in pre-LGM periods. Exploration of the effect of MIS3 ice sheet reconstructions uncertainties on climate models, particularly on model stability, would be valuable. For this purpose, further additional 34 ky / MIS3 ice sheet reconstructions would be very valuable.

3.3 Heinrich-Event preconditioned option

The term “kick” Heinrich event-like was initially introduced by Peltier and Vettoretti (2014) to invoke a ‘kicked’ salt oscillator hypothesis (pseudo-Heinrich type behaviour), to induce D-O type oscillations in an LGM simulation performed with the UofT CCSM4. During the first thousand years of the simulation as the model is spun up and the ocean cools to reach a state consistent with glacial boundary conditions, there are two thermal thresholds during which the strength of the AMOC rapidly reduces (see Figure 2 in Peltier et al., 2020). These abrupt transitions in the AMOC coincide with abrupt reductions in surface temperatures in the North Atlantic and abrupt expansions of sea ice coverage. During the second of these events, the AMOC is reduced to approximately 12 Sv, about half its strength in the pre-industrial control (Peltier et al., 2020). This event may resemble the impact of a Heinrich event-like “kick” to the AMOC though no freshwater perturbation was imposed (Peltier et al., 2020).

In a more recent study, Pedro et al. (2022) examine the CCSM4 simulations that shows unforced D-O type oscillations (Peltier and Vettoretti, 2014; Vettoretti and Peltier, 2018), but with the addition of a (freshwater) H-like event. (Pedro et al.,




Figure 7. MIS3 ice sheet reconstruction from (a,b) Gowan et al. (2021). Also shown for comparison is the (c,d) LGM ICE-6G ice sheet reconstruction from Peltier et al. (2015). Third row shows the differences between the MIS3 ice sheet reconstruction and the LGM ICE-6G reconstruction.

2022). The freshwater flux (0.05 Sv injected into the NA for 500 years) leads to (1) a 5 Sv weaker AMOC compared to the one seen in the unforced model stadial and, (2) a stronger D-O warming transition into the interstadial phase (Pedro et al., 2022).
370 Thus this type of H-E preconditioning can trigger abrupt reductions in the AMOC strength and in NA surface temperatures and sea ice coverage – and it may also help induce a stadial state in other models which is more conducive to unforced (D-O type) oscillations (Pedro et al., 2022).

Given the importance of HEs to starting Bond Cycles of D-O events, an additional experiment to investigate how HE meltwater preconditioning impacts the simulation of D-O like oscillations under MIS3 boundary conditions would be valuable.
375 HE freshwater preconditioning may, as in reality, be more conducive to a (Bon Cycle-like) sequence of spontaneous D-O type oscillations (see Table 1).

The freshwater delivered during Heinrich event iceberg discharge suppresses the AMOC, leading to accumulation of heat in the Southern Hemisphere, and in the North Atlantic subsurface waters (*e.g.* Stocker and Johnsen, 2003). Estimates of the meltwater input into the North Atlantic during Heinrich events range between 2 m and 15 m of sea level equivalent ice volume
380 (Hemming, 2004; Chappell, 2002; Rohling et al., 2004; Roche et al., 2004; Roberts et al., 2014b; Siddall et al., 2008; Grant et al., 2014). It is logical to presume that these freshwater events are important in preconditioning the climate system with respect to D-O behaviour (Peltier and Vettoretti, 2014; Peltier et al., 2020; Pedro et al., 2022). Freshwater perturbations can trigger changes between AMOC states (*e.g.* Ganopolski and Rahmstorf, 2001; Timmermann et al., 2003; Stouffer et al., 2006; Hu et al., 2008; Kageyama et al., 2012; Roberts et al., 2014a; Sime et al., 2019) and a relatively small freshwater flux applied
385 over convection areas can lead to a shutdown of the AMOC (*e.g.* Roche et al., 2010). Studies have shown similarities between observed global features of abrupt D-O changes and the behaviour seen in freshwater forcing experiments (Liu et al., 2009; Menviel et al., 2014). However, the sensitivity of the AMOC to a wide range of freshwater inputs varies according to model, where the meltwater is added, and the background climate state (Ganopolski and Rahmstorf, 2001; Timmermann et al., 2003; Stouffer et al., 2006; Hu et al., 2008; Kageyama et al., 2012; Roberts et al., 2014a; Zhang et al., 2014). Given these various
390 uncertainties, we suggest that it would be useful to run an additional experiment to investigate how preconditioning through a H-like (freshwater) event impacts the simulation of D-O like oscillations under MIS3 boundary conditions.

The range of H-event volumes calculated using ice sheet models varies from 24.2 to 125 x 10⁴km³ (MacAyeal, 1993; Dowdeswell et al., 1995; Marshall and Clarke, 1997; Hulbe, 1997), whilst isotope based estimates and a precipitation balance approach have yielded 86, 649, and 946 x 10⁴km³ of ice volume (Hemming, 2004; Roche et al., 2004; Levine and Bigg, 2008).
395 Roberts et al. (2014b), and using a sediment modelling approach estimated a discharge of 30 to 120 x 10⁴km³ of ice volume. Some of the spread in these estimates could be because the relationship between the oxygen isotope record, sea level, and meltwater volume is not constant when ice is lost from marine basins, such that the use of oxygen isotopes for calculating H-event volumes may produce unrealistically high values (Gasson et al., 2016; Hemming, 2004; Roberts et al., 2014b). There is also some uncertainty about the duration of the H-events, with some previous studies suggesting they could be as short as 250
400 years (Hemming, 2004) and others suggesting a duration of 500 yr is more typical (Roberts et al., 2014b). These considerations suggest that it is possible to justify the use of anywhere between 0.02 - 0.6 Sv freshwater flux over 500 years; or 0.04 - 1.2 Sv over 250 years. More recent estimates of H-event magnitudes tend to favour the lower end of this range. If all forcings are set

to MIS3-cnt values and the H-event freshwater flux is distributed across the North Atlantic this could yield a range of stadial climates (Sime et al., 2019; Zhang and Prange, 2020). After 250-500 years this freshwater forcing should be switched off.

Table 2. Summary of the boundary conditions (BC) and forcings for the MIS3-cnt experiment.

BC/Forcing	Suggested value MIS3-cnt
Atmospheric trace gases	CO_2 : 208 ppm (Bauska et al., 2021)
	CH_4 : 420 ppb (Loulergue et al., 2008)
	N_2O : 204 ppb (Schilt et al., 2010)
Insolation	Eccentricity: 0.01567 Berger et al. (1998)
	Obliquity: 22.6° Berger et al. (1998)
	Precession: -0.016 Berger et al. (1998)
Solar constant	Same as PI control
Ice sheets	35ka ice sheet reconstruction (Gowan et al., 2021); mean global salinity increased by 0.6PSU to account for ice volume
Global freshwater budget	Closed to avoid drifts; Snow should not accumulate over ice sheets and rivers should flow into the ocean.
	Models need to consider lakes when closing the global freshwater budget
Vegetation	Dynamic or fixed as in PI.
	If fixed vegetation: tundra in new land points
Dust	As in PI control
H-kicked variant	initial 0.04 - 1 Sv over 250-500 years
	followed by standard MIS3-cnt simulation

405 4 Conclusions

D-O events are abrupt, large climate changes that punctuated the last glacial period. There is uncertainty whether current IPCC-class models can effectively represent the processes that cause D-O events. We have shown that reduced ice sheets relative to LGM, low obliquity values and low-to-medium MIS3 CO_2 values are more likely to lead to unforced quasi-oscillatory D-O type behaviour. However, the simulations need to be run long enough to allow the strong positive AMOC feedbacks, along
410 with negative feedbacks on long time-scales, which can then lead to D-O type oscillations. Around 40% of the simulations set-up with full LGM or more MIS3-like conditions, have a run length of less than 2000 model years, which makes it difficult to tell whether any of these simulations are capable of, or likely to, exhibit D-O like behaviour. In addition, the vast majority of PMIP4/CMIP6 models have not run LGM or MIS3-like simulations long enough to be sure which models have the capability to oscillate.

415 We have provided boundary conditions for a baseline MIS3-cnt simulation, and a H-event preconditioned variant (freshwater
forced experiment). The MIS3-cnt experiment is centered at 34 ka because it yields the ideal combination of intermediate ice
sheets (smaller in size compared to LGM), low obliquity values and medium-to-low GHG values conducive to oscillatory D-O
type behaviour in models. Ideally, the MIS3 baseline experiment should be run for 5000 years, however, given computational
constraints a minimum duration of 2000 years together with a spin-up of at least 1000 years is a more practical minimum re-
420 quirement. This baseline MIS3-cnt protocol provides a common framework to explore cold-period instabilities using particular
GHG-, insolation-, freshwater-, and NH ice sheet-related forcings, together with diapycnal mixing. More model simulations
run under the here proposed MIS3 DO protocol together with analyses across models, could provide better insights, along the
lines of atmospheric-ice-ocean feedbacks behind DO events. These simulations will allow us to answer questions such as: are
425 current climate models able to reproduce DO-type behaviour under more realistic MIS3 conditions? How well models simu-
late tipping events, abrupt changes? What are the mechanisms that lead to relative levels of model stability? Moreover, a large
number of standardised MIS3 simulations could encourage the creation of new data sets, improving model-data evaluation.

Team list. Ayako Abe-Ouchi; Andreas Born; Nathaëlle Bouttes; Peter Ditlevsen; Michael P. Erb; Georg Feulner; Evan J. Gowan; Lauren Gre-
goire; Chungheng Guo; Sandy P. Harrison; Heather Andres; Masa Kageyama; Marlene Klockmann; Fabrice Lambert; Allegra N. LeGrande;
Ute Merkel; Larissa S. Nazarenko; Kerim H. Nisancioglu; Kevin Oliver; Bette Otto-Bliesner; William R. Peltier; Matthias Prange; Kira
430 Rehfeld; Alexander Robinson; Lev Tarasov, Paul. J Valdes; Guido Vettoretti; Nils Weitzel; Qiong Zhang; Xu Zhang

Author contributions. I.M. compiled all tables. L.C.S. and I.M. wrote the first draft of this manuscript. L.C.S and I.M. produced all figures.
All authors contributed to the final draft.

Competing interests. No competing interests are present

Acknowledgements. This project is TiPES contribution #123: This project has received funding from the European Union's Horizon 2020 re-
435 search and innovation programme under grant agreement No 820970. KR and NW acknowledge funding by the Deutsche Forschungsgemein-
schaft (DFG, German Research Foundation), project no. 395588486 and the PalMod project (www.palmod.de), subproject no. 01LP1926C.
CG acknowledges funding from the Research Council of Norway under grant no. 325333 (ABRUPT).

References

- Abe-Ouchi, A., Saito, F., Kageyama, M., Braconnot, P., Harrison, S. P., Lambeck, K., Otto-Bliesner, B. L., Peltier, W., Tarasov, L., Peter-
440 schmitt, J.-Y., et al.: Ice-sheet configuration in the CMIP5/PMIP3 Last Glacial Maximum experiments, *Geoscientific Model Development*,
8, 3621–3637, 2015.
- Adolphi, F., Ramsey, C. B., Erhardt, T., Edwards, R. L., Cheng, H., Turney, C. S., Cooper, A., Svensson, A., Rasmussen, S. O., Fischer, H.,
et al.: Connecting the Greenland ice-core and U/ Th timescales via cosmogenic radionuclides: testing the synchronicity of Dansgaard–
Oeschger events, *Climate of the Past*, 14, 2018.
- 445 Alley, R., Clark, P., Keigwin, L., Webb, R., et al.: Making sense of millennial-scale climate change, *Geophysical Monograph-American*
Geophysical Union, 112, 385–394, 1999.
- Alley, R. B.: Icing the north Atlantic, *Nature*, 392, 335–337, 1998.
- Andersen, K. K., Azuma, N., Barnola, J.-M., Bigler, M., Biscaye, P., Caillon, N., Chappellaz, J., Clausen, H. B., Dahl-Jensen, D., Fischer,
H., et al.: High-resolution record of the Northern Hemisphere climate extending into the last interglacial period, *Nature*, 431, 147–151,
450 2004.
- Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M., Röthlisberger, R., Ruth, U., Siggaard-Andersen, M.-L., Stef-
fensen, J. P., Dahl-Jensen, D., et al.: The Greenland ice core chronology 2005, 15–42 ka. Part 1: constructing the time scale, *Quaternary*
Science Reviews, 25, 3246–3257, 2006.
- Andrews, J. T. and Voelker, A. H.: “Heinrich events”(& sediments): A history of terminology and recommendations for future usage, *Qua-*
455 *ternary Science Reviews*, 187, 31–40, 2018.
- Armstrong, E., Izumi, K., and Valdes, P.: Identifying the Mechanisms of DO-scale Oscillations in a Gcm: a Salt Oscillator Triggered by the
Laurentide Ice Sheet, 2021.
- Årthun, M., Eldevik, T., Smedsrud, L., Skagseth, Ø., and Ingvaldsen, R.: Quantifying the influence of Atlantic heat on Barents Sea ice
variability and retreat, *Journal of Climate*, 25, 4736–4743, 2012.
- 460 Asmerom, Y., Polyak, V. J., and Burns, S. J.: Variable winter moisture in the southwestern United States linked to rapid glacial climate shifts,
Nature Geoscience, 3, 114–117, 2010.
- Barker, S. and Knorr, G.: Millennial scale feedbacks determine the shape and rapidity of glacial termination, *Nature Communications*, 12,
1–12, 2021.
- Barker, S., Diz, P., Vautravers, M. J., Pike, J., Knorr, G., Hall, I. R., and Broecker, W. S.: Interhemispheric Atlantic seesaw response during
465 the last deglaciation, *Nature*, 457, 1097–1102, 2009.
- Barker, S., Knorr, G., Edwards, R. L., Parrenin, F., Putnam, A. E., Skinner, L. C., Wolff, E., and Ziegler, M.: 800,000 years of abrupt climate
variability, *science*, 334, 347–351, 2011.
- Barton, B. I., Lenn, Y.-D., and Lique, C.: Observed Atlantification of the Barents Sea causes the Polar Front to limit the expansion of winter
sea ice, *Journal of Physical Oceanography*, 48, 1849–1866, 2018.
- 470 Batchelor, C. L., Margold, M., Krapp, M., Murton, D. K., Dalton, A. S., Gibbard, P. L., Stokes, C. R., Murton, J. B., and Manica, A.: The
configuration of Northern Hemisphere ice sheets through the Quaternary, *Nature communications*, 10, 1–10, 2019.
- Baumgartner, M., Kindler, P., Eicher, O., Floch, G., Schilt, A., Schwander, J., Spahni, R., Capron, E., Chappellaz, J., Leuenberger, M., et al.:
NGRIP CH 4 concentration from 120 to 10 kyr before present and its relation to a $\delta^{15}\text{N}$ temperature reconstruction from the same ice
core, *Climate of the Past*, 10, 903–920, 2014.

- 475 Bauska, T. K., Marcott, S. A., and Brook, E. J.: Abrupt changes in the global carbon cycle during the last glacial period, *Nature Geoscience*, 14, 91–96, 2021.
- Berger, A., Loutre, M.-F., and Gallée, H.: Sensitivity of the LLN climate model to the astronomical and CO₂ forcings over the last 200 ky, *Climate Dynamics*, 14, 615–629, 1998.
- Bond, G., Heinrich, H., Broecker, W., Labeyrie, L., McManus, J., Andrews, J., Huon, S., Jantschik, R., Clasen, S., Simet, C., et al.: Evidence
480 for massive discharges of icebergs into the North Atlantic ocean during the last glacial period, *Nature*, 360, 245–249, 1992.
- Broecker, W. and Peteet, D.: Does the ocean-atmosphere system have more than one stable mode of operation?, *Nature*, 315, 21–26, <https://doi.org/10.1038/315021a0>, 1985.
- Broecker, W. S.: Massive iceberg discharges as triggers for global climate change, *Nature*, 372, 421–424, 1994.
- Brovkin, V., Brook, E., Williams, J. W., Bathiany, S., Lenton, T. M., Barton, M., DeConto, R. M., Donges, J. F., Ganopolski, A., McManus,
485 J., et al.: Past abrupt changes, tipping points and cascading impacts in the Earth system, *Nature Geoscience*, 14, 550–558, 2021.
- Brown, N. and Galbraith, E. D.: Hosed vs. unhosed: interruptions of the Atlantic Meridional Overturning Circulation in a global coupled model, with and without freshwater forcing., *Climate of the Past*, 12, 2016.
- Brugger, J., Hofmann, M., Petri, S., and Feulner, G.: On the sensitivity of the Devonian climate to continental configuration, vegetation cover, orbital configuration, CO₂ concentration, and insolation, *Paleoceanography and Paleoclimatology*, 34, 1375–1398, 2019.
- 490 Buckley, M. W. and Marshall, J.: Observations, inferences, and mechanisms of the Atlantic Meridional Overturning Circulation: A review, *Reviews of Geophysics*, 54, 5–63, 2016.
- Chappell, J.: Sea level changes forced ice breakouts in the Last Glacial cycle: new results from coral terraces, *Quaternary Science Reviews*, 21, 1229–1240, 2002.
- Clague, J. J. and Ward, B.: Pleistocene Glaciation of British Columbia, in: *Quaternary Glaciations - Extent and Chronology A Closer Look*, edited by Ehlers, J., Gibbard, P. L., and Hughes, P. D., vol. 15 of *Developments in Quaternary Sciences*, chap. 44, pp. 563 – 573, Elsevier, <https://doi.org/10.1016/B978-0-444-53447-7.00044-1>, 2011.
- Clark, P. U., Pisias, N. G., Stocker, T. F., and Weaver, A. J.: The role of the thermohaline circulation in abrupt climate change, *Nature*, 415, 863–869, 2002.
- Clark, P. U., Hostetler, S. W., Pisias, N. G., Schmittner, A., and Meissner, K. J.: Mechanisms for an 7-kyr climate and sea-level oscillation
500 during marine isotope stage 3, *Geophysical Monograph-American Geophysical Union*, 173, 209, 2007.
- Cook, K. H. and Held, I. M.: Stationary waves of the ice age climate, *Journal of climate*, 1, 807–819, 1988.
- Dalton, A. S., Finkelstein, S. A., Barnett, P. J., and Forman, S. L.: Constraining the Late Pleistocene history of the Laurentide Ice Sheet by dating the Missinaibi Formation, Hudson Bay Lowlands, Canada, *Quaternary Science Reviews*, 146, 288–299, 2016.
- Dalton, A. S., Väiliranta, M., Barnett, P. J., and Finkelstein, S. A.: Pollen and microfossil-inferred palaeoclimate at the Ridge Site, Hudson
505 Bay Lowlands, Canada: evidence for a dry climate and significant recession of the Laurentide Ice Sheet during Marine Isotope Stage 3, *Boreas*, 46, 388–401, 2017.
- Dalton, A. S., Finkelstein, S. A., Forman, S. L., Barnett, P. J., Pico, T., and Mitrovica, J. X.: Was the Laurentide ice sheet significantly reduced during marine isotope stage 3?, *Geology*, 47, 111–114, 2019.
- Dalton, A. S., Pico, T., Gowan, E. J., Clague, J. J., Forman, S. L., McMartin, I., Sarala, P., and Helmens, K. F.: The marine $\delta^{18}\text{O}$ record
510 overestimates continental ice volume during Marine Isotope Stage 3, *Global and Planetary Change*, p. 103814, 2022.
- Dansgaard, W., Clausen, H. B., Gundestrup, N., Hammer, C., Johnsen, S., Kristinsdottir, P., and Reeh, N.: A new Greenland deep ice core, *Science*, 218, 1273–1277, 1982.

- de Lavergne, C., Falahat, S., Madec, G., Roquet, F., Nycander, J., and Vic, C.: Toward global maps of internal tide energy sinks, *Ocean Modelling*, 137, 52–75, 2019.
- 515 Deplazes, G., Lückge, A., Peterson, L. C., Timmermann, A., Hamann, Y., Hughen, K. A., Röhl, U., Laj, C., Cane, M. A., Sigman, D. M., et al.: Links between tropical rainfall and North Atlantic climate during the last glacial period, *Nature Geoscience*, 6, 213–217, 2013.
- Ditlevsen, P. D. and Johnsen, S. J.: Tipping points: early warning and wishful thinking, *Geophysical Research Letters*, 37, 2010.
- Ditlevsen, P. D., Kristensen, M. S., and Andersen, K. K.: The recurrence time of Dansgaard-Oeschger events and limits on the possible periodic component, *J. Climate*, 18, 2594–2603, 2005.
- 520 Ditlevsen, P. D., Andersen, K. K., and Svensson, A.: The DO-climate events are probably noise induced: statistical investigation of the claimed 1470 years cycle, *Climate of the Past*, 3, 129–134, 2007.
- Dokken, T. M., Nisancioglu, K. H., Li, C., Battisti, D. S., and Kissel, C.: Dansgaard-Oeschger cycles: Interactions between ocean and sea ice intrinsic to the Nordic seas, *Paleoceanography*, 28, 491–502, 2013.
- Dowdeswell, J., Maslin, M., Andrews, J., and McCave, I.: Iceberg production, debris rafting, and the extent and thickness of Heinrich layers (H-1, H-2) in North Atlantic sediments, *Geology*, 23, 301–304, 1995.
- 525 Drijfhout, S., Gleeson, E., Dijkstra, H. A., and Livina, V.: Spontaneous abrupt climate change due to an atmospheric blocking–sea-ice–ocean feedback in an unforced climate model simulation, *Proceedings of the National Academy of Sciences*, 110, 19713–19718, 2013.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, 2016.
- 530 Feulner, G., Rahmstorf, S., Levermann, A., and Volkwardt, S.: On the Origin of the Surface Air Temperature Difference between the Hemispheres in Earth’s Present-Day Climate, *Journal of Climate*, 26, 7136–7150, <https://doi.org/10.1175/JCLI-D-12-00636.1>, 2013.
- Galaasen, E. V., Ninnemann, U. S., Irvani, N., Kleiven, H. K. F., Rosenthal, Y., Kissel, C., and Hodell, D. A.: Rapid reductions in North Atlantic Deep Water during the peak of the last interglacial period, *Science*, 343, 1129–1132, 2014.
- Galaasen, E. V., Ninnemann, U. S., Kessler, A., Irvani, N., Rosenthal, Y., Tjiputra, J., Bouttes, N., Roche, D. M., Kleiven, H. K. F., and 535 Hodell, D. A.: Interglacial instability of North Atlantic deep water ventilation, *Science*, 367, 1485–1489, 2020.
- Galbraith, E. and de Lavergne, C.: Response of a comprehensive climate model to a broad range of external forcings: Relevance for deep ocean ventilation and the development of late Cenozoic ice ages, *Climate Dynamics*, 52, 653–679, 2019.
- Ganopolski, A. and Rahmstorf, S.: Rapid changes of glacial climate simulated in a coupled climate model, *Nature*, 409, 153–158, 2001.
- Gasson, E., DeConto, R. M., and Pollard, D.: Modeling the oxygen isotope composition of the Antarctic ice sheet and its significance to 540 Pliocene sea level, *Geology*, 44, 827–830, 2016.
- Gauthier, M. S. and Hodder, T. J.: Surficial geology mapping from Manigotagan to Berens River, southeastern Manitoba (parts of NTS 62P1, 7, 8, 10, 15, 63A2, 7), Report of Activities 2020, Manitoba Agriculture and Resource Development, Manitoba Geological Survey, pp. 41–46, 2020.
- Genty, D., Blamart, D., Ouahdi, R., Gilmour, M., Baker, A., Jouzel, J., and Van-Exter, S.: Precise dating of Dansgaard–Oeschger climate 545 oscillations in western Europe from stalagmite data, *Nature*, 421, 833–837, 2003.
- Goni, M. F. S. and Harrison, S. P.: Millennial-scale climate variability and vegetation changes during the Last Glacial: Concepts and terminology, *Quaternary Science Reviews*, 29, 2823–2827, 2010.
- Gowan, E. J., Zhang, X., Khosravi, S., Rovere, A., Stocchi, P., Hughes, A. L., Gyllencreutz, R., Mangerud, J., Svendsen, J.-I., and Lohmann, G.: A new global ice sheet reconstruction for the past 80 000 years, *Nature Communications*, 12, 1–9, 2021.

- 550 Grant, K., Rohling, E., Ramsey, C. B., Cheng, H., Edwards, R., Florindo, F., Heslop, D., Marra, F., Roberts, A., Tamisiea, M. E., et al.:
Sea-level variability over five glacial cycles, *Nature communications*, 5, 1–9, 2014.
- Guo, C., Bentsen, M., Bethke, I., Ilicak, M., Tjiputra, J., Toniazzo, T., Schwinger, J., and Otterå, O. H.: Description and evaluation of
NorESM1-F: A fast version of the Norwegian Earth System Model (NorESM), *Geoscientific Model Development*, 12, 343–362, 2019a.
- Guo, C., Nisancioglu, K. H., Bentsen, M., Bethke, I., and Zhang, Z.: Equilibrium simulations of Marine Isotope Stage 3 climate, *Climate of*
555 *the Past*, 15, 1133–1151, <https://doi.org/10.5194/cp-15-1133-2019>, <https://cp.copernicus.org/articles/15/1133/2019/>, 2019b.
- Heinrich, H.: Origin and consequences of cyclic ice rafting in the northeast Atlantic Ocean during the past 130,000 years, *Quaternary*
research, 29, 142–152, 1988.
- Helmens, K. F.: The Last Interglacial–Glacial cycle (MIS 5–2) re-examined based on long proxy records from central and northern Europe,
Quaternary Science Reviews, 86, 115–143, 2014.
- 560 Hemming, S. R.: Heinrich events: Massive late Pleistocene detritus layers of the North Atlantic and their global climate imprint, *Reviews of*
Geophysics, 42, 2004.
- Hofer, D., Raible, C., Dehnert, A., and Kuhlemann, J.: The impact of different glacial boundary conditions on atmospheric dynamics and
precipitation in the North Atlantic region, *Climate of the Past*, 8, 935–949, 2012.
- Holden, P. B., Edwards, N., Oliver, K., Lenton, T., and Wilkinson, R.: A probabilistic calibration of climate sensitivity and terrestrial carbon
565 change in GENIE-1, *Climate Dynamics*, 35, 785–806, 2010.
- Hu, A., Otto-Bliesner, B. L., Meehl, G. A., Han, W., Morrill, C., Brady, E. C., and Briegleb, B.: Response of thermohaline circulation to
freshwater forcing under present-day and LGM conditions, *Journal of Climate*, 21, 2239–2258, 2008.
- Huber, C., Leuenberger, M., Spahni, R., Flückiger, J., Schwander, J., Stocker, T. F., Johnsen, S., Landais, A., and Jouzel, J.: Isotope calibrated
Greenland temperature record over Marine Isotope Stage 3 and its relation to CH₄, *Earth and Planetary Science Letters*, 243, 504–519,
570 2006.
- Hughes, A. L., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J., and Svendsen, J. I.: The last Eurasian ice sheets—a chronological database and
time-slice reconstruction, *DATED-1, Boreas*, 45, 1–45, 2016.
- Huisman, S. E., Den Toom, M., Dijkstra, H. A., and Drijfhout, S.: An indicator of the multiple equilibria regime of the Atlantic meridional
overturning circulation, *Journal of Physical Oceanography*, 40, 551–567, 2010.
- 575 Hulbe, C. L.: An ice shelf mechanism for Heinrich layer production, *Paleoceanography*, 12, 711–717, 1997.
- IPCC: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Inter-*
governmental Panel on Climate Change. [Stocker, T.F. and Qin, D and Plattner, G and Tignor, M and Allen, S.K. and Boschung, J and
Nauels, A and Xia, Y and Bex, V and Midgley, P.M (eds.)], Tech. Rep. 5, Intergovernmental Panel on Climate Change, Cambridge, United
Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324>, 2013.
- 580 Jackson, L. and Vellinga, M.: Multidecadal to centennial variability of the AMOC: HadCM3 and a perturbed physics ensemble, *Journal of*
climate, 26, 2390–2407, 2013.
- Jacobel, A., McManus, J., Anderson, R., and Winckler, G.: Climate-related response of dust flux to the central equatorial Pacific over the
past 150 kyr, *Earth and Planetary Science Letters*, 457, 160–172, 2017.
- Johnsen, S., Clausen, H., Dansgaard, W., Fuhrer, K., Gundestrup, N., Hammer, C., Iversen, P., Jouzel, J., Stauffer, B., et al.: Irregular glacial
585 interstadials recorded in a new Greenland ice core, *Nature*, 359, 311–313, 1992.

- Johnsen, S. J., Dahl-Jensen, D., Gundestrup, N., Steffensen, J. P., Clausen, H. B., Miller, H., Masson-Delmotte, V., Sveinbjörnsdóttir, A. E., and White, J.: Oxygen isotope and palaeotemperature records from six Greenland ice-core stations: Camp Century, Dye-3, GRIP, GISP2, Renland and NorthGRIP, *Journal of Quaternary Science: Published for the Quaternary Research Association*, 16, 299–307, 2001.
- 590 Jungclaus, J., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and Von Storch, J.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *Journal of Advances in Modeling Earth Systems*, 5, 422–446, 2013.
- Justino, F. and Peltier, W. R.: The glacial North Atlantic Oscillation, *Geophysical Research Letters*, 32, 2005.
- Kageyama, M., Merkel, U., Otto-Bliesner, B., Prange, M., Abe-Ouchi, A., Lohmann, G., Roche, D., Singarayer, J., Swingedouw, D., and Zhang, X.: Climatic impacts of fresh water hosing under Last Glacial Maximum conditions: a multi-model study., *Climate of the Past Discussions*, 8, 2012.
- 595 Kageyama, M., Albani, S., Braconnot, P., Harrison, S. P., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Marti, O., Peltier, W. R., Peterschmitt, J. Y., et al.: The PMIP4 contribution to CMIP6-Part 4: Scientific objectives and experimental design of the PMIP4-CMIP6 Last Glacial Maximum experiments and PMIP4 sensitivity experiments, *Geoscientific Model Development*, 10, 4035–4055, 2017.
- Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclaus, J. H., Otto-Bliesner, B. L., Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., et al.: The PMIP4 contribution to CMIP6–Part 1: Overview and over-arching analysis plan, *Geoscientific Model Development*, 11, 1033–1057, 2018.
- 600 Kageyama, M., Harrison, S. P., Kapsch, M.-L., Lofverstrom, M., Lora, J. M., Mikolajewicz, U., Sherriff-Tadano, S., Vadsaria, T., Abe-Ouchi, A., Bouttes, N., et al.: The PMIP4 Last Glacial Maximum experiments: preliminary results and comparison with the PMIP3 simulations, *Climate of the Past*, 17, 1065–1089, 2021a.
- 605 Kageyama, M., Sime, L. C., Sicard, M., Guarino, M.-V., de Vernal, A., Stein, R., Schroeder, D., Malmierca-Vallet, I., Abe-Ouchi, A., Bitz, C., et al.: A multi-model CMIP6-PMIP4 study of Arctic sea ice at 127 ka: sea ice data compilation and model differences, *Climate of the Past*, 17, 37–62, 2021b.
- Kawamura, K., Abe-Ouchi, A., Motoyama, H., Ageta, Y., Aoki, S., Azuma, N., Fujii, Y., Fujita, K., Fujita, S., Fukui, K., et al.: State dependence of climatic instability over the past 720,000 years from Antarctic ice cores and climate modeling, *Science advances*, 3, e1600446, 2017.
- 610 Kindler, P., Guillevic, M., Baumgartner, M. F., Schwander, J., Landais, A., and Leuenberger, M.: Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Climate of the Past*, 10, 887–902, 2014.
- Kleman, J., Jansson, K., De Angelis, H., Stroeven, A. P., Hättestrand, C., Alm, G., and Glasser, N.: North American ice sheet build-up during the last glacial cycle, 115–21 kyr, *Quaternary Science Reviews*, 29, 2036–2051, 2010.
- 615 Kleppin, H., Jochum, M., Otto-Bliesner, B., Shields, C. A., and Yeager, S.: Stochastic atmospheric forcing as a cause of Greenland climate transitions, *Journal of Climate*, 28, 7741–7763, 2015.
- Klockmann, M., Mikolajewicz, U., and Marotzke, J.: The effect of greenhouse gas concentrations and ice sheets on the glacial AMOC in a coupled climate model, *Climate of the Past*, 12, 1829–1846, 2016.
- Klockmann, M., Mikolajewicz, U., and Marotzke, J.: Two AMOC states in response to decreasing greenhouse gas concentrations in the coupled climate model MPI-ESM, *Journal of Climate*, 31, 7969–7984, 2018.
- 620 Klockmann, M., Mikolajewicz, U., Kleppin, H., and Marotzke, J.: Coupling of the subpolar gyre and the overturning circulation during abrupt glacial climate transitions, *Geophysical Research Letters*, 47, e2020GL090361, 2020.

- Knorr, G., Barker, S., Zhang, X., Lohmann, G., Gong, X., Gierz, P., Stepanek, C., and Stap, L. B.: A salty deep ocean as a prerequisite for glacial termination, *Nature Geoscience*, 14, 930–936, 2021.
- 625 Kuniyoshi, Y., Abe-Ouchi, A., Sherriff-Tadano, S., Chan, W.-L., and Saito, F.: Effect of Climatic Precession on Dansgaard-Oeschger-Like Oscillations, *Geophysical Research Letters*, 49, e2021GL095695, 2022.
- Lambert, F., Bigler, M., Steffensen, J. P., Hutterli, M., and Fischer, H.: Centennial mineral dust variability in high-resolution ice core data from Dome C, Antarctica, *Climate of the Past*, 8, 609–623, 2012.
- Landais, A., Barnola, J., Masson-Delmotte, V., Jouzel, J., Chappellaz, J., Caillon, N., Huber, C., Leuenberger, M., and Johnsen, S.: A
630 continuous record of temperature evolution over a sequence of Dansgaard-Oeschger events during Marine Isotopic Stage 4 (76 to 62 kyr BP), *Geophysical Research Letters*, 31, 2004.
- Lang, C., Leuenberger, M., Schwander, J., and Johnsen, S.: 16° C rapid temperature variation in central Greenland 70,000 years ago, *Science*, 286, 934–937, 1999.
- Larter, R. D., Anderson, J. B., Graham, A. G., Gohl, K., Hillenbrand, C.-D., Jakobsson, M., Johnson, J. S., Kuhn, G., Nitsche, F. O., Smith,
635 J. A., et al.: Reconstruction of changes in the Amundsen Sea and Bellingshausen sea sector of the West Antarctic ice sheet since the last glacial maximum, *Quaternary Science Reviews*, 100, 55–86, 2014.
- Lenton, T. M., Held, H., Kriegler, E., Hall, J. W., Lucht, W., Rahmstorf, S., and Schellnhuber, H. J.: Tipping elements in the Earth’s climate system, *Proceedings of the national Academy of Sciences*, 105, 1786–1793, 2008.
- Levine, R. C. and Bigg, G. R.: Sensitivity of the glacial ocean to Heinrich events from different iceberg sources, as modeled by a coupled
640 atmosphere-iceberg-ocean model, *Paleoceanography*, 23, 2008.
- Lhardy, F., Bouttes, N., Roche, D. M., Crosta, X., Waelbroeck, C., and Paillard, D.: Impact of Southern Ocean surface conditions on deep ocean circulation during the LGM: a model analysis, *Climate of the Past*, 17, 1139–1159, 2021.
- Li, C. and Battisti, D. S.: Reduced Atlantic storminess during Last Glacial Maximum: Evidence from a coupled climate model, *Journal of Climate*, 21, 3561–3579, 2008.
- 645 Li, C. and Born, A.: Coupled atmosphere-ice-ocean dynamics in Dansgaard-Oeschger events, *Quaternary Science Reviews*, 203, 1–20, 2019.
- Li, C., Battisti, D., Schrag, D., and Tziperman, E.: Abrupt climate shifts in Greenland due to displacements of the sea ice edge, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2005BL023492>, 2005.
- Li, C., Battisti, D., and Bitz, C.: Can North Atlantic sea ice anomalies account for Dansgaard-Oeschger climate signals?, *J. Clim.*, 23, 5457–5475, <https://doi.org/10.1175/2010JCLI3409.1>, 2010.
- 650 Liu, Z., Otto-Bliesner, B., He, F., Brady, E., Tomas, R., Clark, P., A.E., C., Lynch-Stieglitz, J., Curry, W., Brook, E., Erickson, D., Jacob, R., Kutzbach, J., and Cheng, J.: Transient simulation of last deglaciation with a new mechanism for Bolling-Allerod warming, *Science*, 325, 310–314, <https://doi.org/10.1126/science.1171041>, 2009.
- Löfverström, M., Caballero, R., Nilsson, J., and Kleman, J.: Evolution of the large-scale atmospheric circulation in response to changing ice sheets over the last glacial cycle, *Climate of the Past*, 10, 1453–1471, 2014.
- 655 Lohmann, G., Butzin, M., Eissner, N., Shi, X., and Stepanek, C.: Abrupt climate and weather changes across time scales, *Paleoceanography and Paleoclimatology*, 35, e2019PA003782, 2020.
- Lohmann, J. and Ditlevsen, P.: Objective extraction and analysis of statistical features of Dansgaard-Oeschger events, *Climate of the Past*, 15, 1771–1792, 2019.
- Lohmann, J. and Ditlevsen, P. D.: Random and externally controlled occurrences of Dansgaard–Oeschger events, *Climate of the Past*, 14,
660 609–617, 2018.

- Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J.-M., Raynaud, D., Stocker, T. F., and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years, *Nature*, 453, 383–386, 2008.
- Lynch-Stieglitz, J.: The Atlantic meridional overturning circulation and abrupt climate change, *Annual review of marine science*, 9, 83–104, 2017.
- 665 MacAyeal, D.: A low-order model of the Heinrich event cycle, *Paleoceanography*, 8, 767–773, 1993.
- Madonna, E., Li, C., Grams, C. M., and Woollings, T.: The link between eddy-driven jet variability and weather regimes in the North Atlantic-European sector, *Quarterly Journal of the Royal Meteorological Society*, 143, 2960–2972, 2017.
- Manabe, S. and Broccoli, A.: The influence of continental ice sheets on the climate of an ice age, *Journal of Geophysical Research: Atmospheres*, 90, 2167–2190, 1985.
- 670 Marcott, S. A., Clark, P. U., Padman, L., Klinkhammer, G. P., Springer, S. R., Liu, Z., Otto-Bliesner, B. L., Carlson, A. E., Ungerer, A., Padman, J., et al.: Ice-shelf collapse from subsurface warming as a trigger for Heinrich events, *Proceedings of the National Academy of Sciences*, 108, 13415–13419, 2011.
- Marshall, S. J. and Clarke, G. K.: A continuum mixture model of ice stream thermomechanics in the Laurentide Ice Sheet 2. Application to the Hudson Strait Ice Stream, *Journal of Geophysical Research: Solid Earth*, 102, 20615–20637, 1997.
- 675 Martin, T., Park, W., and Latif, M.: Southern Ocean forcing of the North Atlantic at multi-centennial time scales in the Kiel Climate Model, *Deep Sea Research Part II: Topical Studies in Oceanography*, 114, 39–48, 2015.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., et al.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and its response to increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, 2019.
- 680 McMartin, I., Campbell, J. E., and Dredge, L. A.: Middle Wisconsinan marine shells near Repulse Bay, Nunavut, Canada: implications for Marine Isotope Stage 3 ice-free conditions and Laurentide Ice Sheet dynamics in north-west Hudson Bay, *Journal of Quaternary Science*, 34, 64–75, 2019.
- Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J. M., Noda, A., et al.: Global climate projections. Chapter 10, 2007.
- 685 Menviel, L., Timmermann, A., Friedrich, T., and England, M.: Hindcasting the continuum of Dansgaard-Oeschger variability: mechanisms, patterns and timing, *Climate of the Past*, 10, 63–77, <https://doi.org/10.5194/cp-10-63-2014>, 2014.
- Merz, N., Raible, C. C., and Woollings, T.: North Atlantic eddy-driven jet in interglacial and glacial winter climates, *Journal of Climate*, 28, 3977–3997, 2015.
- Mitsui, T. and Crucifix, M.: Influence of external forcings on abrupt millennial-scale climate changes: a statistical modelling study, *Climate Dynamics*, 48, 2729–2749, 2017.
- 690 Montoya, M., Born, A., and Levermann, A.: Reversed North Atlantic gyre dynamics in present and glacial climates, *Climate dynamics*, 36, 1107–1118, 2011.
- Muglia, J. and Schmittner, A.: Glacial Atlantic overturning increased by wind stress in climate models, *Geophysical Research Letters*, 42, 9862–9868, 2015.
- 695 NGRIP Project Members: High-resolution record of Northern Hemisphere climate extending into the last interglacial period, *Nature*, 431, 147–151, <https://doi.org/10.1038/nature02805>, 2004.
- Nilsson, J., Broström, G., and Walin, G.: The thermohaline circulation and vertical mixing: Does weaker density stratification give stronger overturning?, *J. Phys. Oceanogr.*, 33, 2781–2795, 2003.

- Ohgaito, R., Yamamoto, A., Hajima, T., O'ishi, R., Abe, M., Tatebe, H., Abe-Ouchi, A., and Kawamiya, M.: PMIP4 experiments using
700 MIROC-ES2L Earth system model, *Geoscientific Model Development*, 14, 1195–1217, 2021.
- Oliver, K. I. C. and Edwards, N. R.: Location of potential energy sources and the export of dense water from the Atlantic Ocean, *Geophysical Research Letters*, 35, <https://doi.org/https://doi.org/10.1029/2008GL035537>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008GL035537>, 2008.
- Pausata, F. S., Li, C., Wettstein, J., Nisancioglu, K. H., and Battisti, D. S.: Changes in atmospheric variability in a glacial climate and the
705 impacts on proxy data: a model intercomparison, *Climate of the Past*, 5, 489–502, 2009.
- Pausata, F. S., Li, C., Wettstein, J., Kageyama, M., and Nisancioglu, K. H.: The key role of topography in altering North Atlantic atmospheric circulation during the last glacial period, *Climate of the Past*, 7, 1089–1101, 2011.
- Pedro, J., Andersson, C., Vettoretti, G., Voelker, A., Waelbroeck, C., Dokken, T. M., Jensen, M. F., Rasmussen, S., Sessford, E., Jochum, M.,
710 et al.: Dansgaard-Oeschger and Heinrich event temperature anomalies in the North Atlantic set by sea ice, frontal position and thermocline structure, *Quaternary Science Reviews*, 289, 107 599, 2022.
- Peltier, W. R.: Global glacial isostasy and the surface of the ice-age Earth: the ICE-5G (VM2) model and GRACE, *Annu. Rev. Earth Planet. Sci.*, 32, 111–149, 2004.
- Peltier, W. R. and Vettoretti, G.: Dansgaard-Oeschger oscillations predicted in a comprehensive model of glacial climate: A “kicked” salt oscillator in the Atlantic, *Geophysical Research Letters*, 41, 7306–7313, 2014.
- 715 Peltier, W. R., Argus, D., and Drummond, R.: Space geodesy constrains ice age terminal deglaciation: The global ICE-6G_C (VM5a) model, *Journal of Geophysical Research: Solid Earth*, 120, 450–487, 2015.
- Peltier, W. R., Ma, Y., and Chandan, D.: The KPP trigger of rapid AMOC intensification in the nonlinear Dansgaard-Oeschger relaxation oscillation, *Journal of Geophysical Research: Oceans*, 125, e2019JC015 557, 2020.
- Pico, T., Creveling, J., and Mitrovica, J.: Sea-level records from the US mid-Atlantic constrain Laurentide Ice Sheet extent during Marine
720 Isotope Stage 3, *Nature communications*, 8, 1–6, 2017.
- Polyakov, I. V., Pnyushkov, A. V., Alkire, M. B., Ashik, I. M., Baumann, T. M., Carmack, E. C., Goszczko, I., Guthrie, J., Ivanov, V. V., Kanzow, T., et al.: Greater role for Atlantic inflows on sea-ice loss in the Eurasian Basin of the Arctic Ocean, *Science*, 356, 285–291, 2017.
- Rahmstorf, S.: Bifurcations of the Atlantic thermohaline circulation in response to changes in the hydrological cycle, *Nature*, 378, 145–149,
725 1995.
- Rahmstorf, S.: Ocean circulation and climate during the past 120,000 years, *Nature*, 419, 207–214, <https://doi.org/10.1038/nature01090>, 2002.
- Rasmussen, S. O., Bigler, M., Blockley, S. P., Blunier, T., Buchardt, S. L., Clausen, H. B., Cvijanovic, I., Dahl-Jensen, D., Johnsen, S. J., Fischer, H., et al.: A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized
730 Greenland ice-core records: refining and extending the INTIMATE event stratigraphy, *Quaternary Science Reviews*, 106, 14–28, 2014.
- Rasmussen, T. L. and Thomsen, E.: The role of the North Atlantic Drift in the millennial timescale glacial climate fluctuations, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 210, 101–116, 2004.
- Rind, D., Schmidt, G. A., Jonas, J., Miller, R., Nazarenko, L., Kelley, M., and Romanski, J.: Multicentury instability of the Atlantic meridional circulation in rapid warming simulations with GISS ModelE2, *Journal of Geophysical Research: Atmospheres*, 123, 6331–6355, 2018.
- 735 Riviere, G., Laine, A., Lapeyre, G., Salas-Méllia, D., and Kageyama, M.: Links between Rossby wave breaking and the North Atlantic Oscillation–Arctic Oscillation in present-day and Last Glacial Maximum climate simulations, *Journal of Climate*, 23, 2987–3008, 2010.

- Roberts, W. H., Valdes, P. J., and Payne, A. J.: Topography's crucial role in Heinrich Events, *Proceedings of the National Academy of Sciences*, 111, 16688–16693, 2014a.
- Roberts, W. H., Valdes, P. J., and Payne, A. J.: A new constraint on the size of Heinrich Events from an iceberg/sediment model, *Earth and Planetary Science Letters*, 386, 1–9, 2014b.
- 740 Roche, D., Paillard, D., and Cortijo, E.: Constraints on the duration and freshwater release of Heinrich event 4 through isotope modelling, *Nature*, 432, 379–382, 2004.
- Roche, D. M., Wiersma, A. P., and Renssen, H.: A systematic study of the impact of freshwater pulses with respect to different geographical locations, *Climate Dynamics*, 34, 997–1013, 2010.
- 745 Rohling, E. J., Marsh, R., Wells, N. C., Siddall, M., and Edwards, N. R.: Similar meltwater contributions to glacial sea level changes from Antarctic and northern ice sheets, *Nature*, 430, 1016–1021, 2004.
- Romé, Y. M., Ivanovic, R. F., Gregoire, L. J., Sherriff-Tadano, S., and Valdes, P. J.: Millennial-scale climate oscillations triggered by deglacial meltwater discharge in last glacial maximum simulations, *Paleoceanography and Paleoclimatology*, p. e2022PA004451, 2022.
- Rousseau, D.-D., Boers, N., Sima, A., Svensson, A., Bigler, M., Lacroix, F., Taylor, S., and Antoine, P.: (MIS3 & 2) millennial oscillations in Greenland dust and Eurasian aeolian records—A paleosol perspective, *Quaternary Science Reviews*, 169, 99–113, 2017.
- 750 Rousseau, D.-D., Bagniewski, W., and Ghil, M.: Abrupt climate changes and the astronomical theory: are they related?, *Climate of the Past*, 18, 249–271, 2022.
- Samelson, R.: Simple mechanistic models of middepth meridional overturning, *J. Phys. Oceanogr.*, 34, 2096–2103, 2004.
- Sanchez Goñi, M. F. and Harrison, S. P.: Millennial-scale climate variability and vegetation changes during the Last Glacial: Concepts and terminology, *Quaternary Science Reviews*, 29, 2823–2827, 2010.
- 755 Sánchez Goñi, M. F., Desprat, S., Daniau, A.-L., Bassinot, F. C., Polanco-Martínez, J. M., Harrison, S. P., Allen, J. R., Anderson, R. S., Behling, H., Bonnefille, R., et al.: The ACER pollen and charcoal database: a global resource to document vegetation and fire response to abrupt climate changes during the last glacial period, *Earth System Science Data*, 9, 679–695, 2017.
- Schilt, A., Baumgartner, M., Schwander, J., Buiron, D., Capron, E., Chappellaz, J., Loulergue, L., Schüpbach, S., Spahni, R., Fischer, H., 760 et al.: Atmospheric nitrous oxide during the last 140,000 years, *Earth and Planetary Science Letters*, 300, 33–43, 2010.
- Schulz, M., Berger, W. H., Sarnthein, M., and Grootes, P. M.: Amplitude variations of 1470-year climate oscillations during the last 100,000 years linked to fluctuations of continental ice mass, *Geophysical Research Letters*, 26, 3385–3388, 1999.
- Seager, R. and Battisti, D. S.: Challenges to our understanding of the general circulation: Abrupt climate change, *Global Circulation of the Atmosphere*, pp. 331–371, 2007.
- 765 Seierstad, I. K., Abbott, P. M., Bigler, M., Blunier, T., Bourne, A. J., Brook, E., Buchardt, S. L., Buizert, C., Clausen, H. B., Cook, E., et al.: Consistently dated records from the Greenland GRIP, GISP2 and NGRIP ice cores for the past 104 ka reveal regional millennial-scale $\delta^{18}\text{O}$ gradients with possible Heinrich event imprint, *Quaternary Science Reviews*, 106, 29–46, 2014.
- Sepulchre, P., Caubel, A., Ladant, J.-B., Bopp, L., Boucher, O., Braconnot, P., Brockmann, P., Cozic, A., Donnadiou, Y., Dufresne, J.-L., et al.: IPSL-CM5A2—an Earth system model designed for multi-millennial climate simulations, *Geoscientific Model Development*, 13, 770 3011–3053, 2020.
- Sherriff-Tadano, S., Abe-Ouchi, A., Yoshimori, M., Oka, A., and Chan, W.-L.: Influence of glacial ice sheets on the Atlantic meridional overturning circulation through surface wind change, *Climate dynamics*, 50, 2881–2903, 2018.
- Siddall, M., Rohling, E. J., Thompson, W. G., and Waelbroeck, C.: Marine isotope stage 3 sea level fluctuations: Data synthesis and new outlook, *Reviews of Geophysics*, 46, 2008.

- 775 Sidorenko, D., Rackow, T., Jung, T., Semmler, T., Barbi, D., Danilov, S., Dethloff, K., Dorn, W., Fieg, K., Gößling, H. F., et al.: Towards multi-resolution global climate modeling with ECHAM6–FESOM. Part I: model formulation and mean climate, *Climate Dynamics*, 44, 757–780, 2015.
- Sidorenko, D., Goessling, H. F., Koldunov, N., Scholz, P., Danilov, S., Barbi, D., Cabos, W., Gurses, O., Harig, S., Hinrichs, C., et al.: Evaluation of FESOM2. 0 coupled to ECHAM6. 3: preindustrial and HighResMIP simulations, *Journal of Advances in Modeling Earth Systems*, 11, 3794–3815, 2019.
- 780 Sime, L. C., Hopcroft, P. O., and Rhodes, R. H.: Impact of abrupt sea ice loss on Greenland water isotopes during the last glacial period, *PNAS*, 116, 4099–4104, <https://doi.org/10.1073/pnas.1807261116>, <https://www.pnas.org/content/116/10/4099>, 2019.
- Singh, H. A., Battisti, D. S., and Bitz, C. M.: A heuristic model of Dansgaard–Oeschger cycles. Part I: Description, results, and sensitivity studies, *Journal of climate*, 27, 4337–4358, 2014.
- 785 Spolaor, A., Vallelonga, P., Turetta, C., Maffezzoli, N., Cozzi, G., Gabrieli, J., Barbante, C., Goto-Azuma, K., Saiz-Lopez, A., Cuevas, C. A., et al.: Canadian Arctic sea ice reconstructed from bromine in the Greenland NEEM ice core, *Scientific reports*, 6, 33 925, 2016.
- Stocker, T. F. and Johnsen, S. J.: A minimum thermodynamic model for the bipolar seesaw, *Paleoceanography*, 18, 2003.
- Stouffer, R. J., Yin, J., Gregory, J., Dixon, K., Spelman, M., Hurlin, W., Weaver, A., Eby, M., Flato, G., Hasumi, H., et al.: Investigating the causes of the response of the thermohaline circulation to past and future climate changes, *Journal of Climate*, 19, 1365–1387, 2006.
- 790 Tarasov, L., Dyke, A. S., Neal, R. M., and Peltier, W. R.: A data-calibrated distribution of deglacial chronologies for the North American ice complex from glaciological modeling, *Earth and Planetary Science Letters*, 315, 30–40, 2012.
- Tesi, T., Muschitiello, F., Mollenhauer, G., Miserocchi, S., Langone, L., Ceccarelli, C., Panieri, G., Chiggiato, J., Nogarotto, A., Hefter, J., et al.: Rapid Atlantification along the Fram Strait at the beginning of the 20th century, *Science advances*, 7, eabj2946, 2021.
- Thomas, E. R., Wolff, E. W., Mulvaney, R., Johnsen, S. J., Steffensen, J. P., and Arrowsmith, C.: Anatomy of a Dansgaard–Oeschger warming transition: High-resolution analysis of the North Greenland Ice Core Project ice core, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- 795 Tierney, J. E., Zhu, J., King, J., Malevich, S. B., Hakim, G. J., and Poulsen, C. J.: Glacial cooling and climate sensitivity revisited, *Nature*, 584, 569–573, 2020.
- Timmermann, A., Gildor, H., Schulz, M., and Tziperman, E.: Coherent resonant millennial-scale climate oscillations triggered by massive meltwater pulses, *Journal of Climate*, 16, 2569–2585, 2003.
- 800 Turner, J. and Marshall, G. J.: *Climate change in the polar regions*, Cambridge University Press, 2011.
- Tzedakis, P., Drysdale, R. N., Margari, V., Skinner, L. C., Menviel, L., Rhodes, R. H., Taschetto, A. S., Hodell, D. A., Crowhurst, S. J., Hellstrom, J. C., et al.: Enhanced climate instability in the North Atlantic and southern Europe during the Last Interglacial, *Nature communications*, 9, 1–14, 2018.
- 805 Ullman, D., LeGrande, A., Carlson, A. E., Anslow, F., and Licciardi, J.: Assessing the impact of Laurentide Ice Sheet topography on glacial climate, *Climate of the Past*, 10, 487–507, 2014.
- Uriarte, A.: *Historia del Clima de la Tierra*. Servicio Central de Publicaciones del Gobierno Vasco, 2009, 403 p, 2019.
- Valdes, P.: Built for stability, *Nature Geoscience*, 4, 414–416, 2011.
- Valdes, P. J., Armstrong, E., Badger, M. P., Bradshaw, C. D., Bragg, F., Crucifix, M., Davies-Barnard, T., Day, J. J., Farnsworth, A., Gordon, C., et al.: The BRIDGE HadCM3 family of climate models: HadCM3@ Bristol v1. 0, *Geoscientific Model Development*, 10, 3715–3743, 2017.
- 810

- van der Schrier, G., Drijfhout, S., Hazeleger, W., Noulin, L., and van Oldenborgh, G.-J.: Increasing the Atlantic subtropical jet cools the circum-North Atlantic region, in: EGU General Assembly Conference Abstracts, p. 4651, 2010.
- Vanneste, H., De Vleeschouwer, F., Martínez-Cortizas, A., Von Scheffer, C., Piotrowska, N., Coronato, A., and Le Roux, G.: Late-glacial elevated dust deposition linked to westerly wind shifts in southern South America, *Scientific reports*, 5, 1–10, 2015.
- 815 Vettoretti, G. and Peltier, W. R.: Thermohaline instability and the formation of glacial North Atlantic super polynyas at the onset of Dansgaard-Oeschger warming events, *Geophysical Research Letters*, 43, 5336–5344, 2016.
- Vettoretti, G. and Peltier, W. R.: Fast physics and slow physics in the nonlinear Dansgaard–Oeschger relaxation oscillation, *Journal of Climate*, 31, 3423–3449, 2018.
- 820 Vettoretti, G., Ditlevsen, P., Jochum, M., and Rasmussen, S. O.: Atmospheric CO₂ control of spontaneous millennial-scale ice age climate oscillations, *Nature Geoscience*, pp. 1–7, 2022.
- Voelker, A. H. et al.: Global distribution of centennial-scale records for Marine Isotope Stage (MIS) 3: a database, *Quaternary Science Reviews*, 21, 1185–1212, 2002.
- Volodin, E. M., Mortikov, E. V., Kostykin, S. V., Galin, V. Y., Lykossov, V. N., Gritsun, A. S., Diansky, N. A., Gusev, A. V., Iakovlev, N. G., Shestakova, A. A., et al.: Simulation of the modern climate using the INM-CM48 climate model, *Russian Journal of Numerical Analysis and Mathematical Modelling*, 33, 367–374, 2018.
- 825 Wagner, J. D., Cole, J. E., Beck, J. W., Patchett, P. J., Henderson, G. M., and Barnett, H. R.: Moisture variability in the southwestern United States linked to abrupt glacial climate change, *Nature Geoscience*, 3, 110–113, 2010.
- Wang, X., Auler, A. S., Edwards, R. L., Cheng, H., Cristalli, P. S., Smart, P. L., Richards, D. A., and Shen, C.-C.: Wet periods in northeastern Brazil over the past 210 kyr linked to distant climate anomalies, *Nature*, 432, 740–743, 2004.
- 830 Wang, Y., Cheng, H., Edwards, R. L., Kong, X., Shao, X., Chen, S., Wu, J., Jiang, X., Wang, X., and An, Z.: Millennial-and orbital-scale changes in the East Asian monsoon over the past 224,000 years, *Nature*, 451, 1090–1093, 2008.
- Welander, P.: A simple heat-salt oscillator, *Dynamics of Atmospheres and Oceans*, 6, 233–242, 1982.
- Wunsch, C.: Abrupt climate change: An alternative view, *Quaternary Research*, 65, 191–203, 2006.
- 835 Zhang, X. and Prange, M.: Stability of the Atlantic overturning circulation under intermediate (MIS3) and full glacial (LGM) conditions and its relationship with Dansgaard-Oeschger climate variability, *Quaternary Science Reviews*, 242, 106 443, 2020.
- Zhang, X., Lohmann, G., Knorr, G., and Xu, X.: Different ocean states and transient characteristics in Last Glacial Maximum simulations and implications for deglaciation, *Climate of the Past*, 9, 2319–2333, 2013.
- Zhang, X., Lohmann, G., Knorr, G., and Purcell, C.: Abrupt glacial climate shifts controlled by ice sheet changes, *Nature*, 512, 290–294, 840 2014.
- Zhang, X., Knorr, G., Lohmann, G., and Barker, S.: Abrupt North Atlantic circulation changes in response to gradual CO₂ forcing in a glacial climate state, *Nature Geoscience*, 10, 518–523, 2017.
- Zhang, X., Barker, S., Knorr, G., Lohmann, G., Drysdale, R., Sun, Y., Hodell, D., and Chen, F.: Direct astronomical influence on abrupt climate variability, *Nature Geoscience*, 14, 819–826, 2021.

Table A1: Models that exhibit spontaneous oscillations.

Study	Model	Period	GHG	Ice sheet	Insolation	FWF	Run Length	Main findings	Mechanisms for D-O cooling event	Mechanisms for D-O warming event	Similar Mechanism to schematic in Fig.2 or 3
Driffhout et al. (2013)	EC-Earth model	PI	PI	PI	PI	None	1125 yrs	Spontaneous cold event that lasts around 100 years	An anomalous high atmospheric blocking over the eastern subpolar gyre causes the cold event. Ocean currents transport sea ice southwards and there is a shut-down of deep water convection in the Labrador Sea.	No warming event reported.	No. Atmospheric blocking-sea-ice-ocean feedback identified as a main cause behind the cold event.
Kleppin et al. (2015)	CCSM4	PI	PI	PI	PI	None	1000 yrs	Spontaneous cold event that lasts around 200 years	The cooling event has a duration of 200 years and is linked to a weakened state of the SPG and deep water convection in the Labrador Sea.	The warming event is triggered by a stronger Icelandic low and therefore deep water convection recovers and SPG circulation resumes.	No. Stochastic atmospheric forcing identified as a potential cause for sea ice variations.
Sidorenko et al. (2015)	ECHAM6-FESOM	Present day (PD)	PD - 1990	PD	PD - 1990	None	350 yrs	Events of sudden reduction of deep water convection and increase of sea ice cover in the Labrador Sea.	An anomalous inflow of warm and saline water into the deep Labrador Sea causes a weakening of the subpolar gyre and modifies the upper freshwater budget over the Labrador Sea.	No warming event reported.	No. The strong surface winds over the subtropical NA alter the Gibraltar Strait outflow path into the NA and are identified as the main cause behind the saline and warm anomalies in the deep NA.
Martin et al. (2015)	KCM	PD	PD	PD	PD	None	1300 yrs	Centennial-scale variability of the AMOC as well as variations in the NA heat content and subpolar gyre strength.	As the NA Current accelerates, deep convection in the Weddell Sea enables a positive heat content anomaly to propagate northwards in the upper Atlantic Ocean. Eventually, the heat anomaly reaches the northern NA resulting in a reduction in deep water formation there.	The retreat of Antarctic Bottom Water (AABW) leads to enhanced meridional density gradient that results in an increased North Atlantic Deep Water (NADW) cell.	No. Interhemispheric teleconnection; Variability in the Southern Ocean deep water convection identified as the main cause behind AMOC oscillations.
Pelzer and Vettoretti (2014); Vettoretti and Pelzer (2016, 2018)	UoT CCSM4	LGM	LGM - 21ka	LGM - ICE-6G (VM5a)	LGM - 21ka	None	5000 yrs	Spontaneous millennial-scale D-O type oscillations	The continuous flux of sea ice from the Arctic basin into the NA subpolar gyre area across the East Greenland Current, favours melting of sea ice as it moves over the warm ocean surface. This freshwater input restratifies the high-latitude NA and results in a considerable decrease in the rate of NA Deep Water formation.	The initiation of the abrupt warming events is associated with the opening of a large polynya over the Irminger Sea. The stability of the water column is key and depends on transport of salt to the subpolar gyre along the Irminger Current and Denmark Strait in the decades preceding the warming event.	Yes
Vettoretti et al. (2022)	CCSM4	Glacial simulations	CO_2 levels from 190 to 225 ppm	LGM - ICE-6G (VM5a)	LGM	None	8000-10000 yrs	Spontaneous millennial-scale D-O type oscillations within a window of CO_2 levels from 190 to 225 ppm	Old sea ice from the Arctic is exported to the NA, sea-ice growth is favoured through ice-albedo feedback, high-latitude convection is reduced through sea-ice melt, and consequently Antarctica and Greenland cool. The interstadial-to-stadial transition happens with fast NA sea-ice expansion and NADW production collapses.	During a stadial, sea ice thins in the Southern Ocean and Antarctica warms. There are increases in salt convergence in the NA, NADW fluctuations are amplified via salt advection feedback, and the volume of NADW increases, allowed by late-stadial decreases in AABW formation. Late-stadial vertical stratification experiences thermohaline instability and the Nordic and Irminger Seas are destabilised, triggering rapid sea-ice loss in the NA and the transition from stadial to interstadial states.	Yes
Armstrong et al. (2021)	HadCM3B	MIS3 - 30ka	30ka	30ka	30ka	None	6000 yrs	Spontaneous millennial-scale D-O type oscillations	Ocean forcing initiates the stadial phase; The collapse of the salinity gradient between the Northern NA and STG leads to a reduced advection in the Nordic Seas and decreased deep-water formation	The initiation of the interstadial phase is associated with a wind-driven atmospheric forcing in the Nordic Seas due to increased regional temperatures, reduced sea ice cover and increased sea level pressure, which enhances wind stress and convection	Yes, partially. The D-O type oscillations reflect a salt oscillator mechanism in the NA.

Continued on next page

Table A1 – continued from previous page

Study	Model	Period	GHG	Ice sheet	Insolation	FWF	Run Length	Main findings	Mechanisms for D-O cooling event	Mechanisms for D-O warming event	Similar Mechanism to schematic in Fig. 2 or 3
Zhang et al. (2021)	COSMOS	MIS3: 40-32ka	40ka	40ka	One transient simulation of 40-32ka and one 40ka snapshot simulation (with 34 ka orbital parameters)	None	+5000 yrs	Spontaneous millennial-scale D-O type oscillations; Orbitally induced AMOC changes	Transitions from warm interstadial to cold stadial are linked to (1) a precession-controlled rise in low-latitude boreal summer insolation by modifying the NA low-latitude hydroclimate and/or (2) an obliquity-controlled reduction in high-latitude annual insolation by altering high-latitude sea ice-ocean-atmosphere interactions.	While the AMOC is in its weak phase, a gradual increase in subsurface temperature in the subtropical ocean together with enhanced northward transport of salt in the NA, drive the AMOC back to its strong phase.	Some similarities: unforced AMOC oscillations are triggered by either the tropical salt impact (linked to precession-controlled summer insolation) and/or the subtropical thermal impact (linked to obliquity-controlled mean annual insolation).
Brown and Galbraith (2016)	CM2Mc	Mixed forcing	$CO_2 = 180$ ppm	PI	Obliquity; 22°; Precession: 90°	None	more than 8000 yrs	Spontaneous millennial-scale D-O type oscillations	During a weak AMOC phase, NA deep convection is largely reduced and there is an expansion of sea ice in the northeast Atlantic. Heat accumulates at depth in the NA linked to the weak advection of warm waters from the tropics.	During a strong AMOC phase, NA deep convection is intense and there is a retreat of sea ice in the northeast Atlantic	Yes, partially. Salt advection is a key driver of the oscillations, specifically the salt exchange between subtropical and subtropical NA.
Klockmann et al. (2018, 2020)	MPLESM	Mixed forcing	$CO_2 = 195-217$ ppm; $CH_4 = 396-494$ ppb; $N_2O = 209-227$ ppb	PI	LGM – 21 ka	None	8000-12350 yrs	Spontaneous millennial-scale D-O type oscillations	Stadial phases correspond to weak AMOC and strong SPG phases. The extensive SPG results in low northward salt transport and deep convection only occurs sporadically in the Iceland basin. The Nordic Seas are entirely ice covered which results in a weak Icelandic Low and therefore in a weak wind stress curl. Subsurface waters in the Nordic Seas are around 3 K warmer than during interstadial phases.	During interstadial phases, the AMOC is strong and the SPG is contracted and weak. There is a broad inflow of salty subtropical water to the subtropical NA. Changes in the SPG are driven by variations in the cross-gyre density difference. The eastern NA is fully ice-free. Deep convection occurs continuously in the Iceland basin, Irminger Sea and the Nordic Seas.	Yes, partially. The proposed mechanism behind the spontaneous AMOC oscillations compromises three components: (1) oscillations in salinity comparable to Pelier and Vettorelli (2014), (2) a density-driven feedback loop comparable to Montoya et al. (2011), and (3) a wind-driven feedback loop comparable to Drijfhout et al. (2013) and Kleppin et al. (2015).
Kuniyoshi et al. (2022)	MIROC4m	Mid-glacial conditions	$CO_2 = 220$ ppm	LGM - ICE-5G	Obliquity: 22.949°; Eccentricity: 0.04; Perihelion: 270° and 90°	None	6000 yrs	Spontaneous millennial-scale D-O type oscillations	Changes in subsurface ocean temperature in the NA plays an important role in modifying the stratification of the vertical water column and then reversing the AMOC mode (thermohaline oscillator)	The SPG remains weak (strong) when the AMOC is weak (strong), as well as during the transitions between the two AMOC modes. There is a positive feedback between AMOC and SPG, in agreement with (Li and Born, 2019).	Yes, partially. The opening of a big polynya determines the stadial-interstadial transition. Abrupt changes in AMOC lead to changes in salt advection with the NA subtropical gyre and works as a positive feedback.

Table A2: List of simulations run under MIS3/mid-glacial conditions.

Study	Model	Period	GHG	Ice sheet	Insolation	FWF	Run Length	Main findings
Guo et al. (2019b)	NorESM1-F	MIS3 – 38 ka	$CO_2 = 215$ ppm; $CH_4 = 550$ ppb; $N_2O = 260$ ppb	Data-constrained 38 ka	38 ka	None	2500 yrs (recently extended to 6000 model years)	The equilibrium MIS3 simulation does not show spontaneous D-O type oscillations. Attempts at perturbing the system into a cold stadial state, by modifying the height of the LIS and atmospheric CO_2 levels, show that the modelled MIS3 interstadial state is rather stable, and thus questioning the occurrence of spontaneous D-O type oscillations in the lack of interactive ice sheet-meltwater dynamics.
Zhang and Prange (2020)	CCSM3	MIS3 – 38 ka	$CO_2 = 215$ ppm; $CH_4 = 501$ ppb; $N_2O = 234$ ppb	ICE-5G ice sheet configuration (Peltier, 2004).	38 ka	12 hosing/extraction experiments with freshwater fluxes from ± 0.005 Sv to ± 0.2 Sv, injected in the Nordic Seas for 500 years.	2170 yrs	AMOC is more sensitive to meltwater fluxes under MIS3 conditions than under LGM conditions. The lower AMOC stability under MIS3 conditions proposes that D-O type oscillations could have been triggered by small perturbations in the ocean surface meltwater forcing e.g. linked to ice-sheet processes.
Continued on next page								

Table A2 – continued from previous page

Study	Model	Period	GHG	Ice sheet	Insolation	FWF	Run Length	Main findings
Kawamura et al. (2017)	MIROC 4m	Mid-glacial state	$CO_2 = 215$ ppm; $CH_4 = 350$ ppb; $N_2O = 200$ ppb	Intermediate-size ice-sheet configuration (at 15 ka)	15 ka	Hosing experiments with freshwater fluxes of 0.05 Sv and 0.1 Sv, injected in the North Atlantic Ocean (50°N to 70°N) for 500 years.	More than 2000 yrs	The climate response to freshwater perturbations is much lower under LGM conditions than under MIS3 conditions. The unperturbed LGM AMOC is unusually weak (around 6 Sv) and thus could barely be further lessened, such that meltwater hosing does not largely affect the large-scale climate.
Vettoretti et al. (2022)	CCSM4	Glacial run	$CO_2 = 210$ ppm	LGM - ICE-6G (VM5a)	LGM	Hosing experiment (H event-like pulse) with freshwater fluxes of 0.05 Sv for 500 years in two separate stadial periods in a glacial simulation run with CO_2 of 210 ppm. The freshwater flux is injected in the North Atlantic (50°N to 70°N).	8000 yrs	The Heinrich simulation has a large Northern Hemisphere temperature and AMOC overshoot after the Heinrich stadial ends. Nevertheless, this fast AMOC rise above regular interstadial levels is in agreement with observations only for a few H-stadial periods (H4 and H5).

Continued on next page

Table A2 – continued from previous page

Study	Model	Period	GHG	Ice sheet	Insolation	FWF	Run Length	Main findings
Zhang et al. (2014, 2017)	COSMOS (ECHAM5-JSBACH-MPIOM)	Mixed forcing	LGM	Sensitivity experiments applying different heights of the NH ice sheets	LGM	Freshwater fluxes from 0.00 Sv to ± 0.02 Sv, injected in the NA for 100-300 years	Snapshot and transient simulations 250-700 yrs	An AMOC bi-stability regime is found under intermediate CO_2 and ice sheet conditions roughly resembling that of the MIS3 climate. In the bi-stable MIS3 regime, transitions from weak to strong AMOC state and vice versa could be initiated by not only gradual variations in LIS height and atmospheric CO_2 but also freshwater perturbations. Changes in the LIS height can initiate a positive atmosphere-ocean-sea ice feedback leading to D-O type climate shifts. A gradual increase in the ice sheet height results in a northward shift of the winds, and favours a more saline Labrador Sea both by reducing the sea ice/freshwater import from the Arctic and increasing the advection of salt into the area.
Continued on next page								

Table A2 – continued from previous page

Study	Model	Period	GHG	Ice sheet	Insolation	FWF	Run Length	Main findings
Zhang et al. (2014, 2017)	COSMOS (ECHAM5-JSBACH-MPIOM)	Mixed forcing	CO ₂ levels increased from 185 to 205 ppm during 500 years.	40% of LGM ice-sheet configuration	LGM	None	500 yrs	For a moderate ice volume, 0.25-0.45 times the LGM, two stable AMOC modes are identified. A variation of 15 ppm in atmospheric CO ₂ concentration – equivalent to changes during D-O cycles containing HE – is enough to trigger oscillations between a weak stadial state to a strong interstadial circulation state.
Zhang et al. (2014, 2017)	COSMOS (ECHAM5-JSBACH-MPIOM)	Mixed forcing	CO ₂ levels increased from 185 to 239 ppm at a rate of 0.02 ppm per year.	Intermediate ice sheet configuration - ice volume equivalent to approximately a sea-level drop of 40 m	LGM	None	600 yrs	
Zhang et al. (2014, 2017)	COSMOS (ECHAM5-JSBACH-MPIOM)	Mixed forcing	CO ₂ levels increased from 185 ppm and 245 ppm at a rate of 0.05 ppm per year.	LGM ice volume	LGM	Persistent freshwater flux of 0.15 Sv	600 yrs	

Table A3: Possible oscillatory behaviour for other time periods

Study	Model	Period	GHG	Ice sheet	Insolation	FWF	Run Length	Main findings
Brugger et al. (2019)	CLIMBER 3 α	415 Ma and 380 Ma	CO ₂ = 1500 ppm. Few runs with 500 ppm, 800 ppm, 2000 ppm	None	S0=1315 W/m ² (415 Ma), 1319 W/m ² (380 Ma), various obliquity values, eccentricities, precession angles	None	5000 yrs	Decadal to centennial temperature fluctuations at high northern latitudes.
Rind et al. (2018)	GISS E2-R (TCADI)	Future warming simulation (experiment "abrupt 4x CO2" in Rind et al. (2018))	4 x PI CO ₂	PD	PD	None	2500 yrs	Multi-centennial cessation with restoration and rapid overshooting in NADW formation

Continued on next page

Table A3 – continued from previous page

Study	Model	Period	GHG	Ice sheet	Insolation	FWF	Run Length	Main findings
Rind et al. (2018)	GISS E2-R (TCADI)	Future warming simulation (experiment "1 pct CO2" in Rind et al. (2018))	CO ₂ rises at 1% per year until 4 x CO ₂ (after 140 years) and then held constant.	PD	PD	None	2500 yrs	Multi-centennial cessation with restoration and rapid overshooting in NADW formation
Rind et al. (2018)	GISS E2-R (TCADI)	Future warming simulation (experiment "rcp85" in Rind et al. (2018))	rcp85 emissions	PD	PD	None	4300 yrs	Multi-centennial cessation with restoration and rapid overshooting in NADW formation

Appendix B

Table B1. Summary of LGM/MIS3-like simulations discussed in the text. Models that reproduce D-O type oscillations are highlighted in bold.

Study	Model	Period	N° of simulations	Run length
Peltier and Vettoretti (2014)	UofT CCSM4	PMIP4 LGM	1	5000
Lohmann et al. (2020)	AWI-ESM1-1-LR	PMIP4 LGM	1	1300
Sidorenko et al. (2019)	AWI-ESM-2-1-LR	PMIP4 LGM	1	600
Tierney et al. (2020)	CESM1.2	PMIP4 LGM	1	1800
Valdes et al. (2017)	HadCM3B-M2.1aD	PMIP4 LGM	3	400-2900 ^a
Lhardy et al. (2021)	iLOVECLIM1.1.4	PMIP4 LGM	2	5000
Volodin et al. (2018)	INM-CM4-8	PMIP4 LGM	1	50
Sepulchre et al. (2020)	IPSLCM5A2	PMIP4 LGM	1	1200
Ohgaito et al. (2021)	MIROC-ES2L	PMIP4 LGM	1	8960
Mauritsen et al. (2019)	MPI-ESM1.2	PMIP4 LGM	1	3850
Armstrong et al. (2021)	HadCM3B-M2.1aD	MIS3 (30 ka)	1	6000
Zhang et al. (2021)	COSMOS	MIS3 (40-32 ka)	2	5000
Guo et al. (2019b)	NorESM	MIS3 (38 ka)	1	+6000
Zhang and Prange (2020)	CCSM3	MIS3 (38 ka)	1	2170
Kawamura et al. (2017)	MIROC4m	Mid-glacial conditions	1	+2000
Kuniyoshi et al. (2022)	MIROC4m	Mid-glacial conditions	2	6000
Vettoretti et al. (2022)	CCSM4	Glacial conditions	4^b	8000
Brown and Galbraith (2016)	CM2Mc	Mixed forcing	1	+8000
Klockmann et al. (2018)	MPI-ESM	Mixed forcing	3	+8000
Zhang et al. (2014)	COSMOS	Mixed forcing	11 ^c	300-4000 ^d

^a Only one simulation run longer than 2000 model years

^b Four simulations run with CO_2 levels: 200, 210, 220, 225 ppm

^c We do not consider FWF runs nor transient simulations forced with varying CO_2 and/or NH ice sheet height

^d Only two simulations with a duration of +2000 model years

Appendix C

Table C1. Contributing members to PMIP4/CMIP6 that have run simulations under LGM or MIS3 conditions.

Model	Period	Run length
ACCESS-ESM1-5	-	-
AWI-ESM-1-1-LR	LGM	1300
CESM2	-	-
CNRM-CM6-1	-	-
EC-Earth3-LR	-	-
FGOALS-f3-L	-	-
FGOALS-g3	-	-
GISS-E2-1-G	-	-
HadGEM3-GC31-LL	-	-
INM-CM4-8	LGM	50
IPSL-CM6A-LR	-	-
MIROC-ES2L	LGM	+8000
MPI-ESM1-2	LGM	3850
MRI-ESM2-0	-	-
NESM3	-	-
NorESM1-F	MIS3	6000
NorESM2-LM	-	-