

In situ LA-ICPMS U-Pb dating of sulfates: Applicability of carbonate reference materials as matrix-matched standards

Aratz Beranoaguirre^{1,2,3,4}, Iuliana Vasiliev⁵, Axel Gerdes^{2,3}

¹ Institut für Angewandte Geowissenschaften, Karlsruher Institut für Technologie, Adenauerring 20 b, 76131 Karlsruhe, Germany

² Institut für Geowissenschaften, Goethe-Universität Frankfurt, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

³ Frankfurt Isotope and Element Research Center (FIERCE), Goethe-Universität Frankfurt, Frankfurt am Main, Germany

⁴ Geologia Saila, Euskal Herriko Unibertsitatea UPV/EHU, Sarriena z/g, 48940 Leioa, Spain

⁵ Senckenberg Biodiversity and Climate Research Centre (BiK-F), Senckenberganlage 25, 60325 Frankfurt am Main, Germany

10 Correspondence to: Aratz Beranoaguirre (aratz.beranoaguirre@kit.edu)

Abstract. Recent developments in analytical capabilities in the field of in situ laser ablation mass spectrometry (LA-ICPMS) have expanded the applications of U-Pb geochronometers in low-U minerals such as carbonates or garnets. The rapid evolution of the technique relies on well-characterized matrix-matched reference materials. In this article, we explore the suitability of using carbonate as “almost-matrix-matched reference materials” for in situ U-Pb dating of sulfates. For such purpose, we have 15 used the astrochronologically dated gypsum and anhydrite samples deposited during the Messinian Salinity Crisis (5.97 – 5.33 Ma) and compared these dates with the U-Pb ages obtained by LA-ICPMS. Although the majority of the samples failed due to the elevated common Pb content and low $^{238}\text{U}/^{204}\text{Pb}$ ratios, five of the samples showed a higher dispersion on U/Pb ratios. The obtained dates in four of these samples are comparable with the expected ages while another gave an unexpected younger age, each of them with 6–11% of uncertainty. The pit depth of the spots showed that the sulfates ablate similar to carbonates, 20 so the offset due to the crater geometry mismatch or downhole fractionation can be assumed to be negligible. To sum up, the bias between the U-Pb and expected cyclostratigraphic ages, if any, is included in the uncertainty and thus, the results obtained here suggest that carbonate reference material is currently the best option for standardisation of in situ U-Pb sulfate analyses.

1 Introduction

Recent developments in instrumentation and analytical capabilities of LA-ICPMS techniques have greatly expanded the 25 applicability of the U-Pb geochronometer. The high spatial resolution, low cost of analysis and high throughput with relatively good precision (Schaltegger et al., 2015) achievable with the new generation of laser and mass spectrometers favour the study of minerals with low and heterogeneous U concentrations like carbonates or garnets (e.g. Roberts et al., 2020). In fact, carbonate geochronology has gone from scarce publications that involve tedious and long-lasting isotope dilution techniques 30 (e.g., Brannon et al., 1996; Grandia et al., 2000; Woodhead et al. 2006, 2012; Rasbury and Cole, 2009) to a bloom of dozens of publications per year (extensive review in Roberts et al., 2020). Likewise, garnet U-Pb dating is rapidly developing in skarn or metamorphic garnets, with U contents even below 100 parts per billion (e.g., Burisch et al., 2019; Yan et al., 2020; Millonig

et al., 2020). In addition, several laboratories have started to investigate the possibility of measuring other types of minerals: dolomites (Burisch et al., 2017), fluorite (Piccione et al., 2019; Lenoir et al., 2021), nacrite (Piccione et al., 2019) or anatase (Sindern et al., 2019), among others.

35 The rapid evolution of U-Pb dating in low-U phases is closely related to the availability of reference materials (WC-1 carbonate, Roberts et al., 2017; or Mali garnet, Seaman et al., 2017). Well-characterized matrix-matched reference material is essential for U-Pb analyses by ion probe or laser ablation as sample matrix affects the ablation, transport and ionisation (Sylvester, 2008; Yang et al., 2018). Indeed, LA-ICPMS dates could only be as good as the homogeneity of the reference materials, and the accuracy and precision to which such material is known (Schaltegger et al., 2015). Several authors, however,
40 have appraised the suitability of using non-matrix-matched standardisation, with different levels of success. Deng et al. (2017) and Wafforn et al. (2018) used 91500 and GJ1 zircon, respectively, to correct U/Pb fractionation of garnet and assumed they obtained the correct ages, whereas Yang et al. (2018) measured 11 % too old garnet ages using zircon standardisation. Similarly, Parrish et al. (2018) measured Mud Tank zircon within carbonate analyses and reported a bias between zircon and calcite of c. 4.7%. Piccione et al. (2019) used the WC-1 carbonate reference material for fluorite analysis assuming that the
45 bias between calcite and fluorite may likely be less than the one between calcite and zircon.

This study aims to continue opening new possibilities in the field of in-situ U-Pb dating of low-U minerals, by (I) demonstrating that sulfates can be dated by U-Pb and (II) examining the suitability and reliability of using calcite as “almost matrix-matched reference material” for sulfates. Accurate U-Pb dating of sulfates could contribute to a better understanding of their formation and/or transformation (hydration-dehydration) with the potential of dating diagenetic, pedogenic or tectonic processes.
50 Gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) and Anhydrite (CaSO_4) are the two most abundant sulfates of marine and non-marine evaporite deposits (e.g. Murray, 1963; Babel and Schreiber, 2014). Sedimentary gypsum forms by direct precipitation out of water evaporation under arid climatic conditions in hydrologically restricted environments. Under terrestrial evaporitic conditions, gypsum is the dominant primary mineral and anhydrite forms through gypsum dehydration caused during diagenesis. In the presence of water at shallower levels, the anhydrite is rapidly converted back to gypsum (e.g. Conley and Bundy, 1958; Murray,
55 1964; Ossorio et al., 2014; Warren, 2016). Although less frequent, non-evaporitic gypsum formation can also take place (see Van Driessche et al., 2019 and references therein).

In the absence of sulfate matrix-matched reference material, we have assumed that the bias between calcite and sulfate is smaller than with the other available reference materials. Both minerals behave very similarly during ablation (e.g., drill speed, U/Pb downhole fractionation, etc.) and ionization in the plasma (Ca^{2+} as the main cation). For evaluating the suitability of the
60 calcite-based corrections, we have analysed gypsum and anhydrite samples from the Messinian Salinity Crisis (MSC) in the Mediterranean Sea (Roveri et al., 2014a, 2014b; Vasiliev et al., 2017; Grothe et al., 2020; Andreetto et al., 2021) and compared them with their astrochronological data (calibrated with astronomically tuned timescales, such as Milankovic cycles, Laskar et al. 1999). Chronostratigraphy of Late Miocene to Early Pliocene within the MSC is well constrained (CIESM, 2008; Manzi et al., 2013, Roveri et al., 2014a) and thus, makes those samples ideal for comparison purposes.

65 2 Geological Background

The Messinian Salinity Crisis (MSC, 5.97-5.33 Ma) successions record extreme fluctuations in the Mediterranean's paleoceanographic and environmental conditions (e.g. Hsü et al., 1973; Krijgsman et al., 1999; Manzi et al., 2013). At the end of the Miocene, the Mediterranean's connections with the Atlantic Ocean were extremely reduced (e.g. Flecker et al., 2015; Krijgsman et al., 2018) whereas the freshwater supply from the Eastern Paratethys increased (Flecker and Ellam, 2006; 70 Krijgsman et al., 2010). Those paleoceanographic changes lead to the formation of hypersaline water bodies and the deposition of a kilometre-thick evaporite unit (Fig. 1A) (Ryan, 2009). The original definition of the MSC referred to a marked environmental change at the base of the Tripoli diatomite formation (Sicily, Italy) close to the Tortonian/Messinian boundary (Sell, 1960). Astronomical tuning of the pre-evaporitic succession showed that the MSC onset was synchronous throughout 75 the Mediterranean (e.g., Krijgsman et al., 1999; Manzi et al., 2018; Meijlison et al., 2018). According to the shallow water-deep basin model (Hsü et al., 1973; Roveri et al., 2014a), evaporite precipitation was associated with a sea-level drop in the range of 1500 m, up to the almost complete desiccation of the Mediterranean, culminating in halite precipitation and marked by the incisions of deep canyons at the Mediterranean margins. However, a non-evaporitic gypsum formation during MSC has been also described (Hsü et al., 1973). Isotope analyses of gypsum hydration water and the salinity of fluid inclusions in MSC 80 gypsum indicate large freshwater inputs during gypsum formation (Natalicchio et al., 2014; Evans et al., 2015; Costanzo et al., 2019). Additionally, suggestions of the important role of sulfur-oxidizing bacteria in biogeochemically mediated gypsum formation (Grothe et al., 2020) are increasingly used to explain a low salinity, yet high concentrations of Ca^{2+} and SO_4^{2-} (Clauer et al., 2000), during the formation of MSC evaporites.

According to previous publications (Roveri et al., 2008a, 2008b and 2014a), the MSC can be separated into three main stages (Fig. 1B). Stage I (5.97-5.60 Ma), the so-called Primary Lower Gypsum (PLG, Lugli et al., 2010), is defined by the deposition 85 of primary selenite gypsum unit. During the stage II (5.60-5.55 Ma), large evaporite deposits occurred (Resedimented Lower Gypsum unit, RLG), which includes halite, gypsum cumulates and brecciated limestones ('Calcare di Base' type 3, Manzi et al., 2011). Likewise, clastic gypsum derived from the dismantlement of the PLG unit can be found within this stage. Finally, alternating gypsum (mainly bottom grown selenite and cumulate) and fine- to coarse-grained terrigenous deposits form the Upper Gypsum unit (UG, stage III, 5.55-5.33 Ma). There is no outcrop with the complete section of the MSC, but different 90 segments are well exposed throughout the Mediterranean

3 Methodology

U-Pb data was acquired *in situ* in polished mounts and slabs using a RESOLution 193 nm ArF excimer laser (CompexPro 102) equipped with a two-volume ablation cell (Laurin Technic S155) coupled to a (I) single collector (SC)-ICPMS (ElementXR, ThermoScientific) or (II) multicollector (MC)-ICPMS (Neptune Plus, ThermoScientific) at FIERCE (Frankfurt Isotope & 95 Element Research Center), Goethe University Frankfurt. The method is modified after Ring and Gerdes (2016) and Burisch et al. (2017). Samples are pre-screened in order to identify sub-zones with a higher $^{238}\text{U}/^{206}\text{Pb}$ ratio before each analytical session.





Fig. 1: A) Geological sketch of the Messinian evaporite deposits along the Mediterranean Sea (modified after Rouchy and Caruso, 2006). Note that only the successfully dated sample locations are shown. **B)** Chronostratigraphy of Late Miocene to Early Pliocene with MSC events in the Mediterranean (modified from Vasiliev et al., 2017).

The first sessions, between December 2019 and May 2020, were performed with the SC-ICPMS. Prior to the measurements, signal strength was tuned for maximum sensitivity while keeping oxide formation below ~ 0.5 % (UO/U) and element fraction low (e.g. Th/U ~ 0.9). This was done by ablating at 3 $\mu\text{m}/\text{s}$ with a 60 μm spot at 6 Hz and 3.5 J/cm² fluence in the glass SRMNIST 612 (Jochum et al., 2011). The average sensitivity obtained for the line is ca. 100,000 cps per $\mu\text{g/g}$ for ²³⁸U. The 105 detection limits (4 x background signal) of the instrument for ²⁰⁶Pb and ²³⁸U were c. 0.3 and 0.03 ng/g. Data were acquired in fully-automated mode overnight. Each analysis consists of 18 s background acquisition followed by 18 s of sample ablation and 20 s washout. During 36 s data acquisition, the signal of ²⁰⁶Pb, ²⁰⁷Pb, ²⁰⁸Pb, ²³²Th and ²³⁸U were detected by peak jumping in simultaneous analogue and pulse counting mode. Detailed data acquisition parameters are summarised in Table 1.

Due to the low precision obtained in those sessions, where only two samples from a single session can be considered acceptable 110 (see results and discussion), the use of the more sensitive MC-ICPMS (Craig et al., 2018, 2020) was deemed necessary for subsequent measurements. The sessions with the MC-ICPMS were carried out between July 2020 and December 2020. As for the single collector, signal strength was tuned for maximum sensitivity while keeping oxide formation below ~ 0.5 % (UO/U) and element fraction low (e.g. Th/U ~ 0.9). In that case, it was done with a 35 μm , 6 Hz, ca. 3.5 J/cm² fluence and a 3 $\mu\text{m}/\text{s}$ line ablated in the glass SRMNIST 614 (Jochum et al., 2011). The average sensitivity obtained for the line is ca. 120,000 cps 115 per $\mu\text{g/g}$ for ²³⁸U (note the smaller spot size compare to the SC-ICPMS). The detection limits in the multicollector ICPMS were c. 0.3 and 0.01 ng/g for ²⁰⁶Pb and ²³⁸U, respectively. The analyses were done during 31 s (14 s background and 16 s of ablation) in static mode, measuring ²⁰⁶Pb and ²⁰⁷Pb with Secondary Electron Multipliers (SEMs), ²⁰²Hg and ²⁰⁴Pb with Multiple Ion Counters (MICs) and ²³²Th and ²³⁸U on Faraday cups with $10^{13} \Omega$ amplifiers. Faraday signals in V are converted into cps by using a factor of 62,400,000. Detailed data acquisition parameters are summarised in Table 2.

120 In each analytical session, soda-lime glass SRMNIST614 was used as the primary reference material to correct for mass bias (²⁰⁷Pb/²⁰⁶Pb) and the interelement fractionation and instrumental drift (²⁰⁶Pb/²³⁸U) throughout the entire analytical session. Carbonate reference material WC-1 (254 Ma, Roberts et al., 2017) was used to determine the difference of the Pb/U fractionation between carbonate and synthetic glass matrix. Depending on the analytical conditions (i.e, spot size, laser fluence, torch position, sample gas flows, etc.) the matrix effect can vary up to 12 % (FIERCE laboratory observation, e.g. Cruset et 125 al. 2021) and even at similar tuning parameters, two sessions separated by some weeks could result in different Pb/U correction factors. So far, this behaviour is not very well understood and due to its unpredictability, the matrix correction is calculated for each session (see below). Secondary reference calcite materials, ASH-15D calcite (2.965 ± 0.011 Ma, Nuriel et al., 2021), B-6 (42.99 ± 0.99 Ma, only LA-ICPMS data, Pagel et al., 2018) and in-house calcite (reproducible age of ca. 36 Ma) were measured for quality control. Not all the secondary reference materials were used in each session (see information in Tables 1 130 and 2).

Laboratory & Sample Preparation	
Laboratory name	FIERCE, Frankfurt Isotope & Element Research Center Goethe Universität, Frankfurt am Main
Sample type/mineral	Sulphate
Sample preparation	25 mm polished resin mounts
Imaging	Petrographic microscope & 2400 dpi digital scan
Laser ablation system	
Make, Model & type	RESOlution ArF excimer laser (COMplex Pro 102)
Ablation cell	Two-volume ablation cell (Laurin Technic S155)
Laser wavelength	193 nm
Pulse width	20 ns
Fluence	2 J/cm ²
Repetition rate	10 Hz
Pre-ablation	4 pulses (same parameters as main ablation)
Ablation duration	18 s
Ablation rate	~ 0.6 µm/s (in the primary RM), ~ 0.9 µm/s (in carbonate and sulfate)
Spot shape & size	Round, 154 µm (diameter)
Sampling mode	Static spot ablation
Gasses	Sample cell: He. Funnel: He + Ar. Tubbing: He + Ar + N ₂
Gas flows	He (300 ml/min), Ar (1050 ml/min), N ₂ (8 ml/min).
ICP-MS Instrument	
Make, Model & type	ThermoScientific ElementXr sector field ICP-MS
Sample introduction	Ablation aerosol
RF power	1300 W
Detection system	Secondary electron multiplier (with conversion dynode at -8kV). Simultaneous analogue and counting (pulse) modes of detection (conversion factors calculated per mass and applied offline). Magnetic field fixed. Detection by peak jumping with electrostatic analyzer.
Masses measured	206, 207, 232, 238
Dwell times	206: 6.4 ms, 207: 7.5 ms, 232: 2.0 ms, 238: 4.6 ms
Samples per peak/integration type	4 for all masses/average
Total time per run	99 ms
Number of runs/total time	370 / 36.6 s
Acquisition mode	Trigger from laser (20 s after pre-ablation), background: 18 s, ablation: 18 s
Dead time	29 ns
Data Processing	
Gas blank	20 s on-peak zero subtracted.
Calibration strategy	SRMNIST614 as primary RM, WC-1 as offset RM, and ASH15D as validation RM.
Reference Material (RM) information	Soda-lime glass SRMNIST614 (Jochum et al., 2011), WC-1 (Roberts et al., 2017), ASH15D (Nuriel et al., 2021)
Data processing / LIEF correction	In-house VBA spreadsheet program (Gerdes and Zeh, 2006, 2009). Intercept method for LIEF correction, assumes Pbc corrected WC-1 and samples behave identically.
Mass discrimination	²⁰⁷ Pb/ ²⁰⁶ Pb (0.2%) and ²⁰⁶ Pb/ ²³⁸ U (5%) normalised to primary standard
Common-Pb correction	No common-Pb correction applied to the data.
Uncertainty level & propagation	Uncertainties are quoted at 2σ absolute and are propagated by quadratic addition of the within run precision (SD of the mean of ratios in log-ratio space), counting statistics, background, common Pb correction (if applicable) and the excess of scatter (calculated from the primary RM). In addition, an excess of variance of 1.45 % (1σ), calculated from the offset RM, was added quadratically to the ²⁰⁶ Pb/ ²³⁸ U ratios. Systematic uncertainties are reported as an expanded uncertainty, considering long term reproducibility (1.5%, 2σ) and decay constant uncertainties.
Quality control / Validation	WC-1: 254.7 ± 2.3 / 4.4 Ma (2s, MSWD = 1.00, n = 28) ASH15D: 3.004 ± 0.153 / 0.159 Ma (2s, MSWD = 0.85, n = 28) (Ages are the ²⁰⁶ Pb/ ²³⁸ U lower intercept ages of the calculated isochrons with the concordia curve in the Tera-Wasserburg space)

Table 1: LA-SC-ICPMS U-Pb analysis procedure at Goethe University Frankfurt, FIERCE laboratory.

Laboratory & Sample Preparation	
Laboratory name	FIERCE, Frankfurt Isotope & Element Research Center Goethe Universität, Frankfurt am Main
Sample type/mineral	Sulphate
Sample preparation	25 mm polished resin mounts and polished thick sections
Imaging	Petrographic microscope & 2400 dpi digital scan
Laser ablation system	
Make, Model & type	RESolution ArF excimer laser (COMPex Pro 102)
Ablation cell	Two-volume ablation cell (Laurin Technic S155)
Laser wavelength	193 nm
Pulse width	20 ns
Fluence	2 J/cm ²
Repetition rate	10 Hz
Pre-ablation	2 pulses (same parameters as main ablation)
Ablation duration	16 s
Ablation rate	~ 0.6 µm/s (in the primary RM) ~ 0.9 µm/s (in carbonate and sulfate)
Spot shape & size	Round, 130 µm (75 µm for primary RM), except for session 1 (90 µm for all spots)
Sampling mode	Static spot ablation
Gases	Sample cell: He. Funnel: He + Ar. Tubbing: He + Ar + N ₂
Gas flows	He (300 ml/min), Ar (~950 ml/min), N ₂ (5–10 ml/min). Ar and N ₂ are turned each session, so the values can be slightly different
ICP-MS Instrument	
Make, Model & type	ThermoScientific Neptune Plus multi-collector ICP-MS
Sample introduction	Ablation aerosol
RF power	1300 W
Detection system	Simultaneous multi-collection. Secondary Electron Multipliers (SEMs) for ²⁰⁶ Pb and ²⁰⁷ Pb Multiple Ion Counters (MICs) for ²⁰² Hg and ²⁰⁴ Pb Faraday cups with 10 ¹³ Ω amplifiers for ²³² Th and ²³⁸ U
Masses measured	202, 204, 206, 207, 232, 238
Total time per run	131 ms
Number of runs/total time	230 / 30.1 s
Acquisition mode	Trigger from laser (14 s after pre-ablation), background: 15 s, ablation: 16 s
Data Processing	
Gas blank	15 s on-peak zero subtracted.
Calibration strategy	SRMNIST614 as primary RM, WC-1 as offset RM, and ASH15D, B6 & in-house calcite as validation RM.
Reference Material (RM) information	Soda-lime glass SRMNIST614 (Jochum et al., 2011), WC-1 (Roberts et al., 2017), ASH15D (Nuriel et al., 2021), B-6 (Pagel et al., 2018), CalBraun (in-house calcite RM)
Data processing / LIEF correction	In-house VBA spreadsheet program (Gérdes and Zeh, 2006, 2009). Intercept method for LIEF correction, assumes Pbc corrected WC-1 and samples behave identically.
Mass discrimination	²⁰⁷ Pb/ ²⁰⁶ Pb and ²⁰⁶ Pb/ ²³⁸ U normalised to primary standard (variable in each session)
Common-Pb correction	No common-Pb correction applied to the data.
Uncertainty level & propagation	Uncertainties are quoted at 2σ absolute and are propagated by quadratic addition of the within run precision (SD of the mean of ratios in log-ratio space), counting statistics, background, common Pb correction (if applicable) and the excess of scatter (calculated from the primary RM). In addition, an excess of variance calculated for each session from the offset RM, was added quadratically to the ²⁰⁶ Pb/ ²³⁸ U ratios. Systematic uncertainties are reported as an expanded uncertainty, considering long term reproducibility (1.5%, 2σ) and decay constant uncertainties.
Quality control/ Validation	Sequence 1: WC-1: 254.8 ± 1.9 / 4.3 Ma (2s, MSWD = 1.0, n = 12) B-6: 42.73 ± 0.59 / 0.87 Ma (2s, MSWD = 0.84, n = 12) CalBraun: 36.72 ± 1.23 / 1.35 Ma (2s, MSWD = 0.89, n = 12) Sequence 2: WC-1: 254.1 ± 2.0 / 4.4 Ma (2s, MSWD = 1.0, n = 20) B-6: 42.66 ± 0.47 / 0.80 Ma (2s, MSWD = 0.50, n = 22) CalBraun: 36.07 ± 0.65 / 0.85 Ma (2s, MSWD = 0.61, n = 22) Sequence 3: WC-1: 254.5 ± 3.2 / 5.0 Ma (2s, MSWD = 1.0, n = 10) ASH15D: 3.060 ± 0.193 / 0.198 Ma (2s, MSWD = 1.0, n = 10) B-6: 43.54 ± 0.79 / 1.02 Ma (2s, MSWD = 1.13, n = 10) Sequence 4: WC-1: 254.5 ± 1.6 / 4.1 Ma (2s, MSWD = 1.0, n = 20) ASH15D: 3.091 ± 0.102 / 0.112 Ma (2s, MSWD = 0.88, n = 20) B-6: 43.83 ± 0.39 / 0.77 Ma (2s, MSWD = 0.56, n = 20) (Ages are the ²⁰⁶ Pb/ ²³⁸ U lower intercept ages of the calculated isochrons with the concordia curve in the Tern-Wasserburg space. WC-1 RM are anchored at 0.85 value of ²⁰⁷ Pb/ ²⁰⁶ Pb)

Table 2: LA-MC-ICP-MS U-Pb analysis procedure at Goethe University Frankfurt, FIERCE laboratory.

135 Raw data were corrected offline using an in-house VBA spreadsheet program (Gerdes and Zeh, 2006, 2009). Following background and interferences corrections, outliers ($\pm 2\sigma$) were rejected based on the time-resolved $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{206}\text{Pb}/^{238}\text{U}$ ratios and the Pb and U signal. All in all, five sessions were performed, and the matrix Pb/U correction factors (carbonate vs SRMNIST glass) applied to each of them are as follows: 4.5 % for SC-ICPMS session, 8 % for MC-ICPMS session 1 (same spot size for both carbonate and SRMNIST glass, see Table 2), 0.5 % for session 2 (different spot size, Table 2), 0 % for
140 session 3, and 0 % for session 4.. The $^{206}\text{Pb}/^{238}\text{U}$ downhole fractionation during 16/18 s depth profiling was estimated to be 3%, based on the common Pb-corrected WC-1 analyses, and was applied as an external correction to all sulfates analyses and secondary reference materials. Uncertainties for each isotopic ratio are the quadratic addition of the within-run precision, counting statistic uncertainties of each isotope, and the excess of scatter and variance (Horstwood et al., 2016) calculated from the SRMNIST 614 and the WC-1 after drift correction. To account for the long-term reproducibility of the method we added
145 by quadratic addition an expanded uncertainty of 1.5% to the final age of all analysed samples (Montano et al., 2021). This was deducted from repeated analyses of ASH-15D in the FIERCE laboratory between 2017 and 2019. Data were displayed in Tera-Wasserburg plots and ages were calculated as lower concordia-curve intercepts using the same algorithms as Isoplot 4.15 (Ludwig, 2012). All uncertainties are reported at the 2σ level. After the analysis, the depth of the ablation pit was measured in several spots per sample, including the WC-1 and SRMNIST 614 reference materials, using the Keyence VHX 6000 digital
150 microscope.

4 Samples and results


4.1 U-Pb Dating

U-Pb dating was applied to 32 samples from the different locations and all available gypsum/anhydrite varieties (large selenite crystals, banded selenite, gypsum cumulates, anhydrite, halite with gypsum and anhydrite intercalation) across the
155 Mediterranean Sea (Fig. 1), which display variable contents of Pb and U. Only five of them were successfully dated (15 % of success). The undatable samples are characterized by analyses that clustered near the common Pb intercept, disclosing a large amount of common Pb (Fig. 2). This low μ ($^{238}\text{U}/^{204}\text{Pb}$ ratio) makes it impossible to draw any regression line. No link between successful/unsuccessful samples and their texture could have been established and both successful and unsuccessful samples have been found within the same type of gypsum. The successfully dated samples are described below, and their results are
160 presented in Fig. 3 as well as in Tables 3 and 4.

4.1.1 Sample MTO 4-4

The MTO 4-4 sample was collected at the Monte Tondo gypsum quarry, located within the Vena del Gesso basin (along the western Romagna Apennines), and belongs to the PLG (Lugli et al., 2007, 2010; Vasiliev et al., 2017). It is a banded selenite (type F4 of Lugli et al., 2010) and the cyclostratigraphic age is 5.920 Ma, close to the onset of the MSC. The sample was
165 measured in three different sessions. The maximum U and Pb content on the analysed spots are 2.34 $\mu\text{g/g}$ and 3.85 $\mu\text{g/g}$,

respectively, depicting a maximum U/Pb ratio of 98.4 in the best case. The first of the sessions was measured with the SC-ICPMS and the analyses define a regression line with a lower intercept at 6.01 ± 1.19 Ma ($\pm 2\sigma$, MSWD = 1.07, Fig. 3). The other two sessions were measured with the MC-ICPMS and the lower intercept of the regression lines are 5.55 ± 0.61 Ma ($\pm 2\sigma$, MSWD = 1.00, Fig. 3) and 5.73 ± 0.37 Ma ($\pm 2\sigma$, MSWD = 1.13, Fig. 3).



170

Fig. 2: Diagram showing $U_{\text{mean}} / Pb_{\text{mean}}$ content vs. maximum value on $^{238}\text{U}/^{206}\text{Pb}$ axis. The successfully dated samples have a distinctively higher U/Pb heterogeneity.

4.1.2 Sample MTO 11-3

This sample was also collected by Vasiliev et al., (2017) at the Monte Tondo gypsum quarry. It is a massive selenite (F3 of 175 Lugli et al., 2010) and belongs to the younger cycles of the PLG. Its estimated cyclostratigraphic age is 5.701 Ma. MTO 11-3 was measured as well in three different sessions. The maximum U and Pb content on the analysed spots are 5.49 $\mu\text{g/g}$ and 0.97 $\mu\text{g/g}$, respectively, depicting a maximum U/Pb ratio value of 155.2 in the best case. The first of the sessions was measured with the SC-ICPMS and the analyses define a regression line with a lower intercept at 5.40 ± 0.84 Ma ($\pm 2\sigma$, MSWD = 1.13, Fig. 3). The other two sessions were measured with the MC-ICPMS and the lower intercept of the regression lines are $5.46 \pm$ 180 0.44 Ma ($\pm 2\sigma$, MSWD = 1.41, Fig. 3) and 5.55 ± 0.32 Ma ($\pm 2\sigma$, MSWD = 1.03, Fig. 3).

4.1.3 Sample BOX 108

BOX 108 is a halite with anhydrite nodules. It comes from the borehole EMS-4 (Cattolica Eraclea) in the Caltanissetta Basin (southwest of Sicily) and was donated to Prof. Cita (University of Milano). The core was drilled from - 82 m to - 665 m below 185 sea level and the sample was located almost at the bottom (approximately at - 610 m). Cyclostratigraphic ages point to 5.55- 5.60 Ma. The analyses were made both in halite and in anhydrite, but only the anhydrite was successful. It was measured twice with the MC-ICPMS. The maximum U and Pb content on the analysed spots are 5.70 $\mu\text{g/g}$ and 1.67 $\mu\text{g/g}$, respectively, depicting a maximum U/Pb ratio value of 158.0 in the best case. The analyses define a regression line with a lower intercept at 5.55 ± 0.35 Ma ($\pm 2\sigma$, MSWD = 1.01, Fig. 3) in the first of the sessions and 5.54 ± 0.38 Ma ($\pm 2\sigma$, MSWD = 1.49, Fig. 3) in 190 the second.




Fig. 3: Tera-Wasserburg diagram ($^{207}\text{Pb}/^{206}\text{Pb}$ vs. $^{238}\text{U}/^{206}\text{Pb}$) for the samples MTO 4-4, MTO 11-3, BOX 108, Pu 05 and BCR 9644. The blue ellipses and error envelope in samples MTO 4-4 and MTO 11-3 correspond to the analyses with the SC-ICPMS, while the orange and black colours refer to two independent sessions with the MC-ICPMS. Both internal and expanded uncertainties are $\pm 2\sigma$.

4.1.4 Sample BCR9644

The sample BCR9644 was collected from the cores of Deep Sea Drilling Program Site 42A hole 376 cored in 1975 West of Cyprus and stored at Bremen International Ocean Drilling Program repository. BCR9644 was collected from a Gypsum Breccia, at 170.28 m below sea level and has a stratigraphic age of ca. 5.55 - 5.60 Ma. It was measured twice with the MC-ICPMS. The maximum U and Pb content on the analysed spots are 2.31 µg/g and 0.61 µg/g, respectively, although Pb rarely exceeds 0.1 µg/g. The maximum U/Pb ratio obtained in that sample is 577.5 in the best case. The low Pb contents imply large error ellipses, but successful regression lines have been defined, with a lower intercept at 2.98 ± 0.34 Ma ($\pm 2\sigma$, MSWD = 0.79, Fig. 3) in the first of the sessions and 2.98 ± 0.32 Ma ($\pm 2\sigma$, MSWD = 1.40, Fig. 3) in the second.

4.1.5 Sample Pu 05

This sample was collected in the Ploutis region (Central Crete, Greece) and it is a gypsum breccia. The stratigraphic age of these gypsum units is disputed between being part of the PLG (Zachariasse et al., 2008) but the texture is direct capping by Lago Mare deposits strongly suggest that Pu 05 belongs to the UG unit. Its Cyclostratigraphic age is ca. 5.40 Ma. Pu 05 was also measured twice with the MC-ICPMS. The maximum U and Pb content on the analysed spots are 1.44 µg/g and 0.16 µg/g, respectively, depicting a maximum U/Pb ratio value of 158.0 in the best case. Each session defines a regression line with a lower intercept at 5.15 ± 0.42 Ma ($\pm 2\sigma$, MSWD = 0.68, Fig. 4) and 5.54 ± 0.61 Ma ($\pm 2\sigma$, MSWD = 1.02, Fig. 4), respectively.

4.2 Pit depth measurements

After the analyses, pit depths were measured in all the samples as well as in the carbonate reference materials. The measured pit depth averages were used for calculating the U and Pb contents (Tables 3 and 4). The shape and depth of the craters in WC-1 primary carbonate are all similar and their average depth is 15.0 µm (SD = 1.34, n = 16). Few spots corresponding to the secondary reference materials were also checked and they are comparable to those of WC-1. The pits of the SRMNIST 614 are ca. 33% shallower than the ones in the calcite matrix, around 10 µm deep. Regarding the sulfate samples, the pit depth of samples MTO 4-4 and MTO 11-3 is rather homogeneous with mean values of 29.6 µm (SD = 6.2, n = 44) and 18.9 µm (SD = 5.9, n = 37, Fig. 4A), respectively. The samples BCR 9644 and BOX 108 display zones with different heights in some of the ablation holes (Fig. 4B). Although they are exceptional, two ca. 90 µm and two ca. 60 µm pits were measured in BOX 108. Considering them, the average depth is 28.2 µm (SD = 16.4, n=64) whereas excluding those four heights the standard deviation improves substantially (25.0 µm, SD = 8.8, n = 60). The average depth for the sample BCR 9644 is 16.2 µm (SD = 6.7, n=32) excluding two ca. 60 µm spots. On the other hand, the sample PU 05 shows higher variability and larger standard deviation, since the pit depth varies from 29 µm to 107 µm. The calculated average is 62.6 µm (SD = 23.0, n = 48).

5 Discussion

225 5.1. Low success rate

5.1.1. High common Pb content and potential applicability

The majority of the analysed samples, 27 out of 32, were unsuccessful due to the high common Pb content and hence, low or non-existent spread in the $^{238}\text{U}/^{206}\text{Pb}$ axis. Recent studies in the field of environmental hazards have shown that Pb tends to incorporate into sulfates, both gypsum and anhydrite (Astilleros et al., 2010; Morales et al., 2014; Kameda et al., 2017). In fact, in presence of high-Pb fluids, anglesite (PbSO_4) is simultaneously intergrown with those sulfates. The behaviour of uranium remains unknown, although experiments carried out on phosphogypsum, a waste by-product generated from apatite in the production process of phosphoric acid and phosphate fertilizers, suggest that U uptake by gypsum is pH controlled (Lin et al., 2018). Thus, the more alkaline the environment is the higher U concentration could be expected in gypsum. However, the pH of evaporating seawater rarely reaches those values and tends to drop as the evaporation process goes on (Babel and Schreiber, 2014). Considering a low salinity, but high concentrations of Ca^{2+} and SO_4^{2-} (Clauer et al., 2000) during the formation of MSC evaporites, the alkalinity of the depositional environment might have increased. In any case, even the gypsum precipitated in U-rich environments like uranium mine tailings contain a high amount of Pb among other metals (Liu and Hendry, 2011).

The amount of common Pb is a challenge for dating young rocks, as their success strongly depends on the spread in the X-axis ($^{238}\text{U}/^{206}\text{Pb}$). In turn, given the same initial $^{238}\text{U}/^{206}\text{Pb}$ ratio, older samples would have produced sufficient radiogenic Pb, and thus, a certain spread in the Y-axis ($^{207}\text{Pb}/^{206}\text{Pb}$) as to be projected in a more precise regression line. Indeed, older samples are more influenced by the $^{207}\text{Pb}/^{206}\text{Pb}$ ratio and therefore, it is highly likely that the success rate increases with the age of the sample.

5.1.2. SC-ICPMS vs. MC-ICPMS

245 The first set of samples was measured with the SC-ICPMS. The U and Pb contents in the samples were rather low and produce large error ellipses in every single spot. This issue, together with low μ ratios (i.e., spread on $^{238}\text{U}/^{206}\text{Pb}$), produces substantial uncertainties in the final ages (Fig. 3) and a comparison with the depositional ages is poor. In order to achieve better results, we decided to accomplish subsequent measurements with the MC-ICPMS, which provides about three times better sensitivity and simultaneous isotope detection (Craig et al., 2018; 2020). The higher sensitivity implies smaller uncertainties in each spot 250 and hence, more accurate and precise regression lines (i.e., ages) can be depicted.

Indeed, the improvement in age precision is clearly illustrated in Fig. 3. Although the results can be biased because fewer data were acquired during SC-ICPMS analyses, given a similar spread in the $^{238}\text{U}/^{206}\text{Pb}$ axis, the uncertainties of c. 15 % (MTO 11-3) or 20 % (MTO 4-4) obtained with the SC-ICPMS were reduced to 8 % (MTO 11-3, seq 2) and 11 % (MTO 4-4, seq 2) by using the MC-ICPMS (Fig. 3). Furthermore, the re-measurement of these two samples in another independent session in which 255 higher $^{238}\text{U}/^{206}\text{Pb}$ ratios were found, reduced the uncertainties even more down to ca. 6%.




Fig. 4: Pit depth profile of the samples MTO 11-3 (a) and BCR 9644 (b). Whereas the pit shape is roughly homogeneous in the MTO 11-3, the sample BCR 9644 displays deeper areas in some of the pits. The profiles are measured using a Keyence digital microscope VHX-6000.

260 **5.2 U-Pb ages vs cyclostratigraphic ages**

Well-characterized matrix-matched reference material is essential for U-Pb analytical techniques using laser probes as matrix differences between sample and reference standard can cause a significant offset in the obtained ages (Yang et al., 2018; Guillong et al., 2020). However, in the absence of sulfate reference materials, an attempt to use calcite reference materials was carried out, expecting that the offset between both materials was going to be low or negligible. The light absorption observed
265 in calcite and gypsum is similar and they are easily ablated even at low fluence (less than 2 J/cm²). As a comparison, Piccione et al. (2019) obtained analogous ages on contemporary fluorite and nacrite, both corrected to the same calcite reference material, even when the fluorite has different light absorption and higher energy is needed for its ablation (5-6 J/cm²). Due to those reasons, we expected a significantly lower matrix-induced offset than the one observed between calcite and zircon (4.7 %, Parrish et al., 2018).

270 The cyclostratigraphic ages of the MSC samples are well-known (e.g. Vasiliev et al. 2017) and we have used them for testing the suitability of the corrections with respect to carbonate matrix. As pointed out above, the majority of the samples contain a significant amount of common Pb and only five ages were obtained. Although the μ -values of those samples were only moderate, the individual uncertainties range between 6 to 11 % and the ages obtained for the samples MTO 4-4, MTO 11-3, Pu 05 and BOX 108 are in accordance with the cyclostratigraphic ages (e.g. Lugli et al., 2007; Vasiliev et al., 2017). A direct
275 comparison of the U-Pb and cyclostratigraphic ages (Fig. 5), however, points to a slight bias toward younger ages, suggesting a systematic offset between both. Taken singly, each U-Pb date overlaps the cyclostratigraphic age, but a more precise measure is the inverse variance weighted mean of all ten discrepancies between the two ages. The calculated weighted average, i.e. the mean discrepancy, is -0.14 ± 0.14 Ma ($\pm 2\sigma$, MSWD = 0.77). This can now have both an analytical and a geological significance; it can be interpreted as I) matrix mismatch between carbonate and sulfate or II) dating of a subsequent event
280 instead of sedimentation. In fact, the mobilisation of U and Pb during sediment compaction causes some U/Pb heterogeneity, which improves or enables the possibility of dating these sediments by the U-Pb method. The small mean age discrepancy obtained on the sulfate samples is in line with that reported from Montano et al. (2022) on lacustrine carbonates. In this study, although overlapping within uncertainties, a systematic offset was found between U-Pb ages of carbonate cement and that of zircon from ash layers. Thus, U-Pb ages of carbonate and sulfate cement likely dates early diagenesis and not the sediment
285 deposition. This supports our hypothesis that there is no difference in U-Pb fractionation between sulfate and carbonate matrix, although it is not direct evidence.

On the other hand, the sample BCR 9644 resulted in an unexpected younger age of ca. 3 Ma. The brecciated nature of the sample, together with its extremely low Pb content (0.03 µg/g on average) in comparison with surrounding samples suggest a subsequent (re)crystallization and remobilization of U and Pb that could be related to the breccia formation. Warthman et al.
290 (2000) proposed an important bacterial activity after the evaporite formation. For the equivalent in time Site 374, located South-East Sicily, an ~3 m thick dolomitization front in Pliocene hemipelagic succession overlying the UG was identified. Here, a hypothesized role of the deep biosphere, sulfate-reducing bacteria thriving on the dissolution of sulfate-bearing

minerals (WARTHMAN et al., 2000; PETRASH et al., 2017) was suggested. MONTANO et al. (2019, 2021) showed that biological activity may control the U-Pb partitioning on carbonates, so the connection between the bacterial activity and the 3 Ma age could not be discarded. Although gypsum to anhydrite to gypsum (two-step) transformation can be considered as another possible scenario, there is no observation neither in the literature, that supports this hypothesis.




Fig. 5: Comparison of the obtained U-Pb ages and the expected cyclostratigraphic ages. The weighted mean of the offsets between both ages is -0.14 ± 0.14 Ma ($\pm 2\sigma$).

300 5.3 Pit depth profiles

Guillong et al. (2020) showed that different ablation parameters produce distinctive pit profiles (the so-called “aspect ratio” or depth/diameter ratio) and it could result in a noticeable bias in the data. The carbonate reference materials analysed here with a 130 μm spot size, resulted in a depth of ca. 15 μm (aspect ratio of 0.12), whereas the sulfates vary between 16 μm and 63 μm (aspect ratio between 0.12 and 0.48, Fig. 4). The ablation on NIST glass resulted in shallower ca. 10 μm deep holes and an aspect ratio of 0.13, similar to the carbonates. This divergence between the sulfates could be devoted to various non-excluding features such as different textures, particle size, porosity or compaction (Elisha et al., 2021). However, in the cases with an aspect ratio mismatch relative to the primary standard of less than 2, a deviation lower than 5% is anticipated (Guillong et al., 2020), which lies in the final result uncertainty of the majority of the samples analysed here. The larger discrepancy observed in the sample Pu 05 (relative mismatch of 4) could result in age offsets up to 10 % (Guillong et al., 2020, their Figure

310 4). However, Fig. 4b reveals an important heterogeneity in the pit profile in some samples, with a silhouette that resembles pores. Whether they correspond to porosity or chunks released due to a badly coupled laser beam, the signal remained stable. These pit depth issues are also related to the downhole fractionation corrections. Mangenot et al. (2018) claimed that shallow pit depth compared to the spot size could minimize the downhole fractionation. That argument could apply to our reference materials and sulfates with shallower pit depth, but how it affects depths beyond 50-60 μm can be arguable. Lenoir et al.,
315 (2021) obtained coherent regression lines in fluorites even with pit depths (up to 50 % variable) larger than spot sizes. Notwithstanding, the lack of bias between our U-Pb ages and cyclostratigraphic ages suggests that the different downhole fractionation is not noticeable or remains within the uncertainties.

6 Conclusions

In this contribution, we have evaluated the applicability of carbonates as “almost-matrix-matched reference materials” for U-
320 Pb dating of sulfates and for that purpose, gypsum and anhydrite samples from the Messinian Salinity Crisis were analysed. The known cyclostratigraphic ages of these evaporites were compared with the in-situ U-Pb ages obtained. The samples showed a high amount of common Pb and low spread in the U/Pb axis and therefore, only 15 % of the samples were successful. In fact, due to the large uncertainties obtained at the beginning, we were forced to switch from the SC-ICPMS to MC-ICPMS
325 in order to improve the precision of the measurements. Four of the five successfully dated samples were indistinguishable within error from the expected ages, while the other was considerably younger. We assume that all the factors that could produce a bias in the final age, if any, are contained in the uncertainty and therefore, the use of carbonate reference materials could be a trustworthy approach for in situ U-Pb dating of sulfates. We acknowledge that the availability of sulfate reference material in the future will result in an improvement in both reliability and precision.

330

335

340

Table 3: U-Pb data of the LA-SC-ICP-MS measurements.

Analysis number	name / sample	$^{207}\text{Pb}^{\text{a}}$ (cps)	U ^b (ppm)	Pb ^b (ppm)	Th ^b U	$\frac{^{238}\text{U}}{^{206}\text{Pb}}^{\text{c}}$	$\pm 2\sigma$ (%)	$\frac{^{207}\text{Pb}}{^{206}\text{Pb}}^{\text{c}}$	$\pm 2\sigma$ (%)
U436	MTO 4-4	11732	0.075	0.027	0.17	11.66	12	0.8208	3.2
U438		197284	0.26	0.46	0.32	2.043	15	0.8269	0.80
U447		1526928	0.96	3.9	0.18	0.8510	11	0.8307	0.41
U448		28825	0.034	0.066	0.33	1.942	22	0.8350	1.6
U449		25729	0.52	0.058	0.14	34.39	8.4	0.8024	1.6
U450		10535	0.34	0.024	0.11	51.35	7.2	0.8019	1.8
U451		103061	0.97	0.23	0.37	15.87	11	0.8170	0.81
U453		84862	0.67	0.19	0.11	13.91	15	0.8182	0.71
U455		813450	1.1	1.9	0.13	2.276	12	0.8266	0.50
U456		434819	0.82	0.95	0.16	3.529	6.1	0.8333	0.51
U457		334973	0.16	0.82	0.16	0.7034	21	0.8373	0.80
U458		137635	0.92	0.35	0.13	8.870	15	0.8230	0.77
U459		122106	0.41	0.25	0.14	7.284	9.8	0.8248	0.85
U460		66245	1.1	0.16	0.14	24.36	7.2	0.8096	1.2
U461		96692	1.7	0.24	0.13	23.00	7.8	0.8195	0.87
U463		153816	1.3	0.39	0.17	11.87	6.6	0.8224	0.84
U464		533967	1.5	0.96	0.15	9.984	13	0.8292	0.70
U465		198998	1.8	0.45	0.13	15.48	7.4	0.8195	0.70
U466		230504	0.82	0.58	0.22	5.050	7.3	0.8241	0.66
U467		16206	0.006	0.049	1.08	0.4013	25	0.8295	2.1
U468		392950	1.6	0.87	0.10	7.466	9.5	0.8230	0.57
U469		142184	0.56	0.34	0.18	5.895	6.3	0.8285	0.75
U470		105200	1.2	0.25	0.11	17.73	6.2	0.8231	0.76
U471		165896	1.2	0.41	0.14	10.53	13	0.8235	0.82
U472		101404	0.73	0.26	0.36	9.160	12	0.8246	1.2
U473		139917	2.0	0.35	0.11	18.97	6.2	0.8173	0.66
U474		309796	1.7	0.74	0.12	8.422	9.4	0.8193	0.76
U475		190576	2.0	0.48	0.11	14.00	15	0.8172	0.76
U476		489694	1.8	1.2	0.12	5.065	8.3	0.8288	0.51
U477		627160	1.5	1.6	0.16	3.303	19	0.8290	0.53
U479		62648	0.55	0.15	0.22	13.19	7.0	0.8186	0.91
U480		186988	0.59	0.40	0.28	6.266	11	0.8272	0.57
U481		65836	0.25	0.17	0.38	4.834	9.9	0.8266	0.82
U482		211822	1.9	0.53	0.21	12.51	6.0	0.8187	0.70
U576	MTO 11-3	62812	1.2	0.16	0.31	26.79	7.2	0.8138	0.93
U577		71210	1.6	0.18	0.21	29.91	11	0.8051	1.1
U578		27053	1.3	0.063	0.15	82.26	5.5	0.7775	1.2
U579		66175	1.1	0.16	0.33	25.25	5.2	0.8123	0.92
U580		73038	1.3	0.18	0.33	24.35	4.9	0.8092	1.0
U581		116332	0.92	0.31	0.61	9.981	5.6	0.8209	0.71
U582		183789	1.5	0.48	0.22	10.55	7.8	0.8152	0.68
U583		282627	2.0	0.73	0.30	9.601	6.3	0.8217	0.53
U584		216408	3.0	0.51	0.15	23.04	10	0.8181	0.61
U585		53627	0.56	0.12	0.34	20.15	15	0.8144	1.3
U586		93652	0.35	0.25	1.57	4.614	7.8	0.8251	0.71
U587		192319	1.5	0.47	0.24	11.90	14	0.8215	0.74
U588		350075	2.7	0.69	0.23	21.17	16	0.8154	0.70
U589		87555	0.27	0.23	2.51	3.877	6.3	0.8313	0.85
U590		298582	1.7	0.67	0.20	11.20	18	0.8216	0.68
U591		263989	3.5	0.50	0.13	42.55	13	0.7960	0.59
U592		137796	1.2	0.36	0.53	11.01	6.5	0.8182	0.63

^a Within run background-corrected mean ^{207}Pb signal in cps (counts per second).^b U and Pb concentrations and Th/U ratio were calculated relative to the primary reference material.^c Corrected for background, within-run Pb/U fractionation (in case of $^{206}\text{Pb}/^{238}\text{U}$) and

345 **Table 4: U-Pb data of the LA-MC-ICP-MS measurements.**

Analysis number	name / sample	$^{207}\text{Pb}^a$ (cps)	U ^b (ppm)	Pb ^b (ppm)	Th ^b U	$\frac{^{238}\text{U}^c}{^{206}\text{Pb}}$	$\pm 2\sigma$ (%)	$\frac{^{207}\text{Pb}^c}{^{206}\text{Pb}}$	$\pm 2\sigma$ (%)	session
009_U	BOX 108	47348	1.5	0.12	0.072	20.17	5.3	0.8166	0.71	1
010_U		44263	1.1	0.056	0.23	29.83	6.6	0.8160	0.89	1
011_U		8120	0.30	0.019	0.13	24.49	6.7	0.8187	1.4	1
012_U		159615	0.70	0.39	0.22	2.864	15	0.8321	0.56	1
013_U		52758	1.7	0.11	0.15	24.44	7.1	0.8109	0.68	1
015_U		45815	1.2	0.10	0.14	18.66	5.8	0.8205	0.69	1
016_U		210465	0.67	0.50	0.14	2.107	5.8	0.8319	0.50	1
017_U		153200	1.1	0.20	0.17	9.094	11	0.8307	0.56	1
020_U		8660	1.3	0.021	0.088	91.85	5.6	0.7616	1.6	1
020_Ui		54426	1.1	0.11	0.099	15.05	5.7	0.8207	0.68	1
021_U		40138	1.3	0.081	0.11	26.06	13	0.8202	0.75	1
022_U		13276	2.0	0.034	0.056	90.51	4.3	0.7733	1.2	1
024_U		15689	0.50	0.036	0.066	21.67	4.8	0.8219	1.4	1
025_U		24934	0.77	0.061	0.18	20.22	8.6	0.8253	0.90	1
026_U		217333	1.4	0.49	0.14	4.348	7.4	0.8290	0.47	1
027_U		45280	4.3	0.12	0.36	54.80	4.0	0.8014	0.75	1
028_U		106376	4.2	0.26	0.36	25.85	6.8	0.8208	0.63	1
029_U		46138	1.1	0.11	0.52	15.40	9.0	0.8247	0.85	1
030_U		143271	1.3	0.31	0.098	6.515	12	0.8286	0.50	1
031_U		104559	1.0	0.26	0.14	6.336	3.5	0.8281	0.56	1
032_U		66441	0.94	0.15	0.33	9.637	5.2	0.8307	0.78	1
033_U		40961	1.8	0.099	0.12	28.05	4.8	0.8213	0.76	1
034_U		14702	1.1	0.035	0.12	47.15	5.1	0.7992	1.6	1
035_U		28122	1.1	0.067	0.066	24.78	10	0.8136	0.87	1
036_U		42203	0.59	0.10	0.081	9.232	9.2	0.8277	0.71	1
037_U		164255	1.1	0.37	0.14	4.766	15	0.8291	0.53	1
038_U		260017	1.8	0.49	0.098	5.783	11	0.8302	0.47	1
039_U		27954	1.1	0.054	0.24	30.81	5.5	0.8191	0.83	1
040_U		59629	0.96	0.13	0.69	11.90	3.6	0.8280	0.65	1
041_U		20142	1.2	0.047	0.064	39.23	11	0.8030	1.1	1
042_U		172098	0.87	0.46	0.33	2.984	12	0.8315	0.51	1
043_U		26953	0.61	0.040	0.040	24.13	8.6	0.8164	0.98	1
044_U		25995	1.1	0.063	0.15	27.36	5.9	0.8133	1.1	1
045_U		30038	0.94	0.071	0.15	20.81	12	0.8210	1.0	1
046_U		416821	0.98	0.73	0.071	2.138	27	0.8309	0.46	1
047_U		27537	1.6	0.050	0.34	49.55	9.3	0.7956	1.1	1
048_U		34116	1.5	0.083	0.23	28.89	3.3	0.8144	0.76	1
049_U		9837	1.2	0.023	0.10	80.08	3.2	0.7825	1.2	1
050_U		17639	5.7	0.054	0.031	158.0	6.9	0.7188	1.3	1
059_U		60318	1.9	0.15	0.20	20.29	5.9	0.8131	0.82	1
060_U		14510	1.1	0.030	0.10	57.79	5.7	0.7979	1.1	1
061_U		99693	0.90	0.23	0.098	6.175	11	0.8280	0.64	1
062_U		106403	1.4	0.23	0.078	9.364	6.6	0.8275	0.54	1
063_U		33943	1.4	0.049	0.19	43.08	8.4	0.7970	1.1	1
067_U		9178	0.25	0.019	0.19	20.53	4.6	0.8202	1.4	1
068_U		60040	3.4	0.14	0.063	37.77	7.1	0.8103	0.71	1
069_U		42920	3.3	0.11	0.060	47.99	6.5	0.8031	0.78	1
070_U		71312	2.1	0.18	0.069	18.77	8.0	0.8194	0.73	1
071_U		16427	1.6	0.040	0.036	60.56	6.5	0.7900	1.4	1
009_U	BCR 9644	1031	0.061	0.001	0.012	74.49	7.0	0.8324	3.9	2
010_U		87	0.006	0.000	0.0056	68.34	11	0.8105	13	2
011_U		87	0.014	0.000	-0.00112	156.2	10	0.7736	14	2
014_U		110	0.020	0.000	0.0083	243.5	9.5	0.7783	13	2
015_U		343	0.016	0.000	0.0030	108.7	9.3	0.8263	8.4	2
017_U		337	0.036	0.000	-0.00574	130.1	9.7	0.8259	7.1	2
018_U		125	0.025	0.000	-0.00418	220.6	9.0	0.7825	11	2
019_U		275	0.019	0.000	-0.02679	201.2	9.5	0.7862	10	2
021_U		109	0.025	0.000	0.0081	217.8	8.9	0.7236	11	2
022_U		470	0.029	0.000	0.0089	240.5	9.8	0.7104	8.7	2
023_U		108	0.019	0.000	0.0084	206.7	9.9	0.7573	12	2
024_U		141	0.030	0.000	0.0024	246.3	8.1	0.8004	11	2
025_U		720	0.045	0.001	-0.00528	119.5	6.9	0.8306	5.4	2
026_U		347	0.038	0.001	0.046	117.3	8.2	0.7971	6.9	2

Table 4: continue...

027_U	1796	0.57	0.002	0.0067	488.8	19	0.6571	4.2	2	
028_U	2103	0.020	0.003	0.032	9.585	9.5	0.8217	2.9	2	
029_U	2505	0.62	0.002	0.0018	577.5	21	0.6308	5.7	2	
031_U	285	0.030	0.000	0.010	109.3	6.5	0.7944	7.0	2	
032_U	1114	0.19	0.001	0.0056	247.8	6.6	0.7212	4.3	2	
033_U	148	0.059	0.000	0.0095	360.8	8.7	0.6898	11	2	
034_U	180	0.017	0.000	0.014	106.2	9.4	0.7711	9.8	2	
035_U	73633	0.30	0.025	0.0011	19.18	36	0.8220	0.93	2	
036_U	200	0.006	0.000	0.043	30.98	11	0.8088	8.4	2	
037_U	925	0.10	0.002	0.00099	73.86	21	0.8161	5.5	2	
039_U	406	0.068	0.001	0.0050	164.1	8.8	0.7734	6.3	2	
350_U	9048	0.48	0.016	0.019	46.44	7.7	0.8196	1.5	2	
359_U	50690	0.64	0.079	0.057	12.70	11	0.8247	0.66	2	
360_U	5648	0.62	0.009	0.0088	109.5	5.9	0.7752	1.9	2	
361_U	1218	0.11	0.002	0.0097	82.26	8.4	0.7972	4.1	2	
365_U	49349	0.54	0.061	0.039	14.14	9.0	0.8213	0.72	2	
366_U	4943	0.040	0.007	0.018	8.691	21	0.8239	2.5	2	
368_U	1226	0.021	0.002	0.058	17.45	8.1	0.8227	3.5	2	
369_U	65958	0.25	0.12	0.11	3.320	13	0.8291	0.62	2	
374_U	305	0.032	0.001	-0.00810	92.10	7.1	0.7948	7.8	2	
375_U	2308	0.087	0.003	0.014	52.37	12	0.8094	4.9	2	
376_U	27934	0.78	0.032	0.020	38.35	11	0.8195	0.89	2	
377_U	17604	1.0	0.035	0.0051	44.70	10	0.8127	1.0	2	
378_U	58189	0.63	0.082	0.012	12.09	15	0.8195	0.62	2	
382_U	9122	0.93	0.015	0.0072	96.07	5.6	0.7875	1.5	2	
384_U	882	0.019	0.001	0.099	21.28	18	0.8240	4.5	2	
387_U	6049	0.32	0.008	0.010	61.42	13	0.8097	1.8	2	
388_U	6781	0.52	0.010	0.0076	79.65	11	0.8000	1.6	2	
389_U	6780	1.2	0.010	0.0023	182.9	11	0.7553	2.3	2	
390_U	17237	0.90	0.028	0.015	50.27	6.2	0.8145	1.1	2	
391_U	1518	0.11	0.002	0.0077	74.06	10	0.8176	3.3	2	
392_U	6719	0.40	0.008	0.0075	82.20	4.7	0.7937	1.7	2	
160_U	MTO 4-4	42459	0.071	0.033	0.12	3.455	16	0.8322	0.76	2
161_U		36404	0.100	0.024	0.086	6.599	10	0.8274	0.80	2
162_U		29300	0.31	0.025	0.12	19.64	15	0.8157	0.81	2
164_U		107586	0.65	0.064	0.11	16.10	12	0.8229	0.62	2
166_U		62776	1.1	0.055	0.13	32.05	4.3	0.8194	1.1	2
167_U		111666	0.47	0.12	0.16	6.312	15	0.8264	0.55	2
168_U		86330	0.82	0.076	0.12	17.08	7.0	0.8159	0.59	2
169_U		361703	0.88	0.24	0.10	5.739	15	0.8268	0.43	2
170_U		635386	0.50	0.56	0.14	1.426	11	0.8318	0.49	2
171_U		124867	0.27	0.083	0.11	5.252	16	0.8296	0.55	2
174_U		39215	0.67	0.032	0.075	33.18	9.4	0.8066	0.78	2
177_U		51601	0.29	0.049	0.11	9.368	8.3	0.8236	0.72	2
178_U		109476	1.3	0.079	0.093	26.16	11	0.8123	0.60	2
179_U		239957	0.80	0.19	0.12	6.474	9.5	0.8249	0.47	2
180_U		296212	0.93	0.23	0.13	6.498	15	0.8244	0.45	2
181_U		361123	0.98	0.35	0.082	4.497	13	0.8251	0.43	2
182_U		178465	0.91	0.12	0.14	12.41	5.1	0.8252	0.51	2
183_U		39048	0.49	0.037	0.067	21.37	8.1	0.8182	0.96	2
184_U		1048150	0.65	0.94	0.046	1.089	3.9	0.8279	0.39	2
185_U		102247	0.84	0.090	0.10	14.76	8.1	0.8206	0.61	2
186_U		121692	1.0	0.11	0.10	14.40	5.0	0.8202	0.50	2
187_U		198711	0.62	0.14	2.61	6.826	7.4	0.8273	0.47	2
188_U		291589	1.2	0.27	0.11	6.743	6.3	0.8238	0.43	2
189_U		231299	1.1	0.18	0.14	9.687	8.5	0.8269	0.45	2
190_U		413220	0.81	0.35	0.12	3.665	5.6	0.8296	0.42	2
191_U		11917	0.074	0.008	0.14	13.84	9.4	0.8215	1.4	2
192_U		126152	0.96	0.12	0.092	12.84	11	0.8217	0.62	2
193_U		73154	0.75	0.062	0.29	19.21	10	0.8187	0.86	2
194_U		549529	0.61	0.49	0.13	1.982	12	0.8351	0.45	2
197_U		51681	0.30	0.046	0.32	10.41	9.6	0.8230	0.72	2
198_U		273244	0.68	0.23	0.20	4.658	9.0	0.8330	0.45	2
209_U		477618	0.84	0.36	0.12	3.646	17	0.8277	0.41	2
210_U		412526	1.0	0.30	0.10	5.471	9.2	0.8287	0.43	2

Table 4: continue...

211_U	293172	1.0	0.18	0.13	8.843	11	0.8263	0.45	2	
212_U	555903	1.1	0.38	0.088	4.670	16	0.8287	0.41	2	
213_U	76097	0.92	0.047	0.084	31.17	8.0	0.8095	0.68	2	
214_U	346163	1.1	0.31	0.23	5.896	6.1	0.8265	0.43	2	
215_U	120901	0.90	0.10	0.11	14.03	14	0.8228	0.53	2	
217_U	262806	1.3	0.35	0.13	5.921	8.3	0.8314	0.44	2	
219_U	127957	1.0	0.10	0.13	16.17	13	0.8172	0.58	2	
220_U	7919	0.11	0.004	0.16	42.58	11	0.8090	1.8	2	
221_U	148260	1.6	0.13	0.24	19.25	5.8	0.8175	0.48	2	
222_U	128181	1.3	0.094	0.089	20.96	13	0.8179	0.55	2	
223_U	307635	1.5	0.26	0.14	9.247	4.3	0.8249	0.44	2	
225_U	254729	1.6	0.23	0.11	11.19	8.6	0.8251	0.48	2	
226_U	355548	1.3	0.24	0.11	8.578	9.1	0.8241	0.43	2	
227_U	156455	1.1	0.14	0.098	12.89	7.5	0.8222	0.50	2	
228_U	681461	0.67	0.64	0.095	1.651	10	0.8295	0.45	2	
229_U	95834	1.0	0.084	0.10	19.44	7.7	0.8180	0.57	2	
231_U	157487	1.3	0.14	0.081	13.86	12	0.8220	0.50	2	
232_U	3039	0.066	0.003	0.16	33.96	7.1	0.8091	3.0	2	
233_U	118467	2.0	0.10	0.060	30.51	7.1	0.8138	0.51	2	
234_U	133809	1.6	0.11	0.12	23.19	8.0	0.8166	0.52	2	
235_U	168534	1.8	0.15	0.10	19.46	6.0	0.8155	0.52	2	
236_U	220800	1.8	0.19	0.18	14.73	11	0.8218	0.47	2	
237_U	246643	1.3	0.21	0.084	9.590	7.9	0.8252	0.46	2	
238_U	379877	0.62	0.29	0.089	3.429	32	0.8280	0.44	2	
239_U	217983	0.78	0.19	0.11	6.434	14	0.8287	0.49	2	
240_U	412692	0.73	0.38	0.066	3.048	12	0.8311	0.45	2	
241_U	48316	0.51	0.040	0.15	20.14	7.7	0.8204	0.64	2	
244_U	MTO 11-3	31010	0.020	0.020	2.18	1.605	14	0.8367	1.0	2
245_U	24767	0.047	0.030	0.53	2.522	22	0.8298	0.98	2	
247_U	63055	0.28	0.079	0.38	5.659	7.2	0.8300	0.68	2	
248_U	99922	0.26	0.13	0.53	3.038	5.8	0.8329	0.55	2	
249_U	16054	0.064	0.024	0.59	4.306	7.1	0.8338	1.1	2	
250_U	95971	0.20	0.094	0.47	3.360	5.4	0.8268	0.57	2	
259_U	20407	0.83	0.026	0.11	50.50	12	0.7972	1.1	2	
260_U	1786	0.011	0.003	0.74	6.092	8.0	0.8376	3.1	2	
265_U	124037	0.14	0.10	0.95	2.202	8.5	0.8317	0.55	2	
266_U	19405	0.088	0.021	0.47	6.703	14	0.8330	1.0	2	
268_U	145975	1.7	0.21	0.21	12.39	7.1	0.8255	0.58	2	
269_U	136367	0.12	0.19	1.64	1.023	5.3	0.8312	0.51	2	
271_U	12070	0.034	0.018	1.15	2.977	14	0.8323	1.2	2	
272_U	55571	0.22	0.075	0.45	4.556	5.9	0.8267	0.65	2	
274_U	9126	0.059	0.012	0.55	7.695	11	0.8312	2.2	2	
275_U	11007	0.054	0.017	1.23	5.116	6.3	0.8262	1.2	2	
276_U	58251	0.092	0.080	0.54	1.841	16	0.8373	0.67	2	
277_U	112321	0.29	0.13	1.17	3.413	8.0	0.8274	0.55	2	
278_U	18070	0.048	0.028	0.75	2.698	7.3	0.8292	0.93	2	
279_U	105781	0.37	0.15	0.69	3.774	6.9	0.8304	0.54	2	
281_U	153346	4.0	0.22	0.081	28.52	7.9	0.8107	0.51	2	
282_U	141144	3.6	0.20	0.069	27.90	8.6	0.8121	0.52	2	
283_U	70729	0.71	0.11	0.15	9.890	4.2	0.8229	0.59	2	
284_U	57878	0.28	0.065	0.32	6.812	14	0.8344	0.67	2	
285_U	179536	0.73	0.19	0.34	6.075	4.7	0.8349	0.48	2	
286_U	201969	1.3	0.32	0.31	6.386	4.9	0.8261	0.44	2	
287_U	374142	2.5	0.56	0.055	7.103	7.0	0.8247	0.42	2	
288_U	128171	3.2	0.18	0.087	28.71	5.0	0.8112	0.58	2	
289_U	136966	4.1	0.18	0.082	35.20	5.3	0.8090	0.53	2	
290_U	107867	4.9	0.16	0.060	48.17	7.7	0.8005	0.57	2	
291_U	106011	1.9	0.17	0.43	17.90	18	0.8193	0.65	2	
292_U	61587	4.5	0.11	0.075	63.75	15	0.7965	1.1	2	
293_U	91650	3.8	0.14	0.043	42.71	5.7	0.8084	0.58	2	
294_U	80760	3.8	0.12	0.075	47.86	5.8	0.8000	0.66	2	
295_U	67617	4.3	0.11	0.043	62.75	5.6	0.7921	0.67	2	
436_U	Pu 05	1336	0.008	0.001	0.018	17.15	12	0.7968	3.4	2
440_U	3541	0.013	0.002	0.047	9.396	12	0.7993	3.3	2	
441_U	5779	0.013	0.001	0.089	18.43	9.4	0.7958	2.4	2	

Table 4: continue...

442_U	819	0.010	0.000	0.0016	36.21	12	0.7929	4.3	2	
444_U	3503	0.002	0.001	0.058	5.613	15	0.7931	2.7	2	
445_U	1398	0.006	0.001	0.043	12.30	8.4	0.7884	3.4	2	
463_U	2058	0.019	0.001	0.043	19.92	10	0.7828	3.4	2	
464_U	2093	0.019	0.001	0.059	23.30	7.5	0.7863	2.9	2	
465_U	1188	0.007	0.001	0.12	16.62	13	0.7885	4.6	2	
466_U	7720	0.049	0.005	0.043	14.59	4.8	0.7909	1.6	2	
467_U	2652	0.032	0.002	0.032	26.36	15	0.7822	3.6	2	
468_U	51097	0.31	0.030	0.045	15.94	4.4	0.7918	0.65	2	
469_U	39483	0.30	0.024	0.13	19.22	4.7	0.7889	0.77	2	
470_U	18990	0.23	0.011	0.031	33.65	6.1	0.7832	0.96	2	
472_U	77384	0.56	0.046	0.022	18.78	5.0	0.7925	0.58	2	
473_U	42753	1.2	0.029	0.014	62.46	7.5	0.7622	0.78	2	
474_U	38267	0.51	0.026	0.017	29.85	12	0.7871	0.98	2	
475_U	6462	0.25	0.004	0.0099	92.20	9.6	0.7538	2.3	2	
476_U	55482	0.93	0.039	0.092	36.70	9.8	0.7884	0.89	2	
477_U	24674	0.11	0.016	0.047	11.19	4.7	0.7968	1.0	2	
478_U	2847	0.075	0.002	0.0068	74.37	12	0.7544	2.4	2	
479_U	70165	0.52	0.045	0.079	17.80	6.7	0.7906	0.60	2	
480_U	184870	0.88	0.12	0.069	11.87	3.8	0.7968	0.48	2	
481_U	11662	0.62	0.008	0.0093	115.9	7.6	0.7352	1.2	2	
483_U	1829	0.014	0.001	0.030	15.84	11	0.7971	3.2	2	
486_U	652	0.003	0.001	0.032	8.749	22	0.7941	5.5	2	
487_U	97239	1.4	0.060	0.019	36.73	4.6	0.7742	0.57	2	
488_U	12860	0.70	0.009	0.0061	122.2	8.9	0.7343	1.4	2	
489_U	22723	0.84	0.014	0.011	89.94	4.9	0.7468	0.93	2	
490_U	39877	0.51	0.026	0.013	30.42	6.1	0.7844	0.81	2	
491_U	30830	1.0	0.014	0.016	114.4	16	0.7368	0.91	2	
492_U	19082	1.0	0.014	0.013	106.9	10	0.7335	1.1	2	
009_U	MTO 11-3	90036	2.9	0.29	0.14	31.54	10	0.8054	0.65	3
010_U		60711	4.8	0.19	0.058	80.09	4.6	0.7784	0.73	3
011_U		321269	3.1	0.97	0.36	10.25	6.6	0.8207	0.40	3
012_U		18890	1.8	0.061	0.081	92.39	6.5	0.7715	1.1	3
013_U		104308	2.0	0.32	0.12	20.01	8.7	0.8134	0.49	3
014_U		49427	2.4	0.13	0.076	72.22	11	0.7788	0.99	3
015_U		140824	4.3	0.43	0.14	32.07	4.3	0.8054	0.50	3
017_U		107005	3.6	0.34	0.083	34.10	5.5	0.8103	0.54	3
018_U		63650	0.60	0.19	0.30	10.74	7.3	0.8207	0.60	3
024_U		112992	0.55	0.35	0.92	5.077	9.5	0.8279	0.48	3
025_U		148314	0.57	0.45	0.77	4.294	9.0	0.8277	0.44	3
028_U		148368	3.1	0.40	0.11	29.45	11	0.8088	0.60	3
031_U		54347	2.1	0.17	0.075	42.55	8.5	0.7968	0.76	3
033_U		158602	3.3	0.50	0.23	21.72	4.9	0.8142	0.45	3
034_U		47147	0.31	0.14	0.38	7.791	5.9	0.8239	0.66	3
035_U		161917	3.2	0.50	0.34	20.44	5.5	0.8167	0.42	3
037_U		245790	3.2	0.68	0.23	17.08	6.7	0.8149	0.47	3
038_U		152560	4.6	0.48	0.14	30.05	6.2	0.8048	0.54	3
040_U		45567	2.2	0.14	0.11	50.70	5.5	0.7961	0.68	3
041_U		43416	0.35	0.14	0.43	8.464	8.0	0.8208	0.94	3
042_U		41614	0.27	0.13	0.39	7.039	7.3	0.8276	0.77	3
043_U		51784	0.32	0.16	0.43	6.705	5.5	0.8184	0.67	3
044_U		185745	4.0	0.53	0.19	26.15	8.7	0.8085	0.46	3
045_U		197622	1.6	0.58	0.12	9.503	18	0.8224	0.45	3
046_U		163159	3.2	0.52	0.15	19.59	7.1	0.8129	0.52	3
047_U		137303	3.2	0.45	0.17	21.78	6.0	0.8140	0.54	3
048_U		74586	3.5	0.24	0.098	47.14	5.7	0.7992	0.63	3
049_U		142496	4.4	0.45	0.14	31.29	4.5	0.8067	0.43	3
050_U		148349	4.8	0.45	0.14	36.35	6.6	0.8066	0.51	3
059_U		102822	3.0	0.32	0.18	30.04	6.2	0.8036	0.51	3
061_U		125428	3.2	0.40	0.16	25.59	6.9	0.8108	0.47	3
062_U		37071	1.1	0.13	0.13	23.38	9.8	0.8098	0.76	3
063_U		48032	0.78	0.16	0.12	15.90	6.3	0.8180	0.71	3
064_U		69242	0.51	0.21	0.58	8.442	13	0.8203	0.63	3
065_U		165434	4.1	0.51	0.060	26.34	9.2	0.8096	0.53	3
066_U		134214	5.5	0.42	0.085	41.66	9.2	0.7982	0.55	3

Table 4: continue...

067_U	35032	3.5	0.12	0.051	91.67	4.7	0.7642	0.82	3	
068_U	197102	2.8	0.65	0.17	13.21	5.2	0.8204	0.40	3	
069_U	152700	2.9	0.46	0.12	21.30	17	0.8147	0.57	3	
071_U	108748	0.34	0.35	1.35	3.081	9.5	0.8266	0.52	3	
072_U	170691	0.85	0.55	0.59	4.946	11	0.8292	0.52	3	
073_U	9935	1.7	0.032	0.068	155.2	7.0	0.7228	1.5	3	
074_U	92588	5.5	0.30	0.059	58.10	9.9	0.7929	0.77	3	
009_U	Pu 05	2167	0.030	0.002	0.0045	64.40	17	0.7620	3.1	4
011_U	642	0.003	0.001	0.018	12.15	16	0.8304	6.2	4	
013_U	2053	0.002	0.002	0.062	4.612	6.6	0.8177	2.8	4	
014_U	550	0.011	0.000	0.0052	75.72	18	0.7705	7.0	4	
015_U	1351	0.027	0.001	0.00067	67.07	8.5	0.7705	3.5	4	
016_U	7081	0.10	0.006	0.015	56.46	7.6	0.7632	1.6	4	
017_U	2027	0.064	0.002	0.0088	132.8	9.8	0.7411	2.8	4	
018_U	1355	0.051	0.001	0.0042	127.1	15	0.7475	3.5	4	
019_U	9372	0.10	0.007	0.018	63.09	8.6	0.7824	2.2	4	
020_U	2028	0.015	0.002	0.013	27.00	6.9	0.7825	3.0	4	
021_U	3008	0.011	0.002	0.0037	18.46	5.1	0.8120	2.8	4	
022_U	1140	0.005	0.001	0.0098	13.19	19	0.7949	4.4	4	
023_U	1051	0.033	0.001	0.0029	94.71	16	0.7526	4.8	4	
025_U	4316	0.059	0.004	0.0071	51.96	14	0.7794	1.9	4	
026_U	7481	0.050	0.007	0.022	23.39	9.4	0.7891	1.6	4	
027_U	6387	0.028	0.006	0.010	12.93	18	0.7951	1.7	4	
028_U	12431	0.023	0.011	0.0018	6.389	14	0.8304	1.3	4	
029_U	1118	0.023	0.001	0.0070	73.49	9.6	0.7553	4.8	4	
030_U	1099	0.004	0.001	0.00034	6.039	35	0.8159	3.8	4	
031_U	2456	0.003	0.002	0.029	4.920	11	0.8120	2.5	4	
032_U	8610	0.057	0.007	0.042	29.39	9.7	0.7989	1.8	4	
033_U	2450	0.006	0.002	0.086	8.355	7.5	0.8029	3.9	4	
034_U	1943	0.010	0.001	0.029	22.89	6.7	0.8047	3.3	4	
037_U	1471	0.010	0.001	0.014	23.40	12	0.8005	4.0	4	
038_U	4241	0.057	0.003	0.0077	79.29	9.5	0.7796	2.4	4	
040_U	5569	0.039	0.005	0.049	25.95	6.2	0.8039	1.8	4	
041_U	2789	0.008	0.002	0.015	10.48	20	0.8099	2.3	4	
042_U	2272	0.022	0.002	0.0071	35.64	20	0.8083	2.9	4	
043_U	845	0.006	0.001	0.017	16.92	20	0.8020	4.5	4	
044_U	1053	0.002	0.001	0.046	8.587	15	0.8135	3.7	4	
045_U	2058	0.004	0.002	0.023	6.070	8.1	0.8143	2.8	4	
046_U	1004	0.003	0.001	0.018	14.56	9.9	0.7973	3.8	4	
047_U	1329	0.016	0.001	0.00077	33.81	31	0.8012	3.7	4	
048_U	1422	0.021	0.001	0.0060	45.24	15	0.7810	3.1	4	
049_U	276573	1.2	0.16	0.049	37.36	12	0.7884	0.56	4	
060_U	1411	0.006	0.001	0.0053	15.00	12	0.8117	3.0	4	
061_U	7356	0.090	0.005	0.084	81.01	27	0.7538	2.1	4	
062_U	1435	0.043	0.001	0.0064	117.0	14	0.7277	3.2	4	
064_U	633	0.056	0.001	0.0010	219.2	23	0.6783	5.8	4	
067_U	108576	0.66	0.094	0.019	21.24	4.2	0.8007	0.55	4	
068_U	22860	0.23	0.019	0.017	37.25	4.5	0.7906	0.91	4	
073_U	2091	0.14	0.002	0.0021	199.1	9.7	0.6805	3.3	4	
075_U	748	0.007	0.001	0.0033	29.09	18	0.8043	4.8	4	
078_U	1316	0.061	0.001	0.0023	150.7	9.0	0.7338	3.7	4	
079_U	2405	0.044	0.002	0.0056	60.40	15	0.7854	2.5	4	
080_U	2285	0.032	0.002	0.0047	46.32	13	0.7933	2.8	4	
081_U	2198	0.040	0.002	0.011	78.19	6.2	0.7541	2.5	4	
088_U	MTO 4-4	154270	1.5	0.23	0.24	22.94	8.6	0.8128	0.51	4
089_U	66593	1.0	0.11	0.075	31.35	8.3	0.8050	0.64	4	
092_U	25818	0.94	0.043	0.14	72.14	9.5	0.7867	1.3	4	
093_U	188855	1.5	0.29	0.15	18.28	12	0.8136	0.45	4	
094_U	70717	1.5	0.12	0.099	42.66	3.3	0.7971	0.68	4	
095_U	392652	1.3	0.68	0.14	5.799	17	0.8244	0.40	4	
096_U	39333	0.54	0.065	0.14	27.02	9.7	0.8107	0.72	4	
098_U	343014	0.64	0.58	0.33	3.486	9.0	0.8302	0.43	4	
099_U	15397	0.083	0.027	0.36	9.619	12	0.8195	1.2	4	
100_U	35522	0.27	0.060	0.24	14.65	4.5	0.8160	0.78	4	
109_U	202639	1.6	0.34	0.081	14.63	19	0.8141	0.57	4	

355 **Table 4: continue...**

110_U	42283	0.70	0.069	0.083	33.87	5.6	0.8050	0.75	4
111_U	112583	1.5	0.19	0.099	25.52	9.2	0.8088	0.50	4
112_U	120520	0.70	0.21	0.043	10.44	16	0.8226	0.54	4
113_U	4129	0.17	0.007	0.067	75.44	9.7	0.7836	2.3	4
114_U	64831	0.23	0.12	0.051	6.233	6.3	0.8270	0.63	4
115_U	386587	2.3	0.63	0.061	11.80	8.6	0.8204	0.36	4
116_U	101833	0.059	0.18	0.59	1.008	20	0.8328	0.51	4
117_U	36664	0.023	0.066	0.93	1.063	21	0.8262	0.80	4
118_U	263914	1.5	0.45	0.11	10.72	10	0.8204	0.42	4
119_U	198490	1.0	0.33	0.16	10.32	10	0.8199	0.46	4
120_U	321214	1.3	0.54	0.099	7.662	5.1	0.8241	0.38	4
121_U	77903	0.53	0.13	0.046	12.56	6.3	0.8208	0.64	4
122_U	477392	1.1	0.81	0.13	4.040	11	0.8258	0.43	4
123_U	62459	0.84	0.11	0.089	25.60	12	0.8143	0.66	4
124_U	83446	0.45	0.14	0.13	9.875	14	0.8151	0.56	4
125_U	328913	0.86	0.53	0.11	5.216	11	0.8267	0.38	4
126_U	51257	1.2	0.080	0.082	53.19	6.5	0.7948	0.88	4
127_U	165366	1.2	0.28	0.094	13.05	6.3	0.8199	0.44	4
128_U	128085	0.78	0.24	0.14	9.825	15	0.8209	0.48	4
129_U	40122	0.48	0.068	0.076	22.54	8.6	0.8123	0.72	4
130_U	243432	1.6	0.40	0.11	12.91	8.9	0.8182	0.41	4
131_U	410018	1.0	0.63	0.098	5.566	9.6	0.8252	0.38	4
132_U	453513	1.3	0.75	0.11	5.295	13	0.8225	0.39	4
133_U	140762	1.0	0.23	0.10	14.54	10	0.8205	0.45	4
134_U	171451	1.8	0.25	0.11	27.07	5.9	0.8110	0.49	4
135_U	299485	1.3	0.50	0.088	8.646	7.3	0.8253	0.40	4
136_U	273271	0.95	0.46	0.13	6.742	7.5	0.8255	0.38	4
137_U	261422	1.00	0.44	0.17	7.251	8.8	0.8219	0.40	4
139_U	336303	1.2	0.56	0.20	6.730	14	0.8277	0.38	4
140_U	91664	0.34	0.15	0.11	7.230	15	0.8242	0.51	4
141_U	2833	0.13	0.005	0.16	98.39	7.1	0.7727	3.4	4
143_U	178759	0.87	0.30	0.089	9.142	10	0.8233	0.44	4
144_U	175078	1.1	0.30	0.10	11.67	18	0.8186	0.50	4
145_U	190119	1.2	0.30	0.11	13.46	8.7	0.8178	0.46	4
146_U	85820	1.4	0.15	0.22	30.20	3.5	0.8081	0.62	4
147_U	329223	0.75	0.55	0.14	4.281	9.5	0.8254	0.40	4
148_U	114269	1.1	0.20	0.094	18.16	8.5	0.8142	0.53	4
149_U	248241	1.1	0.42	0.10	8.312	8.3	0.8215	0.41	4
150_U	181792	0.92	0.28	0.100	11.66	10.0	0.8192	0.47	4
159_U	122725	0.83	0.21	0.17	12.59	5.4	0.8211	0.48	4
160_U	126548	0.90	0.22	0.10	12.68	8.3	0.8214	0.50	4
161_U	110684	1.9	0.17	0.086	39.06	4.6	0.8017	0.58	4
162_U	78280	1.2	0.13	0.092	30.26	7.1	0.8049	0.71	4
165_U	195480	1.0	0.34	0.092	9.686	9.0	0.8209	0.44	4
166_U	303135	1.3	0.49	0.11	8.744	7.7	0.8241	0.40	4
169_U	62163	0.87	0.11	0.079	22.90	9.7	0.8160	0.62	4
170_U	124538	1.1	0.21	0.093	17.94	7.9	0.8174	0.45	4
171_U	156199	1.3	0.27	0.089	15.78	13	0.8168	0.46	4
172_U	58090	0.92	0.094	0.090	33.53	7.6	0.8049	0.63	4
173_U	88433	1.5	0.14	0.10	40.49	7.9	0.7982	0.53	4
175_U	70717	1.1	0.11	0.100	34.96	6.1	0.8048	0.71	4
176_U	115260	1.2	0.15	0.099	39.42	5.0	0.8018	0.75	4
177_U	35514	0.82	0.058	0.079	47.42	5.3	0.7983	0.80	4
178_U	60006	1.5	0.090	0.095	59.06	8.5	0.7877	0.77	4
179_U	263568	1.7	0.45	0.091	11.81	9.8	0.8172	0.51	4
180_U	237512	1.2	0.38	0.089	10.36	26	0.8202	0.53	4
181_U	79587	1.0	0.14	0.12	24.39	5.7	0.8099	0.56	4
182_U	181676	1.5	0.31	0.16	15.23	8.6	0.8179	0.49	4
183_U	148496	0.62	0.39	0.15	3.828	6.4	0.8256	0.42	4
184_U	338583	1.7	0.59	0.067	9.117	8.8	0.8256	0.39	4
185_U	79508	1.1	0.13	0.065	26.54	4.1	0.8143	0.56	4
186_U	85236	0.73	0.15	0.089	15.83	9.9	0.8170	0.54	4
187_U	307851	1.0	0.53	0.26	5.904	13	0.8191	0.43	4
188_U	116771	1.1	0.21	0.087	16.25	11	0.8141	0.53	4
280_U	BOX 108	3723	0.027	0.015	4.210	34	0.8405	2.5	4

Table 4: continue...

283_U	215245	2.5	0.52	0.12	19.76	7.4	0.8240	0.47	4
284_U	95417	1.1	0.28	0.14	13.04	4.9	0.8259	0.51	4
287_U	9849	1.3	0.030	0.033	133.6	4.0	0.7471	1.5	4
288_U	11288	1.5	0.035	0.042	124.2	4.9	0.7493	1.6	4
289_U	34194	0.096	0.099	0.22	3.185	13	0.8344	0.81	4
290_U	242588	2.0	0.72	0.18	8.867	7.6	0.8307	0.42	4
291_U	142226	1.3	0.40	0.11	10.56	12	0.8321	0.45	4
292_U	205563	1.8	0.52	0.11	13.32	8.8	0.8231	0.44	4
293_U	174825	1.0	0.41	0.054	11.00	18	0.8305	0.45	4
294_U	155002	1.8	0.44	0.15	14.07	3.5	0.8248	0.46	4
296_U	135623	1.4	0.34	0.25	16.69	9.9	0.8252	0.47	4
297_U	431598	1.3	1.1	0.34	4.623	9.7	0.8293	0.37	4
298_U	73380	1.8	0.22	0.027	26.06	9.7	0.8186	0.58	4
299_U	285598	2.7	0.76	0.012	12.45	7.6	0.8291	0.43	4
300_U	177111	1.4	0.51	0.14	8.972	13	0.8245	0.46	4
309_U	20078	0.082	0.061	0.23	4.368	34	0.8415	1.4	4
311_U	162504	1.2	0.49	0.14	7.592	13	0.8274	0.50	4
312_U	300617	2.1	0.89	0.22	7.673	8.5	0.8305	0.42	4
313_U	4524	0.035	0.013	0.061	8.581	14	0.8286	2.2	4
314_U	149794	0.98	0.42	0.16	8.132	5.8	0.8278	0.46	4
315_U	151082	2.8	0.45	0.15	20.25	3.9	0.8208	0.45	4
316_U	403090	5.0	1.2	0.066	13.02	7.7	0.8299	0.41	4
317_U	66022	1.6	0.20	0.15	25.40	4.4	0.8132	0.62	4
318_U	69901	1.3	0.21	0.10	21.32	15	0.8190	0.62	4
319_U	43921	2.7	0.13	0.062	68.54	9.7	0.7921	0.79	4
320_U	95589	0.99	0.30	0.26	10.46	3.5	0.8274	0.49	4
321_U	86168	0.84	0.26	0.21	10.38	11	0.8324	0.54	4
322_U	45837	0.39	0.13	0.079	11.04	8.6	0.8248	0.70	4
323_U	81816	1.1	0.25	0.14	14.89	3.8	0.8234	0.61	4
324_U	291563	2.1	0.88	0.11	7.571	16	0.8275	0.44	4
325_U	54488	2.5	0.17	0.066	46.74	9.2	0.8017	0.66	4
326_U	166955	1.6	0.51	0.035	10.17	5.0	0.8268	0.48	4
327_U	378811	0.73	1.1	0.18	2.132	9.8	0.8380	0.40	4
328_U	125856	2.6	0.40	0.035	19.49	20	0.8257	0.64	4
330_U	157915	2.0	0.45	0.074	15.73	9.3	0.8244	0.48	4
331_U	245854	1.8	0.66	0.29	9.850	7.9	0.8210	0.58	4
332_U	47905	0.41	0.12	0.15	12.59	9.9	0.8276	0.76	4
333_U	566996	4.6	1.7	0.12	8.640	9.0	0.8295	0.38	4
334_U	84562	1.2	0.25	0.096	16.17	3.2	0.8216	0.55	4
335_U	66581	1.8	0.19	0.12	31.10	8.8	0.8126	0.64	4
336_U	192058	3.3	0.54	0.056	21.03	5.8	0.8238	0.44	4
337_U	82708	1.7	0.25	0.14	21.94	5.4	0.8239	0.50	4
338_U	99827	2.1	0.30	0.19	22.02	6.9	0.8194	0.51	4
339_U	49540	1.6	0.15	0.091	33.08	11	0.8110	0.65	4
340_U	209779	1.5	0.61	0.14	8.328	28	0.8299	0.49	4
341_U	93794	0.90	0.26	0.086	11.96	6.9	0.8256	0.51	4
342_U	44297	1.6	0.12	0.051	50.34	5.0	0.7976	0.81	4
343_U	16242	0.025	0.049	0.36	1.664	30	0.8431	1.3	4
344_U	74235	0.35	0.21	0.30	5.544	4.8	0.8306	0.54	4
345_U	195664	1.3	0.54	0.11	8.889	6.4	0.8261	0.44	4
346_U	48143	0.61	0.12	0.15	22.81	3.8	0.8228	0.73	4
348_U	71996	0.59	0.20	0.094	10.79	4.2	0.8295	0.57	4
349_U	102507	1.1	0.30	0.26	12.24	15	0.8253	0.50	4
350_U	198463	2.5	0.51	0.035	18.76	9.3	0.8261	0.44	4
359_U	8939	0.043	0.024	0.45	6.144	27	0.8324	1.6	4
360_U	57033	0.98	0.17	0.24	18.08	9.6	0.8228	0.61	4
361_U	203520	4.3	0.62	0.069	22.01	12	0.8204	0.40	4
362_U	35763	2.9	0.11	0.038	80.60	6.4	0.7848	0.83	4
364_U	79345	1.4	0.25	0.083	17.82	4.7	0.8228	0.53	4
365_U	39697	2.2	0.12	0.096	59.48	4.7	0.7974	0.71	4
366_U	144201	0.82	0.43	0.12	6.093	16	0.8321	0.46	4
367_U	53472	0.69	0.16	0.088	13.48	8.2	0.8254	0.67	4
369_U	50693	2.5	0.16	0.10	49.25	7.4	0.7975	0.74	4
370_U	52833	2.4	0.17	0.076	44.69	6.3	0.8078	0.59	4
371_U	51179	0.82	0.16	0.11	16.64	14	0.8172	0.80	4

Table 4: continue...

372_U	14847	0.28	0.040	0.050	25.52	12	0.8150	1.3	4	
373_U	67995	0.55	0.20	0.083	8.701	13	0.8239	0.64	4	
374_U	169346	1.3	0.47	0.11	9.985	12	0.8279	0.45	4	
375_U	73258	1.4	0.23	0.11	19.10	10	0.8199	0.56	4	
376_U	30039	0.42	0.097	0.077	13.11	14	0.8220	0.77	4	
377_U	243980	1.9	0.75	0.15	8.114	4.6	0.8310	0.40	4	
378_U	28838	1.6	0.089	0.085	58.11	5.3	0.7966	0.95	4	
379_U	106946	2.6	0.33	0.039	24.87	5.1	0.8131	0.56	4	
380_U	64981	2.6	0.19	0.10	44.16	5.7	0.8001	0.62	4	
381_U	82318	2.1	0.21	0.091	39.33	13	0.8072	0.72	4	
382_U	78516	1.3	0.24	0.098	17.00	12	0.8192	0.59	4	
383_U	BCR 9644	22865	0.86	0.064	0.024	47.56	5.1	0.8056	0.97	4
384_U	6644	0.84	0.019	0.0040	146.9	11	0.7724	2.0	4	
386_U	77128	0.27	0.22	0.20	4.265	4.1	0.8256	0.55	4	
388_U	3416	0.41	0.011	0.016	114.6	4.8	0.7689	2.2	4	
390_U	18463	0.33	0.058	0.075	17.53	12	0.8061	1.0	4	
392_U	4305	0.14	0.008	0.0033	180.6	13	0.7574	3.8	4	
393_U	14388	1.3	0.044	0.016	91.44	5.8	0.7816	1.2	4	
394_U	18496	0.92	0.054	0.019	57.05	5.9	0.7943	1.1	4	
395_U	4428	0.20	0.012	0.041	58.79	5.9	0.8002	2.0	4	
396_U	12559	0.81	0.032	0.016	104.3	5.4	0.7788	1.5	4	
397_U	1342	0.59	0.004	0.0025	376.0	9.0	0.6875	4.7	4	
399_U	146754	0.20	0.32	0.11	3.469	6.1	0.8260	0.55	4	
409_U	11621	0.41	0.035	0.018	39.82	11	0.8117	1.5	4	
411_U	5627	0.48	0.018	0.021	84.35	7.9	0.7917	1.8	4	
412_U	23357	1.1	0.066	0.038	60.11	6.7	0.8029	0.97	4	
413_U	14314	0.22	0.041	0.061	19.04	15	0.8151	1.2	4	
415_U	6924	0.056	0.014	0.15	29.96	8.1	0.8014	4.3	4	
416_U	6828	0.17	0.020	0.026	28.80	8.2	0.8266	1.7	4	
423_U	4952	0.090	0.013	0.027	28.72	6.2	0.8009	2.2	4	
427_U	962	0.047	0.003	0.10	42.18	11	0.8056	4.0	4	
433_U	733	0.30	0.002	0.0013	444.8	12	0.6898	6.1	4	
435_U	3290	0.025	0.006	0.0021	35.94	17	0.8262	4.6	4	
438_U	1004	0.076	0.003	0.035	74.37	9.7	0.7965	4.1	4	
440_U	891	0.007	0.003	0.025	8.704	10	0.8189	4.4	4	
441_U	3306	0.47	0.010	0.0019	139.0	5.6	0.7832	2.1	4	
442_U	1782	0.18	0.005	0.0024	103.9	4.6	0.7908	3.0	4	
443_U	1876	0.10	0.005	0.0018	68.27	5.5	0.8084	2.9	4	
445_U	892	0.009	0.003	0.74	7.259	21	0.8357	4.3	4	
446_U	5383	0.20	0.017	0.0096	39.50	8.5	0.8110	1.6	4	
447_U	5316	0.36	0.016	0.0056	72.00	3.8	0.8105	1.7	4	
448_U	4019	0.44	0.013	0.0056	107.8	7.5	0.7851	3.3	4	
449_U	839	0.16	0.002	0.0045	235.9	9.6	0.7546	4.5	4	
450_U	27342	1.5	0.084	0.040	56.17	8.4	0.8088	1.0	4	
459_U	5280	0.17	0.017	0.016	31.91	19	0.8161	1.7	4	
462_U	37636	0.062	0.11	0.23	1.778	9.2	0.8254	0.80	4	
464_U	200339	2.3	0.61	0.050	11.77	6.9	0.8140	0.43	4	
467_U	676	0.005	0.002	0.74	9.950	9.0	0.8126	5.2	4	
468_U	1808	0.060	0.006	0.013	33.81	11	0.8128	3.0	4	
469_U	3923	0.13	0.013	0.17	29.07	15	0.8056	2.1	4	

^a Within run background-corrected mean ²⁰⁷Pb signal in cps (counts per second).^b U and Pb concentrations and Th/U ratio were calculated relative to the primary reference material.^c Corrected for background, within-run Pb/U fractionation (in case of ²⁰⁶Pb/²³⁸U) and subsequently normalised to the primary reference material (ID-TIMS value/measured value).

References

- Andreetto, F., Matsubara, K., Beets, C.J., Fortuin, A.R., Flecker, R., and Krijgsman, W.: High Mediterranean water level during the Lago-Mare phase of the Messinian Salinity Crisis: insights from the Sr isotope records of Spanish marginal basins (SE Spain), *Paleogeogr. Paleoclimatol. Paleoecol.*, 562, 110139, <https://doi.org/10.1016/j.palaeo.2020.110139>, 2021.
- 365 Astilleros, J.M., Godelitsas, A., Rodríguez-Blanco, J.D., Fernández-Díaz, L., Prieto, M., Lagoyannis, A., and Harissopoulos, S.: Interaction of gypsum with Pb - bearing aqueous solutions, *Appl. Geochem.*, 25, 1008-1016. <https://doi.org/10.1016/j.apgeochem.2010.04.007>, 2010.
- Babel, M., and Schreiber, B.C.: Geochemistry of evaporites and evolution of seawater, in: Treatise on Geochemistry (2nd edition), edited by: Turekian, K., and Holland, H., Elsevier, Oxford, UK, 483-560, <https://doi.org/10.1016/B978-0-08-095975-7.00718-X>, 2014.
- 370 Brannon, J. C., Cole, S. C., Podosek, F. A., Ragan, V. M., Coveney, R. M., Wallace, M. W., and Bradley, A. J.: Th-Pb and U-Pb dating of ore-stage calcite and Paleozoic fluid flow, *Science*, 271, 491-493, <https://doi.org/10.1126/science.271.5248.491>, 1996.
- Burisch, M., Walter, B. F. and Markl, G.: Silicification of Hydrothermal Gangue Minerals in Pb-Zn-Cu-Fluorite-Quartz-595 Baryte Veins, *The Canadian Mineralogist*, 55(3), 501-514, <https://doi.org/10.3749/canmin.1700005>, 2017.
- Burisch, M., Gerdes, A., Meinert, L., Albert, R., Seifert, T., and Gutzmer, J.: The essence of time - fertile skarn formation in the Variscan Orogenic Belt, *Earth Planet. Sci. Lett.*, 519, 165-170, <https://doi.org/10.1016/j.epsl.2019.05.015>, 2019.
- CIESM: The Messinian salinity crisis from mega-deposits to microbiology, in: A consensus report. 33ème CIESM Workshop Monographs 33, edited by: Briand, F., CIESM Publisher, Monaco, 91-96, 2008.
- 380 Clauer, N., Chaudhuri, S., Toulkeridis, T., and Blanc, G.: Fluctuations of Caspian Sea level: beyond climatic variations? *Geology*, 28, 1015–1018, [https://doi.org/10.1130/0091-7613\(2000\)28<1015:FOCSLB>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<1015:FOCSLB>2.0.CO;2), 2000.
- Conley, R.F., and Bundy W.M.: Mechanism of gypsumification, *Geochim. Cosmochim. Acta*, 15, 57-72, [https://doi.org/10.1016/0016-7037\(58\)90010-3](https://doi.org/10.1016/0016-7037(58)90010-3), 1958.
- Costanzo, A., Cipriani, M., Feely, M., Cianfione, G., and Dominici, R.: Messinian twinned selenite from the Catanzaro Trough, 385 Calabria, Southern Italy: field, petrographic and fluid inclusion perspectives, *Carbonates and Evaporites*, 34, 743–756. <https://doi.org/10.1007/s13146-019-00516-0>, 2019.
- Craig, G., Managh A.J., Stremtan, C., Lloyd, N.S., and Horstwood, M.S.A.: Doubling Sensitivity in Multicollector ICPMS Using High-Efficiency, Rapid Response Laser Ablation Technology, *Anal. Chem.*, 90, 11564-11571, <https://doi.org/10.1021/acs.analchem.8b02896>, 2018.
- 390 Craig, G., Bracciali, L., and Lloyd, N.: LA-ICP-MS for U-(Th)-Pb geochronology: Which analytical capability is right for my laboratory? *Thermo Fish. Sci. Smart. Note* 30581, 2020.
- Cruset, D., Verges, J., Rodrigues, N., Belenguer, J., Pascual-Cebrian, E., Almar, Y., Perez-Caceres, I., Macchiavelli, C., Trave, A., Beranoaguirre, A., Albert, R., Gerdes, A., and Messager, G.: U-Pb dating of carbonate veins constraining timing of beef

- growth and oil generation within Vaca Muerta Formation and compression history in the Neuquen Basin along the Andean
395 fold and thrust belt, Mar. Petrol. Geol., 132, 10520, <https://doi.org/10.1016/j.marpetgeo.2021.105204>, 2021.
- Deng, X.D., Li, J.W., Luo, T., and Wang, H.Q.: Dating magmatic and hydrothermal processes using andradite-rich garnet U-Pb geochronometry, Contrib Mineral Petr., 172, 71, <https://doi.org/10.1007/s00410-017-1389-2>, 2017.
- Elisha, B., Nuriel, P., Kylander-Clark, A., and Weinberger, R.: Towards in-situ U-Pb dating of dolomites, Geochronology 3, 337-349, <https://doi.org/10.5194/gchron-3-337-2021>, 2021.
- 400 Evans, N.P., Turchyn, A.V., Gázquez, F., Bontognali, R.R., Chapman, H.J., and Hodell, D.A.: Coupled measurements of $\delta^{18}\text{O}$ and δD of hydration water and salinity of fluid inclusions in gypsum from the Messinian Yesares Member, Sorbas Basin (SE Spain), Earth Planet. Sci. Lett., 430, 499–510, <https://doi.org/10.1016/j.epsl.2015.07.071>, 2015.
- Flecker, R., and Ellam, R.M.: Identifying Late Miocene episodes of connection and isolation in the Mediterranean - Paratethyan realm using Sr isotopes, Sediment. Geol., 188-189, 189-203. <https://doi.org/10.1016/j.sedgeo.2006.03.005>, 1999
- 405 Flecker, R., Krijgsman, W., Capella, W., de Castro Martíns, C., Dmitrieva, E., Mayser, J.P., Marzocchi, A., Modestu, S., Lozano, D.O., Simon, D., Tulbure, M., van den Berg, B., van der Schee, M., de Lange, G., Ellam, R., Govers, R., Gutjahr, M., Hilgen, F., Kouwenhoven, T., Lofi, J., Meijer, P., Sierro, F.J., Bachiri, N., Barhoum, N., Alami, A.C., Chacon, B., Flores, Jose A., Gregory, J., Howard, J., Lunt, D., Ochoa, M., Pancost, R., Vincent, S., and Yousfi, M.Z.: Evolution of the Late Miocene Mediterranean Atlantic gateways and their impact on regional and global environmental change, Earth-Sci. Rev., 150, 365-410 392. <https://doi.org/10.1016/j.earscirev.2015.08.007>, 2015
- Gerdes, A., and Zeh, A.: Combined U-Pb and Hf isotope LA-(MC-)ICP-MS analyses of detrital zircons: comparison with SHRIMP and new constraints for the provenance and age of an Armorican metasediment in Central Germany, Earth Planet. Sci. Lett., 249, 47-61, <https://doi.org/10.1016/j.epsl.2006.06.039>, 2006.
- 415 Gerdes, A., and Zeh, A.: Zircon formation versus zircon alteration — new insights from combined U-Pb and Lu-Hf in-situ LA-ICP-MS analyses, and consequences for the interpretation of Archean zircon from the Central Zone of the Limpopo Belt, Chem. Geol., 261 (3-4), 230-243, <https://doi.org/10.1016/j.chemgeo.2008.03.005>, 2009.
- Grandia, F., Asmerom, Y., Getty, S., Cardellach, E., and Canals, A.: U-Pb dating of MVT ore-stage calcite: implications for fluid flow in a Mesozoic extensional basin from Iberian Peninsula, J. Geochem. Explor., 69, 377-380, [https://doi.org/10.1016/S0375-6742\(00\)00030-3](https://doi.org/10.1016/S0375-6742(00)00030-3), 2000.
- 420 Grothe, A., Andreetto, F., Reichart, G.J., Wolthers, M., Van Baak, C.G., Vasiliev, I., Stoica, M., Sangiorgi, F., Middelburg, J.J., Davies, G.R., and Krijgsman, W.: Paratethys pacing of the Messinian Salinity Crisis: low salinity waters contributing to gypsum precipitation?, Earth Planet. Sci. Lett., 532, 116029, <https://doi.org/10.1016/j.epsl.2019.116029>, 2020.
- Guillong, M., Wotzlaw, J.-F., Looser, N., and Laurent, O.: Evaluating the reliability of U-Pb laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) carbonate geochronology: matrix issues and a potential calcite validation reference material, Geochronology, 2, 155-167, <https://doi.org/10.5194/gchron-2-155-2020>, 2020.
- 425 Horstwood, M. S. A., Košler, J., Gehrels, G., Jackson, S. E., McLean, N. M., Paton, C., Pearson, N. J., Sircombe, K., Sylvester, P., Vermeesch, P., and Bowring, J. F.: Community-derived standards for LA-ICP-MS U-(Th-) Pb geochronology-Uncertainty

- propagation, age interpretation and data reporting, *Geostand. Geoanal. Res.*, 40, 311-332, <https://doi.org/10.1111/j.1751-908X.2016.00379.x>, 2016.
- 430 Hsü, K.J., Ryan, W.B.F., and Cita, M.B.: Late Miocene desiccation of the Mediterranean, *Nature*, 242, 240-244, <https://doi.org/10.1038/242240a0>, 1973.
- Hsü, K. J., Montadert, L., Ross, D.A., and Neprochnov, Y.P.: Annotated record of the detailed examination of Mn deposits from DSDP Leg 42 (Holes 372 and 379A). *Pangaea*, <https://doi.org/10.1594/PANGAEA.871889>, 1978. Supplement to: Worstell, P.J., Mélières, F., Bernoulli, D., Erickson, A.J., Wright, R., Bizon, G., Cita, M.B., Müller, C., Kidd, R.B., Fabricius, 435 F.H., Garrison, R.E., Hsü, K.J., and Montadert, L.: Initial Reports of the Deep Sea Drilling Project. Initial Reports of the Deep Sea Drilling Project, U.S. Government Printing Office, XLII Pt. 1, 1249 pp + 1244 pp, <https://doi.org/10.2973/dsdp.proc.42-1.1978>, 1978.
- Jochum, K. P., Weis, U., Stoll, B., Kuzmin, D., Yang, Q., Raczek, I., Jacob, D.E., Stracke, A., Birbaum, K., Frick, D.A., Günther, D., and Enzweiler, J.: Determination of reference values for NIST SRM 610–617 glasses following ISO guidelines, 440 *Geostand. Geoanal. Res.*, 35, 97–429, <https://doi.org/10.1111/j.1751-908X.2011.00120.x>, 2011.
- Kameda, K., Hashimoto, Y., Wang, S.-L., Hirai, Y., and Miyahara, H.: Simultaneous and continuous stabilization of As and Pb in contaminated solution and soil by a ferrihydrite-gypsum sorbent, *J. Hazard. Mater.*, 327, 171-179, <https://doi.org/10.1016/j.jhazmat.2016.12.039>, 2017.
- Krijgsman, W., Hilgen, F. J., Raffi, I., Sierra, F. J., and Wilson, D. S.: Chronology, causes and progression of the Messinian 445 Salinity Crisis, *Nature*, 400, 652-655, <https://doi.org/10.1038/23231>, 1999.
- Krijgsman, W., Stoica, M., Vasiliev, I., and Popov, V. V.: Rise and fall of the Paratethys Sea during the Messinian Salinity Crisis, *Earth Planet. Sci. Lett.*, 290 (1-2), 183-191, <https://doi.org/10.1016/j.epsl.2009.12.020>, 2010.
- Krijgsman, W., Capella, W., Simon, D., Hilgen, F.J., Kouwenhoven, T.J., Meijer, P.T., Sierra, F.J., Tulbure, M.A., van den Berg, B.C.J., van der Schee, M., and Flecker, R.: The Gibraltar Corridor: watergate of the Messinian Salinity Crisis, *Mar. 450 Geol.*, 403, 238-246, <https://doi.org/10.1016/j.margeo.2018.06.008>, 2018.
- Laskar, J.: The limits of Earth orbital calculations for geological time-scale use, in: Astronomical (Milankovitch) Calibration of the Geological Time-Scale, edited by: Shackleton, N.J., McCave, I.N., and Graham, P.W., *Philos. Trans. R. Soc., Ser. A.*, 357(1757), 1735–1759. <https://doi.org/10.1098/rsta.1999.0399>, 1999.
- Geochemistry of evaporites and evolution of seawater, in: Treatise on Geochemistry (2nd edition), edited by: Turekian, K., 455 and Holland, H., Elsevier, Oxford, UK, 483-560, <https://doi.org/10.1016/B978-0-08-095975-7.00718-X>, 2014.
- Lenoir, L., Blaise, T., Somogyi, A., Brigaud, B., Barbarand, J., Boukari, C., Nouet, J., and Pagel, M.: Uranium incorporation in fluorite and exploration of U-Pb dating, *Geochronology*, 3, 197-227, <https://doi.org/10.5194/gchron-3-199-2021>, 2021.
- Lin, J., Sun, W., Desmarais, J., Chen, N., Feng, R., Zhang, P., Li, D., Lieu, A., Tse, J.S., and Pan, Y.: Uptake and speciation of uranium in synthetic gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$): applications to radioactive mine tailings, *J. Environ. Radioact.*, 181, 8-17, 460 <https://doi.org/10.1016/j.jenvrad.2017.10.010>, 2018.

- Liu, D.J., and Hendry, M.J.: Controls on ^{226}Ra during raffinate neutralization at the Key Lake uranium mill, Saskatchewan, Canada, App. Geochem., 26, 2113-2120, <https://doi.org/10.1016/j.apgeochem.2011.07.009>, 2011.
- Ludwig, K.R.: User's Manual for Isoplot Version 3.75-4.15: a Geochronological Toolkit for Microsoft Excel, Berkeley Geochronological Center Special Publication, 5, 2012.
- 465 Lugli, S., Bassetti, M.A., Manzi, V., Barbieri, M., Longinelli, A., and Roveri, M.: The Messinian 'Vena del Gesso' evaporites revisited: characterization of isotopic composition and organic matter, J. Geol. Soc. Lond., 285, 179-190, <http://dx.doi.org/10.1144/SP285.11>, 2007.
- Lugli, S., Manzi, V., Roveri, M., and Schreiber, B.C.: The primary Lower Gypsum in the Mediterranean: a new facies interpretation for the first stage of the Messinian salinity crisis, Palaeogeogr. Palaeoclimatol. Palaeoecol., 297, 83-99,
- 470 <https://doi.org/10.1016/j.palaeo.2010.07.017>, 2010.
- Mangenot, X., Gasparrini, M., Rouchon, V. and Bonifacie, M.: Basin-scale thermal and fluid flow histories revealed by carbonate clumped isotopes ($\Delta 47$) - Middle Jurassic carbonates of the Paris Basin depocentre, Sedimentology, 65(1), 123-150, <https://doi.org/10.1111/sed.12427>, 2018.
- Manzi, V., Gennari, R., Lugli, S., Roveri, M., and Schreiber, B.C.: The Messinian "Calcare di Base" (Sicily, Italy) revisited,
- 475 Geol. Soc. Am. Bull., 123, 347-370, <https://doi.org/10.1130/B30262.1>, 2011.
- Manzi, V., Gennari, R., Hilgen, F., Krijgsman, W., Lugli, S., Roveri, M., and Sierro, F.J.: Age refinement of the Messinian salinity crisis onset in the Mediterranean, Terra Nova, 315-322, <https://doi.org/10.1111/ter.12038>, 2013.
- Manzi, V., Gennari, R., Lugli, S., Persico, D., Reghizzi, M., Roveri, M., Schreiber, B.C., Calvo, R., Gavrieli, I., and Gvirtzman, Z.: The onset of the Messinian salinity crisis in the deep Eastern Mediterranean basin, Terra Nova 30 (3), 189-198,
- 480 <https://doi.org/10.1111/ter.12325>, 2018.
- Meijlison, A., Hilgen, F., Sepúlveda, J., Steinberg, J., Fairbank, V., Flecker, R., Waldmann, N.D., Spaulding, S.A., Bialik, O.M., and Boudinot, F.G.: Chronology with a pinch of salt: integrated stratigraphy of Messinian evaporites in the deep Eastern Mediterranean reveals long-lasting halite deposition during Atlantic connectivity, Earth-Sci. Rev. 194, 374-398. <https://doi.org/10.1016/j.earscirev.2019.05.011>, 2019
- 485 Millonig, L.J., Albert, R., Gerdes, A., Avigad, D., and Dietsch, C.: Exploring laser ablation U-Pb dating of regional metamorphic garnet - The Straits Schist, Connecticut, USA, Earth Planet. Sci. Lett., 552, 116589, <https://doi.org/10.1016/j.epsl.2020.116589>, 2020.
- Montano, D., Gasparrini, M., Gerdes, A., Albert, R., Rohais, S., and Della Porta, G.: In-situ carbonate U-Pb analysis by LA-ICP-MS: From absolute dating to understanding the U-Pb partitioning in lacustrine systems, Goldschmidt 2019 Abstracts,
- 490 2323, 2019.
- Montano, D., Gasparrini, M., Gerdes, A., Della Porta, G., and Albert, R.: In-situ U-Pb dating of Ries Crater lacustrine carbonates (Miocene, South-West Germany): implications for continental carbonate chronostratigraphy, Earth Planet. Sci. Lett., 568, 117011, <https://doi.org/10.1016/j.epsl.2021.117011>, 2021.

- Montano, D., Gasparrini, M., Rohais, S., Albert, R., and Gerdes, A.: Depositional age models in lacustrine systems from zircon and carbonate U-Pb geochronology, *Sedimentology*, in press, <https://doi.org/10.1111/sed.13000>, 2022.
- Morales, J., Astilleros, J.M., Jiménez, A., Göttlicher, J., Steininger, R., and Fernández-Díaz, L.: Uptake of dissolved lead by anhydrite surfaces, *Appl. Geochem.*, 40, 89-96, <https://doi.org/10.1016/j.apgeochem.2013.11.002>, 2014.
- Murray, R.C.: Origin and diagenesis of gypsum and anhydrite, *J. Sedimen. Res.*, 34, 512-523, <https://doi.org/10.1306/74D710D2-2B21-11D7-8648000102C1865D>, 1964.
- Natalicchio, M., Dela Pierre, F., Lugli, S., Lowenstein, T.K., Feiner, S.J., Ferrando, S., Manzi, V., Roveri, M., and Clari, P.: Did Late Miocene (Messinian) gypsum precipitate from evaporated marine brines? Insights from the Piedmont Basin (Italy), *Geology*, 42, 179–182, <https://doi.org/10.1130/G34986.1>, 2014.
- Nuriel, P., Wotzlaw, J.F., Ovtcharova, M., Vaks, A., Stremtan, C., Šala, M., Roberts, N.M.W., and Kylander-Clark, A.R.C.: The use of ASH-15 flowstone as a matrix-matched reference material for laser-ablation U – Pb geochronology of calcite, *Geochronology*, 3, 35-47, <https://doi.org/10.5194/gchron-3-35-2021>, 2021.
- Ossorio, M., Van Driessche, A.E.S., Pérez, P., and García-Ruiz, J.M.: The gypsum-anhydrite paradox revisited, *Chem. Geol.*, 386, 16-21, <https://doi.org/10.1016/j.chemgeo.2014.07.026>, 2014.
- Pagel, M., Bonifacie, M., Schneider, D. A., Gautheron, C., Brigaud, B., Calmels, D., Cros, A., Saint-Bezar, B., Landrein, P., Sutcliffe, C., and Davis, D.: Improving paleohydrological and diagenetic reconstructions in calcite veins and breccia of a sedimentary basin by combining $\Delta 47$ temperature, $\delta^{18}\text{O}$ water and U-Pb age, *Chem. Geol.*, 481, 1-17, <https://doi.org/10.1016/j.chemgeo.2017.12.026>, 2018.
- Parrish, R. R., Parrish, C. M., and Lasalle, S.: Vein calcite dating reveals Pyrenean orogen as cause of Paleogene deformation in southern England, *J. Geol. Soc.*, 175, 425-442, <https://doi.org/10.1144/jgs2017-107>, 2018.
- Pettrash, D.A., Bialik, O.M., Bontognali, T.R.R., Vasconcelos, C., Roberts, J.A., McKenzie, J.A., and Konhauser, K.O.: Microbially catalyzed dolomite formation: From near-surface to burial, *Earth-Sci. Rev.*, 171, 558-582, <https://doi.org/10.1016/j.earscirev.2017.06.015>, 2017.
- Piccione, G., Rasbury, E.T., Elliott, B.A., Kyle, J.R., Jaret, S.J., Acerbo, A.S., Lanzirotti, A., Northrup, P., Wootton, K., and Parrish, R.R.: Vein fluorite U-Pb dating demonstrates post-6.2 Ma rare-earth element mobilization associated with Rio Grande rifting, *Geosphere*, 15, 6, 1958– 1972, <https://doi.org/10.1130/GES02139.1>, 2019.
- Rasbury, E. T., and Cole, J. M.: Directly dating geologic events: U-Pb dating of carbonates, *Reviews of Geophysics*, 47(3), <https://doi.org/10.1029/2007RG000246>, 2009.
- Rouchy, J.M., and Caruso, A.: The Messinian salinity crisis in the Mediterranean basin: a reassessment of the data and an integrated scenario, *Sediment. Geol.*, 188–189, p. 35–67, <https://doi.org/10.1016/j.sedgeo.2006.02.005>, 2006.
- Ring, U. and Gerdes, A.: Kinematics of the Alpenrhein-Bodensee graben system in the Central Alps: Oligocene/Miocene transtension due to formation of the Western Alps arc, *Tectonics*, 35, 1367-1391, <https://doi.org/10.1002/2015TC004085>, 2016.

Roberts, N. M. W., Rasbury, E. T., Parrish, R. R., Smith, C. J., Horstwood, M. S. A., and Condon, D. J.: A calcite reference material for LA-ICP-MS U-Pb geochronology, *Geochem. Geophys. Geosy.*, 18, 2807-2814, <https://doi.org/10.1002/2016GC006784>, 2017.

530 Roberts, N. M. W., Drost, K., Horstwood, M. S. A., Condon, D. J., Chew, D., Drake, H., Milodowski, A. E., McLean, N. M., Smye, A. J., Walker, R. J., Haslam, R., Hodson, K., Imber, J., Beaudoin, N., and Lee, J. K.: Laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) U-Pb carbonate geochronology: strategies, progress, and limitations, *Geochronology*, 2, 33-61, <https://doi.org/10.5194/gchron-2-33-2020>, 2020.

Roveri, M., Lugli, S., Manzi, V., and Schreiber, B.C.: The Messinian Sicilian stratigraphy revisited: toward a new scenario for the Messinian salinity crisis, *Terra Nova*, 20, 483-488, <https://doi.org/10.1111/j.1365-3121.2008.00842.x>, 2008a.

Roveri, M., Bertini, A., Cosentino, D., Di Stefano, A., Gennari, R., Glioza, E., Grossi, F., Iaccarino, S.M., Lugli, S., Manzi, V., and Taviani, M.: A high-resolution stratigraphic framework for the latest Messinian events in the Mediterranean area, *Stratigraphy*, 5, 323-342, 2008b.

Roveri, M., Flecker, R., Krijgsman, W., Lofi, J., Lugli, S., Manzi, V., Sierro, F.J., Bertini, A., Camerlenghi, A., De Lange, G., 540 Govers, R., Hilgen, F.J., Hübscher, C., Meijer, P.T., and Stoica, M.: The Messinian Salinity Crisis: past and future of a great challenge for marine sciences, *Mar. Geol.*, 352, 25-58, <https://doi.org/10.1016/j.margeo.2014.02.002>, 2014a.

Roveri, M., Lugli, S., Manzi, V., Gennari, R., and Schreiber, B.C.: High-resolution strontium isotope stratigraphy of the Messinian deep Mediterranean basins: implications for marginal to central basins correlation, *Mar. Geol.*, 349, 113-125, <https://doi.org/10.1016/j.margeo.2014.01.002>, 2014b.

545 Ryan, W.B.: Decoding the Mediterranean salinity crisis, *Sedimentology*, 56 (1), 95-136, <https://doi.org/10.1111/j.1365-3091.2008.01031.x>, 2009

Schaltegger, U., Schmitt, A.K., and Horstwood, M.S.A.: U-Th-Pb zircon geochronology by ID-TIMS, SIMS, and laser ablation ICP-MS: Recipes, interpretations, and opportunities, *Chem. Geol.*, 402, 89-110, <https://doi.org/10.1016/j.chemgeo.2015.02.028>, 2015.

550 Sell, R.: Il Messiniano Mayer-Eymar 1867. Proposta di un neostratotipo, *Giornale di Geologia*, 28, 1-33, 1960.

Seman, S., Stockli, D.F., and McLean, N.M.: U-Pb geochronology of grossular-andradite garnet, *Chem. Geol.*, 460, 106-116, <https://doi.org/10.1016/j.chemgeo.2017.04.020>, 2017.

Sindern, S., Havenith, V., Gerdes, A., Meyer, F.M., Adelmann, D., and Hellmann, A.: Dating of anatase-forming diagenetic reactions in Rotliegend sandstones of the North German Basin, *Int. J. Earth Sci.*, 108, 1275-1292, 555 <https://doi.org/10.1007/s00531-019-01705-x>, 2019.

Sylvester, P.: Matrix effects in Laser ablation-ICP-MS, in: *Laser Ablation-ICP-MS in the Earth Sciences: Current Practices and Outstanding Issues*, edited by: Sylvester, P., Mineralogical association of Canada, 67-78, 2008.

Van Driessche, A.E.S., Stawski, T., and Kellermeier, M.: Calcium sulfate precipitation pathways in natural and engineering environments, *Chem. Geol.*, 530, 119274. <https://doi.org/10.1016/j.chemgeo.2019.119274>, 2019.

- 560 Vasiliev, I., Mezger, E.M., Lugli, S., Reichart, G.J., Manzi, V., and Roveri, M.: How dry was the Mediterranean during the Messinian salinity crisis?, *Paleogeogr. Paleoclimatol. Paleoecol.*, 471, 120-133, <https://doi.org/10.1016/j.palaeo.2017.01.032>, 2017.
- Wafforn, S., Seman, S., Kyle, J.R., Stockli, D., Leys, C., Sonbait, D., and Cloos, M.: Andradite garnet U-Pb geochronology of the big Gossan skarn, Ertsberg-Grasberg mining district, Indonesia, *Econ. Geol.*, 113, 769-778, 565 <https://doi.org/10.5382/econgeo.2018.4569>, 2018.
- Warren, J.K.: *Evaporites: A Geological Compendium*, Springer, Berlin, 1813 p, <https://doi.org/10.1007/978-3-319-13512-0>, 2016.
- 570 Warthmann, R., van Lith, Y., Vasconcelos, C., McKenzie, J.A., and Karpoff, A.M.: Bacterially induced dolomite precipitation in anoxic culture experiments, *Geology*, 28, 1091-1094, [https://doi.org/10.1130/0091-7613\(2000\)28<1091:BIDPIA>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<1091:BIDPIA>2.0.CO;2), 2000.
- Woodhead, J., Hellstrom, J., Maas, R., Drysdale, R., Zanchetta, G., Devine, P., and Taylor, E.: U-Pb geochronology of speleothems by MC-ICPMS, *Quat. Geochronol.*, 1, 208-221, <https://doi.org/10.1016/j.quageo.2006.08.002>, 2006.
- Woodhead, J., Hellstrom, J., Pickering, R., Drysdale, R., Paul, B., and Bajo, P.: U and Pb variability in older speleothems and strategies for their chronology, *Quat. Geochronol.*, 14, 105-113, <https://doi.org/10.1016/j.quageo.2012.02.028>, 2012.
- 575 Yan, S., Zhou, R.J., Niu, H.C., Feng, Y.X., Nguyen, A.D., Zhao, Z.H., Yang, W.B., Qian, D., and Zhao, J.X.: LA-MC-ICP-MS U-Pb dating of low-U garnets reveals multiple episodes of skarn formation in the volcanic-hosted iron mineralization system, Awulale belt, Central Asia, *GSA Bulletin*, 132 (5-6), 1031-1045, <https://doi.org/10.1130/B35214.1>, 2020
- Yang, Y.H., Wu, F.Y., Yang, J.H., Mitchell, R.H., Zhao, Z.F., Xie, L.W., Huang, C., Ma, Q., Yang, M., and Zhao, H.: U-Pb age determination of schorlomite garnet by laser ablation inductively coupled plasma mass spectrometry, *J. Anal. At. Spectrom.*, 33, 231-239, <https://doi.org/10.1039/c7ja00315c>, 2018.
- 580 Zachariasse, W.J., van Hinsbergen, D.J.J., and Fortuin, A.R.: Mass wasting and uplift on Crete and Karpathos during the early Pliocene related to initiation of south Aegean left-lateral, strike-slip tectonics, *Geol. Soc. Am. Bull.*, 120, 976–993, <https://doi.org/10.1130/B26175.1>, 2008.