



## Shallow marine carbonates as recorders of orbitally induced past climate changes – example from the Oxfordian of the Swiss Jura Mountains

## 5 André Strasser

Department of Geosciences, University of Fribourg, Fribourg, 1700, Switzerland

Correspondence to: André Strasser (andreas.strasser@unifr.ch)

10

Aby Tay. The detailed analysis of shallow marine carbonate sections in the Swiss Jura Mountains allows interpreting the climate changes that affected the sedimentary system during the middle to late Oxfordian Jurassic). The sediments formed on a subtropical platform where carbonate-producing organisms proliferated, and oncoids and oncoids were common. The sections are composed of hierarchically stacked elementary, small-scale, and medium scale denositional sequences where facing changes inply deepening challowing trends. The

- 15 scale, and medium-scale depositional sequences where facies changes imply deepening-shallowing trends. The major sequence boundaries Ox 6, Ox 7, and Ox 8 can be correlated with those of other European basins and place the studied sections in a broader framework. The chronostratigraphic tie points imply that the medium- and small-scale sequences formed in tune with the orbital eccentricity cycles of 405 and 100 kyr, respectively, and the elementary sequences with the precession cycle of 20 kyr. Orbitally controlled insolation changes at the top
- 20 of the atmosphere translated into climate changes: low insolation generally resulted in low sea level amplitudes at the 20 kyr frequency, and in a cool and humid climate at the palaeolatitude of the Jura platform. Terrigenous material was eroded from the hinterland and distributed over the platform. High insolation led to sea level rise to create accommodation, and to warm and semiarid to arid conditions in which coral reefs could grow. However, nutrient input favoured growth of microbialites that encrusted the corals. The reconstruction of high-frequency
- 25 sea level fluctuations based on facies analysis compares well with the curve of insolation changes calculated for the past 500 kyr. It is therefore assumed that sea level fluctuations were mainly due to thermal expansion and retraction of ocean surface water. Two models are presented that explain the formation of elementary sequences, one for low and one for high insolation. Despite the important lateral facies variations typical of a shallow marine platform, and despite the uncertainties in the reconstruction of sea level changes, this study demonstrates
- 30 the potential of carbonate ecosystems to record past climate changes at a time resolution of 20'000 years. Relatively short time windows can thus be opened in the deep geologic past, and processes and products there can be compared with those of the Holocene and the Recent. For example, it appears that today's anthropogenically induced sea level rise is more than ten times faster than the fastest rise reconstructed for the Oxfordian.
- 35

#### **1** Introduction

- 40 In today's world of global climate change and its impact on *Homo sapiens*, research into past climate variability as recorded in sediments becomes particularly interesting. Rapid and profound climate-induced changes in the palaeoecosystems can thus be documented and may serve as example of changes we may experience in the near future.
- 45 Orbital cycles (Milankovitch cycles) are an important factor steering global climate as they control the amount of solar energy received by the Earth. The quasi-periodicities of these cycles (precession, obliquity, short and long eccentricity) have been calculated from today back to the Precambrian (e.g., Berger et al., 1992; Laskar et al., 2011; Hinnov, 2018). They are used to calibrate geological time scales (e.g., Gradstein et al., 2020) and to reconstruct past climate changes with a high time resolution. At the equator, the daily insolation fluctuates
- around 400 W/m<sup>2</sup>, with the precession cycle modulated by the eccentricity cycles. At higher latitudes the insolation values are higher, with higher amplitudes, and the obliquity signal is better expressed (Fig. 1). Today, the frequency peaks of precession are at 21 and 23 kyr, the obliquity at 41 kyr, the short eccentricity at 100 kyr, and the long eccentricity at 405 kyr. While the eccentricity cycles stay constant through time, precession and obliquity were shorter in the geological past (Berger et al., 1992; Hinnov, 2018). In the Oxfordian (Late Jurassic)
- 55 considered here, the precession cycle had a value of about 20 kyr.

In deep water sedimentary records, the effects of the orbital cycles on sedimentation have been well documented for many geological periods. For example, Amies et al. (2019) explained how sapropel formation in the





Mediterranean basin reflects a northward migration of the monsoon rain belt over North Africa during the last interglacial, and this migration was controlled by the precession cycle (Lourens et al., 1996). Another example comes from the Valanginian and Hauterivian in the Vocontian basin analyzed by Martinez (2018), where precession, obliquity, and the two eccentricity cycles are held responsible for the formation of repetitive, hemipelagic limestone-marl alternations, and which help refining the duration of these two stages. In the Kimmeridge Clay Formation of the U.K., Armstrong et al. (2016) documented the short and long eccentricity

65 cycles that periodically induced extremely humid conditions leading to high burial of organic matter. In the Paleogene-Neogene Ebro Basin, Valero et al. (2014) described a 20 Myr lacustrine sedimentary record that formed in tune with the orbital eccentricity cycles.



70

Figure 1. Mean daily insolation values for June 21 (northern summer solstice) at different latitudes, calculated for the last million years (modifed from Hinnov, 2018).

75

In deep marine or deep lacustrine basins, there is a good chance that sedimentation was relatively continuous and interrupted only by tectonic pulses or gravitational flows. Sea level falls have an indirect influence by forcing progradation and thus an increase in turbidity currents and debris flows. On shallow platforms and ramps, however, already minor sea level drops can lead to subaerial exposure and thus to hiatuses in the sedimentary record (e.g., Sadler, 1994; Strasser, 2015). Also, the depositional environment is much less homogenous than in

80 record (e.g., Sadler, 1994; Strasser, 2015). Also, the depositional environment is much less homogenous than in deep water settings: facies changes may occur over short (100 meter scale) distances, creating facies mosaics (e.g., Rankey and Reeder, 2010). Nevertheless, also shallow marine carbonates have a potential to record climate changes, and this at a (geologically speaking) high time resolution.

#### 85

100

#### 2 Climatic controls on shallow marine carbonate systems



Carbonate-producing marine organisms such as corals, calcareous algae, bivalves, gastropods, echinoderms, foraminifera, or serpulids are sensitive to water depth, water temperature, water energy, water chemistry, turbidity, nutrients, and substrate, all factors that are directly or indirectly controlled by climate. Carbonate particles such as ooids or oncoids commonly form under the influence of microbes, and calcifying microbes can build up stromatolites or thrombolites. The microbial communities again are dependent on multiple ecological factors and thus on climate.

Orbital cycles induce insolation changes at the top of the atmosphere, dependent on the latitude (Fig. 1). The translation of insolation changes into climate changes that ultimately influence the shallow marine carbonate platform or ramp is complex and passes through atmospheric and oceanic circulation (Fig. 2), which are themselves dependent on latitude, orography, and land-ocean distribution, as well as on shape and position of the oceanic basins (e.g., Feng and Poulsen, 2014). Multiple feedback processes occur, with different frequencies and





amplitudes. For example, vegetation or ice cover will modify the albedo, or sea level change, ocean dynamics, and nutrient cycling may impact on atmospheric CO<sub>2</sub> levels and water chemistry (e.g., Wallmann et al., 2016). This will finally result in the regional climate that influences the carbonate platform under investigation (air and water temperature, rainfall, wind, and seasonality). However, it also has to be considered that if the seasonal insolation change is below a certain threshold it may not induce a significant climate change and will not be

105 insolation change is below a certain threshold it may not induce a significant climate change and will not be recorded in the sedimentary system (Hinnov, 2018). In this paper, only subtropical carbonate systems are considered, and the following case study deals with such sediments. In fact, carbonates are produced in different "factories", from tropical to cool water, and from shallow to deep (Flügel, 2004; Reijmer, 2021).



**Figure 2.** Chain of factors that influence a hypothetical, subtropical, shallow marine carbonate platform, including feedback processes. Amplitudes and frequencies of these factors may vary through time. The idealized cross section includes typical carbonate-producing environments (modified from Strasser, 2018).

Sea level is an important factor that controls water depth. Water depth influences the growth of calcareous algae that use photosynthesis, and of carbonate-producing organisms that contain light-dependent symbionts such as corals. Wave action in shallow water reworks the sediment and winnows out the mud fraction, while in deeper water carbonate mud can accumulate. The formation of ooid shoals is most efficient in a regime of tidal current which are strongest in shallow channels. Sea level also controls accommodation, i.e. the potential to accumulate

the sediment. Eustatic sea level changes were highly asymmetrical in icehouse worlds through slow buildup of polar icecaps and mountain glaciers, followed by rapid melting. The amplitudes were in the order of tens of meters (120 m during the last glacial-interglacial cycle; Shackleton, 1987). In greenhouse worlds such as the

Late Jurassic (e.g., Sellwood and Valdes, 2008), the sea level changes were meter scale only, caused by changes in ice volume of small polar ice caps and glaciers, by thermally controlled volume changes of the ocean surface waters, by density controlled volume changes of deep ocean waters, and/or by changes in the volume of freshwater sequestered on the continents (Schulz and Schäfer-Neth, 1997; Church and Clark, 2013; Sames et al.,

130 2016; Ray et al., 2019).

If it can be shown that the physical and ecological parameters reigning over a sedimentary system were controlled by climate, and assuming that climate was at least partly controlled by orbital cycles, then a time resolution in the order of the shortest such cycles, the precession cycle of about 20 kyr, becomes possible. In the following, the potential of shallow marine carbonates as recorders of orbitally controlled climate changes will be

135 following, the potential of shallow marine carbonates as recorders of orbitally ce illustrated with an example from the Oxfordian of the Swiss Jura Mountains.

115





#### 3 Oxfordian case study

#### 3.1 Stratigraphy and palaeogeography

145

The Oxfordian sedimentary rocks in the Swiss Jura Mountains have been well studied by Gygi (1995, 2000). He established a lithostratigraphic scheme and dated it biostratigraphically by numerous ammonites (Fig. 3). In addition, Gygi et al. (1998) defined sequence boundaries that are correlated with the boundaries found in the chart of Hardenbol et al. (1998) that summarizes the sequence stratigraphy of European basins. In the present study, the focus is set on the interval between sequence boundaries Ox 6 and Ox 8 (Fig. 3).

150

Chro stro	no- at.	Biostra ammonite zones	itigraphy subzones	Sequence boundaries	Lithostratigraphy	
Kimm.		Hypselocyclum		— Kim 2 —	Reuchenette Fm	
		Platynota		— Kim 1 —		
	fordian late	Planula	Galar Planula	— Ox 8 — — Ox 7 —	Courgenay Porrentruy Mb Verena Mb	
		Bimammatum	Hauffianum		Fm La May Mb O Holznun Laufen Mb Mb	
			Bimammatum		Bure Mb Oolithe rousse	
L LE			Berrense Hypselum		Hauptmumienbank Mb	
fordi		Bifurcatus	Grossouvrei	— Ox 6 —	Vellerat Röschenz Mb Günsberg OO Effingen Mb	
l ă	OX middle	Transversarium	Rotoides / Schilli	— Ox 5 —	Vorbourg Mb Fm Wildegg Fm	
			Luciaef. / Parandieri		Birmenstorf Mb	
		Plicatilis	Antecedens	Ov 4	St. Ursanne Fm OO / Pichoux Fm Birmenstorf Mb	
			Antecedens	0.4	Condensed)	
			Densiplicatum / Vertebr.	— Ox 3 —	O O Bärschwil Fm	

155 Figure 3. Stratigraphic scheme of the middle to late Oxfordian. Lithostratigraphy and biostratigraphy (white circles marking biostratigraphically significant ammonites) according to Gygi (1995, 2000). Sequence boundaries according to Hardenbol et al. (1998) and Gygi et al. (1998). Fm: formation; Mb: member; Kimm: Kimmeridgian. The focus of this study is indicated in gray.

160

The numerical ages of sequences and ammonite zones used in the chart of Hardenbol et al. (1998) are taken from the work of Berggren et al. (1995). The base of the Oxfordian is dated at  $159.4 \pm 3.6$  Ma, its top at  $154.1 \pm 3.2$ Ma, implying a duration of about 5.3 Myr, albeit with wide error margins (Fig. 4). The ages of ammonite zones and sequence boundaries are interpolations between these values. Strasser et al. (2000) and Strasser (2007)

- 165 performed a cyclostratigraphic analysis of Oxfordian platform sections and noticed that the sequences identified by Hardenbol et al. (1998) in many European basins actually correspond to sequences that formed in tune with the 405 kyr long eccentricity cycle, or multiples thereof (Fig. 4). In the Geologic Time Scale (GTS) of 2012, Ogg et al. revised the duration of the Oxfordian, making it last 6.2 Myr. In the GTS of 2020, Hesselbo et al. attributed a still longer time interval of 6.75 Myr to the Oxfordian. An floating time scale is suggested by Huang
- 170 (2018) who counted three 405 kyr cycles between sequence boundaries (SB) Ox 6 and Ox 8, based on the analysis of several deep water sections. The discrepancies in radiometric dating are quite large and preclude a precise attribution of ages to the interval studied here. Furthermore, in the GTS 2020 the sequences boundaries are not dated, and there are differences in the position of sequence boundaries with respect to ammonite zones between the charts of Hardenbol et al. (1998) and Ogg et al. (2012). Nevertheless, the timing of the interval
- 175 between SB Ox 6 and Ox 8 appears to be relatively stable: Hardenbol et al. (1998) attributed 155.81 Ma to SB Ox 6 and 154.63 to Ox 8, implying a duration of 1.18 Myr. Basically the same value (1.2 Myr) was found by Strasser (2007) and Huang (2018). Ogg et al. (2012) dated SB Ox 6 at 159.4 Ma and Ox 8 at 158.2, making the interval also 1.2 Myr long. Consequently, this duration will be applied in the present study.







180

**Figure 4.** Comparison of three different time scales for the Oxfordian, and cyclostratigraphic interpretation of Strasser (2007) and Huang (2018). Note that Strasser (2007) simplified the duration of the long eccentricity cycle to 400 kyr, while Huang (2018) used 405 kyr. For explanation refer to text.

185

In the Oxfordian, the Jura platform was part of the wide passive margin to the north of the Tethys ocean. This margin was a complex array of shallow and deeper marine areas, with several emergent Hercynian massifs that were surrounded by siliciclastic aprons (Carpentier et al., 2006; Fig. 5). The platform was structured by synsedimentary faults (Allenbach, 2001; Strasser et al., 2015). Palaeolatitude was 26 to 27°N (Dercourt et al., 1993).

190 199

The average global climate during Oxfordian times was that of a greenhouse world, i.e. large polar icecaps were absent. Consequently, orbitally induced sea level changes were of low amplitude (see Chapter 1). Late Jurassic climate models have been proposed by, e.g., Weissert and Mohr (1996), Rais et al. (2007), or Louis-Schmid et

195 al. (2007), and a general circulation model has been developed by Sellwood and Valdes (2008). These models, however, only indicate million year trends and do not consider high-frequency climate fluctuations in a restricted palaeogeographic area.







200

(2.00)

Palaeogeographic setting of the Jura platform during the Oxfordian. Modified from Carpentier et al. Figure :

#### 205 3.2 Definition and correlation of depositional sequences

Depositional sequences are here defined in terms of sequence stratigraphy (e.g., Catuneanu et al., 2009). Their formation is controlled by sea level changes. Sequence boundaries (SB) form when sea level drops rapidly, which, on a shallow platform, may lead to subaerial exposure. Lowstand deposits commonly are very thin or 210 absent on the platform because there is no room to accommodate the sediment. A transgressive surface (TS) forms when sea level rise leads to flooding of the platform, and the sediments above this surface then display a deepening up trend (transgressive deposits). The fastest rise of sea level is indicated by the relatively dep

- water fauna and/or condensed deposits if sediment accumulation cannot keep up with sea level rise (M maximum flooding interval). If a hardground develops or a rapid facies change occurs, a maximum flooding 215 surface (MFS) can be defined. Slowing sea level rise then allows the sediment to fill the space created during maximum flooding, and a shallowing up facies evolution is recorded (highstand deposits). A new sea level drop then terminates the sequence. These definitions are independent of the size of the sequences and the time
- involved in their formation (Mitchum and Van Wagoner, 1991; Strasser et al., 1999). Sea level changes may be local or regional (caused by tectonic uplift or subsidence) or global. The latter, eustatic, sea level fluctuations 220 may be long term (million-year scale caused by volume changes in ocean basins as a result of plate tectonics and basalt injections) or short term (caused by orbital cycles; see Chapter 1).

The detailed analysis of facies evolution in the Oxfordian sections studied in the Swiss Jura Mountains allows defining depositional sequences of three different orders, which can be correlated between the sections. An 225 exhaustive sequence- and cyclostratigraphic analysis based on 19 sections covering the middle Oxfordian to late Kimmeridgian interval was carried out by Strasser (100). Here, only three selected sections are presented (Fig.



230



6). The Péry-Reuchenette section was logged by Pittet (1996) and Hug (2003) in a quarry south of the village of Péry-Reuchenette, at the exit of the canyon cutting through the first Jura anticline north of the city of Bienne (Swiss coordinates 585'800/226'250). The Gorge de Court section, studied by Hug (2003), follows a footpath along the Birs River, north of the Village of Court and south of the town of Moutier (Swiss coordinates

- 592'900/223'000). The Hautes-Roches section was analysed by Dupraz (1999) and Stienne (2010). It runs along a forest trail southwest of the Village of Hautes-Roches, north of Moutier (Swiss coordinates 594'950/238'250).
- The biostratigraphy established by Gygi (1995, 2000; see Fig. 3) allows correlating three of the five prominent sequence boundaries shown in Fig. 6 to the boundaries defined in European basins by Hardenbol et al. (1998), i.e. Ox 6, Ox 7, and Ox 8. Assuming that the time interval between Ox 6 and Ox 8 represents about 1.2 Myr (Fig. 4), it is suggested that each of the four prominent sequences represents 400 kyr (termed medium-scale sequences; Stracer, 2007). Each 400 kyr sequence is composed of four small-scale sequences, implying that these lasted 10 (ky) Considering the hierarchical stacking pattern of these sequences and the timing, it is
- 240 proposed that they formed in tune with orbital cycles: 405 kyr for the long eccentricity cycle and 100 kyr for the short one (note that, for simplification, the value of 400 kyr is used for the medium-scale sequences, although they formed in tune with the 405-kyr long eccentricity cycle). These long eccentricity cycles have been found to be dominant also in the early to middle Oxfordian Terres Noires Formation in southeastern France (Boulila et al., 2010). The small-scale sequences are composed of individual beds, the number of which varies between 2
- 245 and 15 per small-scale sequence (Fig. 6). These individual beds are more difficult to attribute to an orbital frequency (obliquity or precession) because, on the shallow platform, also local processes such as lateral migration of sediment bodies could have produced beds, independent of orbitally controlled sea level changes (e.g., Pratt and James, 1986; Strasser, 1991). Also, hiatuses were frequent, hampering a straightforward interpretation of such high-frequency cyclicity (e.g., Sadler, 1994; Strasser, 2015). However, many individual
- 250 beds show a deepening-shallowing facies evolution that can be interpreted as resulting from a sea level cy (e.g., Strasser et al., 1999). Many bed limits are underlined by thin marly layers, which reflect a more humid climate and/or a sea level that washed clays into the sedimentary system. If, in addition, a small-scale sequence is composed of beds, these can be attributed to the 20 kyr precession cycle and are called elementary sequences. Sub-Milankovitch cycles, i.e. cycles with frequencies shorter than those of the precession
- 255 cycle, have been described and interpreted for shallow and deep marine successions (e.g., Zühlke, 2004; de Winter et al., 2014), but there is no evidence for such cycles in the studied Oxfordian sections. In the following, three examples chosen from the sections in Fig. 6 will be discussed in detail.







**Figure 6.** Facies evolution and correlation of three sections logged in the Swiss Jura Mountains (based on Strasser, 2007). Texture terminology according to Dunham (1962) and Embry and Klovan (1971). For explanations refer to text.





265

275

280

### 3.3 Description and interpretation of characteristic sect



#### 270 3.3.1 Gorges de Court

This detailed section (Fig. 7) starts with wackestones of a low energy, protected lagoon, followed by a boundstone composed of corals and oncoids in the Hauptmumienbank (meaning "main mummy bed", referring to the oncoids). Corals indicate warm, oligotrophic conditions, and both corals and echinoderms point to normal marine salinity. Oncoids form in lagoons with high microbial activity. The top of this bed is slightly dolomitized,

suggesting shallow water and semiarid conditions. This is where a first sequence boundary can be placed.

The following part of the section is characterized by terrigenous components (up to 5% quartz, feldspar, and terrestrial organic matter). At the top of the fourth thin bed, birdseyes point to a tidal flat, and a second SB can be placed there. The facies is reddish, and the ooids explain the name of Oolithe rousse Member. This whole interval points to generally low sea level, albeit with high-frequency fluctuations in water depth to explain the

passage from tidal flat to ooid bar. The siliciclastics originate from erosion in the hinterland under a humid climate, and low sea level favoured their progradation into the lagoon. The terrigenous organic matter equally suggests a humid climate allowing for growth of vegetation, in contrast to the dolomite just below.

285

Above follow facies varying between bioturbated packstones (low water energy) and grainstones (high energy). The main components, ooids, oncoids, bivalves, and serpulids, all imply lagoonal conditions. Corals, commonly encrusted by microbialites, become dominant in the middle part of the section, where the individual beds are also thickest (up to 1.8 m). Nevertheless, several bed tops in this interval show dolomitization, suggesting rapid

- 290 changes between sea level rise to accommodate the coral patch reefs under fully marine conditions, and by sea level drops to allow for penecontemporaneous dolomitization on tida . Between meters 15 and 16, even some pseudomorphs after anhydrite are found, pointing to an arid clipped.
- The facies above the coral-rich interval are dominated by oncoids, which formed in protected, low energy 295 lagoons (wackestones, packstones, and floatstones). Tidal flats still occur in the lower part of this interval. Echinoderms and brachiopods point to normal marine salinity. The strongly bioturbated surface at meter 20.6 is interpreted as a maximum flooding surface: no sedimentation for a certain time and several generations of organisms that burrowed through the seafloor. An important change occurs at meter 21.3 of this section: after an intensely dolomitized tidal flat, two 2 to 3 cm thick layers rich in terrigenous organic matter imply the formation
- 300 of marshes and of ponds, in which a marginal marine vegetation could grow. Climate must have changed rapidly from arid or semiarid to humid. Sea level was low and allowed the formation of a major sequence boundary. After this important sequence boundary, sea level rose to accommodate a fully marine lagoon with brachiopods and echinoderms, and then for thick bedded, high energy ooid and peloid bars.
- 305 According to Gygi (1995), the Oolithe rousse Member belongs to the Bimammatum subzone (Fig. 3) and thus also contains SB Ox 7 of Hardenbol et al. (1998). The organic-rich layers at meter 21.3 occur in the upper part of the La May Member, which corresponds to the limit between the Bimammatum and Planut es and thus to SB Ox 8 (Fig. 3). The interval between SB Ox 7 and Ox 8 corresponds to about 400 kyr (F the value of the long eccentricity cycle (405 kyr), and it can thus be postulated that this major sequence observed 310 at Gorges de Court formed in tune with this orbital cycle.

While SB Ox 8 is represented by two thin layers rich in terrestrial organic material testifying to a sea level drop, SB Ox 7 is less well defined. It has been interpreted at the top of the bed with birdseyes (indicating a tidal flat) at meter 6.2, but the entire silicilastic-rich interval implies low sea level. This is why a sequence boundary zone is

315 indicated in Figure 7, corresponding to a time interval that favoured the formation of surfaces that qualify as sequence boundaries (Montañez and Osleger, 1993; Strasser et al., 1999). The same holds for the maximum flooding zone in Fig. 7: thickest beds and most open marine facies define an interval where sea level rose rapidly to create accommodation and fully marine conditions. SB Ox 7 and Ox 8 as well as the maximum flooding zone correlate well with other sections on the Swiss Jura platform (Strasser, 2007; Fig. 6).







325

**Figure 7.** Detail of the Gorges de Court section. Log and facies according to Hug (2003), sequence- and cyclostratigraphic interpretation based on Strasser et al. (2000). Ool. rousse: Oolithe rousse Member. For discussion refer to text.





Between SB Ox 7 and Ox 8, 22 beds are counted (Fig. 7). These beds are separated by thin layers of marls, the clays having been washed into the system during high-frequency sea level drops and/or during phases of more
rainfall in the hinterland. Thus, they qualify as elementary sequences, even if their facies not always display deepening then shallowing up trends (Strasser et al., 1999). 405 kyr divided by 22 results in 18.4 kyr, which is close to the 20 kyr duration of the orbital precession cycle in the Oxfordian (Berger et al., 1992). Considering the complexity of the translation of orbitally induced insolation cycles into the sedimentary system (Fig. 2), and considering the possibility of local processes potentially creating beds, it is not astonishing that the fit is not perfect.

The identification of the small-scale sequences reflecting the short eccentricity cycle of 100 kyr in this outcrop is not easy. Their boundaries are not well developed and are mainly inferred from correlation with other sections (Fig. 6; Strasser, 2007). In Fig. 7, they correspond to rapid changes from high energy grainstone or floatstone

345 facies implying shallower water with waves and currents to low energy wackestone or packstone facies formed below wave base (thus implying an increase in water depth). Following these criteria, 4 such sequences are counted within the 400-kyr sequence. The small-scale sequences are composed of 4 to 6 individual beds. Again, the perfect fit would have been 5 beds per sequence but, again, the sedimentary system was complex and did not always translate the orbital cyclicity one-to-one.

### 350

#### 3.3.2 Hautes-Roches A

- In the Hautes-Roches section, two coral-rich intervals are encountered (Fig. 6; Dupraz, 1999). The lower one (Fig. 8) corresponds to a small-scale sequence that formed in tune with the 100 kyr eccentricity cycle. In this outcrop, the boundaries of the sequence are defined by sharp transgressive surfaces (TS), which superpose crossstratified ooid grainstones over lagoonal sediments (the sequence boundaries are hidden in the sediments just below the TS, or amalgamated to them; Strasser et al., 1999). This suggests that, at the beginning of a sea level rise, ooid shoals migrated over lagoons. The bed above the first ooid shoal still contains ooids but also coral
- 360 debris, suggesting that a nearby coral reef was being dismantled by storms. Healthy corals then grew on top of the rubble, implying increasing water depth. Later, the corals adopted a flat shape, possibly because they had to increase their surface to catch enough light for the photosynthetic zooxanthellae, which again points to increasing water depth. Coral diversity was high (mainly *Microsolena, Heliocoenia, Stylina, Isastraea*, and *Calamophylliopsis*; Dupraz, 1999). The corals gradually became covered by microbialites, which implies
- 365 increased nutrient input and eutrophication of the system (Hallock and Schlager, 1986; Dupraz and Strasser, 2002). A sharp surface terminates this bed. Above, coral growth sets in again, but the corals are covered by microbialites, and oncoids are present. Especially the Microsolenids are associated with microbialites, suggesting that this family supported some degree of eutrophication. A washover deposit rich in echinoderms interrupted the growth of this reef, implying that a major storm passed over it. Above a last, large coral head
- 370 (probably not in situ but toppled) and flat corals, only microbialites are present; they have a thrombolite texture and include some sponges. This sequence, from ooid shoal through coral reef to thrombolite, suggests that water energy decreased through time, that the water became deeper and more turbid to hamper the photosynthesis of the coral symbionts, and that nutrient excess favoured microbial growth that eventually suffocated the corals.
- 375 This evolution ends with a bioturbated, nodular layer capped by a bored hardground, indicating that the sea floor was exposed over some time to become cemented and perforated by lithophaga bivalves. The hardground also marks the end of a deepening up trend and is thus considered as a maximum flooding surface (MFS). Above the hardground, brachiopods indicate normal marine salinity. Quartz sand and (in the upper part of the bed) plant debris testify to terrigenous input and progradation of lagoonal sediments. This is interpreted as shallowing up
- 380 highstand deposits. A thin bed with echinoderms (possibly a tempestite) and a new ooid shoal then represent the beginning of a new sequence.







385

Figure 8. Detail A of the Hautes-Roches section. Log and facies based on Dupraz (1999). For discussion refer to text.

- 390 The sequence is composed of 7 beds. One, however, is an event deposit (the washover), and one belongs to the composite hardground covering the thrombolite thus suggested that the formation of each bed was controlled by the orbital precession cycle of 20 The bed limits are transgressive surfaces or maximum flooding surfaces; the sequence boundaries are not developed (Strasser et al., 1999). The washover deposit that served as a substratum for renewed coral growth is attributed to elementary sequence 4, but the storm event may
- 395 also have occurred at the end of sequence 3. During the first four precession cycles, conditions were those of a gradual transgression, which ended in the hardground. The highstand deposits took only 20 kyr to form, making the record of this 100 kyr cycle highly asymmetrical.





#### 3.3.3 Hautes-Roches B 405

The second interval at Hautes-Roches shown here is dominated by lagoonal marls. The section illustrated in Fig. 9 starts with a firmground that covers lacustrine mud- to wackestones with charophytes. Above, a first thin bed still has lacustrine facies. The following thin bed contains charophytes as well as marine fossils, implying a marine, transgressive pulse that reworked lacustrine elements. A second transgressive surface then leads to a

- 410 strongly bioturbated lagoon. The next two meters contain echinoderms and brachiopods, implying normal marine salinity. Three nodular beds are found in this interval. They are thickening up, and quartz content increases upwards to up to 20% (Stienne, 2010). This evolution is typical of a shallowing upwards highstand deposit. In this case, the maximum flooding interval can be placed at the strongly bioturbated bed, while a sequence boundary is identified at the base of the third nodular bed (Fig. 9). Five elementary sequences are
- 415 identified, leading to the hypothes for the correspond to the 20 kyr precession cycle, and the small-scale sequence they make up to the 100 for the ccentricity cycle. Each of the elementary sequences has a lower, carbonate-rich part interpreted as transgressive deposits and a siliciclastic-rich upper part that represents the highstand deposits.
- 420 The following 1.2 meters are composed of lagoonal marls with a fully marine fauna. They are covered by a sharp transgressive surface, above which grew a reef composed of small corals. This transgressive surface can be followed over the entire Swiss Jura platform (Védrine and Strasser, 2009; Fig. 6). It also corresponds to the turn-around from a long-term regressive trend that began in the early Oxfordian to a long-term transgressive trend that ended in the late Kimmeridgian (Hardenbol et al., 1998). This TS marks the base of the coral- and ooid-rich
- 425 Steinebach Member, respectively of the oncoid-bearing Hauptmumienbank Member (Fig. 3). The corresponding sequence boundary is hidden in the marls below the TS, or amalgamated to it. Elementary sequences cannot be recognized.
- Reef development in this section was short-lived. Above the thin reefal boundstone with low coral diversity (mainly *Isastrea* and *Microsolena*; Dupraz, 1999), only reworked coral clasts appear that, however, suggest that reef growth continued in the neighbourhood. The phase of reef development was then interrupted by ooid shoals, one of which displays sigmoidal cross-beds formed by tidal currents.





	Hautes-Roches B			
Sequence- and cyclostratigraphy		Facies	Depositional environment	
3		ooid grainstone with sigmoidal structures	ooid shoal with tidal sandwave	
		ooid grainstone		
2		reworked corals (rudstone)	storm and wave reworking	
1		reworked corals (rudstone)	storm and wave reworking	
тс	The Weller	small corals (boundstone)	reef installation	
?		lagoonal marls with diversified fauna	lagoon with clay input	
SB ——		mudstone with quartz, echinoderms, and forams more diversified, open-	sandy lagoon	
5	⊕ ⊕ ★	(brachiopods)		Corals
4	*	lagoonal marls with agglutinated foraminifera	lagoon with	<ul> <li>coral clasts</li> <li>ooids</li> <li>brachiopods</li> </ul>
_	*	and ostracodes	clay input	★ echinoderms
	8 8 8 8			<ul> <li>bivalves</li> <li>asstrenads</li> </ul>
3	2 1 2 2 2 1 m		bioturbated lagoon	gastropous     senthic foraminifera
MF		lagoonal wackestone with Thalassinoides burrows		O charophytes
2		marls with mixed lacustrine and marine fossils		ని bioturbation
1				marls and quartz sand
SB/TS —	0 ( 200000	Fe-rich surface	firmground	TS transgressive surface
	0	wackestone	lake	SB sequence boundary

#### 435

Figure 9. Detail B of the Hautes-Roches section. Log and facies based on Dupraz (1999) and Stienne (2010). For discussion refer to text.

#### 440

#### 4 Discussion

445 Although the climate factors influencing a shallow marine carbonate platform are intimately linked (Fig. 2), they will be discussed here separately.

# 4.1 Water temperature 450

In ancient carbonates, oxygen isotope ratios are often used as a proxy for the temperature of the water in which they formed. This, however, requires that the original mineralogy be preserved in the fossil record. In shallow marine carbonates, this is rarely the case: early freshwater diagenesis dissolves the aragonite of corals, gastropods, and green algae, and the high-Mg calcite of echinoderms is transformed intervelop. An

- gastropods, and green algae, and the high-Mg calcite of echinoderms is transformed in the bw-Mg calcite. An
   exception are brachiopod or bivalve shells, which have a relatively stable low-Mg calcite. An
   found in the studied sections. However, the δ<sup>18</sup>O values that are recorded also depend on the geochemistry of the
   platform waters, which may vary considerably across a complexly structured platform such as one in the Jura
   Mountains (e.g., Immenhauser et al., 2003; Colombié et al., 2011).
- 460 Among the tropical and subtropical carbonate-producing organisms, corals are the most sensitive to temperature changes. Today, reef-building zooxanthellate corals do not support water temperatures below 18°C and above 34°C (Wood, 1999). For short periods, however, they tolerate fluctuating and even higher temperatures (e.g.,



465



Schoepf et al., 2015). Aragonite saturation in seawater, which is important for efficient coral skeleton formation, closely follows water temperature (Wood, 1999). If it is assumed that Oxfordian corals had the same requirements regarding temperature as the recent ones, a high-diversity reef would be a proxy for water temperatures ranging from about 23 to 29°C. In the middle and late Oxfordian, coral reefs were widely distributed between the palaeolatitudes of 40°N in Japan to 45°S in the Neuquén Basin (Leinfelder et al., 2002) where these water temperatures were common. The reef in the Hautes-Roches section (Fig. 8) with a relatively high coral diversity may thus indicate water temperatures between 23 and 29°C. Low diversity, however, may

- 470 not only be an effect of temperature but also of other unfavourable conditions such as too shallow or too deep water, high sediment input that makes the water turbid and thus hinders photosynthesis of the symbionts, nutrient excess that favours microbialites and fleshy algae that compete with the corals, or water acidification that impacts on skeletal growth.
- 475 Based on a mean value of -3.50%  $\delta^{18}$ O in micritic whole rock (which averages the isotopic signal of the different components) from middle Oxfordian sections in the Swiss Jura, Plunkett (1997) calculated ambient water temperatures of 26 to 27°C (following Anderson and Arthur, 1983). This compares well with Frakes et al. (1992) who indicate up to 27°C for late Oxfordian ocean surface temperatures, according to oxygen-isotope ratios measured on planktonic foraminifera and belemnites. Martin-Garin et al. (2010) measured oxygen-isotope
- 480 ratios on brachiopods and oysters from outcrops in the Swiss Jura. They calculated water temperatures of below 20 °C for the low-diversity coral reefs (15 genera) in the Liesberg Member (Fig. 3), of about 25°C for the highdiversity reefs (37 genera) in the St. Ursanne Formation, and of about 20°C in the Günsberg Member where coral diversity declined (13 genera). The reef complex shown in Fig. 8 is situated in this latter member.
- 485 For the carbonate platform in the French Jura, Olivier et al. (2015) interpreted a climate evolution going from warm and humid to warm and arid in the course of the Bimammatum zone. This corresponds to the time interval between sequence boundaries Ox 6 and Ox 8 discussed here (Fig. 3). In the Paris basin, Brigaud et al. (2014) cite sea surface temperatures of 20 to 29°C for the Late Jurassic, with an optimum during the middle Oxfordian. At the onset of the late Oxfordian, water temperatures dropped by 6 to 7°C, and rainfall increased. Carpentier et al.
- 490 (2010) state that, in the eastern Paris Basin, after a warm and arid interval favouring carbonate production in the Transversarium zone, climate became cooler and more humid in the Rotoides and Stenocyloides subzones. In the southern North Sea, Abbink et al. (2001) analysed sporomorphs as palaeoclimate proxies and showed that there was a temperature drop between the boreal Serratum and Regulare ammonite zones. This corresponds to the Semimammatum subzone in the Tethyan realm (Fig. 3), and thus to the marl-rich interval above sequence 495
- boundary Ox 6 (Fig. 6).

#### 4.2 Sea level

- 500 Sea level changes in the Milankovitch frequency band are an important proxy as they reflect changes in global air temperature. During the Quaternary, glaciation cycles controlled by the 100 kyr short eccentricity cycle led to important, asymmetrical sea level fluctuations due to slow buildup of ice caps followed by rapid melting (Shackleton, 1987; Fig. 11). In greenhouse worlds, i.e. when major ice caps were absent, the translation of orbitally controlled insolation changes into eustatic sea level changes was more direct. Thermal expansion and
- 505 retraction of ocean water was an important factor controlling sea level fluctuations in greenhouse worlds, in tune with the eccentricity cycles of 100 and 405 kyr (Sames et al., 2014). Schulz and Schäfer-Neth (1997) proposed that thermally induced volume changes in deep-water circulation resulted in sea level changes. Water storage and release on the continents (aquifer eustasy; Jacobs and Sahagian, 1993; Sames et al., 2014; Wendler and Wendler, 2016) depends on the climate over the land areas: sea level rise is linked to a dryer climate with less
- 510 precipitation, when aquifers empty more rapidly than they fill up. Consequently, aquifer eustasy is dependent on the latitudes where the land areas occur and on the atmospheric circulation cells, i.e. it is controlled also by the obliquity and precession cycles. Obliquity is dominant at high latitudes and cannot be identified in the studied sections. The tyr precession cycles, on the other hand, are recorded in the elementary sequences forming individual b
- 515

For Cretaceous high-frequency sea level cycles, Davies et al. (2020) proposed the combined effects of thermal expansion, glacioeustasy, and aquifer eustasy to produce the 65, 30, and 50 m amplitudes observed in the Valanginian, Turonian, and Maastrichtian, respectively. Aquifer eustasy taken alone would be responsible for amplitudes of less than 5 m, which is in the range of the reconstructed Oxfordian amplitudes (see below).

Wendler et al. (2016) suggested that aquifer eustasy could drive sea level changes even in the order of 30 to 40 520 m, and this at the frequency of the long eccentricity cycle that shifted the Hadley circulation and the width of the equatorial humid belt.





For the part of the Gorges de Court section shown in Fig. 7, Strasser (2018) reconstructed a sea level curve that explains the observed stacking pattern of beds. For this, the section was first decompacted according to facies (Goldhammer, 1997; Strasser and Samankassou, 2003). The following decompaction factors are applied: 1.2 for grainstones, 1.5 for packstones, 2 for wackestones, 2.5 for mudstones, and 3 for marls. Boundstones have resisted compaction and a factor of 1.2 is used. Floatstones with their micritic matrix are treated like packstones. Estimation of water depth for the different facies of course is not easy. For this exercise, the following values

- 530 have been assumed: Tidal flats are set at zero meters. Grainstones often result from winnowing by waves and currents, and a water depth of 1 m is attributed. Packstones and floatstones the abundant fauna are set at 2 m. Coral patch reefs potentially can grow up to sea level but are assumed to have been assumed at least 2 m water depth to develop. When an open marine fauna (e.g., brachiopods) is present, a minimum water depth of 3 m is assumed. To accumulate the sediment pile between Ox 7 and Ox 8 (15.3 m today, 26 m when decompacted; Strasser,
- 535 2018), a gain in accommodation of 26 m is needed (Fig. 10). This is created by subsidence and long-term sea level rise. This long-term trend can be subdivided into 4 portions, according to the space needed to accumulate the individual small-scale, 100 kyr sequences. Deducing these trends from the reconstructed sea level curve, the deviation of sea level change from the average accommodation gain can be visualized for each 100 kyr sequence. It is interesting to note that, for the medium-scale sequence, two candidates for a maximum flooding
- 540 zone can be proposed, both situated in the interval of high accommodation gain (Fig. 10). However, when comparing with other sections in the Swiss Jura, it is rather the lower one that is recognized all over the platform (Strasser, 2007; Fig. 6 with the example of the Péry-Reuchenette section).
- 545 The small-scale sequences 3 and 4 are well defined and each contains 5 elementary, 20 kyr sequences. Sequence 545 boundary Ox 7, which defines the limit between small-scale sequences 0 and 1, is placed at the tidal flat. This results in only 4 elementary sequences for small-scale sequence 0, and 6 for small-scale sequence 1 (Fig. 10). This misfit may be explained by a wrong definition of the elementary sequences, or else by other factors but sea level that created a topographic high to install the tidal flat (differential subsidence, lateral migration of sediment bodies). The top of the second small-scale sequence (elementary sequence 14 in Fig. 10) shows corals,
- 550 microbialites, and lithophaga borings but also evaporites, pointing to rapid changes in water depth. It is possible that the 5<sup>th</sup> elementary sequence was not recorded due to lack of accommodation.

The next step is to reconstruct the amplitudes of the highest-frequency sea level fluctuations corresponding to the 20 kyr precession cycle by subtracting the 100 kyr trends (Fig. 11). When compared to the envelope curve based on the calculated insolation changes at the equator and for the last 500 kyr, a partial correspondence becomes

- 555 on the calculated insolation changes at the equator and for the last 500 kyr, a partial correspondence becomes visible: stronger insolation due to the 100 and 405 kyr cyclicity generally translates into higher sea level amplitudes for the 20 kyr cycles. Even considering the complex facies distribution on a shallow marine platform and accommodation changes due to differential subsidence, and even considering the uncertainties in estimating decompaction factors and water depth, it can be concluded that, in the Oxfordian case discussed here, orbitally
- 560 controlled insolation changes translated into sea level changes. The tively good correspondence seen in Fig. 11 suggests that the translation was mainly via thermal expansion betraction of the ocean water. Aquifer eustasy and low-amplitude glacioeustasy certainly were present but are difficult to quantify in the present study.







565

570

**Figure 10.** Reconstruction of high-frequency sea level changes for the Gorges de Court section (modified from Strasser, 2018). The section (Fig. 7) is first decompacted according to facies, and for each facies water depth is estimated. Elementary sequences are assumed to correspond to the 20 kyr precession cycle, small-scale sequences to the 100 kyr one. Average accommodation gain needed to record the small-scale sequences varies through time. By deducing these trends, a sea level curve is reconstructed that serves as input to Fig. 11. Symbols and abbreviations as in Fig. 6.







575

**Figure 11**. Comparison between the high-frequency, meter-scale sea level changes reconstructed for the Gorges de Court section with the insolation curve calculated for the equator (Hinnov, 2018). The envelopes of the two curves (green lines) reflect the 100 and 405 kyr eccentricity cycles, suggesting that the sedimentary record at Gorges de Court was at least partly controlled by insolation cycles. During icehouse times, the sea level changes had a much higher amplitude and were asymmetrical ( $\delta^{18}$ O values serving as proxy for sea level changes; simplified from Railsback et al., 2015; MIS: marine isotope stage). For discussion refer to text.

580

It has to be kept in mind that the precession cycle today has a 21/23 kyr periodicity, while in the Oxfordian the periodicity was at about 20 kyr (Berger et al., 1992; Hinnov, 2018). The number of high-frequency cycles in Fig. 11 therefore is not the same for the Quaternary and Gorges de Court. On the other hand, the 100 and 405 kyr cycles are stable through time and can be directly compared. A synthesis of 12 Oxfordian sections in the Swiss Jura showed that many small-scale sequences controlled by the 100 kyr short eccentricity cycle are composed of a lower, transgressive part dominated by high energy, carbonate dominated facies (e.g., ooid or bioclastic shoals,

coral reefs), while the upper, regressive part displays low energy carbonate facies (e.g., tidal flats) or marls (Strasser et al., 2015). The conclusion is that during a warming climate sea level rise pushed back the terrigenous delivery, and clear, oligotrophic, warm and deepening waters allowed for carbonate production and accumulation. During a cooler climate, sea level dropped, siliciclastics prograded, water depth diminished, and carbonate production was reduced. Cooler climate coincided with more rainfall (see below), thus reinforcing erosion on the land areas.

Pittet and Gorin (1997) analysed the palynofacies in the Vellerat Formation (Fig. 3), They found that the ratio between land-derived particulate organic matter (plant debris, pollen and spores) and marine particles (phytoplankton, foraminifer linings) reflects proximal to distal trends, which depended on the land-ocean configuration but clear on each local barries of the Courts action (Fig. 7), rateful arthur to a set of the courts of the courts

600 configuration but also on sea level changes. In the Gorges de Court section (Fig. 7), reefal carbonates are also associated with high eccentricity, albeit at the 405 kyr scale (Fig. 11). The analysis of associations of benthic foraminfera at the Oxfordian-Kimmeridgian boundary in the Paris Basin by Lefort et al. (2011) revealed a correlation with sea level changes: a highly diversified association composed of small agglutinated and calcitic forms characterizes high sea level stands, while an association richer in large agglutinated foraminifera is solured with each level level level that there is no indication of the sea level and calcited by the sea level level level that the sea level stands.

605 abundant when sea level was low. Unfortunately, there is no indication of the duration of the sea level cycles.



610



One degree Celsius of global warming today leads to 20 to 60 cm of sea level rise through thermal expansion of the ocean water (Church and Clark, 2013). If the same values are assumed for Oxfordian times, and if thermal expansion is considered as the main factor, a sea level rise of 2 m would suggest an atmospheric warming of 3 to 10 degrees. Insolation by itself would not have allowed for this warming, but its input was enhanced by feedbacks due to greenhouse gases. For example, Ikeda et al. (2020) suggested that ca. 10 Myr orbital variations controlled summer monsoon dynamics, which changed terrestrial weatherability, affected atmospheric *p*CO<sub>2</sub>,

and eventually led to variations in sea surface temperatures in the order of 3 to 7°C. At the scale of obliquity cycles, Laurin et al. (2015) explained the  $\delta^{13}C_{earb}$  signature in the Late Cretaceous English Chalk by recycling of carbon through formation and decay of reservoirs of organic matter and/or methane on land, in lakes, and in euxinic marine settings.

It is interesting to compare rates of sea level rise between the Oxfordian and today. Taking two examples from Fig. 10, it can be estimated that, to accommodate elementary sequence 5, sea level must have risen by about 2 m within 10'000 years (half a precession cycle), and by about 3 m for elementary sequence 9. This amounts to

- 620 within 10'000 years (half a precession cycle), and by about 3 m for elementary sequence 9. This amounts to values of 0.2 mm per year for the first case, and of 0.3 mm/y for the second one (under the condition that sea level change was mainly controlled by thermal expansion of the ocean water and thus followed a symmetrical curve). Based on the detailed analysis of one elementary sequence in the Röschenz Member, Strasser et al. (2012) estimated a rise of 0.3 mm/y during maximum flooding, i.e. during the fastest rise of the sea level cycle.
- Today, due to anthropogenically induced global warming, global mean sea level rose by 3.7 mm/y between 2006 and 2018 (Masson-Delmotte et al., 2022), i.e. more than ten times faster than in the Oxfordian.

#### 630 4.3 Rainfall

Terrigenous material (clays, quartz, feldspar, terrestrial organic matter) appear at regular intervals in the studied sections. Thin, only mm- or cm-thick marl layers commonly the limits between individual beds, and several meter-thick intervals around sequence boundaries Ox 7 are dominated by siliciclastics and

- 635 contain plant fragments. The maximum flooding zone at meter 40 in the Pery-Reuchenette and Gorges de Court sections also contains clays and 1% of quartz (Fig. 6; Hug, 2003). According to the sequence stratigraphic interpretation, these intervals commonly correspond to low sea level stands, with the exception remaining flooding at meter 40. Falling sea level causes erosion of the hinterland, i.e. of the Hercynian m (Fig. 5). Deltas prograde, and terrigenous material is being delivered into the lagoons. A second factor of course is
- 640 rainfall in the hinterland, which activates the rivers that transport the siliciclastics and plant fragments towards the ocean. Input of aeolian quartz under an arid climate cannot be excluded, but transport by rivers is considered to have been the main process (Gygi and Persoz, 1986; Hug, 2003).

From Fig. 6 it is clear that the terrigenous material is not recorded equally in the studied sections. This can be explained by the variable platform morphology including highs where the siliciclastics bypassed and lows where they were ponded (e.g., Strasser, 2018). Furthermore, currents picked up the clays and redistributed them over the platform.

- At a palaeolatitude of 26 to 27°N (Dercourt et al., 1993), the studied platform was situated in the subtropical zone, which is confirmed by the diversity and type of carbonate-producing organisms typical of such platforms (Reijmer, 2021). At this latitude and at the insolation maximum, atmospheric circulation was controlled by the Hadley cell, while at the insolation minimum, the Ferrel cell was active (e.g., Matthews and Perlmutter, 1994). Orbital cyclicity thus caused latitudinal shifts of atmospheric cells and shifts between high pressure and low pressure zones. For example, Khon et al. (2012) showed how insolation changes at the 100 kyr scale between the
- 655 last interglacial (Eemian; MIS 5e) and the early Holocene controlled feedback loops between sea surface temperatures, easterly winds, water evaporation, shifting of the Intertropical Convergence Zone, and circulation patterns of the Hadley and Walker cells. These insolation changes thus indirectly controlled rainfall at a given latitude. Cecca et al. (2005) documented latitudinal shifts of climate belts based on the distribution of coral reefs and ammonites during the Oxfordian. Following the General Circulation Model developed by Sellwood et al.
- 660 (2000), they concluded that the late Oxfordian was characterized by low rainfall in the Tethyan Mediterranean area (0.5 to 2 mm per day) and slightly higher rainfall (2 to 4 mm per day) in central and northern Europe.

Orographic effects modifying the atmospheric circulation were probably minor as there was no high mountain range near the Jura platform but only widely dispersed, low-relief massifs (Fig. 5). Insolation minima





665 corresponded to low sea levels (Fig. 10), and this explains that many sequences boundaries in the studied sections contain terrigenous material that was eroded from the hinterland under humid conditions.

More arid conditions prevailed during transgression and maximum flooding, when climate was warmer and sea level rose, allowing for the growth of coral reefs under oligotrophic conditions. However, nutrients were washed

- 670 into the system, which stimulated microbial growth that eventually smothered the corals, as illustrated in Fig. 8. While corals could still grow at the beginning of the second 20 kyr sequence, they were suffocated by microbialites at the of this sequence when eutrophication increased. Towards the end of the transgressive part of the 100 kyr ence, eutrophication was such that corals could not grow anymore. This evolution suggests that nutrients were washed in from the hinterland and affected the reef during transgression and
- 675 maximum flooding, before the siliciclastics arrived and covered the coral-microbial buildup. Eutrophication can also be caused by increased upwelling bringing nutrients from deep to shallow waters. However, considering the position of the Jura platform far from the open Tethys ocean (Fig. 5), this scenario is less likely.
- Védrine et al. (2007) showed that the morphology of oncoids in the Hauptmumienbank Member varied according to water depth, water energy, and terrigenous nutrient influx, all of these related to climate changes induced by the 100 and 20 kyr orbital cycles. Analysing coral-microbialite relationships such as those illustrated in Fig. 8, Dupraz (1999) and Dupraz and Strasser (2002) came to the same conclusion. On a ramp at the southern margin of the French Jura platform, Olivier et al. (2011) described abundant microbial carbonates (oncoids and coral-microbialite reefs) that formed during the Bimammatum zone, i.e. during to the interval shown in Fig. 6.
- This is explained by a humid and relatively cool climate, by terrigenous nutrient input, and by mesotrophic conditions.
- On the Aalenian to Oxfordian platform in western France, Andrieu et al. (2016) found that high carbonate productivity correlated with high eccentricity intervals, which promoted a dry climate and oligotrophic conditions for the carbonate-producing organisms. However, they referred to the ca. 9 Myr eccentricity cycle evidenced by Martinez and Dera (2015) in Jurassic to Early Cretaceous Tethyan sections.

A rapid shift from a cool and humid climate to warmer and more arid conditions is recorded in the North Sea at the end of the early Oxfordian (Densiplicatum subzone) and may have been of global extent, related to

- hydrothermalism at the onset of rifting in the North and South Atlantic (Abbink et al., 2001). From the middle Oxfordian to the Kimmeridgian, the general evolution was towards warmer and more arid conditions (Olivier et al., 2015; Carpentier et al., 2010). However, at the beginning of the late Oxfordian, rainfall increased in the Paris basin (Brigaud et al., 2014). In the Swiss Jura, this corresponds to the interval around sequence boundary Ox 6 where siliciclastics and terrigenous organic matter are present (Fig. 6). Superimposed on these general trends, rainfall patterns were controlled by the orbital cycles. Above sequence boundary Ox 6, siliciclastics
- 700 rainfall patterns were controlled by the orbital cycles. Above sequence boundary Ox 6, siliciclastics predominate. A well developed medium-scale sequence controlled by the 405 kyr long eccentricity cycle then leads to sequence boundary zone Ox 7, which again is rich in siliciclastics. Arid conditions are indicated by evaporites in the middle of the next medium-scale sequence (Fig. 7). SB Ox 8 contains only a thin interval of terrigenous material, thus confirming the general trend towards arid conditions in the early Kimmeridgian
- 705 (Abbink et al., 2001). An interesting issue is the presence of siliciclastics at the maximum flooding of the medium-scale sequences (e.g., at the top of the Steinebach Member at meter 40; Fig. 6). Either there was a pulse of rainfall related to the northward expansion of the Intertropical Convergence Zone (e.g., Khon et al., 2012), or else sea level was high enough to reach deep into the land areas and mobilize terrigenous material there.
- 710 In the Hautes-Roches and Gorges de Court sections, siliciclastics and plant material are present below sequence boundary Ox 6 (Fig. 6). Marls become important above Ox 6 in all three sections shown in the figure, up to the major transgressive surface marking the base of the Steinebach Member. This implies rainfall over the emerged land areas. The same interval also corresponds to the end of the large-scale regressive trend documented in European basins by Hardenbol et al. (1998), meaning that sea level was generally low, that more land was
- 715 exposed to erosion, and that terrigenous material was forced to prograde into the ocean. More exposed land areas also led to more continental weathering, potentially drawing down atmospheric CO<sub>2</sub> levels. This has been modelled for million-year time scales by, e.g., Goddéris et al. (2012), and the feedbacks between weathering and atmospheric CO<sub>2</sub> have also been discussed for future climate change (e.g., Beaulieu et al., 2012). In hemipelagic carbonates in the Jura Mountains and the Vocontian basin, Padden et al. (2002) found negative δ<sup>13</sup>C excursions
- 720 in the Transversarium and Bifurcatus ammonite zones (i.e. below SB Ox 6; Fig. 3), which are explained by release of methane introducing light <sup>12</sup>C into the marine carbon reservoir. If methane was released also into the atmosphere, it may have contributed to greenhouse effects that led to warming and increased precipitation over the land areas, and thus to more siliciclastic input. The general cooling trend due to CO<sub>2</sub> drawdown may thus have been punctuated by short warming periods, but humidity was maintained.





725

Clay minerals also hold palaeoclimatic information, and the ratio between kaolinite and illite is commonly used to define humid versus arid conditions during soil formation: a high K/I ratio generally indicates a warmer and more humid climate (Curtis, 1990; Thiry, 2000). On the Jura platform, kaolinite becomes dominant over illite above sequence boundary Ox 6 (Gygi and Persoz, 1986). However, rates of soil formation are slow, and the time

- 730 resolution of the clays deposited in the ocean therefore is considered to be low (1 to 2 Myr; Thiry, 2000). Furthermore, clays may be stored in deltas, ponded in lagoons, and distributed across the platform at different times. Thus, the climate information they contain may not correspond to the time at which they were finally recorded at a given position on the platform. Nevertheless, Védrine (2007) and Strasser et al. (2012) found that in the Hauptmunienbank Member the K/I ratio increases slightly around sequence boundaries and maximum
- 735 flooding intervals of the small-scale, 100 kyr sequences. At the sequence boundaries, kaolinite is associated with quartz. This implies that climate was humid during low sea level, and quartz and clays were eroded from the hinterland. At maximum flooding, with the ocean encroaching on the land areas, only clays were mobilized and transported across the platform. As discussed above, there might also have been a humid phase during maximum flooding, related to the northward expansion of the Intertropical Convergence Zone.

740

The importance of the palaeolatitude in the distribution of rainfall patterns is illustrated by the comparison of Oxfordian sequences in the Swiss Jura and in the Soria region of Spain (Pittet and Strasser, 1998). In the small-scale, 100 kyr sequences in the Swiss Jura, siliciclastics are commonly concentrated around the sequence boundaries, while in Spain they occur preferentially in the maximum flooding intervals. The sections studied in

745 Spain were at a palaeolatitude of 23 to 24°N (Dercourt et al., 1993), i.e. 3 to 4° farther south than the Jura sections. This latitudinal difference may explain that the humid periods during a 100 kyr cycle did not occur at the same time.

#### 750

#### 4.4 Wind and storms

During the Oxfordian, the Jura Platform was situated in the subtropical zone and under the influence of trade winds. These would have forced currents into a westerly direction, but the presence of several emergent land masses certainly deviated these currents. Nevertheless, considering the palaeogeography shown in Fig. 5, it is probable that the siliciclastics found on the Jura platform stemmed mainly from the Bohemian Massif. This massif lies on a more northern palaeolatitude, which implies that rainfall could have increased weathering and erosion there, while on the Jura platform more arid conditions prevailed. This may explain the direct superposition of marly layers on a dolomitized carbonate facies (Fig. 7). Nutrients in a dissolved state travelled faster across the platform and nourished the microbialites before the siliciclastics arrived (Fig. 8).

The cross-bedded ooid and bioclastic shoals encountered in the studied outcrops were mainly controlled by tidal currents, the direction of which varied according to the platform morphology.

- 765 Storms episodically affected the platform. Being far away from the open ocean, fetch was reduced. Nevertheless, the presence of reef rubble and washovers (Fig. 8) testifies to currents and waves strong enough to dismantle a reef and throw sediment over topographic highs. As it is the case today with global warming, storm frequency and intensity were likely to increase when water temperature increased (Masson-Delmotte et al., 2022). Pittet and Gorin (1997) noted an increase of tempestites and washovers in the marly interval above sequence boundary
- 770 Ox 6, which corresponds to the highstand of the corresponding 405 kyr sequence (Fig. 6). The abundance of siliciclastics in the interval, however, implies humid conditions. Possibly, a preservational bias has to be considered: in very shallow water or in the intertidal zone, only thin layers of sediment are deposited during a storm (as for example on the tidal flats of the Bahamas; Rankey et al., 2004). In deeper water, however, there is room to accommodate the reworked sediment. In the Kimmeridgian of Aquitaine (France), limestone-marl
- 775 alternations are regularly interrupted by thin tempestite layers (Colombié et al., 2018). These authors show that tempestite occurrence (and thus storm frequency) was highest during maximum flooding intervals of the sequences controlled by the 405 kyr long eccentricity cycle. Maximum flooding relates to rapid sea level rise forced by a warming of the climate.





#### 785 4.5 Combined climatic effects at orbital scale

From the discussion above it becomes clear that the sedimentary record of shallow marine carbonate systems is controlled by several, interdependent climatic processes. For the Oxfordian in the Swiss Jura, two models are proposed to explain the interactions of climate drivers and the resulting sedimentary record. The shortest

- 790 decipherable time scale is chosen, i.e. a sea level cycle related to the 20 kyr orbital precession cycle. To simplify, the cycle is assumed to have been symmetrical, but asymmetries due to processes mentioned above certainly were present. Accommodation gain is the sum of subsidence and long-term sea level rise. Subsidence in the Late Jurassic of the Jura platform was about 3 m/100 kyr (Wildi et al., 1989). Long-term sea level rise is the sum of the example in Fig. 10: 405 kyr to accumulate 26 m of decompacted sediment results in 6.4 m/100 kyr fotal
- 795 accommodation gain. This value minus the subsidence gives 3.4 m/100 kyr long-term sea level rise. This is consistent with the major transgressive evolution in European basins from the Oxfordian to the Kimmeridgian, which started in the Semimammatum subzone with the major transgressive surface above sequence boundary Ox 6 (Hardenbol et al., 1998).
- 800 Model A (Fig. 12) describes the hypothetical formation of an elementary sequence controlled by the 20 kyr precession cycle. This cycle is situated in the middle of a 100 kyr and/or 405 kyr cycle when insolation was high (Fig. 1). It is thus expected that sea level amplitudes on the 20 kyr scale were relatively high, in the order of a few meters (Figs 10, 11). For the model, an amplitude of 3 m is chosen, and it is assumed that base and top of the elementary sequence are at zero sea level. At the beginning of the cycle, climate at the given subtropical
- 805 latitude is humid, sea level is low, siliciclastics are eroded from the emergent massifs, transported across the platform, and accumulated at the base of the sequence. Accommodation is low, and only a thin sediment layer can be recorded. With increasing insolation, the climate becomes progressively dry, the water becomes warmer and expands to rise sea level, coastlines retreat, rivers are less active, and siliciclastic input is reduced. At still low sea level, a tidal flat forms and can keep up for a while with rising sea level before it is drowned. With
- 810 increasing water depth, conditions become ideal for carbonate production, and high accommodation gives room for the accumulation of ooid shoals and the growth of coral reefs. Of course, depending on the position on the platform, lateral facies variations are considerable: while at one point reef growth can keep up with rising sea level, at another point a deep lagoon develops (Fig. 2). The demise of the hypothetical coral reef could be due to eutrophication like in the example of Fig. 8, or to smothering by sediment that progrades because sea level rise
- 815 slows down. Still under a warm and arid climate, sea level rise slows down and allows the sediment to fill in the available space. If sea level then drops below the sediment surface and climate becomes humid again, a karst develops. If the sediment surface stays in the intertidal or supratidal zone as in Fig. 12A, evaporites form in a sabkha under an arid climate. When insolation decreases in the second part of the 20 kyr cycle, climate at this given latitude becomes cooler and more humid, and the cycle starts over again. It is clear from Fig. 12A that the
- time interval of vigorous sedimentation covers only a short part of the 20'000 years and that much time is represented by the sequence boundaries (e.g., Strasser, 2015).

Model B (Fig. 12) illustrates the situation during generally low insolation at the 100 kyr and/or 405 kyr scale (Fig. 1). Sea level amplitudes at the 20 kyr scale are low, and an amplitude of 1 m is chosen for the model. The climate is rather cool and humid. Consequently, more land areas are exposed, rainfall favours their erosion, rivers are active, and deltas prograde. Consequently, the elementary sequence will be dominated by siliciclastics. A relatively short time interval of less humid conditions and warmer temperatures allows for some carbonate production, but siliciclastics are still present. Again, much time of this cycle is spent in intertidal to supratidal

830

conditions.

From the graphs in Fig. 12 sedimentation rates can be deduced. At the fastest rise of the sea floor during about 2000 years (corresponding to the fastest rise of sea level) about 1 m of coral reef conclusion number of 0.5 mm/yr. In model B, representing a siliciclast comminated scenario, only 30 cm of lagoonal facies accumulate in 2000 years, giving a sedimentation rate of 0.15 mm/yr. Holocene rates for

835 these environments reach 14 mm/y for a healthy reef and between 1 and 2 mm/yr for a tidal flat (Enos, 1991; Strasser and Samankassou, 2003). This discrepancy does not necessarily imply that sedimentation rates in the Oxfordian were generally lower than in the Holocene but rather that the time of non-deposition, reworking, and/or condensation is underestimated in the models (Sadler, 1994; Strasser, 2015).







- 845 **Figure 12.** Conceptual models that explain the formation of elementary sequences as a function of climate changes in tune with the orbital precession cycle. A: high insolation within a 100 and/or 405 kyr eccentricity cycle; B: low insolation. For discussion refer to text. Symbols and abbreviations as in Fig. 6.
- 850 These two models must undergo many modifications depending on the position on the platform, on the currents distributing nutrients and siliciclastics, and on the ecology of the carbonate-producing organisms. Also, they are idealized cases where facies evolution exhibits deepening-shallowing trends within an elementary sequence. In the studied sections, this is not always the case, and facies may stay the same throughout the sequence (with the exception of the marly layers at the sequence boundaries). This may be due to bioturbation that homogenized the sediment in a lagoon, or to constant reworking by waves and currents in the case of an ooid shoal. Nevertheless,
- 855 sediment in a lagoon, or to constant reworking by waves and currents in the case of an ooid shoal. Nevertheless, the models illustrate the potential of shallow marine carbonates to record climate changes at a relatively high time resolution.





#### 5 Conclusions

- Orbital cycles controlling the energy received by the Earth from the sun are important drivers of environmental changes that occur on land and in the ocean. In the case study presented here, it is shown that, during the Oxfordian (Late Jurassic), the shallow marine carbonate systems on the platform of the Swiss Jura Mountains reacted to these changes at the frequencies of the long of the controlling factors were meter-scale sea level changes and climate changes between humid and arid. Insolation changes translated more or less
- 870 directly into eustatic sea level changes through thermal expansion and retraction of the ocean surface waters. In addition, glacioeustasy and aquifer eustasy certainly occurred but are difficult to quantify. At all frequencies, humid and cool periods corresponded to low insolation and low sea level, while arid and warm periods prevailed during intervals of high insolation and high sea level. At a subtropical latitude on the shallow marine platform, many carbonate-producing organisms (e.g., algae, foraminifera, gastropods, bivalves, serpulids) were present,
- and especially coral reefs were sensitive to environmental changes. Ooids, oncoids, and microbialites also contributed to the sediment.

An interval of 1. An interval of 1. An interval of a discussed, calibrated by bio-, chrono-, and sequence stratigraphy. Three detailed sections are chosen to illustrate the evolution of facies and stacking pattern of the depositional sequences within this interval. During high insolation in the middle of a 405 kyr cycle, high amplitudes of sea level fluctuations and arid conditions allowed for the development of ooid shoals and coral reefs. However, the corals were regularly encrusted by microbialites, suggesting nutrient input even under an arid climate. At low insolation, low sea level amplitudes, and more humid and cooler conditions, siliciclastics were washed into the system and hampered the organic carbonate production.

885

890

880

The orbital frequencies are reflected in the hierarchical stacking of sequences: medium- and small-scale sequences formed in tune with the 405 and 100 kyr eccentricity cycles, respectively, while the 20 kyr precession cycles controlled the formation of elementary sequences. The obliquity cycle was not detected in the studied outcrops. After decompaction of the sections and estimation of the water depths in which the different facies evolved, a composite sea level curve has been reconstructed. It compares well with the curve of insolation changes over the last 500 kyr.

Based on the studied sections and the facies encountered, two models are presented. Model A describes the growth and demise of a coral reef in warm waters and a high-amplitude sea level rise to accommodate the reef. Arid conditions are exemplified by evaporites. Model B simulates a low-amplitude sea level cycle and rainfall in the hinterland that washed siliciclastics and plant matter onto the platform. The sedimentation rates deduced

- from the models are below those of Holocene sedimentary environments, implying that the time of nondeposition in the models probably is underestimated.
- 900 Despite the high lateral and vertical facies variability on the Oxfordian carbonate platform, despite the uncertainties in attributing water depth and compaction factors to the different facies, and despite the multiple feedback loops between insolation at the top of the atmosphere and the depositional environment, this study demonstrates the potential of shallow marine carbonates to record climate changes at the frequencies of orbital cycles. The time resolution of 20'000 years (the precession cycle) offers the opportunity to estimate rates of sea
- 905 level change, of ecological processes, and of sediment accumulation in the geologic past. The interpretation of the evolution of ancient sedimentary systems can thus be refined and better compared to today's changes in ecosystems. Concerning the rate of climate change, this study implies that anthropogenically induced global warming and subsequent sea level rise today occurs more than ten times faster than the fastest rise reconstructed for the Oxfordian.

Competing interests: The author declares that he has no conflict of interest.

915

#### Acknowledgements

I thank editor Denis-Didier Rousseau for having invited me to write this contribution. It is based on field work and research carried out by many MSc and PhD students at the University of Fribourg, and was financed mainly by the Swiss National Science Foundation. Their intellectual, institutional, and financial support is gratefully acknowledged. I also thank the reviewers ..... whose constructive criticism greatly improved the manuscript.

<sup>910</sup> 





#### References

925	Abbink, O., Targarona, J., Brinkhuis, H., and Visscher, H.: Late Jurassic to earliest Cretaceous palaeoclimatic
923	8181(01)00101-1, 2001. Allenback B. P. Synsedimentary tectonics in an encontinental sea: a new interpretation of the Ovfordian basing
	of northern Switzerland, Eclogae Geol. Helv., 94, 265-287, 2001.
930	Andrieu, S., Brigaud, B., Barbarand, J., Lasseur, E., and Saucède, T.: Disentangling the control of tectonics, eustasy, trophic conditions and climate on shallow-marine carbonate production during the Aalenian– Oxfordian interval: From the western France platform to the western Tethyan domain, Sed. Geol., 345, 54-84. https://doi.org/10.1016/i.sedgeo.2016.09.005, 2016.
935	Amies, J. D., Rohling, E. J., Grant, K. M., Rodriguez-Sanz, and L., Marino, G.: Quantification of African monsoon runoff during last interglacial sapropel S5, Paleoceanogr. Paleoclim., 34, 1487–1516, https://doi.org/10.1029/2019PA003652, 2019.
	Anderson, T. F. and Arthur, M. A.: Stable isotopes of oxygen and carbon and their application to sedimentology and paleoenvironmental problems, SEPM Short Course, 10, 151 p., 1983.
940	Armstrong, H. A., Wagner, T., Herrinshaw, L. G., Farnsworth, A. J., Lunt, D. J., Harland, M., Imber, J., Loptson, C., and Atar, E. F. L.: Hadley circulation and precipitation changes controlling black shale deposition in the Late Jurassic Boreal Seaway, Paleoceanography, 31, 1041-1053, https://doi.org/10.1002/2015PA002911, 2016.
	Beaulieu, E., Goddéris, Y., Donnadieu, Y., Labat, D., and Roelandt, C.: High sensitivity of the continental- weathering carbon dioxide sink to future climate change, Nature Clim. Change, 2, 346-349, https://doi.org/10.1038/NCLIMATE1419, 2012.
945	Berger, A., Loutre, MF., and Laskar, J.: Stability of the astronomical frequencies over the Earth's history for paleoclimatic studies. Science. 255, 560-566. https://doi.org/10.1126/science.255.5044.560, 1992.
	Berggren, W. A., Kent, D. V., Aubry, M. P., and Hardenbol, J. (Eds.): Geochronology, Time Scales and Global Stratigraphic Correlation SERM Spec Publ. 54, 386 pp. 1995
950	<ul> <li>Boulila, S., Galbrun, B., Hinnov, L. A., Collin, PY., Ogg, J. G., Fortwengler, D., and Marchand, D.:</li> <li>Milankovitch and sub-Milankovitch forcing of the Oxfordian (Late Jurassic) Terres Noires Formation (SE France) and global implications, Basin Res., 22, 717-732, https://doi.org/10.1111/j.1365-2117.2000.00420 #: 2010.</li> </ul>
955	<ul> <li>Brigaud, B., Vincent, B., Carpentier, C., Robin, C., Guillocheau, F., Yven, B., and Huret, E.: Growth and demise of the Jurassic carbonate platform in the intracratonic Paris Basin (France): Interplay of climate change, eustasy and tectonics, Marine Petrol. Geol., 53, 3-29, https://doi.org/10.1016/j.marpetgeo.2013.09.008, 2014</li> </ul>
	Carpentier, C., Lathuilière, B., and Ferry, S.: Sequential and climatic framework of the growth and demise of a
960	carbonate platform: implications for the peritidal cycles (Late Jurassic, north-eastern France), Sedimentology, 57, 985-1020, https://doi.org/10.1111/j.1365-3091.2009.01128.x, 2010.
200	climatic implications in the eastern Paris Basin (France), Terra Nova, 18, 191-201, https://doi.org/10.1111/ji.1365-3121.2006.00679.x. 2006
	Catuneanu, O., Abreu, V., Bhattacharya J. P., Blum, M. D., Dalrymple, R. W., Eriksson, P. G., Fielding, C. R., Ficher, W. L., Golloway, W. F., Gibling, M. P., Giles, K. A., Holbreck, J. M., Jordan, P., Kondell, C. G.
965	<ul> <li>St. C., Macurda, B., Martinsen, O. J., Miall, A. D., Neal, J. E., Nummedal, D., Pomar L., Posamentier H.</li> <li>W., Pratt, B. R., Sarg, J. F., Shanley, K. W., Steel, R. J., Strasser, A., Tucker, M. E., and Winker, C.:</li> <li>Towards the standardization of sequence stratigraphy, Earth-Sci. Rev., 92, 1-33,</li> </ul>
	Cecca, F., Martin-Garin, B., Marchand, D., Lathuilière, B., and Bartolini, A.: Paleoclimatic control of
970	biogeographic and sedimentary events in Tethyan and peri-Tethyan areas during the Oxfordian (Late Jurassic), Palaeogeo., Palaeoclim., Palaeoeco., 222, 10-32, https://doi.org/10.1016/j.palaeo.2005.03.009,
	2005. Church, J. A. and Clark, P. U. (coordinating lead authors): Sea Level Change, in: Climate Change 2013: The
975	Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Interrovernmental Panel on Climate Change, Cambridge Univ. Press, 1137-1216, 2013.
715	Colombié, C., Carcel, D., Lécuyer, C., Ruffel, A., and Schnyder, J.: Temperature and cyclone frequency in Kimmeridgian Greenhouse period (late Jurassic), Global Planet. Change, 170, 126-145, https://doi.org/10.1016/j.clogaloche.2018.08.005.2018
	Colombié, C., Lécuyer, C., and Strasser, A.: Carbon- and oxygen-isotope records of palaeoenvironmental and
980	carbonate production changes in shallow-marine carbonates (Kimmeridgian, Swiss Jura), Geol. Mag., 148, 133-153, https://doi.org/10.1017/S0016756810000518, 2011.



990



- Curtis, C. D.: Aspects of climate influence on the clay mineralogy and geochemistry of soils, paleosols and clastic sedimentary rocks, J. Geol. Soc. London, 147, 351-357, https://doi.org/10.1144/gsjgs.147.2.0351, 1990.
- 985 Davies, A., Gréselle, B., Hunter, S. J., Baines, G., Robson, C., Haywood, A., David C. Ray, D. C., Simmons, M. D., and van Buchem, F. S. P.: Assessing the impact of aquifer-eustasy on short-term Cretaceous sea-level, Cretaceous Res., 112. 104445, https://doi.org/10.1016/j.cretres.2020.104445, 2020.
  - de Winter, N. J., Zeeden, C., and Hilgen, F. J.: Low-latitude climate variability in the Heinrich frequency band of the Late Cretaceous greenhouse world, Clim. Past, 10, 1001-1015, https://doi.org/10.5194/cp-10-1001-2014, 2014.
    - Dercourt, J., Ricou, L. E., and Vrielynck, B. (eds): Atlas: Tethys Palaeoenvironmental Maps, Gauthier-Villars, Paris, 1993.

Dunham, R. J.: Classification of carbonate rocks according to depositional texture, AAPG Mem., 1, 108–121, 1962.

995 Dupraz, C.: Paléontologie, paléoecologie et évolution des faciès récifaux de l'Oxfordien moyen-supérieur (Jura suisse et français), Ph.D. thesis, GeoFocus, 2, Fribourg, Switzerland, 247 pp., 1999.

Dupraz, C. and Strasser, A.: Nutritional modes in coral-microbialite reefs (Jurassic, Oxfordian, Switzerland): evolution of trophic structure as a response to environmental change, Palaios, 17, 449-471, https://doi.org/10.1669/0883-1351(2002)017<0449:NMICMR>2.0.CO;2, 2002.

- 1000 Embry, A. F. and Klovan, J. E.: A late Devonian reef tract on northeastern Banks Island, N.W.T., Bull. Canadian Petrol. Geol., 19, 730-781, 1971.
  - Enos, P.: Sedimentary parameters for computer modeling, Kansas Geol. Survey Bull., 233, 63-99, 1991.
    - Feng, R., and Poulsen, C. J.: Andean elevation control on tropical Pacific climate and ENSO, Paleoceanography, 29, 795-809, https://doi.org/10.1002/2014PA002640, 2014.
- 1005 Flügel, E.: Microfacies of Carbonate Rocks, Springer, Berlin, Germany, 976 pp., 2004. Frakes, L. A., Francis, J. E., and Syktus, J. I.: Climate Modes of the Phanerozoic, Cambridge Univ. Press, 274 pp., 1992.
  - Goddéris, Y., Donnadieu, Y., Lefebvre, V., Le Hir, G., and Nardin, E..: Tectonic control of continental weathering, atmospheric CO<sub>2</sub>, and climate over Phanerozoic times, C. R. Geoscience, 344, 652-662, https://doi.org/10.1016/j.crte.2012.08.009, 2012.
- 1010
   https://doi.org/10.1016/j.crte.2012.08.009, 2012.

   Goldhammer, R. K.: Compaction and decompaction algorithms for sedimentary carbonates, J. Sed. Res., 67, 26-35, https://doi.org/10.1306/D42684E1-2B26-11D7-8648000102C1865D, 1997.
  - Gradstein, F. M., Ogg, J. G., Schmitz, M. D., and Ogg, G. (Eds.): Geologic Time Scale 2020, vol.1, Elsevier, Amsterdam, The Netherlands, 1357 pp., 2020.
- 1015 Gygi, R. A.: Datierung von Seichtwassersedimenten des Späten Jura in der Nordwestschweiz mit Ammoniten, Eclogae Geol. Helv., 88, 1-58, 1995.
  - Gygi, R. A.: Integrated stratigraphy of the Oxfordian and Kimmeridgian (Late Jurassic) in northern Switzerland and adjacent southern Germany, Mem. Swiss Acad. Sci., 104, 152 pp., 2000.
- Gygi, R. A., Coe, A. L., and Vail, P. R.: Sequence stratigraphy of the Oxfordian and Kimmeridgian stages (Late
   Jurassic) in northern Switzerland, SEPM Spec. Publ., 60, 527-544, 1998.
  - Gygi, R. A. and Persoz, F.: Mineralostratigraphy, litho- and biostratigraphy combined in correlation of the Oxfordian (Late Jurassic) formations of the Swiss Jura range, Eclogae Geol. Helv., 79, 385-454, 1986.
     Hallock, P. and Schlager, W.: Nutrient excess and the demise of coral reefs and carbonate platforms, Palaios, 1,
- 389-398, 1986.
   Hardenbol, J., Thierry, J., Farley, M. B., Jacquin, T., de Graciansky, P.-C., and Vail, P. R.: Jurassic sequence chronostratigraphy, in: Mesozoic and Cenozoic Sequence Stratigraphy of European Basins, edited by: de Graciansky, P.-C., Hardenbol, J., Jacquin, T., and Vail, P. R, SEPM Spec. Publ., 60, 1998.
  - Hesselbo, S. P., Ogg, J. G., and Ruhl, M.: The Jurassic Period, in: Geologic Time Scale 2020, vol. 1, edited by: Gradstein, F. M., Ogg, J. G., Schmitz, M. D., and Ogg, G., Elsevier, Amsterdam, The Netherlands, 955-1021, 2020.
- 1030 1021, 2020.
   Hinnov, L. A.: Cyclostratigraphy and astrochronology in 2018, in: Stratigraphy and Timescales, vol. 3, edited by: Montenari, M., Elsevier, Amsterdam, The Netherlands, 1-80, https://doi.org/10.1016/bs.sats.2018.08.004, 2018.
- Huang, C.: Astronomical time scale for the Mesozoic, in: Stratigraphy and Timescales, vol. 3, edited by: Montenari, M., Elsevier, Amsterdam, The Netherlands, 81-150, https://doi.org/10.1016/bs.sats.2018.08.005, 2018.
  - Hug, W.: Sequenzielle Faziesentwicklung der Karbonatplattform des Schweizer Jura im späten Oxford und frühesten Kimmeridge, Ph.D. thesis, GeoFocus 7, Univ. Fribourg, Switzerland, 156 pp., 2003.
- Ikeda, M., Ozaki, K., and Legrand, J.: Impact of 10-Myr scale monsoon dynamics on Mesozoic climate and<br/>ecosystems, Nature Sci. Rep., 10, 11984, https://doi.org/10.1038/s41598-020-68542-w, 2020.





	Immenhauser, A., Della Porta, G., Kenter, J. A. M., and Bahamonde, J. R.: An alternative model for positive shifts in shallow-marine carbonate δ <sup>13</sup> C and δ <sup>18</sup> O, Sedimentology, 50, 953–959, https://doi.org/10.1046/j.1365-3091.2003.00590.x, 2003.
	Jacobs, D. K. and Sahagian, D. L.: Climate-induced fluctuations in sea level during non-glacial times, Nature,
1045	361, 710-712, https://doi.org/10.1038/361710a0, 1993.
	Khon, V. C., Park, W., Latif, M., Mokhov, I. I., and Schneider, B.: Tropical circulation and hydrological cycle
	response to orbital forcing, Geophys. Res. Lett., 39, L15708, https://doi.org/10.1029/2012GL052482,
	2012. Laskar I. Fienga A. Gastineau M. and Manche H.: La2010: a new orbital solution for the long term motion
1050	of the Earth. Astron. Astronbys. 532, A89, https://doi.org/10.1051/0004-6361/201116836, 2011.
1020	Laurin, J., Meyers, S. R., Uličný, D., Jarvis, I., and Sageman, B. B.: Axial obliquity control on the greenhouse
	carbon budget through middle- to high-latitude reservoirs, Paleoceanography, 30, 133-149,
	https://doi.org/10.1002/2014PA002736, 2015.
1055	Lefort, A., Lathulliere, B., Carpentier, C., and Huault, V.: Microfossil assemblages and relative sea-level fluctuations in a lagoon at the Oxfordian/Kimmeridgian boundary (Upper Jurassic) in the eastern part of the Paris Basin, Facies, 57, 649-662, https://doi.org/10.1007/s10347-010-0259-4, 2011.
	Leinfelder, R. R., Schmid, D. U., Nose, M., and Werner, W.: Jurassic reef patterns - the expression of a
	changing globe, SEPM Spec. Publ., 72, 465-520, 2002.
1060	in pCO <sub>2</sub> and climate in the Oxfordian (Late Jurassic): Carbon isotope and modeling evidence, Earth Planet, Sci. Lett., 258, 44-60, https://doi.org/10.1016/j.ensl.2007.03.014, 2007.
	Lourens, L. J., Hilgen, F. J., Raffi, I., and Vergnaud-Grazzini, C.: Early Pleistocene chronology of the Vrica
	Section (Calabria, Italy), Paleoceanography, 11, 797-812, https://doi.org/10.1029/96PA02691, 1996.
	Martin-Garin, B., Lathuilière, B., Geister, J., and Ramseyer, K.: Oxygen isotopes and climatic control of
1065	Oxfordian coral reefs (Jurassic, Tethys), Palaios, 25, 721-729, https://doi.org/10.2110/palo.2010.p10-
	02/r, 2010. Martinez, M. Mashaniama of measuration of the accountribity and langar term Milankavitah avalas in detrital
	supply and carbonate production in hemipelagic marl-limestone alternations, in: Stratigraphy and
	Timescales, vol. 3, edited by: Montenari, M., Elsevier, Amsterdam, The Netherlands, 189-218,
1070	https://doi.org/10.1016/bs.sats.2018.08.002, 2018.
	Martinez, M. and Dera, G.: Orbital pacing of carbon fluxes by a ~9-My eccentricity cycle during the Mesozoic,
	PNAS, 112, 12604–12609, https://doi.org/10.10/3/pnas.1419946112, 2015. Masson Dalmotta V. Zhoi P. Dirani A. Connors S. L. Dáan C. Dargar S. Caud N. Chan V. Goldfark I.
	Gomis, M. L. Huang, M., Leitzell, K., Lonnov, E., Matthews, J. B. R., Mavcock, T. K., Waterfield, T.,
1075	Yelekçi, O., Yu, R., and B. Zhou, B. (Eds.): Climate Change 2021: The Physical Science Basis.
	Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on
	Climate Change, Cambridge Univ. Press, 2022. Mattheway M. D. and Backmuttan M. A.: Clabal avalation targen privation to the Econor Crean Biver
	basin in: Orbital Forcing and Cyclic Sequences, edited by de Boer P and Smith D G IAS Spec. Publ
1080	19, 459-481, 1994.
	Mitchum Jr., R. M. and Van Wagoner, J. C.: High-frequency sequences and their stacking patterns: sequence-
	stratigraphic evidence of high-frequency eustatic cycles, Sed. Geol., 70, 131-160,
	https://doi.org/10.1016/003/-0/38(91)90139-5, 1991. Montañez I A and Osleger D A : Parasequence stacking patterns, third-order accommodation events, and
1085	sequence stratigraphy of Middle to Upper Cambrian platform carbonates. Bonanza King Formation.
	southern Great Basin, AAPG Mem., 57, 305-326, 1993.
	Ogg, J. G., Hinnov, L. A., and Huang, C.: Jurassic, in: The Geologic Time Scale 2012, edited by: Gradstein, F.
	M., Ogg, J. G., Schmitz, M. D., and Ogg, G., Elsevier, Amsterdam, The Netherlands, 731-791, 2012.
1090	platform, French Jura Mountains, Swiss J. Geosci., 108, 273-288, https://doi.org/10.1007/s00015-015-
	0189-9, 2015.
	Olivier, N., Colombié, C., Pittet, B., and Lathuilière, B.: Microbial carbonates and corals on the marginal French
	Jura platform (Late Oxfordian, Molinges section), Facies, 57, 469-492, https://doi.org/10.1007/s10347-
1095	010-0240-9, 2011. Padden, M., Weissert, H., Funk, H., Schneider, S., and Gansner, C.: Late Iurassic lithological evolution and
	carbon-isotope stratigraphy of the western Tethys, Eclogae geol. Helv., 95, 333-346, 2002.
	Pittet, B.: Contrôles climatiques, eustatiques et tectoniques sur des systèmes mixtes carbonates-siliciclastiques de
	plate-forme: exemples de l'Oxfordien (Jura suisse, Normandie, Espagne), Ph.D. thesis, Univ. Fribourg,
	Switzerland, 258 pp., 1996.





1100	Pittet, B. and Gorin, G. E.: Distribution of sedimentary organic matter in a carbonate-siliciclastic platform environment: Oxfordian in the Swiss Jura Mountains, Sedimentology, 44, 915-937,
	<ul> <li>Pittet, B. and Strasser, A.: Long-distance correlations by sequence stratigraphy and cyclostratigraphy: examples and implications (Oxfordian from the Swiss Jura, Spain, and Normandy), Geol. Rundsch., 86, 852-874,</li> </ul>
1105	https://doi.org/10.1007/s005310050181, 1998. Plunkett, J. M.: Early Diagenesis of shallow platform carbonates in the Oxfordian of the Swiss Jura Mountains,
	PhD thesis, Univ. Fribourg, Switzerland, 155 pp., 1997.
1110	Pratt, B. R. and James, N. P.: The St George Group (Lower Ordovician) of western Newfoundland: tidal flat island model for carbonate sedimentation in shallow epeiric seas, Sedimentology, 33, 313–343,
1110	Railsback, L. B., Gibbard, P. L., Head, M. J., Voarintsoa, N. R. G., and Toucanne, S.: An optimized scheme of
	lettered marine isotope substages for the last 1.0 million years, and the climatostratigraphic nature of
1115	isotope, Quat. Sci. Rev., 111, 94-106, https://doi.org/10.1016/j.quascirev.2015.01.012, 2015. Rais, P., Louis-Schmid, B., Bernasconi, S. M., and Weissert, H.: Palaeoceanographic and palaeoclimatic reorganization around the Middle Late Iurgesic transition Palaeogea. Palaeoceanographic and palaeoceanographic
1115	546. https://doi.org/10.1016/i.palaeo.2007.05.008, 2007.
	Rankey, E. C., Enos, P., Steffen, K., and Druke, D.: Lack of impact of Hurricane Michelle on tidal flats, Andros
	Island, Bahamas: Integrated remote sensing and field observations, J. sed. Res., 74, 654-661, https://doi.org/10.1306/021704740654, 2004.
1120	Rankey, E. C., and Reeder, S. L.: Controls on platform-scale patterns of surface sediments, shallow Holocene platforms. Bahamas, Sedimentology, 57, 1545-1565, https://doi.org/10.1111/j.1365-3091.2010.01155.x, 2010.
	Ray, D. C., van Buchem, F. S. P., Baines, G., Davies, A., Gréselle, B., Simmons, M. D., and Robson, C.: The
1125	magnitude and cause of short-term eustatic Cretaceous sea-level change: a synthesis, Earth-Sci. Rev., 197, 102001, https://doi.org/10.1016/j.earscirey.2019.102001, 2019
1125	Reijmer, J. J. G.: Marine carbonate factories: review and update, Sedimentology, 68, 1729-1796,
	https://doi.org/10.1111/sed.12878, 2021.
	Sadler, P. M.: The expected duration of upward-shallowing peritidal carbonate cycles and their terminal hiatuses, Bull Geol Soc Am 106 791-802 https://doi.org/10.1130/0016-
1130	7606(1994)106<0791:TEDOUS>2.3.CO;2, 1994.
	Sames, B., Wagreich, M., Wendler, J. E., Haq, B. U., Conrad, C. P., Melinte-Dobrinescu, M. C., Hu, X.,
	Wendler, I., Wolfgring, E., Yilmaz, I., and Zorina, S. O.: Review: Short-term sea-level changes in a greenhouse world – a view from the Cretaceous. Palaeogeo, Palaeoclim., Palaeocco., 441, 393-411.
	https://doi.org/10.1016/j.palaeo.2015.10.045, 2016.
1135	Schoepf, V., Falter, J. L., and McCulloch, M. T.: Limits to the thermal tolerance of corals adapted to a highly
	fluctuating, naturally extreme temperature environment, Sci. Rep., 5, 1/639, https://doi.org/10.1038/srep17639_2015
	Schulz, M. and Schäfer-Neth, C.: Translating Milankovitch climate forcing into eustatic fluctuations via thermal
	deep water expansion: a conceptual link, Terra Nova, 9, 228-231, https://doi.org/10.1111/j.1365-
1140	3121.1997.tb00018.x, 1997.
	Shackleton, N. J.: Oxygen isotopes, ice volume and sea level, Quat. Sci. Rev., 6, 183–190, https://doi.org/10.1016/0277-3791(87)90003-5, 1987.
	Sellwood, B. W. and Valdes, P. J.: Jurassic climates, Proc. Geol. Assoc., 119, 5-17,
1145	Sellwood, B. W., Valdes, P. J., and Price, G. D.: Geological evaluation of multiple general circulation model
	simulations of Late Jurassic palaeoclimate, Palaeogeo. Palaeoclim. Palaeoeco., 156, 147-160,
	https://doi.org/10.1016/S0031-0182(99)00138-8, 2000.
	Stienne, N.: Paléoécologie et taphonomie comparative en milieux carbonatés peu profonds (Oxfordien du Jura
1150	Strasser, A.: Lagoonal-peritidal sequences in carbonate environments: autocyclic and allocyclic processes, in:
	Cycles and Events in Stratigraphy, edited by: Einsele, G., Ricken, W., and Seilacher, A., Springer, Heidelberg, Germany, 709-721, 1991.
	Strasser, A.: Astronomical time scale for the Middle Oxfordian to Late Kimmeridgian in the Swiss and French
1155	Strasser, A.: Hiatuses and condensation: an estimation of time lost on a shallow carbonate platform. Depositional
	Record, 1, 91-117, https://doi.org/10.1002/dep2.9, 2015.
	Strasser, A.: Cyclostratigraphy of shallow-marine carbonates – limitations and opportunities, in: Stratigraphy
	https://doi.org/10.1016/bs.sats.2018.07.001, 2018.





<ul> <li>cyclicity, Terra Nova, 12, 303-311, https://doi.org/10.1046/j.1365-3121.2000.00315.x, 2000.</li> <li>Strasser, A., Pittet, B., Hillgärtner, H., Pasquier, JB., 1999. Depositional sequences in shallow carbonate dominated sedimentary systems: concepts for a high-resolution analysis, Sed. Geol., 128, 201-221, https://doi.org/10.1016/S0037-0738(99)00070-6, 1999.</li> <li>Strasser, A., Pittet, B., and Hug, W.: Palaeogeography of a shallow carbonate platform: The case of the Microsoft Strasser, A., Pitter, B., and Hug, W.: Palaeogeography of a shallow carbonate platform.</li> </ul>	iddle
<ul> <li>Strasser, A., Pittet, B., Hillgärtner, H., Pasquier, JB., 1999. Depositional sequences in shallow carbonate dominated sedimentary systems: concepts for a high-resolution analysis, Sed. Geol., 128, 201-221, https://doi.org/10.1016/S0037-0738(99)00070-6, 1999.</li> <li>Strasser, A., Pittet, B., and Hug, W.: Palaeogeography of a shallow carbonate platform: The case of the M.</li> </ul>	iddle
dominated sedimentary systems: concepts for a high-resolution analysis, Sed. Geol., 128, 201-221, https://doi.org/10.1016/S0037-0738(99)00070-6, 1999.	iddle
https://doi.org/10.1016/S0037-0738(99)00070-6, 1999. 1165 Strasser, A., Pittet, B., and Hug, W.: Palaeogeography of a shallow carbonate platform. The case of the M	iddle
1165 Strasser, A., Pittet, B., and Hug, W.: Palaeogeography of a shallow carbonate platform: The case of the M	iddle
to Late Oxfordian in the Swiss Jura Mountains, J. Palaeogeo., 4, 251-268,	
https://doi.org/10.1016/j.jop.2015.08.005, 2015.	
Strasser, A. and Samankassou, E.: Carbonate sedimentation rates today and in the past: Holocene of Florid	la Bay,
Bahamas, and Bermuda vs. Upper Jurassic and Lower Cretaceous of the Jura Mountains (Switzerla	ind and
1170 France), Geol. Croatica, 56, 1-18, 2005.	
Strasser, A., Vedrine, S., and Stienne, N.: Kate and synchronicity of environmental changes on a shallow	
transition bill to the organization of the org	
Thiry, M. Palaeolimatic interpretation of clay minerals in marine denosits: an outlook from the continent	al
1175 right added match method and real matching and real service and real	ai
Valero, L., Garcés, M., Cabrera, L., Costa, E., and Sáez, A.: 20 Myr of eccentricity paced lacustrine cycles	in the
Cenozoic Ebro Basin, Earth Planet. Sci. Lett., 408, 183-193, https://doi.org/10.1016/j.epsl.2014.10.	007.
2014.	,
Védrine, S.: High-frequency palaeoenvironmental changes in mixed carbonate-siliciclastic sedimentary sy	stems
1180 (Late Oxfordian, Switzerland, France, and southern Germany), Ph.D. thesis, GeoFocus, 19, Univ.	
Fribourg, Switzerland, 216 pp., 2007.	
Védrine, S. and Strasser, A.: High-frequency palaeoenvironmental changes on a shallow carbonate platfor	m
during a marine transgression (Late Oxfordian, Swiss Jura Mountains), Swiss J. Geosci., 102, 24/-:	270,
https://doi.org/10.100//S00015-009-1326-0, 2009.	and
1165 Vedrine, S., Strässer, A., and Hug, W.: Oncold growth and distribution controlled by sea-level induitations adjuncted (Lato Orfordian Swing Jurg Mountaine) Equips 52 525 555 [https://doi.org/10.1007/s102	s and 17
007-0114-2007	+/-
Wallmann K. Schneider B. and Samthein M. Effects of custatic sea-level change ocean dynamics and	1
$\alpha$ minimized, $\beta$ , $\beta$ many	nodel
1190 study, Clim. Past, 12, 339-375, https://doi.org/10.5194/cp-12-339-2016, 2016.	
Weissert, H. and Mohr, H.: Late Jurassic climate and its impact on carbon cycling, Palaeogeo. Palaeoclim	,
Palaeoeco., 122, 27-43, https://doi.org/10.1016/0031-0182(95)00088-7, 1996.	
Wendler, J. E. and Wendler, I.: What drove sea-level fluctuations during the mid-Cretaceous greenhouse	
climate? Palaeogeo. Palaeoclim., Palaeoeco., 441, 412-419, https://doi.org/10.1016/j.palaeo.2015.0	8.029,
1195 2016.	
Wendler, J. E., Wendler, I., Vogt, C., and Kuss, J.: Link between cyclic eustatic sea-level change and cont	inental
weathering: evidence for aquifer-eustasy in the Cretaceous, Palaeogeo. Palaeoclim., Palaeoeco.,441	, 430-
437, https://doi.org/10.1016/j.palaeo.2015.08.014, 2016.	
Wildi, W., Funk, H., Loup, B., Amato, E., and Huggenberger, P.: Mesozoic subsidence history of the Euro	pean
1200 marginal shelves of the Alpine Tethys (Helvetic realm, Swiss Plateau and Jura), Eclogae Geol. Hel	v., 82,
01/-040, 1969. Wood $P = 1000$ Reaf Evolution Oxford Univ. Press 414 pp. 1000	
Zühlke R · Integrated cyclostratigraphy of a model Mesozoic carbonate platform – the Latemar (Middle	
Triassic. Italy). SEPM Spec. Publ., 81, 183-211, 2004.	
1205	