



Assessing agriculture's vulnerability to drought in European pre-Alpine regions

Ruth Stephan¹, Stefano Terzi^{2,3,4}, Mathilde Erfurt¹, Silvia Cocuccioni², Kerstin Stahl¹, and Marc Zebisch²

¹Environmental Hydrological Systems, University of Freiburg, Freiburg i. Br., 79085, Germany

²Institute for Earth Observation, Eurac Research, Viale Druso 1, 39100, Bolzano, Italy

³Center for Global Mountain Safeguard Research, Eurac Research, Viale Druso 1, 39100, Bolzano, Italy

⁴United Nations University Institute for Environment and Human Security (UNU-EHS), Platz der Vereinten Nationen 1, 53113 Bonn, Germany

Correspondence: Ruth Stephan (ruth.stephan@hydrology.uni-freiburg.de)

Abstract. Droughts are natural hazards that lead to severe impacts in the agricultural sector. Mountain regions are thought to have abundant water, but have experienced unprecedented drought conditions as climate change is affecting their environments more rapidly than other places. The effect radiates by reducing water availability well beyond the mountains' geographical locations. This study aims to improve the understanding of agriculture's vulnerability to drought in Europe's pre-Alpine region, considering two case studies that have been severely impacted in the past. We applied a mixed-method approach combining the knowledge of regional experts with quantitative data analyses in order to define region-specific vulnerability based on experts' identified factors. We implemented two aggregation methods by combining the vulnerability factors that could be supported with subregional data. Whereas the equal weighting method combines all factors with the same weight, the expert weighting method combines the factors with varying weight based on expert opinion. These two methods resulted in vulnerability maps with the expert weighting showing in general higher vulnerability, and partly relocating the medium and lower vulnerabilities to other subregions within the case study regions. In general, the experts confirmed the resulting subregions with higher vulnerability. They also acknowledged the value of mapping vulnerability by adopting different aggregation methods confirming that this can serve as a sensitivity analysis. The identified factors contributing most to the regions' vulnerability point to the potential of adaptation strategies decreasing the agriculture's vulnerability to drought that could enable better preparedness. Apart from region-specific differences, in both study regions the presence of irrigation infrastructure and soil texture are among the most important conditions that could be managed to some extent in order to decrease the regions' vulnerability. Throughout the analyses, the study benefited from the exchange with the experts by getting an in-depth understanding of the regional context with feedback-relations between the factors contributing to vulnerability. Qualitative narratives provided during the semi-structured interviews supported a better characterization of local vulnerability conditions and helped to better identify quantitative indicators as proxies to describe the factors. Thus, we recommend to apply this mixed-method approach to close the gap between science and practitioners.



1 Introduction

Past and recent droughts have led to severe environmental, social and economic impacts in many regions of the world. The combination of climate change exacerbation and the arising pressures on water demand from socio-economic activities affect the intensity and frequency of drought conditions (van Loon et al., 2016). This is particularly relevant in mountain regions where climate change effects are occurring more rapidly than in other places with consequences on their water tower role and water provision to downstream areas (Beniston and Stoffel, 2014; Immerzeel et al., 2020; Terzi et al., 2021). Recent drought events highlighted the vulnerability of the European Alps and areas dependent on water from the Alps such as pre-alpine regions to unexpected conditions of reduced water availability (Hanel et al., 2018; Laaha et al., 2017). Stephan et al. (2021) showed that within the European Alps a wide range of drought impacts occurred in different socio-economic sectors, with agriculture and the public water supply most impacted.

Drought impacts are triggered by the natural hazard itself, such as the intensity, duration, frequency and extent of water deficits, but local exposure and vulnerability characteristics shape them (Hagenlocher et al., 2019). While the drought hazard components have been investigated and a set of indices are already established and available to describe hydroclimatic variations in terms of precipitation (e.g. SPI) and evapotranspiration (e.g. SPEI) in the Alpine region (e.g. Haslinger and Blöschl (2017)), the characterization of drought vulnerability and exposure still remains a challenge. Multiple conceptual frameworks of vulnerability in the context of natural hazards assessments have been developed (Birkmann et al., 2013; González Tánago et al., 2016). The Intergovernmental Panel on Climate Change (IPCC) provides the risk concept definition, where exposure refers to 'the presence of people, livelihoods, species or ecosystems, environmental functions, services, and resources, infrastructure, or economic, social, or cultural assets in places and settings that could be adversely affected' by a drought hazard' and vulnerability refers to the 'the propensity or predisposition to be adversely affected' (IPCC, p. 1048) due to the system's sensitivity or susceptibility combined with a lack of short-term coping capacity and long-term adaptive capacity. In case of drought risk assessments, an example of a short-term coping capacity is the existence of an irrigation system to reduce agricultural impacts, while the development of an agricultural system increasing the water use efficiency refers to a long-term adaptive capacity. Although exposure and vulnerability are internationally recognised as important drivers of drought risk processes and final impacts, their operational assessment is still discussed and in development. In particular, drought vulnerability studies for areas such as the heterogeneous European Alpine and pre-alpine region are rare. So far studies exist on the aspects of vulnerability in forest growth and for impacts on pasture (e.g. Hartl-Meier et al. (2014)). Melkonyan (2014) Melkonyan (2014) carried out a study to assess socio-economic vulnerability of the agricultural sector in the mountainous region of Armenia.

Moreover, various approaches have been applied often relying on either quantitative or qualitative data and hence covering only specific aspects of drought risk processes. The selection of the underlying approach often depends on the study's spatial scale, as well as on the data availability. Most studies that adopt a quantitative approach focus on large scale (e.g. national or global level, Meza et al. (2020); Carrão et al. (2016)), where datasets on socio-economic conditions are freely available. Additionally, data on past drought impacts are also often only available on a large scale, but necessary for an effective validation of the assessment. Regional datasets on socio-economic conditions are indeed often lacking, leading to an underrepresentation



or omission of regional risk conditions. Therefore, regional or local studies often follow a qualitative approach to assess vulnerability, e.g., by involving the local communities in the process of understanding the vulnerability dimensions (Ayantunde et al., 2015; Birhanu et al., 2017; Martin et al., 2016). The results of these two approaches have rarely been combined or compared.

60 For these reasons, this study considers the Impact Chains (IC), a mixed-method approach recognized as a valuable methodology to integrate both quantitative and qualitative information into the description of the hazard, exposure and vulnerability components for advancing the assessments of drought risk conditions (Schneiderbauer et al., 2020; Zebisch et al., 2021). The IC provides a guideline to systematically identify, select and assess relevant factors involved in risk processes through the combination of quantitative data with information coming from local experts and stakeholders (GIZ, 2017). This process is particularly important for evaluations of the vulnerability and exposure components since no standard set of factors exists to identify and characterise vulnerability to drought in agriculture.

Besides the identification of vulnerability factors, expert knowledge is often used to weigh different factors in the mapping of an overall vulnerability index (Zebisch et al., 2021). This is often adopted when certain factors are perceived to be more important than others and thus have a greater (or lesser) influence on the overall vulnerability. In the context of impact chains, equal weighting is usually applied more often than other weighting methods. Moreover, a comparison of different weighting methods in order to analyse the effect of such choice on the final vulnerability assessment, has rarely been done.

This study applies a vulnerability assessment approach in two case studies located in the European pre-Alpine region. The case study regions 'Thurgau' in Switzerland and 'Podravska' in Slovenia have experienced severe drought impacts in recent years and need to improve their resilience (Zorn and Hrvatin (2015); Komac et al. (2019), DROUGHT-CH). The overarching objective is to systematically identify vulnerability factors linked to agricultural production that potentially lead to agricultural impacts during a drought to assess how they vary spatially.

Subordinate methodological research questions are:

- To what degree can we characterise vulnerability to drought in the two case studies by combining experts' opinions with the openly accessible data?
- 80 – How sensitive is the final vulnerability assessment to different weighting methods for the vulnerability factors?

2 Case study areas

We selected the two study regions 'Thurgau' in Switzerland and 'Podravska' in Slovenia due to the increasing number of reported drought impacts in agriculture in both regions, despite their proximity to the water-rich conditions in the European mountainous regions (Stephan et al., 2021). Most impacts archived in the Alpine Drought Impact report Inventory (EDII_{ALPS}) report consequences on agriculture and livestock farming highlighting the agricultural sector of both regions vulnerable to drought in the past. According to EDII_{ALPS}, in Thurgau the most reported impacts occurred in 2015 and 2018, whereas in Podravska most impacts occurred in 2003 and 2017. In Thurgau, reports claimed a reduced productivity of annual and permanent



crop cultivation and shortages of feed and water for livestock as the majority of the regions' drought impacts. For instance EDII_{ALPS} reports a comment of a local farmer from Stettfurt about the considerable harvest loss: 'Production was down by
90 30 to 40 %. Especially iceberg lettuce had to be irrigated, because they were sensitive to the drought due to their small root system. Irrigation had to be done even at night, so water from the Stettfurt public utility company had to be used in part, which was much more expensive than the water from the Lauche River that he usually used'. In Podravska the reports are clearly dominated by reduced productivity of annual crop cultivation, often with yield losses $\geq 30\%$. For example, the Drought Management Centre for Southeastern Europe stated in 2017, 'Agricultural drought [...] worsened over June and July. Most affected
95 were regions of northeastern and southern half of Slovenia where maize completely stopped. [...] yield was reduced by 30-50 %. Hay production was seriously affected as well.' (DMCSEE, 2017).

Both the Thurgau and Podravska regions are characterised by extensive areas of agricultural land (Fig. 1) that make them particularly exposed to drought conditions, while showing different societal and economic conditions influencing the vulnerability of the agricultural sector. Moreover, they are also part of the Alpine Drought Observatory (ADO, <https://www.alpine-space.org/projects/ado>), an Interreg project aiming to improve the understanding of drought processes and impacts towards increasing
100 levels of drought preparedness through monitoring. Thurgau is known for its agriculturally shaped landscape reaching from specialty crops on the coast of Lake Constance to high elevation pastures (<https://www.landschaftsqualitaet-tg.ch/>). This study region covers 991.77 km², 54 % of which are covered by agricultural land (Fig. 1, CLC, 2018), with a total of 2531 farms in 2019. The main agricultural type (51 %) consists of natural meadows and pastures, followed by cropland (33 %), managed
105 meadows (9 %), and vegetables, fruit, vines, berries (6.5 %) (LID, 2007). Thurgau had 279,547 inhabitants in 2019 (SFSO, 2022) resulting in a population density of 324 inhabitants per km². Podravska's landscape varies from hilly lowlands mainly along the river Drava up to 1517 masl in the western mountainous part. The region covers an area of 2170 km², more than double the size of Thurgau. In 2019, the region had 324,875 inhabitants and thus a lower population density (149.7 inhabitants per km²) than Thurgau ('population' from SURS, 2022). In Podravska more than 30 % of the total area is utilised as agricul-
110 tural land with a total of 10,990 agricultural holdings in 2016, and almost one third of these are located in mountainous areas ('agriculture, forestry and fishery' from SURS, 2022).

3 Data and methods

This study applies a mixed-method approach based on qualitative and quantitative information for the assessment of the agriculture's vulnerability to drought conditions in the two case studies regions Thurgau and Podravska. Following the guidelines
115 of the vulnerability sourcebook (Fritzsche et al., 2014) our study approach was based on an initial exploratory phase during the ADO project meeting held online on the 24th September 2020. During the group discussion, experts from the two case study areas provided a first regional overview of the drought issues through the creation of impact chains following the model by Zebisch et al. (2021). For this study we then considered these impact chains as context for further discussions and refinements focusing on the vulnerability factors and their characteristics. Starting from this context, the methodological approach
120 presented here is composed of five consequential phases for the assessment of vulnerability considering its spatial variabil-




Figure 1. The two study regions a) Thurgau and b) Podravška within the European mountain region. The political border of the study regions is marked in red and the LAU2 regions are marked with white. The black labels present the most important cities and common known subareas (italic). The blue labels present the largest important rivers and lakes. The colours present the regional land cover adapted and modified by the Corine Land Cover (CLC, 2018). The land cover's share is shown in the histograms.

ity within both case study regions: (1) identification of vulnerability factors, (2) data acquisition and indicators selection, (3) pre-processing, normalisation and direction, (4) weighting and aggregating, and (5) participatory validation (Fig. 2).




Figure 2. Conceptual overview of the methodological steps for the drought vulnerability assessment. The grey box represents the preliminary group discussion on agricultural drought while green boxes refer to the five consequential phases for a spatial assessment of drought vulnerability.

3.1 Identification of vulnerability factors

The set of factors used to describe vulnerability to drought in agriculture was generated through semi-structured interviews with nine high-level experts identified as key people with extensive expertise and knowledge on the two case studies and they were held from the 24th August to 9th September 2021. Each discussion followed a flexible interactive structure (reported in S1) allowing to integrate the established questions with further information on the context and expertise from each participant. The interviews aimed to identify the perceived factors making the regions' agriculture vulnerable to drought and whether these factors increase or decrease the vulnerability. Further, the participants assigned an importance to each of the identified factors from a low-medium-high scale of relevance. Finally, at the end of each interview we gathered information from the participants on which quantitative indicators from local biophysical and socio-economic datasets they deemed suitable to describe the spatial characteristics of the identified factors.

3.2 Data acquisition and indicator selection

In order to quantitatively describe the identified vulnerability factors, we scanned the datasets proposed by the interviewed participants to select suitable indicators matching the meaning of each factor. In addition, we searched databases from authorities at regional (e.g Amt für Geoinformation Thurgau), national (e.g. Swiss Federal Statistical Office (SFSO), Statistical Office of the Republic of Slovenia (SUR)) to the European scale (e.g. European Environment Agency (EEA), European Soil Data



Centre (ESDAC)). Data availability and accessibility proved highly variable. If two different sources offered the same dataset to describe suitable indicators we prioritised the most recent and local dataset, as we aimed for current data with a spatial resolution as high as possible and at least with a subregional resolution of LAU1 regions.

3.3 Pre-processing, normalisation and direction

Depending on the type of data and data availability we gathered continuous and categorical data for both study regions. We transformed the categorical data into (a) continuous data by ordering the categories, or into (b) presence-absence data with the presence of the most important category according to the responses of the interview participants. In order to facilitate the analyses all available datasets were rasterized to grid data with the boundaries of Thurgau and Podravska and their region-specific projection. Thus, we gathered one raster layer for each indicator (r_i) representing the factor quantitatively, if subregional data is available.

In the next step, we directed r_i according to the increasing and decreasing effect on the regions' vulnerability resulting from the interviews with the experts, similar to Meza et al. (2020). We transformed indicator layers with presence-absence data to values of '1', if presence has an increasing effect and to values of '0', if absence has an decreasing effect. Regarding the other indicator layers, we multiplied r_i with '+1', if the indicator increases the vulnerability or we multiplied the values of r_i with '-1', if the indicator decreases the vulnerability. Then, we normalised the directed indicator layers to account for different value ranges and units. Both steps are applied as follows

$$r_{i,d,norm} \in [0,1] = r_i * d \in \{-1,1\} * \frac{r_i - \min_{r_i}}{\max_{r_i} - \min_{r_i}} \quad (1)$$

where $r_{i,d,norm}$ represents the directed and normalised indicator layer, r_i represents the indicator layer with the original values from its datasource(s), d represents the direction factor in order to assign an increasing '+1' or decreasing '-1' effect, \min_{r_i} represents the minimum value across r_i , \max_{r_i} represents the maximum value across r_i . That means $r_{i,d,norm}$ presents with '0' the lowest possible vulnerability according to the indicator layer r_i , and with '1' the highest possible vulnerability according to the indicator layer r_i . All other values are transformed in-between with increasing values representing increasing vulnerability.

3.4 Weighting and aggregation

We applied two different weighting methods to calculate the vulnerability index V : the equal-based weighting method (V_{eq}), and the expert-based weighting method (V_{ex}). For each method the assigned weights to the indicators sum up to 1, and thus can be seen as proportional weights dependent on the method. We aggregated the weighted indicators by summing them up. Practically, we added the weighted raster layers corresponding to the indicators for each method. According to Fig. 1 the agricultural land is very heterogeneously distributed, wherefore we masked the summed indicators with the region-specific agricultural land by setting to zero the indicators' values in non-agricultural areas. By doing so, we accounted for the density of agricultural areas over the whole area within each LAU2 region in the next step. We aggregated the summed indicators at



each LAU2 region within Thurgau and Podravka calculating mean values of vulnerability for each LAU2 region. The final
 170 maps enable a comparison of the two weighting methods for both study regions. The above described weighting methods are
 specified in the following.

For V_{eq} we assigned to each indicator i the same weight $w_{i_{eq}}$ as follows

$$V_{eq} = \sum_{i \in \{1, 2, \dots, n\}} r_{i_{d, norm}} * w_{i_{eq}}, \text{ with } \sum_{i \in \{1, 2, \dots, n\}} w_{i_{eq}} = 1 \quad (2)$$

$$V_{eq} = \sum_{i \in \{1, 2, \dots, n\}} r_{i_{d, norm}} * \frac{1}{n} \quad (3)$$

175 where n represents the total number of all factors, $r_{i_{d, norm}}$ represents the directed and normalised indicator layer (see. eq.
 1), and $w_{i_{eq}}$ represents the weight of indicator i that equals in this method to $\frac{1}{n}$. In eq. 3 we aggregated the weighted rasters by
 summing them up to the equal-based vulnerability method V_{eq} .

To gather V_{ex} , we used the importance rating Imp_r by the interview participants from the low-medium-high scale and the
 frequency, how often a factor was named. We weighted each corresponding indicator i with the expert-weight $w_{i_{ex}}$ as follows

$$180 \quad w_{i_{ex}} = \sum_{i \in \{1, 2, 3\}} i * Imp_r * \frac{p_{i, r}}{W_{tot}} \quad (4)$$

$$W_{tot} = \sum_{i \in \{1, 2, \dots, n\}} \sum_{i \in \{1, 2, 3\}} i * Imp_r * p_{i, r} \quad (5)$$

where $w_{i_{ex}}$ represents the experts' weight of indicator i , Imp_r represents the importance rate on the low-medium-high scale
 translated into the integers 1-2-3, $p_{i, r}$ represents the number of the experts who weighted indicator i with the weight r .
 W_{tot} represents the total weight assigned to all indicators, and n represents the total number of all indicators. Analogue to the
 185 equal weighting method, we weighted each indicator layer $r_{i_{d, norm}}$ with the corresponding weight $w_{i_{ex}}$ and summed up these
 weighted raster layers to calculate V_{ex} as follows

$$V_{ex} = \sum_{i \in \{1, 2, \dots, n\}} r_{i_{d, norm}} * w_{i_{ex}}, \text{ with } \sum_{i \in \{1, 2, \dots, n\}} w_{i_{ex}} = 1 \quad (6)$$

3.5 Participatory validation

In order to validate the results, we conducted one group interview involving the previously consulted experts for each case
 190 study. We held the group interviews between the 10th and 21st June 2022 following the same structure (reported in S2). Firstly,
 we asked the experts to identify the most and least vulnerable subregions across Thurgau and Podravka based on their past
 experiences in dealing with drought events. Secondly, we presented the maps displaying the most important vulnerability fac-
 tors and asked their opinions on the differences displayed across the study regions. Thirdly, we presented the final (aggregated)
 vulnerability maps according to the two weighting methods and we asked for feedback on both the most and least vulnerable
 195 regions. Finally, we asked them to provide their opinion on the differences between the maps based on the two weighting
 methods and which one provides a better description of real drought vulnerability conditions in agriculture.



4 Results

4.1 Factors and indicators describing the region specific vulnerability

During the semi-structured interviews the experts named in total 31 unique factors describing vulnerability to agricultural drought impacts (see Table 1). The experts identified 10 common factors for both study regions, whereas they identified 6 factors solely for Thurgau and 13 factors solely for Podravska (see Table 1). The common factors were altitudes, distance to large water bodies, share of drought resistant crop types, farm size, humus content, presence of irrigation infrastructure, slope, soil texture, topsoil depth and water holding capacity. The unique factors for Thurgau were south facing area, share of intensive livestock, share of pastures, political conservative vote, share of specialty crops and type of irrigation infrastructure, whereas the unique factors for Podravska were distance to mountains, landscape diversity, water permits, accessibility to local food market, farm diversification, intensity of farming, absence of drought policies, food price, agro-technical measures, clear landownership, compensations, farmers' age and farmers' education. Thus, the identified factors describe various aspects of vulnerability, such as geographic conditions (e.g. elevation, south facing), hydrological characteristics (e.g. distance to large water bodies), soil characteristics (e.g. water holding capacity), agricultural practices (e.g. intensive farming), agricultural infrastructure (e.g. irrigation infrastructure), farmers' education, and policies (e.g. compensations). In comparison, the experts from Podravska claimed more and a wider range of factors describing the regions' vulnerability than the experts from Thurgau.

For both regions we found suitable indicators with subregional data from various sources to cover a majority of the identified vulnerability factors (see Table 1, Fig. S1). However, we were not able to support with subregional data 6 out of 18 factors in Thurgau and 11 out of 25 factors in Podravska. For both regions we used data from the European Environment Agency (EEA, 2022) and from the European Soil Data Centre (ESDAC, 2022). Regarding Thurgau, we used data from national sources as the Swiss Federal Statistical Office (SFSO, 2022) and the Swiss Federal Office of Environment (FOEN, 2022), and data from regional sources the Office for Geoinformation by Thurgau (Amt für Geoinformation Thurgau, 2022). Regarding Podravska, we used data from the Statistical Office of the Republic of Slovenia (SURSTAT, 2022) and from the INSPIRE Slovenian Data Portal (INSPIRE, 2022).

Some factors could be easily described through indicators supported by data available at the subregional scale. In these cases the factors, originally described by the experts, were fully represented by the selected indicators. For example, with the digital elevation model by the EEA we could represent the factor slope with the indicator slope measured in radians. In addition, with a combination of shapefiles by the EU-Hydro - River Network Database respectively (EU-Hydro) for large rivers, reservoirs and lakes we could represent the factor distance to large water bodies by calculating this distance for each location in both case studies. For other factors we were forced to select proxy indicators which were either more specific or only covering a partial aspect of the overall concept provided by the experts during the interviews. For example, for the factor presence of irrigation infrastructure we found information about the presence or absence of permanently irrigated agricultural land across both study regions and used this as a representative indicator. For example, in Thurgau we used the indicator dominant soil texture with classes from clay to sand to describe the factor soil texture as a whole, and we used the indicator livestock units for the factor




Figure 3. The identified vulnerability factors for (a) Thurgau and (b) Podravka that could be supported with subregional data. The size of each factor within the bars corresponds to the weight comparing the equal weighting method with the expert weighting method. For an interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.

230 share of intensive livestock. This way, we could support a majority of 12 factors in Thurgau and a majority of 14 factors in Podravka with data from indicators (see Table 1).

For some indicators the data is static over time, such as elevation. For indicators that show a temporal development we were able to find recent data, such as number of farms > 30 ha (from 2019 in Thurgau) and average utilised agricultural area per agricultural holding (from 2010 in Podravka). Regarding the available spatial resolution in Thurgau, we found shapefiles with different spatial resolution for five indicators, raster files with a resolution at least of 25 hectares for four indicators, and data for LAU1 regions for three indicators. In Podravka, we found raster files with a resolution of at least 25 hectares for six indicators, data for LAU2 regions for four indicators, with different spatial resolution for three indicators, and point data for one indicator. Most of the factors are represented by indicators with continuous data (respectively $n = 6$ and $n = 10$), followed by indicators with categorical data ($n = 4$ and $n = 3$) and by indicators with binary data ($n = 2$ and $n = 1$).

240 4.2 Vulnerability Thurgau

Thurgau's vulnerability is calculated with 12 factors for both the equal weighting method and the expert weighting method (see Fig. 3). The following five factors were attributed a greater importance by the experts and were therefore assigned with higher weights and accounting for a total of 71 % of the total vulnerability: distance to large water bodies, humus content, presence of irrigation infrastructure, soil texture and water holding capacity, whereas the other factors lost weight in the expert weighting method, specifically the factor slope.



Table 1. Set of factors the interview participants used to describe vulnerability to agricultural drought impacts and that could be supported with data in Thurgau and Podravaska. The data source, data type, data range, the direction, the latest update and the spatial resolution of the data is shown.

Thurgau					
Factor¹	Indicator	Direction	Latest update	Data type	Data source
Altitudes	Altitudes [masl]	↘	2016	Continuous, Raster	EU-DEM
Distance to large water bodies	Distance to large water bodies [km]	↗	2020	Continuous, Raster	EU-Hydro and FOEN
Farm size	No. of farms \geq 30 ha [no.]	↘	2019	Continuous, Vector	SFSO
Humus content	Topsoil organic carbon content [%]	↘	2006	Classes, Vector	ESDAC
Presence of irrigation infrastructure	Permanently irrigated agricultural land	↘	2018	Binary, Raster	CLC
Share of intensive livestock*	Livestock units [LU]	↗	2020	Continuous, Vector	SFSO
Share of pastures*	No. of farms specialized for pasture farming [no.]	↘	2020	Continuous, Vector	SFSO
Slope	Slope [rad]	↗	2014	Continuous, Raster	CLC
Soil texture	Dominant soil texture	↗	2005	Binary, Vector	Amt für Geoinformation Thurgau
Southfacing area*	Southfacing	↗	2005	Binary, Vector	Amt für Geoinformation Thurgau
Topsoil depth	Dominant topsoil depth	↗	2005	Classes, Vector	Amt für Geoinformation Thurgau
Water holding capacity	Topsoil available water capacity [mm]	↘	2006	Classes, Vector	ESDAC
Podravaska					
Factor¹	Indicator	Direction	Latest update	Data type	Data source
Access to local food market*	Agricultural holdings with main destination for sale [%]	↘	2010	Continuous, Vector	SURS
Altitudes	Altitudes [masl]	↗	2017	Continuous, Raster	INSPIRE
Distance to large water bodies	Distance to large water bodies [km]	↗	2020	Continuous, Raster	EU-Hydro and INSPIRE
Distance to mountains*	Distance to mountains [m]	↗	2008	Continuous, Raster	European Mountain Areas
Farm diversification*	Average no. of permanent beds per agricultural holding [no.]	↘	2009/2010	Continuous, Vector	SURS
Farm size	Average utilised agricultural area per agricultural holding [ha]	↘	2010	Continuous, Vector	SURS
Humus content	Topsoil organic carbon content [%]	↘	2006	Classes, Vector	ESDAC
Intensity of farming*	Average total yield per municipality [t]	↗	2010/2020	Continuous, Vector	SURS and INSPIRE
Landscape diversity*	Landscape diversity (Shannon's evenness index)	↘	2018	Continuous, Raster	CLC
Presence of irrigation infrastructure	Permanently irrigated agricultural land	↘	2018	Binary, Raster	CLC
Slope	Slope [rad]	↗	2017	Continuous, Raster	INSPIRE
Soil texture	Subsoil textural class	↘	2006	Classes, Vector	ESDAC
Water holding capacity	Topsoil available water capacity [mm]	↘	2006	Classes, Vector	ESDAC
Water permits*	Discharge from permits [l/y]	↘	2012	Continuous, Point data	INSPIRE

¹marked with '*' for factors only mentioned for either Thurgau or Podravaska, the remaining were mentioned for both study regions.

Each mapped factor shows higher and lower vulnerabilities across the region according to the indicator values (see Fig. S2). The increasing and decreasing effect of the factor depends on the direction defined by the experts (see Table 1). The factor with the highest weight according to the experts is the presence of irrigation infrastructure which accounts for 19 % of the total vulnerability. Permanently irrigated land is relatively equally distributed across Thurgau. The LAU2 regions covered most



250 with irrigation infrastructure are in the central North of Thurgau (Homburg, Kemmental). Regions least covered with irrigation
infrastructure are in the higher elevation-regions of the South (Fischingen) and east of the city of Constance and along the coast
(Berlingen, Gottlieben). The second most important factor is distance to large water bodies accounting for 17 % of the total
vulnerability. This mapped factor presents heterogeneous distances across the region, with lower vulnerability along the coast,
the rivers Rhine, Thur and Murg, and around larger water reservoirs and lakes mostly located in the Northeast. Subsequently, the
255 highest vulnerabilities related to this factor occur in the LAU2 regions in-between the rivers and lakes (Homburg, Raperswilen,
Wäldi) and in the mountainous South (Fischingen). The third most important factor in Thurgau is humus content accounting
for 13 % of the total vulnerability. The best fitting indicator supporting this factor is called Topsoil organic carbon content [%]
and offered by the European Soil Database (ESDAC, 2022). This categorical indicator presents higher carbon content and thus
a lower vulnerability in the LAU2 regions located along the river Thur (Hüttlingen, Fellben-Wellhausen, Hohentannen). On
260 the other hand, most areas across Thurgau present very low carbon content leading to 21 LAU2 regions with carbon content <
1 % spread across the study region. The fourth and fifth most important factors are soil texture and water holding capacity. For
soil texture we used data from the Amt für Geoinformationen, which presents the dominant topsoil texture with five classes
from clay to clay rich sand that we transformed in three classes with sand as the most vulnerable texture and clay as the
least vulnerable texture, as described by the experts (see Table 1). Only a few areas spread across Thurgau are covered with
265 sandy soils and are thus most vulnerable (Basadingen-Schlattingen, Felben-Wellhausen, Lommis), whereas Thurgau is mostly
covered by clay and silt decreasing the vulnerability. For the factor Water holding capacity we used the indicator Topsoil
available water capacity from the European Soil Database (ESDAC, 2022). This indicator presents a very high capacity (>190
mm) along the river Thur (Märstetten, Felben-Wellhausen, Wigoltingen). In contrast, lowest capacities (<99 mm) are presented
for several LAU2 regions along the coastline and close to the river Rhine. Accordingly, the LAU2 regions in the South are less
270 vulnerable and the LAU2 regions in the North are more vulnerable with respect to this factor.

In order to represent the overall vulnerability, we summed all factor raster layers and set to zero all non-agricultural areas in
order to compute a mean value of vulnerability for each LAU2 region according to the equal (V_{eq}) and expert (V_{ex}) weighting
methods (Fig. 4). To identify the agricultural areas we used regional data from the Office for Geoinformation by Thurgau (Amt
für Geoinformation Thurgau, 2022).

275 The five highest V_{eq} towards agricultural drought impacts occur in the central north and northeast part of Thurgau with V_{eq}
> 0.25. The highest V_{eq} is indicated for Raperswilen ($V_{eq} = 0.29$), Berg ($V_{eq} = 0.27$), Dozwil ($V_{eq} = 0.26$), Wigoltingen ($V_{eq} =$
0.26), and Lommis ($V_{eq} = 0.25$). All these LAU2 regions show a high share of agricultural land over the total region land, hence
making them more exposed to possible drought impacts. In particular, although Raperswilen Wigoltingen and Lommis show
a larger presence of irrigation infrastructure, many other factors, such as distance to large water bodies, humus content and
280 water holding capacity, contribute to an increased vulnerability. Differently from Raperswilen Wigoltingen and Lommis, the
LAU2 region of Dozwil and Berg show lower presence of irrigation infrastructure, but low humus content and water holding
capacity. The five lowest V_{eq} are located in the central and southern part of Thurgau with $V_{eq} < 0.07$. The lowest V_{eq} values
are reported for Fischingen ($V_{eq} = 0.05$), Bichelsee-Balterswil ($V_{eq} = 0.05$), Bettwiesen ($V_{eq} = 0.06$), Wilen ($V_{eq} = 0.06$),
Wuppenau ($V_{eq} = 0.07$). For all these regions, almost all underlying factors show moderate to low values besides presence of



285 irrigation infrastructure and share of intensive livestock having higher values. Additionally, a low share of agricultural land
drives the low vulnerability. The highest vulnerability calculated with the expert-based method shows some differences in the
spatial distribution of LAU2 regions with the northwest increasing in vulnerability. In particular, the region of Raperswil
is again ranked as first ($V_{ex} = 0.35$), followed by Herdern ($V_{ex} = 0.31$), Homburg ($V_{ex} = 0.30$), Wäldi ($V_{ex} = 0.30$), and
Uesslingen-Buch ($V_{ex} = 0.28$). For all these regions, the factors water holding capacity, humus content and distance to large
290 water bodies present high vulnerability, apart from Uesslingen-Buch with relatively short distance to the river Thur. In contrast,
the factor soil texture presents these regions covered mostly by clay, and the factor presence of irrigation infrastructure displays
almost all regions covered with irrigation systems, both decreasing vulnerability. However, the combined effect of a large share
of agricultural land and visible patches of high vulnerability values across the factors distance to large water bodies, humus
content and water holding capacity contribute to the final high vulnerability values. The LAU2 regions with the five lowest
295 V_{ex} are almost the same as for the equal-weights method with slightly different changes. The lowest values are reported for
Fischingen ($V_{ex} = 0.04$), Bichelsee-Balterswil ($V_{ex} = 0.05$), Wilen ($V_{ex} = 0.06$), Bettwiesen ($V_{ex} = 0.06$) and Berlingen (V_{ex}
 $= 0.06$). This latter region differs from the areas with lowest values for the equal weighting scheme and although having a low
presence of irrigation infrastructure, it shows, among others, low share of intensive livestock, high values of farm size as well
as clay rich sand as dominant soil texture.

300 From a comparative point of view, (Fig. 5, bottom row) results from Vex point to the northwest part of Thurgau with higher
vulnerability to drought in agriculture driven by the spatial cluster from the factors distance to large water bodies, water holding
capacity, and soil texture. The LAU2 regions showing higher vulnerability with V_{ex} (+0.04) are mostly located between the
river Rhine and Lake Constance (Homburg, Raperswil, Herdern, Basadingen-Schlattigen, Wäldi). In contrast, regions with
decreased vulnerability with V_{ex} (-0.02) are located in the Northeast (Dozwil, Salmsach, Egnach) and North (Gottlieben), and
305 in central Thurgau (Weinfelden).

During the participatory validation with Thurgau's experts the most important factors (see Fig. 4) have been discussed
regarding their correctness. The expert agreed that in general all these factors present plausible patterns across the region.
However, they expected more irrigation in the 'Thur valley', the valley along the river Thur (see Fig. 1). They explained
that this valley is intensively used for agricultural purposes, and therefore expected more irrigation infrastructure. As well,
310 they expected less clay-rich soil with clay in this area, because of the high amount of gravel in the valley, typically a sign
for sandy soils. They concluded that due to the intensive use for agriculture in the Thur valley, the soil conditions could
be modified due to adapted cultivation techniques. Regarding the factor soil texture, they confirmed the clay-rich, but sandy
soils in 'Unterthurgau' covering the regions Diessenhofen, Schlatt, and Basadingen-Schlattigen in northwestern Thurgau,
and the 'Lauche valley', the valley along the river Lauche (see Fig. 1). Further they confirmed the clay-rich soils along the
315 so-called 'Seerücken', a hill range up to 723 masl between Lake Constance and the river Thur (see Fig. 1, https://peter-hug.ch/lexikon/1888_bild/45_0487). Regarding the factor humus content, they questioned the low amount of topsoil organic carbon
content along the Seerücken. Regarding the mapped vulnerability of agriculture to drought, the experts confirmed that both
maps present reasonable differences across Thurgau with higher vulnerability in the North compared to the South. However,
the experts highlighted the vulnerability map based on the expert weighting method, because of the better presentation of



320 known hotspot regions. In specific, the experts pointed to hotspot regions along the Seerücken, and to the hotspot Unterthurgau
(see Fig. 1). The expert weighting method presents all these hotspot regions with higher vulnerability compared to the equal
weighting method (see Fig. 5, Table S5).

The reasons for, as well as the differences between the hotspot regions with high vulnerability have been discussed during
the interview. The experts report the Seerücken and Unterthurgau as impacted regularly in the past. However, they explained
325 that Unterthurgau has access to (irrigation) water from the river Rhine and from a large groundwater aquifer filled by the Lake
and the Rhine. Thus drought typically leads to impacts on agriculture, if soil moisture is abnormally low. This is different
regarding the Seerücken, during drought conditions typically characterised by low soil moisture, but additionally by low river
discharges leading to impacts on agriculture. The experts pointed to the Lauche valley that is specifically better highlighted by
the expert weighting method as the regions Stettfurt and Lommis are presented more vulnerable (+ 0.04, + 0.02) compared to
330 other regions (Table S5). The experts presented the Lauche as a medium-sized river delivering water for several uses that is
quickly overused during drought conditions. Subsequently, in the past user conflicts on irrigation occurred regularly first in the
Lauche valley, which is also the case in the current dry situation (21st June, 2022).

4.3 Vulnerability Podravška

Podravška's vulnerability is calculated considering 14 factors for both the equal weighting and the expert weighting method
335 (see Fig. 3). Within them, the four factors access to local food market, farm diversification, presence of irrigation infrastructures
and soil texture increased their weights in the expert weighting scheme accounting for a total of 56 % of the overall vulnerability
weight.

According to the experts, the most influential factor is the presence of irrigation infrastructure, which accounts for 15 % of the
total vulnerability. Irrigated areas are mainly located in the flat areas along the Drava and Polskava rivers which are crossing
340 from North-West and South-West to East the Podravška region (Miklavž na Dravskem Polju, Starše, Kidričevo, Hajdina,
Markovci, Gorišnica). Areas covered the least with irrigation infrastructure are mainly located in the outer parts of Podravška
and at higher elevation. The second and third most important factors are soil texture and farm diversification accounting for
14 % of the total vulnerability. For soil texture we use data on the subsoil textural classes provided by the European Soil
Database (ESDAC, 2022) transformed in three classes with sand as the most vulnerable texture and clay as the least vulnerable
345 texture according to the expert's opinion (see Table 1 1, Fig. 6). Although most of the areas in Podravška show coarser soil
texture values with higher contribution to vulnerability, some clay and silt soils patches are located in the central southern area
(municipalities of Slovenska Bistrica, Rače–Fram, Kidričevo, Majšperk and Makole), along the Pesnica river in the north east
part of Podravška and also close to the municipalities of Ormož and Središče ob Dravi. The factor farm diversification refers to
the presence of additional incomes for farmers, particularly focussing on the possibility of hosting tourists, as highlighted by
350 the experts. As for the factor soil texture, the factor farm diversification, shows a general condition of homogeneous low values.
Only the municipalities of Maribor, Šentilj, Cerkvenjak and Ormož showed higher values in the number of permanent beds
per agricultural holding (respectively of 4.86, 8.84, 7.40 and 8.55, see Table S6) which is making them less dependent on the
agricultural income, hence more diversified and less vulnerable to potential drought impacts. The fourth most important factor




Figure 4. Thurgau’s most important factors (bold title) masked with agricultural used land and describing the regions’ vulnerability according to the experts (see Fig. 3) supported by data of the indicators (see legend). The factor’s increasing or decreasing effect on the vulnerability is indicated by the arrow in the map (bottom right) and by the colour choice (the darker the colour, the higher the vulnerability). LAU2 regions are indicated by black borders and labelled when showing relatively high or low vulnerability. For an interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.

355 access to local food market showed a more heterogeneous situation with lower values mainly in the northern part (Šentilj) and in the southern part (Poljčane, Makole, Žetale, Podlehnik, Cirkulane and Zavrč). The highest values were found for the municipality of Kidričevo where more than 66 % of the agricultural holdings have 'for sale' as their main destination of their products, followed by Gorišnica with 58.9 %, Rače–Fram with 58.7 % and Hoče – Slivnica with 56.8 %. The lowest values




Figure 5. The mapped results of equal weighting, expert weighting methods, and their difference across Podravška. On the left, masked with agricultural used land and before aggregation on LAU2 regions (black borders). On the right, after aggregation for each LAU2 region. LAU2 regions are labelled when showing relatively high or low vulnerability, or relevant differences between the weighting methods. For an interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.

were found for Podlehnik 18.8 % and Cirkulane 14.4 % in the southern part, where the agricultural holding products are mainly intended for own consumption.

360 Similarly to the Thurgau case study, we summed all factor raster layers and set to zero all non-agricultural areas in order to compute a mean value of vulnerability for each LAU2 region according to the equal (V_{eq}) and expert (V_{ex}) weighting methods



(Fig. 7). To identify the agricultural areas we used regional data on agriculture parcels with declared crop from the INSPIRE Slovenian Data Portal.

The five highest V_{eq} towards agricultural drought impacts occur in the northeast part of Podravska with $V_{eq} > 0.26$. The
365 highest V_{eq} is indicated for Trnovska vas ($V_{eq} = 0.29$), followed by Sveti Jurij v Slov. goricah ($V_{eq} = 0.28$), Destrnik ($V_{eq} =$
0.27), Benedikt ($V_{eq} = 0.27$) and Sveti Andraž v Slovenskih goricah ($V_{eq} = 0.26$). Moreover, other LAU2 regions with high
values are still located in the northeast of Podravska. These regions show high share of agricultural areas for each LAU2 region
and a combination of low values in terms of presence of irrigation infrastructure, soil texture, farm diversification and access
to local food market (Fig. 6), but also for other factors with lower weight on the overall vulnerability such as distance to
370 mountains, intensity of farming and humus content (Fig. S3).

The five lowest V_{eq} towards agricultural drought impacts are located in the northwest and southwest parts of Podravska with
 $V_{eq} < 0.1$. In particular, Ruše shown the lowest V_{eq} value ($V_{eq} = 0.03$), followed by Lovrenc na Pohorju ($V_{eq} = 0.04$), Selnica
ob Dravi ($V_{eq} = 0.07$), Maribor ($V_{eq} = 0.08$) and Poljčane ($V_{eq} = 0.1$). These regions show a low share of agricultural areas for
each LAU2 region since they are located close to mountains with forest as the main land cover type or with large urban areas
375 as for the case of Maribor. Within these regions, the agricultural areas are located along the Drava river and benefit by the low
distance to the main urban areas in terms of farm diversification.

The expert weighting method shows the highest values of vulnerability in the northeast part of Podravska with the five
highest LAU2 regions being almost the same as for the equal weighting and only changing their rank as follows: Sveti Jurij v
Slov. goricah ($V_{ex} = 0.35$), Benedikt ($V_{ex} = 0.32$), Trnovska vas ($V_{ex} = 0.31$), Destrnik ($V_{ex} = 0.30$) and Pesnica ($V_{ex} = 0.29$,
380 Fig. 7b). The shift in highest values of LAU2 regions is visible in the difference between the two maps at the bottom of Fig. 7
with Sveti Jurij v Slov. goricah showed the highest increase (+0.07) driven among others by the large share of agricultural areas
without the presence of irrigation, a coarse soil texture and low values of farm diversification.

The lowest values of vulnerability from the expert weighting method are also still located in the northwest and southwest
part of Podravska without changes in the rank of the LAU2 regions going from Ruše ($V_{ex} = 0.04$), Lovrenc na Pohorju ($V_{ex} =$
385 0.05), Selnica ob Dravi ($V_{ex} = 0.1$), Maribor ($V_{ex} = 0.1$) and Poljane ($V_{ex} = 0.14$).

When comparing the expert weighting results with the equal weighting results (Fig. 7, at the bottom) there are only positive
variations coming from higher weights by the experts compared to the equal weighting scheme. This means that there is a
generalised worsening of vulnerability conditions as represented by the experts in Podravska. The LAU2 regions showing
higher vulnerability compared to $V_{ex} (\geq +0.05)$ are mostly located in the North and central South of Podravska: Sveti Jurij v
390 Slov. goricah, Gorišnica, Kungota, Benedikt, and Starše. In contrast the regions with least changes between the two methods
are spread across Podravska (Središče ob Dravi, Kidričevo, Ruše, Lovrenc na Pohorju, Sveti Andraž v Slovenskih goricah).

During the participatory validation with the experts from Podravska, we discussed each map of the most important vulnera-
bility factors (see Fig. 6). The experts agreed that most shallow soils with coarse texture and subsequently higher vulnerability
are close to rivers and springs. Therefore, they pointed to the so-called 'Drava valley' along the river Drava and the lowlands
395 that are presented with coarse soils by the factor Soil texture in Fig. 6. Regarding the presence of irrigation infrastructure, the
expert confirmed that still a large part of Podravska is not irrigated yet, which is causing problems in the North, Northeast, and



South. In contrast, they expected the East (especially Ormož) to be more covered with irrigation infrastructure, and questioned if the data source from the EEA Copernicus Land Monitoring Service published in 2018 is correctly displaying the current situation. The experts agreed on the spatial distribution of the factor farm diversification with the regions Maribor, Ormož and Šentilj showing farms with a higher touristic share due to the presence of vineyards and orchards often offering touristic opportunities. Regarding the factor Access to local food market, they agreed that farms around cities usually have higher access to food markets, as they can sell their products, which is partly displayed around Maribor. However, they expected the East part of Podravska (e.g. Ormož) to have less access to food markets, as the markets there are known to be less developed.

Regarding the final vulnerability maps of agriculture to drought, the experts did not prefer one map out of the two methods, as they both show the main vulnerable regions across Podravska with higher vulnerability in the Northeast, in the South and in the centre of Podravska. They pointed to the fact that subjectivity is less prominent with the equal weighting method, and that the expert weighting method should be validated in depth with local farmers working on the field. In particular, they pointed to the Drava valley and the lowlands called 'Ptuj field' and 'Drava field' (see Fig. 1), as these regions experienced drought impacts, as they are characterised by shallow soils with low water holding capacity and subsequently more vulnerable to drought. Nevertheless, the experts agreed that the Drava valley is among the most vulnerable regions in Podravska, because of the intensive production, whereas the vulnerability maps display the North and South more vulnerable. Focusing on the Northeast and the South the experts pointed to the fact that these regions are more hilly, and additionally the water distribution is not organised in order to enable irrigation, wherefore these regions are correctly displayed with higher vulnerability. The experts agreed that the Western part of Podravska is less vulnerable, as this mountainous region is not intensively used for agriculture, but more covered with forests. They also confirmed that the East is less vulnerable (Ormož), although more recent data should be considered to capture the local characteristics of lower access to local food market and higher values of presence of irrigation infrastructure.

5 Discussion

5.1 Sensitivity of region-specific vulnerability

Results from the vulnerability assessment highlighted how the type of weighting methods can affect each region's specific vulnerability. The implementation of different weighting methods allowed the creation of possible 'scenarios' of vulnerability conditions. Through the equal weighting we provided a neutral description of the geographical variation of vulnerability conditions. Comparably, the expert weighting method showed similar spatial patterns with a higher vulnerability. Comparing the outcome of the two different methods provided a wider perspective on potential conditions to address, an advantage acknowledged by the experts during the group interview. In particular, keeping such a perspective helps to inform stakeholders and decision makers when results are particularly difficult to validate through quantitative data only.

The equal weighting method is a simple and easily computable method (Kienberger et al., 2016; Becker et al., 2014; Schneiderbauer et al., 2020) showing some limitations. In particular, the weight of each factor is dependent on the number of factors included in the analyses. This means, the more factors are considered the less weight is assigned to each factor. This reduces




Figure 6. Podravskas’s most important factors (bold title) masked with agricultural used land and describing the regions’ vulnerability according to the experts (see Fig. 3) supported by data of the indicators (see legend). The factor’s increasing or decreasing effect on the vulnerability is indicated by the arrow in the map (bottom right) and by the colour choice (the darker the colour, the higher the vulnerability). LAU2 regions are indicated by black borders and labelled when showing relatively high or low vulnerability. For an interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.

430 the effect of each factor on the final vulnerability map. Subsequently factors with a wider range get more influence. In our study regions, these weights were rather similar, as we could identify and supply 12 factors with data in Thurgau meaning each factor weighs $112 \approx 0.08$, and 14 factors in Podravka meaning each factor weighs $114 \approx 0.07$.

The expert weighting method uses the experts’ opinion built on specific knowledge of regional drought leading to agricultural impacts in the study regions. For example, the factor distance to large water bodies received substantially higher weight in both study regions than the equal weights. Thus, the expert weighting method benefits by integrating region-specifics that might be missed by the other method. Moreover, this approach guarantees a higher level of involvement of local stakeholders (Menk et al., 2022).

Looking at the vulnerability factors identified by the experts, our study presents several common factors between both study regions. Nevertheless, the majority of the identified factors were solely mentioned for one of the regions. This demonstrates the region-specifics and accordingly the limitation to extrapolate the expert weights or factor selection to other regions. In




Figure 7. The mapped results of equal weighting, expert weighting methods, and their difference across Podravska. On the left, masked with agricultural used land and before aggregation on LAU2 regions (black borders). On the right, after aggregation for each LAU2 region. LAU2 regions are labelled when showing relatively high or low vulnerability, or relevant differences between the weighting methods. For an interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.

this study, only for one out of ten common factors the assigned direction by the experts differed between the two case studies (i.e. altitudes) showing similar underlying processes leading to drought impacts. Regarding the common factors, in both study



regions the importance rating differed slightly. The factors presence of irrigation infrastructure and soil texture played a major role in both regions and, in conclusion, can be considered generally important for agriculture's vulnerability to drought in
445 pre-Alpine regions.

5.2 Strengths and weaknesses of the mixed method approach

This study applied a mixed-method approach based on qualitative and quantitative information for the assessment of agriculture's vulnerability to drought. First of all, the implementation of a mixed-method approach supports a comprehensive understanding of the region-specifics, as the factors driving or mitigating the regions vulnerability have been discussed during
450 several interviews and analysed quantitatively with the help of indicators. This way, misconceptions and false conclusions are clarified during the discussion with and between the regional experts.

One of the main limitations when performing local analysis is the availability of suitable quantitative regional and local data to support the identified factors. For this reason, the involvement and exchange of information with local experts provided a solid baseline of references and datasets for spatial and temporal data. Nevertheless, for Thurgau we were not able to find
455 subregional data for six out of 18 factors, and for Podraska for 11 out of 25 (Fig. S1). For Thurgau this included e.g., data for the type of irrigation infrastructure, the share of drought prone and drought resistant species, presence of drought management strategies. For Podravska also e.g., food price, farmers age and education, clear landownership, agro-technical measures, and absence of drought policy. Those and more were named by the experts as factors influencing drought vulnerability. Subsequently, they are required to project and better understand potential future impacts. In addition, not all data supporting the
460 factors has proven to be fully suitable to describe the factors highlighted by the experts. For example, the no. of beds per agricultural holding representing the factor farm diversification in Podravska simplifies the initial meaning that farms with complementary activities are less vulnerable to drought. The here selected indicators partly described the corresponding factors. They were included in the assessment as no better data was accessible and with the agreement of the experts throughout the assessment steps. Overall, the collection of local data remains a time and cost consuming process that can be supported by
465 the application of mixed-method approaches to overcome some of this limitation.

When quantitative data was available, we applied data from different time scales. While this introduced some degree of uncertainty to map the current condition, it represented the best possible way to combine multiple information. Moreover, most of the selected factors either are static in time (e.g. altitudes) or require years to show changes at a regional level (e.g. presence of irrigation infrastructures). Moreover, the data was only accessible with different spatial scales. As the aim was to assess and
470 map differences across the study regions, we combined data with different resolutions in a pragmatic way.

For 3 out of 26 mapped factors, we downscaled the coarse resolution to enable data aggregation. This suggests similar conditions in regions that cannot be proven, wherefore we based our final results on mean values in LAU2 regions avoiding overinterpretation. During the participatory validation the experts confirmed the need of higher-resolution vulnerability maps as a starting point for identifying and implementing more local planning and adaptation strategies.

475 Furthermore, we faced the typical challenge of vulnerability assessment to unify various units and value ranges in order to aggregate the data. The literature presents several methods to do so, ranging between building classes and normalising the



values (OECD and JRC, 2008). We normalised the data leading to stronger effects on the final vulnerability the greater the value range. However, this method avoids additional subjectivity on how to define the classes for each factor making them easy to interpret.

480 As this study tailors the vulnerability to the agricultural sector, we therefore applied an agricultural mask; respectively we integrated the land not agriculturally used as zeros in the calculation of the mean values of the LAU2-regions (Section 3.4). This way we excluded land that is not agriculturally used and considered density of agricultural land as one main aspect driving each region's vulnerability with agreement of the experts. Nevertheless, the experts in Thurgau questioned the mask in some smaller regions, as they expected slightly more agricultural used land in the mountainous region of the South and slightly less
485 agricultural used land along the Seerücken in northern Thurgau. The experts of both regions agreed to include the agricultural mask to enable interpretable results that are specific for the agricultural sector.

The implementation of a participatory approach throughout this study enabled a continuous exchange with local experts. At each step of the analysis this exchange provided additional information, especially on local features that cannot or are not fully described by the mapped data. In particular, qualitative narratives, as provided during the semi-structured interviews and
490 the group discussions, integrated and supported a better characterization of local vulnerability conditions. Besides the experts' knowledge of possible data sources, their region-specific knowledge of drought leading to agricultural impacts in the study regions substantially improved the study. In particular, the regional experts can bring their knowledge that external people might not have (as in the case of access to local food markets in Podravska and share of pastures in Thurgau) or report general assumptions that might not be valid for the specific study region.

495 Nevertheless, the vulnerability maps are dependent on the selected experts that participated. Regarding drought leading to agricultural impacts, the perception of important vulnerability factors might vary substantially between the interviewed persons and their background, such as politicians, water suppliers or farmers (van Duinen et al., 2015). To avoid biased ratings and the omission of important factors one option is to reach a saturation point by interviewing as many persons as possible and use a statistical measure to receive a final weight (Glaser et al., 1968; Guest et al., 2006). However, this is time-costly
500 and might display the current opinion and not the best knowledge based on the experience with drought conditions. Another option is to question multiple experts from different fields in order to cover specific possible perceptions (Bogner and Menz, 2009). According to this, we asked regional experts with agricultural, political and scientific background to best cover various aspects (i.e. Department of agriculture and infrastructure in rural areas, Department of environment, water construction and hydrometry, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Slovenian Environment Agency ARSO,
505 Chamber of Agriculture and Forestry KGZS).

Furthermore this approach guarantees the involvement of local stakeholders through a 'bottom-up and participatory appraisal' (Zebisch et al., 2021). The interaction with stakeholders increases the legitimacy of results while supporting a clear communication and bridging the gap between science and society.



5.3 Towards adaptation strategies to decrease vulnerability

510 During the vulnerability assessment, we identified a range of factors characterising vulnerability conditions to drought in agriculture in both regions (Section 4.1). Some factors represent a sensitivity to static physical terrain conditions (e.g. altitude) while others were associated with conditions that can be changed in time, such as presence of infrastructures (e.g. irrigation) or farming features (e.g. livestock presence and farm size). Within this context, it is crucial to understand what are the factors to leverage the maximum decrease in vulnerability through tailored adaptation strategies. In particular, in both regions
515 the involved experts weighted the presence of irrigation infrastructure as the factor with the highest importance. This result provided information at a spatial level (Fig. 4 and 6) on the areas with the highest potential to decrease vulnerability through the implementation of efficient irrigation systems (e.g. drop irrigation).

In Thurgau, humus content, water holding capacity and soil texture are the other most important factors. Their conditions might be somewhat improved by the type and extent of agricultural practices, such as diversified crop rotations, providing
520 organic matters to the soil (e.g. plant residues, organic fertiliser), keep the soil properly limed, and apply site-specific cultivation practices to avoid erosion and compaction (Wezel et al., 2014; Hobbs et al., 2008). While some of these practices might already be locally implemented, a regional variation of these factors' values require widespread practices implemented over years.

In Podravska, farm diversification and access to local food markets presented high potential to decrease vulnerability. In case of farm diversification, farmers could diversify their intake, for example creating or increasing their touristic offers in order to
525 be less dependent on the pure agricultural production. According to the experts, the owners of vineyards and orchards already offer overnight stays in apartments or touristic tours, a strategy that other farms could implement. Moreover, to increase the access to local food markets, farmers rely on the existence of food markets nearby. This is typically the case for farms close to large cities such as Maribor or Ptuj, but not in rural areas distant from urban centres. Improving road connectivity in rural areas to urban centres or developing cooperative partnerships could enable access to food markets with a higher variety of
530 agricultural products, as farms could combine their products and share transportation costs (Mather and Preston, 1980).

While some of these adaptation strategies are already implemented in parts of both regions, our results show which factor to prioritise and where to intervene in order to trigger the largest decrease in agricultural vulnerability to drought conditions. The presented strategies range from land-use planning, over farm cultivation techniques, towards policies that strengthen the agricultural sector to better cope with drought. Further adaptation planning should integrate them with the available local
535 infrastructural and financial resources through efficient communication among the stakeholders from agriculture, policy and science.

6 Conclusions

Drought vulnerability is defined by various regional and local factors, which differ depending on the region's conditions and on the regions' adaptation due to experiences of past drought impacts. In this study, we identified a wide range of vulnerability
540 factors with differences and similarities among the two case study regions highlighting the complexity of vulnerability and the difficulty to upscale the results to other, respectively larger regions, such as the whole Alpine Space. For both case study



regions we could not support all vulnerability factors with data, a restriction that can be interpreted as uncertainty of the presented results. Additionally, the different weighting methods aggregating vulnerability factors can serve as a measure of sensitivity towards the calculation approach. Being aware of the range of sensitivity and uncertainty, the final results can be embedded better in the region specific context, which is essential for any planning and adaptation strategies. Even though this comes along with limitations in the quantitative part of the mixed-method approach, the resulting vulnerability maps were in general confirmed by the regional experts showing that we might not need to draw the complete picture of all facets defining vulnerability. Other regions may benefit from the approach presented here and tailor the participatory approach with their regional experts to validate the quantitative results with region-specific knowledge. This highlights the benefits of the mixed-method approach combining quantitative with qualitative analyses. The results of this vulnerability assessment identified a range of adaptation strategies dependent on regional resources and efficient communication between the agricultural, political and scientific professionals. Many facets of the described adaptation strategies to decrease vulnerability to drought are in accordance with sustainability goals and climate change adaptation demonstrating the need to move from emergency actions to better preparedness. In order to better understand and quantitatively describe feedback relations and interactions between vulnerability factors the development of drought vulnerability models integrating non-linearities is required, a field still highly underexplored.

Author contributions. Ruth Stephan: Conceptualization, Methodology, Data Curation, Formal analysis, Visualisation, Writing - Original Draft, Writing - Review & Editing. Stefano Terzi: Conceptualization, Methodology, Data Curation, Formal analysis, Visualisation, Writing - Original Draft, Writing - Review & Editing. Mathilde Erfurt: Conceptualization, Methodology, Formal analysis, Writing - Original Draft, Writing - Review & Editing. Silvia Cocuccioni: Conceptualization, Methodology, Formal analysis, Writing - Original Draft, Writing - Review & Editing. Kerstin Stahl: Conceptualization, Supervision, Writing - Review & Editing. Marc Zebisch: Conceptualization, Supervision, Writing - Review & Editing.

Competing interests. The authors declare that they have no conflict of interest.

Disclaimer. This research has been supported by the EU Interreg. Alpine Space Programme project ADO (Alpine Space Observatory; grant no. ASP940). Additional funding was provided by the Provincia autonoma di Bolzano – Alto Adige within the AquaMount project (grant no. D59C20000160003)

Acknowledgements. This research was funded by the EU Interreg. Alpine Space Programme project ADO (Alpine Space Observatory; grant no. ASP940). We acknowledge all the experts involved in this study for their time and support. We thank our ADO partners from Podravska



and Thurgau for their effort and support in data collection. The authors acknowledge the Provincia autonoma di Bolzano – Alto Adige
570 Ripartizione Innovazione, Ricerca e Università as a financing institution of the AquaMount project.



References

- Amt für Geoinformation Thurgau: Geoinformation for the Kanton Thurgau, <https://geoinformation.tg.ch/>, last access: 15 March 2022.
- Ayantunde, A. A., Turner, M. D., and Kalilou, A.: Participatory analysis of vulnerability to drought in three agro-pastoral communities in the West African Sahel, *Pastoralism*, 5, 1–11, 2015.
- 575 Becker, D., Renner, K., and Schneiderbauer, S.: Assessing and Mapping Climate Change Vulnerability with the Help of GIS: Example of Burundi: *GI_Forum, Geospatial Innovation for Society*, pp. 101–104, http://austriaca.at/0xc1aa5576_0x0030d408.pdf, 2014.
- Beniston, M. and Stoffel, M.: Assessing the impacts of climatic change on mountain water resources, *The Science of the total environment*, 493, 1129–1137, <https://doi.org/10.1016/j.scitotenv.2013.11.122>, 2014.
- Birhanu, Z., Ambelu, A., Berhanu, N., Tesfaye, A., and Woldemichael, K.: Understanding resilience dimensions and adaptive strategies
580 to the impact of recurrent droughts in Borana Zone, Oromia Region, Ethiopia: A grounded theory approach, *International journal of environmental research and public health*, 14, 118, 2017.
- Birkmann, J., Cardona, O. D., Carreño, M. L., Barbat, A. H., Pelling, M., Schneiderbauer, S., Kienberger, S., Keiler, M., Alexander, D., and Zeil, P.: Framing vulnerability, risk and societal responses: the MOVE framework, *Natural Hazards*, 67, 193–211, 2013.
- Bogner, A. and Menz, W.: The theory-generating expert interview: epistemological interest, forms of knowledge, interaction, in: *Interviewing experts*, pp. 43–80, Springer, 2009.
- 585 Carrão, H., Naumann, G., and Barbosa, P.: Mapping global patterns of drought risk: An empirical framework based on sub-national estimates of hazard, exposure and vulnerability, *Global Environmental Change*, 39, 108–124, <https://doi.org/10.1016/j.gloenvcha.2016.04.012>, 2016.
- CLC: Copernicus Land Monitoring Service - Corine Land Cover, <https://land.copernicus.eu/pan-european/corine-land-cover/>, last access:
590 15 June 2022.
- DMCSEE: Drought Mitigation Centre for Southeastern Europe, *Drought monitoring bulletin: Overview from February to September 2017*, http://www.dmcsee.org/uploads/file/430_dmcsee_bulletin_season2017.pdf, 2017.
- DROUGHT-CH: Early recognition of critical drought and low flow conditions in Switzerland, www.drought.ch, last access: 20 July 2022.
- EEA: European Environment Agency, <https://www.eea.europa.eu/>, last access: 15 March 2022.
- 595 ESDAC: European Soil Data Centre: Thematic centre for soil related data in Europe, <https://esdac.jrc.ec.europa.eu/>, last access: 15 March 2022.
- EU-DEM: Copernicus Land Monitoring Service - EU-DEM, <https://land.copernicus.eu/imagery-in-situ/eu-dem/eu-dem-v1.1/view>, last access: 15 March 2022.
- EU-Hydro: EU-Hydro – River Network Database, Version 1.3, <https://land.copernicus.eu/imagery-in-situ/eu-hydro/eu-hydro-river-network-database?tab=mapview>, last access: 15 March 2022.
- 600 European Mountain Areas: European Mountain Areas - Version 1, <https://www.eea.europa.eu/data-and-maps/data/european-mountain-areas>, last access: 15 March 2022.
- FOEN: Federal Office for the Environment: Federal Centre of Excellence for the Environment, <https://www.bafu.admin.ch/bafu/en/home.html>, last access: 15 March 2022.
- 605 GIZ, E.: Risk supplement to the vulnerability sourcebook, *Guidance on How to Apply the Vulnerability Sourcebook's Approach with the New IPCC AR5 Concept of Climate Risk*, 2017.



- Glaser, B. G., Strauss, A. L., and Strutzel, E.: The discovery of grounded theory; strategies for qualitative research, *Nursing research*, 17, 364, 1968.
- González Tánago, I., Urquijo, J., Blauhut, V., Villarroya, F., and de Stefano, L.: Learning from experience: a systematic review of assessments
610 of vulnerability to drought, *Natural Hazards*, 80, 951–973, <https://doi.org/10.1007/s11069-015-2006-1>, 2016.
- Guest, G., Bunce, A., and Johnson, L.: How many interviews are enough? An experiment with data saturation and variability, *Field methods*, 18, 59–82, 2006.
- Hagenlocher, M., Meza, I., Anderson, C. C., Min, A., Renaud, F. G., Walz, Y., Siebert, S., and Sebesvari, Z.: Drought vulnerability and risk assessments: state of the art, persistent gaps, and research agenda, *Environmental Research Letters*, 14, 083002,
615 <https://doi.org/10.1088/1748-9326/ab225d>, 2019.
- Hanel, M., Rakovec, O., Markonis, Y., Máca, P., Samaniego, L., Kyselý, J., and Kumar, R.: Revisiting the recent European droughts from a long-term perspective, *Scientific reports*, 8, 1–11, 2018.
- Hartl-Meier, C., Zang, C., Dittmar, C., Esper, J., Göttlein, A., and Rothe, A.: Vulnerability of Norway spruce to climate change in mountain forests of the European Alps, *Climate Research*, 60, 119–132, 2014.
- 620 Haslinger, K. and Blöschl, G.: Space-Time Patterns of Meteorological Drought Events in the European Greater Alpine Region Over the Past 210 Years, *Water Resources Research*, 53, 9807–9823, <https://doi.org/10.1002/2017WR020797>, 2017.
- Hobbs, P. R., Sayre, K., and Gupta, R.: The role of conservation agriculture in sustainable agriculture, *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 363, 543–555, <https://doi.org/10.1098/rstb.2007.2169>, 2008.
- Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B. J., Elmore, A. C., Emmer,
625 A., Feng, M., Fernández, A., Haritashya, U., Kargel, J. S., Koppes, M., Kraaijenbrink, P. D. A., Kulkarni, A. V., Mayewski, P. A., Nepal, S., Pacheco, P., Painter, T. H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A. B., Viviroli, D., Wada, Y., Xiao, C., Yao, T., and Baillie, J. E. M.: Importance and vulnerability of the world’s water towers, *Nature*, 577, 364–369, <https://doi.org/10.1038/s41586-019-1822-y>, 2020.
- INSPIRE: Slovenian Data Portal, <https://eprstor.gov.si>, last access: 15 March 2022.
- 630 IPCC: Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation: A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change.
- Kienberger, S., Borderon, M., Bollin, C., and Jell, B.: Climate change vulnerability assessment in Mauritania: Reflections on data quality, spatial scales, aggregation and visualizations, *GI_Forum Journal*, 1, 167–175, 2016.
- Komac, B., Ferik, M., Pipan, P., Tičar, J., and Zorn, M.: Natural Hazards in Slovenia, *The Geography of Slovenia. World Regional Geography Book Series*, pp. 259–277, https://doi.org/10.1007/978-3-030-14066-3_17, 2019.
- 635 Laaha, G., Gauster, T., Tallaksen, L. M., Vidal, J.-P., Stahl, K., Prudhomme, C., Heudorfer, B., Vlnas, R., Ionita, M., van Lanen, H. A. J., Adler, M.-J., Caillouet, L., Delus, C., Fendekova, M., Gailliez, S., Hannaford, J., Kingston, D., van Loon, A. F., Mediero, L., Osuch, M., Romanowicz, R., Sauquet, E., Stagge, J. H., and Wong, W. K.: The European 2015 drought from a hydrological perspective, *Hydrology and Earth System Sciences*, 21, 3001–3024, <https://doi.org/10.5194/hess-21-3001-2017>, 2017.
- 640 LID: Landwirtschaftlicher Informationsdienst: Thurgauer Landwirtschaft Zum Anfassen, https://www.lid.ch/fileadmin/lid/Produkte/Kantonsbroschueren/55263_Broschuere_Thurgauer-Landwirtschaft.pdf, 2007.
- Martin, R., Linstädter, A., Frank, K., and Müller, B.: Livelihood security in face of drought—assessing the vulnerability of pastoral households, *Environmental Modelling & Software*, 75, 414–423, 2016.



- Mather, J. W. and Preston, H. J.: Cooperative benefits and limitations, Cooperative information report; no. 1, sec. 3, <https://www.rd.usda.gov/files/cir1sec3.pdf>, 1980.
- 645
- Melkonyan, A.: Environmental and socio-economic vulnerability of agricultural sector in Armenia, *Science of the Total Environment*, 488, 333–342, 2014.
- Menk, L., Terzi, S., Zebisch, M., Rome, E., Lückerrath, D., Milde, K., and Kienberger, S.: Climate Change Impact Chains: A Review of Applications, Challenges, and Opportunities for Climate Risk and Vulnerability Assessments, *Weather, Climate, and Society*, 2022.
- 650
- Meza, I., Siebert, S., Döll, P., Kusche, J., Herbert, C., Eyshi Rezaei, E., Nouri, H., Gerdener, H., Popat, E., Frischen, J., Naumann, G., Vogt, J. V., Walz, Y., Sebesvari, Z., and Hagenlocher, M.: Global-scale drought risk assessment for agricultural systems, *Natural Hazards and Earth System Sciences*, 20, 695–712, <https://doi.org/10.5194/nhess-20-695-2020>, 2020.
- OECD and JRC: Handbook on constructing composite indicators: methodology and user guide: Methodology and user guide, https://knowledge4policy.ec.europa.eu/sites/default/files/jrc47008_handbook_final.pdf, 2008.
- 655
- Schneiderbauer, S., Baunach, D., Pedoth, L., Renner, K., Fritzsche, K., Bollin, C., Pregolato, M., Zebisch, M., Liersch, S., Del Rivas López, M. R., and Ruzima, S.: Spatial-Explicit Climate Change Vulnerability Assessments Based on Impact Chains. Findings from a Case Study in Burundi, *Sustainability*, 12, 6354, <https://doi.org/10.3390/su12166354>, 2020.
- SFSO: Swiss Federal Statistical Office: Statistical data on Switzerland, <https://www.bfs.admin.ch>, last access: 15 March 2022.
- Stephan, R., Erfurt, M., Terzi, S., Žun, M., Kristan, B., Haslinger, K., and Stahl, K.: An inventory of Alpine drought impact reports to explore past droughts in a mountain region, *Natural Hazards and Earth System Sciences Discussions*, 21, 2485–2501, <https://doi.org/10.5194/nhess-21-2485-2021>, 2021.
- 660
- SURS: Statistical Office of the Republic of Slovenia: Statistical data provided by SURS, <https://www.stat.si/StatWeb/>, last access: 15 March 2022.
- Terzi, S., Sušnik, J., Schneiderbauer, S., Torresan, S., and Critto, A.: Stochastic system dynamics modelling for climate change water scarcity assessment of a reservoir in the Italian Alps, *Natural Hazards and Earth System Sciences Discussions*, 21, 3519–3537, 2021.
- 665
- van Loon, A. F., Stahl, K., Di Baldassarre, G., Clark, J., Rangelcroft, S., Wanders, N., Gleeson, T., van Dijk, A. I. J. M., Tallaksen, L. M., Hannaford, J., Uijlenhoet, R., Teuling, A. J., Hannah, D. M., Sheffield, J., Svoboda, M., Verbeiren, B., Wagener, T., and van Lanen, H. A. J.: Drought in a human-modified world: reframing drought definitions, understanding, and analysis approaches, *Hydrology and Earth System Sciences*, 20, 3631–3650, <https://doi.org/10.5194/hess-20-3631-2016>, 2016.
- 670
- Wezel, A., Casagrande, M., Celette, F., Vian, J.-F., Ferrer, A., and Peigné, J.: Agroecological practices for sustainable agriculture. A review, *Agronomy for Sustainable Development*, 34, 1–20, 2014.
- Zebisch, M., Schneiderbauer, S., Fritzsche, K., Bubeck, P., Kienberger, S., Kahlenborn, W., Schwan, S., and Below, T.: The vulnerability sourcebook and climate impact chains – a standardised framework for a climate vulnerability and risk assessment, *International Journal of Climate Change Strategies and Management*, ahead-of-print, 2021.
- 675
- Zorn, M. and Hrvatin, M.: Damage caused by Natural Disasters in Slovenia between 1991 and 2008, *Acta Geobalcanica*, 1, 33–43, <http://dx.doi.org/10.18509/AGB.2015.04>, 2015.