



Dynamical Perturbation of the Stratosphere by a Pyrocumulonimbus Injection of Carbonaceous Aerosols

Giorgio Doglioni^{1,2}, Valentina Aquila³, Sampa Das^{4,5}, Peter R. Colarco⁵, and Dino Zardi^{1,2}

¹University of Trento, Center Agriculture, Food, Environment.

²University of Trento, Department of Civil, Environmental and Mechanical Engineering.

³American University, Washington DC, USA.

⁴Earth System Science Interdisciplinary Center (ESSIC), University of Maryland, MD, USA.

⁵NASA Goddard Space Flight Center, Greenbelt, Maryland, USA.

Correspondence: Giorgio Doglioni (giorgio.doglioni@unitn.it)

Abstract. The Pacific Northwest pyrocumulonimbus Event (PNE) took place in British Columbia during the nighttime hours between the 12th and 13th of August 2017. Several pyroconvective clouds erupted in this occasion, and released in the upper troposphere/lower stratosphere unprecedented amounts of carbonaceous aerosols (300 ktn). Only a few years later, an even larger pyroCb injection took place over Australia. This event, named the Australian New Year (ANY) event, injected up to 5 1100 ktn of aerosol between December 29th 2019 and January 4th 2020. Such large injections of carbonaceous aerosol modify the stratospheric radiative budgets, locally perturbing stratospheric temperatures and winds. In this study, we use the Goddard Earth Observing System Chemistry Climate Model (GEOS CCM) to study the perturbations on the stratospheric meteorology induced by an aerosol injection of the magnitude of the PNE. Our simulations include the radiative interactions of aerosols, so that their impact on temperatures and winds are explicitly simulated. We show how the presence of the carbonaceous aerosols 10 from the pyroCb causes the formation and maintenance of a synoptic scale stratospheric anticyclone. We follow this disturbance considering the potential vorticity anomaly and the brown carbon aerosol loading and we describe its dynamical and thermodynamical structure and its evolution in time. The analysis presented here shows that the simulated anticyclone undergoes daily expansion-compression cycles governed by the radiative heating, which are directly related to the vertical motion of the plume, and that the aerosol radiative heating is essential in maintaining the anticyclone itself.

15



1 Introduction

Wildfires release large quantities of water vapour and aerosols in the atmosphere, as well as carbon oxides and other gases (Wiedinmyer et al., 2011). Under favourable atmospheric conditions, the rising of plumes from the wildfire can reach the stratosphere, as shown by Peterson et al. (2018) and Fromm et al. (2000, 2010). This lofting to the stratosphere is driven by different mechanisms. First, as soon as the plume reaches its lifting condensation level, the release of latent heat from the condensation of water vapor results in its further buoyant rise. This process is referred to as *pyroconvection* and produces pyroCumulus (pyroCu) clouds. Then, if the state of the troposphere is conducive to the formation of dry thunderstorms (Peterson et al., 2017), pyroCu clouds can further develop vertically, reaching the tropopause and evolving into pyroCumulonimbus (pyroCb) clouds, i.e. wildfire-driven thunderstorms loaded with smoke from the wildfire (Fromm et al., 2010).

A distinctive feature of pyroCb clouds is to transport of wildfire by-products to the Upper Troposphere Lower Stratosphere (UTLS): these thunderstorms act as chimneys, promoting the accumulation of aerosols and other gases in the UTLS. Once in the stratosphere, the main depletion mechanisms for aerosols (i.e. wet and dry deposition) are greatly reduced compared to the troposphere, and as a consequence aerosols have a significantly longer lifetime than in the troposphere (Yu et al., 2019). In the UTLS, the optically thick carbonaceous aerosols from wildfires absorb radiation, causing the temperature of the smoke plume to rise. This warming results in a gradual diabatic lofting of the plume itself (Torres et al., 2020; Das et al., 2021). This process is more pronounced in the days following the aerosol injection into the UTLS, when the concentration of aerosol is higher (de Laat et al., 2012; Ditas et al., 2018).

The second largest pyroCb aerosol injection ever recorded is the 2017 Pacific Northwest Event (PNE). The PNE was characterized by an outbreak of five pyrocumulonimbus clouds in British Columbia (Canada) and the state of Washington (USA) during the evening of the 12th of August 2017 and the first hours of the following day (Peterson et al., 2018). Overall, the five pyroCbs injected into the stratosphere between 0.1 Tg and 0.35 Tg of carbonaceous aerosols (Peterson et al., 2018; Torres et al., 2020). This record was surpassed by the injection from the December 2019/January 2020 Australian New Year event (ANY), which resulted in the stratospheric injection of up to 1.1 Tg of aerosol (Peterson et al., 2021).

Several studies (Das et al., 2021; Bourassa et al., 2019; Kloss et al., 2019; Torres et al., 2020; Yu et al., 2019; Christian et al., 2019; Baars et al., 2019) have characterized the aerosol plume from the PNE event and its interaction with large scale stratospheric features such as the Asiatic Summer Monsoon Anticyclone. In the days following the PNE, the smoke gradually rose into the stratosphere due to diabatic heating, with peak ascent rates of 2-3 km/day (Das et al., 2021; Torres et al., 2020). The plume eventually reached 22 km in height around 20 days after the injection, as detected in OMPS LP (Ozone Mapping and Profiling Suite, Limb Profiler) and SAGE III (Stratospheric Aerosol and Gas Experiment) observations (Torres et al., 2020; Bourassa et al., 2019). The plume was also detected by ground based lidars, as presented in Khaykin et al. (2018). The smoke remained in the atmosphere for several months, with an estimated half-life of 5 months (Christian et al., 2019; Yu et al., 2019); during this period it perturbed the radiative balance of the atmosphere, causing global mean radiative forcing anomalies at the surface of the order of $-0.12 \pm 0.03 \text{ W m}^{-2}$ over September 2017 (Das et al., 2021). Lestrelin et al. (2021) analyzed the dynamical features of the PNE plume by inspecting ERA5 reanalysis fields and CALIOP (Cloud-Aerosol Lidar with Orthogonal



50 Polarization) measurements, finding that it evolved into stratospheric anticyclones that persisted for almost two months. These smoke-induced stratospheric anticyclones were first reported following the ANY event (Allen et al., 2020; Kablick III et al., 2020; Khaykin et al., 2020; Lestrelin et al., 2021) and have been named SWIRLs (Smoke With Induced Rotation and Lofting) by Allen et al. (2020). These studies underline the stability of the SWIRLs and their resilience against the large scale shear characterizing the background wind field. The main signatures of SWIRLs are a deep potential vorticity anomaly with respect to the zonal mean (negative in the northern hemisphere, positive in the southern) and an associated anticyclonic motion, enhanced optical thicknesses and a vertical temperature anomaly dipole. Also, the SWIRL encases air from the UTLS with low ozone content and this is reflected in a characteristic negative ozone concentration anomaly as the SWIRL moves upwards. As pointed out by Khaykin et al. (2020), the SWIRL effectively traps the carbonaceous aerosol, with the result of efficiently transporting it to higher altitudes. The cited studies underline how the presence of carbonaceous aerosol is the necessary condition in the formation and maintenance of SWIRLs and how the coupling between aerosol, radiation, and dynamics is necessary to reproduce a SWIRL in model simulations (Khaykin et al., 2020; Allen et al., 2020).

In this work we use a chemistry climate model to simulate the impact on the stratosphere of a pyroCb plume from an event of the magnitude of the 2017 Pacific Northwest Event (PNE), as well as the effect of the aerosol radiative interaction in the development of the plume itself. As in Das et al. (2021), we use the Goddard Earth Observing System Chemistry-Climate Model (GEOS CCM), focusing in this study on the GEOS Atmospheric General Circulation Model configuration with prognostic aerosols from the GOCART (Goddard Chemistry, Aerosol, Radiation, and Transport) module to simulate the long term transport of the plume and its impact on the radiative budget of the atmosphere. While Das et al. (2021) used GEOS CCM in a relaxed replay mode, i.e. driven by observations in the troposphere, we present here free running simulations (i.e. weather forecasts). In previous work by Khaykin et al. (2020), Allen et al. (2020) and Lestrelin et al. (2021), the structure of the SWIRL was reproduced and maintained thanks to the assimilation of the observations and not by the local radiative heating caused by the carbonaceous aerosols. Here, we show how the GEOS CCM is capable of forming and maintaining a SWIRL following a stratospheric aerosol injection such as the one from the PNE event, thanks to its representation of the radiative impact of the aerosol on the dynamics.

The article is structured as follows. In section 2 we briefly describe the GEOS model as well as the setup used for the simulation. In section 3 we present the results from the simulations, analyzing the dynamical signatures and thermodynamical characteristics of the reproduced SWIRL and comparing them to previous works; more specifically, we will analyze the anomalies of potential vorticity, wind, temperature, and geopotential that describe the SWIRL as well as their evolution during the SWIRL life; moreover, we will show how the diabatic heating provided by the presence of the aerosol is correlated to the dynamics of the SWIRL itself, and how the SWIRL undergoes a daily expansion/compression cycle following this diabatic heating. Also, the geometrical characteristics of the SWIRL will be pointed out, and we will compare the resulting dimensions with the SWIRLs observed in previous works. In section 4 we will present some considerations about the reproduced SWIRLs and draw some conclusions about the novelty and limitation of this work in comparison to other studies.



2 Methods

2.1 Model Description and simulation setup

85 The Goddard Earth Observing System (GEOS) can simulate atmospheric circulation and chemistry, oceanic circulation and biogeochemistry, land surface processes and data assimilation at horizontal resolutions as small as 12 km (Molod et al., 2015; Rienecker et al., 2008). In this study we use GEOS as an atmospheric general circulation model, resolving the atmospheric circulation and composition but using prescribed sea surface temperatures and sea ice fractions. GEOS can be used in *replay* and *free running* configurations. In the replay mode, simulations are constrained by the MERRA-2 reanalysis fields, while
90 in the free running mode the evolution of the atmosphere is not constrained to the observations. The results shown in this study were produced in free running simulations with the Icarus 3.3 version of GEOS. The meteorological fields for the model initialization were obtained from version 2 of the Modern-Era Retrospective analysis for Research and Applications (Gelaro et al., 2017).

Simulations were run on a cubed-sphere horizontal grid with horizontal resolution of ~ 50 km with 72 hybrid vertical sigma
95 levels extending from the surface to 0.01 hPa (Rienecker et al., 2008). Aerosol concentrations and aerosol processes are simulated with the Goddard Chemistry, Aerosol, Radiation, and Transport module (GOCART), a bulk aerosol module simulating the evolution of black carbon (BC), brown carbon (BrC), organic carbon (OC), nitrates (NO_3), sulfates (SO_4), dust, and sea salt (Chin et al., 2002, 2009; Colarco et al., 2010, 2017). In our simulations, aerosols are radiatively coupled to the dynamics and have a direct impact on the meteorological forecast. The optical properties of all the species, except dust and BrC are deter-
100 mined by the OPAC data set (Hess et al., 1998), while dust optics are treated as described in Colarco et al. (2014). As in Das et al. (2021), wildfire emissions are assigned to BC and BrC, which is more absorbing than OC in the near UV. The simulations also include the stratospheric chemistry module StratChem (Consideine et al., 2000; Douglass and Kawa, 1999), which allows for the inclusion of background stratospheric sulfate from the oxidation of carbonyl sulfide. Stratospheric chemical reaction rates are impacted by the pyroCb smoke through temperature changes, but not via changes in heterogeneous chemistry.

105 The PNE plume is represented as a 0.3 Tg injection of carbonaceous aerosol, which is consistent with the observational findings from Peterson et al. (2018) and Torres et al. (2020). Of this, 97.5 % is BrC and the remaining 2.5% is BC. The BC/BrC ratio was set by Das et al. (2021) to reproduce the observed diabatic lofting of the plume, which is sensible to the overall loading of the strongly absorbing BC. The 0.3 Tg are injected in three different locations in British Columbia, in correspondence to the most pronounced three pyroCb clouds of the PNE. The aerosol have been initialized as smeared in regions centered on these
110 locations, spanning horizontally 2° longitude by 2.5° latitude and vertically from 10 to 12 km. The aerosol injections occur during the first hours of the 13th of September, with the first two puffs occurring from 0 to 3 UTC (0.19 Tg, locations 1A and 1B, Fig. 1) and the third from 4 UTC and 6 UTC (0.11 Tg, location 2, Fig. 1). These time intervals have been obtained using Cloud Top brightness temperature measurements by Das et al. (2021). The modal radius of the BrC aerosol in the model was set to $0.035 \mu\text{m}$ to achieve an Angstrom Exponent of 1.3, following SAGE III measurements of the event (Das et al. (2021)).
115 In this work we present a simulation with the PNE injection, starting on 13th August 2017 and ending on 30th September 2017, that shows the formation of a SWIRL.




Figure 1. Locations of the carbonaceous injections included in the GEOS simulations on the 13th of August.

2.2 Definition of SWIRL boundaries and data analysis

We define the boundaries of the SWIRL based on the local anomaly of Ertel's potential vorticity (PV) (Ertel, 1942) with respect to the zonal mean and the brown carbon concentration. We identify the SWIRL as the region where the Ertel's PV anomaly is smaller than -10% of the zonal mean value and the mass mixing ratio of brown carbon is higher than $1.25 \mu\text{g kg}^{-1}$. These criteria do not select PV anomalies that are not connected to the presence of high carbonaceous aerosol concentrations. In order to obtain smooth SWIRL contours, both the carbonaceous aerosol concentration fields and the PV anomaly fields have been filtered horizontally using a gaussian filter with a standard deviation of 1 degree in the latitudinal direction and 1.5 in the longitudinal direction. We concentrate on model levels higher or equal to 100 hPa, so as to ensure that the detected SWIRL is above the tropopause. The analysis proposed here to track the SWIRL follows the approach of Lestrelin et al. (2021), which uses the Lait Potential Vorticity anomaly (Lait, 1994), along with the ozone concentration anomaly, to locate the SWIRL from ERA5 reanalysis fields. A similar approach was also used by Khaykin et al. (2020) to track the SWIRL from the ANY event. To investigate the characteristics of the SWIRL, we analyzed the following diagnostic quantities: Ertel's potential vorticity (PV), temperature (T), potential temperature (θ), density (ρ), zonal wind speed (U), meridional wind speed (V), vertical wind



130 speed (W) and carbonaceous aerosol (BC and BrC) concentrations. All these quantities are saved on model levels every 6
hours, and will be presented as absolute fields or as anomalies with respect to their zonal means at the same model level and
latitude unless specified otherwise. The geopotential of the model levels that is present in our discussion is approximated by the
mid-layer geometrical height multiplied by the gravitational acceleration g . In our analysis we have also used the temperature
tendencies due to the diabatic processes resolved by the model (radiation, moist physics, friction, turbulence, and gravity wave
135 drag) and to the dynamics. Additionally, the model calculates the contribution of the aerosol on the temperature tendency due
to radiation through two calls to the radiation model, with and without the inclusion of aerosols. In order to better capture the
relationship between the SWIRL and its closest surroundings, we restricted our analysis to a region centered on the SWIRL
spanning 43.75° longitude by 35° latitude, and calculate the diagnostic variables as horizontal averages over the portions of
this region inside and outside the SWIRL. The SWIRL geometry was described considering its volume, which is calculated by
140 summing the volumes of the grid cells inside the SWIRL contours over the height of the SWIRL, and its average area, which
is calculated by vertically averaging the areas of the SWIRL computed for every layer by summing the areas of the cells lower
boundaries inside the SWIRL selection. The thickness was obtained by considering the height difference between the highest
and lowest levels at which the SWIRL was detected.

To visualize and describe the evolution of the bulk SWIRL properties, we considered averages and extremes over the SWIRL
145 volume of our meteorological fields and temperature tendencies.

3 Results

3.1 SWIRL Tracking

In the days following the PNE aerosol injection in UTLS, the smoke plume drifted eastward and the diabatic self-lofting
150 promoted its vertical movement. A brief description of the smoke transport is available in appendix A, where we also include
a comparison of the plume transport in the free running and replay configurations of the GEOS model. In our free running
simulations the plume reached 100 hPa on the 16th of August, consistent with OMPS LP observations (Das et al., 2021),
marking the start of the SWIRL detection using the algorithm described in Sec. 2.2. By the end of the 18th of August the
detection algorithm revealed that the bulk of the SWIRL was above the 100 hPa limit, spanning between the levels 100 hPa and
155 61 hPa. For our analysis, we mark the 18th of August as the first day of the stratospheric SWIRL. Before this date some signs
of SWIRL formation were visible at levels lower than 100 hPa and above the simulated tropopause height (about 150-200 hPa
in the days following the injection in the regions corresponding to the plume). We have chosen not to include this initial part
in the analysis, since we are interested in the life of the SWIRL when it is well into the stratosphere.

The SWIRL is first detected by the algorithm over the Atlantic on the 18th of August (figure 2) and traveled southeastward
160 until it reached the northern Atlantic. Here it stalled for 7 days (days 6-12 after the PNE, 19th-25th August) before starting its
westward movement, which eventually brought it to the Arabic peninsula (day 28 after the PNE, 10th of September). The next
day (11th of September, day 29), the SWIRL started losing compactness and that marked the end of its stratospheric lifetime.




Figure 2. Time evolution of simulated SWIRL starting on the 18th August. The contours show the locations where the SWIRL is detected in at least two vertical layers. Due to the overlapping of the contours, we show here the days: 18th-19th-23rd-27th of August and then every 2 days until the 16th of September

After this date, the detection algorithm was still able to detect its fragmented remains until at least the 16th of September, when they reached the Caribbeans. In our simulations, the SWIRL lasted overall 25 days, which is shorter than in observations
 165 Lestrelin et al. (2021). Lestrelin et al. (2021) reports the splitting of the initial SWIRL and the generation of offspring anticyclones between the 22nd of August and the 1st of September. In our simulations a similar event takes place over the Atlantic ocean during the stalling of the SWIRL, but apart from the main SWIRL no offspring circulation is generated by the resulting aerosol plumes.

3.2 Analysis of the SWIRL on the 23rd of August

170 In this section we will describe the structure of the simulated SWIRL on August 23rd. On this day the vortex is well into the stratosphere (detected in the model levels between 72 and 43 hPa), with tangential wind speed anomalies of 10 ms^{-1} , and well organized, with a clear anticyclonic circulation closed around its center. Inside the borders of the SWIRL lies the deep negative PV anomaly, which indicates the presence of an anticyclone (Figs. 3a-d and 4a-d). Indeed, the horizontal wind anomaly structure of the SWIRL shows an anticyclonic circulation centered over the PV anomaly minimum, with tangential wind mag-
 175 nitudes of about 10 m/s (Figs. 3q-t and 4q-t). The magnitude of the tangential wind speed in the SWIRL here simulated is comparable to the ones presented in literature for the SWIRL developed after the Australian New Year event (Kablick III et al., 2020; Allen et al., 2020). The analysis of the geopotential shows a positive anomaly in correspondence of the SWIRL (Figs. 3q-t and 4q-t): this indicates that the effect of the aerosol is to trigger an outward pressure gradient force that, together with Coriolis acceleration produces the anticyclonic motion of the SWIRL.

180 The pyroCb aerosol causes a strong internal heating of the SWIRL during daytime (Fig. 3). During nighttime the aerosol does




Figure 3. Horizontal sections at different heights of the SWIRL on the 23rd of August, 18z, when the maximum heating occurs. The first to the fourth columns show 43, 52, 61, and 72 hPa altitudes, respectively. The first row (a-d) shows the potential vorticity anomaly [%], the second (e-h) the vertical velocity, the third (i-l) the potential temperature anomaly, the fourth (m-p) the temperature tendency due to radiative heating/cooling of the aerosol, and the fifth row (q-t) the geopotential anomaly (color shades) and the wind field anomaly (white arrows). All anomalies are calculated with respect to the zonal mean. The geopotential anomaly was made positive in the considered region by summing the absolute value of the minimum of the geopotential anomaly to the entire field. This was done for representation purposes. The black contours represent the boundary of the detected SWIRL.




Figure 4. Same as 3, but for the 23rd August 00 z, when the aerosol is cooling due to the lack of incoming solar radiation.

not absorb SW radiation but emits LW radiation, inducing a net cooling of the plume (Fig 4m-p). Vertical velocities within the SWIRL are comparable to those outside the SWIRL (Fig 4e-h). From this analysis there is no evidence of an organized structure in the vertical wind field in correspondence of the SWIRL and there is no sign of its upward movement. This however is expected considering that the vertical displacement of the SWIRL, which can be quantified in hundreds of meters per day, should be driven by a vertical velocity of the order of fractions of cm s^{-1} , which cannot be visible in the noisy vertical velocity

185



background. The lofting of the SWIRL will be made evident later on, considering the average of the vertical velocity in the SWIRL (Fig. 7).

190 While the higher levels of the SWIRL are approximately elliptic, the lowermost levels present a peculiar shape, consisting of an ellipsoid to which a protrusion is attached in the downwind direction. This protrusion can be seen as the tail of the SWIRL, which gradually loses material in time, leaving a trail of air with negative PV anomaly and high aerosol concentration that is classified as SWIRL by our detection method. Such tails have been also observed by Lestrelin et al. (2021).

195 Another characteristic feature of SWIRLs is the vertical potential temperature anomaly dipole. This is visible considering the maps of potential temperature anomaly (Figs. 3i-l and 4i-l), which indicate a negative anomaly above the center of the SWIRL and a positive anomaly below. As stated by Allen et al. (2020), this signature in the potential temperature is consistent with the vertical expansion of the plume due to the diabatic heating of the aerosol.

200 In the vertical sections of the SWIRL (Figs. 5-6e) a strong anticyclonic motion is visible as well as the vertical structure of the diabatic heating provided by the aerosol (Fig. 5-6c), which is particularly intense in the middle of the SWIRL. The analysis of the meridional wind anomaly curtain (Fig. 6e) reveals also the tilting of the SWIRL, whose axis lies at an angle with respect to the vertical. As described by Allen et al. (2020), the tilting of the vortex is given by the wind shear in the vertical direction. The characteristic vertical temperature and potential temperature anomaly dipoles that characterize SWIRLs are also visible in Figs. 5b-f and 6b-f, with negative anomalies in the upper part of the SWIRL and positive anomalies below. The magnitude of the temperature dipole (10 K) is comparable to that reported for the ANY case (Allen et al., 2020), which reached up to 15 K. The temperature and potential temperature anomalies extend well below and above the potential vorticity anomaly, as is also reported in Allen et al. (2020); Lestrelin et al. (2021) and Kablick III et al. (2020). As in the horizontal sections, the vertical velocity field does not exhibit any detectable difference between the interior of the SWIRL and its surroundings.




Figure 5. Vertical cross section at fixed longitude (-59° W) of the SWIRL on the 23rd of August, 18z. (a) shows the percent relative potential vorticity anomaly, (b) the temperature anomaly, (c) the temperature tendency due to the aerosol heating, (d) the vertical velocity, (e) the zonal wind anomaly, and (f) the potential temperature anomaly. All anomalies are calculated with respect to the zonal mean between -180° and $+180^\circ$ longitude.

The temperature anomaly dipole showed in Figs. 5b and 6b is reflected in a reduced lapse rate inside the SWIRL compared to the surroundings (Fig. 7a), with the internal temperature profile intersecting the external one roughly in the middle of the SWIRL. The same can be observed in the potential temperature anomaly (Fig. 7d). Fig. 7h shows the diurnal heating and nocturnal cooling of the SWIRL due to aerosols and Fig. 7b its resulting vertical motion. The difference in the average vertical velocity between the interior and exterior of the SWIRL is crucial, since it explains how the SWIRL actually moves upwards despite being immersed in a noisy vertical velocity field.

The profile of the horizontal wind speed anomaly (Fig. 7e) shows stronger wind speeds in the middle part of the SWIRL; this was visible also in Fig 3. The reason for this is that the concentrated heating that takes place inside the SWIRL produces a strong rotation which does not interact with the background flow. Indeed, the similarly strong heating taking place at the bottom




Figure 6. Same as fig.5 but for a fixed latitude (36° N). Panel (e) contains the meridional velocity anomaly.

of the SWIRL (Fig. 7h) does not result in a similarly strong rotation due to the SWIRL interaction with the background flow. This fact is also supported by the shape of the SWIRL, which presents tails in its lower levels caused by the dispersive action of the background flow.

220 The radiative heating/cooling due to the aerosol is the main factor among the physical diabatic processes (Fig. 7g-i). During the day, the SWIRL is subject to strong diabatic heating, that promotes the vertical motion from buoyancy. The vertical motion is accompanied by the expansion of the SWIRL, which partially compensates for this heating; this is evident in the temperature tendency due to the dynamics (Fig. 7b and i, yellow curves). During the night, the situation is the opposite. The diabatic cooling drives a downward vertical motion of the SWIRL and its subsequent heating due to expansion (Fig. 7b and i, blue curves). The
 225 nocturnal cooling is less intense in absolute value than the diurnal heating. This points toward a net ascent of the SWIRL, given the enhanced intensity of the diurnal lofting with respect to the nocturnal descent (Fig. 7h).




Figure 7. Vertical profiles at 4 different times (0-6-12-18 z) for the 23rd of August. Full lines represent vertical profiles of the horizontal averages inside the SWIRL selection, while dotted lines represent vertical profiles of the horizontal averages outside the SWIRL selection in a rectangular region 43.75° longitude by 35° latitude centered on the SWIRL. The panels show (a) absolute temperature, (b) vertical velocity, (c) Ertel’s potential vorticity, (d) potential temperature anomaly with respect to the zonal mean, (e) horizontal speed anomaly with respect to the zonal mean, (f) density anomaly, (g) temperature tendency due to diabatic processes, (h) temperature tendency due to the aerosol, and (i) temperature tendency due to the dynamics.

3.3 Evolution in time of the SWIRL properties




Figure 8. Evolution in time of the SWIRL (a) volume, (b) mean radius, (c) thickness, (d) average potential temperature, (e) carbonaceous aerosol loading and (f) absolute temperature. In (d) the red line shows the result of the linear fit between day 5 and 11 of the potential temperature, while the blue line between day 11 and day 30. The steepness of the red line is 10.6 K day^{-1} , while for the blue line it is 6.2 K day^{-1} .

The volume of the SWIRL increases until day 10-11 (22nd-23rd August) and then decreases steadily in time (Fig. 8a). The daily modulation of the volume, corresponding to the expansion/compression cycle caused by the diurnal radiative cycle, is visible especially in the first 11-12 days. Accordingly, the calculated radius (Fig. 8b) shows the same time evolution, peaking around day 11 at 700 km. The peak radius of the simulated SWIRL is comparable to the maximum radius of the SWIRL observed following ANY (peaking at 750 km, Allen et al. (2020)); instead, the simulated SWIRL is consistently larger in comparison to the main SWIRL observed following PNE, whose horizontal dimensions (diameter) were estimated at 700 km from composite analysis Lestrelin et al. (2021). After day 28 (11th September) the SWIRL volume increases significantly, following its dispersion and fragmentation. The detection algorithm still categorizes the remains of the plume as SWIRL even




Figure 9. Evolution in time of (a) the minimum percent relative Ertel’s PV anomaly with respect to the zonal mean inside the SWIRL; (b) the average vertical velocity in the SWIRL (black line) and outside the SWIRL (green line); (c) Average temperature tendency in the SWIRL due to diabatic processes (red line), average temperature tendency due to aerosol inside the SWIRL (blue line), average temperature tendency outside the SWIRL due to diabatic processes (green line); (d) Maximum (red line), minimum (blue line) and average (black line) speed anomalies with respect to the zonal mean inside the SWIRL (red line); (e) Maximum (red line) and minimum (blue line) temperature anomalies inside the SWIRL with respect to the zonal mean; The black line represents the magnitude of the temperature anomaly dipole computed as the distance between the red and blue lines; (f) Averages inside the SWIRL of dynamical (black line) and total diabatic (red line) temperature tendencies. The green lines have been computed as follows: a region centered in the SWIRL of 43.75° longitude by 35° latitude was selected and the averages were carried out over the portion of this region at the same height of the SWIRL but outside the SWIRL.

thought they lack the characteristic compactness of SWIRLs. This process is also visible in the calculated average radius, with a steep increase after day 28. Moreover, the SWIRL thickness (Fig. 8c) indicates that the SWIRL reaches a maximum thickness of 5 km between day 11 (24th August) and 16 (29th August); before and after this period, the thickness value fluctuates between 3 and 4 km; the magnitude of the SWIRL thickness is comparable to the one observed in the ANY SWIRL, that peaked at 6 km in the moment of its maximum expansion (Allen et al., 2020), and it is also close to the thickness of 3 km of the PNE SWIRL



obtained from the composite analysis of Lestrelin et al. (2021). The step-like fluctuations of the SWIRL thickness is due to the vertical resolution of the model at the considered levels, which is around 1 km. The ascent of the SWIRL can be divided in two phases: the steep initial ascent, during which the volume of the SWIRL increases, and a second slower ascent phase, characterized by a decrease in volume of the SWIRL. The potential temperature θ increases approximately linearly (Fig. 8d) during both ascent phases, with a slope of about 10 K day^{-1} during the first phase and of 6 K day^{-1} during the second. The ascent speeds are consistent with the observed ones during the ANY SWIRL: in Allen et al. (2020) the ascent speed during the first 27 days of the SWIRL life is estimated at 7.8 K day^{-1} and 5.5 K day^{-1} afterwards. Lestrelin et al. (2021) show that the SWIRL generated by the PNE plume that reached subtropical latitudes penetrated the stratosphere until reaching 570 K in potential temperature 60 days after the PNE. The SWIRL presented in this work instead reaches a potential temperature of 590 K before its disruption 29 days after the PNE. While the two figures are of similar magnitude, the simulated SWIRL reaches it in half the time with respect to the observed one; however, it is important to remark that the observed SWIRL following the PNE had already reached potential temperatures of the order of 520-540 K a month after the injection (Lestrelin et al., 2021), so the ascent velocities of the SWIRLS are actually comparable. Lastly, the average absolute temperature T within the SWIRL (Fig. 8f) shows little variation in time, with values between 206 and 209 K.

The magnitude of the temperature anomaly dipole in the SWIRL is approximately stable until day 10-11 (22nd-23rd August), and then gradually decreases as the SWIRL ages (Fig. 9e). The simulated temperature dipole (peaking at 7 K) is significantly more pronounced than the one calculated for the main PNE SWIRL via composite analysis by Lestrelin et al. (2021) (peaking at 2 K). The maximum horizontal wind speed anomaly in the SWIRL (Fig. 9d) first decreases for three days, in the same period in which the radius of the SWIRL increases significantly; then it stabilizes around $12\text{-}13 \text{ m s}^{-1}$ before decreasing again after day 15 (28th August) and reaching figures between $6\text{-}8 \text{ m s}^{-1}$. The same behaviour is observed in the average horizontal velocity anomaly that decreases in the first days from 10 m s^{-1} to 5 m s^{-1} , then remains constant at 5 m s^{-1} until day 15 and subsequently decreases gradually reaching 3 m s^{-1} at the end of the SWIRL lifetime.

The minimum of the PV anomaly (Fig. 9a) abruptly increases at day 6-7 from about -140% to about -90% and remains roughly stable until day 11, when it starts increasing at a slower rate. The average vertical velocity of the SWIRL (Fig. 9b) is positive (indicating upwelling) at the end of the daily heating period and negative during the night, with maximum values peaking at 0.7 cm s^{-1} . The average vertical velocity of the SWIRL reaches more pronounced peaks than those of the surroundings, indicated by a green line in Fig. 9b.

The diabatic temperature tendency (Fig. 9c) is initially dominated by the radiative heating of the aerosol. The mean aerosol heating rate within the SWIRL is significantly larger than the average background diabatic temperature tendency until day 22-23 (4th-5th September). After this day, the diabatic temperature tendency inside the SWIRL becomes progressively closer to the diabatic temperature tendency outside. The quick loss of coherence of the SWIRL that takes place after day 28 is driven by the loss of this differential heating between its inner region and the surroundings. The dynamical temperature tendency is anticorrelated with the diabatic temperature tendency (9f), which is dominated by the aerosol heating, and is directly linked to the vertical velocity of the SWIRL: when the SWIRL is heated by radiation, it rises and expands, meaning that its average vertical velocity is positive and the dynamical temperature tendency is negative. During the night this tendency is reversed



along with an average negative vertical velocity. This means that while the SWIRL is cooling, it moves downward while being subject to compression.

4 Conclusions

280 In this work, we present how the GEOS CCM is able to reproduce a SWIRL, i.e. a stratospheric aerosol-induced anticyclone. Previous studies used reanalyses and observations to characterize the SWIRLs generated by the PNE (Lestrelin et al., 2021) and ANY (Allen et al., 2020; Kablick III et al., 2020; Khaykin et al., 2020) events. The simulations presented here reproduce several of their findings, and in particular the structures of the anomalies characterizing the SWIRLs. The use of a free-running model with aerosol-radiation coupling allows us for the first time to resolve the role that the aerosol radiative interaction plays
285 in the development of the SWIRL, as well as to characterize the diurnal cycle of the SWIRL and associated transport effects. The presence of the heating of the aerosol is crucial, and Khaykin et al. (2020) showed that, without this contribution, SWIRLs dissipate in around 6-7 days.

In this work, the simulated SWIRL has been first recognized by using a simple detection algorithm, based on the Ertel's PV anomaly and on the brown carbon concentration. The analysis of the SWIRL has been carried out by considering its structure
290 on the 23rd of August and then by evaluating how its bulk properties evolved during its lifetime. This analysis shows that the simulated anticyclonic disturbance reproduces the magnitudes of the anomalies of the observed ones; for instance, the magnitude of the flow of the anticyclone is about 10 m s^{-1} , which is close to the magnitude the flow in the SWIRL observed following ANY (Allen et al., 2020). The same is true for the magnitude of the temperature anomaly dipole, which is observed together with a steeper lapse rate inside the SWIRL than in the surroundings. We also showed that the anticyclonic circulation is
295 triggered by a pressure gradient force, triggered by a positive anomaly of the geopotential of the isobaric levels. This enhanced pressure in the center of the SWIRL is given by the diabatic heating of the aerosol, that also leads the expansion/compression cycle that the SWIRL undergoes and its consequent net diabatic lofting.

The diurnal cycle of the SWIRL is modulated by the daily radiation cycle. During daytime, the aerosol heats up the atmosphere through the absorption of solar radiation. This triggers a radial expansion of the plume and a consequent vertical motion due to
300 buoyancy. This vertical motion, despite small, triggers an adiabatic temperature tendency that counteracts the radiative heating. During the night, when the plume cools down radiatively, the process is inverted, and the SWIRL moves downward. The upward daytime motion is more pronounced than the nighttime one, leading to a net ascension of the plume. The analysis of the evolution of the radius/volume of the SWIRL during its lifetime reveals this daily expansion-compression cycle of the SWIRL. While the simulated SWIRL intensity and characteristics are in agreement with the ones detected in the past (Lestrelin et al.,
305 2021; Allen et al., 2020), GEOS maintains the SWIRL for a shorter time. We have not investigated the reason for this discrepancy in details, but a key point in maintaining the SWIRL appears to be the difference between the diabatic heating within the SWIRL and the surroundings. Therefore, a role in determining the development of the SWIRL must be played by the specific meteorological conditions after the event, which are not expected to be reproduced in a free-running simulation.

Despite being able to reproduce many characteristics observed in SWIRLs, the GEOS model has limitations that might impact



310 the results. First, the vertical resolutions at the stratospheric altitudes here analyzed is about 1 km. A finer resolution would
provide a better description of the vertical ascent, as clear in the step-wise behaviour of the SWIRL thickness visible in Fig. 8.
Additionally, GOCART, the aerosol module in the GEOS configuration used here, is a bulk model, i.e. it does not represent the
microphysical evolution of aerosols. GOCART does include a parameterization of the transformation of carbonaceous particles
from hydrophobic to hydrophilic and the subsequent hygroscopic growth of hydrophilic particles, but does not include changes
315 in optical properties due to coagulation of aerosols, condensation of gaseous material, or the transfer from external to internal
mixture. Lastly, GOCART assumes that black and brown carbon aerosols are spherical. Changes in depolarization ratio (both
in space and time) as shown in Christian et al. (2019), or fractal structures such as in Yu et al. (2019) are not simulated. Since
(Yu et al., 2019) showed that fractal structures are more absorbing at visible wavelengths, the radiative impact that we simulate
might be underestimated. This might explain the shorter SWIRL lifetime in our simulations than in observations. This work
320 only described the SWIRL during the period in which it was relatively stable and well-formed in the stratosphere. We did not
investigate the SWIRL characteristics during its formation in the UTLS, or during its dissipation in the stratosphere. A whole
ensemble of free-running simulations should be used to analyze the importance of specific background meteorological condi-
tions on the formation and dissipation of the SWIRL, as well as the interaction of aerosol heating and background conditions
in the initial and final stage of the SWIRL lifetime.

325 *Code availability.* The GEOS model is available from an externally accessible subversion software repository, whose details are provided at
https://gmao.gsfc.nasa.gov/GEOS_systems/geos5_access.php (NASA, 2020).

Data availability. The dataset used to obtain the results presented in this article are available for download in the online repository "GEOS
CCM free-running simulation data of the Pacific-Northwest pyrocumulonimbus Event-like aerosol injection, SWIRL selection", url: <https://doi.org/10.5281/zenodo.6366106>, (Doglioni, 2022).



330 Appendix: Aerosol Transport




Figure A1. Comparison of the zonal mean stratospheric aerosol optical thickness (SAOD) at 550 nm from the free running and replay simulations for four different days in the month following the PNE.

Following the injection from the PNE, the aerosol released in UTLS started its journey in the stratosphere. The diabatic self lofting promoted the gradual ascension of the plume while it was advected and dispersed by the large scale circulation. The horizontal transport of aerosol is visualized considering maps of SAOD (Stratospheric Aerosol Optical Depth) at 550 nm (Fig. A1). The SAOD is calculated summing the simulated extinction coefficients fields due to the presence of Black and Brown carbon, and vertically integrating them from the tropopause up to the model's top of the atmosphere. The computation of the SAOD has been carried out both for the free running and replay simulations. As shown in Das et al. (2021), the latter well capture the aerosol transport as observed by OMPS LP; thus, the replay simulation is used here as a reference of the actual aerosol transport.

In the first 2-3 days following the injection, the aerosol is advected and gradually dispersed by the wind field in the UTLS, that moves the plume towards the Atlantic both in the free running and replay simulations (Fig. A1 (a)); at this stage the difference



is limited between the different simulations.

As time passes, the free running simulation begins to significantly differ from the one in replay configuration (fig. A1, (d)). Of particular interest during this phase is the splitting of a portion of the plume that can be observed in the free running simulation around day 11 (24th August 2017) above the northern Atlantic ocean (fig. A1, compare (b) with (c)). After this event, a section of the plume reaches lower latitudes and starts moving towards the Gulf of Mexico.

This southern part of the aerosol plume exhibits interesting properties such as the confinement of the plume itself and its resilience against the disruption operated by the background flow. This portion of the plume generated the SWIRL, which is presented in this article. The splitting of the plume does not occur in the replay configuration, in which the bulk of the aerosol reaches Europe.




Figure A2. Time evolution of the horizontal average over the northern hemisphere of the 500 nm extinction coefficient of carbonaceous aerosols.

In order to compare the vertical displacement in time of the plume, we consider the time evolution of the horizontal average over the northern hemisphere of the extinction coefficient due to carbonaceous aerosols at 550 nm (Fig. A2), which reveals a fast ascent of the aerosol plume in the first days following the injection. Comparing the free running with the replay configurations it is evident that in the former the vertical transport is more pronounced: in the free running simulation the aerosol that reaches 100 hPa in 5 days is consistently more than in the replay simulation. Only from days 9-10 we find comparable extinction coefficients



355 above 100 hPa in the replay simulation. This difference in the ascent rate in the replay and free running configurations could
be given by the different background meteorology. Also, in the free running configuration the model's physics is free from the
observational constraints, so that the aerosol diabatic heating can have a direct impact on it; instead in the replay configuration
the action of the aerosol in the dynamics might be dampened by the replay procedure itself. However, caution is needed when
exploring this argument, since we have a single free running simulation; the study of an ensemble of free running simulations
360 might be the correct instrument to better quantify and explain the differences in the diabatic self-lofting between the replay and
free running simulations.

Author contributions. PRC and SD designed the modeling approach and performed the simulations. GD performed the data analysis and
analyzed the results under the supervision of VA and DZ. GD wrote the manuscript with VA and DZ. All the authors reviewed the manuscript
before the submission.

365 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. We would like to acknowledge the NASA Earth Science Division and GEOS model developmental efforts at GMAO for
their support and the NASA MAP funding under the GEOS-CCM project (CCM Workpackage; program manager David Considine). Sampa
Das's research at NASA GSFC for this effort was supported by an appointment to the NASA Postdoctoral Program (NPP), administered by
the Universities Space Research Association under contract with NASA. The computing resources supporting the simulations shown in this
370 work were provided by the NASA High-End Computing (HEC) Program through the NASA Center for Climate Simulation (NCCS) at the
Goddard Space Flight Center.

We would also like to acknowledge the Department of Civil, Mechanical and Environmental Engineering of the University of Trento for the
financial support given to Giorgio Doglioni during the development of this paper as a graduate student grant.



References

- 375 Allen, D. R., Fromm, M. D., III, G. P. K., and Nedoluha, G. E.: Smoke with Induced Rotation and Lofting (SWIRL) in the Stratosphere, *Journal of the Atmospheric Sciences*, 77, 4297 – 4316, <https://doi.org/10.1175/JAS-D-20-0131.1>, 2020.
- Baars, H., Ansmann, A., Ohneiser, K., Haarig, M., Engelmann, R., Althausen, D., Hanssen, I., Gausa, M., Pietruczuk, A., Szkop, A., Stachlewska, I. S., Wang, D., Reichardt, J., Skupin, A., Mattis, I., Trickl, T., Vogelmann, H., Navas-Guzmán, F., Haeffele, A., Acheson, K., Ruth, A. A., Tatarov, B., Müller, D., Hu, Q., Podvin, T., Goloub, P., Veselovskii, I., Pietras, C., Haeffelin, M., Fréville, P., Sicard, M., Comerón, A., Fernández García, A. J., Molero Menéndez, F., Córdoba-Jabonero, C., Guerrero-Rascado, J. L., Alados-Arboledas, L., Bortoli, D., 380 Costa, M. J., Dionisi, D., Liberti, G. L., Wang, X., Sannino, A., Papagiannopoulos, N., Boselli, A., Mona, L., D’Amico, G., Romano, S., Perrone, M. R., Belegante, L., Nicolae, D., Grigorov, I., Gialitaki, A., Amiridis, V., Soupiona, O., Papayannis, A., Mamouri, R.-E., Nisantzi, A., Heese, B., Hofer, J., Schechner, Y. Y., Wandinger, U., and Pappalardo, G.: The unprecedented 2017–2018 stratospheric smoke event: decay phase and aerosol properties observed with the EARLINET, *Atmospheric Chemistry and Physics*, 19, 15 183–15 198, 385 <https://doi.org/10.5194/acp-19-15183-2019>, 2019.
- Bourassa, A. E., Rieger, L. A., Zawada, D. J., Khaykin, S., Thomason, L. W., and Degenstein, D. A.: Satellite Limb Observations of Unprecedented Forest Fire Aerosol in the Stratosphere, *Journal of Geophysical Research: Atmospheres*, 124, 9510–9519, <https://doi.org/https://doi.org/10.1029/2019JD030607>, 2019.
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: 390 Tropospheric Aerosol Optical Thickness from the GOCART Model and Comparisons with Satellite and Sun Photometer Measurements, *Journal of the Atmospheric Sciences*, 59, 461 – 483, [https://doi.org/10.1175/1520-0469\(2002\)059<0461:TAOTFT>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<0461:TAOTFT>2.0.CO;2), 2002.
- Chin, M., Diehl, T., Dubovik, O., Eck, T. F., Holben, B. N., Sinyuk, A., and Streets, D. G.: Light absorption by pollution, dust, and biomass burning aerosols: a global model study and evaluation with AERONET measurements, *Annales Geophysicae*, 27, 3439–3464, <https://doi.org/10.5194/angeo-27-3439-2009>, 2009.
- 395 Christian, K., Wang, J., Ge, C., Peterson, D., Hyer, E., Yorks, J., and McGill, M.: Radiative Forcing and Stratospheric Warming of Pyrocumulonimbus Smoke Aerosols: First Modeling Results With Multisensor (EPIC, CALIPSO, and CATS) Views from Space, *Geophysical Research Letters*, 46, 10 061–10 071, <https://doi.org/https://doi.org/10.1029/2019GL082360>, 2019.
- Colarco, P., da Silva, A., Chin, M., and Diehl, T.: Online simulations of global aerosol distributions in the NASA GEOS-4 model and comparisons to satellite and ground-based aerosol optical depth, *Journal of Geophysical Research: Atmospheres*, 115, 400 <https://doi.org/https://doi.org/10.1029/2009JD012820>, 2010.
- Colarco, P. R., Nowotnick, E. P., Randles, C. A., Yi, B., Yang, P., Kim, K.-M., Smith, J. A., and Bardeen, C. G.: Impact of radiatively interactive dust aerosols in the NASA GEOS-5 climate model: Sensitivity to dust particle shape and refractive index, *Journal of Geophysical Research: Atmospheres*, 119, 753–786, <https://doi.org/https://doi.org/10.1002/2013JD020046>, 2014.
- Colarco, P. R., Gassó, S., Ahn, C., Buchard, V., da Silva, A. M., and Torres, O.: Simulation of the Ozone Monitoring Instrument aerosol index 405 using the NASA Goddard Earth Observing System aerosol reanalysis products, *Atmospheric Measurement Techniques*, 10, 4121–4134, <https://doi.org/10.5194/amt-10-4121-2017>, 2017.
- Considine, D. B., Douglass, A. R., Connell, P. S., Kinnison, D. E., and Rotman, D. A.: A polar stratospheric cloud parameterization for the global modeling initiative three-dimensional model and its response to stratospheric aircraft, *Journal of Geophysical Research: Atmospheres*, 105, 3955–3973, <https://doi.org/https://doi.org/10.1029/1999JD900932>, 2000.



- 410 Das, S., Colarco, P. R., Oman, L. D., Taha, G., and Torres, O.: Pyrocumulonimbus Events over British Columbia in 2017: The Long-term Transport and Radiative Impacts of Smoke Aerosols in the Stratosphere, *Atmospheric Chemistry and Physics Discussions*, 2020, 1–35, <https://doi.org/10.5194/acp-2020-1240>, 2021.
- de Laat, A. T. J., Stein Zweers, D. C., Boers, R., and Tuinder, O. N. E.: A solar escalator: Observational evidence of the self-lifting of smoke and aerosols by absorption of solar radiation in the February 2009 Australian Black Saturday plume, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD017016>, 2012.
- 415 Ditas, J., Ma, N., Zhang, Y., Assmann, D., Neumaier, M., Riede, H., Karu, E., Williams, J., Scharffe, D., Wang, Q., Saturno, J., Schwarz, J. P., Katich, J. M., McMeeking, G. R., Zahn, A., Hermann, M., Brenninkmeijer, C. A. M., Andreae, M. O., Pöschl, U., Su, H., and Cheng, Y.: Strong impact of wildfires on the abundance and aging of black carbon in the lowermost stratosphere, *Proceedings of the National Academy of Sciences*, 115, E11 595–E11 603, <https://doi.org/10.1073/pnas.1806868115>, 2018.
- 420 Doglioni, G.: GEOS CCM free-running simulation data of the Pacific-Northwest pyrocumulonimbus Event-like aerosol injection, SWIRL selection, <https://doi.org/10.5281/zenodo.6366106>, 2022.
- Douglass, A. R. and Kawa, S. R.: Contrast between 1992 and 1997 high-latitude spring Halogen Occultation Experiment observations of lower stratospheric HCl, *Journal of Geophysical Research: Atmospheres*, 104, 18 739–18 754, <https://doi.org/10.1029/1999JD900281>, 1999.
- 425 Ertel, H.: Ein neuer hydrodynamischer Erhaltungssatz, *Naturwissenschaften*, 30, 543–544, <https://doi.org/10.1007/BF01475602>, 1942.
- Fromm, M., Alfred, J., Hoppel, K., Hornstein, J., Bevilacqua, R., Shettle, E., Servranckx, R., Li, Z., and Stocks, B.: Observations of boreal forest fire smoke in the stratosphere by POAM III, SAGE II, and lidar in 1998, *Geophysical Research Letters*, 27, 1407–1410, <https://doi.org/10.1029/1999GL011200>, 2000.
- Fromm, M., Lindsey, D., Servranckx, R., Yue, G., Trickl, T., Sica, R., Doucet, P., and Godin-Beekmann, S.: The Untold Story of Pyrocumulonimbus, *Bulletin of the American Meteorological Society*, 91, <https://doi.org/10.1175/2010BAMS3004.1>, 2010.
- 430 Gelaro, R., McCarty, W., Suarez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419 – 5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 435 Hess, M., Koepke, P., and Schult, I.: Optical Properties of Aerosols and Clouds: The Software Package OPAC, *Bulletin of the American Meteorological Society*, 79, 831 – 844, [https://doi.org/10.1175/1520-0477\(1998\)079<0831:OPOAAC>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2), 1998.
- Kablick III, G. P., Allen, D. R., Fromm, M. D., and Nedoluha, G. E.: Australian PyroCb Smoke Generates Synoptic-Scale Stratospheric Anticyclones, *Geophysical Research Letters*, 47, e2020GL088 101, <https://doi.org/10.1029/2020GL088101>, e2020GL088101
- 440 10.1029/2020GL088101, 2020.
- Khaykin, S., Legras, B., Bucci, S., Sellitto, P., Isaksen, I., Tencé, F., Bekki, S., Bourassa, A., Rieger, L., Zawada, D., Jumelet, J., and Godin-Beekmann, S.: The 2019/20 Australian wildfires generated a persistent smoke-charged vortex rising up to 35 km altitude, *Communications Earth & Environment*, 1, 22, <https://doi.org/10.1038/s43247-020-00022-5>, 2020.
- 445 Khaykin, S. M., Godin-Beekmann, S., Hauchecorne, A., Pelon, J., Ravetta, F., and Keckhut, P.: Stratospheric Smoke With Unprecedentedly High Backscatter Observed by Lidars Above Southern France, *Geophysical Research Letters*, 45, 1639–1646, <https://doi.org/10.1002/2017GL076763>, 2018.



- Kloss, C., Berthet, G., Sellitto, P., Ploeger, F., Bucci, S., Khaykin, S., Jégou, F., Taha, G., Thomason, L. W., Barret, B., Le Flochmoen, E., von Hobe, M., Bossolasco, A., Bègue, N., and Legras, B.: Transport of the 2017 Canadian wildfire plume to the tropics via the Asian monsoon circulation, *Atmospheric Chemistry and Physics*, 19, 13 547–13 567, <https://doi.org/10.5194/acp-19-13547-2019>, 2019.
- 450 Lait, L. R.: An Alternative Form for Potential Vorticity, *Journal of Atmospheric Sciences*, 51, 1754 – 1759, [https://doi.org/10.1175/1520-0469\(1994\)051<1754:AAFFPV>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<1754:AAFFPV>2.0.CO;2), 1994.
- Lestrelin, H., Legras, B., Podglajen, A., and Salihoglu, M.: Smoke-charged vortices in the stratosphere generated by wildfires and their behaviour in both hemispheres: comparing Australia 2020 to Canada 2017, *Atmospheric Chemistry and Physics*, 21, 7113–7134, <https://doi.org/10.5194/acp-21-7113-2021>, 2021.
- 455 Molod, A., Takacs, L., Suarez, M., and Bacmeister, J.: Development of the GEOS-5 atmospheric general circulation model: evolution from MERRA to MERRA2, *Geoscientific Model Development*, 8, 1339–1356, <https://doi.org/10.5194/gmd-8-1339-2015>, 2015.
- Peterson, D., Campbell, J., and Hyer, E. e. a.: Wildfire-driven thunderstorms cause a volcano-like stratospheric injection of smoke, *npj Clim Atmos Sci* 1, 30, 587–590, <https://doi.org/10.1038/s41612-018-0039-3>, 2018.
- Peterson, D. A., Hyer, E. J., Campbell, J. R., Solbrig, J. E., and Fromm, M. D.: A Conceptual Model for Development of Intense Pyrocumulonimbus in Western North America, *Monthly Weather Review*, 145, 2235 – 2255, <https://doi.org/10.1175/MWR-D-16-0232.1>, 2017.
- 460 Peterson, D. A., Fromm, M. D., McRae, R. H. D., Campbell, J. R., Hyer, E. J., Taha, G., Camacho, C. P., Kablick, G. P., Schmidt, C. C., and DeLand, M. T.: Australia’s Black Summer pyrocumulonimbus super outbreak reveals potential for increasingly extreme stratospheric smoke events, *npj Climate and Atmospheric Science*, 4, 38, <https://doi.org/10.1038/s41612-021-00192-9>, 2021.
- Rienecker, M., Suarez, M., Todling, R., Bacmeister, J., Takacs, L., Liu, H., Gu, W., Sienkiewicz, M., Koster, R., Gelaro, R., Stajner, I., and 465 Nielsen, J.: The GEOS-5 Data Assimilation System— Documentation of Versions 5.0.1, 5.1.0, and 5.2.0, Tech. rep., Global Modelling and Assimilation Office (GMAO), NASA, <https://gmao.gsfc.nasa.gov/pubs/docs/tm27.pdf>, 2008.
- Torres, O., Bhartia, P. K., Taha, G., Jethva, H., Das, S., Colarco, P., Krotkov, N., Omar, A., and Ahn, C.: Stratospheric Injection of Massive Smoke Plume From Canadian Boreal Fires in 2017 as Seen by DSCOVR-EPIC, CALIOP, and OMPS-LP Observations, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032 579, <https://doi.org/https://doi.org/10.1029/2020JD032579>, e2020JD032579 470 2020JD032579, 2020.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geoscientific Model Development*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.
- 475 Yu, P., Toon, O. B., Bardeen, C. G., Zhu, Y., Rosenlof, K. H., Portmann, R. W., Thornberry, T. D., Gao, R.-S., Davis, S. M., Wolf, E. T., de Gouw, J., Peterson, D. A., Fromm, M. D., and Robock, A.: Black carbon lofts wildfire smoke high into the stratosphere to form a persistent plume, *Science*, 365, 587–590, <https://doi.org/10.1126/science.aax1748>, 2019.