

# Pathfinder v1.0.1: a Bayesian-inferred simple carbon-climate model to explore climate change scenarios

Thomas Bossy<sup>1,2,\*</sup>, Thomas Gasser<sup>1,\*</sup>, and Philippe Ciais<sup>2</sup>

<sup>1</sup>International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

<sup>2</sup>Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Gif-sur-Yvette, France

\*These authors contributed equally to this work.

**Correspondence:** Thomas Gasser (gasser@iiasa.ac.at)

**Abstract.** The Pathfinder model was developed to fill a perceived gap within the range of existing simple climate models. Pathfinder is a compilation of existing formulations describing the climate and carbon cycle systems, chosen for their balance between mathematical simplicity and physical accuracy. The resulting model is simple enough to be used with Bayesian inference algorithms for calibration, which enables assimilation of the latest data from complex Earth system models and the IPCC 6th assessment report, as well as a yearly update based on observations of global temperature and atmospheric CO<sub>2</sub>. The model's simplicity also enables coupling with integrated assessment models and their optimization algorithms, or running the model in a backward temperature-driven fashion. In spite of this simplicity, the model accurately reproduces behaviours and results from complex models – including [several](#) uncertainty ranges – when [running](#) following standardized diagnostic experiments. Pathfinder is open-source, and this is its first comprehensive description.

## 1 Introduction

Simple climate models (SCMs) [typically simulate global mean temperature change caused by either atmospheric concentration changes or anthropogenic emissions, of CO<sub>2</sub> and other climatically active species](#). They are most often composed of ad hoc parametric laws that emulate the behaviour of more complex Earth system models (ESMs). The emulation allows simulating large ensembles of experiments that would be too costly to compute with ESMs. However, the SCM denomination refers to a fairly broad range of models whose complexity can go from a couple of boxes that only emulate one part of the climate system (e.g. a global temperature impulse response function; Geoffroy et al., 2013b) to hundreds of state variables representing the different cycles of greenhouse gases and their effect on climate change (e.g. the compact Earth system model OSCAR; Gasser et al., 2017). Simpler models are easier and faster to solve, but they may not [be](#) adequate for all usages. Therefore, finding the “simplest but not simpler” model depends on a study's precise goals.

In our recent research, we have perceived a deficiency within the existing offer of SCMs, in spite of their large and growing number (Nicholls et al., 2020). We have therefore developed the Pathfinder model to fill this gap: it is a parsimonious [CO<sub>2</sub>-only](#) model that carefully balances simplicity and accuracy of representation of physical processes. Pathfinder was designed to fulfil three key requirements: 1. the capacity to be calibrated using Bayesian inference, 2. the capacity to be coupled with integrated

assessment models (IAMs), and 3. the capacity to explore a very large number of climate scenarios to narrow down those compatible with limiting climate impacts. The latter motivated the model's name.

While these three requirements clearly call for the simplest model possible, as they all need a fast solving model, they also imply a certain degree of complexity. The Bayesian calibration requires an explicit representation of the processes (i.e. the variables) that are used to constrain the model. Coupling with IAMs requires accurately embedding the latest advances of climate sciences to be policy relevant (National Academies of Sciences and Medicine, 2017). And exploring future climate impacts requires the flexibility to link additional (and potentially regional) impact variables to the core carbon-climate equations.

The Pathfinder model [is essentially an integration of existing formulations, adapted to our modelling framework and goals.](#) [It](#) is calibrated on Earth system models that contributed to the Coupled Model Intercomparison Project phase 6 (CMIP6), on additional data from the 6th assessment report of the IPCC (AR6), and on observations of global Earth properties up to the year 2021. [The calibration philosophy of Pathfinder is to use complex models as prior information, and only real-world observations and assessments combining many lines of evidence as constraints.](#)

[Compared to other SCMs \(Nicholls et al., 2020\), Pathfinder is much simpler than models like MAGICC \(Meinshausen et al., 2011\), OSCAR \(Gasser et al., 2017\) or even HECTOR \(Hartin et al., 2015\). It is comparable in complexity to FaIR \(Smith et al., 2018\) or BernSCM \(Strassmann and Joos, 2018\), although it is closer to the latter as it trades off an explicit representation of non-CO<sub>2</sub> species for one of the carbon cycle's main components. This choice was made to help calibration, keep the model invertible, and be compatible with IAMs such as DICE \(Nordhaus, 2017\). While most SCMs are calibrated using procedures that resemble Bayesian inference \(Nicholls et al., 2021\), Pathfinder relies on an established algorithm whose implementation is fully tractable, and that allows for an annual update as observations of atmospheric CO<sub>2</sub> and global temperature become available.](#)

Here, we present the first public release of Pathfinder [and its source code](#). We first provide a detailed description of the model's equations. We then describe the Bayesian setup used for calibration, the sources of prior information for it, and the resulting posterior configuration. We end with a validation of the model using standard diagnostic simulations and quantitative metrics for the climate system and carbon cycle.

## 2 Equations

An overview of Pathfinder is presented in Figure 1. The model is composed of a climate module, of three separate modules for the carbon cycle (ocean, land without land use and land permafrost), and of two additional modules describing global impacts: sea level rise (SLR), and surface ocean acidification. We do not emulate cycles of other non-CO<sub>2</sub> gases. Mathematically, the model is driven by prescribing time series of any combination of two of four variables: global mean surface temperature (GMST) anomaly (noted  $T$ ), global atmospheric CO<sub>2</sub> concentration ( $C$ ), global non-CO<sub>2</sub> effective radiative forcing ( $R_x$ ), and global anthropogenic emissions of CO<sub>2</sub> ( $E_{\text{CO}_2}$ ). The model can therefore be run in the traditional emission-driven and concentration-driven modes, but also in a temperature-driven mode [\(in terms of code, implemented as separate versions of the model\)](#). This is notably important for the calibration, during which it is driven by observations of GMST and atmospheric CO<sub>2</sub>.

The following presents all equations of the models. Variables are noted using Roman letters, and compiled in Tables B1 and B2. With a few exceptions, parameters are noted using Greek letters, and summarized in Tables B3 and B4. The model has 21 state variables that follow a first-order differential equations in time. The time variable is noted  $t$  and kept implicit unless  
60 required.

## 2.1 Climate

The GMST change ( $T$ ) induced by effective radiative forcing (ERF;  $R$ ) is represented using a widely used two-box energy balance model with deep ocean heat uptake efficacy (Geoffroy et al., 2013a; Armour, 2017). The first box represents the Earth surface's temperature (including atmosphere, land and surface ocean), and the other one is the deep ocean's temperature ( $T_d$ ).  
65 Their time-differential equations are:

$$\Theta_s \frac{dT}{dt} = R - \frac{\phi \ln(2)}{T_{2\times}} T - \epsilon_{\text{heat}} \theta (T - T_d) \quad (1)$$

and

$$\Theta_d \frac{dT_d}{dt} = \theta (T - T_d) \quad (2)$$

where  $\phi$  is the radiative parameter of CO<sub>2</sub>,  $T_{2\times}$  is the equilibrium climate sensitivity (ECS) at CO<sub>2</sub> doubling,  $\Theta_s$  is the heat capacity of the surface,  $\Theta_d$  is the heat capacity of the deep ocean,  $\theta$  is the heat exchange coefficient, and  $\epsilon_{\text{heat}}$  is the deep ocean heat uptake efficacy.  
70

The global ERF is simply the sum of the CO<sub>2</sub> contribution ( $R_{\text{CO}_2}$ ), expressed using the IPCC AR5 formula (Myhre et al., 2013), and that of non-CO<sub>2</sub> climate forcers ( $R_x$ ):

$$R = R_{\text{CO}_2} + R_x \quad (3)$$

75 with

$$R_{\text{CO}_2} = \phi \ln \left( \frac{C}{C_{\text{pi}}} \right) \quad (4)$$

where  $C_{\text{pi}}$  is the preindustrial atmospheric CO<sub>2</sub> concentration.

The above energy balance model naturally provides the ocean heat content (OHC;  $U_{\text{ohc}}$ ) as:

$$U_{\text{ohc}} = \alpha_{\text{ohc}} (\Theta_s T + \Theta_d T_d) \quad (5)$$

80 and the ocean heat uptake (OHU) as:

$$\frac{dU_{\text{ohc}}}{dt} = \alpha_{\text{ohc}} \left( \Theta_s \frac{dT}{dt} + \Theta_d \frac{dT_d}{dt} \right) \quad (6)$$

where  $\alpha_{\text{ohc}}$  is the fraction of energy used to warm the ocean (i.e. excluding the energy needed to heat up the atmosphere and land, and to melt ice).

## 2.2 Sea level rise

85 Global SLR has been implemented in Pathfinder as a variable of interest to model climate change impacts. In this version, it is firstly a proof of concept, modelled in a simple yet sensible manner. The total sea level rise ( $H_{\text{tot}}$ ) is the sum of contributions from thermal expansion ( $H_{\text{thx}}$ ), Greenland ice sheet (GIS;  $H_{\text{gis}}$ ), Antarctica ice sheet (AIS;  $H_{\text{ais}}$ ), and glaciers ( $H_{\text{gla}}$ ):

$$H_{\text{tot}} = H_{\text{thx}} + H_{\text{gis}} + H_{\text{ais}} + H_{\text{gla}} \quad (7)$$

The thermal expansion contribution scales linearly with the OHC (Goodwin et al., 2017; Fox-Kemper et al., 2021):

$$90 \quad H_{\text{thx}} = \Lambda_{\text{thx}} U_{\text{ohc}} \quad (8)$$

where  $\Lambda_{\text{thx}}$  is the scaling factor of the thermosteric contribution to SLR. Note, however, that the thermal capacity of the climate module does not match that of the real-world ocean (Geoffroy et al., 2013b), and so this equation cannot describe equilibrium SLR over millennial timescales.

To model contributions from ice sheets and glaciers, we followed the general approach of Mengel et al. (2016). The SLR caused by GIS follows a first-order differential equation with its specific timescale, and the equilibrium SLR from GIS is assumed to be a cubic function of GMST:

$$\frac{dH_{\text{gis}}}{dt} = \lambda_{\text{gis}} + \frac{1}{\tau_{\text{gis}}} (\Lambda_{\text{gis1}} T + \Lambda_{\text{gis3}} T^3 - H_{\text{gis}}) \quad (9)$$

where  $\lambda_{\text{gis}}$  is an offset parameter introduced because GIS was not in a steady state at the end of the preindustrial era,  $\Lambda_{\text{gis1}}$  is the linear term of equilibrium of GIS SLR,  $\Lambda_{\text{gis3}}$  is the cubic term of equilibrium of GIS SLR, and  $\tau_{\text{gis}}$  is the timescale of the GIS contribution. The motivation for replacing the quadratic term of Mengel et al. (2016) by a cubic one is the oddness of the cubic function that leads to negative (and not positive) SLR for negative  $T$  (which happens during the earlier years of the calibration run).

The contribution from glaciers is also a first-order differential equation with an equilibrium inspired by Mengel et al. (2016). We expanded it with a cubic term to account for the fact that we aggregate all glaciers together and allow more skewness in the curve describing the equilibrium SLR as a function of  $T$ . In addition, we added an exponential sensitivity to speed up the convergence to equilibrium under warmer climate:

$$\frac{dH_{\text{gla}}}{dt} = \lambda_{\text{gla}} + \frac{\exp(\gamma_{\text{gla}} T)}{\tau_{\text{gla}}} (\Lambda_{\text{gla}} (1 - \exp(-\Gamma_{\text{gla1}} T - \Gamma_{\text{gla3}} T^3)) - H_{\text{gla}}) \quad (10)$$

where  $\lambda_{\text{gla}}$  is an offset parameter accounting for the lack of initial steady-state,  $\Lambda_{\text{gla}}$  is the SLR potential if all glaciers melted,  $\Gamma_{\text{gla1}}$  is the linear sensitivity of glaciers' equilibrium to climate change,  $\Gamma_{\text{gla3}}$  is the cubic sensitivity of glaciers' equilibrium to climate change,  $\tau_{\text{gla}}$  is the timescale of the glaciers contribution, and  $\gamma_{\text{gla}}$  is the sensitivity of glaciers' timescale to climate change.

Following Mengel et al. (2016), the contribution from AIS is further divided in two terms, one for surface mass balance (SMB;  $H_{\text{ais,smb}}$ ) and one for solid ice discharge (SID;  $H_{\text{ais,sid}}$ ), so that  $H_{\text{ais}} = H_{\text{ais,smb}} + H_{\text{ais,sid}}$ . It is expected that precipitation will increase over Antarctica under higher GMST, leading to increase in SMB and to a negative sea level rise contribution

115 modeled as:

$$\frac{dH_{\text{ais,smb}}}{dt} = -\Lambda_{\text{ais,smb}} T \quad (11)$$

where  $\Lambda_{\text{ais,smb}}$  is the AIS SMB sensitivity to climate change (expressed in sea level equivalent). At the same time, increasing surface ocean temperatures will cause more SID through basal melting, which we model using a first-order differential equation assumed to be independent of the SMB effect, and with a term that speeds up the effect the more SID happened:

$$120 \quad \frac{dH_{\text{ais,sid}}}{dt} = \lambda_{\text{ais}} + \frac{1 + \alpha_{\text{ais}} H_{\text{ais,sid}}}{\tau_{\text{ais}}} (\Lambda_{\text{ais}} T - H_{\text{ais,sid}}) \quad (12)$$

where  $\lambda_{\text{ais}}$  is an offset parameter accounting for the lack of initial steady-state,  $\Lambda_{\text{ais}}$  is the SLR equilibrium of AIS SID,  $\tau_{\text{ais}}$  is the timescale of the AIS SID contribution, and  $\alpha_{\text{ais}}$  is the sensitivity of the timescale to past SID. In the model's code, however, we directly solve for the total AIS contribution as:

$$\frac{dH_{\text{ais}}}{dt} = -\Lambda_{\text{ais,smb}} T + \lambda_{\text{ais}} + \frac{1 + \alpha_{\text{ais}} (H_{\text{ais}} - H_{\text{ais,smb}})}{\tau_{\text{ais}}} (\Lambda_{\text{ais}} T - (H_{\text{ais}} - H_{\text{ais,smb}})) \quad (13)$$

## 125 **2.3 Ocean carbon**

To calculate the ocean carbon sink, we use the [elassie](#) mixed-layer impulse response function model from Joos et al. (1996), updated to the equivalent box-model formulation of Strassmann and Joos (2018), and extended in places to introduce parameter adjustments for calibration. In the model, the mixed layer is split into 5 boxes (subscript  $j$ ), as represented in Figure 2, so that the total carbon in the mixed layer pool ( $C_o$ ) is:

$$130 \quad C_o = \sum_j C_{o,j} \quad (14)$$

This total carbon mass is converted into a molar concentration of dissolved inorganic carbon (DIC;  $c_{\text{dic}}$ ) following:

$$c_{\text{dic}} = \frac{\alpha_{\text{dic}}}{\beta_{\text{dic}}} C_o \quad (15)$$

where  $\alpha_{\text{dic}}$  is a fixed conversion factor, and  $\beta_{\text{dic}}$  is a scaling factor for the conversion. (The latter can be seen as a factor multiplying the mixed layer depth: it is 1 if the depth is unchanged from the original Strassmann and Joos (2018) model.)

135 The non-linear carbonate chemistry in the mixed layer is emulated in two steps. First, the model's original polynomial function is used to determine the partial pressure of CO<sub>2</sub> affected by changes in DIC only ( $p_{\text{dic}}$ ):

$$\begin{aligned} p_{\text{dic}} = & (1.5568 - 0.013993 T_o) c_{\text{dic}} \\ & + (7.4706 - 0.20207 T_o) 10^{-3} c_{\text{dic}}^2 \\ & - (1.2748 - 0.12015 T_o) 10^{-5} c_{\text{dic}}^3 \\ & + (2.4491 - 0.12639 T_o) 10^{-7} c_{\text{dic}}^4 \\ & - (1.5768 - 0.15326 T_o) 10^{-10} c_{\text{dic}}^5 \end{aligned} \quad (16)$$

where  $T_o$  is the preindustrial surface ocean temperature. Second, the actual partial pressure of CO2 ( $p_{CO2}$ ) is calculated using  
 140 an exponential climate sensitivity (Takahashi et al., 1993; Joos et al., 2001):

$$p_{CO2} = (p_{dic} + C_{pi}) \exp(\gamma_{dic} T) \quad (17)$$

where  $\gamma_{dic}$  is the sensitivity of  $p_{CO2}$  to climate change.

The flux of carbon between the atmosphere and the ocean ( $F_{ocean}$ , defined positively if it is a carbon sink) is caused by the difference in partial pressure of CO2 in the atmosphere and at the oceanic surface, following an exchange rate that varies  
 145 linearly with GMST, that is here used as a proxy for wind changes:

$$F_{ocean} = \nu_{gx} (1 + \gamma_{gx} T) (C - p_{CO2}) \quad (18)$$

where  $\nu_{gx}$  is the preindustrial gas-exchange rate, and  $\gamma_{gx}$  is its sensitivity to climate change.

This flux of carbon entering the ocean is split between the mixed layer carbon subpools, and this added carbon is subsequently transported towards the deep ocean at a rate specific to each subpool. This leads to the following differential equations:

$$150 \quad \frac{dC_{o,j}}{dt} = -\frac{C_{o,j}}{\kappa_{\tau_o} \tau_{o,j}} + \alpha_{o,j} F_{ocean}, \quad \forall j \quad (19)$$

where  $\alpha_{o,j}$  are the subpools' splitting shares (with  $\sum_j \alpha_{o,j} = 1$ ),  $\tau_{o,j}$  are the subpools' timescales for transport to the deep ocean, and  $\kappa_{\tau_o}$  is a scaling factor applied to all subpools. Finally, the deep ocean carbon pool ( $C_d$ ) is obtained through mass balance:

$$\frac{dC_d}{dt} = \sum_j \frac{C_{o,j}}{\kappa_{\tau_o} \tau_{o,j}} \quad (20)$$

## 155 2.4 Ocean acidification

While in the real world, ocean acidification is directly related to the carbonate chemistry and the ocean [uptake of anthropogenic carbonsink](#), we do not have a simple formulation at our disposal that could link it to our ocean carbon cycle module. We therefore use a readily available emulation of the surface ocean acidification (pH) that links it directly to the atmospheric concentration of CO2 (Bernie et al., 2010) with the following polynomial approximation:

$$160 \quad pH = \kappa_{pH} (8.5541 - 0.00173 C + 1.3264 \cdot 10^{-6} C^2 - 4.4943 \cdot 10^{-10} C^3) \quad (21)$$

where  $\kappa_{pH}$  is a scaling factor (that defaults to 1). We note that this approach is reasonable for the surface ocean, as it quickly equilibrates with the atmosphere (but it would not work for the deep ocean).

## 2.5 Land carbon

The land carbon module of Pathfinder is a simplified version of the one in OSCAR (Gasser et al., 2017, 2020). It is shrunk  
 165 down to four global carbon pools: vegetation, litter, active and passive soil (see Figure 3). All terrestrial biomes are lumped

together, and there is therefore no accounting of the impact of land use change on the land carbon cycle in this version of Pathfinder. This is an extreme assumption – although very common in SCMs – motivated by simplicity, and it implies that CO2 emissions from fossil fuel burning and land use change are assumed to behave in the exact same way, in spite of their not doing so in reality (Gitz and Ciais, 2003; Gasser and Ciais, 2013).

170 The vegetation carbon pool ( $C_v$ ) results from the balance between net primary productivity (NPP;  $F_{\text{npp}}$ ), emission from wildfires ( $E_{\text{fire}}$ ), emission from harvest and grazing ( $E_{\text{harv}}$ ), and loss of carbon from biomass mortality ( $F_{\text{mort}}$ ):

$$\frac{dC_v}{dt} = F_{\text{npp}} - E_{\text{fire}} - E_{\text{harv}} - F_{\text{mort}} \quad (22)$$

NPP is expressed as its own preindustrial value multiplied by a function of CO2 and of GMST ( $r_{\text{npp}}$ ). This function thus embeds the so-called CO2-fertilisation effect, [whereby NPP increases with atmospheric CO2](#), described using a generalised logarithmic functional form:

$$F_{\text{npp}} = F_{\text{npp0}} r_{\text{npp}} \quad (23)$$

with

$$r_{\text{npp}} = \left( 1 + \frac{\beta_{\text{npp}}}{\alpha_{\text{npp}}} \left( 1 - \left( \frac{C}{C_{\text{pi}}} \right)^{-\alpha_{\text{npp}}} \right) \right) (1 + \gamma_{\text{npp}} T) \quad (24)$$

where  $F_{\text{npp0}}$  is the preindustrial NPP,  $\beta_{\text{npp}}$  is the CO2-fertilisation sensitivity,  $\alpha_{\text{npp}}$  is the CO2-fertilisation shape parameter for saturation, and  $\gamma_{\text{npp}}$  is the sensitivity of NPP to climate change ([that can be positive or negative](#)). [The generalised logarithmic functional form implies that:  \$r\_{\text{npp}} \rightarrow \(1 + \beta\_{\text{npp}} \ln\(C/C\_{\text{pi}}\)\)\(1 + \gamma\_{\text{npp}} T\)\$  as  \$\alpha\_{\text{npp}} \rightarrow 0^+\$ .](#)

Harvesting and mortality fluxes are taken proportional to the carbon pool itself even though in reality the mortality fluxes are climate dependent. For simplicity we assume a constant mortality following the equations in OSCAR (Gasser et al., 2017):

$$185 \quad E_{\text{harv}} = \nu_{\text{harv}} C_v \quad (25)$$

and

$$F_{\text{mort}} = \nu_{\text{mort}} C_v \quad (26)$$

where  $\nu_{\text{harv}}$  is the harvesting/grazing rate, and  $\nu_{\text{mort}}$  is the mortality rate.

Wildfires emissions are also assumed proportional to the vegetation carbon pool, but with an additional linear dependency of the emission rate on CO2 ([as a proxy of changes in leaf area index and evapotranspiration](#)) and GMST ( $r_{\text{fire}}$ ):

$$E_{\text{fire}} = \nu_{\text{fire}} r_{\text{fire}} C_v \quad (27)$$

with

$$r_{\text{fire}} = \left( 1 + \beta_{\text{fire}} \left( \frac{C}{C_{\text{pi}}} - 1 \right) \right) (1 + \gamma_{\text{fire}} T) \quad (28)$$

where  $\nu_{\text{fire}}$  is the wildfires rate,  $\beta_{\text{fire}}$  is the sensitivity of wildfires to CO<sub>2</sub>, and  $\gamma_{\text{fire}}$  is their sensitivity to climate change.

195 Soil carbon is divided into three pools. The litter carbon pool ( $C_{s1}$ ) receives the mortality flux as sole input, it emits part of its carbon through heterotrophic respiration ( $E_{\text{rh1}}$ ), and it transfers another part to the next pool through stabilization ( $F_{\text{stab}}$ ):

$$\frac{dC_{s1}}{dt} = F_{\text{mort}} - F_{\text{stab}} - E_{\text{rh1}} \quad (29)$$

Similarly, the active soil carbon pool ( $C_{s2}$ ) receives the stabilization flux, is respired ( $E_{\text{rh2}}$ ), and transfers carbon to the last pool through passivization ( $F_{\text{pass}}$ ):

$$200 \quad \frac{dC_{s2}}{dt} = F_{\text{stab}} - F_{\text{pass}} - E_{\text{rh2}} \quad (30)$$

The passive carbon pool ( $C_{s3}$ ) receives this final input flux and is respired ( $E_{\text{rh3}}$ ):

$$\frac{dC_{s3}}{dt} = F_{\text{pass}} - E_{\text{rh3}} \quad (31)$$

Although information pertaining to this fourth pool is not commonly provided by ESMs, it was introduced in Pathfinder to adjust the complex models' turnover time of soil carbon to better match isotopic data (He et al., 2016). For completeness, we  
 205 note that the total heterotrophic respiration is  $E_{\text{rh}} = E_{\text{rh1}} + E_{\text{rh2}} + E_{\text{rh3}}$ , and the total soil carbon pool is  $C_s = C_{s1} + C_{s2} + C_{s3}$ .

All soil-originating fluxes are taken proportional to their pool of origin, and multiplied by a function ( $r_{\text{rh}}$ ) explained hereafter. For the litter pool, this gives:

$$E_{\text{rh1}} = \nu_{\text{rh1}} r_{\text{rh}} C_{s1} \quad (32)$$

and

$$210 \quad F_{\text{stab}} = \nu_{\text{stab}} r_{\text{rh}} C_{s1} \quad (33)$$

where  $\nu_{\text{rh1}}$  is the litter respiration rate, and  $\nu_{\text{stab}}$  is the stabilization rate. For the active soil pool, we have:

$$E_{\text{rh2}} = \frac{\nu_{\text{rh23}} - \nu_{\text{rh3}} \alpha_{\text{pass}}}{1 - \alpha_{\text{pass}}} r_{\text{rh}} C_{s2} \quad (34)$$

and

$$F_{\text{pass}} = \nu_{\text{rh3}} \frac{\alpha_{\text{pass}}}{1 - \alpha_{\text{pass}}} r_{\text{rh}} C_{s2} \quad (35)$$

215 and for the passive soil pool:

$$E_{\text{rh3}} = \nu_{\text{rh3}} r_{\text{rh}} C_{s3} \quad (36)$$

where  $\nu_{\text{rh23}}$  is the soil respiration rate (averaged over active and passive pools),  $\nu_{\text{rh3}}$  is the passive soil respiration rate, and  $\alpha_{\text{pass}}$  is the fraction of passive carbon (over active+passive soil carbon). This slightly convoluted formulation is motivated by the lack of information regarding the active/passive split in ESMs, which we alleviate using additional data during calibration.

220 In addition, the function  $r_{rh}$ , describing the dependency of respiration (and related fluxes) on temperature and on the availability of fresh organic matter to be decomposed, is defined as:

$$r_{rh} = \underbrace{\left(1 + \beta_{rh} \left( \frac{C_{s1}}{C_{s1} + C_{s2} + C_{s3}} \left(1 + \frac{\nu_{stab}}{\nu_{rh23}}\right) - 1 \right)\right)}_{\left(1 + \beta_{rh} \left( \frac{C_{s1}}{C_s} \frac{C_s(t_0)}{C_{s1}(t_0)} - 1 \right)\right)} \exp(\gamma_{rh} T) \quad (37)$$

where  $\beta_{rh}$  is the sensitivity of the respiration to fresh organic matter availability (expressed here as the relative change in the  $C_{s1}/C_s$  ratio with regard to preindustrial times), and  $\gamma_{rh}$  is its sensitivity to climate change (equivalent to a " $Q_{10}$ " formulation with  $Q_{10} = \exp(10 \gamma_{rh})$ ).

Finally, the net carbon flux from the atmosphere to the land ( $F_{land}$ , defined positively if it is a carbon sink) is obtained as the net budget of all pools combined:

$$F_{land} = F_{npp} - E_{fire} - E_{harv} - E_{rh} \quad (38)$$

and this system of equations leads to the following preindustrial steady-state:

$$230 \quad \begin{cases} C_v(t_0) = \frac{F_{npp0}}{\nu_{fire} + \nu_{harv} + \nu_{mort}} \\ C_{s1}(t_0) = C_v(t_0) \frac{\nu_{mort}}{\nu_{rh1} + \nu_{stab}} \\ C_{s2}(t_0) = C_{s1}(t_0) \frac{\nu_{stab}}{\nu_{rh23}} (1 - \alpha_{pass}) \\ C_{s3}(t_0) = C_{s1}(t_0) \frac{\nu_{stab}}{\nu_{rh23}} \alpha_{pass} \end{cases} \quad (39)$$

## 2.6 Permafrost carbon

As the land carbon cycle described in the previous section does not account for permafrost carbon, we implemented this feedback using the emulator developed by Gasser et al. (2018) but aggregated into a unique global region. [Figure 4 gives a representation of the permafrost module as described in the following.](#) The emulation starts with a theoretical thawed fraction ( $\bar{a}$ ) that represents the fraction of thawed carbon under steady-state for a certain level of local warming. It is formulated with a sigmoid function (that equals 0 at preindustrial and 1 under very high GMST):

$$\bar{a} = -a_{min} + \frac{(1 + a_{min})}{\left(1 + \left(\left(1 + \frac{1}{a_{min}}\right)^{\kappa_a} - 1\right) \exp(-\gamma_a \kappa_a \alpha_{lst} T)\right)^{\frac{1}{\kappa_a}}} \quad (40)$$

where  $-a_{min}$  is the minimum thawed fraction (corresponding to 100% frozen soil carbon),  $\kappa_a$  is a shape parameter determining the asymmetry of the function,  $\gamma_a$  is the sensitivity of the theoretical thawed fraction to local climate change, and  $\alpha_{lst}$  is the proportionality factor between local and global climate change.

The actual thawed fraction ( $a$ ) then moves towards its theoretical value at a speed that depends on whether it is thawing (i.e.  $a < \bar{a}$ ) or freezing (i.e.  $a > \bar{a}$ ). This is written as a non-linear differential equation:

$$\frac{da}{dt} = 0.5 (\nu_{thaw} + \nu_{froz}) (\bar{a} - a) + 0.5 |(\nu_{thaw} - \nu_{froz}) (\bar{a} - a)| \quad (41)$$

where  $\nu_{\text{thaw}}$  is the rate of thawing, and  $\nu_{\text{froz}}$  is the rate of freezing. Because  $\nu_{\text{thaw}} > \nu_{\text{froz}}$ , the absolute value in the equation  
 245 leads to the right-hand side being  $\nu_{\text{thaw}}(\bar{a} - a)$  if  $a < \bar{a}$ , or  $\nu_{\text{froz}}(\bar{a} - a)$  if  $a > \bar{a}$ . The change in the pool of frozen carbon ( $C_{\text{fr}}$ ) naturally follows:

$$\frac{dC_{\text{fr}}}{dt} = -\frac{da}{dt} C_{\text{fr}0} \quad (42)$$

where  $C_{\text{fr}0}$  is the amount of frozen carbon at preindustrial times.

Thawed carbon is not directly emitted to the atmosphere: it is split into three thawed carbon subpools ( $C_{\text{th},j}$ ) that have their  
 250 own decay time, but are all affected by an additional function ( $r_{\text{rt}}$ ). This leads to the following budget equations:

$$\frac{dC_{\text{th},j}}{dt} = -\alpha_{\text{th},j} \frac{dC_{\text{fr}}}{dt} - \frac{C_{\text{th},j}}{\kappa_{\tau_{\text{th}}} \tau_{\text{th},j}} r_{\text{rt}}, \quad \forall j \quad (43)$$

where  $\alpha_{\text{th},j}$  are the subpools' splitting shares (with  $\sum_j \alpha_{\text{th},j} = 1$ ),  $\tau_{\text{th},j}$  are the subpools' decay times, and  $\kappa_{\tau_{\text{th}}}$  is a scaling factor applied to all subpools. The additional  $r_{\text{rt}}$  function describes the sensitivity of heterotrophic respiration to climate change in boreal regions, using a Gaussian formula:

$$255 \quad r_{\text{rt}} = \exp(\kappa_{\text{rt}} \gamma_{\text{rt}1} \alpha_{\text{lst}} T - \kappa_{\text{rt}} \gamma_{\text{rt}2} (\alpha_{\text{lst}} T)^2) \quad (44)$$

where  $\kappa_{\text{rt}}$  is a factor scaling the sensitivity of thawed carbon against that of regular soil carbon,  $\gamma_{\text{rt}1}$  is the sensitivity to local temperature change (i.e. a Q10), and  $\gamma_{\text{rt}2}$  is the quadratic term in the latter sensitivity that represents a saturation effect. Noting that all the emitted carbon is assumed to be CO<sub>2</sub>, the global emission from permafrost ( $E_{\text{pf}}$ ) is thus:

$$E_{\text{pf}} = \sum_j \frac{C_{\text{th},j}}{\kappa_{\tau_{\text{th}}} \tau_{\text{th},j}} r_{\text{rt}} \quad (45)$$

## 260 2.7 Atmospheric CO<sub>2</sub>

The change in atmospheric concentration of CO<sub>2</sub> is the budget of all carbon cycle fluxes to which we add the exogenous anthropogenic emissions ( $E_{\text{CO}_2}$ ):

$$\alpha_C \frac{dC}{dt} = E_{\text{CO}_2} + E_{\text{pf}} - F_{\text{land}} - F_{\text{ocean}} \quad (46)$$

where  $\alpha_C$  is the conversion factor from volume fraction to mass for CO<sub>2</sub>.

## 265 3 Bayesian calibration

### 3.1 Principle

Bayesian inference is a powerful tool for assimilating observational data into reduced-complexity models such as Pathfinder (Ricciuto et al., 2008). The approach consists in deducing **joint** probability distributions of parameters from a priori knowledge on those distributions and on distributions of observations of some of the model's **state** variables, using Bayes' theorem (Bayes,

270 1763). [Summarily, the Bayesian calibration updates the joint distribution of parameters to make it as compatible with the constraints as possible given their prior estimates, which increases internal coherence of Pathfinder by excluding combination of parameters that are unlikely.](#)

[Such a Bayesian calibration is vulnerable to the possibility that the priors draw on the same information as the constraints. However, given that Pathfinder is a patchwork of emulators whose parameters are obtained independently from one another](#)  
 275 [and following differing experimental setups, we expect that the coherence of information contained within the priors and the constraints is very low. Our choice of using only complex models as prior information and only observations and assessments as constraints also aims at limiting this vulnerability.](#)

Concretely, the posterior probability  $\mathcal{P}_{\text{post}}$  of a sample  $k$  from the joint parameters distribution  $\xi_k$ , conditional to a set of observations  $\mathbf{x}$ , is proportional (symbol  $\propto$ ) to its own prior probability  $\mathcal{P}_{\text{pre}}$  and to the likelihood  $\mathcal{L}$  of the model simulating  $\mathbf{x}$   
 280 given  $\xi_k$ :

$$\mathcal{P}_{\text{post}}(\xi_k|\mathbf{x}) \propto \mathcal{L}(\mathbf{x}|\xi_k) \mathcal{P}_{\text{pre}}(\xi_k) \quad (47)$$

Here, we assume all observations are independently and identically distributed following a normal distribution (with mean values  $\mu_{\mathbf{x}}$ , and standard deviations  $\sigma_{\mathbf{x}}$  expressed in real physical units), which leads to the following likelihood:

$$\mathcal{L}(\mathbf{x}|\xi_k) = \prod_{i=1}^{n_x} \frac{1}{\sigma_{x,i} \sqrt{2\pi}} \exp\left(-\frac{(\mathcal{F}_i(\xi_k) - \mu_{x,i})^2}{2\sigma_{x,i}^2}\right) \quad (48)$$

285 where  $\mathcal{F}_i(\xi_k)$  is the model's output for the  $i$ -th observable (out of  $n_x$ ) with input parameters  $\xi_k$ .

### 3.2 [Implementation](#)

The Pathfinder model is a set of [\(discretized\)](#) differential equations with a number of input parameters, of which  $n_{\xi}$  are calibrated through Bayesian inference, and an additional two input variables provided as time series (i.e. one value per time step required). While the two input time series can be any combination of two out of four variables (anthropogenic CO2  
 290 emissions, non-CO2 ERF, atmospheric CO2 concentration, or GMST), for calibration we use the two most well constrained variables that are direct physical observations of the global Earth system: atmospheric CO2 and GMST. These input time series cover the historical period from 1751 to 2020<sup>4</sup>. Therefore, the  $\xi_k$  vector is:

$$\xi_k = \left\{ \{\xi_j\}_{j=1}^{n_{\xi}}, \{C(t)\}_{t=1751}^{2021}, \{T(t)\}_{t=1751}^{2021} \right\}_k \quad (49)$$

However, to ease the computation by reducing the dimension of the system, we do not use annual time series of observations  
 295 as inputs, but we assume that each input time series (for variable  $X$  being  $C$  or  $T$ ) follows:

$$X(t) = X_{\mu}(t) + \tilde{\sigma}_X X_{\sigma}(t) + \epsilon_X \text{AR1}(t; \rho_X) \quad (50)$$

where  $X_{\mu}$  and  $X_{\sigma}$  are fixed exogenous annual time series (i.e. structural parameters),  $\tilde{\sigma}_X$  is the relative standard deviation of the time series (without noise),  $\epsilon_X$  is the noise intensity, and AR1 is an autoregressive process of order 1 and autocorrelation

parameter  $\rho_X$ . This assumption leads to the final expression of the  $\xi_k$  vector:

$$\xi_k = \left\{ \left\{ \xi_j \right\}_{j=1}^{n_\xi}, \tilde{\sigma}_C, \epsilon_C, \rho_C, \tilde{\sigma}_T, \epsilon_T, \rho_T, \right\}_k \quad (51)$$

During this Bayesian assimilation, the Pathfinder model is run solely over the historical period (from 1750 to 2021), as the constraints concern only preindustrial or historical years. For the computation, the time-differential system of Pathfinder is solved using an implicit-explicit numerical scheme (also called IMEX), with a time step of one quarter of a year. This solving scheme relies on: first, writing the differential equations of all state variables  $X$  as:

$$\frac{dX}{dt} = -\nu X + \mathcal{R} \quad (52)$$

where  $\nu$  is the constant speed of the linear part of the differential equation, and  $\mathcal{R}$  is its non-linear part; second, discretizing these equations as:

$$\frac{X(t + \delta t) - X(t)}{\delta t} = -\nu X(t + \delta t) + \mathcal{R}(t) \quad (53)$$

where  $\delta t$  is the solving time step (which is  $\frac{1}{n_t}$  times the annual time step of the model's inputs and outputs, here  $n_t = 4$ ); and third, explicitly solving for all  $X(t + \delta t)$ . We note this is also the default solving scheme for regular simulations with the model, although the value of  $n_t$  can be altered and alternative schemes are available.

The Bayesian procedure itself is implemented using the Python computer language, and specifically the PyMC3 package (Salvatier et al., 2016). The solving of equation 47 and its normalization distribution sampling and parameter distributions estimation are done using the package's through a full-rank Automatic Differentiation Variational Inference (ADVI) algorithm (Kucukelbir et al., 2017), with 100,000 iterations (and default algorithm options). The choice of variational inference instead of Markov chain Monte Carlo is motivated by the significant size our model (Blei et al., 2017) and the speed of ADVI. An additional strength of the full-rank version of the ADVI algorithm is its speed, and its ability to generate correlated (i.e. joint) posterior distributions even if the prior ones are uncorrelated. Convergence of the algorithm was controlled through convergence of the ELBO metric (Kucukelbir et al., 2017). All results presented hereafter are obtained through drawing 2000 sets of parameters – that we call configurations – from the posterior or prior distributions.

### 3.3 Constraints

We use a set of 19 constraints related to all aspects of the model that correspond to the set of observations  $x$  in the Bayesian calibration. Many of the constraints are observations, but some are ranges assessed by expert panels such as the Global Carbon Project or the IPCC. They cover either a recent point in time or an assumed preindustrial equilibrium, and they are typically taken over a period of at least a few years to reduce the effect of natural variability. During the Bayesian assimilation, the Pathfinder model is run solely over the historical period (from 1750 to 2021).

Table 1 summarises these constraints, the periods over which they are considered, and their distributions. The following subsections provide further details on the constraints, and the constraints distributions are shown in Figure 6.

### 3.3.1 Climate system

330 To constrain the temperature response, we use the same five data sets of observed GMST as in Section 3.4.8, to derive average and standard deviation of two constraints: the average GMST change, and the average GMST yearly trend obtained through second-order accuracy gradient (Fornberg, 1988), both over the latest 20 years of data (2002–2021). Because this data is already used as input to the Bayesian setup, albeit in a different way, it does not provide much of a constraint, and is used mostly to ensure the  $\tilde{\sigma}_T$  and  $\epsilon_T$  parameters remain within sensible range.

335 To further constrain the climate system, we use the mean OHU assessed by the IPCC AR6 over 2006–2018 (Gulev et al., 2021, Table 2.7), and the non-CO2 ERF (averaged over 2010–2019) also estimated for the AR6. The central value of the latter is taken from Dentener et al. (2021, Table AIII.3, and corresponding GitHub repository), and its uncertainty is constructed using data from Forster et al. (2021, Table 7.8) and assuming the ERF of all species are normally distributed and uncorrelated, but fully correlated in time for each separate species (which likely overestimates the uncertainty).

340 To better align with the IPCC AR6, we also constrain the ECS of our model (i.e. the  $T_{2\times}$  parameters). To do so, because the distribution of ECS cannot be assumed normal, we follow the framework of Roe and Baker (2007) who define the climate feedback factor  $ff$  so that  $T_{2\times} = T_{2\times}^* / (1 - ff)$ , where  $T_{2\times}^*$  is the minimal ECS value (roughly corresponding to the Planck feedback). We assume this feedback factor follows a logit-normal distribution, which implies  $\text{logit}(ff) = \ln(ff / (1 - ff)) = \ln(T_{2\times} / T_{2\times}^* - 1)$  follows a normal distribution. We therefore constrain  $\text{logit}(ff)$ , using distribution parameters and a value of

345  $T_{2\times}^*$  calibrated to fit the probabilistic ranges of ECS provided by the AR6. This fit of the ECS distribution is illustrated in Figure B9.

### 3.3.2 Carbon cycle

Similarly to what is done with GMST, we constrain the atmospheric CO2 level over the latest 10 years of data (2012–2021) using the NOAA/ESRL data (Tans and Keeling, 2010). The rest of the global CO2 budget is constrained using the 2021 Global

350 Carbon Budget (GCB; Friedlingstein et al., 2022). We use namely the net atmospheric CO2 growth and total anthropogenic emissions (fossil and land use) over the [last](#) 10 years, and the ocean and land carbon sinks accumulated since the beginning of the instrumental measurement period (1960–2020). Note that our definition of the land carbon sink ignoring land use change is consistent with that of the GCB.

Given its number of parameters and their inconsistent sources, we further constrain the land carbon module by considering

355 present-day (mean over 1998–2002) NPP (Ciais et al., 2013; Zhao et al., 2005), and preindustrial vegetation and soil carbon pools. These preindustrial pools are taken from the AR6 for the central value (Canadell et al., 2021, Figure 5.12), but their relative uncertainty is taken from the AR5 (Ciais et al., 2013, Figure 6.1) since it is lacking in the AR6. In addition, the soil carbon pool constraint is corrected downward by estimates of peatland carbon (Yu et al., 2010, Table 1), since it is an ecosystem missing in TRENDY models (and in ours) but not in the IPCC assessments.

### 360 3.3.3 Sea level rise

To constrain the separate SLR contributions from thermal expansion, GIS, AIS and glaciers, we use the model-based SLR speed estimates over the recent past (averaged over 2006–2018) reported in the AR6 (Fox-Kemper et al., 2021, Table 9.5). To constrain the total contribution, we also use the historical (1901–1990) sea level rise inferred from tide gauges from the same source, although the value is corrected upward for the missed impact of uncharted glaciers (Parkes and Marzeion, 2018).

365 Contrarily to all other modules, the SLR module is not assumed to start at steady-state in 1750, which is represented through the  $\lambda_{\text{ice}}$  ( $\text{ice} \in [\text{gla}, \text{gis}, \text{ais}]$ ) parameters. We assume this is entirely due to the so-called little ice age (LIA) relaxation, which we assume can be simply modeled in Pathfinder through exponential decay of our three ice-related contributions since  $t_0 = 1750$ . This gives a net LIA contribution of  $H_{\text{lia}} = \sum_{\text{ice}} \lambda_{\text{ice}} \tau_{\text{ice}} \exp\left(-\frac{t-t_0}{\tau_{\text{ice}}}\right)$ . We constrain this diagnostic variable using the global SLR reported by Slangen et al. (2016) over 1900–2005 for their control experiment.

### 370 3.4 Parameters (prior distributions)

Out of the model’s 77 parameters, 33 are assumed to be fixed (i.e. they are structural parameters), and the remaining  $n_{\xi} = 44$  parameters are estimated through Bayesian inference. Prior distributions of the  $\xi_j$  parameters are assumed log-normal if the physical parameter must be defined positive, logit-normal if it must be between 0 and 1, and normal otherwise. To avoid extreme parameter values that could make the model diverge during calibration, the posterior distributions are bound to  
375  $\mu_{\xi,j} \pm 5\sigma_{\xi,j}$ , where  $\mu_{\xi,j}$  and  $\sigma_{\xi,j}$  are the mean and standard deviation of the  $j$ -th parameter’s prior distribution. These two values are taken from the literature, deduced from multi-model ensembles, or in a few instances arbitrarily set, as described in the following subsections. Note that when parameters are deduced from multi-model ensembles, there are effectively two rounds of calibration: first, a calibration on individual models using ordinary least square regressions to obtain prior distributions, and second, the Bayesian calibration itself that leads to the posterior distributions. In addition, the prior distributions of  $\tilde{\sigma}_X$ ,  $\epsilon_X$  and  
380  $\rho_X$  are assumed normal, half-normal and uniform, respectively (see Figure 5). All prior distributions are assumed independent, so that the prior joint distribution  $\xi$  does not exhibit any covariance.

All parameters are summarised in Tables B5 and B6 along with their properties and values. [The following subsections further explain how the prior distributions of the parameters are established, and these distributions are shown in Figure 5](#)

#### 3.4.1 Climate

385 All the parameters of the climate module are calibrated. The prior [distribution of the](#) radiative parameter  $\phi$  is taken from the AR5 (Myhre et al., 2013, Table 8.SM.1). All other prior [distributions of the](#) parameters of the climate module (i.e.  $T_{2\times}$ ,  $\Theta_s$ ,  $\Theta_d$ ,  $\theta$  and  $\epsilon_{\text{heat}}$ ) are taken from 35 CMIP6 models whose climate responses were derived for the AR6 using the *abrupt-4xCO2* experiment (Smith et al., 2021, Section 7.SM.2.1, and corresponding GitHub repository). Here,  $T_{2\times}$  is simply assumed to be half the reported equilibrium temperature at quadrupled CO2. In addition, the prior [distribution of the](#) ocean warming fraction  
390  $\alpha_{\text{ohc}}$  is taken from the AR6 (Forster et al., 2021, Table 7.1).

### 3.4.2 Sea level rise

Some parameters from the SLR module are structural: the maximum SLR contribution from glaciers ( $\Lambda_{\text{gla}}$ ) is taken from Fox-Kemper et al. (2021, Section 9.6.3.2), the equilibrium AIS SLR ( $\Lambda_{\text{ais}}$ ) is from (Church et al., 2013, Figure 13.14), and the  $\tau_{\text{gis}}$ ,  $\tau_{\text{gla}}$  and  $\tau_{\text{ais}}$  timescales are the mean values from Mengel et al. (2016, Table S1) (assuming they provide the 90%-range of a log-normal distribution). All other parameters are calibrated. The prior [distribution of the](#) thermosteric parameter  $\Lambda_{\text{thx}}$  is taken from the AR6 (Fox-Kemper et al., 2021, Section 9.2.4.1), as are the prior [distributions of the](#) preindustrial offset parameters  $\lambda_{\text{gis}}$ ,  $\lambda_{\text{gla}}$  and  $\lambda_{\text{ais}}$  (Fox-Kemper et al., 2021, earliest period of Table 9.5). For the remaining parameters, we derive prior distributions using SLR projections compiled by Edwards et al. (2021) for a number of ice sheets and glaciers models, over various RCP and SSP scenarios. Using the models' outputs, we apply equation 9 to estimate the  $\Lambda_{\text{gis1}}$  and  $\Lambda_{\text{gis3}}$  parameters, equation 10 for the  $\Gamma_{\text{gla1}}$ ,  $\Gamma_{\text{gla3}}$  and  $\gamma_{\text{gla}}$  parameters, and equation 12 for the  $\Lambda_{\text{ais,smb}}$  and  $\alpha_{\text{ais}}$  parameters. During these fits, all other parameters are assumed to take their default value if structural, and their best-guess value otherwise. Results of this calibration on the individual models compiled by Edwards et al. (2021) are shown for each SLR contribution in Figures B6, B7 and B8.

### 3.4.3 Ocean carbon

The ocean carbon cycle module has a number of structural parameters:  $\alpha_{\text{dic}}$ , all  $\alpha_{o,j}$  and all  $\tau_{o,j}$  are taken from Strassmann and Joos (2018, Tables A2 and A3, based on the Princeton model). The prior [distribution of the](#) adjustment factor  $\kappa_{\tau_o}$  is arbitrarily taken to [apply](#) add a 20% uncertainty on the oceanic transport timescales. All other prior [distributions](#) for this module's parameters are derived from 12 CMIP6 models with interactive carbon cycle that contributed to C4MIP (Arora et al., 2020).  $T_o$  is taken [on average over](#) from the *piControl* simulation.  $\nu_{\text{gx}}$  and  $\gamma_{\text{gx}}$  are calibrated by applying equation 18 to the models' outputs for the *IpctCO2*, *IpctCO2-rad* and *IpctCO2-bgc* experiments, while  $\beta_{\text{dic}}$  and  $\gamma_{\text{dic}}$  are calibrated by applying equations 14-17 and 19. Results of this calibration on the individual CMIP6 models is shown in Figures B1 and B2.

### 3.4.4 Ocean acidification

In this version of Pathfinder,  $\kappa_{\text{pH}}$  is a structural parameter set to 1.

### 3.4.5 Land carbon

Parameters related to the passive soil carbon pool are taken from He et al. (2016, Table S5):  $\nu_{\text{rh3}}$  is structural, while  $\alpha_{\text{pass}}$  is not. All the prior [distribution of the](#) parameters related to the preindustrial steady-state of the land carbon (i.e.  $F_{\text{npp0}}$ ,  $\nu_{\text{fire}}$ ,  $\nu_{\text{harv}}$ ,  $\nu_{\text{mort}}$ ,  $\nu_{\text{stab}}$ ,  $\nu_{\text{rh1}}$  and  $\nu_{\text{cs}}$ ) are derived from 11 TRENDYv7 models (Sitch et al., 2015; Le Quéré et al., 2018), exactly as for OSCAR v3.1 (Gasser et al., 2020) except that all biomes and regions are lumped together. The prior [distribution of the](#) remaining parameters are derived from 12 CMIP6 models that contributed to C4MIP (Arora et al., 2020). Using the models' outputs for the *IpctCO2*, *IpctCO2-rad* and *IpctCO2-bgc* experiments, we calibrated  $\beta_{\text{npp}}$ ,  $\alpha_{\text{npp}}$  and  $\gamma_{\text{npp}}$  through equation 24,

$\beta_{\text{fire}}$  and  $\gamma_{\text{fire}}$  through equation 28, and  $\beta_{\text{rh}}$  and  $\gamma_{\text{rh}}$  through equation 37. Results of this calibration on the individual CMIP6 models is shown in Figure B3, B4 and B5.

### 3.4.6 Permafrost carbon

The permafrost module's ~~prior and structural~~ parameters are recalibrated using the same algorithm as used by Gasser et al. (2018), but adapted to the global formulation of Pathfinder. First, the algorithm is run once to obtain a set of parameters reproducing the behavior of the ~~multi-model~~ global average of five permafrost models (with data from UVic (MacDougall, 2021) added to the four original models). This gives the values of the structural parameters (i.e.  $\alpha_{\text{lst}}$ ,  $\gamma_{\text{rt1}}$ ,  $\gamma_{\text{rt2}}$ ,  $\kappa_{\text{rt}}$ ,  $a_{\text{min}}$ , all  $\alpha_{\text{th},j}$ , all  $\tau_{\text{th},j}$ ,  $\nu_{\text{thaw}}$  and  $\nu_{\text{froz}}$ ). Second, the algorithm is run five additional times, for each of the five permafrost models separately, with the structural parameters established in the first step, to obtain prior distributions ~~offer~~ for the remaining parameters (i.e.  $C_{\text{fr0}}$ ,  $\kappa_a$ ,  $\gamma_a$  and  $\kappa_{\tau_{\text{th}}}$ ).

### 3.4.7 Atmospheric CO2

The conversion factor  $\alpha_C$  is a structural parameter whose value is taken from the latest GCBs (e.g. Le Quéré et al., 2018). The prior distribution of preindustrial CO2 concentration ( $C_{\text{pi}}$ ) is taken from the AR6 (Gulev et al., 2021, Section 2.2.3.2.1), assuming the difference between minimum and maximum over the 0–1850 period is representative of the 90% uncertainty range.

### 3.4.8 Historical CO2 and GMST

The structural  $X_\mu$  and  $X_\sigma$  time series are taken from the latest observations, as follows.  $T_\mu$  and  $T_\sigma$  are taken as the average and standard deviation of 5 observational GMST data sets: HadCRUT5 (Morice et al., 2021), Berkeley Earth (Rohde et al., 2013; Rohde, 2013), GISTEMP (Hansen et al., 2010), NOAAGlobalTemp (Huang et al., 2020), and JMA. We use the 1850–1900 period to define our preindustrial baseline, and GMST ~~change~~ is assumed to be zero before the earliest date available in each data set. Regarding atmospheric CO2,  $C_\mu$  is taken as the global value reported by NOAA/ESRL (Tans and Keeling, 2010) and  $C_\sigma$  as a constant  $\pm 1$  ppm uncertainty, for 1980 onward (this uncertainty is arbitrarily taken higher than the actual uncertainty estimated through instrumental measures to ~~increase have more degrees of~~ freedom in the calibration). Before that ~~period~~,  $C_\mu$  comes from the IPCC AR6 (Dentener et al., 2021, Table AIII.1a), and  $C_\sigma$  is linearly interpolated backwards from the instrumental uncertainty in 1980 to the preindustrial one (Gulev et al., 2021) in 1750. Finally, the prior distribution of  $\rho_X$  is set to Uniform over  $[0,1]$ , that of  $\tilde{\sigma}_X$  is a unit Normal distribution, and that of  $\epsilon_X$  is set arbitrarily to a Half-Normal of parameter 0.05 K for GMST and 0.5 ppm for CO2.

## 3.5 Results (posterior distributions)

The following subsections discuss the adjustments between the prior and posterior parameters that are the results of the Bayesian calibration, as well as the matching of the constraints. These sections constantly refer to Figure 5 that shows the

prior and posterior distributions of the model’s parameters, Figure 6 that shows those of the constrained variables, and Figure 7 that displays the correlation matrix of the posterior parameters. (There is no correlation among the prior parameters.) Prior and posterior values of the parameters can also be retrieved from Table B6.

### 3.5.1 Climate system

455 Our climate-related constraints lead to adjusting all the parameters of the climate module. As explained in Section 3.4.8, the constraints for present-day GMST change and its derivative are met by construction.

The ECS ( $T_{2\times}$ ) is the parameter with the strongest adjustment, since it is directly constrained. Its precise value is discussed hereafter in Section 4.2, but we note that it is unsurprisingly decreased, as the CMIP6 model ensemble tends to overestimate the ECS compared to the IPCC assessed value. Consequently, our posterior  $\text{logit}(\text{ff})$  matches well the constraint. The adjustment  
460 of the ECS significantly reduces the gap between our posterior distribution of the non-CO2 ERF and its constraint, although the posterior central value remains 41% lower (but well within uncertainty range).

Among the dynamic parameters that are adjusted, we note that the deep ocean heat capacity  $\Theta_d$  is somewhat increased compared to the prior, and the heat exchange coefficient  $\theta$  is also increased. These dynamic parameters are likely adjusted through our OHU constraint that is corrected in the posterior so the difference in the central values is lowered from 22% to  
465 14%, which remains well within the uncertainty range.

In addition, a number of weak but physically meaningful correlations across the climate module’s parameters are found, such as a positive correlation between  $T_{2\times}$  and  $\epsilon_{\text{heat}}$  (see e.g. Geoffroy et al., 2013a), a positive correlation between  $T_{2\times}$  and  $\Theta_d$  (that tends to exclude configuration that would warm fast and high), and a negative correlation between  $T_{2\times}$  and  $\phi$  (to match the GMST and ERF constraints together).

### 470 3.5.2 Carbon cycle

Similarly to GMST, the posterior distribution of atmospheric CO2 concentration matches the constraint by construction. Its derivative, however, is (slightly overly) corrected to match the GCB estimate. Global anthropogenic CO2 emissions are significantly increased to get closer to the GCB constraint, but [their central value](#) remains 9% too low. Since these emissions are determined through mass balance and the atmospheric CO2 matches observations, this implies that the total carbon sinks (i.e.  
475  $F_{\text{land}} + F_{\text{ocean}}$ ) must be weaker.

This is confirmed for the ocean sink, as the posterior [central value of](#)  $F_{\text{ocean}}$  is 8% lower than the constraint, but still noticeably corrected if compared to the prior. This correction is explained by small adjustments in some parameters of the ocean carbon module. The mixed layer depth is slightly increased through  $\beta_{\text{dic}}$ . All other parameters remain mostly unaffected by the calibration, and only minor correlations are found. These results, along with the fact that our prior distribution spans  
480 only about half of the constraint’s distribution, suggest that there is a structural limitation in our ocean carbon module that warrants further investigation.

It is also confirmed that the posterior land sink is weaker than the constraint, by 15% [for the central value](#), which is nevertheless a significant reduction of the prior gap of 34%. To explain this adjustment, we observe that the CO2-fertilization

sensitivities ( $\beta_{\text{npp}}$  and  $\gamma_{\text{npp}}$ ) are adjusted upwards. However our constraint on present-day NPP prevents these adjustments  
 485 ~~from being too~~ important, as the posterior distribution of this variable is similar to the prior and ~~its central value~~ remains  
 8–9% higher than its constraint. An increased preindustrial NPP mechanically leads to an increase in preindustrial carbon  
 pools, but these require further adjustments of the land carbon turnover rates, and most notably the mortality rate  $\nu_{\text{mort}}$  and the  
 passive carbon fraction  $\alpha_{\text{pass}}$ , to better match their constraints (of which the one on total soil carbon is perfectly met).

The land carbon module exhibits significant correlations among posterior parameters. This is likely a consequence of all  
 490 the constraints combined as they dictate both the preindustrial steady-state of the module and its transient response over the  
 historical period. Eliminated configurations are those, for instance, that would show high initial carbon pools that are very  
 sensitive to climate change (as these would lead to a very weak land sink), or that would exhibit a weak CO<sub>2</sub>-fertilization effect  
 associated with a fast turnover time (that would also lead to a weak sink).

### 3.5.3 Sea level rise

495 The prior parameters of the SLR module are the least informed of ~~our~~this Bayesian setup. The model initially underestimates  
 the thermal expansion, as well as the GIS and AIS SLR rates. The calibration brings the posterior distributions closer to their  
 respective constraints but it always remain in the lower end of the uncertainty range. The correction is done by adjusting many  
 of the module's parameters (most notably  $\Lambda_{\text{gis1}}$ ,  $\Lambda_{\text{ais, smb}}$ ,  $\lambda_{\text{ais}}$  or  $\lambda_{\text{gla}}$ ), and by finding strong correlations among them (thus  
 excluding physically unrealistic combinations).

500 The historical SLR is markedly corrected by the constraint: from a 19% gap between ~~the central values of~~ the constraint and  
 the prior estimate, to only 7% after calibration. Here, we also note that the ~~sum~~some of individual contributions to historical  
 SLR reported in AR6 do not match that total SLR (Fox-Kemper et al., 2021, Table 9.5), which likely has some impact on the  
 consistency between our constraints. Finally, although the LIA relaxation contribution is not altered by the calibration, ~~as its~~  
 505 ~~central value~~and remains 50% too high, it is the likely source of the strong correlations found among the parameters of this  
 module, because it straightforwardly links the individual SLR contributions together.

## 4 ~~Model diagnosis~~Diagnostics

### 4.1 Historical period

Because in the Bayesian setup we do not use annual time series of observations as constraints, the posterior distributions given  
 in Figure 6 do not inform on the whole dynamic of the model over the historical period. To further diagnose the model's  
 510 behavior, Figure 8 gives the time series from 1900 to 2021 of six key variables. GMST and atmospheric CO<sub>2</sub> match very  
 well the historical observations, by construction of these input time series. The non-CO<sub>2</sub> ERF exhibits a very high variability,  
 owing to our temperature-driven setup and the natural variability in the GMST input. Beyond that, the ERF time series is  
 consistent with the AR6 estimates (Smith et al., 2021), albeit somewhat lower on average in the recent past, as seen with the  
 posterior distribution. Consistently with the interpretation of carbon cycle variables in the calibration results, anthropogenic

515 CO<sub>2</sub> emissions, and the ocean and land carbon sinks are slightly underestimated compared to the GCB estimates (Friedlingstein et al., 2022). Several reasons could explain this discrepancy, from the lack of land use change in Pathfinder to the inconsistency of the GCB figures (that do not close the budget, while ours do). Nevertheless, the interest of the calibration is clearly illustrated, as the posterior uncertainty range covers observations much better than the prior one.

## 4.2 Idealized simulations

520 To complete the diagnosis of our model with common metrics used with climate and carbon models, we ran a set of standard idealized experiments, corresponding to the CMIP6 *abrupt-2xCO<sub>2</sub>*, *1pctCO<sub>2</sub>*, *1pctCO<sub>2</sub>-bgc* and *1pctCO<sub>2</sub>-rad*. A summary of these metrics' values is given in Table 2, and the resulting time series are shown in Figure 9.

The *abrupt-2xCO<sub>2</sub>* experiment sees an abrupt doubling of atmospheric CO<sub>2</sub>, and it is used to diagnose the model's ECS that is defined as the equilibrium temperature for a doubling of the preindustrial atmospheric concentration of CO<sub>2</sub> (we acknowl-  
525 edge that it is superfluous with this version of Pathfinder since it is also a parameter). Using the GMST anomaly at the end of 1500 years of this experiment leads to an unconstrained estimate of ECS of  $4.1 \pm 1.3$  K and a constrained estimate of  $3.3 \pm 0.7$  K. Consistently, the latter value is between the ECS value extracted from CMIP6 models (Meehl et al., 2020) that is higher ( $3.7 \pm 1.1$  K) and the final value assessed in the AR6 that is lower (3.0 K, with a 67% confidence interval between 2.5 and 4.0 K).

530 Using the *1pctCO<sub>2</sub>* experiment that sees a 1% yearly increase in atmospheric CO<sub>2</sub>, we can estimate the model's transient climate response (TCR) that is defined as the GMST change after 70 years, when atmospheric concentration CO<sub>2</sub> has just doubled. The CMIP6 models have a TCR of  $2.0 \pm 0.4$  K (Meehl et al., 2020). Pathfinder's unconstrained value is higher, at  $2.2 \pm 0.5$  K, while the constrained one goes down to  $1.9 \pm 0.3$  K. If we divide the TCR by the cumulative anthropogenic CO<sub>2</sub> emissions compatible with the atmospheric CO<sub>2</sub> increase in this experiment, we obtain an estimate of the transient climate  
535 response to emissions (TCRE). Similarly to the TCR, it is higher in the unconstrained ensemble and lower in the constrained one, when compared to CMIP6 models (Arora et al., 2020). Both downward adjustments of the TCR and TCRE are consistent with that of ECS, with the posterior TCRE matching very well the AR6 assessed range (Canadell et al., 2021).

To look more closely at the carbon cycle, we perform two variants of the latter experiment: in *1pctCO<sub>2</sub>-rad*, atmospheric CO<sub>2</sub> only has a radiative effect, as it is kept at preindustrial level for the carbon cycle; whereas in *1pctCO<sub>2</sub>-bgc*, atmospheric  
540 CO<sub>2</sub> only has a biogeochemical effect, as the climate system sees only preindustrial level. These three experiments are used to calculate the carbon-concentration ( $\beta$ ) and carbon-climate ( $\gamma$ ) feedback metrics that measure the carbon sinks' sensitivities to changes in atmospheric CO<sub>2</sub> and GMST, respectively. We apply the same method as Arora et al. (2020) to calculate these, which leads to metrics at the time of CO<sub>2</sub> doubling that are in line with CMIP6 models (Arora et al., 2020). As both carbon sinks were adjusted upwards by the Bayesian calibration, the constraints logically increased both  $\beta_{\text{ocean}}$  and  $\beta_{\text{land}}$ , to values  
545 fairly close to those of the complex models. The  $\gamma_{\text{ocean}}$  is not affected by the calibration, and remains 45% too low, which again suggests a structural limitation in our formulation of the ocean sink. This is however compensated during calibration by the  $\gamma_{\text{land}}$  being 26% higher than in complex models.

### 4.3 Scenarios

To further validate Pathfinder, we run the five SSP scenarios (Riahi et al., 2017) for which climate and carbon cycle projections were reported by a large-enough number of models in the AR6 (namely, *ssp119*, *ssp126*, *ssp245*, *ssp370* and *ssp585*). These simulations are run with prescribed CO<sub>2</sub> concentration and non-CO<sub>2</sub> ERF (the latter is taken from Smith et al. (2021)). Time series of GMST and cumulative land and ocean sinks are shown on Figure 9. Table 3 shows a comparison of the projected changes in GMST to the CMIP6 estimates (Lee et al., 2021, Table 4.2), and of carbon pools to Liddicoat et al. (2021) (since this was not directly reported in the AR6).

Be it on short-, mid- or long-term, Pathfinder’s projections of GMST are very much in line with the one assessed by the IPCC in the AR6 based on multiple lines of evidence (Lee et al., 2021, Table 4.5). The only significant difference is a smaller uncertainty range in our projections for the longer-term periods. Although this is the result of the efficiency of the Bayesian calibration, one might wonder whether the climate module is over-constrained (or equivalently, too limited in its number of parameters).

The ocean carbon storage appears overestimated by 5% to 20% by Pathfinder across SSP scenarios. This is consistent with the upward adjustment of the ocean carbon sink stemming from our Bayesian calibration. To compare the land carbon storage with CMIP6 models, because our land carbon module does not include land use change processes, we correct the value reported by complex models by the cumulative land use change emissions of each scenario (Riahi et al., 2017; Gidden et al., 2019). While the land carbon storage of Pathfinder is well in line under *ssp126* (a scenario consistent with the 2 °C target), it is underestimated in *ssp119* (consistent with the 1.5 °C target), and increasingly overestimated in higher warming scenarios. A likely explanation is that the climate-carbon feedback on land is underestimated in Pathfinder, as suggested by the  $\gamma$  metric seen in Section 4.2. Alternatively (or concurrently), the absence of loss of sink capacity caused by land cover change (Gasser and Ciais, 2013; Gasser et al., 2020) can explain the overestimation of the land sink under high CO<sub>2</sub>. The Pathfinder model’s estimates of both sinks remain nonetheless well within the CMIP6 models’ uncertainty ranges.

Our SLR emulator gives estimates (Table 4) that are always on the lower of the range reported in the AR6 (Fox-Kemper et al., 2021, Table 9.9). This can be explained by the fact that our individual SLR rate estimates are on the lower end of their respective constraints (see Section 3.5.3). This discrepancy also highlights potential structural limitations in the SLR module (e.g. too few separate contributions), and the difficulty of calibrating the module given the short time period of data available, both from complex models (that cover the 21st century only) and observations, compared to the long time scale of the SLR processes. Nevertheless, our estimates remain within uncertainty range of the IPCC assessment, especially as we do not account for contribution from land water storage that causes an additional 0.03 [0.01, 0.04] m of SLR in all scenarios in 2100 (Fox-Kemper et al., 2021).

## 5 Concluding remarks

In this paper, we have presented the Pathfinder model: a simple global carbon-climate model with selected impact variables, carefully designed to balance accuracy of representation and simplicity of formulation, and calibrated through Bayesian in-

ference on the latest data from Earth system models and observations. Pathfinder has been shown to perform very well in comparison to complex models, although there remains room for further improvement of the model and its calibration setup. We identify four main avenues to improve the model.

585 First, some parts of the model may well lean too much on the complexity side of the simplicity–accuracy balance we aimed to strike, owing to the creation process of Pathfinder that mostly compiled existing formulations. Future development should therefore strive to reduce complexity wherever possible. The ocean carbon sub-pools and perhaps the land carbon pools are potential leads in this respect.

Second~~First~~, the ocean carbon module alone appears to be limited by its structure inherited from a 25-year-old (yet seminal) article (Joos et al., 1996). Although it is undoubtedly a significant undertaking, developing an alternative formulation of the 590 ocean carbon dynamic, calibrated on state-of-the-art ocean models and properly connected to ocean pH and~~interacting with~~ the ocean of the climate module, would benefit more than just the SCM community.

Third~~Second~~, integration of land use and land cover change in such a model appears warranted, despite the difficulty of doing so in a physically sensible yet simple manner. Given our expertise with the OSCAR model and its bookkeeping module (Gasser et al., 2020), we are confident that this can be done, although it will demand extra care to keep the model compatible 595 with the IAMs it is also meant to be linked to.

Fourth~~Third~~, the Bayesian setup can be extended, notably by including more time periods for the existing constraints, but also by introducing and constraining entirely new variables such as isotopic ratios (Hellevang and Aagaard, 2015) or inter-hemispheric gradients (Ciais et al., 2019); although a balance must be struck~~stricken~~ with respect to the calibration’s computation time. Here, we caution against including complex models’ results as constraints in the Bayesian calibration, as 600 was done for the IPCC AR6 (Smith et al., 2021; Nicholls et al., 2021), as it goes against the philosophy of Pathfinder that is to use complex models’ results as prior information ~~only and only real-world observations (or assessment combining many lines of evidence) as constraints.~~

Fifth~~Fourth~~, although our model is restricted to CO<sub>2</sub> by design because of how IAMs like DICE (Nordhaus, 2017) are also limited to CO<sub>2</sub> emissions, we can imagine many reasons why one would want to add non-CO<sub>2</sub> climate forcers into Pathfinder. 605 We would suggest doing so by following the model’s philosophy: that is, by taking existing reduced-complexity formulations such as something between FaIR (Leach et al., 2020) and OSCAR (Gasser et al., 2017), and adding the new parameters into the Bayesian setup with the relevant observational constraints.

In spite of these few shortcomings and potential development leads, Pathfinder v1.0.1 is a powerful tool that fits perfectly the niche it has been created for. We will further demonstrate the strengths and flexibility of Pathfinder in other publications. 610 Meanwhile, we invite the community to seize this open source model, and use it in any study that could benefit from a simple, fast and accurate carbon-climate model, aligned with the latest climate science.

## References

- Japan Meteorological Agency, Global Average Surface Temperature Anomalies, [https://ds.data.jma.go.jp/tcc/tcc/products/gwp/temp/ann\\_wld.html](https://ds.data.jma.go.jp/tcc/tcc/products/gwp/temp/ann_wld.html), accessed: 2022-07-28.
- 615 Armour, K. C.: Energy budget constraints on climate sensitivity in light of inconstant climate feedbacks, *Nature Climate Change*, 7, 331–335, 2017.
- Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P., et al.: Carbon–concentration and carbon–climate feedbacks in CMIP6 models and their comparison to CMIP5 models, *Biogeosciences*, 17, 4173–4222, 2020.
- 620 Bayes, T.: LII. An essay towards solving a problem in the doctrine of chances. By the late Rev. Mr. Bayes, FRS communicated by Mr. Price, in a letter to John Canton, AMFR S, *Philosophical transactions of the Royal Society of London*, pp. 370–418, 1763.
- Bernie, D., Lowe, J., Tyrrell, T., and Legge, O.: Influence of mitigation policy on ocean acidification, *Geophysical Research Letters*, 37, 1–5, <https://doi.org/10.1029/2010GL043181>, 2010.
- Blei, D. M., Kucukelbir, A., and McAuliffe, J. D.: Variational inference: A review for statisticians, *Journal of the American statistical Association*, 112, 859–877, 2017.
- 625 Bossy, T., Gasser, T., and Ciais, P.: Pathfinder v1. 0: a Bayesian-inferred simple carbon-climate model to explore climate change scenarios, *EGUsphere*, pp. 1–52, 2022.
- Canadell, J. G., Monteiro, P. M. S., Costa, M. H., Cotrim da Cunha, L., Cox, P. M., Eliseev, A. V., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P. K., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global Carbon and other Biogeochemical
- 630 Cycles and Feedbacks, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 2021.
- Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., et al.: Sea level change, Tech. rep., PM Cambridge University Press, 2013.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R., Piao, S., and Thornton, P.: AR5-Working Group 1, Chapter 6: Carbon and Other Biogeochemical Cycles 6 - Contribution
- 635 of Working Group I, *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, 2013.
- Ciais, P., Tan, J., Wang, X., Roedenbeck, C., Chevallier, F., Piao, S.-L., Moriarty, R., Broquet, G., Le Quéré, C., Canadell, J., et al.: Five decades of northern land carbon uptake revealed by the interhemispheric CO<sub>2</sub> gradient, *Nature*, 568, 221–225, 2019.
- 640 Cowtan, K. and Way, R. G.: Coverage bias in the HadCRUT4 temperature series and its impact on recent temperature trends, *Quarterly Journal of the Royal Meteorological Society*, 140, 1935–1944, 2014.
- Dentener, F. J., Hall, B., and Smith, C.: Annex III: Tables of historical and projected well-mixed greenhouse gas mixing ratios and effective radiative forcing of all climate forcers, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 2021.
- 645 Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E., Smith, C. J., et al.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74–82, 2021.
- Fornberg, B.: Generation of finite difference formulas on arbitrarily spaced grids, *Mathematics of computation*, 51, 699–706, 1988.

- Forster, P., Storelvmo, T., Armour, K., Collins, T., Dufresne, J. L., Frame, D., Lunt, D. J., Mauritsen, T., Palmer, M. D., Watanabe, M., Wild, M., and Zhang, H.: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity, *Climate Change 2021: The Physical Science Basis*. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, 2021.
- 650 Fox-Kemper, B., Hewitt, H. T., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S. S., , Edwards, T. L., Golledge, N. R., Hemer, M., Kopp, R. E., Krinner, S. S., Mix, A., Notz, D., Nowicki, S., Nurhati, I. S., Ruiz, L., Sallée, J.-B., and Slangen, A. B. A., a. Y. Y.: Ocean, Cryosphere and Sea Level Change, *Climate Change 2021: The Physical Science Basis*. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, 2021.
- 655 Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Bakker, D. C., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., et al.: Global carbon budget 2021, *Earth System Science Data*, 14, 1917–2005, 2022.
- Gasser, T.: tgasser/Pathfinder: v1.0, <https://doi.org/10.5281/zenodo.7003849>, 2022.
- Gasser, T. and Ciais, P.: A theoretical framework for the net land-to-atmosphere CO<sub>2</sub> flux and its implications in the definition of "emissions from land-use change", *Earth System Dynamics*, 4, 171–186, <https://doi.org/10.5194/esd-4-171-2013>, 2013.
- 660 Gasser, T., Ciais, P., Boucher, O., Quilcaille, Y., Tortora, M., Bopp, L., and Hauglustaine, D.: The compact Earth system model OSCAR v2.2: Description and first results, *Geoscientific Model Development*, 10, 271–319, <https://doi.org/10.5194/gmd-10-271-2017>, 2017.
- Gasser, T., Kechiar, M., Ciais, P., Burke, E. J., Kleinen, T., Zhu, D., Huang, Y., Ekici, A., and Obersteiner, M.: Path-dependent reductions in CO<sub>2</sub> emission budgets caused by permafrost carbon release, *Nature Geoscience*, 11, 830–835, <https://doi.org/10.1038/s41561-018-0227-0>, 2018.
- 665 Gasser, T., Crepin, L., Quilcaille, Y., Houghton, R. A., Ciais, P., and Obersteiner, M.: Historical CO<sub>2</sub> emissions from land use and land cover change and their uncertainty, *Biogeosciences*, 17, 4075–4101, 2020.
- Geoffroy, O., Saint-Martin, D., Bellon, G., Voldoire, A., Olivie, D. J., and Tyteca, S.: Transient climate response in a two-layer energy-balance model. Part II: Representation of the efficacy of deep-ocean heat uptake and validation for CMIP5 AOGCMs, *Journal of Climate*, 26, 1859–1876, <https://doi.org/10.1175/JCLI-D-12-00196.1>, 2013a.
- 670 Geoffroy, O., Saint-Martin, D., Olivie, D. J. L., Voldoire, A., Bellon, G., and Tyteca, S.: Transient Climate Response in a Two-Layer Energy-Balance Model. Part I: Analytical Solution and Parameter Calibration Using CMIP5 AOGCM Experiments, *Journal of Climate*, 26, 1841–1857, <https://doi.org/10.1175/JCLI-D-12-00195.1>, 2013b.
- Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., Van Vuuren, D. P., Van Den Berg, M., Feng, L., Klein, D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O., Harmsen, M., Hasegawa, T., Havlik, P., Hilaire, J., Hoesly, R., Horing, J., Popp, A., Stehfest, E., and Takahashi, K.: Global emissions pathways under different socioeconomic scenarios for use in CMIP6: A dataset of harmonized emissions trajectories through the end of the century, *Geoscientific Model Development*, 12, 1443–1475, <https://doi.org/10.5194/gmd-12-1443-2019>, 2019.
- 675 Gitz, V. and Ciais, P.: Amplifying effects of land-use change on future atmospheric CO<sub>2</sub> levels, *Global Biogeochemical Cycles*, 17, 2003.
- Goodwin, P., Haigh, I. D., Rohling, E. J., and Slangen, A.: A new approach to projecting 21st century sea-level changes and extremes, *Earth's Future*, 5, 240–253, <https://doi.org/10.1002/2016EF000508>, 2017.
- 680 Gulev, S. K., Thorne, P. W., Ahn, J., Dentener, F. J., Domingues, C. M., Gerland, S., Gong, D., Kaufman, D. S., Nnamchi, H. C., Quaas, J., Rivera, J. A., Sathyendranath, S., Smith, S. L., Trewin, B., von Shuckmann, K., and Vose, R.: Changing State of the Climate System, *Climate Change 2021: The Physical Science Basis*. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, 2021.

- 685 Hansen, J., Ruedy, R., Sato, M., and Lo, K.: Global Surface Temperature Change, *Review of Geophysics*, 5, 1–29,  
<https://doi.org/10.1029/2010RG000345.1>.INTRODUCTION, 2010.
- Harris, C. R., Millman, K. J., Van Der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J.,  
 et al.: Array programming with NumPy, *Nature*, 585, 357–362, 2020.
- Hartin, C. A., Patel, P., Schwarber, A., Link, R. P., and Bond-Lamberty, B.: A simple object-oriented and open-source model for scientific  
 690 and policy analyses of the global climate system–Hector v1. 0, *Geoscientific Model Development*, 8, 939–955, 2015.
- He, Y., Trumbore, S. E., Torn, M. S., Harden, J. W., Vaughn, L. J., Allison, S. D., and Randerson, J. T.: Radiocarbon constraints imply  
 reduced carbon uptake by soils during the 21st century, *Science*, 353, 1419–1424, 2016.
- Hellevang, H. and Aagaard, P.: Constraints on natural global atmospheric CO<sub>2</sub> fluxes from 1860 to 2010 using a simplified explicit forward  
 model, *Scientific reports*, 5, 1–12, 2015.
- 695 Hoyer, S. and Hamman, J.: xarray: N-D labeled arrays and datasets in Python, *Journal of Open Research Software*, 5,  
<https://doi.org/10.5334/jors.148>, 2017.
- Huang, B., Menne, M. J., Boyer, T., Freeman, E., Gleason, B. E., Lawrimore, J. H., Liu, C., Rennie, J. J., Schreck III, C. J., Sun, F., et al.:  
 Uncertainty estimates for sea surface temperature and land surface air temperature in NOAA GlobalTemp version 5, *Journal of Climate*,  
 33, 1351–1379, 2020.
- 700 Joos, F., Bruno, M., Fink, R., Siegenthaler, U., Stocker, T. F., Le Quéré, C., and Sarmiento, J. L.: An Efficient and accurate representation of  
 complex oceanic and biospheric models of anthropogenic carbon uptake, 1996.
- Joos, F., Prentice, I. C., Sitch, S., Meyer, R., Hooss, G., Plattner, G.-K., Gerber, S., and Hasselmann, K.: Global warming feedbacks on ter-  
 restrial carbon uptake under the Intergovernmental Panel on Climate Change (IPCC) emission scenarios, *Global Biogeochemical Cycles*,  
 15, 891–907, 2001.
- 705 Kucukelbir, A., Tran, D., Ranganath, R., Gelman, A., and Blei, D. M.: Automatic differentiation variational inference, *Journal of machine  
 learning research*, 2017.
- Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell,  
 J. G., Arneeth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., and Zheng, B.: Global Carbon Budget 2018, *Earth System Science Data*,  
 pp. 2141–2194, <https://doi.org/10.5194/essd-8-605-2016>, 2018.
- 710 Leach, N. J., Nicholls, Z., Jenkins, S., Smith, C. J., Lynch, J., Cain, M., Wu, B., Tsutsui, J., and Allen, M. R.: GIR v1. 0.0: a generalised  
 impulse-response model for climate uncertainty and future scenario exploration, *Geoscientific Model Development Discussions*, pp. 1–29,  
 2020.
- Lee, J. Y., Marotzke, J., Bala, G., Cao, L., Corti, S., Dunne, J. P., Engelbrecht, F., Fischer, E., Fyfe, J. C., Jones, C., Maycock, A., Mutemi, J.,  
 Ndiaye, O., Panickal, S., and Zhou, T.: Future Global Climate: Scenario-Based Projections and Near-Term Informations, *Climate Change*  
 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on  
 715 Climate Change, 2021.
- Liddicoat, S. K., Wiltshire, A. J., Jones, C. D., Arora, V. K., Brovkin, V., Cadule, P., Hajima, T., Lawrence, D. M., Pongratz, J., Schwinger,  
 J., et al.: Compatible Fossil Fuel CO<sub>2</sub> Emissions in the CMIP6 Earth System Models’ Historical and Shared Socioeconomic Pathway  
 Experiments of the Twenty-First Century, *Journal of Climate*, 34, 2853–2875, 2021.
- 720 MacDougall, A. H.: Estimated effect of the permafrost carbon feedback on the zero emissions commitment to climate change, *Biogeo-  
 sciences*, 18, 4937–4952, 2021.

- Meehl, G. A., Senior, C. A., Eyring, V., Flato, G., Lamarque, J.-F., Stouffer, R. J., Taylor, K. E., and Schlund, M.: Context for interpreting equilibrium climate sensitivity and transient climate response from the CMIP6 Earth system models, *Science Advances*, 6, eaba1981, 2020.
- 725 Meinshausen, M., Raper, S. C., and Wigley, T. M.: Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, *MAGICC6–Part 1: Model description and calibration*, *Atmospheric Chemistry and Physics*, 11, 1417–1456, 2011.
- Mengel, M., Levermann, A., Frieler, K., Robinson, A., Marzeion, B., and Winkelmann, R.: Future sea level rise constrained by observations and long-term commitment, *Proceedings of the National Academy of Sciences*, 113, 2597–2602, 2016.
- Morice, C. P., Kennedy, J. J., Rayner, N. A., Winn, J., Hogan, E., Killick, R., Dunn, R., Osborn, T., Jones, P., and Simpson, I.: An updated  
730 assessment of near-surface temperature change from 1850: The HadCRUT5 data set, *Journal of Geophysical Research: Atmospheres*, 126, e2019JD032361, 2021.
- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: AR5 - Working Group 1, Chapter 8 : Anthropogenic and Natural Radiative Forcing - Contribution of Working Group I, Cambridge University Press, 23, 56, 2013.
- 735 National Academies of Sciences, E. and Medicine: Valuing climate damages: updating estimation of the social cost of carbon dioxide, National Academies Press, 2017.
- Nicholls, Z., Meinshausen, M., Lewis, J., Corradi, M. R., Dorheim, K., Gasser, T., Gieseke, R., Hope, A. P., Leach, N., McBride, L. A., et al.: Reduced complexity Model Intercomparison Project Phase 2: Synthesizing Earth system knowledge for probabilistic climate projections, *Earth's Future*, 9, e2020EF001900, 2021.
- 740 Nicholls, Z. R., Meinshausen, M., Lewis, J., Gieseke, R., Dommenges, D., Dorheim, K., Fan, C.-S., Fuglestad, J. S., Gasser, T., Golücke, U., et al.: Reduced Complexity Model Intercomparison Project Phase 1: introduction and evaluation of global-mean temperature response, *Geoscientific Model Development*, 13, 5175–5190, 2020.
- Nordhaus, W. D.: Revisiting the social cost of carbon, *Proceedings of the National Academy of Sciences*, 114, 1518–1523, <https://doi.org/10.1073/pnas.1609244114>, 2017.
- 745 Parkes, D. and Marzeion, B.: Twentieth-century contribution to sea-level rise from uncharted glaciers, *Nature*, 563, 551–554, 2018.
- Pérez, F. and Granger, B. E.: IPython: a System for Interactive Scientific Computing, *Computing in Science and Engineering*, 9, 21–29, <https://doi.org/10.1109/MCSE.2007.53>, 2007.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P.,  
750 Humenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Streffer, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environmental Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- 755 Ricciuto, D. M., Davis, K. J., and Keller, K.: A Bayesian calibration of a simple carbon cycle model: The role of observations in estimating and reducing uncertainty, *Global biogeochemical cycles*, 22, 2008.
- Roe, G. H. and Baker, M. B.: Why is climate sensitivity so unpredictable?, *Science*, 318, 629–632, 2007.
- Rohde, R.: Comparison of Berkeley Earth, NASA GISS, and Hadley CRU averaging techniques on ideal synthetic data, *Berkeley Earth Memo*, January, 2, 013, 2013.


- 760 Rohde, R., Muller, R., Jacobsen, R., Perlmutter, S., Rosenfeld, A., Wurtele, J., Curry, J., Wickham, C., and Mosher, S.: Berkeley Earth temperature averaging process, *Geoinformatics & Geostatistics: An Overview*, 1, 1–13, 2013.
- Salvatier, J., Wiecki, T. V., and Fonnesbeck, C.: Probabilistic programming in Python using PyMC3, *PeerJ Computer Science*, 2, e55, 2016.
- Sitch, S., Friedlingstein, P., Gruber, N., Jones, S. D., Murray-Tortarolo, G., Ahlström, A., Doney, S. C., Graven, H., Heinze, C., Huntingford, C., et al.: Recent trends and drivers of regional sources and sinks of carbon dioxide, *Biogeosciences*, 12, 653–679, 2015.
- 765 Slangen, A., Church, J. A., Agosta, C., Fettweis, X., Marzeion, B., and Richter, K.: Anthropogenic forcing dominates global mean sea-level rise since 1970, *Nature Climate Change*, 6, 701–705, 2016.
- Smith, C., Nicholls, Z. R. J., Armour, K., Collins, W., Forster, P., M. M., Palmer, M. D., and Watanabe, M.: The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity Supplementary Material, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 2021.
- 770 Smith, C. J., Forster, P. M., Allen, M., Leach, N., Millar, R. J., Passerello, G. A., and Regayre, L. A.: FAIR v1. 3: a simple emissions-based impulse response and carbon cycle model, *Geoscientific Model Development*, 11, 2273–2297, 2018.
- Strassmann, K. M. and Joos, F.: The Bern Simple Climate Model (BernSCM) v1.0: An extensible and fully documented open-source re-implementation of the Bern reduced-form model for global carbon cycle-climate simulations, *Geoscientific Model Development*, 11, 1887–1908, <https://doi.org/10.5194/gmd-11-1887-2018>, 2018.
- 775 Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., and Sutherland, S.: Seasonal variation of CO<sub>2</sub> and nutrients in the high-latitude surface oceans: A comparative study, *Global Biogeochemical Cycles*, 7, 843–878, 1993.
- Tans, P. and Keeling, R.: NOAA, ESRL ([www.esrl.noaa.gov/gmd/ccgg/trends/](http://www.esrl.noaa.gov/gmd/ccgg/trends/)), 2010.
- Theano Development Team: Theano: A Python framework for fast computation of mathematical expressions, *arXiv e-prints*, abs/1605.02688, <http://arxiv.org/abs/1605.02688>, 2016.
- 780 Van Rossum, G. and Drake, F. L.: *Python 3 Reference Manual*, CreateSpace, Scotts Valley, CA, 2009.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors: SciPy 1.0: Fundamental Algorithms for Scientific
- 785 Computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- Vose, R. S., Arndt, D., Banzon, V. F., Easterling, D. R., Gleason, B., Huang, B., Kearns, E., Lawrimore, J. H., Menne, M. J., Peterson, T. C., et al.: NOAA’s merged land–ocean surface temperature analysis, *Bulletin of the American Meteorological Society*, 93, 1677–1685, 2012.
- Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophysical research letters*, 37, 2010.
- 790 Zhao, M., Heinsch, F. A., Nemani, R. R., and Running, S. W.: Improvements of the MODIS terrestrial gross and net primary production global data set, *Remote sensing of Environment*, 95, 164–176, 2005.

*Code and data availability.* The source code of Pathfinder is openly available at: <https://github.com/tgasser/Pathfinder>. [A frozen version of the code and data as developed in the paper can be found on Zenodo: https://doi.org/10.5281/zenodo.7003848](https://doi.org/10.5281/zenodo.7003848) (Gasser, 2022).


*Author contributions.* TG developed the model, with contributions from TB on sea level rise and PC on land carbon cycle. TG coded the  
795 model and its calibration. TB ran the diagnostic simulations and made the final figures. TB and TG wrote the manuscript with input from PC.

*Competing interests.* The authors declare they have no competing interests.


*Acknowledgements.* Development of Pathfinder was supported by the European Union's Horizon 2020 research and innovation programme under grant agreement #820829 (CONSTRAIN project), and by the Austrian Science Fund (FWF) under grant agreement P31796-N29 (ERM project). We thank A. H. MacDougall for sharing permafrost simulations made with UVic.




**Figure 1.** Pathfinder in a nutshell: Green blocks represent the carbon cycle, and red blocks the climate response. Blue blocks with dotted arrows are impacts that can be derived with the model. Grey blocks are variables that are directly related to anthropogenic activity. Possible inputs of the model are distinguishable through the bold contours of the blocks. In this scheme, arrows correspond to a forward mode where inputs would be  $E_{CO_2}$  and  $R_x$




**Figure 2.** The ocean sink model in Pathfinder follows the structure of the mixed-layer pulse response function introduced by Joos et al. (1996). The mix-layer is represented through five subpools which each has a different timescale for transport to the deep ocean carbon pool




**Figure 3.** The land sink model in Pathfinder is derived from OSCAR (Gasser et al., 2017) and represents the biosphere as a set of four carbon pools: vegetation, litter, active soil and passive soil. These pools exchange carbon through fluxes whose direction is given by the arrows.



**Figure 4.** The permafrost carbon model in Pathfinder is taken from Gasser et al. (2018). The frozen pool dynamic lags behind a theoretical value that is determined by the temperature anomaly. Thawed carbon is then split between three pools that are emitted to the atmosphere at different rates.




**Figure 5.** Parameters distributions before (black lines) and after (blue lines) the Bayesian calibration. Parameters are noted under their text notation, and Tables B3 and B4 provide the corresponding notation in code.




**Figure 6.** Distributions of the constrained variables. Dashed lines give the distributions used to constrain. Black lines give the distribution before calibration while blue lines give the distribution posterior to calibration. Under a variable's name, we give the period over which the constraint is estimated, and the data processing method: mean over the period, difference between last and first year, or sum of all the years over the period. (1750 is the preindustrial.) Variables are noted under their text notation, and Tables B1 and B2 provide the corresponding notation in code.





**Figure 8.** Historical time series of key variables from Pathfinder. Red lines are observations, black lines are the model's outputs before calibration, and blue lines are the same after calibration. Shaded areas and vertical bars correspond to the  $1\sigma$  uncertainty range. Temperature observations are taken from HadCRUT5 (Morice et al., 2021), Cowtan and Way (2014), Berkeley Earth (Rohde et al., 2013; Rohde, 2013), GISTEMP (Hansen et al., 2010), and NOAA/MLOST (Vose et al., 2012). Other sources are NOAA/ESRL (Tans and Keeling, 2010), GCB 2021 (Friedlingstein et al., 2022), and AR6 (Smith et al., 2021).



**Figure 9.** Time series of GMST change, [integrated](#) land carbon uptake and [integrated](#) ocean carbon uptake for idealized experiments (*abrupt-2xCO2*, *1pctCO2*, *1pctCO2-bgc* and *1pctCO2-rad*), and projections according to SSP scenarios in Pathfinder. Shaded areas give the 1 $\sigma$  uncertainty range.

Variable	Period	Method	Prior	Posterior	Constraints	Unit
$E_{\text{CO}_2}$	2011-2020	Mean	$9.1 \pm 1.3$	$10.0 \pm 0.7$	$11.0 \pm 0.9$	$\text{PgC yr}^{-1}$
$\frac{dC}{dt}$	2011-2020	Mean	$2.41 \pm 0.06$	$2.40 \pm 0.01$	$2.40 \pm 0.01$	$\text{ppm yr}^{-1}$
$F_{\text{land}}$	1960-2020	Sum	$95 \pm 52$	$123 \pm 26$	$145 \pm 35$	PgC
$F_{\text{ocean}}$	1960-2020	Sum	$89 \pm 12$	$97 \pm 13$	$105 \pm 20$	PgC
$C_v$	1750	Mean	$407 \pm 54$	$407 \pm 37$	$450 \pm 50$	PgC
$C_s$	1750	Mean	$1181 \pm 735$	$1086 \pm 284$	$1088 \pm 249$	PgC
$F_{\text{NPP}}$	1998-2002	Mean	$60.0 \pm 7.9$	$59.5 \pm 3.9$	$55.0 \pm 5.0$	$\text{PgC yr}^{-1}$
$C$	2012-2021	Mean	$403.6 \pm 0.3$	$403.6 \pm 0.1$	$401.2 \pm 0.1$	ppm
$R_x$	2010-2019	Mean	$0.01 \pm 0.47$	$0.33 \pm 0.37$	$0.56 \pm 0.53$	$\text{W m}^{-2}$
$T$	2001-2020	Mean	$0.96 \pm 0.08$	$0.97 \pm 0.06$	$1.00 \pm 0.07$	K
$\frac{dT}{dt}$	2000-2019	Mean	$0.028 \pm 0.003$	$0.028 \pm 0.002$	$0.029 \pm 0.002$	$\text{K yr}^{-1}$
$\frac{dU_{\text{ohc}}}{dt}$	2006-2018	Mean	$0.56 \pm 0.10$	$0.62 \pm 0.09$	$0.72 \pm 0.17$	$\text{W m}^{-2}$
$\frac{dH_{\text{thx}}}{dt}$	2006-2018	Mean	$1.02 \pm 0.22$	$1.14 \pm 0.21$	$1.39 \pm 0.40$	$\text{mm yr}^{-1}$
$\frac{dH_{\text{glia}}}{dt}$	2006-2018	Mean	$0.63 \pm 0.24$	$0.62 \pm 0.04$	$0.62 \pm 0.03$	$\text{mm yr}^{-1}$
$\frac{dH_{\text{ais}}}{dt}$	2006-2018	Mean	$-0.02 \pm 0.23$	$0.30 \pm 0.10$	$0.37 \pm 0.08$	$\text{mm yr}^{-1}$
$\frac{dH_{\text{gis}}}{dt}$	2006-2018	Mean	$0.36 \pm 0.12$	$0.57 \pm 0.10$	$0.63 \pm 0.07$	$\text{mm yr}^{-1}$
$H_{\text{tot}}$	1901-1990	Difference	$72 \pm 17$	$83 \pm 10$	$89 \pm 32$	mm
$H_{\text{lia}}$	1750	Mean	$45 \pm 17$	$45 \pm 11$	$30 \pm 13$	mm
logit(ff)	1750	Mean	$1.69 \pm 0.38$	$1.47 \pm 0.28$	$1.38 \pm 0.37$	1

**Table 1.** Constrained variables in Pathfinder, with values before and after calibration. Variables are noted under their text notation, and Tables B1 and B2 provide the corresponding notation in code. The uncertainty correspond to the  $1 \sigma$  uncertainty range.

$2 \times \text{CO}_2$	Pathfinder unconstrained	Pathfinder constrained	CMIP6	AR6
ECS (K)	$4.1 \pm 1.3$	$3.3 \pm 0.7$	$3.7 \pm 1.1$	3.0 (2.0, 4.5)
TCR (K)	$2.2 \pm 0.5$	$1.9 \pm 0.3$	$2.0 \pm 0.4$	1.8 (1.4, 2.2)
TCRE (K EgC $^{-1}$ )	$2.20 \pm 0.63$	$1.65 \pm 0.32$	$1.77 \pm 0.37$	1.65 (1.0, 2.3)
$\beta_{\text{ocean}}$ (PgC ppm $^{-1}$ )	$0.81 \pm 0.10$	$0.87 \pm 0.11$	$0.91 \pm 0.09$	
$\gamma_{\text{ocean}}$ (PgC K $^{-1}$ )	$-12.9 \pm 5.4$	$-12.5 \pm 6.0$	$-8.6 \pm 2.9$	
$\beta_{\text{land}}$ (PgC ppm $^{-1}$ )	$1.05 \pm 0.5$	$1.26 \pm 0.30$	$1.22 \pm 0.40$	
$\gamma_{\text{land}}$ (PgC K $^{-1}$ )	$-33.2 \pm 26.6$	$-25.3 \pm 24.2$	$-34.1 \pm 38.4$	

**Table 2.** Diagnostics of climate and carbon-cycle responses in Pathfinder before and after Bayesian calibration. Comparison with AR6 (Forster et al., 2021; Canadell et al., 2021) and CMIP6 data (Arora et al., 2020; Meehl et al., 2020) is shown. For AR6 data we give the median and the 90% confidence interval while for every other values we give the mean  $\pm 1 \sigma$

Experiment	Model	GMST (K)	GMST (K)	GMST (K)	Ocean Carbon Storage (PgC)	Land Carbon Storage (PgC)
		2021–2040	2041–2060	2081–2100	2015–2100	2015–2100
ssp119	Pathfinder	1.5 (1.3, 1.8)	1.6 (1.4, 1.9)	1.5 (1.2, 1.7)	132 ± 21	49 ± 33
ssp119	AR6 or CMIP6	1.5 (1.2, 1.7)	1.6 (1.2, 2.0)	1.4 (1.0, 1.8)	111 ± 11	73 ± 33
ssp126	Pathfinder	1.5 (1.3, 1.8)	1.8 (1.6, 2.1)	1.9 (1.6, 2.2)	179 ± 28	109 ± 45
ssp126	AR6 or CMIP6	1.5 (1.2, 1.8)	1.7 (1.3, 2.2)	1.8 (1.3, 2.4)	162 ± 8	120 ± 50
ssp245	Pathfinder	1.6 (1.4, 1.8)	2.1 (1.8, 2.4)	2.8 (2.4, 3.3)	265 ± 41	225 ± 76
ssp245	AR6 or CMIP6	1.5 (1.2, 1.8)	2.0 (1.6, 2.5)	2.7 (2.1, 3.5)	252 ± 11	178 ± 76
ssp370	Pathfinder	1.6 (1.4, 1.8)	2.2 (1.9, 2.6)	3.7 (3.2, 4.3)	354 ± 53	330 ± 112
ssp370	AR6 or CMIP6	1.5 (1.2, 1.8)	2.1 (1.7, 2.6)	3.6 (2.8, 4.6)	338 ± 15	269 ± 124
ssp585	Pathfinder	1.7 (1.5, 1.9)	2.5 (2.2, 2.9)	4.4 (3.8, 5.2)	420 ± 63	409 ± 148
ssp585	AR6 or CMIP6	1.6 (1.3, 1.9)	2.4 (1.9, 3.0)	4.4 (3.3, 5.7)	398 ± 17	311 ± 162

**Table 3.** Comparison of SSP scenarios for GMST change projections (w.r.t. 1850–1900) to AR6 (Lee et al., 2021, Table 4.5), and for ocean and land carbon storage projections to CMIP6 (Liddicoat et al., 2021). Land carbon storage projections were corrected using the land use change emissions data from SSPs (Riahi et al., 2017; Gidden et al., 2019). For GMST data we give the median and the 90% confidence interval while for every other values we give the mean  $\pm 1 \sigma$

Experiment	Model	SLR (m)	SLR (m)	SLR rate (mm yr <sup>-1</sup> )	SLR rate (mm yr <sup>-1</sup> )
		2050	2100	2040–2060	2080–2100
ssp119	Pathfinder	0.15 (0.13, 0.18)	0.30 (0.25, 0.36)	3.5 (2.9, 4.2)	2.7 (2.2, 3.4)
ssp119	AR6	0.18 (0.15, 0.23)	0.38 (0.28, 0.55)	4.1 (2.8, 6.0)	4.2 (2.4, 6.6)
ssp126	Pathfinder	0.16 (0.14, 0.19)	0.35 (0.30, 0.43)	4.1 (3.5, 5.0)	3.6 (2.9, 4.5)
ssp126	AR6	0.19 (0.16, 0.25)	0.44 (0.32, 0.62)	4.8 (3.5, 6.8)	5.2 (3.2, 8.0)
ssp245	Pathfinder	0.17 (0.15, 0.20)	0.46 (0.39, 0.56)	5.0 (4.2, 6.0)	6.2 (5.1, 8.0)
ssp245	AR6	0.20 (0.17, 0.26)	0.56 (0.44, 0.76)	5.8 (4.4, 8.0)	7.7 (5.2, 11.6)
ssp370	Pathfinder	0.18 (0.15, 0.21)	0.56 (0.47, 0.69)	5.5 (4.7, 6.7)	9.1 (7.4, 11.7)
ssp370	AR6	0.22 (0.18, 0.27)	0.68 (0.55, 0.90)	6.4 (5.0, 8.7)	10.4 (7.4, 14.8)
ssp585	Pathfinder	0.19 (0.17, 0.23)	0.67 (0.56, 0.82)	6.4 (5.4, 7.8)	11.4 (9.1, 15.0)
ssp585	AR6	0.23 (0.20, 0.29)	0.77 (0.63, 1.01)	7.2 (5.6, 9.7)	12.1 (8.6, 17.6)

**Table 4.** Comparison of SSP scenarios between Pathfinder and AR6 for SLR (w.r.t. 1995–2014) and SLR speed projections (Fox-Kemper et al., 2021, Table 9.9). We give the median value and the 90% confidence interval in parentheses

## 800 **Appendix A: Additional information on the model**

### **A1** [Technical requirements](#)

Pathfinder has been developed and run in Python (v3.7.6) (Van Rossum and Drake, 2009), preferentially using IPython (v7.19.0) (Pérez and Granger, 2007). Currently, packages required to run it are NumPy (v1.19.2) (Harris et al., 2020), SciPy (v1.5.2) (Virtanen et al., 2020) and Xarray (v0.16.0) (Hoyer and Hamman, 2017), and it has hard-coded dependencies on  
805 PyMC3 (v3.8) (Salvatier et al., 2016) and Theano (v1.0.4) (Theano Development Team, 2016) that are in fact used only for calibration. Other versions of Python or these packages were not tested.

The calibration procedure takes about 9 hours to run on a desktop computer (with a base speed of 3.4 GHz). Simple use of the model is much faster: the idealized experiments and SSP scenarios for this description paper, which represent 2984 simulated years, were run in about 20 minutes for all 2000 configurations and on a single core. A single simulated year takes  
810 a few tenth of a second, although a number of options in the model can drastically alter this performance. Note also that this scales sub-linearly with the amount of configurations or scenarios because of the internal workings of the Xarray package, albeit at the cost of increased demand in random-access memory.

### **A2** [Known issues](#)

Two relatively benign issues that have been identified during development remain unsolved. First, the model requires a high  
815 number of sub-time steps (i.e. high  $n_t$ ) to remain stable under high CO<sub>2</sub>, because of the ocean carbon cycle. Second, the version of the model that is driven by  $T$  and  $R_x$  time series is extremely sensitive to its inputs, because mathematically it requires the first two derivatives of  $T$  and the first derivative of  $R_x$ .


### **A3** [Changelog](#)

Brief description of the successive versions of Pathfinder:


820 **v1.0.1.** Exact same physical equations and numerical values as v1.0. Added best-guess parameters calculated as the average of the posterior distribution, and corresponding historical outputs, for single-configuration runs. Improved README and MANUAL files.

**v1.0.** First release. Exact model described in the preprint version of this very paper (Bossy et al., 2022).


## **Appendix B: Additional figures and tables**




**Figure B1.** Calibration to estimate prior  $\nu_{gx}$  and  $\gamma_{gx}$  from CMIP6 time series of  $F_{ocean}$ . We fit our equation on the results of the +1%  $CO_2$  (*IpctCO2*) experiment (in blue) and its variants *IpctCO2-rad* (in green) and *IpctCO2-bgc* (in orange). Coloured markers are CMIP6 models data while the solid black lines show the fit from Pathfinder. Panels without black line indicate that at least one of the required variables was not reported by the complex model.




**Figure B2.** Calibration to estimate prior  $\beta_{\text{dic}}$  and  $\gamma_{\text{dic}}$  from CMIP6 time series of  $p_{\text{CO}_2}$ . We fit our equation on the results of the +1%  $\text{CO}_2$  (*1pctCO2*) experiment (in blue) and its variants *1pctCO2-rad* (in green) and *1pctCO2-bgc* (in orange). Coloured markers are CMIP6 models data while the solid black lines show the fit from Pathfinder. Panels without black line indicate that at least one of the required variables was not reported by the complex model.




**Figure B3.** Calibration to estimate prior  $\beta_{ef}$  and  $\gamma_{ef}$  from CMIP6 time series of  $r_{fire}$ , shown as a ratio over its preindustrial value. We fit our equation on the results of the +1%  $CO_2$  ( $IpctCO_2$ ) experiment (in blue) and its variants  $IpctCO_2-rad$  (in green) and  $IpctCO_2-bgc$  (in orange). Coloured markers are CMIP6 models data while the solid black lines show the fit from Pathfinder. Panels without black line indicate that at least one of the required variables was not reported by the complex model.




**Figure B4.** Calibration to estimate prior  $\beta_{rh}$  and  $\gamma_{rh}$  from CMIP6 time series of  $r_{rh}$ , shown as a ratio over its preindustrial value. We fit our equation on the results of the +1%  $\text{CO}_2$  (*IpctCO2*) experiment (in blue) and its variants *IpctCO2-rad* (in green) and *IpctCO2-bgc* (in orange). Coloured markers are CMIP6 models data while the solid black lines show the fit from Pathfinder. Panels without black line indicate that at least one of the required variables was not reported by the complex model.




**Figure B5.** Calibration to estimate prior  $\beta_{npp}$ ,  $\alpha_{npp}$  and  $\gamma_{npp}$  from CMIP6 time series of  $r_{npp}$ , shown as a ratio over its preindustrial value. We fit our equation on the results of the +1%  $\text{CO}_2$  (*IpctCO2*) experiment (in blue) and its variants *IpctCO2-rad* (in green) and *IpctCO2-bgc* (in orange). Coloured markers are CMIP6 models data while the solid black lines show the fit from Pathfinder. Panels without black line indicate that at least one of the required variables was not reported by the complex model.




**Figure B6.** Calibration to estimate the prior of GIS SLR module parameters ( $\Lambda_{gis1}$ ,  $\Lambda_{gis3}$  and  $\lambda_{gis0}$ ). We fit our equation on the compiled outputs from Edwards et al. (2021) for which there is more than one scenario available. Each panel's title displays the name of the institute, model and configuration used. Coloured markers are the models data while the solid black lines show the fit from Pathfinder.



**Figure B7.** Calibration to estimate the prior of Glaciers SLR module parameters ( $\Gamma_{gla1}$ ,  $\Gamma_{gla3}$ ,  $\gamma_{gla}$  and  $\lambda_{gla0}$ ). We fit our equation on the compiled outputs from Edwards et al. (2021) for which there is more than one scenario available. Each panel's title displays the name of the model or study used. Coloured markers are the models data while the solid black lines show the fit from Pathfinder.



**Figure B8.** Calibration to estimate the prior of AIS SLR module parameters ( $\Lambda_{ais,smb}$ ,  $\alpha_{ais}$  and  $\lambda_{ais0}$ ). We fit our equation on the compiled outputs from Edwards et al. (2021) for which there is more than one scenario available. Each panel's title displays the name of the institute, model and configuration used. Coloured markers are the models data while the solid black lines show the fit from Pathfinder.



**Figure B9.** Distribution of the logit of the ECS ( $T_{2x}$ ) inferred from AR6. Blue points are the assessments from AR6, the plain line is the CDF fitted on those assessment and the dashed line is the PDF associated with the CDF (arbitrary scale). The value of the fitted parameters is given above the plot.

In manual	In code	Description	Units
$R_c$	RFco2	CO2 (effective) radiative forcing	$\text{W m}^{-2}$
$R_x$	ERFx	Non-CO2 effective radiative forcing	$\text{W m}^{-2}$
$R$	ERF	Effective radiative forcing	$\text{W m}^{-2}$
$T$	T	Global surface temperature anomaly	K
$T_d$	Td	Deep ocean temperature anomaly	K
logit(ff)	logit_ff	Logit of the climate feedback factor (for calib.)	1
$U_{\text{ohc}}$	OHC	Ocean heat content (anomaly)	$\text{W yr m}^{-2}$
$H_{\text{thx}}$	Hthx	Thermosteric sea level rise	mm
$H_{\text{gla}}$	Hgla	Glaciers' contribution to sea level rise	mm
$H_{\text{gis}}$	Hgis	Grenland ice sheet's contribution to sea level rise	mm
$H_{\text{ais,smb}}$	Hais_smb	Surface mass balance component of Hais	mm
$H_{\text{ais}}$	Hais	Antartica ice sheet's contribution to sea level rise	mm
$H_{\text{tot}}$	Htot	Total sea level rise	mm
$H_{\text{lia}}$	Hlia	Sea level rise from relaxation after LIA between 1900 and 2005 (for calib.)	mm
$C_{o,j}$	Co_j	Change in surface ocean carbon subpools	PgC $j \in \llbracket 1, 5 \rrbracket$
$C_o$	Co	Change in surface ocean carbon pool	PgC
$C_d$	Cd	Change in deep ocean carbon pool	
$c_{\text{dic}}$	dic	Change in surface DIC	$\mu\text{mol kg}^{-1}$
$p_{\text{dic}}$	pdic	Subcomponent of pCO2	ppm
$p_{\text{CO2}}$	pCO2	CO2 partial pressure at the ocean surface	ppm
$F_{\text{ocean}}$	Focean	Ocean carbon sink	$\text{PgC yr}^{-1}$

**Table B1.** Summary of PathFinder's equation variables in climate, sea level and ocean carbon modules. [Prog?](#) is for prognostic variables

In manual	In code	Description	Units
$r_{\text{npp}}$	r_npp	Relative change in NPP	1
$r_{\text{fire}}$	r_fire	Relative change in wildfire intensity	1
$r_{\text{rh}}$	r_rh	Relative change in heterotrophic respiration rate	1
$E_{\text{npp}}$	NPP	Net primary productivity	PgC yr <sup>-1</sup>
$E_{\text{fire}}$	Efire	Emissions from wildfire	PgC yr <sup>-1</sup>
$E_{\text{harv}}$	Eharv	Emissions from harvest and grazing	PgC yr <sup>-1</sup>
$F_{\text{mort}}$	Fmort	Mortality flux	PgC yr <sup>-1</sup>
$E_{\text{rh1}}$	RH1	Litter heterotrophic respiration	PgC yr <sup>-1</sup>
$F_{\text{stab}}$	Fstab	Stabilization flux	PgC yr <sup>-1</sup>
$E_{\text{rh2}}$	RH2	Active soil heterotrophic respiration	PgC yr <sup>-1</sup>
$F_{\text{pass}}$	Fpass	Passivization flux	PgC yr <sup>-1</sup>
$E_{\text{rh3}}$	RH3	Passive soil heterotrophic respiration	PgC yr <sup>-1</sup>
$F_{\text{land}}$	Fland	Land carbon sink	PgC yr <sup>-1</sup>
$E_{\text{rh}}$	RH	Heterotrophic respiration	PgC yr <sup>-1</sup>
$C_v$	Cv	Vegetation carbon pool	PgC
$C_{s1}$	Cs1	Litter carbon pool	PgC
$C_{s2}$	Cs2	Active soil carbon pool	PgC
$C_{s3}$	Cs3	Passive soil carbon pool	PgC
$C_s$	Cs	Total soil carbon pool	PgC
$r_{\text{rt}}$	r_rt	Relative change in permafrost respiration rate	1
$\bar{a}$	abar	Theoretical thawed fraction	1
$a$	a	Actual thawed fraction	1
$E_{\text{pf}}$	Epf	Emissions from permafrost	PgC yr <sup>-1</sup>
$C_{\text{th},j}$	Cth_j	Thawed permafrost carbon subpools	PgC $j \in \llbracket 1, 3 \rrbracket$
$C_{\text{fr}}$	Cfr	Frozen permafrost carbon pool	PgC
$E_{\text{CO2}}$	Eco2	Anthropogenic CO2 emissions	PgC yr <sup>-1</sup>
$C$	CO2	Atmospheric CO2 concentration	ppm
pH	pH	Surface ocean pH	1

**Table B2.** Summary of PathFinder’s equation variables for land carbon, permafrost and atmospheric modules.

In manual	In code	Description	Units
$\phi$	phi	Radiative parameter of CO2	$\text{W m}^{-2}$
$T_{2\times}$	T2x	Equilibrium climate sensitivity	K
$\Theta_s$	THs	Heat capacity of the surface	$\text{W yr m}^{-2} \text{K}^{-1}$
$\Theta_d$	THd	Heat capacity of the deep ocean	$\text{W yr m}^{-2} \text{K}^{-1}$
$\theta$	th	Heat exchange coefficient	$\text{W m}^{-2} \text{K}^{-1}$
$\epsilon_{\text{heat}}$	eheat	Deep ocean heat uptake efficacy	1
$T_{2\times}^*$	T2x0	Minimal value of the ECS distribution (for calib.)	K
$\alpha_{\text{ohc}}$	aOHC	Fraction of energy warming the ocean	1
$\Lambda_{\text{thx}}$	Lthx	Proportionality factor of thermosteric SLR	$\text{mm m}^2 \text{W}^{-1} \text{yr}^{-1}$
$\lambda_{\text{gla}}$	lgla0	Initial imbalance in SLR from Glaciers	$\text{mm yr}^{-1}$
$\Lambda_{\text{gla}}$	Lgla	Maximum contribution to SLR from Glaciers	mm
$\Gamma_{\text{gla1}}$	Ggla1	Linear sensitivity of steady-state Glaciers SLR to climate	$\text{K}^{-1}$
$\Gamma_{\text{gla3}}$	Ggla3	Cubic sensitivity of steady-state Glaciers SLR to climate	$\text{K}^{-3}$
$\tau_{\text{gla}}$	tgla	Timescale of Glaciers' contribution to SLR	yr
$\gamma_{\text{gla}}$	ggla	Sensitivity of Glaciers' timescale to climate	$\text{K}^{-1}$
$\lambda_{\text{gis}}$	lgis0	Initial imbalance in SLR from GIS	$\text{mm yr}^{-1}$
$\Lambda_{\text{gis1}}$	Lgis1	Linear sensitivity of steady-state GIS SLR to climate	$\text{mm K}^{-1}$
$\Lambda_{\text{gis3}}$	Lgis3	Cubic sensitivity of steady-state GIS SLR to climate	$\text{mm K}^{-3}$
$\tau_{\text{gis}}$	tgis	Timescale of GIS contribution to SLR	yr
$\Lambda_{\text{ais,smb}}$	Lais_smb	Sensitivity of AIS SMB increase due to climate	$\text{mm yr}^{-1} \text{K}^{-1}$
$\lambda_{\text{ais}}$	lais	Initial imbalance in SLR from AIS	$\text{mm yr}^{-1}$
$\Lambda_{\text{ais}}$	Lais	Sensitivity of steady-state AIS SLR to climate	$\text{mm K}^{-1}$
$\tau_{\text{ais}}$	tais	Timescale of AIS contribution to SLR	yr
$\alpha_{\text{ais}}$	aais	Sensitivity of AIS timescale to AIS SLR	$\text{mm}^{-1}$
$\alpha_{\text{dic}}$	adic	Conversion factor for DIC	$\mu\text{mol kg}^{-1} \text{PgC}^{-1}$
$\beta_{\text{dic}}$	bdic	Inverse-scaling factor for DIC	1
$\gamma_{\text{dic}}$	gdic	Sensitivity of pCO2 to climate	$\text{K}^{-1}$
$T_o$	To	Preindustrial surface ocean temperature	$^{\circ}\text{C}$
$\nu_{\text{gx}}$	vgx	Surface ocean gas exchange rate	$\text{yr}^{-1}$
$\gamma_{\text{gx}}$	ggx	Sensitivity of gas exchange to climate	$\text{K}^{-1}$
$\alpha_{o,j}$	aoc_j	Surface ocean subpools fractions	$1 \ j \in \llbracket 1, 5 \rrbracket$
$\tau_{o,j}$	toc_j	Timescales of surface ocean subpools	$\text{yr } j \in \llbracket 1, 5 \rrbracket$
$\kappa_{\tau_o}$	k_toc	Scaling factor for timescales of surface ocean subpools	1

**Table B3.** Parameters used in the climate, ocean carbon and sea level modules.

In manual	In code	Description	Units
$\beta_{\text{npp}}$	bnpp	Sensitivity of NPP to CO2 (= fertilization effect)	1
$\alpha_{\text{npp}}$	anpp	Shape parameter for fertilization effect	1
$\gamma_{\text{npp}}$	gnpp	Sensitivity of NPP to climate	$\text{K}^{-1}$
$\beta_{\text{fire}}$	bfire	Sensitivity of wildfire intensity to CO2	1
$\gamma_{\text{fire}}$	gfire	Sensitivity of wildfire intensity to climate	$\text{K}^{-1}$
$\beta_{\text{rh}}$	brh	Sensitivity of heterotrophic respiration to fresh organic matter	1
$\gamma_{\text{rh}}$	grh	Sensitivity of heterotrophic respiration to climate	$\text{K}^{-1}$
$F_{\text{npp},0}$	npp0	Preindustrial NPP	$\text{PgC yr}^{-1}$
$\nu_{\text{fire}}$	vfire	Wildfire intensity	$\text{yr}^{-1}$
$\nu_{\text{harv}}$	vharv	Harvest and grazing rate	$\text{yr}^{-1}$
$\nu_{\text{mort}}$	vmort	Mortality rate	$\text{yr}^{-1}$
$\nu_{\text{stab}}$	vstab	Stabilization rate	$\text{yr}^{-1}$
$\nu_{\text{rh1}}$	vrh1	Litter heterotrophic respiration rate	$\text{yr}^{-1}$
$\nu_{\text{rh23}}$	vrh23	Soil (active and passive) respiration rate	$\text{yr}^{-1}$
$\nu_{\text{rh3}}$	vrh3	Passive soil respiration rate	$\text{yr}^{-1}$
$\alpha_{\text{pass}}$	apass	Fraction of passive soil	1
$\alpha_{\text{lst}}$	aLST	Climate scaling factor over permafrost regions	1
$\gamma_{\text{rt1}}$	grt1	Sensitivity of (boreal) heterotrophic respiration to climate	$\text{K}^{-1}$
$\gamma_{\text{rt2}}$	grt2	Sensitivity of (boreal) heterotrophic respiration to climate (quadratic)	$\text{K}^{-2}$
$\kappa_{\text{rt}}$	krt	Scaling factor for sensitivity of permafrost respiration to climate	1
$a_{\text{min}}$	amin	Minimal thawed fraction	1
$\kappa_a$	ka	Shape parameter for theoretical thawed fraction	1
$\gamma_a$	ga	Sensitivity of theoretical thawed fraction to climate	$\text{K}^{-1}$
$\nu_{\text{thaw}}$	vthaw	Thawing rate	$\text{yr}^{-1}$
$\nu_{\text{froz}}$	vfroz	Freezing rate	$\text{yr}^{-1}$
$\alpha_{\text{th},j}$	ath_j	Thawed permafrost carbon subpools fractions	1 $j \in \llbracket 1, 3 \rrbracket$
$\tau_{\text{th},j}$	tth_j	Timescales of thawed permafrost carbon subpools	$\text{yr}$ $j \in \llbracket 1, 3 \rrbracket$
$\kappa_{\tau_{\text{th}}}$	k_tth	Scaling factor for timescales of surface ocean subpools	1
$C_{\text{fr},0}$	Cfr0	Preindustrial frozen permafrost carbon pool	$\text{PgC}$
$\alpha_C$	aCO2	Conversion factor for atmospheric CO2	$\text{PgC ppm}^{-1}$
$C_{\text{pi}}$	CO2pi	Preindustrial CO2 concentration	ppm
$\kappa_{\text{pH}}$	k_pH	Scaling factor for surface ocean pH	1
$\tilde{\sigma}_C$	std_CO2	Relative standard deviation of the historical CO2 time series (for calib.)	1
$\epsilon_C$	ampl_CO2	Noise amplitude of the historical CO2 time series (for calib.)	ppm
$\rho_C$	corr_CO2	Autocorrelation of the historical CO2 time series (for calib.)	1
$\tilde{\sigma}_T$	std_T	Relative standard deviation of the historical T time series (for calib.)	1
$\epsilon_T$	ampl_T	Noise amplitude of the historical T time series (for calib.)	K
$\rho_T$	corr_T	Autocorrelation of the historical T time series (for calib.)	1

**Table B4.** Parameters used in the permafrost, land carbon modules **ag3** for calibration.

Parameters	Prior	Unit
Lgla	380	mm
Lais	1200	mm
tgla	190	yr
tgis	481	yr
tais	2093	yr
T2x0	0.61	K
adic	4.49	$\mu\text{mol kg}^{-1} \text{PgC}^{-1}$
aoc_1	0.87	1
aoc_2	0.06	1
aoc_3	0.04	1
aoc_4	0.02	1
aoc_5	0.01	1
toc_1	1.3	yr
toc_2	16.7	yr
toc_3	65	yr
toc_4	348	yr
toc_5	$10^9$	yr
vrh3	$8.27 \cdot 10^{-5}$	$\text{yr}^{-1}$
aLST	1.87	1
grt1	0.12	$\text{K}^{-1}$
grt2	0.0029	$\text{K}^{-2}$
krt	1.34	1
amin	0.98	1
vthaw	0.14	$\text{yr}^{-1}$
vfroz	0.011	$\text{yr}^{-1}$
ath_1	0.05	1
ath_2	0.12	1
ath_3	0.83	1
tth_1	18.2	yr
tth_2	252	yr
tth_3	3490	yr
aCO2	2.12	$\text{PgC ppm}^{-1}$
k_pH	1	1

**Table B5.** Structural parameters values. Parameters are noted under their code notation, and Tables B3 and B4 provide the corresponding notation in text.

Parameters	Prior	Posterior	Unit	Parameters	Prior	Posterior	Unit
phi	$5.35 \pm 0.54$	$5.29 \pm 0.54$	$\text{W m}^{-2}$	T2x	$4.13 \pm 1.37$	$3.37 \pm 0.77$	K
THs	$8.14 \pm 0.99$	$8.21 \pm 1.06$	$\text{W yr m}^{-2} \text{K}^{-1}$	THd	$108.6 \pm 61.8$	$123.8 \pm 57.8$	$\text{W yr m}^{-2} \text{K}^{-1}$
th	$0.61 \pm 0.13$	$0.67 \pm 0.12$	$\text{W m}^{-2} \text{K}^{-1}$	eheat	$1.35 \pm 0.40$	$1.41 \pm 0.43$	1
aOHC	$0.91 \pm 0.02$	$0.91 \pm 0.02$	1	Lthx	$1.82 \pm 0.21$	$1.85 \pm 0.23$	$\text{mm m}^2 \text{W}^{-1} \text{yr}^{-1}$
lgla0	$0.59 \pm 0.24$	$0.40 \pm 0.21$	$\text{mm yr}^{-1}$	Ggla1	$0.34 \pm 0.18$	$0.34 \pm 0.05$	$\text{mm K}^{-1}$
Ggla3	$0.022 \pm 0.013$	$0.022 \pm 0.013$	$\text{mm K}^{-3}$	ggla	$0.12 \pm 0.09$	$0.11 \pm 0.07$	$\text{K}^{-1}$
Lgis1	$82 \pm 45$	$189 \pm 55$	$\text{mm K}^{-1}$	Lgis3	$5.7 \pm 1.4$	$5.8 \pm 1.5$	$\text{mm K}^{-3}$
Lais_smb	$0.61 \pm 0.19$	$0.40 \pm 0.10$	$\text{mm K}^{-1} \text{yr}^{-1}$	lais0	$0.00 \pm 0.11$	$0.07 \pm 0.08$	$\text{mm yr}^{-1}$
aaais	$0.002 \pm 0.003$	$0.004 \pm 0.003$	$\text{mm}^{-1}$	lgis0	$0.33 \pm 0.14$	$0.35 \pm 0.14$	$\text{mm yr}^{-1}$
k_toc	$1.00 \pm 0.20$	$0.91 \pm 0.18$	1	vgx	$0.19 \pm 0.06$	$0.20 \pm 0.07$	$\text{PgC ppm}^{-1} \text{yr}^{-1}$
ggx	$0.018 \pm 0.029$	$0.019 \pm 0.033$	$\text{K}^{-1}$	To	$18.0 \pm 0.5$	$18.0 \pm 0.5$	K
bdic	$0.87 \pm 0.08$	$0.90 \pm 0.09$	1	gdic	$0.04 \pm 0.02$	$0.04 \pm 0.02$	$\text{K}^{-1}$
npp0	$48.2 \pm 5.1$	$46.5 \pm 3.3$	$\text{PgC yr}^{-1}$	vfire	$0.006 \pm 0.003$	$0.006 \pm 0.002$	$\text{yr}^{-1}$
vharv	$0.003 \pm 0.003$	$0.003 \pm 0.002$	$\text{yr}^{-1}$	vmort	$0.11 \pm 0.01$	$0.11 \pm 0.01$	$\text{yr}^{-1}$
vstab	$0.32 \pm 0.28$	$0.30 \pm 0.22$	$\text{yr}^{-1}$	vrh1	$0.33 \pm 0.29$	$0.27 \pm 0.20$	$\text{yr}^{-1}$
vrh23	$0.024 \pm 0.009$	$0.024 \pm 0.008$	$\text{yr}^{-1}$	bnpp	$0.93 \pm 0.37$	$1.09 \pm 0.25$	1
anpp	$0.48 \pm 0.57$	$0.36 \pm 0.38$	1	gnpp	$-0.014 \pm 0.023$	$-0.005 \pm 0.024$	$\text{K}^{-1}$
bfire	$-0.05 \pm 0.12$	$-0.06 \pm 0.14$	1	gfire	$0.052 \pm 0.072$	$0.044 \pm 0.088$	$\text{K}^{-1}$
brh	$1.06 \pm 0.43$	$1.01 \pm 0.41$	1	grh	$0.056 \pm 0.053$	$0.042 \pm 0.035$	$\text{K}^{-1}$
apass	$0.69 \pm 0.19$	$0.63 \pm 0.20$	1	ga	$0.13 \pm 0.04$	$0.13 \pm 0.04$	$\text{K}^{-1}$
ka	$2.6 \pm 2.0$	$2.4 \pm 1.8$	1	k_tth	$0.96 \pm 0.93$	$1.00 \pm 0.87$	1
Cfr0	$546 \pm 120$	$538 \pm 122$	PgC	CO2pi	$278 \pm 3$	$279 \pm 3$	ppm

**Table B6.** Calibrated parameters values before and after Bayesian calibration. Parameters are noted under their code notation, and Tables B3 and B4 provide the corresponding notation in text. The uncertainty correspond to the  $1 \sigma$  uncertainty range.